


THE
CANADIAN
FORMULARY
OF
UNOFFICIAL PREPARATIONS.

1908



SMEGTK

RS 125
.05
1908

THE ONTARIO COLLEGE OF PHARMACY

TORONTO, ONT.

Ca

THE

T

U. W. O. MEDICAL LIBRARY

THE
Canadian Formulary
OF
UNOFFICIAL PREPARATIONS.

BY AUTHORITY OF

THE ONTARIO COLLEGE OF PHARMACY

PRICE, ■ 50 CENTS

PUBLISHED BY

THE ONTARIO COLLEGE OF PHARMACY

TORONTO, ONT.

1908

T
Coll
purp
and

acce
for t
phy
exp
Dom
Ont
kno
supp
Rec
assi
mos

artic
prep
fanc
indi
fanc
prof

meth
pers
care
foun
read
unde
kno
effici
test

inclu
supp
publ
the
phar
in F

63725

PREFACE.

THIS second edition of the Canadian Formulary is published under the authority of the Ontario College of Pharmacy, by committees from the Colleges of Pharmacy, of the Provinces of Ontario and Quebec, appointed for the purpose of investigating and approving of formulas believed to be appropriate and suitable for the purpose and object for which the publication is authorized.

Recognition of certain formulas bearing a semi-official title prepared according to the formulas prevailing in localities, has demonstrated the necessity for the adoption of some uniform system of authoritative formulas, whereby the physician can intelligently prescribe and the pharmacist dispense, and the result expected and obtained be uniform and identical throughout the whole of the Dominion of Canada. This was the desire and intention of the Council of the Ontario College of Pharmacy, when the work was inaugurated, with the full knowledge and belief, that only by and with the general co-operation and support of the pharmacists of the entire Dominion can success be achieved. Recent developments point to such encouragement for a wider co-operation and assistance from the pharmacists, from which we feel justified in concluding, that, most valuable and material benefits will accrue.

The establishment of uniform and authoritative standards for medicinal articles, to meet the demands upon the medical and pharmaceutical professions for preparations brought to the attention of the prescriber, under various and fanciful coined names, with very extravagant claims for medicinal virtues, indicated as possessed only by the one special preparation and marketed at fanciful trade prices, is essentially a step in the interests of the public and the professions.

The best means of introducing the preparations, or the most successful method of obtaining due recognition of the preparations, should be through personal introduction to the physician by the pharmacist. If the pharmacist will carefully examine the various formulas, it is believed, that many of them will be found particularly applicable to the requirements in his locality, they can be readily prepared by any qualified pharmaceutical chemist, and with an intelligent understanding of the medicinal properties of the preparation, as well as a knowledge of any extravagant claims for competitive proprietary articles, the efficient pharmacist should be able to impress the physician and induce him to test the reliability of the articles presented.

Attention is particularly directed to the fact that many formulas are included in the book, for the express purpose of enabling the pharmacist to supply the popular demand for preparations on the market for which the formulas published will produce an article of identical properties, and that in some cases the formula is not to be considered or recommended as a truly scientific pharmaceutical exhibit of the ingredients contained in the preparation, (as shown in Formula numbers 33 and 35).

The formulas are largely selected and compiled from a careful survey and investigation of many recognized authorities, with the intention on the part of the committee of allowing due credit in each case to the source from which it is obtained. Valuable assistance was given the work by many pharmacists in Ontario and Quebec, also by Prof. Chas. F. Heebner, Dean of the Ontario College of Pharmacy, J. E. Morrison, Montreal College of Pharmacy, and Fred. W. Flett, Toronto, who are worthy of special mention, and to whom a large share of credit is due. Criticisms and suggestions on all formulas will be cheerfully received by the committees, and recommendations for new formulas eligible for inclusion in subsequent editions, will materially advance the scope and usefulness of the work.

Both Imperial and Metric weights and measures are given throughout the Formulary. It has been somewhat difficult, in the course of a single paragraph embodying formulas involving definite quantities of materials, to give precise directions for their employment in *two different systems* of weights and measures, hence those who use the Formulary are requested to avoid the assumption that Imperial and Metric quantities thus placed in juxtaposition are necessarily equivalent to one another. The intention has been to furnish formulas that will yield liquid products measuring twenty fluidounces (or a convenient multiple of that volume) or one thousand cubic centimeters. Except for wholly insignificant fractional differences, a preparation made according to either system will contain the same proportions of ingredients; but the two systems cannot both be used in the same operation, and are therefore not interchangeable.

The term 'Diluted Alcohol' which occurs throughout the text, refers to a mixture of equal volumes of commercial (95%) Alcohol and Distilled Water.

RESEARCH COMMITTEES.

ONTARIO COLLEGE OF PHARMACY.—John Hargreaves, *Chairman*;
Henry Watters, R. A. Harrison, J. H. H. Jury, and E. R. Wigle.

QUEBEC PHARMACEUTICAL ASSOCIATION.—W. H. Chapman, *Chairman*;
J. E. Morrison, A. B. J. Moore, H. R. Gray, A. E. Duberger, A. J. Laurence,
H. Lanctot, J. E. W. Lécours.

O.C.P. COUNCIL, 1907-1908.

GEO. E. GIBBARD, *President*. JOHN HARGREAVES, *Vice-President*.

Henry Watters, E. W. Case, John H. H. Jury, J. R. Y. Broughton, Alex. Stewart, Henry Southcott, R. A. Harrison, W. A. Karn, J. F. Roberts, Ebon R. Wigle, A. J. Johnston.

W. B. GRAHAM,
Registrar-Treasurer.

Toronto, 1st, March 1908.

Wate
Dilut
to pe
of ic
funn
and,
wash
cold
tests
test s
Mix t
on a
Alcoh
suffic
Pres
:

Alcoh
Powd
Powd
Spirit
—
N

Alcoh
aside

T
Purifie
tinued

1. ACIDUM HYPOPHOSPHOROSUM.

Hypophosphorus Acid.

(N.F. 1900)

Hypophosphite of Potassium	483 parts.
Tartaric Acid	682 parts.
Distilled Water	500 parts.
Diluted Alcohol ("45% Alcohol")	1000 parts.

Dissolve the Potassium Hypophosphite in 500 parts of Distilled Water, previously warmed, and the Tartaric Acid in 1000 parts of Diluted Alcohol. Mix the solutions in a flask of sufficient capacity to permit agitation, cork and shake well and set the flask in a bath of ice water for 12 hours. Then carefully pour the mixture into a funnel, the neck of which has been closed with a pledget of cotton, and, when all the liquid has been drained off, rinse the flask, and wash the crystalline precipitate in the funnel with small portions of cold Diluted Alcohol until the washings no longer respond to the tests for Hypophosphorus Acid (black precipitate by Silver Nitrate test solution or white precipitate by Mercuric Chloride test solution). Mix the original filtrate and the washings and evaporate the whole on a water-bath at a temperature not exceeding 140° F. until all the Alcohol has been dissipated. Allow the liquid to cool and add sufficient Distilled Water to bring the weight up to 1000 parts. Preserve the product in well stoppered bottles.

NOTE.—This should contain 30 per cent. of absolute Hypophosphorus Acid.

2. ALCOHOL DEODORATUM.

Deodorized Alcohol.

(N.F. 1896)

Alcohol (95 per cent.)	160 fluidounces	5000 Cc.
Powdered Quicklime	300 grains	20 Gm.
Powdered Alum	150 grains	10 Gm.
Spirit of Nitrous Ether	1¼ fluidrachm	4·5 Cc.

Mix the Lime and Alum intimately by trituration; add to the Alcohol and shake well, then add the Spirit of Nitrous Ether, set aside for seven days and filter through powdered Animal Charcoal.

3. AQUA OLEI ROSÆ.

Rose Water.

Oil of Rose	1 Cc.
Calcium Phosphate or Purified Talcum	2 Gm.
Distilled Water	500 Cc.

Triturate the Oil of Rose with the Phosphate of Calcium (or the Purified Talcum), gradually add the Distilled Water, with continued trituration, and filter.

NOTE.—The following Medicated Waters may be made in the same manner as Rose Water, and used in the place of the corresponding *Aquæ* of the text of the B. P. :—

Aqua Olei Anethi.
 Aqua Olei Anisi.
 Aqua Olei Carui.
 Aqua Olei Cinnamomi.
 Aqua Olei Fœniculi.
 Aqua Olei Menthæ Viridis.
 Aqua Olei Menthæ Piperitæ.
 Aqua Olei Pimentæ.

4. CAPSULÆ APIOL ET ERGOTINI.

Capsules of Apiol and Ergotin.

Each capsule to contain Apiol five minims. (0.30 Cc.) and Ergotin two grains (0.13 Gm.).

5. CAPSULÆ COLCHICINÆ ET METHYL SALICYLATIS.

Capsules of Salicylates of Colchicine and Methyl.

Colchicine Salicylate.....	1 grain	0.65 Gm.
Methyl Salicylate.....	1250 minims.	74 Cc.

Dissolve and fill into 250 capsules.

Each capsule contains Colchicine 1-250th grain (0.00025 Gm.), and Methyl Salicylate five minims (0.30 Cc.). Dose, one capsule.

6. CATAPLASMA KAOLINI.

Cataplasm of Kaolin.

(U.S.P. 1905)

Kaolin, in very fine powder.....	11½ ounces	577 Gm.
Boric Acid, in very fine powder.....	395 grains	45 Gm.
Thymol.....	5 grains	0.5 Gm.
Methyl Salicylate (Synthetic Oil of Winter-green).....	20 grains	2 Gm.
Oil of Peppermint.....	5 grains	0.5 Gm.
Glycerin.....	7½ ounces	375 Gm.

Heat the Kaolin in a suitable vessel at 212° F. with occasional stirring, for one hour. Heat the Glycerin at the same temperature for half an hour, and dissolve in it the Boric Acid, and incorporate the hot Kaolin with this liquid. Dissolve the Thymol in the Methyl Salicylate and the Oil of Peppermint, and mix with the above to form a homogeneous mass. The product should be kept in an air-tight container.

Liqu
Whi
Spe
Bor
Oil
Dist

Bees
pour
Bora
finall

to 5)

Chlor
Camp

1
mort

Camp
Chlor

I

Iodofc
Flexib

D
dry bo

No

7. CERATUM GALENI.

Galen's Cerate.

Cold Cream.

Liquid Paraffin	16 fluidounces	160·0 Cc.
White Beeswax	4 ounces	40·0 Gm.
Spermaceti.....	1 ounce	10·0 Gm.
Borax.....	30 grains	0·6 Gm.
Oil of Rose.....	10 minims.	0·2 Cc.
Distilled Water.....	8 fluidounces	80·0 Cc.

Dissolve the Borax in the Distilled Water; melt the White Beeswax and Spermaceti with the Liquid Paraffin at a gentle heat; pour the mixture into a warmed mortar and add while yet hot the Borax solution (previously warmed) with constant trituration, and finally the Oil of Rose, and continue the trituration until cold.

In hot weather the quantity of White Beeswax may be increased to 5½ ounces (53 Gm.) and the Spermaceti to 2 ounces (20 Gm.).

8. CHLORAL CAMPHORATUM.

Camphorated Chloral.

Chloral	2 ounces	50 Gm.
Camphor	2 ounces	50 Gm.

Mix them by agitation in a bottle, or by trituration in a warm mortar until liquified and combined.

9. CHLOROFORMUM CAMPHORATUM.

Camphorated Chloroform.

Camphor.....	2 ounces	200 Gm.
Chloroform.....	1 fluidounce	100 Cc.

Dissolve the Camphor in the Chloroform by agitation.

10. COLLODIUM IODOFORMATUM.

Iodoform Collodion.

(N.F. 1906)

Iodoform, in fine powder.....	5 parts
Flexible Collodion.....	95 parts

Dissolve the Iodoform in the Flexible Collodion contained in a dry bottle, by agitation.

NOTE.—This preparation should be made extemporaneously.

11. ELIXIR ACETANILIDI COMPOSITUM.

Compound Elixir of Acetanilide.

Acetanilide	400 grains	22.75 Gm.
Phenacetin	320 grains	18.3 Gm.
Sodium Bromide	3 ounces 288 grains	91.5 Gm.
Caffeine Citrate	160 grains	9.15 Gm.
Tartaric Acid	80 grains	4.58 Gm.
Sodium Bicarbonate	1 ounce 32 grains	27.5 Gm.
Aromatic Elixir sufficient to make.....	40 fluidounces	1000 Cc.

Mix the Acetanilide, Tartaric Acid and Sodium Bicarbonate and dissolve in 20 fluidounces (500 Cc.) of Aromatic Elixir. To this solution add the Sodium Bromide and Caffeine Citrate; then add sufficient Aromatic Elixir to make 40 fluidounces (1000 Cc.), and filter if necessary.

12. ELIXIR ADJUVANS.

Adjuvant Elixir.

(U.S.P. 1905)

Fluid Extract of Glycyrrhiza	2½ fluidounces	120 Cc.
Aromatic Elixir	17½ fluidounces	880 Cc.

Mix, and filter if necessary.

13. ELIXIR AMMONII BROMIDI.

Elixir of Ammonium Bromide.

Ammonium Bromide	1600 grains	91.5 Gm.
Citric Acid	70 grains	4 Gm.
Aromatic Elixir, sufficient to make.....	40 fluidounces	1000 Cc.

Dissolve the Ammonium Bromide and Citric Acid in about 20 fluidounces (500 Cc.) of Aromatic Elixir, by agitation. Then add enough Aromatic Elixir to make 40 fluidounces (1000 Cc.) and filter, if necessary.

Each fluidrachm contains 5 grains (0.32 Cc.) of Ammonium Bromide.

14. ELIXIR ANISI.

Elixir of Anise.

Aniseed Cordial.

Anethol	65 minims.	3.5 Cc.
Oil of Fennel	10 minims.	0.5 Cc.
Spirit of Bitter Almond	4 fluidrachms	12 Cc.
Deodorized Alcohol	9½ fluidounces	240 Cc.
Syrup	25 fluidounces	625 Cc.
Water	5 fluidounces	125 Cc.
Magnesium Carbonate	4½ drachms	15 Gm.

the
ture
Car
pori

Con
Syr
Pre
Deo

Acc
syr
war
Wat
then
fils
fiter
part:

Spiri
Deo
Simp
Disti
Tale

ally :

Pota
Sodit
Amm
Calci
Lithi
Tinct
Arom
I
of Ca

Mix the Anethol, the Oil and the Spirit of Bitter Almond with the Deodorized Alcohol, add the Syrup and Water and set the mixture aside for 12 hours. Then mix it intimately with the Magnesium Carbonate and filter through a wetted filter, returning the first portions of the filtrate until it passes perfectly clear.

15. ELIXIR AROMATICUM.*Aromatic Elixir.*

Compound Spirit of Orange 230 minims. 12 Cc.
 Syrup 14 fluidounces 375 Cc.
 Precipitated Calcium Phosphate..... 285 grains 15 Gm.
 Deodorized Alcohol and Distilled Water,
 a sufficient quantity of each to make... 40 fluidounces 1000 Cc.

To the Compound Spirit of Orange add enough Deodorized Alcohol to make 10 fluidounces (250 Cc.). To this solution add the syrup in several portions, agitating after each addition, and afterwards add in the same manner 14 fluidounces (375 Cc.) of Distilled Water. Mix the Calcium Phosphate intimately with the liquid and then filter through a wetted filter, returning the first portions of the filtrate until a transparent liquid is obtained. Lastly, wash the filter with a mixture of one part of Deodorized Alcohol to three parts of Water, until the product measures 40 fluidounces (1000 Cc.).

16. ELIXIR AURANTII.*Elixir of Orange.**Simple Elixir.*

Spirit of Orange..... 4 fluidounces 4 Cc.
 Deodorized Alcohol 25 fluidounces 25 Cc.
 Simple Syrup 40 fluidounces 40 Cc.
 Distilled Water..... 31 fluidounces 31 Cc.
 Talcum, a sufficient quantity.

Mix the several ingredients in the order named; shake occasionally and filter through Talcum, until the filtrate passes perfectly clear.

17. ELIXIR QUINQUE BROMIDORUM.*Elixir of Five Bromides.*

Potassium Bromide 1600 grains 91.5 Gm.
 Sodium Bromide 1600 grains 91.5 Gm.
 Ammonium Bromide 960 grains 55. Gm.
 Calcium Bromide 480 grains 27.45 Gm.
 Lithium Bromide 160 grains 9.15 Gm.
 Tincture of Cannabis Indica 2 fluidounces 50. Cc.
 Aromatic Elixir, sufficient to make..... 40 fluidounces 1000 Cc.

Dissolve the Bromides in the Aromatic Elixir, add the Tincture of Cannabis Indica and filter if necessary.

18. ELIXIR BUCHU ET HYOSCYAMI COMPOSITUM.

Compound Elixir of Buchu and Hyoscyamus.

Fluid Extract Buchu	3 fluidounces	75· Cc.
Fluid Extract Uva Ursi	1½ fluidounces	37·5 Cc.
Fluid Extract Pareira	1½ fluidounces	37·5 Cc.
Fluid Extract Hyoscyamus	1½ fluidounces	37·5 Cc.
Fluid Extract Hops	1½ fluidounces	37·5 Cc.
Potassium Acetate	2 ounces	291 grains
Spirit of Nitrous Ether	4½ fluidounces	112·5 Cc.
Aromatic Elixir, sufficient to make	40 fluidounces	1000 Cc.

Mix and set aside for two days. Filter if necessary.

19. ELIXIR CALCII ET SODII GLYCEROPHOSPHATIS.

Elixir of Glycerophosphate of Calcium and Sodium.

Calcium Glycerophosphate	320 grains	18·3 Gm.
Sodium Glycerophosphate	160 grains	9·15 Gm.
Gluside	5 grains	0·286 Gm.
Concentrated Phosphoric Acid	150 grains	8·58 Gm.
Tincture of Fresh Sweet-Orange Peel	1¼ fluidounce	31·25 Cc.
Glycerin	7½ fluidounces	187·5 Cc.
Sherry Wine	10 fluidounces	250· Cc.
Distilled Water, sufficient to make	40 fluidounces	1000 Cc.

Dissolve the Glycerophosphates of Calcium and Sodium in ten fluidounces (250 Cc.) of Distilled Water with which the Concentrated Phosphoric Acid has been previously mixed. Then add the Glycerin, Sherry and the Gluside dissolved in the Tincture of Orange and enough Distilled Water to make the finished Elixir measure 40 fluid ounces (1000 Cc.). Filter through paper sprinkled with Talcum.

NOTE.—Each fluidrachm contains Glycero-Phosphate of Calcium, 1 grain (0·065 Gm.) and Glycerophosphate of Sodium, ½ grain (0·325 Gm.).

20. ELIXIR CINCHONÆ.

Elixir of Cinchona. Elixir of Calisaya.

*(Elixir of Cinchona from "Alkaloids").**Compound Elixir of Quinine.*

(N.F. 1906)

Quinine Sulphate	30 grains	2 Gm.
Cinchonidine Sulphate	15 grains	1 Gm.
Cinchonine Sulphate	15 grains	1 Gm.
Compound Tincture of Cudbear	1½ fluidounces	50 Cc.
Purified Talcum	240 grains	15 Gm.
Aromatic Elixir, sufficient to make	32 fluidounces	1000 Cc.

Aron
cient
tritur
stand
throu
perfe
)
½ grSolut
Wate
Elixir

add

Digit
Solut
Solut
Aromuntil
Elixir
menteach
O
crystaSodiu
Sodiu
Fluid
Tinct
Solut
Arom

Dissolve the Alkaloid Salts in 30 fluidounces (900 Cc.) of Aromatic Elixir, add the Compound Tincture of Cudbear and sufficient Aromatic Elixir to make 32 fluidounces (1000 Cc.) and triturate the Purified Talcum with the mixture. Allow the mixture to stand several hours, if convenient, occasionally shaking, then filter through paper, returning the first portions until the filtrate passes perfectly clear.

Each fluidounce contains Quinine Sulphate about 1 grain, and $\frac{1}{2}$ grain each of Cinchonidine and Cinchonine Sulphates.

21. ELIXIR CINCHONÆ ET FERRI.

Elixir Cinchona and Iron.

Ferrated Elixir of Cinchona.

(N.F. 1906)

Soluble Ferric Phosphate.....	640 grains	36.6 Cc.
Water (boiling).....	5 fluidounces	125 Cc.
Elixir of Cinchona, sufficient to make 40 fluidounces		1000 Cc.

Dissolve the Soluble Ferric Phosphate in the boiling water, then add Elixir of Cinchona sufficient to make 40 fluidounces (1000 Cc.).

22. ELIXIR DIGITALINI COMPOSITUM.

Compound Elixir of Digitalin.

Digitalin (amorphous).....	$1\frac{1}{2}$ grains	171 milligrams
Solution of Strychnine, B.P....	6 fluidrachms	37.3 Cc.
Solution of Trinitrin.....	3 fluidrachms	18.65 Cc.
Aromatic Elixir, sufficient to make 20 fluidounces		500 Cc.

Triturate the amorphous Digitalin with a portion of the Elixir until a solution results. Then add to the remainder of the Aromatic Elixir the Strychnine, Trinitrin and Digitalin solutions, in the order mentioned, mixing thoroughly after each addition.

NOTE.—Each fluidrachm of this Elixir contains approximately $\frac{1}{15}$ grain each of Digitalin and Trinitrin, and $\frac{1}{35}$ grain of Strychnine Hydrochloride.

Only amorphous Digitalin should be used in making this preparation, as the crystalline variety is believed to be five times as strong as the amorphous.

23. ELIXIR EUPHORBIAE COMPOSITUM.

Compound Elixir of Euphorbia.

(Anti-Asthmatic Elixir.)

Sodium Iodide	640 grains	36.6 Cc.
Sodium Bromide.....	640 grains	36.6 Cc.
Fluid Extract of Euphorbia.....	2 fluidounces	50 Cc.
Tincture of Lobelia.....	$1\frac{1}{4}$ fluidounces	31.5 Cc.
Solution of Trinitrin (B.P.).....	3 fluidrachms	9 Cc.
Aromatic Elixir, sufficient to make....	40 fluidounces	1000 Cc.

Dissolve the Sodium Iodide and Bromide in 20 fluidounces (500 Cc.) of Aromatic Elixir, add the remaining ingredients, and, lastly, sufficient Aromatic Elixir to make 40 fluidounces (1000 Cc.). Filter if necessary.

24. ELIXIR FERRI PYROPHOSPHATIS CUM QUININA ET STRYCHNINA.

Elixir of Pyrophosphate of Iron with Quinine and Strychnine.

Quinine Sulphate	160 grains	9.15 Gm.
Sodium Citrate	150 grains	8.6 Gm.
Solution of Strychnine, B.P.	500 minims	26. Cc.
Iron Pyrophosphate, soluble	600 grains	34. Gm.
Alcohol (95 %)	5 fluidounces	125. Cc.
Glycerin	6 fluidounces	150. Cc.
Distilled Water	2 fluidounces	50. Cc.
Simple Elixir, sufficient to make	40 fluidounces	1000 Cc.

Dissolve the Quinine in the Alcohol and 6 fluidounces (150 Cc.) of Simple Elixir, using gentle heat if necessary, and add the Solution of Strychnine. Dissolve the Pyrophosphate of Iron in the Water previously warmed, and 2 fluidounces (50 Cc.) of Simple Elixir, and add it to the Solution of Quinine and Strychnine. Dissolve the Sodium Citrate in the Glycerin, mix the solutions and add sufficient Simple Elixir to make 40 fluidounces (1000 Cc.).

25. ELIXIR FERRI, QUININÆ ET STRYCHNINÆ.

Elixir of Iron, Quinine and Strychnine.

(N.F. 1906)

Tincture of Ferric Citro-Chloride	5 fluidounces	125. Cc.
Quinine Hydrochloride	156 grains	8.75 Gm.
Strychnine Sulphate	3½ grains	0.175 Gm.
Alcohol (95%)	1 fluidounce	25. Cc.
Aromatic Elixir, sufficient to make	40 fluidounces	1000 Cc.

Dissolve the alkaloidal salts in 32 fluidounces (750 Cc.) of the Elixir, then add the Tincture and Alcohol, and finally enough Aromatic Elixir to make 40 fluidounces (1000 Cc.). Filter if necessary.

Each fluidrachm contains $\frac{1}{100}$ grain of Strychnine Sulphate.

26. ELIXIR FORMINI.

Elixir of Formin.

Elixir Hexamethylene-tetramine.

Formin	600 grains	34.125 Gm.
Tincture of Cudbear	5 fluidrachms	15 Cc.
Aromatic Elixir, sufficient to make	40 fluidounces	1000 Cc.

Dissolve the Formin in the Elixir, add the Tincture of Cudbear and filter if necessary.

Flui
Aro

Calc
Sodi
Iron
Pota
Citri
Tinc
Glus
Glyc
Sher
Dist

ounc
add
viou
mak
with

Calc
Sodi
Pota
Mag
Iron
Quir
Stry
Citri
Glus
Tinc
Alco
Glyc
Dist

27. ELIXIR GLYCYRRHIZÆ.

Elixir of Glycyrrhiza.

Elixir of Licorice.

(N.F. 1906)

Fluid Extract of Licorice.....	5 fluidounces	125 Cc.
Aromatic Elixir.....	35 fluidounces	875 Cc.

Filter if necessary.

28. ELIXIR GLYCEROPHOSPHATUM COMPOSITUM.

Compound Elixir of Glycerophosphates.

Calcium Glycerophosphate.....	160 grains	9.2 Gm.
Sodium Glycerophosphate.....	160 grains	9.2 Gm.
Iron Glycerophosphate (Scale).....	80 grains	4.6 Gm.
Potassium Glycerophosphate.....	80 grains	4.6 Gm.
Citric Acid.....	76 grains	4.5 Gm.
Tincture of Sweet Orange Peel.....	50 fluidrachms	15 Cc.
Gluside.....	4 grains	0.25 Gm.
Glycerin.....	6 fluidounces	150 Cc.
Sherry Wine.....	10 fluidounces	250 Cc.
Distilled Water, sufficient to make....	40 fluidounces	1000 Cc.

Dissolve the Glycerophosphates and Citric Acid in 12 fluidounces (300 Cc.) of warm water, add the Glycerin, and when cool add the Tincture of Orange in which the Gluside has been previously dissolved, then the Sherry Wine, and sufficient water to make 40 fluidounces (1000 Cc.). Filter through paper sprinkled with Talcum, returning the filtrate until it passes perfectly clear.

Dose, 2 fluidrachms.

29. ELIXIR GLYCEROPHOSPHATUM CUM QUININA ET STRYCHNINA.

Elixir of Glycerophosphates with Quinine and Strychnine.

Calcium Glycerophosphate.....	240 grains	13.8 Gm.
Sodium Glycerophosphate.....	160 grains	9.2 Gm.
Potassium Glycerophosphate.....	160 grains	9.2 Gm.
Magnesium Glycerophosphate.....	160 grains	9.2 Gm.
Iron Glycerophosphate (Scale).....	80 grains	4.58 Gm.
Quinine Hydrochloride.....	20 grains	1.15 Gm.
Strychnine.....	4 grains	0.25 Gm.
Citric Acid.....	60 grains	3.5 Gm.
Gluside.....	22 grains	1.25 Gm.
Tincture of Sweet Orange Peel.....	2 fluidounces	50 Cc.
Alcohol (95%).....	2 fluidounces	50 Cc.
Glycerin.....	10 fluidounces	250 Cc.
Distilled Water, sufficient to make....	40 fluidounces	1000 Cc.

Dissolve the Glycerophosphates of Quinine and Strychnine and the Citric Acid in 20 fluidounces (500 Cc.) of warm water mixed with the Glycerin, and when cold add the Tincture of Orange Peel and Alcohol, in which the Gluside has been previously dissolved. Filter through paper sprinkled with Talcum and pass sufficient Distilled Water through the filter to make 40 fluidounces (1000 Cc.).

Each fluidrachm contains $\frac{1}{80}$ grain Strychnine Hydrochloride.
Dose, 1 to 2 fluidrachms.

30. ELIXIR LITHII ET HYDRANGÆÆ.

Elixir of Lithium and Hydrangea.

Lithium Salicylate.....	600 grains	34 Gm.
Lithium Benzoate.....	300 grains	17 Gm.
Fluid Extract of Hydrangea.....	$7\frac{1}{2}$ fluidounces	187 Cc.
Alcohol.....	$7\frac{1}{2}$ fluidounces	187 Cc.
Aromatic Elixir, sufficient to make.....	40 fluidounces	1000 Cc.

Dissolve the Lithium salts in 25 fluidounces (625 Cc.) Aromatic Elixir, add the Alcohol to the Fluid Extract of Hydrangea and mix all together. Let the mixture stand for twenty-four hours and filter if necessary.

31. ELIXIR LITHII SALICYLATIS.

Elixir of Lithium Salicylate.

(N.F. 1906)

Lithium Salicylate.....	1600 grains	91.5Gm.
Aromatic Elixir, sufficient to make....	40 fluidounces	1000 Cc.

Dissolve the Lithium Salicylate in about 36 fluidounces (900 Cc.) of Aromatic Elixir by agitation. Then add enough Aromatic Elixir to make 40 fluidounces (1000 Cc.) and filter.

Each fluidrachm contains 5 grains (0.325 Gm.) Lithium Salicylate.

32. ELIXIR PAPAINI.

Elixir of Papain.

Papain.....	640 grains	36.5 Cc.
Diluted Hydrochloric Acid.....	150 minims.	8 Cc.
Distilled Water.....	6 fluidounces	150 Cc.
Glycerin.....	6 fluidounces	150 Cc.
Sherry Wine.....	6 fluidounces	150 Cc.
Gluside.....	20 grains	1.15 Gm.
Aromatic Elixir, sufficient to make....	40 fluidounces	1000 Cc.

Macerate the Papain in the Acid and Water for four days, with occasional agitation. Dissolve the Gluside in the Wine and Elixir, add the Glycerin, mix with Papain mixture and filter; then add Aromatic Elixir sufficient to make 40 fluidounces (1000 Cc.).

Peps
Panc
Dias
Lacti
Hydr
Glyc
Wat
Tinct
Purif
Aron

Panc
occa:
the
40 fl
thore

popul
consis
retain
The l
amou

3.

Pepsi
Panc
Dias
Glyc
Lacti
Hydr
Glyc
Disti
Tinct
Purif
Aron

33. ELIXIR PEPSINI COMPOSITUM.

Compound Elixir of Pepsin.

*Elixir Digestivum Compositum. Elixir of Digestive Ferments.
Elixir of Lactated Pepsin.*

(N.F. 1906)

Pepsin	175 grains	10	Gm.
Pancreatin	17½ grains	1	Gm.
Diastase	17½ grains	1	Gm.
Lactic Acid	½ fluidrachm	1·5	Cc.
Hydrochloric Acid	1 fluidrachm	3	Cc.
Glycerin	10 fluidounces	250	Cc.
Water	5 fluidounces	125	Cc.
Tincture of Cudbear ..	5 fluidounces	125	Cc.
Purified Talcum	1 Av. ounce	30	Gm.
Aromatic Elixir, sufficient to make	40 fluidounces	1000	Cc.

Mix the Acids with the Glycerin and Water, add the Pepsin, Pancreatin and Diastase to this mixture, and macerate with occasional agitation until solution is apparently effected. Then add the Tincture of Cudbear and enough Aromatic Elixir to make 40 fluidounces (1000 Cc.). Incorporate the Purified Talcum thoroughly with the mixture and filter.

NOTE.—This preparation is included to enable the Pharmacist to meet a popular demand for an Elixir of this name. It is not, however, presented as a consistent scientific exhibit of the ingredients specified, it being impossible to retain all three digestive ferments in soluble active condition in the same medium. The best commercial variety of Diastase capable of converting the largest amount of Starch into Dextrin and Glucose, should be used.

34. ELIXIR PEPSINI CUM BISMUTHO COMPOSITUM.

Compound Elixir of Pepsin with Bismuth.

Elixir Lactated Pepsin with Bismuth.

Pepsin (1 in 3000)	175 grains	10	Gm.
Pancreatin	17½ grains	1	Gm.
Diastase	17½ grains	1	Gm.
Glycerin of Bismuth	5 fluidounces	125	Cc.
Lactic Acid	30 minims.	1·5	Cc.
Hydrochloric Acid	1 fluidrachm	3	Cc.
Glycerin	2 fluidounces	50	Cc.
Distilled Water	5 fluidounces	125	Cc.
Tincture of Cudbear	2 fluidounces	50	Cc.
Purified Talcum	1 fluidounce	25	Gm.
Aromatic Elixir, sufficient to make	40 fluidounces	1000	Cc.

Mix the Acids with the Glycerin and Water, add the Pepsin, Pancreatin and Diastase to the mixture and macerate with frequent agitation until solution is apparently effected. Then add the Glycerin of Bismuth and Tincture of Cudbear and sufficient Aromatic Elixir to make 40 fluidounces. (1000 Cc.). Thoroughly incorporate the Purified Talcum and filter.

See note under formula No. 33.

35. ELIXIR PEPSINI CUM QUININA ET FERRO ET STRYCHNINA.

Elixir of Pepsin, with Quinine, Iron and Strychnine.

Elixir of Lactated Pepsin, with Quinine, Iron and Strychnine.

Pepsin....(1 in 3000)	175 grains	10 Gm.
Pancreatin	17½ grains	1 Gm.
Diastase	17½ grains	1 Gm.
Quinine	120 grains	6·9 Gm.
Solution of Ferric Chloride, (B.P.)....	2 fluidounces	50 Cc.
Purified Talcum.....	1 fluidounce	25 Cc.
Solution of Strychnine (B.P.)	1 fluidounce	25 Cc.
Lactic Acid.....	30 minims.	1·5 Cc.
Hydrochloric Acid	1 fluidrachm	3 Cc.
Glycerin	4 fluidounces	100 Cc.
Distilled Water.....	5 fluidounces	125 Cc.
Gluside	20 grains	0·2 Gm.
Aromatic Elixir, sufficient to make....	40 fluidounces	1000 Cc.

Mix the Acids with the Glycerin and Water, add the Pepsin, Pancreatin and Diastase to the mixture and macerate with frequent agitation until dissolved. Dissolve the Gluside in 1 fluidounce (25 Cc.) of Distilled Water. Dissolve the Quinine in the Solution of Iron, add the Solution of Strychnine, the Glycerin and 2 fluidounces of Aromatic Elixir. Mix all together and lastly add Aromatic Elixir sufficient to make 40 fluidounces (1000 Cc.). Thoroughly incorporate with the Purified Talcum and filter.

Each fluidounce contains 3 grains of Quinine and ⅓ grain of Strychnine. See note under formula No. 33.

36. ELIXIR PEPSINI ET BISMUTHI.

Elixir of Pepsin and Bismuth.

(N.F. 1900)

Pepsin(1 in 3000).....	160 grains	9 Gm.
Glycerin	4 fluidounces	100 Cc.
Glycerin of Bismuth.....	5 fluidounces	125 Cc.
Distilled Water	10 fluidounces	250 Cc.
Aromatic Elixir, sufficient to make	40 fluidounces	1000 Cc.

I
add 1
thoro
I
Bism

Potas
Distil
Solut
Elixir

I
about
add tl
40 flu
E

Brom

Calcii
Acetic
Fluid
Arom

I
with t
Acid,
add tl
ounce

I
nesium

Fluid
Fluid
Fluid
Glyce
Arom

Dissolve the Pepsin in the Glycerin and Distilled Water, then add the Glycerin of Bismuth and Aromatic Elixir and mix thoroughly.

Each fluidrachm contains $\frac{1}{2}$ grain of Pepsin and 2 grains of Bismuth and Ammonium Citrate.

37. ELIXIR POTASSII BROMIDI.

Elixir of Potassium Bromide

Potassium Bromide	7 ounces. 138 grains	183 Gm.
Distilled Water	7 fluidounces	175 Cc.
Solution of Carmine	35 minims	2 Cc.
Elixir of Orange, sufficient to make	40 fluidounces	1000 Cc.

Dissolve the Potassium Bromide in the Distilled Water and about 25 fluidounces (625 Cc.) of the Elixir of Orange by agitation; add the Solution of Carmine and sufficient Elixir of Orange to make 40 fluidounces (1000 Cc.). Let stand a few hours and filter.

Each fluidrachm contains 10 grains (0.65 Gm.) of Potassium Bromide.

38. ELIXIR RHEI ET MAGNESII ACETATIS.

Elixir of Rhubarb and Magnesium Acetate.

(N.F. 1906)

Calcined Magnesia	355 grains	20 Gm.
Acetic Acid	a sufficient quantity	
Fluid Extract of Rhubarb	5 fluidounces	125 Cc.
Aromatic Elixir, sufficient to make	40 fluidounces	1000 Cc.

Dissolve the Magnesia in 6 fluidounces (150 Cc.) of Acetic Acid with the aid of a gentle heat, adding, if necessary a little more Acetic Acid, drop by drop until the solution is neutral to test paper. Then add the Fluid Extract and enough Aromatic Elixir to make 40 fluid ounces (1000 Cc.) and filter.

Each fluidrachm represents about 4 grains (0.25 Gm.) of Magnesium Acetate and $7\frac{1}{2}$ grains (0.5 Gm.) of Rhubarb.

39. ELIXIR SERENOÆ COMPOSITUM.

Compound Elixir of Saw Palmetto.

Fluid Extract Saw Palmetto	2 fluidounces	50 Cc.
Fluid Extract Sandalwood	2 fluidounces	50 Cc.
Fluid Extract Couch Grass.	2 fluidounces	50 Cc.
Fluid Extract Corn Silk	2 fluidounces	50 Cc.
Glycerin	2 fluidounces	50 Cc.
Aromatic Elixir, sufficient to make	40 fluidounces	1000 Cc.

Mix and let stand for four days, then filter if necessary.

40. ELIXIR SEX IODORUM.

Elixir of Six Iodides.

Arsenic Iodide	2 grains	0.12 Gm.	Terp
Mercuric Iodide	2 grains	0.12 Gm.	Heri
Manganese Iodide	32 grains	1.85 Gm.	Glus
Sodium Iodide	320 grains	18.5 Gm.	Tinc
Potassium Iodide	320 grains	18.5 Gm.	Brar
Glycerin of Ferrous Iodide	30 minims	1.5 Cc.	Alco
Sodium Hypophosphite, a sufficient quantity			Glyc
Aromatic Elixir, sufficient to make	40 fluidounces	1000 Cc.	Dist

Add the six Iodides to the Elixir, dissolve by agitation, and add sufficient Sodium Hypophosphite to decolorize the liquid. Filter.

41. ELIXIR SODII SALICYLATIS COMPOSITUM.

Compound Elixir of Sodium Salicylate.

Sodium Sulphate	10 ounces	250 Gm.	
Sodium Salicylate (from Oil of Winter-green)	800 grains	45.5 Gm.	
Magnesium Sulphate	9 ounces	225 Gm.	Fluid
Lithium Benzoate	400 grains	22.75 Gm.	Fluid
Tincture of Nux Vomica	2 fluidounces	50 Cc.	Fluid
Solution of Carmine	6 fluidrachms	18 Cc.	Fluid
Simple Elixir sufficient to make	40 fluidounces	1000 Cc.	Arom

Dissolve the salts in 24 fluidounces (600 Cc.) of Simple Elixir by trituration, add the Tincture of Nux Vomica and Solution of Carmine and sufficient Simple Elixir to make 40 fluidounces. (1000 Cc.) Filter if necessary.

42. ELIXIR TERPINI HYDRATIS ET CODEINÆ.

Elixir of Terpin Hydrate and Codeine.

Terpin Hydrate, in fine powder	320 grains	18.3 Gm.	
Codeine Phosphate	40 grains	2.3 Gm.	
Gluside	10 grains	0.6 Gm.	R
Tincture of Fresh Sweet Orange Peel	1 1/4 fluidounces	31.5 Cc.	the W
Alcohol (95 per cent.)	13 fluidounces	325 Cc.	
Glycerin	20 fluidounces	500 Cc.	
Elixir of Orange, sufficient to make	40 fluidounces	1000 Cc.	

Dissolve the Terpin Hydrate, Codeine Phosphate and Gluside in the Alcohol with a gentle heat, add the Tincture of Orange, Glycerin, and enough Elixir of Orange to make 40 fluidounces (1000 Cc.).

Each fluidrachm contains Terpin Hydrate 1 grain (0.065 Gm). and Codeine Phosphate 1/8 grain (0.008 Gm.).

Terp
Heri
Glus
Tinc
Brar
Alco
Glyc
Dist

Glus
Vani
mak

and 1

Fluid
Fluid
Fluid
Fluid
Arom
Arom
1
and a
sprin

R
the W

Cod I
Acacis
Soluti
or Tol
Flavoi
Distill

43. ELIXIR TERPINI HYDRATIS ET HEROINÆ.

Elixir of Terpin Hydrate and Heroine.

0.12 Gm.	Terpin Hydrate	320 grains	18.3 Gm.
0.12 Gm.	Heroine Hydrochloride	13 1/3 grains	0.75 Gm.
0.85 Gm.	Gluside	10 grains	0.6 Gm.
0.5 Gm.	Tincture of Vanilla (1 in 10).	1 1/4 fluidrachms	4 Cc.
0.5 Gm.	Brandy	5 fluidrachms	16 Cc.
0.5 Cc.	Alcohol (95 per cent.).	15 fluidounces	375 Cc.
	Glycerin	20 fluidounces	500 Cc.
000 Cc.	Distilled Water, sufficient to make	40 fluidounces	1000 Cc.

Dissolve the Terpin Hydrate, Heroine Hydrochloride and the Gluside in the Alcohol with a gentle heat; add the Tincture of Vanilla, Brandy, Glycerin and lastly, enough Distilled Water to make 40 fluidounces (1000 Cc.).

Each fluidrachm contains Terpin Hydrate 1 grain (0.065 Gm.) and Heroine Hydrochloride 1/4 grain (0.0025 Gm.).

44. ELIXIR VIBURNI COMPOSITUM.

Compound Elixir of Crampbark.

0.5 Gm.	Fluid Extract of Hydrastis	1 1/2 fluidounces	37.5 Cc.
0.75 Gm.	Fluid Extract of Viburnum Opulus	3 fluidounces	75 Cc.
Cc.	Fluid Extract of Scutellaria	1 1/2 fluidounces	37.5 Cc.
Cc.	Fluid Extract of Mitchella Repens	1 1/2 fluidounces	37.5 Cc.
Cc.	Aromatic Syrup of Licorice	6 fluidounces	150 Cc.
e Elixir	Aromatic Elixir, sufficient to make	40 fluidounces	1000 Cc.

Mix the fluid Extracts, then add the Aromatic Syrup of Licorice and agitate, then add the Aromatic Elixir. Filter through paper sprinkled with Talcum, if necessary.

45. EMULSIO IODOFORMI.

Emulsion of Iodoform.

3 Gm.	Iodoform	10 parts.
1 Gm.	Glycerin	70 parts.
1 Gm.	Distilled Water	20 parts.

Rub the Iodoform to a smooth paste with the Glycerin, then add the Water and continue stirring until a uniform product results.

46. EMULSIO OLEI MORRHUÆ.

Emulsion of Cod Liver Oil.

side	Cod Liver Oil	10 fluidounces	500 Cc.
nge,	Acacia, in fine powder,	2 1/2 fluidounces	125 Gm.
nces	Solution of Gluside	1 fluidrachm	7 Cc.
	or Tolu Syrup.	2 fluidounces	100 Cc.
7m).	Flavoring to suit (see No. 108).		
	Distilled Water, sufficient to make	20 fluidounces	1000 Cc.

Prepare a Primary Emulsion in the following manner:

Place the Powdered Acacia into the centre of a perfectly dry and clean wedgwood mortar, then add the Oil (pouring it *upon* the Acacia rather than *about* it) and triturate until a homogenous mixture results. Then add, *all at once*, twice as much Water as Acacia, (the water having a temperature of not less than 90° F.) and stir briskly with the pestle until a thick creamy emulsion results.

To the Primary Emulsion thus prepared, add the desired flavoring materials, also the Solution of Gluside, or Tolu Syrup, under constant stirring, and likewise enough water to make 20 fluidounces (1000 Cc.).

NOTE.—The Hypophosphites of Calcium or Sodium or other water-soluble salts can be included in this Emulsion, by dissolving them in the water used in diluting the Primary Emulsion.

47. EMULSIO OLEI MORRHUÆ CUM FERRI PHOSPHATO.

Emulsion of Cod Liver Oil with Phosphate of Iron.

Cod Liver Oil	20 fluidounces	500 Cc.
Soluble Ferric Phosphate	240 grains	13·8 Gm.
Powdered Acacia	5 ounces	125 Gm.
Syrup of Orange	2½ fluidounces	62·5 Cc.
Syrup of Tolu	2½ fluidounces	62·5 Cc.
Distilled Water, sufficient to make	40 fluidounces	1000 Cc.

Prepare a Primary Emulsion of the Cod Liver Oil, as directed under Emulsio Olei Morrhuae, then add the Syrups, under constant stirring. Dissolve the Soluble Ferric Phosphate in 3½ fluidounces (87·5 Cc.) of Water, and add this, under stirring, to the mixture, and follow with sufficient Water to make 40 fluidounces (1000 Cc.).

Each fluidounce contains 6 grains of Ferric Phosphate.

48. EMULSIO OLEI MORRHUÆ CUM PEPSINO.

Emulsion of Cod Liver Oil with Pepsin. Phosphatic Emulsion.

Cod Liver Oil	144 fluidounces	4082·40 Cc.
The Yolks of Twenty-four Eggs		
Glycerin	24 fluidounces	680·40 Gm.
White Sugar	40 ounces	1174 Gm.
Compound Powder of Acacia	4½ ounces	127·8 Gm.
Lime Water	75 fluidounces	2041·20 Cc.
Diluted Phosphoric Acid	9 fluidounces	255·15 Cc.
Essence of Pepsin	24 fluidounces	608·40 Cc.
Flavor, as required (See No. 108)	3 fluidrachms	9·15 Cc.

Rub the Yolks of Eggs in a Mortar (Whites of half this number of Eggs may be added with advantage) until a smooth paste results; add the Glycerin and stir briskly. Add the Compound Powder of Acacia, then the Cod Liver Oil in portions of about 8 fluidounces (230 Cc.) at a time. When the oil is emulsified add

the L
then
Pepsin
to sti

Oil of
Lemo
Alcohol
Distill
Magn

A
Carbo
Water
macer
Alcohol

Glycer
Sherry
Glycer
Alcohol
Tinctu
Distill

M

Phenol
Essence

Di:
necessa

the Lime Water containing the Sugar in solution and stir vigorously ; then add the Diluted Phosphoric Acid and finally the Essence of Pepsin, and stir vigorously for fifteen minutes. Allow the Emulsion to stand for two hours, and strain through dairy cloth.

49. ESSENTIA LIMONIS.

Essence of Lemon.

Oil of Lemon (fresh).....	1 fluidounce	25 Cc.
Lemon Peel (freshly grated).....	1 Av. ounce	25 Cc.
Alcohol	28 fluidounces	700 Cc.
Distilled Water	12 fluidounces	300 Cc.
Magnesium Carbonate	4 drachms	12.5 Gm.

Mix the Oil of Lemon and the Lemon Peel with the Magnesium Carbonate. Triturate well, then slowly add the Alcohol and Distilled Water, previously mixed, meanwhile continuing the trituration ; macerate for 24 hours, then filter and add sufficient of the mixed Alcohol and Water to make 40 fluidounces (1000 Cc.).

50. ESSENTIA PEPSINI.

Essence of Pepsin.

Glycerin of Pepsin, B.P.....	4 fluidounces	100 Cc.
Sherry	5 fluidounces	125 Cc.
Glycerin	5 fluidounces	125 Cc.
Alcohol (95 per cent.).....	3½ fluidounces	87.5 Cc.
Tincture of Fresh Sweet-Orange Peel..	5 fluidrachms	15.5 Cc.
Distilled Water, sufficient to make....	40 fluidounces	1000. Cc.

Mix and filter through paper sprinkled with Talcum.

51. ESSENTIA PEPSINI PHENOLATA.

Phenolated Essence of Pepsin.

Phenol (Absolute).....	25 grains	1.4 Gm.
Essence of Pepsin, sufficient to make.....	40 fluidounces	1000 Cc.

Dissolve the Phenol in the Essence of Pepsin, and filter if necessary.

52. ESSENTIA VANILLINI COMPOSITA.

Compound Essence of Vanillin.

Compound Tincture of Vanillin.

(N. F. 1906)

Vanillin	110 grains	6.2 Gm.
Cumarin	8 grains	0.5 Gm.
Alcohol	7½ fluidounces	187.5 Cc.
Glycerin	5 fluidounces	125 Cc.
Syrup	5 fluidounces	125 Cc.
Compound Tincture of Cudbear	5 fluidrachms	17.5 Cc.
Distilled Water, sufficient to make	40 fluidounces	1000 Cc.

Dissolve the Vanillin and Cumarin in the Alcohol, add the Glycerin, Syrup and Tincture of Cudbear and finally sufficient Distilled Water to make 40 fluidounces (1000 Cc.).

53. EXTRACTUM BUCHU FLUIDUM.

Fluid Extract of Buchu.

(U. S. P. 1905)

Buchu Leaves (60 powder)	40 ounces	1000 Gms.
Alcohol 95%, and Water of each, sufficient to make	40 fluidounces	1000 Cc.

Mix 30 fluidounces (750 Cc.) of Alcohol with 10 fluidounces (250 Cc.) of Water, and having moistened the powder with 16 fluidounces (400 Cc.) of this menstruum, pack it firmly in a cylindrical percolator, then add enough menstruum to saturate the powder and leave a stratum above it. Macerate for 48 hours, and continue the percolation process in the usual manner, reserving the first 34 fluidounces (850 Cc.) of the percolate, evaporating the remainder to a soft extract, and dissolving this in the reserved percolate, and adding sufficient menstruum to make 40 fluidounces (1000 Cc.) of Fluid Extract.

54. EXTRACTUM CASCARÆ SAGRADÆ AROMATICUM FLUIDUM.

Aromatic Fluid Extract of Cascara.

Cascara Bark (in coarse powder)	60 ounces	1704 Gm.
Licorice Root (in coarse powder)	10 ounces	284 Gm.
Calcined Magnesia	12 ounces	340 Gm.
Gluside	40 grains	2.3 Gm.
Sodium Bicarbonate	10 grains	.65 Gm.
Oil of Coriander	15 minims.	1 Cc.
Oil of Aniseed	20 minims.	1.25 Cc.
Alcohol, 95%	1 ounce	28.4 Cc.
Glycerin	24 ounces	682 Cc.
Distilled Water (boiling)	100 ounces	2840 Cc.

moist
in a s
pack
water
(or st
the C
Sodiu
both
and s

Seneg
Solut
Alcoh
t
y
(600
contin
Buchu
when

Powd
Hydro
Distill

A
bottle
in por

Green
Boilin
Glycer
R
the bo
suffici

Mix the Cascara, Licorice and Magnesia thoroughly, and moisten thoroughly with the water, stirring well. Place the mixture in a suitable, well-covered container, and macerate for 24 hours, then pack moderately tight in a percolator, and percolate with boiling water until exhausted. Evaporate the percolate over a water-bath (or steam-bath) until it measures 54 fluidounces (1535 Cc.). Dissolve the Gluside in 1 fluidounce (28·4 Cc.) of water with the aid of the Sodium Bicarbonate. Dissolve the Oils in the Alcohol and mix both solutions with the Glycerin, then add the concentrated percolate and shake thoroughly.

55. EXTRACTUM SENEÆ FLUIDUM.

Fluid Extract of Senega.

(U.S.P. 1905)

Senega (No. 40 powder)	40 ounces	1000 Gm.
Solution of Potassium Hydroxide	1¼ fluidounces	30 Cc.
Alcohol 95%, and Water, of each sufficient to make	40 fluidounces	1000 Cc.

Mix the Solution of Potassium Hydroxide with 24 fluidounces (600 Cc.) of Alcohol and 12 fluidounces (300 Cc.) of Water, and continue with the percolation process as given under Fluid Extract of Buchu, using a mixture of Alcohol, two parts, with Water, one part, when all of the alkaline menstruum has been used.

56. GARGARISMA CHLORI.

Chlorine Gargle.

Powdered Potassium Chlorate	4 drachms	2·75 Gm.
Hydrochloric Acid	120 minims.	13 Cc.
Distilled Water, sufficient to make	20 fluidounces	1000 Cc.

Add the Hydrochloric Acid to the Potassium Chlorate in a large bottle; when the gas given off has displaced the air, add the water in portions, corking and shaking the bottle after each addition.

57. GLYCERINUM BELLADONNÆ.

Glycerin of Belladonna.

Green Extract of Belladonna	1 ounce	28·4 Gm.
Boiling Distilled Water	1 fluidrachm	3·5 Cc.
Glycerin, sufficient quantity to make	2 fluidounces	56·8 Cc.

Rub together in a warm mortar the Extract of Belladonna and the boiling Distilled Water to produce a smooth paste; then add sufficient Glycerin to make 2 fluidounces (56·8 Cc.)

58. GLYCERINUM BISMUTHI.

Glycerin of Bismuth.

Bismuth and Ammonium Citrate. 11 ounces 308 grains 332.46 Gm.
 Glycerin 10 fluidounces 250 Cc.
 Strong Solution of Ammonia, a sufficient quantity.
 Distilled Water, sufficient to make..... 40 fluidounces 1000 Cc.

Triturate the Bismuth and Ammonium Citrate with 8 fluidounces (200 Cc.) of Distilled Water, and 3 fluidounces (150 Cc.) Glycerin, and gradually add to it just enough Strong Solution of Ammonia to dissolve the salt and to produce a neutral solution. Then add the remainder of the Glycerin and sufficient Distilled Water to make 40 fluidounces (1000 Cc.)

Each fluidrachm contains 16 grains of Bismuth and Ammonium Citrate.

59. GLYCERINUM FERRI IODIDI.

Glycerin of Ferrous Iodide.

Iron (in wire) 2½ ounces 125 Gm.
 Iodine 6 ounces 405 grains 415 Gm.
 Glycerin 10 fluidounces 500 Cc.
 Sulphurous Acid, B.P. 125 minims. 13 Cc.
 Distilled Water, sufficient to make 20 fluidounces 1000 Cc.

Mix the Iron and Iodine in a flask with 8 fluidounces (200 Cc.) of Distilled Water. Shake the mixture occasionally, checking the reaction, if necessary, by the affusion of cold water, and when the solution has acquired a greenish color and has lost the odor of Iodine, heat it gently to the boiling point, and add at once 2 fluidounces (100 Cc.) of Glycerin and filter the solution into the remainder of the Glycerin. Then add the Sulphurous Acid and sufficient Glycerin to make 20 fluidounces (1000 Cc.), and mix thoroughly.

NOTE.—This preparation should be kept in small, well-filled, well-corked, colorless glass bottles, in a place accessible to light. Each fluidounce contains 220 grains Ferrous Iodide. 1 Volume mixed with 4 Volumes of Simple Syrup will furnish a preparation similar to Syrup of Ferrous Iodide B.P.

**60. GLYCERINUM FERRI PHOSPHATIS CUM
QUININA ET STRYCHNINA.**

Glycerin of Phosphate of Iron with Quinine and Strychnine.

Iron Wire..... 750 grains 45 Gm.
 Concentrated Phosphoric Acid B.P. 12½ fluidounces 31.25 Cc.
 Strychnine..... 50 grains 2.8 Gm.
 Quinine Sulphate..... 1300 grains 73 Gm.
 Glycerin 24 fluidounces 600 Cc.
 Distilled Water, sufficient to make 40 fluidounces 1000 Cc.

with
 cotton
 result
 Glyce
 make
 N
 will fu

Hero
 Amm
 Fluid
 Fluid
 Solul
 Glyce
 Syru
 Cinn
 Glyce
 Wate
 Wild
 40 fl

repe

Buch
 Tinc
 Alco
 Dilut
 Infus

Place the Iron Wire, and the Phosphoric Acid (previously diluted with 4 fluidounces of Distilled Water) in a flask, plug the neck with cotton wool, and heat gently till the Iron is dissolved; in the resulting solution dissolve the Quinine and Strychnine, filter into the Glycerin, and pass sufficient Distilled Water through the filter to make 40 fluidounces (1000 Cc.).

NOTE.—1 Volume of this Glycerin mixed with 4 Volumes of Simple Syrup will furnish a preparation similar to Syrup Triple Phosphates, B.P.

61. GLYCERINUM HEROINI COMPOSITUM.

Compound Glycerin of Heroin.

Heroin	20 grains	1.15 Gm.
Ammonium Hypophosphite	640 grains	36.6 Gm.
Fluid Extract of Hyoscyamus	320 minims.	18.3 Cc.
Fluid Extract of White Pine	2 2/3 fluidounces	66.5 Cc.
Soluble Tincture of Tolu	2 fluidounces	50 Cc.
Glycerin	10 fluidounces	250 Cc.
Syrup of Wild Cherry Bark	6 fluidounces	150 Cc.
Cinnamon Water, sufficient to make . .	40 fluidounces	1000 Cc.

Dissolve the Heroin and the Ammonium Hypophosphite in the Glycerin, previously mixed with 8 fluidounces (200 Cc.) of Cinnamon Water; add the Fluid Extracts, Tincture of Tolu and Syrup of Wild Cherry, and finally add sufficient Cinnamon Water to make 40 fluidounces (1000 Cc.).

62. GLYCERINUM IODI.

Glycerin of Iodine.

Iodine, resublimed	1 part
Glycerin	50 parts

Dissolve the Iodine in the Glycerin with the aid of a gentle heat.

NOTE.—This forms a useful pigment, the skin does not get hardened by repeated applications and does not peel off.

63. INFUSUM BUCHU CONCENTRATUM.

Concentrated Infusion of Buchu.

(B.P.C. 1907)

Buchu Leaves	40 parts
Tincture of Buchu	22.5 parts
Alcohol	10 parts
Dilute Chloroform Water (1 in 1000) sufficient to make	100 parts

Prepare by the Macero-Expression Process for Concentrated Infusions (B.P.C.)

Dose, 1 to 2 fluidrachms.

64. LAC FERMENTATUM.

Fermented Milk.

"Kumyss."

(N.F. 1906)

Cows Milk (fresh)	20 fluidounces	100 Cc.
Yeast (semi-liquid)	45 minims.	5 Cc.
Sugar	5 drachms	35 Gm.

Dissolve the Sugar in the Milk contained in a strong bottle, add the yeast, cork the bottle securely, and keep it at a temperature between 74° and 90° F. for six hours, then transfer it to a cold place.

NOTE.—24 grains of Compressed Yeast, triturated with a little milk, may be used in place of the semi-liquid yeast.

65. LAC HUMANISATUM.

Humanized Milk.

(N.F. 1906)

Milk Powder	100 grains	6.5 Gm.
Cows' Milk (Fresh)	2 fluidounces	56.8 Cc.
Sweet Cream (Fresh)	½ fluidounce	14 Cc.
Distilled Water	2 fluidounces	56.8 Cc.

Triturate the Milk Powder with the water, transfer the mixture into a bottle containing the Milk and Cream, and immerse the bottle in water heated to 100° F. for fifteen minutes; then pour the mixture into a vessel, in which heat it quickly to boiling, and then immediately allow it to cool to the body temperature.

NOTE.—Should be freshly prepared. If directions are carefully followed, the milk will be well peptonized and the pancreatin of the Milk Powder rendered sterile.

66. LINIMENTUM ALBUM.

White Liniment. Stokes' Liniment.

Acetic Turpentine Liniment.

(N.F. 1906)

Oil of Turpentine	3 fluidounces	85.2 Cc.
Fresh Egg	1 (one)	1 (one)
Oil of Lemon	60 minims.	3.5 Cc.
Acetic Acid	300 minims.	17.5 Cc.
Rose Water	2½ fluidounces	71 Cc.

Triturate or beat the contents of the Fresh Egg with the Oils in a mortar until they are thoroughly mixed. Then incorporate the Acetic Acid and Rose Water. Shake the mixture, whenever any of it is to be dispensed.

Strot
Tinct
Glyc
Tinc

depos

Men
Chlo
Olivuse o
Offic
Oil stMen
LiniAmr
for :

67. LINIMENTUM AMMONII IODIDI.

Liniment of Ammonium Iodide.

Strong Solution of Ammonia	5 fluidounces	50 Cc.
Tincture of Iodine	5 fluidounces	50 Cc.
Glycerin	5 fluidounces	50 Cc.
Tincture of Camphor	5 fluidounces	50 Cc.

Mix and agitate.

NOTE.—On standing, the liquid will become colorless, usually with a slight deposit, which may be separated by filtration.

68. LINIMENTUM MENTHOLIS.

Menthol Liniment.

Menthol	2 ounces	50 Gm.
Chloroform	8 fluidounces	200 Cc.
Olive Oil, sufficient to make	40 fluidounces	1000 Cc.

Mix and agitate until the Menthol is dissolved.

NOTE.—The Colonial Addendum of the British Pharmacopœia permits the use of Oleum Sesami (Oil of Benne) in North American Colonies, in making the Official Liniments, Ointments and Plasters, for which the B.P. orders that Olive Oil shall be used.

69. LINIMENTUM MENTHOLIS COMPOSITUM.

Compound Menthol Liniment.

Menthol	1 ounce	10 Gm.
Liniment of Ammonium Iodide	49 fluidounces	490 Cc.

Mix and agitate until the Menthol is dissolved.

70. LIQUOR AMMONII VALERIANATIS.

Solution of Valerianate of Ammonium.

Valerianic Acid	3 parts
Ammonium Carbonate, a sufficient quantity.	
Alcoholic Extract of Valerian	2 parts
Distilled Water, sufficient to make	100 parts

Add the Acid to the Water and neutralize carefully with Ammonium Carbonate, add the Extract of Valerian, and let it stand for 24 hours, then filter.

Dose, 10 to 30 drops in sweetened water.

71. LIQUOR ANTISEPTICUS.

Antiseptic Solution.

(U.S.P. 1905)

Boric Acid	352 grains	20	Gm.
Benzoic Acid	18 grains	1	Gm.
Thymol	18 grains	1	Gm.
Eucalyptol	4 minims.	0.25	Cc.
Oil of Peppermint	8 minims.	0.50	Cc.
Oil of Gaultheria	4 minims.	0.25	Cc.
Oil of Thyme	1½ minims.	0.10	Cc.
Alcohol (95 %)	10 fluidounces	250	Cc.
Purified Talcum	352 grains	20	Gm.
Water, sufficient to make	40 fluidounces	1000	Cc.

Dissolve the Boric Acid in 24 fluidounces (600 Cc.) of Water and the Benzoic Acid in 6 fluidounces (150 Cc.) of Alcohol, and pour the aqueous solution into the alcoholic solution, then dissolve (in a mortar) the Thymol in the Eucalyptol and Oils of Peppermint, Gaultheria and Thyme; thoroughly incorporate the Purified Talcum, and add with constant trituration to the solution first prepared. Allow the mixture to stand with occasional agitation, during forty-eight hours, filter, add 4 fluidounces (100 Cc.) of Alcohol to the clear filtrate, and a sufficient quantity of Water to make the finished product measure 40 fluidounces (1000 Cc.).

72. LIQUOR ANTISEPTICUS ALKALINUS.

Alkaline Antiseptic Solution.

(N.F. 1905)

Potassium Bicarbonate	600 grains	32	Gm.
Sodium Benzoate	600 grains	32	Gm.
Sodium Biborate	310 grains	8	Gm.
Thymol	4 grains	0.2	Gm.
Eucalyptol	4 minims.	0.2	Cc.
Oil of Peppermint	4 minims.	0.2	Cc.
Oil of Wintergreen	7 minims.	0.4	Cc.
Tincture of Cudbear	300 minims.	16	Cc.
Alcohol (95 %)	2½ fluidounces	62.5	Cc.
Glycerin	10 fluidounces	250	Cc.
Purified Talcum	185 grains	10	Gm.
Water, a sufficient quantity to make	40 fluidounces	1000	Cc.

Dissolve the salts in 23 fluidounces (575 Cc.) of water, and the Thymol, Eucalyptol and Oils in the Alcohol. Mix the alcoholic solution with the Glycerin and Tincture of Cudbear, add the solution of the salts and enough water to make 40 fluidounces (1000 Cc.). Add the Talcum, shake occasionally during a few days, then filter.

Lactone

Glycerin Thymol

Arsen
Gold
Brom
a
I
Cc.)
Brom
at a t
does i
a por
gentl
and d
to ma
I
of Tr
of Tr
times
the W
I

Bora:
Sodi:
Carb:
Glyce
Wate
I
and t
by w
(1000

Chlor
Pota:
Tinct
Tinct

73. LIQUOR AURI ET ARSENI BROMIDI.

Solution of Bromide of Gold and Arsenic.

(N.F. 1906)

Arsenous Acid	10 grains	2.5 Gm.
Gold Tribromide	13 grains	3.25 Gm.
Bromine Water, Distilled Water, of each		
a sufficient quantity to make	10 fluidounces	1000 Cc.

Introduce the Arsenous Acid and about $1\frac{1}{3}$ fluidounces (135 Cc.) of Bromine Water in a flask and heat gently until all free Bromine has disappeared. Then add Bromine Water, 20 to 30 drops at a time, until it will be present in slight excess, or until the solution does not become colorless after some time. Transfer the solution to a porcelain capsule, expel the excess of Bromine with the aid of gentle heat, dilute it with Water to about 9 fluidounces (900 Cc.), and dissolve in this the Tribromide of Gold, adding enough Water to make 10 fluidounces (1000 Cc.).

Ten (10) minims. of this solution contain $\frac{3}{32}$ grain (.002 Gm.) of Tribromide of Gold and the equivalent of $\frac{1}{18}$ grain (0.004 Gm.) of Tribromide of Arsenic.

NOTE.—Bromine Water is made by shaking Bromine with about thirty times its weight of Water, occasionally during several hours, and decanting the Water from the undissolved Bromine.

Average dose, 3 minims.

74. LIQUOR BORACIS COMPOSITUS.

Compound Solution of Borax.

Dobell's Solution.

(N.F. 1906)

Borax	130 grains	15 Gm.
Sodium Bicarbonate	130 grains	15 Gm.
Carbolic Acid	25 grains	3 Gm.
Glycerin	$5\frac{1}{2}$ fluidrachms	35 Cc.
Water, sufficient to make	20 fluidounces	1000 Cc.

Dissolve the salts in about 10 fluid ounces (500 Cc.) of Water, and then add the Glycerin and the Carbolic Acid, previously liquified by warming, and lastly enough Water to make 20 fluidounces (1000 Cc.)

75. LIQUOR BROMO-CHLORAL COMPOSITUS

Compound Solution of Bromo-Chloral.

Chloral Hydrate	$3\frac{1}{2}$ ounces	182.75 Gm.
Potassium Bromide	$3\frac{1}{2}$ ounces	182.75 Gm.
Tincture of Indian Hemp	6 fluidrachms	41.65 Cc.
Tincture of Orange Peel	6 fluidrachms	41.65 Cc.

Henbane Juice	3 fluidounces	165.55 Cc.
Syrup	3 $\frac{3}{4}$ fluidounces	187.5 Cc.
Fluid Extract of Licorice	$\frac{1}{2}$ fluidounce	25 Cc.
Distilled Water, sufficient to make	20 fluidounces	1000 Cc.

Dissolve the Potassium Bromide in 7 fluidounces (330 Cc.) of Water, and mix all the other ingredients (except the Water) together. Add the solution of Potassium Bromide, filter and wash the filtrate with enough Distilled Water to make 20 fluidounces (1000 Cc.).

Dose, $\frac{1}{2}$ to 2 fluidrachms.

76. LIQUOR CARMINI.

Solution of Carmine.

(N.F. 1906)

Carmine	1 ounce	87 grains	60 Gm.
Solution of Ammonia	7 fluidounces		350 Cc.
Glycerin	7 fluidounces		350 Cc.
Water, a sufficient quantity to make	20 fluidounces		1000 Cc.

Triturate the Carmine to a fine powder in a wedgwood mortar, gradually add the Solution of Ammonia, and afterwards the Glycerin under constant trituration. Transfer the mixture to a porcelain capsule and heat it upon a water-bath, constantly stirring, until the liquid is free from ammoniacal odor. Then cool and add enough water to make 20 fluidounces. (1000 Cc.).

77. LIQUOR CREOSOTI ET IODI.

Solution of Creosote and Iodine.

Iodine, Resublimed	2 ounces	56.8 Gm.
Creosote	3 fluidounces	85.2 Cc.

Triturate the Iodine in a glass mortar to a fine powder, add the Creosote slowly and continue trituration until solution is effected.

NOTE.—This preparation is intended for dental use.

78. LIQUOR CRESOLIS.

Solution of Cresol.

Cresylic Acid (Cresol)	25 fluidounces	625. Cc.
Resin	5 ounces	125. Gm.
Potassium Hydroxide	350 grains	25. Gm.
Distilled Water, sufficient to make	40 fluidounces	1000 Cc.

Dissolve the Resin in the Cresylic Acid with the aid of heat. Make a solution of the Potassium Hydrate by dissolving in two fluidounces of Distilled Water. Mix the two solutions, and heat until saponification takes place. Set aside to cool, and make up to 40 fluidounces (1000 Cc.), with water.

Pepto
Brand
Alcohol
Syrup
Tinctu
Tinctu
Comp
Distill
E
Water
the re

Glucid
Sodium
Alcohol
Water
D
ounce
the fil
fluidou
E

Potass
Calciu
Sodium
Iron F
Mangr
Potass
Citric
Quinir
Strych
Hypop
Oil of
Alcohol
Glucid
Glycer
Distill

79. LIQUOR FERRI ET MANGANI PEPTONATIS.

Solution of Peptonate of Iron and Manganese.

Peptonate of Iron and Manganese	728 grains	475	Gm.
Brandy	4 fluidounces	100	Cc.
Alcohol	3 fluidounces	75	Cc.
Syrup	5 fluidounces	125	Cc.
Tincture of Orange	2 fluidrachms	6.25	Cc.
Tincture of Cinnamon	2 fluidrachms	6.25	Cc.
Compound Essence of Vanillin	2 fluidrachms	6.25	Cc.
Distilled Water, sufficient to make	40 fluidounces	1000	Cc.

Dissolve the Peptonate in 24 fluidounces (300 Cc.) of Distilled Water, with a gentle heat, then allow the solution to cool and add the remaining ingredients.

80. LIQUOR GLUSIDI.

Solution of Gluside.

Solution of Saccharin.

(N.F. 1906).

Gluside	1 ounce	202 grains	73 Gm.
Sodium Bicarbonate	29 grains	29 grains	33 Gm.
Alcohol	5 fluidounces	250	Cc.
Water, a sufficient quantity to make	20 fluidounces	1000	Cc.

Dissolve the Gluside and the Sodium Bicarbonate in 13 fluidounces (650 Cc.) of Water, filter the solution, add the Alcohol to the filtrate and pass enough Water through the filter to make 20 fluidounces (1000 Cc.).

Each fluidrachm represents 4 grains (0.26 Gm.) of Gluside.

81. LIQUOR HYPOPHOSPHITUM COMPOSITUM SINE SACCHARO.

Compound Solution of Hypophosphites, without Sugar.

Potassium Hypophosphite	320 grains	9.15	Gm.
Calcium Hypophosphite	320 grains	9.15	Gm.
Sodium Hypophosphite	80 grains	2.3	Gm.
Iron Hypophosphite	160 grains	4.58	Gm.
Manganese Hypophosphite	80 grains	2.3	Gm.
Potassium Citrate	300 grains	8.6	Gm.
Citric Acid	100 grains	2.86	Gm.
Quinine (Alkaloid)	80 grains	2.3	Gm.
Strychnine (Alkaloid)	2¼ grains	.064	Gm.
Hypophosphorous Acid (10%)	a sufficient quantity		
Oil of Sweet Orange	12 minims.	0.4	Cc.
Alcohol	10 fluidrachms	15.	Cc.
Gluside	25 grains	0.7	Gm.
Glycerin	20 fluidounces	250	Cc.
Distilled Water, sufficient to make	80 fluidounces	1000	Cc.

Dissolve the Hypophosphites of Potassium, Calcium and Sodium in 28 fluidounces (350 Cc.) of boiling Distilled Water. Dissolve the Hypophosphites of Iron and Manganese, the Citrate of Potassium and Citric Acid, in 8 fluidounces (100 Cc.) of Water with a gentle heat. Dissolve the Alkaloids in a little Water with a sufficient quantity of Hypophosphorous Acids. Mix these solutions and add the Glycerin. Dissolve the Gluside and the Oil of Orange in the Alcohol with gentle heat, and mix with the foregoing solution, then add sufficient Distilled Water to make 80 fluidounces (1000 Cc.).

82. LIQUOR IODI DILUTUS.

Dilute Solution of Iodine.

Iodine.....	440 grains	50 Gm.
Potassium Iodide	600 grains	67.5 Gm.
Distilled Water, sufficient to make	20 fluidounces	1000 Cc.

Dissolve.

83. LIQUOR OPII SEDATIVUS.

Sedative Solution of Opium.

Sedative Liquid.

Extract of Opium	2 ounces 405 grains	72.8 Gm.
Alcohol	6¼ fluidounces	156 Cc.
Water, sufficient to make.....	40 fluidounces	1000 Cc.

Dissolve the Extract of Opium in 16 fluidounces (400 Cc.) of boiling Water. Cool the solution, add the Alcohol and cold Water, filter and add sufficient Water to make 40 fluidounces.

NOTE.—Each fluidrachm represents 4 grains of Extract of Opium.

84. LIQUOR POTASSII CITRATIS.

Solution of Potassium Citrate.

(U.S.P. 1905)

Potassium Bicarbonate	124 grains	40 Gm.
Citric Acid	93 grains	30 Gm.
Distilled Water, sufficient to make.....	3½ fluidounces	500 Cc.

Dissolve the Potassium Bicarbonate and the Citric Acid each in 10 fluidrachms (150 Cc.) of Distilled Water. Filter the solutions separately and wash the filters with enough Distilled Water, to obtain in each case, 15 fluidrachms (225 Cc.). Finally mix the two solutions, and when effervescence has nearly ceased, transfer the liquid to a bottle.

Dose, 4 fluidrachms.

NOTE.—This preparation should be freshly made when wanted.

Oleic
Potassi
Alcohol
Oil of
Ether,

Mi
of Pot
indicate
Ether to

Oil of
Oil of C
Oil of F
Oil of C
Tincture
Concent
Alcohol
Solution
Magnesi
Distilled

Boil
for two
saponifie
Cool and
lastly th
and filter

Diss
preserve

85. LIQUOR SAPONIS ANTISEPTICUS.*Antiseptic Soap Solution*

Oleic Acid	14 fluidounces	350 Cc.
Potassium Hydroxide in solution, (1 in 1) a sufficient quantity.		
Alcohol	6 fluidounces	150 Cc.
Oil of Lavender.....	40 minims.	2·3 Cc.
Ether, sufficient to make.....	40 fluidounces	1000 Cc.

Mix the Oleic Acid and Alcohol and neutralize with the solution of Potassium Hydroxide, using Phenolphthalein Solution as an indicator. Cool and add the Oil of Lavender, then add sufficient Ether to make 40 fluidounces (1000 Cc.).

86. LIQUOR SANTALI FLAVI COMPOSITUS.*Compound Solution of Sandal Oil.*

Oil of Sandal.....	2 fluidounces	50 Cc.
Oil of Cubebs.....	1 fluidounce	25 Cc.
Oil of Copaiba.....	6 fluidrachms	18·75 Cc.
Oil of Pimenta.....	30 minims.	1·5 Cc.
Oil of Cassia.....	30 minims.	1·5 Cc.
Tincture of Buchu.....	6 fluidounces	150 Cc.
Concentrated Infusion of Buchu.....	6 fluidounces	150 Cc.
Alcohol.....	8 fluidounces	200 Cc.
Solution of Potassium Hydroxide.....	6 fluidounces	150 Cc.
Magnesium Carbonate.....	1 Av. ounce	25 Gm.
Distilled Water.....	3 fluidounces	75 Cc.

Boil the Solution of Potash and mix with the Oils; let stand for two days, add the Distilled Water, and shake well (if not saponified boil with the addition of a little more Solution of Potash). Cool and add the Tincture and Infusion of Buchu, the Alcohol, and lastly the Magnesium Carbonate. Mix well, let stand for 24 hours and filter.

87. LIQUOR SODII HYDROXIDI.*Solution of Sodium Hydroxide.**Solution of Soda.*

(U.S.P. 1905)

Sodium Hydroxide.....	5 parts
Distilled Water.....	95 parts

Dissolve the Sodium Hydroxide in the Distilled Water and preserve in a well-stoppered, green glass bottle.

88. LIQUOR ZINGIBERIS.

Solution of Ginger.

Strong Tincture of Ginger (1 in 2).....	10 fluidounces	500 Cc.
Purified Talcum.....	6 $\frac{3}{4}$ ounces	335 Gm.
White Sugar.....	6 $\frac{3}{4}$ ounces	335 Gm.
Distilled Water, sufficient to make.....	20 fluidounces	1000 Cc.

Triturate the Tincture of Ginger with the Sugar and Purified Talcum, add the Distilled Water, shake and filter, returning the first portions of filtrate to the filter, until a clear liquid is obtained.

89. LOTIO CALAMINÆ.

Calamine Lotion.

Levigated Calamine.....	40 grains	2.6 Gm.
Zinc Oxide.....	20 grains	1.3 Gm.
Glycerin.....	20 minims.	1.25 Cc.
Lime Water, sufficient to make.....	1 fluidounce	28.4 Cc.

Elutriate the Calamine and Zinc Oxide by triturating them in a mortar with successive portions of the Water and decanting from the silicious matter, then add the Glycerin.

90. LOTIO CALCIS SULPHURATÆ.

Sulphurated Lime Lotion.

Vleming's Solution.

(N.F. 1906)

Slaked Lime.....	3 ounces 132 grains	165 Gm.
Sublimed Sulphur.....	5 Av. ounces	250 Gm.
Distilled Water, sufficient to make.....	20 fluidounces	1000 Gm.

Mix the Slaked Lime with the Sulphur and add the mixture gradually to 33 fluidounces (1650 Cc.) of boiling water. Then boil the whole under constant stirring until it measures 20 fluidounces (1000 Cc.), strain, and having allowed the solution to become clear by standing in a well-stoppered bottle, decant the clear brown liquid and keep it in completely filled and well stoppered-bottles.

91. MAGMA MAGNESIA.

Magnesia Magma.

(Milk of Magnesia).

(N.F. 1906)

Magnesium Sulphate.....	10 ounces	250 Gm.
Sodium Hydroxide.....	3 ounces	81 Gm.
Distilled Water, sufficient to make.....	40 fluidounces	1000 Cc.

Di
of wat
ounces
Sodium
Sulpha
subside
times w
saline t
drain w
add sul
thoroug
Ea
Magnes
Not
or the ma

Butyl-Cl
Glycerin
Distilled
Mix

Gluside
Sodium I
Chlorofor
Oil of Pi
Oil of Ca
Oil of Cl
Castor Oi
Diss
Bicarbona
and shake

Starch, in
Glycerin .
Water...
Diluted Sc
Boil tl
add the sol

Dissolve the Magnesium Sulphate in 160 fluidounces (4000 Cc.) of water, and the Sodium Hydroxide in another portion of 160 fluidounces (4000 Cc.) of Water, and filter the solutions. Pour the Sodium Hydroxide slowly in a thin stream into the Magnesium Sulphate solution with constant stirring. Allow the precipitate to subside and decant the clear liquid. Wash the Magma several times with water by decantation until the washings are free from saline taste. Transfer the magma to a muslin strainer and allow to drain without pressing. Then re-transfer it to suitable vessels and add sufficient water to make 40 fluidounces (1000 Cc.) and mix thoroughly by stirring.

Each fluidrachm contains about three grains (0.195 Gm.) of Magnesium Hydroxide.

NOTE.—The water used in this preparation must be free from organic matter or the magma will become discolored.

92. MISTURA BUTYL-CHLORAL.

Mixture of Butyl-Chloral.

Butyl-Chloral Hydrate.....	80 grains	9.1 Gm.
Glycerin	5 fluidrachms	35. Cc.
Distilled Water, sufficient to make.....	20 fluidounces	1000 Cc.

Mix and dissolve. Dose, one fluidounce.

93. OLEUM RICINI AROMATICUM.

Aromatic Castor Oil.

"Sweet Castor Oil."

Gluside	7½ grains	0.4 Gm.
Sodium Bicarbonate.....	7½ grains	0.4 Gm.
Chloroform	150 minims.	8. Cc.
Oil of Pimenta	75 minims.	4. Cc.
Oil of Cassia	75 minims.	4. Cc.
Oil of Cloves	75 minims.	4. Cc.
Castor Oil, sufficient to make.....	40 fluidounces	1000 Cc.

Dissolve the Gluside in the Chloroform; add the Sodium Bicarbonate, then add the oils (which have been previously mixed) and shake vigorously.

94. PASTA IODI ET AMYLI.

Iodine and Starch Paste.

Starch, in powder	1 ounce	10 Gm.
Glycerin	2 ounces	20 Gm.
Water.....	6 ounces	60 Gm.
Diluted Solution of Iodine.....	1 ounce	10 Gm.

Boil the Starch in the Glycerin and Water, and when nearly cold, add the solution of Iodine and mix thoroughly.

95. PEPSINUM SACCHARATUM.

Saccharated Pepsin.

(U.S.P. 1905)

Pepsin	10 parts	:
Sugar of Milk, recently dried	90 parts	(

Triturate the Pepsin with the Sugar of Milk to a fine, uniform powder and keep the product in well-stoppered bottles.

Mix t

96. PIGMENTUM IODI COMPOSITUM.

Compound Iodine Paint.

Mandl's Solution.

Iodine	5 grains	0.3 Gm.
Menthol	5 grains	0.3 Gm.
Potassium Iodide	15 grains	1 Gm.
Glycerin	1 fluidounce	28.4 Cc.

Triturate until a perfect solution is obtained.

Salicylic A
Carbolic A
Eucalyptol
Menthol.
Thymol ..
Zinc Sulph
Boric Acid

Tritur
powder, ad
and continu
at a time, u

97. PULVIS ACACIÆ COMPOSITUS.

Compound Acacia Powder.

Powdered Acacia	5 parts
Powdered Tragacanth	5 parts
Powdered Starch	5 parts
Powdered Sugar	5 parts
Powdered Boric Acid	1 part

Triturate the powders together until thoroughly mixed.

NOTE.—Recommended as an emulsifying agent.

Be
Po
Po

Triturat
mortar for t
warmed, and
Dose, ½

98. PULVIS ACETANILIDI COMPOSITUS.

Compound Powder of Acetanilide.

(N.F. 1896)

Acetanilide	7 ounces	70 Gm.
Caffeine	1 ounce	10 Gm.
Sodium Bicarbonate	2 ounces	20 Gm.

Reduce the ingredients separately to a fine powder and mix them thoroughly.

Dose, 3 to 5 grains.

Con
Sug
Mix inti

NOTE.—TI
Humanized Milk

99. PULVIS ALOES ET CANELLÆ.

Powder of Aloes and Canella.

Hiera Picra.

Socotrine Aloes, in fine powder 4 parts

Canella, in fine powder 1 part

Mix them intimately.

100. PULVIS ANTISEPTICUS SOLUBILIS.

Soluble Antiseptic Powder.

(N.F. 1906)

Salicylic Acid75	grains	5 Gm.
Carbolic Acid15	grains	1 Gm.
Eucalyptol15	grains	1 Gm.
Menthol15	grains	1 Gm.
Thymol15	grains	1 Gm.
Zinc Sulphate	4	ounces	125 Gm.
Boric Acid30	ounces	866 Gm.

Triturate the Salicylic Acid and Zinc Sulphate to a very fine powder, add the Carbolic Acid, Eucalyptol, Menthol and Thymol, and continue the trituration, adding the Boric Acid, in small portions at a time, until a uniform impalpable powder is obtained.

101. PULVIS BENZOATIS COMPOSITUS.

Compound Benzoate Powder.

Skeenes' Mixture.

Benzoic Acid 1 part

Potassium Bicarbonate 3 parts

Powdered Sugar 12 parts

Triturate the Benzoic Acid and Potassium Bicarbonate in a hot mortar for ten minutes, then add the Powdered Sugar, previously warmed, and triturate all together, keeping mortar continuously hot.

Dose, $\frac{1}{2}$ to 1 drachm.**102. PULVIS LACTIS COMPOSITUS.**

Compound Milk Powder.

Humanizing Milk Powder.

(N.F. 1906)

Compound Pancreatic Powder 35 parts

Sugar of Milk, in fine powder 965 parts

Mix intimately.

NOTE.—This preparation is intended for convenient use in preparing Humanized Milk. A teaspoonful approximates to about 100 grains (6.5 Gms.).

103. PULVIS PANCREATICUS COMPOSITUS.

Peptonizing Powder.

(N.F. 1906)

Pancreatin	20 parts
Sodium Bicarbonate	80 parts

Mix them by trituration.

NOTE.—To peptonize 16 fluidounces of Fresh Cows' Milk, add 25 grains of Compound Pancreatic Powder to four fluidounces of Tepid Water contained in a suitable flask, and afterwards add 16 fluidounces of Fresh Cows' Milk, previously heated to 100 F. Maintain the mixture at this temperature for thirty minutes, then transfer the flask to a cold place. Milk thus peptonized should not be used when it has been kept over 24 hours, or when it has developed a bitter taste.

104. PULVIS PEPSINI COMPOSITUS.

Compound Powder of Pepsin.

Pulvis Digestivus.

(N.F. 1906)

Saccharated Pepsin	225 grains.	15 Gm.
Pancreatin	225 grains.	15 Gm.
Diastase	15 grains.	1 Gm.
Lactic Acid	15 minims.	1 Cc.
Hydrochloric Acid	30 minims.	2 Cc.
Sugar of Milk, in powder	960 grains.	66 Gm.

Add the Acids gradually to the Sugar of Milk, and triturate until thoroughly mixed. Mix the Pepsin, Pancreatin and Diastase, and then incorporate this mixture by trituration with the Sugar of Milk. Finally rub the mixture through a hair-sieve, and preserve the powder in bottles.

105. PULVIS PRO MISTURA CRETÆ.

Powder for Chalk Mixture.

Prepared Chalk	50 grains.	5.0 Gm.
Powdered Tragacanth	7 grains.	0.7 Gm.
Powdered Sugar	100 grains.	10.0 Gm.

Mix the powders and keep in a well-stoppered bottle.

When required for making Chalk Mixture use 40 grains (2.6 Gm.) of the powder to each fluidounce (28.4 Cc.) of Cinnamon Water.

Santonin
Sub-Chlor
Rhubarb,
Sugar ...
Oil of Pep

Mix i

NOTE.
17 grains of

D
Pc
Sc
Sc

Tritura
powder. T
the crystalli

NOTE.—
that is similar

The qu
(1000 Cc.) of

1. Oil of Ga
2. Oil of Ga
Oil of Sa
3. Compound
4. Oil of Gau
Oil of Bitt
Oil of Cori
5. Oil of Gau
Oil of Sas
Oil of Bitt
6. Oil of Gau
Oil of Bitt

106. PULVIS SANTONINI COMPOSITUS.

Compound Powder of Santonin.

Santonin	125 grains	1.25 Gm.
Sub-Chloride of Mercury	125 grains	1.25 Gm.
Rhubarb, in fine powder	200 grains	2.00 Gm.
Sugar	50 grains	0.5 Gm.
Oil of Peppermint	15 minims.	1 Cc.

Mix intimately.

NOTE.—Each 4 grains contains 1 grain each of Santonin and Calomel, and 1½ grains of Rhubarb.

107. SAL CAROLINUM FACTITIUM.

Artificial Carlsbad Salt.

(N.F. 1906)

Dried Sodium Sulphate.....	44 parts
Potassium Sulphate.....	2 parts
Sodium Chloride, purified	18 parts
Sodium Bicarbonate	36 parts

Triturate the ingredients, previously well dried to a fine, uniform powder. The dried Sodium Sulphate is prepared by slowly drying the crystalline salt until it has lost one-half its weight.

NOTE.—Fifty-three grains dissolved in one pint of water gives a solution, that is similar to Carlsbad (Sprudel) Water, in its essential constituents.

108. SAPORES PRO EMULSIONIBUS.

Flavours for Emulsions.

(N.F. 1906)

The quantities given below are intended for 40 fluidounces (1000 Cc.) of finished Emulsion of Cod Liver Oil.

1. Oil of Gaultheria.....	78 minims.	4.0 Cc.
2. Oil of Gaultheria	40 minims.	2.0 Cc.
Oil of Sassafras	40 minims.	2.0 Cc.
3. Compound Spirit of Orange.....	30 minims.	1.5 Cc.
4. Oil of Gaultheria	40 minims.	2.0 Cc.
Oil of Bitter Almond.....	4 minims.	0.25 Cc.
Oil of Coriander.....	4 minims.	0.25 Cc.
5. Oil of Gaultheria	30 minims.	1.5 Cc.
Oil of Sassafras	30 minims.	1.5 Cc.
Oil of Bitter Almond.....	4 minims.	0.25 Cc.
6. Oil of Gaultheria	48 minims.	2.5 Cc.
Oil of Bitter Almond.....	48 minims.	2.5 Cc.

109. SPIRITUS ASPARAGI COMPOSITUS.

Compound Spirit of Asparagus.

Gum Guaiacum,	1 ounce	28.4 Gm.
Asparagus Seed	1 ounce	28.4 Gm.
Parsley Seed	1 ounce	28.4 Gm.
Black Haw	2¼ ounces	64 Gm.
Henbane Leaves	100 grains	6.5 Gm.
Compound Spirit of Orange	4 fluidrachms	15 Cc.

Diluted Alcohol, a sufficient quantity.

Reduce the drugs to a powder and percolate with Diluted Alcohol to make 15½ fluidounces (425 Cc.) to which add the Compound Spirit of Orange.

110. SPIRITUS AMYGDALÆ AMARÆ.

Spirit of Bitter Almond.

(U.S.P. 1905)

Oil of Bitter Almond	70 minims.	10 Cc.
Alcohol (95%)	16 fluidounces	800 Cc.
Distilled Water, sufficient to make	20 fluidounces	1000 Cc.

Dissolve the Oil in the Alcohol, and add enough Water to make 20 fluidounces (1000 Cc.).

111. SPIRITUS AURANTII.

Spirit of Orange.

Fresh Oil of Sweet-Orange Peel	1 fluidounce	10 Cc.
Deodorized Alcohol	9 fluidounces	90 Cc.

Mix.

112. SPIRITUS AURANTII COMPOSITUS.

Compound Spirit of Orange.

(U.S.P. 1905)

Oil of Orange Peel	4 fluidounces	200 Cc.
Oil of Lemon	1 fluidounce	50 Cc.
Oil of Coriander	3½ fluidrachms	20 Cc.
Oil of Anise,	48 minims.	5 Cc.
Deodorized Alcohol, sufficient to make	20 fluidounces	1000 Cc.

Mix them. Keep in completely filled, well-stoppered bottles, in a cool, dark place.

Mucilag
Simple
Mi:

Codeine
Alcohol
Distilled
Syrup,
Dis:
add the

NOTE
The streng

Fluid Ex
Fluid Ex
Fluid Ex
Fluid Ex
Ammonit
Magnesi
Compoun
Sugar ..
Water, si

Tritu
with the
Water, a
moistene
16 fluidoi
and Amm
fluidounce

113. SYRUPUS ACACIÆ.

Syrup of Acacia.

Mucilage of Acacia	1 fluidounce	25 Cc.
Simple Syrup.....	3 fluidounces	75 Cc.

Mix.

114. SYRUPUS CODEINÆ PHOSPHATIS.

Syrup of Codeine Phosphate.

Codeine Phosphate	40 grains	4.57 Gm.
Alcohol (95%)	7 fluidrachms	47.5 Cc.
Distilled Water.....	3 fluidrachms	18.75 Cc.
Syrup, a sufficient quantity to make.	20 fluidounces	1000 Cc.

Dissolve the Codeine Phosphate in the Water and Alcohol, then add the Syrup.

NOTE.—Recommended as more stable than the official Syrup of Codeine. The strength is identical with Syrupus Codeinæ, P.B.

115. SYRUPUS EUCALYPTI COMPOSITUS.

Compound Syrup of Eucalyptus.

Fluid Extract of Eucalyptus.....	5 fluidounces	125 Cc.
Fluid Extract of Horehound	2 fluidounces	50 Cc.
Fluid Extract of Elecampane	2 fluidounces	50 Cc.
Fluid Extract of Licorice	2 fluidounces	50 Cc.
Fluid Extract of Comfrey	2 fluidounces	50 Cc.
Ammonium Chloride.....	480 grains	28 Gm.
Magnesium Carbonate.....	240 grains	14 Gm.
Compound Spirit of Orange.....	4 fluidrachms	12.5 Cc.
Sugar	20 ounces	500 Gm.
Water, sufficient to make.....	40 fluidounces	1000 Cc.

Triturate the Fluid Extracts and Compound Spirit of Orange with the Magnesium Carbonate, and 8 fluidounces (200 Cc.) of Water, and let stand two hours. Filter through a previously moistened filter, passing enough Water through the filter to make 16 fluidounces (400 Cc.) of filtrate, in which dissolve the Sugar and Ammonium Chloride, then add sufficient Water to make 40 fluidounces (1000 Cc.).

116. SYRUPUS FERRI ET MANGANI IODIDI.

Syrup of Iodide of Iron and Manganese.

(N.F. 1906)

Iodine	3 ozs. 172 grains	81.5 Gm.
Iron Wire, fine, bright, and finely cut $1\frac{1}{4}$ ounce		28.7 Gm.
Manganese Sulphate	1 oz. 48 grains	26.5 Gm.
Potassium Iodide	1 oz. 137 grains	31.5 Gm.
Diluted Alcohol, a sufficient quantity.		
Distilled Water, sufficient to make 40 fluidounces or 1000 Cc.		

Mix the Iron with 10 fluidounces (250 Cc.) of Distilled Water in a flask, add the Iodine, and prepare a solution of Ferrous Iodide, in the usual manner, aiding the process, if necessary, by heating the contents of the flask, at first gently, and finally to the boiling point. Filter the liquid through a small filter, directly upon the Sugar contained in a suitable bottle. Dissolve the Manganese Sulphate in 5 fluidounces (125 Cc.) of Distilled Water, and the Potassium Iodide in 4 fluidounces (100 Cc.) of Diluted Alcohol. Mix the two solutions, and filter into the same bottle which contains the Sugar and the Iron Solution. Wash the filter with 10 fluidrachms (32 Cc.) of cold Distilled Water, receiving the washings in the same bottle. Agitate until the Sugar is dissolved, and if necessary, strain. Finally, make up the volume with Distilled Water to 40 fluidounces (or 1000 Cc.).

NOTE.—Each fluidrachm contains about 6 (0.4 Gm.) grains of Ferrous Iodide and 3 (0.2 Gm.) grains of Manganese Iodide.

Average dose, 15 minims. (1 Cc.).

117. SYRUPUS FERRI PHOSPHATIS COMPOSITUS.

Compound Syrup of Phosphates of Iron.

"Parrish's Chemical Food"

(N.F. 1906)

Precipitated Calcium Carbonate	1 oz. 200 grains	35 Gm.
Soluble Ferric Phosphate	320 grains	17.5 Gm.
Ammonium Phosphate	320 grains	17.5 Gm.
Potassium Bicarbonate	75 grains	4 Gm.
Sodium Bicarbonate	75 grains	4 Gm.
Citric Acid	$3\frac{1}{8}$ ounces	82 Gm.
Glycerin	15 fluidounces	375 Cc.
Phosphoric Acid (B.P.)	$3\frac{3}{4}$ ounces	90.7
Orange Flower Water	5 fluidounces	125 Cc.
Tincture of Cudbear	5 fluidrachms	16 Cc.
Sugar	$12\frac{1}{2}$ ounces	300 Gm.
Water, sufficient to make 40 fluidounces 1000 Cc.		

Tritu
and Sodii
Flower W
solution h
Ammoniu
and add tl
through a
and receiv
Agitate ur
bear, and

NOTE
Phosphate,
and smaller

Avera

11

Licorice R
Solution o
Oil of Cori
Oil of Clov
Alcohol (9
Granulatec
Water, sul

Macer
Distilled V
Ammonia,
colature.
menstruum
reserving
colatures
concentrate
To the fil
and dissolv
add enough

If pref
foregoing :

Fluid Extr:
Mixture
Oil of Cori
Oil of Clov
Alcohol (95
Granulated
Water, suff

Triturate the Precipitated Calcium Carbonate with the Potassium and Sodium Bicarbonates, the Citric Acid, Glycerin and Orange Flower Water, and gradually add the Phosphoric Acid, stirring until solution has been effected. Dissolve the Ferric Phosphate and the Ammonium Phosphate in 10 fluidounces (250 Cc.) of Hot Water, cool and add the solution to that previously prepared. Filter the whole through a pellet of absorbent cotton placed in the neck of a funnel, and receive the filtrate in a graduated bottle containing the Sugar. Agitate until the latter is dissolved, then add the Tincture of Cudbear, and lastly, enough Water to make 40 fluidounces (1000 Cc.).

NOTE.—Each fluidrachm contains about 2 grains (0.13 Gm.) of Calcium Phosphate, 1 grain (0.065 Gm.) each of Phosphates of Iron and of Ammonium and smaller quantities of Sodium and Potassium Phosphates.

Average dose, 1 fluidrachm (4Cc.).

118. SYRUPUS GLYCYRRHIZÆ AROMATICUS.

Aromatic Syrup of Licorice.

Licorice Root, cut small	8 ounces	200	Gm.
Solution of Ammonia	1 fluidounce	25	Cc.
Oil of Coriander	20 minims.	1	Cc.
Oil of Cloves	10 minims.	0.5	Cc.
Alcohol (95%)	2 fluidounces	50	Cc.
Granulated Sugar	27 ounces	675	Gm.
Water, sufficient to make	40 fluidounces	1000	Cc.

Macerate the Licorice Root with 16 fluidounces (400 Cc.) of Distilled Water mixed with 160 minims. (8.5 Cc.) of Solution of Ammonia, for twelve hours; strain and express, reserving the colature. Repeat this operation with the pressed marc and new menstruum of Ammonia and Water twice, straining, pressing and reserving the colature after each maceration. Mix the several colatures and evaporate over a water-bath until the liquid is concentrated to 16 fluidounces (400 Cc.), then cool and filter. To the filtrate add the Oils, previously dissolved in the Alcohol, and dissolve the Sugar, by percolation, in the mixed liquids, then add enough water to make 40 fluidounces (1000 Cc.).

If preferred, the following formula may be substituted for the foregoing :

Fluid Extract of Licorice (for Quinine Mixtures)	8 fluidounces	200	Cc.
Oil of Coriander	20 minims.	1	Cc.
Oil of Cloves	10 minims.	0.5	Cc.
Alcohol (95 per cent.)	2 fluidounces	50	Cc.
Granulated Sugar	27 ounces.	675	Gm.
Water, sufficient to make	40 fluidounces	1000	Cc.

Mix the Fluid Extract with the Alcohol, in which the Oils have been previously dissolved, and 8 fluidounces (200 Cc.) of Distilled Water. Dissolve the Sugar in this liquid and add enough Water to make 40 fluidounces (1000 Cc.).

119. SYRUPUS HYPOPHOSPHITUM COMPOSITUS

Compound Syrup of Hypophosphites.

Calcium Hypophosphite	2¼ oz.	58 grains	29.5 Gm.
Sodium Hypophosphite	2¾ oz.	80 grains	36.5 Gm.
Potassium Hypophosphite	1¼ oz.	94 grains	18.3 Gm.
Manganese Hypophosphite		80 grains	2.3 Gm.
Quinine		40 grains	1.15 Gm.
Strychnine		10 grains	0.28 Gm.
Ferrous Sulphate, in crystals		120 grains	3.45 Gm.
Dilute Hypophosphorous Acid, a sufficient quantity.			
Concentrated Phosphoric Acid		45 minims.	1.15 Cc.
Granulated Sugar		65 ounces	813 Gm.
Distilled Water, sufficient to make		80 fluidounces	1000 Cc.

Dissolve the Sodium and Potassium Hypophosphites and 960 grains (27.2 Gm.) of Calcium Hypophosphite in 35 fluidounces (437.5 Cc.) of boiling Distilled Water. Dissolve the Manganese Hypophosphite in 5 fluidounces (62.5 Cc.) of hot Distilled Water, then dissolve the Alkaloids in this solution, with the aid of a minimum quantity of Dilute Hypophosphorous Acid. Mix the two solutions and filter, if necessary. Make a syrup by dissolving the Sugar in the filtrate by percolation. Dissolve the Ferrous Sulphate in 6 fluidrachms (9.5 Cc.) of Water, previously mixed with the Concentrated Phosphoric Acid. Also dissolve 82 grains (2.3 Gm.) of Calcium Hypophosphite in 6 fluidrachms (9.5 Cc.) of Water; mix this solution with the Ferrous solution, let the mixture stand for twelve hours and filter out the precipitate. (The filtrate will contain approximately 80 grains of Ferrous Hypophosphite.) Mix the filtrate with the Syrup and pass enough Water through the percolator to make the finished product measure 80 fluidounces (1000 Cc.).

NOTE.—Each fluidounce of this Syrup contains Sodium Hypophosphite, 16 grains; Calcium Hypophosphite, 12 grains; Potassium Hypophosphite, 8 grains; Manganese and Ferrous Hypophosphites, 1 grain each; Quinine, ½ grain, and Strychnine, ½ grain.

Dose, one to two fluidrachms.

Tar.....
White Sand.
Glycerin...
Sugar.....
Water, suffi

Mix the
ouces (200
pour off the
of boiling
for 15 min
occasionally
Dissolve the
pass enough
(1000 Cc.).

121.

Potassium E
Magnesium
Precipitated
Quinine Hy
Strychnine
Orange Flo
centrate
Phosphoric
Soluble Ferr
Water.....
Syrup, suffic

Dissolv
Phosphoric
Then dissolv
warmed, and
Syrup to ma

120. SYRUPUS PICIS LIQUIDÆ.

Syrup of Tar.

Tar.....	3 ounces	75 Gm.
White Sand.....	4 ounces	100 Gm.
Glycerin.....	4 fluidounces	100 Cc.
Sugar.....	32 ounces	800 Gm.
Water, sufficient to make.....	40 fluidounces	1000 Cc.

Mix the Tar intimately with the White Sand, pour on 8 fluidounces (200 Cc.) of Water, and stir frequently for 12 hours, then pour off the Water and throw it away. Pour 16 fluidounces (400 Cc.) of boiling Water upon the residue, stir well and frequently for 15 minutes, add the Glycerin and set aside for 24 hours, occasionally stirring, and decant the clear solution and filter. Dissolve the Sugar in the filtrate, with gentle heat, cool, strain and pass enough Water through the strainer to make 40 fluidounces (1000 Cc.).

121. SYRUPUS QUININÆ PHOSPHO-MURIATIS.

Syrup of Phospho-Muriate of Quinine.

Potassium Bicarbonate.....	616 grains	35 Gm.
Magnesium Carbonate.....	352 grains	20 Gm.
Precipitated Calcium Carbonate.....	352 grains	20 Gm.
Quinine Hydrochloride.....	70 grains	4 Gm.
Strychnine Hydrochloride.....	2½ grains	0·14 Gm.
Orange Flower Water, natural, concentrated.....	5 fluidounces	125 Cc.
Phosphoric Acid, 85%.....	5 fluidounces	125 Cc.
Soluble Ferric Phosphate.....	282 grains	16 Gm.
Water.....	310 minims.	16 Cc.
Syrup, sufficient to make.....	40 fluidounces	1000 Cc.

Dissolve the several Carbonates and the Alkaloidal Salts in the Phosphoric Acid, previously diluted with the Orange Flower Water. Then dissolve the Soluble Ferric Phosphate in the Water, previously warmed, and add it to the foregoing solution, and lastly add sufficient Syrup to make 40 fluidounces (1000 Cc.).

122. SYRUPUS RUBI AROMATICUS

Aromatic Syrup of Blackberry.

(N.F. 1906)

Blackberry Root Bark	5 ounces	125 Gm.
Cinnamon Bark	262 grains	15 Gm.
Nutmeg	262 grains	15 Gm.
Cloves	140 grains	8 Gm.
Allspice	140 grains	8 Gm.
Granulated Sugar	26 ounces	650 Gm.
Alcohol (95%)		
Water, Blackberry Juice (fresh) of each, sufficient to make	40 fluidounces	1000 Cc.

Reduce the Blackberry Root Bark and the Aromatics to a moderately coarse (No. 40) powder and percolate in the usual manner with a menstruum of equal volumes of Alcohol and Water, until 10 fluidounces (250 Cc.) of percolate are obtained. To this, add 18 fluidounces (450 Cc.) of Blackberry Juice, and dissolve the Sugar in the liquid by agitation. Lastly, add enough Blackberry Juice to make 40 fluidounces (1000 Cc.).

123. SYRUPUS SARSÆ COMPOSITUS.

Compound Syrup of Sarsaparilla.

(U.S.P. 1905)

Fluid Extract of Sarsaparilla	8 fluidounces	200 Cc.
Fluid Extract of Glycyrrhiza	5 fluidrachms	15 Cc.
Fluid Extract of Senna	5 fluidrachms	15 Cc.
Sugar	26 ounces	650 Gm.
Oil of Sassafras	2 minims.	0.2 Cc.
Oil of Anise	2 minims.	0.2 Cc.
Oil of Gaultheria	2 minims.	0.2 Cc.
Water, sufficient to make	40 fluidounces	1000 Cc.

Add the Oils to the mixed Fluid Extracts, and shake the liquid thoroughly, then add Water enough to make up the volume to 24 fluidounces (600 Cc.), and mix well. Set the mixture aside for one hour, and then filter it. Dissolve the Sugar in the filtrate with the aid of a gentle heat. Cool, strain, and add enough water through the strainer to make the finished product measure 40 fluidounces (1000 Cc.).

124. SYRUPUS SENEGÆ.

Syrup of Senega.

Fluid Extract of Senega	8 ounces	200 Cc.
Glycerin	2 ounces	50 Cc.
Sugar	40 ounces	1000 Gm.
Magnesium Carbonate	360 grains	20.5 Gm.
Distilled Water, sufficient to make	40 fluidounces	1000 Cc.

Mix the Fluid Extract and Glycerin, then triturate with the

Magnesium Carbonate, and dually add 10 fluidounces of Sugar in the form of a solution, and add Water, if necessary, to make 40 fluidounces (1000 Cc.).

121

Fluid Extract
Fluid Extract
Fluid Extract
Fluid Extract
Fluid Extract
Fluid Extract
Potassium Iodide
Sugar
Water, sufficient to make
Mix the Fluid Extract and Water, let stand for 24 hours, and then filter it. Potassium Iodide
Water to make

Solution
Simple
Mix.

NOTE.—This is a Solution of Ginger.

Potassium Iodide
Iodine, resublimed
Sodium Hydroxide
Thymol, in crys-
tals
Distilled Water

Dissolve the Iodine in 28.4 Cc. of Distilled Water and a half fluidounce of Distilled Water. Dissolve the Sodium Hydroxide in Distilled Water and dilute with Water. Mix the solutions slowly and wash the residue with Distilled Water.

Magnesium Carbonate and 4 ounces (100 Gm.) of Sugar, then gradually add 10 ounces (250 Cc.) of Water and filter. Dissolve the Sugar in the remainder of the filtrate by the percolation method, and add Water, if necessary, to make 40 fluidounces (1000 Cc.).

125. SYRUPUS TRIFOLII COMPOSITUS.

Compound Syrup of Trifolium.

Compound Syrup of Red Clover Blossoms.

Fluid Extract of Red Clover Blossoms...	20 fluidrachms	64	Cc.
Fluid Extract of Burdock	10 fluidrachms	32	Cc.
Fluid Extract of Berberis Aquifolium...	10 fluidrachms	32	Cc.
Fluid Extract of Stillingia	10 fluidrachms	32	Cc.
Fluid Extract of Poke Root.....	10 fluidrachms	32	Cc.
Fluid Extract of Cascara Amarga.....	10 fluidrachms	32	Cc.
Fluid Extract of Prickly Ash Bark.....	2½ fluidrachm	8	Cc.
Potassium Iodide	320 grains	18.25	Gm.
Sugar.....	40 ounces	1000	Gm.
Water, sufficient to make	40 fluidounces	1000	Cc.

Mix the Fluid Extracts with 12½ fluidounces (312.5 Cc.) of Water, let stand for a few hours, filter, and dissolve the Sugar and Potassium Iodide in the filtrate, and strain; then add sufficient Water to make 40 fluidounces (1000 Cc.).

126. SYRUPUS ZINGIBERIS.

Syrup of Ginger.

Solution of Ginger
 1 | volume |

Simple Syrup
 9 | volumes |

Mix.

NOTE.—This preparation is similar in strength to the official Syrup of Ginger.

127. THYMOLIS IODIDUM.

Thymol Iodide.

Dithymol-Diiodide.

Potassium Iodide.....	124	grains	8	Gm.
Iodine, resublimed	93	grains	6	Gm.
Sodium Hydroxide	27½	grains	1.8	Gm.
Thymol, in crystals	27	grains	1.7	Gm.
Distilled Water, a sufficient quantity.				

Dissolve the Iodine and Potassium Iodide in one fluidounce (28.4 Cc.) of Distilled Water and add Distilled Water to make one and a half fluidounces (42.6 Cc.)

Dissolve the Sodium Hydroxide in 1 fluidounce (42.6 Cc.) of Distilled Water, and in this solution dissolve the Thymol, and dilute with Water to make 1½ fluidounces (42.6 Cc.) Into this solution slowly pour the Iodine Solution under constant stirring, and wash the resulting precipitate, by alternate affusion and decantation with Distilled Water, then drain and dry carefully.

128. TINCTURA AURANTII CORTICIS DULCIS RECENTIS.

Tincture of Fresh Sweet-Orange Peel.

Fresh Sweet-Orange Peel	5 ounces	250 Gm.
Rectified Spirit, sufficient to make	20 fluidounces	1000 Cc.

Prepare by the maceration process.

229. TINCTURA CARMINATIVA.

Carminative Tincture.

Spirit of Chloroform	5 fluidounces	250 Cc.
Compound Tincture of Cardamon	5 fluidounces	250 Cc.
Compound Tincture of Lavender	5 fluidounces	250 Cc.
Aromatic Spirit of Ammonia	5 fluidounces	250 Cc.

Mix.

Adult dose, 1 fluidrachm (4 Cc.).

130. TINCTURA FERRI CITRO-CHLORIDI.

Tincture of Citro-Chloride of Iron.

Tasteless Tincture of Iron.

Strong Solution of Ferric Chloride, B.P.	5 fluidounces	125 Cc.
Citric Acid, in powder	6 ounces	200 grains 160.5 Gm.
Sodium Bicarbonate	7 ounces	300 grains 190.75 Gm.
Alcohol, (95%)	3 fluidounces	75 Cc.
Distilled Water, sufficient to make	20 fluidounces	500 Cc.

Heat $7\frac{1}{2}$ fluidounces (187.5 Cc.) of Water to the boiling point, and dissolve in it the Citric Acid, continuing the heat while adding the Sodium Bicarbonate in portions, and stirring with a glass rod, if necessary; when effervescence has ceased, add the strong solution of Ferric Chloride and cool the mixture. Then add the Alcohol and sufficient Distilled Water to make 20 fluidounces (500 Cc.)

NOTE.—This preparation is practically identical in strength of Iron, but not in the quantity of Alcohol, with Tinctura Ferri Perchloridi P.B.

131. TINCTURA IGNATIÆ ALKALINA.

Alkaline Tincture of Ignatia.

Gouttes Amères de Baume.

St. Ignatius' Bean	20 ounces	500 Gm.
Potassium Carbonate	90 grains	5.2 Gm.
Alcohol (60%), sufficient to make	40 fluidounces	1000 Cc.

Macerate for ten days and filter.

Dose, 5 to 20 minims (0.3 to 1.2 Cc.).

Iodine, resublim
Potassium Iod
Water
Alcohol (95%),

Dissolve 1
Iodine, and las
pleted, measur

Iodine, resublim
Strong Solution
Rectified Spirit
Dissolve 1
of Ammonia.

Cudbear
Alcohol (95%).
Water, of each,

Prepare by
and Water, to
obtained.

NOTE.—This
tint or color is to b

135.

Cudbear
Caramel
Alcohol, (95%),
Water, of each,

Mix Alcoho
the Cudbear wit
twenty-four hou
add the Carame
Water. Then
through the fil
(1000 Cc.).

NOTE.—This
brownish-red tint or

132. TINCTURA IODI, CHURCHILL.

Churchill's Tincture of Iodine.

(N.F. 1906)

Iodine, resublimed.....	3 ozs. 131 grains	165 Gm.
Potassium Iodide.....	289 grains	33 Gm.
Water.....	5 fluidounces	250 Cc.
Alcohol (95%), sufficient to make.....	20 fluidounces	1000 Cc.

Dissolve the Potassium Iodide in the Water, then add the Iodine, and lastly, enough Alcohol to make the Tincture, when completed, measure 20 fluidounces (1000 Cc.).

133. TINCTURA IODI DECOLORATA.

Decolorized Tincture of Iodine.

Iodine, resublimed.....	250 grains	26.0 Gm.
Strong Solution of Ammonia.....	10 fluidrachms	62.5 Gm.
Rectified Spirit, sufficient to make...	20 fluidounces	1000 Cc.

Dissolve the Iodine in the Alcohol and add the Strong Solution of Ammonia. Keep the mixture in a warm place until decolorized.

134. TINCTURA PERSIONIS.

Tincture of Cudbear.

Cudbear.....	5 ounces	125 Gm.
Alcohol (95%).		
Water, of each, sufficient to make.....	40 fluidounces	1000 Cc.

Prepare by maceration with a mixture of Alcohol, one volume, and Water, two volumes, until 40 fluidounces (1000 Cc.) are obtained.

NOTE.—This preparation is intended as a coloring agent when a *bright-red* tint or color is to be produced, particularly in acid liquids.

135. TINCTURA PERSIONIS COMPOSITA.

Compound Tincture of Cudbear.

(N.F. 1906)

Cudbear.....	300 grains	17 Gm.
Caramel.....	4 ounces	100 Gm.
Alcohol, (95%), and		
Water, of each, sufficient to make.....	40 fluidounces	1000 Cc.

Mix Alcohol, one volume, with Water, two volumes. Macerate the Cudbear with 30 fluidounces (750 Cc.) of the menstruum during twenty-four hours, agitating occasionally; filter through paper and add the Caramel, previously dissolved in 5 fluidounces (125 Cc.) of Water. Then pass sufficient of the before-mentioned menstruum through the filter to make the whole measure 40 fluidounces (1000 Cc.).

NOTE.—This preparation is intended as a coloring agent, when a *brownish-red* tint or color is to be produced.

136. TINCTURA TOLUTANA SOLUBILIS.

Soluble Tincture of Tolu.

(N.F. 1906)

Tolu Balsam	3½ ounces	100 Gm.
Magnesium Carbonate	200 grains	12 Gm.
Glycerin	16 fluidounces	400 Cc.
Water, and Alcohol (95%), of each, sufficient to make..	40 fluidounces	1000 Cc.

Mix 8 fluidounces (200 Cc.) of Alcohol with the Glycerin, and dissolve the Tolu Balsam in the mixture, with the aid of heat, avoiding loss by evaporation. Then add 15 fluidounces (375 Cc.) of Water and allow the mixture to cool. Pour off the milky liquid from the resinous precipitate (which latter is to be rejected), mix it with the Magnesium Carbonate, by trituration, and filter. Lastly, pass enough of a mixture of Alcohol, *one volume*, and Water, *two volumes*, through the filter, to make the whole filtrate measure 40 fluidounces (1000 Cc.).

137. UNGUENTUM ACIDI CARBOLICI COMPOSITUM.

Compound Ointment of Carbolic Acid.

Mercuric Nitrate Ointment, B.P.	4 ounces	40 Gm.
Sublimed Sulphur	1 ounce	10 Gm.
Phenol	2 ounces	20 Gm.
Olive Oil	2 ounces	20 Gm.
Yellow Wax	2 ounces	20 Gm.

Dissolve the Sulphur in the previously heated Olive Oil and melt the Wax in this solution with a gentle heat. Stir while cooling, and when nearly cold, add the Phenol, and stir until dissolved. Rub the Mercuric Nitrate Ointment in a mortar until smooth; then incorporate with it the mixture previously prepared.

138. UNGUENTUM ICHTHYOLIS COMPOSITUM.

Compound Ichthyol Ointment.

Ichthyol	1 ounce	20 Gm.
Solution of Lime	4 fluidounces	80 Cc.
Anhydrous Wool-Fat	2½ ounces	50 Gm.
Soft Paraffin	5 ounces	100 Gm.
Zinc Ointment	2½ ounces	50 Gm.

Triturate the Ichthyol with the Lime Water; add the Wool Fat gradually, under constant trituration, and then the other ingredients in a similar manner.

136

Iodine

Soft Paraffin ..

Finely po

add the Powde

fully combined

140

Lead Plaster

Soft Paraffin.

Oil of Bergam

Melt the L

approaches the

until it conge

141. U

Phenol.

Campho

Hydro:

Yellow |

Yellow |

Liquify the

gentle heat, ar

Phenol and Cau

142. L

Resorcin ...

Zinc Oxide ...

Bismuth Subni

Oil of Cade....

White Paraffin

Hydrous Wool-

Triturate th

Ether, and allo

incorporate with

NOTE.—Darl

tightly covered co

139. UNGUENTUM IODI DENIGRESCENS.

Stainless Iodine Ointment.

Iodine	1 ounce	10 Gm.
Soft Paraffin	19 ounces	190 Gm.

Finely powder the Iodine ; heat the Paraffin until liquefied, then add the Powdered Iodine, continuing a gentle heat, and stirring until fully combined, then remove from heat and stir until it congeals.

140. UNGUENTUM EMPLASTRI PLUMBI.

Ointment of Lead Plaster.

Diachylon Ointment.

Lead Plaster	1 ounce	110 Gm.
Soft Paraffin	1 ounce	110 Gm.
Oil of Bergamot	4 minims.	1 Cc.

Melt the Lead Plaster and Paraffin together ; when the mixture approaches the temperature of 160° or 170° F., add the Oil and stir until it congeals.

141. UNGUENTUM PHENOLIS CAMPHORATUM.

Camphorated Phenol Ointment.

Phenol	15 parts
Camphor	30 parts
Hydrous Wool-Fat	60 parts
Yellow Beeswax	40 parts
Yellow Soft Paraffin	300 parts

Liquify the Paraffin, Beeswax and Wool-Fat, by the aid of a gentle heat, and while the mixture is still warm, dissolve in it the Phenol and Camphor, and stir until it congeals.

142. UNGUENTUM RESORCINI COMPOSITUM.

Compound Resorcin Ointment.

(N.F. 1906)

Resorcin	1 ounce	85 grains	60 Gm.
Zinc Oxide	1 ounce	85 grains	60 Gm.
Bismuth Subnitrate	1 ounce	85 grains	60 Gm.
Oil of Cade	2 3/8	fluidounces	120 Cc.
White Paraffin Ointment	7 ounces		350 Gm.
Hydrous Wool-Fat	7 ounces		350 Gm.

Triturate the Resorcin to a fine powder with the aid of a little Ether, and allow the Ether to evaporate ; then mix the powders and incorporate with the remainder of the ingredients

NOTE.—Darkens on exposure to air and light, and should be kept in tightly covered containers.

143. UNGUENTUM SULPHURIS COMPOSITUM.

Compound Sulphur Ointment.

Wilkinson's Ointment—Hebra's Itch Ointment.

Precipitated Calcium Carbonate.....	1 ounce	10 Gm.
Sublimed Sulphur.....	1½ ounce	15 Gm.
Oil of Cade.....	1½ ounce	15 Gm.
Soft Soap.....	3 ounces	30 Gm.
Lard.....	3 ounces	30 Gm.

Mix the Lard with the Soft Soap and Oil of Cade. Then gradually incorporate the Sublimed Sulphur and Precipitated Calcium Carbonate.

144. UNGUENTUM SULPHURIS ET RUSCI COMPOSITUM.

Compound Ointment of Sulphur and Birch Tar.

Sublimed Sulphur, sifted.....	32 parts
Potassium Carbonate.....	2 parts
Oil of Birch Tar, Russian.(Oleum Rusci)....	2 parts
Zinc Ointment.....	16 parts
Benzoated Lard.....	32 parts

Mix intimately by trituration, in order to produce a smooth and homogeneous ointment.

145. UNGUENTUM SULPHURIS CINEREI COMPOSITUM.

Compound Grey-Sulphur Ointment.

(Edinburgh)

Grey Sulphur.(Sulphur Vivum).....	8 ounces	227 Gm.
Potassium Nitrate.....	60 grains	3.9 Gm.
Powdered White Hellebore.....	1 ounce	28.4 Gm.
Green Soap.....	3 ounces	84 Gm.
Phenol.....	120 grains	7.8 Gm.
Oil of Bergamot.....	30 minims.	2 Gm.
Lard.....	24 ounces	682 Gm.
Water, a sufficient quantity.		

Mix the Lard and Soap, and incorporate the Grey Sulphur and Powdered Hellebore with the mixture. Then add the Potassium Nitrate (previously dissolved in a little water), and the Oil of Bergamot, and lastly the Phenol.

Zinc Stearate,
White ParaffinLiquify the
the Zinc Stear
smooth, then sFluid Extract c
Alcohol (95%)
Sugar.....
Red Wine, suDissolve t
add the Alcoh
to make the li
mixture aside f
Dose, 4 flPepsin.....
Hydrochloric A
Glycerin.....
Sherry, a suffici
Dissolve thMorrhual (Gad
Fluid Extract o
Glycerin.....
Syrup of Wild
Liquid Extract
Compound Syr
Fuller's Earth,
Sherry Wine, sMix the M
Fuller's Earth,
allow it to stanc
add the Syrup c
to make 40 flui

146. UNGUENTUM ZINCI STEARATIS.

Ointment of Zinc Stearate.

Zinc Stearate, in fine powder	1 ounce.	50 Gm.
White Paraffin Ointment	1 ounce.	50 Gm.

Liquify the Paraffin Ointment by the aid of a water bath, add the Zinc Stearate, continuing the heat until the mixture becomes smooth, then stir while cooling, until it congeals.

147. VINUM COCÆ.

Wine of Coca.

(U.S.P. 1905)

Fluid Extract of Coca	2 $\frac{3}{4}$ fluidounces	65 Cc.
Alcohol (95%)	3 fluidounces	75 Cc.
Sugar	3 ounces	75 Gm.
Red Wine, sufficient to make	40 fluidounces	1000 Cc.

Dissolve the sugar in 20 fluidounces (500 Cc.) of Red Wine, add the Alcohol and Fluid Extract of Coca, and enough Red Wine to make the liquid measure 40 fluidounces (1000 Cc.). Set the mixture aside for two days, then filter.

Dose, 4 fluidrachms (16 Cc.)

148. VINUM PEPSINI.

Wine of Pepsin.

Pepsin	320 grains	36.5 Gm.
Hydrochloric Acid	2 fluidrachms	12.5 Cc.
Glycerin	1 fluidounce	50 Cc.
Sherry, a sufficient quantity to make	20 fluidounces	1000 Cc.

Dissolve the Pepsin in the liquids, previously mixed.

149. VINUM OLEI MORRHUÆ.

Wine of Cod Liver Oil.

Morrhual (Gaduol)	80 grains	4.6 Gm.
Fluid Extract of Licorice	3 fluidounces	75 Cc.
Glycerin	2 fluidounces	50 Cc.
Syrup of Wild Cherry	4 fluidounces	100 Cc.
Liquid Extract of Malt	8 fluidounces	200 Cc.
Compound Syrup of Hypophosphites	4 fluidounces	100 Cc.
Fuller's Earth, in powder	240 grains	15 Gm.
Sherry Wine, sufficient to make	40 fluidounces	1000 Cc.

Mix the Morrhual with the Glycerin and triturate with the Fuller's Earth, add the Fluid Extracts and Syrup of Wild Cherry, allow it to stand for 24 hours, agitating occasionally, then filter and add the Syrup of Hypophosphites; lastly add sufficient Sherry Wine to make 40 fluidounces (1000 Cc.).

ERRATA

- PAGE 6, Formula 11—Read “Mix the *Phenacetine*, Acetanilide” etc.
- 9, Formula 21—Read “Elixir of Cinchona and Iron.”
- 14, Formula 35—Insert “*Compositum*” after Strychnina and read “*Compound*” Elixir of Cinchona and Iron.”
- 20, Formula 52—Read “Alcohol, 95%”
- 29, Formula 79—Read “Alcohol, 95%”
- Formula 81—Read in title, “*Hypophosphites*” for Hypophosphites.
- 31, Formula 86—After “Liquor” read “*Olei*”
- 32, Formula 89—In last line but one read “*Lime Water*” for “Water”
- 40, Formula 117—Read in title “*Phosphate*” for Phosphates.

Acetic Turpenti
 Acidum Hypop
 Adjuvant Elixir
 Alcohol Deodor
 Alkaline Antise
 Tincture of
 Allspice Water
 Aniseed Cordia
 Water . . .
 Anti-Asthmatic
 Antiseptic Pow
 Soap, Solu
 Solution . .
 Aqua Olei Aneth
 Anisi .
 Carui .
 Cinnam
 Fœnic
 Menth
 Menth
 Rosæ .
 Artificial Carls
 Aromatic Casto
 Elixir
 Fluid Extr
 Syrup of L
 Syrup of Bl

Calamine Lotio
 Camphorated C
 Chloroform
 Phenol Oin
 Capsulæ Apiole
 Colchicinæ
 cylatis . .
 Capsules of Apio
 Salicylates
 and Meth
 Carlsbad Salt, /
 Carminative Tir
 Carraway Water
 Cataplasma Ka
 Cataplasm of K

INDEX.

A

	PAGE		PAGE
Acetic Turpentine Liniment	24	Ceratum Galeni	5
Acidum Hypophosphorosum	3	Chemical Food, Parrish's	40
Adjuvant Elixir	6	Churchill's Tincture of Iodine	47
Alcohol Deodoratum	3	Chloral Camphoratum	5
Alkaline Antiseptic Solution	26	Chlorine Gargle	21
Tincture of Ignatia	46	Chloroformum Camphoratum	5
Allspice Water	4	Cinnamon Water	4
Aniseed Cordial	6	Cold Cream	5
Water	4	Collodium Iodoformatum	5
Anti-Asthmatic Elixir	9	Compound Acacia Powder	34
Antiseptic Powder, Soluble	35	Benzoate Powder	35
Soap, Solution of	31	Digestive Elixir	9
Solution	26	Elixir of Acetanilide	6
Aqua Olei Anethi	4	Buchu and Hyos-	
Anisi	4	cyamus	8
Carui	4	Cramp Bark	17
Cinnamomi	4	Digitalin	9
Fœniculi	4	Euphorbia	9
Menthæ Piperitæ	4	Glycerophosphates	11
Menthæ Viridis	4	" with Quinine	
Rosæ	3	and Strychnine	11
Artificial Carlsbad Salt	37	Pepsin	13
Aromatic Castor Oil	33	" and Bismuth	13
Elixir	7	" with Quinine,	
Fluid Extract of Cascara	20	Iron and Strych-	
Syrup of Licorice	41	nine	14
Syrup of Blackberry	44	Quinine	8

C

Calamine Lotion	32		
Camphorated Chloral	5	Essence of Vanillin	20
Chloroform	5	Glycerin of Heroin	23
Phenol Ointment	49	Iodine Paint	34
Capsulæ Apiol et Ergotini	4	Menthol Liniment	25
Colchicinæ et Methyl Salicy-		Milk Powder	35
latis	4	Powder of Acetanilide	34
Capsules of Apiol and Ergotin	4	Pepsin	36
Salicylates of Colchicine		Santonin	37
and Methyl	4	Compound Ointment of	
Carlsbad Salt, Artificial	37	Carbolic Acid	48
Carminative Tincture	46	Grey Sulphur	50
Carraway Water	4	Ichthyol	48
Cataplasma Kaolini	4	Resorcin	49
Cataplasm of Kaolin	4	Sulphur and	
		Birch Tar	50

	PAGE		PAGE
Compound Solution of Borax	27	Elixir Ferri Quininae et Strychninae	10
Solution of Bromo-Chloral	27	Formini	10
Hypophosphites without sugar	29	Glycerophosphatum Compositum	11
Sandal Oil	31	Glycerophosphatum cum Quinina et Strychnina	11
Spirit of Asparagus	38	Glycyrrhizae	11
Orange	38	Lithii et Hydrangeae	12
Syrup of Eucalyptus	39	Lithii Salicyatis	12
Hyphosphites	42	Papaini	12
Phosphate of Iron	40	Pepsini Compositum	13
Red Clover Blossoms	45	cum Bismutho	13
Sarsaparilla	44	cum Quinina et Ferro et Strychnina	14
Trifolium	45	Potassii Bromidi	15
Tincture of Cudbear	47	Quinque Bromidorum	7
Vanillin	20	Rhei et Magnesii Acetatis	15
D		Serenozæ Compositum	15
Decolorized Tincture of Iodine	47	Sex Iodorum	16
Deodorized Alcohol	3	Sodii Salicylatis Compositum	16
Diachylon Ointment	49	Terpini Hydratis et Codeinæ	16
Digestive Powder	36	Terpini Hydratis et Heroinæ	17
Dill Water	4	Viburni Compositum	17
Dilute Solution of Iodine	30	Elixir of Ammonium Bromide	6
Dithymol Diiodide	45	Calisaya	8
Dobell's Solution	27	Cinchona	8
E		Cinchona from Alkaloids	8
Elixir Acetanilidi Compositum	6	Cinchona and Iron	8
Adjuvans	6	Digestive Ferments	13
Ammonii Bromidi	6	Five Bromides	7
Anisi	6	Formin	10
Aromaticum	7	Glycerophates with Quinine and Strychnine	8
Aurantii	7	Hexamethylenetetramine	10
Buchu et Hyoscyami Compositum	8	Iron Pyrophosphate with Quinine and Strychnine	10
Calcii et Sodii Glycerophosphatis	8	Lactated Pepsin	13
Cinchonæ	8	Lactated Pepsin with Bismuth	13
Cinchonæ et Ferri	9	Elixir Lactated	13
Digestivum Compositum	9	Quinine, Iron	13
Digitalini Compositum	9	Licorice	13
Euphorbiæ Compositum	9	Lithium and Lithium Sal	13
Ferri Pyrophosphatis cum Quinina et Strychnina	10	Papain	13
		Potassium Rhubarb and Acetate	13
		Six Iodide	13
		Terpin	13
		Codeine	13
		Terpin	13
		Heroin	13
		Emulsio Iodof	13
		Olei Mori	13
		Olei Mori Phosph	13
		Olei Mori sino	13
		Emulsion of	13
		Cod Liver	13
		Cod Liver phosphate	13
		Iodoform	13
		Errata	13
		Essentia Lin	13
		Pepsini	13
		Pepsini	13
		Vanillin	13
		Essence of	13
		Pepsin	13
		Pepsin	13
		Extractum	13
		Cascar	13
		mati	13
		Senega	13
		Fennel Wa	13
		Ferrated E	13
		Flavours f	13
		Fluid Extr	13
		Fluid Extr	13
		Fermented	13
		Galen's Co	13
		Gargarism	13

PAGE	PAGE
Elixir Lactated Pepsin with	Gargle, Chlorine 21
Quinine, Iron and Strych-	Glycerinum Belladonnæ 21
nine 13	Bismuthi 22
Licorice 11	Ferri Iodidi 22
Lithium and Hydrangea. 12	Glycerinum Ferri Phosphatis
Lithium Salicylate 12	cum Quinina et Strychnina 22
Papain 12	Heroini Compositum 23
Potassium Bromide 15	Iodi 23
Rhubarb and Magnesium	Glycerin of Belladonna 21
Acetate 15	Bismuth 22
Six Iodides 16	Ferrous Iodide 22
Terpin Hydrate and	Iodine 23
Codeine 16	Iron Phosphate with
Terpin Hydrate and	Quinine and Strychnine 22
Heroin 17	Triple Phosphates 22
Emulsio Iodoformi 17	<i>Gouttes Ameres de Baume</i> 46
Olei Morrhuæ 17	H
Olei Morrhuæ cum Ferri	Hebra's Itch Ointment 50
Phosphato 18	Hiera Picra 35
Olei Morrhuæ cum Pep-	Humanized Milk 24
sino 18	Humanizing Milk Powder 35
Emulsion of Cod Liver Oil 17	Hypophosphorus Acid 3
Cod Liver Oil with Pepsin	I
Cod Liver Oil with Phos-	Infusum Buchu Concentratum 23
phate of Iron 18	Iodine and Starch Paste 33
Iodoform 17	Iodoform Collodium 5
Errata 52	Itch Ointment (Hebra's) 50
Essentia Limonis 19	K
Pepsini 19	Kumyss 24
Pepsini Phenolata 19	Kaolin Cataplasm 4
Vanillini Composita 20	L
Essence of Lemon 19	Lac Fermentatum 24
Pepsin 19	Lac Humanisatum 24
Pepsin Phenolated 19	Linimentum Album 24
Extractum Buchu Fluidum 20	Ammonii Iodidi 25
Cascaræ Sagradæ Aro-	Mentholis 25
maticum Fluidum 20	Mentholis Compositum 25
Senegæ Fluidum 21	Liniment of Ammonium Iodide 25
F	Liquor Ammonii Valerianatis. 25
Fennel Water 4	Antisepticus 26
Ferrated Elixir of Cinchona. 9	Antisepticus Alkalinus 26
Flavours for Emulsions 37	Auri et Arsenii Bromidi. 27
Fluid Extract of Buchu 20	Boracis Compositus 27
Fluid Extract of Senega 21	Bromo-Chloral Composi-
Fermented Milk 24	tus 27
G	Carmini 28
Galen's Cerate 5	Creosoti et Iodi 28
Gargarisma Chlori 21	

	PAGE		PAGE
Liquor Cresolis	28	Pulvis Pepsini Compositus . .	36
Ferri et Mangani Pepton-		Pro Mistura Cretæ . .	36
atis	29	Santonini Compositus. .	37
Glusidi	29		
Hypophosphitum Com-		R	
positum sine Saccharo	29	Rose Water	3
Iodi Dilutus	30		
Opii Sedativus	30	S	
Potassii Citratis	30	Saccharated Pepsin	34
Saponis Antisepticus . .	31	Sal Carolinum Factitium . . .	37
Olei Santali Flavi Com-		Sapores Pro Emulsionibus . .	37
positus	31	Sedative Liquid	30
Sodii Hydroxidi	31	Sedative Solution of Opium .	30
Zingiberis	32	Simple Elixir	7
M		Skeenes' Mixture	35
Magma Magnesia	32	Soluble Antiseptic Powder . .	35
Magnesia Magma	32	Soluble Tincture of Tolu . . .	48
Mandl's Solution	34	Solution of Ammonium Valer-	
Menthol Liniment	25	ianate	25
Milk of Magnesia	32	Carmine	28
Mistura Butyl-Chloral	33	Cresol	28
Mixture of Butyl-Chloral . . .	33	Creosote and Iodine . . .	28
Mixture, Skeenes'	35	Ginger	32
O		Gluside	29
Oleum Ricini Aromaticum . . .	33	Iron and Manganese	
Ointment of Lead Plaster . .	49	Peptonate	29
Zinc Stearate	51	Iodine (dilute)	30
P		Peptonate of Iron	
Parrish's Chemical Food . . .	40	and Manganese	29
Pasta Iodi et Amyli	33	Potassium Citrate . . .	30
Peppermint Water	4	Saccharin	29
Pepsinum Saccharatum	34	Soda	31
Peptonizing Powder	36	Sodium Hydroxide . . .	31
Phenolated Essence of Pepsin	19	Mandl's	34
Phosphatic Emulsion	18	Spearmint Water	4
Pigmentum Iodi Compositum	34	Spiritus Asparagi Compositus.	38
Powder for Chalk Mixture . .	36	Amygdalæ Amaræ	38
Powder of Aloes and Canella.	35	Aurantii	38
Preface	1	Aurantii Compositus . . .	38
Pulvis Acaciæ Compositus . . .	34	Spirit of Bitter Almond . . .	38
Acetanilidi Compositus	34	Orange	38
Aloes et Canellæ	35	Stainless Iodine Ointment . .	49
Antisepticus Solubilis . .	35	Stokes' Liniment	24
Benzoatis Compositus . . .	35	Sulphurated Lime Lotion . . .	32
Digestivus	36	Sweet Castor Oil	33
Lactis Compositus	35	Syrupus Acaciæ	39
Pancreaticus Composi-		Codeinæ Phosphatis	39
tus	36	Eucalypti Compositus . . .	39
		Ferri et Mangani Iodidi .	40

F
C
F
I
Syrup
C
F
S
S
I
Z
Syrup

Tastel
Thym
Thym
Tinctu

	PAGE
Ferri Phosphatis Compositus	40
Glycyrrhizæ Aromaticus.	41
Hypophosphitum Compositus	42
Lotio Calaminæ.....	32
Calcis Sulphuratæ....	32
Syrupus Picis Liquidæ.....	43
Quininæ Phospho-Muriatis	43
Rubi Aromaticus	44
Sarsæ Compositus.....	44
Senegæ	44
Trifolii Compositus.....	45
Zingiberis.....	45
Syrup of Acacia.....	39
Codeine Phosphate.	39
Iodide of Iron and Manganese.....	40
Tar	43
Phospho-Muriate of Quinine.....	43
Senega	44
Ginger.....	45

T

Tasteless Tincture of Iron ..	46
Thymolis Iodidum.....	45
Thymol Iodide	45
Tinctura Aurantii Corticis	
Dulcis Recentis....	46
Carminativa	46
Ferri Citro-Chloridi.	46
Ignatiæ Alkalina ..	46
Iodi (Churchill)	47
Iodi Decolorata....	47

	PAGE
Tinctura Persionis.....	47
Personis Composita.	47
Tolutana Solubilis..	48
Tincture of Citro-Chloride of	
Iron	46
Cudbear	47
Fresh Sweet Orange Peel.....	46

U

Unguentum	
Acidi Carbolici Compositum	48
Ichthyolis Compositum..	48
Iodi Denigrescens.....	49
Emplastri Plumbi	49
Phenolis Camphoratum.	49
Resorcini Compositum..	48
Sulphuris Compositum...	50
Sulphuris et Rusci Compositum	50
Sulphuris Cinerei Compositum	50
Zinci Stearatis	51

V

Vleminc's Solution.....	32
Vinum Cocæ	51
Pepsini.....	51
Olei Morrhuæ	51

W

White Liniment.....	24
Wilkinson's Ointment.....	50
Wine of Coca	51
Pepsin	51
Cod Liver Oil.....	51

Chlor 1:19.