





IMAGE EVALUATION TEST TARGET (MT-3)



APPLIED IMAGE, Inc
 1653 East Main Street
 Rochester, NY 14609 USA
 Phone: 716/482-0300
 Fax: 716/288-5989

© 1993, Applied Image, Inc., All Rights Reserved

**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1993

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

Coloured covers/
Couverture de couleur

Coloured pages/
Pages de couleur

Covers damaged/
Couverture endommagée

Pages damaged/
Pages endommagées

Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée

Pages restored and/or laminated/
Pages restaurées et/ou pelliculées

Cover title missing/
Le titre de couverture manque

Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées

Coloured maps/
Cartes géographiques en couleur

Pages detached/
Pages détachées

Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)

Showthrough/
Transparence

Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur

Quality of print varies/
Qualité inégale de l'impression

Bound with other material/
Relié avec d'autres documents

Continuous pagination/
Pagination continue

Tight binding may cause shadows or distortion along interior margin/
La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure

Includes index(es)/
Comprend un (des) index

Title on header taken from: /
Le titre de l'en-tête provient:

Blank leaves added during restoration may appear within the text. Whenever possible, these have been omitted from filming/
il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.

Title page of issue/
Page de titre de la livraison

Caption of issue/
Titre de départ de la livraison

Masthead/
Générique (périodiques) de la livraison

Additional comments: /
Commentaires supplémentaires:

This item is filmed at the reduction ratio checked below /
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	12X	14X	16X	18X	20X	22X	24X	26X	28X	30X	32X
<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>									

The copy filmed here has been reproduced thanks to the generosity of:



Library
Agriculture Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol \rightarrow (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

Bibliothèque
Agriculture Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole \rightarrow signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

ONTARIO AGRICULTURAL COLLEGE

EXPERIMENT STATION

BULLETIN LXXII

ROOTS, POTATOES AND FODDER CORN.

BY THOMAS SHAW, PROFESSOR OF AGRICULTURE, AND
C. A. ZAVITZ, B. S., EXPERIMENTALIST.

PUBLISHED BY THE DEPARTMENT OF AGRICULTURE
February 29, 1892

TORONTO

PRINTED BY WARWICK & SONS

MINISTER OF AGRICULTURE

HON. JOHN DRYDEN, TORONTO.

Ontario Agricultural College and Experimental Farm, Guelph,
under control of the Minister of Agriculture.



JAMES MILLS, M.A. President.
THOMAS SHAW Professor of Agriculture and Farm Superintendent.
A. E. SHUTTLEWORTH, B.A. Sc. Professor of Chemistry.
J. HOTES PANTON, M.A., F.G.S. Professor of Natural History and Geology.
F. C. GREENSIDE, V.S. Professor of Veterinary Science.
H. H. DEAN, B.S.A. Professor of Dairy Husbandry.
E. LAWRENCE HUNT, B.A. Assistant Resident and Mathematical Master.
CAPTAIN WALTER CLARKE Instructor in Drill and Gymnastics.
C. A. ZAVITS, B.S.A. Experimentalist.
H. B. SHARMAN, B.S.A. Assistant Chemist.
A. MCCALLUM, Bursar.

ADVISORY BOARD.

C. C. JAMES, M.A., *Secretary* Deputy Minister of Agriculture, Toronto.
JOHN I. HOBSON, *Chairman* Mosborough, County of Wellington.

BULLETIN LXXII.

ROOTS. POTATOES AND FODDER CORN.

Experience has taught our farmers that there is a tendency with all kinds of grain to deteriorate when grown for several years in the same locality, hence the necessity for the introduction of new varieties from time to time. The same holds true of field roots, potatoes and corn, hence the necessity will always exist for conducting experiments which will enable us to determine which of these varieties are worthy of the attention of the agriculturist.

The following include some of the principal objects of this bulletin, viz. :—(1) To give the comparative results obtained from growing leading varieties of roots, potatoes and fodder corn side by side, and under the same conditions, for one year, and in the case of some of them for a longer term ; and (2) to give the comparative results from growing roots (a) upon ridges as against flat cultivation, and (b) upon rows at different distances apart.

In 1891, 57 varieties of turnips, 36 of mangels and 4 of sugar beets were grown at this station and upon plots side by side. Each plot consisted of two drills 6 rods in length. The soil was a clay loam of mild texture. It had produced grain crops for a number of years previously, was manured with a fair dressing of stable manure in the spring of 1890, and produced a crop of rape the same autumn which was pastured off on the land. The seeds of nearly all the varieties of roots tested were obtained from seedsmen in Ontario and the United States, but a few were also obtained from England.

Swede Turnips. 42 varieties were grown. The drills were 27.2 inches apart. In one drill the plants were thinned to a distance of 12 inches and in the other to 16 inches. They were sown June 19th. The following table gives the comparative results from the 6 varieties which gave the highest yields :

Varieties.	Description of roots.		Average weight per root of two seedings.	Average yield per acre.
	Shape.	Firmness of flesh.		
Hartley's Bronze top.....	long ..	good	lb. 2.91	tons. 23.93
Marshall's Purple top.....	globe...	good	2.51	23.77
Sutton's Champion	globe...	medium..	2.95	23.54
Hazard's Swede turnip.....	long ..	good	2.91	23.12
Our selected Purple top.....	long ..	poor	2.89	22.95
Skirving's Swede	globe...	good	2.76	22.73

The following varieties have given the best yields for two years, viz: Royal Norfolk Purple Top, 21.15 tons per acre, Hazard's Swede 20.92, and Carter's Elephant 19.87 tons. The average yield of all the Swedish varieties grown in 1891 was 19.7 tons.

White Fleshed and Yellow Fleshed Turnips. 15 varieties were grown. The drills were 27.2 inches apart. In one drill the plants were thinned to 16 inches and in the other to 13 inches in the row. They were sown June 19th. These varieties are all white fleshed. The comparative results from the six varieties which gave the highest yields are shown below.

Varieties.	Shape of roots.	Average weight per root of two seedings.	Average yield per acre.
		lb.	tons.
Grey Stone Improved	4.29	29.28
Red Globe Norfolk.....	globe.....	3.67	28.69
Jersey Navet.....	".....	3.74	28.23
Grey Stone.....	".....	3.56	28.06
Purple Top Main.....	slightly flat.....	3.48	28.09
Red Top Strap Leaf.....	flat.....	3.15	25.97

The following varieties have given the best yields for two years, viz: Red Globe Norfolk, 27.6 tons per acre, Red Top Strap Leaf, 23.3 tons, and Pomeranian White Globe, 22 tons. These also are all white fleshed varieties. The 11 white fleshed varieties averaged 25.6 tons per acre, and the four yellow fleshed 20.22 tons.

Mangels. 36 varieties were grown. The drills were 22½ inches apart. In one drill the plants were thinned to 7 inches and in the other to 14 inches. They were sown May 23rd. The next table gives the comparative results from the six varieties which gave the highest yields.

Varieties.	Shape.	Uniformity.	Average weight per root of two seedings.	Average yield per acre.
			lb.	tons.
Improved Mammoth Prize, long red	long.....	even.....	2.55	35.56
Carter's Champion.....	half long.....	".....	2.77	34.09
Mammoth Sawlog.....	long.....	".....	2.68	34.00
Steele Bros. Long Red Selected....	".....	medium..	2.61	33.33
Norbiban Giant.....	".....	".....	2.59	33.07
Eiffel Tower.....	half long.....	".....	2.46	32.04

It
variety
The 1
per a
follow
viz: 1
Ward

Sug
apart.
row, t

It w
been g
list in
19.86

Pot
the tw
the ma
taining
years,
in the
180 set
ft. long
Resu

Empir
Summi
Early
Foot
Thorbo

It will be observed that these do not comprise any of the globe varieties. The highest yielding globe variety came 18th in the list. The 14 distinctively long varieties yielded an average of 28.57 tons per acre and the six globe varieties an average of 22.42 tons. The following varieties have given the best yields per acre for two years, viz: Norbitan Giant 22.9 tons, Long Red 19.3 tons and Carter's Warden Orange 18.3 tons.

Sugar Beets. 4 varieties were grown. The drills were 22½ inches apart. In one drill the plants were thinned to 6 inches in the row, and in the other to 12 inches. They were sown May 23rd.

Varieties.	Average weight per root.	Yield per acre.
	lb.	tons.
Vilmorin's Improved White..	2.02	21.90
German White	1.84	20.71
Silesian	1.53	20.59
French White	1.42	18.87

It will be observed that the German White, the kind which has been grown most extensively over the province, comes second on the list in point of yield. The average yield of the four varieties is 19.86 tons per acre.

Potatoes. 74 varieties were grown in 1891, and 25 varieties in the two years past. The seed was obtained from the United States, the maritime provinces and Ontario. The soil was a clay loam containing a large amount of vegetable matter. It lay in sod for five years, was turned over last autumn and was thoroughly cultivated in the spring. One peck of the tubers of each variety was cut into 180 sets and planted in drills 28½ inches apart. The rows were 18½ ft. long.

Results from the six varieties which gave the highest yields :

Varieties.	Maturity. 1891.	Uniformity of tubers. 1891.	Average yield per acre. 1890-91.
Empire State	medium	good	bush. 192.9
Summit	late	good	147.5
Early Maine	early	medium	140.9
Clark No. 1	medium	good	138.0
Pootaluck	medium	good	130.3
Thorburn	medium	good	129.2

By actual test we have found all these varieties to possess good edible qualities with the exception of the Pootaluck, which probably should not be classed higher than medium.

Fodder Corn. 84 varieties were grown in 1891. The seed was obtained chiefly from the northern States, but some of it came from the southern States and some was obtained from Ontario. All these varieties were grown in duplicate in the first instance upon sod which had not been plowed for six years, and in the second on land which had grown different kinds of grain for four years previously. The sod was turned under in the autumn of 1890. No manure was applied nor had it received any since 1886. The stubble land was also plowed the previous autumn, and had received a fair application of farmyard manure in the spring of 1890. The soil was a mild clay loam and rather low lying. The corn was planted in hills, 5 links or 39.6 inches apart each way. But one row of each variety was planted on each kind of soil. The rows were 165 ft. long on the sod, and 99 ft. long on the stubble ground. But four plants were left in each hill. Thorough shallow cultivation was given the rows both ways. Following are the results:

Varieties.	Class of corn.	Maturity.	Weight of green ears per acre, 1891.	Average yield per acre of the whole crop, 1889-90-91.
			lb.	tons.
Sheep's Tooth	White dent ..	Late	1,836	16.63
Chester County Mammoth..	Yellow dent ..	Late	1,800	16.41
Calico Dent	Mixed dent ..	Medium	4,333	15.35
Leaming Dent	Yellow dent ..	Medium late..	3,866	15.33
Oranberry White Dent.....	White dent ..	Late	900	15.10
Compton's Early	Yellow flint..	Early	6,000	15.03
Wisconsin Yellow Dent	Yellow dent ..	Medium late..	3,166	14.94
Mammoth Southern Sweet..	White dent ..	Late	1,060	14.86
South Western	White dent ..	Late	1,700	14.92
Wisconsin White Flint	White flint..	Medium	5,735	14.55

It should be remembered that these varieties were grown for fodder purposes whether in the green or dried state or cured in the silo. Eight of the ten varieties were dent and two were flint. It has been found somewhat difficult hitherto to get flint-varieties which produce sufficient weights per acre, but this experiment shows that the two flint varieties named in the table weigh well and also produce a large quantity of grain. Of the other varieties the most suitable probably for the silo in temperatures the same as at this station are Compton's Early, Wisconsin White Flint, Leaming Yellow, and Wisconsin Yellow Dent.

The
ing S
again
ayers

Flat of
Ridge

It w
obtain
marked

The
Swede
28 inch
inches
inches.
the tur

Distan

16 inch
24 inch
28 inch

It wil
them ga
be requi
with saf

The fo
experime

1. The
white-fl
yellow-f

The next table gives the comparative results obtained from growing Swede turnips, mangels, and sugar beets upon ridged drills as against flat cultivation. Each result mentioned in the table is the average of triplicate tests.

Modes of cultivation.	Average yield per acre (tons).		
	Swede turnips.	Mangels.	Sugar beets.
Flat cultivation	26.06	21.55	13.03
Ridged cultivation	25.83	19.53	12.91

It will be observed that the largest yield in each instance was obtained from the roots grown upon the level surface, and it is most marked in the case of the mangels.

The next table shows the comparative average yields per acre of Swede turnips, mangels and sugar beets grown in rows, 16, 24 and 28 inches apart. The turnips were thinned to the distance of 14 inches in the rows, the mangels 14 inches, and the sugar beets 7 inches. The mangels and sugar beets were sown on May 20th, and the turnips on June 22nd.

Distance between rows.	Average yield per acre (tons).		
	Swede turnips.	Mangels.	Sugar beets.
16 inches.....	27.39	24.05	14.03
24 inches.....	26.66	21.15	13.36
28 inches.....	23.78	16.42	11.51

It will be observed that those with the shortest distances between them gave the highest yields in every instance, but further tests will be required before conclusions can be drawn which can be followed with safety.

Conclusions.

The following conclusions may be safely drawn from the different experiments summarised in the Bulletin :

1. That while the average yield of the 11 varieties of white-fleshed turnips per acre was 853 bushels, that of the 4 yellow-fleshed varieties was but 673.8 bushels.

2. That while the average yield of the 14 distinctively long varieties of mangels was 952 bushels per acre, that of the 6 globe varieties was but 747 bushels.

3. That the best yielding varieties of potatoes tested at this station for the past two years are also possessed of good edible qualities.

4. That for quantity and quality of fodder and earliness in maturing the Compton's Early, Wisconsin White Flint, Leaming Yellow and Wisconsin Yellow Dent are probably the most suitable for the silo among the 35 varieties tested here for the past three years.

5. That in this test the mangels grown on the level gave 10.4 per cent. more of an increase than when grown on ridges, while the difference with the turnips and sugar beets was but slight.

Distribution of Seed. Parties desirous of obtaining seed of any of the leading varieties of field roots and corn grown at this station, for purposes of testing, should apply for circular, which contains all necessary information in reference to this department of the work of our Station.

