





**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**Photographic  
Sciences  
Corporation**

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503

**CIHM/ICMH  
Microfiche  
Series.**

**CIHM/ICMH  
Collection de  
microfiches.**



**Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques**

**© 1983**

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- |  |  |
|--|--|
| <input checked="" type="checkbox"/> Coloured covers/<br>Couverture de couleur  | <input type="checkbox"/> Coloured pages/<br>Pages de couleur   |
| <input type="checkbox"/> Covers damaged/<br>Couverture endommagée  | <input type="checkbox"/> Pages damaged/<br>Pages endommagées   |
| <input type="checkbox"/> Covers restored and/or laminated/<br>Couverture restaurée et/ou pelliculée  | <input type="checkbox"/> Pages restored and/or laminated/<br>Pages restaurées et/ou pelliculées  |
| <input type="checkbox"/> Cover title missing/<br>Le titre de couverture manque   | <input type="checkbox"/> Pages discoloured, stained or foxed/<br>Pages décolorées, tachetées ou piquées  |
| <input type="checkbox"/> Coloured maps/<br>Cartes géographiques en couleur   | <input type="checkbox"/> Pages detached/<br>Pages détachées  |
| <input type="checkbox"/> Coloured ink (i.e. other than blue or black)/<br>Encre de couleur (i.e. autre que bleue ou noire)   | <input type="checkbox"/> Showthrough/<br>Transparence  |
| <input type="checkbox"/> Coloured plates and/or illustrations/<br>Planches et/ou illustrations en couleur  | <input type="checkbox"/> Quality of print varies/<br>Qualité inégale de l'impression   |
| <input type="checkbox"/> Bound with other material/<br>Relié avec d'autres documents   | <input type="checkbox"/> Includes supplementary material/<br>Comprend du matériel supplémentaire   |
| <input type="checkbox"/> Tight binding may cause shadows or distortion<br>along interior margin/<br>La reliure serrée peut causer de l'ombre ou de la<br>distortion le long de la marge intérieure   | <input type="checkbox"/> Only edition available/<br>Seule édition disponible   |
| <input type="checkbox"/> Blank leaves added during restoration may<br>appear within the text. Whenever possible, these<br>have been omitted from filming/<br>Il se peut que certaines pages blanches ajoutées<br>lors d'une restauration apparaissent dans le texte,<br>mais, lorsque cela était possible, ces pages n'ont<br>pas été filmées. | <input type="checkbox"/> Pages wholly or partially obscured by errata<br>slips, tissues, etc., have been refilmed to<br>ensure the best possible image/<br>Les pages totalement ou partiellement<br>obscurcies par un feuillet d'errata, une pelure,<br>etc., ont été filmées à nouveau de façon à<br>obtenir la meilleure image possible. |
| <input type="checkbox"/> Additional comments:/<br>Commentaires supplémentaires:  |  |

This item is filmed at the reduction ratio checked below/  
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
12X	16X	20X	24X	28X	32X

The copy filmed here has been reproduced thanks to the generosity of:




Library of Congress  
Photoduplication Service

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol  $\rightarrow$  (meaning "CONTINUED"), or the symbol  $\nabla$  (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:


Library of Congress  
Photoduplication Service

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole  $\rightarrow$  signifie "A SUIVRE", le symbole  $\nabla$  signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.



Pl 130  
U6 W65

---

NOTES  
ON  
RECENT CANADIAN UNIONIDÆ.

BY J. F. WHITEAVES.

*Reprinted from the Canadian Record of Science, January, 1895.*

---



328 AK

3528 A.K.

"Reprinted from the Canadian Record of Science, January, 1895."

### NOTES ON RECENT CANADIAN UNIONIDÆ.

By J. F. WHITEAVES.<sup>1</sup>

The present paper is intended as a contribution to our knowledge of the geographical distribution of the Unionidæ in North America. It consists of a list of all the species from Canadian localities that are now represented in the museum of the Geological Survey at Ottawa, and is based almost exclusively upon specimens that were either collected by members of the Survey staff or presented by friends interested in its museum. So far as the writer is aware, however, the *Unio tenuissimus* of Lea, which was collected by Dr. G. M. Dawson in 1873, in the Souris River, Manitoba, is the only species of Unionidæ known to occur in Canada that is not represented in the Survey museum. Specimens of most of the nominal species of *Anodonta* and of a few of the more difficult species of *Unio* enumerated in this list have been kindly compared by Mr. Charles T. Simpson, of the United States National Museum, with Dr. Lea's types of North American Unionidæ now preserved in that institution, and identified as correctly as the small number of shells sent from each locality and the incompleteness of his studies of the family would permit. The nomenclature employed throughout this list is that which is now in general use among students of this group in North America, as it is still quite uncertain which of the earlier names of Rafinesque,

<sup>1</sup> Communicated by permission of the Director of the Geological Survey Department.



Lamarck and others, will ultimately have to be retained for some of these shells.

ANODONTA, Lamarck, 1879.

ANODONTA BENEDICTII, Lea.

Specimens which appear to have been identified with this species by Dr. Lea have already been recorded by Dr. R. Bell<sup>1</sup> as having been collected by himself, in 1860, at Batch-ah-wah-nah Bay, Lake Superior; in the St. Mary River, near Sugar Island, and on the north shore of Lake Huron, at Lacloche Island. Professor Macoun has recently (1894) collected it at Rondeau, near Point aux Pins, on the Ontario side of Lake Erie, and a few specimens, which Mr. Simpson thinks are probably referable to *A. Benedictii*, were collected by Dr. R. Bell, in 1883, at Lake Winnipeg, between Fort Alexander and Elk Island. Mr. Simpson is inclined to believe that *A. Benedictii* may be only a variety of *A. ovata*, Lea.

ANODONTA DECORA, Lea.

Eight full grown specimens and one immature shell of a very large *Anodonta*, which Mr. Simpson refers to *A. decora*, were collected by Mr. Law, of Chatham, at Rondeau, Ontario, and presented by him to the Museum of the Survey, through Professor Macoun, in 1884. One of the adult shells from this locality, a fairly average specimen, measures 6.6 inches in length, 4 inches in height and 3.1 inches in breadth or thickness. The umbones of each are remarkably ventricose and prominent. The test is rather thick, the hinge line short, and the cardinal angles are rounded in front and obtusely angular behind. The writer has long been under the impression that these shells could be identified with the typical form of *A. grandis*, Say, as they do not correspond at all well with Lea's figures or measurements of *A. decora*,

<sup>1</sup> In Canad. Nat. and Geol., Vol. VI., p. 269.

the "breadth" or, as it would now be called, the length of which is stated to be 3.9 inches. The recent receipt from Mr. Simpson of outline drawings of specimens from Dr. Lea's collections, labelled "*A. decora*, from the canal at Cincinnati, Ohio," has, however, convinced the writer of the correctness of Mr. Simpson's determination, though it is very generally believed that *A. decora* is not more than a mere variety of *A. grandis*.

ANODONTA EDENTULA, Say. (*A. undulata*, Lea, et auct., but possibly not of Say; *A. Pennsylvanica*, Lamark, and *A. areolata*, Swainson.)

Dr. R. Ellsworth Call has expressed the opinion that *A. edentula*, Say, is peculiar to the Mississippi drainage system, and *A. undulata*, Say., to those waters that drain into the Atlantic, but the writer has never been able to see any tangible difference between these two shells. In a recent letter to the writer, Mr. Simpson says, "*Anodonta undulata* is no doubt the small form which we have here in the Potomac. Though Say gives no locality, he speaks of it as 'thin and fragile, length near half an inch; breadth seven-tenths.' The figure fairly well represents our shell. This may run into *A. edentula*, but I have never yet been able to connect it with that. The material in Lea's collection, under the name of *A. undulata*, Say, is merely a form or forms of *A. edentula*."

Under one or the other of these names this shell has previously been recorded as having been collected in Lake Matapedia, P.Q., by Dr. R. Bell in 1857; in a small lake in the valley of the Riviere Rouge, P.Q., by W. S. M. D'Urban, in 1858; in the St. Charles River, near Quebec city, by the writer, in 1861, and at Brome Lake, P.Q., by Mr. R. J. Fowler, in 1862.

More recently, it has been collected by Dr. R. Bell in 1883, at Lake Winnipeg, between Forts Alexander and Simpson, and by Professor Macoun, in 1894, in Ontario,

at Rondeau, on Lake Erie, and in the east and west branches of the Grand River at Galt and Ayr.

In another letter to the writer, Mr. Simpson makes the following remarks upon this species, "The so-called Anodontas of which this is the type, have more or less perfect cardinals and occasional vestiges of laterals. They group with *Margaritana Elliotti*, *M. Spillmani*, *M. Ravenliana*, etc. The genus *Margaritana* is a medley of forms, which, for the most part, are more nearly related to various groups of *Unio* than to each other. I believe that *Margaritana* should be merged into *Unio*, and with it the Anodontas of the *edentula* group."

ANODONTA FERUSSACIANA, Lea.

L'Orignal Creek, Ottawa River, Dr. R. Bell, 1855 (as *A. pavonia*, Lea). Ponds at the Mile End, Montreal, Dr. R. Bell, 1858, and J. F. Whiteaves, 1862.

ANODONTA FLUVIATILIS, Dillwyn. Sp. (*A. cataracta*, Say.)

Several specimens of this common eastern species, which has previously been recorded as occurring at many localities in the Province of Quebec and neighbourhood of Ottawa, were collected by Dr. R. Bell, in 1883, at Flying Post Route, 100 miles north-east of Michipicoten, and, in 1889, from a small lake near Proudfoot's north and south line, in the Sudbury district of Ontario. A single specimen, which may be referable to this species, was collected by Professor Macoun, in 1884, at White Fish River, north of Lake Superior.

ANODONTA FOOTIANA, Lea.

Specimens which are said to have been identified with this species by Dr. Lea were collected by Mr. W. M. S. D'Urban, in 1858, from three small lakes tributary to the Riviere Rouge, P.Q. Since then, specimens, which Mr. Simpson refers to *A. Footiana*, have been collected in

Ontario, by Professor Macoun, in 1884, at White Fish River, north of Lake Superior, and at Lake Hannah, on the Nepigon River; by Dr. A. C. Lawson, in 1886, at Rainy Lake; by Mr. W. Spreadborough, in 1894, from the Muskoka River, near Georgian Bay; and in Manitoba, by Dr. R. Bell, in 1883, at Shoal Lake, Red River. Mr. Simpson also is of opinion that specimens collected by Mr. R. J. Fowler in the Lachine Canal at Montreal, in 1863, and referred by the writer to *A. Lewisii*, Lea, are young shells of *A. Footiana*.

ANODONTA FRAGILIS, Lamark. (*A. lacustris*, Lea.)

This shell was apparently first collected in Canada by Mr. D'Urban in 1858, associated with *A. Footiana*, in three small lakes in the valley of the Riviere Rouge, and identified shortly afterwards by the late Dr. Isaac Lea with the *A. fragilis* of Lamarck. Specimens collected by Professor Macoun in 1885, from a lake six miles up the Becscie River, Anticosti, were identified with *A. fragilis* by Mr. F. R. Latchford, of Ottawa, and similar shells have long been known to occur at Meach's Lake, near Ottawa. Some of these Anticosti specimens were sent to Mr. Simpson, who thinks that they are essentially similar to shells labelled *A. fragilis* in Dr. Lea's collection, but cannot see how these latter are to be distinguished from *A. lacustris*, Lea, and does not pretend to be always able to separate *A. fragilis* from *A. fluviatilis*.

ANODONTA IMPLICATA, Say.

Lake Winnipeg, between Fort Alexander and Elk Island, Dr. R. Bell, 1883; and Souris River, near Roche Percée, Dr. A. R. C. Selwyn, 1890; a few specimens from each of these localities, which have been identified with this species by Mr. Simpson. It had previously been recorded as occurring in the St. Charles River, near Quebec, where it was collected by the writer in 1861.

## ANODONTA MARRYATTANA, Lea.

Lake Hannah, Nipigon River, and east side of Lake Nipigon, Ontario, Professor Macoun, 1884; and Fairford River, Manitoba, J. F. Whiteaves, 1888; as identified by Mr. Simpson.

ANODONTA NUTTALLIANA, Lea. (*A. Oregonensis*, Lea.)

Okanagan Lake, B.C., A. J. Hill, 1882; two specimens of the variety *Oregonensis*. Near Victoria, V. I., James Fletcher, 1885, and Rev. G. W. Taylor, 1889. Nicola Lake, B.C., Dr. G. M. Dawson, 1889; three specimens of the typical form and one of the variety *Oregonensis*. Salmon Arm, Shuswap Lake, B.C., Dr. Dawson, 1894; several examples of both forms of the species. Stream entering Clayoquot Sound, V. I., at Stubbs Island, W. Spreadborough, 1894.

## ANODONTA OVATA, Lea.

Coulée No. 5, Vermilion River, Alberta, J. B. Tyrrell, 1886.

## ANODONTA PEPINIANA, Lea.

Specimens which Mr. Simpson refers to this species were collected by Dr. R. Bell, in 1883, from the Winnipeg River, Manitoba, and in 1886, from the Attawapishkat River, in the Severn district, which now forms the eastern part of Keewatin. Two left valves of a shell which may be referable to this species were collected by Mr. J. B. Tyrrell, in 1884, at the Lake of the Woods. Mr. Simpson is of the opinion that *A. Pepiniana* may be merely a variety of *A. Simpsoniana*, Lea.

## ANODONTA SIMPSONIANA, Lea.

In Ontario this species was collected by Dr. A. R. C. Selwyn in 1883, at Black Bay, Lake Superior; by Prof. Macoun, in 1884, at the north end of Lake Nipigon, in

1885, at Port Dover, Lake Erie, and in 1890, at Port Colborne, on the same lake.

In Manitoba it was collected by Dr. R. Bell in 1878, at the outlet of Lake Winnipeg and from Lake Winnipeg between Fort Alexander and Elk Island. It occurs, associated with *A. Marrayattana*, Lea, in the Fairford River, and is the only species of *Anodonta* that the writer was able to find in Lake Manitoba (in 1888).

In the district of Saskatchewan one perfect specimen was collected by Dr. R. Bell, in 1882, at Buffalo Lake, near Methy Portage.

Mr. Simpson, to whom the writer is indebted for the identification of specimens from most of these localities, is convinced that *A. Dallasiana* and *A. Kennicotti*, of Lea, are both synonyms of *A. Simpsoniana*.

#### ANODONTA SUBCYLINDRACEA, Lea.

Widely distributed in the provinces of Quebec and Ontario, from Lakes Metapedia and St. John to the eastward, to creeks, rivers and bays at the east end of Lake Superior and north side of Lake Erie to the westward. Mr. Simpson, however, regards *A. subcylindracea* as a mere synonym of *A. Ferrussaciana*, Lea.

#### MARGARITANA, Schumacher, 1819.

##### MARGARITANA CALCEOLA, Lea. (*M. deltoidea*, Lea.)

Lake Erie, at Fort Dover, Professor Macoun, 1890.  
Grand River, at Belwood, Ontario, J. Townsend, 1892.  
East and west branches of the Grand River at Galt and Ayr, Professor Macoun, 1894.

##### MARGARITANA COMPLANATA, Barnes.

Manitoba. Upper Assiniboine River, Dr. R. Bell, 1874;  
Souris River, Dr. A. R. C. Selwyn, 1882 and 1884; Shoal River and near Elk Island, Lake Winnipeg, Dr. R. Bell,

1883; Swan River, J. B. Tyrrell, 1887, and Assiniboine River, J. B. Tyrrell, 1884.

Keewatin. Nelson River, Dr. R. Bell, 1878.

Saskatchewan. Shell River (township 50, range 2 and 3, west of third Initial Meridian) north of the north Saskatchewan, O. J. Klotz, 1890.

Alberta. Battle River, three miles above Grattan Lake, J. B. Tyrrell, 1885.

MARGARITANA MARGARITIFERA, L.

From the Province of Quebec this species has already been recorded as having been collected by Dr. R. Bell (in 1857) in the Green and Rimouski rivers, at Lake St. John and both the Metapedia Lakes, and by the writer, (in 1861) in the River St. Charles, near Quebec City. More recently it has been collected in that province by Dr. H. M. Ami, in 1883, in the Assumption River, near Rawdon; by N. J. Giroux, in 1892, at the Lac de la Ferme, Riviere du Loup, en haut, and in that river; also by A. P. Low, in 1894, in the Romaine River.

In British Columbia, small and thin but characteristic specimens were found by Dr. G. M. Dawson, in 1885, in small streams entering Malaspina Strait, on the mainland side; also, in 1890, in Kakwous Lake, the source of the Bonaparte River, at an altitude of about 4,000 feet.

MARGARITANA MARGINATA, Say.

The small and typical eastern form of this shell is common in the province of Quebec and in eastern Ontario. A few specimens of the large western variety known to students of the Unionidae as *M. truncata*, Say (M. S.) were collected by Professor Macoun, in 1894, at Galt and Ayr, from the east and west branches of the Grand River.

MARGARITANA RUGOSA, Barnes. (? = *M. costata*, Rafinesque, sp.)

This species is widely distributed in the provinces of

Quebec and Ontario. In the latter province unusually large and thick specimens, measuring five inches and a half in length by three inches in height, were collected by Prof. Macoun, in 1894, in the east and west branches of the Grand River, at Galt and Ayr. The species has been recorded by Dr. G. M. Dawson as occurring, though rarely, in the Roseau River, Manitoba.

MARGARITANA UNDULATA, Say.

St. Lawrence River, at Montreal and Quebec, J. F. Whiteaves, 1861. Near Ottawa City, G. C. Heron, 1879.

UNIO, Philipsson, 1788.

UNIO ALATUS, Say.

Widely distributed throughout Ontario. The most easterly locality at which it has been collected is the Ottawa River at L'Orignal, as recorded by Dr. R. Bell, in the Canadian Naturalist and Geologist for June, 1859 (Vol. IV., p. 219). In Manitoba it has been collected in the Red River by Dr. G. M. Dawson, in 1873, and by T. C. Weston, in 1884.

UNIO BOREALIS, A. F. Gray.

A pair of specimens of this species, from the Ottawa River, at Duck Island, the typical locality, was presented to the museum of the Survey by Mr. F. R. Latchford, of Ottawa, in 1886.

UNIO CANADENSIS, Lea.

Two specimens, from the Ottawa River, near Ottawa, which are believed by the donor to be referable to this enigmatical species, were presented to the Museum of the Survey by Mr. Latchford, in 1893.

UNIO CIRCULUS, Lea. (?= *U. subrotundus*, Rafinesque.)

Lake Erie, at Kingsville, Ontario, J. McQueen, 1880,



two specimens. Thames River, at Chatham (several specimens) and Detroit River, below Sandwich, Ontario (one specimen), Professor Macoun, 1894.

UNIO COCCINEUS, Lea.

Grand River, Cayuga, Ontario, Professor Macoun; one "fairly typical specimen," (C. T. Simpson).

UNIO COMPLANATUS (Solander ?) Lea. (*U. purpureus*, Say.)

Abundant in Nova Scotia, New Brunswick, Quebec and Eastern Ontario. Collected by Dr. R. Bell, in 1859, in creeks, rivers and bays on the north shore at the east end of Lake Superior, along the entire north shore of Lake Huron, also in the St. Mary River. Lake Nipissing, Dr. A. R. C. Selwyn, 1884 (whence it had previously been recorded by Dr. Bell, in 1859). Montreal River, Lake Temiscaming, Ontario, Dr. R. Bell, 1887.

UNIO CORNUTUS, Barnes. (? = *U. reflexus*, Rafinesque.)

Grand River, Cayuga, Ontario, Professor Macoun, 1890; a perfect and fresh left valve.

UNIO ELEGANS, Lea. (*U. truncatus*, as of Rafinesque.)

Thames River, at Chatham (Ontario), Professor Macoun, 1894; one dead but perfect specimen.

UNIO ELLIPSIS, Lea. (? = *U. olivarius*, Rafinesque.)

Ottawa River, opposite L'Orignal, R. Bell, 1854, and near Ottawa, G. C. Heron, 1879 (as *U. olivarius*, Rafinesque). St. Lawrence River, at Montreal, R. Bell, 1858, and near Quebec, J. F. Whiteaves, 1861. Missisquoi River, on the north shore of Lake Huron, Dr. R. Bell, 1860. Lake Erie, at Port Colborne, and Detroit River, near Windsor, Professor Macoun, 1885.

UNIO GIBBOSUS, Barnes. (? = *U. dilatatus*, Rafinesque.)

This species, which has long been known to be abundant

in the St. Lawrence and Ottawa rivers, has recently been collected by Professor Macoun in Lake Erie, at Port Colborne, in the Grand River at Cayuga, and its two branches at Galt and Ayr, also in the Detroit River, at Windsor.

UNIO GRACILIS, Barnes. (? = *U. fragilis*, Rafinesque.)

Collected by Professor Macoun, in 1885, from Lake Erie, at Port Colborne, and the Grand River, at Cayuga; in 1890, at Port Dover, Ontario, and in 1894, in the River Thames, at Chatham.

UNIO LACHRYMOSUS, Lea. (Probably = *U. quadrulus*, Rafinesque.)

In Ontario, Professor Macoun collected specimens of this species in the Grand River at Cayuga, in 1885, and in the Thames River, at Chatham, in 1894.

In Manitoba it was found to be abundant in the Red River, by Dr. G. M. Dawson, in 1873, and Professor J. Fowler has presented to the museum of the Survey a specimen, which he collected at Emerson in 1887.

UNIO LIGAMENTINUS, Lamarck.

Grand River, at Caledonia, Ontario, J. Townsend, 1885, and at Cayuga, Professor Macoun, 1890. Thames River, at Chatham, Professor Macoun, 1894. Roseau River, Manitoba, Dr. G. M. Dawson, 1873, and Assiniboine River, at Mil'wood, J. B. Tyrrell, 1888.

UNIO LUTEOLUS, Lamarck.

Common almost everywhere in Canada east of the Rocky Mountains, though its exact range east of Ontario is a little uncertain, owing to its close resemblance to *U. radiatus*. Dr. Lea, in 1862, records it as occurring in Great Slave Lake, Lake Athabasca, and near the mouth of Moose River, Hudson's Bay. In Manitoba it was collected by Mr. J. B. Tyrrell, in 1887, from the Swan River;

in 1888, from the Assiniboine, and in 1889, from the Red Deer River. It appears to be the only *Unio* in Lake Manitoba, where it was collected by the writer in 1888, and from the Fairford River. In Alberta, Mr. Tyrrell collected it, in 1885, in the Blind Man, Battle and Medicine Rivers.

UNIO NASUTUS, Say.

Two fine specimens of this species, from Toronto Bay, were presented to the museum of the Survey, by Mr. Latchford, in 1886, and since then numerous specimens of it were obtained by Professor Macoun (in 1894) at Rondeau, on Lake Erie.

UNIO NOVI-EBORACI, Lea. (Perhaps = *U. iris*, Lea.)

Grand River, at Cayuga, Professor Macoun, 1890; one perfect specimen. Thames River, at Chatham (two specimens) and Detroit River, below Sandwich (one specimen), Professor Macoun, 1894.

UNIO PHASEOLUS, Hildreth. (? = *U. fasciolaris*, Rafinesque.)

Detroit River, at Windsor (one specimen) and Lake Erie, at Port Colborne (two specimens), Professor Macoun, 1885. Lake Erie, at Kingsville, Ontario (one specimen), J. T. McQueen, 1890, and Thames River, at Chatham (one specimen), Professor Macoun, 1894.

UNIO PRESSUS, Lea.

Boulder River, one of the upper branches of the Attawapishkat River, west of James Bay (in lat. 52° 30' and long. 87° 30'), Dr. R. Bell, 1886; a perfect and fresh right valve. West branch of the Grand River, at Ayr, Ontario, Professor Macoun, 1894, a slightly distorted but living shell. This species has long been known to be common in the Rideau Canal and river, near Ottawa, where it was first noticed by the late E. Billings, about the year 1856 or 1857.

UNIO PUSTULOSUS, Lea. (? = *U. bullatus*, Rafinesque.)

Grand River, Caledonia, Ontario, J. Townsend, 1885; one specimen. Thames River, at Chatham, Professor Macoun, 1894; two specimens.

## UNIO RADIATUS (Gmelin), Lamarck.

No new localities are to be recorded for this common eastern species, which has long been known to range from Nova Scotia to at least as far to the westward as Ottawa.

UNIO RANGIANUS, Lea. (Perhaps a var. of *U. perplexus*, Lea.)

Lake Erie, at Kingsville, Ontario, J. T. McQueen, 1890; one perfect specimen of the shell of the female.

## UNIO RECTUS, Lamarck.

Common in the St. Lawrence and Ottawa rivers, and in western Ontario. In Manitoba, it was collected by Dr. G. M. Dawson, in 1873, from the Roseau River, and by Mr. J. B. Tyrrell, in 1888, in the Assiniboine River at Millwood.

UNIO RUBIGINOSUS, Lea. (? = *U. flavus*, Rafinesque.)

In Ontario this shell has been collected by Professor Macoun, in 1890, in the Grand River at Cayuga, and in 1894, in the Thames River, at Chatham. In Manitoba, it was found by Dr. G. M. Dawson, in 1873, in the Red and Roseau Rivers, and by Dr. R. Bell, in 1883, in Lake Winnipeg, between Fort Alexander and Elk Island.

## UNIO SUBROTUNDUS, Lea.

Grand River, Caledonia, J. Townsend, 1885, one specimen, which "approaches *U. ebennus*" (C. T. Simpson). Port Dover, Lake Erie, a specimen "which approaches *U. solidus*, Lea," (C. T. Simpson), and Rondeau, Lake Erie, one specimen, Professor Macoun, 1894.

UNIO TRIANGULARIS, Barnes. (? = *U. triqueter*, Rafinesque.)

Collected by Professor Macoun, in 1885, at Port Colborne, Ontario, and in 1894, at Rondeau and in the Thames River at Chatham.

UNIO TRIGONUS, Lea. (? = *U. undatus*, Barnes.)

Port Dover, Lake Erie, Professor Macoun, 1890, two perfect but worn specimens, which were identified with this species by Mr. Simpson.

UNIO UNDULATUS, Barnes. (? = *U. costatus*, Rafinesque.)

Ontario. Sable River, at Thedford, Mr. Bissell, 1883, per Dr. H. Ami. Grand River, Caledonia, J. Townsend, 1885. Lake Erie, at Port Colborne, and Detroit River, at Windsor, Professor Macoun, 1885. Grand River, at Cayuga, Professor Macoun, 1890, and Thames River, at Chatham, Professor Macoun, 1894.

Manitoba. Black River, Lake Winnipeg, Dr. R. Bell, 1883, two specimens, with the umbonal regions much eroded. Emerson, Professor J. Fowler, one specimen of a small form which approaches *U. plicatus* (Le Sueur, M.S.S.) Say.

UNIO VENTRICOSUS, Barnes. (*U. occidens*, Lea, female, and *U. subovatus*, Lea, male: ? = *U. cardium*, Rafinesque.)

Common in the St. Lawrence and Ottawa rivers and throughout Ontario. In Manitoba it has been collected in the Red and Roseau Rivers by Dr. G. M. Dawson, in 1873, and at Lake Winnipeg, between Fort Alexander and Elk Island, by Dr. R. Bell, in 1883.

OTTAWA, November 30th, 1894.

"Reprinted from the Canadian Record of Science, April, 1895."

ADDITIONAL NOTES ON RECENT CANADIAN  
UNIONIDÆ.

By J. F. WHITEAVES.

UNIO CANADENSIS, Lea.

In a letter to the writer, dated June 18th, 1895, Mr. Simpson says, "I think there can be little doubt, from examining the type of *U. Canadensis*, that it is a somewhat injured specimen of the male of *U. ventricosus*, Barnes."

Mr. Bryant Walker, of Detroit, informs the writer that he has, in his cabinet, specimens of each of the following species, from the Detroit River:

MARGARITANA HILDRETHIANA, Lea.

Main channel of the Detroit River off Belle Isle, collected by the Michigan Fish Commission in 1895.

UNIO LEIBII, Lea.

Detroit River, at the upper end of Fighting Island, collected by Mr. Walker in 1873 or 1874, and identified by the late Dr. James Lewis.

UNIO MULTIRADIATUS, Lea.

Same locality, collector and date as for the preceding species; also, Thames River, Ontario, from the collection of the late Dr. George A. Lathrop.

UNIO SULCATUS, Lea. (= *U. perplexus*, var. *perobliquus*, Conrad. Types from Detroit River, and Wabash R., Indiana.)

Collected by Mr. Walker in the Detroit River at the upper end of Fighting Island, in 1873 or 1874; at the:

upper end of Belle Isle in 1894; and in the same river, at the locality first mentioned, by the Michigan Fish Commission, in 1895.

UNIO VERRUCOSUS, Barnes.

Main channel of the Detroit River off Belle Isle, collected by the Michigan Fish Commission in 1895, and Detroit River opposite Grassy Island, collected by Mr. Walker in 1895.

Mr. Walker also states that he has, in his collection, twenty-six species of Unionidæ from the Detroit River and Lake St. Clair, viz., *Anodonta Benedictii*, *A. Footiana*, *A. fragilis* and *A. subcylindracea*; *Margaritana deltoidea*, *M. Hildrethiana*, *M. marginata*, and *M. rugosa*; *Unio alatus*, *U. circulus*, *U. coccineus*, *U. ellipsis*, *U. gibbosus*, *U. gracilis*, *U. Leibii*, *U. luteolus*, *U. multiradiatus*, *U. nasutus*, *N. Uovi-Eboraci*, *U. phaseolus*, *U. pressus*, *U. Rangianus*, *U. rectus*, *U. triangularis*, *U. ventricosus* and *U. verrucosus*.

OTTAWA, July 9th, 1895.

the same river,  
Michigan Fish

off Belle Isle,  
in 1895, and  
collected by Mr.

in his collection,  
the Detroit River  
*A. Footiana*,  
*Maritana deltoidea*,  
*U. rugosa*; *Urio*  
*is*, *U. gibbosus*, *U.*  
*diatus*, *U. nasutus*,  
*is*, *U. Rangianus*,  
and *U. verrucosus*.



