


IMAGE EVALUATION TEST TARGET (MT-3)



Photographic Sciences Corporation

23 WEST MAIN STREET WEBSTER, N.Y. 14580 (716) 872-4503

STATE OF THE STATE

W Res Res

CIHM/ICMH Microfiche Series. CIHM/ICMH Collection de microfiches.



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques



(C) 1985

### Technical and Bibliographic Notes/Notes techniques et bibliographiques

The to

The por of film

Ori be the sio oth firs sio or

The she Till

Ma diff ent beg rig red me

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.		qu'il de c poin une mod	stitut a mid lui a été p et exempli t de vue b image rep ification d indiqués d	ossible de aire qui so ibliograph roduite, o ans la mé	e se proce ont peut-é nique, qui u qui peu ethode no	tre uniq peuven vent exi	détails jues du t modifier iger une		
	Coloured covers/ Couverture de coule	ur			Coloured Pages de				
	Covers damaged/ Couverture endomm	agée			Pages da Pages en	maged/ dommagé	ies		
	Covers restored and Couverture restaurée					stored and staurées e			
	Cover title missing/ Le titre de couvertur	e manque				coloured, colorées,			
	Coloured maps/ Cartes géographique	s en couleur			Pages de Pages dé				
	Coloured ink (i.e. otl Encre de couleur (i.e				Showthre Transpare				
	Coloured plates and/ Planches et/ou illust					f print val négale de		ion	
	Bound with other ma Relié avec d'autres d					suppleme d du maté			ire
	Tight binding may co along interior margir Lareliure serrée peu distortion le long de	n/ t causer de l'ombre			Seule édi	ion availation dispo	onible	scured b	ov errata
	Blank leaves added appear within the te have been omitted fill se peut que certailors d'une restaurationais, lorsque cela ét pas été filmées.	xt. Whenever possil rom filming/ nes pages blanches on apparaissent dan	ole, these ajoutées is le texte,		slips, tiss ensure th Les pages obscurcie etc., ont	ues, etc., e best poi totaleme	have bee ssible ima ent ou par feuillet d'e s à nouve	n refilm ige/ rtielleme errata, u eau de f	ed to ent ine pelure,
	Additional comment Commentaires suppl	7.11							
_	item is filmed at the ocument est filmé au			Ssous.					
10X	14X	18X		22X		26X	<del></del>	30X	
	124	164	20 Y		24Y		28X		32X

The copy filmed here has been reproduced thanks to the generosity of:

**National Library of Canada** 

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol → (meaning "CONTINUED"), or the symbol ▼ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

L'exemplaire filmé fut reproduit grâce à la générosité de:

Bibliothèque nationale du Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole → signifie "A SUIVRE", le symbole ▼ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents.
Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

1	2	3

1	
2	
3	

1	2	3
4	5	6

rrata o

tails du odifier


une

mage

pelure, 1 à

32Y

32X



# NATIONAL ARITHMETIC,

IM

## THEORY AND PRACTICE;

DESIGNED FOR THE USE OF

## CANADIAN SCHOOLS.

The medical land of the Art of the breefine M. Ferthern int.

By JOHN HERBERT SANGSTER, Eso.

MATHEMATICAL MASTER AND LECTURER IN CHEMISTRY AND NATURAL PHILOSOPHY IN THE NORMAL SCHOOL FOR UPPER CANADA.

Sanctioned by the Council of Public Instruction for Apper Canala.

THIRD EDITION-CAREFULLY REVISED AND CORRECTED.

Montreal :

PRINTED AND PUBLISHED BY JOHN LOVELL;
AND SOLD BY B. & A. MILLER.

Toronto :

R. & A. MILLER, 62 KING STREET EAST. 1862.

## JAMOUTAA

## OLUMNITHA

## HOLDS THE THE TENTOLICE

THE COURSE STREET, STREET, STR

## CANADIAN SURFOLES

Entered, according to the Act of the Provincial Parliament, in the year one thousand eight hundred and fifty-nine, by John Levell, in the Office of the Registrar of the Province of Canada.

Anterior on the Cos of of Physics and and a Costage Costage

the still and tak tourish out terrible score in the

Director and Identification of the second

er i setteren rener reger bet

### PREFACE.

In preparing the following work (undertaken at the suggestion of the Chief Superintendent of Education for Upper Canada), it has been the constant aim of the Author to present it to Canadian teachers and students as a thoroughly reliable Treatise on the Theory and Practice of Numbers, and as an Arithmetic, in some degree, commensurate with the higher qualifications of teachers and the improved methods of instruction now generally

found in our schools:

The Arithmetic now offered to the public is based upon the Irish National Treatise;—in fact, it was at first intended merely to adapt that work to the decimal currency, and to abbreviate the somewhat tedious reasons there given for the various rules. So many alterations and improvements suggested themselves, however, that the original design was speedily abandoned, and, with the exception of the first ten or fifteen pages, which are taken entire from the work in question, the Treatise, as at present issued, is, in all essential respects, an entirely new book. Nevertheless, as it was the sole object of the Author to prepare a complete text-book on the subject of Arithmetic, he has not hesitated to adopt whatever he considered good, either in the Irish National or in the numerous other excellent works on the subject.

By far the greater number of the problems are original; and it is hoped that the practical manner in which many of them are put, will tend to render the study of Arithmetic more interesting and useful than it has hitherto been. It will be observed, that a thorough series of review examples has been given at the close of each of the sections up to the seventh, and a very extensive set at the end of the book. This is deemed an important feature in the present work, as in some degree insisting upon that careful revision of what has been learned from time to time, without which, the pupil arrives at the end of the book with all the rules and principles so confounded with one another, as to render his

knowledge in a great measure worthless.

Since the only difference between simple and denominate numbers is that the one increase and decrease according to the scale of tens and the other according to different scales, there is no reason why the rules relating to them should be separated; and therefore in the following pages no distinction is made between simple and compound rules. A somewhat extended

rent set for compound numbers.

It will be observed that towards the end of the Treatise the rules are mainly deduced algebraically. Some teachers may not, at first, be disposed to regard this as an improvement, but it was not adopted until after careful deliberation and consultation with many of the most successful teachers of Arithmetic in the Province. It is generally conceded that a pupil should commence, in some sort, the study of Algebra as soon as he has progressed through Proportion in Arithmetic. In schools in which this view is adopted by the teacher, no difficulty can be experienced, as, even in the deduction of the rules, the algebraic principles used are of the simplest possible character.

As some teachers, however, prefer always giving the rule in a purely arithmetical form, this has invariably been appended in

all the cases usually treated of in Common Arithmetic.

Vith regard generally to algebraic formulæ, it may be further arbed, that an algebraic formula is simply the most abbrelated form in which it is possible to express a rule or principle. Que the pupil is properly taught their use, he is in a manner independent of mere memory, since from a very few general inciples he is able, without any reference to a text-book, to educe for himself the whole series of rules for Simple and Comaund Interest. Discount. Annuities. Progression, and Position. ven when the pupil is merely required to commit the rules to semory, it is obvious that he can do so much more readily when y are given to him in the shape of algebraic formulæ than in worded paragraphs. Let any one, for instance, compare the ork necessary for committing the eleven rules for Simple Interthat required to commit the corresponding formulæ, and the result will be a thorough conviction of the superiority the latter mode of giving the rules. In short, every expelenced teacher will admit, that even while the pupil remains at shoel it is next to impossible to make him remember all the different rules for Interest, Progression, and Annuities; and that frectly he leaves the school to enter upon the business of life, these rules are either altogether forgotten or are so confounded with one another as to become mere useless mental lumber.

After many years' trial, the Author is persuaded that the only successful mode of treating the rules in question, is to enable the pupil to deduce them algebraically, and then to interpret and y the resulting formulæ.

The attention of the teacher is respectfully directed to the Recapitulation at the end of the first section, where, it is thought,

the definition and essential principles of Notation and Numeration are so concisely worded that they may be advantageously

committed to memory by the pupil,

erest

pre-

diffe-

e the

not,

1.the

com-

ls in

n be

praic

le in

id in

ther

bre-

iple.

nner

eral

, to

om-

ion.

s to hen

n in

ter-

læ,

rity

pe-

at

the

fe.

led

er.

nd

The examination questions throughout the work have been carefully prepared, and are designed both to enable the self-taught student to test, at each section, the extent and thereaghness of his knowledge of the principles therein contained, and also to guide the pupil as to what principles and definitions are of such importance that they require to be committed to memory. This latter object is further secured by the arrangement of type,—all the definitions and leading principles being printed in large type, the explanations, reasons, and remarks in small type, and the problems in a size intermediate to the two.

Great pains have been taken to render the wording of the rules as perfect as possible; and it will be observed that, in order to catch the eye when glancing over the page, they are

invariably printed in Italics.

It is believed that the sections on Proportion, Fractions, Interest, &c., contain a larger amount of information and a better selection of examples than are commonly given; and that the section on the Properties of Numbers and the different scales of Notation will tend very materially to enlarge the pupil's acquaintance with the general principles of the science of Arithmetic.

Although the Preface is not the proper place for discussing methods of teaching Arithmetic, the Author cannot refrain from

urging upon his fellow-teachers the following points:

1st. The pupil should be thoroughly drilled upon the use of the signs and symbols of Arithmetic, because these constitute the language proper to the subject.

2nd. He should be required to commit to memory all essential definitions, and also the tables of money, weights, and measures. The teacher would do well to examine his pupils on these tables once a month or oftener, since if the pupil has to turn back to his book for each table as it is required, it is not to be expected that his progress will be very rapid or thorough. It may be fairly questioned, whether more than half the difficulty and obscurity that cling to the subject of Arithmetic does not arise from the fact that the pupil is not familiar with the signs, the tables, and the principles of notation.

3rd. The teacher should give his class, from time to time, questions of his own construction, either to solve at home or as ordinary school-room work, and the pupils should be encouraged and required to write questions themselves under each rule. This is an important exercise, and no teacher who once adopts it will ever throw it aside.

4th. In all operations in which there are both multiplication and division, the pupil should be taught to first indicate the processes by their appropriate signs and then cancel as far as possible.

5th. The teacher is respectfully reminded, that without frequent and thorough reviews there can be no real progress. Experience has shown that from one-third to one-half of the time devoted to Arithmetic can be profitably devoted to revision and recapitulation.

6th. The teacher should require from his pupil the absolutely correct answer to each question. 'Near enough' is productive of great mischief to the pupil, as it encourages a habit of such carelessness in his operations, that no confidence can be placed on his results. It is not enough that the pupil understands the principles,—although this of course is important. It is possible so to train the pupil that his operations in Arithmetic shall be at once rapid and accurate, and this should be the aim of the teacher.

with an illustrate the same of the particle of

Com and the think of the second of the secon

The activate of some the control of the solution of the soluti

good to the second state of the second of th

And a relative for the distribute functioners on a state that

the state of the state of

WE WATER TO AN A STATE OF THE S

The transfer of the format france

Toronto, December, 1859.

# quierrant for the second edition.

entropy against the first seed of the real forms of the seed of

and the state of t

1 3 3 17

Aufgreiten V. - Officer in der Steiner und ber der Steine Steine

Change of the grate. The Author embraces the opportunity afforded by the issue of a Second Edition, both to thank his fellow-teachers in Canada for the kind and flattering reception they have given his work, and to offer a few words of explanation on what, as far as he can learn, is the only feature that does not meet with very general approval. He refers to the union of the Compound with the Simple Rules. It has been objected to the arrangement adopted in the National Arithmetic, that a pupil must know the Simple Rules before he can work problems in Reduction or in the Compound Rules. Now this is undoubtedly true, and would be a fatal objection to any such arrangement in an Elementary or Primary Arithmetic. The National is, however, an advanced or second book on arithmetic, and the pupil is assumed to have progressed through an elementary text-book before he enters it. If the National Arithmetic were designed for beginners, where would be the necessity for a First or Elementary book on Arithmetic? The objections have arisen altogether from a misconception of the design of the book. The pupil is supposed to have worked through some elementary text-book on arithmetic, and to have acquired a certain amount of practical skill in arithmetical operations. He then commences the National, and, in progressing through it, not only meets with additional and more advanced practical exercises, but also learns the reasons and the mutual relations of the several rules. In the Elementary he is taught how to multiply an abstract by an abstract number, or an applicate by an abstract number. In the National he is shown that these operations, though differing in detail, are essentially the same in principle; and he is thus enabled to generalize and classify.

ation e the

t fre-Extime n and

P. William

THAU

3113,8

 Another objection urged is, that if the National Arithmetic be designed for a second book on the science, the simple problems given at the commencement of each rule, and indeed the earlier rules themselves, should not be inserted. This is also a mistake. The object has been to exhibit a gradual progression from the simple to the more difficult,—to shew that the most simple and the most complicated problems depend essentially upon the same principles. Indeed, were the National Arithmetic intended merely as a second practical work on arithmetic, three fourths of it might have been omitted, and nothing given but the few rules omitted in the Elementary.

The wiften bear the fit is a second of the second of the

But my of the fifth of the last to the same of free or

riginal Mering of the factor of the section of the

with result of the superior or will be and the superior of the

The first transfer of the first transfer of

Nota Ara Rom

Exer Exer Den

> Redu Redu Reco

Misco Exam

Fund Addi Proo Appl Beca Exam

Subt Proo Appl Reca Exer

Mult Mult To M

nt no this base the entire of the first of the surface of Chiaring with this see . I have a small for military vistas in the state of the stat Appropriation agreement, it is a selection of cultime of and the fred of began a still, and being allowed in a facel transfer इत का उन्नी स्टिन्ट स्ट्रांट वर्ग के नाम महिन्दी के लेकिन के जिल्ला है Bound the see Made and rich to water the sea on the majorate a golden of most of ments for the state of the opening was a state of the or all which are the logic to garden execution of copies of the cor is fea of the never of the in the Applific operations Note the there is twice or to be made our was a war in which the leafer the discuss to effect to the tree in it is even to be of the contract of the contra and their inchance succeedings the their compact and their best take as the ha received the contract of the contract of the contract of the contract of the national per the little and a list section with the the the section of the real is a series of the series appel Howards co. These states an about one is now the first of t the property of the factor a the society is on other many or our is now a that there are and adverse distribution is distributed the same in provide a new his is it is entirely to The state of the same of the

# CONTENTS.

go calletty the spinger

etic"

10 &

ion

ithtic,

11 76

10

1 1

374

it }

<sup>ME</sup> 3

SECTION I.	A. THE
Definitions.	
Notation and Numeration.	
Arabic Notation	10
Roman Notation.	
Exercises in Notation.	89
Exercises in Numeration.	
Denomination of Numbers.	
Tables of Money, Weights, and Measures	
Reduction Descending	K0
Reduction Ascending	' KI
Recapitulation to Section I	N. A.
Miscellaneous Exercises on Section I	1 Kg 60
Examination Questions on Section I	
TANIMITIES COLD CALLED ON DECEMBER 12	, J
SECTION II.	P
	-157
Fundamental Rules	. 62
Addition	63
Proof of Addition	. 67
Application	71
Recapitulation	114
Examination Questions on Addition	1.77 75
Subtraction.	. 76
Proof of Subtraction	. 79
Application	. 82
Examination Questions on Addition. Subtraction. Proof of Subtraction. Application. Recapitulation.	. 83
Examination Questions on Subtraction	. 4 08
Multiplication	. 85
mutuplication Table	. V . 04
To Multiply by a Composite Number	. 90

N G D B T T T T T

TOTRY AASMTOTMEMOTTT

	PAGE
To Multiply when the Multiplier contains Decimals	94
Proof of Multiplication	95
Proof of Multiplication	97
Exercises in Multiplication	100
Examination Questions on Multiplication	
Division	102
General Rule for Division	105
General Principles	
Proof of Division	
General Principles	
To Divide by a Composite Number	
To Divide when both Divisor and Dividend are Denomin	
Numbers	
To Divide when the Divisor or Dividend or both cont	ain
Decimals	114
Contractions in Division	
Exercises in Division	116
Examination Questions on Section II	117
Miscellaneous Exercises on Sections I and II	
SECTION III.	Sangial Control
Properties of Numbers	120
Table of Prime Numbers	
To Resolve a Number into its Prime Factors	126
To Find all the Divisors of a Number	127
Number of Divisors	128
To Find a Common Divisor of two or more Numbers	129
To Find the Greatest Common Measure of two Number	· grain
To Find the Greatest Common Measure of more than Numbers	two 130
Second Method of finding the Greatest Common Measur	2 m m m arth F
Least Common Multiple	132
Scales of Notation	136
To Reduce a Number from One Scale to Another	137
To Reduce a Number from Any Scale into the Decimal.	139
Fundamental Rules in Different Scales.	141
Fundamental Rules in Different Scales,  Duodecimal Multiplication	143

### CONTENTS.

PAGE 94 95 97 100 101 .... 102 .... 105 ....108 ....109 inate ......113 ntain with .... 114 g. V. 115 .... 116 .... 117 Brand Ast

.... 120 .... 125 .... 126 .... 127 .... 128 .... 129 ers.. 129 two

130 re. 131

П	PAGE	
П	Examination Questions on Section III	
L	Miscellaneous Exercises on Sections I-III	)
П	SECTION IV.	
Ш	With the second	No.
П	Vulgar and Decimal Fractions 150	
П	General Principles	L
	Definitions of Fractions	
П	Reduction of Fractions 154	
	To Reduce a Mixed Number to a Fraction	
1	To Reduce an Improper Fraction to a Mixed Number 156	
	To Reduce a Fraction to its Lowest Terms 156	
	To Reduce several Fractions to a Common Denominator 15	7
Н	To Reduce several Fractions to their Least Common Deno-	
п	minator 156	-
	To Reduce a Compound Fraction to a Simple One 15:	
П	Cancellation	
	To Reduce a Complex Fraction to a Simple One 16	
	Reduction of Denominate Fractions 16	2
-	To Reduce one Denominate Number to the Fraction of Ano-	
	ther.	
4	Addition of Fractions	Ε,
	Addition of Denominate Fractions	
	Subtraction of Fractions 16	
	Multiplication of Fractions	
	To Multiply a Denominate Number by a Fraction 17	
L	Division of Fractions	
	To Divide a Denominate Number by a Fraction	
н	Multiplication and Division of Complex Fractions 17	
	Examination Questions on Vulgar Fractions	
1	Decimals and Decimal Fractions	
п	To Reduce a Decimal Fraction to its Corresponding Decimal 18	
	To Reduce a Decimal to a Decimal Fraction 18	
П	To Reduce a Vulgar Fraction to a Decimal	
	To Reduce a Denominate Number of Several Denominations	
	to an Equivalent Decimal of a Given Denomination 18	
	To Find the Value of a Decimal of a Denominate Number. 18	
	TO LINE OF ASIER OF S DECIMAL OF S DEPONDED AND AND AND AND AND AND AND AND AND AN	7

PAGE	
Circulating or Repeating Decimals	
To Determine the Number of Places in the Decimal corres-	
ponding to a Given Vulgar Fraction	
To Reduce a Pure Repetend to a Vulgar Fraction 190	
To Reduce a Mixed Repetend to a Vulgar Fraction 191	
Addition of Circulating Decimals 194	
Subtraction of Circulating Decimals	
Multiplication of Circulating Decimals	
Division of Circulating Decimals	J
Miscellaneous Exercises in Decimals	
Examination Questions on Section IV 197	
Miscellaneous Exercises on Sections I-IV	
the state of the s	
SECTION V.	
Ratio 200	
Proportion.       206         Simple Proportion.       208         Compound Proportion.       213         Conjoined Proportion.       218	4
Simple Proportion	60
Compound Proportion	4/2
Conjoined Proportion	· tro
Examination Questions on Section V	1
Miscellaneous Exercises on Sections I-V	40
SECTION VI.	
Practice.	ľ
Table of Aliquot Parts	1
Bills of Parcels.	
Tare and Tret. 230	
Tare and Tret.  230 Examination Questions.	-
Missellaneons Evergines on Sections L-VI	
The second in the second of th	
SECTION VII.	
Percentage	d.
To Find the Percentage of any Given Number	
Commission	
Brokerage	

SIT CSI AAE Indie Gp Sp Co Tail Edition Ex

Pr Ba Al Ex Fo

### CONTENTS.

PAGE	1 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
186	PAGE
10 Et . 1925 pa	To compute Commission or Brokerage when it is to be
189	deducted in advance from a given amount and the
190	balance invested
191	Stock 237
194	Insurance
194	To compute the sum for which property must be insured in
195	order to cover both its value and the premium paid 240
196	Custom-House Business
196	Specific Duties
197	Au valorem Dunes 233
198	Assessment of Taxes
2	Examination Questions on Section VII 245
°, 4	SECTION VIII.
	A comment of the comm
200	Interest
206	Simple Interest
208	Deduction of Rules for Simple Interest
213	Special Rules for 6 per cent
218	Special Rules for other rates
220	Partial Payments 256
222	Compound Interest
4.574	Table of Amounts of \$1 or £1 at Compound Interest 260
	Discounted A of Tributes the web broom 262
100	Bank Discount 264
224	Equation of Payments
224	Simple Partnership
アルではまったと	Compound Partnership 270
230	Examination Questions on Section VIII 272
. 231	Extended to the control of the contr
ZOI -	SECTION IX.
APPAR TO THE	Profit and Loss 274
1 1 1 1 1 1	Barter 278
. 232	Alligation
. 283	Exchange of Currencies.,.,
234	Foreign Moneys of Account
235	Table showing the Value of the Foreign Coins most fre-
5	quently met with 287
7 .4	

PSDCAAAT

To Est As Ta As

林的

to to

	D
PAGE 1	PAGE
Canadian and United States Currencies	288
To Reduce Dollars and Cents to Old Canadian Or or to the Currency of Any State	irrency
or to the University of Any State	288
To Reduce Old Canadian Currency or Any State Cu	The state of the s
to Dollars and Cents	
To Reduce Dollars and Cents to Sterling Money	290
To Reduce Sterling Money to Dollars and Cents	290
Exchange	290
Arbitration of Exchange	294
Examination Questions on Section IX	298
A Commence of the commence of	
SECTION X.	St. Turket V man 1 15
Involution	207
Evolution	299
Extraction of the Square Root	
Application of Square Root,	
Extraction of the Cube Root	
Horner's Method	0
Application of Cube Root	
Extraction of the Roots of Higher Orders	
Logarithms	
To Find the Logarithm of a Number	
To Find the Natural Number corresponding to a	
Logarithm	
Logarithmic Arithmetic	
Multiplication of Numbers by their Logarithms	324
Division of One Number by Another by means of Loga	
To Raise a Number to Any Power by means of Loga	
Extraction of Roots by means of Logarithms	
Examination Questions on Section X	
	2.
SECTION XI.	10 - 1 - 1 - 1 - 1
Arithmetical Progression	
Table of Rules	
Geometrical Progression.	
Table of Rules.	Ake a Ake
+ mbro or Todiobee de consission de consission de contrata de consission de contrata de co	1 540

### CONTENTS.

Top HTE	7,64
PAGE	PAGE
288	Position. 345
rency	Single Position 346
288	Single Position
rency	Compound Interest
289	Annuities
290	Annuities at Simple Interest
290	Annuities at Compound Interest
290	Table showing the Amount of an Annuity of \$1 or £1 for
294	any number of payments 362
296	Table showing the Present Value of an Annuity of \$1 or £1 363
to made with	Examination Questions on Section XI
	Examination Problems
297	Arithmetical Recreations
299	Tables of Logarithms
300	Tables of Squares, Cubes and Roots
305	Answers to Miscellaneous Exercises
308	Answers to Examination Problems
313	cor has defined their it to all more that he has the
314	The second of th
315	and effected of the state of reference that the death to death the
316	and the second of the second o
318	the state of the s
Given S	of book proving and the fine field of creat to be suffered.
322	opening Test 14; and is read 5 to 2 to a. 7 is a 1.
324	the the reflect size. By the is it is not fight the trans
324	on of a chi, it appropries odd, it station, it was thour among
ithms 325	Troop of S. T. V. Sales the Transaction of the Section of the sales
ithms 326	Ten root of A. jer Mr. doi
326	MOTHER RELIEF OF THE FEMALES
329	the first the spirit of could be the second of the second of the second at
5 3	oughness wheels with the gut or war it should be a change to be a
Contract of	the second of th
331	were a least to be the same of
933	tion of the grante of the control of the control of the control of
337	के मार्ग हुन ने अपने कार्य है। है है है है है है है है
340	
41.670	and the state of t

### SIGNS USED IN THIS TREATISE.

money or account or company of the of the off

+ the sign of addition; as 5+7, or 5 to be added to 7.

the sign of subtraction; as 4-3, or 3 to be subtracted from 4.

× the sign of multiplication: as 8×9, or 8 to be mul-

tiplied by 9.

+the sign of division; as 18+6, or 18 to be divided

by 6.

() which is used to show that all the quantities united by it are to be considered as but one. Thus  $(4+3-7)\times 6$  means 4 to be added to 3, 7 to be taken from the sum, and 6 to be multiplied into the remainder. The latter is equivalent to the whole quantity within the brackets.

= the sign of equality; as 5+6=11, or 5 added to 6,

is equal to 11.

 $\frac{2}{4}$ , and  $\frac{2}{3}$ < $\frac{2}{5}$ , mean that  $\frac{3}{4}$  is greater than  $\frac{1}{2}$ , and that  $\frac{3}{4}$  is less than  $\frac{3}{5}$ .

: is the sign of ratio or relation; thus 5 : 6, means the

ratio of 5 to 6, and is read 5 is to 6.

means that there is the same relation between 5 and 10 as between 7 and 14; and is read 5 is to 10 as 7 is to 14.

 $\checkmark$  the radical sign. By itself, it is the sign of the square root; as  $\checkmark$ 5, which is the same as  $5^{\frac{1}{2}}$ , the square root of 5.  $\checkmark$ 3, is the cube root of 3, or  $3^{\frac{1}{2}}$ .  $\checkmark$ 4 is the 7th root of 4, or  $4^{\frac{1}{2}}$ , &c.

EXAMPLE. [ $\sqrt{(8-3+7)\times 4\div 6}+31]\times\sqrt[3]{9}\div 10\frac{1}{2}\times 5^2=556\cdot 25$ , &c., may be read thus: take 3 from 8, add 7 to the difference, multiply the result by 4, divide the product by 6, take the square root of the quotient and to it add 31, then multiply the sum by the cube root of 9, divide the product by the square root of 10, multiply the quotient by the square of 5, and the product will be equal to 556·25, &c.

These signs are fully explained in their proper places.

# ARITHMETIC

. I a son o . - m i salish don you

## LA A A A TOWN THE WAY TO SEE THE SEE T SECTION I.

## DEFINITIONS. The first side as

with the start of the start of the start of the start of the

1. Science is a collection of the general principles or leading truths relating to any branch of knowledge, arranged in systematic order so as to be readily remembered, referred to, and applied. - It s with it shift done many with

2. Art is a collection of rules serving to facilitate the performance of certain operations. The rules of Art are

based upon the principles of Science.

3. Arithmetic is both a Science and an Art.

4. As a Science, Arithmetic treats of the nature and properties of numbers,; as an Art, it teaches the mode of applying this knowledge to practical purposes. The former may be called Theoretical, and the latter Practical Arithmetic. To Practical Arithmetic belong all the operations we perform upon numbers, as addition, subtraction, multiplication, division, the extraction of roots, &c. The discussion of the principles upon which these operations are founded, constitutes the theory of Arithmetic.

5. Any single thing, as a horse, an apple, a day, an inch, is called a unit or one.

6. Numbers are expressions for one or more units, Thus, the words one, two, three, four, five, &c., or the characters 1, 2, 3, 4, 5, &c., are expressions by which we indicate how many single things or units are to be taken.

7. Numbers are divided into two classes:

1. Abstract numbers. - we will have the warm white our

2. Applicate, Concrete, or Denominate numbers.

dded to 7. subtracted

to be mul-

be divided

ties united -3-7)×6 the sum, he latter is ckets. dded to 6,

, and that

means the

10::7:14. and 10 as s to 14. ign of the

the square 1/4 is the

 $9 \div 10 \frac{1}{2} \times$ B. add 7 to e product it add 31,. divide the quotient equal to

er places.

C

pe

gr

of

fir

no

nu

of

8. If the units referred to by a number have reference to particular objects, as seven days, nine inches, &c., it is called an applied, applicate, concrete, or denominate number. If the units represented by a number have no reference to any particular object, as when we say twice eight are sixteen, or seven and two are nine, it is called an abstract number.

#### NOTATION AND NUMERATION.

9. To avail ourselves of the properties of numbers, we must be able both to form an idea of them ourselves, and to convey this idea to others by spoken and by written language—that is, by the voice, and by characters.

The expression of number by characters, is called notation; the reading of these, numeration. Notation, therefore, and numeration, bear the same relation to each other as writing and reading, and, though often confounded, they are in reality perfectly distinct.

10. It is obvious that, for the purposes of Arithmetic, we require the power of designating all possible numbers; it is equally obvious that we cannot give a different name, or character to each, as their variety is boundless. We must, therefore, by some means or another, make a limited system of words and signs suffice to express an unlimited amount of numerical quantities. With what beautiful simplicity and clearness this is effected, we shall better understand presently.

11. Two modes of attaining such an object present themselves; the one, that of combining words or characters already in use, to indicate new quantities; the other, that of representing a variety of different quantities by a single word or character, the danger of mistake at the same time being prevented. The Romans simplified their system of notation by adopting the principle of combination; but the still greater perfection of ours is due also to the expression of many numbers by the same character.

12. It will be useful, and not at all difficult, to explain to the pupil the mode by which, as we may suppose, an idea of considerable numbers was originally acquired, and of which, indeed, although unconsciously, we still avail ourselves; we shall see, at the same time, how methods of simplifying both numeration and notation were naturally suggested.

Let us suppose no system of numbers to be as yet constructed, and that a heap, for example, of pebbles, is placed before us that we may discover their amount. If this is considerable, we cannot ascertain it by looking at them altogether, nor even by separately inspecting them; we must, therefore, have recourse to

ARTS. 8-17.]

eference to, it is number. rence to are sixabstract

we must convey —that is,

notation; , and nuriting and ality per-

ic, we reis equally aracter to e, by some and signs quantities. s effected,

emselves; in use, to a variety he danger mans simle of comue also to

ain to the ea of conh, indeed, shall see, umeration

even by

that contrivance which the mind always uses when it desires to grasp what, taken as a whole, is too great for its powers. If we examine an extensive landscape, as the eye cannot take it all in at one view, we look successively at its different portions, and form our judgment on them in detail. We must act similarly with reference to large numbers; since we cannot comprehend them at a single glance, we must divide them into a sufficient number of parts, and, examining these in succession, acquire an indirect, but accurate idea of the whole. This process becomes by habit so rapid, that it seems, if carelessly observed, but one act, though it is made up of many; it is indispensable, whenever we desire to have a clear idea of numbers—which is not, however, every time they are mentioned.

13. Had we, then, to form for ourselves a numerical system, we should naturally divide the individuals to be reckoned into equal groups, each group consisting of some number quite within the limit of our comprehension; if the groups were few, our object would be attained without any further effort, since we should have acquired an accurate knowledge of the number of groups, and of the number of individuals in each group, and therefore a satisfactory, although indirect estimate of the whole.

We ought to remark that different persons have very different limits to their perfect comprehension of number. The intelligent can conceive with ease a comparatively large one; there are savages so rude as to be incapable of forming an idea of one that is extremely small.

constitute a group, the ratio; it is evident that the larger the ratio, the smaller the number of groups; and the smaller the ratio, the larger the number of groups.

15. If the groups into which we have divided the objects to be reckoned, exceed in amount that number of which we have a perfect idea, we must continue the process, and, considering the groups themselves as individuals, must form with them new groups of a higher order. We must thus proceed until the number of our highest group is sufficiently small.

16. The ratio used for groups of the second and higher orders, would naturally, but not necessarily, be the same as that adopted for the lowest; that is, if seven individuals constitute a group of the first order, we should probably make seven groups of the first order constitute a group of the second also; and so on.

17. It might, and very likely would happen, that we should not have so many objects as would exactly form a certain number of groups of the highest order—some of the next lower might be left. The same might occur in forming one or more of the other groups. We might, for example, in reckening a heap of pebbles, have two groups of the fourth order, three of

in

ha

po

ST.

the third, none of the second, five of the first, and seven individuals or simple units.

18. If we had made each of the first order of groups consist of ten peobles, and each of the second order consist of ten of the first, each group of the third of ten of the second, and so on with the rest, we had selected the decimal system, or that which is not only used at present, but which was adopted by the Helrawa, Greeks, Romans, &c. It is remarkable that the language of every civilized nation gives names to the different groups of this, but not to those of any other numerical system. Its very general diffusion, even among rude and barbarous people, has most probably arisen from the habit of counting on the fingers, which is not altogether abandoned, even by us.

19. It was not indispensable that we should have used the same ratio for the groups of all the different orders. We might, for example, have made four pebbles form a group of the first order, twelve groups of the first order a group of the second, and twenty groups of the second a group of the third order. In such a case we had adopted a system exactly like that to be found in the table of sterling money, in which four farthings make a group of the order of pence, twelve pence a group of the order of shillings, twenty shillings a group of the order of pounds. While it must be admitted that the use of the same system for applicate, as for abstract numbers, would greatly simplify our arithmetical processes—as will be evident hereafter—a glance at the tables given further on, and those set down in treating of enchange, will show that a great variety of systems have actually been constructed.

20. When we use the same ratio for the groups of all the orders, we term it a common ratio. There appears to be no particular reason why ten should have been selected as a "common ratio" in the system of numbers ordinarily used, except that it was suggested, as already remarked, by the mode of counting on the fingers; and that it is neither so low as unnecessarily to increase the number of orders of groups, nor so high as to exceed the conception of any one for whom the system was intended. (See Section III.)

21. A system of numbers is called binary, ternary, quaternary, quinary, senary, septenary, octenary, nonary, denary, undenary or duodenary, according as two, three, four, five, six, seven, eight, nine, ten, eleven, or twelve, is the common ratio. The denary and duodenary systems are more commonly known as the decimal and duodecimal systems. Ours is therefore a decimal or denary system of numbers.

If the common ratio were sixty, it would be a sexegesimal system. Such a one was formerly used, and is still, to some extent, retained—as will be porceived by the tables hereafter given

(Spot, 1.

en indi-

s consist m of the d se on at which d by the the landifferent t system. Is people, ag on the

ve might,
f the first
cond, and
rder. In
hat to be
farthings
oup of the
of pounds.
system for
nplify our
a glance
reating of

of all the pe no par-"common opt that it counting ecessarily high as to stem was

e actually

taternary, denary or ven, eight, he denary the decilecimal or

xagesimal some exfter given for the measurement of arcs and angles, and of time. A duodecimal system would have twelve for its "common ratio"; a vigesimal, twenty, &c.

22. A little reflection will show that it was useless to give different names and characters to any numbers except to those which are less than that which constitutes the lowest group, and to the different orders of groups; because all possible numbers must consist of individuals, or of groups, or of both individuals and groups. In neither case would it be required to specify more than the number of individuals, and the number of each species of group, none of which numbers—as is evident—can be greater than the common ratio. This is precisely what we have done in our numerical system, except that we have formed the name of some of the groups by combining those already used. Thus, "tens of thousands," the group next higher than thousands, is designated by a combination of words already applied to express other groups—which tends still further to simplification.

23. Arabic system of Notation:-

Units of Comparison, or simple units.

First group, or units of the second order, Second group, or units of the third order, Third group, or units of the fourth order, Pourth group, or units of the fifth order, Pifth group, or units of the sixth order, Sixth group, or units of the seventh order,

One
Two
Three
Pour
Pive
Six
Seven
High
Nine
Ten
Hundred
Thousand
Ten Thousand
1,000
Ten Thousand
10,000
Million
1,000,000

Names Cheracters.

24. The characters which express the first nine numbers are the only ones used. They are called digits, from the custom of counting them on the fingers, already noticed,—"digitus" meaning in Latin a finger, and they have also been called significant figures, to distinguish them from the cipher, or 0, which has no value when standing alone, and which is used merely to give the digits their proper position with reference to the decimal point.

25. The decimal point is a point or dot used to indicate the

position of the simple units.

The pupil will distinctly remember that the place where the "simple units" are to be found is that immediately to the left-hand of this point, which, if not expressed, is supposed to stand at the right-hand side of all the digits. Thus, in 462-78 the \$ expresses "simple units," being to the left of the decimal point;

26. We find by the table just given, that, after the first nine numbers, the same digits are constantly repeated, their positions with being, however, changed; to indicate succeeding groups, the digit is moved, by means of a cipher, one place farther to the left. Any one of the digits me need to express its respective number of any of the groups:—thus a would be eight "simple units"; 80, eight groups of the first order, or eight "tens" of simple units; 800, eight groups of the second, or units of the third order; and so on. We might use any of the digits with different groups; thus, for example, 5 for groups of the third order, 3 for those of the second, 7 for those of the first, and 8 for the "simple units," then the whole set down in full would be 5000, 300, 70, 8, or, for brevity's sake, 5378. For we never use a cipher, when the place it would occupy may be filled up by a digit; and it is evident that in 5378 the 378 keeps the 5 four places from the decimal point (understood), just as well as ciphers would have done also the 78 keeps the 3 in the third, and the 8 keeps the 7 in the second place.

27. It is important to remember that each digit has two values, an absolute and a relative. The absolute value is the number of units it expresses, whatever these units may be, and is unchangeable; thus 6 always means six; sometimes, indeed, six tens; at other times six hundreds, &c. The relative value depends on the order of units indicated, and on the nature of the "simple unit."

What has been said on this very important subject is intended principally for the teacher, though an ordinary amount of industry and intelligence will be quite sufficient for the purpose of explaining it, even to a child, particularly if each point is illustrated by an appropriate example; the pupil may be made, for instance, to arrange a number of pebbles in groups, sometimes of one, sometimes of another, and sometimes of several orders, and shan be desired to express them by characters—the "unit of comparison" being occasionally changed from individuals, suppose to tens, or hundreds, or to scores, or dozens, &c. Indeed the pupils must be well acquainted with these introductory matters, otherwise they will content the habit of answering without any very definite ideas of many things they may be called upon to explain, and which they should be expected particular to understand. Any trouble bestowed by the teacher at this period will be well rapaid by the ease and rapidity with which the learner will afterwards advance. To be assured of this, he has only to recollect that most of his future reasonings will be derived from, and his explanations grounded on the very principles we have endeavoured to unfold. It may be taken as a truth, that what is all learns without understanding, he will acquire with disgust, and will see to remember; for it is with children as with persons of more that year — when we appeal successfully to their understandings, the particular what is and pleasure they feel in the attainment of knowledge, cause the law and the wear ness which it costs to be undervalued or forgotten.

Pebbles will answer well for examples—indeed, their use in computing

y , ...\*

1: 1000

6 . 1/1. Om # d

979328

at being

rst nine
positions
hanged;
byed, by
ne of the
t groups
0, eight
1 so on.
s; thus,
se of the
units,"
3, or, for
the place

values, umber of nchangetens; at cends on "simple

evident decimal e done;

ed princind intellieven to a
example;
pebbles in
of several
"unit of
se to tens,
st be well
i content
ines the
parifecta,
on will oc

fterwards ost of his unded on aken as a uire with n as with to their nment of be under-

omputing

### ROMAN SYSTEM OF NOTATION.

28. Our ordinary numerical characters have not been always, or everywhere, used to express numbers; the letters of the alphabet naturally presented themselves for the purpose, as being already familiar, and, accordingly, were very generally adopted—for example, by the Hebrews, Greeks, Romans, &c., each, of course, using their own alphabet. The pupil should be acquainted with the Roman notation on account of its beautiful simplicity, and its being still employed in inscriptions, &c.: it is found in the following table:—

Characte	rs. Numbers Expressed.
· · · · · · · · · · · · · · · · · · ·	One.
11.	<b>Two.</b>
II.	
Anticipated change II	
Change	Five.
V]	Six.
V1	II Seven.
	II Eight.
Anticipated change IX	
Change . X.	. Ten.
X	
	II. Twelve,
( ) ( ) ( ) ( <b>X</b> ]	III Thirteen.
X 1 1 1 1	V. Fourteen.
	V. Fifteen.
	VI. Sixteen.
	VII. Seventeen.
** ** ** * * * * * * * * * * * * * * *	VIII. Eighteen.

XIX.:

has given rise to the term calculation, "calculus" being, in Latin, a pebble; but while the teacher illustrates what he says by groups of particular objects, he must take care to notice that his remarks would be equally true of any others. He must also point out the difference between a group and its equivalent unit, which, from their perfect equality, are generally confounded. Thus, he may show that a penny, while equal to, is not identical with four farthings. This seemingly unimportant remark will be better appreciated hereafter; at the same time, without inaccuracy of result, we may, if we please, consider any group either as a unit of the order to which it belongs, or so many of the next lower as are equivalent.

. Nineteen.

. Twenty.

XXX., &c. . Thirty, &c.

### Characters. Numbers Expressed.

Anticipated chang	zeXL.	. Forty.
Change of the contract of		. Fifty.
man in the state of the state o	LX., &c.,	Sixty, &o. a tre
Anticipated change	geXC.	. Ninety.
Change	<b>C.</b>	. One hundred.
hel care on he ages	CC., &c.,	. Two hundred, &c.
Anticipated chan	geCD.	. Four hundred.
Change		. Five hundred, &c.
Anticipated chan	geCM.	. Nine hundred.
Change	M. or CIO.	. One thousand, &c.
• • •	V. or IDD.	. Five thousand.
	X.orCCIAA	. Ten thousand, &c.
37	Ingo.	. Fifty thousand, &c.
	ccciooo	. One hundred thousand,&c

29. Thus we find that the Romans used very few characters—fewer indeed than we do, although our system is still more simple and effective from our applying the principle of "position," unknown to them.

They expressed all numbers by the following symbols, or combinations of them: I. V. X. L. C. D. or IO. M., or CIO. In constructing their system, they evidently had a quinary in view; that is, as we have said, one in which five would be the common ratio; for we find that they changed their character, not only at ten, ten times ten, &c.; but also at five, ten times five, &c. A purely decimal system would suggest a change only at ten, ten times ten, &c.; a purely quinary, only at five, five times five, &c. As far as notation was concerned, what they adopted was neither a decimal ner a quinary system, nor even a combination of both; they appear to have supposed two primary groups, one of five, the other of ten "units of comparison"; and to have formed all the other groups from these, by using ten as the common ratio of each resulting series.

30. They anticipated a change of character,—one unit before it would naturally occur; that is, not one "simple unit," but one of the units under consideration. In this point of view, four is one unit before five; ferty, one unit before fifty—tens being now the units under consideration; four hundred, one unit before five hundred—hundreds having become the units contemplated.

31. From the table (28) it will be seen that as often as any letter is repeated, so many times is its value repeated. Thus I, standing alone, denotes one, II denotes two, &c. So X denotes ten, XX twenty, &c.

When a letter of less value is placed before a letter of greater value, it takes away its own value from the greater; but when placed after it, it adds its own value to the greater. Thus V denotes five, IV denotes four, and VI six; so X denotes ten, IX nine, and XI eleven, &c.

A line or bar placed over any letter increases its value a thousand-fold. Thus V denotes five, V denotes five thousand; X denotes ten, X denotes ten thousand, &c.

32. To express a number by the Roman method of notation:

RULE.—Find the highest number within the given one, that is expressed by a single character, or the "anticipation" of one (28); set down that character, or anticipation, as the case may be, and take its value from the given number. Find what highest number less than the remainder is expressed by a single character, or "anticipation"; put that character or "anticipation" to the right hand of what is already written, and take its value from the last remainder; proceed thus until nothing is left.

EXAMPLE.—Set down the number eighteen hundred and forty-four, in Roman characters. One thousand expressed by M. is the highest number within the given one, indicated by one character or by an "anticipation"; we put down

and take one thousand from the given number, which leaves eight hundred and forty-four. Five hundred, D, is the highest number within the last remainder (eight hundred and forty-four) expressed by one character, or an "anticipation"; we set down D to the right hand of M,

MD.

and take its value from eight hundred and forty-four, which leaves three hundred and forty-four. In this the highest number expressed by a single character, or an "anticipation," is one hundred, indicated by C: which we set down, and for the same reason two other C's.

#### MDCCC.

This leaves only forty-four, the highest number within which, expressed by a single character or an "anticipation" is forty, KL,—an "anticipation" we set this down also,

MDCCCXL.

Four, expressed by IV, still remains; which, being also added, the whole is as follows:—

MGCCCXLIV.

C.

cc.

&c. usand,&c

haracters still more of "posi-

s, or com-CIO. In y in view; le common ot only at t, &c. A it ten, ten s five, &c. pted was mbination oups, one it to have

nit before le unit," t of view, fty—tens lred, one the units

### EXERCISE. 1.

### 33. Express the following numbers in the Roman notation:-1. Twenty-five.

- 2. Forty-three.

- 3. Sixty-seven.4. Eighty-nine.5. Ninety-eight.
- 6. One hundred and thirty-seven.
- 7. Three hundred and seventy-one.
- 8. Four hundred and two.

- 9. Six hundred and seventeen.
- 10. Nine hundred and ninety-nine.
- 11. One thousand four hundred and forty-six.
- 12. Three thousand eight hundred and five.
- 13. Eight thousand six hundred and seventy.
- 14. Twelve thousand one hundred and sixty-nine.
- 15. Four hundred and ninety-seven thousand, six hundred and eighty-two.

## Answers.

- भ भारती वार वार । ५ ५% १० १ 1. XXV. Asis some 2. XLIII.
- 4. LXXXIX. 5. XCVIII. 6. CXXXVII.

the state of the s

- 10. OMXCIX. 11. MCDXLVI. 12. MMMDCCOV.
- 13. VMMMDCLXX. 14. XMMCLXIX.

## 15. ODXOVMMDOLXXXII.

### EXERCISE 2.

### 34. Read the following expressions:-

- 2. COLXXII. 3. DOLXVIII.
- 1. XCVII.
- 5. XV. 6. VMMMXXXIII. 4. CMIX.
- 7. XVDCCCLXXXVIII. 8. DCXLVMCMIV. 9. XXVXXV.
- 1. Ninety-seven.
- 2. Two hundred and seventy-two.
- 3. Six hundred and sixty-eight.
- 4. Nine hundred and nine.
- 5. Fifteen thousand.
- 6. Eight thousand and thirty-three.
- 7. Fifteen thousand eight hundred and eighty-eight.
- 8. Six hundred and forty-six thousand nine hundred and four.
- 9. Twenty-five thousand and twenty-five,

### ARABIC SYSTEM OF NOTATION.

- 35. In the Common or Arabic system of Notation the same character may have different values, according to the place it holds with reference to the decimal point (25), or perhaps more strictly to the simple units. This is the principle of position.
- 36. The places occupied by the units of the different orders (23), may be described as follows:—simple units, one place to the left of the decimal point, expressed, or understood; tens, two places; hundreds, three places, &c.
- 37. When, therefore, we are desired to write any number, we have merely to put down the digits expressing the amounts of the different units in their proper places, according to the order to which each belongs. If, in the given number, there is any "place" in which there is no digit, a cipher must be set down in that place, when required to keep another digit in its own position.—But a cipher produces no effect, when it is not between one or more digits and the decimal point; thus, 0536, 536-0, and 536 would mean the same thing—the first is, however, incorrect. 536 and 5360 are different; in the latter case the cipher affects the value, because it alters the position of the digits.

EXAMPLE.—Let it be required to set down six hundred and two. The six must be in the third, and the two in the first place; for this purpose we are to put a cipher between the 6 and 2—thus 602. Without a cipher the six would be in the second place—thus, 62; and would mean, not six hundreds, but six tens.

38. In numerating, we begin with the digits of the highest order, and proceed downwards, stating the number which belongs to each order.

To facilitate notation and numeration, it is usual to divide the places occupied by the different orders of units into periods. For a certain distance, the English and French methods of division agree; the English billion is, however, a thousand times greater than the French. This discrepancy is not of much importance, since we are rarely obliged to use so high a number; —we shall prefer the French method. To give some idea of the amount of a billion, it is only necessary to remark, that, according to the English method of notation, there has not been one billion of seconds since the birth of Christ. Indeed, to reckon even a million, counting on an average three per second for eight hours a day, would require nearly 12 days. The following are the two methods;

dred and

gov.

VIII. XXXIII. XV.

nd four.

FRENCH METHOD.	Bundreds of Octillions.   Fens of Octillions.   Cotillions.   Hundreds of Septillions.   Fens of Septillions.   Septillions.	333333	333333	c.   Tens of Elikons.   c.   Billions.   c.   Billions.   c.   Tens of Millions.   c.   Millions.   c.   Millions.   c.   Hundreds of Thousands.   c.   Tens of Thousands.   Tens of Thousands.	
Brelibe Method.	Hunds, of Thous, of Quadrillions. Tens of Thous, of Quadrillions. Thousands of Quadrillions. Hundreds of Quadrillions. Tens of Quadrillions. Ouadrillions.	Hunds, of Thous, of Trilliens, Tens of Thousands of Trillions, Thousands of Trillions, Hundreds of Trillions, Tens of Trillions,	Hunds, of Thous, of Billions. Tens of Thous, of Billions. Thousands of Billions. Hundreds of Billions. Tens of Billions. Billions. Hunds, of Thous, of Millions.	Tens of Thousands of Mrillions Thousands of Millions Hundreds of Millions Tens of Millions Millions Hundreds of Thousands Tens of Thousands	Thousands. Hundreds, Tens. Usite

39. Use of Periods.—For the purpose of reading or writing numbers, we divide them by separating points, into periods—the first separating point being the decimal point, expressed or understood, and the other separating points being placed after every third digit, or place, to the right and left of the decimal point. Each period has three places—of which one or more may be occupied by digits. The lowest place in every period—or that to the right hand, is the "units'" place of that period: and the highest, the "hundreds'" place. And this is true, whether the period is to the left or to the right of the decimal point.

40. The period to the left of the decimal point contains the simple units. The first period to the left of the units' period, contains the thousands; and the first period to the right of it, the thousandths. The second period to the left of the units' period, contains the millions; and the second to the right of it, the millionths. The third period to the left of the units' period, contains the billions; and the third to the right of it the billionths. The fourth period to the left of the units' period, contains the trillions; and the fourth to the right of it, the trillionths. The fifth period to the left of the units' period, con-

ARTS: 39, 40.7

r writing periodspressed or aced after e decimal or more periodat period: is true, e decimal

tains the s' period, ght of it, the units' ght of it, s' period, of it the s' period, of it, the riod, con-

tains the quadrillions; and the fifth to the right of it, the quadrillionths. The sixth period to the left of the units' period, contains the quintillions; and the sixth to the right of it, the quintillionths. The seventh period to the left of the units' period, contains the sextillions; and the seventh to the right of it, the The eighth period to the left of the units' period, contains the septillions; and the eighth to the right of it, the septillionths. The ninth period to the left of the units' period, contains the octillions; and the ninth to the right of it, the octillionths. The tenth period to the left of the units' period, contains the nonillions; and the tenth to the right of it, the nonillionths.

The pupil should be made perfectly familiar with the names of the periods and of the places in each period—so as to be able, without the slightest hesitation, to name the period and place to which any digit belongs, or into which it ought to be put. When he can read or write any one digit, belonging to any period and place, he should be taught to read and write a number consisting of two, three, four, &c., digits, whether they are close tegether, or separated by any number of ciphers.

The whole of what has been said above will become more evident from an attentive consideration of the following table:

,	Sof Onedrillions			of Trillions.		1	of Billiens.			of Millions.			of Thousands.	₹. •		of Units.	, i		of Thousandins.	31	1	or Millionths.			Sol Dilliontills.	· · ·	Cot Teffisonthe		#1 #3 **	CofOnedrillionths			of Quintillionths.	A
W Handreds	OTens	Cellinita	THE drede	Z-Tens	*Units	or Hundreds	@Tens	e Units	oo Hundreds	@Tens	2Units	*Hundreds	&Tens	co-Units	o Hundreds	Tens	e Units	& Hundreds	9 Tens		7 Handreds	STens	& Units	Handreds		o United	Thomas	Lillinite		-Traine	Triffe	- Hundreds	o Tens	. Units
2	ath Deriod 20	8	1	5th Period 2	-	5	4th Period. \ @	2,	8	3rd Period. < %	7.,	4	2nd Period. < &	3	_	1st Period.	2,		1st Period. < &	4	7	2nd Period. < co	8,	·	Srd Period. 4 5	5,	Ath Dowload	T CE TONE		Eth Dowing ) 8	-	3,	6th Period.	9

EXAMPLES.—Let it be required to read off the following number, 576934. We put a point to the left of the 9, and find that there are exactly two periods—thus, 576,934; this does not always occur, as the highest or lowest period is often imperfect, consisting only of one or two digits. Dividing the number thus into parts, shows at once that 5 is in the third place of the second period—that is, in the Hundreds' place of the Thousands' period; and therefore, that it expresses five hundred thousands; that the 7, being in the second place of the same period indicates tens of thousands; and the 6, being in the first indicates thousands. The 9, being in the

fir le

od

be

py CU

CO

to

gr th If th 88 qu it ' it

> an th th

de

po

le

pu

to

m

al

si

third place of the first period, indicates hundreds of units: the 3, being in the second place of the same period, indicates tens of units; and the 4, being in the first, indicates units ("of comparison," or "simple units"). The number, therefore, may be read as follows—"five hundreds of thousands, seven tens of thousands, and six thousands; nine hundreds of units, three tens of units, and four units"; or more briefly, "five hundred and seventy-six thousand nine hundred and thirty-four."

41. To prevent the separating point or that which divides into periods, from being mistaken for the decimal point, the former should be a comma (,)—the latter a full stop (.) Without this distinction, two numbers which are very different might be confounded: thus, 498,763, and 498,763, one of which is a thousand times greater than the other. After a while we may dispense with the separating point, though it is convenient to retain it with large numbers, as they are then read with greater case.

42. To write down any integral or whole number, it is merely necessary to remember the order of the periods, and that every period contains three places, each of which must be filled, either by a digit or a cipher. The one, two, or three digits, belonging to the highest period are first written in their appropriate places; then the next lower period is filled with the digits, or ciphers belonging to it; afterwards the next; and so on, till the whole number is set down.

EXAMPLE.—Let it be required to write the number is set down.

EXAMPLE.—Let it be required to write the number seventy-three trillions two hundred and nine billions eighteen thousand and six. The highest period here mentioned is that of trillions, which we know to be the fifth to the left of the decimal point (40). We therefore set down the digits 73, bearing in mind that we are to put in four complete periods, or twelve places between the 3 and the decimal point. The next period we have is that of billions, which we fill with digits 209 (two hundred and nine). The next period, that of millions, has no significant figures, and we accordingly fill it thus, 600. We now come to the period of thousands, in which we have the digits 18, but, inasmuch as the third place of this period must also be filled, we insert there a cipher, and the full period becomes 018. Lastly, the lowest period, or that of units, is to contain only the digit 6,—the other two places being filled with ciphers, the complete period is written 606. Now setting these periods one after the other in their proper order, we obtain for the entire number the expression, 78,209, 000,018,006.

43. To write down any decimal number we proceed very much in the same way. We have to remark, that in any decimal the last digit to the right gives the denomination to the number. Thus, 68 is read sixty-eight hundredths; 4078 is read four thousand and seventy-eight tenths of thousandths, &c.

Now, when we wish to write any decimal, we first ascertain how many places the proposed denomination or order is to the right of the decimal point; and then, if the given digits will not bring the number to its proper position, we insert between these digits and the decimal point the requisite number of ciphers.

EXAMPLE. 1.—Let it be required to write the number, seven hundred and sixteen thousand and eighty-nine billionths. Now we know (40) that billionths occupy the 9th place to the right of the decimal point. Were we to place the decimal point immediately before the digits themselves, thus, '716089, they would express not so many billionths but so many millionths: since millionths occupy the 6th and billionths the 9th place. It is obvious, then, that to give the digits their proper value, we must insert three ciphers between them and the decimal point, and the number is then correctly written '600.716.089. then correctly written '000,716,089.

s, being in and the 4, le units"). s of thous of units, ndred and

o periods, a comma ers which 763, one of le we may o retain it

is merely
y period
y a digit
e highest
the next
ig to it;
et down.
hree tril
The highethe fith
digits 73,
or twelve

or twelve ve have is and nine). and we acsands, in dis period becomes only the complete other in n, 73,209,

d very lecimal umber. r thou-

tin how e right t bring digits

undred 60) that Were uselves, ny milace. It ust inmber is EXAMPLE 2.—Write the number six thousand two hundred and one hundredths of trillionths. From (40) we know that hundredths of trillionths occupy the 14th place. The given digits (6201) being only four in number, require the aid of ten ciphers in order to fill the 14 places, and the number is thus written, '000,000,000,003,01.

EXAMPLE 3.—Write the number, six millions seven hundred and twenty-seven thousand and twelve tenths of billionths. The given digits, 6727012, are only seven in number, while the denomination tenths of billionths implies that ten places must be filled. We have, therefore, to insert three ciphers between the given digits and the decimal point, and the resulting expression, 000,672,701,2, represents the given number.

- 44. The simple units are, as we have said, always found in the first period to the left of the decimal point. The digits to the left hand, progressively increase in a tenfold degree—those occupying the first place to the left of the simple units being ten times greater than the simple units; those occupying the second place, ten times greater than those which occupy the first, and one hundred times greater than the units of comparison themselves; and so on. Moving a digit one place to the left, multiplies it by ten—that is, makes it ten times greater; moving it two places, multiplies it by one hundred—that is, makes it one hundred times greater; and so of the rest. If all the digits of a quantity be moved one, two, &c., places to the left, the whole is increased ten, one hundred, &c., times—as the case may be. On the other hand moving a digit, or a quantity one place to the right, divides it by ten, that is makes it ten times smaller than before; moving it two places divides it by one hundred, or makes it one hundred times smaller, &c.
- 45. We possess this power of easily increasing, or diminishing, any number in a tenfold, &c., degree, whether the digits are all at the right, or all at the left of the decimal point; or partly at the right or partly at the left. And the pupil must remember that the quantities increase in a tenfold degree to the left, and decrease in the same degree to the right wherever the decimal point may happen to be. We therefore put quantities ten times less than simple units one place to the right of them, just as we put those which are ten times less than hundreds, &c., one place to the right of hundreds, &c. Quantities to the left of the decimal point are called integers because none of them is less than a whole simple "unit"; and those to the right of it, decimals. When there are decimals in a given number, the decimal point is always expressed, and is found at the right-hand side of the simple units.
- 46. The periods to the left of the decimal point may be called the ascending, and those to the right of it the descending series:

  —taken together, however, they constitute but one series, which is an ascending or rescending series, according as it is read from right to left or from left to right. Periods that are equally distant from the units of comparison bear a very close relation to

each other, which is indicated even by the similarity of their names; the only difference being in the terminations (40). We have seen also, that when we divide integers into periods (40), the first separating point must be put to the right of the thousands. In dividing decimals into periods, the first point must be put to the right of the thousandths also.

ne

ar

to

be

to

Th

me

1.

3. 3.

4. 5.

6.

9.

10.

11.

12.

13.

14.

15.

4.

- 47. Care must be taken not to confound what we now call "decimals," with what we shall hereafter designate "decimal fractions"; for they express equal, but not identically the same quantities—the decimals being what shall be termed the "quotients" of the corresponding decimal fractions. This remark is made here to anticipate any inaccurate idea on the subject, in those who already know something of arithmetic.
- 48. There is no reason for treating integers and decimals by different rules, and at different times, since they follow precisely the same laws, and constitute parts of the very same series of numbers (46). Besides, any quantity may, as far as the decimal point is concerned, be expressed in different ways; for this purpose we have merely to change the unit of comparison. Thus, let it be required to set down a number indicating five hundred and seventy-four men. If the unit be one man, the quantity would stand as follows, 574. If a band of ten men, it would become 57-4-for as each man would then constitute only the tenth part of the "unit of comparison," four men would be only four tenths, or 0.4; and since ten men would form but one unit, seventy men would be merely seven simple units, or 7, &c. Again if it were a band of one hundred men, the number must be written 5.74; and lastly, if a band of a thousand men, it would be 0.574. Should the "unit" be a band of a dozen, or a score of men, the change would be still more complicated; as, not only the position of a decimal point, but the very digits also, would be altered.

49. It is not necessary to remark that moving the decimal point so many places to the *left*, or the digits an equal number of places to the *right*, amounts to the same thing.

Sometimes in changing the decimal point, one or more ciphers are to be added; thus, when we move 42.6 three places to the left, it becomes 42600; when we move 27 five places to the right it is 0.00027, &c.

50. It follows from what we have said, that a decimal, though less than what constitutes the unit of comparison, may itself consist of not only one, but several individuals. Of course it will often be necessary to indicate the nature of the "simple units;" as 3 scores, 5 dozen, 6 men, 7 companies, 8 regiments, &c. But its nature does not affect the abstract properties of numbers; for it is true to say that seven and five, when added

y of their 40). We iods (40), the thouat must be

2 2 7 2 1/21/02

now call cimal fracame quanquotients" k is made t, in those

ecimals by w precisely e series of he decimal or this puron. Thus, re hundred tity would ald become tenth part our tenths, venty men if it were 5.74; and 4. Should the change position of

he decimal al number

altered.

ore ciphers aces to the o the right

al, though may itself f course it ne "simple regiments, pperties of hen added

together, make twelve, whatever the unit of comparison may be :- provided, however, that the same standard be applied to both; thus 7 men and 5 men are 12 men; but 7 men and 5 horses are neither 12 men nor 12 horses; 7 men and 5 dozen men are neither 12 men nor 12 dozen men. When, therefore, numbers are to be compared, &c., they must have the same unit of comparison :- or without altering their value, they must be reduced to those which have. Thus we may consider 5 tens of men to become 50 individual men-the unit being altered from ten men to one man, without the value of the quantity being changed. This principle must be kept in mind from the very commencement, but its utility will become more obvious hereafter.

### Exercise 3.

51. Write down the following Numbers:-

1. One hundred and ninety-four. 2. One thousand and seventy-six.

- 3. Twenty thousand five hundred and eight. 4. Two hundred and one thousand and three. 5. Eighty millions four thousand and thirty-three.
- 6. Sixteen quadrillions five hundred and ninety-seven trillions three billions forty-four millions and ninety-one. ी । रहिता भिर हैरे

7. Ninety-seven hundredths.

8. Six hundred and forty-three thousandths.

9. One hundred and twenty-two thousand and eighty-nine expected or to the extent millionths.

10. Thirty-nine tenths of millionths,

- 11. Sixty-three hundredths of trillionths. 12. Seventeen billions four thousand and one, and nine hundred and sixty-seven billionths. If pro to the for the second
- 13. Seven trillions eight hundred and two billions twenty-three thousand and eleven, and nine thousand nine hundred and ninety-nine billionths.

One quadrillion one trillion one billion one million one thousand one hundred and one, and one trillionth.

Eight hundred and ninty-six trillions and two, and nine 15. hundred and four hundredths of millionths.

### Answers.

 1. 194.
 2. 1076.
 3. 20508.

 4. 201003.
 5. 80004033.
 6. 16597003044000091.

 7. 97.
 8. 643.
 9. 122089.

10. 0000039. 11. 0000000000063. 12. 1700004001.00000967. 13. 7802000023011.000009999. 14. 1001001001101.600000000001.

### EXERCISE 4.

# 52. Read the following numbers :-

1. 904.			
1 004			
			٠.

2. 7060.

3. 90004.

4. 40300201.

5. 7060504030.

6. 70003000000400.

7. 604.03.

8. 90767.004003.

9. 9001.00070306.

10. 1237-9134671342913.

11. .00100100100101.

12. 100-2003004005006007.

#### Answers.

- 1. Nine hundred and four.
- 2. Seven thousand and sixty.
- 3. Ninety thousand and four.
- 4. Forty millions three hundred thousand two hundred and one.
- 5. Seven billions sixty millions five hundred and four thousand and thirty.
- 6. Seventy trillions three billions and four hundred.
- 7. Six hundred and four, and three hundredths.
- 8. Ninety thousand seven hundred and sixty-seven, and four thousand and three millionths.
- 9. Nine thousand and one, and seventy thousand three hundred and six hundredths of millionths.
- 10. One thousand two hundred and thirty seven, and nine trillion, one hundred and thirty-four billion six hundred and seventy-one million three hundred and forty-two thousand nine hundred and thirteen tenths of trillionths.
- 11. One hundred billion one hundred million one hundred thousand one hundred and one hundredths of trillionths.
- 12. One hundred, and two quadrillion three trillion four billion five million six thousand and seven tenths of quadrillionths.

#### ON THE DENOMINATION OF NUMBERS.

53. When two numbers have the same unit they are said to be of the same denomination; when the units are not the same, they are said to be of different denominations. For example, 16 shillings and 28 shillings are two numbers of the same denomination; but 23 shillings and three farthings are not of the same denomination, the unit of 23 shillings being one shilling, and of three farthings, one farthing. The kind of unit always expresses the denomination.

Even in abstract or simple numbers, different names are given to the units as we proceed to the right or left of the decimal point, viz., simple units or units of the first order; tens, or units of the second order; hundreds, or units of the third order, &c. Considered in this relation to each other, these units may be regarded as denominate numbers.

The following Tables show the various kinds of denominate numbers in general use, and also the relative values of their different units.

# TABLES OF MONEY, WEIGHTS, AND MEASURES.

#### STERLING MONEY.

54. The denominations are pounds, shillings, pence, and farthings.

TABLE.

	farthir pence		r.) m	ake	1 pen	ny m	arke	d d.
20	shillin	gs	-0		1 pou		11 66	6.1 <b>(£</b> )
er er	qr. 4	p.2.	<i>d</i> . 1		· · · · · · · · · · · · · · · · · · ·	ام کسر ما ما اما	A Francisco	Taras in the same of the same
	48 960	=	12 240		1 20	=	£ 1	1 1 1 1

Other English coins, some of them now out of use:

Moidore	= :	27s.		Noble	,= y	6s. 8d.
Guinea	= -	21s.		Crown	· == `\	58.
Pistole	=	168.	10d.	Angel	= "	10s.
Mark or Merk	=	13s.	4d.	Groat	=	4d.

The letters  $\mathcal{L}$  s. d. and qr. are the initials of the Latin words, libra, solidus, denarius, and quadrans, which respectively signify a pound, a shilling, a penny, and a farthing, or quarter. The mark  $\nearrow$ , which sometimes separates the shillings and pence, is a corruption of the long f (s), arising from the rapidity with which it is made.

Sterling money is supposed to have received its name from the *Esterlines* or German traders in England, by whom it is said to have been first coined.

The pound is so called, because in ancient times it was equal to a pound Troy of silver.—Its present value in Canada is \$4'8666, and hence the value of an English shilling is \$4'9 cents. The guinea was so called from being originally coined from gold brought from Guinea, on the coast of Africa.

dred and

thousand

and four

nine trilndred and
wo thousiths.
dred thou-

nths. ur billion quadrill-

are said are not inations. wo numnd three unit of ngs, one denomThe present standard gold coin of Great Britain consists of 22 parts pure sold and 3 parts of copper. The standard silver coin consists of 37 parts sore eller and 3 parts copper. In copper coin 24 peace weigh a pound woirdupois.

## FEDERAL MONEY.

55. Federal money is the currency of the United States. The denominations are eagles, dollars, dimes, cents and mills.

10	mills	(m.) 1	make	1	cent,	marl	ked ct.	
	cents	` '	"	1	dime,		d.	
10	dimes	3 :	"		dollar		8	
10	dollar	9	"		eagle,		E.	
m.		ct.	1 2		,			
10	=	1			d.			
100	=	10	=		1	\$		
1000	=	100	=	1	0 =	1	, E	
0000	=	1000	=	10	0 =	10	= 1.	

The sign \$ is the symbol for the old Spanish coin of 8 reals. On one side of the Spanish real the pillars of Hercules were represented supporting the world—on the piece of eight reals the pillars were retained and the 8 written over them—thus \$. Many however consider the sign \$ a contraction of the letters U. S., the initials of United States made by dropping the curve of the U and writing the 8 over it.

The present standard for both gold and silver coin in the United States is 950 parts of pure metal and 100 parts of alloy. The alloy for gold is silver and copper, of which not more than one half must be silver; that for silver is pure copper.

The gold coins are the Eagle, the Double Eagle, Half Eagle, Quarter Eagle, and Dollar; the silver coins are the Dollar, Half Dollar, Quarter Dollar, Dime, Half Dime and three cent piece; the copper coins are the Cent and the Half Cent; Mills are never coined.

## OLD CANADIAN MONEY.

56. The denominations are pounds, dollars, shillings, pence, and farthings.

> TABLE. 4 farthings make 1 penny, marked d. " 1 shilling, 12 pence 5 shillings " 1 dollar. " 1 pound; 4 dollars qr., ... d. 4 = pilliria: S. 7 48 J= 12 0= 11 0 A 10 8 11 "By 14 1 mg 240 = 60 = 5 = 1 960 = 240 = 20 = 4 = 1.

eta pure 87 parte a mound

MECT. 1.

States.

oue side rting the and the a conby drop-

ed States l is silver t for sil-

Quarter Quarter are the

illings,

Norm.—Every 3d. of the old coinage is equal to 5 cents of the new. The York shilling is equal to the eighth part of a \$, or to 7\frac{1}{2}d. or to 12\frac{1}{2} cents.

# NEW CANADIAN OR DECIMAL MONEY.

57. The denominations are dollars and cents.

The coins are cents, five-cent pieces, ten-cent pieces, and twenty-cent pieces.

100 cents (c) make 1 dollar, marked \$

## AVOIRDUPOIS WEIGHT.

58. Is used in weighing heavy articles. Its name is derived from French—and ultimately from Latin words signifying "to have weight." Its denominations are tons, hundredweights, quarters, pounds, ounces, and drams.

#### TABLE

16 drams 1	make	1 ounce,	marked	OZ.
16 ounces	66	1 pound,	• 66	lb.
25 pounds	"	1 quarter,	"	qr.
4 quarters	".	1 hundredwe	ight."	cwt.
20 cwt.		1 ton,	""	t.
d. o	z.	• • •		
16 = 1		lb.		
256 = 16	=	1 . 1 . q		
6400 = 400	=	25 = 1	. cw	t.
25600 = 1600	=	100 = 4	= 1	t
512000 = 32000	=	2000 = 80	= 20	= 1

It was formerly the custom to allow 28 lbs. to the quarter, 112 lbs. to the hundredweight; and 2240 to the ton. This has now fallen into disuse; and among merchants in Canada the qr., cwt., and ton are universally considered as respectively equal to 25 lbs., 100 lbs., and 2000lbs. The Custom Houses continue to regard the cwt. as equal to 112 lbs., and some few articles are still weighed by the old cwt. by farmers and others. The English cwt. is 112 lbs.

#### TROY WEIGHT.

59. The denominations of Troy Weight are pounds, ounces, pennyweights, and grains.

#### TABLE.

24	grains	(grs.)	make	1	penny	weight,	mar	ked	dwt.
20	pennyv	veight	s. " "	1	ounce,	19		•	OZ.
12	ounces	5			pound				lb.

This weight was introduced into Europe from Cairo, in Egypt, and was first adopted in Troyes, a city of France—whence its name. It is used in philosophy, in weighing gold, precious stones, &c.

philosophy, in weighing gold, precious stones, &c.

Note.—The origin of all weights used in England, was a grain of wheat taken from the middle of the ear and well dried. A weight equal to 32 of these grains was called a pennyweight, being equal to the weight of a silver penny then in use; 20 of these pennyweights constituted an ounce, which was the 12th part of a pound (Lat. "uncia," a 12th part—compare "inch." the twelfth part of a foot.) In later times the pennyweight came to be divided into 24 equal parts instead of 32, but these still retain the name of grains.

The "Carat," which is equal to about four grains (somewhat less than Troy grains), is used in weighing diamonds. The term carat is also applied in estimating the fineness of gold: the latter, when perfectly pure, is said to be "24 carats fine." If there are 23 parts gold, and one part some other material, the mixture is said to be "23 carats fine"; if 22 parts out of the 24 are gold, it is, "22 carats fine," &c. The whole mass is, in all cases supposed to be divided into 24 parts, of which the number consisting of gold is specified. Our gold coin is 22 carats fine; pure gold, being very soft, would too spon wear out. The degree of fineness of gold articles is marked upon them at the Goldsmiths' Hall; thus we generally perceive "18" on the cases of gold watches: this indicates that they are "18 carats fine"—the lowest degree of purity which is stamped. degree of purity which is stamped.

,			grs.
A Troy ounce contains	 		480
An Avoirdupois ounce	 		4371
A Troy pound	 • • • •	1	5.760
An Avoirdupois pound	 	••	7,000

A Troy pound is equal to 372.965 French grammes. 175 Troy pounds are equal to 144 avoirdupois; 175 Troy are equal to 192 avoirdupois ounces.

# APOTHECARIES' WEIGHT.

60. The denominations of Apothecaries' Weight are pounds, ounces, drams, scruples, and grains.

	1	TA.	BLM.			200
20 grains (grs.)	make	1	scruple,	marked	sc. or	<b>D</b> '
3 scruples	. "	1	dram,		dr. or	
8 drams	"		ounce,	b	oz. or	
12 ounces	"		pound,	. 66	lb.	3
grs. $\ni$				٠.,		i,
	*,e	3	1	** ***		15
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	=	1	3		PROFE.	(1
5760 = 288	= = :	96	= 12	- 10. - 1.	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 1

s used in

of wheat at to 32 of of a silver ce, which e "inch," be divid-of grains. less than to applied re, is said ome other out of the cases sup-

Apothecaries mix their medicines by this weight, but buy and sell by avoirdupois.

The pound and ounce of this weight are the same as in Troy weight.

# LONG MEASURE.

61. The denominations of Long Measure are leagues, miles, furlongs, rods, yards, feet, inches, and lines.

		T	ABLE.	- ;	11
12 lines (l.)	make	1	inch,	marked	in.
12 inches	"		foot,		ft. X
3 feet	cc		yard,	. 66	yd.
5½ yards	**		rod, pole,	or perch.	
40 rods or perches	, cc	1	furlong,	* "	fur.
8 furlongs	66,		mile,	ίι , ,	m.
3 miles	"		league,	, ((	lea.
691 miles (nearly)	"	1	degree or earth's	360th pr	rt of the
in 1	Pt.				San St.

in.		ft.					. 5.45
12	=	1		yd.			
36	=	3	=	1		rd.	
198	=	161	=	51	=	1. 1	fur.
7920	=		=	220	=	40 =	1 . m.
63360	=	5280	=		=	320 =	8 = 1.

100 links, 4 rods, or 22 yards, make 1 Gunter's chain. Each link therefore is equal to  $7_{100}^{99}$  inches.

Eleven Irish are equal to 14 English miles. The Paris foot is equal to 12.792 English inches, the Roman foot to 11.604 English inches, and the French metre to 39.383 English inches.

4 inches make 1 hand (used in measuring horses).

3 inches " 1 palm.
18 inches " 1 cubit.

3 feet " a common pace.

5 feet " a Roman pace.

6 feet " a fathom.

120 fathoms " a cable's length.

# SQUARE MEASURE.

62. This measure is used for estimating artificers' work, such as flooring, plastering, painting, paving, &c., and, in short, any kind of work where surface alone is concerned. It is always employed in measuring land, and hence it is frequently called Land Measure.

of gold is oft, would ked upon the cases the lowest

Troy are

ght are

D 3 33

A square is a four sided figure having all of its sides equal and perpendicular one to another. If the length of each side be an inch, a foot, or a yard, &c., the square is called a square inch, a square foot, or a square yard, &c. It will be observed from the adjacent figure that a square foot contains 12× 12 or 144 square inches. and similarly a square contain 3×3 or 9 square feet.

				-		4	_			-	Ċ.	
	-	,		-			_	_	-	_	_	
-	-	. 0		,	٠,	7	1	ŧ,	ν,	9.5	1.	
		, )			1		۸.,	1		质析		800
	-	_	_	-	_	_		_	_	_		-
_			-	_	_	_	7 (7	_	· ·	11.	1.	
_				_						٨,	• '	_
-	-	_	-	_		<del></del>	. "	-		?	e apri	· •
, -	7	_	-	-	-	-	-	1		1.	ŧ,	-

The denominations of Square Measure are square miles, acres, roods, square perches, square yards, square feet, and square inches.

#### TABLE.

144 square inches make 1 square foot, marked sq. ft. 9 square feet " 1 square yard, sq. yc. 301 square yards 1 square rod, sq. rd. 1 rood, 40 square rods r. 1 acre, 66 4 roods 11 1 square mile, 640 acres

sq. i	n.	sq. ft.			1
144	=	1	sq. yd.		1, 1.
1296	-	• 9 =	1	sq. rd.	1.
39204	=	$272\frac{1}{4} =$	301 =	. 1	r.
1568160	=	10890 =	1210 =	40 =	1 : acre.
6272640	=	43560 =	4840 =	160 =	4 = 1,

68. In measuring land, Gunter's chain is used. It is divided into 100 links.

79% inches	make ·	1 link,	marked	1.
100 links or 4 roo	s. "	1 chain,	agent of the	C
on 80 chains at la c	166	1 mile,	W W G	m.
10000 square links	I Saint	1 square	chain,"	eq. c
10 square chain	5 66	1 acre.	TOLL WAS	. 2.

e miles et, and

q. ft. y. yc. . rd.

m.


re.

It is

# SOLID OR CUBIC MEASURE.

64. This measure is used for finding the solid contents of timber, stone, &c. A cube is a solid bounded by six equal surfaces or squares, and having eight equal edges. It is called a cubic inch, a cubic foot, or a cubic yard, according as each of these edges is an inch, a foot, or a yard in length.

The accompanying figure represents a cubic yard—each edge being 3 feet in length. The top, which is equal to the base, contains 3×3 or 9 square feet; hence, if it were only one foot in height it would contain 9 cubic feet; but it is 3 feet in height, and must therefore contain 9×3 or 27 cubic feet. A cubic yard then contains 3×3×3 or 27 cubic feet.



Similarly it may be shown that a cubic foot contains  $12 \times 12 \times 12$  or 1728 cubic inches.

The denominations of Cubic Measure are cords, tons. cubic feet, and cubic inches.

make 1 c. ft. marked c. ft. 1728 cubic inches " I cubic yd. " c. yd. 27 cubic feet \*40 c. ft. of round timber, or 1 " 1 ton, " ton.

50 c. ft. of sq. or hewn timber

128 cubic feet make 1 cord of firewood, marked c.

c. in. c. ft.

A pile of cord-wood 4 feet high, 4 feet wide, and 8 feet long, contains 128 cubic feet or one cord. One foot in length of such a pile is called a cord-foot. It is equal to 16 solid feet, and is consequently equivalent to the eighth part of a cord.

# CLOTH MEASURE.

65. The denominations of Cloth Measure are French ells, English ells, Flemish ells, quarters, nails, and inches.

<sup>\*</sup> A ton of round timber is that quantity of timber which, when hewn, will make 40 cubic feet.

hd

100 201 403

806 T avo and I bee equ cuk for E

an

We

21 inches (in.) make	1 nail, marked na.
4 nails "	1 quarter . " qr.
3 quarters "	1 Flemish ell, "Fl. e.
4 quarters "	1 yard, '" yd.
5 quarters "	1 English ell, "E. e.
6 quarters "	1 French ell, "F. e.
in. na.	The second secon
2\frac{1}{9} = \frac{1}{4} = \frac{1}{1}	701
$\frac{9}{27} = \frac{4}{12} = \frac{1}{3} = \frac{1}{3}$	Fl. e. 1 yd.
36 = 16 = 4 =	
45 = 20 = 5 =	$1\frac{3}{3} = 1\frac{1}{2} = 1$ Fr. e.
54 = 24 = 6 =	$2 = 1\frac{1}{2} = 1\frac{1}{2} = 1.$

Note. The Scotch ell contains 4 quarters 11 inch.

### DRY MEASURE.

66. By this are measured all dry wares, as grain, beans, coal, oysters, &c.

The denominations of Dry Measure are chaldrons, bushels, pecks, gallons, quarts, and pints.

4 quarts	" "	. 1 gallon,	"	gal.
4 quarts 2 gallons	"	1 peck,	"	pk.
4 pecks	"	1 bushel,	"	bu.
36 bushels	. "	1 chaldron,	"	ch.
pt	qt.	1) to		
	1 .	gal.		
8 = 16 =	4 = 8 =	$\begin{array}{ccc} 1 & & \text{pk.} \\ 2 & = & 1 \end{array}$	bu.	
. 64 =	32 - =	8 = 4 =	= 1	ch.

= 1152 = 288 = 144 =

2 pints (pt.) make 1 quart, marked qt.

Our Standard of Dry Measure is the Winchester bushel. This is an upright cylinder whose internal diameter is 19½ inches and depth 8 inches. It contains 2150'4 cubic inches of 77'627 lbs. Avoirdupois of pure distilled water at 62° Fahr, and 30 in. barometer. The standard unit of Dry Measure in the United States is also the Winchester bushel, so called because the standard measure was formerly kept at Winchester, England. The standard unit of Dry Measure in Great Britain is the Imperial bushel, which is an upright cylinder whose internal diameter is 18'789 inches and depth 8 inches. It contains [2218'192 cubic inches or 80 lbs. Avoirdupois of pure distilled water at 62° Fahr, and 30 in. barometer.

Grain is often bought and sold by weight, allowing for a bushel, 60 lbs. of wheat, 53 lbs. of rye, 56 lbs. of Indian corn, 48 lbs. of barley, 34 lbs. of cets, 60 lbs of peas, 50 lbs. of beans, 40 lbs. of buckwheat, 60 lbs. of timethy or red clover seed.

red clover seed.

### AND MEASURES.

## LIQUID MEASURE.

67. Liquid Measure is used for measuring all liquids.

The denominations of Liquid Measure are tuns, pipes, hogsheads, barrels, gallons, quarts, pints, and gills.

#### TABLE.

4 gills (g.)	make	1	pint,	marked	pt.
2 pints	"	1	quart,		qt.
4 quarts	"	1	gallon,	. "	gal.
314 gallons	"		barrel,	"	bar.
2 barrels			hogshead		hhd.
2 hogsheads	"		pipe,	"	pi.
2 pipes	"	1	tun,		tun.

g:	pt.	at.			11	
8 =	2 =	1	gal.			
32 =	8 =	4 =	1 .	bar.	,	
1008 =	252 =	126 =	$31\frac{1}{4} =$	1	hhd.	
2016 =	504 =	252 =	63 =	2 =	1	pi.
4032 =	1008 =	504 =	126 ==	4 =	. 2 =	1 tun.
8064 =	2016 =	1008 =	252 =	8 =	4 =	2 = 1

The English Imperial gallon contains 277 274 cubic inches or 10 lbs. avoirdupois of pure distilled water, weighed at a temperature of 62° Fahr. and under a barometric pressure of 30 inches.

In the United States the wine gallon contains 231 cubic inches, and the beer gallon 283 cubic inches. The gallon of Great Britain is therefore about equal to 1.2 gallons United States Wine Measure.

By an Act of the Imperial Parliament, 1826, the Imperial gallon of 277°274 cubic inches, was adopted as the only gallon, and is therefore the standard for both liquid and dry measure.

Beer is sold usually by the gallon; sometimes, however, in casks of 5 gals., 10 gals., 20 gals., &c. The beer barrel contains 36 gallons, and the hogshead 54 gallons.

## TIME MEASURE.

68. Time is naturally divided into days and years—the former measured by the revolution of the earth on its axis, and the latter by the revolution of the earth round the sun.

The denominations of Time Measure are years, months, weeks, days, hours, minutes, and seconds.

n, beans,

haldrons,

This is an h 8 inches, re distilled Dry Mea-ed because land. The shel, which nd depth 8 of of pure

l, 60 lbs. of bs. of oats, imothy or

#### TABLE.

60	seconds (sec.) m	ake 1	minute,	marked	min.	
	minutes	1	hour,	. "	h	101
24	hours	" :1	day,	66	d	
	days	" 1	week,	" (( ,	wk.	4 ,
	weeks		lunar month	, "	mo.	
	lunar months or	. )		4 1.	A.9	
12	calendar months	or >	make 1 civil	year, m	arked	yr.
	days (nearly)	. )	. ,			110
-		•	• • •	17		10

80	c. mi	n.			* 7.
60	= 1	h		:	
3600	= 60	= 1	da.		,
86400	= 1440	= 24		wk.	
604800	= 10080	= 168	= 7 =	: 1.	yr.
31557600	==525960	=8766	= 3651 =	52 28	= 1.

The twelve calendar months, into which the civil or legal year is divided, and the number of days in each, are as follows:

First month, January, has 31 days. Second "February," 28 "Third "March" 31 " Third March, 31 " 80 .. Fourth April; May, . 31 \*\* Fifth . \*\* " Sixth June, 30 " 31 \*\* Seventh " July, . 66 " Eighth August, 31 September, " 41 Ninth 30 " October, Tenth 31 November, " December, " Eleventh" Twelfth "

The number of days in the respective months may be recalled by recollecting the following well-known lines:

Thirty days hath September, April, June, and November, February has twenty-eight alone, And all the rest have thirty-one; But leap-year coming once in four, February then has one day more.

The number of days in each month may also be recollected by counting the months on the four fingers and three intervening spaces. Thus, January on the first finger; February in space between first and second fingers; March on second finger; April in second space; May on third finger; June in third space; July on fourth finger: August on first finger (since there are no more spaces); September in first space, &c. Now, when counted thus, all the months having 31 days come on the fingers, and all having 30 and the spaces.

thus, all the months having of days continued the spaces.

The solar year is the time elapsing from the passage of the sun from either solstice back to the same again, and is equal to 365d. 5h. 48m. 48sec.

The sidereal year is the time between two successive conjunctions of the sun with some star, and is equal to 365d. 6h. 9m. 14isec.

The civilor legal year is that in common use among different nations and is

ARTS.

civil sixth the 2 from the L Wen This the ci

> The solar 11m. error the Ju longer XIII., days in Hence

This tember to con done p Date

Style; To el and if

25th M Russ amoun

> 69. Div

Leap elapse

Thu it is 3 be Le

To t years, fourth Julian Thus r Janua

To r year; leap ye 1700, 1 years, hand. fore le years 104.

in.

ked vr.

is divided,

by recol-

counting us, Janud fingers; ger; June nce there n counted having 30

om either ons of the

ons, and is

This additional day is given to every fourth year, in order to make the civil year agree with the solar. It was originally added by repeating the sixth of the calends of March in the Roman calendar—corresponding with the 24th of February with us. The day was called the intercalary day, from the Latin intercalo, to insert; and the year was called biesewite, from the Latin bis, twice, and sextills, sixth (i.e., sixth calend, taken twice). We now call it Leap Year, because it leaps a day more than a common year. This correction was made by Julius Casar, emperor of Rome, and hence the civil year is often called the Julian year.

The addition of one day every four years would be strictly correct, if the solar year contained 365d. 6h.; but it only contains 365d. 5h. 48m. 48s., or 11m. 12s. less than 365d. 6h. Adding 1 day every 4 years, gives us then an error of excess of 44m. 46s., or about 3 days for every 400 years. Thus the Julian calendar was behind the solar time, since the Julian year was longer than the natural year. This error, at the time of Pope Gregory XIII., amounted to 10 days, which he corrected in 1562 by suppressing 10 days in the month of October, the day after the 4th being called the 15th. Hence this calendar is sometimes called the Gregorian colendar.

This correction was not adopted in England till 1752, when the error amounted to 11 days. By Act of Parliament, 11 days after the 2d of September were therefore omitted. The civil year, by the same act, was made to commence on the 1st of January, instead of the 25th of March, as it had done previously.

Dates reckoned by the old method or Julian calendar, are called Old Style; and those reckoned by the new method are called New Style.

To change any date from Old to New Style, we must add 11 days to it; and if the given date in Old Style is between the 1st of January and the 25th March, we must add 1 to the year in New Style.

Russia still reckons dates according to Old Style. The difference now amounts to 12 days.

69. To ascertain whether a year is LEAP YEAR.

Divide the given year by 4, and if there is no remainder it is Leap Year. The remainder, if any, shows how many years have elapsed since a Leap Year occurred.

Thus, dividing the year 1847 by 4, the remainder is 3; hence it is 3 years since the last Leap Year, and the ensuing year will be Leap Year.

To this rule there is an exception; for we have seen that a color year is 11m. 12s. less than a Julian year, which is 365t days. This error, in 400 years, amounts to about 3 days; consequently, if a day is added every fourth year, that is, if we have 100 leap years in 400 years, according to the Julian calendar, the reckoning would fall 3 days behind the color time. Thus reckoning from the commencement of the Christian era, when it was January 1st, 401, by the Julian time, it was January 4th by the solar time.

To remedy this error, only 1 centennial year in 4 is regarded as leap year; or, which is the same in effect, whenever the centennial year, or the number expressing the century, is not divisible by 4, that year is not a leap year, while the other centennial years are. Thus, 17, 18, 19, denoting 1700, 1800, and 1900, are not divisible by 4, consequently they are not leap years, though according to the rule above they would be; ou the other hand, 16 and 20, denoting 1600 and 2000, are divisible by 4, and are thezefore leap years. There is still a slight error; but it is so small that ir, 5000 years it scarcely amounts to a day.

Ju

17

the

whi

pos

sola

sur vey lon

deg

70.—TABLE SHOWING THE NUMBER OF DAYS FROM ANY DAY OF ONE MONTH TO THE SAME DAY OF ANY OTHER MONTH IN THE SAME YEAR.

From any	. "	•		T	the th	e sai	me d	lay	of			
day of	·Jan.	Pob.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
January	365	-31	59	90	120	151	181	212	243	273	304	334
February	334	365	28	59	89	120	150	181	212	242	273	303
March						92	122	153	184	214	245	275
April	275	306	334	365	30	61	91	122	153	183	214	244
May	245	276	304	335	365	31	61	92	123	153	184	214
June	214	245	273	304	334	365	30	61	92	122	153	183
July	184	215	243	274	304	335	365	31	62	92	123	153
August	153	184	212	243	273	304	334	365	31	61	92	
September.	122	153	181	212	242	273	303	334	365	30	61	91
October												
November .					181							
December.	31				151							

The months counted from any day of, are arranged in the left-hand vertical column; those counted to the same day of, are in the upper horizontal line; the days between these periods are found in the angle of intersection, in the same way as in a common table of multiplication. If the end of February be included between the two points of time, a day must be added in leap years.

EXAMPLE 1.—How many days are there from the 15th of March to the 4th of October? Looking down the vertical row of numbers at the head of which October is placed, and at the same time along the horizontal row at the left hand side of which is March, we perceive in their intersection the number 214: so many days, therefore, intervene between the 15th of March and the 15th of October. But the 4th of October is 11 days earlier than the 15th: we therefore subtract 11 from 214, and obtain 203, the number required.

EXAMPLE 2.—How many days are there between the 3rd of January and the 19th of -May? Looking as before in the table, we find that 120 days intervene between the 3rd of January and the 3rd of May; but as the 19th is 16 days later than the 3rd, we add 16 to 120, and obtain 136, the number required.

Since February is in this case included, if it were a leap year, as that month would then contain 29 days, we should add 1 to the 136, and 137 would be the answer.

#### EXAMPLES.

- 1. How many days from May 3d to the 4th of next July?

  Ans. 62 days.
- 2. How many days from July 4th to the 25th of next December?

  Ans. 174 days.
- 3. How many days from March 21st to the 23rd of the next September?

OF ONE

> 91 61

> 30

eft-hand he upper d in the of mul-

the two

ch to the the head ntal row ersection a 15th of ys earlier 203, the

uary and 120 days the 19th number

as that and 137

y? lays. ember? lays. ne next lays. 4. How many days from September 23rd to the 21st of the next March?

Ans. 179 days.

5. How many days from June 21st to the 22nd of the next December?

Ans. 184 days.

6. How many days from December 22nd to the 21st of the next June?

Ans. 181 days.

7. How many days from March 21st to the 21st of the next June?

Ans. 92 days.

8. How many days from January 13th, 1848, to September 17th of the same year?

Ans. 248 days.

71. The unit of time is the basis of that of Length, Mass, and Pressure: the connectious being as follows:—

A Pound Pressure means that amount of pressure which is exerted towards the earth, at the level of the see, by the quantity of matter called a pound.

A Pound of Matter means a quartity equal to that quantity of pure water which, at the temperature of 62° Fahr., would occupy 27°272 cubic inches.

A cubic inch is that cube whose side, taken 39 1393 times, would measure the effective length of a London seconds-pendulum.

A London seconds-pendulum is that which, by the unassisted and unopposed effect of its own gravity, would make 86400 vibrations in an artificial solar day, or 86163'09 in a natural sidereal day.

## CIRCULAR MEASURE.

72. Circular Measure, sometimes called Angular Measure, is chiefly used by astronomers, navigators, and surveyors, for measuring angles and for reckoning latitude and longitude, and the motion of the heavenly bodies.

The Denominations of Circular Measure are signs, degrees, minutes, and seconds.

#### TARLE

60 seconds (") make 1 minute, marked '
60 minutes " 1 degree, " 9
30 degrees " 1 sign, " s.
12 signs or 360 deg. 1 circle, " c.

 The circumserence of every circle is supposed to be divided into 360 equal parts called degrees, as in the subjoined figure. Since a degree is simply the yet part of the circum-ference of a circle, it is obvious that its length must depend upon the size of the circle. If the circumference be 360 miles in length, then a degree of that circle will be one mile long; if the circle be 360 inches in circumference, then a degree will be one inch, &c.

The divisions of the circumference of the circle into 360 equal parts took its origin from the length of the year, which, in round numbers, was supposed to contain 360 days, or 12 months of 30 days each. The 12 signs cor-

sepond to the 12 months.

The term minute is from the Latin minutum "a small part." The term econds is \$11 abbreviated expression for second minutes, or minutes of the

#### MISCELLANEOUS TABLE.

73. 12 individual things m	nake 1 dozen.
12 dozen	" 1 gross.
12 gross	" 1 great gross.
20 individual things	" 1 score.
24 sheets of paper	" 1 quire.
20 quires	" 1 ream.
112 pounds	" 1. quintal
200 "	" 1 barrel of pork or beef.
196 ,	" 1 barrel of flour.
14 24 70	" 1 stone."

## BOOKS.

A sheet folded into two leaves is called a folio.

- folded into four leaves is called a quarto, or 4to.
- folded into eight leaves is called an octavo, or Svo. folded into twelve leaves is called a duodecimo, or 12mo.
- folded into eighteen leaves is called an 18mo

74. When figures are written by the side of each other, thus.

2587931272

the language implies that the unit in each place is equivalent to ten units of the place next to the right; or that ten units of any particular place are equivalent to one unit of the place immediately to the left.

the tio thi

> the tion of f the six

pri nex of t

nur valı 7 0 mu whi we ply

or a

dre

ord we whi thi

twe

a k

4to.

r 8vo.

cimo,

ther,

75. When figures are written thus,

d.

the language implies that 10 units of the lowest denomination make one of the second; ten of the second, one of the third; and ten of the third, one of the fourth.

76. When figures are written thus,

gr. lb. cwt. 21 11

the language implies that 16 units of the lowest denomination make one of the second; 16 units of the second, one of the third; 25 units of the third, one of the fourth; 4 of the fourth, one of the fifth; and 20 of the fifth, one of the sixth.

All other denominate numbers are formed on the same principle; and in all of them we pass from a lower to the next higher denomination by considering how many units of the one make one unit of the other.

## REDUCTION.

77. Reduction is the changing the denomination of a number from one unit to another, without altering the value of the number. For example, if we desire to reduce 7 of the order of hundreds to a lower denomination, we multiply the 7 by 10, and thus obtain 70 of the order tens. which are equal to 7 of the third order or hundreds. If we wish to reduce to a still lower denomination, we multiply the tens by ten, and this gives us 700 of the first order or simple units, which are just equal to 70 tens or 7 hundreds.

If, on the contrary, we wish to reduce 900 of the first order or simple units, to units of the third order or hundreds. we divide by 10, and thus obtain 90 of the second order. which we again divide by 10 and obtain 9 units of the third order or hundreds.

Hence reduction of denominate numbers is divided into two parts:-

1st. To reduce a number from a higher denomination to a lower: this is called Reduction Descending.

2nd. To reduce a number from a lower denomination to a higher: this is called Reduction Ascending.

## REDUCTION DESCENDING.

#### EXAMPLE.

78. Reduce £6 16s. 01d. to farthings.

£ 8. d. 6 16 01

20

136 shillings = £6 16s.

1632 pence = £6 16s. 0d.

6529 farthings = £6 16s. 01d.

EXPLANATION.—In this example we multiply the 26 by 20, because each pound is equal to 20 shillings; 6 pounds are therefore equal to 120 shillings, and the 16 shillings given in the question make 136 shillings. Then we multiply the number of shillings by 12, because each shilling is equal to 12 pence, and, since there are no pence in the question, we simply set down the result, 1632 pence. Lastly, we multiply the 1632 pence by 4, because each penny is equal to 4 farthings, and to the result we add the one farthing given in the question.

From the above example and solution we deduce the following

#### RULE.

Multiply the highest given denomination by that quantity which expresses the number of the next lower contained in one of its units; and add to the product that number of the next lower denomination which is found in the quantity to be reduced.

Proceed in the same way with the result; and continue the process until the required denomination is obtained.

#### EXERCISE 5.

1.	How ma	ny farthings in 23828 pence	? Ans. 93312.
		ny shillings in £348?	Ans. 6960.
3.	How ma	ny pence in £38, 10s. ?	Ans. 9240.
4.	How ma	ny pence in £58 13s.?	Ans. 14076.
5:	How ma	ny farthings in £58 13s.?	Ans. 56304.
6.	How ma	ny farthings in £59 13s. 616	1.? Ans. 57291.
		ny pence in £63 0s. 9d.?	Ans. 15129.
80	How ma	ny pounds in 16 cwt., 2 qrs.,	16 lb. ? Ans. 1666.
9.	How ma	ny pounds in 14 cwt., 3 qrs.	16 lb.? Ans. 1491.
10,	How ma	ny grains in 3 lb., 5 oz., 12 d	lwts., 16 grains?
- dat	a probably to be	high in the resident the second in the second in the second	Ans. 19984.

CT. L

n to

e each Then qual to down ecause ne far-

which units; ration rocess

e the

3312. 6960. 9240. 4076 6304. 7291. 5129. 1666.

491. 9984.

11. How many grains in 7 lb., 11 oz., 15 dwt., 14 grains? Ans. 45974.

12. How many hours in 20 (common) years? Ans. 175200. Ans. 5280.

.13. How many feet in 1 mile? 14. How many minutes in 46 years, 21 days, 8 hours, 56 minutes (not taking leap-year into account)? Ans. 24208376.

15. How many square yards in 74 square perches?

Ans. 2238.5 (2238 and a half). 16. How many square yards in 46 acres, 3 roods, 12 perches? Ans. 226638.

17. How many square acres in 767 square miles? Ans. 490880.

18. How many cubic inches in 767 cubic feet? Ans. 1325376.

19. How many quarts in 767 pecks? Ans. 6136.

20. How many pints in 797 pecks?

Ans. 12752.

## REDUCTION ASCENDING.

79. Example.—Reduce 856347 farthings to pounds, &c.

4)856347.

12)2140864d.

20)17840s, 61d.

£892 0s. 61d. = 856347 farthings.

EXPLANATION.—We divide the farthings by 4, because every four farthings are equal to one penny, and it is evident that what remains after taking away four farthings as often as possible from the farthings must be farthings. We thus obtain 856347 farthings, equal to 214086 pence and 3 farthings. Then we divide the pence by 12, because every 12 pence are equivalent to one shilling, and what remains after taking 12 pence as often as possible from the pence must be pence. We thus ascertain that 214086 pence and 3 farthings are equal to 17840 shillings and 6 pence 3 farthings. Lastly we divide 17840 shillings by 20, because every 20 shillings are equal to one pound.

By this process we have reduced 856347 farthings to £892 0s. 63d.

From the above example and solution we deduce the following-

Divide the given number by that number which it takes of the given denomination to make one of the next higher. Set down the remainder, if any, and proceed in the same manner with each successive denomination till you come to the one required. The last quotient, with the several remainders annexed, will be the answer required.

EXERCISE 6.

1. Reduce 32756 farthings to pounds, shillings, and pence. Ans. £34 2s. 5d.

2. Reduce 23547 troy grains to pounds, &c. 7

Ans. 4 lb. 1 oz. 1 dwt. 3 grs.

3.	Reduce 397024 yards to miles, furlongs, &c.
	Ans. 225 m. 4 fur. 26 r. 1 yd.
4.	How many hours are there in 28635 seconds?
	Ans. 7 h. 57 min. 15 sec.
5.	How many cwt., qrs., and pounds in 1666 pounds?
	Ans. 16 cwt. 2 grs. 16 lb.
R	How many cwt., &c. in 1491 pounds?
2	Ans. 14 cwt. 3 qrs. 16 lb.
17 .	How many pounds troy in 115200 grains? Ans. 20.
	How many pounds in 107520 oz. avoirdupois? Ans. 6720.
0.	now many pounds in 101020 oz. avoirdupois 1 Jus. 6120.
9.	How many cubic feet, &c. in 1674674 cubic inches?
	Ans. 969 feet, 242 inches.
10.	How many yards in 767 Flemish ells?
	Ans. 575 yards, 1 quarter.
11.	How many leagues in 183810 feet?
	Ans. 11 lea. 1 m. 6 fur. 20 rd.
12.	How many cubic yards in 138297 cubic inches?
	Ans. 2 c. yds. 26 ft. 57 in.
12	How many cords of wood are there in 67893 cubic feet?
10.	Ans. 530 cords, 53 cub. ft.
14	In 3561829 seconds, how many weeks?
13.	Ans. 5 wks. 6 dvs. 5 h. 23 min 49 gec

15. In 1597 quarts, how many bushels?

Ans. 49 bushels, 3 pks. 1 gal. 1 qt. 16. In 1000 cord-feet of wood, how many cords? Ans. 125 cords.

17. In 10,000" how many degrees? Ans. 2° 46' 40"

18. In 70,000 square links, how many square chains?

Ans. 7 square chains.

19. In 11521 grains apothecaries' weight, how many pounds? Ans. 2 lbs. 0 3 0 3 0 9 1 gr.

20. In 26025 square feet, how many roods? Ans. 2 r. 15 sq. p. 17 sq. yds. 8 sq. ft. 36 sq. in.

# REDUCTION OF THE OLD CANADIAN CURRENCY TO THE NEW OR DECIMAL CURRENCY.

80. Example.—Reduce £76 14s. 103d. to cents.

£76×400 30400 cents. 280 " 148.× 20

EXPLANATION.—We multi-

14s.× 20 = 280 "

10id.=43 far. × 5 - 12 = 17ii"

276 14s. 10id. = 30697ii cts. | by 20, because each shillings by 5 and divide the result by 12, because each shillings is equal to 20 cents; and lastly we multiply the number of farthings by 5 and divide the result by 12, because each shillings is equal to 20 cents; and lastly we multiply the number of farthings in the pence and farthings by 5 and divide the result by 12, because each shillings in equal to 20 cents. farthing is equal to A of a cent.

That each farthing is equal to A of a cent is evident from the fact that

48 farthings (or one shilling) are equal to 20 cents; or 12 farthings equal 5 cents, or one farthing equal  $\frac{A}{A}$  of a cent.

From the above example and solution we deduce the following—

RULE.

Multiply the pounds by 400, the shillings by 20, and take fivetwelfths of the number expressing how many farthings there are in the given pence and farthings. Add the three results together and their sum will be the number of cents required.

Consider the last two figures as cents, and the result will be

dollars and cents.

Note.—We take five-twelfths of the farthings by multiplying them by five and dividing the result by twelve.

#### EXERCISE 7.

1. How many cts. are there in £3 7s. 11d.? Ans. 13421 cts.

2. How many dollars are there in £29 18s. 3 d.?

- Ans. 11965 cents, or \$119:65 cents. are there in 111d.?

  Ans. 181 cents.
- 3. How many cents are there in 11½d.? Ans. 18½ cents
  4. How many dollars and cents are there in £69 15s. 6d.?
- Ans. 27910 cents, or \$279.10.

  5. How many dollars and cents in 18s. 84d.? Ans. \$3.741

6. How many dollars and cents in £17 16s. 52d.?

- Ans. \$71.297.
- 7. How many dollars and cents in £87? Ans. \$348.00.
- 8. How many dollars and cents in 15s. 11 $\frac{1}{4}$ d.? Ans. \$3·19 $\frac{7}{\sqrt{4}}$ s. How many dollars and cents in £16 6s. 2d.? Ans. \$65·23 $\frac{1}{4}$ .
- 10. Reduce £2 9s. 11d. to dollars and cent3. Ans. \$9.981.

## RECAPITULATION.

I. Science is a collection of the general principles or leading truths of any branch of knowledge systematically arranged.

II. Art is a collection of rules serving to facilitate the

performance of certain operations.

III. The rules of art are based upon the principles of science.

IV. Arithmetic is both a science and an art.

V. The science of arithmetic discusses the properties of numbers and the principles upon which the elementary operations of arithmetic are founded.

VI. The science of arithmetic is called Theoretical

Arithmetic.

VII. The art of arithmetic is called Rractical Arithmetic.

VIII. Practical Arithmetic is the application of rules based upon the science of numbers, to practical purposes, as the solution of problems, &c.

IX. Numbers are expressions for one or more things of

the same kind.

X. Unity, or the unit of a number, is one of the equal

things which the number expresses.

XI. Numbers are divided into two classes, viz.: simple or abstract numbers; and applicate, concrete, or denominate numbers.

XII. An applicate, concrete, or denominate number is a number whose unit indicates some particular object or thing.

XIII. A simple or abstract number is a number whose unit indicates no particular object or thing.

XIV. Numbers may be expressed either by words or by

characters.

XV. The expression of numbers by characters is called Notation.

XVI. The reading of numbers, expressed by characters, is called *Numeration*.

XVII. The characters we use to express numbers are either letters or figures.

XVIII. The expression of numbers by letters is called

Roman Notation.

XIX. The expression of numbers by figures is called Arabic Notation.

XX. In the Roman Notation only seven numeral letters

are used, viz.: I, V, X, L, C, D, M.

XXI. When these letters stand alone, I denotes one, V five, X ten, L fifty, O one hundred, D five hundred, M one thousand.

XXII. All other numbers are expressed by repetitions and combinations of these letters.

XXIII. In combinations of these numerical letters, every time a letter is repeated its value is repeated; also when a letter of a lower value stands before one of a higher, its value is to be subtracted; but when a letter of a lower comes directly after one of a higher value, its value is to be added.

ules ses,

r. I.

s of

ual

ple mi-

s a

ose

by led

ers,

are

led

led

ers

M,

ns

ry a ts er be XXIV. A bar or dash written over a letter or combination of letters, multiplies the value by one thousand. As we have already a character for one thousand, viz., M, and car, by repeating it, express two or three thousand, we do not dash the I, or combinations into which it enters,

XXV. Anciently, IV was written IIII; IX was written VIIII; XL was written XXXX, &c.; D was written ID, and M was written CID. Affixing C to ID increases its value ten times—thus ID=500; IDD=5000; IDD=50000, &c. Prefixing C and affixing D to CID increases its value also ten times, thus CID=1000; CCIDD=10000; CCCIDD=100,000, &c.

XXVI. The figures or characters used in the Arabic or common system of notation are 1, 2, 3, 4, 5, 6, 7, 8, 9, 0, one, two, three, four, five, six, seven, eight, nine, zero.

XX. The first nine of these characters are called significating gures, because each one has always some value or denotes some number. They are also called digits (Lat. digitus, "a finger"), from the almost universal habit of counting on the fingers.

XXVIII. The last or zero is called a cipher or naught, because it is valueless, that is, stands for nothing. It is not, however, useless, since it serves to give the significant

figures their appropriate places.

XXIX. When the 0 stands to the left of an integral number, or to the right of a decimal, i. e. when it does not come between the decimal point and some significant figure, it is both valueless and useless.

XXX. The digits 1, 2, 3, &c. standing immediately to the left of the decimal point expressed or understood, are called simple units, or units of the first order.

XXXI. The decimal point is a small dot or point, used

to indicate the position of the simple units.

XXXII. The digits 1, 2, 3, &c. standing one place to the left of the simple units, are called tens, or units of the second order to the left. When they stand one place to the right of the simple unit, they are called tenths, or units of the second order to the right.

XXXIII. The digits 1, 2, 3, &c. when standing two places to the left of the simple unit, are called hundreds, or units of the third order to the left. When standing two places to the right, they are called hundredths, or units of the third order to the right, &c.

XXXIV. Commencing at the simple units and proceeding to the left, we have units of the first order or simple units; next, units of the second order or tens; next, units of the third order or hundreds; next, units of the fourth order or thousands; next, units of the fifth

order or tens of thousands, &c.

XXXV. Commencing at the simple units and proceeding to the right, we have units of the first order or simple units; next, units of the second order or tenths; next, units of the third order or hundredths; next, units of the fourth order or thousandths; next, units of the fifth order or tenths of thousandths, &c.

XXXVI. Each digit has two values, viz.: a simple or

absolute value, and a local or relative value.

XXXVII. The simple or absolute value of a digit is the value it expresses when simply considered as representing a certain number of repetitions of the digit one.

XXXVIII. The local or relative value of a digit is the value it expresses when considered as occupying a certain

position with reference to the decimal point.

XXXIX. The ratio of one number to another is the relation which one bears to the other with respect to magnitude, when the comparison is made by considering, not by how much the one is greater or less than the other, but what number of times it contains it, or is contained in it.

XL. When several numbers, or groups of units, are so arranged that the second and third have the same ratio to one another as the first and second, and the third and fourth the same ratio as the second and third, &c.,—they (the numbers or groups of units) are said to have a common ratio.

XLI. The common ratio of our system of numbers is 10—by saying which we merely mean that the different orders increase or decrease from one another in a ten-fold

two eds.

ling nits

pror or

ens;

nits fifth

ling

its; the

rder

s of

e or

the

ting

the tain

ore-

ide,

row

hat

80

b to

rth the

on

is

nt old

Ł

SECT. I.]

proportion. i. e. that 10 units of any one order make one unit or the next higher, and vice versa.

XLII. A system of numbers is called a binary, ternary, quaternary, quinary, senary, septenary, octenary, nonary, denary, &c. system, according as two, three, four, five, six, seven, eight, nine, or ten is the common ratio of the orders. Ours is a denary or decimal system.

XLIII. To facilitate the reading of a number we divide it into periods of three places each, by placing separating points after every third figure right and left of the decimal point.

XLIV. The periods to the left of the decimal point are units, thousands, millions, billions, trillions, &c. periods to the right of the decimal point are thousandths. millionths, billionths, trillionths, &c.

XLV. The lowest order used in any reading, whether it be thousands, units, hundredths, tenths of thousandths. hundredths of millionths, &c., gives the name or denomination to the part or whole of the number used in the read-

XLVI. Numbers to the left of the decimal point are integers or whole numbers; those to the right of the decimal point are called decimals.

XLVII. A number is multiplied by 10 every time the decimal point is moved one place to the right, and divided by 10 every time the decimal point is moved one place to the left. Thus, moving the decimal point two, four or six places, either multiplies or divides the number by 100, 10,000, or 1,000,000, according as we move it to the right or to the left.

XLVIII. A number may be read in several ways by changing the nature of the simple unit. Thus the number 576.24 may be read:

<sup>1</sup>st. Five hundreds, seven tens, six units, two tenths, and four hundredths.
2nd. Fifty-seven tens, six units, two tenths, and four hundredths.
3rd. Five hundred and seventy-six units, two tenths, and four hundredths.
4th. Five thousand, seven hundred and sixty-two tenths, and four

<sup>5</sup>th. Fifty-seven thousand, six hundred and twenty-four hundredths. 6th. Five hundred, and seven thousand, six hundred and twenty-four hundredths.

7th. Fifty-seven tens, and six hundred and twen four hundredths.
8th. Five hundred and seventy-six units, and twenty-four hundredths.
9th. Fifty-seven tens, sixty-two tenths, and four hundredths.
10th. Five hundreds, seven hundred and sixty-two tenths, and four hundredths. &c.

#### Exercise 8.

## MISCELLANEOUS PROBLEMS.

- 1. Reduce 6789634 links to acres, and prove by reducing the result to links.
  - 2. Read 67845398678904 and 5900704080040000.00060604.
  - 3. Set down 4769 in Roman numerals.
  - 4. Make 42986 ten thousand times greater.
  - 5. Reduce £16 18s. 61d. Old Canadian Currency to Dollars and Cents.
  - 6. Read LXXVMMOMXCI.
  - 7. Write down, in Arabic numerals, six hundred and five billions, seventy thousand and sixteen, and nine millionths.
    - 9. Make 469789 one hundred times greater.
  - 7. Read the number 6798 in all the ways it can be read. (See Recapitulation XLVIII.)
    - 10. Divide 69800463 by one million.
    - 11. Divide 8439 by ten thousand.
    - 12. Multiply 6789 by one hundred thousand.
    - 13. Multiply 60432986 by ten millions.
  - 14. Write down one quadrillion one billion one thousand and one, and one trillionth.
  - 15. Write down seven thousand six hundred and nine tenths of millionths.
    - 16. Read 90807060504030 and

# 4004040400400000060432.01010203040508

- 17. Reduce 6789463 inches to acres, and prove by reducing the result to inches.
  - 18. Reduce 617 cord-feet of wood to cords.
  - 19. Reduce 91867 cubic feet of wood to cords.

.

to 2

and befo

2

30

2

the the of t call deri

yard
is m
sions
Wha
Impe
our

that

poun barle Wha inche poun Unit

year, grees and quire there

-of

four

the

ollars

d five s.

(See

usand

tenths

ducing

20. Write down 718, 614, 499, 999, 8643, 96149, 163986, and 444444 in Roman numerals.

- 21. Read CCCXXXIII, MCMLXXXIX, and MI.
- 22. Read 6129 in as many ways as it can be read.
- 23. Give all the readings of 634986.
- 24. Give all the readings of 19.639.

SECT. I.1

- 25. Reduce 18s. 9\d.; £6 2s. 11d.; 3s. 7d.; and £189 7s. 4\d. to dollars and cents.
- 26. Give all the readings of the number \$69.863 Federal money.
  - 26. Give all the readings of 9 bush, 3 pk. 1 gal. 3 qts. 1 pt.
- 28. Were the years 1693, 1856, 1728, 1549, 867, 444, 1600, and 927, leap years or not? If not, how many years after or before leap year?
  - 29. How many days from this to the 17th of next March?
- 30. Answer the following questions: What is the meaning of the symbols £ s. d. and q.? In the expression "16," what does the long mark (/) represent? What is the derivation of the word sterling? Why are the pound and guines so called? What is the derivation of the sign \$? What is the derivation of the words "grain," pennyweight," "ounce," and "inch"? What is a "carat"? What is a square? thow that a square yard contains 9 square feet. Show that a cubic yard contains 27 cubic feet. What is a cubic yard? What is meant by a ton of round timber? What must be the dimensions of a pile of wood in order that it shall contain a cori? What is meant by a cord-foot? What are the dimensions of the Imperial bushel?—of the Winchester bushel? Which of these is our standard? Which that of the United States? How many pounds of wheat go to the bushel?--of rye?--of oats?--of barley?—of peas?—of beans?—of buckwheat?—of Indian corn? What is our standard for liquid measure? How many cubic inches of water are there in the Imperial gillon? How many pounds Avoirdupois? What are the standard gallons of the United States? Explain why a day is added to every fourth year. What is the origin of the divisions of the circle into degrees and signs? What is the derivation of the terms "minute" and "second"? How many sheets of paper are there in a quire? How many quires in a ream? How many pounds are there in a barrel of flour? What is the recening of folio?-of 4.0 or quarto?—of 8vo or octavo?—of 12mo or duodecimo? -of 16mo?-of 18mo?

# QUESTIONS TO BE ANSWERED BY THE PUPIL.

NOTE.—Numbers in Roman numerals, thus, XVI, refer to the articles in the recapitulation: those in Arabic numerals, thus, 16, refer to the numbered articles of the Section.

2. What is art? (II.)

(VIII.)

Is arithmetic a science or an art?

(IV.) What is the science of arithm tie

called? (VI.)
What is practical arithmetic?

10. What is the unit of a number ?(X)12. What are applicate or denominate numbers? (XII.)

14. By how many methods may numbers be expressed? (XIV.)

What is science? (I.)
 Upon what are the rules of art based? (III.)

based? (III.)

5. What are the objects of the science of arithmetic? (V.)

7. What name is given to the art of arithmetic? (VII.)

9. What are numbers? (IX.)

11. How many classes of numbers are there? (XI.)

18. What are simple or abstract numbers? (XIII.)

15. What is Notation? (XV.)

16. What is Numeration? (XVI.)

17. What characters do we use to exp

16. What is Numeration? (XVI.)
17. What characters do we use to express numbers? (XVII.)
18. What is Roman Notation? (XVIII.)
19. What is Arabic Notation? (XIX.)
20. What numeral letters are used in Roman Notation? (XX.)
21. What is the value of each of these letters when standing alone? (XXI.) 23. How are all other numbers expressed in Roman Notation? (XXIL.)
23. In combination, when a letter is repeated, what does it indicate?

(XXIII.)
When a letter of a lower is placed before one of a higher value, what

35. When a letter of a lower is placed after one of a higher value, what does it indicate? (XXIII.)
36. What effect has a bar or dash written over a letter or expression? (XXIV.)

27. How do we always write 1000, 2000, 3000? (XXIV.

28. Why do we not dash the I or expressions into which it enters? (XXIV.)
29. How were four, nine, forty, &c., anciently written? (XXV.)
30. How were 500 and 1000 anciently written? (XXV.)

31. How were the expressions Io and CIo increased in value in ten-fold proportion? (XXV.)

32. What are the characters used in Arabic or Common Notation? (XXVI.)
33. What are significant figures, and why are they so called? (XXVII.)
34. What are digits, and why are they so called? (XXVII.)
35. Why is 0 called "cipher" or "naught"? (XXVIII.)
36. Is the cipher of any value? Is it of any use? (XXVIII.)
37. When is the cipher or 0 both valueless and useless? (XXIX.)
38. When are digits called simple vents or units of the first order? (XXVI.)

38. When are digits called simple units or units of the first order? (XXX.)

39. What is the decimal point? (XXXI.)
40. When are digits called tens or units of the second order to the left? (XXXII.)

11. When are digits called tenths or units of the second order to the right? (XXXII.)

When are digits called hundreds, thousands, hundredths, thousandths,

hisme the different orders to the left of the decimal point, to the

Anne the different orders to the left of the decimal point, the them. (XXXIV.) (XXXV.)

44. How many values has each digit? What are they? (XXXVI.)

45. What is the simple or absolute value of a digit? (XXXVII.)

46. What is the local or relative value of a digit? (XXXVIII.)

47. What is meant by the ratio one number bears to another? (XXXIX.)

48. What is meant by a common ratio? (XL.)

SECT

49. W 50.

51.

55. H

58.

59. 60.

ng Inw Gids 64. 63. 64.

> pī th equ

1/7 68. 69.

the 70. Ho res 72. Ho

CW Ho 74. 75. Ho res

res to WE 77. Ho

qu

(60 Wh Ho 79. an

80. W 81. Ho

Ho 82.

[SECT. I.

PIL.

erticles in the num-

or an art?

rithm tie ithmetic ?

mber?(X)r denomi-

may num-

e? (XXI.) indicate ?

alue, what alue, what

19(XXIV.)

(XXIV.)

n ten-fold (XXVI.) XVII.)

? (XXX.)

the left ? he right?

isandths, -to the

XXIX.)

49. What is meant by saying that 10 is the common ratio of our system of numbers? (XLI.)
50. What name is given to a system having 10 for its common ratio?—to one having 6?—to one having 3?—to one having 12?—to one having 7? (XLII.)

51. Why are periods used? How many places are there in each period? (XLIII.)

52. Name the periods right and left of the decimal point. (XLIV.)
53. What order gives the name or denomination to the number read?
(XLV.)
54. What are integers? What are decimals? (XLVI.)
55. How does it affect a number to remove the decimal point to the right?
How to remove it to the left? (XI.VII.)

56. How may a number be read in several ways? (XLVIII.)
57. When figures are written thus, 673:32 what does the notation imply?

(74.)
58. When figures are written thus, 6d. 23h. 16 min. 37 sec., what does the notation imply? (75 and 76.)
59. What is Reduction? (77.)
60. Into what two parts is Reduction divided? (77.)
6'. What is Reduction Descending? Give an example. (77.)
6'. What is Reduction Descending. (78.)
6'. What is Reduction Descending. (78.)
6'. What is Reduction Ascending. (78.)
6'. What are the denominations of Sterling money? Give the table: (54.)
66. How are pounds, shillings, and pence reduced to farthings? Give the process and the reason for each step. (54 and 78) (Answer this and similar succeeding questions after the fo'lowing model.) We multiply the pounds by twenty, and add in the shillings because each pound is equal to twenty shillings. We multiply the shillings by twelve and add in the pence, because each shilling is equal to twelve pence. And lastly, we multiply the pence by four and add in the farthings, because each penny is equal to four farthings.
67. What are the denominations of Federal money? Give the table. (55.)
68. What are the denominations of Canadian money, old currency? Give the table. (55.)

the table. (56.)
69. What are the denominations of Canadian money, ne a currency? Give

the table. (57.)
70. How is Old Canadian Currency reduced to New? Give the process and

71. What are the denominations of Avoirdupois weight? Give the table. (58)
72. How many pounds are there in the new cwt.? How many in the old cwt.? (58.)

cwt.? (58.)
73. How are tons reduced to drams? (58 and 78.)
74. What are the denominations of Troy weight? Give the table. (59.)
75. How are grains Troy reduced to pounds Troy? Give the process and reason for each step. (59 and 79.) (Answer this and succeeding similar questions after the following model.) We divide the grains by 24, because every 24 grains are equal to one pennyweight. We divide the resulting pennyweights by 20, because every 20 pennyweights are equal to one ounce. And lastly, we divide the resulting ounces by 12, because every 12 ounces are equal to one pound.
76. What are the denominations of Apothecaries weight? Give the table (60.)

76. What are the denominations of Apothecaries' weight? Give the table. (60.)
77. How are pounds, ounces, &c., Apothecaries' weight reduced to grains?
(60 and 78.) Answer as in question 66.
78. What are the denominations of Long measure? Give the table. (61.)
79. How are lines reduced to leagues? (61 and 79): Answer after model in question 75.

80. What are the denominations of Square measure? Give the table. (62.)
81. How are square miles reduced to square inches? (62 and 78). Answer

82. How are links reduced to acres? (63 and 79.) Answer after model,

d

01

- 83. What are the denominations of Solid measure? Give the table (64.)
  84. How are cubic inches reduced to cubic feet? (64 and 79.)
  85. How are cubic feet of wood reduced to cords? (64 and 79.)
  86. What is a cord-foo! (64.)
  87. What are the denominations of Cloth measure? Give the table. (65.)
  88. How are English ells reduced to inches? (65 and 78.) Answer after model.
  89. What are the denominations of Dry measure? Give the table. (66.)
  90. How are pints reduced to chaldrons? (66 and 79.) Answer after model.
  91. What are the denominations of Liquid measure? Give the table. (67.)
  92. How are tuns reduced to gills? (67 and 78.) Answer after model.
  93. What are the denominations of Time measure? Give the table. (68.)
  94. How are econds reduced to years? (68 and 79.) Answer after model.
  95. Name the months and the number of days in each. (68.)
  96. What is the Solar year and its length?—the Sidereal year and its length?—the Civil year and its length?—the Sidereal year and its length?—the Civil year and its length? (68.)
  97. How can we ascertain whether any given year be Leap year? (69.)
  98. Show that the unit of time is the basis of the units of length, mass or especity, and weight. (71.)
  99. What are the denominations of Circular measure? Give the table. (72.)
  100. Upon what does the length of a degree depend? (72.) How are degrees reduced to seconds? (72 and 78.)

## SECTION II.

## FUNDAMENTAL RULES.

1. Arithmetic may be divided into four parts:-

1st. The Arithmetic of Whole Numbers, or that which treats of the properties of entire units.

2nd. The Arithmetic of Fractions, or that which treats

of the parts of units.

3rd. The Arithmetic of Ratios, which treats of the relations of numbers, whether integral or fractional, to each other and to the unit 1.

4th. The Application of Arithmetic to practical and

useful purposes.

2. The Arithmetic of Whole numbers includes Addition, Subtraction, Multiplication, Division, Involution, Evolution, &c.

3. The Arithmetic of Fractions may be divided into

1st. Vulgar or Common Fractions, in which the unit is divided into any number of equal parts.

2nd. Decimal Fractions in which the unit is divided

according to the scale of ten.

4. The Arithmetic of Batios relates to the comparison of numbers with respect to their quotients, and embraces Proportion and Progression.

le." (65.)

BOTI II.

(64.)

le. (65.) r model. (66.) r model. ole. (67.) lel. model.

69.) mass or

blę.(72.) degrees

which treats

he relato each

al and

Addilution,

d into

unit is

livided

ison of braces

5. Addition, Subtraction, Multiplication, Division, are called the fundamental rules, or ground rules of Arithmetic, because all the other operations of Arithmetic are performed by means of them.

Whatever operations we may perform upon a number, we can only either increase it or diminish it. If we increase it, the process belongs to addition; if we diminish it, to subtraction. All the rules of Arithmetic are therefore resolvable into these two. Multiplication is only a short method of performing a peculiar kind of addition, in which the addends are all the same; and division is merely an abridged method of performing a particular kind of subtraction, in which the same quantity is to be taken away from

a given number as often as possible.

When any number of quantities, either different, or repetitions of the same, are united together so as to form but one, we term the process, simply, "Addition." When the quantities to be added are the rame, but we may have as many of them as we please, it is called "Multiplication;" when they are not only the same, but their number is indicated by one of them, the process belongs to "Involution." That is, addition restricts us neither as to the kind, nor the number of the quantities to be added; multiplication restricts us as to the kind, but not the number; involution restricts us both as to the kind and number. All, however, are really comprehended under the same rule—addition.

# ADDITION.

7. The sum of two or more numbers is a number which contains as many units, and no more, as are found in all the given numbers.

8. Addition is the process of finding the sum of two or

more numbers.

9. The quantities to be added together are called addends, and the result of the addition is called the sum of the addends.

10. Only those quantities can be added which have the same unit, or, in other words, which are of the same denomination.

Thus it is evident that 6 days and 7 miles cannot be added, since the result would neither be 13 days nor 13 miles; nor can 5 shillings and 3 pence be added, as the result would neither be shillings nor pence. Similarly, we cannot add units and tens, or tenths and hundredths, or units and sevenths, &c.

11. Hence, in writing down the addends preparatory to adding, we must be careful to set units of the same denomination in the same vertical column, i.e. units under units, tens under tens, hundreds under hundreds, &c.; shillings under shillings, pence under pence, &c.; miles under miles, furlongs under furlongs, rods under rods, &c.

EXERCISE 9. Apples. Shillings. Addends 3 Addends Sum of Addends 7 Sum of Addends 24 (3) Addends Sum of Addends 30 (5) (6) (8) (9) (10) pence, sevenths, horses. tens. millionths. 8 23 30 35 12. Let it be required to add together 987 and 689. Í. III. IV. 987 987 987 987 987 689 689 689 689 689 1500 160 16 16 160 1500 160 16 16 1500 70 1000 600 600 70

600

1000

1000

1676

tory ame ader ahil-

ader

(11)

iles.

7

1234

r. II:

EXPLANATION.—We place the given numbers, 987 and 680, under each other, according to (11) and draw a line to separate the addends from the

It is manifest that so long as we add the units of the several orders it is quite immaterial whether we commence at the highest, at the lowest, or at an intermediate denomination.

In the first of the above operations we have commenced continually at the highest or left-hand order. The hundreds added make 15 hundreds or one thousand and five hundred, which we set down; the tens added make 16 tens, equal to 1 hundred and 6 tens, and the units added, make 16 units, equal to 1 ten and 6 units, all of which we set down in their appropriate columns.

Next considering the partial sums 1500, 160, and 16, as so many new addends, we proceed similarly with them and obtain a new set of partial sums, viz: 1000, 600, 70, and 6. But, from the principles of notation (Sec. 1). these last numbers (i. c. 1000, 600, 70 and 6) may be written in one line, thus, 1676, which therefore is the sum of the addends 967 and 669.

In (II), (III), (IV), (V) the same result is obtained by a slightly different process.

In (II), (IV), (V) the same result is obtained by a singitly different process.

In (II) we have commenced at the tens, and in (III), (IV), and (V) at the units or lowest order. (IV) is simply (III) with the unnecessary o's omitted.

(V) is (IV) somewhat modified as follows:—9 units and 7 units make 18 units, equal to 6 units, which we set down, and one ten which we carry to the next column or column of tens; 1 ten and 8 tens make 9 tens, and 8 tens make 17 tens, equal to 7 tens, which we set down, and 1 hundred, which we carry to the column of hundreds; 1 hundred and 6 hundreds make 7 hundreds; and 9 hundreds make 16 hundreds, equal to 6 hundreds and 1 thousand, both of which we set down.

13. From (I), (II), and (III), it is manifest that it is as legitimate to commence at the lowest denomination as at the highest: and from (IV) and (V), that it is most convenient to commence at the lowest denomination.

14. From (V) we learn that when we have obtained the sum of the units, in any column, we reduce it to the next higher denomination, and, setting down the remainder under the column added, carry the units of the next higher denomination to their proper column.

15. The reasoning in (12), (13) and (14) applies to any numbers whatever, whether abstract or denominate, and from it, for addition, we deduce the following general-

Write down the numbers so that units of the same denomination shall fall in the same column (Arts. 10 and 11).

Draw a line beneath the addends (Art. 12).

Add up the units of the lowest denomination and divide their sum by so many as make one of the denomination next higher (Arts. 13 and 14).

Set down the remainder and carry the quotient to the next higher denomination (Art. 14),

Proceed in the same manner through all the denominations to the

16. We commence at the lowest order or tenths of thousandths. There being nothing to add to the 9 tenths of thousandths we simply set down the 9 in its appropriate column. Next we add the thousandths, thus:—I thousandths and 8476 6 thousandths are 8 thousandths and 4 thousandths and 98462 1 hundredth. The 2 thousandths we write down in its own column and carry the hundredth to the column of hundredths. Next we add the column of hundredths in a 1881 9839 in a 1 hundredth (carried) and 6 hundredths make 7 hundredths and 9 hundredths make 16 hundredths which are equal to 8 hundredths and 6 hundredths make 28 hundredths which are equal to 8 hundredths and two tenths. We set down the 6 hundredths and carry the two enths to the next column or column of tenths. Adding the finths we find their sum to be 39 tenths, equal to 9 tenths, which we set down and 3 units which we carry. The simple units added make 41 units, equal to 1 unit, which we set down and 4 tens which we carry; the tens added make 38 tens, equal to 8 tens and 3 hundreds; the hundreds added (with the three hundreds we carry) make 18 hundreds, or 8 hundreds, and 1 thousand, both of which we set down in their proper columns. columns.

TT. We commence as in (16) with the lowest denomination, which, in KAMPLE. this example, is cents. 89 cents and 42 cents and 56 cents and 39 cents, added, make 276 cents. But every 1136 160 cents make one dollar, 276 cents at the seferog equal to 3 dollars and 76 cents. The 76 cents we set down in 1189 their proper place and carry the 2 dollars to the column of EXAMPLE.

Druss Grad : 5 die 18. Example.-Add together £52 17s. 34d., £47 5s. 61d., and £66 14s. 21d.

2166 17 Of rum.

and i make three farthings, which, with 1, make 6 farthings; these are equivalent to one of the next denomination, or that of pence, to be carried, and two of the present, or one half-penny, to be set down. I penny (earried) and 2 are 3, and 6 are 9, and 3 are 12 pence—equal to one of the next denomination, or that of shillings, to be carried, and no pence to be set down; we therefore put a cipher in the pence place of the sum. I shilling (carried) and 14 are 15, and 5 are 20, and 17 are 37 shillings—equal to one of the next denomination, or that of pounds, to be carried, and 17 of the present, or that of shillings, to be set down. I pound and 6 are 7, and 7 are 14, and 2 are 16 pounds,—equal to 6 units of pounds, to be set down, and 1 ten of pounds to be carried; I ten and 6 are 7 and 4 are 11 and 5 are 16 tens of pounds, to be set down.

When the addends are very numerous, we may divide them into two or more parts by horizontal lines, and, adding each part separately, may afterwards find the amount of all the sums.

ARTS. 16-19.]

ons to the

ths. There housendths to column and the said and the are and the are and the and lown in its he column undredths, the make 7 redths, and hundredths the 6 hon-n of tenths. o 9 tenths, mits added which we dreds; the hundreds,

ats and 56 But every efore equal et down in e column of

heir proper

59. 61d.,

of the next to be set l to one of of the pre-7, and 7 set down, and 5 are

ato two:or may after-

Or, in adding each column, we may put down an asteriak, thus, as often as we come to a quantity which is at least equal to that number of the denomination added which is required to make one of the next—carrying forward what is above this number, if anything, and putting the last remainder, or —when there is nothing left at the end—a cypher under the column;—we carry to the next column one for every asteriak. Using the same example.

.21 404 10

2 pence and 4 are 6, and 2 are 8, and care 17 pence—equal to 1 shilling and 5 pence; we put down a dot or an asterisk and carey 5, 5 and 2 are 7, and 4 are 11, and 9 are 20 pence—equal to 1 shilling and 8 pence; we put down a dot or an asterisk and carry 8. 8 and 2 are 10 and 6 are 16 pence equal to 1 shilling and 4 pence; we put down a dot and carry 4. 4 and 4 are 8 and 2 are 10—which being less than 1 shilling, we set down under column of pence to which it belongs, &c. We find on adding them up, that there are three dots; we therefore carry 3 to the column of shillings. 8 shillings and 8 are 11, and 4 are 15, and 4 are 19, and 3 are 22 shillings—equal to 1 pound and 2 shillings: we put down a dot and carry 2. 2 and 17 are 19, &c.

Care is necessary, lest the dots, not being distinctly marked, may be considered as either too few or too many. This method though now but little used, seems a convenient one.

little used, seems a convenient one.

#### PROOF OF ADDITION.

19. FIRST METHOD.—Go through the process again, beginning at the top and adding downwards.

This method of proof is merely doing the same work twice, in

a slightly different manner.

SECOND METHOD.—Separate the addends into two parts. Add each part separately, in the usual touy, and then add their sums. If the last sum is the same as that found by the first addition, the work may be presumed to be correct.

This method of proof is founded on the aziom that "the

whole is equal to the sum of all its parts,"

Example.—Find the sum of 509267, 235809, 72910, and 83925.

OPERATIO	on. Proof by 8	ECOND METHOD.
509267	509267,	. , 72910
235809	235809	83925
72910		
83925	Partial sums 745076	156835
· · · · · · · · · · · · · · · · · · ·	First partial sum	745076
Sum 901911	Second partial sum	156835

Proof.... 901911

. An e		. 4. 6.			
(1)	(2)	(3)	(4)	(5)	(6)
Dollars.	Bushels.	Days.	Acres.	Dollars.	Pounds
15	. 76	765	392	5832	98764
26	48	381	446	- 8907	8753
18	59	872	872	4671	76
61	81	315	969	6789	9889
1			(		
120	264	2333	2679	26199	117482
1		17-	-30 /		

The sum of the numbers in each row of the following table, whether taken vertically or horizontally, or from corner to corner, is 24156. Let the pupil be required to make these 24 distinct additions.\*.

2016	4212	1656	3852	1296	3492	936	3132	576	2772	216
252	2052	4248	1692	3888	1332	3528	972	3168	612	2412
2448	288	2088	4284	1728	3924	1368	3564	1008	2808	648
684	2484	324	2124	4320	1764	3960	1404	3204	1044	2844
2880	720	2520	360	2160	4356	1800	3600	1440	3240	1080
1116	2916	756	2556	396	2196	3996	1836	3636	1476	3276
3312	1152	2952	792	2592	. 36	2232	4032	1872	3672	1512
1548	3348	1188	2988	432	2628	72	2268	4068	1908	3708
3744	1584	3384	828	3024	468	2664	108	2304	4104	1944
1980	3780	1224	3420	864	3080	504	2700	144	2340	4140
4176	1620	3816	1260	3456	900	3096	540	2736	180	2376

This table is formed by multiplying the numbers in the magic square of 11 by 36.

Œ.	
ъ.	7
	K

(36) 0·87

5.273

8-127

25.63

1	
[SEC	r.II.
nd 83	925.
.0 .5	×1
5	
žņ.	1 .
(6) unds	š
8764	20
8753 76	
	. 1
9889 7482	
7482	able,
7482	able, ener, inct
7482 ing to to con 4 dist	able, ener, inct
7482 ing to to con 4 dist	able, mer, inct
7482 ing to to con 4 dist	able, mer, sinct
7482 ing to cor 4 dist 216 2412 648	able, ener, inct
7482 ng ta to con 4 dist  216  2412  648  2844	able, ener, sinct
7482 ng ta to con 4 dist  216  2412  648  2844	able, ener, iinct
7482 ing to con 4 dist 216 2412 648 2844 1080	able, mer, sinct
7482 ing to con 4 dist 216 2412 648 2844 1080	able, ener, sinct
7482 ing to cor 4 dist 216 2412 648	able, ener, iinct

2376 gic square ART. 19.]

(31)

7674

376

6:

(37)

3.785

20.766

0.253

10.004

34.808

74564

82620

(32) 5676

1567

6767

63

(38)

6034.82

712.52

85.742

57.8563

576.03 5000.0 0.37 4000.005 4712.5 427.0 8458.302 213.5 6.53712 3.7.12 0.007 2753.0   MONEY.  (45) (46) (47) (48) £ s. d. £ s. d. £ s. d. £ s. d. 4567 14 6½ 76 14 7 3767 13 11 5674 17 6½ 776 15 7½ 667 13 6 4678 14 10 4767 16 11½ 76 17 9¾ 67 15 7 767 12 9 3466 17 10¾ 51 0 10¼ 5 4 2 10 11 5 5984 2 2½ 44 5 6 3 4 3 4 11 8762 9 9   AVOIRDUPOIS WEIGHT.  (49) (50) (51) (52) cwt. qrs. lb. cwt. qrs. lb. cwt. qrs. rs. 76 3 14 476 1 24½ 447 1 7 14 2 37 2 15 756 3 21½ 576 1 6 3 3 14 1 11 767 1 16 467 1 7½ 2 128 3 15 973 1 12 428 0 0½	4	\$ 8 E.	€.	6.34			(43) (3456·5			0.0	235		• .
4712·5	-1	2	100										
6·53712 3:·12 0·007 2753·0  MONEY.  (45) (46) (47) (48)  £ s. d. £ s. d. £ s. d. £ s. d.  4567 14 6½ 76 14 7 3767 13 11 5674 17 6½  776 15 7½ 667 13 6 4678 14 10 4767 16 11½  76 17 9½ 67 15 7 767 12 9 3466 17 10½  51 0 10½ 5 4 2 10 11 5 5984 2 2½  44 5 6 3 4 3 4 11 8762 9 9   AVOIRDUPOIS WEIGHT.  (49) (50) (51) (52)  cwt. qrs. lb. cwt. qrs. lb. cwt. qrs. lb. cwt. qrs.  76 3 14 476 1 24½ 447 1 7 14 2  37 2 15 756 3 21½ 576 1 6 3 3  14 1 11 767 1 16 467 1 7½ 2  567 2 15 563 1 6 7 0		1-	en var										
MONEY.  (45) (46) (47) (48)  £ s. d. £ s. d. £ s. d. £ s. d.  4567 14 61 76 14 7 3767 13 11 5674 17 61  776 15 71 667 13 6 4678 14 10 4767 16 111  76 17 92 67 15 7 767 12 9 3466 17 102  51 0 101 5 4 2 10 11 5 5984 2 21  44 5 6 3 4 3 4 11 8762 9 9   AVOIRDUPOIS WEIGHT.  (49) (50) (51) (52)  cwt. qrs. lb. cwt. qrs. lb. cwt. qrs.  76 3 14 476 1 241 447 1 7 14 2  37 2 15 756 3 211 576 1 6 3 3  14 1 11 767 1 16 467 1 71 2  567 2 15 563 1 6 7 0		**************************************	1 22										8
(45) (46) (47) (48)  £ s. d. £ s. d. £ s. d. £ s. d.  4567 14 6½ 76 14 7 3767 13 11 5674 17 6½  776 15 7½ 667 13 6 4678 14 10 4767 16 11½  76 17 9½ 67 15 7 767 12 9 3466 17 10½  51 0 10½ 5 4 2 10 11 5 5984 2 2½  44 5 6 3 4 3 4 11 8762 9 9   AVOIRDUPOIS WEIGHT.  (49) (50) (51) (52)  cwt. qrs. lb. cwt. qrs. lb. cwt. qrs.  76 3 14 476 1 24½ 447 1 7 14 2  37 2 15 756 3 21½ 576 1 6 3 3  14 1 11 767 1 16 467 1 7½ 2  567 2 15 563 1 6 7 0				, '				9-		1	9062	76.0	53
£ s. d. £ s. d. £ s. d. £ s. d. 4567 14 6½ 76 14 7 3767 13 11 5674 17 6½ 776 15 7½ 667 13 6 4678 14 10 4767 16 11½ 76 17 9¾ 67 15 7 767 12 9 3466 17 10¾ 51 0 10¼ 5 4 2 10 11 5 5984 2 2½ 44 5 6 3 4 3 4 11 8762 9 9 AVOIRDUPOIS WEIGHT.  (49) (50) (51) (52) (52) (54) (54) (55) (55) (55) (55) (55) (55				66			NEY.	MC	V.	,			11.
4567 14 6½ 76 14 7 3767 13 11 5674 17 6½ 776 15 7½ 667 13 6 4678 14 10 4767 16 11½ 76 17 9¾ 67 15 7 767 12 9 3466 17 10¾ 51 0 10½ 5 4 2 10 11 5 5984 2 2½ 44 5 6 3 4 3 4 11 8762 9 9   AVOIRDUPOIS WEIGHT.   (49) (50) (51) (52)  cwt. qrs. lb. cwt. qrs. lb. cwt. qrs. lb. cwt. qrs. 76 3 14 476 1 24½ 447 1 7 14 2 37 2 15 756 3 21½ 576 1 6 3 3 14 1 11 767 1 16 467 1 7½ 2 567 2 15 563 1 6 7 0			8)				(47)	,	46)	. (	5) _	(48	T.
4567 14 6½ 76 14 7 3767 13 11 5674 17 6½ 776 15 7½ 667 13 6 4678 14 10 4767 16 11½ 76 17 9¾ 67 15 7 767 12 9 3466 17 10¾ 51 0 10½ 5 4 2 10 11 5 5984 2 2½ 44 5 6 3 4 3 4 11 8762 9 9   AVOIRDUPOIS WEIGHT.   (49) (50) (51) (52)  cwt. qrs. lb. cwt. qrs. lb. cwt. qrs. lb. cwt. qrs. row		d.	8:4	£	1.	d	£ s		S.	£	d.	. s.	£
76 17 9\(\frac{1}{2}\) 67 15 7 767 12 9 3466 17 10\(\frac{1}{2}\) 51 0 10\(\frac{1}{2}\) 5 4 2 10 11 5 5984 2 2\(\frac{1}{2}\) 44 5 6 3 4 3 4 11 8762 9 9   SECULTARIZED STATE OF THE STATE	1	61	17	5674	1	1	3767 13	'7 *	14	76	4 61	37 1	456
51 0 10½ 5 4 2 10 11 5 5984 2 2½ 44 5 6 3 4 3 4 11 8762 9 9  5516 14 3½  AVOIRDUPOIS WEIGHT.  (49) (50) (51) (52)  cwt. qrs. lb.	101	114	16	4767	0	1	4678 14				5 71	16 1	77
44 5 6 3 4 3 4 11 8762 9 9  5516 14 3½  AVOIRDUPOIS WEIGHT.  (49) (50) (51) (52)  wt. qrs. lb. cwt. qrs. lb. cwt. qrs. lb. cwt. qrs. 76 3 14 476 1 24½ 447 1 7 14 2 37 2 15 756 3 21½ 576 1 6 3 3 14 1 11 767 1 16 467 1 7½ 2 567 2 15 563 1 6 7 0			17		9	1							
AVOIRDUPOIS WEIGHT.  (49) (50) (51) (52)  cwt. qrs. lb. cw		2}							4	5	0 101	51 " (	5
AVOIRDUPOIS WEIGHT.  (49) (50) (51) (52)  cwt. qrs. lb. cw	14	9	9	8762	1	1	3 4	4	3		5 6	14	4
cwt. qrs. lb.     cwt.				,	IT.	IG E	DIS WE	DUP	OIR	AV	4 31	16 1	551
cwt. qrs. lb.     cwt.	*		52):	(		): }	(51		(0)	(8	2.1	(49)	
76 3 14 476 1 241 447 1 7 14 2 37 2 15 756 3 211 576 1 6 3 3 14 1 11 767 1 16 467 1 71 2 2 567 2 15 563 1 6 7 0	lb	grs.	wt.	` C1	: lb.	qrs.	cwt.				lb.	qrs.	wt.
37 2 15 756 3 21½ 576 1 6 3 3 3 14 1 11 767 1 16 467 1 7½ 2 15 563 1 6 7 0	12					1	447	241	1	476			
567 2 15 563 1 6 7 0	7			- Pa		1	576	211	3		15	<b>2</b>	37
	15	2	5 3		71	1	467	16	1	767	11	1	14
128 3 15 973 1 12 428 0 04	3	0	7	2 1 12	6	1			-				-
	14	- 4.	- A	,	.01	0	428	12	1	973	15	3	128
	e 2	-	-	h : 4	<del></del>	-	-	······································			64		
					1								
· · · · · · · · · · · · · · · · · · ·									٠.	•			

ADDITION.

(39)

0.00007

0.06236

0.0572

0.21

(34)

67674

75670

36

77

(35)

42.37

56·84 27·92

62.41

(40) 5471·3

563·47 21·502

0.0007

(33)

76746

71207

-100

56

Sta SN

# TROY WEIGHT.

(53)	(54)	(55)
lb. oz. dwt. grs.	lb. oz. dwt. grs.	lb. oz. dwt. grs.
7 0 5 9	57 9 12 14	87 3 7 12
5 6 6 7	67 9 11 11	11 12 3
9 5 6 8	66 8 - 10 5	16 14
	74 6 5 3	44 12 10 13
21 11 18 0,	12 3 5 4	67 8 9 10

# TIME.

(56)	(57)	(58)
yrs. ds. hrs. ms.	yrs. ds. hrs. ms.	yrs. ds. hrs. ms.
99 359 9 56 88 0 8 57	60, 90, 0, 50, 6, 76, 1, 57,	50 127, 7, 50 120, 9, 44
77 120 7 49	3 58	76 121 11 44
265 115 2 42	6 1, 2 0	6 47 3 41 8 9 11 17

# CLOTH MEASURE.

(59)	(60)	(61)	(62)
vds. qrs. nls.	yds, qrs. nls.	yds. qrs. nls.	yds. qrs. nls.
567 3 2 476 1 0	147 3 3 173 1 0	157 2 1 143 3 2	156 1 1 176 3 1
72 3 3	148 274	1 2	54 1 0
5 2 1	92 3 2	54 0 3	573 2 3
1122 2 2	. 11. 34 5.		44 (1

# CANADIAN MONEY:

(63) \$978:63	(64) \$ 69·42	(65) \$719:43	(66) \$9868:47
492.29	189:87	912:99	986 10
83.43	674.29	68:68	91.89
729.47	86:43	50:00	7.45
9.00	982:78	9:73	.98
\$2292:82	\$ 75.6	\$ 77.7	\$

13

10

rs. ms.

50

44

44

17

s. nls.

- 67. 0.4+74.47+37.007+75.05+747.077=934.004.
- 68.56.05+4.75+0.007+36.14+4.672 = 101.619.
- 69. 0.78 + 0.0076 + 76 + 0.5 + 5 + 0.05 = 82.3176.
- 70. 0.5+0.005+5+50+500 = 555.505.
- 71. 0.367 + 56.7 + 762 + 97.6 + 471 = 1387.667.

72. Add eight hundred and fifty-six thousand, nine hundred and thirty-three; one million, nine hundred and seventy-six thousand, eight hundred and fifty-nine; two hundred and three millions, eight hundred and ninety-five thousand, seven hundred and fifty-tyo.

Ans. 206729544.

73. Add three millions, and seventy-one thousand; four millions, and eighty six thousand; two millions, and fifty-one thousand; one million; twenty-five millions, and six; seventeen millions, and one; ten millions, and two; twelve millions, and twenty-three; four hundred and seventy-two thousand, nine hundred and twenty-three; one hundred and forty-three thousand; one hundred and forty-three millions. Ans. 217823955.

dred and seventy thousand; thirty-three thousand; eight hundred and forty-seven thousand; thirty-three thousand; eight hundred and seventy-six thousand; four hundred and ninety one thousand.

16. Add together one hundred and sixty-seven thousand; three hundred and sixty-seven thousand; nine hundred and six thousand; two hundred and forty-seven thousand; ten thousand; seven hundred thousand; nine hundred and seventy-six thousand; one hundred and ninety-sive thousand; ninety-seven thousand; and seventy-six thousand; one hundred and ninety-sive thousand; ninety-seven thousand.

#### APPLICATIONS.

1. How many miles is it from the lower end of Lake Huron to the Gulf of St. Lawrence, passing through the River St. Clair, 25 miles long; Lake St. Clair, 20 miles; River Detroit, 23 miles; Lake Eric, 250 miles; Niagara River, 34 miles; Lake Ontario, 180 miles; and the River St. Lawrence, 750 miles long?

Ans. 1282 miles.

2. The city of Toronto has a population of about 50000; Hamilton, 25000; Kingston, 15000; London, 10000; Ottowa, 10000; Montreal, 75000; and Quebec, 45000. What is the population of these seven cities taken together? Ans. 230000.

3. In the year 1856 Canada exported :—Produce of the mine, \$165000; produce of the sea, \$500000; produce of the forest, \$10000000; animals and their produce, \$2500000; agricultural products, \$15000000; manufactures and ships, \$1600000; and various other products to the amount of \$2235000. What was the total value of Canadian exports for that year?

4. A wholesale merchant sells, during the year, goods to the amount of \$11080 in Toronto; \$9427 in Galt; \$1798 in Berlin; \$16422 in Tamilton; \$7496 in Guelph; \$6429 in Woodstock; \$5297 in Chatham; and \$8426 in Goderich. Required the amount of the year's sales.

Ans. \$66376.

5. The Grand Trunk Eailway is 962 miles long, and cost \$60000000; the Great Western is 229 miles long, and cost \$14000000; the Ontario, Simcoe, and Huron is 95 miles long, and cost \$3300000; the Toronto and Hamilton is 35 miles long, and cost \$2000000. What is the aggregate length and cost of these four roads? Ans. Length, 1324 miles, and cost \$79300000.

6. The circulation of promissory notes for the four weeks ending February 3, 1844, was as follows:—Bank of England, about £21228000; private banks of England and Wales, £3480000; Joint Stock Banks of England and Wales, £3446000; all the banks of Scotland, £2791000; Bank of Ireland, £3881000; all the other banks of Ireland, £2429000; what was the total circulation?

Ans. £38455000.

occurred 4604 years before Christ; the deluge, 2348; the call of Abraham, 1921; the departure of the Israelites from Egypt, 1491; the foundation of Solomon's temple, 1012; the end of the captivity, 536. This being the year 1859, how long is it since each of these exercises.

each of these events?

2ns. From the creation, 5863 years; from the deluge, 4207; from the call of Abraham, 3780; from the departure of the Israelites, 3350; from the foundation of the temple, 2371; and from the end of the captivity, 2395.

8. Add together the following:—2d., about the value of the man sestertius; 7id., that of the denarius; 1id., a Greek obolus; 9d., a drachma; £3 15s., a mina; £225, a talent; 1s. 7d., the Jewish shekel; and £342 3s. 9d., the Jewish talent. Ans. £571 2s.

3. Add together 2 dwt. 16 grains, the Greek drachma; 1 lb.

1 oz. 1 dwt., the mina: 67 lb. 7 oz. 5 dwt., the talent.

Ans. 68 lb. 8 oz. 8 dwt. 16 grains.

10. What was the population of the British provinces in North
America in 1834, the population of Lower Canada being stated
at 549005, of Upper Canada, 336461; of New Brunswick, 152156;
of Nova Scotia and Cape Breton, 142548; of Prince Edward's

Island, 3229?; of Newfoundland, 75000?

Ans. 1287462.

11. A owes to B £567 16s. 7½d.; to C £47 16s.; and to D £56 0s. 1d. How much does he owe in all? Ans. £67. 12s. 8½d.

12. A man has owing to him the following sums: 10s. 7d.; £46 0s. 7dd.; and £52 14s. 6d. How much is the wage?

13. A merchant sends off the following quantities of butter:—47 cwt. 2 qrs. 7 lb.; 38 cwt. 3 qrs. 8 lb.; and 16 cvt. 2 qrs. 20 lb. How much did he send off in all?

Ans. 103 cwt. 10 lb.

ods to the in Berlin; oodstock; nired the s. \$66376. and cost sand cost riles long. illes long, nd cost of 29300000. our weeks England, d Wales. 3448000; f Irela 00; what 38455000. the world ; the calf

ige, 4207; parture of e temple,

om Egypt,

end of the

is it since

lue of the reek obolis. 7d., the £571 2s. ma; 1 lb.

16 grains.

s in North
ing stated
c, 152156;
Edward's
1287462.
and to D
128. 81d.
108. 7d.;
ce?
2 58. 81d.
butter:—
qrs. 20 lb.

7t. 10 lb.

14. A merchant receives the following quantities of tallow, viz:—13 cwt. 1 qr. 6 lb.; 10 cwt. 3 qrs. 10 lb.; and 9 cwt. 1 qr. 15 lb. How much has he received in all?

Ans. 33 cwt. 2 qrs. 6 lb.

15. A silversmith has 7 lb. 8 oz. 16 dwts.; 9 lb. 7 oz. 3 dwts.; and 4 lb. 1 dwt. What quantity has he?

Ans. 21 lb. 4 oz.

16. A merchant sells to A, 76 yards 3 quarters 2 nails; to B, 90 yards 3 quarters 3 nails; and to C, 190 yards 1 nail. How much has he sold in all?

Ans. 357 yards 3 quarters 2 nails.

17. A merchant in Toronto sells goods to the following amounts during the week, viz:—Monday, \$429.38; Tuesday, \$711.43; Wednesday, \$419.87; Thursday, \$1080.42; Friday, \$1304.65; Saturday, \$2498.91. Required the whole amount of the week's sales.

Ans. \$6444.66.

18. Looking over my last month's expenditure, I find that I have paid the following sums, viz:—Baker's bill, \$5.73; Butcher's bill, \$20.91; Groceries, \$12.75; Fruit, \$3.29; Rent, \$16.25; Servants' wages, \$10; Tailor's account, \$17.87; Shoemaker's bill, \$11.63; and sundries, \$9.47. Required how much I paid in all.

Ans. \$107.90.

19. Add together \$607.19; \$298.97; \$789.87; \$1723.10; and \$123.00.

Ans. \$3542.13.

20. A farmer sells seven loads of wheat, the first containing 1763 lbs., the second 1827 lbs., the third 1329 lbs., the fourth 1901 lbs., the fifth 1666 lbs., the sixth 1879 lbs., and the seventh 1185 lbs. What was the aggregate weight of the seven loads and how many bushels did they contain?

Ans. 11550 lbs. or 1921 bushels.

Note.—The bushels are found by dividing the aggregate weight by 60

lbs., the weight of one bushel.

21. Having effected an insurance on my household furniture, &c., I am required to make a detailed statement of its value. I find this to be as follows:—Carpets \$250.00, table and bed linen \$90.88, beds and bedding \$173.60, furniture \$791.23, pictures and engravings \$207.18, books \$1649.19, plate and plated ware \$307.18. Required the total value of my household furniture.

Ans. \$3469.26.

22. Toronto has a population of 45000, Hamilton 20000, Brockville 4000, Prescott 2500, Kingston 15000, Ottawa City 10000, Chatham 4000, Goderich 2000, London 10000, Port Hope 4000, Cohourg 5000, Montreal 70000, and Quebec 50000. What is the entire population of these 13 cities and towns?

Ans. 241500.

20. The pupil should not be allowed to leave addition until he can read up the columns without hesitation. For instance, in the following questions, which are inserted for the sake of practice in rapid addition, he should not be permitted to spell the columns thus, 6 and 4 are 10, and 4 are 14, and 4 are 18, and 5 are 23

100

49 (13)

m hi

de ru so

tic

&c., but should be required to read them, i. e., simply touch each digit with his pencil and name the sum, thus:—6, 10, 14, 18, 23, 31, 32, 35, 42, 43, 44, 49, 53, &c., &c.

I.	II.	III.	IV.
244658	275634	135790	123456
492327	386731	246824	786123
635425	987654	135790	456789
321465	321456	864212	123456
732849	989123	579246	788123
376731	456789	835792	459789
935746	123456	468357	123456
847963	789123	924689	789123
745143	456789	753246	456789
234561	123456	835792	123456
746874	789123	468357	789123
934746	456789	924683	456789
872345	123459	579246	123456
934756	789123	835798	789123
842345	456789	642875	456789
873456	123456	334683	123456
864530	789123	579864	789123
234673	456789	297531	456789
325871	246842	135795	871178
479234	357931	246834	936639
845645	642248	824248	248842
823456	756139	357964	525255
245734	246842	872278	736376
872475	657931	375946	875578
896731	642248	624862	473468
456841	753139	375937	934579
314567	246842	872459	894645
814563	357931	837645	123875
427881	642248	644875	767457
932768	753913	472963	875345
456345	375913	875847	874563
345634	426428	864314	375534
734734	573931	734561	937565
784564	624824	273475	475734
834756	735813	845675	698945
	*	B C and a The Co.	

## RECAPITULATION.

I. Addition is the process of finding the sum of two or more numbers.

II. The numbers to be added are called Addends.

III. The result of the addition is called the sum of the addends.

touch 0, 14,

IV. In writing numbers down preparatory to adding them, we write units under units, tens under tens, &c., because it is more convenient, since only like quantities, i. e., quantities of the same name, can be added together.

V. We draw a line under the addends in order to sepa-

rate them from the sum.

VI. We begin the addition at the column containing the lowest denomination, and work from right to left, because, by so doing, we are enabled to corry, from the column added, the number of units of the rext higher denomination it contains, to their appropriate column, and thus perform the work by one addition, which would otherwise require two or more.

VII. We divide the sum of the units of any one denomination by the number required to make one of the next higher, in order to know how many we are to carry to the

next higher.

SECT. II.]

VIII. The addition of simple numbers was formerly called Simple Addition; and the addition of compound or denominate numbers, Compound Addition. As the same rule applies to the addition of all numbers, there is no cason why, in a second course, we should treat of the addition of simple and denominate numbers separately.

#### QUESTIONS.

NOTE.—Arabic numerals, thus (14), refer to the articles of the Section, and Roman numerals, thus (VI.) to the Recapitulation.

1. Into what parts may Arithmetic be divided? (1)
2. Of what does the Arithmetic of whole numbers treat? (1)
3. What rules are included in the Arithmetic of Whole Numbers? (2)
4. Of what does the Arithmetic of Fractions treat? (1)
5. How is the Arithmetic of Fractions treat? (1)
6. How is the unit divided in Vulgar or Common Fractions? (3)
7. How is the unit divided in Decimal Fractions? (3)
8. Of what does the Arithmetic of Ratios treat? (1)
9. What rules of Arithmetic are embraced in the Arithmetic of Ratios? (4)
10. What are the fundamental rules of Arithmetic? (5)
11. Why are they so called? (5)
12. Upon what rules do all the operations of Arithmetic ultimately depend? (6)
13. What is the sum of two numbers? (7)
14. What is Addition? (3 or I.)
15. What are plends? (9 or II.)
16. What has a plends? (9 or II.)
17. What has a plends? (9 or II.)
18. Why input we piece units of the same denomination in the same vertical column? (FV.)

- 19. Why do we draw a line under the addends? (V.)
  20. Why do we begin to add at the lowest denominations? (VI.)
  21. Why do we divide the sum of the units of any one denomination by as many as make one of the next higher? (VII.)
  22. How do we prove addition? (19.)
  23. Upon what axiom is the 2nd method of proof founded? (19)
  24. So far as the result is concarned, does it make any difference where we commence to addition? (12.)
  25. Exhibit the work when we commence adding at the left-hand side, or highest denomination. (12.)
  26. When the addends are very numerous, what plans may we adopt? (18)
  27. Upon what principle does the former of these plans proceed? (19)
  28. What different rules were formerly made in addition? (VIII.)
  29. Is this distinction necessary? Why not? (VIII).
  30. Illustrate the difference between spelling and reading in addition. (20)

#### SUBTRACTION:

- 21. Subtraction is the process of finding the difference between two numbers.
- 23. The greater of the two given numbers, or that which is to be lessened, is called the Minuend (Lat. Minuendus, "to be lessened"); the smaller, or that which is to be subtracted, the Subtrahend (Lat. Subtrahendus, "to be subtracted").
- 23. If anything is left after making the subtraction, it is called the remainder, difference, or excess.
- 24. Only quantities of the same denomination (i. e. which have the same unit) can be subtracted the one from the other.
- 25. Subtraction is indicated by —, called the minus, or negative sign: Thus 5—4=1, read five minus four equal to one, indicates that if 4 is subtracted from 5, unity is left.

Quantities connected by the negative sign cannot be taken, indifferently, in any order; because, for example, 5-4 is not the name as 4-5. In the former case the positive quantity is the greater, and 1 (which means + 1) is left; in the latter, the negative quantity is the greater, and -1, or one to be subtracted, still remains. To illustrate ye further the use and nature of the signs, let us suppose that we have five pounds and owe four; the five pounds we have wil. we represented by 5, and our debt by -4; taking the 4 from the 5, we shall have 1 pound (+1) remaining. Next, let us suppose that we have only four pounds and owe five; if we take the 5 from the 4 (that is, if we pay as far as we can) a debt of one pound, represented by - 1, will still remain; consequently 5-4=1; but 4-5=-1,

tion by

where nd side,

erence

which endus. be sub-

(i. e. e from

nus, or r equal is left.

taken. not the is the er, the racted, ture of

e four; ur debt +1) repounds

pt? (18) ?\_(19) ion. (20)

e subion, it

pay as

26. When several numbers, connected by the signs x and are placed within brackets, thus, (7+4-6-3+9,) the whole expression is to be considered as one quantity. The negative sign before such an expression indicates that the value of the whole expression within the brackets, is to be subtracted, or, what amounts to the same thing, that the numbers having the sign+before them are to be subtracted, and those having the sign, added. Hence a minus sign before a bracket, has the effect of changing the signs of all the quantities within the brackets, when the brackets are removed. So, also, when we desire to place a quantity within brackets, we must change its sign, if the sign preceding the first bracket be minus.

The following examples will show how the brackets affect numbers, according as we make them include an additive, or a

subtractive quantity:

subtractive quantity. 27-4+7-3=27 27-(4+7-3)=19But 27-(4-7+3)=27. [changing all the signs of the original quantities, but the first.]

Again 48+7-8-8+7-2=49; what is in the brackets being additive, it is not necessary to change any signs. 48+7-(3+8-7+2)=49; it is now necessary to change all the signs in the brackets. 48+7-3-(8-7+2)=49; it is necessary in this case, also, to change the signs.

27. When the numbers are small they can be subtracted mentally, thus: from 6 shillings take 4 shillings, and the result is evidently 2 shillings; from 9 pounds take 4 pounds, and the remainder is 5 pounds; from 16 days, take 9 days, and the remainder is 7 days; from 14 sixteenths take 5 sixteenths, and the remainder is 9 sixteenths, &c.

When the numbers are too large to be conveniently retained in the mind, they may be written as in addition.

EXAMPLE 1.—From 97 take 43, that is, from 9 tens and 7 units take 4 tens and 3 units.

OPERATION. 90+7 or 97 = Minuend. 40+3 or 43 = Subtrahend. EXPLANATION.—3 units from 7 units leaves 4 units, and 40 units or 4 tens from 90 units or 9 tens, leave 50 units or 5 tens.

50+4 or 54 = Remainder. Example 2.—Let it be required to subtract 746 from 978, or from 900+70+8 to take 700+40+6.

OPERATION. 2 2 EXPLANATION.—6 units from 8 units, and 2 units 700+70+8 or 7 4 6 remain; 40 units or 4 tens from 70 units or 7 tens, and 30 units or 3 tens remain; and 700 units or 7 hundreds, from 900 units or 9 hundreds, and 200 units, or 2 hundreds remain.

Brangle 3 .- From 842 take 661.

HEFFACE S.—From 542 take 601.

REFFACE ATION.—In placing the subtrahend under the minuend, in this example, we find that, while we can subtract it.

III. tract the units from the units, we cannot subtract the tens from the tens, since we cannot or 650-50-1 or 650-60-1 have 6 tens in the subtrached and only a tens in the minuend. We get over this in the minuend. We get over this in the minuend. We get over this difficulty by considering the minuend to be not 500-60-10-2, but 760-140-2, or in other words, we borrow one of the order of hundreds and reduce it to tens. Now we have 1 unit from 2 units and 1 units or 8 tens remain; 600 units or 6 tens from 140 units or 14 tens, and 80 units or 8 tens remain; 600 units or 6 hundreds, from 700 units or 7 hundreds, and 100 units or 1 hundred remain.

Example 4.—Let it be required to subtract 3 cwt. 2 qrs. 7 lbs. from 9 cwt. 1 qr. 8 lbs.

EXPLANATION.—As we cannot subtract 2 qrs. from 1 qr. we borrow 1 cwt. and reduce it to quarters. The 9 cwt. cwt. qrs. lb. cwt. qrs. lb. 1 qr. 8 lb. we then consider as 8 cwt. 5 qrs. 8 lb. and from it subtract the 5 cwt. 2 qrs. 8 lb. and from it subtract the 5 cwt. 2 qrs. 7 lb. Thus, 7 lbs. from 8 lbs. and 1 lb. remains; 2 qrs. from 5 qrs. and 3 qrs. remain; and 3 cwt. from 8 cwt. and 5 cwt. remain;

28. Hence, to find the difference between two numbers, we deduce the following:—

#### RULE.

Write the subtrahend under the minuend, so that units of the same denomination may be in the same vertical column. (24) Draw a line under the subtrahend to separate it from the remainder. Subtract each digit in the subtrahend from the one over it in the minuend beginning at the lowest denomination.

When the units of any one denomination of the minuend fall short of those of the same denomination in the subtrahend, borrow one of the next higher denomination in the minuend, reduce it to its equivalent units of the required denomination, add them to the units of that denomination given in the minuend, and from their sum subtract the units of that denomination given in the subtrahend.

29. The following is the complete work of a question in Subtraction:

EXAMPLE 5.—From 6400 lbs. 0 oz. 0 dwt. 7.0006 grs. take 987 lbs. 3 oz. 17 dwt. 22.6349 grs.

18 14. 31	.1	OPERA'		
(10) 9 9	11	19	24 9 9 9	
5 3 10 10	12	20	6.701010(10	
4 4 0 0 lbs			. TO 0 0 6	grs. Minuend.
987	1 3	17	22.6 3 4 9	Subtrahend.
ter merete flett fem			rene & con John	1

in this en subince we l only 4 ver this uend to e of the 2 units and 80 7 hun-

7 1bs.

rrow 1 9 cwt. t. 5 grs. lb. reemain : nain. nbers.

of the (24)under. in the

short ow one to its units ir sum

on in

take

iend. end.

der.

EXPLARATION.—Here, as we cannot take 9 tenths of thousandths of a grain from 6 tenths of thousandths of a grain, we borrow one grain, there being no tenths, hundredths, or thousandths in the minuemd. Now this one grain is equivalent to ten of the order of tenths of grains. Borrow one tenth and there remain 9 tenths, and the one tenth we borrowed is equal to 10 hundredths. Borrow 1 hundredth, there remain 9 hundredths, and the one hundredth we borrowed is equal to 10 thousandths. Borrow 1 thousandth, there remain 9, and the 1 thousandth is equal to 10 of the order of tenths of thousandths—the order for which it was necessary to borrow. 10 of the order of tenths of thousandths of grains, make 16, from which take 9 of the order of tenths of thousandths of grains, and there remain 7 of the order of tenths of thousandths of grains; 4 of the order of thousandths from 9 of the order of hundredths remain; 3 of the order of hundredths from 9 of the order of hundredths and 5 of the order of hundredths and 6 hundredths remain; 6 tenths from 9 tenths and 3 tenths remain.

undustricts and 5 of the order of thousandths remain; 3 of the order of hundredths from 9 of the order of hundredths and 6 hundredths remain; 6 tenths from 9 tenths and 8 tenths remain.

Again, as we cannot take 22 grains from 6 grains, we borrow from the next available higher order, which, in this case, is hundreds of pounds. 1 of the order of hundreds of pounds reduced, as above, to its equivalent lower denomination, is equal to 9 tens of 1bs., 9 units of 1bs. 11 oz. 19 dwt. 24 grs. 24 grains, added to 6, make 30 grains, and 22 grains from 30 grains, leave 8 grains; 17 dwt. from 19 dwt. leave 2 dwt; 3 oz. from 11 oz. leave 8 oz.; 7 units of 1bs. from 9 units of 1bs., leave 2 units of 1bs.; 8 tens of 1bs. from 9 tens of 1bs. from 9 units of 1bs. leave 2 units of 1bs.; 8 tens of 1bs., from 3 hundreds of 1bs., so we are compelled to borrow 1 of the order of thousands of 1bs., which is equal to 10 hundreds of 1bs., and 8 hundreds of 1bs., and 4 hundreds of 1bs. remain; 0 thousands of 1bs. from 8 hundreds of 1bs. and 5 thousands of 1bs. remain; 0 thousands of 1bs. from 5 thousands of 1bs. and 5 thousands of 1bs. remain.

30. If any digit of the minuend be smaller than the corresponding digit of the subtrahend, practically, we can proceed in either of two ways. First, we may increase that denomination of the minuend which is too small, by borrowing one from the next higher, (considered as so many of the lower denomination, or that which is to be increased,) and adding it to those of the lower, already in the minuend. In this case we alter the form, but not the value of the minuend; which, in the example given below, would become—

become-

hundreds. tens. units. 12 = 792, the minuend. 7 = 427, the subtrahend.

Or, secondly, we may add equal quantities to both minuerid and subtrahend, which will not alter the difference; then we would have hundreds. tens, units.

2+10 = 792 + 10, the minuend + 10. 7 = 427 + 10, the subtrahend + 10. 2+1 7

3 6 5 = 385 + 0, the same difference.
In this mode of proceeding we do not use the given minuend and subtrahend, but others which produce the same remainder. PROOF OF SUBTRACTION.

31. FIRST METHOD.—Add together the remainder and subtrahend; the sum should be equal to the minuend.

For the remainder expresses by how much the subtrahend is smaller than the minuend; adding therefore, the remainder to the subtrahend, should make it equal to the minuend; thus,

8754 minuend. subtrahend. 5839 2915 difference.

Sum of difference and subtrahend, 8754 = minuend.

Tal

SECOND METHOD.—Subtract the remainder from the minuend, and what is left should be equal to the subtrahend.

For the remainder is the excess of the minuend over the subtrahend; therefore, taking away this excess should leave both equal; thus

8634 minuend.

PROOF: 8634 minuend. 649 remainder.

649 remainder. New remainder, 7985 = subtrahend. In practice, it is sufficient to set down the quantities once; thus

8684 minuend. 7985 subtrahend.

649 remainder.

Difference between remainder and minuend, 7985 = subtrahend

#### EXERCISE 11.

They &	153.2		* 2* 3		
3	(1) 11000000 9919919			00 800000	
	1080081				
From Take	(6) 85·73 42·16	(7) 864·5 73·2	(8) 594·763 85·6	(9) 47.630 0.078	(10) 52·137 20·005
	43.57	;- <del></del>			
	(11)	(12)	(13)	(14)	(15)
	0.00063 0.00048	874·32 5·637			400·3270 5003 0·006
,	0.00015		. , ,	,	
7. 56	5676—56 6789— 7 1000—	5674= 4	91115.	27. 97777- 28. 60000- 29. 75477-	- 1= 59999

16.	7465676-	67456=	898220.	27.	97777-	4=	97773.
17.	566789-	75674=	491115.	28.	60000-	1=	59999.
18.	941000-	5007=	935993.	29.	75477-	76=	75401.
19.	97001-	50077=	46924.	30.	7.97-	1.05=	6 92.
20.	76734	977=	75757.	31.	1.75-0	·074==	1.676.
21.	56400-	100=	56300.	32.	97.07-4	.769=	92.301.
22.	700000-	99=	699901.	33.	7.05-4	.776=	2.274.
23.	5700—	500=	5200.	34.	10.761-9	·001=	1.76.
24.	9777-	89=	9688.	35.	·10009—7	1-121=	<b>₹</b> .97909.
25.	76000-	1 to 1=	75999.	36.	176 1	-700	176.093.
26.	90017-	3=	90014.	37.	15.06	·863=	7.179,

nuend,

rahend;

(15) 0·3270 0·006

> 97773. 59999. 75401.

6 92. 1 676. 92 301. 2 274. 1 76. 97909. 76 093. 7 179, 20207

#### ABT, OL.

## MONEY.

From Take	\$9876·43 987·49	(39) \$427.63 197.21	\$721.73 \$16.25 91.00 9.75
. 4 0 p	\$8888.94	\$230.42	\$ 1,6 \$ 5
From Take	(42) \$1234.50 999.96	(43) \$671.98 99.67	(44) (45) \$286·29 \$7·19 611·89 1·86
3,71	\$234.54	\$572.31	8 9 . S. 1 . S. E

*	, ,	(46)	)	(4	7)	81	P.C.	(48	)	1	(49)	(50)	
From Take	£ 1098 434		d. 6	£ 76.1	s. 14	d. 8	*£	s. 15	d. 6	£ 47.	s. d. 16.7	£ 8. 6 97.14 6 15	1.
- 74	FEES	16	10	-	, ,4,4						. Wegett	79.) med. 5	200

(51) (52) (53) (54) (55) £ s. d. From 98 14 2 47 14 6 97 16 6 147 14 4 560 15 6 Take 77 15 3 38 19 9 88 17 7 120 10 8 477 17 7

#### AVOIRDUPOIS WEIGHT.

b e	* * * * * * * * * * * * * * * * * * *	(56)		~	(57)	2.	, (58)	1. 30	i o	59)	
From Take	cwt. 200	qrs.	lb. 24		2	15	cwt. qrs. 9664 2 9073 0	23	554	0 0	
. 10	100	3	9			_		1 1- 1	L	1 1	-

#### TROY WEIGHT

*			(60)	7	, 1	. (6	31)					)
				grs.						02.		
From						-	10	-	-	7 0		•
Take	97	, 100	16	10		٠	- 17	23	798	3, 0	18	17
	457	9	2	13				, (		6 1.A	P 30	

TA

#### TIME.

			(6	3)			(64)			(68	5)	
	From Take	767 476	ds. 131 110	hrs. 6 14	ms. 30 13	yrs. 475 160	ds. 14 16	hrs. 13	ms. 16 17	yrs. ds. 567 126 400 0	hrs. 14 15	ms. 12 0
4	*** **********************************	001			<del></del>				•	**		

#### APPLICATIONS.

1. A shopkeeper bought a piece of cloth containing 42 yards for £22 10s., of which he sells 27 yards for £15 15s.; how many yards has he left, and what have they cost him?

Ans. 15 yards; and they cost him £6 15s. 2. A merchant bought 234 tons, 17 cwt., 1 quarter, 23 lb., and sold 147 tons, 18 cwt., 2 quarters, 24lb.; how much re-

Ans. 86 tons, 18 cwt. 2qrs. 24lb. mained unsold?

3. In 1356 the revenue of Canada was as follows :- customs, \$4500000; public works, \$500000; crown lands, \$500000; and casual, \$320000. For the same year the expenditure was as follows :- interest on public debt, &c., \$1000000; civil government, \$225000; legislation, \$450000; administration of justice, \$450000; education, \$380000: collection of revenue, \$940000; public works, &c., \$1755000. How much did the total revenue of that year exceed the total expenditure?

Ans. \$620000. 4. The census of 1852 gives the population of Upper Canada as 962004, and that of Lower Canada as 890261. By how much did the population of the former exceed that of the latter?

Ans. 71743. 5. Upper Canada contains 147832 square miles; Lower Canada, 209990 square miles; Nova Scotia and Cape Breton, 18746 square miles; New Brunswick, 27620 square miles; Prince Edward's Island, 2173 square miles; Newfoundland, 36000 square miles; and Hudson's Bay Territory, 2436000 square miles. By how much does the aggregate extent of these British North American Provinces fall short of the total area of the

Ans. 57755 square miles. 6. A merchant has 209 casks of butter, weighing 400 cwt. 2 grs. 14lb.; and ships off 173 casks, weighing 213 cwt. 2 qrs. 24lb. How many casks has he left; and what is their weight?

United States—the latter being 2936116 square miles?

Ans. 36 casks, weighing 186 cwt. 3 grs. 15lb. 7. If from a piece of cloth containing 496 yards, 3 quarters, and 3 nails, I cut 247 yards, 2 qrs., 2 nails, what is the length of the remainder.

Ans. 249 yards, 1 quarter, 1 nail. 8. A field contains 769 acres, 3 roods, and 20 perches, of

I. betwe II

thi Yo

exc

doe

cor

\$94

cus

WOI

11 11.

1

1

32

13-

take

with

from

lowi

be a

4 fro

be re

10, 1

nueno

SECT. II.

42 yards low many

er, 23 lb., much reeqrs. 24lb.
—customs,
\$500000;
diture was
000; civil
inistration
of revenue,
h did the
ere?
; \$620000.

er Canada
By how
the latter?
Ins. 71743.
Lower Caton, 18746
s; Prince
Ind. 36000
00 square
lese British
area of the
? \*

are miles.
400 cwt. 2
cwt. 2 qrs.
weight?
grs. 15lb.
quarters,
hat is the
ter, 1 nail.
perches, of

which 576 acres, 2 roods, 23 perches are tilled; how much remains untilled?

Ans. 193 acres, 37 perches.

9. I owed my friend a bill of £76 16s. 91d., out of which I paid £59 17s. 101d.; how much remained due?

Ans. £16 188. 103d.

10. The population of London is 2363141, and that of Paris is 1053262. How much does the population of London exceed that of Paris?

Ans. 1309879.

11. The population of Liverpool is 384265, and that of New York 515547. How much does the population of New York exceed that of Liverpool?

Ans. 131282.

12. Lake Huron contains 20000 square miles: by how much does it exceed the area of Lakes Erie and Ontario—the former containing 11000 square miles, and the latter 7000 sq. miles?

Ans. 2000 square miles.

13. A merchant has \$6947.87 in bank; \$4789.63 in stock; \$9491.11 in property; and \$14167.93 on his books against his customers: his debts amount to \$19478.25. How much is he worth after paying what he owes?

Ans. \$15918.29.

14. What is the value of 6-3+15-4?

Ans. 14.

15. Of 43+(7-3-14)?

Ans. 33.

16. Of 47·6—(2+1—24+16—0·34)?

Ans. 52.94.

17. What is the difference between 15+13-6-81 and 15+13-(6-81+62)?

Ans. 100.

32. Before the pupil leaves subtraction he should be able to take any of the nine digits, continually, from a given number, without stopping or hesitating, thus, in subtracting 7 continually from 94, he should say, 94, 87, 80, 73, 66, 59, &c. In the following examples, which are inserted for practice, he should not be allowed to spell the subtraction, thus, 6 from 9 and 3 remain, 4 from 2, we can't, but 4 from 12 and 8 remain, &c.; but should be required to read as follows:—6, 9..3; 4, 12..8; 9, 13..4; 10, 11..1; 10, 18..8, &c.

(18)

9800046043019181697800041981329 191347813191681473199916199846

(19)

74321913047123098706540456007139 1342345678912345678912345678912

#### RECAPITULATION.

I. Subtraction is the process of finding the difference between two numbers.

II. The greater of the two numbers is called the minuend.

III. The smaller of the two numbers is called the subtrahend.

IV. What is left after making the subtraction is called the remainder or difference.

V. Only quantities of the same denomination can be subtracted.

VI. Subtraction is indicated by the sign —; which is called minus, or the negative sign.

VII. When several numbers are inclosed in brackets, they are to be considered as constituting only one quantity.

VIII. When a negative sign precedes the first bracket it indicates that all the quantities within the brackets are to have their signs changed when the brackets are removed.

IX. When quantities are removed into brackets, precoded by the negative sign, all their signs must be changed.

X. We begin subtraction at the lowest denomination, because it is sometimes necessary to borrow from the higher denominations and reduce.

XI. Instead of thus borrowing and reducing, we may consider any denomination in the minuend increased by as many units of that denomination as make one of the next higher, and then add one to the next higher denomination in the subtrahend. This is merely adding the same quantity under different forms to both minuend and subtrahend, and consequently cannot affect the value of the remainder. (30.)

#### QUESTIONS TO BE ANSWERED BY THE PUPIL.

Note.—Numbers in Roman numerals, thus (V), refer to the Recapitulation; those in Arabic numerals, thus (25), refer to the articles of the Section.

1. What is Subtraction? (I.)

2. What is the minuend? (II.)

3. What is the derivation of the word minuend? (22)

4. What is the oubtrahend? (III.)

5. What is the derivation of the word subtrahend? (22)

6. What is the remainder? (IV.)

6. What is the remainder? (IV.)

6. What is the remainder? (1V.)
7. What kind of quantities can be subtracted? (V.)
8. How is subtraction indicated? (VI.)
9. When several numbers are inclosed together in brackets, how are they to be taken? (VII and 26.)
10. What effect has a negative sign preceding brackets? (VIII and 26.)
11. When quantities are removed into brackets, preceded by the sign—what must be done with them? (IX and 26.)
12. That is the rule for subtraction? (28.)

12. What is the rule for subtraction? (28.)

AR

ber mul

mul plica

we n

3 man the p TI

of th facto 37 divid

38 integr comp

38 taking in the 1st.

be equ 2nd

will be multip 3rd. d the sub-

n is called

on can be

-; which is

brackets, e quantity. irst bracket rackets are re removed.

ackets, prebe changed. nomination, n the higher

ing, we may reased by as of the next enomination same quansubtrahend, e remainder.

PUPIL.

he Recapitulaarticles of the

s, how are they II and 26.) l by the sign-

13. Why must we put units of the same denomination in the same vertical column? (24)
14. When a digit in the subtrahand is greater than the corresponding digit in the minuend, what is done? (27 Example 3, or 29)
15. What other plan may be adopted? (30)
16. Upon what principle does this plan proceed? (XI.)
17. Why do we begin to subtract at the right-hand side? (X.)
18. How do we prove subtraction? (31)
19. Upon what principles are these methods of proof founded? (31)
20. Illustrate the difference between spelling and reading in subtraction. (32)

tion. (32)

#### MULTIPLICATION.

33. Multiplication is a short process of taking one number as many times as there are units in another. Hence multiplication is a short method of performing addition.

34. The number to be taken or multiplied is called the multiplicand, and in addition would be called an addend.

35. The number denoting how many times the multiplicand is to be taken, or, in other words, that by which we multiply, is called the multiplier.

36. The number arising from taking the multiplicand as many times as there are units in the multiplier, is called the product, and corresponds to the sum of the addends in addition.

The multiplicand and multiplier are called the factors of the product because they make or produce it, (Lat. factor, "a maker, agent, or producer.")

37. A prime number is one which cannot be exactly divided by any whole number, except the unit one and itself.

38. A composite number is the product of two or more integral factors; neither of which is unity. Thus 16 is a composite number, and its factors are 8 and 2, or 4 and 4.

39. Since the product is the result which arises from taking the multiplicand as many times as there are units in the multiplier, it follows:

1st. If the multiplier be equal to unity, the product will

be equal to the multiplicand.

2nd. If the multiplier be greater than unity, the product will be as many times greater than the multiplicand as the multiplier is greater than unity.

3rd. If the multiplier be less than unity, that is, if it be

10

11 12 It

two N for t

profi

lows, dera woul 13 ti

2 are

expr by 9

a proper fraction, the product will be as many times less than the multiplicand as the multiplier is less than unity.

40. Let it be required to multiply any two numbers' together, say 7 and 6.

If we make in a horizontal live as many stars as there are units in the multiplicand, and make as many such lines of stars as there are units in the multiplier, it is manifest that the entire number of stars will represent the number of units which result from taking the multiplicand as many times 6 as there are units in the multiplier.

But it is evident that we may consider the 42 stars in the above figure, either as 7 stars taken 6 times, or as 6 stars times, that is, 6 × 7 = 42 = 7 × 6.

Hence either of the factors may be used as multiplier without altering the product.

41. Let it be required to multiply the number 8 by the composite number 6, or which the factors are 3 and 2.

If we write 8 stars in a horizontal line and make 6 such lines, we shall evidently have in all 8×6 = 48, the number of units in all the lines.

But we may consider the 6 lines as 2 sets of 3 lines each, and in each set of 3 lines there are 6×3 = 24 units. Therefore in the 2 sets there are 24×2 =48 unit. Again we may consider the 6 lines as 3 zets of 2 lines each, and in each sat of 2 lines there are 8×2 = 16 units. Therefore in 3 such sets there are 16×8 = 48 units.

Hence 8×6 = 48 8×3 = 24 and 24×2 = 48 = 8×6

 $8\times2=16$  and  $16\times5=48=8\times6$ .

And as the same may be shewn for any other composite number as well as for 6, we may conclude that,

When the multiplier is a composite number we may multiply by each of the factors in succession, and the last product will be the entire product sought.

42. As the multiplication of the higher numbers may be resolved into the multiplication of one digit by another, the pupil should make himself perfectly familiar with the following table:

This table is called the Multiplication Table, and was calculated by: Pythagoras, a celebrated Greek philosopher who flourished about 500 years before Christ, It was calculated after the following manner:—Pand 2 are 2 are 4; 3 and 3 are 6; twice 3 are 6; 5 and 4 are 8—twice 4 are 8, &c.

nes less unity.

umbers

1 0

ultiplier

site numi

2 = 16 3 = 48

, we shall nesch set rare 24×2 each, and such sets

per as well

we may

may been the pupil ng table:

rulated by: t 500 years and 2 area are 8, &c.

#### MULTIPLICATION TABLE

		TII BION		Territoria Calculation	10-20-0
Twice,	3 times	4 times	5 times	6 times	7 times
1 are : 2.	1 are 3	1 are 4	1 are 5	lare 6	, 1 are; 7
2 - 4	2 6	2 8	2 10	2 - 12	2 - 14
3 - 6	3 - 9				
	4 - 12				
	1 5 - 15				
	m6 - 18.			6 - 36	
	7. — 21.			7 - 42	7 - 49
	8 - 24				
	9 - 27				
	10 - 30				
	11 - 33				
	12 — 36				
	9 times				
lare: 8			10 1		1 are: 12.
	7 2 — 1				
3 24	3 - 2	7 3			3 - 36
4 - 32	4 - 3	6 4 —	40 4	- 44	4. 48.
5 - 40	5 - 4	5 5	50 . 5.		5 60
6 - 48				66	6 - 72
7:56	7 - 6	3 7 -			7 - 84
8 64			80: 8		8 - 96
9 - 72					9 - 108
10 - 80					0 - 120
					1 - 132
11 88	<del></del>				
11 — 88 12 — 96					2 - 144

It appears from this table, that the multiplication of the same two numbers in whatever order taken, produce the same product.

NGTE.—Though the part of the multiplication table given above is enough for the pupil to commit to memory at first; yet, after he has made some proficiency in a chimetic, he may find it advantageous to commit what follows, as it will enable him, in many cases, to shorten his work in a considerable degree. The labour of committing a still more extended table would be scarcely compensated by the advantage resulting.

13 times	14;times:	15 times	16 times	17 times	18 times	19 times
2 are: 26	2 are 28	2 are 30	2 are 32	2 are 34	2 are 36	2 are 38
3 - 39	8 - 42	3 - 45	3 - 48.	3 - 51	3 - 54	8 - 57
4 - 521	4 - 56	4 60	4 - 64	4 - 68	4 - 72	4 - 78
5 65	5 - 70	5 — 75	5 - 80	5 - 85	5 90	5 20
6 - 78	6 - 84	6 - 90	6 - 98	6 - 102	6 - 108	6 - 114
7 - 91	7 - 98	7 - 105	7 — 112	7 — 119	7 - 126	7 - 183
8 — 104	8 — 112	8 - 120	8 — 128	8 136	8 - 144	8 - 152
9 - 117	9 126	9 - 135	19 - 144	9 158	9 - 162	19 - 171

43. The multiplication of one quantity by another is expressed by  $\times$ ; thus  $7 \times 9 = 63$ , means that 7 multiplied by 9 is equal to 63.

44. Quantities connected by the sign of multiplication are multiplied by any number, if we multiply any one of the factors by that number; thus  $(9 \times 10 \times 2) \times 27 = 9 \times 10 \times 54$ , or  $9 \times 270 \times 2$ ; that is, if we multiply the factor 2 or the factor 10 by 27, we, in

effect, multiply the whole number  $(9 \times 10 \times 2)$  by 27.

45. When a quantity within brackets, consisting of several terms connected by the signs + and -, is to be multiplied by any number, each of its parts or terms must be multiplied. This arises from the fact that we consider the several terms within the bracket as constituting but one quantity, and to multiply the whole, we must multiply each of its parts. Thus  $(7+8-3) \times 3$  $=7\times3+8\times3-3\times3$ ; and  $(8+7-5)\times(13-2)$  means that each of the terms within the former bracket is to be multiplied by each of the terms within the latter, or by their difference.

46. Let it be required to multiply 768 by 9.

Now  $768\times9=(700+60+8)\times9=700\times9+60\times9+8\times9(Art.45)$ . Hence, to far as the result is concerned, it matters not whether we commence multiplying at the lowest or at the highest denomination;  $700\times9+60\times9+8\times9$  being evidently equal to  $8\times9+60\times9+700\times9$ .

Commencing the multiplication at the left-hand side, or highest denomination; the work is as follows:

768 which may 9 be thus abbreviated.	768	EXPLANATION.—7 hundreds; taken 9 times, are 63 hundreds; by 9, are 54 tens; and 8 units m 72 units. 63 hundreds, 54 tens, a together, make 6912. The secon the only abbreviation possible w at the highest denomination.	6 tens multiplied nultiplied by 9, are nd 72 units, added ad operation shows
TITLE TO MAKE A		at the lightest denomination.	n 1

Let us now take the same question and commence at the right-hand or lowest denomination.

	PERAT			
I. which may 768 be thus ab- 9 breviated.	768 9	and thus still farther abbre- viated.	111. 768 9	
540 6800	72 54 63	· ·	6912	
6912	6912			

EXPLANATION.—No. II. differs from No. I. only in having the unnecessary 0's omitted. In No. III. the principle of carrying is taken advantage of, thus—8 units, multiplied by 9, are 72 units, equal to 2 units and 7 tens to

6912 6912 carry—6 tens, multiplied by 9, are 54 tens, and 7 tens, make 61 tens, equal to 1 ten, and 6 hundreds to carry; 7 hundreds, multiplied by 9, are 63 hundreds, and 6 hundreds, make 69 hundreds, equal to 6 thousands and 9 hundreds.

Hence, in order that we may be enabled to take advantage of the principle of CARRYING, we commence the multiplication at the right-hand or lowest denomination.

47. From the last article (46), for multiplying by any integral multiplier, not exceeding 12, (or 20 if the extended Multiplication Table be used) we deduce the following :-

Multiply every order of units in the multiplicand in succes-

n are ctors 0×2; we, in

cr. II

veral ed by This. ithin ly the 3) ×3 t each ed by

o far as iplying being enomi-

by 9, or Itiplied y 9, are , added nmence

and or

only in ary 0's I. the staken units, tens to ied by tens, ds, and

of the t-hand tegral

plica-

ucces-

sion beginning with the lowest, by the multiplier, and divide each product, so formed, by the number of that denomination which makes one unit of the next higher; write down each remainder under units of its own order, and carry the quotient to the next product.

Example 1.—Multiply \$7896.43 by 11.

PERATION.

27896:43

plied by 11, make 33 hundredths, equal to 3 hundredths, to set down, and 3 tenths to carry; 4 tenths of dollars, or tens of cents, multiplied by 11, make 44 tenths of dollars, or tens of tenths we carried, make 47 tenths, equal to 7 tenths and 4 units to carry; 6 units, multiplied by 11, make 66 units, and 4 units we carried, make 70 units, equal to 0 units to set down and 7 tens to carry; 9 tens, multiplied by 11, make 99 tens, and 7 tens, make 106 tens, equal to 6 tens and 10 hundreds; 8 hundreds, multiplied by 11, make 86 hundreds, and 10, make 98 hundreds, equal to 8 hundreds and 9 thousands; 7 thousands, multiplied by 11 make 77 thousands, and 9, make 86 thousands, equal to 6 thousands and 8 tens of thousands.

Example 2.—Multiply 3 cwt. 2 qrs. 11 lbs. 7 oz. 6 drs. by 7.

OPERATION.

EXPLANATION.-7 times 6 drams are 42 drams 

		Exercise 12	2. * - ,	t neg n ship
Multiply By	· (1) 48960 5	75460 9	678000 8	57800 6
	244800	9		· · ·
Multiply	(5) 5·2736	• (6) 8 • 7563	(7) 0·21375	(8) ····································
Ву	10.5479	4	6	8
· (*)	10.5472			
Multiply By	(9) \$767·62 2	(10) \$672·56 2	(11) \$789·76 6	(12) \$573·46 5
,	\$1535.24			
Multiply	(13) 866342	(14) 738579	(15) 4716375	(16) 8429763
Ву	11	12	11	12

- 17. Multiply £32 8s. 6id. by 5.
  - Ans. £162 28. 81d.
- 18. Multiply £43 11s. 41d. by 8.
- Ans. £348 11s. 2d. Ans. £1507 16s. 3d.
- 19. Multiply £126 138, 01d. by 12.
- 20. Multiply 10 cwt. 3 qrs. 6 lbs. by 3. Ans. 32 cwt. 1 qr. 15 lbs. 21. Multiply 7 yds. 3 qrs. 1 na. by 7. Ans. 54 yds. 2 qrs. 3 na.
- 22. Multiply 11 oz. 10 dwt. 19 grs. by 12.

Ans. 11 lbs. 6 oz. 9 dwt. 12 gr.

48. When the multiplier is a composite number, and can be resolved into two or more factors, neither of which is greater than 12, we deduce from (41) the following:—

Multiply by each of the factors in succession and the last product will be the entire product sought.

Example 1.—Multiply 3 hrs. 7 min. 14 sec. by 64.

OPERATION. hrs. min. sec.×64=8×8 EXPLANATION.—Multiplying 3 hrs. 7 min. 14 sec. by 8, we obtain 1 day 6 hrs. 57 min. 52 sec., which we again multiply by 8, and obtain 8 days 7 hrs. 42 min. 56 sec., which is the product of the product duct of 3 hrs. 7 min. 14 sec., by 8 times 8 or 64.

7 .. 48 ... 56 Ans.

1 0 57

EXAMPLE 2.—Multiply 796:437 by:132.

OPERATION. 796:437×132=11×12

EXPLANATION .- We first multiply the 796'437×132=11>12 given number by 11, or, in other words, take it 11 times, and then take this result 12 times, which is evidently 8760'807=11 times multiplicand. equivalent to taking the given number 12 times 11 or 132 times.

105129'684=12 times 11 times multiplicand.

Example 3.—Multiply 16 cwt. 3 qrs. 11 lb. by 270.

OPERATION.

EXPLANATION.—270=10 times 27 or 10×3×9.

If, therefore, we take the given multiplicand 3 times, and then this product 9 times, and then this second product 10 times, it is evident we shall have, in effect, taken the given multiplicand  $3\times9\times10$  or 270 times.

EXERCISE 13.

- 1. Multiply \$169.78 by 36.
- 2. Multiply 796342:3 by 121.
- 3. Multiply \$33460 by 144.
- 4. Multiply 735 by 648.
- 5. Multiply £3 7s. 6d. by 18.

Ans. \$6112.08.

Ans. 96357418:3. Ans. 34818240.

Ans. 476280.

Ans. £60 16s. 0d.

II.

d.

d.

A.

a:

Tw

ıd

sh

ct!

n. 52 in

.0-

9.ddti

Ans. £125 19s. 11d. 6. Multiply £5 14s. 61d. by 22.

7. Multiply £3:48. 7d. by 810. Ans.: £2615 12s. 6d.

8. Multiply 11 cwt. 3 qrs. 14 lb. 7 oz. by 54.

Ans. 642 cwt. 1 qr. 4 lbs. 10 oz.

9. Multiply 26 bush. 3 pks. 1 gal. 1 qt. 1 pt. by 49.

Ans. 1319 bush. 0 pks. 1 gal. 1 qt. 1 pt.

10. Multiply 2 yds. 2 qrs. 2 na. 2 in. by 63.

Ans. 168 yds. 3 grs. 2 na. 0 in.

11. Multiply 5 days 17 hrs. 33

Ans. 16 10

11 % ... by 288. 15 hrs. 16 min. 48 sec.

49. When the multiplication and the multiplier is greater number, we proceed according to

denominate number but not a composite ollowing :-

Take the nearest composite number to the given multiplier, multiply successively by its factors and add to or subtract from the product so many times the multiplicand as the assumed composite number is less or greater than the given multiplier.

EXAMPLE 1.-Multiply £62 12s. 6d. by 76.

£		rion. d. 6 8	
- 501	0	0	
-	4	1411	

EXPLANATION.—We take 76=
9×6+4, and thus we get 72 times
the multiplicand, and to it adding
4 times the multiplicand, obtain
the desired product, viz., 76 times
the multiplicand.


4509 0 0 = 72 times multiplicand. 250 10 0 = 4 times multiplicand.


£4759 10 0 = 76 times multiplicand.

Instead of multiplying as above, we might have multiplied by 7 and 10 and increased the result by 6 times the multiplicand or we might have multiplied by 7 and 11, and decreased the result by once the multiplicand.


Example 2.—Multiply 17 lbs. 3 oz. 7 dr. 2 scr. 16 grs. by 789.

lb. 170		dr.	SCT.	grs.	×'9'=	9 ti	mes m	ultipli	cand.
173	3	7	1.	10	× 8 =	80 ti	mes m	ultipli	cand.
1788	8 1	111	1.	0,7	t,		ł .,	11 178	* 90
12132 1386 155	7:3	.1. 12dr	1: 2:: 1	0	= 700 = 80	time	mult	iplican iplican	dalar
13675	5	3	1	4	<del> 789</del> 1	.∫ times		nlican	





# IMAGE EVALUATION TEST TARGET (MT-3)



Photographic Sciences Corporation

23 WEST MAIN STREET WEBSTER, N.Y. 14580 (716) 872-4503

STATE OF THE STATE



EXPLANATION.—We divide the given multiplier into 700+50+0, and ob-ain the 3 partial products, which we add together, for the entire product.

Example 3.—Multiply 3 wks. 6 days 17 hrs. 21 min. 12 sec. by 4736.

#### OPERATION.

wks. ds. h. min. sec. 3 6 17 21 12×6=	wks. ds. h. min. sec. 23 5 8 7 12 =	6 times multiplicand.
10		and the state of the state of
30 4 5 33 0×3=	118 5 16 36 0=	30 times multiplicand.
and the same of th	TO SEE SEE SEE SEE SEE SEE SEE SEE SEE SE	200 Atm

0 = 700 times multiplicand. 0×4=15841 0 = 4000 times multiplicand.

> Ans. 18756 4 9 23 12 = 4786 times multiplicand.

EXAMPLE 4.—Multiply £47 16s. 2d. by 5783.

#### 1783=5×1000+7×100+8×10+8.

#### OPERATION.

143 8 6=product by units of the multiplicand.

8×8= 3824 13 4= product by tens of the multiplicand.

180 16 8×7 = 33465 16 8 = product by hundreds of the multiplicand.

6 8×5 = 239041 13 4= product by thousands of the multiplicand.

## Ass. 276475 11 10 = product by entire multiplier.

## EXERCISE 14. 14 MA MILE AND AND ADDRESS AN

- 1. Multiply £12 2s. 4d. by 83. med and Ans. £1005 13s. 8d.
- Ans. 962040 2s. 51d. 2. Multiply £963 0s. 03d. by 999.
- 3. Multiply £3 6s. 51d. by 3178. Ans. £10556 18s. 41d.
- 4. Multiply 16 bush. 3 pks. 1 gal. by 678.

Ans. 11441 bush. 1 pk. 0 gal.

5. Multiply 23 m. 6 fur. 33 rds. 4 yds. by 247.

Ans. 5892 m. 2 fur. 10 rds. 34 yds.

6. Multiply 38. 16° 30′ 45" by 721. Ans. 2559S. 25° 30' 45"

50. It may be proper here to caution the pupil against the absurd attempt to multiply one denominate number by another. Multiplication is merely a particular kind of addition, and when we are required to multiply a quantity by any number, we are simply required to repeat it as many times as there are units in the multiplier. It is evident, then, that to talk of multiplying \$19 19s. 11fd., by £19 19s. 11fd., or, in other words, of adding or repeating \$19 19s. 11fd. £19 19s. 11fd. times is simply ridiculous. Nevertheless, great pains have been taken to show that 2s. 6d. may be multiplied by 2s. 6d., and that the product will be either 3fd. or 6s. 3d !! Undoubtedly 2s. 6d. can be taken 2f times, and the result will be 6s. 3d.; or it can be taken one-eighth

of a time, and the result will be 3id., but this is a very different thing fretaking it 2s. 6d. times. In fact it is quite as nonsensical to talk of taking 2s. 6d. 2s. 6d. times as it would be to talk of taking 6 lbs. of beef 6 lbs. beef times; or, 7 bars of music 7 bars of music times, &c. Daudscim multiplication, which is sometimes adduced, as a proof that one denoming number can be multiplied by another, affords no support whatever to fatheory, as will be fully shown hereafter. (See Sec. III.)

51. Let it be required to multiply 729 by 478.

OPERATION. EXPLANATION.—From the preceding examples it is evided that when units are multiplied into any order whatever, the product will always be of that order. Here, then, we first multiply by the 8 units, as in (47). Next we multiply by the 5832 tens, thus:—9 units, multiplied by 7 tens, give 65 tens, equal to 3 tens, which we set down in the column of tens, and hundreds which we carry; 2 tens, multiplied by 7 tens, give 5103 to 3 tens, which we carry; 2 tens, multiplied by 7 tens, give 52 tens, and 6 hundreds which we carried, make 20 hundreds; equal to 6 hundreds to set down and 2 thousands to carry, &c. Next we multiply by the 4 hundreds as follows:

9 units, multiplied by 4 hundreds, give 36 hundreds, equal to 6 hundreds set down in the hundreds column, and 3 thousands to carry, &c. Lasti we add the several partial products together.

Hence, when the multiplicand is an abstract number, the multiplier being greater than 12 and not a composite number, we have the following:-

L. L. BULB. 2 . Of Multiply the multiplicand by each figure of the multiplier separately, beginning with the lowest, and write the partial products in separate lines, placing the first figure of each line directly under the figure by which you multiply, and, lastly, add the several partial products together.

EXAMPLE. Multiply 7423 by 6709.

EXPLANATION.—Here, as there are no tens in the multiple we may either proceed directly to the hundreds after mul OPERATION. 7428 plying by the units, or we may set down a 0 under the and then write the product by the hundreds in the same always remembering to place the first digit of the partial duct under the figure by which we are multiplying in othat all the digits of the same order may come in the 6709 66807 519610 44538 vertical column.

19800907

d.

	10,1, 6	LAX	RUIDHE	10.	4	1.55
	(1)	.2: (2)	• ,	(3)	(4) 997	(5)
Multiply	325	765	1/2	732	997	667
- <b>By</b>	95.	765	i ve List	456	345	347
		20.00	_			

6. Multiply 7071 by 556. Ans. 3931476. Ans. 48288058. 7. Multiply 15607 by 3094. 8. Multiply 39948123 by 6007. Ans. 239968374861.

Ans. 27416327796.

9. Multiply 2778588 by 9867.

52. Let it be required to multiply 63.5 by .97.

AND ATION. EXPLANATION.—Since (51) any order, multiplied by units, will give that erder—tenths, multiplied by tenths tenths. Hence it is obvious that tenths multiplied by tenths will give the next lower order, or hundredths, and also that tenths, multiplied by hundredths, will give the next lower order, or hundredths, and also that tenths, multiplied by hundredths, will give the next lower order again, or thousandths. In the above crample, therefore, we proceed thus:—5 tenths, multiplied by 7 hundredths, give 55 themsandths, equal to 5 thundredths, into the tenths, give 10 hundredths, and 5 hundredths we carried, make 24 hundredths, give 11 hundredths, give 45 tenths, and 5 tenths we carried, make 44 tenths, equal to 4 hundredths, give 45 tenths, and 5 tenths, multiplied by 7 hundredths, equal to 5 hundredths, equal to 6 tenths, equal to 6 hundredths, equal to 5 hundredths, equal to 6 hundredths equal to 6 hundredths equal to 6 hundredths equal to 6 hundredths equal to 6 hu

53. Strictly speaking, all examples in multiplication of decimals should be worked according to the above method. An attentive consideration of the reasonings in (52) will. however, show that the lowest digit of the product of any two numbers containing decimals, must always be a number of places to the right of the decimal point, equal to the sum of the decimal places, in both multiplicand and multiplier.

Hence, when the multiplicand or multiplier, or both, contain decimals, we deduce the following-

Multiply as though there were no decimals, and then remove the decimal point in the product as many places to the left as there are decimals in both the multiplicand and the multiplier.

EXAMPLE 1 .- Multiply 5:63 by 0:00005.

OFFICATION.

BYPLANATION.—We multiply 561 and remove the decimal point access places to the sale since there are for decimal places in the multiplier and two in the multiplierand, that is, we have taken a number a hundred times too great a hundred thousand times too often, and the product 2815 is therefore ten million times too great, and to make it what it should be, we divide it by ten millions; or, in other words, remove the decimal point seven places to the

EXAMPLE 2 .- Multiply 2.073-by 5.12.

EXPLANATION.—We multiply as though both were whole numbers, and out off five decimals, since there are three in the multiplicand and two in the multiplier. OPERATION. 2073

4146

fore,

d to 4 hunl to 4 hun-

n of

hod.

will.

any

the

mul-

ooth,

e the

e are mul-

ndred

, and

o the

vhole

ARTS, 52-54.7

#### EXERCISE 16.

N 80 91 19 1	. 1.7 s.gre(1) as	(2)	5 10 5 m m m (3)	Line 19 9 9 1 .
Mul	tiply ::003296	41.78		
By	5-782	. w. 4)2. 4:4 •08	29 2.00	06
	or de la Carlo	Territoria	24 2 317	23 h 23
Pro	duct -0190574	72 2.627	962	3 88 37
4. Multi	ply 3.2517 by	023.	Ans	0747891.
5. Multi	ply 64.001 by	340.	Ans	21760-34.
6. Multi	ply 482000 by	•37.		w. 178340.
7. Multi	ply 3782.4 by	00917.	Ans.	34.684608.
8. Multi	ply 87.96 by 2	20.	i, An	s. 19351·2.
( )	.55005	AT 94500 HVD 10		to the same

#### PROOF OF MULTIPLICATION

54. If the multiplier is not greater than 12, multiply the multiplicand by the multiplier, minus one, and add the multiplicand to the product. The sum should be the same as the product of the multiplicand by the whole multiplier.

If the multiplier be greater than 12 and the multiplicand an abstract number:—

FIRST METHOD.—Multiply the multiplier by the multiplicand, and if the product thus obtained agree with the other the work may be considered correct.

This method of proof depends upon the principle (40) that the product of two numbers is the same whichever is taken as multiplier.

SECOND METHOD.—Divide the product by one of the factors, and if the quotient thus obtained is equal to the other factor, the work is correct.

This is simply reversing the operation, i. e., breaking up the product into its factors.

TEIRD MATHOD.—Divide the sum of the digits of the multiplicane by 9 and set down the remainder; divide also the sum of the digits of the multiplier by 9 and set down the remainder; multiply these two remainders together, divide the sum of the digits in their product by 9, and if the remainder thus obtained is equal to the remainder abtained by dividing the sum of the digits in the product of the multiplicand and the multiplier by 9, the work is generally correct: if these two last remainders are different, it must be wrong.

EXAMPLE 1.—Let the quantities multiplied be 9426 and 3785.

Taking the nines from 8436, we get 3 as remainder.

And from \$785, we get 5.

47180 75408 \$×5=15,from which 9 being taken, 6 are laft. 65982

Taking the nines from 85677410, 6 are left.

The remainders being equal, we are to presume the multiplication is correct. The same result, however, would have been obtained even if we had displaced digits, added or omitted cyphers, or fallen into errors which had counteracted each other; but, with ordinary care, none of these are likely to occur.

EXAMPLE 2.—Let the numbers be 76542 and 8436.

Taking the nines from 76542, the remainder is 6.
Taking them from 8486, it is 3.

6×3=18, the remainder from which is 0. 612336

Taking the nines from 645706812 also, the remainder is 0.

The remainders being the same, the multiplication may be considered

NOTE.—This proof applies, whatever may be the position of the decimal point in either of the given numbers.

EXAMPLE 3.—Let the numbers be 4.63 and 5.4.

From 4'63, the remainder is 4. From 5'4, it is 0.

1852 4×0=0, from which the remainder is 0.

From 25.002 the remainder is 0.

55. The principle on which this process depends is, that if any number is divided by 9, and the sum of its digits also be divided by 9, the remainders, are, in both cases, the same.

Thus taking the number 7825, we have.

$$7835 = 7000 + 800 + 20 + 5 = 7000 + 800 + 20 + 5 = 7 \times 1000 + 8 \times 100 + 2 \times 10 + 5 = 7 \times (111 + \frac{1}{2}) + 8 \times (11 + \frac{1}{2}) + 2 \times (1 + \frac{1}{2}) + 5 = 777 + \frac{7}{4} + 88 + \frac{9}{4} + \frac{2}{3} + \frac{2}{3} + \frac{2}{3} = 777 + 88 + 2 + \frac{7}{4} + \frac{3}{3} + \frac{3}{4} + \frac{2}{3} = 777 + 88 + 2 + \frac{7+8+2+3}{4}$$

Hence the remainder arising from the division of 7825 by 9 is evidently the same as that arising from dividing 7+8+2+5 or 22, which is the sum of its digits, by 9.

56. Casting the nines from the factors, multiplying the resulting remainders, and casting the nines from the product, will leave the same remainder as if the nines were cast from the product of the factors—provided the multiplication has been correctly performed.

Thus, let the factors be 578 and 464.

Casting the nines from 5+7+3 (which we have just seen is the same as casting the nines from 573), we obtain 6 as remainder. Casting the nines from 4+6+4, we get 5 as remainder. Multiplying 6 and 5 we obtain 80 as product, which, when the nines are taken away, will give 3 as a remainder.

h is 0.

idered ecimal

that if also be

by 9 is 2+5 or

resultet, will he proen cor-

the nines

We see show that 3 will be the restaining, flor. If we cost the pines from the product of the factors; which is effected by setting down this product of the first that are equal to it—as follows:

573×654—(the product of the factors).

=(0×100+7×10+5)×(0×100+6×10+6)

= {5× (0+1)+7×(0+1)+5} × {4× (0+1)+6× (0+1)+6}

= 5×30+5+7×3+7+5)×(4×30+4+3×2+6+4)

5×30 expresses a number of nines; it will continue to do so when multiplied by all the quantities within the second brackets, and is, therefore, to be east out; and, for a similar reason, 7×2. Again 4×30 supresses a number of nines; it will continue to do so when multiplied by the quantities within the first brackets, and is therefore to be cast out; and for a similar reason, 6×9. There will show he left only (4+7+3)×(4+6+4)—from which the nines are still to be cast out, the reasonders to be multiplied together, and the nines so the cast from their product;—but we have done all this already, and obtained 8 as remainder.

#### CONTRACTIONS IN MULTIPETORTION.

57. I. To multiply by 5:

Affix a 0 to the maltiplicand and divide the feralt by 2.

Reason 5 = 40 Tomber 1... will a sile of he make flight

II. To multiply by 15: 1 to ough these s t us which the

Affin a 0 to the multiplicand and to the result add half of title. Reason 15 = 10 + 14.

III. To multiply by 25:

Affix two 0s to the multiplicand and divide the result by 4.

Reason 25 = 191.

IV. To multiply by 126 that then to it william a final man

Affix three 0s to the multiplicand and divide the result by 8.

19 8 19 21 (47 0) 21 1 2 2 1 2 1 2 2

Reason 125 = 1900.

V. To multiply by 75:

. To multiply by 75:

Affix two 0s to the multiplicand and from the result take one fourth of itself. 

VI. To multiply by 175:

Affix two 0s-multiply the result by 7 and divide by 4.

VII. To multiply by 275:

Affix two 0s-multiply the result by 11 and divide by 4.

Reason 275 = 1100.

VIII. To multiply by 13, 14, 15, &c., or by 1 with sither of the other digits afficed to it; out to site

The state of the s

EXAMPLE. Multiply by the unite' figure of the multiplier, 2325 × 13 and write each figure of the partial product one place to the right of that from which it arises; finally, add the partial product to the multiplicand, and the result will be the answer required.

REASON.—This is the same in effect as if we actually multiplied by the common method. We merely make the multiplicand serve for the second partial product.

IX. To multiply by 21, 31, 41, &c., or by 1 with either of the other significant figures prefixed to it:

BEAMPLE. Multiply by the tend figure of the multiplier,

365×21 and write the first figure of the partial product in

the tend place; finally, add this partial product to

the multiplicand, and the result will be the answer

required.

RMASON.—The reason of this method of contraction is substantially the same as that of the preceding.

X. To multiply by 101, 102, 103, 104, &c., or by 10 with either of the other digits affixed to it:

Multiply by the units' figure of the multiplier and write the partial product, thus obtained, two places to the right of the multiplicand—finally, add the partial product to the multiplicand.

REASON.—Substantially the same as No. 8.

XI. To multiply by any number of nines:

Remove the decimal point of the multiplicand so many places to the right (by affixing 0's if necessary) as there are nines in the multiplier; and subtract the multiplicand from the result.

EXAMPLE 1.—Multiply 7347 by 999.

7347×999=7347000-7347=7339653.

We, in such a case, merely multiply by the next higher convenient composite number, and subtract the multiplicand as many times as we have taken it too often; thus, in the example just given—7347×999=7847×(1000-1)=7347000-7347=7339653.

EXAMPLE 2.—Multiply 678943 by 999999.

678943×1000000 = 678943000000 678948×1 = 678948

078943×990009 == 678942321.057

EXAMPLE 3.—Multiply 78.9645 by 99993.

78·9645×100000=7896450 78·9645× 7= 552·7515

78-9645×99993 = 7895827-2485

XII. When it is not necessary to have as many decimal places in the product, as are in both multiplicand and multiplier—

ART. 57.7

other site

olier, t one rises : cand.

by the econd

ither

iplier. uct in luct to mewer.

lly the by 10

ite the multi-

laces to in the

nt comwe have

A300 : 151

ecimal d and

Reverse the multiplier, putting its unite' place under the place of that denomination in the multiplicand, which is the lowest of the required product.

Multiply by each digit of the multiplier beginning with the denomination over it in the multiplicand; but adding what would have been obtained, on multiplying the preceding digit of the multiplicand—unity, if the number obtained would be between 5 and 15; 2, if between 15 and 25; 3, if between 25 and 35, &c.

Let the lowest denominations of the products, arising from the different digits of the multiplicand, stand in the same vertical column.

Add up all the products for the total product; from which cut off the required number of decimal places.

Example 1.—Multiply 5.6784 by 9.7324, so as to have four decimals in the product.

Short method. 56784 42879	the contraction	Ordinary Method. 5.6784 9.7324
511058	an qui	22(7136
89749 1708	战三海 机单位	1708 58
118		89748 8 511056
mil : RKYGAQ . Chat yo	b mile it idea.	KN-9844 6016

9 in the multiplier expresses units; it is therefore put under the fourth decimal place of the multiplicand—that being the place of the lowest decimal required in the product.

In multiplying by each succeeding digit of the multiplier we neglect an additional digit of the multiplicand; because, as the multiplier decreases, the number multiplied must increase—to keep the lowest denomination of the different products, the same as the lowest denomination required in the total product. In the example given, 7 (the second digit of the multiplier) multiplied by 8 (the second digit of the multiplicand) will evidently produce the same denomination as 9 (one denomination higher than the 7), multiplied by 4 (one denomination lower than the 8). Were we to multiply the lowest denomination of the multiplicand by 7, we should get (53) a result in the fifth place to the right of the decimal point; which is a denomination supposed to be, in the present instance, too inconsiderable for notice—since we are to have only four decimals in the product. But we add unity for every ten that would arise, from the multiplication of an additional digit of the multiplicand; since every such fon constitutes one in the lowest denomination of the required product. When the multiplication of an additional digit of the multiplicand would give more than 5, and less than 15, it is nearer to the truth to suppose we have ten than either 0 or 20; and therefore it is more correct to add 2 than 1 or 3; do. We may consider 5 either as 0 or 10; 15 either as 10 or 20; do.

On inspecting the results obtained by the abridged, and ordinary methods, the difference is perceived to be inconsiderable. When greater accuracy is desired, we should proceed as if we intended to have more decimals in the product, and afterwards reject those that are unnecessary.

to the state of men services.

Example 2.—Multiply 8.76539 by 0.5765, so as to have three decimal places.

870888 A for a finished and a finish

There are no units in the multiplier; but, as the rule directs, we put its units place under the third decimal place of the multiplicand. In multiplying by a since there is no digit over it in the multiplicand, we marrily set down what would have resulted from the multiplying the preceding denomination of the multiplicand.

Example 3.—Multiply 0.23257 by 0.243, so as to have four

decimal places.

We are obliged to place a cipher in the product to make up the required number of decimals.

#### EXERCISE 17.

1. The canals in Canada amount to 216 miles in length, and their average cost was \$83469 per mile. What was the total cost of the canals of Canada?

2. The Great Western Railroad is 229 miles in length, and its cost was about \$61135.37 per mile. What was the total

cost of this road?

3. The Austrian empire contains 255226 square miles, and the population averages 143 per square mile. What is the entire population of the Austrian empire?

4. France contains 203736 square miles, and the population averages 176 per square mile. What is the entire population of

France ?

5. Great Britain contains 116700 square miles, and the population averages 235 per square mile. What is the entire po-

pulation of Great Britain?

6. The total number of Common Schools in operation in Canada West, during the year 1857, was 3721; allowing an average of 73 pupils to each, how many children were in attendance at the Common Schools?

7. 32000 seeds have been counted in a single poppy; how many

would be found in 297 of these?

8. 9344000 eggs have been found in a single cod fish; how many would there be in 35 such?

100

OHE

ired

and tal and tal

and ire ion

of

popq-

> in an

ny

9. Multiply 128 lbs. 4 os. 7 drs. 2 ser. 17 gr. by 149.

10. Multiply 1698732 by 999998.

11. Multiply 128 bush. 1 pk. 1 gal. 1 gt. 1 pt. by 640.

12. What will be the cost of a cheet of tes containing 89

lbs. at 78 cents per lb.?

13. How much cloth will it take to make the clothes for a regiment of soldiers containing 1148 men, if each suit requires

7 yds. 3 grs. 2 ns. 1 in.? 14. Multiply 1634 5789 by 635000.

15. A person dying bequeathed the whole of his property to his three sons. To the youngest he gave \$968.49; to the second, 3.4 times as much as the youngest; and to the eldest 3.7 times. as much as to the second. Required the value of his property.

### QUESTIONS TO BE ANSWERED BY THE PUPIL.

Norn.—The numbers after the questions refer to the articles of the section.

- 1. What is multiplication? (33) 9. What is the multiplicand? (36)
  3. What is the multiplier? (35) 4. What is the product? (36)
  5. Why are the multiplier and multiplicand called the factors of the pro-

- 3. Why are the multiplier and multiplierad called the factors of the product (36).

  8. What is a composite number? (37).

  7. What is a composite number? (38).

  8. If the multiplier be greater than unity, how will the product compare with the multiplierand? (39).

  9. If the multiplier be equal to unity, how will the product compare with the multiplierad? (39).

  10. If the multiplier be less than unity, how will the product compare with the multiplicand? (39).

  11. Show that either of the factors may be used as multiplier without altering the value of the product. (40).

  12. Show that when the multiplier is a composite number we may estain the entire product by multiplying by each of the factors in succession. (41).

  13. By whom was the multiplication table calculated? (48).

- 18. By whom was the multiplication table calculated? (49)
  14. How was it calculated? (48)
  15. What is the sign of multiplication? (48)
  16. How do we multiply a quantity consisting of several factors connected by the sign of multiplication? (44)
- 17. How do we multiply a quantity consisting of several terms, connected by the signs + and enclosed within a bracket? (45)

  18. What is meant by (7+3-2+5) × (2+3-7)? (45)

  19. Why do we begin multiplying a number at the right-hand side? (46)

  20. What is the rule for multiplication when the multiplier is not greater

- than 12? (47)

  21. What is the rule when the multiplier is a composite number, none of its factors being greater than 12? (48)

  23. What is the rule when the multiplicand is a denominate number, and the multiplier greater than 13; but not a composite number? (49)

  23. Show the absurdity of attempting to multiply one denominate number.

- by another. (50)

  24. When the multiplicand is an abstract number, and the multiplier conter than 13, but not a composite number, what is the rule? (51)

  25. When the multiplicand or multiplier, or both, contain decimals, what is the rule? (51)

- 6. Give the reason of this rule. (53 and 53)
  7. How do we prove multiplication when the multiplier is less than 13?(54)
  8. How do we prove multiplication when the multiplicand is an abstract number and the multiplier is greater than 13? (54)
  9. Upon what does the proof by casting out the nines depend? (55)
  10. Prove this principle. (55)
  11. Prove that casting the nines from the factors, multiplying the resulting remainders, and casting the nines from the product, will leave the same remainder as if the nines were cast from the product of the

- factors. (56)

  35. What short methods have we for multiplying by 5, 25 and 125? (57)

  36. How may we multiply by 175? How by 275? (57)

  36. How may we multiply by 13, 14, 15, &c.? (57)

  36. How may we multiply by 13, 14, 15, &c.? (57)

  37. How may we multiply by any number of nines? (57)

  38. How may we contract the work when we require only a limited number of decimals? (57)

## DIVISION.

- 58. Division is the process of finding how many times one number is contained in another.
  - 59. The number by which we divide is called the divisor.
  - 60. The number to be divided is called the dividend.
- 61. The number obtained by division, that is, the number which shows how many times the divisor is contained in the dividend is called the quotient (Lat. quoties, " how many times.")
- 62. If the divisor be less than the dividend, the quotient will be greater than unity.

If the divisor be equal to the dividend, the quotient will

be equal to unity.

If the divisor be greater than the dividend, the quotient will be less than unity.

68. It is sometimes found that the dividend does not contain the divisor an exact number of times; in such cases the quantity left after the division is called the remainder.

The remainder, being a part of the dividend, is, of course,

of the same denomination.

The remainder must be less than the divisor—otherwise the divisor would be contained once more in the dividend.

64. Division is merely a short method of performing a particular kind of subtraction (Art. 6, Sec. II.) The dividend corresponds to the minuend, the divisor to the

?(54)

(57)

ober .

nos

or.

d. 3.

ım-

red

OW

ent

vill

ent

tot

368

er.

se,

180

nd.:

2 8 H

VI-

he

subtrahend, and the remainder to the difference. quotient has no corresponding quantity in subtractionsince it simply tells how many times the divisor can be subtracted from the dividend.

It will help us to understand how greatly division abbreviates subtraction, if we consider how long a process would be required to discover—by actually subtracting it—how often 7 is contained in 8563495724, while as we shall find, the same thing can be effected by division in less than a minute.

65. Since the quotient shows how many times the dividend contains the divisor, it follows that the divisor and quotient are the factors of the dividend. Hence if the divisor and quotient be multiplied together, and the remainder, if any, added to the product, the result will be equal to the dividend.

66. We have three ways of expressing the division of one quantity by another:-

1st. By the sign: + written between them; thus, 15+ 3 = 5.

2nd. By the sign: written between them; thus, 15:3-5. 3rd. By writing the dividend above and the divisor below a horisontal line; thus, 4 = 5.

Two quantities written thus 1<sup>4</sup> constitute what is called a fraction, and the expression is read six-elevenths.

It is usual and proper to write the remainder obtained in division, in the form of a fraction; thus 17+3 gives 5 as a quotient and 2 as a remainder. Now the remainder, 2, is written above the line; and divisor 3 below the line; the whole quotient being expressed thus 5\frac{1}{2}\$ (read five and two-thirds); the meaning of which is, that 3 is contained in 17, 5 times and \frac{1}{2}\$ of a time.

- 67. When a quantity consisting of several terms connected by the sign of multiplication is to be divided, dividing any one of the factors will be the same as dividing the product; thus 5×10×25÷5=4×10×25, for each is equal to 250.
- 68. When a quantity consisting of several terms connected by the signs + and -, contained within brackets, is to be divided, it is necessary, on removing the brackets, to put the divisor under each of the terms of the quantity;

6+3-7+9 thus  $(6+3-7+9) \div 3$ , or we do not divide the whole unless we divide all its parts.

69. It will be seen from (68) that the horizontal line

OP

tho

12)

i. e.

of t in a sen 71d

spe

which separates the dividend from the divisor assumes the place of a pair of brackets when the dividend consists of several terms; and, therefore, when the quantity to be divided is subtractive, it will sometimes be necessary to change the signs, as already directed (26); thus:

15-6+9 27-15+6-9 27 3 6 + 13 - 3

70. Example 1. Let it be required to divide 798 by 3.

OPERATION. EXPLANATION.—Place the divisor a little to the left of the dividend and separate them by a short ourve line. Also draw a straight line beneath the dividend. 2âñ

600 180 18 = 200 600+180+18 798 700-1-90-1-8 600+190+8 +60+6=266 (See 68).

Instead of going through this long operation it is evident that we may proceed as follows: 3 units into 7 hundreds will go 2 (hundreds) times and leave a remainder 1, which being of the order of hundreds, is equal to 10 tens: 10 tens and 9 tens make 19 tens, and 3 into 19 goes 6 (tens) times and leaves a remailider 1, which, being of the order of tens is equal to 10 tinits; 10 units and 8 units make 18 units, and 3 units into 18 units goes 6 (units) times.

EXAMPLE 2. Let it be required to divide 917 lb. 13 oz. 12 dr.

OPERATION. EXPLANATION.—Placing the dividend and divisor as before,
1b. os. dr.
4)917 16 18
18 (tens) times and 1 over; 1 hundred, equal to 10 tens, and 1 ten make 11 tens; 4 in 11, 2
1919 7 7 hundred, equal to 10 tens, and 1 ten make 11 tens; 4 in 11, 2
1920 7 7 hundred, equal to 10 tens, and 1 ten make 11 tens; 4 in 11, 2
1920 7 7 hundred, equal to 10 tens, and 1 ten make 11 tens; 4 in 11, 2
1920 7 7 hundred, equal to 30 units, and 7 units
10 make 37 units; 4 in 37, 9 times and 1 over, which is 1 os., since the 20 are 0s.;
10 make 39 os., 4 in 39, 7 times and 1 over, which is 1 os., since the 20 are 0s.;
10 make 39 os., 4 in 39, 7 times and 1 over, which is 1 os., since the 20 are 0s.;
10 make 39 os., 4 in 39, 7 times and 1 over, which is 1 os., since the 20 are 0s.;
10 make 37 units, 4 in 30, 9 times and 1 over; which is 1 os., since the 20 are 0s.;
10 make 39 os., 4 in 39, 7 times and 1 over; which is 1 os., since the 20 are 0s.;
10 make 37 units, 4 in 37, 9 times and 1 over; which is 1 os., since the 20 are 0s.;
10 make 37 units, 4 in 37, 9 times and 1 over; which is 1 os., since the 20 are 0s.;
10 make 37 units, 4 in 37, 9 times and 1 over; which is 1 os., since the 20 are 0s.;
10 make 37 units private the 30 units, and 7 units

Example 3. Let it be required to divide 9789 by 26.

which we do by placing the 26 under the 13. as is explained in (Art. 66).

The complete quotient is therefore 3765 read 376 and thirteen-twentysixths or 876 and 18 divided by 26.

the

to of o be

ry to

Asset) of the · c. 2 (10%) 5

= 200

May

tens; COVES

inita)

dr.

efore.

er 1 11,3 unita

1 16.

1303.

05. es. ient.

visor

f in the 3 ply-sub-er 19

nd 8 ulti-sub-

ens.

169

not

71. From the preceding illustration and examples we deduce, for the division of numbers, the following general

BULE.

Beginning with the highest order of units in the dividend, pass on to the lower orders until the Jewest number of Agures be found that will contain the divisor; divide these Agures by it, for the first figure of the quotient; this figure will be of the same order as that of the lowest used in the partial dividend.

Multiply the divisor by the quotient figure so found, and subtract the product from the dividend, being careful to place units of the same order in the same vertical column. Reduce the remainder to units of the next lower order, and add in the units of that order found in the dividend: this will furnish a new dividend.

Proceed in a similar manner until units of every order shall have

been divided.

EXAMPLE 1 .- Divide 98765 by 7.

OPERATION. BEPLATATION.—Here we say 7 in 9, 1 and 2 over; in 28
7)98765 4 and 0 over; in 7, 1 and 0 over; in 6, 0 times and 6 over; in
65, 9 and 2 over; Beneath this 2 we write the divisor 7 to inlates division. We may, however, carry on the division by considering the 2 units reduced to tenths, &c., and the quotient becomes 14109 2857.

Thus 2 units, equal to 20 tenths, 7 in 20, 2 and 6 over; 6 tenths are equal to 60 hundredths, 7 in 60, 5 times, and 4 over; 4 hundredths are equal to 40 thousandths, 7 in 40, 5 and 5 over; 5 thousandths are equal to 50 tenths of thousandths, &c.

Example 2.—Divide 124789 by 12.

EXPLANATION.—Here again we may either stop at the units and write the remainder 1 over the divisor 12, or we may reduce the 1 unit to tenths, &c., as in the second ope-OPERATION. 12) 194789 10309 1 ration.

Triff of to give the of the bus his grant was

Ord 12) 124780

EXAMPLE 3.—Divide £1986 14s. 71d. by 9.

OPERATION. EXPLANATION.—2 in 19, 2 and 1 over; 9 in 18, 2 and 0
9) £1986 14 7 over; 9 in 6, 0 and 6 over; £6 are equal to 120c. and
14c. make 134c.; 9 in 134 14, and 8 over; 8c. are equal
£220 14 11 to 96d. and 7d. make 103d.; 9 in 103. 11 times and 4 over;
4d. are equal to 16 farthings and 2 farthings make 18 farthings; 9 in 18, 3,
i. e. one minth of 18 farthings is 2 farthings, written thus 1d.

LA Clieby Fry Corelling Copper

72. In example 3, we are, in reality, required to find one-ninth of the dividend. The obvious meaning is, not that 9 is contained in £1986 14s. 74d. £220 14s. 114d. times, which would be nonsense, but that £220 14s, 111d, is the ninth part of £1986 14s. 71d.: so also in all similar questions.

Notwithstanding this, all such examples are reducible to a species of subtraction. Thus, in the above example, we for the

78 dete nece

pe m divis mult

grea

the

the 1 quot

moment, consider the divisor 9 to be of the same denomination as the dividend, and ascertain how many times £9 will go into (i. e., can be subtracted from) £1988. We get, as a result, 220 times, and a remainder of £6. Then we argue, from the principles already established, that since £9 is contained in £1986 220 times, with a remainder of £6; £220 is contained in £1986. 9 times, with a remainder of £6; that is, that the ninth part of £1986 is £220, with a remainder £6. Next reducing this £6 to shillings, and adding in the 14s., we obtain a total of 134s., and we find that 9s. is contained in 134s. 14 times, with a remainder of 8s., whence we conclude that 14s. is contained in 134s. 9 times, with a remainder of 8s., that is, that the ninth part of 184s. is 14s. with a remainder of 8s., or that the ninth part of £1986 14s. is £220 14s., with 8s. still undivided, &c.

EXAMPLE 4.—Divide 978964 by 8429.

DIPERATION.

EXPLANATION.—3489 into 9789 (the smallest numproperty of figures that will contain the divisor) goes 3
times, we therefore put 3 in the quotient. Hultiplying 3489 by 2, we get 6888, which we subtract from
20316 5789; and obtain as remainder 2031, which we reduce
27408 to the next lower order (tens) and add in the 6 tens,
3489 into 20316 goes 8 times. We therefore place 8
in the quotient. Multiplying 3429 by 8 we get 97433,
17145 which we subtract from 20316, and obtain 1834 as a
remainder. Reducing to units and adding in the 4,
or what amounts to the same thing, bringing down
the 4 and writing it after the 1884 we get 18344; and
the divisor 3439.

the divisor 8490

78. When the dividend is an abstract number, it is evident that bringing down the next figure and writing it to the right of the remainder, is the same in effect as reducing the remainder to the next lower denomination and adding in the units of that order found in the dividend. Thus, in the last example, bringing down the 6 and writing it directly to the right of the first remainder, 2931, makes the next partial dividend 29316, which is the same as reducing the 2931 to the next lower order and adding to the result the 6 of that order found in the dividend.

EXAMPLE 5.—Divide 6421284 by 642,

OPERATION: EXPLANATION.—642 goes once into 648, and leaves no remainder. Bringing down the next digit of the dividend gives no digit in the quotient, in which, therefore, we put a cipher after the 1. The next digit of the dividend, in the same way, gives no digit in the quotient, in which, consequently, we put another cipher, and, for similar reasons, another in bringing down the next; but the next digit makes the quantity brought down 1884, which contains the divisor twice, and gives no remainder:—we put 2 in the quotient.

Norn.—After the first quotient figure is obtained, for each figure of the dividend which is brought down, either a significant figure, or a cipher, must be put in the quotient, without his believed ARTS. 78-75.]

The Think of the Control of the Cont

D. MILES

4月 194以 数如此病!

A Transport LA

ation into t, 220 prin-£1986 £1986 art of £6 to and inder

times,

6 148.

t numgoes 2
Multict from
reduce 6
tens,
place 8
t 97433,
884 as a
n the 4,
g down
ld: and

vident ght of ainder of that bringne first which er and idend.

aves no be diviserefore, te divitent, in and, for tt; but ins the

each ificant 74. Why there is a remainder, we may continue the division, adding decimes to the quotient, as follows—
EXAM: 6.—Divide 796347 by 847, and the result by 7234.

847)708847 (940:197166, &c.
7628

3404
3588

1670
847

8230
7623

6070
5929

1410
847

5880
5088

7234)940·1971#6(0·129969, &c.

216·79 144·68 78·117

THE THE PARTY EN TO THE A

41、果然而是 140 元 一种国际

7:0111 6:5106 -50056 -48404

66526 65106

1420, &c.

75. When the divisor is large, the pupil will find assistance in determining the quotient figure, by finding how many times the first figure of the divisor is contained in the first figure, or, if necessary, the first two figures of the dividend. This will give pretty nearly the right figure. Some allowance, must, however, be made for carrying from the product of the other figures of the divisor, to the product of the first into the quotient figure. After multiplying the divisor by the quotient figure, if the product is greater than the corresponding partial dividend, this shows the quotient was taken too great, and must be diminished. If the remainder, after subtraction, is greater than the divisor, the quotient was taken too small, and must be increased.

mi in

na

vide

Th

34m

obtai Sin

# EXAMPLE 7.—Divide 279 cwt. 3 grs. 14 ib. 9 os. by 129.

16 = drams in oz.

1184 = drams. 1116

23 remainder.

i. e., the 129th part of 279 cw., is 2 cwt., with a remainder of 21 cwt. With a remainder of 21 cwt. With a remainder of 21 cwt. We reduce to quarters by multiplying by 4 and adding in the 3 qrs. The 129th part of 87 qrs. is equal to 0 qr. and we therefore place a 0 in the quarters' place of the quotient. We next reduce qrs. to lbs. by multiplying by 25 and adding in the 14lbs. of the dividend. We thus obtain 2189 lbs., of which the 129th part is 16 lb., with an undivided remainder of 125 lbs. Reducing 125 lbs. to 02., and adding in the 9 cs., we obtain 2009 oz., of which the 129th part is 15 os., with an undivided remainder of 74 oz. to drams, we obtain 1184 drams, of which the 129th part is 2 drams, with an undivided remainder of 23 drams, under which we place the divisor 129 to indicate its division. Thus we find the total quotient to be 2 cwt. 9 qr. 16 lb. 15 oz. 9,23 drs.

76. The general principles on which the operations in division depend are:

1st. The quotient arising from the division of the whole dividend by the divisor, is equal to the sum of the quotients arising from the division of the several parts of the dividend by the divisor. (68)

2nd. The divisor and quotient are the factors of the dividend. (65)

and. The product of the divisor, by the entire quotient, is equal to the sum of the products of the divisor by the several parts of the quotient. (45)

6 in 189, 279 cw., ainder of we reduce plying by qrs. The i. is equal fore place place of

qrs. The
l. is equal
fore place
place of
next remultiplygs in the
end. We
of which
ilb., with
ainder of
25 lbs. to
the 9 oz.,
of which
i oz., with
nder of 74
74 oz. to
84 drams,
part is 9
ivided rens, under
e divisor
divisor
divisor,

otal quoor. 16 lb.

tions in

e whole rotients he divi-

the di-

ptient,

YYA TOTAL

We said how many tieses the divisor is contained in a part of the divisor dend, and thus a part of the quotient is found; the product of the divisor by this part is taken from the dividend, showing how much of the latter remains undivided; then a part of the remaining divident is taken and another part of the quotient is found, and the product of the divisor, by it, is taken away from what before remained; and thus the operation are ceeds till the whole of the dividend is divided, or till the remained; it less than the divisor.

77. We begin at the left-hand side, because what remains of the higher denomination may still give a quotient in a lower; and the question is, how often the divisor will go into the dividend—its different denominations being taken in any convenient way. We cannot know how many of the higher we shall have to add to the lower denominanations, unless we begin with the higher.

#### PROOF OF DIVISION.

78. FIRST MATHOD. Multiply the quotient by the divisor, and to the product add the remainder, if any; the result should be equal to the dividend. (65)

EXAMPLE 8.-Divide £5681 18s. 4d. by 700.

A PARTY A	s. d. PROOF	20 18 36 1	"科特"的线。
700)5881 13 4 (8	Total Burning Barrell B. d		ercon publishment frame-subservation with
1 1 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	46 4 4 2 5 B	estroi :	130203
(C.Com	(1)	0	1.46)
it will a so that we will are	5 .8 1 81 8	A STORY	200 8
1688	7 m / 6 1/2 1 1	M. Garage	Drotter or
1400	811 18	Andrew description to the articles	मान्येका इत्यान्त्रे सम्बन्धानामा । अन्यमान्य वृत्यानीकृतिः
600	To all by	Classified and	43 44 41 12
233	5681 18-	28 26. 4d ×7	oosdividend.
101 33 4 3		- TIT 1818	La Maratant
2800		1. 1. 80 M.	statistic re

SHOOND MATHOD.—Subtract the remainder, if any, from the dividend, divide the dividend, thus diminished, by the quotient; and if the result is equal to the given divisor, the work is right.

This is merely doing the same work by a different method.

THIRD METHOD.—Cast the nines out of the divisor and quotient, and thattiply the remainders together; and to their product the remainder, if any, after division, and cast the nines out of this sum; the remainder thus obtained should be equal to the remainder obtained by casting the nines out of the dividend.

Since the divisor and quotient answer to the multiplier and multiplicand, and the dividend to the product, it is evident that the principle of casting out the Ga will apply to the proof of division as well as to that of multiplication.

FOURTH MATHON .- Add the remainder and the respective products

ni di th

th

of the divisor into each quotient figure together; and if the sum is equal to the dividend, the work is right.

This mode of proof depends upon the principle that the whole of a quantity is squal to the sum of all its parts.

BEAMPLE 9 .- Divide 147856 by 97.

la Grano a a lichide ce co

the wair's a come of.

willed and maintain to the

in a word of made, many

waterio need to respect out of is

it the water summers and so it

		97)147856(1524		1 4 1 ALM 18 1 1 1
vi. min	the freezewant	97)147856(1524	i dut si	They have the

( ) T#1000(T092		-4 . 3 . 51	1/3	
13a 970 m	lit in th	West 194	4 19 224 4 12 12	
485°	selmi	with the	er ihr i	
, .k	9/1/25	1 4 10 %	o' mai	
285	The party	Mr. Y	Tri to	
M . and the .	Ty 18	16 P. BUS	1 160 313 }	
416 388	1 3 1/2	· Apple	131 Ty 13.	
14.1 1.1 28° (1) A	1 111	Markett &	41 1 1 1	
147856				

Norm.—The asterisks show the lines to be added.

. 19 1001 1

# Exercise 18

(1) 12)876967	7)891023	(3) 9)76345		8)654	(4) 132·8	78	See that
73080 <sub>7</sub> <sup>2</sup> s (5) \$ cts. 9)6789·60	127289 (6) \$ cts. 11)4298·76	8482 (7) £ s. 4)19 6	84 d.	wks. 9)69	(8 ds.		min.
\$754.40	\$390.79	4 16	7	7	5.	4	50
<ol> <li>Divide £1</li> <li>Divide 56</li> <li>Divide 67</li> <li>Divide £4</li> <li>Divide \$5</li> </ol>	8965 by 6423. 176 14s. 6d. by 789 by 741. 85158 by 7894 1728 16s. 2d. b 97896 64 by 42 0763 by 6.	12. y 317.	innet.	Committee of the State of the S	£14 Ans 14 1 ns, \$	1 14s. Ins. 7 2. 859 8s. 4	514d. 9318.
16. Divide 71 17. Divide 97 18. Divide 72	234 by 9. 7076 by 47600 80 lbs. 6 oz. 4	drs. 2 sci	. 13 7 oz.	grs. by 5 dr. 0	Ans 498.	. 20 <del>2</del>	5145 5878.
20. Divide 78	67674 by 9712 2 m. 3 fur. 38	il-to him:	W. L. 200	A. An	Ans	. 810	954

fa quan-

1 112 My 1

of the in

the sad to sit

# alarmon brodieso General Principles. Compa i Can

79. If a given divisor is contained in a given dividend a certain number of times, the same divisor will be contained in double that dividend twice as many times; in three times that dividend thrice as many times, &c. Hence,

When the divisor remains the same, multiplying the dividend by any number has the effect of multiplying the quotient by the same number.

Thus  $9 \div 3 = 3$ ;  $9 \times 2$  or  $18 \div 3 = 6 = 3 \times 2$ ,  $9 \times 5$  or  $45 \div 3 = 15 = 3 \times 5$ , &c.

80. If a given divisor is contained in a given dividend a certain number of times, the same divisor will be contained in half that dividend half as many times; in one-third of that dividend one-third as many times, &c. Hence,

When the divisor remains the same, dividing the dividend by any number, has the effect of dividing the quotient by the same number.

Thus 48:3 = 16; 44:3 or 24:3 = 8 = 46; 44:3 or 6:3 = 2 = 46, &c.

81. If a given divisor is contained in a given dividend a certain number of times, half that divisor will be contained in the same dividend twice as many times, one-third of that divisor thrice as many times, &c. Hence,

When the dividend remains the same, dividing the divisor by any number has the effect of multiplying the quotient by that number.

Thus 48:6=8; 48:3 or 48:3=16=8×2; 48:3 or 48:2=24=8×8, &c. 82. If a given divisor is contained in a given dividend a certain number of times, twice that divisor will be contained in the same dividend only half as many times, three times that divisor only one-third as many times, &c. Hence,

When the dividend remains the same, multiplying the divisor by any number has the effect of dividing the quotient by the same number.

Thus 48:2 = 24; 48: twice 2 or 48:4 = 12 = half of 24.
48: eight times 2 or 48:16 = 3 = one-eighth of 24, &c.

83. If a given divisor is contained in a given dividend a certain number of times, twice that divisor is contained in twice that dividend the same number of times; thrice that divisor in thrice that dividend the same number of times, &c. Hence,

When the divisor and dividend are both multiplied by the same number, the quotient will remain unchanged.

Thus 12:43; 24 or twice 12:8 or twice 4=3; 72 or thrice 24:24 or thrice 3=3, &c.

78 2225

hrs. min. 19 30

4 50 1248713. 14s. 61d. ns. 76473. . 8594813. 8s. 4517d.

228 19<del>11</del>3 3 8333+ ns. 7914<del>§</del>

2024878

12433 gr. 53d. 487. 8103714. 1r. 14 rds. the offer

my simi

Banay time, 20.

Ar mad W

र याच्या १ वे पार्थन

data fix

fin

W

ed

abi

84. If a given divisor is contained in a given dividend a certain number of times, half that divisor is contained in half that dividend the same number of times; one-third that divisor in one-

When the divisor and dividend are both divided by the same number, the quotient will remain unchanged.

Thus 40+34=3; 36 or belf of 48+18 or half of 24=3, 40,

# TO DIVIDE BY A COMPOSITE NUMBER

#### BULE.

85.—Divide the dividend by one of the factors of the divisor; then the resulting quotient by another factor; and so on till all the factors are used. The last quotient will be the passer.

Multiply each remainder by all the preceding divisors and addition produces to the first remainder. If any, for the true remainder.

When the divisor is separated into only two factors, the rule for finding the true remainder may be thus expressed

Multiply the last remainder by the Arit divisor, and to their product and the Arit remainder, if any; the recilit will be the true

EXAMPLE. Divide 718 lbs. by 72.

3)795 Elvis (a	lit rema	mder	li mba	ni.
4)250-1	and rema	inder=5×8	TOTAL S	16,
6)59-8	ard rema	inder—5×4	× 8= 60	lb.

true remainder to a to Wilk . Ast 979

That dividing by the factors of a number will give the same quotient as dividing by the number, itself, follows directly from Art. 84.

In the last example, dividing by 3 distributes the 718 lbs. into 250 parcels of 8 lbs. each, and leaves a remainder of 1 lbf.; dividing next by 4 distributes the 230 parcels into 50 atill larger percels, each containing 4 of the supplier of 3 lbs. percels, and leaves a remainder 5, which is not 5 lbs. but 5 percels, each of 5 lbs. lbs. but 5 percels, each of 5 lbs. lbs. but 5 percels, each of 5 lbs. cach, and leaves a remainder 5, which is, of course, 5 of the 13 lb. parcels. Hence the reason of the rule for anding the true remainder.

TOTAL OF OR A	Sand the second of the second
1. Divide 3766 by 25. 2. Divide 26406 by 42	62816. The ven divisor is contained in a gi
3. Divide 25431 by 98	the control description will be a second to the control of the con
4. Divide £24 178. 6d	
5. Divide £740 13s. 4	d. by 49
6. Divide £847 120. 4	
7. Divide 6788436 by	36 line duchous out and And 19398344
8. Divide 753293 by 1	47 (=7×7×3) Ans. 5124 4.

Ans. 22 lbs. 2 oz. 9 dwt. 04f grs.

A Only I minet

with the divisor

to prepare the

Jung of the original said

h enthy there.

I TO THE WHITE THE THE

Children of my of the particle for the WEARING TO STANKE THE

3. 4.00.30 + 28.004 A (8.7.1) It is gridly much the to gright

di shiming decimile, it r delvib on continuos

rtain

31445

word in

86. When both the divisor and the dividend are denominate numbers ... o for Lara of the action of

Reduce both the divisor and the dividend to the lowest denomis nation contained in either, and then proceed as in Art. 71.

EXAMPLE 1.—Divide £37 5s. 91d. by 3s. 61d.

		distance Staffing			11	
	8 61	2 87.	5 94 1431	12 0	A 17	88.
ř	18	13. (51) 20	. (11 6	Alta nh	HAT OF THE	147 6
	42 2 75%	TT 70 10 745	ni di	S. Stanto	was den	1 00 5 221
,	Late & South	] _nife_1	DI FILE	divib v	il coldi	i din i
	170 farthi	ngs. 8940				

ton promet and held it. mily sign. "his how is the his two 170 might that give in all of the

(4) is anterbur, and cherolites. Her floren es a hiertebe construit. This reception the ducional plant were at eater for it, in his buch 87. In the above and all similar questions we are required to find what fraction the divisor is of the dividend; or, in other words, how often the divisor is contained in, or can be subtracted from, the dividend, and the quotient must necessarily be an abstract number. . . been write at to the s we done and

EXAMPLE 2.—Divide 729 cwt. 3 qrs. 16 lb. by 3 qrs. 9 lb. 7 os.

3 9 25		729 8	à 16
===	-		1.3 3
84		919	
16	, 3 0	, 25	1 3 tale
511	14		LED THE
84	No.		ntonwaya. Novi tetk
351 oz			11° 19
			distribute in
et	1 . V 435	946	
	th 720		

I wire with the food of the first and be delined to the court of the court of

metrical co 8708 0 - 190-00 organic a deline of - coodspers = P8108 of figores / f. 8006 ASTUTE A CHICAGO 112 COUNTY 11 TO THE 9. 0878-03-1-0-7071 cm 00 not 700 4

### Bankoise 20.

- 1. Divide £8968 13s. 7jd. by £491 12s. 0jd. And 1811499
- 2. Divide 1027 m. 1 fur. 6rds. by 17 m. 5 fur. 27 rds. Ans. 58. 8. Divide £171 ls. 104d. by £57 0s. 74d.
- 4. Divide 9lb. 9 oz. 8 dwts. 12 grs. by 5 dwts. 9 grs. das. 436.
- 5. Divide 2866 acres 8 roods 36rds, by 91 acres 6 rds. Ans. 26.
- 88. When the dividend alone contains decimal places, the preceding rules are sufficient; but when the divisor contains decimals, it becomes necessary to prepare the quantities for division according to the following—

#### BULE.

Remove the decimal point as many places to the right in both the dividend and the divisor, as there are decimals in the divisor, and then proceed as in Art. 71.

This is simply multiplying both dividend and divisor by the same number, and therefore (Art.83) does not affect the quotient. Thus removing the decimal point one place to the right, in both dividend and divisor, is equivalent to multiplying each by 10; two places, the same as multiplying each by 100; three places, by 1000, &c.

EXAMPLE 1.—Divide 87-6 by .0009

Multiplying each by 10000, or, in other words, removing the decimal mint four places to the right, in each, (since there are four decimals in the divisor,) gives us 876000-19, and this (Art. 88) must give the same quotient as 87.6. 0009, therefore 87.6. 0009 = 876000 - 9 = 9783888, &c.

EXAMPLE 2.—Divide .06 by 8.934.

·06 ÷ 8·934 = 60 ÷ 8934.

8984)60.000(0.0067, &c.

53.604 6.8960 6.2538

1400

Removing the decimal point three places to the right, in each, we get 60:-8934, and we then proceed thus: 8934 into 60 (units), 0 (units) times; set down 0 with the decimal point after it; 9894 into 600 (tenths), 0 times; into 6000 (thionandths), 6 (thousandths) times, &c.

EXAMPLE 3.—Prepare 93.004 - 0000069 for division.

Ans. 93.004 - 0000069 = 930040000 - 69.

# Exercism 21.

- 1. 43 ÷ 0006947 = 430000000 ÷ 6947.
- 2.  $9378 \cdot 92 \div 9 \cdot 7891 = 93789200 \div 97891$ .
- 3.  $4.96723 \div 23.934 = 4967.23 \div 23984$
- 4. ·793 ÷ ·49 = 79·3 ÷ 49.

3.

136.

26.

ces.

isor

the

th the

, and

5. ·001 + 674·937=1 + 674937.

8. Divide 47.955 by 4.5.

ARTS. 88-94.7

7. Divide 756.98 by 76.73612.

8. Divide 47.5782975 by 28.175.

9. Divide 1 by 7.6345.

10. Divide 75.347 by 0.3829.

11. Divide .0002 by .000000008

Ans. 10-50.

Ans. 9.864-4. Ans. 1.8177.

Ame. 0.130

Ans. 198.7791 Ans. 25000.

### CONTRACTIONS IN DIVISION.

89. To divide by 10, 100, 1000, &c.

Remove the decimal point as many places to the left in the dividend as there are 0s in the divisor.

**90.** To divide by 25.

Multiply by 4 and divide by 100.

Reason 25 = 190.

91. To divide by 15, 35, 45, or 55.

Double the dividend, and divide the product by 30, 70, 90, or 110, as the case may be.

REASON.—This method is simply doubling both the divisor and dividen We must therefore divide the remainder, if any, by 2, for the free mainder.

To divide by 125.

Multiply the dividend by 8, and divide the product by 1000.

REASON.—This contraction is multiplying both the dividend and divisor by 8. For the true remainder, therefore, we must divide the remainder, if any, by 8.

93. To divide by 75, 175, 225, or 275.

Multiply the dividend by 4, and divide the product by 300, 700, 900, or 1100, as the case may be.

REASON.—75 = 390 175 = 190, &c. For the true remainder, divide the remainder, if any thus found, by 4.

94. When there are many decimals in the dividend and but few are required in the quotient, we may abbreviate the division by the following-

Proceed as in Art. 71 till the decimal point is placed in th quotient, and then cut off a digit to the right hand of the divisor each new digit of the quotient; remembering to carry what have been obtained by the multiplication of the digit nagle unity if this multiplication would have produced more than less than 15; 2 if more than 15, and less than 25, &c.

v the tient. both v.10;

laces,

ecimal in the uotient

we get times; times; andths)

- 69

### -LB.-Divide 764 337385 by 61.347.

Method. 7)75-487-885(12-296 61847				00 618	ntracted 17)754837 61347	Method. 885(12:296	1
	14006	7. 6.		,	140867 : 122694 :	es e e	
ø	1817 1226	8·8 9·4		` .	18173	al atmir	
		3-98 1-25	- ,	(11)	5904° 5521		
		2.755 3.082	nin c	30	383		
	1	4:6780		. \	15	7- 1/2	

According as the denominations of the quotient become small, their products by the lower denomination of the divisor become inconsiderable, and may be neglected, and consequently, the portions of the dividend from which they would have been subtracted. What should have been carried from the multiplication of the digit neglected—since it belongs to a higher denomination than what is neglected—must still be retained.

#### EXERCISE 22.

1. The Ontario, Simcoe, and Huron Railway is 95 miles in length, and cost \$3300000. What was the cost per mile?

2. The Rideau Canal is 126 miles in length, and cost \$3860000.

What was the average cost per mile?

3. The distance of the earth from the sun is 95270400 miles; how long would it take a cannon ball, going at the rate of 28800 miles per day, to reach the sun?

4. The national debt of France is 1145012096 dollars, and the number of inhabitants is 35781628; what is the amount of

indebtedness of each individual?

5. The national debt of Great Britain is 3764112127 dollárs, and the number of inhabitants is 27475271; what is the amount of indebtedness of each individual?

6. What is the ninth part of \$972?

7. What is each man's part, if \$972 be divided equally among

8. Divide a legacy of \$8526 equally between 294 persons.
9. Divide 340480 ounces of bread equally between 792 per-

sons. A cubic foot of distilled water weighs 1000 ounces; what

the the weight of one cubic inch?

The many Sabbath days' journeys (each 1155 yards) in the law is day's journey, which was equal to 33 miles and 2 follows: English?

12. How many pounds of butter, 19 cents per lb., would pur-

muse a cow, the price of which is \$47.50?

13, Divide 978.634 by 96.34762,

n 0.

of

hd

of

nt

hg

in

14. Divide 729 bush. 1 pk. 1 gal. 1 qt. 1 pt. by 297.

15. Divide 179 cwt, 3 qr. 4 lb. 16 os. by 9 lb. 7 os. 8 drs.

16. The circumference of the earth is about 25000 miles; if a vessel sails 93 m. 4 fur. 7 rds. a day, how long will it require to sail round the earth?

# QUESTIONS TO BE ANSWERED BY THE PUPIL.

NOTE.—The numbers after the questions refer to the articles of the

Note.—The numbers eyes section.

1. What is the division? (56)

2. What is the dividend? (50)

3. What is the dividend? (90)

4. What is the quotient? What is the derivation of the word 'quotient' (61)?

5. Explain when the quotient will be equal to unity, and when greater or less than unity. (62)

6. Under what circumstances does a remainder arise in division? (62)

7. What is the denomination of the remainder? (63)

8. Why can it never be as great as the divisor? (63)

9. What is the correspondence between the minuend and the subtrahend in subtraction and the divisor and the dividend in division? (64)

10. What may we consider as the factors of the dividend? (65)

11. How many ways have we of expressing the division of one quantity by another? What are they? (66)

12. When a quantity consisting of several terms, connected by the signs is to be divided by any number, how may the work be performed? (13). When a quantity consisting of several terms, connected by the signs or —, contained within brackets, is to be divided, what must be done upon removing the brackets? (68)

or —, contained within brackets, is to be divided, what must be done upon removing the brackets? (68)

14. Give the general rule for division. (71)

15. In the question "Divide 11 m. 7 fur. 20 per. 3 yds. by 279," emakin what is really required. (72) Show that all such questions are reducible to a species of subtraction. (73)

16. In dividing abstract numbers, explain what bringing down the next figure of the dividend is equivalent to. (73)

17. When there is a remainder, how is it to be written? (71, Rample 1)

18. What are the three general principles upon which the operations of division depend? (76)

division depend? (76) Why do we begin dividing at the left-hand side? (77)

19. Why do we begin dividing at the total and the second s

The dividend remaining unchanged, what effect has dividing the divisor

 The dividend remaining unchanged, what effect has dividing the divisor by any number? (81)
 The dividend remaining unchanged, what effect has multiplying the divisor by any number? (82)
 What is the effect upon the quotient when the divisor and the dividend are both multiplied by the same number? (83)
 What is the effect upon the quotient when the divisor and the dividend are both divided by the same number? (84)
 How do we divide by a composite number? (85)
 When we divide by the divisors of a composite divisor, how do we obtain the correct remainder? (85)
 When the divisor is separated into only two factors, how may the rule for obtaining the correct remainder be worded? (85)
 When the divisor and the dividend are both denominate numbers, what is the rule? (85)
 When one denominate number is divided by another, what kind of a number must the quotient always be? (87) number must the quotient always be? (87)

- 82. In the question." Divide 37 lb. 2 oz. 15 dr. by 1 lb. 9 oz. 11 dr.," what are we in reality required to do? (\$7).
  83. When the divisor contains decimals, how do we proceed? (\$8) Upon

what principle do we do this? (89)

34. How do we divide by 1, followed by any number of 00? (89)

35. How do we contract the work when dividing by 25? How by 15, 35, 45,

or 55? (90, 91)

36. How do we divide by 125? How by 75, 175, 225, or 275? (92, 93)

37. How do we abbreviate the work when there are many decimals in the dividend and but few are required in the quotient? (94)

#### EXERCISE 23.

### MISCELLANEOUS EXERCISE. (On preceding rules.)

1. Multiply 789643 by 999998.

2. Read the following numbers: 67813420.021030046. 72000000.000000072, 1001000100.0010000010000001.

3. Express 709, 4376, 9999, 86004, and 3947596 in Roman numerals.

4. Multiply 749 lb. 10 oz. avoirdupois by 72.

- 5. What is the price of 17 pairs of gloves at 4s. 72d per pair? 6. The planet Neptune is 2850 millions of miles from the sun :
- how long would it take a locomotive to travel from the sun to Neptune, at the rate of 30 miles an hour?

7. Reduce £729 17s. 61d. to dollars and cents.

8. From \$10000 subtract \$9876.23.

9. Write down five hundred and twenty billions, six millions, two thousand and forty-three, and five thousand and sixteen trillionths.

10. Reduce 7964327 inches to acres, roods, &c.

11. Add together the following quantities: \$729.43, \$16.70, \$976.81, \$9987.17, \$429.00, \$129.19.

12. Multiply 6 weeks 4 days 3 hours 17 minutes by 429.

13. Take the number 741, and, by removing the decimal point: (1) multiply it by 1000000; (2) divide it by 100000; (3) make it millions; (4) make it billionths; (5) make it trillionths; (6) make it hundredths of thousandths; (7) make it tenths.

14. Multiply 78.96 by .00042.

15. How many hogsheads of sugar, each containing 13 cwt. 2 qrs. 14 lbs., may be put on board a ship of 324 tons burden?

16. A farmer's yearly income was 9237 dollars. He paid for repairing his house 136 dollars, for hired help on his farm 4 times as much lacking 95 dollars, and for other expenses 1902 dollars; how much does he save yearly?

17. How many suits of clothes can be made from a piece of cloth containing 39 yds. 2 qrs. 3 nls.; each suit requiring 3 yds.

1 qr. 2 nls.?

18. There is a farm consisting of 732 acres; 25 acres of which is planted with corn and potatoes; 197 acres sown with rye; 156 with oats; 97 with wheat; 199 is pastured; and the remainder is meadow. How many acres of meadow?

19. Bought 96 acres 3 roods 17 perches of land, for which I

pay \$7764; what did I pay for it per perch?

20. A lady, having 312 dollars, paid for a bonnet 20 dollars, for a shawl 75 dollars, for a silk dress 97 dollars, and for some delaines 83 dollars; how much had she remaining?

21. A silversmith received 36 lb. 8 oz. 14 dwt. 16 grs. of silver to make 12 tankards; what would the weight of each tankard be?

22. I bought four fields; in the first there were 8 acres 3 rds. 12 perches; in the second, 7 acres 2 roods; in the third, 9 acres and 13 perches; in the fourth, 5 acres 2 roods 36 perches. How much in all?

23. A merchant expended 294 dollars for broadcloth, consisting of three different kinds; the first at 5 dollars a yard; the second at 7 dollars; and the third at 9 dollars a yard. He had as many yards of one kind as of another—how many yards of

each kind did he buy?

24. A silversmith made three dozen spoons, weighing 5 lb. 9 oz. 8 dwt.; a tea-pot, weighing 3 lb. 2 oz. 16 dwt. 16 grs.; two pair of silver candlesticks, weighing 4 lb. 6 oz. 17 dwt.; a dozen silver forks, weighing 1 lb. 8 oz. 19 dwt. 22 grs.; what was the weight of all the articles?

25. Reduce £972 11s. 111d. to dollars and cents.

26. Reduce 179 lbs. 3 oz. 3 dr. 1 scr. 14 grs. to grains.

27. There is a house 56 feet long, and each of the two sides of the roof is 25 feet wide; how many shingles will it take to cover

it, if it require 6 shingles to cover a square foot?

28. A merchant bought 4 bales of cotton; the first contained 6 cwt. 2 qr. 11 lb.; the second, 5 cwt. 3 qr. 16 lb.; the third, 8 cwt. 0 qr. 7 lb.; the fourth, 3 cwt. 1 qr. 17 lb. He sold the whole at 15 cents a pound; what did it amount to?

29. A merchant has 29 bales of cotton cloth, each bale containing 57 yards; what is the value of the whole at 15 cents a

yard?

30. A man willed an estate of \$370129 to his two children and wife, as follows: to his son, \$139468; to his daughter, \$98579; and to his wife the remainder. How much did he will to his wife?

31. Divide £1694 16s. 011d. by £9 19s. 111d. 32. Reduce £19 19s. 111d. to dollars and cents.

33. A merchant having purchased 12 cwt. of sugar, sold at one time 3 cwt. 2 qrs. 11 lb., and at another time he sold 4 cwt. 1 qr. 15 lb.; what is the remainder worth, at 15 cents per pound?

34. Bought 4 chests of hyson tea; the weight of the first was 2 cwt. 0 qr. 17 lb.; the second 3 cwt. 2 qrs. 15 lb.; the third, 2 cwt. 1 qr. 20 lb.; the fourth, 5 cwt. 3 qr. 17 lb.; what is the value of the whole at 371 cents a pound?

35. Express 100200300709 in Roman numerals.

36. Divide 43-2 by 76-8437.

37. Divide 123.4 by .000000066. Collaboration : att Range

38. From \$2789.27 take 17 times \$63.29.

39. Add together \$278.43, \$417.16, \$11.27, \$2110.40, \$723.15,

and £29 6s. 112d. and divide the sum by 173.

40. In 1857 the total number of volumes in the Common School and other Public Libraries of Canada West was estimated at 491544 and the number of libraries at 2076. How many volumes were there upon an average to each library?

# steller . Mille Burney SECTION ., III.

PROPERTIES OF NUMBERS, PRIME NUMBERS, MEASURES,
GREATEST COMMON MEASURE, LEAST COMMON
MULTIPLE, SCALES OF NOTATION, AND APPLICATION OF THE FUNDAMENTAL RULES TO DIFFERENT
SCALES. DUODECIMALS.

1. A divisor, or measure of a number, is a number which will divide it exactly; that is, leaving no remainder.

2. A multiple of a number is a number of which the given number is a divisor.

3. An integer, or integral number, is a whole number.

4. Integers are either prime or composite, odd or even.

5. An Even Number is that of which 2 is a divisor.

6. An Odd Number is that of which 2 is not a divisor.

A Prime Number is one which has no integral divisor except unity and itself, thus 2, 3, 5, 7, 11, 13, 17, 19, 23,

29. &c., are primes.

8. A Composite Number is a number which is not prime; or is a number which has other *integral* divisors besides unity and itself, thus 4, 6, 9, 10, 12, 14, 15, 16, 21, &c., are composite numbers.

9. The Factors of a number are those numbers which,

when multiplied together, produce or make it.

or aliquot parts.

11. A Common Measure of two or more numbers, is a number which will divide each of them without a remainder; thus 7 is a common measure of 14, 35, and 63.

12. Two or more numbers are prime to one another when they have no common divisor except unity; thus, 9 and 14 are "prime to each other."

15,

aoel

l at

mes

Hence all prime numbers are prime to each other; but composite numbers may or may not be prime to one another.

13. Commensurable Numbers are those which have some common divisor.

Thus 55 and 35 are commensurable, the common divisor being 11.

14. Incommensurable Numbers are those which are prime to one another.

Thus 55 and 34 are incommensurable.

15. A Square Number is one which is composed of two equal factors.

Thus 25=5×5 is a square number: so also 64=8×8, &c.

16. A Cube Number is one which is composed of three equal factors.

Thus 348=7×7×7 is a cube number: so also 27=3×3×8, &c.

17. A Perfect Number is one which is exactly equal to the sum of all its divisors.

Thus, 6=1+2+3 is a perfect number; so also 28=1+2+4+7+14 is a perfect number.

All the numbers known to which this property really belongs, are the eight following: 6; 28; 496; 8128; 33550336; 8589669056; 137438691328; and 2305843008189952128.

Note.—All perfect numbers terminate with 6, or 28.

18. Amicable Numbers are such pairs of integers that each of them is exactly equal to the sum of all the divisors of the other.

Thus, 220 and 284 are amicable; for, 220=1+2+4+71+142, which are all the divisors of 284, and 284=1+2+5+11+4+10+22+20+44+55+110, which are all divisors of 220.

Other amicable numbers are 17296 and 18416; also 9363583 and 9437056.

19. By the term properties of numbers, is meant those qualities or elements which are inseparable from them. Some of the most important properties of numbers are the following:

I. The sum of two or more even numbers is an even

number.

II. The difference of two even numbers is an even number.

III. The sum or difference of two odd numbers is an even number.

IV. The sum of three, five, seven, &c., odd numbers, is an odd number.

ICA-ENT aber

RES.

MON

der. the

or. visor 23,

en.

me; ides &c.,

ich, les,

is a ain-

her

V. The sum of two, four, six, eight, &c., odd numbers, is an even number.

VI. The sum or difference of an even and an odd num-

ber, is an odd number.

VII. The product of two even numbers, or of an even and an odd number, is an even number.

VIII. If an even number be divisible by an odd num-

ber, the quotient will be an even number.

IX. The product of any number of factors will be even if one of the factors be even.

X. An odd number is not divisible by any even number. XI. The product of any number of factors is odd if they are all odd.

XII. If an odd number divide an even number, it will

also divide half of it.

XIII. Any number that measures two others must like wise measure their sum, their difference, and their product,

Thus, if 6 goes into 24 four times, and into 18 three times, it will go into 34+18 or 42, three plus four, or seven times.

Also, if 6 goes into 24 four times, and into 42 seven times, it will go into 42—24 or 18, seven minus four, or three times.

Lastly, if 6 goes into 24 four times, and into 12 twice, it will evidently go into 12 times 24, twelve times 4 times, or 48 times.

XIV. If one number measure another, it must likewise measure any multiple of that other.

Thus, if 7 measures 21, it must evidently measure 6 times 21, or 11 times 21, or 17 times 21, ac.

XV. Any number, expressed by the decimal notation, divided by 9, will leave the same remainder as the sum of its digits divided by 9. (See Art. 55, Sec. II.)

This property of the number 9 affords an ingenious method of proving each of the fundamental rules. The same property belongs to the anuaber 3; for 3 is a measure of 9, and will therefore be contained an exact number of times in any number of 9s. But it belongs to no other digit. The preceding is not a necessary but an incidental property of the number 9. It arises from the law of increase in the decimal notation. If the radix of the system were 8, it would belong to 7; if the radix were 12, it would belong to 11; and, universally, it belongs to the number that is one less than the radix of the system of notation.

XVI. If the number 9 be multiplied by any single digit, the sum of the figures composing the product will make 9.

Thus,  $9 \times 4 = 36$ , and 3 + 6 = 9; so also  $8 \times 9 = 72$  and 7 + 2 = 9.

XVII. If we take any two numbers whatever; then one of them, or their sum, or their difference, is divisible by 3.

Th sum.

reme place alter Cessa

Tak sum o 4+5+ vided

not m the re X

divisi X X

XX the w XX

by 8,  $\mathbf{X}\mathbf{X}$ visible

XX sible b XX

even p odd pl Thus to

The n

So also in the oc places. XX

prime For sin

factors t these fa other fac are prim abers,

num-

a even

l num-

e even

umber. if they

it will

ust like product. ill go into

ill go into idently go

likewise

r 11 times

notation. sum of

of proving he number ct number

f the num-on. If the were 12, it that is one

gle digit, make 9.

then one ble by 3. Thus, take 11 and 17; though neither the numbers themselves, nor their sum, is divisible by 3, yet their difference is, for it is 6.

XVIII. Any number divided by 11, will leave the same remainder as the sum of its alternate digits in the even places, reckoning from the right, taken from the sum of its alternate digits in the odd places, increased by 11, if ne-CESSATY.

Take any number as \$8405605, and mark the alternate figures. Now the sum of those marked, viz: 8+0-16+3=17. The sum of the others, viz: 8+4-15+0=12. And 17-12=5, the remainder sought. That is; \$8405603 divided by 11, will leave 5 remainder.

Again, take \$847302, the sum of the marked figures is 14; the sum of those not marked is 31. Now 21 taken from 25, (i.e. 14 increased by 11) leaves 4, the remainder sought—remainder obtained by dividing 5847302 by 11.

XIX. Any number ending in 0, or an even number, is divisible by 2.

XX. Any number ending in 5 or 0 is divisible by 5. XXI. Any number ending in 0 is divisible by 10.

XXII. When two right-hand figures are divisible by 4, the whole is divisible by 4.

XXIII. When the three right-hand figures are divisible by 8, the whole number is divisible by 8.

XXIV. When the sum of the digits of a number is divisible by 9, the number itself is divisible by 9.

XXV. When the sum of the digits of a number is divi-

sible by 3, the number itself is divisible by 3.

XXVI. When the sum of the digits, standing in the even places, is equal to the sum of the digits standing in the odd places, the number is divisible by 11.

Thus to illustrate the last five properties.

The number 7416 is divisible by 4, because 16, the last two digits, is divisible by 4.

divisible by 4.

—is divisible by 8, because 416, its last three digits, is divisible by 8,

—is divisible by 9, because the sum of its digits, 7+4+1

+6=18, is divisible by 9,

—is divisible by 3, because the sum of its digits, 7+4+1

+6=18, is divisible by 3.

So also the number 4567321 is divisible by 11, since the sum of the digits in the odd places, 1+3+6+4=14=2+7+5, the sum of the digits in the even

XXVII. Every composite number may be resolved into prime factors.

For since a composite number is produced by multiplying two or more factors together, it may evidently be resolved into those factors; and if these factors themselves are composite, they also may be resolved into other factors, and thus the analysis may be continued until all the factors are prime numbers.

XXVIII. The least divisor of any number is a prime number.

For every whole number is either prime or composite (Art. 4); but a composite number can be resolved into factors (XXVII): consequently, the least divisor of any number must be a prime number.

XXIX. Every prime number, except 2, if increased or diminished by 1 is divisible by 4. (See table of prime numbers on next page).

XXX. Every prime number except 2, is odd; and

therefore terminates in an odd digit.

NOTE.—It must not be inferred from this that all odd numbers are prime.

XXXI. All prime numbers, except 2 and 5, must terminate with 1, 3, 7, or 9. Every number that ends in any other digit than 1, 3, 7, or 9, is a composite number.

For all prime numbers, except 2, must end in an odd digit (XXX), and all numbers ending in 5 are divisible by 5.

XXXII. Every prime number, except 2 and 3, if increased or diminished by 1, is divisible by 6.

20. To find the prime numbers between any given

limits-

Write down all the odd numbers, 1, 3, 5, 7, 9, &c. Over every third from 3 write 3; over every fifth from 5 write 5; over every seventh from 7 write 7; over every eleventh from 11 write 11; and

Then all the numbers which are thus marked are composite; and

the others, together with 2, are prime.

Also the figures thus placed over, are factors of the numbers over which they stand.

#### EXAMPLE.

Find all the prime numbers less than 100.

17.8	1 1	Part IV		8			3.2	
1	3	5 .	7	9	11	13	/ 15	17
1 11	- 8.7	5 ,	5	8			3.11	5.7
19	- 21	23	25	3 27	29	31	33	35
	8.13			- 3.2	Dr. aug a	7	8.17	
37	39	41	43	45	47	49	51	. 53
5'11	3.19		9 41	8.7	5.13		3'23	i t
55	57	59	61	63	- 65	67	69	71
٦	3.2	7.11		8		5'17	3.29	3415 4
73	75	77-	79	81	83	85	87	89
7'18	3.81	5'19	4 A 16 M	811	1 1 1	1 44 1,	4 10 11	
91	93	95	97	99			2 11	

ART. 2

Hen prime 43, 47 numb

This primes Eratos scribed numbe what r

When first no number must en table, co divide o of the ta is a prime

rt. 4); but a jequently, the

e of prime

odd; and

oers are prime. 5, must terends in any mber.

and 3, if in-

any given

5; over every write 11; and

omposite; and

e numbers over

3.5 1.5 3.11 3.11 5.7 3.3 3.5 3.17 5.1 5.3 3.23 6.9 7.1 3.29 8.7 8.9 Hence, rejecting all the numbers which have superiors, the primes less than 100 are 1, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 37, 41, 43, 47, 53, 59, 61, 67, 71, 73, 79, 83, 89, 97, together with the number 2.

This process may be extended indefinitely, and is the method by which primes are found even by modern computators. It was invented by Eratosthenes, a learned librarian at Alexandria (Born B. C. 275). He inscribed the series of odd numbers upon parchment, then cutting out such numbers as he found to be composite, his parchment with its holes somewhat resembled a sieve: hence, this method is called 'Eratosthenes' Sieve.'

### TABLE OF PRIME NUMBERS FROM 1 TO 3407.

1	173	409	659	941	1223	1511	1811	2129	2423	2741	8079
2	179	419	661	947	1229	1523	1823	2131	2437	2740	8088
3	181	421	678	953	1281	1531	1881	2137	2441	2753	3089
5	. 191	431	677	967	1237	1543	1847	2141	2447	2767	3109
7	193	433	683	971	1249	1549	1861	2143	2459	2777	3119
11	197	439	691	977	1259	1553	1867	2153	2467	2789	3121
13	199	443	701	983	1277	1559	1871	2161	2473	2791	3137
17	211	449	709	991	1279	1567	1873	2179	2477	2797	3163
19	223	457	719	997	1283	1571	1877	2203	2503	2801	3167
23	227	461	727	1009	1289	1579	1879	2207	2521	2803	3169
29	229	463	783	1018	1291	1583	1889	2213	2531	2819	3181
81	233	467	739	1019	1297	1597	1901	2221	2539	2833	3187
37	239	479	743	1021	1301	1601	1907	2237	2543	2837	3191
41	241	487	751	1031	1303	1607	1913	2239	2549	2843	3203
43	251	491	757	1033	1307	1609	1981	2243	2551	2851	3209
17	257	409	761	1039	1319	1618	1933	2251	2557	2857	3217
53	263	503	769	1049	1321	1619	1949	2267	2579	2861	3221
59	269	509	778	1051	1327	1621	1951	2269	2591	2879	3229
81	271	521	787	1061	1361	1627	1973	2273	2593	2887	3251
67	277	523	797	1063	1867	1637	1979	2281	2609	2897	8253
71	281	541	800	1069	1373	1657	1987	2287	2617	2903	8257
78	283	547	811	1087	1381	1668	1993	2293	2621	2909	3259
79	293	557	821	1091	1399	1667	1997	2297	2633	2917	3271
83	307	563	823	1093	1409	1669	1999	2309	2647	2927	3299
89	311	569	827	1097	1423	1693	2003	2311	2657	2939	3301
97	313	571	829	1103	1427	1697	2011	2333	2659	2953	3307
01	317	577	839	1109	1429	1699	2017	2339	2663	2957	3313
03	331	587	853	1117	1433	1709	2027	2841	2671	2963	3319
07	837	593	857	1123	1439	1721	2029	2347	2677	2969	3325
09	847	599	859	1129	1447	1723	2039	2351	2683	2971	3329
18	349	601	863	1151	1451	1733	2053	2357	2687	2999	333
27	353	607	877	1158	1453	1741	2063	2371	2689	8001	834
181	359	613	881	1163	1459	1747	2069	2377	2693	3011	834
137	367	617	883	1171	1471	1753	2081	2381	2699	3019	335
189	373	619	887	1181	1481	1759	2083	2383	2707	3023	336
149	379	631	907	1187	1483	1777	2087	2389	2711	3037	337
151	383	641	911	1193	1487	1783	2089	2393	2718	8041	337
157	389	643	919	1201	1489	1787	2099	2399	2719	3040	338
163	397	647	929	1213	1493	1789	2111	2411	2729	3061	339
167	401	653	937	1217	1499	1801	2113	2417	2731	3067	340

When it is required to determine whether a given number is a prime, we first notice the terminating figure; if it is different from 1, 3, 7, or 9, the number is composite; but if it terminate with one of the above digits, we must endeavour to divide it with some one of the primes, as found in the table, commencing with 3. There is no necessity for trying 2, for 2 will divide only the even numbers. If we proceed to try all the successive primes of the table until we reach a prime which is not less than the square-root

of the number, without finding a divisor, we may conclude with certainty that the number is a prime.

The reason why we need not try any primes greater than the square-root of the number, is drawn from the following consideration: If a composite number is resolved into two factors, one of which is less than the square-root of the number, the other must be greater than the square-root.

The square of the last prime given in our table is 11607649; hence, this table is sufficiently extended to enable us to determine whether any number not exceeding 11607649 is a prime. It is obvious that numbers may be proposed which would require by this method very great labor to determine whether they are primes, still this is the only sure and general method as yet discovered.

21. To resolve a Composite Number into its Prime Factors.

Divide the given number by the smallest number which will divide it without a remainder; then divide the quotient in the same way, and thus continue the operation till a quotient is obtained which can be divided by no number greater than 1. The several divisors with the last quotient, will be the prime factors required. (19—

XXVII.)

REASON.—Every division of a number, it is plain, resolves it into two factors, viz. the divisor and the quotient. But according to the rule, the divisors, in every case, are the smallest numbers that will divide the given number or the successive quotients without a remainder, consequently they are all prime numbers. (19-XXVIII.) And since the division is continued till a quotient is obtained, which cannot be divided by any number but unity or itself, it follows that the last quotient must also be a prime number; for, a prime number is one which cannot be exactly divided by any whole number except unity and itself. (Art. 7.)

Note.—Since the least divisor of every hunder is a prime number, it is evident that a composite number may be resolved into its prime factors by dividing it continually by any prime number that will divide the given number and the successive quotients without a remainder. Hence,

A composite number can be divided by any of its prime factors without a remainder, and by the product of any two or more of them, but by no other number.

Thus, the prime factors of 43 are 2.3, and 7. Now 42 can be divided by 2.3, and 7; also by 2×3, 2×7, 3×7, and 2×3×7; but it can be divided by no other number.

EXAMPLE 1.—Resolve 210 into its prime factors.

OPBRATION. 2)210

8)105 6)85

We first divide the given number by 2, which is the least number that will divide it without a remainder, and which is also a prime number. We next divide by 3, then by 5. The several divisors and the last quotient are the prime factors required.

Ans. 2, 3, 5, and 7.

PROOF.—2×3×5×7=210. EXAMPLE 2.—Resolve 728 into its prime factors.

OPERATION. 2)728

1 72 1 . 1 2) 364 & 70 1

or of the state of 

#### with certainty

he square-root If a consposite an the square-e-root. 807849; hence, e whether any numbers may eat labor to de-ure and general

ME FACTORS.

ch will divide he same way, tained which veral divisors equired. (19-

evolves it into ling to the rule, will divide the nainder, conse-e the division is ed by any numalso be a prime

me number, it is orime factors by livide the given

Hence, factors without, but by no other n be divided by be divided by no

y 2, which is the ut a remainder, e next divide by

×2×2×7×18, on the prime factor

#### EXERCISE 24.

3. Resolve 11368 into its prime factors.	Anis. \$ 23 × 72 × 29.
4. What are the prime factors of 2934?	Ans. 2×3°×163.
5. What are the prime factors of 1011?	Ans. 8×337
6. What are the prime factors of 1000?	Ans. 23×53.
7. What are the prime factors of 1024?	Ans. 210.
8. What are the prime factors of 32320?	Ans. 26×5×101.
9. What are the prime factors of 707?	Ans. 7×101.
10. What are the prime factors of 1118?	Ans. 2 × 18 × 43.
and they they they had the to a to	

# DIVISORS.

22. From Art. 21, Note, for finding all the divisors of any number, we deduce the following-

#### RULE.

Resolve the number into its prime factors; form as many series of terms as there are prime factors, by making 1 the first term of each series, the first power of one of the prime factors for the second term, the second power of this factor for the third term, and so on, until we reach the highest that occurred in the decomposition. Then multiply these series together, and the partial products thus obtained will be the divisors sought.

### Example 1.—What are the divisors of 48?

Here we find  $48=34\times3$ . Therefore our series of terms will be  $1\cdot 2\cdot 4\cdot 8\cdot 16$  and  $1\cdot 3$ ; multiplying these together.  $1\cdot 2\cdot 4\cdot 8\cdot 16$ 

1 · 2 · 4 · 8 · 16 · 3 · 6 · 12 · 24 · 48

Therefore the divisors of 40 are 1, 2, 3, 4, 6, 8, 12, 10, 24, and 48.

We begin each series with 1, because, were we not to do so, the different powers of the prime factors would not themselves appear among the partial products.

EXAMPLE 2.—What are the divisors of 360.

The prime factors of 360 are 23×32×5 and therfore the series are 1.2. 4:8:1.3.9 and 1.5.

#### OPERATION.

1 -2 -4 -8 1.3.9

-2 - 4 - 8 - 3 - 6 - 12 - 24 - 9 - 18 - 36 - 72 = products of 1st and 2nd series

1 - 2 - 4 - 8 - 3 - 6 - 12 - 24 - 9 - 18 - 36 - 72 - 5 - 10 - 20 - 40 - 15 - 30 - 60 -

120 · 45 · 90 · 180 · 360.

Therefore the divisors of 360 are 1, 2, 3, 4, 5, 6, 8, 9, 10, 12, 15, 18, 20, 24, 30, 36, 40, 45, 60, 73, 90, 130, 180, 360.

The small fleures written to the right of the factors and above the line, are called exponents, and show how often the digit is taken as factor.

mai

will

num

K 424

#### EXERCISE 25.

1. What are the divisors of 100?

Ans. 1, 2, 4, 5, 10, 20, 25, 50, 100.

2. What are the divisors of 810?

Ans. (1, 2, 3, 5, 6, 9, 10, 15, 18, 27, 30, 45, 54, 81, 90, 135, 162, 270, 405, 810.

3. What are the divisors of 920?

Ans. 1, 2, 4, 5, 8, 10, 20, 23, 40, 46, 92, 115, 184, 230, 460, 920.

4. What are the divisors of 25000?

§ 1, 2, 4, 5, 8, 10, 20, 25, 40, 50, 100, 125, 200, 250, 500, 625, 1000, 1250, 2500, 3125, 5000, 6250, 12500, 25000.

#### NUMBER OF DIVISORS.

23. Since the series of terms which we multiplied together, by the last rule, to obtain the divisors of any number commenced with 1, it follows that the number of terms in each series will be one more than the units in the exponent of the factors used.

Hence, to find the number of divisors of any number, without actually setting them down, we have the following-

Resolve the number into its prime fuctors and express them as in examples 3, 4, and 6, in Art. 21. Increase each exponent by unity and multiply the resulting numbers together. The product will be the number of divisors.

# EXAMPLE.—How many divisors has 4320?

 $4320=2^{5}\times8^{8}\times5$ . Here the exponents are 5,8, and 1: each of which being increased by one, we obtain 6, 4, and 2, the continued product of which is  $6\times4\times2=48$ —the number of divisors sought.

#### EXERCISE 26.

1.	How many	livisors h	as 88200 ?	master on the	Ans. 108.
2:	How many	divisors b	as 3500 ?	3 10 4	Ans. 24.
3.	How many	divisors b	as 6336 ?		Ans. 42.
4.	How many d	livisors h	as 824 ?		Ans. 8.
5.	How many	livisors b	as 49000?	, .	Ans. 48.
6.	How many	divisors b	as 81000 ?		Ans. 80.
7.	How many	divisors l	has 75600?		Ans. 120.
8.	How many	divisors l	has 25600 ?	A km ! HI	Ans. 33.

# AGO ..... GREATEST COMMON MEASURE.

24. The greatest common measure, or greatest common divisor of two or more numbers, is the greatest number that will divide each of them without a remainder,

([ografi

mon mber

25. To find a common divisor or common measure of two or more numbers:-

Resolve the given numbers into their prime factors, then if any factor be common to all, it would be a common measure.

If the given numbers have not a common factor they cannot have a common measure greater than unity, and consequently are either prime numbers or are prime to each other. (Arts. 7 and 12.)

EXAMPLE.—Find a common divisor of 14, 35, and 63.

14=2×7: 35=5×7, and 63=3×3×7. The factor 7 is common to all the given numbers, and is therefore a common measure of them.

#### Evenoren 27

						Edda 2	24.5 5	4 - 52	
. 171 3		32-1	-201	10 04		3 0			
1. Find a	common	GIVISOR	01 31,	10, 41	and 30.	4 14. T	1	Ans.	3.
ALLE TO THE PARTY OF THE PARTY			10100			1			

26. To find the greatest common measure of two quantities :-

#### BULE.

Divide the larger by the smaller; then the divisor by the remainder; next the preceding divisor by the new remainder:continue this process until nothing remains, and the last divisor will be the greatest common measure. If this be unity, the given numbers are prime to each other.

Example.—Find the greatest common measure of 3252 and 4248

3252)4248(1

996)\$252(8

o. Wat the par took come 8802

204)264(1

204

60)204(3

24)60(2

12)24(2

996, the first remainder, becomes the second divisor; 264, the second remainder, becomes the third divisor, &c. 12, the last divisor, is the required greatest common measure.

From Proces.—In order to establish the truth of this rule, it is necessary to remember (19-XIII. and XIV.) that if one number measure another it will likewise measure any integral multiple of that other; and if one number measure two others, it will also measure their sum or their difference.

First, then, 13 is a common measures 23, it also measures 24, a multiple of the process; because 12 measures 12, it also measures 24, a multiple of 14, because 13 measures 13 and also 45, it measures 60, a multiple of 24, because 13 measures 14, a multiple of 15, a measures 15, and also 24, it measures 150, and also 24, it measures 150, and also 24, it measures 150, and likewise 60, it measures their sum, which is 204, because 13 measures 204, it measures 792, a multiple of 264; and because 13 measures 793, and also 204, it measures their sum, which is 224. Therefore, measures 205, it measures 2060, a multiple of 265; and because 13 measures 2053, and also 204, it measures their sum, which is 2248. 12, therefore, measures each of the given numbers, and is a common measure; next it is their greatest common measure.

For, if not, let some other as 18, be greater. Then, (beginning now at the top of the process) because 15 measures 2353, and also 234, it measures their difference, which is 234; because 13 measures 204, it measures their difference, which is 234; because 15 measures 204, it measures 208, a multiple of 204; and because 18 measures 252, and also 204, it measures their difference, which is 234; because 18 measures 264, it also measures 208, a multiple of 264; and because 18 measures 253, and also 204, it measures their difference, which is 234; because 18 measures 264, it also measures 208, a multiple of 264; and because 18 measures 255, and also 204, it measures their difference, which is 236; because 18 measures 264, it measures 264, it measures 265, a multiple of 264; and because 18 measures 265, and also 204, it measures 265, a multiple of 264; and because 18 measures 264, it measures 265, a multiple of 264;

manner it may be shown that no number greater than 13 is a common measure. Therefore 13 is the greatest common measure.

As the rule might be proved for any other example equally well, it is

true in all cases.

# Exercise 28.

- 1. What is the greatest common measure of 296 and 407?
- Ans. 37. 2. What is the greatest common measure of 506 and 308?
- Ans. 22. 3. What is the greatest common measure of 74 and 84? Ans. 2.
- 4. What is the greatest common measure of 1825 and 2555?
- Ans. 365. 5. What is the greatest common measure of 556 and 672?
- 27. To find the greatest common measure of more than two numbers :---

#### RULE.

Find the greatest common measure of two of them; then, of this common measure and a third; next of this last common measure and a fourth, &c. The last common measure found will be the greatest common measure of all the given numbers.

EXAMPLE 1.—Find the greatest common measure of 679, 5901, and 6734.

7 An The 0 and Th

AR

mon of th Th comm

1. W 2. W

3. W 4. WI

28 or gre be the mon : that a

Her more r

Resol product greatest

Ans. 4.

EXAM and 198

Hence, factors,

her it will e number

rence. ning at the a multiple because 12 because 12 ree 180, and es 104, and it measures it measures 986, a mul-sures their sures their n numbers.

sasure. ing now at it measures sures 1988, a so me it measures also 204, it it measures, it measures sasures their reater num-

in a similar s common ly well, it is

4077 Ans. 37.

308 ? Ans. 22. 14? Ans. 2. 1 2555?

Ans. 365. 672 ?

Ans. 4.

more than

hen, of this on measure will be the

679, 5901,

By the last rule we find that 7 is the greatest common measure of 679 and 5901; and by the same rule that it is the greatest common measure of 7 and 6784 (the remaining number), for 6784—7=962, with no remainder. Therefore 7 is the required number.

EXAMPLE 2.—Find the greatest common measure of 986, 736, and 142.

The greatest common measure of 936 and 736 is 8, and the greatest com-

The greatest common measure of use and 750 is 5, and the greatest common measure of 8 and 142 is 3; therefore 8 is the greatest common measure of the given numbers.

This rule may be shown to be correct in the same way as the last; except that in proving the number found to be a common measure, we are to begin at the end of all the processes, and go through all of them in succession; and in proving that it is the greatest common measure, we are to begin at the commencement of the first process, or that used to find the common measure of the two first numbers, and proceed successively through

#### EXERCISE 29.

- 1. What is the greatest common measure of 110, 140, and 680? Ans. 10.
- 2. What is the greatest common measure of 1326, 3094, and Ans. 442.
- 3. What is the greatest common measure of 468, 922, and 375? Ans. They have none.
- 4. What is the greatest common measure of 204, 1190, 1445, and 2006? Ans. 17.

### SECOND METHOD.

28. It is manifest that the greatest common measure or greatest common divisor of two or more numbers, must be their greatest common factor, and that this greatest common factor must be the product of all the prime factors that are common to all the given numbers.

Hence to find the greatest common measure of two or

more numbers, we have the following:—

Resolve each of the given numbers into its prime factors; and the product of those factors, which are common to all, will be the greatest common measure.

Example 1.—What is the greatest common measure of 1365 and 1995?

3)1365		i	3)1995
5)455	10		5)665
7)91			7)183
3			

Hence, 3, 5, 7, and 18 are the prime

Hence, 3, 5, 7, and 19 are the prime factors.

And the factors that are common to both are 3, 5, 7. Hence 3×5×7=105 =greatest common measure.

EXAMPLE 2.—What is the greatest common measure of 108, 126, and 162?

 $108=2^2\times3^3$ ,  $126=2\times3^2\times7$ , and  $162=2\times3^4$ . Hence, the factors that are common are 2 and  $3^2$ , and the greatest common measure=2×32==18.

EXERCISE 30.

1. Work by this method all the preceding examples.

2. What is the greatest common measure of 56, 84, 140, 168? Ans. 28.

3. What is the greatest common measure of 241920, 380160, 69120, 103680? Ans. 34560.

4. What is the greatest common measure of 10800, 28040, and 2160 ? Ans. 40.

### LEAST COMMON MULTIPLE.

29. One number is a common multiple of two or more others when it can be divided by each of them without a remainder.

30. One number is the least common multiple (l. c. m.) of two or more others when it is the least number that can

be divided by each of them without a remainder.

31. It is evident that a dividend will contain a divisor an exact number of times, when it contains, as factors, every factor of that divisor; and hence, the question of finding the *least* common multiple of several numbers is reduced to finding a number which shall contain all the prime factors of each number and none others. If the numbers have no common prime factor, their product will be their least common multiple.

Suppose we wish to see what is the least common multiple of 9, 12, 16, 20, and 35. Resolving these into their prime factors, we obtain  $9=3^2$ ,  $12=2^2\times3$ ,  $16=2^4$ ,  $20=2^2\times5$ , and  $35=7\times5$ . Now it is plain that  $2^4$  must enter into the least common multiple as a factor, and, since  $2^4$  is a multiple of  $2^2$ , we do not consider  $2^2$  also a factor of the least common multiple. So sleep  $2^2$ not consider 22 also a factor of the least common multiple. So also 32 must be a factor of the least common multiple; and since it contains 3, we do not again multiply by 3. Lastly, 5 and 7 must enter into the least common multiple.

The factors of the least common multiple are then 24, 32, 5 and 7; and these, multiplied together, give  $2^4 \times 3^2 \times 5 \times 7 = 5040 = \text{least common multiple}$ .

Hence, to find the least common multiple of two or more numbers, we have the following:-

Resolve the numbers into their prime factors (Art. 21), select all the different factors which occur, observing when the same factor as different powers, to take the highest power. The continued proct of the factors thus selected will be the least common multiple.

1. W

Here 25, mu

W

3. WH

4. WH

5. Wh

6. Wb and

32. two or

Writ Divide them w number Divi

the open any nui The .

line wil EXAM and 10

The le multiple the num will be th line; the (SECT. III.

3×5×7=105

ure of 108,

reatest com-

, 140, 168?
 Ans. 28.
20, 380160,
 Ans. 34560.
28040, and

wo or more without a

Ans. 40.

le (l. c. m.) er that can

in a divisor as factors, question of numbers is ain all the s. If the roduct will

of 9, 12, 16, 20, 3<sup>2</sup>, 12=2<sup>2</sup> × 3, enter into the of 2<sup>2</sup>, we do s. So also 3<sup>2</sup> contains 3, we the least com-

5 and 7; and mon multiple. of two or

1), select all same factor ntinued proon multiple.

#### EXERCISE 31.

1. What is the least common multiple of 8, 9, 10, 12, 25, 32, 75, and 80?

Here  $8 = 2^3$ ,  $9 = 3^2$ ,  $10 = 2 \times 5$ ,  $12 = 2^2 \times 3$ ,  $25 = 5^2$ ,  $32 = 2^5$ ,  $75 = 5^2 \times 3$ ,  $80 = 2^4 \times 5$ . Therefore the least common multiple  $= 2^5 \times 3^2 \times 5^2 = 70200$ .

2. What is the least common staltiple of 6, 7, 42, 9, 10, and 630 ?

Ans.  $2 \times 3^2 \times 5 \times 7 = 630$ .

3. What is the least common multiple of the nine digits?

Ans.  $2^3 \times 3^2 \times 5 \times 7 = 2520$ . 4. What is the least common multiple of 6, 9, 12, 15, 18, 21, and 30?

Ans. 1260.

5. What is the least common multiple of 670, 100, 335, and 25?

6. What is the least common multiple of 8, 10, 18, 27, 36, 44, and 396?

Ans. 11880.

### SECOND METHOD.

32. We may also find the least common multiple of two or more numbers by the following:—

#### RITE.

Write the given numbers in a line, with two points between them. Divide by the LEAST number which will divide any two or more of them without a remainder, and set the quotients and the undivided numbers in a line below.

Divide this line and set down the results as before; thus continue the operation till there are no two numbers which can be divided by any number greater than 1.

The continued product of the divisors and the numbers in the last line will be the least common multiple sought.

EXAMPLE 1.—What is the least common multiple of 16, 48, and 108?

2)16 .. 48 ..108 2) 8 .. 24 .. 54 2)4 .. 12 .. 27 2)2 .. 6 .. 27 3)1 .. 3 .. 27 1 .. 1 .. 9

Ans.  $2\times2\times2\times2\times3\times9=432=$  least common multiple.

The least common multiple of 1, 1, and 9 is 9, and the least common multiple of 1, 1, and  $9\times3$ , will be the least common multiple of 1, 3, and 27, the numbers of the fifth line; the least common multiple of 1,3 and 27,  $\times$  2, will be the least common multiple of 2, 6, and 27, the numbers of the fourth line; the least common, multiple of 2, 6, and 27,  $\times$  2, will be the least com-

mon multiple of 4, 12, and 27, the numbers in the third line; the least common multiple of 4, 12, and 27, $\times$ 2, will be the least common multiple of 8, 24, and 54, the numbers in the second line; and the least common multiple of 8, 24, and 54, $\times$ 2, will be the least common multiple of 16, 48, and 140, the given numbers.

The reason of the preceding rule depends upon the principle; that the least common multiple of two or more numbers, is composed of all the prime factors of the given numbers, each taken the greatest number of times it is found in either of the given numbers.

Note.—In finding the least common multiple by this method, it is necessary to divide by the *smallest* number, which will divide two or more of them without a remainder, because the divisor may otherwise be a composite number (Art. 21), and have a factor *common* to it, and one of the quotients in the last line. Consequently the continued product of the divisors and these quotients or undivided numbers in the last line, would be too great for the least common multiple.

Thus in the third of the following operations the divisor 9 is a composite number, containing the factor 3, common to it and the 3 in the quotients consequently the product is three times too large. In the second operation the divisor 12 is a composite number, and contains the factor 6 common to it, and the 6 in the quotient; therefore the product is six times too large.

it, and the 6 in the quotient: therefore the product is six times too large.

The object of arranging the given numbers in a line, is that all of them may be resolved into their prime factors at the same time; and also to present at a glance the factors that compose the least common multiple required.

EXAMPLE 2.—What is the least common multiple of 12, 18, 36?

2)12 18 36	12)12 18 36	2)12 . 18 . 36
2)6 918	3)118 3	2)6 9 . 18
3)3 9 9	161	9)3 9 9
8)1 3 3	12×3×6=216	$ \begin{array}{c} 3 \dots 1 \dots 1 \\ 2 \times 2 \times 9 \times 3 = 108. \end{array} $
1 1 1 9×9×9×9 = 98 = 1	lom ? . Illa .	s who e is

#### EXERCISE 32.

- 1. Find the least common multiple of 12, 20, and 24. Ans. 120.
- 2. Find the least common multiple of 14, 21, 3, 2, and 63.
- Ans. 126.
  3. Find the least common multiple of 18, 12, 39, 216, and 234.
- 4. Find the least common multiple of 8, 18, 15, 20, and 70.

  Ans. 2520.
- 5. Find the least common multiple of 24, 16, 18, and 20.
- 6. Find the least common multiple of 60, 50, 144, 35, and 18.
- Ans. 25200.
  7. Find the least common multiple of 27, 54, 81, 14, and 63.

Ans. 1134.

mos

and comm An

unca exac of th and P

> M mul

60,

5, and duce Westo 11 the sa fact which it is the

the seco

39,

beca

S, a

#### the least ultiple of mon mul-3, and 16

SECT. III.

rinciple abers, is rs, each or of the

l, it is ner more of composite quotients risors and too great

a compoquotient: operation mmon to too large. ll of them id also to multiple :

, 18, 36? 36 18

= 108.

Ins. 120.

63. ns. 126. nd 234. s. 2808.

70. s. 2520.

ns. 720. and 18.

25200. id 63.

s. 1134.

#### THIRD METHOD.

33. The least common multiple of several numbers is most expeditiously found by the following:

Write the given numbers in a line. Take any one of them as divisor, and strike out of each of the given numbers all the factors that are common to it and the assumed number.

Arrange the uncancelled factors of the given numbers, and the uncancelled numbers in a line, take the least other number which exactly contains one or more of them, and strike out all the factors of the numbers in the second line which are common to any of them and the second assumed number.

Proceed thus until the assumed numbers cancel all the factors

of the given numbers.

Multiply all the assumed numbers together for the least common multiple of the given numbers.

Example 1.—What is the least common multiple of 16, 27, 45, 60, 88, 96, 100.

Assume 100 | 16 .. 27 .. 45 .. 60 .. 88 .. 96 .. 100 4 .. 27 .. 9 .. 8 .. 22 .. 24 Assume 24 9 · · · B · · · Assume 99  $100 \times 24 \times 99 = 237600 = 1. c. m.$ 

EXPLANATION.—4, a factor of 100, reduces 16 to 4, 88 to 22, and 96 to 24; 5, another factor of 100, reduces 45 to 9; and 20, another factor of 100, reduces 60 to 3. The numbers in the second line then are 4, 27, 9, 3, 22, and 26. We assume 24 of which a factor, 4, cancels 4; another factor 2 reduces 25 to 11; and another factor, 3, reduces 27 to 9 and 9 to 8. The numbers in the third line them are 9, 3, and 11. For this line we assumed 29, of which a factor, 3, cancels 3; another factor, 9, cancels 9; and a third, 11, cancels 11. Now since the least common multiple of a series of numbers is a number which still contains all the prime factors of each number, and none others, it is manifest that the least common multiple of the given numbers will be the same as the least common multiple of 100, and 4, 27, 9, 3, 22, and 24, because only those factors, which were common to the given numbers and 100 were struck out.

100 were struck out.

Similarly, the least common multiple of 100, 24, and 9, 3, and 11, will be the same as the least common multiple of 100, and the numbers in the second line, since only those factors which were common to 34 and the num-

bers of the second line are struck out.

Finally the least common multiple of 100, 24, and 99, is equal to the least common multiple of the given numbers.

EXAMPLE 2.—What is the least common multiple of 120, 12, 39, 65, 88, and 16?

Assume 120 | 120 - 40 - 32 - 65 - 88 - 16 Assume . 13 Assume  $120 \times 13 \times 22 = 34320 = 1.$  c. m.

EXPLANATION.—We first assume 120. Now this cancels 120 and 40. Also, 3, a factor of 130, reduces 30 to 13, and 5, another factor, reduces 65 to 13. Also 8, another factor, reduces 88 to 11 and 16 to 2. Next assume 13, this cancels 15 and 18. Next assume 22, of which 11, one factor, cancels the 11, and one than factor 8 cancels 15. and another factor 2, cancels 2.

till

EXAMPLE 3.—Find the least common multiple of 12, 16, 20, 24, 30, 48, 56, and 64.

#### Exercism 33.

What is the least common multiple of 300, 200, 150, 50, 60, 75, and 125?

2. What is the least common multiple of 20, 60, 15, 165, 210, 63, and 27?

3. What is the least common multiple of 12, 132, 144, 60, 96, and 1728?

Ans. 95040.

Work also by this method all the preceding questions in least common multiple.

# DIFFERENT SCALES OF NOTATION.

84. The radix or base of a scale of notation is its common ratio. Thus in our system the radix is 10; in the duodecimal system the radix is 12, &c.

35. If the expression 12345 represents a number in the common or decimal scale of notation, we read it twelve thousand three hundred and forty-five; but if it expresses a number in any other scale, we cannot so read it, because the names thousands, hundreds, &c., belong only to the decimal scale. In order to read it properly in any other scale we should have to invent names for the different orders. In place, however, of doing this, we simply read over the digits and indicate the scale. For example, if the expression 24678 be a number in the nonary scale, we read it thus—two, four, six, seven, eight in the nonary scale.

36. We may express the number 4578 (decimal scale) by writing the order of each digit beneath it, thus,

4578

10 10 10 ...

and then read it 8 units, 7 of the order of tens, 5 of the order of hundreds or tens squared, or second order of tens, 4 of the third order of tens, &c. Similarly if 4578 express a number in the *nonary* scale, we may write it.

4578

the soft it is

6, 20,

0, 60, 3000. 210, 1580. 0, 96,

5040.

com- : the

a the relve esses ause the ther

lers. the pres-

ale)

the ens.

688

and read it 8 units, 7 nines, 5 of the second order of nines, 4 of the third order of nines. &c.

- 37. The expression 10 always represents the radix of the scale. In the decimal scale 10 is equal ten; in the binary scale 10 is equal two; in the undenary scale 10 is equal eleven. &c.
- 38. It is obvious that, in any scale, the highest digit used must be one less than the radix. Thus, in the decimal scale, the highest digit is 9; in the ternary, 2; in the octenary, 7, &c. In writing numbers in the duodenary scale we use the letter t to represent ten, and e, eleven, and in the undenary scale t likewise represents ten.
- 39. Let it be required to reduce 337 from the decimal to the octenary scale.
- BATION.

  BENDLANATION.—If we divide 337 by 8, we distribute it into 8)337

  42 groups of 8 each, and have a remainder of 1 unit. If now we divide these groups of 8 by 8, we obtain 5 groups of a still higher order, each containing 8 of the former groups, with a remainder of 2 of these groups.

  5-2

  337, in the decimal scale, is therefore equal to 531 in the octonary scale; i. e. the successive remainders written in order constitute the equivalent expression in the required scale. OPERATION.

Hence, to reduce a number from one scale to another, we have the following: " " the transfer of the

#### RULE.

Divide the number continually by the radix of the proposed scale, till the quotient is less than the radix.

Write all the remainders, thus obtained, in regular order from left to right, beginning with the last, and placing 0s where there are no remainders. The result will be the required number.

EXAMPLE 1.—Reduce 7342 from the common to the quinary

PERATION. 5)7842	alon Print				40
5)1468—2				Brook in	(.:
5)2933	Therefore 784		21,8882	quina	ry
2-1	3577 + 64 (1) 5-577	1 1100 Te	Auto A		

EXAMPLE 2.—Express nine millions, three hundred and fortytwo thousand and twenty-seven, in the duodenary scale.

OF PANOT. The right want of working as the of the house it outs to be the real 12)64875-2 Therefore 9342027 denary = 3166323 duodenary. The selver and iven at the 12)450-6 we want be blow he then be come 12)87-6 est is . A at a specific to their outs of string, make of

### Exercise 34.

1. Change 592835 from the decimal to the duodenary scale. Ans. 24 10te.

2. Express the common number 3700 in the quinary scale.

Ans. 104300.

Ans. 7571. 3. Express 10000 in the undenary scale.

4. Express a million in the senary scale. Ans. 33233344. 5. Express 10000 in the octenary scale. Ans. 23420.

Transform 12345654321 into the duodenary scale.

Ans. 248664et69. 7. Express 10000 in the nonary scale. Ans. 14641.

8. Transform 300 from the common to the binary scale. Ans. 100101100.

EXAMPLE 1.—Transform 2313042 from the quinary to the octenary scale. Test the sails to be for the sail also sails

> EXPLANATION.—We divide here as before, bearing in mind, however, that the ratio is no longer ten, but five. We proceed thus.—8 in 2, no times; twice five (the radix) is ten and 3 make thirteen; 8 in 18, 1 and 5 over; 5 times 5 are 25, and 1 make 26; 8 in 26, 3 times and 2 over; twice 5 are 10, and 3 make 18, 8 in 18, once and 5 over, &c. OPBRATION. 8)2313042 8)181310-7 8)10100-8)311-2 8)20-1 Therefore 2313042 quinary = 121257 octenary,

e a suite se tree to the tree and a finished

Note.—The Roman Numeral written over the number indicates the radix of the scale.

A. L'111

18 MA F.

Ote.

300.

571.

344.

420.

2169. 641.

100.

the

efore, is no in 2, and 8

mes 5

once

sary.

indi-

31, 11

EXAMPLE 2.—Transform S78t18 from the undenary to the duodenary scale.

Observe the first two figures here are not thirty-seven, but 3×11+7=0. We say 13 into 40, 3 times and 4 over; next, 13 into 4×11+8 or 53, &c. OPERATION. 12)878#18 12)34456-8 (or it is a amitio) a 12)3132-4 12)294-0 878£13 undenary = 240948, duodenary, Ans. 12)26-9 12)2-4

EXAMPLE 3.—Transform #423# from the duodenary to the nonary scale. The second of the of sinits and least - then tries

Observe, here we say 9 into \$ tea, 1 and 1 over; 9 into 16, (1×12+4) 1 and 7 over; 9 into 86, (7×12+2) 9 and 5 over; 9 into 63, (5×12+3) 7; 9 into 6, 1 and 1 over. OPERATION. XII. 9) t423t And we proceed in the other lines in the same 9)11971-1 9)1649-4 t423t duodenary = 356341 nonary. 9)206-3 9)28-6 " be stone the will be so to 3-5

#### Examples 35.

- 1. Transform 37704 from the nonary to the octenary scale.
- 2. Transform 444 and 4321 from the quinary to the septenary 1. Ans. 285 and 1465.
- 3. Transform 1212201 from the quaternary to the nonary scale.

40. A number may be transformed from any scale to the decimal by the preceding rule, but the following is more convenient.

Multiply the left hand figure by the given radix, and to the product add the next figure.

Then multiply this sum by the radix and add the next figure. Continue this process until all the figures have been used. the last product will be the number in the decimal scale.

Norm.—Both this and the preceding rule are the same in principle as reducing denominate numbers from one denomination to another. wing the first more than the first of the first

EXAMPLE 1.—Reduce 76345 from the octenary scale to the decimal scale.

OPERATION. VIII. 62 of the fourth order. 409 of the third order. 3996 of the second order. 31978 units = required number in decimal scale.

EXAMPLE 2.—Transform ettete from the duodenary to the common or decimal scale.

XII 142 = number of fifth order. 1714 = number of fourth order. 20579 = number of third order. 246958 = number of second order. 2963507 = units = required number in decimal scale. .oloog grand tout of gat Exercise 36. A AVE TO A TO LEGIST TO

- 1. Change 20212331 from the quaternary into the decimal scale. 10 1903 Change L. 1 19 19 19 Ans. 35261.
- 2. Change 101202220 from the ternary into the decimal scale." Ans. 7854.
- 3. Transform 1522365 from the nonary into the decimal scale. Ans. 841568.
- 4. Transform 33233344 from the senary into the decimal scale. . 1000000.

Example 5 .- Transform 2784, octenary seale, into the undenary, septenary, and quinary scales, and prove the results by reducing all four numbers to the decimal scale.

VIII.	Dani T	VIII.	Lilly H .VIII.
11)9784	· .	7)2784	5)2784
11)810-4		7)826-2	5)454-0
11)14-4	11 10	7)86-4	5)74-0
1-1	7.8.00	4-2	5)14-0

herefore 2734	octenary=1144 undenary=48	enary=22000 quinary
8	11	1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1
23 8	12 80 7	18
187	196 196	4 To diller
8	ra ministration of the state of	25

1500 denary. 1500 denary. 1500 denary. 1500 denary. Since the results all agree when reduced to the denary scale, we conclude the work is correct.

6. Transform 132'13 nonary, into the ternary, duodenary, and octenary scales, and prove the results by reducing all four numbers to the denary scale.

7. Transform t2t290 duodenary, into the nonary, senary, quaternary, and binary scales, and prove the result by reducing

all five numbers to the decimal scale.

# FUNDAMENTAL RULES.

41. The fundamental rules of arithmetic are carried on in the different scales as with numbers in the ordinary or decimal scale; observing that, when we wish to find what to carry in addition, subtraction, multiplication, &c., we divide, not by ten, but by the radix of the particular scale used.

EXAMPLE 1.—Add together 34120, 3121, 13102, 31410, 12314, 112243 and 444444 in the senary scale.

OPERATION. Observe the sum of the first line is 14, which, divided by 6, VI. the radix of the scale, gives us 2 to set down and 2 to carry; 34120 the sum of the second line is 16, which, divided by the radix, 3121 6, gives us 4 to set down and 2 to carry, &c.

e.

0.

8. e.

0.

1144042 Ans.

EXAMPLE 2.—From 43t'16 take 9t09, in the underary scale.

OPREATION. Observe, here we say 9 from 6, we cannot, but 9 from 17 (I

XI. borrowed=11 and 6) and 8 remains, &c.

9609

35068

EXAMPLE 3,-Multiply 3426 by 567, in the octenary scale.

OPERATION.

VIII. 8496 567

Observe, we say 7 times 6 are 42, 8 (the radix) into 42 5 to carry and 2 to set down, 7 times 2 are 14 and 5 make 19, equal to 8 to set down and 2 to carry, &c.

21556

2400472 Ans.

EXAMPLE 4.—Divide 671384 by 7876, in the nonary scale.

OPERATION.

7876)671884(75<del>749}</del> Ans. 61786

the particular

Here 7876 will go into 67138 7 times (observe it would go 8 times in the decimal scale); and 7876 multiplied by 7 gives 61766, this being subtracted, gives a remainder, 5242, to which we bring down the next digit, 4, and proceed as in common division.

well all the the training

Nors.—After the units' figure is brought down, we may either write the remainder in the form of a fraction, as in example 29, or we may place a point, and annexing 0s, continue the division as in the following example.

Observe, this point is called the decimal or denary point only in the decimal system. In every other scale of notation it takes its name from the system—thus, in the duodenary or duodecimal system it is called the duodenary or duodecimal point, in the senary system, the senary point, &c.

EXAMPLE 5.—Divide t134567 by e473, in the duodenary scale.

r diving house of

15 ,0175868 . July 1 1 1 . .... 1 4 . . 1/4" 67829

47 7 HAR THAT . 97897 6 1 FAR 1 Th at the transfer of the serve 9500 and ont?

647'3

645'30 452'79

#### EXERCISE 37.

- 1. Multiply 252 by 252, in the senary scale. Ans. 122024.
- 2. Divide 32e75721 by 62te, in the duodenary scale. Ans. 62te.
- 3. From 201210 take 102221, in the ternary scale. Ans. 21212.
- 4. Multiply 57264 by 675, in the octenary scale. Ans. 51117344.
- 5. Add together 101, 1001, 1111, 1011, 1000, 1111, and 10101, in the binary scale. Ans. 1010100.

to 42

make

6. Divide 142613 by 2143, in the ceptenery scale.

Ans 50.5254+. 7. Add together 65432, 43210, 1444, 65001, and 54321, in the septenary scale.

8. From 71348 take 5e814, in the disordenary scale. Ans. 11864.

9. Multiply 3417 by 6666, in the duodenary scale.

Ans. 1136e296.

10. Divide 1010100001 by 100101, in the binary scale.

Ans. 10010 100101.

42. All the methods of proof given in Sec. II., for the fundamental rules in the common scale, apply to the various other scales; but it must be remembered that, in using the principle of the proof by nines for multiplication and division, we use, not nine, but a number one less than the radix of the scale.

Thus, in applying this principle to the proof in Example 4, eccess cast out of 57364, give a remainder 3; sevens cast out of 676, give a remainder 4, 4×3, and eccess cast out, give a remainder 5; sevens cast out of 51117846, give a remainder 5.

If the radix be 15, we cast out the 11s; if the radix be 6, we cast out the

48. Numbers containing digits to the right of the separating point, are dealt with according to the rules given in Arts. 53 and 88, Sec. II.

EXAMPLE .- Multiply 37-1443 by 6-1et in the duodenary scale. OPERATION. We place the separating point in the product so as to have XII. seven digits to the right of it, because there are four to the \$71448 right of the point in the multiplicand and three in the multiplicand three in the multiplicand and th

Chairman of the contract of the contract of the

1/11/086836

# DUODECIMAL MULTIPLICATION.

J Las mine to be in

44. The term duodecimal is commonly applied to a set of denominate fractions having 1 foot (linear, square, or cubic measure) for their unit.

The foot is supposed to be divided into 12 equal parts, called primes; each of which is divided into 12 equal parts, called seconds, &c.

> TABLE. 12 fourths" make 1 third, marked" 12 thirds " 1 second, 1 prime, 12 seconds 12 primes .... " of 1 foot,

serve and which

ither e 29, ision

only akes decia the

ale.

024. 32te. 212.

101, 100.

45. The term "inch," sometimes used in this table, is objectionable, corresponding to "prime" only when the unit is a linear foot. When the unit is a square foot, the prime is 1/2 of a square foot, or is a surface 12 inches long and 1 inch wide; when the unit is a cubic foot, the prime is 12 of a cubic foot, or is a solid 12 inches long, 12 inches wide, and 1 inch thick.

46. Let AEHG represent the surface of a rectangular table four feet in length and three in breadth. Now, if AE be divided into four equal parts, and AH into three equal parts, each of these parts, Ab, bc, fl, &c., will be 1 foot long, and if lines bk, ce, dm are drawn through b, c, and d, parallel to AH, and lines fp, lo through f and l, parallel to AE, they will divide the whole surface into the small figures, Abef, berc, &c.

And, since Ab=1 foot, and Af=1 foot, Afso is a square foot, so likewise is each of the other figures, berc, crxd, &c.

Now it is evident that there are as many vertical rows of these square feet as there are linear feet in AE, and as many squares in each row as there are linear feet in AH, that is in this case the number of square feet in the surface—4×8=12.

At the same method of proof would apply in any similar case, it appears



91

01

As the same method of proof would apply in any similar case, it appears.

that-

The area of any rectangular surface is found in square feet, and fractions of a square foot, by multiplying the number expressing how many linear feet, &c., there are in the length, by the number expressing how many linear feet, &c., there are in the breadth.

NOTE.—In linear measure, primes are linear inches; in square measure, seconds are square inches; and in cubic measure, thirds are cubic inches.

47. The example under Section 43, page 143, is, in effect, equivalent to finding the area of a rectangle, one side of which is 43 feet 1' 4" 10" and 3"" long, and the other 6 ft. 1' 11" 10" long. The answer may be translated 265 sq. ft. 10' 0" 8" 11" 8"" 3"" and 6""".

NOTE.—161, the number to the left of the separating point, is a number in the duodenary scale. In order to read it in common terms, we convert it to an equivalent number in the decimal scale (Art. 40), and thus obtain 265. It is obvious that, since the orders primes, seconds, thirds, &c., form a series of numbers descending in a 12-fold proportion from left to right, we must allow the digits to the right of the point to remain as they are.

EXAMPLE.—Find the area of a rectangular ceiling 43 ft. 4'

7" long by 20 ft. 11' 10" wide. 13 14

Here, since 45 and 20 are numbers in the common scale, we must reduce them to the duodenary scale before attaching them by the point to the other parts of the numbers. We thus obtain for the first, 37, and for the second, 18. After multiplying and pointing off four places in the product, we find 63t to the right of the point; this, reduced to an equivalent number in the common scale, gives us 910, to which we attach the other four digits, with their indices, as below. OPBRATION. XII. 87.47 18'et 30196

SECT. III.

is table, y when are foot, 2 inches bot, the long, 12

od E

t e m G likewise is ese square was there feet in the

it appears.

are feet, expressthe numbreadth.

measure, c inches. B, is, in

one side e other ted 265

a number convert it btain 265. to right, ey are. 43 ft. 4'

scale, we attaching ers. We 8. After duct, we n equiva-which we below.

48. The common arithmetical rule for duodecimal multiplication is as follows:—

#### RULE.

Write the multiplier under the multiplicand having quantities of the same denomination under each other.

Multiply each term of the multiplicand by each term of the mul-

tiplier separately.

Write the partial products under one another, so as to have quantities of the same name in the same vertical column, and add the several partial products together.

Nors.—Considering the foot to have no index, the denomination of the product of any two factors is found by adding their indices.

Thus 3"×2" give 6""; 4 ft.×7"" give 28""; 2 ft.×3 ft. give 6 ft.; 9'×11 give 99", &c.

This is commonly expressed, for the sake of brevity, by saying—feet into feet produce feet, feet into primes produce primes, &c., primes into feet produce primes, primes into primes produce seconds, &c., seconds into seconds produce forths, &c.

EXAMPLE 1.—Multiply 43 ft. 4' 7" by 20 ft. 11' 10".

OPERATION. 20 11 10 8 0 1 9" 10" Here 7 and 10, multiplied together, give us 70, and adding their indices, we see that the product is so many fourths—70", are equal to 10" to set down and 5" to carry. Next 4'×10"=40" and 5" make 45"=3" 9", &c.

910 5' 0" 2" 10""

49. In comparing this example with the previous number it will be seen that the two methods very closely agree—the only difference being that, in the latter method, upon reaching the units or feet, we drop the duodecimal scale and carry on the process in the decimal scale, while, in the former, we carry on the whole process in the duodecimal scale, and afterwards reduce that part of the expression to the left of the separating point to the common or decimal scale.

50. Provided we multiply every part of the multiplicand by every part of the multiplier, it is perfectly immaterial where we commence the process. It is customary, however, to commence, not as we have done in the last example, with the lowest denomination of both multiplier and multiplicand, but with the highest of the multiplier and the lowest of the multiplicand. Hence duodecimal multiplication is frequently called Cross Multiplication.

Example 2.—Multiply 8 ft. 2' 7" 4" by 1' 3" 7"

3 ft.		2'1	OPERATION AND STREET			OM.	ON.		
		.8	9	7 7 10	10	0""" 8	 Amin	ij.	
		4	-		8''''			Ans.	

1. Multiply 4 ft. 7' 6" 10" by 9 ft. 7' 11" 11".

Ans. 44 sq. ft. 9 1" 8" (ym 5" 5"

2. Multiply 19 ft. 10' 3" by 11 ft. 2' 4".

Ans. 222 sq. ft. 8' 0" 5" 9"".

3. Multiply 9" 7" 4"" by 7" 3"" 11""

Ans. 5" 10" 4" 11" 11" 8" 11"

4. How many square inches, &c., are there in a sheet of paper 92 inches and 5 inches 7" 4" wide?

Ans. 4' 6" 8" 6"" or 5447 sq. inches.

5. What is the superficial contents of a sheet of glass whose length is lift. M' 11" and breadth Sft. 2' 2"?"

Ass. 23 sq. ft. 6' 9" 7" 10"".

51. The solid contents are found by multiplying together the length, breadth, and thickness.

EXAMPLE.—How many cords of wood are there in a pile 79 ft. 8 inches long, 4 ft. 2 inches wide, and 7 ft. 11 inches high?

> OPERATION. SECOND METHOD. FIRST METHOD. 237:04 214348 741774 2627 10' 8" 8"'÷128. (number of ft. in cord)

No. of ft..in cord = t8)162e t88(18.64469 duodenary 20140 fb cords. Ans. 2927828 com. scale. 714

<sup>\* 14 12</sup>x + 17xx, &c. of a square foot.

un gum

5/11 9/1/

ווווווווון

of paper . inches.

7" 10"".

together

ile 79 ft. igh?

ETHOD.

t. in cord) cords. Ans.

### ARTS. 51, 52.]

#### EXERCISE 39:

1. Multiply together 15 ft., 1 ft., 1 ft. 2', and 8'.

Ans. 11 cubic ft. 8'=11 cubic ft. 1152 cubic in.

2. Multiply together 53 ft. 6 in., 10 ft. 3 in., and 2 ft.

Ans. 1096 cubic ft. 9'.

3. How many cords of wood in a pile 10 ft. long, 5 ft. high, and Ans. 2 cords 94 cubic ft.

4. How many cords of wood are there in a pile 4 ft. wide, 5 ft. 3 in. high, and 70 ft. long?

5. What are the exact cubic contents of a block of marble 4 ft.

7'8" long by 9 ft. 6' wide and 2 ft. 11' thick?

Ans. 128 cubic ft. 6' 5" 2".

6. How many bricks, 8 inches long, 4 inches wide, and 2 inches thick, will it require to make a wall 25 ft. long, 20 ft. high, and 2 ft. 6 inches thick? Ans. 33750 bricks.

52. It is sometimes asked how we can multiply feet, inches, &c., by feet, inches, &c., while we cannot multiply pounds, shillings and pence. The answer is very simple.

1st. When we say that feet multiplied by feet give square feet, we merely use, as we have seen, (Art. 46), an abbreviated form of expression for the following, viz: that "the number of square feet contained in any rectangular surface, is equal to the product of two numbers, one of which represents the number of linear feet in one side; and the other the number of linear feet in the adjacent side."

2nd. When we are multiplying together prines, seconds, &c., we are merely multiplying together a set of factors having 12 or powers of 12 for denominators; and when we say that seconds multiplied by fourths, give staths; primes, multiplied by seconds, give thirds, &c., we simply mean that the product of any two of these fractions is a fraction having for its denominator a power of 12, which power is indicated by the sum of the indices of the factors. indices of the factors.

It is hence obvious that duodecimal multiplication affords no support whatever to the idea that money may be multiplied by money.

#### QUESTIONS TO BE ANSWERED BY THE PUPIL.

Norn.—The numbers after the guestions refer to the articles of the Section.

- 1. What is the measure of a number? (1)
  2. What is the multiple of a number? (2)
  3. What is an integer? (6)
  4. Of how many kinds are integers? (4)
  5. What is an even number? (5)
  6. What is an odd number? (6)

- 6. What is an odd number? (6)
  7. What is a prime number? (7)
  8. What is a composite number? (8)
  9. What are the factors of a number? (9)
  10. By what other names are factors known? (10)
  11. What is a common measure of two or more numbers? (11)
  12. When are two or more numbers prime to each other? (12)
  13. Are all prime numbers prime to each other? (12)
  14. Are all composite numbers prime to each other? (12)
  15. What are commensurable numbers? (13)
  16. What is a grane number? (15)

18. What is a cube number? (16)

19. What is a perfect number? (17)

- 20. Mention some perfect numbers. How do all perfect numbers terminate? (17)
- 21. What are *omicable* numbers? Mention some amicable numbers. (18)
  22. What is meant by the *properties of numbers*? (19)
  23. What is the sum of two or more even numbers? (19-I.)

24. What is the difference of two even numbers? (19-II.)
25. What is the sum of 3, 5, 7, &c., odd numbers? (19-IV.)
26. What is the sum of 2, 4, 6, 8, &c., odd numbers? (19-V.)
27. What is the sum or difference of an odd and an even number? (19-VI.)

When is the product of any number of factors even? (19-IX.)
When is the product of any number of factors odd? (19-XI.)
When will a number measure the sum, difference and product of two numbers? (19-XII.)

numbers? (19-XIII.)

31. If the number 9 be multiplied by any single digit to what is the sum of the digits in the product equal? (19-XVI.)

32. By what is any number ending in 0 divisible? (19-XX.)

33. By what is any number ending in 2 divisible: (19-XXX.)

34. By what is any number ending in 2 divisible? (19-XIX.)

35. When is a number divisible by 4? (19-XXII.)

36. When is a number divisible by 9? (19-XXIV.)

38. When is a number divisible by 9? (19-XXV.)

39. When is a number divisible by 11? (19-XXV.)

39. When is a number divisible by 11? (19-XXVI.)

40. Show that every composite number may be resolved into prime factors. (19-XXVII.)
41. Show that the least divisor of any number is a prime number.

(19-XXVIII.)

42. With what digits must all prime numbers except 2 and 5 terminate? (19-XXXI.)

13. How do you find the prime numbers between any limits? (20)

When it is required to ascertain whether a given number is prime or not, what is the first thing we do? (20)
When we try the primes of the table as divisors, which is the highest

we need use? (20)

7. Why is it unnecessary to try any divisor greater than the square root of the number? (20)
16. How do we resolve a composite number into its prime factors? (21)

19. By what numbers can a composite number he divided? (21-Note.)

What is the rule for finding all the divisors of a number ? (22

51. How do we find simply how many divisors a number has? (23)
52. What is the greatest common measure of two or more numbers? (24)
53. How do we find a common measure of two or more numbers? (25)

54. How do we find the greatest common measure of two numbers? (26)
55. Prove the rule in Art. 26.
56. How do we find the G. C. M. of three or more numbers? (27)
57. What is the second method of finding the G. C. M.? (28)

Upon what principle does this method rest? (28)

What is a common multiple of two or more numbers? (29)
What is the least common mutiple of two or more numbers? (30)
Give the first rule for finding the l. c. m. of two or more numbers. (31)
Give the second rule. (32). What is the reason of this rule? (32)

Give the most convenient and expeditious rule for finding the l. c. m. of several numbers. (33)
What is meant by the radix or base of a system of notation? (34)

How do we read numbers in different scales ? (35)

Express the number 234213 quinary as in Art. 36.

What does the expression 10 always represent? (37)
What is the highest digit used in any scale? (38)
How do we reduce a number from one scale to another? (39)

rmin-(18)

9-VI.)

of two

e sum

ae fac-

mber.

inate?

ime or

aighest

re root

(24)

(26)

rs. (31)

l. c. m.

- 70. What is the rule for transforming a number from any scale into the
- decimal? (40)
  71. How are the fundamental operations carried on in the different scales? (41).
  72. How is the separating point named in the different scales? (41-Note.)
  73. How are operations in the different scales proved? (42)
  74. What are duodecimals? (44)

- 75. Give the table of duodecimals. (44)
  76. What is a prime? (45)
  77. How is the area of a rectangular surface found? (46)
  78. What is the rule for duodecimal multiplication? (48)
  79. How may the rule for fluding the denomination of the product be concisely worded? (48)
  80. How are solid contents found? (51)
  - 81. Show that duodecimal multiplication affords no support to the idea that money may be multiplied by money, &c. (52)

# Exercise 40.

### MISCELLANEOUS EXERCISE.

(On preceding rules!) All the service of the servic

- 1. Add together \$729.18, \$710.50, \$166.78, £9314s. 71d., £276 198. 101d., \$497.81 and £275 4s. 112d. All Control
- 2. Multiply 47 miles, 6 fur. 17 per. 4 yds. 2 ft. 7 in. by 576.
- 3. How many divisors has the number 243000?
- 4. From 713427 octenary take 4234434 quinary and give the answer in both scales.
- 5. Divide 79.342 by .00006378. Appear to strate ye as we
- 6. Express 79423 and 234567 in Roman numerals.
- 7. What is the l. c. m. of 5, 7, 9, 11, 15, 18, 20, 21, 22, 24, 28, 30, 33, 35, 36, 40, 42, 44, 45, 48, and 50.
- 8. Give all the readings of 376.342.
- 9. Multiply 64276.3427 by 9999993000.
- 10. Transform 78263 nonary into the quinary and undenary scales and prove the results by reducing all the numbers to the septenary scale.
- 11. Form a table of all the prime numbers less than 200.
- 12. Reduce £672 7s. 7d. to dollars and cents.
- 13. What is the G. C. M. of 243000, 891, 37800 and 35100.
- 14. Give all the readings of 6 yards 3 qrs. 3 nails 2 inches.
- 15. Write down as one number, seven hundred and forty-two quintillions, nine hundred and five billions, seventy-eight thousand and fourteen, and eighty-seven million, two hundred thousand and eleven tenths of trillionths.
- 16. Read the following numbers:

71300100200401.000000070402

134900101000100100.000200020002 47000000000007·000000000000278

- 17. Add together £178 16s. 43d., £97 15s. 114d., £693 19s. 113d., £216-11s. 9\d., £678 14s. 7\d., £197 13s. 11\d., £117 6s. 5d., and £91 1s. 13d.
- 18. What are the prime factors of 276000?

ev

m

th

de

th

m

nu

di

21

2n

3r

5t

- 18. Multiply 6.ft. 2 7"-9" 10" by 13 ft. 11" 11" 11" 11"
- 20. Divide 7te9:047 by 713196 in the duodenary neals.
- 21. What number in the common scale is the greatest that can be expressed by seven figures in the quaternary scale?
- 22. What number in the common scale is the least that can be expressed as an integral number by five figures in the octenary scale?
- 23. Reduce 74002702 square inches to acres;
- 24. What is the least common multiple of 240, 780, 1620, and 1728?
- 25. Divide \$7894 16 among 3 men, 4 women and 6 children, so that each woman shall have twice as much as a child and each man 5 times as much as a woman. What is the share of each?
- 26. What are the greatest and least integral numbers in the common scale that can be expressed by 10 figures in the binary scale?
- 27. Divide 729 yds. 3 qrs. 3 na. 1 in by 7(yds: 1 qr. 1 na. 1 in.
- 28. Multiply 762.4978 by 63.423:
- 29. From 723426 take 938:9126141.
- 30. From 129 lb. take 63 lb. 4 oz. 7 drs. 2 ser.
- 31. What are the divisors of 1064?
- 32. How many yards of carpet 2 ft. 7/in. wide, will be required to cover a floor 30 ft. 6 in. long and 20 ft. 11 in. wide?

# SECTION IV.

# VULGAR AND DECIMAL FRACTIONS, &c.

- 1. A fraction is an expression, representing one or more of the equal parts into which any quantity may be divided.
- 2. If a quantity be divided into 2, 5, 9, or 34, &c., equal parts, then one of these parts is called one half, one fifth, one-ninth, or one-thirty-fourth, &c., as the case may be.

and half is written 1	one-ninth is written
	one-hundredth is written 180
one founth is written	one-sixty-eighth is written-
one of this written	oleran comente antha in smitter
	eleven-seventeenths is written

3. The division of one number by another may be in-

directed in three different ways, vin : by using the full sign of division, -- or either of its parts, --, or :

Thus we may indicate the division of 17 by 8, by writing them thus 17-48, or thus 17: 8, or thus 17:

Now the last of these, vis: W is a fraction, and so in every other case, a fraction indicates the division of one number, called the numerator, by another number, called the denominator.

4. In a fraction the number below the line is called the denominator, because it indicates into how many equal parts the unit is divided,—i. e., it tells the denomination of the parts. The number above the line is called the numerator, because it numerates or tells how many of these equal parts are to be taken. (Art. 2)

5. The numerator and denominator are called the terms

of the fraction.

6. Since every fraction expresses the division of the numerator by the denominator, it follows that—

The value of the fraction is the quotient obtained by

dividing the numerator by the denominator.

7. Hence, 1st. When the numerator is less than the denominator, the value of the fraction is less than 1.

2nd: When the numerator is equal to the denominator the value of the fraction is equal to 1.

3rd. When the numerator is greater than the denominator the value of the fraction is greater than 1.

8. From (Art. 6) and (Arts. 79—84, Sec. II.) it is manifest that

1st. Multiplying the numerator of a fraction by any number multiplies the fraction by that number.

2nd. Multiplying the denominator of a fraction by any number divides the fraction by that number.

3rd. Multiplying both numerator and denominator of a fraction by the same number does not affect the value of the fraction.

4th. Dividing the numerator of a fraction by any number divides the fraction by that number.

5th. Dividing the denominator of a fraction by any number multiplies the fraction by that number.

uired ?

can

n be:

the.

and

n, 80

and

hare

a the

a the

1 in.

more ided.

may

1 1 d o

itten

be in-

6th. Dividing both n merator and denominator of a fraction by the same number does not affect its value.

9. Fractions are divided into two classes : - vulgar and decimal.

10. A Decimal Fraction is a fraction in which the denominator is 1, followed by one or more 0s. want doing we have

11. All other fractions are called Vulgar or Common

Fractions.

Note. The word vulgar is here used in the sense of common.

12: There are six kinds of vulgar fractions proper, improper, mixed, simple, compound, and complex.

13. A Proper Fraction is one in which the denominator

is greater than the numerator.

A Proper Fraction may also be defined to be a fraction whose value is less than 1.

Thus  $\frac{1}{13}$ ,  $\frac{4}{5}$ ,  $\frac{1}{15}$ ,


The following diagrams represent unity, seven-sevenths, and the proper fraction, five-sevenths.



The very faint lines indicate what 4 wants to make it equal to unity and identical with I. In the diagrams which are to follow, we shall, in this manner, generally subjoin the difference between the fraction and unity.

The teacher should impress on the mind of the pupil that he might have chosen any other unity to exemplify the nature of a fraction.

14. The following will show that 4 may be considered as either the \$ of 1 or the \$ of 5, both—though not identical-being perfectly equal.



In one case we may suppose that the five parts belong to but 1 unit; in the other, that each of the five belongs to different units of the same kind.

frao-

REF CLES

and

odemon

oper, or sq nator

whose

enths,

F1 15.0


ity and

in this

unity.

idered dentiARTS. 9-19.]

Lastly, 4 may be supposed as the 2 of one unit five simes as large as the former; thus ( ) did the real ( ) at the supposed of ( )



15. An Improper Fraction is a fraction whose denominator is not greater than its numerator.


An Improper Fraction may also be defined to be a fraction whose value is equal to or greater than 1.

Thus, 2, 16, 7; 11, 262, 113, 3, 28, &c., are improper fractions.

16. A Mixed Number is a number made up of a whole number and a fraction.

Thus, 163, 1934, 113, 9991, 6,3, 21, &c., are mixed numbers.

17. An Improper Fraction is always equal either to a whole number or to a mixed number. The following will exemplify an improper fraction, and its equivalent mixed number:



18. A Simple Fraction expresses one or more equal parts of unity.

Thus, 4, 8, 6, 11, 4, 187, &c., are simple fractions.

19. A Compound Fraction expresses one or more equal parts of a fraction; or in other words, is a fraction of a fraction.

a rollife outer of the last make how appet you

Thus, \$ of \$, \$ of \$ of 1\$ of \$ of 1\$2, &c., are compound fractions.

init: in

Jirli.

ARTI

min

3.

to

gin

OP

180: 1 of 3 means, not the four-nintles of unity, but the four-nintles of the three-rourths of unity:—that is, unity being divided into such three of these are to be divided into nine parts and then four of these nine are to be taken; thus—



Note.—The word "of," placed between the several parts of a con pound fraction, is equal to end may be replaced by x, the sign of multiplication.

21. A Complex Fraction is one having a fraction on a mixed number in its numerator or denominator, or in both.

Thus, 
$$\frac{2}{3}$$
,  $\frac{2}{7}$ ,  $\frac{3}{7}$ ,  $\frac{4}{7}$ ,  $\frac{91}{11}$ ,  $\frac{4}{1873}$ ,  $\frac{71}{213}$ , &c. are complex fractions.

NOTE. 

means, that we are to take the fourth part, not of unity, but of the populary. This will/be exemplified by—



22. Since fractions, like integers, are capable of being increased or diminished, they may be added, subtracted, &c.

23. Every integer may be considered as a fraction having unity, for its denominator.

Thus, 13 may be written 1,2; 6, 4; 29, 29, &c.

# REDUCTION OF FRACTIONS.

24. Since (Art 8) multiplying both numerator and denominator by the same number does not alter the value of the fraction, we may reduce an integer to a fraction having any proposed denominator, by the following:—

#### A DEPOS LEGICAL.

Write the integral number in the form of a fraction having 1

for its denominator. (Art. 23.)

And multiply both numerator and denominator of the resulting expression by the proposed denominator. (Art. 8.)

EXAMPLE 1.—Reduce 16 to a fraction having 11 for its denominator.

EXAMPLE 2.—Reduce 173 to a fraction having 31 for its denominator.

### Exercise 41.

- 1. Reduce 29 to a fraction having 12 for its denominator.
- Ans. 344. 2. Reduce 243 to a fraction having 3 for its denominator.
- 3. Reduce 7, 23, and 101 to fractions having 13 for denominator.
- Ans. 91, 292, 1913. 4. Reduce 4, 37, 126, 73, and 1007 to fractions having 101 for denominator.
- 5. Reduce 204, 7011, and 1999 to fractions having 207 for denominator.

25. Let it be required to reduce the mixed number 817 to an improper fraction.  $8\frac{7}{11}$  is equal to the whole number 8, and the fraction  $\frac{7}{11}$ , and by (Art. 24.)  $8 = \frac{6}{11}$ , therefore  $8\frac{7}{11} = \frac{6}{11} + \frac{7}{11} = \frac{6}{11}$ .

Hence, to reduce a mixed number to an improper fraction, we deduce the following:

Multiplying the whole number by the denominator of the fraction, to the product add the given numerator and place the sum over the given denominator.

BEAMPLE 1.-Reduce 78% to an improper fraction.

EXPLANATION.—We multiply the whole number, 78, by 9 and add in the numerator, 4. This gives us 661, which we write over the given denominator, 2, and the resulting fraction, 181, is the improper fraction sought. OPERATION. 734 ngl Ans.

EXAMPLE 2.—Reduce 27617 to an improper fraction.

D ar

oth.

ions.

hut

eing &c. hav-

lidegeref aing

rec lon rec

nu

eq

for

ne

### EXERCISE: 42.

1. Reduce the mixed numbers, 7314, 1814, and 12818 to improper

fractions.

2. Reduce the mixed numbers 384f, 673 4, 4792 6, and 568 7. to improper fractions.

Ans. 3461, 2767, 119801, and 16474.

26. Since every fraction indicates the division of the numerator by the denominator—to reduce an improper fraction to a mixed number, we have the following:

Divide the numerator by the denominator and the quotient will be the required mixed number.

Example 1.—Reduce 294 to a mixed number. 204 = 204 : 7 = 291 Ans.

EXAMPLE 2.—Reduce 20047 to a mixed number. 20047-11 = 1822-4 Ans.

#### The market ten FEXERCISE: 43.

Reduce the improper fractions 407, 2082, and 1847.5 0 mixed numbers.
 Ans. 31, 47, 47, 43, and 16, 12, 17.
 Reduce the improper fractions 2847, 2841, and 2841 to mixed numbers.
 Ans. 8834, 15845, and 78.

27. To reduce a fraction to its lowest terms:

Divide both terms by their greatest common measure.

This is simply dividing both terms by the same number—which does not affect the value of the fraction. (Art. 8.)

The greatest common measure may be found by (Art. 26, Sect. III.) or, very frequently, by inspection.

Example 1.—Reduce 50 to its lowest terms.

Greatest common measure = 25. Dividing both terms by 25;  $40 = \frac{2}{3}$  Ans.

EXAMPLE 2.—Reduce 136 to its lowest terms.

Greatest common measure of 126 and 162 = 18.

Dividing both terms by 18 we get  $\frac{126}{61} = \frac{7}{9}$  Ans.

### EXERCISE 44.

- 1. Reduce #180 to its lowest terms.
- 2. Reduce \$7378 to its lowest terms

Ans. Tto.

Ans. 1591.

proper 1988 568

OF, IV.

f the proper

nt will

43 - 13

4 13 .5 17,0 .0. 61217. and to nd 78.

- 22 - 3

loes not

(Art.

Ans.

3. Reduce \$4944 and \$14 to their lowest terms. Ans. 1 and 1. 4. Reduce 1948, Ala and 1111 to their lowest terms.

Ans. 12, 47, and 7388. 28 Instead of dividing both terms by their greatest common measure we may divide both by any common measure. We thus reduce the fraction to lower terms, and, continuing the division as long as the terms have a common measure, we shall finally have reduced the fraction to its lowest terms.

Note.—It is advisable to commit to memory the properties of numbers given in Art. 19, Sec. III from XVIII to XXIV. EXAMPLE 21.—Reduce 329480 to its lowest terms.

282489 dividing by 10. (XXI. of Art. 19, Sec. III.) = 111 dividing by 8. (XXIII. of Art. 19, Sec. III.)
= 111 dividing by 9. (XXIV. of Art. 19, Sec. III.) 18 dividing by 3. (XXV. of Art. 19, Sec. III.) 183 Ans.

EXAMPLE 22.—Reduce 2374 to its lowest terms.

38 44 dividing by S. (XX. in Art. 19, Sec. III.) = 18 dividing by 9. (XXIV. in Art. 19, Sec. III.) 4+ dividing by 8. (XXV. in Art. 19, Sec. III.)

Ans.

#### Exercise 45.

1. Reduce 3ff to its lowest terms. Ans. 17. Reduce 134500 to its lowest terms.
 Reduce 1375000 to its lowest terms.
 Reduce 15750 to its lowest terms. Ans. 3110. Ans. 7.

5. Reduce 308, 5149 and 16238 to their lowest terms. Ans. 11, 2381, and 181.

Ans. Agg.

11 25 11

29. To reduce fractions of different denominators to equivalent fractions having the same denominator:-

Multiply each numerator by all the denominators except its own for a new numerator, and all the denominators together for a new denominator.

This is merely multiplying both numerator and denominator of each fraction by the same quantity, vis: the product of all the other denominators, and consequently (Art. 8.) it does not alter the value of the fraction.

Example 1.—Reduce 3, 7 and 4 to a common denominator.

8×11× 9 = 297 = 1st numerator. 7× 4× 9 = 352 = 2nd numerator. 5× 4×11 = 220 = 3rd numerator. 4×11× 9 = 396 = common denominator.

Therefore the equivalent fractions are \$97, 148, and \$49

EXAMPLE 2.—Reduce 1, 1, 4, and 7 to equivalent fractions having a common denominator.

1×5×7×11=885=1st numerator.

5×6×7×11=485=3nd numerator.

4×5×5×11=440=3nd numerator.

9×5×5×7=680=46h numerator.

9×5×7×11=776=800mmon denominator.

And the equivalent fractions are 348, 448, 448 and 778.

#### EXPROISE 46.

- 1. Reduce 3, 4, 5, and 7, to equivalent fractions having a common denominator.
- 2. Reduce \$1, 12, and \$2 to fractions having a common denominator.
- 3. Reduce 4, 4, 4, and to fractions having a common denominator.
  - Ans. +28+2, 1600's, 14014, 14014, and 74014.
- 4. Reduce 14, 4, and 15 to a common denominator.
- 5. Reduce \$, \$, \$, and \$\frac{1}{2}\$ to a common denominator.
- 6. Reduce 1, 3, 3, and 4 to a common denominator.

  Ans. 195, 146, 175, and 200.
- 30. To reduce fractions to equivalent fractions having their least common denominator:—

#### RULE.

Find the least common multiple of all the denominators. (Arr. 33, Sec. III.)

Multiply both terms of each fraction by the quotient obtained by dividing this least common multiple by the denominator of that fraction.

This is merely multiplying both terms by the same quantity, as in Art.

EXAMPLE 1.—Reduce 1, 7, 3, and 5 to their least common denominator.

The least common multiple of 4, 12,3, and 15, is 48.

Multiplying both terms of the 1st fraction by 12 (i.e. 48) it becomes \$8.

Said by 18 (i.e. 48) it becomes \$8.

"" Said by 18 (i.e. 48) it becomes \$8.

actions

EXAMPS 2.—Reduce 4, 11, 18, 11, 18, and 2 to their least common denominator.

The least common multiple of 6, 11, 20, 44, 55, and 4, is 220. The multiplier for both terms of the first fraction is  $s \neq 0 = 44$ , for second. 320; for the third, 320 = I1; for the fourth, 320 = 5; for the fifth,

Multiplying by these numbers, we obtain 178, 188, 188, 180, 180,

and 199 for the required fractions.

### EXERCISE 47.

1. Reduce 4, 1, 1, 1, and 76 to their least common denominator. Ans. 140, 140, 130, and 100.

Aus. \$78, \$48, 188, and 500.

6. Reduce 1, 2, 1, 5, 7, 11, 16, and 11 to their least common denominator.

7. Reduce 4, 12, 18, 27, 28, and 15 to their least common denominator.

Ans. 4888, 4888, 4888, 4888, 4888, 4888, and 4888.

8. Reduce 44, 4, 4, 14, 27, 48, 4, and 48 to their least common denominator.

Ans. 8614, 1926, 1920, 8478, 5948, 8778, 5218, and 7828.

31. Let it be required to reduce 15 of 15 to a simple fraction.

13 of 15 means 12 times 17 of 11. We get 17 of 17, i. e. divide 17 by 17, when we multiply the denominator 11 by 17 (Art. 8). Therefore 17 of 16 = 18 17, and to multiply this result by is, we multiply the numerator, 8, by 18, (Art. 8.)

Therefore  $\frac{1}{13}$  of  $\frac{6}{15} = \frac{6 \times 12}{11 \times 17} = \frac{73}{187}$ .

Hence to reduce a compound fraction to a simple one we deduce the following:

Multiply all the numerators together for a new numerator, and all the denominators together for a new denominator.

aving a

n deno-1. 708g. common

1 7007

nd polor.

nd

and for.

having

. (Arr.

tained by of that

as in Art.

common

ecomes 13. somes ? ecomes ? ecomes 37 inator, are

de

# EXAMPLE 3.—Reduce \ of \ of \ to a simple fraction.

 $\frac{2\times4\times5}{3} = \frac{140}{189}$ 

NOTE.—In all cases the answer must be reduced to its lowest terms.

#### Exercise 48.

1. Reduce \$ of 3 of 1 of 75 to a simple fraction.

2. Reduce \$ of \$ of \$ of \$ of \$\frac{3}{100}\$ of \$\frac{3}{2}\$ to a simple fraction.

Ans. \$\frac{3}{2}\$.

3. Reduce \$\frac{3}{2}\$ of \$\frac{3}{10}\$ of \$\frac{3}{2}\$ to a simple fraction.

Ans. \$\frac{7}{2}\$.

4. Reduce \$\frac{3}{2}\$ of \$\frac{4}{2}\$ of \$\frac{1}{3}\$ to a simple fraction.

Ans. \$\frac{7}{6}\$. Ans. 3126.

32. Since the several numerators of the compound fraction form the factors of the numerator of the simple fraction, and also the several denominators of the compound fraction, the factors of the denominator of the simple frac-

Before applying the rule in (Art. 31) we may cast out or cancel all the factors that are common to a numerator and a denominator of the compound fraction.

Example 1.—Reduce of of \$ of \$ of \$ of \$6 to a simple fraction.

CANCELLED.

$$\frac{6}{11} \text{ of } \frac{4}{7} \text{ of } \frac{3}{5} \text{ of } \frac{29}{27} \text{ of } \frac{35}{16} = \frac{6 \times 4 \times 3 \times 22 \times 35}{11 \times 7 \times 5 \times 27 \times 16} = \frac{6}{11 \times 7 \times 5} \times \frac{32}{27 \times 16} \times \frac{32}{27 \times 16} = \frac{1}{3} \text{ Ans.}$$

Here 6 and 27 contain a common factor, 3, which is cast out, and these numbers thus reduced to 2 and 9. Next this 2 reduces 16 to 8, and the 9 is reduced to 3 by the third numerator, which is thus cancelled. Again, 11 cancels 11 (the first denominator) and reduces 22 to 2, and this 2 reduces the 8, before obtained from the 16, to 4. Next, this 4 is cancelled by the 4 in the numerator. Again, 7 cancels the 7 in the denominator and reduces the 35, in the numerator, to 5, and this 5 cancels the 5 in the denominator. All the numerators are now reduced to unity, as also all the denominators but the fourth, which is 3. The resulting fraction is therefore  $\frac{1\times i\times 1\times 1\times 1}{1\times 1\times 1\times 3\times 1}$ but this is simply 1.

EXAMPLE 2.—Reduce 1 of \$ of \$ of \$ to a simple fraction.

STATEMENT. CANCELLED.

$$\frac{7}{11} \text{ of } \frac{4}{6} \text{ of } \frac{3}{5} \text{ of } \frac{55}{20} = \frac{7 \times 4 \times 3 \times 55}{11 \times 6 \times 5 \times 20} = \frac{7 \times 4 \times 3 \times 55}{11 \times 6 \times 5 \times 20} = \frac{7}{11 \times 6 \times 5 \times 20} = \frac{7}{2 \times 5} = \frac{7}{2 \times 5} = \frac{7}{10} \text{ Ans}$$

Note.—If any of the terms of the compound fraction are whole or mixed numbers, they must be reduced to fractions (Arts. 23 and 25).

The process of cancelling exemplified above should always be adopted when possible.

#### EXERCISE 49.

1. Reduce 5 of 6 of 3 of 16 to a simple fraction. Ans. 5%.

2. Reduce \( \frac{3}{2} \) of \( \frac{1}{3} \) of

3. Reduce \$ of 4 of 51 to a simple fraction.

Ans. 4.

4. Reduce \(\frac{1}{3}\) of \(\

5. Reduce of of of of of 24 of 34 of 64 to a simple fraction.

Ans. Zs.

6. Reduce 4 of 3 of 154 to a simple fraction.

Ares. 24.

33. Let it be required to reduce the complex fraction -- to a simple fraction.

Since (Art. 8) we may multiply both numerator and denominator of a fraction by the same number, without altering its value—we may multiply both terms of the given fraction by 4, i. e., by the denominator with its terms inverted, without altering its value.

Therefore 
$$\frac{4}{3} = \frac{4 \times 4}{3 \times 4} = \frac{4 \times 4}{1} = 4 \times 4 = \frac{6 \times 4}{7 \times 8}$$

Hence, to reduce a complex fraction to a simple one, we deduce the following:

Reduce the expression (Arts. 23 and 25) to the form of fraction

i. e., reduce both numerator and denominator to simple fractions. Then multiply the extremes or outside numbers together for a new numerator, and the means or intermediate numbers together for a new denominator.

EXAMPLE 1.—Reduce 41 to a simple fraction.

Note.—Factors that are common to one of the extremes and one of the means, are to be struck out or cancelled. (Art. 33).

EXAMPLE 2.—Reduce  $\frac{711}{111}$  to a simple fraction.

ns. TI-

ns. 70.

pound simple pound

e fracr cancel minator

simple

nd these d the 9 is Again, 11 2 reduces by the 4 reduces minator. XIXIX1 X1X8X1

10 Ans.

raction.

or mixed adopted

### EXERCISE 50.

1. Reduce 
$$\frac{\frac{1}{4}\frac{5}{6}}{\frac{1}{2}\frac{1}{6}}$$
 to a simple fraction. Ans.  $\frac{5}{27}$ .

2. Reduce 
$$\frac{11}{7\frac{1}{16}}$$
 to a simple fraction. Ans.  $\frac{2}{6}$ .

4. Reduce 
$$\frac{11\frac{3}{3}}{12\frac{3}{5}}$$
,  $\frac{3\frac{1}{4}}{9}$  and  $\frac{3}{3}$  to simple fractions.

Ans. 
$$\frac{175}{202}$$
,  $\frac{13}{36}$ , and  $\frac{19}{29}$ .

5. Reduce 
$$\frac{12}{153}$$
,  $\frac{53}{76}$  and  $\frac{23}{33}$  to simple fractions.

6. Reduce 
$$\frac{16\frac{2}{3}}{11\frac{2}{3}}$$
,  $\frac{6\frac{1}{6}}{13}$ ,  $\frac{17}{18\frac{1}{3}}$ ,  $\frac{21\frac{1}{6}}{10\frac{2}{3}}$ , and  $\frac{1}{4\frac{3}{6}}$  to simple fractions.

Ans.  $1^2$ ,  $\frac{3}{6}$ ,  $\frac{5}{6}$ ,  $\frac{5}{6}$ ,  $\frac{2}{10}$ , and  $\frac{5}{6}$ .

# 1 34. A denominate fraction is a fraction of a denominate number.

Thus, 4 of a lb., 7 of a mile, 3 of a day, &c., are denominate fractions.

25. Reduction of denominate fractions consists in changing them from one denomination to another without altering their values.

36. Let it be required to reduce \$\displays \text{ of a pint to the fraction of a bushel.}

Since 1 qt. = 2 pints, \$\displays \text{ of a pint} = \displays \text{ of a quart.}

Also because 1 gal. = 4 qts. 4 of a pint = 2 of 1 of a gal.

Similarly 4 of a pint = 1 of 1 of 1 of 4 of a bushel = 11 = 1 bushel.

Hence to reduce a denominate fraction from a lower to a higher denomination, we deduce the following:

#### RULE.

Take the number expressing how many of the given denomination are required to make one of the next higher; also the number expressing how many of this denomination are required to make one of the next higher again, and so on until the required denomination be reached.

Write the fractions formed by these numbers as denominators, with 1 as numerator and the given fraction in the form of a compound fraction, which reduce to a simple fraction. (Art. 31.)

RCT, IV.

Example 1.—Reduce 3 of a minute to the fraction of a week. Ans. 3 of do of de of de anden of a week.

Example 2.—Reduce \$4 of a grain troy, to the fraction of an ounce.

64 of 14 of 10 ale of an oz. Troy.

### EXERCISE 51.

- 1. Reduce 4 of an oz. to the fraction of a pound, avoirdupois. Ans. Jo lb.
- 2. Reduce of of a penny to the fraction of a pound. Ans. £ BAG.
- Ans. A wk. 3. Reduce 3 of 83 days to the fraction of a week.
- 4. Reduce of of 16 nails to the fraction of an English ell. Ans. T. E.e.
- 5. Reduce 3 of 4 of a yard to the fraction of a perch. the state and a day and Ans. An per.
- 6. Reduce 3 of 4 of  $21_{14}$  of a cord foot to the fraction of a cord. Ans. 1 cord. The same of the sa
- 7. Reduce 19 of 17 of 91 square perches to the fraction of an for to the secretary screen

37. Let it be required to reduce 1 of a day to the fraction of a minute. Since there are 24 hours in a day and 60 minutes in an hour;

\$ of a day will be 24 times \$ of an hour and 60 times 24 times \$ of a minute; that is,  $\frac{1}{2}$  of a day is equal to  $\frac{1}{2} \times \frac{24}{24} \times \frac{60}{24}$  of a minute.

Therefore  $\frac{1}{2}$  of a day  $\frac{1}{2}$  of  $\frac{1}{2}$  of a minute  $\frac{1}{2}$  minute.

Hence, to reduce a denominate fraction from a higher to a lower denomination, we have the following:

Take the number expressing how many of the next lower denomination make one of the given denomination; also, the number, expressing how many of the next lower again make one of this denomination, and so on till the required denomination be reached.

Write the fractions formed by these numbers as numerators, with 1 as denominator, as the given fraction in the form of a compound

fraction, which reduce to a simple fraction. (Art. 31.)

EXAMPLE 1.—Reduce } of a £ to the fraction of a penny. g of ep of 49 = 100 pence.

EXAMPLE 2.—Reduce 1 of 5 of 14 of a furlong to the fraction

. 18 1 3 of 4 of 18 of 14 of 1 of 3 = 800 ft. Ans.

and if.

and  $J_0$ . tions.

and 46.

minate

ctions. chang-

ushel. THALA.

bushel. wer to

ination ber exzke one ination

nators. a com-

# A Transit & Exercise 52. Transit - Taylor, T.

1. Reduce 14 of a bushel to the fraction of a quart.

2. Reduce & of a gal. to the fraction of & of & of a gill.

Ans. 150.

3. Reduce 7 of 2 pecks to the fraction of 1 of 3 of a pint.

4. Reduce 17 of a lb. to the fraction of a scruple.

- 5. Reduce soon of 3 of 2 of 11 of 29 of a lb. avoirdupois to the Ans. 1376 dr. fraction of a dram.
- 38. To find the value of a denominate fraction in terms of a lower denomination:

# Antes who mercaning bulk. how him is in a company

Divide the numerator by the denominator according to the rule riven in Art. 71, Sec. II.

This is only actually performing the work which the fraction indicates. "in the life " the distance in the

EXAMPLE.—What is the value of 11 of a mile?

#### 11 miles : 13

88 = number of furlongs.

or in the tree of the property and the delication

1007 100 40 = perches in furlong.

400 = perches. W. Traffic Black

han the state of = yards in a perch.

THE STATE OF STAN BUFF, LESS and the state of yards and to and amount content to and the state of t

with the state of the tree of a grant - and the state of

When the french one former to there and & one has reintered and the

# Exercise 53. 'At an and see a constant of the

1. What is the value of 3 of a bushel and also of 4 of a lb. avoirdupeis? Aus. 1 pk. 0 gal. 0 qt. 1 pt. and 13 oz. 11 drams.

2. What is the value of 13 of a yard of cloth?

Ans. 2 grs. 0 na. 115 inches.

8. What is the value of § of a lb. troy; and also of 11 sq. mile?

Ans. 10 oz. 13 dwt. 8 grs.; and 62 acres, 1 rood, 8 sq. per. 4 sq. yds. 2 ft. 79 1 in.

4. What is the value of \$ of a furlong; and of \$ of a £? Ans. 35 rds. 3 yds. 0 ft. 2 in.; and 11s. 5.d. 40. Addition of the thousand present of this is

A che to the 39. Let it be required to reduce 2s. 72d. to the fraction of 27. 18s.

28. 7\frac{7}{20. 7\frac{7}{2}d.} = \frac{127}{7584} \frac{\text{farthings.}}{\text{farthings.}} \text{Therefore 2s. 7\frac{7}{2}d.} = \frac{127}{7584} \text{ of \$27\$ 18.}

Hence, to reduce one denominate number to the fraction of another, we deduce the following:-

# . The later of being the RULE.

Reduce both quantities to the lowest denomination contained in either.

Then place that quantity which is to be the fraction of the other as numerator and the remaining quantity as denominator.

EXAMPLE 1 .- Reduce 3 days 4 hours to the fraction of a week.

3 days 4 hours = 76 hours.

1 week = 168 hours.

And the required fraction is  $768 = \frac{19}{48}$  Ans.

Example 2.—What fraction is 3 lb. 4 oz. 2 dr. 2 scr. 7 gra. of 63 lb. 4 oz. 7 dr. Apothecaries' weight?

> 3 lb. 4 oz. 2 dr. 2 scr. 7 grs. = 19367 grs. 68 lb. 4 oz. 7 dr. == 365220 grs. And the fraction is 19307 Ans.

#### Exercise 54.

- 1. What fraction is 6 bush. 1 pk. 1 gal. 1 qt. 1 pt. of 50 bush.? or only and distance best of the continuence Ans. 4100.
- 2. What fraction is 35 per. 9 ft. 2 in. of a furlong?
- 3. What fraction is 7 h. 12 m. of a day?
- 4. What fraction is 2 sq. yds. 2 ft. 120 in. of 3 sq. per. 131 yds. 1 ft. 72 in.? Ans. 45.
- 5. What fraction is 7 oz. 7 dr. 2 scr. 14 grs. of 21 lbs. Apoth.?
- 6. Reduce 9 min. 48 sec. to the fraction of a day. Ans. 7100.
- 7. Reduce 16 bush. 1 pk. 1 pt. to the fraction of 69 bush.
- Ans. 1474. 8. Reduce 3 grs. 31 na. to the fraction of an ell Eng. Ans. 14.
- 9. What part of a lb. Troy is 13 dwt. 7 grs.? Ans. 2180.
- 10. What part of 54 cords of wood is 4800 cubic feet? Ans. 14.

# ADDITION OF VULGAR FRACTIONS.

40. Addition of fractions is the process of finding a single fraction which shall express the value of all the fractions added.


Addition may be illustrated as follows.



41. In order that fractions may be added they must have a common denominator

Thus 144 make neither 4 nor 4; but if we reduce them to equivalent frections having a common denominator, as and and we are enabled so add them and thus obtain for their sum 11.

Me so Pachiens; before and after they rective a dommon denegrinator, will be represented as follows:-



We have increased the number of the parts just as much as we have diminished their size

42. For the addition of fractions we have therefore the following:

Reduce compound and complex fractions to simple ones, and all to a common denominator. (Arts. 29, 31, and 33.)

Add all the numerators together, and beneath their sum place

the common denominator.

Reduce the resulting fraction, when an improper fraction, to a mixed number. (Art. 26.)

Now ... If mixed numbers occur as mg the addends, the integral portions are to be added separately and their sum added to the sum of the fractions.

ng a the

have

Ment bled

z de-

h as

v the

d all dace

n, to

gral sum

EXAMPLE 1.-Add together 1, 17, 10, 17 and 17.

Here, since the fractions have already a common denominator, we have mphy to will the numerators and place 11, the common denominator, simply to said the bene in their sum.

Thus 14 + 37 + 37 + 17 + 19 = 4+3+8+7+10 = 34 = 2 14 And

Example 2.—Add together \$, \$, \$, \$ and \$\frac{1}{4}.

These tractions reduced to their least common denominator by Art. 30, become 1, 84, 88, 48, 18.

= 138 = 13 = 3 Ans. And 35+184+88+88+88

Example 3.—Add together 1, 1, 2 and 1 of 1 of 17 of 19 of 51. 35 of 3 of 3 of 52 of 51 is equal to 7 (Art. 31).

The fractions to be sided are therefore \$ + 11-2-17: These reduced to a common denominator (Art. 30); become

Example 4.—Add together 91, 112, 161, 431, and 71

Here the last fraction is a complex fraction and is equal to \$ 91+112+162 +441+1=5+11+16+48+(1+2+2+14+14);

And 9+11+16+45=79. Also 1+2+3+3+4= 188+116+118+118+1165= 1000 = 3100.

Therefore the sum of the given quantities is 79+3,19 = 82,166.

Harris 5. Add together \$ 3 and 51

Here adding the three fractions together we obtain 1344 for their sum to which we add the integral number 5 and thus obtain the child sum 67 14.

# EXERCISE 55

Ans. 33 = 275. 1. Add together +1, +3 and 13.

2. Add together 13, 13, 13, 14, 14 and 15. Ans. 71 = 11 = 31.

3. Add together 43, 114, 163, 212 and 194

 $A_{70}$  71+12 = 784.

4. Add together 1625, 1117, 1845, 1713 and 11244. Ans. 17714.

5. Add together 4; 1; and The combined Ans. 671.

Ans. 643. 6. Add together 1, 1, 1, 1, 1, 1, 1, 1 and 1. Ans. 24 7. Add together \$, \$, and \$.

Ans. 3547 8. Add together i, i. ; i and i. 9. Add together in i. i. i. i and i.

Ans. 1

10. Add together 16,3, 47, 2117, 7 and 191. Ans. 10414. 11. Add together 174, 434, 1684, 207, and 506144. Ans. 94347. 12. Add together 63, 114, 26, 16, 7, 1, 5, and 1711. Ans. 53143.

13. Add together 1, 2, 7 and 681. Ans. 69444.

14. Add together 173, 3, 84 and 9111. Ans. 273284.

15. Add together 114, 223, 324 and 428. Ans. 13338. 16. Add together 1, 13, 48, 34, 16, 3, 1 and 4. Ans. 31.

17. Add together 7, 111, 18, 263 and 79 A

168

Ans. 14245.

18. Add together 3, 7 and 4 of 3 of 101. Ans. 11 74. - 207

19. Add together 7, 1 of 3, of 4 of 23, and 7.

20. Add together 34, 111 and 1438. Ans. 2913.

21. Add together 1 of 2, 3 of 4, 3 of 5, 3 of 1, and 41 of 1 of 1 of 1 of 1.

22. Add together 411, 1052, 3002, 2412 and 4721.

Ans. 116132.

152

value

of a

1. W

2. Ad

3. Ad

4. W

5. W

6. Ad 7. W

44.

ing th Weh

non de

they m case ala

No

Ex

23. Add together 92,5, 37,5 and 74. Ans. 137764.

24. Add together 211, 351, 23 and 3 of 7.

25. Add together 23 of 33, 11, 24 of 41 of 15, and 43 of 2 of 21 of 13. All now bushed Ans. 341138.

43. In order to add denominate fractions they must not only have a common denominator, but they must be fractions of the same

unit, i. e., must be of the same denomination.

Thus 23, 38. and 3d. cannot be added together, as the result would be

neither & of a pound; & of a shilling, nor & of a penny.

But if we reduce them all to the fraction of a pound, or all to the fraction of a shilling, or all to the fraction of a penny, it is obvious that we may then add the resulting fractions, having first reduced them to a common denominator.

Hence, for the addition of denominate fractions, we have the following :-

Reduce all the fractions to the same denomination (Arts. 36 and 37). Reduce the resulting fractions to a common denominator (Arts. 29 and 30). Add (as in Art. 42) and find the value of the resulting fraction (Art. 38).

IV.

EXAMPLE 1.—Add together \$ of a day and \$ of an hour.

 $\frac{1}{3}$ h.  $+\frac{1}{4}$ h.  $=\frac{1}{3}$  $\frac{1}{4}$ +  $\frac{1}{4}$  $\frac{1}{4}$ +  $\frac{1}{4}$ +

EXAMPLE 2.—Add together  $T_f$  of a pound,  $\frac{1}{2}$  of a shilling, and  $\frac{1}{2}$  of a penny.

 $\frac{7}{11}$  of a  $e = \frac{7}{11}$  of  $\frac{9}{11}$  of a penny = 152 A pence.  $\frac{9}{11}$  of a shilling =  $\frac{9}{11}$  of  $\frac{1}{11}$  of a penny =  $\frac{1}{11}$  pence.

 $152\frac{4}{17} + 4\frac{1}{2} + \frac{2}{7} = 156 + \frac{280 + 508 + 155}{385} = 157\frac{2}{3}\frac{2}{3}\frac{2}{3}$  pence = 13s.  $1\frac{2}{3}\frac{2}{3}\frac{2}{3}$ d.

Note.—In place of proceeding as above, we may find the value of each fraction separately (Art. 38) and add the results.

EXAMPLE 3.—Add together 2 of a bushel, 7 of a peck, and 10 of a gal.

f of a bushel = 3 pks. 0 gal. 1 qt. 1 pts. 0 of a peck = 1 gal. 3 qts. 1 pts. 1 of a gal. = 1 pts. 1 pts.

Sum=1 bush, 0 pks. 0 gal. 1 qt. 024 pts. Ans.

### EXERCISE 56.

1. What is the sum of Alb. Apothecaries' weight, \$ oz. Ar. and \$ scr.? Ans. 4 oz. 6 drs. 2 scrs. 1811 grs.

2. Add together # yd. 4 ell Eng. and 4 qr.

Ans. 3 grs. 3 z. 1132 in.

3. Add together 1 of a yard, 1 of a foot, and 4 of an in.

Ans. 7 inches.

4. What is the sum of 77 of a mile, 43 of a furlong, and 22 of a yard? Ans. 5 fur. 16 rds. 0 yds. 0 ft. 3 22 in.

5. What is the sum of 1 wk. 1 day, 1 h.?

Ans. 2 days 2 h. 12 m.

6. Add together £1, \$s., and \$\frac{1}{4}d\$.
7. What is the sum of \$\frac{1}{4}\$ of \$21s\$. \$\frac{1}{4}\$ of \$5s\$. \$\frac{1}{4}\$ of \$21s\$. \$\frac{1}{4}\$ of \$

#### SUBTRACTION OF VULGAR FRACTIONS.

44. Subtraction of vulgar fractions is the process of finding the difference between two fractions.

We have seen that before fractions can be added they must have a common denominator, and that when denominate fractions are to be added they must be also of the same denomination, and this is manifestly the case also in the subtraction of fractions.

Hence, for the subtraction of fractions, we have the following:—

RULE.

Reduce compound and complex fractions to simple ones and all to the while destruction, if not already such.

Reduce value of the resulting fractions to a common denominator.

Subtract the numerator of the subtracted from the numerator of the middlend, and belieath the difference write the common denominator.

Norm.—In the case of mixed numbers it frequently happens that the fractional part of the subtrained is greater than the fractional part of the minuend. When this occas, instead of reducing both quantities to improper fractions and then applying the rule, it is much better to borrow using from the integral part of the minuend and considering it as a fraction, having the common denominator, add it to the fractional part of the minuend. (See 3rd, 4th and 5th Examples below.)

EXAMPLE 1.-From ? take %.

10. 11.

13.

15.

16.

45

He No

time

If, in or Bu

and .

Ther

mult

inato

dedi

R

and 23 a Co

nomi

Here reducing \$ and 14 to a common denominator they become 151 and 14.

Example 2.—From  $\frac{1}{2}$  of  $\frac{1}{2}$  of  $\frac{1}{4}$  of

And 37 of 1 of 1 = 1. And 2 - 1 = 13 - 16 - 5. Ans.

Example 3. - From 1921 take 1618.

 $\frac{3}{10}$  and  $\frac{1}{10}$  reduced to a common denominator become  $\frac{3}{10}$ , and  $\frac{1}{10}$  and

Here, since we come subtract 195 from 32, we have to borrow 1 from the ting-rad part of the minuend and densidering it as 114 add it to 125.

We have reduce 188. to 1918 1 and then make the subtraction.

Example 4. From 29 7 take 164.

 $29_{11}^{4} - 164 = 29\frac{1}{4} - 1644 = 28 + 1\frac{1}{4} - 1644 = 28\frac{1}{4} - 1644 = 124$ = 124 Ans.

EXAMPLE . Wie m 1177 take 67 19

1173 6744 = 11 188 - 67798 = 116+1+38 - 67798 = 116998 - 67798 = 49478 Ahr.

EXAMPLE 6.—What is the difference between 1 of 5 of 5 of 23

of 2 of of 27 days 8 of a day 5 of 2 of an hour 12 hours 17 hours; and 2 of 3 of 5 hours 18 hours 17 hours

And 17 h.-136 h. = 175 -135 163 hours. Ans.

# Examples: 57.

1. From ! take 70.	· 1: ( .4.10)	a state to	Ans. 2
2. From 145 17 of	12 of 19 tal	83	And On

Ans. 952-1770. 3. From 98217 take 2918.

4. What is the difference between 69 7 and 18745? Ans. 501923:

5. What is the difference between 100; and 9; ? Ans. 9070

6. What is the difference between 64 and 1 of 94? Ang., 11. 7. From 611 for take 610 f \$ . 8. From \$ of 2 take \$ of \$ +++. Ans. A

9. From i of a lb. avoirdupois take i of a dram.

Ans. 10 oz. 97 drs. 10. What is the difference between 24 and 217 ? .dns. 2117.

11. What is the difference beween I of a mile, and Tr of a fur-Ans. 1 fur. 5 rd. 3 yds. 1 ft. 10 in.

12. Find the value of & of 115 - 1 of 281.

13. Find the value of 12,346 + 1 of 1 of 3 of 81 of

14. Find the value of 312+82-313-25+51+61-161. Ans. 23

15. From 1 of an acre take for sperch.

Ans. 1 rood 17 p. 22 yds. 2 ft. 108 in.

16. From 161 take 914; and from 16917; take 831%. Ans. 6-34 and 85-330.

# MULTIPLICATION OF YULGAR PHACTIONS.

45. Let it be required to multiply  $\frac{3}{4}$  by  $\frac{7}{4}$ . Here we are required to multiply  $\frac{3}{4}$  by  $\frac{7}{4}$ .

Here we are required to multiply to 15 that is by 1 of 7.

Now if we multiply 1 by 7 we shall have multiplied by a quantity 8 times too great, and the product will be 8 times too great,

If, therefore, we multiply in by 7 we shall have to divide the result by 8 in order to get the product of  $A_1 \times 7$ .

But (Art. 8) we multiply 1 by 7, when we multiply the numerator by 7. and we divide the result by 8 when we multiply the denominator by 8.

Therefore, 11 × 11×8 that is to multiply fractions together, we multiply the numerators together for a new numerator, and the denominators together for a new denominator.

Hence, for the multiplication of vulgar fractions we deduce the following:-

Reduce compound and complex fractions to simple ones (Arts. 31. and 33) and whole and mixed members to improper fractions (Arts. 23 and 25)

Cancel any factors that are common to a nun rator and a denominator of the resulting fractions (Art. 32).

and all

ET. IV.

re fol-

inator. perator n deno-

appens in the tead of applyategral ing the minu-

ome 119

91798

w 1 from t 10 7%.

- 1644

f 4 of 24

bours -

Multiply all the reduced numerators together for a new numerator, and all the reduced denominators together for a new denominator.

Reduce the result, if necessary, to a mixed number.

EXAMPLE 1 .- Multiply # by | 9.

Here we cancel the first denominator and reduce the second numerator to 3.

EXAMPLE 2.—Multiply together 71, 3, 31 and 11.

STATEMENT.

CANCELLED.

$$f_1 \times f \times f \times f = \frac{q}{11} \times \frac{q}{4} \times \frac{q}{4} \times \frac{q}{96} = \frac{1}{1} Ans.$$

Example 3.—Multiply together \$, \$\frac{1}{17}, 6\frac{2}{7}, 9\frac{2}{7}, and 63.

STATEMENT.

CANCELLED.

$$\frac{2}{4} \times \frac{3}{11} \times \frac{44}{7} \times \frac{48}{8} \times \frac{8}{2} \times \frac{8}{1} = \frac{2 \times 3 \times 4 \times 48}{1} = 1152 \text{ Ans.}$$

Example 4.—Multiply together  $\frac{1}{179}$ ,  $18_{1}^{7}$ ,  $9\frac{1}{5}$ ,  $\frac{1}{2}$  of  $\frac{3}{4}$  of

STATEMENT.

CANCELLED.

$$\frac{1}{179} \times \frac{205}{11} \times \frac{48}{5} \times \frac{21}{8} \times \frac{185}{8} = \frac{205 \times 3 \times 8 \times 3}{179} = \frac{5535}{179} = 30\frac{195}{179} \text{ Ans.}$$

EXAMPLE 5. Multiply together 7, 3,1, 41, 8, 61 and 518.

CANCELLED. 15.00 http://or off. dec. of

with the the water of the west ( Lett. 23)

$$\frac{x}{x} \times \frac{247}{81} \times \frac{x}{2} \times \frac{x}{5} \times \frac{43}{5} \times \frac{77}{15} = \frac{247 \times 43 \times 77}{81 \times 5 \times 15} = \frac{817817}{6075} = 1348792.$$

AR

2.

5.

7.

9. I

12. N

18. M

15. F

17. M

18. Fi

19. M

20. Fi

X

46.

Multi and div

Normhaving 1

EXAM

Exam:

物質消

### Exercise 58.

.IV.

mer-

7, and

	. What is the product of 17 × 4?	ans.	
	. What is the product of †×1?	das.	
	. What is the product of TXTE?	Ante-	
	Multiply together I, and A.	And Ann	
	. Multiply together 14, 15 and 34.	Ans.	
6.	. Multiply together & 82, 4 and 11.	Ans.	
7.	. Required the product of \$, 17, 17, 18	and f. Ans.	
8.	. Required the product of \$, \(\frac{1}{2}\), \$\(\frac{1}{2}\), 21	, and 5. Ans.	
9.	. Required the product of \$, \$, 11, 14	nd 209. Ans.	<b>)</b> }.
	Find the value of $61 \times 111 \times 167 \times 13$		
11.	. Find the value of $\dagger$ of $\dagger_1$ of $\dagger_6$ of 77	X of 1 of 91 X 6	13.
1-77	mortes of School of the College	m 1	1271.
	7, 7, 42	ulaman't motera	. 1 .
12.	Multiply together 8, 91, 4, 7,1, 27,	and 14. Ans. 7	T:
	. Multiply 1 of 8 by 7 of 19.	Ans. 10	05.
	. Multiply to of 7 by the of 873.	Ans. 40	
	. Find the value of 61×1×1×4.	Ane. 24	-10 - 4
	Find the value of 31×41×15.	de la la Ans. 26	
	Multiply tof 8t of to of 91 by 814)		
	151 of 1111.	Ans. 472	
	27 878 7 81		374.
18.	Pind the value of	and the same of th	-75
19.	Multiply \$877 by + of + of +7.	Ans. \$	
20	Find the value of 108 4 of 82 X 78 of	28 × 17 × 1 × 143 ×	100
40.	Find the value of $\frac{75\frac{1}{6}}{6\frac{1}{11}} \times \frac{\frac{3}{4} \text{ of } 8\frac{1}{4} \times \frac{1}{16} \text{ of } 6\frac{1}{8} \times \frac{1}{17} \text{ of } 6\frac{1}{8} \times \frac{1} \text{ of } 6\frac{1}{8} \times \frac{1}{17} \text{ of } 6\frac{1}{8} \times \frac{1} \text{ of } 6\frac$	24×15×7×144	121
6111	4 4	with my me that the state of the second	. 4 11 4
4.1	×51×9.	10.1 72 1 & Ans. 1'	1988.
	16.1	1 . 2 . 4 . 9 4 . 200 . 2 . 2	9 9 8 11 5

46. To multiply an integral denominate number by a fraction, we have the following:

### entil athailahalbat 1.85

Multiply the denominate number by the numerator of the fraction and divide the result by the denominator.

Note.—This is merely considering the denominate number as a fraction having 1 for its denominator (Art. 23), and applying the preceding rule.

Example 1.—How much is \$ of \$129:63.

3 of \$120-63\_\$120-63×4\_\$618-53\_\$57-613. Ane.

EXAMPLE 2.—How much is 77 of 1 of 10 lb. 6 oz. 4 dr. Avoir?

77 of 2 of 10 lb, 6 oz. 4 dr.—75 of 10 lb. 6 oz. 4 dr.—10 lb. 6 oz. 4 dr.×7

23

3 lbs. 4 oz. 1414 drams, Ans.

gi

ms

ber

ver

·T

the

2. D

7. D

.1

### Exercise 59.

- How much is 1% of 4 days 5 h.? Ans. 5 days 38 m. 20 sec.
   How much is 1% of £29? Ans. 5 days 38 m. 20 sec.
- 3. How much is 7 of 186 acres 3 roods? Ans, 145 acres 1 rood.
- 4. How much is 14 of 9 of 10 of 231 times 24 h. 30 m.?

Ans. 1 hour 38 min.

- 5. How much is 3 of 4 of 10 of 33 bush. 2 pk. 1. gal. 7 Ans. 2 bush. 2 pk. 0 gal. 3 qt. 117 pt.
- 47. From the principles already established, it is evident that

1st. When the multiplier is less than unity, the product is less than the multiplicand.

2nd. To multiply a fraction by a whole number, we may either multiply the numerator of the fraction or divide the denominator by that number. (Art. 8).

3rd. To multiply a whole number by any fraction having unity for its numerator, we simply divide the whole number by the denominator.

Thus, to multiply by 1, 1, 1, 1, 1, 2c., we divide by 2, 3, 4, 7, 11, &c.

4th. When multiplying by a mixed number of which the fractional part has unity for its numerator, it is better to multiply by the integral part of the multiplier first and then by the fractional part, afterwards adding the two partial products together.

### DIVISION OF VULGAR FRACTIONS.

The same of the same of the

48. Let it be required to divide by fr.

Here we are required to divide 3 by A that is, by A of 5.

Now if we divide 3 by 5, we use a divisor 11 times too great, and the quotient is 11 times less than the required quotient.

Therefore, to obtain the correct quotient of 9- 4, after dividing ; by 5, we shall have to multiply the result by 11.

But (Art. 8) we divide the fraction \$ by 5, when we multiply the denominator 7 by 5, and we multiply the result by 11 when we multiply the numerator 3 by 11.

Therefore = X 1 = dividend × divisor with its terms in-, at Till of the selection of the land of

Hence for the division of meetions we have the following:-

1

ct. IV.

20 sec. s. 67d. 1 rood.

8 min. 147 pt.

vident

product we may

ide the

le num-

of which is better first and the two

st, and the

the denomiply the nu-

ts terms in-

ne follow-

### 

Reduce compound and complex fractions to simple ones; whole and mixed numbers to improper fractions.

Invert the terms of the divisor and proceed as in multiplication.

In addition to the foregoing analysis, the following may be given as a proof of the truth of this rule.

 $\frac{3}{4} \cdot \frac{5}{11} = \frac{7}{11}$  because the dividend of any question in division may be made the numerator and the divisor the denominator of a fraction.

Now since we may multiply both terms of the fraction in the fraction by the number, we may multiply them by 11, i. e., the denominator with its terms inverted.

Therefore  $\frac{3}{11} = \frac{\frac{3}{7} \times \frac{11}{6}}{\frac{1}{7} \times \frac{11}{3}} = \frac{\frac{2}{7} \times \frac{11}{6}}{1}$  (because  $\frac{5}{17} \times \frac{11}{6} = 1$ )= $\frac{3}{7} \times \frac{11}{6}$ : whence the truth of the rule.

EXAMPLE 1.—Divide 13 by 11.

13 11 13 × 11 = 33 Ans.

EXAMPLE 2.—Divide \(\frac{1}{1-11}\) of \(\frac{1}{1-12}\) of \(\frac{1}{1-12}\) of \(\frac{1}{1-12}\) of \(\frac{1}{1-12}\) \(\frac{1}{1-12}\) of \(\frac{1}{1-12}\) \(\frac{1}{1-12}\) of \(\frac{1}{1-12}\) \(\frac{1}{1-12}\) of \(\frac{1}{1-12}\) of \(\frac{1}{1-12}\) \(\frac{1}{1-12}\) of \(\frac{1}{1-12}

Example 3.—Divide 3. of 1. of 3. ×11=6×1. 46 = 211 Ans.

Example 4.—Divide 3. of 1. of 3. ×31 by 1. of 54 ×43.

STATEMENT. THREE OF DIVISOR INVERTED.

### (i 1970 fire. 1. gii 113 a **Exercise 60.** file (i i a bankivii)

1. Divide a of a by a of 84.	J1160 T7E.
2. Divide 14 by 15 and divide the result by fr.	Ans. 8.
3. Divide 82 14 by 2844. Ale i so not estimental 2013	Ans. Byffs.
4. Divide 81 by 4 bet as as to so where as in	Ans. olyge
5. Divide 12 by + of 22 of 16 of 82 of 15.	Ans. 211.99
6. Divide 21 by (3 - x of 9.)	Ans. 780.
7. Divide 484 by \$ 4 100 6. James and a maine	Ans. 1995.
8. Divide 6 by 8 of 46 4 fraise see a 10 von	Ans. 6874.

	Divide 41 of 31 by 21 of		Ans. 118.
10.	Divide $\frac{73}{118}$ by $\frac{7}{41}$ .	कारी इता अहा त्याहरी, ताही सम्बद्धाः कृताः कृति । वहीति । वहीति	Ans. 6176.
11.	Divide & of 7 h by 1 of	177.	Ans, 149.
12.	Divide 117 of 19 of 2 of	18 by 8 of 18 of 2 of 5.	Ans. 3#7.
13.	Divide $\frac{1\frac{1}{4}}{4\frac{1}{4}}$ by $\frac{2\frac{1}{3}}{2\frac{1}{4}}$ .	ette sa isterio med ne i del	Ans. 3.
	many or letter and 44 to the loss		
			- 4
15.	Divide 14 of by 7 of 87	of 198 1 1 35 208 24 24	Ans. 12889.

16. Divide 151 of  $\frac{3}{4}$  of  $\frac{7}{3}$  of  $\frac{70}{3}$  by  $\frac{44}{7}$  of  $\frac{3}{42}$  of  $\frac{1}{31}$  of  $\frac{23}{4}$ .

Ans.  $28_{2455}^{1425}$ .

49. To divide an integral denominate number by a fraction:—

#### RULE

Multiply it by the denominator and divide the result by the numerator of the fraction.

Norm.—This is, in effect, merely considering the denominate number as a fraction having 1 for its denominator (Art. 23) and applying the foregoing rule.

EXAMPLE.—Divide 6 days 17 hours 11 minutes by 11.

6 days 17h. 11m.  $\div \frac{5}{11} = 6$  days 17h. 11m.  $\times \frac{11}{5} = \frac{6 \text{ days } 17\text{h. } 11\text{m.} \times 11}{5}$ = 14 days 18h. 36m. 12 sec. Ans.

### Exercise 61.

- 1. Divide £8 14s. 6\(\frac{1}{2}\)d. by \(\frac{1\frac{1}{11}}{1\frac{3}{2}}\). Ans. £8 8s. 5\(\frac{1}{2}\)d.
- 2. Divide 1m. 5 fur. 91 yds. 2 feet by 27 of 17.

Ans. 2 far. 124 yds. 2 ft.

3. Divide 3 acres, 3 roods and 3 perches by 3.

Ans. 6 acres 1 rood 5 per.

4. Divide £7 16s. 2d. by 4. Ans. £17 11s. 41d.

50. To reduce a fraction having a complex fraction in its numerator or denominator or both to a simple fraction we have simply to apply as often as necessary the rule given in Art. 33.

Norm.—Particular attention must be paid to the relative...
length and heaviness of the separating lines as they determine the various numerators and denominators.

1 ts. 6170.

T. 1V:

3<del>87</del>.

2889.

by a

2266

or as a egoing

a. × 11

. 2 ft.

5 per.

on in action rule

lative ...

EXAMPLE 1.—Simplify 
$$\frac{1}{\frac{3}{4}}$$

OPERATION.

 $\frac{3\frac{1}{4}}{\frac{3}{4}}$ 
 $\frac{3\frac{1}{4}}{\frac{3}{4}}$ 
 $\frac{3\frac{1}{4}}{\frac{3}{4}}$ 

EXAMPLE 2.—Simplify  $\frac{3\frac{1}{4}}{\frac{3}{4}}$ 
 $\frac{3\frac{1}{4}}{\frac{3}{4}}$ 

OPERATION.

 $\frac{3\frac{1}{4}}{\frac{3}{4}}$ 
 $\frac{3\frac{1}{4}}{\frac{3}{4}}$ 
 $\frac{3\frac{1}{4}}{\frac{3}{4}}$ 

OPERATION.

 $\frac{3\frac{1}{4}}{\frac{3}{4}}$ 
 $\frac{3\frac{1}{4}}{\frac{3}{4}}$ 

OPERATION.

 $\frac{3\frac{1}{4}}{\frac{3}{4}}$ 
 $\frac{3\frac{1}{4}}{\frac{3}{4}}$ 

OPERATION.

 $\frac{3\frac{1}{4}}{\frac{3}{4}}$ 
 $\frac{3\frac{1}{4}}{\frac{3}}$ 
 $\frac{3\frac{1}{4}}{\frac{3}{4}}$ 
 $\frac{3\frac{1}{4}}{\frac{3}}{\frac{3}}{\frac{3}{4}}$ 

13×13×17

M

		TIXERUISE O	4.	
	12½ 7		3 of 32	i agrice
1. Multiply	9	by .	91	.Ans. 2117.
- (5, -,-)	3 3.	s. *, , , , , , , , , , , , , , , , , , ,	31	F
, y is to linear the to be	41	95,		
¥ .	7			e
2. Divide	6½ 9½	b <b>y</b>	7	Ans. 1237.
	3	d	°, ,	(4 4 1 2)
•	121		2 <del>1</del> 5	
3. Divide	32	by:	43	Ans. 3\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\
2	51	N	163	

51. From what has already been said, the truth of the following principles is evident.

1st. When the dividend is equal to the divisor, the quotient will be 1.

2nd. When the dividend is greater than the divisor, the quotient will be greater than 1.

3rd. When the dividend is less than the divisor, the quotient will be less than 1.

of the

or, the

sor, the

or, the

4th. The quotient will be as many times greater or less than 1 as the dividend is greater or less than the divisor.

5th. To divide a fraction by a whole number, we may either divide the numerator or multiply the denominator by that number.

6th. To divide a whole number by a fraction having 1 for its numerator, we simply multiply the whole number by the denominator of the fraction.

Thus, to divide by  $\frac{1}{2}$ ,  $\frac{1}{3}$ ,  $\frac{1}{5}$ ,  $\frac{1}{7}$ , &c., we multiply by 2, 3, 5, 7, &c.

### QUESTIONS TO BE ANSWERED BY THE PUPIL.

NOTE.—The numerals after the Questions refer to the numbered articles of the Section.

- 1. What is a fraction? (1 and 3)
  2. What does every fraction indicate? (3)
  3. What is the denominator of a fraction, and why is it called so? (4)
  4. What is the numerator of a fraction, and why is it so called? (4)
  5. What are the terms of a fraction? (6)
  6. How is the value of a fraction obtained? (6)
  7. When is the fraction equal to 1, and when greater or less than 1.9.

- 7. When is the fraction equal to 1, and when greater or less than 1 ? (7) 8. What effect has multiplying the numerator of a fraction by any number ? (8)
- 9. How does multiplying the denominator of a fraction by any number affect the value of the fraction? (8)

  10. How does multiplying both terms of a fraction by the same number affect its value?(8)

- 11. How does dividing the numerator by any number affect the value of the fraction ?(8)

  12. How does dividing the denominator by any number affect the value of the fraction? (8)
- 18. How does dividing both numerator and denominator by the same num-
- ber affect the value? (8)

  14. Into what classes are fractions divided? (9)

  15. What is the distinction between vulgar and decimal fractions? (10 and 11.)
- 16. What is the meaning of the word "vulgar" as applied to fractions? (11)
  17. Enumer, to the six different kinds of vulgar fractions. (12)
  18. What is a proper fraction? (13)
  19. What is an improper fraction? (15)
  20. What is a mixed number? (16)

- 21: To what must an improper fraction always be equal? (17)
  22. What is a simple fraction? (18)
  23: What is a compound fraction? (19)
  24: What is a complex fraction? (21)
- 25. How may we convert an integer into a fraction? (23)
  26. How may we reduce a whole number to a fraction having a given denominator? (24)
  27. How is a mixed number reduced to an improper fraction? (23)
- 28. How is an improper fraction reduced to a mixed number?
- 29. How is a fraction reduced to its lowest terms? (27 and 28)
  30. How are fractions reduced to a common denominator? (29)
  31. How are fractions reduced to their least common denominator? (30)
  32. How is a compound fraction reduced to a simple one? (51)

33. What is meant by cancelling? (32)

34. Upon what principle may we cancel factors common to numerator and denominator? (32 and 3)

35. How do we reduce complex fractions to simple ones? (33)

36. What is a denominate fraction? (34)

37. In what does reduction of denominate fractions consist? (35)

38. How do we reduce a denominate fraction from a lower to a higher denomination? (36)
39. How do we reduce a denominate fraction from a higher to a lower denomination? (37)
40. How do we find the value of a denominate fraction? (38)
41. How do we reduce one denominate number to the fraction of another?

(39)

What is addition of fractions? (40)

48. What kind of fractions only can be added? (41)
44. What is the rule for addition of fractions? (42)
45. What is the rule for the addition of denominate fractions? (42, note)
46. What is the rule for the addition of denominate fractions? (43)

What is the rule for the subtraction of fractions? (44)
What is the rule for multiplication of fractions? (45)
Give a proof of the truth of this rule. (45)

66. Give a proof of the truth of this rule. (45)
50. How do we multiply an integral denominate number by a fraction? (48)
51. How may we multiply a fraction by a whole number? (47)
52. How do we multiply a whole number by a fraction having 1 for numerator? (47)
53. How do we multiply a whole number by a mixed number, the fractional part of which has 1 for numerator? (47)
54. What is the rule for division of fractions? (48)
55. Give a proof of the truth of this rule, (48)
56. How do we divide an integral denominate number by a fraction? (49)
57. How do we divide a fraction by a whole number? (51)
58. How do we divide a whole number by a fraction having 1 for its numerator? (51)

numerator? (51)

#### EXERCISE 63.

### MISCELLANEOUS EXERCISE ON VULGAR FRACTIONS.

- 1. The Ottawa River is 800 miles long; the Gatineau 420 miles, the Chaudière 100 miles, the Richelieu 160 miles, and the Niagara 35 miles. The entire length of the St. Lawrence. from the upper end of Lake Superior to the Sea is 2000 miles. How will the lengths of these different rivers be expressed as fractions of that of the St. Lawrence?
- The population of Goderich is ? of that of Peterborough. the population of Peterborough is 11 of that of Breckville. the population of Brockville is 13 of that of Prescott, the population of Prescott is 1 of that of Ottawa City, the population of Cttawa City is 21 of that of Port Hope, and the population of Port Hope is 4 of that of Toronto. What fraction is the population of Goderich of that of Toronto?
- 3. What will 67 pounds of tea cost, at 652 cents per lb.?
- 4. Suppose I have ? of a ship, and that I buy it more; what is my entire share?

5. A boy divided his marbles in the following manner; he gave to A = 0 of them, to B = 0, to C = 0, and to D = 0, keeping the rest to himself; how many did he give away, and how many did he keep?

6. Find the value of  $\frac{5\frac{2}{3}-2\frac{1}{5}}{3\frac{2}{5}+\frac{9}{5}}$  of  $\frac{4\frac{1}{5}+5\frac{1}{5}}{4\frac{1}{5}}$  of  $\frac{2\frac{2}{5}+1\frac{3}{5}}{7+\frac{9}{5}-2\frac{1}{5}}$ .

7. What cost 1670 72 pounds of coffee at 121 cents per pound?

8. A tree whose length was 136 feet, was broken into two pieces by falling; } of the length of the longer piece equalled 2 of the length of the shorter. What was the

length of the two pieces respectively?

9. A farmer bought at one time 971 acres of land, for 1000 dollars; at another, 1273 acres, for 13751 dollars; at another, 500% acres for 68314 dollars; and at another, 3334 acres for 401378 dollars. What was the whole quantity of land that he purchased, and the sum that he paid for it?

10. Find the value of  $(126 - 81 - 1_{10} + 1_{6}) \times 41 \times (74 - 61)$ ,

and also of  $(\frac{3}{3} \div 1\frac{4}{7}) - (\frac{4}{3} \div 3\frac{4}{11})$ .

11. What is the value of 19% barrels of flour, at \$6% a barrel?

12. What is the value of 376% acres of land, at \$75% per acre?

13. Bought at one time 1473 bushels of coal, and at another time 320% bushels. Having consumed 156% bushels, I desire to know what quantity of the coal purchased is still on hand?

14. Divide  $\frac{7(1\frac{1}{3} \text{ of } \frac{3}{4})}{\frac{1}{6}(\frac{3}{3\frac{1}{4}} \text{ of } 7)}$  by  $7\frac{7}{8}$ ; and find the value of  $\frac{1}{\frac{1}{24} + \frac{1}{34} + \frac{1}{44}}$ 

15. If 174 bushels of wheat sow 74 acres how many bushels

will it require to sow one acre?

16. Multiply the sum of 33, 41, and 45, by the difference of 75 and 5%; and divide the product by the sum of 941 and

17. Divide 2 by the sum of 23, 4, and 4; add 13-7 to the quotient; and multiply the result by the difference of 51 and

18. Find the value of  $(\frac{1}{2}+\frac{1}{2})\times(\frac{1}{2}+\frac{1}{2})\times(\frac{1}{2}-\frac{1}{2})\times(\frac{1}{2}-\frac{1}{2})$ ;

and also of (12+21)+(51+31).

19. A person dies worth \$40000, and leaves 1 of his property to his wife, i to his son, and the rest to his daughter. The wife at her death leaves ; of her legacy to the son, and the rest to her daughter; but the son adds his fortune to his sister's and gives her 1 of the whole. How much will the sister gain by this; and what fraction will her gain be of the whole?

THE THE TREE TREET OFFICE

(46)

te)

er?

nd .

ımeional

(49)or its

ONS.

niles. l the ence, 2000 rs be

bugh, ville, , the

, the . and

onto. at of

what

### DECIMALS AND DECIMAL FRACTIONS.

52. A decimal fraction is a fraction having unity with one or more 0s to the right of it for denominator:

Thus 1000, 100, for 100000, &c. are decimal fractions.

58. A decimal fraction is reduced to its corresponding decimal by dividing the numerator by the denominator; but since (Art. 52) this denominator is unity followed by one or more 0s, we divide the numerator by the denominator when we move the decimal point as many places to the left in the numerator, as there are 0s in the denominator.

EXAMPLE 1. Reduce Total to a decimal.

Ans. .743.

2. Reduce Thoughton to a decimal.

Ans. :00092376.

#### EXERCISE 64.

1. Reduce 1000, 100000 and 10 to decimals.

Ans.:567,:00098 and .7.

2. Reduce 10000000 and 10000000 to decimals.

Ans. :0000023 and :0000176.

3. Reduce To 378 84800 to a decimal.

Ans. .000278643.

- 54. It is as inaccurate to confound a decimal fraction with its corresponding decimal as to confound a vulgar fraction with its quotient: Thus the value of  $\frac{1}{2}$  is .75, so also the value of  $\frac{1}{100}$  is .75 but .75 and  $\frac{1}{100}$  are no more identical than are  $\frac{1}{2}$  and .75.
- 55. To reduce a decimal to its corresponding decimal

#### RULE.

Consider the significant part of the decimal as numerator and beneath it write for denominator 1 followed by as many 0s as there are places in the decimal.

### Exercise 65.

1. Reduce 73, 092 and 0003 to decimal fractions.

Ans. 736, 1860, and 10800.

2. Reduce :137 and :000006943 to decimal fractions.

Ans. 1000, and 10088680000.

3. Reduce 13578967 and 023004003 to decimal fractions.

Ans. 135778967σ, and 1378984883σ.

IV.

ith

ing

or;

by

mi-

to

or.

76.

76.

43.

its

ion

the

ical

mal

and

here

30.

.000

<u>β3</u>π•

56. Decimal fractions follow exactly the same rules as vulgar fractions.—It is, however, generally more convenient to obtain their quotients, and then perform on them the required processes of addition, &c., by the methods already described (Sect. II).

To reduce a vulgar fraction to a decimal or to a decimal fraction :--

B'ILE.

Divide the numerator by the denominator and the quotient will be the required "decimal"; the latter may be changed to its corresponding decimal fraction by (Art. 55).

This is merely actually performing the division which the fraction indi-

cates.

EXAMPLE 1. Reduce 3 to a decimal and also to a decimal fraction. 8)7. 5" 's \_ 'P" . 'ja 3not \_\_

.875 Ans. = 1000 Ans.

2. Reduce 16 to a decimal.

·5625 Ans.

### EXERCISE 66.

- 1. Reduce 1 and 3 to decimals.
- Ans. 5 and 375.
- 2. Reduce 26 and 4 to decimal fractions. Ans. 76 and 1866.
- 3. Reduce 78, 171, and 15 to decimals.

Ans. 9733 +, 4.666 + and .44117 +.\*

4. Reduce 9, fe, and 5 to decimals.

Ans. . 857142 +, .4166 + and .44444 +.

5. Reduce 1 and 1 to decimals.

Ans. 15178571428 + and .554012 +.

57. Let it be required to reduce £3 7s. 63d. to the decimal of a pound.

OPERATION. id=75d hence 6id = 675d. If now we divide this by 12 we shall have its value as the coimal of a shilling.
6id=575d=5625s, hence 7s 6id=7 5625s.

Next if we divide this by 20 we shall have its value as a desimal of a pound.

7s. 61d=7\*5625s=£\*378125. Therefore £3 7s 61d=£\*378125.

Hence to reduce a denominate number of different denominations to an equivalent decimal of a given denomination we deduce the following:

<sup>\*</sup>The sign + written after these answers simply indicates that there is still a remainder consequently that the division may be carried on

10.

11.

de

An

tor fol

low

and

nat

nat

Divide the lowest denomination named by that number which makes one of the next higher denomination.

Annex this quotient to the number of the next higher denomination given and divide as before.

Proceed thus through all the denominations to the one required, and the last result will be the one sought.

EXAMPLE 1. Reduce 3 days, 12 hours, 3 minutes, 30 seconds, to the decimal of a week.

60)30=sec.=30 sec.

60)3'5=decimal of a minute=3 min. 30 sec.

24)12'0588 decimal of an hour=12 h. 3 m. 30 sec.

7)3'5024505=decimal of a day=3 days 12h. 3m. 30 sec.

Ane. '5003472=decimal of a week=3 days 12h. 3m. 80 sec.

EXAMPLE 2. Reduce 187 lb. 13 oz. 11 drams to the decimal of a ton.

OPERATION.

60)11' drains.

16)18.6875 ounces.

3000)187 85546875 lbs.

Here we divide the 11 drams by 16 operation.

In the we divide the 11 drams by 16 and thus obtain '8875 to which we prefix the given 13 oz. Next we divide this by 16 and obtain '85548875 to which we bring down the 187 lb. and divide the result by 2000, the number of lbs. in a ton.

Norm.—To divide by 2000 remove the decimal point three places to the loss and divide by 2; similarly to divide by 60, 20, &c., remove the decimal point one place to the left and divide by 6, 2, &c.

### Exercise 67.

- 1. Reduce 3 yds 2 ft. 1 in. to the decimal of a furlong. Ans. 01679+.
- 2. Reduce 3 dwt. 17 grs. Troy, to the decimal of a pound. Ans. 01545138+.
- 3. Reduce 2 ser. 7 grs. to the decimal of a pound, Apoth. Ans. .0081597+.
- 4. Reduce 5 fur. 35 per. 2 yd. 2 ft. 9 in. to the decimal of a mile. Ans. . 73603+.
- 5. Reduce 8 qr. 2 na. to the decimal of a yard.

Ans. . 875.

6. Reduce 5s. to the decimal of 13s. 4d.

Ans. . 375.\*

Reduce 5s. first to the fraction of 18s. 4d. and then reduce the regulting fraction to a decimal.

Thus 5s. reduced to the fraction of 18s. 4d.  $=\frac{60}{150}$   $=\frac{3}{3}$  = 375.

hich

IV.

omired.

, to

imal by 16

pre-ivide 5 to mber

e the

19+. 8+.

74. nile. 3十.

875. B75.\*

alting

7. Reduce 12 h. 55 min. 21 sec. to the decimal of a day. Ans. . 5384375.

8. Reduce 4 of 1 of 61d. to the decimal of £1.

Ans. .012053+. 9. Reduce 1 of 1 of a mile to the decimal of 31 inches.

Ans. 3620.571428-

10. Reduce 1 of 3 of 31 lb. Avoir. to the decimal of 2 of an oz. Ans. 9.2444+.

11. Reduce 3 g k. and 1 g 1 pt. to the decimal of a bushel. Ans. . 921876

58. Let it be denomination of

OPERATION.

to and the value in terms of a lower yard.

7825 41700

EXPLANATION.—Since there are 8 feet in a yard, it is evident that any decimal of a yard is three times as great a decimal of a foot. Hence to reduce the decimal of a yard to a decimal of a foot we multiply it by 8. This gives us two feet and '8475 of a foot. Similarly multiplying the decimal of a foot by 12 reduces it to an equivalent decimal of an inch. We thus find '3475 of a foot equal to 4 inches and '17 of an inch. Again, multiplyinches and '17 of an inche. Again, multiplying this last by 12 reduces it to the decimal
of a line, and we thus find the whole quantity

Ans. 2 ft. 4 in. 2 04 lines. 7825 of a yard equal to 2 ft. 4 in. 2 04 lines.

Norz.—In these multiplications we only multiply the number to the

right of the separating point.

Hence, to find the value of a denominate number in terios of integers of a lower denomination we have the following:-

Multiply the given decimal by the number of units of the next lower denomination that make one of the given denomination.

Point off as many decimal places as there were in the multiplier, and the integral portion, if any, will be units of that lower denomination; the decimal part may be reduced to a still lower denomination, and so on.

EXAMPLE 1.—Find the value of £.97875.


OPERATION. 97875 20 19°575006.


18 6:90000d. 2 10 7 6 Sar.

Ans. 19s. 62d. + 3 of a farthing. 2 "32" while to use adoptional some and are


Commence of the second

8'0000of.






# IMAGE EVALUATION TEST TARGET (MT-3)



Photographic Sciences Corporation

23 WEST MAIN STREET WEBSTER, N.Y. 14580 (716) 872-4503



-4.4.2.1. C.S.

đị

de

w

be

Example 2.—Find the value of 7863625 of a pound Apothecarios weight.

OPPEATION. Andrew of a standard standard and a standard and the by a to lace or

Ans. 9 oz. 3 dr. 1 scr. 9.448 grains.

design of the street of the of the Butter a six I sale has been a sometimes the

1'4734000 per. is det. Love in du requires to that the galed of transfer a forten

### EXERCISE 68.

1. Find the value of 0.3945 of a day.

Ans. 9 hours 28 min. 4.8 sec.

2. Find the value of 0 3965 of a mile.

THE THE PARTY OF THE PARTY OF THE PARTY OF THE

. devous horizon of 1835 of a rard.

WASHINGTON BOTTON

dis. 3 für. 6 per. 4 yds. 2 ft. 6 24 in.

4. Find the value of 22 75 of £2 2s. 6d. Ans. £48 6s. 101d. 6. Find the value of 11 17825 of 7 bush. 1 pk. 1 gal. 1 qt. Ans. 82 bush. 8 pks. 0 gal. 1 qt. 0 4805 pt.

6. Find the value of :2057 of a lb. Troy.

Ans. 2 os. 9 dwt. 8-832 grains. 7. Find the value of 176 of 1 for. 36 per. 2 yds. 5 inc.

Ans. 13 per. 2 yds. 1 ft. 4 in. 8. Find the value of .625 of a league. Ans. 1 mile 7 fur.

9. What is the value of .015625 of a bushel? Ans. I pint.

10. What is the value of .9378 of an acre?

to the 3 roods 80 per 1 yd. 4 ft. 9 % inches.

11. Find the value of 2775 of lag. yd. 3 ft. 72 in. Male to at a special and a second and and Ans. 3 ag. ft. 671 in.

### it want to not hat you to the it is CIRCULATING OR REPEATING DECIMALS.

59. Let it be required to reduce and f to decimals

OPERATION.

555555, &c. 7)6

857149857148857142, &c.

If the given quantity be expressed in more than one denomination it should be reduced to one before applying the rule. Thus in this example 7 bush. 1 pk. 1 gal. 1 qt. = 207 qts. and 11 17825 × 237 = 2049 24526 qts. = 82 bush. 3 pks. 0 gal. 1 qt. 0'4905 pints.

In these and many other eases the division does not terminate, and the value of the fraction can only be approximately expressed. In the former of the above examples the figure 5 is constantly repeated, and in the latter the series of figures 857142.

- 60. Decimals which do not terminate, i. c., which consist of the same digit or set of digits constantly repeated, are called Repeating or Circulating Decimals.
- 61. The digit or set of digits, which repeats, is called a repetend, period or circle.

Nors.—The terms period and circle are commonly used only when the repetend contains two or more digits.

62. A Single Repetend is one in which only a single digit repeats,

Thus 3338 &c. ; 7777 &c. ; 38888 &c. arc single repetends.

63. A Single Repetend is expressed by writing the digit that repeats with a dot over it,

Thus, '333 &c. is written '3; '777 &c. is written '7.

64. A Circulating Decimal or Compound Repetend is one in which more than one digit repeats,

Thus, 34547347 &c.; 202020 &c.; 123412341234, &c., are Circulating Decimals or Compound Repetends.

- 65. A Circulating Decimal is expressed by writing the recurring period once with a dot over its first and last digits.

  Thus, 347347 &c. is written 347; 2000 &c. 20; 1254434 &c. is written 1824.
- 66. A Pure Repetend or Circulating Decimal is one in which the repetend commences immediately after the decimal point.
- 67. A Mixed Repetend or Circulating Decimal is one which contains one or more ciphers or significant figures between the repetend and the decimal point.

Thus, i, i, are Pure Repetends.

'78917, '0578, '002 are Mixed Repetends." 23 Village

'72, '045, '91876 are Pure Cinculating Decimals.

1378, 073200, 0717800 are Mixed Circulating Decimals

tion it

othe-

24 in.

101d.

5.pt.

raine.

in.

pint.

68. Similar Repetends are those which commence at the same number of places from the decimal point,

Thus, 71846, 912786 and 00071846 are Similar Repetends.

69. Dissimilar Repetends are those which commence at a different number of places from the decimal point,

Thus, 7848, 928687 and 9184878 are Dissimilar Repetends.

70. Coterminous Repetends are those which terminate at the same number of places from the decimal point,

Thus. 7457, '6348 and '1347 are Coterminous Repetends.

71. Similar and Coterminous Repetends are those which both commence and end at the same distance from the decimal point,

Thus, 734207, 16-471312, 198-161341 are Similar & Coterminous Repetends.

73. In reducing a fraction to a decimal we place a point after the numerator, and annex to it until it is exactly divisible by the denominator. But since the point does not affect the division, merely determining the place of the point in the resulting quotient, it is manifest that we may leave it altogether out of consideration, so that annexing to to the numerator becomes in effect multiplying it by such a power of 10 as will make it contain the denominator. Now if the fraction, before proceeding to the division, be reduced to its lowest terms, the denominator can have no factor in common with the numerator; and if the denominator be exactly contained in the numerator with the 0s annexed, it can only be from its being contained in that power of 10 by which the original numerator was multiplied. But since 10 contains only the factors 2 and 5, any power of 10 can contain only the factors 2 and 5; and hence, in order that the denominator may be exactly contained in numerator with 0s annexed, it must contain only the factors 2 and 5.

Hence, when a vulgar fraction is reduced to its lowest terms, if the denominator contain no factors other than 2 and 5, the corresponding decimal will be finite; but if the denominator contain any other factor than 2 and 5, as 3, 7, 11, &c., the corresponding decimal will be infinite, i. o., will be a repetend.

EXAMPLE.—Can 76, 16, 18 and 176 be exactly expressed as decimals?

16, the denominator of the first,  $=2\times2\times2\times2$ , (i. e. contains no prime factor other than 2 or 5) therefore it can be exactly expressed by a decimal.

25=5 × 5 (i. e. no prime factor other than 2 or 5) therefore

 $\frac{11}{12}$  can be exactly expressed by a decimal.  $12=2\times2\times3$  (i. e. does contain a factor other than 2 or 5)

therefore  $\frac{1}{12}$  cannot be exactly decimated.

125 = 5 × 5 × 5 (i.e. no factor other than 2 or 5) therefore can be exactly decimated.

nce at E TON THE TOTAL

BOT. VI.

TOWNSON'S ence at or mailten

minate the sta

which m the

petends. e numeminator. ning the we may vill mak ng to the ator was

lowest than 2 if the , as 3, e, i. e.,

Tringe !! ssed as ontains

exactly erefore

2 or 5)

erefore

### EXERCISE 69.

Of the following fractions, which can and which cannot be exactly decimated, i. e., reduced to equivalent decimals?

- 1. 7, 17, 13, 1817, and 173.
  - 2. 178, 4, 25, 800, 111
  - 3. 11, 17, 18, 3, and 1980.
- 78. We may determine the number of places in the decimal or finite part of the decimal corresponding to a vulgar fraction by the following:

Reduce the fraction to its lowest terms, and decompose the denominator into its prime factors.

If the denominator contains no factors other than 2 or 5, or

powers of 2 or 5, the whole decimal is finite.

If the denominator does not contain 2 or 5 as factor, the decimal contains no finite part.

The highest exponent of 2 or 5 will indicate the number of decimal places in the finite part of the corresponding decimal.

EXAMPLE 1.—How many decimal places will be required to express 1847 7 Min a pade par season of an a

Here,  $$195 = 5 \times 5 \times 5 \times 5 \times 5 = 5^5$ . Therefore the equivalent decimal will

EXAMPLE 2.—How many decimal places will be required to express 1400 ? Is casafferst sinder ovier post eff on a

Here,  $1600 = 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 5 \times 5 = 26 \times 5^3$ . Hence 6 is the highest exponent, and the number of decimal places will therefore be 6.

### EXERCISE 70.

- 1. How many decimal places will be required to express the following fractions, viz :- 16, 20, 1000 and 1016 to 1000 advantaged traditions to the Ans. 4, 3, 6 and 10.
  - 2. How many places will there be in the finite part of the decimals corresponding to 78, 196, 18750 and 1747 13 30 9 9 7 a dne 5, 7, 4 and 11.
  - 74. In decimating vulgar fractions where many places are required in the decimal, the method of continually dividing becomes very tedious. In such cases we may sometimes shorten the work as follows:—

EXAMPLE.—What decimal is equivalent to the vulgar fraction 22.3

"ora of down beauton and

thought consenouted to b

i alcuirolo inoloxiano co

TIACOST IA

Resident and the

OPERATION. 29)1-00(0-08448

	7	1	210	,
	9		4	
	1	÷	-	Harry .
	1	40	3	10
	-	24		
Mahmu,		91	t	Ind
0	3 7	立意	8	

= 0.03448 . Therefore = 0.27586 and substituting this

value for  $f_0$  we get:—  $f_0 = 0.0344827586 f_0$ . Hence  $f_0 = 0.2068965517 f_0$  and substituting this for  $f_0$  we get:—  $f_0 = 0.03448275862068965517 f_0$ . Hence  $f_0 = 0.241379810844$ -

8275862022 and substituting this value for we get :-= 0.0344827586206896591724137931. Ans.

75. The number of places in a period cannot exceed the units in the denominator minus one.

This is manifest from the fact that all the remainders that occur must be denominated, and their number earned in greater than the denominator, minus one; because we carry on the division by affixing a and it follows that whenever we obtain a remainder like one that has averday occurred, the digits of the decimal will begin to repeat.

Thus 4 = 0 657142; where the small figures above the line represent the successive remainders, none of which, of course, can be as great as 7, the divisor,—the next remainder after the 6 would be 4; and consequently the digits would commence to repeat.

76. These repetends that have as many places, minus one, as there are units in the denominators of their equivalent vulgar fractions are sometimes called perfect repetends.

The following are the only fractions having a denominator less than 100 that give perfect repetends when decimated :-

1, 17, 19, 95, 99, 47, 89, 81 and 97.

77. To reduce a pure repetend to an equivalent vulgar Toy aomeser!

that which the work study

Put the period for numerator, and as many nines as there are places in the period for denominator.

ing this

substi-

810844get:--

exceed

cour must then than a flixing that has

line relecourse, der after ald com-

minus equivapetends.

minator :--

vulgar

here are

Example - What vulgar fractions are equivalent to :7, :93,

Ans. 7 = 1; 93 = 11 = 11; 704 = 111; 007043 = 11115.

Reason 1 = 1 therefore 1, 1, 1, 4c., = 2 3, 4, 4c., benefit 2, 3, 4c., =

Similarly 1 = 01, therefore 2 = 07; 22 = 15; 12 = 75; &c.

Hence '01; = 3; '07 = 3; '38 = 1; '17 = 1; '40. 3001 and

no also ga - dei : ag - dei ; ag - dei ; ag - dei ; ag c

Hence '001 = 113; 245 = 343; &c., whence the reason of the rule is ordent.

### Exercise 71.

- 1. Reduce 8, 05, 342, 7004 and 002003 to equivalent vulgar fractions.
- 2. Reduce 19, 1067, 11115 and '704103 to equivalent vulgar fractions.

  Ans. 19, 1081 = 37, 11115 = 1744 and 100100 = 200181.
- 3. Reduce 102, '0013, '00007108, '01020804 and '907084031 to equivalent vulgar fractions.

  Ans. 114, vista vulgar, fractions.

78. To reduce a mixed repetend to an equivalent valgar

#### RULE

Subtract the finite part from the whole and set down the difference for the numerator.

For denominator put as many 9s as there are places in the infinite part followed by as many 0s as there are places in the finite part.

EXAMPLE.—Reduce ·73, ·1234 and ·7132092 to their equivalent vulgar fractions.

73- 7 = 60 = numerator of first fraction.

715206-713-7131070 = "third."

\$0 = let Dénominator, since the repetend contains one place in the finite, and one place in the infinite part.

\$0002-2506 Benduitinton/concert the repetend contains two places in the Anisomer and two his he infinite part.

in

of

88

ha

of

pla re

las

m

th pli

contains four places and the finite part free places.

Hence, '73=16=11, '1234=1888=111, and '7132092=1111878.

REASON.—Let it be required to reduce 978734 to an equiva-

Let 
$$x = .978734$$
 (I)

Then 
$$100x = 97.8734$$
 (II)

And 1000000x = 978734.8734 (III); subtracting (II) from (III) gives 999900x = 978734 - 97.

5 15 LEAR TOTAL 978784-197 THE - = Whole repetend minus the finite part

for a numerator; and as many 9s as there are places in infinite part, followed by as many 0s as there are places in finite part for denominator.

The rule may also be explained as follows:-

Taking the same example 978734 and multiplying it by 100, we get

 $978736 \times 100 = 978734 = 97 + 8734 = 97 + 8734 = 97 + 8734 = 100 times too great. There-$ 

fore 978734 100 + 2578 00 and to add these fractions we must reduce them to a com

97×10000-97 + 6784 = 900000 97×(10000-1)+ 8784

= Whole repetend whose finite part for numerator; and as any to as there are places in infinite part, followed by as many to as sere are places in finite part for denominator.

Whence the truth for the rule is manifest.

# Exercise 72. Therefore to the work a par-

- 1. Reduce :8325, :147658, and :4320075 to their equivalent vulgar fractions. Ans. \$368 = 2121, 147611 and 4318868 = 1422481.
- 2. Reduce 875 4965 and 301 82756 to their equivalent mixed numbers. Ans. 8751278 and 3011488.
- duce '083, '0714285, and '123456' to their equivalent vulgar fractions.

cimal,

B, IV.

H858.

quiva-

(III)

te part infinite te part

we get

Therereduce

9784 999900 and as

ivalent

mixed

ivalent

ivalent

79. There are several properties belonging to repetends which it is necessary to remember. They are as follows:

1st. Any finite decimal may be regarded as a repetend if we make the 0s recur:

Thus, '27 = '270 = '2700 = '27000 = '2700000, &c.

2nd. A repetend having any number of places may be reduced to one having twice, thrice, &c., that number of places.

Thus a repetend having 2 places may be reduced to one having 4, 6, 8, 10, 12, &c., places.

For example, 372 = 37272 = 3727272, &c.

·232134 = ·2321342134 = ·23213421348134, &c.

3rd. Two or more repetends, having a different number of places in each, may be reduced to others having the same number of places in each, by the following:—

BITLE.

Take the numbers indicating how many places there are in each repetend, and find their least common multiple. Reduce each repetend to that number of places.

Thus, let it be required to reduce :147, :932, :8417, to repetends having the same number of places.

Here the numbers of places are 1, 2, and 3, and the least common multiple of 1, 2 and 3 is 6, and hence each new repetend must have 6 places.

Therefore 147 = 14777777, 932 = 9323232, and 8417 = 8417417.

4th. Any repetend may be transformed into another having a finite part and an influe part containing as many places as the original repetend; and hence any two or more repetends may be made similar.

Thus, 4123 = 41231 = 412312, &c.

7.654321 = 7.6543216 = 7.65432165, &c. 7.654321

5th. Having made two or more repetends similar by the last article, they may be made coterminous by the preceding one, and hence two or more repetends may always be made similar and coterminous.

6th. If several repetends of equal places be added together their sum will be a repetend of the same number of places; since every set of periods will give the same sum.

### ADDITION OF CIRCULATING DECIMALS.

80. To add circulating decimals :-

# HE THE SEE DOLLE IN SULL.

Make the repetends similar and coterminous and write them under one another, so as to have the units of the same order in the same vertical column.

Add, beginning at the right hand side and carrying what would have been obtained if the decimale had been carried out two or three places further.

Example—Add together .788, .927, .421 and 9-128456.

issimilar.	Similar.	. Similar and Cote	rminous.
783 =	788	= '788888888888	14 14 19 19 19 19 19 19 19 19 19 19 19 19 19
1007		9272727272727	i mi
41 <b>41</b> 34 3 7 7	49142 :	= 4914914914914	
P198466 =	9-123456	91234563456345	i carried.

Sum. = 11.95548389768904

Exercise 78.

1. Add together .9, 6.327, 19.43, 27.0278 and .0347123.

Ans. 53.8198688274.

2. Add together 7:427, 9:1234, 17:2987643 and 18:67.

Ans. 52.526228203901471.

3. Add together 4.95, 7.164, 4.7123 and .97317.

Ans. 17.8092502138.

4. Add together 1.5, 99.083, 162, 814, 2.93, 3.769230, 97.26 and 134.09. Ans. 339.626177443.

SUBTRACTION OF CIRCULATING DECIMALS.

81. To subtract one repetend from another:

#### RULE.

Make the repetends similar and coterminous, and write one be-

the

38.

1.26

ben

neath the other, so as to have units of the same order in the same vertical column.

. Subtract as in whole numbers, taking notice whether one would have been borrowed if the periods had been extended.

### Example.—From 97.03429 take 11.03876.

	Dissimilar.	Similar.	Similar and Coterminous
	97.08439	97.08420	97.084302020
7	11.08876	11.088768	11:088768768 FELTINA

### True difference. 85 995524160

If the periods had been extended, we would have had to borrow one from the last figure of the minuend period; and bearing this in mind, we say a from 8, 0, &c.

### EXERCISE 74.

2.				
1.	From	729-3427	take 93·126.	Ans. 636-216742.
2	From	1.437201	take :00713	Ans 1-4301600K97894

3. From 1.2754 take .47384. Ans. .65370016280907.

4. From 42·18763 take 17·0000008482. Ans. 25·1876824900.

### MULTIPLICATION OF CIRCULATING DECIMALS.

82. To multiply one repetend by another or by a finite decimal:—

#### RIILE.

Change the decimals into their equivalent vulgar fractions (Arts. 77 and 78), multiply these together, and reduce the product to its equivalent decimal.

EXAMPLE 1.- Multiply .3 by .78.

$$3 = 3 = \frac{1}{2}$$
 and  $78 = \frac{14}{2} = \frac{14}{2}$ .

Therefore,  $3 \times 78 = \frac{1}{4} \times \frac{14}{14} = \frac{14}{14} = 26 \text{ Ans.}$ 

Example 2.—Multiply .318 by .7432.

$$\cdot 318 = \frac{7}{24}$$
 and  $\cdot 7432 = \frac{44}{2}$ .

Therefore,  $318 \times 7432 = 7 \times 44 = 34 = 23648$ .

1. Multiply 7.25 by 2.9.

Am. 1911-750 Antillusia.

2. Multiply 297 by 7-72.

And 9:40542

3. Multiply .818 by .77.

- fne. 2 68.

4. Multiply 1.735 by .47058.

Ans. 81654168350.

5. Multiply 4.722 by .198.

Ans. 985.

### DIVISION OF CIRCULATING DECIMALS.

88. To divide one repetend by another or by a finite dening the state of the state o

Change the decimals into their equivalent vulgar fractions. divide as in Art. 48, and reduce the result to its corresponding

Example. - Divide :427 by .818.

.427 = 170 and .818 = 17.

### EXERCISE 76.

1. Divide .082 by .123.

Ans. '6.

and a me is a 2. Divide 389-185 by 15.7.

Ans. 24.6.

8. Divide .81654168350 by .47053.

Ans. 1-785

4. Divide :45 by :118881. Ans. 3.8235294117647058.

### EXERCISE 77.

### MISCELLANEOUS EXERCISE ON DECIMALS.

- 1. Reduce 1 of 7 of 14 to its equivalent decimal.
- 2. Multiply .67 by 2.13.
- 3. Find the value of .678125 of a week.
- 4. Reduce 92437 to its equivalent fraction.
- 5. Add together 67.234,98.713, and 91.03471234, and from their To Them it was the sum take 100·123456789.
- 6. Reduce 5 fur. 36 rife. 2 yds. 2 ft. 9 in, to the decimal of

6.

6

- 7. Find the difference between 17:428571 sq. ft. and 100:8 sq. in.
- 8. What is the value of 91789772 of two some?
- 9. Reduce 11-287 and 1-0428571 to yulgar fractions.
- 10. Divide 47-845 by 1.76.
- 11. From 85.62 take 18.76482.
- 12. What is the difference between .784 of a lb. and .198 of an os. avoirdupois?
- 18. How many yards of carpet 2 ft. 51 in, wide will be required. to cover a floor 27.3 ft, long and 20.16 ft. wide.
- 14. Multiply 3-145 by 4-29%.
- 15. How many finite places are there in the decimals corresponding to 30, 30, 16, tot, 16, and 1886?
- 16. Add together 813, 61-126, 32833; and 5:624.
- 2-8 of 2-27 to a sim-4.4-2.83 of 6.8 of 8 17. Reduce. ple quantity.

### QUESTIONS TO BE ANSWERED BY THE PUPIL.

NOTE.—The numbers after the questions refer to the articles of the

- Norm.—The numbers after the questions refer to the articles of the Bection.

  1. What is a decimal fraction? (52)
  2. What is the distinction between a decimal and its corresponding decimal fraction? (54-and Art. 47 Sect. I.)
  2. How is a decimal reduced to its corresponding decimal fraction? (55)
  3. How would you reduce 4 os. 17 dwt. 18 ers. to the decimal fraction? (55)
  4. How would you find the value of '71345 of a French elle? (58)
  7. What is an ent by repeating or circulating decimals? (49)
  8. What is a repetend, period, or circle? (81)
  9. What is a single repetend, and how is it expressed? (62 & 63).
  10. What is a circulating decimal or compound repetend, and how is it expressed? (64 & 65)
  11. What is a pure repetend? (67)
  12. What is a pure repetend? (67)
  13. What are similar repetends? Give example. (69)
  14. What are distimilar repetends? Give example. (69)
  15. What are observations repetends? Give examples. (70)
  16. When are repetends mid to be both similar and cotemninque? Give examples. (71)
  17. When can a vulgar fraction be exactly expressed by a decimal? (72)
  18. Show that this must hecosiarily be the case. (76)
  19. How can we assorted the number of pictors in the finite part of the decimal corresponding to any vulgar fraction contain are petend. What is the greatest number of pictors in the finite part of the decimal corresponding to any vulgar fraction contain are petend. What is the greatest number of pictors in the finite part of the Matt are particle repetends? (70)
  18. Show that this must necessarily be the case.

  25. What are particle repetends? (70)
  26. How is a pure repetend reduced to a vulgar fraction? (77)
  27. How is a pure repetend reduced to a vulgar fraction? (77)

- 24. How is a mixed repetend reduced to a vulgar fraction? (78)
  25. Show the truth of this rule. (75)
  26. Show that any finite decimal may be made into a repetend. (79)
  27. Show that any repetend may be reduced to another having twice, thrice, &c., as many places. (79).
  28. Show that any number of repetends may be made to have the same number of places, and give the rule. (79)
- Show that any pure repetend may be transformed into a mixed repetend. (79). Show that two or more repetends may be made similar and cotermi-

- nous. (70)

  31. How are circulating decimals added? (80)

  32. How are circulating decimals subtracted? (81)

  33. How do we multiply circulating decimals together? (82)

  34. How do we divide one circulating decimal by another? (83)

### TREE TENEDIST 78.

### MISCELLANEOUS EXERCISE.

### (On preceding Rules.)

- 1. Transform 4312131 quinary, into the nonary, ternary, and octenary scales, and prove the results by reducing all four numbers to the decimal scale.
- 2. Write down seven hundred and two trillions seven millions thirty thousand and seventeen, and four millions and seventysix tenths of quadrillionths.
- 3. Divide 976.432 by .00000096.
- (21+·5625-1·5++k)-1 4. What is the value of (1 1 × 1 × 296 × 101 :

- 5. Divide 97 lb. 3 oz. 4 dr. 1 sor. 17 grs. by 9 lb. 7 oz. 7 dr. 2 scr.
  6. A wall is to be built 15 yards long, 7 feet high, and 13 in. thick, with a doorway 6 ft. high and 4 ft. wide: how many bricks will it require, the solid contents of each being 108 cubic inches?
- 7. Multiply 9 ft. 6' 4" 7" by 11 ft. 7' 9" 11"
- 8. Find the value of 1 of 13+6 of 4.
- 9. Reduce 782436 pints to bushels, &c.
- 10. Find the least common multiple of 77, 42, 27, 21,33, 14, 7, 11, 63, and 30.
- 11. Divide 36187942 by 28e4 in the duodecimal scale. Also change 3762814 from the nonary to the decimal scale.
- 12. How many divisors has the number 150528?
- 13. Find the value of 1234625 of 2 weeks and 2 days.
- 14. Multiply 27 lb. 4 oz. 3dr., avoirdupois, by 7281.
- 15. Add together \$98.17, \$42.29, £16 3s. 82d., \$97.19,\$127.874, and from their sum subtract £67 17s. 71d
- 16. Reduce .8, .76, .9123, and .003327 to their equivalent vulgar

17. Take the number 704 and by removing the decimal point, (1) Make it 10000 times greater; (2) make it 10000000 times less; (3) make it billions; (4) make it hundredths of billionths; (5) make it tenths of millionths; (6) make it hundredths.

> $[\{(2\frac{1}{2} \times \cdot 5 \text{ of } 1\frac{1}{7}) + 9\frac{1}{7} + \cdot 09 + \frac{9}{4}\} - 11\frac{1}{7} + (\frac{1}{7}) + (\frac{1}{7})$ [(.7632763×11)×+ of +8+]×(+ of .2 of .3 of .25 of 96)+.2

- 18. Reduce 1 of .6732467 +1.
- 19. Divide £550 3s. 11d. among 4 men, 6 women, and 8 children, giving to each man double of a woman's share; and to each woman triple of a child's.
- 20, Add together 16,7, 194, 284, and 1294. 21. Write down all the divisors of 8100.
- 22. Find the G. C. M. of 2691, 11817 and 9828.
  23. Find the exact length of the lunar month which contains
- 2551448 seconds, and of the solar year, which contains 31555928 seconds.
- 24. How many times will a carriage wheel turn in going from Toronto to Hamilton, a distance of 38 miles, the circumserence of the wheel being 14 feet 11 inches?
- 25. What is the weight of the water contained in a rectangular cistern 11 feet wide, 13 feet long, and 15 feet deep, and how many gallons of water does it contain?

NOTE.—A cubic foot of water weighs 625 lbs, and a gallon weighs 10 lbs.

- 26. Reduce £73 17s. 11ad. to dollars and cents.
- 27. From 93 take 76 3 and divide the result by
- 28. Find the value of  $\frac{1}{1}$  of  $\frac{2}{3}$   $\div$   $10\frac{1}{3}$   $\times$   $\frac{1}{3}$  of  $\frac{1}{3}$  of  $\frac{5}{3}$ .
- 29. Transform 91342 undenary into the quinary, duodenary and binary scales and prove the results by reducing all four numbers to the decimal scale.
- 30. What are the prime factors of 7680?
- 31. Reduce 72 miles, 3 fur., 7 per., 2 yds., 1 ft., 7 in. to lines.
- 32. Find the price of 97 pairs of gloves at 47 cents per pair.
- 33. What is the worth of a pile of cord wood 73 feet long, 4 feet wide and 11 feet high, at \$3.62} per cord?
- 34. Divide 93.723 by 29.4173.
- 35. How many bushels of oats are there in 73429 lbs?
- 36. What is the worth of 719630 lbs. of wheat at \$1.80 per bushel?
- 37. Add together \$72.14 and \$93.76; multiply the sum by 9.47 and divide the product equally among 11 persons.
- 38. Find the G. C. M. of 21389 and 180781.

<sup>\*</sup> These questions though apparently difficult are not so in reality—they are designed for exercise in cancelling, and do not require much work.

15 t

the

the

and

por

39. Reduce 17, \$, \$, \$, \$1, 70, and 1 to equivalent fractions, having a common denominator.

40. Purchased 17 yards of cotton at 11 cents per yard, 19 yards of ribbon at 37½ cents a yard, 14½ yards of silk at \$2.17 a yard, a parasol \$4.75, a bonnet \$11.50, 67 yards of sheeting at 27 cents a yard, 15 yards of French merino at \$1.37½ a yard, and trimmings \$7.93. Required the amount of my bill.

# SECTION V.

### RATIO AND PROPORTION.

1. Two numbers having the same unit may be compared with one another in two ways.

1st. By considering how much greater or less one is than

the other; and

2nd. By considering how many times one contains the

2. Ratio is the relation which one number bears to another with respect to magnitude, when the numbers are compared by considering, not how much greater or less one is than the other, but how many times or parts of a time one contains the other. Hence:

The ratio of two numbers is the quotient arising from the

division of one by the other.

Thus the ratio of 18 to 6 is 3, since 18 +6=3, the ratio of 7 to 21 is \( \frac{1}{2} \), since

7+21=7,=1.

- 3. The ratio of one number to another, when measured with respect to their difference, is sometimes called arithmetical ratio, to distinguish it from the ratio considered as in (Art. 2), which is called geometrical ratio. In the following pages, whenever the term ratio is used, geometrical ratio is meant; we shall use the term difference in place of arithmetical ratio.
- 4. Since ratio simply expresses the quotient arising from the division of one number by another, and since (Art. 66, Sect. II.) we have three ways of indicating division, it follows that we have three ways of expressing the ratio of one number to another.

Thus the ratio of 9 to 4 is expressed either by 9÷4, or by  $\frac{9}{4}$ , or by 9:4. The ratio of 7 to 13 is indicated either by 7÷13, or by  $\frac{7}{13}$ , or by 7:13.

5. Ratio can exist only between numbers of the same kind.

T. Y.

ons. ards 17 2

ting

.371

my

ared

than

the

s to

are

the

since

ct to

sh it

ratio

sing

ince

livi-

the

ame

Thus it is obvious that no comparison with respect to magnitude can be made between 6 hours and 11 pounds, or between 19 days and 16 index, ac. i.e., these numbers are not of the same kind, and therefore no ratio can exist between them.

KATIQ.

6. Numbers are of the same kind when they are of the same denomination, or when they have the same unit. or when one can be multiplied so as to exceed the other.

7. The two given numbers which constitute the ratio are called the terms of the ratio; when spoken of together they are called a couplet.

8. The first term of a couplet is called the antecedent;

the last term, the consequent.

When the ratio is expressed in the form of a fraction, the numerator is the antecedent and the denominator the consequent.

9. Ratio is either direct or inverse, simple or compound.

10. A Direct Ratio is that which arises from the division of the antecedent by the consequent.

11. An Inverse or Inverted Ratio is that which arises from the division of the consequent by the antecedent,

Thus the inverse ratio of 15 to 3 is 3:15 or 3, or 3:15, or 1.

12. An Inverse Ratio is semetimes called a reciprocal ratio.

Thus the reciprocal ratio of 15 to 8 is 3:15 or  $\frac{3}{16} = \frac{1}{2} =$ inverse ratio of 15 to 8.

13. The reciprocal of a quantity is unity divided by that quantity.

Thus the reciprocal of 8 is 1; of 11, 17; of 3, 7; of 13; of 1, 9; of

- 14. When the direct ratio of two numbers is expressed by points, the inverse or reciprocal ratio is expressed by inverting the order of the terms; when by a fraction, by inverting the fraction.
- 15. A Simple Ratio is one that has but one antecedent and one consequent.

Thus 9:3, 7:11, 18:2, &c., are simple ratios.

16. A Compound Ratio is a ratio produced by compounding or multiplying together the corresponding terms of two or more simple ratios.

17. It must be distinctly remembered that a compound ratio is of the same nature as any other ratio, and, like a simple ratio country of one antecedent and one consequent. The term compound ratio is used merely to indicate the origin of the ratio in particular cases.

18. Ratios are compounded by multiplying together all the antecedents for a new antecedent, and all the consequents for a new consequent.

Thus, the ratio compounded of 2:7, 2:3, 5:11, and 4:3 is  $2 \times 2 \times 5 \times 4:7 \times 3 \times 11 \times 3$  or 80:963.

### EXERCISE 79. 4 17 to the

1. What is the ratio of 27 to 3?	Ans. 9.
2. What is the ratio of 7 to 11?	Ans. 7r.
3. What is the ratio of 9 to 27?	Ans. f.
4. What is the ratio of 42 to 5?	Ans. 8%.
5. What is the ratio of 72 to 6?	Ans. 12.

### Required the ratio of the following numbers:

	13. \$17 to \$8.50. Ans. 2.
	14. \$93 to \$31. Ans. 3.
	15. 14 bus. to 2 pks. Ans. 28. 16. 40 m. to 12 fur. Ans. 26.
	17. 24 lb. to 12 oz.
11. 23 to 299.	18. 17 shillings to £51.
	19. 16 acres to 30 sq. per.
to the same of the	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1

### Required the inverse ratio of the following numbers:-

20. 7 to 21. Ans. 3	3. 27. 6 days to 4 weeks. Ans. 43.
21. 12 to 2. Ans.	28. 11 min. to 30 sec. Ans. 32.
	. 29. 4 lbs. to 12 oz. Ans. 36.
23. 9 to 36. Ans.	4. 30. 3 qts. to 43 gals. Ans. 571.
24. 19 to 57.	31. 70 per. to 2 miles.
25. 81 to 9.	32. 7 Flem. ells to 9 Eng. ells.
26. 187 to 17.	33. 11 oz. to 68 scruples.

## Required the reciprocal ratio of the following numbers:-

34. 7 to 42. Ans. $+:\frac{1}{42}=6$ .	39.	1 to 36. 1 1 1 Ans. 3.
35. \( \) to \( \) \( \) Ans. \( 8:2=4. \) 36. \( 42 \) to \( 28 \) Ans. \( \)		
		1 to 1.

### Required the ratios compounded of the following ratios:-

	44. 2	to 3	. 5 to.	7 and 1	to 7.		A .	Ans	. 10 to 1	147.
	45. 8	to 6	and 1	7 to 3.	1 1 11	1 016			. 136 to	
						3 and		Ans	. 2520 : 8	364.
T.	47. 1	to 7	, 1 to	3, 3 to	1 and	5 to 1	186 - 1		Ans. 15:	21.
4	48. 2	to 5,	3 to 3	, 4 to	5, 21 t	o 2 and	1 to 9.	Ans	.504:31	150.

19. Since the antecedent of a couplet is a dividend, the consequent a divisor, and the ratio the quotient, it follows from the principles established in Arts. 79-84, Sect. II., that:—

1s ing t that

> Thu The 2n

that I

3rd seque the v

20 gether ed by Sect.

Who fore m that is

Ex

18 11 84

Exa follow

\_ **4**₹ 8 **4** 

14 2

77

ns. 44. ns. 22.

ns. 16. s. 571. ells.

Ans. 3. Ins. 1. ns. 76.

0 147. to 18. : 864. 5:21.

3150. d. the llows . II.,

1st. Multiplying the antecedent of a couplet or dividing the consequent by any number multiplies the ratio by that number.

Thus the ratio of 28 to  $112=\frac{1}{2}$ . The ratio of 28  $\times$  8 to  $112=\frac{1}{2}=\frac{1}{2}\times 3=$  three times the ratio of 28 to 113. 2nd. Dividing the antecedent of a couplet or multiplying the consequent by any number divides the ratio by that number.

Thus the ratio of 64 to 16 = 4. The ratio of 64  $\div$  2 to 16 = 32 : 16 = 2 = 4  $\div$  2 = half the ratio of 64 to 16.

3rd. Multiplying or dividing both antecedent and consequent of a couplet by the same number does not alter the value of the ratio.

Thus the ratio of 18 to 6 is 8, The ratio of 18  $\times$  7 : 6  $\times$  7 = 126 : 42 = 8 = ratio of 18  $\div$  2 : 6  $\div$  2 = 9 : 8.

20. Since any number of ratios to be compounded together may be expressed as fractions and then compounded by the rule for multiplication of fractions (Art. 45, Sect. IV.) it follows that:

When several ratios are to be compounded together we may, before multiplying the corresponding terms together, cancel any factor that is common to an antecedent and a consequent.

EXAMPLE 1.—Compound together 4: 17, 34:55, 11:2, 13:7, and 21:65.

5. 55 .11  $=4\times3:5\times5$ 18 5

EXPLANATION.—17 cancels 17 and reduces 34 to 2 and this 2 cancels 2, the third consequent; 11 reduces 55 to 5; 13 reduces 65 to 5 and 7 reduces 21 to 3.

×3:5×5 The only antecedents now left are 4 and 3 which multiplied together make 12, and the only remaining consequents are 5 and 12:25 Ans. 5 which multiplied together make 25.

The ratio 12 to 25 is therefore the ratio compounded of all the given ratios.

203

EXAMPLE 2.—Compound the following ratios: OPERATION.

Y : 16 9: 49 or 18 : 13 Ans. 14 : 11 23 : 13

EXAMPLE 3.—Find the ratio compounded of the following ratios:

OPERATION. 2B 16 23 . Zg 98 : 319 =1: 4 Ans. 328 : 64

### Exercise 80.

1. Find the ratio compounded of 9: 16, 25: 31; 341: 18 and 48: 100.

2. Find the ratio compounded of 18: 25, 7: 9, 11: 12, and 91: 49.

3. Find the ratio compounded of 1:2, 2:3, 3:4, 4:5; 5:6:6 and 7:11.

4. Find the ratio compounded of 2: 5, 8: 11, 14: 17 and 187: 112.

Find the ratio compounded of 8: 5, 7: 9, 11: 13, 15: 17
 and 19: 21.

21. If the antecedent of a couplet be equal to the consequent, the ratio is equal to 1 and is called a ratio of equality.

If the antecedent be greater than the consequent the ratio is greater than I and is called a natio of greater incommittee.

If the entecedent be less than the consequent the ratio is less than 1, and is called a ratio of less inequality.

Thus the ratio of 7: 7 = 1 is a ratio of equality.

The ratio of 7: 2 = 3; is a ratio of greater inequality.

The ratio of 7: 14 = ; is a ratio of less inequality.

### Exercise 81.

In examples 1-43 of Exercise 79 point out which are ratios of greater and which ratios of less inequality.

32. Ratios are compared with one another by expressing them in the form of fractions—reducing these to their equivalent fractions having a common denominator and comparing the numerators.

Ratios may also be compared by actually dividing the antecedent by the consequent and thus ascertaining which gives the greatest quotient.

North-The latter method is usually the more convenient.

Example 1.—Which is the greatest and which the least of the following ratios, viz: 8:4, 7:8, and 9:107

By 1st Rule 7:  $8 = \frac{3}{1} = \frac{30}{10}$  Hence 9: 10 is greatest and 3: 4 least.

By 2nd Rule 7: 8=7: 8= 875 Hence 9: 10 is, greatest 9: 10=9: 10=9: 4 least,

EXAMPLE 2. Compare together the following ration, 7:8, 2:3 and 11:13 and 5:6.

1/A

By

Ву

1. Po

2. Po

3. Po

same have

Thus, and the That

other resulti

Thus, ratio of —2 = 85

25.

Thus, The re

the sa

Thus.

The r. 8:46.

Carona VI

: 18 and e. 28 : 6. 12, and 43 : 150. : 5; 5 : 6 s. 7 : 66. : 17 and

no. 2:5. 15:17 09:663. the conratio of

eater in-

the mtio

ratios of

g them in fractions

ntecedent greatest

least of

and 3:4

greatest

7. 1. 8;

7: 8= 1=11 By 1st Rule 2: 8= 3co 11 | Hence 7: 8 is the greatest and 2: 3 is the least.

7: 6= 7: 8=675
2: 3= 2: 3=6

By 2nd Method 11: 13=11:13=846153
5: 6= 5: 6=83

Hence 7:8 is the greatestand 2:8 the least.

### Extrores 82.

1. Point out which is greatest and which least of the ratios 7:4, 6:3, 17:8, and 11:5.

Ans. 11: 5 is greatest and 7: 4 least.

2. Point out which is greatest and which least of the ratios 16:9, 10:3, 7:2, and 8:3.

Ans. 7: 2 is greatest and 16: 9 least.

3. Point out which is greatest and which least of the ratios 7: 33, 11: 49, 16: 71, and 21: 106.

Ans. 16: 71 is the greatest and 21: 106 least.

23. If the terms of two or more couplets, having the same ratio, be added together, the resulting couplet will have the same ratio.

Thus, the ratio of 6:2=3, the ratio of 21:7=3, and the ratio of 38:11=3, and the ratio 6+21+33 to 2+7+11, that is, of 60 to 20 is also 3.

That is, if 6:2=21:7=36:11, then 6+21+33:2+7+11=6:2.

24. If from the terms of any couplet the terms of another couplet having the same ratio be subtracted, then the resulting couplet will have the same ratio.

Thus, the ratio of 35 to 5 is 7, and the ratio of 14 to 2 is 7. So also the ratio of 35—14:3 2, then 35—14:5 — 2 = 35:5.

25. A ratio of greater inequality is diminished by adding the same number to both terms.

Thus, the ratio of 48+12:8+12 or 60:20=8 which is less than ratio 48:8.

26. A ratio of less inequality is increased by adding the same number to both terms.

Thus, the ratio of the Ly the land of

The ratio of 5+15:45+15 or 30:50 = which is greater than ratio of 8:46.

quo

quoi

be f

181

21

370

4th

E

E

4th t

are §

3rd

### PROPORTION.

27. Proportion is an equality of ratios.

Thus, the ratios 15: 3 and 25: 5 constitute a proportion, since 15: 3 = 5 =

28. The terms of the two couplets are called proportionals.

29. Proportion may be expressed in two ways,

1st. By placing =, the sign of equality, between the ratios.

2nd. By placing four points, thus :: , between the two

Thus, we may express the proportion existing between 15, 8, 25, and 5 by 15:8=25:5, or by 15:3::25:5.

We read either of them by saying the ratio of 15 to 8 equals the ratio of 25 to 5; or simply 15 is to 3 as 25 is to 5.

Note.—The sign: is supposed to be derived from —, the sign of equality, the four points being merely the extremities of the lines.

30. In every proportion there must be four terms, since there must be two complets, and each couplet consists of two terms.

31. When three numbers constitute a proportion, one of them is repeated so as to form two terms.

Thus, if 18, 6, and 2 are proportionals.

In this case the 6, i. e., the term repeated, is called the *middle* term or a mean proportional between the other two numbers.

The 3 is called the *third* term or a *third* proportional to the other two

numbers.

32. It is important to remember the distinction between ratio and proportion.

A ratio consists of two terms, an antecedent and a consequent.

A proportion consists of two couplets or four terms.

One ratio may be greater or less than another. One proportion cannot be greater or less than another, since equality does not admit of degrees.

33. The outer terms of a proportion are called the extremes, and the two intermediate ones, the means.

Thus, in the proportion 3:17::21:119.
3 and 119 are the extremes.

17 and 21 are the means.

34. If four quantities be proportionals, the product of the extremes is equal to the product of the means.

6:11::18:33. Then 6 × 35 = 11 × 18.

:8=5=

propor-

een the

the two

s, and 5 by

he ratio of of equality,

ms, since nsists of

ion, one

e term or a

veen ratio

nsequent.

her, since

d the ex-

roduct of

S. SAR MA

This may be established in the following manner:—6: 11= $\frac{1}{4}$  and 18: 88= $\frac{1}{4}$ , and since 6:11::18:88,  $\frac{1}{4}$ = $\frac{1}{4}$ § (Art. 27.) Now, since multiplying equals by the same number does not destroy their equality, if we multiply these fractions by 11 we get  $6=\frac{16\times11}{38}$ ; and multiplying each of these by 38, we have  $6\times 23=18\times 11$ ; but 6 and 33 are the extremes and 18 and 11 are the means; therefore in any geometrical proportion the product of the extremes equals the product of the means.

The same fact may be established more generally as follows:

Let a, b, c and d be any four proportionals whatever. Then a: b:: c: d

But  $a: b = \frac{a}{b}$  and  $c: d = \frac{c}{d}$ 

Therefore  $\frac{a}{b} = \frac{\sigma}{d}$  — Multiplying each of these equals by  $b \times d$ , we have  $a \times d = b \times c$ . But a and d are the extremes and b and c are the means. Therefore, &c.

35. This principle then may be considered the test of a geometrical proportion. If the product of the extremes equals the product of the means, the four quantities are proportional; if the products are not equal, the numbers are not proportional.

### 86. It follows from Art. 34 that :-

1st. If the product of the means be divided by one extreme, the quotient will be the other extreme.

2nd. If the product of the extremes be divided by one mean, the quotient will be the other mean.

and hence,

3rd. If any three terms of a proportion be given, the fourth may be found thus:

Example 1.—What is the fourth proportional to 7, 11 and 35?

4th term =  $\frac{2nd \text{ term} \times 3rd \text{ term}}{1st \text{ term}} = \frac{11 \times 35}{7} = 55 \text{ Ans.}$ 

EXAMPLE 2.—The first, second and fourth terms of a proportion are 9, 16 and 128. Required the third term.

$$8rd term = \frac{1st \times 4th}{2nd} = \frac{9 \times 128}{16} = 72 Anc.$$

### Exercise 83.

1. The second, third and	fourth terms of	a pro	portion	are 17,
11; and 93j. What is	the first term?	- 11ti		Ans. 2.

2. The first, third, and fourth terms of a proportion are 21, 68 and 39. Required the second term.

3. The first three terms of a proportion are 2, 3 and 7. What is the fourth term? Ans. 101.

4. The last three terms of a proportion are 91, 88 and 104. Required the first term. Ans. 77.

E THE THE TOUTH Proportional to		seemed in
5. 4 yds. 18 yds. and \$96.	Ans	\$432.
6. 5 lb. 2 lb. and \$3.75.	Ans.	\$1-00.
7. 1 cwt. 215 cwt. and \$7.50.	Ans. \$10	312.50.
8. 6 miles, 1 mile and 27 shillings.	Ans.	
9. 10 lb 150 lb and £6 8s. 9d.	Ang £92 1	

10. 4 days, 27 days and \$100. Ans. 3675.

37. It will be useful to remember the following properties of decimetrical proportion. As the proofs are given in every common work on Algebra, it has not been thought advisable to insert them here. a, b, c and d stand for any four proportionals whatever.

	If a: b::e:d Or i	if 15:6:;10:4 15:10::6:4
	Inversely b:a::d:c	6:-16::4:10
	By Composition $a+b:b::c+d:d$ By Division $a-b:b::c-d:d$ By Conversion $a:a+b::c:c+d$	15-6:6::10-4:4, or 9:6::0:4 15:15+6::10:10+4, or 15:21::10:14
14	Ora:a-b::a:a-d	15:15 - 6::10:15-4. or 15:9::10:6

38. Proportion in Arithmetic is usually divided into simple, compound and conjoined.

### SIMPLE PROPORTION.

39. Simple Proportion is frequently called the Rule of Three, because when three terms are given, by means of them a fourth may be found. It is also sometimes called the Golden Rule from its extensive utility.

40. Example.—If 16 barrels of flour cost \$112, what will 129 barrels cost?

In this and every other question in Shaple Proportion there are two ratios, one of which is perfect (i.e. has both terms given) and the other imperfect and from the nature of proportion we know that these two ratios must be both of the same kind, that is, they must be both ratios of greater inequality or both ratios of less inequality.

Now in the above example, the ratio of \$112 to the answer is a ratio of less inequality since it is evident that, if 16 barrels cost \$112, 129 barrels will cost more. Therefore the other ratio is also a ratio of less inequality and must be written 16: 199.

and must be written 16: 129.

re 17, ne. 2.

11, 63 s. 13.

What 101.

104. u. 77.

\$432.

51-00. 12.50.

ls. 6d.

3s. 3d. \$675."

ties of every ble to

tionals

:: 10:14 :10:6 d into

lule of fthem ed the

ill 129

retwo her imo ratios greater

ratio of barrels quality And since the ratios are equal.

dollars. 16:129::112: Ans.

Also (Art. 36) Ans. = 112×129 = \$908.

PROOF.—Set 903 in the fourth place, thus:
i6:129::113:903
and see if the product of extremes = product of means (Art. 85.)
16×903 = 14448 = 129×112.

From the preceding illustrations and principles we deduce for Simple Proportion the following general

#### RULE.

Set the given term of the imperfect ratio in the third place, and

the letter x, to represent the answer, in the fourth.

Then, if, by the nature of the question, the ratio of the third term to the answer is a ratio of greater inequality, make the remaining ratio a ratio of greater inequality also; but if the ratio of the third term to the answer be a ratio of less inequality, make the other ratio a ratio of less inequality also.

Lastly, (Art. 36,) multiply the second and third terms together, divide the product by the first term, and the quotient will be the

answer in the same denomination as the third term,

PROOF.—Multiply the first term and the answer together, and, if the product is equal to the product of the second and third terms, the work is correct. (Art. 35.)

Example 1.—If a man can walk 155 miles in 12 days, how many miles can he walk in 60 days?

Here the imperfect ratio is 155 miles to x, and, in order to ascertain whether it is a ratio of greater or less inequality, we have merely to ask the following simple question—If a man can walk 155 miles in 12 days, can be walk more or less in 60 days? Evidently more. Therefore the ratio of 155:x is a ratio of less inequality, or, in other words, the antecedent must be the least of the two numbers, and the statement is

days. miles. 12:60::155:#,

Whence the answer  $=\frac{60\times155}{19}=775$  miles.

11. Since the second and third terms multiplied together, constitute a dividend, and the first term is a divisor, it is manifest, from the principles of division (Arts. 79-84, Sect. II.), that we may cancel any factor that is common to the first term and either of the other terms.

Thus in the last example we have 12:60::155:x and, dividing the first and second by 12, we get 1:5::155:x and  $155\times 5=775$  Ans.

EXAMPLE 2.—If 96 bushels of wheat cost \$128, what will 15 bushels cost ?

As the answer to the question must be in dollars, the imperfect ratio is \$128: x, and from the nature of the question, we know that 15 bushels will

cost less than 96 bushels; we therefore place 15, the smaller of the remaining terms, is the second place, and the other term, 96, in the first place. Hence the statement is 96: 15 bushels;: \$125; \( \alpha\_i^{-1} \)

OPERATION.

\$\\ \frac{1}{6} \times 4 = \$20 Ans.

Here 82 reduces 96 to 8 and 128 to 4, and 3 cancels, 3 and reduces 15 to 5.

The teacher would do well to insist upon his pupils performing all questions in Proportion by analysis,

Thus, to solve the last question, we begin as follows: If 96 bushels cost \$125, 1 bushel will cost \$125, or \$133. Then if 1 bushel cost \$133. 15 bushels will cost 15 times as much, which is \$20.

Example 3.—If 27 men can mow 60 acres of grass in a day, how many acres can 93 men mow?

men. acres. 21, 198: 60: 20
3 81 × 20
3 3 7 7 2063 acres Ans.

Here the imperfect ratio is 60: acres, and since 96 men will evidently mow more than 27 men, we make 93 the second term and 27 the first. Hence the statement is 27: 93:: 60: a. Then 3 reduces 27 to 9 and 93 to 31, and 8 again reduces 9 to 3 and 60 to 20, and the answer is equal to 31 multiplied by 20, and divided by 5.

This question may be thus performed by analysis:

If 27 man mow 00 acres a day, 1 man will mow  $\frac{1}{47}$  of 60 acres, or  $2\frac{2}{9}$  acres; so men will therefore mow 95 times  $2\frac{2}{3}$  acres =  $206\frac{2}{3}$  Are.

#### Exercise 84.

- 1. If 11 baskets of peaches cost \$13.42, what will 87 baskets cost? Ans. \$106.14.
  - 2. If 28 cords of wood cost \$266, what will 25 cords cost?

    Ans. \$237.50.
  - 3. If a man receives \$29.20 for 16 days' work, for how many days should he work for \$83.60?

    Ans. 4543 days.
  - 4. If 16 bags potatoes are sold for \$12.80, what will 156 hags bring?

    Ans. \$124.80.
  - 5. If a stick 7 feet long cast a shadow of 5 feet, what will be the height of a tree which casts a hadow of 112 feet long?
  - 8. If a stack of hay will feed 27 cows for 99 days, how long will it feed 55 cows?
  - 7. If 9 bushels of peas sow 5 acres, how many bushels will be required to sow 48 acres?

    Ans. 867 bushels.
- J. If 3 men put up 73 perches of fencing in 2 days, how long will they take to put up 803 perches?

  Ans. 22 days.
- 9. If 1/6 pails of maple sap make 100 lbs. of sugar, how much sugar will 1128 pails make?

  Ans. 640 ft lbs.
- 10. If it cost \$20:88 to weave 108 yards of cloth, what will it cost to weave 465 yards?

ining

and 3

per-

s cont 1'85 jr.

day,

the same and the same are same

acres;

askets 06·14. st? 137·50. many days.

11 156 124.80. vill be long?

days.
will be ushels.
w long days.
much

17 1bs. will it \$89.90. 11. If \$16 pay for the carriage of 72 barrels of flour, for the carriage of how many barrels will \$1278 pay? Ans. 5751 barrels.

12. If 11 men plough 165 acres in a week, how many acres would 3 men plough in the same time?

Ans. 45 acres.

13. If 4 barrels flour make 250 four-pound loaves of bread, how many such loaves will 67 barrels make?

Ans. 4187 loaves.

14. If 190 bushels of apples make 16 barrels of cider, how many barrels of cider will 38 bushels of apples make?

Ans. 31 barrels.

15. If 90 men can build a wall in 12 days, how many men could build it in 15 days?

Ans. 72 men.

16. If 17 days' work pay for 2 barrels of flour, for how many barrels will 279 days' work pay?

Ans. 3244 barrels.

17. If a train travel 27 miles per hour, how far will it avel in 24 hours?

18. If 7 cows make 30,lbs. of butter a week, how much may be expected from 23 cows?

Ans. 98† lbs.

42. If any of the terms contain fractions or mixed numbers, apply the rules in Section IV.

EXAMPLE 1.—If \$ of a basket of peaches cost \$ of a dot ar, how much will 1 of a basket of peaches cost?

#### OPERATION.

 $\frac{3}{3}$ :  $\frac{3}{1}$ : :  $\frac{3}{3}$ : x. Therefore answer =  $\frac{3}{3}$  ×  $\frac{3}{1}$ :  $\frac{3}{3}$  =  $\frac{3}{3}$  ×  $\frac{3}{1}$  ×  $\frac{3}{1$ 

EXAMPLE 2.—If 18 of a bushel cost 14 of a pound, what will 14 of a bushel cost ?

#### OPERATION.

 $r_6:\frac{1}{12}::\mathcal{L}_{\uparrow\uparrow}:x$ . Therefore answer  $=\frac{1}{12}\times\frac{1}{12}:\frac{9}{16}=\frac{1}{12}\times\frac{1}{12}\times\frac{1}{12}\times\frac{1}{16}=\frac{1}{12}\times\frac{1}{12}\times\frac{1}{12}\times\frac{1}{16}=\frac{1}{12}\times\frac{1}{12}$ 

Norm.—If the first term be a fraction, invert it and connect it to the others by the sign of multiplication.

#### Exercise 85.

1. If 16 of a ship cost \$9750, what will 16 cost? Ans. \$42000.

2. How much will 1 of a yard come to if 1 of a yard cost 2 of a shilling?

3. If \$7.49 pay for \$\frac{7}{2}\$ of a ton of coals, what will \$\frac{3}{2}\$ tons cost?

Ans. \$80.25.

4. If 54 yards of broadcloth cost \$28.42, what will 4 of a yard come to?

5. If \(\frac{1}{2}\) of a dollar pay for \(\frac{1}{2}\) of a bag of apples, for what part of a bag will \(\frac{7}{2}\) of a dollar pay?
Ans. \(\frac{7}{2}\) of a bag.

6. If \$100 stock is worth \$981; what will \$472 H stock be worth? and disable in the bare will approximate Ans. \$467-123.

eral top significant

- 7. If 17 \$\frac{2}{3}\$ tons of hay last a certain number of horses 107-\$\frac{2}{3}\$, how many days will 11\frac{1}{2}\$ tons last the same number of horses?

  Ans. 70\frac{2}{3}\frac{4}{3}\text{ days.}
- 8. If 22‡ cords of wood last as long as  $15\frac{7}{13}$  tons of coal, how many cords of wood will last as long as  $11\frac{2}{6}$  tons of coal?

  Ans.  $16\frac{7}{18}$  cords of wood.
- 9. If 1 of 3 of  $3\frac{1}{3}$  yards of broadcloth cost 3 of  $3\frac{1}{3}$  of \$43, what will  $\frac{3}{3}$  of  $\frac{1}{3}$  of  $\frac{5}{3}$  of a yard cost?

  Ans.  $\$_{2}^{1}\$_{4}$  or \$0.0669.
- 43. When the first and second terms are not of the same denomination or contain different denominations—

#### RULE.

Reduce both to the lowest denomination contained in either, and then apply the rule in Art. 40.

EXAMPLE.—If 11 bushels 2 pks. 1 gal. cost \$74, what will 76 bushels 1 pk. 1 gal. 1 qt. 1 pt. cost?

#### OPERATION.

The lowest denomination contained in either is pints.

11 bush. 2 pks. 1 gal.: 76 bush. 1 pk. 1 gal. 1 qt. 1 pt.:: \$74:x; this reduced becomes 744: 4891:: \$74:x.

Ans.  $\frac{$74 \times 4891}{744} = $486.47 +$ 

In this example 11 bush. 2 pk. 1 gal. == 744 pints and 76 bush. 1 pk. 1 gal. 1 pt. == 4891 pints.

#### EXERCISE 86.

- 1. What will 37 sq. yds. 4 ft. 120 in. of painting cost, if 9 sq. yds. 2 ft. cost \$3.50?

  Ans. \$14.245.
- 2. How much will 12 lb. 10 oz. of silver come to at \$1.25 per oz.?

  Ans. \$192.50.
- 3. If 10 yards of ribbon cost \$3.40, what will 3 yds. 2 qrs. cost?

  Ans. \$1.19.
- 4. If 15 oz. 12 dwt. 16 grs. cost \$3.80, what will 13 oz. 14 grs. cost?

  Ans. \$3.167.
- 5. What will 3 lb. 1 oz. 11 dwt. cost, if 12 lb. 6 oz. 4 dwt. cost \$600?

  Ans. \$150.
- 6. If a man can pump 54 barrels of water in 2 hrs. 46 min. 30 sec., in what time will he pump 24 barrels?
- Ans. 1 h. 14 min.
  7. What will 73 yds. 3 qrs. 2 na. 1 in. of velvet cost, if 3 Flem.
  ells 2 qrs. 1 na. cost £4 17s. 8½d?

  Ans. £128 6s. 10½dd.
- 8. If 4\( \frac{1}{2}\) oz. avoirdupois cost 8\( \frac{1}{2}\) shillings, what will 8\( \frac{1}{2}\) ibs. cost?

  Ans. £13 9s. 0\( \frac{1}{2}\) a.
- 9. In the copy of a work containing 327 pages, a remarkable passage commences at the end of the 156th page. On what page might it be expected to begin in a copy containing 400 pages?

  Ans. On the 191st page.

6

q. 5.

er

0. s. 9.

s. 7.

> st 0.

80

10. If the rent of 46 acres, 3 roods, and 14 perches be £100, what will be the rent of 35 acres, 2 roods and 10 perches?

Ans. £75 188. 63199d.

11. When A had travelled 68 days at the rate of 12 miles a day,
B, who had travelled 48 days, overtook him. How many
miles a day did B travel, allowing both to have started
from the same place?

Ans. 17.

12. If 21 shillings pay for 16 lbs. of prunes, how many pounds can be bought for 32 shillings?

Ans. 24 4 15 lbs.

13. A ton of coal yields about 9000 cubic feet of gas; a street lamp consumes about 5, and an argand burner (one in which the air passes through the centre of the flame) 4 cubic feet in an hour. How many tons of coal would be required to keep 17493 street lamps, and 192724 argand burners in shops, &c., lighted for 1000 hours? Ans. 953734.

14. The gas consumed in London requires about 50000 tons of coal per annum. For how long a time would the gas this quantity may be supposed to produce (at the rate of 9000 cubic feet per ton), keep one argand light, (consuming 4

cubic feet per hour) constantly burning?

Ans. 12842 years and 170 days.

15. Suppose 11270 lbs. of beef for a ship's use were to be cut up in pieces of 4 lb., 3 lb., 2 lb., 1 lb., and ½ lb.—there being an equal number of each. How many pieces would therebe of each?

Ans. 1073; and 3½ lb. left.

16. The sloth does not advance more than 100 yards in a day.

How long would it take to crawl from Toronto to Kingston,

allowing the distance to be 180 miles?

Ans. 3168 days, or about 82 years.

17. Suppose that a greyhound makes 27 springs while a hare makes
25, and that their springs are of equal length. How many
springs must the hound make to overtake the hare, if the
latter has a start of 50 springs?

Ans. 675.

### COMPOUND PROPORTION.

44. Compound Proportion is an equality between a compound ratio and a simple ratio.

Thus 7:11 compounded with 22:21::34:51, is a compound ratio.
Or 7×22:11×21::34:51, and applying Art. 40 we have 7×22×51=11×21
×34.

45. Compound Proportion is also called the Double Rule of Three. It enables us to obtain the answer by a single statement, although two or more proportions are contained in the question.

46. In Compound Proportion there are three or more ratios, one of which is imperfect and all the others perfect.

47. Let it be required to solve the following question: If 18 men dig a trench 30 yards long, in 24 days, by working 8 hours a day, how many men will dig a trench 60 yards long, in 64 days, working 6 hours a day?

Let us suppose the time to be the same in both cases, and this question becomes the same as the following:

If 18 men dig 30 yards of trench, how many men will dig 60 yards?

Here it is evident the answer will be the same fraction of 18 that 60 yards is of 30 yards; or, in other words, the required number of men=3% of 18

Next let us take into account the number of days; but suppose they work the same number of hours per day in both cases.

The question then becomes: If 50 of 18 men require 24 days to dig a

trench, how many men will dig it in 6s days?

In this case it is plain that the answer will be the same fraction of \$6. of 18 men that 24 days is of 64 days; that is, the required number of men=

testly, let us take into consideration the time worked each day. The question then becomes: If 24 of 30 of 18 men dig a trench in a certain number of days, working 8 hours per day, how many men will dig it working 6 hours per day?

In this case the answer is obviously—

6 of 34 of 30 of 18 men, or dividing

Asswer these equals by 18. = 8×81×98.

Or taking the reciprocals  $\frac{10}{4nswer} = 8 \times 44 \times 50$ .

That is the ratio compounded of 6:8, 64:24, and 30:60 = ratio of

18: Answer, or, 64: 24 :: 18: Answer.

The answer is equal to the continued product of the third term, and all the second terms, divided by the continued product of all the first terms.

From the preceding principles and illustrations, we deduce the following general

#### RULE FOR COMPOUND PROPORTION.

Place that number which is of the same kind as the answer in the third term, and the letter x to represent the answer in the fourth term.

Then take the other numbers in pairs, or two of a kind, and

arrange them as in simple proportion.

Finally multiply together all the second terms and the third term, divide the result by the product of the first term, and the quotient will be the fourth term or answer required.

Norr.—Since the third term and second terms multiplied together constitute a dividend, and the first terms multiplied together a divisor, we may (Arts. 79-84, Sect. II) cancel any factors that are common to any of the first terms and to the third term or any of the second terms.

EXAMPLE 1.—If 5 compositors, in 16 days, 11 hours long, can compose 25 sheets of 24 pages in each sheet, 44 lines in each page, and 40 letters in a line; in how many days, each 10 hours long, may 9 compositors compose a volume, to be printed in the same letter, consisting of 36 sheets, 16 pages to a sheet, 50 lines to a page, and 45 letters to a line?

# STATEMENT. 9 comp.: 5 comp. 10 hours: 11 hours. 25 sheets: 36 sheets. 24 pages: 16 pages. 44 lines: 50 lines. 40 letters: 45 letters. SAME CANCELLED. 9: 5 10: 11 20: 11 21: 12 22: 16 23: 24: 16 24: 25 26: 25 2

EXPLANATION.—The imperfect ratio is that of 16 days to an unknown number of days. We place this ratio to the right hand-side, as in Simple Proportion. Now we compare each pair, of terms with this ratio, in order to decide whether they constitute a ratio of greater or less inequality. Thus, if 5 compositors require 16 days, will 9 compositors require more or less? Evidently less: therefore it is a ratio of greater inequality, and we must write it 9:5. Next, if 11 hours to the day require 16 days, will 10 hours to the day require more or less?—more; therefore we must write 10:11. Next, if 25 sheets require 16 days, will 36 days require more or less?—more: therefore we write 25:36. Next, if 44 lines to a page require 16 days, will 50. Lastly, if 40 letters to a line require 16 days, will 45 letters to a line require more or less?—more; therefore we write 44:50. Lastly, if 40 letters to a line require more or less?—more; therefore we write 40:45.

The statement is now complete, and we cancel as follows; 5 cancels 5, the first consequent, and reduces 25, the fifth consequent, to 10, and 10 dancels this 10 and 10, the second antecedent. Again, 9 cancels the interest of the second antecedent.

The statement is now complete, and we cancel as follows; 5 cancels 5, the first consequent, and reduces 25, the third antecedent to 5, and 5 cancels this 5, and reduces 50, the fifth consequent, to 10, and 10 cancels this 10 and 10, the second antecedent. Again, 9 cancels the first antecedent and reduces 36, the third consequent, to 4, and 4 cancels this 4 and reduces 44, the fifth antecedent, to 11, and 11 cancels this 11 and 11, the second consequent. Again, 8 reduces 24 to 3 and 16 to 2, 3 carbells this 3 and reduces 45 to 15. 2 cancels the 2 resulting from the 16 and reduces 40 to 20, and 5 reduces this 20 to 4 and the 15 resulting from 45 to 3. Lastly, 4 cancels this 4 and reduces 16, the third term, to 4. There remain but 3 and 4 which multiplied together make 12.

EXAMPLE 2.—If 24 men can saw 90 cords of woods in 6 days when the days are 9 hours long, how many cords can 8 men saw in 36 days, when they are 12 hours long?

1 - 1 -	SAME C	ANCELLED.	
	424 : 8 <sup>2</sup> 6	Ans. 10 × 2×12=	=
	:: 90 : x.	cords. 24:82:36:36:36	:: 90 : x.

Here the imperfect ratio is 90: Ass. If 24 men saw 90 cords, will 8 men saw more or less?—less: therefore it is a ratio of greater inequality, and we write 24: B. Next, if 6 days saw 90 cords of wood, will 36 days saw more or less?—more; therefore it is a ratio of less inequality, and we write 6: 36. Lastly, if 9 hours per day saw 90 cords, will 12 hours per day saw more or less?—more; therefore it is a ratio of less inequality, and we write 9: 12,

a 64

rore

fect.

f 18

ours

of 18

work dig a

of §6 nen=

a cerdig it

tio of

nd all rms. e de-

per in

fourth and

third d the

d tod tol any third

10.

11.

12.

13.

15. ]

16. I

17. In

18. I

19. If

fe

in

80

m

cl

ps

40

EXAMPLE 3.—If 248 men, in 51 days, of 11 hours each, dig a trench of 7 degrees of hardness, 2321 yards long, 33 wide, and 21 deep; in how many days, of 9 hours long, will 24 men dig a trench of 4 degrees of hardness, 3371 yards long, 53 wide, and 31 deep?

### STATEMENT.

The answer will be  $(2^{16} \times \frac{1}{4} \times 1 \times \frac{6}{4} \times \frac{6}{4} \times \frac{7}{4} \times \frac{1}{4} \times \frac$ 

#### CANCELLED.

$$\begin{array}{c} \frac{3}{48} \times \frac{11}{1} \times \frac{4}{1} \times \frac{15}{1} \times \frac{4}{1} \times \frac{28}{5} \times \frac{28}{5} \times \frac{7}{2} \times \frac{11}{2} \times \frac{1}{24} \times \frac{1}{9} \times \frac{1}{7} \times \frac{2}{485} \times \frac{8}{11} \times \frac{3}{7} \\ = 4 \times 3 \times 11 = 132 \text{ days.} \end{array}$$

### Exercise 87.

- 1. If 120 bushels of corn last 14 horses 56 days, how many days will 90 bushels last 6 horses?

  Ans. 98 days.
- 2. If a wall of 28 feet high were built in 15 days by 63 men, how many men would build a wall 32 feet high in 8 days?

  Ans. 135 m.sn.
- 3. If 1 lb. of thread make 3 yards of linen of 1½ yards wide, how many pounds of thread would be required to make a piece of linen of 45 yards long and 1 yard wide? Ans. 12lb.
- 4. If 3 lb. of worsted make 10 yards of stuff of 1½ yards broad, how many pounds would make a piece 100 yards long and 1½ broad?

  Ans. 25 lb.
- 5. If 12 horses in 5 days draw 44 tons of stones, how many horses would draw 132 tons the same distance in 18 days?

  Ans. 10 horses.
- 6. If 27s. are the wages of 4 men for 7 days, what will be the wages of 14 men for 10 days?

  Ans. £6 15s.
- 7. 3 mesters, who have each 8 apprentices, earn \$144 in 5 weeks—each consisting of 6 working days. How much would 5 masters, each having 10 apprentices, earn in 8 weeks, working 51 days per week—the wages being in both cases the same?

  Ans. \$440.

8. If 6 shoemakers, in 4 weeks, make 36 pair of men's and 24 pair of women's shoes, how many pair of each kind would 18 shoemakers make in 5 weeks?

Ans. 135 pair of men's and 90 pair of women's shoes.

wall is to be built of the height of 27 feet; and 9 feet high

it are built by 12 men in 6 days. How many men must

be employed to finish the remainder in 4 days?

Ans. 36.

- 10. If a footman travels 130 miles in 3 days, when the days are 14 hours long, in how many days of 7 hours each will be travel 390 miles?

  Ans. 18.
- 11. If the price of 10 oz. of bread, when the flour is 1s. 101d. per stone, is 1d., what must be paid for 3lb. 12 oz. when the flour is 2s. 6d. per stone?

  Ans. 8d.
- 12. If 5 compositors in 16 days of 14 hours long, can compose 20 sheets of 24 pages in each sheet, 50 lines in a page, and 40 letters in a line; in how many days of 7 hours long may 10 compositors compose a volume to be printed in the same letter, containing 40 sheets, 16 pages in a sheet, 60 lines in a page, and 50 letters in a line?

  Ans. 32 days.
- 13. If 336 men, in 5 days of ten hours each, dig a trench of 5 degrees of hardness, 70 yards long, 3 wide and 2 deep, what length of trench of 6 degrees of hardness, 5 yards wide, and 3 deep, may be dug by 240 men in 9 days of 12 hours each?

  Ans. 36 yards.
- 14. If a pasture of 16 acres will feed 6 horses for 4 months, how many acres will feed 12 horses for 9 months?
- 15. If 25 persons consume 300 bushels of corn in one year, how much will 139 persons consume in 7 years at the same rate?

  Ans. 11676 bushels.
- 16. If 32 men build a wall 36 feet long, 8 feet high, and 4 feet wide, in 4 days; in what time will 48 men build a wall 864 feet long, 5 feet high, and 3 feet wide?
- Ans. 30 days.

  17. If a regiment of 679 soldiers consume 702 bushels of wheat in 336 days, how many bushels will an army of 22407 soldiers consume in 112 days?

  Ans. 7722 bushels.
- 18. If 12 tailors in 27 days can finish 13 suits of clothes, how many tailors in 19 days of the same length, can finish the clothes of a regiment of soldiers consisting of 494 men.

  Ans. 648 tailors.
- 19. If 17 head of cattle consume 5 acres 2 roods 10 perches of pasture in 30 days, how many acres would be consumed by 40 head in 51 days?

Ans, 22 acres 1 rood.

⟨<del>3</del>

ays.
ien,
ys?
ide,
te a
21b.
ad,

lb.
any
ys?
ses.
the
5s.
eks
d 5

the

40.

20. If 180 bricks, 8 inches long, and 2 wide, are required for a walk 20 feet long, and 6 feet wide, how many bricks will be required for a walk 100 feet long and 4 feet wide?

Ans. 600 bricks.

### CONJOINED PROPORTION.

- 48. Conjoined Proportion is a kind of Compound Proportion, in which the ratio of one of the terms to its corresponding term is made to depend on equivalencies among the intermediate terms of the proportion.
- 49. Conjoined Proportion is sometimes called the Chain Rule from the peculiar manner in which the different pairs of terms are linked, as it were, together. It relates principally to exchanges between different countries, in respect to specie, weights, and measures, but is applicable to common business transactions.
- 50. EXAMPLE 1.—Suppose 7 yards of velvet in Toronto cost as much as 9 in Montreal, and 16 in Montreal as much as 24 in Paris, how many yards in Toronto will cost as much as 54 in Paris.

EXPLANATION.—This question may be stated as a problem in Compound Proportion as follows:

The imperfect ratio is 7 yards Toronto to an unknown 9:16 24:54 2:17:x pay for 7 yards Toronto. Then, if 9 yards Montreal, pay for 7 yards Toronto, will 16 yards pay for more or less?—more; therefore we write 9:16. Next if 24 yards Paris pay for a certain number  $\left(\frac{16\times7}{9}\right)$  yards Toronto, will 54 yards Paris

pay for more or less?—more; therefore we write the ratio 24:54. Now (Art. 47) the answer  $\frac{16\times54\times7}{9\times24}$ ; and it is evident that we may consider

all the factors of the numerator as antecedents, and all the factors of the denominator as consequents, and then make the statement thus:

7 yds. Toronto = 9 yds. Montreal. 16 " Montreal = 24 " Paris. 54 " Paris = x " Toronto.

Since the left-hand numbers constitute a dividend and the right hand numbers a divisor, we may cancel factors that are common. Merely writing the numbers and doing this we have—

SAME CANCELLED.

From the preceding principles and illustrations we deduce the following;

sign alwa Mi divid be the

ARTS.

W

Ex goats how

25 sb 33 gc 38 cc x ho

> Her term, Not equal Ex

in Ha 30 lb Bosto much are w

19 G 7 H 30 Q 81 B 10 L

ous x H

1. I

2. I

equired bricks ride? bricks.

d Proits coramong

e Chain ifferent relates ries, in plicable

nto cost as 24 in as 54 in

Compound

unknown Montreal, or more or if 24 yards ards Paris

: 54. Now y consider

tors of the

right hand erely writ-

ns we de-

### RULE FOR CONJOINED PROPORTION.

Write the equivalent terms, as they occur, right and left of the sign of equality, taking care that terms of the same name shall always be on opposite sides.

Multiply all the terms on the same side as the odd term for a dividend and all on the other side for a divisor. The quotient will

be the required term.

Example 2.—If 25 sheep eat as much hay as 19 goats, and 33 goats as much as 10 cows, and 38 cows as much as 22 horses, how many horses will eat as much as 60 sheep?

STATEMENT. SAME CANCELLED. 25 sheep = 19 goats Or writing the 33 goats = 10 cows numbers merely, 38 cows = 22 horses cancelling and apx horses=60 sheep plying the rule. \$ 25 =19<sub>2</sub> 8 88 =1011 4 3 88 = 32 x =6020

Ans. 4×2=8 horses. Here, since the term 25 sheep is on the left hand-side, we put the odd term, 60 sheep, on the right-hand side.

Note.—The sign=in such questions, merely means equal in value, or equal in time, or equal in time, or equal in time.

EXAMPLE 3.—If 19 lbs. of tea in Guelph cost as much as 20 lbs. in Hamilton, and 7 in Hamilton as much as 91 lbs. in Quebec, and 30 lbs. in Quebec as much as 291 lbs. in Boston, and 81 lbs. in Boston as much as 51 lbs. in London, and 10 lbs. in London as much as 57 lbs. in Hong Kong; how many lbs. in Hong Kong are worth 100 lbs. in Guelph?

STATEMENT. SAME CANCELLED. 19 Guelph = 20 Hamilton 19 = 20= 91 Quebec 7 Hamilton X = 91 8 86 = 383,41 30 Quebec = 29 Boston = 51 London = 57 Hong Kong 2 81 = 51 19 81 Boston 10 London 10 = 24  $x = 100^{10}$ and x Hong Kong= 100 Guelph

Ans.  $10 \times 91 \times 51 = 5063$  lbs.

### EXERCISE 88.

1. If 17 cords of wood are equivalent to 116 lbs. of tea, and 87 lbs. of tea to 23 barrels of flour, and 19 barrels of flour to 34 days' work, and 92 days' work to 57 baskets of peaches, and 31 baskets of peaches to 24 dollars, and 12 dollars to 2 tons of coal; how many cords of wood may be purchased for 35 tons of coal?

2. If 6 lbs. of tea are worth 29 lbs. of sugar, and 17 lbs. of sugar pay for 1 bushel of wheat, and 27 bushels of wheat are equivalent to 4 tons of coal, and 34 tons of coal purchase 15 cows, and 29 cows cost \$1160; how many pounds of tea can be purchased for \$20?

SECT. V

16. Wh 17. Wh

18. Sinc

19. Hov

20. Ho

21. Ho

22. How

28. How

25. Whe

26. Whe

29. Wha 81

31. Wh

32. How 33. Wha

85. Whe

36. Point 37. Wha

37. Wha 38. Prov

39. Wha

40. Dedu

41. If r: 2n 42. Wha 43. Wha

44. Give 45. Give

46. Show If an

48. If th

49. Wha

51. How

52. In st

53. How ine

54. Whe:

55. Show COL

aldesuns

Wha

th

thi

the

mg

Whe

How

Wh

How

in

3. If 11 bushels of barley pay for 21 bushels of potatoes, and 19 bushels of potatoes for 29 bushels of oats, and 115 bushels of oats for 44 bushels of wheat, and 14 bushels of wheat for 38 bushels of peas, and 60 bushels of peas for 55 bushels of rye, and 75 bushels of rye for 11 bushels of clover seed; for how many bushels of barley will 36 bushels of clover seed pay? Ans. 8794.

4. If 16 baskets of pears pay for 29 turkeys, and 17 turkeys for 7 days' work, and 71 days' work for 187 loaves of bread, and 31 loaves of bread cost as much as 4 lbs. of veal, and veal is 11 cents per pound, and \$7.92 pay for 63 lbs. of sugar; how many pounds of sugar will 21 baskets of pears purchase? Ans. 4041.

5. Suppose A can do as much work in 7 days as B can in 11 days, and B as much in 5 days as C can in 8 days, and C as much in 15 days as D can in 21 days, and D as much in 11 days as E can in 5 days; in how many days would A do as much work as E can do in 42 days? Ans. 261.

6. If 7 barrels of flour pay for 23 cords of wood, and 6 cords of wood pay for 11 cwt. of beef, and 46 cwt. of beef cost £28, and £77 pay for 9 sheep, and 5 sheep are worth as much as 8 tons of coal; how many barrels of flour may be purchased for 9 tons of coal? Ans. 131.

If 15s. in N. England be the same in value as 20s in N. York. and 24s. in N. York the same as 22s. 6d. in N. Jersey, and 30s. in N. Jersey the same as 20s. in Canada; how many pounds in N. England are the same in value as £240 7s. 6d. in Canada? Ans. £288 9s.

### QUESTIONS TO BE ANSWERED BY THE PUPIL.

NOTE.—The numbers following the questions refer to the numbered articles of the section.

- 1. In how many ways may one number be compared with another with respect to magnitude? (1)
- 2. What is ratio? (2) What is the difference between the Geometrical and the Arithmetical ratio of numbers? (3)
- 4. How many ways have we of expressing the ratio of one number to another? (4)
- 5. Between what kind of quantities only can ratio exist? (5) When are quantities said to be of the same kind? (6)
- 6. When are quality? (7)
  7. What is a couplet? (7) 8. What is the antecedent?—the consequent? (8)
- 9. How many kinds of ratio are there? (9)

- 10. What is a direct ratio? (10)
  11. What is an inverse ratio? (11)
  12. What is the reciprocal of a quantity? (13)
  13. What is a reciprocal ratio? (12)
- 14. How is the reciprocal ratio of two numbers expressed? (14) 15. Show that "reciprocal ratio" and "inverse ratio" are interchangeable terms? (12)

es, and bushels f wheat bushels

er seed ; f clover ns. 8744. keys for ead, and

and veal f sugar; ars purns. 4041. can in 11 and C as

uch in 11 A do as Ans. 261. 1 6 cords cost £28.

much as ourchased Ans. 131. N. York, rsey, and ow many

40 7s. 6d. £288 9s.

numbered

other with rithmetical

number to

16. What is a simple ratio? (15) What is a compound ratio? (16)

18. Since a compound ratio does not differ in nature from a simple ratio, why is the term used? (17)
19. How are ratios compounded together? (18)

How are ratios compounded together? (18)
 How does multiplying the antecedent or dividing the consequent of a couplet by any number, affect the ratio? (19)
 How does dividing the antecedent or multiplying the consequent of a couplet by any number, affect the ratio? Why? (19)
 How does multiplying or dividing both antecedent and consequent of a couplet by any number, affect the ratio? Why? (19)
 How does it happen that we may cancel any factors common to an antecedent and a consequent, before compounding ratios together? (20)
 When is a ratio called a ratio of equality? (21)
 When is a ratio called a ratio of greater inequality? (21)
 How are ratios compared with one another? (22)
 When equal ratios are added together, what is the nature of the resulting ratio? (23)
 What effect has adding the same number to both terms of a ratio? (25)

29. What effect has adding the same number to both terms of a ratio? (25 and 26)

What is Proportion? (27)

31. What are the terms of the two equal ratios called? (28)
32. How many ways are there of expressing Proportion? (29) 33. What is the supposed derivation of the sign::? (29—Note)

33. What is the supposed derivation of the sign:: (22-Note)
34. How many terms must there be in every proportion? (30)
35. When three numbers constitute a proportion, what is the repeated term called?—What is the last term called? (31)
36. Point out the distinctions between ratio and proportion. (32)
37. What are "extremes" and "means"? (33)
38. Prove that if four quantities are proportional, the product of the extremes is equal to the product of the means. (34)
39. What is the test of geometrical ratio? (35)
40. Deduce from this principle a rule for finding any one of the terms when

40. Deduce from this principle a rule for finding any one of the terms when

the other three are given. (36)
41. If r: w:: x: y, what does the proportion become? 1st, by composition, 2nd, alternately; 3rd, by conversion; 4th, by division; 5th, inversely. (37)

42. What are the different kinds of Proportion? (38)
43. What other names has Simple Proportion?—Why so called? (39)
44. Give the rule for making the statement in Simple Proportion. (40)
45. Give the rule for finding the unknown quantity after the statement is made. (40.)

46. Show that we may cancel any factors that are common to the first term

and either of the others, before applying the rule. (41)
47. If any of the terms contain fractions, what is done? (42)
48. If the first and second terms are not of the same denomination, what is the rule? (43)

Aldespies was a

49. What is Compound Proportion? (44)
50. What other name has Compound Proportion? (45)

50. What other name has Compound Proportion (90)
51. How many ratios are there in Compound Proportion, and how many of them are perfect? (46)

The proportion in Compound Proportion, what do you make the

52. In stating a question in Compound Proportion, what do you make the third term? (47)

53. How do you know whether the other ratios are ratios of greater or less inequality? (47)
54. When the statement is made, how is the answer obtained? (47)

55. Show that before applying the rule we may cancel any factors, that are common to any of the first terms, and to the second and third terms. (47-Note) In h . h . h

The said of the said of the said

changeable

20.

21.

22. ] 23. I

25. I

26. E

27. M

28. D

29.: V

30. 1

31. If

34. W

35. A

36. R

37. A

38. If

tl

56. What is Conjoined Proportion? (48)
57. Why is it sometimes called the Chain Rule? (49)
58. Give the rule for Conjoined Proportion. (59)
59. In what sense is the sign = taken in these statements? (5?)

### EXERCISE 89.

### MISCELLANEOUS EXERCISE.

### (On preceding Rules).

1. What is the ratio compounded of the ratios 7:8, 17:11, 23:29, 319:119, and 16:69?

2. Reduce £119 16s. 6 d. to dollars and cents.

3. How many days are there from 12th March to the 17th of the following February?

4. Compare together the following ratios, and point out which is greatest and which least, 9: 13, 21: 27, 7:10, and 11:15.

5. From 76.23478 take 19.1342291.

6. Multiply 71324t undenary by 23421 quinary and divide the result by 14e7 duodenary. Give the answer in each scale.

7. If 5.63 cubic inches of water weigh 3.254 ounces avoirdnpois, what will be the weight of 7.9 cubic inches of nitric acid having a specific gravity of 1.220?

8. Divide 63 yds. 3 grs. 2 na. 1 in. of ribbon equally among 17

What is the value of 913625 of an acre at 67 cents per sq.

10. Multiply 1 of 3 of 7 of 20 bushels by 5×6×7.

11. Of the ratios 6: 7, 17: 8, 23: 11, and 88: 176, point out (1) which is the greatest, (2) which is the least, (3) which are ratios of greater inequality, (4) which are ratios of less inequality, (5) what is the ratio compounded of these ratios.

12. The population in Canada in 1851 was 1842265, and in 1857 it was estimated at 2571437. What was the rate per

cent. of increase?

13. From one-half of two-thirds of eighteen twenty-ninths subtract one-eighth of two-thirds of five-sevenths.

14. Deduct 7 per cent. from 11 feet.

15. What is the value of 79 lbs. of tea at £ 00163 per ounce?

16. If 3 men in 21 days, working 12 hours a day, can cradle a field of wheat containing 20 acres, in how many days can 4 men, working 10 hours a day, cradle a field of wheat containing 35 acres?

17. Find the value of (\$ of 10 × 02× 456) + (17 of \$ of \$ of 51).

18. A certain number is divided by 5, the result is divided by 4, this result by 1,3, and this last result by \$. The last quotient is 2; what was the original number?

1, 23:29,

th of the

ut which 1 11 : 15.

ivide the h scale. irdupois, tric acid

among 17

ts per sq.

nt out (1) which are f less inse ratios. , and in rate per

nths sub-

ounce? cradle a days can heat con-

f t of 51). ided by 4, t quotient 19. If 50 barrels of flour in Toronto are worth 125 yards of cloth in New York, and 80 yards of cloth in New York 6 bales of cotton in Charleston, and 13 bales of cotton in Charleston 31 hogsheads of sugar in New Orleans; how many hogsheads of sugar in New Orleans are worth 1000 barrels of flour in Toronto?

20. Multiply 73.47 by .0063, and divide the result by 17.2345.

21. Reduce 2 roods 7 per. 4 yds. 3 ft. 117 in. to the decimal of

22. Deduct .73 of 11 furlongs from \$ of \$ of \$ of 70 miles.

23. From 274312 nonary take 1101011010 binary, and multiply the result by 5555 septenary. Give the answer in all three scales.

24. Find the 1. c. m. of 44, 275, 18, 190, 209, and 225.

25. If 60 men in 6 weeks of 5 working days, of 10 hours each, build an embankment 800 yards in length, 18 feet in mean breadth and 11 ft. in mean height, how many men will make an embankment 8742 feet long, 20 feet wide and 8 ft. high, in 10 weeks, of 6 days each, and 11 working hours to each day?

26. How many divisors has the number 172000?

27. Multiply 42.7 by 9.7123.

28. Deduct 27 per cent. from \$73.42.

29. What are all the divisors of 6300?

30. If \$ of \$ of 31 lbs, of coffee cost \$ of \$ of \$4 of 1 of a dollar;

what will 3 of 7 of 6 of 31 of 90 lbs. cost?

31. If \$2739.18 be divided among 7 men, 2 women, and 11 children, so that each child shall have # of a woman's share, and each woman in of a man's share, what will be the amount received by each?

32. What is the reciprocal ratio of \(^2: \frac{1}{3};\) the direct ratio of

93: 17, and the inverse ratio of  $\frac{2}{3}$  of  $\frac{7}{3}$ ?

33. Add together f of 61 yards, 3 of 4 of 82 ft., and 3 of 37 of 7% inches. 34. What is the ratio compounded of 23: 7, 4: 11, 6:5, 13:111;

and 381: 3?

35. A pint contains 9000 grains of barley, and each grain is one third of an inch long. How far would the grains in 23 bush. 2 pks. 1 gal. 1 qt. 1 pt. reach if placed one after another?

36. Reduce 1436 to its lowest terms.

37. Add together 1, 3, 3 and 7 in the octenary scale.

38. If 17 sheep eat as much grass as 6 cows, and 26 cows require 272 acres, and 12 acres supply 13 horses, and 11 horses est as much as 28 goats, how many goats will eat as much as 68 sheep?

25 of

State 100 lb

39. Suppose that 50 men, by working 5 hours each day, can dig, in 54 days, 24 cellars, which are each 36 feet long, 21 feet wide and 10 feet deep, how many men would be required to dig, in 27 days, 18 cellars, which are each 48 feet long, 28 feet wide, and 9 feet deep, provided they work only 3 hours each day?

### SECTION VI.

### PRACTICE.

1. Practice is so called from its being the method of calculation practised by mercantile men; it is an abridged mode of performing processes dependent on the Rule of Three—particularly when one of the terms is unity.

The statement of a question in practice, in general terms, would be—
One quantity of goods: another quantity of goods:: price of former: price
of latter.

- 2. The simplification of the Rule of Three by means of practice, is principally effected, either by dividing the given quantity into "parts," and finding the sum of the prices of these parts; or by dividing the price into "parts," and finding the sum of the prices of each of these parts; in either case, as is evident, we obtain the required price.
  - 3. An Aliquot Part is an exact or even part.

Thus, 2 shillings is an aliquot part of a pound; 12½ cents is an aliquot part of a dollar; 6 months, 4 months; 3 months, 2 months, 1½ months are aliquot parts of a year, &c.

TABLE OF ALIQUOT PARTS.

Parts of \$1.	Parts of a year.	Parts of a month.	Parts of £1.	Parts of 18.	Parts of a ewt. of 112 lbs.
50 ots.= 1	0 m'ths= 1	15 days== 1	10s = }	6d = 1	56 lb = }
$33\frac{1}{25} = \frac{1}{3}$	4 = 3	$10 = \frac{1}{3}$	88 8d = 3	1d = 3	28 lb =
	$3 = \frac{1}{4}$	71 = 1	58 = 1	$3d = \frac{1}{4}$	16 lb + = 4
20 = 1	3 = 8	8 = 1	4s = 1	2d = 1	14 lb =
161 = 1	11 = 1	5 = 1	3s 4d = 6	11 = 1	8 lb = 1
121 = 1	1 =18	3 = 10	2s 6d == 8	1d =12	71b =
$10 = \frac{1}{10}$		2 =18	2s = 10	1100	parts of a qr.
81 = Ty	1 th 14	1 =30	1s 8d =14	1 1	of 28 lbs.
61 = 1	i in the	H	18 4d =18	(A)	14 lb =
5 = T	si a libina .	No - 12mg P IS	1s 3d =16	4 46 3	7 lb - =
4 = 1	17		ls ===	. 4.	31 lb =
8 = 80	T C	4			14 lb 11 =

\*Although we allow but 100 lbs. to the cwt. in Canada, it is often necessary to make calculations with the old cwt., of 112 lbs. This arises from the

n dig. 1 feet quired long. only 3

DT. VI.

hod of oridged Rule of

d beer: price

neans of he given orices of s," and arts; in rice.

an aliquot onths are

of a cwt." rts of a qr.

> ften neceses from the

EXAMPLE 1.—Find the price of 2783 yards of silk at \$3.371 per yard. OPPRATION.

The cost of 2783 yards at \$3'371 = cost at \$3 + cost at ... 0. 11 19783 375 cents.

2783 yds. at \$3 comes to 8 times as much as at \$1 ; i.e., to 8 times \$2783, or \$8349.

375 cents, hence, 2783 yds. at 375 cents = price at 25 cents + 8340 194 C. 1

132 c. | | 695.75 cents, hence, 2783 yds. at 37½ cents == price at 25 cents + 347.87½ price at 13½ cents.

Since 2783 yards at \$1 comes to \$2783, and 25 cents == ½

Ans. \$9392.62½ of a dollar; 2783 yards at 25 cents come to ½ of \$2783,
i.e., to \$695.75. Again, because 2783 yards at 25 cents

come to \$695.75 and 12½ cents equals ½ of 25 cents, 2783 yards at 12½ cents

will come to ½ of \$695.75; i.e., to \$347.87½.

Then 2783 yards at \$3.37½ == price at \$3 + price at 25 cents + price at 12½

cents = \$83.40 + \$695.75 + \$347.87½ = \$9392.62½.

EXAMPLE 2.—What is the cost of 972 oz. of gold dust at £3 14s. 8id. per oz. ?

OPERATION. 104. |} = cost at £3 0 22916 86. 4d. 10d. 486 162 = cost at 0 = cost at 0 0 10 5d. 40 10s. = cost at 0 0 10 20 = cost at 1 8d. = cost at £3629 16 3 = cost at £3 14 81

EXAMPLE 3 .- Find the price of 729 days' work at £1 7s. 11d. per day.

OPERATION. 2729 0 0 = price at £1 182 5 0 = price at 0 1s. 8d. 5d. 60 15 0 = price at 15 3 9 = price at 15 21 = price at 0 0 01 £987 18 111 = price at £1 7 11

EXAMPLE 4.—What is the cost of 624 bush. 1 pk. 1 gal. 3 gt. of wheat at \$2.874 per bushel?

OPERATION. 50 cts. |1 \$1248 = price of 624 bush, at \$2.00 25 ots. 312 = price at 156 = price 121 cts. at 78 = price 121 \$1794 = price of 624 bush. at \$2.871

fact that the latter is still in common use in Great Britain, several of the States of the American Union, &c. The aliquot parts of the new owt.. of 100 lbs. are the same as the aliquot parts of \$1.

10

11

12.

13,

14

15.

16.

17.

18.

19.

20.

21.

22.

23.

24.

25.

```
1 pk. | 1 | 32.87
               = price of 1 bush.
          '717 = price of 1 pk.
2 qt.
          '35 | 6 = price of 1 gal.
          '1732 = price of 2 qt.
1 qt. 2
          0863 = \text{price of 1 qt.}
```

\$1.3449 = price of 1 pk. 1 gal. 3 qt. n \$1794 = price of 624 bushels at \$2.871 per bushel, 1.3412 = price of 1 pk. 1 gal. 3 qt. at \$2.871 per bush. Then \$1794

\$1795.3449 = price of 624 bush. 1 pk. 1 gal. 3 qt. at \$2.871 per bush.

EXAMPLE 5.—What is the price of 96 acres 1 rood 141 per. at £7 11s. 5id. per acre?

£726 18 = price of 96 acres at £7 11 5}

```
1 rood | 1 | £7 11 5
            1 17 101+1
                           = price of 1 rood.
               = price of 10 perches.
= price of 4 perches.
 per.
                  51 + 320 = price of 1 perch.
```

22 11 7 + 360 f. = price of 1 rd. 142. per. at 27 118.54d. per ac. £726 18 = price of 96 acres.

Ans. 2729 9s. 7d.  $+\frac{1}{320}$  f. = price of 96 acres 1 rood 144 per.

EXAMPLE 6.—What is the cost of 9641 square yards of plastering at 221 cents per square yard?

```
20 cts. | }
                        964
                    $192.80 = cost of 964 yds. at 20 cts. 225×11 = 16; cents. 24.10 = cost of 964 yds. at 2; cts. 225×15 half of the cost of 964 yds. at 2; cts.
21 cts. 1
```

\$216.90 = cost of 964 yds, at 221 cts. '16' = cost of + of a yd. at 22' cts

Ans. \$217'06; = cost of 964; yds. at 22; cts. per yd.

### EXERCISE 90.

1. Required the value of 92647 lbs. of tea at 35 cents per lb. Ans. \$32426.45.

the set when I for the thing is the fact that the

2. What is the cost of 94937 pails at 1s. 5d. each? with the transfer of the Ans. £6724 148, 1d.

ART. 8.]

ECT. VI.

3. What is the worth of 95972 boxes at 71 cents?

Ans. \$7197.90.

4. What is the cost of 62 acres at \$28.80 per acre?

Ans. \$1785.60.

- 5. Find the price of 2310 lbs. at 321 cents per lb. Ans. \$750.75.
- 6. Find the price of 2117 bags at 371 cents each. Ans. \$793.871.
- 7. Find the price of 7506 pair of shoes at 1s. 93d. a pair.

  Ans. £680 4s. 74d.
- 8. What is the value of 1217 lbs. of coffee at 171 cents. per lb?

  Ans. \$212.971.
- 9. Find the price of 2103 cords of wood at \$3.071 per cord.
- 10. What is the cost of 2096 oz. of gold dust at £3 18s. 101d. per oz.?

  Ans. £8266 2s. 0d.
- 11. Required the value of 6 oz. 18 dwt. 20 grs. of silver at \$1.55 per oz.

  Ans. \$10.7523.
- 12. What is the cost of 98 yds. 3 qrs. 1 na. of cloth at £1 15s. per yard?

  Ans. £172 18s. 5\fmathref{1}d.
- 13. What is the rent of 344 acres 3 roods 15 per. at £4 1s. 1d. per acre?

  Ans. £1398 1s. 034d.
- 14. What is the price of 5 oz. 6 dwt. 17 grs. of mercury at 5s. 10d. per oz.?

  Ans. £1 11s. 123d.
- 15. Find the price of 4 yards 2 qrs. 3 nails of satin at £1 2s. 4d. per yard.

  Ans. £5 4s. 84d.
- 16. Find the price of 32 acres 1 rood 14 perches at £1 16s. per acre.

  Ans. £58 4s. 1 d.
- 17. Find the price of 3 gals. 5 pts. of spirits of wine at 7s. 6d. per gallon.

  Ans. £1 7s. 21d.
- 18. How much will 724 bushels of apples come to at \$1.671 per bushel?

  Ans. \$1212.70.
- 19. What is the cost of 721 bush, of wheat at \$1.93‡ per bush.?

  Ans. \$1396.93‡.
- 20. What is the cost of 4514 rods of fencing at £2 17s. 7id. per rod?

  Ans. £13005 19s. 3d.
- 21. What is the price of 37493 acres at £3 15s. 6d. per acre?

  Ans. £14153 17s. 92d.

Allowing 112 lbs. to the cwt., find the value of-

22. 17 cwt. 1 qr. 17 lbs. at £1 4s. 9d. per cwt.

Ans. £21 10s. 837 d.

- 23. 75 cwt. 3 qrs. 12 lbs. at \$11.55 per cwt. Ans. \$910.80.
- 24. 20 tons 19 cwt. 3 qrs. 27 lbs. at £10 10s. per ton?

  Ans. £220 9s. 11 d. nearly.

25. 219 tons 16 cwt. 3 qrs. at \$45.50 per ton. Ans. \$10002.60

per bush.

per. at

d. per ac.

of plas-

16} eents.

er lb. 2426·45.

14s, 1d.

Mı

19

19

8

5

21

22

Mr.

9

13 114 14

121 118

#### EXERCISE 91.

### BILLS OF PARCELS.

(No. 1.)

QUEBEO, 16th April, 1859.

The state of the fifth of

Mr. JOHN DAY.

Bought of RICHARD JONES. 301

s. d. £ s. d. 6 per yard, 10 2 6 15 yards of fine broadcloth, at..... 13 24 yards of superfine ditto, at ..... 18 9 22 10 0 66 27 yards of yard wide ditto, at..... 8 4 11 5 70 44

5 0 D 46 12 yards of serge, at ..... 2 10 1 14 " 2 13 32 yards of shalloon, at..... 1 8

Ans. £53 4 10

### (No. 2.)

MONTREAL, 24th June, 1859.

Mr. JAMES PAUL,

Bought of THOMAS NORTON.

9 pair of worsted stockings, at.... 4 6 per pair, a barrie 

19 yards of flannel, at ...... 1 71 per yard,

Ans. £23 15

Salas (No. 3.)

TORONTO, 10th July, 1859.

Mr. WM. FILBERT.

A SANGIETONIA

Bought of GRORGE PRICE.

75] lbs. of sugar, at..... 7] cents per lb., W. L. L. W. 93 63 lbs. of tea, at ..... 126 lbs. of butter, at ..... 13 354 lbs. of raisins, at..... 17 lbs. of sago, at...... ·9 : 17 / 16:

23 lbs. of rice, at..... 581 lbs. of starch, at.....

Ans. \$105.02}

Mr J

1859.

4 10

1859.

## 5 tes (No. 4.)

HAMILTON, 12th August, 1859.

Mr. John James,

### Bought of James THOMAS.

11	S ots.	7.
198 Sangster's National Arithmetic, at	0.60	+
197 Robertson's Philosophy of Grammar, at	0.50	
83 Hodgins' Geography, at	1.00	THE L
57 Sangster's Algebraic Formula, at	0.124	, 2 7 1
217 Strachan's Canadian Penmanship, at	0.371	543
143 Hodgins' Geography of British Provinces, at	0.45	-
227 Sangster's Elementary Arithmetic, at	0.30	

Ans. \$521.25

### (No. 6.)

NIAGABA, 17th September, 1859.

### Mr. ALEX. LEITH,

### Bought of LAWRENCE MERCER.

01		2.70	0.	. 3
94 yards of flow	, atvered ditto, at	18	a per yaru	7 2
114 yards of lust	tring, at	6 1	0 4 44	4
	cade, at		3 - " "	- Ann
	n, at		8 . W. 48 Sin.	1
	ret, at		O High	

Ans. £44 15 10

### (No. 6.)

KINGSTON, 11th July, 1859.

### Dr. ALEX. HAMILTON,

是是"五人"

### Bought of TIMOTHY PASTLE.

14	OE.	ipecacuanna, at			. DO.04
23	"	laudanum, at			0.89
17	88	emetic tartar, at			1.25
25	66	cantharides, at		1. 180	2.17
27	"	gum mastic, at			0.61
56	. 66	gum camphor at	- da-1*	20 38	. 0.27

Ano. \$136.04

5.02

1859.

6. W 9. V 10. V 11. W

10.-I

11. I

12. T

### (No. 7.)

LONDON, C. W., 1st May, 1859.

### Mr. JAS. GREY.

### Bought of MICHAEL LEWIS.

THE TANK OF THE PARTY OF THE PA	8 0	
151 lbs. of currants, at	0 4	per lb.,
174 lbs. of Malaga raisins, at		
193 lbs. of sun raisins, at		
17 lbs. of rice, at		A alleria
81 lbs. of pepper, at	1	3 1
3 loaves of sugar, weight 321	1bs., at 0 8	4
13 oz. of cloves, at		per oz.

Ans. £3 13

### TARE AND TRET.

4. Tare and Tret is the name given to a rule by means of which merchants calculate the amount of certain allowances which were formerly made in buying and selling goods by weight in large quantities. They were as follows:

Tret, an allowance for waste in weighing.

2. Tare, an allowance for the actual or supposed weight of the box, bag, barrel, &c., containing the goods. And

3. Cloff, an allowance of 2 lbs. in every 336 for the

turn of the scale in retailing goods.

Of these the only one known in Canada is Tare; and as this is always set down in full in the invoice, Tare and Tret, as a rule, has no existence in Canadian mercantile transactions, and has therefore been altogether omitted.

### QUESTIONS TO BE ANSWERED BY THE PUPIL.

NOTE.—The numbers after the questions refer to the articles of the section.

1. What is Practice? (1)
2. Why is it so called? (1)
3. Of what rule is Practice merely a modification? (1)
4. What would be the general statement of a question in Practice? (1)
5. How is the process for finding the price of a number of articles simplified by Practice? (2)
6. What is an aliquot part? (3)
7. What are the aliquot parts of a dollar? (3)

strought har bush and asset where 6. What are the aliquot parts of a year? (3)
9. What are the aliquot parts of a month? (3)
10. What are the aliquot parts of a £? (3)
11. What are the aliquot parts of a shilling? (3)

12. What are the aliquot parts of a cwt. (112 lbs.) ? (3)

### EXERCISE 92.

#### MISCELLANEOUS EXERCISE.

(On preceding Rules.)

1. Take the number 70204, and by removing the decimal point of (1) multiply it by 100000; (2) divide it by 10000; (3) make it thousandths; (4) make it tenths of billionths; (5) make it tenths; and (6) make it hundredths of billionths.

2. Divide 427·1 by .0000637.

- 3. What will 19 tons 19 cwt. 3 qrs. 271 lbs. of hops cost, at £19 19s. 112d per ton?
- 4. Add together 73.723, 11.342, 16.713, 19.034, 713.213437, and

12.345678.

5. Of the ratios 5: 7, 9: 12, 12: 17, and 7: 10, point out (1) which is greatest, (2) which is least, (3) what is the ratio compounded of these?

6. If 1 acre of land cost \$80.50, what will 25 acres, 2 roods,

35 rods cost?

7. What is the G. C. M. of 144, 485, and 63.

8. What is the price of 7439 cords of wood at \$3.682 a cord?

9. Reduce 135775, 714335, 183376, and 1933 to their lowest terms.

10. If 341 bushels of turnips are worth 17 bushels of potatoes, and 9 bushels of potatoes 591 lbs. of tea, and 6 lbs. of tea 111 stone of flour, and 13 stone of flour \$3.60, and 38 cents pay for 12 lbs. of bread; how many bushels of turnips are worth 119 lbs of bread?

11. If 27 men in 7 days, working 8 hours a day, paint 42 floors, each 20 feet long and 16 feet wide, with 3 coats of paint to each; in how many days, of 11 hours each, will 54 men paint 77 floors, each 24 feet long and 22 feet wide, giving

each 5 coats of paint?

12. Take the number 7449164 and by removing the decimal point, make it (1) One hundred thousand times greater.

(2) One million times less.

(3) Hundredths of quadrillionths.

(4) Thousandths.

(5) Tenths of billionths.

(6) Tenths.

13. Reduce 72342 nonary to equivalent expressions in the duodenary, senary, and ternary scales, and prove the results by reducing all four numbers to the decimal scale.

eans

859.

rtain sells fol-

posed the

the

nd as Tret, rans-

of the

impli-

- 14. Express in the decimal scale the greatest and least numbers that can be formed with six digits in the binary, quaternary, senary, octenary, and duodenary scales.
- 15. Write down all the divisors of 1728.
- 16. What is the l. c. m. of the first fifteen even numbers, 2, 4, 6, 8, &c.?
- 17. From 97.91342 take 18.1234567.
- 18. What would be the cost of painting a ceiling 20 ft. 7 in. long and 19 ft. 5in. 7" wide, at \$2.87\frac{1}{2} per square yard?
- Divide 916 acres, 3 roods, 17 per., 7 yards, by 43 acres, 1 rood,
   2 per., 17 yds.

### SECTION VII.

# PERCENTAGE, COMMISSSION, BROKERAGE, STOCKS, INSURANCE, CUSTOM-HOUSE BUSINESS, ASSESSMENT.

1. The term Per Cent. is derived from the Latin word per, "by" or "for" and centum, "a hundred," and means "for a hundred." The term is usually employed to indicate the allowance paid for the use of money, but may also be used to express so much the hundred units of any other quantity.

Thus, the term 5 per cent. on so many dollars, gallons, miles, days, &c., signifies \$5 on every \$100, or 5 gallons on every 100 gallons, or 5 miles on every 100 miles, or 5 days on every 100 days, &c.

2. When the rate per cent. is known, the rate per unit is easily obtained by dividing the rate per cent. by 100.

Thus, 1 per cent. is equal to  $\frac{1}{100}$  or '01 per unit.

2 per cent. is equal to  $\frac{1}{100}$  or '02 per unit.

7 per cent. is equal to  $\frac{1}{100}$  or '07 per unit.

9 per cent. is equal to  $\frac{1}{100}$  or '09 per unit.

10 per cent, is equal to  $\frac{1}{100}$  or '10 per unit.

18 per cent. is equal to  $\frac{1}{100}$  or '18 per unit.

39 per cent. is equal to  $\frac{1}{100}$  or '39 per unit.

95 per cent. is equal to  $\frac{1}{100}$  or '95 per unit.

125 per cent. is equal to  $\frac{1}{100}$  or '78 per unit.

bers rary.

VII.

4, 6,

7 in. ood,

KS,

rord ansi ndialso

ther

A Alle &c. ZAX

unit

per cent, is equal to  $\frac{1}{100}$  or '005 per unit.  $\frac{1}{2}$  per cent. is equal to  $\frac{3}{100}$  or '0025 per unit.  $\frac{2}{3}$  per cent, is equal to  $\frac{2}{100}$  or '0075 per unit. per cent. is equal to 100 or 00125 per unit. 6t per cent, is equal to 6t or '065 per unit, &c.

1. What rate per unit is equivalent to 1.6 per cent., 11 per cent., 17 per cent., 63 per cent.?

What rate per unit is equivalent to 6 per cent., 25 per cent.

137 per cent.?

3. What rate per unit is equivalent to 81 per cent., 91 per cent.,

4. What rate per unit is equivalent to } per cent., } per cent.,

82 per cent. ?

6. At 61 per cent., how much is it for 1? Ans. . 0625. 6. At 183 per cent., how much is it for 17 Ans. . 186. 7. At 235 per cent., how much is it for 17 Ans. .23625.

8. At 2.734 per cent., how much is it for 17 Jins. .02734. 9. At 82.7 per cent.; how much is it for 1? Ans. 827.

10. At 19 per cent., how much is it for 1?

Ans. . 193.

### 8. To find the percentage of any given number-

### 

Multiply the given number by the rate per unit expressed decimally, and point off the product as directed in Art. 63, Sec. II.

EXAMPLE 1.-What is 7 per cent. on \$673.93?

#### OPERATIONAL PLANS AND

EXPLANATION.—7 per cent. is equivalent to '07 per unit; or, in other words, the percentage on each dollar is 7 cents. It is obvious then that the percentage on the whole sum will be as many times 7 cents as the sum contains dollars; that is '07×673-93.

Example 2.—What is 61 per cent. on \$2934?

Ans. \$2934×.065=\$190.71.

Example 3.—What is 47? per cent. on 7893 gallons of molasses? Ans. 7893 gal. x 4775=3768:9075 gallons.

#### EXERCISE 94.

1. What is 5 per cent. of \$742.10? 2. What is 11 per cent. of \$1000?

Ane. \$37.104. Ans. \$110.

3. Fow much is 10 per cent. of \$734-19?

Ano. 673-419.

per

- 4. How much is 874 per cent. of \$1624.50 ? Ans. \$1421.4375.
- 5. What is 121 per cent. on \$994.70? Ans.\$124.3375.
- 6. What is 82 per cent. on \$777.50? Ans. \$68.03.
- 7. What is 21 per cent. of \$7135.80? Ans. \$160.5555.
- 8. A merchant imports 2740 boxes of oranges, and finds, upon receiving them, that 20 per cent. of the whole quantity are decayed. To how many boxes was his loss equivalent?

  Ans. 548 boxes.
- 9. A gentleman purchases a farm for \$7490, agreeing to pay 10 per cent. down, 17 per cent. at the end of the first year, 27 per cent. at the end of the second year, and 46 per cent. at the end of the third year. What is the amount of each payment?

  Ans. \$749 down.

\$1273 30 at the end of 1st year. \$2022 30 at the end of 2nd year. \$3445 40 at the end of 3rd year.

- 10. What is the difference between 41 per cent. of \$740 and 21; per cent. of \$1680?
- 11. If I purchase 729 gallons of brandy and lose 11 per cent. by leakage, &c., how much have I remaining?
- 12. Add together 25 per cent. of \$763.22, 16 per cent. of \$847.16, and 64 per cent. of \$1284.17.
- 13. A person dying leaves an estate worth \$17429.40 to be divided among his three sons. The eldest is to receive 43 per cent. of the whole, the second 37 per cent. of the whole, and the youngest son the remainder; what is the share of each?

Ans. The eldest receives \$7494.64, the second \$6448.87;

- 14. A merchant purchases vinegar to the amount of 63978 gallons, and finds, upon receiving it, that 36 per cent. had leaked away. What was his loss?

  20. 24832 08 gallons.
- 15. A brick kiln contains 29800 bricks, and it is found after burning that 17 per cent. of the entire quantity are worthless; how many good bricks were there in the kiln?

men wire on a liveron i ampili man te con

### COMMISSION.

4. Commission is the percentage charged by agents, or commission merchants, for their services in purchasing or selling goods, collecting bills, &c.

The person who buys or sells goods for another is called an Agent, a Commission Merchant, a Factor, or a Correspondent.

5.

5. 3. 5.

n

re

s.

ay ir, it.

ch

ar.

Ir.

21

10.

by.

18.

16,

5.

be .

43

he

he

14,

18.

ed

er

h-

s,

g

ir.

5. To find the commission on any sum at a given rate per cent. is simply to find the percentage on that sum, and the rul imployed is the same as that in Art. 3, viz:

Multiply the given amount by the rate per unit expressed deci-

EXAMPLE 1.—What is the commission on \$790.80 at 3 per cent.?

Ans.  $$790.80 \times .03 = $23.724$ .

EXAMPLE 2.—A commission merchant sells goods to the amount of \$7982.75; what is his commission at 21 per cent.?

Ans. \$7982.75 × .0275 = \$219.525625.

### I had govern of the Exercise 95. 1 . . Price will .0

- 1. What is the commission on \$1000 at 41 per cent.? Ans. \$45.
- 2. What is the commission on \$1678.30 at 21 per cent.?

  Ans. \$37.76175.
- 3. What is the commission on \$7531:19 at 32 per cent.?; 600013
- 4. Find the commission on \$508.60 at 11 per cent. The inner of the state of the sta
- 5. Find the commission on \$7863.50 at 12 per cent.? And the
- Ans. \$137.61125.

  6. An agent collects debts to the amount of \$878.30; what is his commission at 21 per cent.?

  Ans. \$21.9578.
- 7. A correspondent purchases teas for me to the amount of \$7193.16; what have I to pay him for commission at 31 per cent.?

  Ans. \$224.78625.
- 8. A commission merchant sells goods to the amount of \$6734.10; what is his commission at 17 per cent.?

  Ans. \$1144.797.
- 9. An agent sells 718 barrels of flour at \$7.13 a barrel; what is his commission at 41 per cent.?

  Ans. \$217:57195,
- 10. A commission merchant disposes of 8243 bushels of wheat at \$1.85 per bushel; what is the amount of his commission at 57 per cent.?

  Ans. \$857.7871875.

### BROKERAGE.

6. Brokerage is the percentage charged by money dealers, called Brokers, for negotiating notes, mortgages, bills of exchange, &c., or for buying or selling stocks, &c.

7. Brokerage is merely another name for commission, and is computed by the same rule.

£

pa

in

### Exercise 96.

- 1. What is the brokerage on \$7893-87 at 2 per cent.?

  Ans. \$167-8774.
- 2. What is the brokerage on \$8000 at } per cent.? Ans. \$70.
- What is the brokerage on \$8643-22 at 11 per cent.?
   Ans. \$108-04025.
- 4. What is the brokerage on \$78963.80 at I per cent.?

  Ans. \$690.93325.
- 5. What is the brokerage on \$1987.27 at 31 per cent?
  Ans. \$74.522625.
- 8. Commission and Brokerage should both be computed on the amount of money collected or invested.

For example: If I receive \$10000 to invest and charge 5 percent, my brokerage would be \$500 if I invested the whole \$10000; but if, as is usually the case, I am requested to deduct, from the amount sent, my brokerage or commission, and invest the remainder, it would obviously be unjust to charge commission on the whole amount,—i. e., on the sum invested and also on the sum I retain for commission. Hence, in all cases, the sum actually expended is the proper basis upon which to compute the commission, brokerage, &c.

9. To compute commission or brokerage when it is to be deducted in advance from a given amount, and the balance invested:

#### BULE.

- 1. Divide the given amount by \$1, plus the commission on \$1, and the result will be the sum to be invested.
- 2. Subtract the part to be invested from the given amount, and the remainder will be the commission or brokerage.

Example.—A correspondent receives \$16782, with instructions to deduct his commission at 3½ per cent., and invest the belance in sugar at 9½ cents per pound. How much sugar does he ship to his employer, and what is his commission?

#### OPERATION,

\$16782 - 1 035 = \$16214'40275 = sum to be invested. \$16782 - \$16214'40275 = \$567.50725 = commission. \$16214'40275 - 91 cents = 170678'871 lbs. Ans.

EXPLANATION.—The commission on \$1, at the rate of \$1 per cent, is \$0.000. Hence, for every time he receives \$1.005, he keeps \$0.000 for commission and invests \$1. It is plain, then, that if we divide the given amount, \$16783, by \$1.085, or in other words, find how often the latter sum is contained in the former, we shall find how often he invests \$1; i.e. how many dollars he invests.

M.

74:

70.

25:

25.

25.

per:

ole

ct,

est

is-

lso a

the.

m-

32 15

to'

he.

BI,

ınd

IC-

the ·

Deg

The work may be proved by finding the commission on the sum invested (Art. 5), and comparing it with the commission as found by deducting the sum invested from the whole sum sent. If these are equal, the work is correct.

### Exercise 97.

- 1. An agent receives \$4000, with instructions to purchase Great Western Railway stock. After deducting his brokerage at 11 per cent., how much money had he to invest and what was his brokerage?

  Ans. Invested \$3950-61728.

  Commission \$49-38271.
- 2. A merchant sends his agent \$7500, with instructions to deduct his commission at 41 per cent., and purchase laces with the remainder. What is the commission, and what sum was expended in laces?

  Ans. Commission \$322.96651.
- Invested \$7177:03349.

  3. A commission merchant receives \$8470, with instructions to purchase the best brand of Canadian superfine flour at \$6.40 per barrel. He is to receive out of this sum 5 per cent. on the amount he invests. How many barrels of flour does he purchase?

  Ans. 1260-5 barrels.
- 4. A broker receives \$11000, with instructions to invest it in Bank stock—deducting his brokerage at 7 per cent. What sum had he to invest?

  Ans. \$10904.584882.
- b. If I remit to my agent \$13000, instructing him to purchase broad cloth at \$3.68 per yard, and he keeps 41 per cent. on the sum invested, for commission; how much cloth does he send me, and what is his commission?

Ans. 3427-0499 yds. of cloth. \$559-8086 commission.

### STOCK.

10. Stock is a term used to denote the Copital of moneyed institutions, as Banks, Railroad Companies, Gas Companies, Insurance Companies, Manufactories, &c.

11. Stock is usually divided into portions of \$100 or £100 each, called shares, and the different individuals owning these are called shareholders or stockholders.

- 12. The Association of Shareholders, is called a Company or Corporation; and the Act of Parliament specifying their corporate powers, rights, and privileges is called a charter.
- 13. The nominal or par value of a share is its original cost of valuation.

14. The market or real value of a share is the sum for which it can be sold.

15. The rise and fall in the value of Stock is reckoned

at a certain per cent. on its nominal or par value.

16. When stocks sell for their original cost or valuation, they are said to be at par; when they sell for more than their original valuation, they are said to be at a premium or advance, or above par; when they do not bring their original cost or valuation, they are said to be at a discount, or below par.

Nors.—Par is a Latin word, and means equal or a state of equality. Stock is at par when a hundred-dollar share sells for \$100; it is above par when it brings more than \$100, and below par when it will not bring as

much as \$100.

17. Persons who deal in stocks are called stock-brokers or stock-jobbers.

18. To find how much stock either above or below par

a given sum will purchase:-

#### BULE.

Divide the given amount by the worth of \$1 stock, and the result will be the stock required.

Example 1.—How much stock at 10 per cent below par can be purchased for \$25000 ? .... Ans. \$25000 ÷ 0.90 = \$27777.773.

BEYLANATION.—When stock is 10 per cent. below par, each share of \$100 sells for only \$90, i. e. \$90 money will purchase \$100 stock, therefore \$0.90 money will purchase \$1 stock and the given sum will purchase \$1 stock as often as it, (the given sum) contains \$0.90.

Example 2.—How much stock at 15 per cent. premium may be purchased for \$7000?

Ans. \$7000 ÷ 1.15 = \$6086.9565.

EXPLANATION.—When stock is 15 per cent. above par, it requires \$115 money to purchase \$100 stock, or \$1.15 money to purchase \$1 stock. Hence if we divide the whole sum to be invested by the value of \$1 stock, it is evident we must get the amount of stock produced.

EXAMPLE 3.—I own \$16400 stock of the Bank of Montreal, and sell out at 13 per cent premium. What do I receive?

Ans. \$16400 × 1.13 = \$18532.

EXPLANATION.—Each \$100 stock brings me \$113 money, or \$1 stock brings \$1:13 money, therefore \$16400 stock must bring \$16400 x 1:13 money.

EXPLANATION.—Each \$100 stock brings me \$113 money.

1. A person has \$9000 which he wishes to invest in Grand Trunk Railway shares, then selling at 17 per cent. discount, what amount of stock can he purchase?

Ans. \$10843.373.

2. If I invest \$8500 in Upper Canada Bank stock, which is selling 11 per cent. above par, what amount of stock do I receive?

for

oned

alua-

more

at a

bring

at a

cality. de par ing as

okers

par

d the

r can

7.778.

hare of

ase \$1

n may

9565.

Hence k, it is

itreal, 18532.

1.stock

money.

Grand

count,

3.373.

is sell-

ceive ?

3. If I remit to my agent \$17500, with instructions to deduct his brokerage at 1½ per cent., and invest the remainder in Great Western Railroad stock, then selling at 7 per cent, premium, what amount of stock do I receive.

Ans. \$16153.22.

4. If I receive \$20000, with instructions to deduct my commission at 12 per cent., and invest the balance in stock, which is then seiling at 3 per cent. discount, what amount of stock do I remit to my employer?

Ans. \$20263-987.

5. Mr. A. owns 200 shares in the Canada Life Assurance Company. The par value is \$100 a share, the stock at a premium of 51 per cent.; if I purchase it through a broker who charges me 7 per cent. for the transaction; how much do my 200 shares cost me.

Ans. \$21284.625.

#### INSURANCE.

- 19. Insurance is a written agreement by which an individual or an incorporated company becomes bound, in consideration of a certain sum paid in advance, to exempt the owners of certain kinds of property, as houses, household furniture, merchandise, ships, &c., from loss by fire, hipwreck, or other calamity.
- 20. The Written Instrument, or contract between the parties, is called a Policy of Insurance.
- 21. The sum paid for the insurance is called the *Premium*, and is usually a certain per cent. on the sum for which the property is insured.
- 22. Houses, merchandize, furniture, &c., are usually insured against risk of fire for the year, or other specified time.

Note.—The rate of insurance on dwelling houses, stores, goods, household furniture, &c., varies from ; to 2 per cent. per annum, on the sum insured according to the character and position of the tenement; vessels are insured for the voyage or the year.

23. To compute the premium for insurance for 1 year, or a specified time, we use the same rule as for Commission or Brokerage.

EXAMPLE.—If I insure my house and furniture for \$7389, at the rate of 1½ per cent. per annum, what premium must I pay yearly?

Ans. \$7389 × 0125 = \$92.3625.

EXPLANATION.—1; per cent., i. e. \$1:25 per \$100, is equal to \$0.0125 per dollar. The premium, therefore will be as many times \$0.0125 as the sum insured contains \$1; i. e. the premium will be 0.0125 × 7389.

tembel of vacioty is before

### than & niver ningua off the L. Brancis 99.

- 1. What is the premium for insurance on \$7500, at 12 per cent.? Ans. \$131.25.
- 2. What is the premium for insurance on \$8375, at 2 per. cent.? dom'n proude n Ans. \$62.8125.
- 3. What is the premium for insurance on \$6000, at 14 per cent.? Ans. \$112.50.
- 4. What is the premium for insurance on \$5000 at \$1.17 per cent. (i. e. per \$100) ? Ans. \$58.50.
- 5. What is the premium for insurance on \$6400, at \$0.90 per go (Scent. ? ... Ans. \$57.60.
- 6. What is the premium for insurance on \$4500, at \$0.35 per Ans. \$15.75.
  - 7. What premium must I pay for insuring a cargo of flour worth \$36000, from Quebec to Liverpool, at \$3 per cent.?
  - Ans. \$1080. 8. A firm, owning four steamers running on lake Ontario, effect an insurance with a company in Toronto to the amount of \$27000 on each, paying \$4.82 per cent. (i. e.  $4\frac{83}{100}$  per cent.) What is the total premium on the four steamers?
  - Ans. \$5205.60. 9. What is the annual premium on an insurance for \$39000, at 21 per cent.?
  - A farmer insures his barns and their contents to the amount of \$17800. What premium does he pay at | per cent.
- 11. A vessel running between Hamilton and Oswego is insured for \$12350, at the rate of 13 per cent. per month. To what loes the premium of insurance amount for 7 months, beginning with the 10th of April and ending with the 10th of Ans. \$1235. November?
- 24. To find what sum must be insured on property so that, if destroyed, its value and the premium may both be recovered: of constraint and antique.

Divide the value of the property by \$1, minus the premium on a \$1 at the given rate per cent.

In case of the property of \$1, minus the premium on a \$1 at the given rate per cent.

Example 1 -A ship-owner wishes to insure a vessel valued at \$17450, so that if it be wrecked he may recover both the value of the vessel and the premium. In order to do so, for what sum must be insure, at \$4.60 per cent.? Ans. \$17450 - 964 = \$18291.40461. REPLANATION.—If I insure goods to the value of \$100, at 46 per cent, and they are destroyed, I receive only \$95'40 towards my loss, since I paid \$4'60 for insurance; that is, for every \$1 of my loss I receive \$0'954. Since, then, the recovery of \$0'954 requires \$1 to be insured, the recovery of \$1'2450 will require as many dollars to: be insured as \$0'954 is contained times in \$17450.

PROOF. \$18301.40461 > '046=\$841.40461 = the premium, and \$18301.46461 = \$841.40461 = \$17450 = value of the vessel.

Example 3.—What sum must be insured on a house valued at \$6000, at 3 per cent so that in case of fire the value of both premium and property may be secured?

Ans. \$6000 ÷ .97 = \$6185.567.

EXPLANATION.—For every dollar I lose (taking premium into account) I receive 97 cents, that is, in order to receive 97 cents, I must insure for \$1, and in order to receive \$6000, without any loss, I must insure for \$6000 ÷ 97 = \$6185.567.

### Exercise 100.

- 1. For what sum must I insure a cargo valued at \$17000, so that in case the whole is lost I may recover both the value of the property and the premium of 31 per cent.?
- 2. For what sum must I insure on \$22750 in order to cover both the premium of 6 per cent. and the value of the property insured?

  Ans. \$24202:127.
- What sum must be insured at 21 per cent. on property worth \$15000 so that the owner may be secured against all loss?

  Ans. \$15345.2685.
- 4. A steamer worth \$33000 is insured at 51 per cent for such a sum, that in case of its becoming a total wreck, the owners recover both the worth of the yessel, and the prenium of insurance. For what sum is it insured?

Ans. \$35013 2625

### CUSTOM HOUSE BUSINESS.

25. All goods coming into Canada from Foreign countries are required by law to be landed at certain places or ports called *Ports of Entry*.

26. At every Port of Entry in Canada, the Government has an establishment called a Custom House, with one or more officers attached to it, called Custom-House Officers.

27. A certain charge called a Duty, fixed by Act of Parliament, is made upon nearly all goods entering Canada from Foreign countries.

28. It is the business of the Custom-House Officers to inspect the cargoes of all vessels entering at any of these

Q

25. t.? 25.

11.7

TI.

it.? 50. per 50.

per 60. per 75.

75. rth

80. ect of

60. 00, 58.

89. red hat

of 35.

be

ued

the for

61.

ports, to examine the invoice of goods, collect the duties,

&c., &c.

29. Besides the duties on merchandize, all vessels engaged in commerce are required to pay certain charges for the privilege of entering the port, &c.; these charges are called harbor dues.

30. The duties levied by law on goods imported into

Canada are of two kinds:

1st. Specific duties. 2nd. Ad Valorem duties.

31. A specific duty is a certain sum levied on the ton. owt., lb., gallon, square yard, &c., of a particular kind of merchandise, as so much per square yard on woollens, fannels or cloths, so much per lb. on tea, so much per gallon on brandy, wine, &c.

32. An ad valorem duty is a certain percentage on the actual cost of the goods in the country in which the

were purchased.

Thus an ad valorem duty of 10 per cent. on satin purchased in France is a charge for duty of 10 per cent. of the sum the invoice of satin cost in

Nore 1.—The term ad valorem is from the Latin; and means according

to the value, i.e., when the value.

Nors 2.—An invoice is a written statement of the goods, showing the quantity of each sort and its value or price.

23. In the United States Custom Houses certain legal allowances are made for draft, tare, leakage, &c., before specific duties are imposed. In Canada, however, as before remarked, (Art. 4, Sect. VI.,) these are not known, the tare being found by actually weighing one or more of the boxes, &c., containing the goods, and the leakage by guaging the cask.

Nors.—At present (1859) the various kinds of spirits are the only articles upon which specific duties are charged by the Canadian Tariff.

184. To calculate the specific duty on an invoice of

n AULD offern mining A

sponet) the tare, leakage, &c., and multiply the remainder by the given duty per gallon, lb., yard, &c.

At 41 cents per lb. what is the specific duty on There of cottes weighing 73 lbs., each, allowing 4 lbs. per 100 or tare?

1000

T

to Van

n,

of

3. 1. 11

no

ev

ade .c

the

gal

ore

28

m,

by

on.

100

of

931

# A CA TO THE OPERATION. THE TENT OF THE PARTY OF

78 × 7 = 511 lbs. = gross weight. 511 × 04 = 2011 lbs. = tare.

40014 = net at 41 cents per lb. = 49014 × 41 = \$20.8468. = 4ms.

Example 2.—What is the specific duty on 10 chests of tea, the net weight 783 lbs., at 11 cents per lb.?

## OPERATION.

 $783 \times 11 = 8613 \text{ cents} = $86.13. Ans.$ 

### Exercise 101.

- 1. What is the specific duty, at 3½ cents per lb., on 5 hhis. of sugar, each weighing 1347 lbs., allowing tare 6 lbs. per 100?

  Ans. \$221.58.
- 2. What is the specific duty, at \$1.20 per 100 lbs., on 11 bags of rice, each weighing 127 lbs., allowing 3 lbs. per 100 for tare?

  Ans. \$16.26.
- 3. What is the specific duty, at 13 cents per gallon, on 129 gallons of oil?

  Ans. \$16.77.
- 4. What is the specific duty, at 52 cents per lb., on 207 drums of figs, each weighing 31 lbs., allowing 21 lbs. a drum for tare?

  Ans. \$342-1968.
- 5. What is the specific duty, at 47 cents per yard, on 214 yards of black silk velvet?
  Ans. \$100.58.
- 35. To find the ad valorem duty on an invoice of merchandise:—

#### RULE

Multiply the value of the goods at the place in which they were purchased by the per cent. charged, expressed decimally, and the result will be the duty required.

Example. 1—What is the ad valorem duty, at 27 per cent. on an invoice of brandy which cost \$7493.70?

#### OPERATION.

# \$7493.70 × .27 = \$2023.299, Ans.

EXAMPLE. 2. What is the ad valorem duty, at 19 per cent. on a quantity of broadcloth which cost \$4116.40 ?

#### OPERATION.

\$4116.40 × ·19 = \$782.116. Ans.

#### EXERCISE 102.

- 1. What is the ad valorem duty, at 21 per cent. on an invoice of silks which cost \$17429.80?

  Ans. \$3660.2580.
- 2. What is the sh valorem duty, at 71 per cent. on 40 boxes of ten which gost \$2920-16?

  Ans. \$219-012.
- 3. What is the ad valorem duty, at 25 p r cent., on an invoice of jewellery which cost \$71342:90?

  Ans. \$17835-725.

4. What is the ad valorem duty, at 20 per cent., on an invoice of boots and shoes which cost \$913.73? Ans. \$182.746.

5. What is the ad valorem duty at 33 per cent, on an invoice of French silks which cost \$14713.19.7 Ans. \$4855.3527.

# ASSESSMENT OF TAXES.

36. A tax is a certain sum required to be raised by a municipality for local improvement, payment of officers, and other general purposes. Theis collected from each

citizen in proportion to the value of his property.

37. In levying taxes the first thing to be done is to make a complete inventory of the value of all the property in the city, town, township, &c., in which the tax is to be raised. This inventory is made by officers called Assessors sondinted by the municipality.

38. To calculate the amount of taxes any one indivi-

dual has to pay:-

Divide the whole sum to be levied by the whole value of rateable property in the town, township, &c.: the quotient will be the sum to be paid on each dollar.

Multiply the rate per dollar by the amount of the person's prop-

erty, and the product will be the amount of his tax.

EXAMPLE—A certain township requires to raise the sum of \$14729.00 for general purposes; the whole amount of rateable property in the municipality being set down at \$2743500, what proportion must I bear if my property is assessed at \$7490.00.

#### OPERATION.

\$14729 ÷ \$2743500 = \$0.005368 = rate per dollar. \$0.005368 × 7490 = \$40.20632. Ahs.

# Exercise 103.

1. The assessment rolls of a town show the value of the rateable property to be \$7142300. A tax of \$23900 is to be levied for general purposes; how much is my proportion, my prop-Ans. \$49.2878. erty being set down at \$14729.50.

2. A tax of \$100000 is to be levied on a county having rateable property to the value of \$5793000; what is the amount

horne by A, whose property is valued at \$18600?

Ans. \$321 9732.

3: In the last examply what would be the amount of B's tax, the value of his property being \$7500? Ans. \$129:465.

4. In the same example what would be the amount of O's tax, his property being assessed at \$11400. And \$186:7868.

ic.

16.

lce

7 8 TS. ch

The Miles to cty be

ors

VI-

ıtethe

op-

um ble hat D. Kalitik

ble

ied op-78. ble unt 32.

SI,

65.

ex, 88.

# QUESTIONS TO BE ANSWERED BY THE PUPIL.

NOTE.—The numerals after the questions refer to the numbered articles of the section.

1. What is the meaning and derivation of the term per cent.? (1)
2. When the rate per cent. is known, how is the rate per unit obtains 1? (2)
3. How do we ascertain the percentage on any given number? (3)

What is commission ? (4)

4. What is commission? (4)
6. What is the person who sells goods for another called? (4)
6. How do we find the commission on any given sum? (5)
7. What is brokerage? (6)
8. How is the brokerage on any sum computed? (7)
9. Upon what sum should commission and brokerage be computed? (8)

10. Explain this by an example.

11. How do we compute commission or brokerage when it is to be deducted in advance from a given amount, and the balance invested? (9)

12. How is this rule proved? (9)

13. What is understood by the term Stock? (10)

14. How is Stock usually divided? (11)

15. What is meant by the terms Shareholders, Corporation, and Charter?

(11 and 12).

16: What do you understand by the nominal or par value of Stock? (18)

17: What is meant by the market or real value of Stock? (14)

18. When is Stock said to be at par? when at a premium or above par?

and when at a discount or below par? (16)

19. What is the meaning of the term par? (16 note.)

20. What are persons who deal in Stocks called? (17)

21. When Stock is either above or below par, how do we find how much of it a given sum will purchase? (18)

22. What is Insurance? (19)
23. What is a Policy of Insurance? (20)
24. What is meant by the Premium of Insurance? (21)

25. For what length of time is property usually insured? (22)

22. For what length of time is property usually insured? (22)
26. How do we compute the premium of insurance on any amount of goods, property, &c.? (23)
27. How do we compute the amount for which we must insure in order to cover both the value of the property and the premium paid? (24)
28. How may the truth of this rule be proved? (24)
29. What are Ports of Entry? (25)
30. What is the duty of Custom-House Officers? (28)
21. What are duties? (27)
32. What are harborduse? (20)

32. What are harbor dues? (29)

33. What different kinds of duties are levied on goods in Canada? (30)

The wine the test of the second relative delice in the

I wish of the real of the state of the state

e natebalkovski skore, i slove se se koje koje se provinski provinski provinski provinski provinski provinski

84. What are specific duties? (31)
35. What is an ad valorem duty? (32)

36. What is the meaning of the term ad valorem? (82)

37. What is an invoice? (82)

38. What is the rule for computing specific duties? (34)
39. What is the rule for calculating ad valorem duties? (35)
40. What is a tax? (36)
41. How are taxes imposed? (9 and 38)

# SECTION VIII.

# INTEREST, DISJOUNT, EQUATION OF PAYMENTS, AND PARTNERSHIP.

1. Interest is the sum allowed for the use of money, and is usually reckoned at a certain rate per cent. per annum; that is, so many pounds for the use of £100 for one year, so many dollars for the use of \$100 for one year, &c.

Note.—The term per cent. means per hundred; per annum means per year.

2. Interest differs from Commission, Brokerage, &c., in that the latter are computed at a certain per cent. without regard to time, while interest is calculated at a certain rate per cent. for one year, and consequently for longer and shorter periods in like proportion.

3. The Principal is the sum lent.

4. The Rate per cent. is the sum paid for the use of each hundred dollars, pounds, &c.

5. The Rate per unit is the sum paid for the use of each dollar, pound, &c.

6. The Interest is the whole sum received for the use

of the principal.

7. The Amount is the sum obtained by adding together the principal and the interest.

Thus, if I lend \$200 for a year, on the agreement that I am to receive interest at the rate of 7 per cent. (per annum, understood), at the end of the year I receive back the \$200, and in addition \$14 for interest. Here, \$200 to the principal.

7.00 is the rate per cent.
0.07 is the rate per unit.
14.00 is the interest.
214.00 is the amount = principal + interest.

8. Interest is either Simple or Compound.

9. Money is lent at Simple Interest when the interest is not added to the principal so as to hear interest.

Thus, if \$100 be lent at simple interest at 5 per cent., the *principal* remains unchanged, being always \$100, and the *interest* for each successive year is \$5.

10. Money is lent at Compound Interest when the interest, as it falls due from time to time, is added to the principal; the sum thus obtained constituting a new principal for the ensuing year, half year, quarter, &c., as the case may be.

Thus: if \$100 oe lent at 5 per cent. per annum compound interest, the principal charges at the end of each year; being \$100 for the first year. 5105 (i. e. former principal-its interest) for the second, \$110°25 for the third, do. The interest is consequently \$5 for the first year, \$5°25 for the second, \$5°5125 for the third, do.

## SIMPLE INTEREST.

11. Questions in Interest are dependent on Proportion, and may all readily be solved by one or more statements in the Bule of Three; but in order to deduce special rules, we shall represent the different quantities by their initial letters, and thus obtain a series of algebraic formulæ, which translated, become the common arithmetical rules for interest.

It is to be presumed that the pupil has made sufficient progress in Algebra before he arrives at this point, to readily understand what follows. The operations involved are of the simplest kind, and may without difficulty be comprehended, even by those wholly ignorant in Algebra. The only part, however, absolutely necessary for working any problem in interest, is the interpretation of the formula, i. e. the arithmetical rule, and this we have always appended. A glance at the formule and the corresponding rules will show how much less labor is necessary to remember the former than the latter;—and indeed the pupil should be required to deduce from time to time ary formula he may find it necessary to use.

Note.—When two or more letters are written together thus, prt. the meaning is that the values of these letters are to be maltiplied together. Thus, Prt means that the value of P is to be multiplied by the value of r, and that by the value of t.

When letters are written in the form of a fraction, thus  $\frac{A-P}{Pr}$  the meaning is the same as in common arithmetical fractions; i. e., that the part constituting the numerator is to be divided by the part constituting the denominator.

Thus,  $\frac{A-P}{Pr}$  means that the value of P is to be subtracted from the value of A, and this difference is to be divided by the value of P multiplied by the value of r.

ney,

ND

for ear.

per

, in tout tain ger

e of

e of

use

her

eive id of lere,

rest

l re-

yes

fine

inte

cent

inte

H  $\mathbf{T}\mathbf{h}$ Th

give

IN

the 1 E

13 p

Th

22 da

give

IN cipa

E

3 pe

He

No

He

12. Let P = Principal, I = Interest, A = Amount, r = rate per unit, and t = time (i.e., number of years).

$$I = Prt (I.)$$

$$P = \frac{I}{rt}(II.)$$

$$t = \frac{I}{Pr}(II.)$$

$$A = P (1+rt) (V.)$$

$$P = \frac{A}{1+rt} (VI.)$$

$$I = \frac{A+P}{1+rt} (VI.)$$

大宗 - (IX:)

 $r = \overline{t_n}(X)$ n=ir+1 (XI.) Then because t interest of 01 for 1 year, and t = number of years, rt = interest of 01 for the given time, and Prt = interest of given principal for given time and at given rate. Therefore I = Prt and dividing each of these equals, 1st by rt, and by Pt, and St by Pr, we get formulas (II.) (III.) and (IV.) in the marrin. (III.) and (IV.) in the margin.

(III) and (IV.) in the margin.

Again, because rb = interest of \$1 at given rate and for given time, 1+rt = the amount of \$1 at given rate and time, and P times 1+rt, that is. P (1+rt) = amount of given principal at the given rate and time. Therefore A = P (1+rt), which is formula (V.) in the margin, and dividing each of these equals by 1+rt, we get formula (VI.) in the margin. Taking (V.) and actually multiplying as indicated, the part within the brackets by P, we get A = P + Prt; and subtracting P from each of these, we get A - P = Prt. Dividing these equals, 1st by Pt and 2nd by Pr, we get formulas (VII.) and (VIII.) in the margin. Taking, if we are required to find in what time any sum of money will amount to any given number of times itself at a given rate per cent, or, in other words, in what time any principal where n simply stands for the required number of times, we have in formula (VIII.) in the margin. A P P P  $t = P_r$ because the amount is to be nP; and dividing both numerator and denomi-

\*\*P; and dividing both numerator and denominator of this fraction by P; we get formula (IX.) in the margin, multiplying (IX.) by t we get tr=
\*\*-1: and dividing these equals by t, we get formula (X.); and, again, adding 1 to each of these same equals, we get formula (XI.)

# APPLICATIONS.

13. When the principal, rate per cent., and time are given, to find the interest-

Rule I = Prt (i.)

INTERPRETATION .- The interest is found by multiplying the principal by the rate per unit, and the resulting product by the time.

Example. What is the interest on \$342.20 for 7 years at 8 per cent. ?

Sparie to to the operation. Here  $P = $342 \cdot 20$ ,  $r = \cdot 08$ , and t = 7. Then  $I = Prt = $342 \cdot 20 \times \cdot 08 \times 7 = $191 \cdot 682$ . Ans.

14. When the interest, rate per cent., and time are given to find the principal

We had of the diese of in RULE. P=rt (ii.)

INTERPRETATION .- The principal is found by dividing the interest by the product of the rate per unit and the time,

III.

mit,

the pal

rt.

ate

l at t is, the

rt), ling

ıula ally

the

ub-

Prt.

gin. ime

4611

nt.,

168,

be

mi-

X.) or.

989

re

the

ze.

8

Example.—What principal will give \$207.50 interest in 64 years at 41 per cent.?

OPERATION.

Here I = \$207.50, t = 6.5, and r = .0475.

Then 
$$P = \frac{I}{rt} = \frac{$207.50}{6.5 \times .0475} = \frac{$207.50}{.30875} = $672.664$$
. Ans.

15. When the interest, principal, and time are given to find the rate per cent.—

子のは、よりないがくまというとなる

Rule. 
$$r = \frac{I}{Pt}$$
 (iii.)

INTERPRETATION.—The rate per unit is found by dividing the interest by the product of the principal and time, and the rate per cent. is found from the rate per unit by multiplying the latter by 100.

EXAMPLE.—At what rate per cent. will \$729.18 give \$109.11 interest in 9 years?

OPERATION.

Here  $P = $729^{\circ}18$ ,  $I = $109^{\circ}11$ , and t = 9. 109.11

109.11 Then  $r = \frac{1}{Pt} = \frac{1}{729.18 \times 9} = \frac{1}{6562.62}$ = 0.01662 = rate per unit.

Therefore the rate per cent. =  $0.01662 \times 100 = 1.662 = 11$  nearly. Ans.

16. When the interest, principal, and rate per cent. are given, to find the time-

Rule. 
$$t = \frac{\pi}{12r}$$
 (iv.)

INTERPRETATION. - The time is found by dividing the interest by the product of the principal and rate per unit.

EXAMPLE.—In what time will \$850 give \$89:75 interest, at 13 per cent.?

Here P = \$850, I = \$89.75, and r = 13. Then  $t = \frac{I}{Pr} = \frac{89.75}{850 \times 13} = \frac{89.75}{110.5} = \frac{897.5}{1105} = 0.812217$  years = 9 months, 22 days.

17. When the principal, rate per cent, and time are given, to find the amount-

Rule. 
$$A = P(1+rt)$$
 (v.)

INTERPRETATION.—The amount is found by multiplying the principal by the amount of \$1 for the given rate and time.

EXAMPLE.—To what sum will \$789.80 amount in 11 years, at 3 per cent.?

OPERATION.

Here P = \$789.80, r = 03, and t = 11. Then  $A = P(1+it) = $789.80 \times 1.38 = 1050.434$ . Ans. Note. -(1+rt) in this question =1+3×11=1+38=1-88.

134 3

any

I

15115 B

elev

per

amo

R

I

fron amo E

to 2

amo

R

I

E in 2

tiply

18. When the amount, rate per cent., and time are given, to find the principal—

Rule. 
$$P = \frac{A}{1+rt}$$
(vi.)

INTERPRETATION.—The principal is found by dividing the given amount by the amount of \$1 for the given time at the given rate.

EXAMPLE.—What principal put to interest at 71 per cent. will amount to \$2000 in 8 years?

Here 
$$A = $2000$$
,  $r = .075$  and  $t = 8$ .  
Then  $P = \frac{A}{1+rt} = \frac{2000}{1.60} = \frac{20000}{16} = $1250$ . Ans.

19. When the amount, principal, and time are given, to

Rule. 
$$r = \frac{A-P}{Pt}$$
 (vii.)

INTERPRETATION.—The rate per unit is found by subtracting the principal from the amount, and dividing the difference by the principal multiplied by the time. The rate per ce t is found by multiplying the rate per unit by 100.

Example.—At what rate per cent. will \$730 amount to \$2783.80 in 23 years?

#### OPERATION.

Here 
$$A = \$2783^{\circ}80$$
,  $P = \$730$  and  $t = 23$ .  
Then  $r = \frac{A-P}{Pt} = \frac{\$2783^{\circ}80 - \$730}{\$730 \times 28} = \frac{\$2053^{\circ}80}{\$16790} = 1223 = \text{rate per unit.}$ 

Hence rate per cent. = 1823 = 121 nearly.

20. When the amount, principal, and rate per cent. are given, to find the time—

Rule. 
$$t = \frac{A-P}{Pr}$$
 (viii.)

INTERPRETATION.—The time is found by subtracting the principal from the amount, and dividing the difference by the principal multiplied by the rate per unit.

EXAMPLE.—In what time will \$666.33 amount to \$983.73 at 12 per cent.?

Here 
$$A = \$963.73$$
,  $P = \$666.33$  and  $r = 12$ .  
Then  $t = \frac{A - P}{Pr} = \frac{983.73 - 866.53}{666.33 \times 13} = \frac{317.40}{79.9596} = \frac{5174000}{79.9596} = 3.9695$  years = 3 years 11 months 19 days. Ans.

BOY. VIII.

me are FILL >

ths given he given

B, and per cent. s sonis

given, to

TOP TOP

btracting ce by the found by

lount to

er unit.

cent. are

he principrincipal

183:73 at

years = 8

21. To find the time in which any sum will amount to any given number of times itself at a given rate per cent-

Rule. 
$$t = \frac{n-1}{r}$$
 (ix.)

/ ARTS. 18-28.]

INTERPRETATION.—To find the time in which a given sum will amount to n times itself at a given rate per cent., subtract 1 from n, and divide the remainder by the rate per unit.

EXAMPLE 1.—In what time will any sum of money amount to eleven times itself at 8 per cent. ?

Here 
$$t = \frac{11}{r}$$
 and  $r = \frac{108}{108} = \frac{1000}{108} = \frac{1000}{125}$  years, 4ns.

Example 2.—In what time will \$67.83 quadruple itself at 41 per cent. ?

OPERATION.

Here n = 4, since the money is to quadruple itself, and r = .0475. Then  $t = \frac{n-1}{r} = \frac{4-1}{.0475} = \frac{8}{.0475} = \frac{30000}{.475} = 63.157$  years. Ans.

22. To find the rate per cent. at which any sum will amount to a given number of times itself in a given time-

Rule. 
$$r = \frac{n-1}{4}$$
 (x.)

INTERPRETATION.—The rate per unit is found by subtracting 1 from n, the number of times itself to which the given principal is to amount, and dividing the remainder by the given number of years.

Example.—At what rate per cent. will a given sum amount to 25 times itself in 72 years?

OPERATION.

Here 
$$n = 25$$
,  $t = 72$ .

Then  $r = \frac{n-1}{t} = \frac{25-1}{72} = \frac{24}{72} = \frac{1}{3} = 33\frac{1}{3} = \text{rate per unit.}$ 

Hence rate per cent. =  $33\frac{1}{3}$ . Ans.

23. To find to how many times itself a given sum will amount in a given time at a given rate per cent.

Rule. 
$$n = tr + 1$$
. (xi.)

INTERPRETATION. - The number of times, or n, is found by multiplying the time by the rate per unit, and adding 1 to the product.

EXAMPLE.—To how many times itself will four cents amount in 20 years at 17 per cent.?

H

annu

ing:

Di

Ex

years

7 year

mont

7 year

T

1. F

2. B

3. I

4. 7

5. Y

6. T

7. H

8. H

9. F

10. F

25.

month cent.

per an

1000

num

Ex

COPERATION. THE STATE OF THE STATE OF Here t=20 and r=17Then'n = tr + 1 = 20 x, 17 + 1 = 34 + 1 = 44 = 47 times itself. Ans. Exercise 104.

1. What is the interest on \$723.19 for 7.32 years at 8.7 per cent.? Ans. \$354.6813086.

2. To what sum will \$857.19 amount in 61 years at 61 per cent? Ans. \$1219.352775.

3. To how many times itself will £2 19s. 9 d. amount in 11 years at 721 per cent.?

Ans. 8.975, or nearly 9 times.

4. In what time will \$654.32 give \$234.56 interest at 7 per

Ans. 5.12112, or 5 years 1 m. 13 days.

5. At what rate per cent. will \$700 amount to \$1200 in 5 years? Ans. 149 per cent.

6 In what time will any sum of money quadruple itself at 23 per cent. Ans. 13 years 15 days.

7. Find the time in which \$270 will give \$87 interest, at 7 per Ans. 4 years 7 months.

8. To what sum will \$680 amount in 111 years, at 11 per cent? Ans. \$1540.20.

9. What principal will amount to \$2000 in 20 years, at 8 per Ans. \$769.23 14.

10. At what rate per cent. will any sum of money amount to 21 times itself in 24 years? Ans. 831 per cent.

11. In what time will a given sum of money amount to 23 times itself, at 16 per cent.? Ans. 1371 years.

12. Find the interest on \$679.18 at 72 per cent., for 11.73 years. Ans. \$617.4255.

At what rate per cent. will \$950 amount to \$1763.42 in 10 Ans. 8:562 per cent., or rather over 81 per cent.

14. In what time will \$666 amount to \$1347.50, at 6 per cent.? Ans. 17:054 + years, or 17 years 19 days.

15. In what time will \$278 give \$100 interest, at 9 per cent.?

Ans. 4 years 25 days. 16. At what rate per cent. will \$476.30 amount to \$500 in 2 years? Ans. 218 per cent.

17. At what rate per cent. will \$749.49 give \$257 interest in 7 Ans. 4.898 per cent.

18. What principal will amount to \$1111 11 in 11 years, at 11 per cent.? Ans. \$502.7647.

19. Find the interest on £187.47, at 11 per cent. for 9 years. Ans. £165 158. 10 88d.

# SPECIAL RULES.

24. The interest of \$100 at 6 per cent, for one year, is \$6; hence the interest on \$1 at 6 percent, for one year, is \$0.06, and for two months it is } of \$0.06; i.e., 1 cent. 77。14年,15年的中华企业中有

Hence, to find the interest of \$1, at 6 per cent. per annum for any number of months, we deduce the following:—

RULE

Divide the number of months by 2, and call the quotient cents.

EXAMPLE 1.—What is the interest of \$1 at 6 per cent. for 7 years and 9 months?

OPERATION.

7 years and 9 months, = 93 months, and 98+2 = 464 cents = \$0'465. Ans.

EXAMPLE 2.—Find the interest on \$72.93 for 7 years and 8 months at 6 per cent.

OPERATION.

7 years 8 mo. = 92 months, half of 92 = 46 cents = interest of \$1 for given rate and time.

Then \$0'48×79'98 = \$33'5478. Ans.

## EXERCISE 105.

1. Find the interest on \$1 for 11 months at 6 per cent.

Aire. 51 cents.

2. Find the interest on \$1 for 16 months at 6 per cent.

Ans. \$0.08, or 8 cents.

3. Find the interest on \$1 for 9 years 8 months at 6 per cent.

Ans. \$0.58.

4. What is the interest on \$1 for 16 yrs. 3 months at 6 per cent.?

Ans. \$0.971.05. Whatis the interest on \$1 for 11 yrs. 7 months at 6 per cent.?

Ans. \$0:695.

6. What is the interest on \$1 for 12 yrs. 5 months at 6 per cent.?

Ans. \$0.745.

7. Find the interest on \$279.40 for 3 yrs. 2 mo's. at 6 per cent.

Ans. \$53.086.

8. Find the interest on \$189.70 for 6 yrs. 7 mo's. at 6 per cent.

Ans. \$74.9315.

9. Find the interest on \$1463 for 3 yrs. 11 mo's. at 6 per cent.

Ans. \$343.805.

10. Find the interest on \$28967.50 for 11 years 1 month at 6 per cent.

Ans. \$19263.3875.

25. Since in computing interest the month is taken as 30 days, two months will contain 60 days, and, by Art. 24, the interest on \$1 at 6 per cent. for 2 months or 60 days is one cent, the interest on \$1 at 6 per cent. per annum, for 6 days, will therefore be 10 of one cent; i. e. one mill or 1000 of \$1.

Hence, to find the interest on \$1 at 6 per cent. per annum for days, we have the following:

er cent.? 3813036.

If. Ano.

er cent? 852775. nt in 11

9 times. at 7 per 13 days. 100 in 5

per cent. elf at 23 15 days.

months. per cent?

at 8 per 169·23 / s. int to 21 per cent.

23 times 71 years. 73 years.

17.4255. 42 in 10 per cent. r cent.?

19 days. cent. ? 25 days. 500 in 2

per cent. rest in 7 per cent.

rs, at 11 02-7647. 7647s.

ce the in-

# S RULE.

Call one-sixth of the number of days mills or thousandths of a dollar.

Example.—What is the interest on \$1 at 6 per cent. for 16 days?

#### OPERATION.

### 16-: 6=2} mills=\$0.0026. Ans.

# EXERCISE 106.

- 1. What is the interest on \$1 for 2 days at 6 per cent.?

  Ans. \$0.0003.
- 2. What is the interest on \$1 for 7 days at 6 per cent.?
- Ans. \$0.0011.
- 3. What is the interest on \$1 for 11 days at 6 per cent.?
- 4. What is the interest on \$1 for 27 days at 6 per cent.?
- Ans. \$0.004}

  5. What is the interest on \$1 for 47 days at 6 per cent.?
- 6. Required the interest on \$1 for 8 months 12 days at 6 per
- cent. So·042.
- 7. Required the interest on \$1 for 66 days at 6 per cent.
- 8. Required the interest on \$1 for 2 years 2 months 19 days
- 9. Find the interest on \$1 for 7 years 8 months 9 days at 6 per cent.

  Ans. \$0.1331...

  Ans. \$0.4611...
- 10. What is the interest on \$1 for 17 years 11 months 23 days at 6 per cent.?
- 11. Required the interest on \$1 for 12 years 7 months 17 days at 6 per cent. Ans. \$0.757%.
- 26. To find the interest on any sum of money at 6 per cent. per annum for any time:—

#### RULI

Find the interest on \$1 for the given time, by Arts. 24 and 25, and multiply this by the given principal.

EXAMPLE.—What is the interest on \$763:20 at 6 per cent. for 6 years 7 months and 26 days?

The

1. Fi

2. Fi

3. R

4. R

6. Fi

7. Fi

8. W m 9. W

10. Fi

11. To at 12. To

27 per ce

đ

Find per cen

Then
part of
per ans

togethe

This is the method in common use for computing interest for days; but, since it considers the year as containing only 360 days instead of 365, the result is too large by \$55, or \$15 of itself. Hence, when perfect accuracy is desired, the interest for the days when obtained by the rule must be diminished by \$15 part of itself.

be retai

VIII.

0003. 8

001∦. ∧

D018.3

004

0074.

6 per

·042.

0.11.0

1332.01

days

at 6

4611.

days

days

7578.

6 per

t. for

BOCHmust

0784.

# s for the Agree of the a coperation.

SIMPLE INTEREST.

Interest on \$1 for 6 years 7 months

Thorefore interest on \$1 for 6 yrs. 7 months 26 days \$0 3991 Then\* \$0'3991 × 763'20=\$304'7712. Ans.

### EXERCISE 107.

- 1. Find the interest on \$917.30 for 7 months 17 days at 6 per Ans. \$34.704516.
- 2. Find the interest on \$842.50 for 3 months 13 days at 6 per Ans. \$14.462916.
- 3. Required the interest on \$573.83 at 6 per cent. for 2 years Ans. \$101.3766. 11 months 10 days.
- 4. Required the interest on \$642.30 at 6 per cent. for 6 years 9 months 19 days. Ans. \$262-16545.
- 5. Required the interest on \$1427.871 at 6 per cent. for 5 years 5 months 7 days. Ans. \$465:7252.
- 6. Find the interest on \$709.63 for 4 years 7 months 16 days at 6 per cent. Ans. \$197.040596.
- 7. Find the amount of \$2463.20 at 6 per cent. for 7 years 7 months 22 days. Ans. \$3592.9877.
- 8. What is the interest on \$999.99 at 6 per cent. for 9 years 9 months 9 days? Ans. \$586.494135.
- 9. What is the interest on \$68.70 for 3 years 4 months 27 days. at 6 per cent. ? Ans. \$14.04915.
- 10. Find the interest on \$742.63 at 6 per cent. for 3 years 28 days. Ans. \$137.139.
- 11. To what sum will \$200 amount in 7 years 4 months 11 days at 6 per cent.? Ans. \$288.366.
- 12. To what sum will \$743.63 amount in 9 years 3 months 9 days at 6 per cent.? Ans. \$1157-460095.
- 27. To find the interest on any sum at any other rate per cent. for any given time:-

Find the interest on the given principal for the given time at 6 per cent, by Art. 26.

Then add to or subtract from this interest such a fractional part of itself as the given rate exceeds or falls short of 6 per cent.

The amount is obtained by adding the interest and the principal together.

<sup>\*</sup> In order to obtain the correct answer, this fraction when it occurs must be retained in the form of a vulgar fraction; and in that case it is better to make the interest of \$1 for the given time the multiplier.

Fo

EXAMPLE.—What is the interest on \$450 for 3 years 6 months 11 days at 8 per cent. ?

# OPERATION.

Interest on \$1 at 6 per cent, for given time=\$0'2116. Interest on \$450 at 6 per cent, for given time=\$0.2112 × 450=\$95.825.

Hence interest on \$450 at 8 per cent, for given time=\$95'325+one third

of \$95.325=\$127.10. Ans.

Note.—Since \$6 = 6 + 2 = 6 + \( \frac{1}{2} \) of 6 we find the interest at 6 per cent., and increase it by one third of itself for the interest at 8 per cent. So for interest at 9 per cent., we should find the interest at 6 per cent., and increase it by one-half of itself; for 7 per cent., increase the interest at 6 per cent by one-sixth; at 14 per cent, double the interest at 6 per cent., and increase it by 1 of the interest at 6 per cent.; at 5 per cent., find the interest at 6 per cent. and deduct one-sixth; at 41 per cent, find the interest at 6 per cent. rest at 6 per cent., and deduct one-fourth, &c., &c.

# EXERCISE 108.

- 1. Required the interest on \$1234.56 for 8 years 9 months 10 days at 7 per cent. Ans. \$758.5685.
- 2. Required the interest on \$9876.54 for 2 years 1 month 11 days at 3 per cent. Ans. \$626 337245.
- 3. Required the interest on \$715.30 for 3 years 7 months 10. days at 8 per cent. Ans. \$206.6422.
- 4. To what sum will \$555.55 amount in 2 years 4 months 8 Ans. \$712.58546. days at 12 per cent.?
- 5. To what sum will \$7766.55 amount in 100 days at 5 per Ans. \$7874.41875.
- 6. To what sum will \$500 amount in 8 years 8 months 8 days at 16 per cent.? Ans. \$1195.111.
- What is the interest on \$576 for 3 years 5 months 7 days. Ans. \$98.96. at 5 per cent.?
- 8: What is the interest on \$2478.91 for 2 years 6 months 11 days at 41 per cent.? Ans. \$282.285.
- 9: What is the interest on \$780 from May 9, to December 11, Ans. \$28.08. at 6 per cent. ?
- 10. What is the interest on a note of \$1830:63 from August 16, 1851, to June 19, 1852, at 7 per cent.? Ans. \$109.63439.
- 11. What is the amount of a note of \$6200 from Sept. 3, 1858, to January 9, 1859, at 6 per cent.? Ans. \$6332.266.

# PARTIAL PAYMENTS.

28. To compute the interest, on notes or bonds, when partial payments have been made:

If the interest be paid by days: Multiply the sum by the number of days which have elapsed before any payment was made. Subtract the first payment, and multiply

VIII.

onths

e third

r cent.. ent er cent., er cent.,

find the he inte-

nths 10 8.5685. onth 11 337245.

nths 10 06.6422. onths 8

2.58546. at 5 per 4.41875.

s 8 days 195.111. s 7 days

\$98.96. onths 11

282.285. mber 11.

\$28.08. gust 16. 9.63439.

3, 1858, 332·266.

is, when

ad before multiply

the remainder by the number of days which passed between the first and second payments. Subtract the second payment, and multiply this remainder by the number of days which passed between the second and third payments. Subtract the third payment; &c.

Add all the products together, and find the interest of their sum

for one day.

If the interest is to be paid by the week or month, substitute weeks or months for days, in the above rule.

EXAMPLE.—How much principal and interest have I to pay on the following note on the 10th November, 1859?

TORONTO, 18th October, 1858.

For value received, I promise to pay to Timothy Thomas, or order, the sum of six hundred and twenty dollars, on demand, with interest at 6 per cent.

THOMAS WILLIAMS.

The following endorsements were made on this note:

1858.—November 25th, there was endorsed \$ 47.50 " December 28th, 166 

1859.—February 11th 44 June 6th, 160.10 September 2nd, " " 183-25

# tono avil- " COPERATION.

From 18th October to 25th November there are 38 days.

" 25th Nov. to 28th December " 33 "
25th Nov. to 28th December " 35 "
28th Dec. to 11th February " 45 "
11th February to 6th June " 115 "
6th June to 2nd September " 88 "
2nd September to 10th Nov. 69 "
Whole sum \$620,00 for 85 days \$23560,00 for 1 day.

First endorsement 47'50

Balance \$572.50 for 33 days \$18892.50 for 1 day. Second endorsement 108'93

Balance \$463.57 for 45 days=\$20860.65 for 1 day. Third endorsement 216'18

Balance \$247:39 for 115 days=\$28449.85 for 1 day. Fourth endorsement

Balance \$18729 for 88 days=\$16481 52 for 1 day. Fifth endorsement 183'25

Balance \$4.04 for 69 days 278.76 for 1 day.

Whole interest = that of \$108523 28, for 1 day. 

Balance on note . ..... 7406 Principal and interest due = \$21.8794

La Track Land Benevier a har

# EXERCISE 109.

1. What principal and interest was due on the following note on the 7th October, 1860?

GUELPH, June 2nd, 1859.

For value received, I promise to pay, on demand, to James George, or order, the sum of twelve hundred and seventeen dollars and thirty cents, with interest from date at 6 per cent.

JOSEPH JOHNS.

On this note there were endorsed the following payments:

1859 July 17th, received	1 \$207:80
Oct. 6th, "	209.60
" Dec. 11th, "	320.90
1860March 29th, "	421.83
Mark Mark Art Control of the Control	Ans. \$98.6816.
it is the set of freezeway the rest the reserve	" M 3 3 4 4 5 5 5 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6

2. What principal and interest was due on the following note on the 1st May, 1863?

PORT HOPE, June 17th, 1860.

For value received, I promise to pay, on demand, to Messrs. Henly & Jobson, or order, the sum of seven thousand, three hundred and forty-eight dollars and twenty-five cents, with interest from date at 8 per cent.

HENRY GOODPAY.

by

On this note there were endorsed the following payments:

1860	-Septembe	r 5th, receive	ed \$2463:8	30 = A = A0
16	Decembe	r 7th, 4	392-2	100 1112
	-June 11th		982.2	10 200
	February		2842-9	0
	December		317-2	13
1.40	P 100			02.1222

# COMPOUND INTEREST.

- 29. In the present article we shall merely take some of the simpler problems in Compound Interest, learning the full discussion of the rule until after the pupil is familiar with the use of Logarithms. (See Sect. XI.)
- 30. We have seen (Art. 10) that when money is lent at compound interest, the interest is added to the principal at the close of each period, and, with it, constitutes a new principal for the next term.

Hence to find the compound interest of any sum for any

given time at a given rate per cent :-

Find the interest on the given principal for one period, i. e., ONE YEAR, HALF YEAR, or QUARTER, as the case may be, and add it to the principal.

Then find the interest on this amount for the NEXT PERIOD and

add it to the principal used for that period, as before.

Proceed in this manner with each successive year or period of

the proposed time.

Then the last result will be the amount of the given principal, at the given rate, for the given time. Subtract the given principal from this, and the remainder will be the Compound Interest required.

Example.—What is the Compound Interest on \$1000 for 4 years at 5 per cent. per annum?

OPERATION.

\$1000.00 Principal. 50.00 Interest for 1st year.

\$1050.00 Amount for 1 year—principal for 2nd year.
52'50 Interest for 2nd year.

\$1102'50 Amount for 2 years—principal for 3rd year. 55'125 Interest for 3rd year.

\$1157.625 Amount for 3 years—principal for 4th year. 57.88125 Interest for 4th year.

\$1215'50625 Amount for 4 years. 1000.00 given Principal.

Ans. \$215.50625=Compound Interest required.

# Exercise 110.

1. What is the Compound Interest of \$1800 for 5 years at 6 per cent. per annum?

Ans. \$608.806.

2. What is the Compound Interest of \$700 for 3} years at 7 per cent. half-yearly?

Ans. \$424-040.

Nore.—Since the payments are made half-yearly, and bear interest at the rate of 7 per cent. per half year, we simply find the amount of the given principal at 7 per cent. for 7 payments.

3. What are the amount and Compound Interest of \$673.40 for 2 years at 3 per cent. quarterly?

Ans. \$353.0429 = Amount \$179.6429 = Interest.

4. What are the amount and Compound Interest of \$860 for 3 years at 4 per cent. half-yearly?

Ans. \$1088:1743 = Amount. \$228:1743 = Interest.

31. Compound Interest is most expeditiously calculated by the following—

ng note

1859. o James een dolent

HNS.

ents:

2 2749 T

ving note

1860. to Messrs.

hree hunth interest

oodpay. payments:

Ans. CLL

1333.

impler proble rule until ect. XI.)

is lent at rincipal at ates a new

m for any

3

OVO

S. 7.

fo

# TABLE

SHEWING THE AMOUNTS OF \$1 OB £1 AT COMPOUND INTEREST, FOR ANY NUMBER OF PAYMENTS FROM 1 TO 50.

Par	per	per cent.	per cent.	per cent.	No of Pay- meats.	per cent.	per cant.	per cent.	per cent.
		1 .04000				8 15659		S 55567	4.54938
		1 '03160				2 28120		8 73346	4 82235
199	3 1.0927		1 15,82			2 28793		3:92013	5.11169
		11 16996				2 35657		4 11614	5 41839
1	5 1.1592	9.122365	1 27028	r 39829	80	2 '42728	3 24340	4.32194	5 74340
	6 1 1940	1 26532	1 34010	1 41852	81		8 2/818	4.58804	6,'08810
Win er		7 1 31505				3 :24908		4:76494	6:45339
		7 1 36857					3 64888	9.00310	6 84059
-		7 1 42331					8 79432	5 25335	7 25102
, 1	0 11 3438	2 1 48024	1 413801	T AAAGS	35	2.31386	S *94609	2.21601	7 '68609
18	1 .0040	3 1 5394	1 .71084	1 .60660	38	0.00000	4 10393	5.79182	8 14725
		6 1 .60103					4 26809	6.08141	
	3 1.4685	3 1 66507	1 :88565	2 18293	38	3.07478		6.38548	
		91 78168					4 61637	6 70475	
		7 1 '80094					4'80102		10 28572
	6 1 6047	1 1 .8729	2 18287	2 84085	41	9 - 85000	4 99306	7-80180	10 90286
		5 1 9479					5 19278		11 '55708
		3 2 '0258					5 40040		12 25045
	9 1 7535	1 2 '1068	2 '52695	3 02560			5 '61651		12 93548
		1 2 1911	3 '65330	3 20718			5 '84118		18:76461
No.	1 10000	o oroni	o Poros	0.000	. 2 .			THE P. L	
		9 2 2787					6 07482		14 59049
		0 2 36925 9 2 46475					8 81782		15 46592
		9 2 5633							16:39387 17:37700
		8 2 6658							18 42515

32. To compute Compound Interest by the above

RULE

Find by the table the amount of \$1 for the given time and at the given rate.

Multiply the sum thus found by the given principal, and the result will be the required amount.

Subtract the principal from this amount, and the remainder will be the Compound Interest.

EXAMPLE.—What are no amount and Compound Interest of \$3400 at 5 per cent. The years?

OPERATION.

By the table the married of \$1 at 5 per cent, for 11 years \$2 07898.

Then 42 07893 × 24th 2 27068 565 = Amount.

2400 = Principal.

18:362 = Interest.

EXAMPLE.—What is the amount and compound interest of

APOUND S FROM

per cent.

67 4.54938 466 4.82235 013 5.11169 014 5.41839 194 5.74349

182 8 14725 141 8 63609 3548 9 15425 1475 9 70351 3999 10 28572

3426 14 59049 0597 15 46592 0127 16 39387 2133 17 37700 6740 18 42515

the above

ne and at the ind the result

mainder will

interest of

=\$2.07893

OPERATION.

We find by the table that

£47 10s. for 6 years at 3 per cent. half yearly?

21.42576 is the amount of 21 for the given time and rate.

267 7236 = 67 14 51 is the required amount.

47 10 0 is the given principal.

And £20 4 5 is the required interest.

# Exercise 111.

1. What are the amount and compound interest on \$875 for 11 and years at 6 per cent?

Ans. Amount = \$1661.0125.

Interest = \$786.0125.
2. What are the amount and compound interest on \$643.98 for

13 years at 4 per cent. half yearly?

Ans. Amount = \$1785.41523.

Interest = \$1141.43253.

3. What are the amount and compound interest of 1 cent at 6 per cent. per annum for 45 years? Ans. Amount = \$137646.

Interest = \$ 127646.

4. What are the amount and compound interest of \$78.20 for 7

1010 years at 3 per cent. quarterly? Ans. Amount = \$178.916.

Interest = \$100.716.

5. What are the amount and compound interest of \$777-77 for 9

Ans. Amount=\$1871.7968.

Interest = \$1094.0268.

6. What are the amount and compound interest of £44 5s. 9d.

over for 11 years at 6 per cent. per annum?

Ans. Amount = £84 1s. 5d. Interest = £39 15s. 8d.

7. What are the amount and compound interest of £32 4s. 91d. for 3 years at 4 per cent. half-yearly?

Ans. Amount = £40 15s. 104d. nearly.

Interest = £ 8 11s. 1d.

33. Given the amount, time and rate—to find the principal; that is, to find the present worth of any sum to be due bereafter—a certain rate of interest being allowed for the money now paid—

RITE.

Find by the Table the amount of \$1 at the given rate and for the given time; and divide it into the given amount. The quotient will be the principal.

AMO

1797

NES

fron

pay

cent

25500

Ter

中的小

PR

E

9 mc

Here

Then

1. W

8:10

4. W

6. If

88

E

EXAMPLE—What principal will amount to \$10000 in 12 years at 6 per cent. compound interest?

#### OPERATION.

Amount of \$1 for 12 years at six per cent. = \$20122. \$10000 ÷ 20122 = \$4969.684. Ans.

### EXERCISE 112.

1. What principal will amount to \$7439.87 in 7 years at 4 per cent. compound interest?

Ans. \$5653.697.

 What principal will amount to \$9193.90 in 20 years at 5 per cent. compound interest?

Ans. \$3465.081.

3. What ready money ought to be paid for a debt of £595 10s.

2 d. to be due 3 years hence, allowing 6 per cent. per annum compound interest?

Ans. £500.

4. What ready money ought to be paid for a debt of \$7111-11.
to be due 7 years hence, allowing 6 per cent. compound interest?

Ans. \$4729-295.

5. What principal, put to interest for 6 years, would amount to £268 0s. 41d. at 5 per cent. per annum?

Ans. £200.

# DISCOUNT.

34. Discount is an allowance made for payment of a debt before it is due.

35. The present worth of a debt payable at some future time, without interest, is that sum of money which, being put out at legal interest, will amount to the debt by the time it becomes due.

Thus, if I owe a man \$100 and give him a note for that amount, payable one year hence without interest, the present value of my note is less than \$100, since \$100 being put out at interest for 1 year at 6 per cent. will amount to \$106...

28. From Art. 18 it is evident that to find the present worth of a note, payable at some future time, without interest, is simply to find what principal, put to interest at the rate specified, will amount to the sum named on the face of the note in the given time; i.e. by the time the note becomes time.

Hence to find the present worth of any sum to be paid at some future time without interest, we have (Art. 18) the following:—

Rule. 
$$P = \frac{A}{1+rt}$$

INTERPRETATION.—The present worth is found by dividing the amount of the note, debt, &c., by the amount of \$1, at the specified rate per cent, for the given time.

1784 . Fare

ears

1 per .697. 5 per

.081. 108. ran-£500.

1.11. ound 293.

int to £200.

nt of

future being y the

nount, of my est for

a note, at prinnamed ecomes

e paid t. 18)

in: the perified

Norn .- The discount is found by deducting the present value from the note, debt, &c.

EXAMPLE 1.—What is the present value of a note for \$860 payable 3 years hence, allowing discount at the rate of 6 per cent. per annum?

OPERATION.

Here A = \$860, r = 08, and t = 3. Whence 1 + rt = 178. 860 =\$728.8124. Ans.

1+rt 1.18 PROOF.—Interest on \$720.813 for 3 years at 6 per cent. = \$131.1835.

Added principel..... = 728.8121.

EXAMPLE 2.—What is the discount on a note for \$728.63 due 9 months hence, allowing discount at 7 per cent. per annum?

OPERATION.

Here A = \$728.63, r = .07, and t = .75 year. Whence 1 + rt = 1.0325.

= \$692.285 present worth. 1+rt 1.0525

> Then amount on face of note...\$728.63 Present value...... 602'285

> > Discount ...... \$ 86'344 Ane.

# Exercise 113. A s. to & Controller . 1

1. What is the present worth of a note for \$962, payable in one year, at 4 per cent. discount?

2. What is the present worth of \$2202, payable in 5 years and 9 months, at 6 per cent. per annum discount?

Ans. \$1637:174. 3. What sum will discharge a debt of \$1003.50; to be due in 8 months hence, allowing 6 per cent. per annum discount? Therefore South Or & up god i the in Model four to Ans. \$964.9038.

4. What ready money will now pay a debt of \$716 due 7 months hence, allowing discount at 8 per cent.? Ana. \$684.0764.

5. What ready money will now may a debt of \$1342.50, due 125 days hence, at 61 per cent.?

Ans. \$1313 266.

6. If a legacy of \$2400 is left, to me on the 3rd of May, to be paid on the Christmas day following, what must I receive as present payment, allowing 5 per cent. per annum discount? augitefeld at Ans. \$2324:84.

7. Find the discount on a bill of \$2202 at 5 per cent., payable 9 months hence. Ans. \$79.59036.

8. What is the present worth of a note for \$4360, payable one year and 5 months hence, at 8 per cent.? Ans. \$4018-43317.

9. What is the present worth of a note for \$1647, due 11 mouths 

- 10. Required the present worth of a note for \$2000 due 3 years 7 months hence, at 6 per cent. Ans. \$1646.09053.
- 11. What is the discount on a note for \$2070.90, payable 1 year 7 months hence, at 5 per cent.?

  Ans. \$151.919.
  - 2. That is the present worth of a note of \$970.63, payable in 1 months at 8 per cent.?

    Ans. \$904.313.

Note.—When the payments are to be made at different times, find the present value of the sums separately; their sum will be the present value of the note, and, as before, this subtracted from the whole amount will give the discount.

- 13. What is the discount on \$3024, the one half payable in 5 and the remainder in 12 months, 7 per cent. per annum being allowed?

  Ans. \$150.0464.
- 14. A merchant owes \$440, payable in 20 months, and \$896, payable in 24 months; the first he pays in 5 months, and the second in one month after that. What did he pay, allowing 8 per cent. per annum?

  Ans. \$1200.

# BANK DISCOUNT.

- 87. Bank Discount is a charge made by a bank for the payment of money on a note before the note is due, and differs materially from discount as commonly calculated.
- 38. Banks consider the discount to be the same as the interest on the whole amount of the note, from the time it is discounted until the time it becomes due. Bank Discount is therefore greater than the true discount by the interest on the discount.
- 39. The three days of grace, which by mercantile usage, are allowed to clapse after a note falls due, before it is payable, are always included by banks in the time for which they calculate the discount,
  - 40. Two kinds of notes are discounted at banks:
- 1st. Business notes or business paper. These are notes actually given by one individual to another for property sold or value received.
- and commodation notes, called also accommodation paper. These are in and for the purpose of borrowing money from the banks.
  - 41. To find the bank discount on a note:-

#### BITT. W

Add 3 days to the time which the note has to run before it becomes due, and calculate the interest for this time at the given rate per cent.

Branch What is the bank discount on a note of \$700, payable in 69 days, allowing discount at 6 per cent.?

3 years

R-09053.

le 1 year 151 919.

ayable in

s, find the sent value nount will

in 5 and

um being

150.0464.

and \$896.

nths, and

#### OPERATION.

Here the time the note has to run is 72 days = 2 months 12 days.

Interest of \$1 at 6 per cent. for 2 months 12 days, is \$0.012.

Interest of \$700 at 6 per cent. for 2 months 12 days \$0.012 ×700=
\$5'40. Ans.

# Exercise 114.

1. What is the bank discount on a note for \$986, having 2 years and 3 months to run, allowing discount at 7 per cent.?

Ans. \$155.8701.

2. If I have a note for \$640, payable in 100 days, and get it discounted at the rate of 8 per cent. per annum, what discount am I charged?

Ans. \$14.6488.

3. I sell a horse and carriage for \$563.80, and receive a note for that sum, payable, without interest, 91 days hence. Now if I get this discounted at the rate of 6 per cent, per annum, what sum do I receive?

Ans. \$554.967.

42. It is often necessary to make a note of which the present value shall be a certain sum.

Thus, suppose I require to receive from the bank \$1000, and wish to give my note, payable in 7 months, at 6 per cent., what amount must I put on the face of the note?

New the interest on \$1 at 6 per cent, for 7 months and 3 days (i.e. days of grace) is \$0.0355, and this will be the bank discount on \$1 for 7 months at 6 per cent.

To get the present value of \$1, we subtract \$0.0355 from \$1, which gives us \$0.9645.

Hence, for every \$0.9645 I receive, I must put \$1 on the face of the note;

and therefore to receive \$1000, I must put 0'9645, i. e. \$1035'806 on the face of the note.

esent value.....\$1000.00

Hence to find the face of a note, due at some future time and discounted at a given rate per cent. per annum, that shall have a known present value, we have the following:—

e pay, als. \$1200.

due, and culated. me as the ne time it

Bank Disut by the

it is payfor which

tually given her. These banks.

th per cent.
of \$700,

ations, b

These examples are worked by the rule given in Arts. 26 and 37. If the absolutely correct answer is required, it must be obtained by desinoting from these results of the interest for the deep used, as before explained. In example 2, it will be observed, this makes a difference of 30 cents.

to

11

me

£5 wb

eit

dis

#### ROWE

Find the present value of \$1 for the same time (adding the three days of grace) and at the same rate; divide the required present value of the note by this, and the quotient will be the face of the note.

BEANPLE.—For what sum must a note be drawn at 8 months 18 days, so that discounted immediately at 6 per cent. it shall produce \$670?

### OPERATION.

Interest on \$1 for 8 months 21 days at 6 per cent. \$0.0435, and this taken from \$1 gives us \$0.9565 present worth of \$1.

Then 0.0565 = \$700.47. Ane.

### EXERCISE 115.

1. What sum must I put on the face of a note payable in 90 days so that I may obtain \$3755 when discounted at a bank at 7 per cent.?

Ans. \$3824-15.

2. For what sum must a note be drawn payable in 6 months in order that its proceeds at 5 per cent. bank discount may be \$1147.80?

Ans. \$1177.734.

3. For what sum must a note be drawn payable in 45 days so that its proceeds at 3½ per cent. bank discount may be \$713.90?

Ans. \$717.2471.

# EQUATION OF PAYMENTS.

48. Equation of Payments is the process of finding the equated or average time when two or more payments, due at different times, may be made at once without loss to either party.

44. The average time for the payment of several sums due at different times is called the mean time or equated

time

45. To find the equated time for any number of payments:—

### RULE.

First multiply each debt by the time before it becomes due; then divide the sum of the products thus obtained by the sum of the payments, and the quotient will be the equated time required.

Work by Arts. 26 and 27.

† This rule is based upon the supposition that what is gained by keeping certain payments after they become due is equal to what is lost by naving other payments before they become due. This, however, is not exactly true; for the gain is the interest, while the loss is equal only to the

resent of the

onthe shall

taken

bank 24.15. onths t may 7.734.

in 90

BYS SO ay be -2471.

g the , due

sums uated

f pay-

: then of the

keeplost by is not to the Nors.—When there are both days and months, they must all be reduced to the same unit; i. e., the payments mast all be reckoned for so many days, or so many months or parts of a month. If one of the payments is due on the day from which the equated time is reckoned, the corresponding product will be nothing; but in finding the sum of the debts, this payment must be added with the others. (See Example 3 below.)

EXAMPLE 1 .- A merchant purchases a vessel for \$7000,\$2000 to be paid in 3 months, \$2000 in 5 months, and the balance in 11 months. Now if he wishes to make the whole in one payment for what time must his note be drawn?

OPERATION. \$2000 × \$=\$ \$000 × 1 2000 × 5= 10000 × 1 3000 × 11= \$3000 × 1

\$2000 \times 3 = \$6000 \times 1

2000 \times 5 = 10000 \times 1

\$2000 \times 5 = 10000 \times 1

\$2000 \times 1 = \$2000 \times 1

\$2000 for three months is equal to the interest of \$6000 for one month. Similarly, the interest of the second payment is equal to the interest of \$10000 for one month, and the interest of the interest of \$3000 for one month. Hence, the interest of the several payments at the siven times will be appeared to the interest of \$3000 for one month.

Hence, the interest of the several payments, at the given times, will be equal to that of \$49000 for one mouth; and if we divide this \$49000 by the sum of the payments, \$7000, we obtain 7 months for the equated time.

That is, \$7000: \$40000::1 month: Ans.= \$7000 =7 months.

EXAMPLE 2 .- A person owes another £20, payable in 6 months; £50, payable in 8 months; and £90, payable in 12 months. At what time may all be paid together, without loss or gain to either party?

> OPERATION.  $20 \times 6 = 120$   $50 \times 8 = 400$  $90 \times 12 = 1080$ 160) 1600 (10 months. Ans.

discount, which (Art. 33) is always less than the interest: but the discrepancy is so trifling as not to make any material difference in the result.

With this exception, the rule is true, and may be demonstrated as follows:—Let p = first payment, and t = the time hefore it becomes due; p' = other payment, and t' = the time before it becomes due; x = equated time, and r = the rate of interest per unit. And since x, the equated time, lies between t and t' the time between t and x is = x - t, and that between t' and x is = t' - x. The interest of p for the time x - t is (from Art. 13) pr(x - t). Also interest of p' for the time t' - x is p'r(t' - x). Hence pr(x - t) = p'r(t' - x).

And  $x = \frac{p\ell + p'\ell'}{p + p'}$ , which is the rule, and may be similarly proved for any number of payments.

lo

w

of

B 17.

TO EU

EXAMPLE 3. A debt of \$450 is to be paid thus: \$100 immediately, \$300 in four, and the rest in 6 months. When should it be paid altogether?

OPPERATION. \$100×0===0 \$00×4==1200 50×6== 300

450 450)1500(3} months. Ans.

# EXERCISE 116.

1. A owes B \$600, of which \$200 is payable in 3 months, \$150 in 4 months, and the rest in 6 months; but it is agreed that the whole sum shall be paid at one payment. When should the payment be made? Ans. In 41 months.

2. A debt is to be discharged in the following manner: 1 at present, and † every three months after until all is paid. What is the equated time?

3. A debt of \$120 will be due as follows: \$50 in 2 months, \$40 in 5, and the rest in 7 months. When may the whole be

paid together?

4. I owe \$1000 to be paid down, \$1500 in 1 month, \$600 in 3 months, \$700 in 5 months, and \$1400 in 7 months. For what time must my note be drawn so that the whole may be paid in one payment? Ans. 35 months.

5. Bought of Messrs. Hendrie & Robarts, goods to the following

amounts, on the credit of six months;

15th of January, a bill of \$3750, 10th of February, a bill of 3000,

6th of March, a bill of 2400, 8th of June, a bill of 2250, I wish on 1st of July to give my note for the amount; at what time must it be made payable? fas 31st August.

# PARTNERSHIP OR FELLOWSHIP.

46. Partnership or Fellowship is the joining together of two or more persons for the transaction of business, agreeing to share the profits and losses in proportion to the amount of money each invests in the business.

47. The persons thus associated are called Partners,

and the association itself a Company or Firm.

48. The money employed is called the Capital or Stock. 49. The gain or loss by hared is called the Dividend.

# SIMPLE PARTNERSHIP.

50. When the partners employ their shares of the capital for the same period of time, the partnership is called Simple Partnership.

It is also called Single Partnership or Partnership without Time.

51. It is evident that the whole stock which suffers the gain or loss must been the same proportion to the stock of each partner that the whole gain or loss bears to his share of the gain or loss.

Hence, for partnership without time, we have the following :-

As the whole stock is to each man's share of the stock, so is the whole gain or loss to each man's share of the gain or loss.

EXAMPLE.—A and B epter into trade with a capital of \$3700, of which A contributes \$2000 and B the remainder. They gain \$1200. What is each man's share of the profits?

### OPERATION.

Whole stock: A's stock:: whole profit: A's profit.

That is, \$3700: \$2000:: \$1200: 3700 = \$648:648 = A'
Again, whole stock: B's stock:: whole profit: B's profit.

That is, \$3700: \$2700: 1900: 1700×1280 = \$648.648 = A's shars.

That is, \$3700; \$1700; 1900: 3700; \$551:351 B's share.

North After A's share has been found, B's share may be obtained by subtracting A's profit from the whole profit.

# vadi Sada i i il ito Mi Exercise 117.

1. Two merchants enter into partnership with a stock of \$4300, of which A contributes \$3000. They gain \$1117; how should this be divided between them?

Ans. A's share = \$779.302.

B's share = \$837.697.

2. Three persons A, B and C, agree to form a company for the manufacture of woollen cloths. A puts in \$6470, B \$3780, and O \$9860. By the end of the year they find that they have gained \$7890. What portion of this profit belongs to ach? Ans. A's share = \$2538.453.

B's share = \$1483.053. O's share = \$3868.493.

- 3. B and C buy certain merchandize, amounting to \$320, of which B pays \$120, and C \$200? and they gain \$80. How Ans. B \$30 and O \$50. is it to be divided?
- 4. B and O gain by trade \$728; B put in \$1200, and C \$1600. Ans. B \$312 and O \$416. What is the gain of each?
- 5. Two persons are to share \$100 in the proportions of 2 to B and I to O. What is the share of each? Ans. B \$66.661 and C[\$38.334.

meld it

\$150 reed Vhen nths.

1 at paid. nths. \$40 le be nths.

in 3 For may nths. wing

what gust.

ether ness. o the

mers,

tock. end.

YA

F 11

bu

mo

C2.

3.

1881

Sir

6. A merchant failing, owes to B £500 and C £900; but has only £1100 to meet these demands. How much should each creditor receive? Ans. B £3929 and C £7071.

Three merchants load a ship with butter; B gives 200 casks, C 300, and D 400; but when they are at sea it is found necessary to throw 180 casks overboard. How much of this loss should fall to the share of each merchant?

Ans. B should lose 40 casks, C 60, and D 80.

8. Three persons are to pay a tax of \$100, according to their estates. B's yearly property is \$800, C's \$600, and D's \$400. How much is each person's share?

Ans. B's \$44.44\frac{1}{2}, C's \$33.33\frac{1}{2}, and D's \$22.22\frac{1}{2}.

9. Divide 120 into three such parts as shall be to each other as 1, 2 and 3. Ans. 20, 40, and 60.

10. A ship worth \$900 is entirely lost; } of it belonged to B, to C, and the rest to D. What should be the loss of each. \$540 being received as insurance?

Ans. B \$45, C \$90 and D \$225.

11. Three persons have gained \$1320; if B were to take \$6, C ought to take \$4, and D \$2. What is each person's share? Ans. B's \$660, C's \$440, and D's \$220.

12. Three persons join; B and C put in a certain stock, and D puts in £1090; they gain £110, of which B takes £35, and O £20. How much did B and C put in; and D's share of the gain? - beatle .. Ans. B put in £829 6s. 1114d., # £687 3s. 513d.,

and D's part of the profit is £46.

# COMPOUND PARTNERSHIP.

52. When the partners employ their capital for different periods of time, the partnership is called Compound Partnership or Compound Fellowship.

It is likewise called Double Partnership, or Partnership With Time.

For stample; Suppose A puts in \$200 for 3 years, and B \$300 for 4 years, and they make a certain gain or loss. This would give a case of Compound Partnership.

In such cases it is plain that each man's share of the profit depends upon two circumstances:

1st. The amount of his stock; and
2nd. The period for which it is continued in the business.
Also that when the times are equal, the shares of the gain or loss are as
the stocks; when the stocks are equal, the shares are as the times; and
when neither the times nor the stocks are equal, the shares are as their products.

Hence, for Compound Partnership we have the following:

Multiply each man's stock by the time he continues it in trade; then say, as the sum of the products is to each particular product, so is the whole gain or loss to each man's share of the gain or loss,

has fould 707}. asks,

VIII.

D 80. their

ch of

other d 60. to B, each,

\$225. \$6, C share? \$220. and D 5, and hare of

iare of 111 d., 513 d., is £46.

fferent Part-

4 years, mpound ds upon

s are as es; and as their

ng :---

trade; roduct, r loss. EXAMPLE.—A contributes \$120 for 6 months, B \$336 for 11 months, and O \$384 for 8 months; and they lose \$56. What is O's share of the loss?

#### OPERATION.

\$120 × 6 = \$720 for one month) \$36 × 11 = \$696 for one month \$84 × 8 = \$072 for one month)

galage for a first of the second

\$3072×56 = \$22974.

EXPLANATION.—It is clear that \$120 contributed for 6 months are, as far as the gain or loss is concerned, the same as 6 times \$120, or \$720, contributed for one month. Hence A's contribution may be taken as \$720 for 1 month; and, for the same reason, B's as \$1696 for the same time; and C's as \$2072, also for the same time. This reduces the question to one in Simple Fellowship.

## Exercise 118.

1. Three merchants enter into partnership; B puts in \$357 for 5 months, O \$371 for 7 months, and D \$154 for 11 months; and they gain \$347.20. What should be each person's share of it?

Ans. B's \$102, O's \$148.40, and D's \$96.80.

2. B, C, and D pay \$160 as the year's rent of a pasture. B puts
40 cows on it for 6 months, U 30 for 5 months, and D 50 for
the rest of the time. How much of the rent should each
person pay? Ans. B \$87.27<sub>1</sub>3<sub>1</sub>, C \$54.54<sub>1</sub>3<sub>1</sub>, and D \$18.18<sub>1</sub>3<sub>1</sub>.

3. Three dealers, A, B, and O, enter into partnership, and in a certain time make £291 13s. 4d. A's stock, £150, was in trade 6 months; B's, £200, 3 months; and U's, £125, 16 months. What is each person's share of the gain?

Ans. A's is £75, B's, £50, and C's, £166 13s. 4d.

4. Three persons have received \$665 interest; B had put in \$4000 for 12 months, C \$3000 for 15 months, and D \$5000 for 8 months. How much is each person's part of the interest?

Ans. B's \$240, C's \$225, and D's \$200.

5. Three troops of horse rent a field, for which they pay \$320; the first sent into it 26 horses for 12 days, the second 64 for 15 days, and the third 80 for 18 days. What must each pay?

Ans. The first must pay \$70,

The second "100, The third 150.

6. Three merchants are concerned in a steam-vessel; the first,
A, puts in \$960 for 6 months; the second, B, a sum unknown
for 12 months; and the third, C, \$640, for a time not known
when the accounts were settled. A received \$1200 for his
stock and profit, B \$1400 for his, and C \$1040 for his: what
was B's stock, and C's time?

Ans. B's stock was \$1600;
and C's time was 15 months.

8

20 21

22 23

25.

26. 27.

29.

30.

31.

83.

85.

36. I

37. H 88. H 89. H

40. W 41. H 42. H

43. W 44. W 45. H 46. W

47. W 49. W

50. Ho

51. Ho

Wh

Wh

52. 53. W1 54. Ho 55. Wh 56. Wi 57. Wh

NOTEIf A gain \$240 in 6 mon	ths, he would	gain \$480 in 1	2 months; tha	t
Norm.—If A gain \$240 in 6 mon is, A's stock and profit at = \$1440.	the end of 1	menths won	d be \$960 1-\$48	0
= arago.	I have to	1 2 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	a reference 11	

Then \$1440 : 2400 :: \$980 : B's stock ; or 3440 =\$1600 E's stock. Again. B's stock : C's stock :: B's profit : C's profit for samé time, vis :

640×800 12 months. That is \$1600 : \$640 :: \$900: 1600 = \$320 = C's profit for 12 months Lastly, C's profit for 12 months : C's given profit :: 12 months; C's

100 × 12 320 =15 mo. = Cs time. time; that is, \$320: \$400:: 12 months:

7. In the foregoing question A's gain was \$240 during 6 months, B's \$800 during 12 months, and C's \$400 during 15 months; and the sum of the products of their stocks and times is 34560. What were their stocks? Ans. A's was \$ 960,

8. In the same question the sum of the stocks is \$3200; A's stock was in trade 6 months, B's 12 months, and C's 15 months; and at the settling of accounts, A is paid \$240 of the gain, B \$800, and C \$400. What was each person's stock? Ans. A's was \$960, B's \$1600, and C's \$640.

# QUESTIONS TO BE ANSWERED BY THE PUPIL.

NOTE.—The numbers following the questions refer to the articles of the

1.5What is interest? (1)

2. What is the meaning of the terms per cent. and per annum? (1) 3. In what respect does interest differ from Commission and Brokerage?

What is the principal? (3)

5. What is meant by the rate per cent? (4)
6. What is meant by the rate per unit? (5)
7. What is the interest? (6)
8. What is the amount? (7)

9: Of how many kinds is interest? (8)

10, Explain the distinction between Simple and Compound Interest. (9

11. In using formulas for interest, what is the meaning of the letters P. A.

12. It, and r? (12).

13. Deduce algebraically a full set of rules for Simple Interest. (12)

13. How is the interest found when the principal, rate per cent, and time

Norm—Answer this and succeeding similar questions by giving the formula.

14. Interpret this formula. (13)

15. When the interest, rate der cent, and time are given what is the continuous similar questions by giving the interest.

14. Interpret this formula. (13)
15. When the interest, rate per cent., and time are given, what is the rule
16. Interpret this formula. (14)
17. How is the rate per cent. found when the interest, principal, and time
18. Interpret this formula. (15)
18. When the interest, principal, and rate are given, how is the time
found? (13)

that

stock.

vis:

pro-

s; C's

s time.

onths.

nths :

nies is

960.

1600.

); A's C's 15

240 of

erson's

\$640.

es of the

kerage!

rest. (9

ers P. A.

nd time

v giving

the rule

nd time

he time

640.

20. Interpret this formula. (16)
21. When the principal, rate, and time are given, hew is the amount found? (17)

22. Interpret this formula. (17)

23. When the amount, rate and time are given, how do we find the prin-

24. Interpret this formula. (18)

23. When the amount, principal, and time are given, how do we find the rate? (19)

26. Interpret this formula. (19)

27. Vi hen the amount, principal, and rate are given, how do we find the time.? (20)

28. Interpret this formula. (20)

29. How do we find the time in which any sum of money will amount to any given number of times itself at a given rate? (21)

30. Interpret this formula. (21)

31. How do we flud the rate at which any sum will amount to a given number of times itself in a given time? (22)

32. Interpret this formula. (22)

83. When the time and rate are given, how do we find to how many times itself a given sum will amount? (23)

34. Interpret this formula. (23)
85. How do we find the interest on \$1 at 6 per cent. per annum for any number of months ? (21)

36. How do we find the interest on \$1 at 6 per cent. for any number of days? (25)

How do we find the interest of any sum for any given time at 6 per cent. ? (26)

38. How may we find the interest at any other rate than 6 per cent. ? (27)

39. How do we compute interest on notes, &c., when partial payments are

40. What is the rule for calculating Compound Interest? (30)

41. How is Compound Interest calculated by the table given in Art. 31? (32)

42. How do we ascertain the present worth of a debt the some given time hence, allowing Compound Interest at a given rate (33).

43. What is Discount? (34)

44. What is meant by the present worth of a debt, note, &c. ? (35)
45. How do we compute the present worth of a debt or note? (36)

What is Bank Discount? (37)
What is the distinction between Bank Discount and True Discount? (38 and 35)

48. What are days of grace? (39)
49. What are the two kinds of notes discounted at banks? (40)
50. How do we calculate the bank discount on notes, &c.? (41)
51. How do we find what amount to put on the face of a note so that its
prevent value shall be a certain sum? (42)

52. What is meant by the Equation of Payments? (43)
53. What is meant by the mean time or equated time of payment? (44)
54. How do we find the equated time of payment? (45)
55. What is Partnership or Fellowship? (46)

What are the persons associated together in partnership called? (47) What is the money employed in the business called? (48)

What is meant by the dividend? (49)
What is the distinction between Simple and Compound Fellowship? (50 and 52)

By what other names is Simple Partnership known? (50) What is the rule for Simple Partnership? (51) What is the rule for Compound Partnership? (52)

pr lo

los

be

dol

wh

# SECTION IX.

PROFIT AND LOSS, BARTER, ALLIGATION, CUR-RENCIES, EXCHANGE, &c.

# PROFIT AND LOSS.

1. Profit and Loss is a rule by which we are enabled to ascertain what we gain or lose in mercantile transactions. It also instructs us how much we must increase or diminish the price of our goods in order that our gain or loss may be so much per cent.

# CASE I.

2. To find the total gain or loss on a certain quantity of goods when the prime cost and selling price are given:

#### FIRST BULE,

Find the price of the goods at prime cost and also at the selling price. The difference will be the whole gain or loss.

EXAMPLE 1.—What do I gain if I buy 207 cords of wood at \$3.78 per cord and sell it at \$4.25?

#### OPERATION.

207 cords @ \$4'25 = \$879'75 = whole sum for which goods were sold.

# Difference = \$97'29 = whole gain = Ans.

EXAMPLE 2.—If I purchase 900 bushels of wheat at \$1.47 per bushel and sell it at \$1.25, what do I lose upon the whole transaction?

#### OPERATION.

900 bushels @ \$1'47 = \$1328 = whole cost, 900 bushels @ \$1'25 = \$1125 = whole sum received for wheat,

\$198= whole loss = Ans.

#### SECOND RULE.

Find the difference between the buying and selling price of a buhsel, lb., yard, &c.

Multiply the gain or loss per bushel, lb., yard, &c., by the number of bushels, lbs., or yards, and the result will be the whole gain or loss.

old.

ng-

Example.—Bought 211 yards of flannel at 371 cents per yard, and sold it at 45 cents; required my total gain?

#### OPERATION.

\$0°375 = buying price. \$0°45 = selling price.

\$0°075 gain per yard \$0°075 × 211 = \$15°825. Ans.

NOTE.—This second rule affords the shorter method of finding the gain or loss.

## EXERCISE 119.

1. Bought 317 lbs. of butter at 9 cents per lb., and sold it at 124 cents; what was my gain on the whole? Ans. \$11.095.

2. Bought 2138 bushels of potatoes at 871 cents per bushel, and sold them at \$1.20; what was my gain on the whole? Ans. \$694.851

3. Bought 13 barrels of sugar, each weighing 317 lbs. net at 15 cents per lb., and sold the whole for \$735; how much did I gain or lose on the transaction? Ans. Gained \$116.85.

4. Bought 17 kegs of wine, each containing 22 gallons, at \$3.15 per gallon, and paid in addition \$26.33 for carriage, &c., and an ad valorem duty of 371 per cent. I sold the whole for \$1625; what was my gain or loss? Ans. Loss \$21.2175.

# CASE II.

3. Let it be required to find for what sum I must sell a house which cost \$2900 so that I may gain 15 per cent.

Here for every \$100 the house cost me I am to receive \$115, or for every 1 cost I am to receive \$1'15.

The selling price must evidently be as many times \$1°15 as the buying price contains \$1; i.e., \$1°15×2900 = \$3335'00. Ans.

Again: If a person buys a horse for \$230, and afterwards sells it so as to lose 11 per cent.; how much does he receive for it?

Here for every \$1 he paid for the horse he receives only \$0°89 (since he loses 11 per cent., i.e. 11 cents on the \$1.)

Then, the selling price will obviously be \$0°89×230=\$204'70. Ans.

Hence, to find at what price an article must be sold so as to gain or lose a specified per centage, the cost price being given :-

Find (Art. 2, Sect. VII.) how much must be received for each dollar of the buying price, and multiply this by the whole buying The result will be the selling price.

EXAMPLE.—Bought a quantity of oatmeal for \$1793.80. For what must I sell it so as to gain 8 per cent.?

### OPERATION.

Here for every \$1 I expend I desire to receive \$108; hence, the selling price will be \$108×1793'80 = \$1937'304. Ans.

EXAMPLE. Bought a lot of sheep for \$7000, and am willing to lose 3 per cent. For what sum must I sell?

Here for every \$1 I expend I am willing to receive \$0.97, and hence the set ing price will be \$0.97×7000 = \$6790. Ans.

### Exercise 120.

- 1. Bought cordwood at \$3.25 per cord; at what rate per cord
- must I sell it in order to gain 30 per cent.?

  Ans. \$4.221.

  2. Bought a stock of goods for \$13420; for how much must it be sold in order to gain 5 per cent.? Ans. \$14091.
- 3. Bought a quantity of wood at 11 cts. a lb., and wish to sell so as to gain 15 per cent.; at what rate per lb. must I sell Ans. 12 13 cents.
- 4. Bought axes at \$15.25 a doz., and desire to sell them so as to gain 23 per cent.; at what rate per doz. must I sell?
- 5. Bought a farm for \$7890, and am willing to lose 11 per cent.; at what price must I sell? Ans. \$7022.10.

# CASE III.

4. Let it be required to find what per cent. of profit a merchant makes by buying tea at 43 cents per lb. and selling it at 67 cts.

Here the gain on each lb. is 24 cents.

That is every 43 cents invested gives a gain of 24 cents.

Therefore every cent invested gains 13 of 24 cents = 23 cents.

And hence, the gain per cent.  $=\frac{24}{13} \times 100 = 2400 = 55$  per cent.

Hence to find the rate per cent. of profit or loss when the prime cost and selling price are given, we have the following !--

Find the difference between the buying and selling price, and hence the gain or loss per unit.

Multiply this by 100, and the result will be the gain or loss per

cent.

EXAMPLE.—A speculator invests \$44400 in stocks, and sells out for \$50000; what per cent. does he make by the operation?

#### OPBRATION.

Here the whole gain is \$50000 \$4400 = \$5600.

That is \$44400 gain \$5600, and therefore \$1 gains  $\frac{5600}{4000} = \frac{14}{111}$  of a dollar Hence gain per cent =  $\frac{11}{111} \times 100 = \frac{140}{110} = 12.6$ . Ans.

Nors.—The above and all similar questions may be solved by Proportion .

Thus this question is, if \$44400 gain \$5000, what will \$100 gain \$2000.

And the statement is \$44400: \$100:: \$5600: Ans.- 44400

# EXERCISE 121.

- 1. Bought tes at 60 cents a lb., and sold it at 871 a lb.; how much did I gain per cent.?
- 2. Bought coffee at 13 cents and sold it at 11 cents a pound; what was my loss per cent.?
- 3. Bought flour at \$6.20 a barrel, and sold it at \$7.80; what was the per cent. of profit?

  Ans. 25; per cent.
- 4. Bought cloth at \$2.75 per yard, and sold it at \$3.10; what was my gain per cent.?

  Ans. 124 per cent.
- 5. Bought oats at \$0.47 per bushel, and sold them at \$0.56; what was my gain per cent.?

  Ans. 19 7 per cent
- 6. Bought meat at 12 cents per lb., and sold it at 10 cents a pound; what was my loss per cent.?

  Ans. 12 per cent.
- 7. Eought a horse for \$93, and sold it for \$127; what per cent. of profit did I make?

  Ans. 3653.
- 8. A man bought a farm for \$6742.50, and sold it for \$6000; what was his loss per cent.?

  Ans 11 1890 per cent.
- 9. If I purchase a house for \$5700, a horse for \$275, and pay \$1987.32 for household furniture and a carriage, and then sell the whole for \$8750, what is my gain or loss per cent?

  Ans. Gain 9.89 or nearly 10 per cent.
- 10. I purchase 723 yards of black si'k velvet in Paris and pay \$4.25 a yard; I further pay 7 per cent. for insurance, \$23.70 for carriage, \$2.70 for harbor dues, \$3.16 for wharfage and storage, and an ad valorem duty of 22 per cent., and then sell the whole for \$5270; what is my gain or loss per cent.?

  Ans. Gein 31.96749 or nearly 32 per cent.

# CASE IV.

5. Let it be required to find the prime cost of cloth which I sold for \$4 and gained 10 per cent. thereby.

Here the gain on \$1 was 10 cents, or what I sold for \$1.10 cost me out

Therefore the cost price will contain \$1 as many times as the selling price contains \$1.10.

That is the cost price = 14.00 = \$3.636. Ans.

Hence, to find the cost price, the selling price and the gain or loss per cent. being given, we have the following

#### RULE

Find the gain or loss per unit, and add it to unity if it be gain, but subtract it from unity if it be loss.

Divide the selling price by the quantity thus obtained, and the result will be the cost price.

Or s \$100 | gain per cent. (or as \$100—loss per cent.) is to \$100 is the selling price to the cost price.

EXAMPLE.—Sold a quantity of coal for \$719, and lost 7 per cent. by the transaction; what was the prime cost?

if Jan J. west OPERATION. Li Ja oblico Ju

lst Rule.—Loss on \$1 is 7 cents, or for every \$1 paid I receive \$0.98. Hence cost = \$7 \frac{1}{2} = \$773 \cdot 118.

BND RULE. - 100: 100: 2719: Ans. = 100×719 = 1773'118

### Exercise 122.

1. For what did I buy a quantity of sugar which I sold for \$24.60, losing 4 per cent.?

2. A gentleman sold his library for \$2360, which was 10 per cent less than dost; what did he give for it? Ans. \$2622.22.

3. A farmer sold his farm for \$7400, gaining 11 per cent. on the prime cost; what did he give for it? ... Ans. \$666.666.

4. A merchant sold a quantity of silk velvet for \$3789 40.

gaining 17 per cent. by the transaction; required the buying price?

Ans. \$3238.803.

5. Sold a lot of cattle for \$2740, losing 13 per cent. by the transaction; what did I give for them? ... Ans. \$3149.425.

### BARTER.

6. Barter signifies an exchange of goods or articles of commerce at prices agreed upon so that neither party in the transaction may sustain loss.

7. The principle of solution depends upon finding the value of the commodity whose price and quantity are given, and thence the equivalent quantity of a second commodity of a given price, or the equivalent price of a given quantity of a second commodity.

EXAMPLE 1.—How much tea at \$1:10 per lb. ought to be given for 712 lb. of sugar at 13 cents per lb.?

### OPERATION.

713 lbs. of sugar at 13 cents per lb. \$92.56, and \$92.56. \$1.10 84.1454 lbs. 84 lbs. 21 oz. Ans.

EXAMPLE 2.—I desire to barter 96 lbs. of sugar, which cost me 8 cents per lb., but which I sell at 13 cents, giving 9 months' credit, for calico which another merchant sells for 17 cents per yard, giving 6 month's credit. How much calico ought I to receive?

### OPERATION.

I first find at what price I could sell my sugar, were I to give

If 9 months give me 5 cents profit, what ought 6 months to give?

9:6::5: 5 = 1 cente.

Hence, were I to give 6 months' credit, I should charge 8+3;=11; centa, per lb. Next-

IX.

per

for

325.

per

.22.

. on

666.

.40.

uy-

803.

the

425.

s of

7 in

e of

the

the

ven

1454

cost

ths'

per

to

nts.

As my selling price is to my buying price, so ought his selling to be to his buying price, both giving the same credit.

The price of my awar, therefore, is 96×8 cents, or \$768; and of his called, 12 cents per yard.

Hence  $\frac{1}{12} = 64$ , is the required number of yards.

### EXERCISE 123.

- 1. A has coffee which he barters at 10 cents the lb. more than
  it against tea which stands B in \$2, leaf which he
  from 65 per lb. How much did the coffee had a thirst?
- 2. A which cost \$2.80 per lb.; B has cloth at \$2.50, which only \$2 the yard. How much must A charge for his ark, to make his profit equal to that of B?
- 3. I have cloth at 8 cents the yard, and in barter charge for it 13 cents, and give 9 months' time for payment; another merchant has goods which cost him 12 cents per lb., and with which he gives 6 months' time for payment. How high must he charge his goods to make an equal barter?
- Ans. At 17 cents.

  4. K and L barter. K has cloth worth \$1.60 the yard, which he barters at \$1.85 with L, for linen cloth at 60 cents per yard, which is worth only 55 cents. Who has the advantage; and how much linen does L give to K for 70 yards of his cloth? Ans. L gives K 2155 yards; and K has the advantage.
- 5. B has five tons of butter, at \$102 per ton, and 101 tons of tallow, at \$135 per ton, which he barters with C; agreeing to receive \$600.30 in ready money, and the rest in beef at \$4.20 per barrel. How many barrels is he to receive?


### Ans. 316.


### ALLIGATION.

8. Alligation is the method of finding the value of a mixture of ingredients of different values, or of forming a compound which shall have a given value.


Norm.—The term alligation is derived from the Latin word allige " to tie or bind," the reference being to the manner of connecting or tring the numbers together in a certain class of questions.

- 9. Alligation is divided into Alligation Medial and Alligation Alternate.
- 10. Alligation Medial (Latin medius, "mean or average,") enables us to find the value of a mixture when the






# IMAGE EVALUATION TEST TARGET (MT-3)



Photographic Sciences Corporation

23 WEST MAIN STREET WEBSTER, N.Y. 14580 (716) 872-4503

STATE OF THE STATE



Me

por

con

thos less

L betu

wri

I

111

who

mix

it is

whi

who

A TO

WOW

ingredients, of which it is composed and their prices are

11. Alligation Alternate enables us to find what proportion must be taken of several ingredients, whose prices are known, in order to form a compound of a given price.

### ALLIGATION MEDIAL.

12. Let it be required to find the price per lb. of a mixture containing 47 lbs. of sugar at 11 cents per lb., 29 lbs. at 13 cents, and 24 lbs. at 17 cents.

### OPERATION.

67 lbs. at 11 cents = 517 cents.

29 lbs, at 18 cents = 377 cents. 24 lbs. at 17 cents = 408 cents.

Then 100 lbs. cost 1802 cents and 1 lb. will cost 1302 = 18 cents.

Hence for Alligation Medial we deduce the following:

Divide the entire cost of the whole mixture by the sum of the ingredients, and the quotient will be the price per unit of the

Example 1.—What will be the price per lb. of a mixture of tea containing 7 lbs. at \$0.50 per lb., 11 lbs. at \$0.80, 19 at \$1.06, and 3 lbs. at \$1.23?

OPERATION.

10 enut 193 fan 17 lbm. @ \$0.50 = \$9500 to saft at each nalescent 11. "an @ \$180 = \$200 mg teld an each at each nalescent 12. " @ \$176 = \$2014 at 2012 at 2012

40 lbs. = sum of ingre- \$36'15 = Total cost.

40)\$38-13(\$0-9018. Ans. 86.0

a lo other extractation bolton self at notes the

EXAMPLE 2.- A goldsmith has 3 lbs. of gold 22 carats fine, and 2 lbs. 21 carats fine. What will be the fineness of the mixture?

In this case the value of each kind of ingredient is represented by a num-

OPERATION.

8 lbs. × 22 = 66 carsts 8 " × 21 = 43 "

Filester 19108 F. Filester apit WE G.

The mixture is 31 carata line,

ices

100.

ture

13

the

of

06,

### Exercise 124.

1. Having melted together 7 os. of gold 22 carats fine, 12} os. 21 carats fine, and 17 oz. 9 carats fine, I wish to know the fineness of each ounce of the mixture?

Ans. 1542 carats.

2. A vintner mixed 2 gallons of wine, at 14s. per gallon, with 1 gallon at 12s., 2 gallons at 9s., and 4 gallons at 8s. What is one gallon of the mixture worth?

Ans. 10s.

3. A farmer mixes 15 bushels of wheat worth \$1.20 with 30 bushels worth \$1.50, and 60 bushels worth \$1.10 and 83 bushels worth \$1.75. What is one bushel of the mixture worth?

4. A grocer mixes together 12 lbs. of tea at 50 cents, 16 lbs. at 72 cents, 12 lbs. at 65 cents, 18 lbs. at 85 cents, and 100 lbs. at 42 cents. How much per lb. is the mixture worth?

Ans. 53 cents.

### ALLIGATION ALTERNATE.

13. Alligation Alternate is the reverse of Alligation Medial, and may be proved by it.

### CASE I.

14. Given the prices of the ingredients, to find the proportion in which they must be mixed in order that the compound may be worth a given price:—

### RULE.

Set down the prices of the ingredients in two columns, placing those greater than the price of the compound to the left, and those less than it to the right.

Between these columns form two others composed of the differences between the prices of the several ingredients and of the compound; writing each difference next to the number by which it was obtained.

Link, by means of a line, the left-hand differences to the right-

hand differences in any order.

Then each difference will express how much of the quantity with whose difference it is connected, should be taken to form the required mixture.

If any difference is connected with more than one other difference, it is to be considered as repeated for each of the differences with which it is connected; und the sum of the differences with which it is connected is to be taken as the required amount of the quantity whose difference it is.

EXAMPLE 1.—How many pounds of tea at 5s. and 8s. per lb., would form a mixture worth 7s. per lb.?

2.

### OF REALTON.

### Prices. Differences Prices 4. I no services 2

It is connected with 2, the difference between the 7, the required price, and 5s. hence there must be 1 h. at 5s. 2 he connected with 1, the difference between 8s, and the required price; hence there must be 3 bs. at 8s. Then 1 lb. of tea at 5s. and 1 lbs. at 8s. per lb. will form a mixture worth 7s. per lb.—as may be proved by the last rule.

"It is evident that any equimoltiples of these quantities would answer equally as well; hence a great number of answers may be given to such a question.

basels weren he tall heart so may realist of conditions Example 2.—How much sugar at 9d., 7d., 5d., and 10d., will d produce sugar at '8d: per lb. ? Lagrating a second

Second and sipple of Direction in and all are 35 12 व धाँडा क्रिक्स विशेषक के जान है। क

8=\{\frac{10-2}{10-2} \frac{1+7}{3+5}\}=8

1 is connected with 1, the difference, between 7d, and the mean, 8; hence there is to be 1 lb. of sugar at 7d. per 1b. 2 is connected with 3, the difference between 5d, and the mean; hence there is to be 2 lbs. at 5d. 1 is connected with 1, the difference between 3d and the mean; hence there are to be 3 lbs. at 10d. per 1b.

Consequently we are to take 1 lb, at 7d. and 2 lbs. at 5d., 1 lb. at 9d. and 3 lbs. at 10d. If we examine the price of the mixture these will give (Art. 12) we shall find it to be the given mean.

Example 3.—What quantities of tee at 4s., 6s., 8s., and 9s. per 1b., will produce amixture worth 8s.?

### OPPRATION.

Differences, Prices,




3, 1, and 4 are connected with 1s., the difference between as and the mean; therefore we are to take 5 lbs. + k lbs. + k lbs. at the mean; therefore we are to take 1 lb. of tea at a., 1 lb. of tea at a., 2 lb. and fis., and the mean; therefore we are to take 1 lb. of tea at a., 1 lb. of tea at a., and 1 lb. at 96. per lb.

EXAMPLE 4.—How much of any thing at 38, 48:, 58., 78, 88., 98., 118., and 128. per lb., would form a mixture worth 68. per lb.?

### OPERATION.

Prices. Differences. Prices.



1 lb, at 2s, 2 lbs, at 4s, 2 lbs, at 7s, 2 lbs, at 6s, 24-5+5; i.s., 14 lbs. at 5s, 1 lb, at 18s, and 1 lb, at 12s, per lb, will form the required

11.1180.75

the differ-blos at 8s.

ould answer

10d., will SELECT ST

NOTE. The principle upon which, this rule proceeds is that the cases of one ingredient above the mean is made to counterbalance which are separately being equal to the mean. This in crample 7, 1 the at separate lib. gives a deficiency of Sa.; but this is corrected by Sa. excess in the 3 lbs. at Sa. per lb. Ilbs. at 8s. per lb. at 7d. gives a deploising of 1d., 1 lb. at 9d. gives an excess of 1d., but the excess of 1d. and the deficiency of 1d. exactly neutralize each other.

Again, it is evident that 2 lbs., at 5d, and 3 lbs. at 10d. are worth just as much as 5 lbs. at 8d,—that is, 8d. will be the average price if we mix 2 lbs. at 5d. with 3 lbs. at 10d.

EXERCISE 125.

1. How much wheat at \$1.80, \$1.40, \$1.10, and \$1 per bushel must be mixed together in order to form a mixture worth \$1.25 per bushel? Give at least two sets of answers.

4ns. \$5 bushels at \$1.10, 15 at \$1.00, 15 at \$1.00, and 35 at \$1.40, and 35 bushels at \$1.00, 15 at \$1.40, 15 at \$1.10, and 25 at \$1.00.

2. How much wine at 60 cents, 50 cents, 42 cents, 38 cents, and 30 cents per quart, will make a mixture worth 45 cents a quart? Ans. 15 qts. at 42 q 5 qts. at 30 c., 3 qts. at 60 c. and 22 qts, at 50 c. and 5 quarts at 38 cents.

3. A merchant has sugar worth 10 cents, 12 cents, 14 cents. 15 cents, 16 cents, 17 cents, and 18 cents per pound, and wishes to form a mixture worth 121 cents a lb. How many pounds of each must he use? Ans. 21 lbs. at 14 c., 11 lbs. at 10 c., 16 lbs. at 12 c. and 1 lb. at each other price.

4. A grocer has sugar at 5d., 7d., 12d., and 13d. per Ib. How much of each kind will form a mixture worth 10d. per lb.? Ans. 2 lbs. at 5d., 3 lbs. at 7d., 5 lbs. at 12d., and 3 lbs. at 13d.

CASE II, TE as and dram were 15. When a given quantity of one of the ingredients is to be taken:

I. Find the proportional quantities of the ingredients as in Case I. II. Then say, as the amount of the ingredient, as thus found is to the given amount of the same ingredient, so is the amount of any other ingredient (found by Case I.) to the required quantity of that

EXAMPLE, 1.—29 lbs of tea at 4s, per lb., is to be mixed with teas at 6s., 8s., and 9s. per lb., so as to produce what will be worth 5s. per lb. What quantities must be used?

### OPERATION.

By Case I we find that 8 lbs. of tea at 4s, and 1 lb. at 6s., 1 lb. at 8s., and 1 lb. at 9s., will make a mixture worth 5s. per lb.

Therefore 8 lbs. (the quantity of tea at 4s. per lb., as found by the rule):
29 lbs. (the given quantity of the same tea)::1 lb. (the quantity of tea at 6s. per lb., as found by the rule:) or 8 lb. = 8s. lbs. 4ss.

We may in the same manner find what quantities of tea at 6s. and 9 per lb., correspond with 29 lb. of tea at 4s. per lb.

n, 8; hence the differ-i. 1 is con-e there is to en Tod; and

at 9d. and Il give (Art.

and 9s.

and the An men lb. f ton at os.

78., 88., per lb. ?

14 lbs. at

g E in 73

6s., 8

11lbs.

lbat

11 :

The

1. A

-xi\$1

\$1

pu \$1 2. 271

eac

gai \$2

a \$9

194174

rice.

Ale B

. 17.

a sum

count

tions o

or mor

19.

kind, 20:

its ma

usage.

TATI DE

18.

3. Ho

EXAMPLE 2.—A refiner has 10 oz. of gold 20 carats fine, and melts it with 16 oz. 18 carats fine. What must be added to make the mixture 22 carats fine?

10 oz. of 20 carats fine = 10×20 = 200 carats. 16 os. of 18 carats fine = 16×18 = 288

26: 1 :: 488 : 1814 carate, the Ane-

has nees of the mixture. a mis

24—22 = 2 carats baser metal, in a mixture 22 carate fine. 24—1849 = 5,3 carats baser metal in a mixture 1849 carate fine. Then 2 carats: 22 carats:; 5.3: 57.2 carats of pure gold—required to change 5,3 carats baser metal into a mixture 22 carats fine. But there are already in the mixture 18+9 carats gold; therefore 57-73 18+9-38+9 carats gold are to be added to every ounce. There are 26 oz.; therefore 26×384 9. = 1008 carats of gold are wanting. There are 24 carats in every os; therefore 1998 carats = 42 oz. of gold must be added. There will then be a mixture containing:

this thinny ion 68: 1 os. :: 1496 : 22 carata, the required finenes

### Exercise 126.

1. How much molasses at 16 cents, at 19 cents, and at 23 cents per quart must be mixed with 87 quarts at 31 cents in order that the mixture may be worth 25 cents per quart?

Ans. 30+2 qts, at each price. 2. How much oats at 37 cents per bushel and barley at 68 cts. per bushel must be mixed with 70 bushels of peas at 80 cts. a bushel so that the mixture may be worth 75 cents per bushel? Ans. 71 bush, at each price.

How much brass at 14d. per lb., and pewter at 101d, per lb., must I melt with 50 lbs. of copper at 16d. per lb., so as to make the mixture worth 1s. per lb.?

Ans. 50 lbs. of brass, and 200 lbs. of pewter. 4. How much gold of 21 and 23 carats fine must be mixed with 30 oz. of 20 carats fine, so that the mixture may be 22 carats and one of an or .all s Ins. 30 of 21, and 90 of 28. fine?

### CASE III.

16. When the quantity of the compound is given as well as the price :-

Find the proportional quantities as in Case I.

II. Then say, as the sum of the proportional quantities is to each proportional quantity, so is the given quantity to the corresponding part of each.

[Spor. IX.

s fine, and added to

a, the Anc-

arats fine.

-required to

sut there are

=361 g carats

ore 26×361 g.

y os; thereen be a mix-

at 23 cents ts in order

ach price.
at 68 cts.
at 80 cts.
cents per
ach price.
101d, per
r lb., so as

of pewter. aixed with 22 carats 1 90 of 23.

given as

is to each

EXAMPLE—What must be the amount of tea at 4s. per lb. in 736 lb. of a mixture worth 5s.per lb., and containing tea at 6s., 8s., and 9s., per lb.?

To produce a mixture worth 5s. per lb., we require 8 lbs. at 4s., 1 at 6s., and 1 at 9s. per lb. (Art. 14). But all of these added together, will make 11lbs. in which there are 8 lbs. at 4s. Therefore

lbe. lbs. lbs. lbs. lbs. os.

11 : 8 :: 786: 8×786 = 535 47 , the required quantity of ten at 4s.

That is, in 736 lbs. of the mixture there will be 535 lbs. 44, or. at 4s. per lb. The amount of each of the other ingredients may be found in the same way.

### Exercise 127.

- 1. A druggist is desirous of producing, from medicine at \$1.00, \$1.20, \$1.60, and \$1.80 per lb., 168lbs. of a mixture worth \$1.40 per lb; how much of each kind must be use for the purpose? Ans. 28lbs. at \$1.00, 56lbs. at \$1.20, 56lbs. at \$1.60, and 28 lbs. at \$1.80 per lb.
- 2. 27lbs. of a mixture worth 4s. 4d. per lb. are required. It is to contain tea at 5s. and at 3s. 6d. per lb.; how much of each must be used? Ans. 15lbs. at 5s., and 12lbs. at 3s. 6d.
- 3. How much brandy at \$2.40, \$2.60, \$2.80, and \$2.90, per gallon, must there be in one hogshead of a mixture worth \$2.70 per gallon? Ans. 18 gals. at \$2.40, 9 gallons at \$2.60, \$9 gals at \$2.80, and 27 gals. at \$2.90 per gallon.

### EXCHANGE OF CURRENCIES.

- 17. Exchange of Currencies is the process of changing a sum of money expressed in the denomination of one country to an equivalent sum expressed in the denominations of another country.
- 18. By the currency of a country is meant the coins, or money, or circulating medium of trade of that country.
- 19. The intrinsic value of a coin is determined by the kind, purity, and quantity of metal it contains.
- 20. The relative value or commercial value of a coin is its market value, and is fixed by law and commercial usage.

in the color that a correction is all in said in the color with the color of the

if the old plate rate area areas core, but is the general rates in and areas

8T. TUS

TUR

Guin CROV

HAL SHIL

DOLL

FIVE LIVE

CROV Loun FLOR GUIL FLOR THAL

RIX-I FLOR MARC FLOR FLOR FLOR RIX-I

SPECI

DOLL MILE

MILE

MILRI REAL

REAL PISTO:

RIAL PIETE CROSS RUBL INP TO

### FOREIGN MONEYS OF ACCOUNT,

WITH THE PAR VALUE OF THE UNIT, AS FIXED BY COMMERCIAL USAGE, EXPRESSED IN DOLLARS AND CENTER

UBAGE, EXPRESSED IN DOLLARS AND UBAGE.
AUSTRIA.—60 kreutsers = 1 florin (silver) =
BREKEN5 schwares =1 grote; 72 grotes =1 rix-dollar(silver)= '787
Burrish India 12 pice 1 anna; 16 annas 1 Company's rupee 445
Burnos Arens.—Srisis—1 dollar currency (variable) mean value.  CARTON, 10 cash t = 1 candarines; 10 cand = 1 mace; 10 mace = 1 tael = 1.48
CAPE OF GOOD HOPE.—6 stivers—1 solding; 8 schilings = 1 rix-dollar 818
Cayton 4 pice = I fanam; 12 fanams = 1 rix-dollar=
CURA, COLOMBIA AND CHILL-8 fials = 1 dollar =
Druggang 12 prenning 1 lakilling; 16 skillings 1 mare; 6 mares
Best AND. 4 farthings 1 penny: 12 pence 1 shilling: 20shil 21 4867
France.—10 centimes—1 decimes 10 decimes—1 franc—
Gminon—100 lepta—1 drachme; 1 drachme (dlver)—
HAMMUNGH.—12 prenning = 1 schiling; 16 sebil. = 1 mare; 8 mares
18. minen brapde at \$2.40, \$2.60, \$2.80, and allower TE
MALTA -20 grains = 1 taro; 12 tari =1 soudo; 24 soudi =1 pezza = 100
Mr.An. 12 denist 1 soldo 20 sold 1 hra
Marroo.—8 rials 21 dollar 2013 10 100 100 Monre Vidno,—100 centesimos =1 rial; 8 rials = 1 dollar =
NAPLES.—10 grani = 1 carlino: 10 carlini = 1 ducat (silver) = 80
Norway. 120 skillings 1 rix-dollar specie (silver) =
PAPAL STATES. 10 bajocchi=1-paolo; 10 paoli=1acudo or crown= 170
Puro e riale 1 dollar (diver) = 1 100 rees = 1 milree or crown = 1 12
Prince A the manage of growth (aliver); 20 growth in 1 thaler or
dolla =
Russia. 100 copecks = 1 ruble (silver) =
SARDINIA 100 centesimi = 1 lira =
Swent 1 addings = 1 rix-dollar specie = 106
Srong - Sgrapi = 1 terp; 20 teri = 1 onds (gold) = 10
8 reals = 1 plastre; 4 plastres;=1 pistole of exchange.
20 reals vellon = 1 Spanish dollar =
The current silver rupes of Bombay, Madras, and Bengal, is worth

The entrent silver rupes of Bombay, Madras, and Bengal, is worth 1044. In India also they use couries for coin. These are small shells found in the Maldives and elsewhere: 2500 cowries make a rupes, and 10000 rupes make a los.

† The cash, made of copper and lead, is said to be the only money coined in China.

The old plate real is not a coin, but is the denomination in which changes are usually made.

BOAL To John T	Toning to the state of the stat	-886 -181/10/1 -90*2/1 -005 ***
828 787	United States of America:—16 mills—1 cent; 10 cents — I dime; 10 dimes—1 dollar—	100 100 II
98.00	21. The following table exhibits the commercial va of the Foreign coins most frequently met with.	ME nonoxi
rix- •818	difficulty and the second of t	10 807 316
1:00 se	HALF-CROWN of England. SHILLING of England. DOLLAR of the United States	10b.j.s
11= 1865 186	FRANC Of France. FIVE-FRANC PIBOR Of France.	·181 ·98 ·184
166 40 Arcs	CROWN OF France.	106 y La
24 = 1 00 20 1 00	FLORIS of the Netherlands.  GUILDES of Southern Germany.	40
888 80	RIX-DOLLAR of Bremen.  FLORIN Of Prussia.  GWA-Francisco 10 Walst A.	·69 ·78‡ ·22‡
100 100 100 112	FLORIN of Austria and city of Angsburg.	85 48 48
er or	FLORIN of Nuremburg, Frankfort, and Creveld.  RIX-DOLLAR of Denmark.  SPECIA-DOLLAR of Denmark.  DOLLAR of Sweden and Norway.  MILER of Portugal.	W V
186 108 240	MILEE of Portugal  MILEE of Azores	12
1:00	REAT VELLOR of Spain  REAT PLATE of Spain  PISTOLE of Spain  8	10 C
al, is worth small shells supes, and	Brat of Spain	11 13
n in which	Protestant Caoss Pist Assess Rubils (cilibraty) of Bussia. Impurial of Bussia.	14 ST

1

_ fe, ett				, 'F	Lat'	10 w 10 1 1	מאלש שלא	. B. a	725 H.
DOUBLOON of	Mexico								172.00
HALF-JOE of	Portugal		• • • • • •		00000				8'58
LIBA of Tues	any and Lo	mbar	ly				1 77.		16
LIBA of Sard	inte	17	4.2	A-7	10 m	1	200	9 6 31 6	183
Ounon of Sic	dly		*****		7.9		1 1087	1.0	2.40
DUCAT Of N	inles							A 14	-80
CROWN of Tu	ibioi	•••••	•••••	*****				g (2	1.06
Florence Liv	mounty	A.	2					A .he	15
									181
Genos "									
Geneva "									21
Leghorn Don	LAB								.80
Swim LIVER.									- 27
SOUDO of Ma	lta		gel.			6 1 . 5	*		. 40
Turkish Pras	TRE		. He		. #		Sec. 4.14.	1 1	-05
PAGODA of I	ndie			.0.		70.01	1.5.5.5	up of the f	1.84
Ruran of Ind	lia.				•••••		128 21 Kg.	1-04	2.441
TABL of Chir		•				*****	traine li	- 1 E	1.49
TABL OF CHI		••••••	•••••			******			TAN DO A
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1			. 1		- 4	हर र जारेक ह	* * * * * * * * * * * * * * * * * * * *	To do the	6 6 35 Oc 8

23. In Canada all accounts were kept in pounds, shillings, pence, and arthings, previous to the adoption of the decimal coinage by Act of Provincial Parliament in 1858. In the United States also accounts were similarly topt prior to the adoption of Federal Money in 1786. In the States, at the time Federal money was adopted, the Colonial currency or bills of credit had become more or less depreciated in value, i. e., a colonial shilling was worth less than a shilling sterling, a.c., and the depreciation in value being greater in the surrencies of some colonies than in others gave rise to the different values of the present old currencies of the different States.

### TABLE OF CURRENCIES

### IN CANADA AND THE UNITED STATES.

In Canada, Nova Scotia, New Brunswick, &c.,	\$1 = 58. or £1.
In N. Y., N. C., Ohio, and Mich.,	\$1 = 8s. or £.
In N. Eng., Va., Ky., Ten., Ia., Ill., Miss.	
Missouri	$1 = 68$ . or $\mathcal{L}_{10}^{s}$ .
	1 = 78. 6d. or £3.
In Georgia and S. C.,	1 = 4s. 8d. or £ 0.
in deorgia and S. C.,	1 - 40. Ou. or -80.

Nors.—The remaining States use the Federal money exclusively.

23. To reduce dollars and cents to old Canadian Currency, or to any State Currency:—

### RULE

Multiply the given sum by the value of \$1 in the required currence expressed as a fraction of a pound. The product will be passed and decimals of a pound.

der (Art. 58, Sect. IV.) decimals to shillings, pence, and

. \$15.60

183

1.06 .181

-21 .90 . 40

1.84 40 17-441 11

ence, and ct of Pro-were simi-he States, or bills of out out of out of n in value ave rise to States (PAGE )

> or £1. or £3.

MI SO STALL IN

STATE IN

or £36 or £1.

y exclu-

an Cur-

ering"

red cur-

EXAMPLE 1 .- Reduce \$498.72 to Old Canadian Currency. OPERATION.

406-71×1= £135-46 = £126 80, 71d. Ans.

EXAMPLE 2 .- Reduce \$749.80 to New England Currency.

OPERATION.

740'80×4,= £224'94 = £224 18s. 97d. Ans.

EXAMPLE 3.-Reduce \$1111.11 to New York Currency.

OPERATION.

1111'11×4 = 2444'444 = 2444 80, 101 4d, Anc.

### EXERCISE 128.

- 1. Reduce \$1974.80 to New Jersey Currency. Ans. £740 11s.
- 2. Reduce \$765.43 to Michigan Currency. Ans. £306 3s. 576d.
- 3. Reduce \$7172.19 to Old Canadian Currency.

Ans. £2043 0s. 11ad.

24. To Reduce Old Canadian Currency or any State Currency to dollars and cents:-I had be the me or a year 1 %.

Express the given sum decimally and divide it by the value of a dollar expressed as a fraction of a pound; the quotient will be dollars, cents, &c.

EXAMPLE 1 .- Reduce £179 18s. 4ad., Old Canadian Currency, to dollars and cents.

2179 18s. 43d. = 2179 9197916 and 179 9197916 + 1 = \$719 67916. Ans. Norm.—Old Canadian Currency may be most expeditiously reduced to dollars and cents by the rule given in Art, 80, Sect. I.

EXAMPLE 2. Reduce £234 18s. 91d., Ohio Currency, to dollars and cents. OPERATION TO SELECT THE PROPERTY OF THE PROPER

£234 18s. 91d.=£234 9385416 and £34 9385416 + 1 = \$587 84635416. Ans.

### EXERCISE 129.

- 1. Reduce £743 18s. 11d., New England Currency, to dollars Ans. \$2479.8194.
- 2. Reduce £119 9s. 8id., Maryland Currency, to dollars and
- 3. Reduce £473 17s, 12d., Georgia Currency, to dollars and

## 25. To reduce dollars and cents to sterling money :-

### BULB.

Divide the given sum by the value of £1: sterling (\$4.8674), the quotient will be pounds sterling and decimals of a pound.

Reduce the decimal part (Art. 58, Soct IV) to shillings and pence.

Example.—Reduce \$749.83 to sterling money.

### OPERATION.

### 740-88 :4-867=2154-0541=2154 18, 81d, Ans.

### Exercise 130.

1. Reduce \$1006.90 to sterling money. Ans. £206 17s. 73d.

2. Reduce \$916.87 to sterling money. Ans. £188 7s. 84d. 8. Reduce \$2114.81 to sterling money. Ans. £434 10s. 42d.

## 26. To reduce sterling money to dollars and cents:

### RULE,

Express the given sum decimally and multiply by the legal value of £1 sterling (\$4.867).

EXAMPLE.—Reduce £78 11s. 47d. to dellars and cents.

### OPERATION.

278 11s. 41d.=278'5697916, and 78'5697916×4'867=\$382'398. Ans.

### 

- 1. Reduce £2043 11s. 3d. sterling to dollars and cents.
- 2. Reduce £777 7s. 7d. sterling to dollars and cents.
- Ans. \$3783.50427.
  3. Reduce £557 19s. 51d. sterling to dollars and cents.

  Ans. \$2715.65418.

### MACHANGE. 1925 .

27. Exchange is a commercial term, denoting the payment of money by a person residing in one place to a person residing in another, by draft or bill of exchange.

28. A bill of exchange is a written order addressed to a person directing him to pay, at a specified time and place, a certain sum of money to another person or his order.

29. The person who signs the bill of exchange is called the drawer or maker of the bill.

374), the

nd pence.

17s. 73d. 7s. 81d. 10s. 47d.

nts:-

legal value

nts. 82:308. Ans.

946-01868.

783.50427. ts.

715:65418.

the payto a pernge.

dressed to time and on or his

e is called

80. The person on whom it is drawn is called the drawee, and, after he has accepted it, the acceptor.

31. The person to whom the money is directed to be

paid is called the payer.

32. The person who purchases the bill of exchange, i. e., the person in whose favor it is drawn, is called the buyer or remitter.

83. The person who has legal possession of the bill is

called the holder.

84. The acceptance of a bill or draft is a promise on the part of the drawee to pay it at maturity or the specified time. The usual mode of accepting a bill is for the drawee to attach his signature to the word "accepted," written either across the face of the note or on its back.

Norm.—A draft or bill of exchange should be presented to the drawer, for his acceptance, immediately on its receipt.

35. If the payee or holder of a bill or draft wishes to sell it or transfer it, he endorses it, i. e., he writes his name on the back.

NOTE.—If the endorser directs the bill to be paid to a particular person, the endorsement is call a special endorsement and the person therein named is called the endorses.

named is called the endorsee.

If the endorser simply writes his name on the back of the bill, the endorsement is called a blank endorsement.

When the endorsement is blank, or when the bill is made payable to bearer, it may be transferred from one to another at pleasure, and the drawee is bound to pay it to the holder at maturity. If the drawee or acceptor of a bill fail to pay it, the endorsers are responsible for the payment.

3.6. When the drawee of a bill refuses acceptance, or, having accepted, this to make payment when it becomes due, the bill is immediately protected.

37. A protest is a formal declaration in writing, made by a public officer called a Notary Public, at the request of the holders of the bill, notifying the drawer, endorsers, &c., of its non-acceptance or non-payment.

Nota.—If the drawer and endorsers are not notified within a reasonable time of the non-acceptance or non-payment of the bill, they are not re-

time of the non-acceptance or non-payment of the bill, they are not responsible for its payment.

When a bill is protested for non-acceptance, the drawer must pay it immediately, even though the specified time has not arrived.

38. The time specified for the payment of a bill varies, and is a matter of agreement between the drawer and buyer. Some are payable at sight, some at a certain number of days or months after sight or after date. In both cases it is customary to allow three days of grace.

39. Bills of Brohange are divided into inland and foreign bills. When both drawer and drawer reside in the same country, they are called inland bills or drafts; when in different countries, foreign bills.

Note,—Three bills are commonly drawn for the same amount, &c., and are called respectively the First, Second, and Third of Brohange, and tegether constitute a set. These are sent by different ships or conveyances; and when the first that arrives is accepted or paid, the others become void. This plan is adopted in order to avoid the delays which might arise from accidents, miscarriage, &c.

### FORM OF AN INLAND BILL OR DRAFT.

\$3000.

TORONTO, 1st July, 1859.

Ten days after sight, pay to the order of George McCallum, Esq., Three Thousand Dollars, value received, and charge the same to

RIDOUT & STEVEN.

Messrs. Hardman & Morris,
Bankers, Hamilton.

FORM OF A FOREIGN BILL OF EXCHANGE.

Exchange 8000 francs.

TORONTO, 17th July, 1859.

At sixty days sight of this first of exchange (the second and third of the same date and tenor unpaid) pay to Edward Atkinson, Esq., or order, the sum of Eight Thousand Francs, with or without further advice.

JOHN HENDERSON.

Messrs. Duhamel & Beauharnois, Bankers, Paris.

40. The par of exchange is that amount of the money of one country actually equal to a given sum of the money of another, and is either intrinsic or commercial.

41. The intrinsic par of exchange is the real value of the money of different countries, as determined by the

weight and purity of their standard coins.

Thus, the English sovereign is intrinsically worth \$4.861 of the gold coin of the United States.

42. The commercial par of exchange is a comparison of the coins of different countries, according to their nominal or market value.

Thus, the English sovereign varies in market value from \$4.83 to \$4.85.

Note.—The intrinsic par is always the same so long as the standard coins are of the same kind, quantity, and quality of metal; the commercial par is determined by commercial usage, and fluctuates, being different at different times.

43. The Course of Exchange signifies the current price paid in one country for bills of exchange drawn on another.

Note.—The course of exchange is constantly fluctuating from various causes. When the exports of a country just equal its imports, the exchange will be at par; when the balauce of trade is against a place, i. e. when its imports exceed its exports, bills on foreign countries will be above par, because there will be a greater demand for them to pay the bills due abroad; when the balance of trade is in favor of a country, i. e. when its exports exceed its imports, bills of exchange on foreign countries will be below pay since fewer of them will be required.

The course of exchange can never very greatly exceed the *intrinsic par value*, because when the premium on bills of exchange becomes great it is less expensive to importers to pay for the insurance and transportation of bullion and coin to meet their payments than to transmit bills of exchange.

44. By an old act of Provincial Parliament it was enacted that £100 sterlings or 100 sovereigns should be equivalent to £1115 Canadian money, i. e. to \$444'444 or Einterling = \$4'444. It was found however that this was very much below the real or intrinsic value of the sterling pound, accordingly, while its legal value was only \$4.444, the market or commercial value varied from \$4.83 to \$4.86. By an act recently passed by the Provincial Parliament, the value of the pound sterling was fixed at \$4.866.

Now the new paris equal to the old par plus nine and a-half per cent. of the old par, that is, \$4.444 | 91 per cent. of \$4.444, which is .422, make \$4.866= the new par. Consequently the rate of exchange between Canada and Great Britain must reach the nominal premium or 91 per cent. before it is at par, according to the new standard.

45. Rates of exchange between Canada and Great Britain are commonly reckoned, at a certain per cent. on the old par of exchange, instead of on the new par.

EXAMPLE 1.—A merchant in Hamilton wishes to remit to London £749 3s. 6d. sterling; exchange being at 10 per cent. premium; how much must he pay for the bill of exchange?

OPERATION.

Old commercial par of £1 sterling = \$4 144

To which add 10 per cent. of itself = '444'

Gives price of £1 = 4'888

Then £740 3s. 6d.=£740 175 × 4.888 = \$3662.631. Ans.

EXAMPLE 2.—A merchant in Toronto wishes to remit 144479 francs to Paris, exchange being at a premium of 2 per cent. What will be the cost of his bill in dollars and cents?

OPERATION.

Commercial value of the franc = 186 cents. Add 2 per cent. \*372

Gives value for remitting = 18.972 " Then 18 972 × 144479 = \$27410 55588. Ans.

EXAMPLE 3.—What sum in dollars and cents will purchase a bill of exchange on Hamburg for 14667 marcs banco, exchange being at 11 per cent. discount?

OPERATION.

Commercial value of the marc banco = 35

Gives value for remitting = 34.475
Then 34.475 cents × 14867 = \$5056.448. Ans.

en.

lum,

e the

59. and. tkin-

ith or

noney

noney

lue of v the

old coin

son of minal

0 \$4'85. rd coins ifferent

other.

various xchange when its abroad exports

20

Di

to

at

and

1.

### Exercise 132.

1. If I wish to remit \$16785.25 to Paris, for how many francs and centimes can I obtain a bill—exchange being 5 francs 4 centimes to the dollar?

Ans. 84597 francs 66 centimes.

2. What is the cost of a bill of exchange for 4000 marcs banco at one per cent. above par?

Ans. \$1414.

3. How much must I give for a draft on New York for \$35678 at 21 per cent. premium?

Ans. \$36480.755.

4. What will a bill of exchange on St. Petersburg for 2560 rubles cost in dollars and cents, at 2 per cent. discount, the par being 75 cents per ruble?

Ans. \$1881-60.

5. What will be the cost of a bill of exchange on Great Britain for £800 sterling at 8 per cent, premium?

Ans. \$3840.00.

### Ale wit to Arbitration of Exchange. "Hornto on

rivital trad for class

46. Arbitration of exchange is the process of changing a given amount of the money of one country into an equivalent sum of the money of another, through the medium of one or more intervening currencies with which the first and last are compared.

NOTE.—Arbitration enables a person to accertain whether it is more advantageous to draw or remit a bill of exchange direct from one country to another or indirectly through other places.

47. When there is but one intervening country, the operation is termed simple arbitration; when there are two or more intervening countries, compound arbitration.

46. All question in arbitration of exchange may be solved by one or more statements in simple proportion; it is more convenient, however, to consider them as problems in Conjoined Proportion, and work them by the rule given in Art. 50, Sec. V.

Note.—Care must be taken to reduce all the money of the same country to the same denomination before linking them as directed in the rule.

EXAMPLE 1.—A merchant in Toronto wishes to remit 2000 marcs banco to Hamburg, and the exchange between Toronto and Hamburg is 35 cents for one marc banco. He finds, however, that the exchange between Toronto and Lisbon is \$1.08 for 1 milree, that between Lisbon and Paris is 6 milrees for 38 francs, and that between Paris and Hamburg is 19 francs for 10 marcs banco. How much will he gain by the circuitous exchange?

प अवार्षकी की रहे

### OPERATION.

STATEMEN	(P. 1 5	RANCE (	CANCELLED.
108 cents	= 1 milree.		Hart See
6 milraes	= 38 francs.	Prof 42341244	188 = 1
19 france	= 10 marcs	banco. 200/ 1	JE 176 200
2000 mares banco	= 4.2.11 - "C" 2"	banco. 200 1	A
THE PARTY OF THE P	= 200 X 3 X	The - bolto.	" . " . " . " . " . " . " . " . " . " .
$3000 \times 35 = $700.00 =$	what he has i	to new by dire	ct exchang

648:00 = what he has to pay by circuitous exchange.

Difference=\$ 52:00 = what he gains by the latter mode.

Example 2.—£824 Flemish being due to me at Amsterdam, it is remitted to France at 16d. Flemish per franc; from France to Venice at 800 france per 60 ducats; from Venice to Hamburg at 100d. per ducat; from Hamburg to Lisbon at 50d. per 400 rees; from Lisbon to England at 5s. 8d. sterling per milree; and from England to Canada at \$4-867 per £1 sterling. Shall I gain; or lose; and how much, the exchange between Canada and Amsterdam being 7s. 1d. Flemish per dollar?

### ्रा स्थानमंत्री र तथा वर्षा वर्षात्री है। कार्यो कार्यार्थ वर्षात्री पत्र अस्ति है है है

O'BEARON.
GAME CANCELLED
in Alice Ali
16d, Flemish = 1 franc. 18 = 1
50 - 50
300 francs = 60 ducats. 300 = 60
1 ducat = 100d. Flemish. $1 = 100$
50d. Flemish = $400$ rees. $50 = 400^{\circ}$
1000 rees = 68d. British, 1000 = 68 17
240d. British = \$4.867. 3296
x = 197760d. Flemish. x = 191760 19176
14 17 × 4967 × 3296 200 13 1 10 10 10 10 10 10 10 10 10 10 10 10 1
2 2797'074 - amount remitted
to off at 12.X DU to a 1 to an all the control of t
Then since exchange between Canada and Amsterdam is 7s. 1d. Flemish
per dollar we have
The state of the s

85d. Flemish == = 197760d. Flemish.

Here  $x = \frac{197760 \times 100}{2}$  = \$232658 = sum I should have received had it

been transmitted direct from Amsterdam to Canada.

Hence by the circuitous exchange I gain the difference between \$2727.07. and \$2326.58 that is \$400.401.

### EXERCISE 133.

1. If London would remit £1000 sterling to Spain, the direct exchange being 42 d. per plastre of 272 maravedie; it is asked whether it will be more profitable to remit directly, or to remit first to Holland at 35s. per pound; thence to France at 193d, per franc; thence to Venice at 300 france per 60 ducate; and thence to Spain at 360 maravedis per ducated Ass. The circular exchange is more advantageous by 103 piastres, 3 reals, 20 maravedis.

ncs les. aco

nes

IX.

678 55. 560

int, tain

ıg a iva-

n of and

more ntry ope-

or ved

nore con-Art.

ame d in

000 onto ver. or 1 ncs. BICE 2. A merchant wishes to remit \$4888.40 from Montreal to London, and the exchange is 10 per cent. He finds that he can remit to Paris at 5 francs 15 centimes to the dollar, and to Hamburg at 35 cents per marc banco. Now, the exchange between Paris and London is 25 francs 80 centimes for £1 sterling, and between Hamburg and London 132 marcs banco for £1 sterling. How had he better remit?

Ans. If he remits direct to London he will obtain a

bill for £1000.

If he remits through Paris he will obtain a bill for only £975 15s. 81d.

If he remits through Hamburg he will obtain a bill for £1015 15s. 5d.

Hence the best way to remit is through Hamburg, and the next best way is direct to London.

3. A merchant in Quebec wishes to remit 1200 marcs banco to Hamburg, and the exchange of Quebec on Hamburg is 35 cents for 1 marc. He finds the exchange of Quebec on Paris is 18 cents for 1 franc; that of Paris on London, is 25 francs for £1 sterling; that of London on Lisbon, is 180 pence for 3 milrees; that of Lisbon on Hamburg, is 5 milrees for 18 marcs banco. How much will he gain by the circuitous exchange?

> Ans. Direct exchange \$420; circuitous exchange \$375; gain \$45.

QUESTIONS TO BE ANSWERED BY THE PUPIL.

NOTE.—The numbers after the questions refer to the numbered articles of the section.

. What is profit and loss? (1)

w nat is profit and loss? (1)
 How do we find the total gain or loss on a quantity of goods when the cost price and selling price are given? (2)
 How do we find at what price an article must be sold so as to gain or lose a specified percentage, the cost price being given? (3)
 How do we find the rate per cent. of profit or loss? (4)
 How do we find the cost price when the selling price and the gain or loss per cent. are given? (5)
 What is barter? (6)
 What is alligation? (8)

7. What is parter? (a)
7. What is alligation? (8)
8. Into what rules is alligation subdivided? (9)
9. What is alligation medial? (10)
10. What is alligation alternate? (11)
11. How is alligation alternate proved? (13)
12. Give the different rules for alligation. (12, 14-16)
13. What is meant by the exchange of currencies? (17)

125. Give the different rules for alligation. (12, 14-16)
13. What is meant by the exchange of currencies? (17)
14. What is meant by the currency of a country? (18)
15. How is the intrinsic value of a coin determined? (19)
16. What fires the commercial value of a coin? (20)
17. How do you account for the fact that the \$\mathbf{s}\$ is of different values in the American States? (22)
18. Give the value of the pound currency in Canada, and in the different States. (22)

19. How do we reduce dollars and cents to old Canadian currency or to any state currency ? (28)

mult Thu

20.

32. E

35. W 36. B

by it powe Thu

3×8= 3.

third twice.

Thus factor, 4.

becau is exp by the to

t he lar, the

mes 137 ?

in a

in a

tain amt to

anco g is

ebec don, n, is

, is 5 n by

ange

ticles

n the

in or

20. 18.45 in or

n the

erent

any

- 20. How do we reduce old Canadian currency or any state currency to
- dollars and cents? (24)
  21. How do we reduce dollars and cents to sterling money? (25)
  22. How do we reduce sterling money to dollars and cents? (25)
  23. What is a bill of Exchange? (28)
- 24. Explain the terms drawer, drawes, acceptor, payes, holder, endorser, and endorses. (29-35)
  25. How is a bill accepted? (34)
  26. What is the difference between a blank endorsement and a special
- endorsement? (35)
- 27. What is meant by protesting a bill? (36, 37)
  28. Explain what is meant by the First, Second, and Third of Exchange.

- 29. What is the par of Exchange? (40)
  30. Explain the difference between the intrinsic par and the commercial par of Exchange. (41, 42)
  31. What is the course of Exchange? (43)
  32. Explain what is meant by saying the par of Exchange between Canada and Britain is 9\frac{1}{2} per cent. (44)
  33. Upon what is the rate of Exchange between Canada and Britain reckoned? (45)
  34. What is arbitration of Exchange? (46)
  35. What is the difference between simple and compound arbitration? (47)
  36. By what rule are questions in arbitration of Exchange worked? (46)

- 36. By what rule are questions in arbitration of Exchange worked? (48)

## SECTION X.

### INVOLUTION, EVOLUTION, LOGARITHMS, AND LOGARITHMIO ARITHMETIC.

1. A power of any number is the product obtained by multiplying that number by itself one or more times.

Thus  $25 = 5 \times 5$  is a power of 5;  $81 = 3 \times 3 \times 3 \times 3$  is a power of 8, &c.

2. The number which, being multiplied once or oftener by itself, produces the power, is called the root of that power.

Thus 5 is the root of 25, since  $5\times5=25$ ; 3 is the root of 81, since  $3\times8\times$  $3 \times 8 = 81.$ 

3. The powers of a number are called the first, second, third, fourth, fifth, &c., according as the root is taken once, twice, thrice, four times, five times, &c., as factor.

Thus, 81 is called the fourth power of 3, because 3 is taken 4 times as factor, in order to produce 81.

4. The second power of a number is also called its square, because a square surface, the length of one of whose sides is expressed by a given number, will have its area expressed by the second power of that number. (See Art. 62, Sec. I.)

5. The third power of a number is also called its cube; because if the length of one side of a cube be expressed by a given number, the solid contents of the cube will be expressed by the third power of that number. (See Art. 64, Sec. I.)

6. The index or exponent of a power is a small figure written to the right, indicating how often the root has to be taken as factor in order to produce the given power.

Thus, 21 = 2 = 2 = First power of 2.

22 = 2 × 2 = 4 = Second, power of 3.

23 = 2 × 2 × 2 = 8 = Third power of 3.

24 = 2 × 2 × 2 × 2 = 16 = Fourth power of 3.

24 = 2 × 2 × 2 × 2 × 2 = 5 = Firth power of 3.

So also 87 means the seventh power of 8; i. e., a number produced by taking 8 seven times as factor, &c.

7. (5+8) means that the sum of 5 and 8 is to be squared as one number and is a very different thing from 52+62, which means the sum of the squares of 5 and 8.

Thus  $(5+8)^2 = 13^2 = 169$ , while  $5^2+8^2 = 25+64 = 89$ .

Therefore  $(5+8)^2 = 25+80+64 = 18t$  part squared, plus twice product of 1st part by 2nd part, plus 2nd part squared.

- 8. The process of finding a power of a given number by multiplying it into itself is called Involution.
  - 9. To involve a number to any required power :-

### RULE.

in Jourisman and Take the given number as factor as many times as there are units in the index of the required power and find the continued product of these factors.

Norn. Fractions are implied by multiplying both numerators and denominators as above, and mixed numbers should be reduced to fractions before applying the rule.

EXAMPLE 1.—What is the fifth power of 7?

### OPERATION.

Here the index of the required power is 5 and hence the given number 7 must be taken 5 times as factor.

7×7×7×7×7=16667 And
EXAMPLE 2.—What is the third power of 1?

Ans. (3)3=3×3×3=64 Ans.

### EXERCISE 134.

- 1. Find the fifth power of 3.

- 3: Required the sixth power of 1:05.
- 4. Find the seventh power of ...
- 5. Find the fifth power of §.
- 6. Required the third power of 11%.
- Ans. 243.
- Aur. 1:340095640625.

- Ans.  $\frac{2187}{73136}$ .
  Ans.  $\frac{3187}{31065}$ .
  Ans.  $\frac{187}{148}$ .  $= 1481 \frac{68}{14}$ .

ARTS.

3.10.

multi

11.

Ther He

power diviso

12.

He multip requir Thus.

1. Wini

2. Divi 3. Find

4. Find

5. Find

13. of a gi

NOTE. find a por find the r

powers another

two me

ibe: l by EX-

gure as to

LAST MY ed by umber queres

roduct

mber

re are

tinued . HATTY PR rators educed

umber

0000.

81 ff.

10. Let it be required to find the product of 4° by 4°.  $4^{2} = 4 \times 4 \times 4$  and  $4^{2} = 4 \times 4$ . Therefore  $4^{3} \times 4^{3} = (4 \times 4 \times 4) \times (4 \times 4)$ =  $4 \times 4 \times 4 \times 4 \times 4 = 4^{3} = 4^{3} + 2$ .

Hence two or more powers of the same number are multiplied together by adding their indices or exponents.

Thus, 6°×6°×6° = 65 + 2 + 3 = 610. 5×5° × 5° × 5° = 51 + 2 + 3 + 7 = 513, 20, 20.

11. Let it be required to divide 35 by 33.

Therefore 85 +82 = 35 8×8×3×3×3 =3×8×8=8 = 86.4.

Hence, to divide one power of a number by another power of the same number, we subtract the index of the divisor from the index of the dividend.

Thus, 75-73-75-3-78 311-34=311-4=37, &c., &c., &c.

12. Let it be required to find the third power of 72.  $(72)^{3}=7^{2}\times7^{2}\times7^{3}=7\times7\times7\times7\times7\times7\times7=7^{6}=7^{2}+3$ 

Hence to find any required power of a given power, we multiply the index of the given power by the index of the required power.

Thus, (24)5=24+5=230; (33)7=33 x 7=314, &c., &c.

## Exprose 135.

1. Multiply together 42,40,46, and 47, and in the same 12.

on throw site i self

3. Find the fifth power of 33.

Ans. 315.

4. Find the value of  $\{(7^4 \times 7^3) \div (7^9 \times 7^2)\}^6$ 

5. Find the value of  $\{(5^3 \times 5^4 \times 5^{11} \times 5^9) \div (5^3 \times 5^2 \times 5^7 \times 5^9)\}^{\frac{3}{2}}$ Twenty of the transaction of the hollow days Ans. 550

### in any of Evolution. The footing of congil

13. Evolution is the process of finding any required root of a given power. White at the bear there

Note.—Evolution is the reverse of involution; the latter teaches how to find a power of a number by multiplying it into itself; the former, how to find the root of a power by resolving it into equal factors. It follows that powers and roots are correlative terms. If one number is a power of another the latter is a root of the former.

14. A root of a number may be indicated by either of two methods.

ART

begin

place the q

left-

to th

11

II

I

V dend. obtai

V

in th

rema

nd IV TRIAL

until

requir

No: canno

Ex

22

16

87)6

948)

9465

many

Bearin

and in that that the

ives 6

bring

Agai double

exclus

in the Mul the ne

is 946 the rig

478 is 1 PRO

1st. By using  $\sqrt{\ }$ , called the radical sign (Lat. radia, a root).

2nd. By using a fractional index having unity for its numerator, and the number expressing the degree of the root for denominator.

Thus. The square root of 7 is expressed either by 4/7 or by 71. The cube root of 6 is The seventh root of 2 is 2 or by 21.

Norm.—The figure placed in the radical sign, or as denominator of the fractional index denotes the root.

A fractional index with numerator greater than one is sometimes used; in such cases the denominator denotes the root, and the numerator the power to be taken.

Thus, 27 means either the cube root of the square of 2 or the square of the cube root of 2.

The radical sign  $\sqrt{a}$  corrupted form of the letter r, the initial letter of the Latin word radix, "a root."

### EXERCISE 136.

1. Express the square root of 17 and the cube root of 11.

Ans. 17 or 17 and 1/11 or 11

THE THE PROPERTY OF A CONTRACT 2. Express the fifth root of 4. Ans. \$\frac{1}{4}\$ or 4\$

3. Express the fourth root of 53 Ans. 1/53 or 54

4. Express the sixth root of  $7^4$ . Ans.  $\sqrt[6]{7^+}$  or  $7^6 = 7^5$ 

5. Express the third power of the fifth root of 1. Ans. (1/2) or 2

6. Express the eleventh power of the tenth root of 161.

Ans. (10/161)11 or 16110

15. Let it be required to extract the fifth root of 315.

The fifth root of  $3^{15}$  is expressed either by  $2/3^{15}$ , or by  $3^{15}$ .

Taking the latter mode, we have  $3^{\frac{15}{8}} = 3^3 = 3^{15} \div 5$ .

Hence, to extract any root of a given power of a number we divide the index of the power by the index of the root

> Thus, The seventh root of 214 is 214-7-22 The fourth root of 212 is 212:4-23, &c., &c.

### EXTRACTION OF THE SQUARE ROOT.

16. To extract the square root of a number, is to find a number which, being multiplied once by itself, will produce the given number.

idia, a

for its

of the

or of the

es used :

rator the

square of

letter of

### BULB.

I. Point off the given number into periods of two figures each, beginning at the decimal point,

II. Find the highest square contained in the left-hand period and place its root to the right of the number, in the place occupied by the quotient in division.

III. Subtract the square of the digit put in the root, from the left-hand period, and to the remainder bring down the next period to the right, for a new dividend.

IV. Double the part of the root already found for a TRIAL DIVISOR. V. Find how many times the trial divisor is contained in the dividend, exclusive of the right-hand digit, and place the figure thus obtained both in the root and also to the right of the trial divisor.

VI. Multiply the divisor thus completed by the digit last put in the root; subtract the product from the dividend, and to the remainder bring down the next period for a new dividend.

VII. Again, double the part of the root already found for a new TRIAL DIVISOR; proceed as in V. and VI., and continue the process until all the periods are brought down.

Note.—If the given number is not a perfect square, its exact square root cannot be found; but by annexing periods of ciphers, we can obtain any required approximation to it.

EXAMPLE 1.—What is the square root of 22420225?

22420225(4785, is the required root.

87)643 609

943)3302 2829

9465)47325

EXPLANATION.—Here 22 is the left hand period, and the highest square in 22 is 16, of which the square root is 4. We place 4 in the root and subtract 16 from 22 This leaves a remainder 6, to which we bring down the next period, 42, and thus obtain 642 for the new dividend. Our next step is to find the trial divisor, which we obtain by doubling the part of the root already found. This gives us 8,

by doubling the part of the root already found. This gives us 8, (= 4 doubled) and we ask how many times 8 will go into 64 (the dividend exclusive of the right hand digit). Bearing in mind that we are to put the digit thus obtained both in the root and in the divisor, and that the completed divisor will be over 80, we find that the required digit is 7, which we accordingly place both in the root and in the divisor. The complete divisor is 87, which multiplied by 7, gives 609, and this subtracted from 642, gives a remainder 33, to which we bring down the next period, 02, and thus get 3302 for the next dividend.

Again, doubling the part of the root already found, we obtain 94 (= 47 doubled) for a trial divisor, and as this will go into 330 (the dividend exclusive of the right hand digit) 3 times, we place 3 both in the root and

in the divisor.

Multiplying the 943 thus obtained by 3; subtracting and bringing down the next period, we get 47325 for the next dividend. The next trial divisor is 946 (=473 doubled) which will go into 4732 (the dividend exclusive of the right hand figure) 5 times; and we therefore place 5 both in the root and in the divisor. Multiplying and subtracting, we find no remainder. 473 is therefore the square root of 22420235.

PROOF,-4735×4735=22420225.

1 or 11

number he root

to find a produce

E give num I. since thou squa high

whic

re n tens

perio

enter there meht III

steps 1

oI.a decim II. decim

II.

### EXPLANATION AND REASON.

17. We may consider every number as consisting of its tone plus its units; has in if the tone be represented by the letter a and the units by the letter b. Number = a+b; and Number squared  $= (a+b)^2 = a^2 + 2ab + b^2$ .

Hence the square of a number is equal to the square of the tens, plus twice the product of the tens by the units. plus the square of the units.

Thus, 69 = 60+9 And (69) 2=(60+9) 3=(60) 3+2+60×9+93=3600+1080+81=4761.

18. Let it now be required to extract the square root of 4761.

It is evident that the square of a number consisting of a single digit can nevel contain more than two digits or less than one; conversely the square root of a number of one or two digits must be a number of one digit. Again the square of a number consisting of two digits can never contain more than four or less than three digits; conversely the square root of a number of three or four digits must be a number consisting of two digits. Similarly, the square of a number consisting of three digits can contain neither more than six nor less than five digits, and conversely, the square root of a number consisting of two or six digits, must be a number of three digits do; that is, one digit in the root is equivalent to two digits in the square, or conversely, two digits in the square are equivalent to one digit in the root.

Hence, if we divide the given number into periods of two figures each beginning at the decimal point, the number of periods will indicate the number of digits in the root

II. Taking the number 4761, we divide it into periods, thus, 4761, and since there are two periods in the square there must be two digits in the root. We thus learn that 4761 is the square of a certain number of tens, plus a certain number of units. Now it is manifest that the square of the tens can only be found in the second period, 47, since tens squared can give no digit of a lower order than hundreds. Also, that no part of the square of the units can be found in the second period, 47, since any single unit what are give no digit of a higher order than tens.

Therefore the square of the units is found only in the first or lowest period, the square of the tens only in the second period, the square of the hundreds only in the third period, &c.

" OPERATION! A !! A !!

things blood to water out to a

derivate it is both

4761(69 = square root. 36 = highest square in 2nd period.

6 teas ×2=12 teas+9 units=129) 1161 = remainder which contains, lat, The same of the sa

III. In extrapting the aguare root of this number, we look first for the digit occupying the place of tens in the root. We know (II.) that the square of tens is contained in the second period, 47, and the highest square contained in 47 must be the square of the highest square in 47 is 36, the square root of which is 6. Placing 36 under the 47, 8 in the root, we subtract and bring down the next period, 61, and thus get a total remainder of 1161. Now

ARTS. 17-19.7

s units; letter b.

are of units.

f 4761. digit can de square Again in more imilarly, er more

the root. of two of pe-

gipe, sho;

4761, and the root. the tens n give no square of ngie unit

first or period,

period.

sins, lat, tens by

the digit quare of outsined nd in the (Art. 17) the whole number 4761 consists of the square of the tens, plus twice the product of the tens by the units, plus the square of the units; and since we have subtracted from it. So, (or if the square of the units; and shoe we have subtracted from it. So, (or if the square of the units; and the tens the remainder, 1161, must contain twice the product of the tens by the units, plus the square of the units; that is, twice 6 tens × by a certain number of units, plus the square of that number of units; and because we do not know as yet wint the units. Bure of the root is, we use twice the tens for a trial divisor.

IV. Since we are now seeking the units digit of the root, and since tens multiplied by units can give no digit of a lower order than tens, the right hand digit of the divided dan form no part of twice that product of the tens by the units, and we have simply to ascertain how often 13 tens by the units, and we have simply to ascertain how often 13 tens of the root, and in the divisor, because the dividend contains not only twice the product of the tens by the units, but also the square of the units. Now when we multiply the 9 by 9 we get the aquare of the units, and when we multiply the 13 tens by 9 units, we get twice the product of the tens of the troot by the units.

Example 2.—Extract the square root of 127449.

OPERATION.

12740 (357

(265)874

\$2 " AL . 16 | 707)4949 .

EXPLINATION AND RHASON.—From the pointing off we learn that the given number is the square of a certain number of hundreds, plus a certain

number of tens, plus a certain number of units.

I. We are first then to look for the digit in the place of hundreds, and

I. We are first then to look for the digit in the place of hundreds, and since hundreds squared can give no digit of a lower order than tens of thousands or of a higher order than hundreds of thousands, we see that the square of the hundreds can be found only in the left hand period. The highest square contained in the left hand period is 9, the square root of which is the left hand digit of the entire root.

II. After subtracting, we bring down the next period only, because we are now looking for the digit in the place of tens in the root. And since tens squared can give no digit of a lower order than hundreds, the lowest period cannot enter into any part of the square of tens, much less can it enter into any part of twice the product of the hundreds by the tens, and therefore when locking for the tens of the root, we pay no attention to the might hand period of the square.

III. The remainder of the process is similar, and the reason for the various steps the same as in example 1.

steps the same as in example 1.

19. To extract the square root of a decimal:

h digar of the car at books

I. Annex one cipher, if necessary, in order that the number of decimal places may be even. Ly ... i v. Hfa o is he.

II. Point off into periods of two figures each, beginning at the decimal point, and extract the square root as in whole numbers.

remembering that the number of decimal places in the root will be equal to the number of periods in the equare.

### Express 137.

	Extract the se			Ans.	
3.	Extract the se	quare root of	954064.	Ans.	
10	81 st + 5 - 4	6 9 2001	12 h 1 n 2		2.28606

5. Extract the square root of 5 true to six decimal places.

6. Extract the square root of 60.487129. Aug. 7.777.

7. Extract the square root of 79792266297612001.

Ans. 282475249.

1. 1 2. I

3. I 4. H 5. H

sequ

ang

20. To extract the square root of a fraction :-

### RULE

I. Reduce mixed numbers to improper fractions, and compound and complex fractions to simple ones, and the resulting fraction to its lowest terms.

II. Extract the square root of both numerator and denominator separately, if they have exact roots; but if they have not both exact roots, reduce the fraction to its corresponding decimal, by Art. 56, Sec. IV., and then extract the root as in Art. 19.

Example 1 .- Extract the square root of 21.

### OPERATION

Ans. 
$$2\frac{1}{4} = \frac{2}{4}$$
 and  $\sqrt{2} = \frac{1}{4} = \frac{1}{4}$ .

EXAMPLE 2.—Extract the square root of 37.

Char to a wast

## in the sect of the first to the contract of th

34 = 3.42857142 and \3.42857142 = 1.8516.

## EXERCISE 138.

1.	Find	the square	root of		THE PRESENTANT	Ans.	16 m.
2.	Find	the square	root of	787.		Ans.	
3.	Find	the square	root of	51.	1 8		267786.
4.	Find	the square	root of	\$ 17. AMERICAN			63509.
5.	Find	the square	root of	131.	· SI S ASSIGNA	Ame 1	1.63318

21. Let it be required to extract the square root of 63513-423 applenary.

ill be

T. X.

06.

06.

15249. 11.

pound tion to

inator h exact ert. 56.

1.8516.

267786. 3509. 83318. 513-423

43)235

466) 4318 4161

5051)128'42 50'51

505 '95)41 '6180 84 '8564

505 '335) 4 '223300

453628

OPERATION.

EXPLANATION.—We point off into periods of two places each, as in the decimal or common configuration.

Then the highest square in 6, the first period, is 4, of which the square root is 2. Subtracting 4 from the 6 and bringing down the next period, 35, we get 235 for the dividend.

Next doubling the 2 we obtain 4, and we find that this will go into 23, the dividend exclusive of the right hand figure, 3 times. Placing this 3 in both root and divisor, multiplying (bearing of the right hand figure, 3 times. Figure this 5 in both root and divisor, multiplying (bearing in mind that 7 is the common ratio of the system) and subtracting, we obtain a remainder of 43, to which we bring down the next period, 18, and thus get 4313 for the next dividend, &c.

Example. Extract the square root of 4731392 und vary true to two places to the right of the separating point.

OPERATION.

4731892(2182.99. Ans.

428)3213

4352) 11592

3594 '64 3594 '64

4355 · 79) 404 · 0700 359 · 5744

55 \*5#67

EXERCISE 139.

- 1. Extract the square root of 11333311 septenary. Ans. 2626.
- 2. Extract the square root of 33233344 senary. 3. Extract the square root of 4234-10123 quinary. Ans. 43-412.
- 4. Extract the square root of 88888 888 nonary. Ans. 888.88
- 5. Extract the square root of 248664et69 duodenary. Ans. 54373.

### APPLICATION OF SQUARE ROOT.

22. A triangle is a figure having three sides, and consequently three angles. When one of the angles is a right angle, like the corner of a square, the triangle is called a right angled triangle.

13

23. In a right angled triangle the side opposite the right angle is called the hypothenuse, and the sides containing the right angle, are called the base and the perpendicular.

24. It is shown by elementary geometry that the square described on the hypothenuse of a right angled triangle is equal to the sum of the squares described on the other two sides.

Or if h be the hypothenuse, b the base, and p the perpendicular; then

$$h^2 = b^2 + p^2$$
, and hence

$$h = \sqrt{b^2 + p^2}$$

$$b = \sqrt{h^2 - p^2}$$

$$p = \sqrt{h^2 - b^2}$$

That is—to find the hypothenuse of a right angled triangle when the other sides are given we add the square of the base to the square of the perpendicular and extract the square root of the sum.

To find the length of the base we subtract the square of the perpendicular from the square of the hypothenuse and extract the square root of the remainder.

To find the length of the perpendicular we subtract the square of the base from the square of the hypothenuse and extract the square root of the remainder.

25. The following principles are also established by geometry:—

Circles are to each other as the squares of their diameters.

If the diameter of a circle be multiplied by 3.1416, the product is the circumference.

If the square of half the diameter of a circle be multiplied by

3.1416, the product is the area.

If the square root of half the square of the diameter of a circle be extracted, it is the side of an inscribed square.

If the area of a circle be divided by 3.1416, the quotient is the equare of half the diameter.

Example 1.—If the hypothenuse of a right angled triangle is 12 feet long and the base 10 feet, how long is the perpendicular?

OPERATION.

 $12^2 = 144$   $10^2 = 100$ 

difference = 44 and  $\sqrt{44}$  = 0.63384. Ans.

EXAMPLE 2.—If the foot of a ladder be placed 20 feet from the side of a house, how long must it be in order to reach to the top of the house, the latter being 46 feet high? ht ng

r. re is

WO.

hen

hen uare

peruare

e of uare

by

ct is

ircle

the le is

lar?

rom

OPERATION. 462 = 2116 202 = 400

sum = 2516 and  $\sqrt{2516}$  = 50.15. Ans.

### Exercise 140.

- 1. Suppose a ladder 100 feet long be placed 60 feet from the foot of a tree; how far up the tree will the top of the ladder reach?

  Ans. 65 feet.
- 2. Two persons start from the same place, and go, the conduction north 50 miles, the other due west 80 miles. How far apart are they?

  Ans. 94-34 miles, nearly.

3. How large a square stick of timber can be hewn from a round stick 24 inches in diameter? Ans. 16.97 in. to the side.

- 4. A man has a ladder 36 feet long, which, when put on the outside of a ditch 20 feet wide, exactly reaches the top of the wall. Required the height of the wall. Ass. 29:933.
- 5. A ladder 40 feet long is placed against a wall 14 feet high, and just reaches the top; it is then turned over and touches the top of another wall 26 feet high. Required the breadth of the street.

  Ans. 22.622 yds.

6. If the area of a circle be 1760 yards, how many feet must there be in the side of a square to contain that quantity?

Ans. 125.857.

7. A certain general has an army of 141376 men. How many must he place in rank and file to form them into a square?

8. What is the distance through the opposite corners of a square yard?

Ans. 4:24264 feet.

9. The distance between the lower ends of two equal rafters, in the different sides of a roof, is 32 feet, and the height of the ridge above the foot of the rafters is 12 feet. What is the length of a rafter?

Ans. 20 feet.

10. What is the distance measured through the centre of a cube from one corner to its opposite corner, the cube being 3 feet, or 1 yard, on a side?

Ans. 5-196 feet.

11. If an iron wire to inch in diameter will sustain a weight of 450 pounds, what weight might be sustained by a wire an inch in diameter?

Ans. 45000 lbs.

12. What length of repe must be tied to a horse's neck, in order that he may feed over an acre?

Ans. 7:136+perches.

13. Four men A, B, C, D, bought a grindstone, the diameter of which was 4 feet; they agreed that A should grind off his share first, and that each man should have it alternately until he had worn off his share; how much did each man grind off?

Norm:—In this question we disregard the thickness of the grindstone.

After the first has ground off his portion, there will remain t of the stone

Then the whole stone: part remaining::square of diameter of whole stone:square of diameter of part remaining. (Art. 25)

That is, 1:  $\frac{1}{4}$ ::  $4^2$ :  $4^2$ :  $4^2$ :  $4^2$ : and hence  $4^2$ : 4

Similarly, after the second has ground off his portion there will remain is of the stone, and after the third has taken his portion, is of the stone.

Hence 1:  $\frac{1}{2}$ ::  $4^2$ ::  $x^2$ , whence x=4  $\sqrt{\frac{1}{2}}=2.828$  ft. = diameter after 2nd has taken his portion.

1:  $\frac{1}{4}$ :  $\frac{4^2}{4^2}$ :  $\frac{4^2}{4^2}$ : whence  $x = 4 \times \sqrt{\frac{1}{4}} = 2$  ft. = diameter after 3rd has taken off his portion.

Hence A takes off  $\frac{4}{4}$ : 464 = 536 ft. = 6.432 inches.

Hence A takes off 4—3·464 = '536 ft. = 6·492 inches.

B " 3·464—2·828 = '636 ft. = 7·632 inches.
C " 2·828—2 = '828 ft. = 9·936 inches.
D " remaining 2 ft. = 24 inches.

### CUBE ROOT.

26. To extract the cube root of a number is to find a number which taken three times as factor will produce the given number:—

### RULE

I. Point off the number into periods of three figures each beginning at the decimal point.

II. Find the highest cube contained in the left hand period and place its root to the right of the number, in the place occupied by the quotient in division.

III. Subtract the cube of the digit put in the root from the left hand period, and to the remainder bring down the next period to the right for a new dividend.

IV. Multiply the square of the part of the root already found by 300 for a TRIAL DIVISOR.

V. Find how many times the trial divisor is contained in the dividend and put the figure thus obtained in the root.

VI. Complete the TRIAL DIVISOR by adding to it:

Griffing Chang.

1st. The part of the root previously found x the last digit put in the root x 30 and

2nd. The square of the last digit put in the root.

VII. Multiply the divisor thus completed by the digit last put in the root; subtract the product from the dividend, and to the remainder bring down the next period for a new dividend.

VIII. Again multiply the square of the part of the root already found by 300 for a new TRIAL DIVISOR, find what digit to place next in the root as in V, complete the divisor by making the two additions to the trial divisor described in VI, multiply, subtract and bring down as directed in VII, and continue the process until all the periods are brought down.

X

ole

n 🛊 🦠 ter en

he

in-

undthe

eft

to

by

the

but

the

dy

ace

wo

rct

### EXAMPLE. - What is the cube root of 429172932007? OPERATION.

	429172932007 7543 Ans.
1st trial divisor = 7° × 300= 14700 1st increment = 7 × 5×30= 1050 2nd " 5° = 25	86172 = 1st dividend.
1st complete divisor = 15775	78875 = product of comp. div.
2nd trial divisor = 75 <sup>2</sup> × 300= 1687500 1st increment= 75×4×30 = 9000 2nd " = 4 <sup>2</sup> = 16	7297932 = 2nd dividend, 2014
and complete divisor = 1696516	6786064—product of comp. div.
3rd trial divisor = 7542 × 800 = 170554800 1st increment = 754 × 3×30 = 67860 2nd " = 82 = 9	511868007 = 3rd dividend.
3rd complete divisor =170622669	511868007 = product of comp. div. by 8.

EXPLANATION.—After pointing off we find that the highest cube number contained in the left hand period is 343, of which the cube root is 7. We therefore place 7 in the root and subtract 343 from the first period. This gives us a remainder of 86, to which we bring down the next period 178, and thus obtain 86172 for a new dividend.

Next we take 7, the part of the root already found, square it and multiply the 49 thus obtained by 300, this gives the first trial divisor 14700 which we find will go into the dividend 86172 (…aking due allowance for the increase of the divisor) 5 times.

Next we complete the divisor by adding to it

1st, 7×5×30=1050, and 2nd, 5<sup>2</sup>=25 which gives us

15775 for a complete divisor. This we multiply by 5, the digit last put in the root, subtract the product 78875 from the 1st dividend, and to the remainder 7297 bring down the next period 932, &c., &c.

27. EXPLANATION AND REASON.—We have seen (Art. 17) that we may consider every number as consisting of its tens plus its units, or if a=tens consider even then and b=units, then Number = a+b; and Number = a+b;

Number cubed =  $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ .

Hence the cube of a number is equal to the cube of the tens, plus three times the product of the tens squared multiplied by the units, plus three times the product of the tens multiplied by the square of the units, plus the cube of the units.

Thus 
$$69 = (60+9)$$
; and  $69^3 = (60+9)^3 = 60^3 + 8 \times 60^2 \times 9 + 8 \times 60 \times 9^2 + 9^3 = 21600 + 97200 + 14580 + 729 = 528509$ .

28. Let it now be required to extract the cube root of 328509.

I. It is manifest that the cube of a single digit can never centain more than three digits or less than one digit, and hence the cube root of a number (i. e., perfect cube) of one, two or three digits must be a number of one digit. Again the cube of a number consisting of two digits can never contain more than six or less than four digits, and conversely the cube root of a perfect cube consisting of four, five or six digits must be a number of two digits. Similarly the cube root of a perfect cube consisting of seven, eight or nine digits must be a number of three digits, &c.

Hence, one digit in the root is equivalent to three digits in the cube, and conversely three digits in the cube are equivalent to one digit in the root, and therefore if we divide the given number into periods of three digits each, beginning at the decimal point, the number of periods will indicate the number of digits in the root.

II. The cube of the units can be found only in the period immediately to the left of the decimal point, since any unit cubed can give no digit of a higher order than hundreds. Also the cube of the tens can be found only in the second period to the left of the decimal point, since tens cubed can give no digit of a higher order than hundreds of thousands, or of a lower order than thousands. Similarly the cube of the hundreds can be found only in the third period to the left of the decimal point, &c.

Hence, counting from the decimal point towards the left, the cube of the units can be found only in the first period, the cube of the tens only in the second period, the cube of the hundreds only in the third period, &c.

III. Taking the number 328509 we divide it into periods, thus 328509, and since there are two periods in the cube there must be two digits in the root. We thus learn that 328509 is

0PERATION.

328509(69
216

62 = 36 × 300 = 10800|112500.

6×9=54 × 30 = 1620|
92 = 81

root. We thus learn that 328509 is the cube of a certain number of tens plus a certain number of units. We first then look for the digit in the place of tens in the root. We know (II) that the cube of the tens is contained in the second period 328, and the highest cube contained in 328 must evidently be the cube of the highest digit that can occupy the place of tens in the root—which digit we are seeking. The highest cube

second period 328, and the highest cube contained in 328 must evidently be the cube of the highest digit that can occupy the place of tens in the root—which digit we are seeking. The highest cube contained in 328 is 216, of which the cube root is 6. We then subtract 216 from 328 and to the remainder bring down 509, the next period, which IV. From the given number we have only subtract.

IV. From the given number we have only subtracted 216 (or if the ciphers be affixed, 216000) the remainder, 112509 therefore consists (Art. 27) of three times the product of the square of the tens by the units, plus three times the product of the tens by the square of the units, plus three times the product of the tens by the square of the units, plus the cube of the units; that is, 112509 consists of (6 tens)  $^2 \times 3 \times a$  certain number of units; (6 tens)  $\times 3 \times$  (that number of units) + (that number of units)  $^3$ ; and because we do not knowns yet what the units figure is, we use (6 tens)  $^2 \times 3$  for a trial divisor.

But  $(6 \text{ tens})^2 \times 3 = (60)^2 \times 3 = (6 \times 10)^2 \times 3 = 6^2 \times 10^2 \times 3 = 6^2 \times 300$ ; or in other words, any number of tens squared, multiplied by 3, is equal to that same number of units squared and multiplied by 300. Hence we obtain the constant multiplier 300.

the constant multiplier 300.

V. 6<sup>2</sup> = 36, and this multiplied by 300 gives us 10800. In asking how often this is contained in 112509 we have to bear in mind that we must increase the trial divisor by the two additions indicated in the sixth section of the rule. Making allowance for these additions, we find the units' figure of the root to be 9.

nore umone ot of

raX.

r of ven, the t to

ımmal gits

tely of a only can wer

the ube eds

and the ) is ens We ace hat the ube the cuich

ube 216 ich ers ree nes the 8 nd X3

ıat in inon ta'

VI. If we were to multiply the 10600 we have obtained as a trial divisor by 9, the units figure of the root, we should only get three times the product of the square of the tens by the units; but we require also three times the product of the tens by the square of the units and lastly the cube of the units. Our complete divisor must therefore evidently consist of-

Three times the square of tens.

2nd. Three times the tens multiplied by the units.

2nd. The square of the units; or representing the tens by a and the units by b, the divisor must =  $3a^2 + 3ab + b^2$ , and this multiplied by b, the digit in the units' place will give  $(3a^2 + 3ab + b^2)b = 3a^2b + 3ab^2 + b^3 = \text{the dividend.}$ 

Now (6 tens)  $\times$  3 = (60)  $\times$  3 = 6  $\times$  10  $\times$  3 = 6  $\times$  30, i.e. the product of any number of tens multiplied by 3 is equal to the product of that same number of units multiplied by 30.

Hence we obtain the constant multiplier 30.

The additions we make then are  $6 \times 30 \times 9 = 1620$ , and  $9^2 = 81$ , and thus we obtain the complete divisor  $12501 = (60)^2 \times 3 + 60 \times 3 \times 9 + 9^2$ , and multiplying this by 9, we get

 $(60)^2 \times 3 + 60 \times 3 \times 9 + 9^2$   $9 = 60^2 \times 3 \times 9 + 60 \times 3 \times 9^2 + 9^3 = three$ times the square of the tens multiplied by the units, plus three times the tens multiplied by the square of the units, plus the cube of the units.

Note.—When there are more than two periods, the reasons are analogous, since we never have to do with more than tens and units of the root at one time; i.e., when we are seeking the second digit of the root, we call the first digit tens and the second, units; when we are seeking the third digit of the root, we consider the first two as so many tens, and the third as units, &c.

The reason for bringing down only one period at a time is similar to the reason for the same step in the extraction of the square root (for which

see Art. 18, Example 2).

## 29. To extract the cube root of a decimal:

I. Annex two ciphers, if necessary, in order to make the last period complete.

II. Point off into periods of three places each, beginning at the decimal point, and extract the cube root as in whole numbers, remembering that the number of decimal places in the root will be equal to the number of periods in the cube.

## EXERCISE 141.

1. What is the cube root of 62712728317?	Ans. 3973.
2. Extract the cube root of 1953125.	Ans. 125.
3. Extract the cube root of 1076890625.	Ans. 1025.
4. What is the cube root of .697864103?	Ans887.
5. What is the cube root of 102503.232?	Ans. 46.8.
6. Find the cube root of 179597.069288.	Ans. 56.42.
7. Find the cube root of 483.736625.	Ans. 7.85.
8. Find the cube root of .636056.	Ans 86.

30. To extract the cube root of a mixed number or a vulgar fraction :-

#### RULE

I. Reduce mixed numbers to improper fractions, and compound or complex fractions to simple ones, and the resulting fraction to its lowest terms.

II. Extract the cube root of both numerator and denominator separately, if they have exact roots; but if they have not both exact roots, reduce the fraction to its corresponding decimal by Art. 56, Sect. IV, and then extract the root as in Art. 29.

EXAMPLE 1.—What is the cube root of 33?

OPERATION.

$$\sqrt[8]{3\frac{3}{8}} = \sqrt[8]{\frac{27}{8}} = \frac{\sqrt[8]{27}}{\sqrt[8]{8}} = \frac{3}{8} = 1\frac{1}{2}$$
. Ans.

EXAMPLE 2.—Extract the cube root of 171.

1. Extract the cube root of 3.

OPERATION.

 $17\frac{1}{2} = 17:125$ , and  $\sqrt[8]{17:125} = 2:5/77$ , nearly.

#### EXERCISE 142.

2. Extract the cube root of 3.	Ans. :5609.
3. Extract the cube root of 1 of 21.	Ans. 941.
4. Extract the cube root of 283.	Ans. 3.063.
5. Extract the cube root of 32-8.	Ans. 3.198.
21 In autmosting the subs root of a numb	A to the

31. In extracting the cube root of a number in any scale, other than the decimal, we proceed in the same manner, pointing off into periods of three figures each, finding a trial divisor and afterwards completing it as in the preceding examples.

Note.—In all scales having a radix higher than 3, the constant multipliers are 300 and 30; but as in the binary and ternary scale we cannot use a digit so high as 3, these multipliers become respectively 1100 and 110 for the binary scale, and 1000 and 100 for the ternary scale.

Example 3.—Extract the cube root of 613412·132 septenary.

#### OPERATION.

```
\begin{array}{c} 6^2 = 51 \times 300 = 21300 \\ 6 \times 30 = 240 \times 5 = 1560 \\ 5^2 = 34 \\ \hline \\ 23224 \\ 152456 \\ \hline \\ 65^2 = 6304 \times 300 \\ 0 = 25215000 \\ 650 \times 30 = 26100 \times 4 \\ 143400 \\ 4^2 = 22 \\ \hline \\ 252323422 \\ \hline \\ 1402 \cdot 630621 \\ \hline \\ 290 \cdot 9013442 \\ \hline \end{array}
```

1. E

2. E

4. E

5. E

root t ing it proce multi

I.

II.
place
root,
plyin
col. I
tract
down

III I., m first visor

> dend, third by th divise the p next

vas j till ti

deci

rund

a to

ator

both

Art.

721. 309.

141.

)63.

198.

any

an-

ing

re-

ılti-

not

110

### EXERCISE 143.

- 1. Express one million in the senary scale and then extract its cube root.

  Ans. 244.
- 2. Extract the cube root of 6131271 octenary. Ans. 165.32.
- 3. Extract the cube root of 10221012:102 ternary.

Ans. 112.012.

- 4. Extract the cube root of teteet in the duodenary scale true to two places to the right of the separating point.

  Ans. e7-t2.
- 5. Extract the cube root of 421030.4412 quinary true to two places to the right of the separating point. Ans. 44.004.
- 32. Since many teachers prefer Horner's method of extracting the cube root to the common method, we shall give it here. Upon closely examining it the student will find that the reasons for the several steps of the process are identical with those given in Arts. 27 and 28. The constant multipliers 300 and 30 are still used, but in a disguised form.

#### RULE.

- I. Point off as in the common method.
- II. Find the greatest cube in the first period on the left hand; place its root, on the right of the number for the first figure of the root, and also in col. I. on the left of the number. Then multiplying this figure into itself, set the product for the first term in col. II.; and multiplying this term by the same figure again, subtract this product from the period, and to the remainder bring down the next period for a dividend.
- III. Adding the figure placed in the root to the first term in col. I., multiply the sum by the same figure, add the product to the first term in col. II., and to this sum annex two ciphers, for a divisor; also add the figure of the root to the second term of col. I.
- IV. Find how many times the divisor is contained in the dividend, and place the result in the root, and also on the right of the third term of col. I. Next multiply the third term thus increased by the figure last placed in the root, and add the product to the divisor; then multiply this sum by the same figure, and subtract the product from the dividend. To the remainder bring down the next period for a new dividend.
- V. Find a new divisor in the same manner that the last divisor was found, then divide, &c., as before; thus continue the operation till the root of all the periods is found.

EXAMPLE. What is the cube root of 783146, true to two decimal places.

No

EX

mu

Re

 $E_{3}$ 

extra

prim

2. W

3. E:

4. E

5. Ez

Thu

Exam

#### OPERATION.

Col. I.	Col. II. 16×4 =	78314'600, (42'78+.
2nd " 8	4800, 1st divisor )	14314
3rd " 122	5044×2 =	10088
4th " 124	529200, 2d divisor )	4226600
5th " 1967	538069×7 =	3766483
oth : 1274	54698700, 8d divisor	460117000
7th " 12818	54801244×8 =	438409952

EXPLANATION.—The cube root of the greatest cube in 78 is 4 which is placed in the root and also in column I, then multiplying this 4 by itself gives us 16 which is the 1st term in column II, and again multiplying this 16 by 4 gives us 64, the number which we are to subtract from the first period 78.

Subtracting and bringing down the next period 314 we get 14314 for the

next dividend.

Now adding 4, the figure placed in the root, to 4 the 1st term in col. I. gives us 8, the 2nd term in col. I, multiplying this 8 by the 4, i. e., the figure in the root, gives us 32 which we add to the 1st term of col. II, and affix two ciphers. We thus obtain 4800 the second term of col. II, which is our

We then find that 4800 goes 2 times in the dividend. This 3 we place in the root and also to the right of the sum of the 1st and 2nd terms of col. I. The 1st and 2nd terms of col. I, added together make 12 and the 3 of the root affixed makes 122, the third term of col. I. Then we multiply this 122 by 2, the last digit put in the root, this gives us 244 which we add to 4800, the second term of col. II. and thus obtain 5044, the 3rd term. Lastly this third term multiplied by 2, gives us the number to subtract, &c.

NOTE.—For examples in this method work any of the preceding questions.

tions.

## APPLICATION OF THE CUBE ROOT.

33. Principles Assumed.—I. Spheres are to one another as the cubes of their diameters.

II. Cubes and all other regular solids are to one another as the cubes of their like dimensions.

## EXERCISE 144.

1. If a cannon ball 3 inches in diameter weighs 8 lbs., what will be the weight of a ball of the same metal 4 inches in diameter?  $3^3:4^3::8$  lbs.: Ans. = 1839 lbs.

2. If a ball 3 inches in diameter weighs 4 lbs., what will be the weight of a ball that is 6 inches in diameter? Ans. 32 lbs.

3. If a globe of gold one inch in diameter be worth \$120, what is the value of a globe 31 inches in diameter? Ans. \$5145.

4. If the weight of a well proportioned man, 5 feet 10 inches in height be 180 pounds, what must have been the weight of Goliath of Gath, who was 10 feet 42 inches in height?

Ans. 1015.1 lbs.

5. A person has a cube of c. whose sides are 973 ft. long; he wishes to take out of the same 5 cubes whose sides are 45 feet, 62 feet, 30 feet, 80 feet, and 20 feet. He requires to know the length of the side of the cube that can be formed out of the remaining clay.

Ans. 972 69 ft.

6. What is the side of a cube which will contain as much as a chest 8 feet 3 inches long, 3 feet wide, and 2 feet 7 inches deep?

Ans. 47 9843 inches.

7. Four ladies purchased a ball of exceeding fine thread, 3 in. in diameter. What portion of the diameter must each wind off so as to share off the thread equally?

Ans. 1st lady must wind off 27432 inches.

3rd " " 49122 " 4th " 1.88988 "

Norm.—This question is solved by a method similar to that adopted in Example 13, Exercise 140.

## EXTRACTION OF THE ROOTS OF HIGHER ORDERS.

34. When the index of the root is a power of 2 or 3, or a multiple of any power of 2 by any power of 3—

#### RULE.

Resolve the given index into its prime factors.

Extract the root denoted by one of these factors, then of this root, extract the root denoted by another factor, and so on till all the prime factors be used.

Thus, for the 4th root extract the square root of the square root.

for the 6th root extract the cube root of the square root.

for the 8th root extract the square root of the square root of the

square root.

for the 13th root extract the cube root of the square root of the square root.

for the 16th root extract the square root four times.

for the 18th root extract the cube root of the cube root of the square root, &c., &c.

#### Exercise 145.

1. What is the fourth root of 19987173376?	11	Ans.	376.
2. What is the sixth root of 308915776?	2	Ans.	26.
3. Extract the ninth root of 40353607.	1	Ans.	7.
4. Extract the eighteenth root of 387420489.	1.10-	Ans.	≥ 3.
K Everent the twenty-seventh root of 124217728	4. Di	Ans	2.

4 which is 4 by itself plying this m the first

m in col. I. , the figure II, and affix hich is our

314 for the

we place in ms of col. I. the 2 of the ply this 122 add to 4800, rm. Lastly et, &c.

her as the

ther as the

lbs., what inches in 1834 lbs. vill be the 22. 32 lbs. 120, what 25. \$5145. inches in weight of height?

## LOGARITHMS.

85. The Logarithm of a number is the index of the power to which it is necessary to raise a given root or base, in order to produce the given number.

86. The Base of a system of logarithms is the fixed number to which all the logarithms of that system belong

as indices.

Thus  $10^3 = 1000$ ; here 3 is called the logarithm of 1000, to the base 10. So also  $2^5 = 32$ ; here 5 is called the logarithm of 32, to the base 2, &c., &c.

37. A System of Logarithms is a collection of the logarithms of a series of numbers corresponding to the same base.

Any number whatever may be taken as the base of the system; but it is obvious that some numbers are much more convenient than others.

38. Two system of logarithms have been constructed and tables calculated with great care. They are,-Ist. The Common System or Briggean System, whose base is 10.

2nd. Napierian System, whose base is 2.71828.

The Narierian System was invented by Baron Napier, and the peculiar base, 2°713. was adopted chiefly because the logarithms having that base are more simply expressed and more easily calculated than any other. It has hence been called the Natural System of Logarithms. These logarithms were also formerly called Hyperbolic logarithms, from certain relations found to exist between them and the asymptotic spaces of the hyperbola, and which were erroneously believed to be peculiar to them.

The Common System was shortly afterwards invented by Briggs and adopted by Baron Napier, and is the system now universally employed for the purposes of calculation.

39. The Characteristic of a logarithm is the part which stands to the left of the decimal point.

40. The Mantissa (handful) is that part of the logar-

ithm which stands to the right of the decimal point.

41. Since 10 is the base of the common system of logarithms and at the same time the radix of our system of notation, we have-

= 105;	whence	log.	100000	= . 5
= 104;	whence	log.	10000	= 4
$= 10^3$ :	whence	log.	1000	= 8
$=10^{2}$ ;	whence	log.	-100	=\2
$= 10^{1}$ ;	whence -	log.	10	= 1
= 10°;	whence	log.	164911	= 0
$= 10^{-1};$	whence	log.	111	= -1.
= 10-9;	whence	log.	*01	= -2
= 10-3	whence	log.	*001	= -8
== 10-4;	whence	log.	: 0001	=-41
	= 10 <sup>4</sup> ; = 10 <sup>3</sup> ; = 10 <sup>2</sup> ; = 10 <sup>1</sup> ; = 10 <sup>-1</sup> ; = 10 <sup>-2</sup> ; = 10 <sup>-3</sup> ;	= 104; whence = 103; whence = 101; whence = 100; whence = 10-1; whence = 10-3; whence = 10-3; whence	= 104; whence log. = 103; whence log. = 101; whence log. = 100; whence log. = 100; whence log. = 100; whence log. = 100; whence log. = 1000; whence log.	$ \begin{array}{c} = 10^4 \  \   \text{whence log.}  10000 \\ = 10^3 \  \   \text{whence log.}  1000 \\ = 10^2 \  \   \text{whence log.}  100 \\ = 10^1 \  \   \text{whence log.}  10 \\ = 10^1 \  \   \text{whence log.}  1 \\ = 10^{-1} \  \   \text{whence log.}  1 \\ = 10^{-3} \  \   \text{whence log.}  01 \\ = 10^{-3} \  \   \text{whence log.}  001 \\ \end{array} $

and 10 mal; so than 1 logarith

He its to rically

Thus. of 3.004 43. between equal to and '01 be -2 p

is neg 0s whisignifi

of the lo of '00027

> Nor mantis indicat charact

> > Wha numbe 1. 723

2. .02

3. 1:1

44. of the vidend the nu

317

ISECT. X.

ex of the n root or

the fixed em belong

base 10. se 2, &c., &c. n of the ng to the

m; but it is thers. nstructed

m, whose

the peculiar ng that base yother. It e logarithms in relations e hyperbola,

Briggs and mployed for

art which

the logarnt.

ystem of system of 42. From this it appears that the logarithm of and 10 will be more than 0 and less than 1; i. e., will a fractio a decimal; so also the logarithm of any number between 10 and 100 will be greater than 1 and less than 2; i. e., will be 1 and a fraction, or a decimal; so also the logarithm of any number between 100 and 1000 will be 2 and a decimal, &c.

Hence, the characteristic of any number containing digits to the left of the decimal point is positive and numerically one less than the number of such digits.

Thus, the characteristic of 7842 is 3; of 978 26 it is 2; of 813426789 it is 8; of 3 00429 it is 0; of 26789 426789 it is 4, &c.

43. It also appears, from Art. 41, that the logarithm of every number between 1 and 1 will be less than 0 and greater than—1; that is, it will be equal to—1, plus some decimal; the logarithm of every number between 1 and 01 will be less than—1 and greater than—2; or, in other words, will be—2 plus some decimal; so also the logarithm of every number between 101 and 1001 will be—3 plus some decimal, &c., &c.

Hence, the characteristic of the logarithm of a decimal is negative and numerically one greater than the number of Os which come between the decimal point and the first significant figure.

Thus, the characteristic of the logarithm of '000001 is 6; the characteristic of the logarithm of '00000000002347 is 11; the characteristic of the logarithm of 000278926345 is 4. &c., &c.

Norn.—The negative sign affects only the characteristic—the mantissa or decimal portion of a logarithm is always positive. To indicate this it is customary to write the negative sign over the characteristic, as in the above examples, and not before it.

## EXERCISE 146.

What are the characteristics of the logarithms of the following numbers:

1. 723, 9126.4, 81234.567, 912678.96124567, 23.912342.

Ans. 2, 3, 4, 5, and 1.

2. .027, .002134, .000000698, .8126714, .0000000002134.

Ans. 2, 3, 7, 1, and 10.

3. 1.1111111; 111111.11, 1000000000, .000000002162, 7, 12.78,

Ans. 0, 5, 9, 9, 0, and 1. 44. Since (Art. 11), to divide one power of a number by another power of the same we subtract the index of the divisor from the index of the dividend, and since common logarithms are indices to the base 10, let us take the number 47280 and successively dividing it by 10, examine the results.

Logarithms. = 4'674677 Numbers. 47280 ..... = 3.674677 4728 4728 = 2.674677 4728 = 1.674677 4.728 = 0.674677 4728 ..... = 1.674677 \*004728 . ..... = 3.674677 A ... ... Here we have simply performed the mane operation by two different methods, let. dividing the wambers by 10, and and, from the logarithms corresponding to the numbers, subtracting 1, the logarithm of 10.

From this illustration it is evident that.—

1st. The characteristic of the logarithm of a number is dependent wholly upon the position of the decimal point in that number, and is not at all affected by the sequence of the digits that compose that number; and

2nd. The Mantissa or decimal part of the logarithm of a number is dependent wholly upon the sequence of the digits that compose that number, and is not at all affected

by the position of the decimal point.

Nors.—It is only common logarithms (i. e., those having 10 for their base) that possess the important property of having the same mantises for the same figure, whether integral or decimal, or both, and it was this property that induced Briggs to adopt that base in preference to the Napierian base 1971388.

base, 371838.

'45. Since the characteristic of the logarithm of any number does not depend upon the value of the digits composing that number, and is so easily found by attention to the rules found in Arts. 42, 43, it is customary to omit it altogather in logarithmic hables, and merely give the mantissa. The annexed tables contain the logarithms of all numbers from I to 1000 calculated to 6 decimal places. When greater accuracy is required, tables calculated to a greater number of places are used. By means of the proportional parts and difference given in the tables, the logarithm corresponding to all numbers whatever, may be found with sufficient accuracy for all intentical numbers. for all practical purposes.

46. To find the logarithm of any number not greater than 100:-

Find on the first page of the table of logarithms, the given number in the column marked No., and directly opposite to it,—in the column marked log., will be found the logarithm.

EXAMPLE 1.—What is the logarithm of 47? Ans. 1.672098.

NOTE.—By maying that 1.672098 is the logarithm of 47, we simply mean that the base 10, raised to the power 1.672098, is equal to 47, or briefly 101.672098 = 47.

EXAMPLE 2.-What is the logarithm of 93? Ans. 1'968483.

47. To find the logarithm of any number consisting of not more than four digits:-

Find, in the column marked N, the first three digits of the given

Then the mantiese will be found in the intersection of the horizontal line containing these three digits and the vertical column at the head of which stands the fourth digit.

To this mantissa attach the characteristic as found by the rules

in Arts. 6, 42, 48.

Ex Loo on p page this h which requir and to Ex

horiso colum (sinco figure

1. W

2: W 3. W

4. W

than

Fn rithm tissa: the o numb in th cut o A

corre tissa L

E:

Th and t o different logarithms

## EXAMPLE 1.—What is the logarithm of 7983?

Looking in the column marked N, we find the first three digits 798, on page 393 in the fourth horisontal division, counting from the top of the page and in the last line but one of that division. Carrying the eye along this horisontal line till we come to the vertical column, at the head of which stands the remaining digit, 3, we obtain for the mantissa of the required logarithm '903166, to which we prefix the characteristic 3 (since there are four digits to the left of the decimal point in the given number), and thus obtain the required logarithm 3'903166.

EXAMPLE 2.—What is the logarithm of '0000001234?

The first three digits, vis: 123, are found in the fourth line of the third horizontal division on page 382, and at the intersection of this line with the column headed 4, is found '091315. To this we attach the characteristic 7. (since there are six 0s, between the decimal point and the first significant figure) and thus obtain the required logarithm, 7:091315.

### EXERCISE 147.

- 1. What are the logarithms of 5794, 57.94, 5794000, and .0005794? Ans. 3.762978, 1.762978, 6.762978, and 4.762978.
- 2: What are the logarithms of 1.169, 11690, and 1000000?
- Ans. 0.067815, 4.067815, and 3.067815. 3. What are the logs. of .734, 7340000000, and .00000000734?
- Ans. 1.865694, 9.865696, and 9.865696. 4. What are the logarithms of 978.4, 9.784, 978400, and .9784? Ans. 2.990516, 0.990516, 5.990516, and 1.990516.
- 48. To find the logarithm of a number containing more than four digits:-

FIRST METHOD.—Find the mantissa corresponding to the logarithm of the first four digits by the last rule. Subtract this mantissa from the next following mantissa in the tables. Multiply the difference thus obtained by the remaining digits of the given number, and cut off from the product as many digits as there were in the multiplier (but at the same time adding unity if the highest cut off be not less than 5).

Add the number thus obtained to the mantissa of the logarithm corresponding to the first four digits, and the result will be the mantissa of the given number.

Lastly, attach the characteristic to this mantissa.

EXAMPLE 1.—What is the logarithm of 58803.2?

#### OPBRATION.

The mantissa of the logarithm of 5880 (the first four digits) is '780782 and the next following mantissa is '730863.

Then from '730863 Subtract '730782

Difference 81; and \$1×32 (remaining digits of given number)

point in uence of rithm of

e of the

affeoted

umber is

0 for their antissa for a this pro-Napierian

or dose not , and is so customary mantissa 1 to 10000 red, tables of the pro--Barroo one t accuracy

greater

ven num-,—in the

672098. iply mean or briefly

'968483.

nsisting

he given

the horilumn at

the rules

20011

it d

rigi cor and rigi

sevi rigi

be 1

dig the

of t

The

FINI

= 2502, from which we cut off two digits, since we multiplied by a number of two digits, and since the highest digit cut off is not less than 5, we add unity to the part retained, which gives us 26.

Then mantissa of logarithm of first four digits '730782

Mantissa of logarithm of given number '730808

To which attach the characteristic 4 and required logarithm = 4730808.

Note.—Except at the beginning of the tables, where the mantissas increase rapidly in magnitude, the difference may be taken from the right hand column, (headed D) and opposite the first three digits of the given number, where the mean difference of the mantissas in that line will be

## EXAMPLE 2.—What is the logarithm of 832.17242?

#### OPERATION.

920176 38

To which we attach the characteristic 2 and required logarithm = 2 920214 49. The difference given in the column headed D in the tables, is that due to an increment of one unit in the fourth figure of natural number, thus 

Difference of natural numbers =1; difference of logarithms = 75

And since it is shown in common works on Algebra that, with small increments in the natural numbers the logarithms corresponding to them increase in arithmetical progression, in order to find the logarithm of any number between those given above, we consider that the increment of the logarithm to be added to 3'758'61, bears the same proportion to 75 (the increment for 1), that the increment of the natural number does to 1.

For example,—Let it be required to find the logarithm of 5738'47.

Here the increment of the given number below 15 and form the present of the given number leaves.

Here the increment of the given number being 47, we form the proportion 1: 47::75: 47×75 == 35°25, the increment to be added to 3 °758761, and this addition having been made, we get 3 °758796 for the logarithm of 5738°47. Similarly, if the increment of the natural number had been '047 or '0047, the corresponding increment of the log. would have been 3 '525 or '3525. These illustrations sufficiently explain the reasons of the last rule.

50. Taking the same number as in the last article and dividing the difference 75 by 10, we obtain 75 the difference corresponding to an increase of one unit in the fifth place of the natural number; the double of this, or 15 for two units, the treble or 22.5 for the three units, and so on; and each of the numbers thus obtained will be the increment of the logarithm corresponding to an increase of that number of units in the fifth place of the natural number. The increments thus obtained, and corresponding to each of the nine digits, are inserted in the left hand column of the tables, headed P. P. (Proportional Parts.)

51. The numbers in the column headed P.P., as arready explained, are the increments in the logarithm for an increase in the fifth place of the natural numbers. They express also the increments for the digits in the sixth, seventh, eighth, ninth, &c., places of the natural number, when they are divided by 10, 100, 1000, &c., as the case may be.

52. Hence to find the logarithm of any number containing more than four digits:-

Water the commence of the contraction of the contra

# a number

SECT. X.

=4.730808.
mantissas
the right
the given
ine will be

.... 920176

.. 38

= 2 920214 bles, is that mber, thus 58761 58836

= 75
with small
ing to them
thm of any

ment of the n to 75 (the es to 1.

proportion 61, and this 65738'47. 047 or '0047, or '3525.

rule.
g the differincrease of
f this, or 15
and each of
thm corresplace of the
ling to each
the tables.

plained, are place of the ligits in the , when they

nber con-

#### RULE.

SECOND METHOD.—Find the mantiesa of the logarithm corres-

ponding to the first four digits of the given number.

Find in the same horizontal division as that in which the mantissa is found, the proportional part in the column headed P. P., corresponding to the digit in the fifth place of the given number, and set it down beneath the part of the mantissa already found, so that their right hand digits may be in the same vertical line. Find the P. P. corresponding to the digit in the sixth place of the given number, and set it down so that its right hand figure may be one place to the right of the last. Find the P. P. corresponding to the digit in the seventh place of the given number and set it down one place to the right of the last, and so on till all the digits of the given number be used.

Add the part of the mantisea already found, and the P. Ps. as written, together, and reject from the result all but the first six digits to the left, adding one to the last retained, if the highest of the rejected digits be not less than 5—the result will be the mantisea of the logarithm of given number.

Lastly, attach the proper characteristic to this mantisea, and the result will be the required logarithm.

EXAMPLE 1.—What is the logarithm of 8372.468?

#### OPERATION.

Manti	ssa of logarithm of	8372 =	922829
P. P.	corresponding to	'4 =	21
P. P.	to to	'06 =	31
P. P.	" anto	'008 ==	77.42

Sum'= '922853| 52

Therfore required mantissa = 922854 and required log. = 3 922854. EXAMPLE 2.—What is the logarithm of 403567?

#### OPERATION.

Mantis	sa of le	ogai	ithm (	of 405	3500=	605844
P. P. co	rresp	ond	ing to	3350	, 60 ==	64
P.P.	13.4	125	nio to	39	77=	, P 75

Sum = .6059155

Therefore required logarithm is 5.605916.

## EXERCISE 148.

FIND THE LOGARITHMS OF THE FOLLOWING NUMBERS BY THE FIRST

- 1. What are the logarithms corresponding to 8193217, 73.9245, and .843742?

  Ans. 6.913455, 1.868789, and 1.926210.
- 2. Find the logarithms corresponding to .000234564 and .001007013. Ans. 4:370261 and 3:003035.

nu

the

rec

lo

Hi

Hi Hi

Th

ist

#### USING THE TABULAR DIFFERENCES.

3. Find the logarithms corresponding to 52.376 and 129.476.

Ans. 1.719133 and 2.112189.

#### USING THE PROPORTIONAL PARTS.

- 4. Find the logarithms corresponding to 000471398 and 9136712.
  - Ans. 4.673387 and 6.960790.
- Find the logarithms corresponding to 4.23429 and 763.12987.
   Ans. 0.626780 and 2.882598.
  - 53. To find the logarithm of a vulgar fraction:

Subtract the logarithm of the denominator from the logarithm of the numerator.

54. To find the logarithm of a mixed number :-

#### RIII.E.

Either reduce the mixed number to a fraction and proceed as in Art. 53, or reduce the fractional part to a decimal, attach it to the whole number and proceed as in Arts. 48-52.

55. To find the natural number corresponding to any given logarithm:—

#### RULE

FIRST METHOD.—Find that logarithm in the table which is next lower than the given one, and the four digits corresponding to it will be the first four digits of the required number.

II. Subtract this logarithm from the given logarithm, to the remainder annex one cipher and divide by the tabular difference corresponding to the four digits already obtained, the quotient will be the fifth digit.

III. To the remainder attach another cipher and again divide by the tabular difference, the quotient will be the sixth digit, and thus proceed till a sufficient number of digits has been obtained.

IV. The characteristic of the logarithm shows where to place the decimal point.

Note.—The number cannot be carried with accuracy to more places than the logarithm has decimal places. (See Art. 56)

Example 1.—Find the number corresponding to the logarithm 4.923267.

#### OPERATION.

Given log. 923267 Next lower in tables, 923244 = log. of 8380.

Difference = 23 Tabular difference = 52.

Then 23000 ÷ 53 gives 442 for digits in 5th, 6th, and 7th places.

129·476. 2·112189.

398 and

6·960790. 63·12987. 2·882598.

logarithm.

oceed as in ch it to the

ng to any

hich is next onding to it

thm, to the r difference uotient will

gain divide digit, and tained. to place the

more places

logarithm

Hence the digits of the natural number are 8380442; and since the characteristic is 4, 4. e. one less than the number of digits to the left of the decimal point, the required number is 83804'42.

SECOND METHOD.—Find the first four digits of the required number and also the difference between the given logarithm and the next lower in the table as in the last rule.

II. Find in the same horizontal division of the table the highest P. P. that does not exceed this difference. Opposite to it in the column headed N. will be found the digit of the fifth place.

III. Subtract this P. P. from the difference, to the remainder annex one cipher and find the highest P. P. not exceeding the number thus formed. Opposite to it in column N. will be found the sixth digit.

IV. Continue this process by the addition of ciphers till the required number of digits be found.

EXAMPLE 2.—Find the natural number corresponding to the logarithm 3.553259.

#### OPBRATION.

Given log. '553259

Next lower in table '553155 = log. of 3574

Highest P. P. not greater than 104 = 98 corresponds to 8 for Afth

Highest P. P. not greater than 60 = 49 corresponds to 4 in sixth.

Highest P. P. not greater than 110 = 110 corresponds to 9 in seventh

Therefore digits of required number are 3574849; and since the characteristic is 3, there must be four digits to the left of the decimal point.

Hence required number is 3574849.

### Exercise 149.

## BY FIRST METHOD.

1. Find the natural numbers corresponding to the logarithms

Ans. 13713.227, 5.225578 and .0004319376.

2. Of what numbers are 2.921686 and '922165 the logarithms?

Ans. 835 and .8359211.

## BY SECOND METHOD.

- 3. Of what numbers are 5.407968, 7.408386 and 3.416369 the logarithms?

  Ans. 255839.4, 25608588 and .0026083.
- 4. What are the natural numbers corresponding to the logarithms 4.877777 and 0.555555?

Ans. 75470.5168 and 3.5938.

A.G. In order to ascertain how many figures of these results may be relied upon as correct, let us take from the tables any logarithm, as 4.235635.

Now the real value of this logarithm if carried to a greater number of places might be anything between 4.2566355 and 4.2566355, and might therefore differ from the given logarithm by very hearly 0000005, which is therefore the extreme limit of the error attached to tables of six places; i.e. any difference less than 0000005 might occur without producing any change in the logarithm as given in the table.

Now it is demonstrated in within the acting of the theory of logarithms.

the logarithm as given in the table.

Now it is demonstrated in works treating of the theory of logarithms that the difference between the logarithms of numbers, which differ only by unity, is less than the modulus of the system divided by the smaller number. The modulus of the common system of logarithms is 4342945, and if we let a represent the smaller number, the difference between the logarithms of a and of a 1 is less than 4342945.

Now we have shown that the difference between the true logarithm and that given in the table to six places, may be nearly equal to 0000005, which

If tables of seven or eight places are used, the result can be depended on to seven or eight places, if the number he less than 868889 or if the mantissa be less than 9378; but if greater, then the result can be relied on only to one less number of figures than the decimals of the logarithm.

## LOGARITHMIC ARITHMETIC.

57. The Arithmetical Complement of a logarithm is the remainder obtained by subtracting the logarithm from 10.

Thus the arithmetical complement of 2.713426 is 10-2.713426 = 7.286574.

## Exercise 150.

- 1. Find the arithmetical complements of 5:631642 and 0:714000. Ans. 4.368358 and 9.286000.
- 2. Find the arithmetical complements of 3.123456 and 7.213149. Ans. 12.876544 and 16.786851.
- 3. Find the arithmetical complements of 6-124357 and 2-000887. 12 16 20 3 h.dns. 3.875643 and 11.999163.
- 58. To multiply two or more numbers together by means of logarithms:-

### The RULE. 1st

I. Add their logarithms and the sum will be the logarithm of their product.

II. Find the natural number corresponding to this logarithm.

NOTE 1.—For reason see Art. 10.

not the state of the state of the

North & The following exercises are all worked by the difference, and not by the proportional parts:

umber of is therechange in

ogarithms differ only he smaller tween the

rithm and 005, which ut 0000005 è less than digite, but e table will

pended on f the manied on only

m is the from 10.

== 7°286574.

0:714000. 286000.

7.213149. 6.786851. 2-000887.

1.999163.

ether by

garithm of

rithm.

ference, and

EXAMPLE. - Multiply 5631 by 47.

Logarithm of 5631 = 8 750586 47 = 1 672098

5 574296847 Tall to south J' all will I'm

5-422590 = logarithm of 264500

Ans. 204657

## 

1. Multiply 61, 22, and 65 together. Ans. 87230.
2. Multiply 52, 734, and 6 together. Ans. 229008.

3. Multiply together 35.86, 2.1046, .8372 and .00294.

Ans. . 185761.

4. Multiply -00008764 by 86359.

ARTS, 55-60.1

Ans. 0000 75685.

59. To divide numbers by means of their logarithms:-

I. Subtract the logarithm of the divisor from the logarithm of the dividend: the result will be the logarithm of the required quotient.

II. Find the natural number corresponding to this.

NOTE.—For reason see Art. 11.

Example 1.—Divide 6732:7 by 478.

OPERATION.

Logarithm of 6782-7 = 3:828180 Logarithm of 478' = 2.679428

Difference = 1'148761

1'148603 = logarithm of 14'0800

Ans. 14.0851

EXAMPLE 2.—Divide .036584 by .00078593.

OPERATION.

Logarithm of 00078598 = Tsi95884

Difference = 1.667907 1'867826 = logarithm of 48'8400 : San Diz Pool to

81 =

Ans. 40'5487

Bli. Instead of subtracting the logarithm of the divisor, we may all its arithmetical complement—the result, with 10 subtracted from the characteristic, will be the logarithm of the quotient.

Thus, in the last example the arithmetical complement of 4.895384 is 13'104616, and this added to 2'563291 gives 11'667907, and subtracting 10 from this characteristic, gives us 1'667907, the same as obtained by the other method.

NOTE.—This method of using the arithmetical complement is very convenient when we have to divide one number by the product of several

#### EXERCISE 152.

1	Dimid	0.794	. h	0009278.	
1.	DIVIG	3 6 7 34	DA .	UUU9278.	

Ans. 725.8033.

log

in

neg

who logi

N

and

3.88

Dei

fac

to

T

11

any 21

logi can

log

120

2. Divide 437.89 by 62.735.

Ans. 6.98.

3. Divide 93.217 by .0007132.

Ans. 130702.4.

4. Divide 983526' by the product of 23, 189 and 2.748.

Ans. 823.339.

## 61. To raise a quantity to any power by means of logarithms:-

ે સાતા લોકેલ કોલો, એક સા**લાકા.** કેલ્કસ અને કે કેલ . છે

I. Multiply the logarithm of the given number by the index of the required power, the result will be the logarithm of the required power.

II. Find the natural number corresponding to this logarithm.

NOTE.—For reason see Art. 12.

EXAMPLE 1.—Find the 10th power of 2.

## OPERATION.

Logarithm of 2 = 0.301030.

 $0.301030 \times 10 = 3.010300 = logarithm of 1024$ . Ans.

EXAMPLE 2.—Find the 7th power of 2.71.

#### OPERATION.

Logarithm of 2.71 = 0.432969.

Then  $0.432969 \times 7 = 3.030783 = logarithm of 1073.45$ . Ans.

NOTE.—In order to obtain the correct result when the characteristic happens to be negative, it must be recollected that the mantissa is always positive.

## EXERCISE 153.

1. What is the 5th power of 5?

Ans. 3125.

- and is a true town this is it

2. What is the 6th power of 1.073?

Ans. 1.5261.

3. What is the 4th power of :0279?

Ans. :00000060592.

4. What is the 11th power of 1.111? Ans. 3.1831.

## 62. To extract any root of a given number by means of logarithms:-

I. Find the logarithm of the given number and divide it by the index of the required root, the result will be the logarithm of the root. the method older of the most than of the order than of the say

5384 is

10 from other

ry conseveral

5.8033.

702.4. 3.339.

ins of

dex of equired

cteristic always

060592.

means

t by the

II. Find the natural number corresponding to this logarithm.

Note.—For reason see Art. 15.

EXAMPLE.—What is the cube root of 12345?

OPERATION.

Logarithm of 12345 = 4.091491. Then 4.091491 ÷ 3 = 1.363830 = logarithm of 23.11159. Ans.

63. To extract any root when the characteristic of the logarithm of the given number is negative:-

I. If the characteristic is exactly divisible by the divisor, divide in the ordinary way, but make the characteristic of the quotient negative.

II. If the negative characteristic is not exactly divisible add what will make it so, both to it and to the decimal part of the

logarithm. Then proceed with the division. EXAMPLE 22.—Extract the fourth root of .0076542.

## OPERATION.

Logarithm of '0076542 = 3.883899.

Now since 3 is not exactly divisible by 4 we add—1 to the characteristic and +1 to the mantissa which gives us 4+1.833899 and this is evidently = 3.883800.

Thin  $\frac{7}{4} + 1.883899 \div 4 = \overline{1.4709747} = \text{logarithm of .295784.}$  Ans. EXERCISE 154.

1. Extract the 7th root of 913426000.

Ans. 19.0588.

2. Extract the 11th root of 1.61342.

Ans. 1.04444.

3 Extract the 5th root of .000007139. Extract the 7th root of .002147.

Ans. .0934817. Ans. .41575.

64. When the logarithms of two or more prime numpers are given, the logarithm of any multiples of these factors by each other can be easily obtained by attention to the foregoing rules.

Thus if the logarithm of 2 and 3 be given:-

1st. We can obtain the logarithm of any power of 2 or 3 by Art. 61, and

any root of 2 or 3 by Art. 62.

2nd. We know the logarithm of 10 to be 1, and hence we can obtain the logarithm of 5 since  $10 \div 2 = 5$  and also of 3.3 since  $10 \div 3 = 3.3$ , hence we can also obtain the logarithm of any power or root of 5 or 3.8.

3rd. By Arts. 58, 59, we can obtain the logarithm of any power or root of

2, 3, 5 and 3'3 multiplied by any power or root of 2, 3, 5 or 3'3.

Example 27.-Given the logarithm of 2 = 0.301030 and the logarithm of 3 = 0.477121. Find the logarithms of 500, 24, 54, 120, 75000, 163, 1, and 13.5.

```
328
                                     LOGARITHMIC ARITHMETIC.
                                                                                                                 [SECT. X.
                                                      OPERATION.
Since 5 = 10 \div 2 the logarithm of 5 = \log. 10 - \log. 2 = 1 - 0.301030 = 0.808970. Then logarithm of 500 = 2.698970. 24 = 8 \times 3 = 2^3 \times 3 \cdot \cdot \log \cdot 24 = (\log \cdot 2) \times 3 + (\log \cdot 3.)
                                                              \log 2 = 0.301030 \times 3 = 0.903090
                                                              log. 3=
                                                                                                       477121
                                                                                        Sum == 1.380211=log.24
                                                          Bouly been
   54 = 27 \times 2 = 3^3 \times 2. \log. 54 = (\log. 3) \times 3 + (\log. 2.)
\log. 3 = 0.477121 \times 3 = 1.431363
\log. 2 = 0.301030
\begin{array}{c} \text{Sum} = 1.782398 = \log. 54. \\ 120 = 4 \times 3 \times 10 = 2^2 \times 3 \times 10 \cdot \cdot \cdot \log. 120 = (\log. 2) \times 2 + (\log. 3) + (\log. 10.) \\ \log. 2 = 0.301030 \times 2 = 0.602090 \\ \log. 3 = 0.477121 \end{array}
                                                                                       Sum =1:782398 = log. 54.
                                                        log. 10=
                                                                                    Sum = 2.079181 = log. 120.
75000 = 25 \times 3 \times 1000 = 5^2 \times 3 \times 1000 \cdot \log_{10} 75000 = (\log_{10} 5) \times 2 + (\log_{10} 3)
           + (log. 1000.)
                                                       5 = 0.698970 \times 2 = 1.397940
                                           log.
                                           log. 1000 =
                                                                             Sum = 4.875061 = log. 75000.
```

163 = 3·3 × 5 ·· logarithm of 163 = (log. 3·3) + (log. 5.)
Since 10÷3=3·3, log. 3·3=log. 10-log. 3=1-0·477121=0·522879
logarithm 5= 0·698970

Sum =1.221849=log. 162.

SR

19:

of ti

9.

11. 12. 13. 14. 15.

16.

18.

19. 20.

21.

 $\frac{1}{2} = 5$  .. by changing only the characteristic =  $\overline{1} \cdot 698970 = \log \operatorname{rithm} 1.$   $13 \cdot 5 = 5 \times 27 = 5 \times 3^3$  .. logarithm  $13 \cdot 5 = (\log_2 3) \times 5 + (\log_2 5)$   $\log \operatorname{rithm} 3 = 0.477121 \times 3 = 1.431863$ 

" . I Washington

logarithm 5 =

1.698970

Sum =1.130333=log. 13.5

would be creek at Exercise 155.

1. Given logarithm 2 = 0.301030 and log. 7 = 0.845098, find the

logarithms of 14000, 4.9, .00196, 1750, 1428.571428, .00000112 and 3.0625.

Ans Log. 14000 = 4-146128. Log. 4-9 = 0-690196.

 $\begin{array}{c} \text{Log. } \cdot 00196 = \overline{3} \cdot 292266. \\ \text{Log. } 1750 = 3 \cdot 243038. \end{array}$ 

Log.  $1428 \cdot 571428 \Rightarrow 3 \cdot 154902$ . Log.  $00000112 \Rightarrow 6 \cdot 049218$ . Log.  $3 \cdot 0626 \Rightarrow 0 \cdot 486076$ .

Norm.—1428-571428 = + × 10000, also 5'0825 = 40 + 16.

SECT. X.1

EXAMPLE 2.—Given logarithm 1 = 1.698970 logarithm 3 = 0.477121 logarithm 11= 1.041393

Find the logarithms of 491, 363, 4.09, 2.4, 392.72, 2933331 and 19.965.

491 = 1.694605. Ans. Logarithm of 363 = 2.559907.Logarithm of Logarithm of 4.09 = 0.611819. Logarithm of 2.4 = 0.388181.Logarithm of 392.72 = 2.594090. Logarithm of 2933331 = 5.467362. Logarithm of 19.965 = 1.300270.

## QUESTIONS TO BE ANSWERED BY THE PUPIL.

NOTE. The numbers after the questions refer to the numbered articles of the section.

1. What is the power of a number? (1)

2. What is a root of a number? (2)
3. Why is the second power of a number called its square? (4)
4. Why is the third power of a number called its cube? (5)

5. What is the index or exponent of a power? (6)
6. What is involution? (8)
7. How do we multiply two or more different powers of the same number

together? (10)
3. How do we divide any power of a number by another power of the same number? (11)
9. How do we find any required power of a given power? (12)
10. What is evolution? (13)
11. By what methods do we indicate a root of a number? (14)

11. By what methods do we indicate a root of a number? (12)

12. How do we extract any root of a given power of a number? (15)

13. What is meant by extracting the square root of a number? (16)

14. What is the first step in extracting the square root of a number? (16)

15. Why do we point off into periods of two figures each? (18-1)

16. What is the second step in the process of extracting the square root?

17. How do we know that the square root of the highest square in the left hand period is the highest digit of the root? (18-II)
18. What is the third step in the process of extracting the square root?

19. Why do we bring down only the next period to the right? (18-II in

20. What is the fourth part of the process for extracting the square root?

21. Why do we double the part of the root already found for a trial divisor? (18-III).
 22. What is the next step in extracting the square root of a number? (16)
 23. Why do we not include the right hand figure of the dividend when seeking how many times the trial divisor is contained in it? (18-IV.)
 24. Why do we piace the digit thus found in both the divisor and the root? (18-V)
 25. What are the other steps used in extracting the square root? (16)
 26. How do we extract the square root of a decimal? (19)

26. How do we extract the square root of a decimal ? (19)

0.698970.

=log.24

= log. 54.

(log. 10.)

log. 120.  $+(\log 3)$ 

g. 75000.

log. 16%.

thm .

=log. 13'5

find the

571428,

146128. 690196.

292286. 243038

154902.

.049218.

486076.

27. How do we extract the square root of a fraction or mixed number? (20)
28. What is a triangle? (22) What is a right-angled triangle? (23)
29. How may any one side of a right-angled triangle be found when the other two are given? (24)
30. What proportion exists between different circles? (25)
31. How may the area of a circle be found when its diameter is known?

33. What is meant by extracting the cube root of a number? (26)
33. Give the different steps of the process of extracting the cube root. (26)
34. If a number consist of a certain number of tens, plus a certain number

34. If a number consist of a certain number of tens, plus a certain number of units, of what does its cube consist? (??)

35. Why do we divide off into periods of three figures each? (28, I.)

36. How do we know that the cube root of the highest cube contained in the left hand period is the highest digit of the root? (28, II.)

37. Whence do we obtain, in the cube root, the constant multipliers 300 and 30. Illustrate by an example. (28 IV, and VI.)

38. Why do we make the two additions, indicated in the rule, to the trial divisor? (28, VI.)

39. How do we extract the cube root of a decimal? (29)

40. How do we extract the cube root of a fraction or mixed number? (30)

40. How do we extract the cube root of a fraction or mixed number? (30)

41. In extracting the cube root of a number in any other scale, what changes must we make in the rule? (31)

42. Give the different steps of Horner's method of extracting the cube root. (32)

43. What proportion exists between the magnitude of similar solids? (83)

43. What proportion exists between the magnitude of similar solids? (33)
44. How do we extract the higher roots when the index is a power of 2 or 8 or a multiple of 2 by 3? (34)
45. What is a logarithm? (35)
46. What is the base of a system of logarithms? (36)
47. What is a system of logarithms? (37)
48. What systems of logarithms have been constructed and how do they differ from the another? (38)
49. What is the characteristic of a logarithm? (39)
50. What is the decimal part of the logarithm called? (40)

50. What is the characteristic of a logarithm called? (40)
51. How do we find the characteristic of a logarithm? (42 and 43)
52. Why is the negative sign written over the characteristic of the logarithm of a decimal? (43, Note.)
53. Show that the characteristic of the logarithm of a number depends only on the position of the decimal point in the number, and the manting poly in the coverne (44).

tissa only in the sequence of figures. (44)

54. Explain clearly what is meant by the numbers in column D of the tables. (49)

55. Explain how the proportional parts in column P. P. are obtained. (50)

56. Explain how the numbers in the column headed P. P. become the incre-

ments to be added to the logarithms for an increase in the sixth, seventh, eighth, &c., place in the natural number. (51)

57. How do we find the logarithm of a vulgar fraction? (53)

58. Explain to how many figures we may rely upon the accuracy of the results obtained by logarithmic tables. (56)

59. What is the arithmetical complement of a logarithm? (57)

60. How do we multiply numbers by means of their logarithms? (58)

61. How do we divide numbers by means of their logarithms? (59, 60)

62. How do we involve and evolve quantities by means of logarithms?

(61, 62, 63)

A fine a new construction of a second

(61, 62, 63)

tentel in the " March in the tente

PRO

when t Thus, ference l

12, 10, being 2.

called

be con

These being gi Cases ar 4. If

we shal preted, 5. Th

where t Thus in number less tha Hence equal to

one less 6. Si in any o

Also Hence But (a-

## r? (20)

CT. X.

hen the

mown?

ot. (26) number

ained in iers 300

he trial

er? (30) le, what he cube

is? (83) of 2 or 8

do they

he logaads only he man-

D of the

red. (50) ie increie sixth.

y of the

ñ8) rithms?

# SECTION XI.

## PROGRESSION, POSITION, COMPOUND INTEREST,! AND ANNUITIES.

## PROGRESSION.

1. Quantities are said to be in Arithmetical Progression when they increase or decrease by a common difference.

Thus, 2, 5, 8, 11, 14, &c., are in arithmetical progression, the common difference being 3.

12, 10, 8, 6, &c., are in arithmetical progression, the common difference being 2.

2. In every progression the first and the last terms are called the extremes, and the intermediate terms the means.

## ARITHMETICAL PROGRESSION.

- 3. In arithmetical progression there are five things to be considered:
  - The first term.
     The last term.
  - 3. The common difference.
  - The number of terms. 5. The sum of the series.

These quantities are so related to one another that any three of them being given the other two can be found, and hence there are 20 distinct cases arising from these combinations.
4. If we represent these five quantities by letters, thus:

a = the first term. l = the last term.

d = the common difference.
n = the number of terms.

s = the sum of the series.

we shall be able easily to deduce algebraic formulæ which, being interpreted, become the common arithmetical rules for arithmetical progression.

5. The general expression for an arithmetical series then becomes

a+(a+d)+(a+2d)+(a+3d)+(a+4d)+(a+5d)+, &c.

where the coefficient of d is always 1 less than the number of the terms. Thus in the third term the coefficient of d is 2, which is 1 less than the number of the term: in the fifth term the coefficient of d is 4, which is 1

less than the number of the term, &c. Hence l = a + (n-1) d; that is, the last term of an arithmetical series is equal to the first term added to the product of the common difference by one less than the number of terms.

6. Since the sum of the series is equal to the sum of all the terms taken in any order whatever, we have

S =  $a+|a+d+|a+2d+|a+3d+| \dots l-3d+|l-2d+|l-d+|l-d+|l-d+|l-2d+|l-3d+| \dots a+3d+|a+2d+|a+d+|a|$ Hence 2s =  $(a+l)+(a+l)+(a+l)+(a+l)+\dots$  to n terms. But (a+l)+(a+l)......to n terms = (a+l)n.

Therefore 2 = (a+i)n, and dividing these equals by 3, we have  $c = (a+i) \frac{\pi}{2}$ . That is, the sum of the strice is found by adding together the first and last terms and multipliying their sum by half the number of terms.

Norm.—In adding the corresponding terms of the foregoing series together the d's people out; thus adding the second terms of the right hand members together we have a+d+l-d, where the d's cancel, and the sum becomes a+l: so also in the third terms we have a+2d+l-2d = a+l, &c.

7. From the formula obtained in Art. 5, we find by transposing the terms

$$\begin{array}{l}
l = a + (n-1)d \\
a = l - (n-1)d \\
d = \frac{b-a}{n-1} \\
n = \frac{l-a}{d} + 1
\end{array}$$

and substituting these values of *l*, *a*, *d*, and *n* in the formula obtained in Art. 6, we find

$$s = \left\{ 2a + (n-1)d \right\} \frac{n}{s}$$

$$s = \left\{ 2l - (n-1)d \right\} \frac{n}{s}$$

$$(l-a)(l+a) + l+a$$

$$s = \frac{n}{s}$$

We thus obtain the five fundamental formulas from which the other afteen are derived by transposing the terms, &c. Thus

$$l = a + (n-1)d$$
 gives formulas for  $i, a, n, d = 4$ 

$$s = (a+l) \frac{n}{2} \qquad s, a, l, n = 4$$

$$s = \left\{2a + (n-1)d\right\} \frac{n}{2} \qquad s, a, n, d = 4$$

$$s = \left\{2l - (n-1)d\right\} \frac{n}{2} \qquad s, l, n, d = 4$$

$$s = \left(\frac{l+a}{2}\right)(\frac{l-a}{2}) + \frac{l+a}{2} \qquad s, a, l, d = 4$$

$$Total 20$$

And the state of t

ABTE,

ARIT

\_

No

I

.

·VI

.0

0

od

10

1

XI

X

X X

XVI XVI

ynd

or. XI.

ries tont hand the sum -1, &c. te terms

ined in

he other

11 1 11 11

(a) 1117

# S. THE EQLIOWING TABLE GIVES THE 20 FOR WILLS FOR ARITHMETICAL PROGRESSION WITH THEIR RELATIONS, &C.

No.	Given	Required.	Formulas.	Whenee derived
	0, d, n	ी र त	1=0+(x-1)d -11	fundamental.
	a, d, s		$l = \frac{1}{2a} + \sqrt{2ds + (a - 1d)}$ $l = \frac{2a}{a} - a$	Tall.
,	a, n, e		* - * ·	101
IA;	a, n, e		$l = \frac{10}{n} + \frac{(n-1)d}{2}$	VII.
V.	a, l, n		$s = (a+1) \frac{n}{3}$	fundamental.
VI.	a, d, n	• )	$a = \{2a + (n-1)d\} \frac{a}{3}$	V. and I.
VIL	4, 1, 4		्= (श्रिः (कारोस्) के	v. and XVII.
1	a, d, 1	130	- ( to) (but)   late	V. and XIII.
	<del>'</del>		26	Grand Charles
IX.	a, n, l		$d = \frac{r - a}{4^{-1}}$	I.
X.	a, 18, 4		$d = \frac{2s - 2an}{n(n-1)}$	<b></b>
XI.	9 1,0	d	$d = \frac{(l+a)(l-a)}{(l-a)}$	VIII.
1)			d = 2nl 30 a2 5	COLING TO
. o consi	2, 10, 0	1	1 0 (3(3-1)	Mills of the State
XIII.	a, d,		$n = \frac{l-a}{d} + 1$	.I. 1994
	a, d,		d-20+1/20 /20-00	11 / [7
1		75	(120 0	is ca <b>yl</b> a,gr
XV.	a, 1, e		1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	n verill
XVI	d, l,		n = 21+3+1 (21+4) 2 2	VII.
			1 20 7	12 m junt "
XYII	d. a.		a = (n-1)d	
XVIII	d, 2,	11111	8 (701)G	, , , VI.
XIX	. 1, n, a	4 -	$a = \frac{2s}{l} - l$	· : V:+ 3
XX	. d, l, s		$a = \frac{1}{2}d + \sqrt{(l+\frac{1}{2}d)^2 - 2de}$	VIII
	, , ,		(.7) (1	(- i) III 5

9. The following examples will enable the student to understand clearly the interpretation and application of these formula.

10. To find the last term of an arithmetical series when the first term, the common difference, and the number of terms are given:—

RULE.

$$l = a + (n-1)d$$
. (1.)

INTERPPETATION.—The last term of a series is found by adding the first term to the product of the common difference by 1 less than the number of terms.

EXAMPLE.—What is the tenth term of the arithmetical series 1, 3, 5, &c.?

Here we have given the first term 1, the common difference 2 and the number of terms 10; to find the tenth or last term.

Then  $l = a + (n-1)d = 1 + (10-1) \times 2 = 1 + 9 \times 2 = 1 + 18 = 19$ . Ans.

11. To find the common difference of an arithmetical series when the first term, the last term, and the number of terms are given:—

RULI

$$d = \frac{l-a}{n-1} \quad \text{(13.)}$$

INTERPRETATION.—To find the common difference of an arithmetical series,—Subtract the first term from the last term and divide the difference thus obtained by one less than the number of terms.

EXAMPLE.—The first term of an arithmetical series is 3, the

#### OPERATION.

Here we have given the first term 3, the last term 55, and the number of terms 13, to find the common difference.

Then 
$$d = \frac{l-a}{n-1} = \frac{55-3}{13-1} = \frac{52}{12} = 41 = Ans.$$

12. To find the sum of an arithmetical series when the first term, the last term, and the number of terms are given:—

RULE.

$$s = (a+l) \frac{n}{2}. \quad (v.)$$

INTERPRETATION.—Add the first and last terms together and multiply their sum by half the number of terms.

EXAMPLE.—Find the sum of an arithmetical series whose first term is 2, last term 50, and number of terms 17.

Her

The

the r

d:

termi series numb will

num

746 w

He

whe

twice different

rend

tern the the

the Wh when

OPERATION.

Here we have given the first term 2, the last term 50 and the number of terms 17 to find s, the sum of the series.

Then 
$$s = (a+1) \frac{n}{2} = (2+50) \times \frac{17}{2} = 52 \times \frac{17}{2} = 26 \times 17 = 442$$
. Ans.

13. To find the common difference when the last term, the number of terms, and the sum of the series are given:

$$d = \frac{2nl-2s}{n(n-1)}$$
. (XII.)

INTERPRETATION .- Take twice the product of the number of terms by the last term, and from it subtract twice the sum of the Divide the resulting difference by the product of the number of terms by 1 less than the number of terms and the quotient will be the common difference.

EXAMPLE.—In an arithmetical series the last term is 80, the number of terms 11 and the sum of the series 746, required the common difference.

OPERATION.

Here we have given l, n, and s to find d and since l = 80, n = 11 and s =746 we have:

$$d = \frac{2nl-2s}{n(n-1)} = \frac{(2 \times 11 \times 80) - (2 \times 746)}{11 \times (11-1)} = \frac{1760 - 1492}{11 \times 10} = \frac{268}{110} = \frac{234}{38}. \text{ Ans.}$$

14. To find the number of terms of an arithmetical series when the first term, the common difference, and the sum of the series are given :-

$$n = \frac{d-2a}{2d} + \sqrt{\frac{2s}{d} + \left(\frac{2a-d}{2d}\right)^2}$$
. (xiv.)

INTERPRETATION. - I. Subtract the common difference from twice the first term, divide the remainder by twice the common difference, square the quotient, add the result to the quotient obtained by dividing twice the sum of the series by the common difference and extract the square root of this sum.

II. Next, from the common difference subtract twice the first term, divide the remainder by twice the common difference, and to the quotient add the square root obtained in I. The sum will be the number of terms.

EXAMPLE—The first term of an arithmetical progression is 7, the common difference 1, and the sum of all the terms 142. What is the number of terms?

adding 1 less

BOT. XI

aber of

l series

and the

19. Ans. metical umber

n aritherm and mber of

s 3, the

umber of

ien the ms are

ter and

ose first

= 16. Ans.

OPERATION.

Here we have given a, d, and e, to find n and since a = 7,  $d = \frac{1}{4}$ , and

$$n = \frac{d-2a}{2d} + \sqrt{\frac{2a}{d}} + \left(\frac{2a-d}{2d}\right)^2 = \frac{\frac{1}{2}-3\times7}{9\times\frac{1}{4}} + \sqrt{\frac{142\times9}{2} + \left(\frac{2\times7-\frac{1}{2}}{9\times\frac{1}{4}}\right)^2} = \frac{\frac{1}{2}-14}{9\times\frac{1}{4}} + \sqrt{\frac{12}{2}+\frac{1}{2}} + \sqrt{\frac{12}{2}+\frac{1}{2}} = -27\frac{1}{2} + 45\frac{1}{2}.$$

EXERCISE 156.

1. In an arithmetical series the first term is 4, the number of terms 17 and the sum of the series 884. What is the last term?

2. The extremes of an arithmetical series are 21, and 497, and the number of terms is 41. What is the common difference?

Ans. 1176.

3. In an arithmetical series, the first term is 12, the last term is 96, and the common difference is 6. Required the number of terms?

4. In an arithmetical series, the last term is an the common difference 1 and the sum of the series 1955 Required the number of terms of the control of the series 1955 Required the number of terms of the control of the series of the control of the control of the series of the control of the

5. The first term of an arithmetical series is 3, the common difference 3, and the sum of the series 1180. What is the last term ?.

6. If the extremes of an arithmetical series are 8 and 170 and the sum of the series 4895, what is the common difference?

7. If the extremes of an arithmetical series are 5 and 271 and the common difference 21, what is the number of terms?

Ans. 11.

8. If the first term of a series is 2, the last term 478 and the number of terms 86, what is the sum of the series?

9. In an arithmetical series the last term is 998, the first term 2 and the common difference 6. What is the sum of the series?

10. In an arithmetical series the first term is 5, the number of terms 11 and the common difference 21. What is the last term?

11. In an arithmetical series the last term is 199, the common difference is 11 and the number of terms 19. Required the sum of the series?

12. The sum of an arithmetical series is 39840, and the extremes are 2 and 478. What is the number of terms? Ans. 166.

13. The sum of an arithmetical spries is \$3500 and the extremes are 998 and 2. Required the common difference? Ans. 6.

15.

16.

17.

18.

19.

20.

21.

whe

or co 100 being

be c

, and

+ 431.

per of g, the 100. 7. and differ-1170.

t term Hum-15. c mmon ed the

15. mmon is the 39}.

70 and differ-3,0 10 71 and

erms? 11. nd the 7

20640. t term of the 83500. nber of is the

271. ommon equired 1900. tremes

1s. 166. tremes Ans. 6. 14. A snail crawls up a flag staff 130 feet high and upon reaching the top begins to descend. In what time will he again reach the ground if he goes 2 feet the first day, 4 feet the second, 6 feet the third, and so on? Ans. 15 days, 15 hours, 10 min. 27-264 sec.

15. The sum of an arithmetical series is 83500, the first term is 2 and the common difference 6, what is the last term? Ans. 998.

16. A person wishes to discharge a debt of \$1125 in 18 annual payments which shall increase in arithmetical progression. How much must his first payment be in order that the last may be \$120? Ans. \$5.

17. In an arithmetical series the extremes are 5 and 271 and the number of terms is 11. What is the common difference? Ans. 21.

18. 220 stones are placed in a straight line exactly 21 yards apart, the first being 21 yards from a basket, how far will a person go whilst picking up the stones, returning with one at a time and depositing it in the basket?

Ans. 69 1 miles. 19. The sum of an arithmetical series is 39840, the number of terms is 166 and the last term is 478. What is the first

20. A person travelled from Toronto to Kingston, in 12 days, walking 4 miles the first day, 6 miles the second, 8 miles the third, and so on. How far is Toronto from Ringston? Ans. 180 miles.

21. The clocks of Venice strike from 1 to 24. How many strokes does one of these clocks make in the day?

## GEOMETRICAL PROGRESSION.

15. Quantities are said to be in Geometrical Progression when they increase or decrease by a common multiplier.

Thus 3, 12, 48, 192, &c., are in geometrical progression, the common ratio or common multiplier being 4. 100, 20, 4, 4, 5, 25, &c., are in geometrical progression, the common ratio

being t.

16. In geometrical progression there are five things to be considered:

The first term.
 The last term.

3. The common ratio.

4. The number of terms. 5. The sum of the series.

1 1 1 10 . 1 a

As in arithmetical progression, these five quantitie and related that any three of them being given the other two can be found, and heare there are 20 dictinot cases arising from their combinations.

17. Representing these five quantities by letters, thus,

a = the first term.

r = the common ratio.

n = the number of terms.

• = the sum of the series.

the general expression for a geometrical series becomes

perion for a Reomentical series becomes

a+ar+ar2+ar2+ar4+ar5+, &c.,

where the index of r is always one less than the number of the term.

Thus in the third term the index of r is 2, which is one less than the number of the term; in the fifth term the index of r is 4, which is one less than the number of the term, &c.

Hence  $l = ar^{-1}$ ; that is, the last term is equal to the first term multiplied by the common ratio raised to that power which is indicated by one less than the number of terms.

18. Since the sum of the series is equal to the sum of all the terms.

$$s = a + ar + ar^{2} + ar^{3} + \dots + ar^{-3} + ar^{-2} + ar^{-1}$$
, multiplying by  $r$  we get  $sr = ar + ar^{2} + ar^{3} + \dots + ar^{-3} + ar^{-2} + ar^{-1} + ar^{-1}$ 

Hence 
$$sr-s = ar-a$$
; or  $s(r-1) = a(r-1)$ , and therefore  $s = \frac{a(r-1)}{r-1}$ 

That is, the sum of the series is found by finding that power of the common ratio which is expressed by the number of terms—subtracting 1 from this, dividing the remainder by one less than the common ratio and multiplying the quotient by the first term.

Note.—The second of the above series is found from the first by multiplying both sides of the equation by r; and in subtracting we take the terms of the upper series from the corresponding terms of the lower. Only the first three or four and the last three or four terms are written and between  $ar^2$  and  $ar^{-2}$  there may be any number of intermediate terms. The  $ar^{-2}$  in the lower series is obtained by multiplying the term before  $ar^{-3}$  in the upper series, which is  $ar^{-4}$ , by r.

19. From the formula obtained in Art. 17 we get by transposing the terms, &c.

$$l = ar^{-1}$$

$$a = \frac{l}{r^{-1}}$$

$$r = \left(\frac{l}{a}\right) \stackrel{!}{=} 1$$

$$r = \frac{\log l - \log a}{\log r} + 1$$

and

An

are t

geome be ok XIV but re ere

Ħ.

us,

a the e less

nultiy bne

m. of we get

(r-1) r-1

multike the before

et by

And substituting these values of l, a, r, n in the formula obtained in Art. 18 we find

$$\delta = \frac{rl - a}{r - 1}$$

$$\delta = \frac{l(r - 1)}{(r - 1)r^{n - 1}}$$

$$\delta = \frac{r^{n - 1} - a^{n - 1}}{\frac{1}{2r - 1} - \frac{1}{2r - 1}}$$

and these together with the two formulas obtained in Arts. 17 and 18.

$$e = \frac{a(r-1)}{r-1}$$

$$l = ar-1$$

are the fundamental formulas of geometrical progression from which the other fifteen are derived by reduction. Thus,

$$s = \frac{rl - a}{r - 1} \text{ gives formulas for s, r, l, and } a = 4$$

$$s = \frac{l(r^{n} - 1)}{(r - 1)r^{n-1}} \quad \text{" s, r, l, and } n = 4$$

$$s = \frac{a}{\frac{1}{p-1} - \frac{1}{a^{n-1}}} \quad \text{" s, r, a, and } a = 4$$

$$t = ar^{n-1} \quad \text{" s, r, a, and } n = 4$$

$$t = ar^{n-1} \quad \text{" l, a, r, and } n = 4$$

$$Total \ 20$$

20. The following table gives the 20 formulas for geometrical progression with their relations, &c. It will be observed that questions involving formulas III, XII, XIV, and XVI cannot be solved by common arithmetic, but require the aid of the higher mathematics. All the formulas for n involve the use of logarithms.

No.	Given.	Required.	Formulas.	Whence derived.
II. III.	a, r, n, a, r, s, a, n, s, r, n, s,	ı	$l = ar^{n-1}$ $l = \frac{a + (r - 1)s}{r}$ $l(s - l)^{n-1} - a(s - a)^{n-1} = 0$ $s = \frac{(r - 1)sr^{n-1}}{r^{n-1}}$	fundamental. VI. VII. VIII.
VI.	$a, r, n, a, r, l, a, \hat{n}, l, r, n, l, l,$		$s = \frac{a(r^{n}-1)}{r^{n}-1}$ $s = \frac{r^{1}-a}{r-1}$ $s = \frac{1}{l^{n}-1} - a^{\frac{1}{l^{n}-1}}$ $s = \frac{l(r^{n}-1)}{(r-1)r^{n}-1}$	fundamental  V. and I  V. and XIII  V. and IX
<b>X</b> :	r, n, l r, n, e r, l, e, n, l, e,	ciffic quis pe	$a = \frac{l}{r^{n-1}}$ $a = \frac{(r-1)s}{r^{n-1}}$ $a = r(l-s) + s$ $a(s-a)^{n-1} - l(s-l)^{n-1} = 0$	v. vi. vii.
XIV:	a, n, l, a, n, s, a, l, s,	1. 18 100 To a	$r = \left(\frac{l}{a}\right)^{\frac{1}{n-1}}$ $r = \frac{s}{a}r + \frac{s-a}{a} = 0$ $r = \frac{s-a}{s-1}$	I. v. v. v.
XVIII.	a, 7/3	Tennis	$r = \frac{s}{s-l}r^{s-1} + \frac{l}{s-l} = 0$ $n = \frac{\log l - \log a}{\log r} + 1$ $n = \frac{\log [a + (r-1)s] - \log a}{\log l - \log a}$ $n = \frac{\log l - \log a}{\log (s-a) - \log (s-l)} + \frac{\log l - \log (r^l - (r-1)s)}{\log r} + \frac{\log l - \log (r^l - (r-1)s)}{\log r} + \frac{\log l - \log (r^l - (r-1)s)}{\log r}$	VII

ART

2 nun

In raise of te

E

He Th 4.2

last

8 ; In the class t

Commall the

Hei The

num

INT
is ind
divide
Lat
the se Ex

sum comm

tel.

ıtal.

d I.

III.

IIX.

## APPLICATIONS.

21. Given the first term, the common ratio, and the number of terms, to find the last term:

$$l = ar^{n-1}$$
. (1.)

INTERPRETATION.—Multiply the first term by the common ratio raised to that power which is indicated by one less than the number of terms. The result will be the last term.

EXAMPLE.—What is the 9th term of the series 7, 21, 63, &c.?

Here 
$$a = 7, r = 3, and n = 9.$$

Then 
$$l = ar^{n-1} = 7 \times 3^{n-1} = 7 \times 3^n = 7 \times 6561 = 45927$$
. Ans.

22. Given the first term, the common ratio, and the last term, to find the sum of the series:—

RULE.

$$s = \frac{rl - a}{r - 1} \quad (VI.)$$

INTERPRETATION.—Subtract the first term from the product of the common ratio by the last term and divide the remainder by one less than the common ratio.

EXAMPLE.—The first term of a geometrical series is 5, the common ratio 4, and the last term 1000000. What is the sum of all the terms?

OPERATION.

Here 
$$a = 5$$
,  $r = 4$ , and  $l = 1000000$ .  
Then  $s = \frac{rl - a}{r - 1} = \frac{4 \times 1000000 - 5}{4 - 1} = \frac{3999995}{3} = 1333331\frac{3}{3}$ . Ans.

23. Given the first term, the common ratio and the number of terms, to find the sum of the series:—

RULE.

$$s = a \left( \frac{r^s - 1}{r - 1} \right) \text{ (v.)}$$

INTERPRETATION.—Find that power of the common ratio which is indicated by the number of terms, subtract one from it, and divide the remainder by one less than the common ratio.

Lastly, multiply the quotient thus obtained by the first term of the series, and the result will be the sum of all the terms.

EXAMPLE.—The first term of a geometrical series is 3, the common ratio is 4, and the number of terms 9. Required the sum of the series.

8.

10.

11.

12.

OPERATION,

Here a = 3, r = 4, and n = 9.

Then 
$$s = a \left( \frac{r-1}{r-1} \right) = 3 \times \frac{49-1}{4-1} = 3 \times \frac{262144-1}{3} = 962145$$
. Ans.

24. To find the common ratio when the first term, the last term, and the sum of the terms are given :--

el laving marking of the contraction of the contrac

Transport of the State of the said

The same of the best of the best of the

teck course indeed by une

INTERPRETATION .- Divide the difference between the first term and the sum by the difference between the last term and the sum; the quotient will be the common ratio.

EXAMPLE.—The first term of a geometrical series is 1, the last term 19683, and the sum of all the terms, 29524. What is the common ratio?

OPERATION.

Here a = 1, l = 19683, and s = 29524.

Then 
$$r = \frac{s-a}{s-l} = \frac{29524-1}{29524-19683} = \frac{29523}{9841} = 3$$
. Ans.

## 

- 1. A nobleman dying left 11 sons, to whom he bequeathed his property as follows: to the youngest he gave £1024; to the next, as much and a half: to the next 11 of the preceding son's share; and so on. What was the eldest son's fortune; and what was the amount of the nobleman's property? Ans. The eldest son received £59049, and the father was worth £175099.
- 2. The first term of a geometrical progression is 7, the last term is 1240029, and the sum of all the terms is 1860040. What is the ratio?
- 3. What debt can be discharged in a year by monthly payments in geometrical progression, the first term being £1, and the last £2048; and what will be the common ratio? Ans. The debt will be £4095; and the ratio 2.
- 4. The ratio of the terms of a geometrical progression is 3, the number of terms is 8, and the last term is 106493. What is the sum of all the terms? Jns. 307#1.
- 5. In a geometrical progression the first term is 1, the number of terms 7, and the common ratio 3, what is the sum of the series? Ans. 1093.

the

the

at is

his

: to

pre-

dest

ble-

049

last

040.

ents

and

io 2.

the

hat

nper

sum

6. The first term of a geometrical progression is 1, the last term is 10077696, and the number of terms is 10. What is the sum of all the terms?

Ans. 12093235.

7. The first term of a geometrical progression is 6, the last term is 3072, and the sum of all the terms is 6138. What is the ratio?

8. The ratio of the terms of a geometrical progression is 2, the number of terms is 11, and the sum of all the terms is 20470. What is the last term?

Ans. 10240.

9. A gentleman married his daughter on New Year's day, and gave her husband 1 shilling towards her portion, and was to double it on the first day of every month during the year. What was her portion?

Ans. £204 15s.

10. What will be the price of a horse sold for 1 farthing for the first nail in his shoes, 2 farthings for the second, 4 for the third, &c., allowing 8 nails in each shoe?

Ans. £4473924 5s. 3\frac{3}{2}d.

11. The first term of a geometrical progression is 4, the last term is 78732 and the number of terms is 10. What is the ratio?

12. A person travelling, goes 5 miles the first day, 10 miles the second day, 20 miles the third day, and so on increasing in geometrical progression. If he continue to travel in this way for 7 days, how far will he go the last day?

Ans. 320 miles.

13. The first term of a geometrical progression is 5, the last term is 327680, and the ratio is 4. What is the sum of all the terms?

Ans. 436905.

14. A king in India, named Sheran, wished (according to the Arabic author Asephad,) that Sessa, the inventor of chess, should himself choose a reward. He requested the king to give him 1 grain of wheat for the first square, 2 grains for the second square, 4 grains for the third square, and so on; reckoning for each of the 64 squares of the board twice as many grains as for the preceding. Sheran was angry at a demand apparently so insignificant; but when it was calculated, to his astonishment it was found to be an enormous quantity. What was the number of grains of wheat and what was its worth at \$1.50 per bushel, reckoning 7680 grains to a pint?

Ans. 18446744073709551615 grains. 37529996894754 bushels. \$56294995342131.

15. The ratio of the terms of a geometrical progression is 3, the number of terms is 10, and the sum of all the terms is 295240. What is the last term?

16. The first term of a geometrical progression is 1, the last term is 2048, and the number of terms is 12. What is the sum of all the terms?

17. The first term of a geometrical progression is 5, the ratio is 4. and the number of terms 9. What is the last term?

25. When the common ratio of a geometrical series is a proper fraction, i.e., less than 1, the series is a descending one, and when the number of terms becomes very large ra becomes very small. In an infinite descending series ra becomes infinitely small, i.e. its value becomes = 0, and therefore are may be neglected and the formula for finding the sum becomes

 $a = \frac{ar^n - a}{r-1} = \frac{-a}{r-1} = \frac{a}{1-r}$ . Hence for finding the sum of any infinite series when r is less than 1:-

$$s = \frac{a}{1-r}(xxi.)$$

INTERPRETATION.—The sum of an infinite series-is found by dividing the first term by unity minus the common ratio.

EXAMPLE 1.—What is the sum of the infinite series 1 + 1 + 26 + 128, &c. ?

OPERATION.

Here a=1 and r=t

~ 1 7 7 11 B.

EXAMPLE 2.—What is the sum of the infinite series .734? to produce of

OPERATION. Here a = 1000 and r = 1000. Then  $s = \frac{a}{1-r} = \frac{7}{1-\frac{340}{1000}} = \frac{734}{\frac{990}{1000}} = 734$ . Ans.

# EXERCISE 158.

1. What is the sum of the infinite series 2, 35, 18, &c.?.

2. What is the sum of the infinite series 4, 2, 1, 1, 1, &c.?

Ans. 8.

3. What is the sum of the infinite series '79? Ans. 78.

4. What is the sum of the infinite series :1234? Ans. 1333.

26. To insert any number of means between two given extremes:

by fo first differ mula ratio term

Ex

Sinc Then o

1st ter And se

EXA

Since th Then r:

Hence 2 And seri

1. Inser

2. Inse 3. Find

4. Find

27. means could n

NOTE. Error,

XI.

last t is

S. .

o is

390.

ion,

er of

fore

nite

. 8.

78.

334.

ven

#### RULE.

If the series is an arithmetical one, find the common difference by formula IX. ART. 8. Then add this common difference to the first term and the result will be the second term; add the common difference to the second and the result will be the third term. &c.

If the series is a geometrical one, find the common ratio by formula XIII. ART. 20. Then multiply the first term by the common ratio and the product will be the second term; multiply the second term by the common ratio and the result will be the third. &c.

EXAMPLE 1.—Insert 7 arithmetical means between 3 and 51,

## OPERATION.

Since there are 7 means and 2 extremes the number of terms is 9.

Then 
$$d = \frac{l-a}{n-1} = \frac{51-3}{9-1} = \frac{48}{8} = 6$$
.

1st term = 3; 2nd = 3 + 6 = 9; 3rd = 9 + 6 = 15; 4th = 15 + 6 = 21; 5th = 21 + 6 = 27; 6th = 27 + 6 = 33, and so on.

And series is 3, 9, 15, 21, 27, 33, 39, 45, 51.

Example 2.—Insert 6 geometrical means between 1 and 128.

#### OPERATION.

Since there are 6 means and 2 extremes the number of terms is 8.

Then 
$$r = \left(\frac{l}{a}\right) \frac{1}{a-1} = \left(\frac{128}{1}\right) \frac{1}{a-1} = (128) \frac{1}{2} = 2.$$

Hence 2nd term =  $1 \times 2 = 2$ ; 3rd term =  $2 \times 2 = 4$ ; 4th =  $4 \times 2 = 8$ , &c. And series is 1, 2, 4, 8, 16, 32, 64, 128.

### EXERCISE 159.

- 1. Insert 9 arithmetical means between 2 and 92.
  - Ans. 2, 11, 20, 29, 38, 47, 56, 65, 74, 83, 92,
- 2. Insert 4 arithmetical means between 7 and 50.
  - Ans. 7, 153, 241, 324, 413, 50.
- 3. Find 8 geometrical means between 4096 and 8.
  - Ans. 2048, 1024, 512, 256, 128, 64, 32, and 16.
- 4. Find 7 geometrical means between 14 and 23514624.
  - Ans. 84, 504, 3024, 18144, 108864, 653184, and 3919104.

### POSITION.

- 27. Position is a rule which enables us to solve, by means of assumed numbers, a class of problems which we could not otherwise solve without the aid of algebra.
- Norm.—Position is also called the Rule of False, or the Rule of Trial and Error,

repli

will

\*All frequent upon t

examp

age is equal t

B's =

EXA

28. Position is divided into:-

1st. Single Position—when only one assumed number is used.

2nd. Double Position—when two assumed numbers are used.

29. Single position is employed in the solution of those problems in which the required number is increased or decreased in any given ratio, i. e., when it is increased or diminished by any part of itself, or when it is multiplied or divided by any given number.

30. Double Position is employed in the solution of those problems in which the result found by increasing or decreasing the required number in any given ratio, is itself increased or diminished by some other number which is no known part or multiple of the required number.

## SINGLE POSITION.

31. Single Position proceeds upon the principle that the results are proportional to the numbers used, and is employed in all cases when the problem can be stated algebraically in the form of ax = b, where x = the required number, a the given multiplier, integral or fractional, and b the given result.

32. Let it be required to find a value of x such that ax = b. Suppose x' to be this value, and instead of b we obtain b' for the result. Then we have ax = b and ax' = b', and dividing we get  $\frac{ax'}{ax} = \frac{b'}{b}$  or  $\frac{x'}{x} = \frac{b'}{b}$  whence b':

Hence for single position we deduce the following:—

Assume a number, and perform with it the operations described in the question; then say, as the result obtained is to the number used, so is the true or given result to the number required.

EXAMPLE 1.—What number is that which being increased by its fourth part and diminished by its fifth part gives 63 for the result?

Assume any number, 40. Then one-fourth of number = 10, and one-fifth is

<sup>\*</sup>For the sake of convenience we assume a number of which we can take the required parts without using fractions.

um-

bers

hose

or

a of g or

tself s no

that

d is

ated

Te-

frac-

have

b':

11 .

ribed

mber

d by

r the

files

40 + 10 - 8 = 48, which by the question should have been 48.

Than—Result obtained; Result required; Number used; Number required.

Or, 48; 68:: 40; 68×40 = 60. And,

PROOF, -60 + 1 of 60 -1 of 60 = 68.

EXAMPLE 2.—A teacher being asked how many pupils he had, replied, if you add 1, 1, and 1 of the number together, the sum will be 18; what was their number?

### OPERATION.

Assume 60 to be the number of pupils,
Then one-third of 60 = 20
one-fourth of 60 = 15
one-sixth of 60 = 10

Bum = 45, but it should, by question, equal 18,

Then 45: 18:: 60: 45 - 24, Anc.

Proop. 45 of 24 × 1 of 24 = 18.

# It ou ti flin out Bregiorent 160. "

1. A gentleman distributed 78 pence among a number of poor persons, common of men, women, and children; to each man he gave 6d., to each women 4d., to each child 2d.; there were twice as many women as men, and three times as many children as women. How many were there of each?

Ans. 3 men, 6 women, and 18 children.

2. A person bought a chaise, horse, and harness, for £60; the horse came to twice the price of the harness, and the chaise to twice the price of the horse and harness. What did he give for each? Ans. He gave for the harness, £6 13s. 4d.; for the horse, £13 6s. 8d.; and for the chaise, £40.

3. A's age is double that of B's; B's is treble that of C's; and the sum of all their ages is 140. What is the age of each?

Ans. A's is 84, B's 42, and C's 14.

4. After paying away 1 of my money; and then 1 of the remainder, I had 72 guineas left. What had I at first?

Ans. 120 guineas.

\*All questions in position may be solved by simple analysis, and very frequently this is the better method, and indeed the teacher should insist upon the pupil thus solving each problem. The following will serve as examples of the mode of solution.

EXAMPLE 5.—Since 140 is equal to A's age, + B's age, + C's age, and B's age is equal to three times C's, and A's to 6 times C's, it follows that 140 is equal to 1+8+6=10 times C's age, and hence C's age is  $\frac{1}{10}$  of 140 = 14; B's = 14 × 3 = 42; and A's = 14 × 6 = 84.

tion

In thes resu

Th

Bu

Su

DI

proc

IF 01

the

will

is as

as h

duri

to the

- 5. A can do a piece of work in seven days; B can do the same in 5 days; and O in 6 days. In what time will all of them execute it?
  Ans. In 1484 days.
- execute it?

  6. A and B can do a piece of work in 10 days; A by himself can do it in 15 days. In what time will B do it?

  Ans. In 30 days.
- 7. A cistern has three pipes; when the first is opened all the water runs out in one hour; when the second is opened, it runs out in two hours; and when the third is opened, in three hours. In what time will it run out, if all the pipes are kept open together?

8. What is that number whose \$, \$ and \$ parts, taken together, make 27 ?

9. There are 5 mills; the first grinds 7 bushels of corn in 1 hour, the second 5 in the same time, the third 4, the fourth 3, and the fifth 1. In what time will the five grind 500 bushels, if they work together?

Ans. In 25 hours.

10. There is a cistern which can be filled by a pipe in 12 hours; it has another pipe in the bottom, by which it can be emptied in 18 hours. In what time will it be filled, if both are left open?

Ans. In 36 hours.

# DOUBLE POSITION.

38. When the number sought is to be increased or diminished by some absolute number, which is not a known multiple, or part of it—or when two propositions, neither of which can be banished, are contained in the problem, we use doubte position, assuming two numbers. If the number sought is, during the process indicated by the question, to be involved or evolved, we obtain only an approximation to the quantity required. In other words doubte position is employed in all cases in which the problem stated algebraically would take the form of

where  $\alpha$  is the number sought,  $\alpha$  the given multiplier, integral or fractional, b the given increment, and c the given result.

EXAMPLE 7. BY ANALYSIS.—Since A can do the whole work in 7 days, in 1 day he will do \(\frac{1}{2}\) of the whole work, similarly in 1 day B will do \(\frac{1}{2}\), and C \(\frac{1}{2}\) of the whole work. Therefore working together they will do \(\frac{1}{2}\) + \(\frac{1}{2}\) + \(\frac{1}{2}\) of the whole work, and they will require as many days to do the whole work as \(\frac{1}{2}\) is contained times in 1, i. e., \(1 + \frac{1}{2}\) = \(\frac{1}{2}\) \(\frac{1}{2}\) days. Anc.

ne in

34. Let it be required to find a value for a such as to satisfy the equation, ax + b = c.

In such a case assume any two known numbers n and n' and perform on these the operations indicated in the question, and let the errors in the result be s and s', both suppose in excess.

Then an + b = o + o (1) and an' + b = o + o' (11), and, by the question, an + b = o' (111).

Subtracting III from I we get an -aa = e, or a(n-a) = e(IV).

Subtracting III from II we get an'-an = d, or a (n'-s) = d' (V.)

Dividing IV by V we get  $\frac{a(n-a)}{a(n'-a)} = \frac{e}{a'}$  or  $\frac{n-a}{n'-a} = \frac{e}{a'}$ .

And reducing this we get  $a = \frac{n'e - ne'}{e - e'}$ .

Hence for double position we deduce the following:-

### RULH

I. Assume two convenient numbers, and perform upon them the processes supposed by the question, marking the error derived from each with form, according as it is an error of excess, or of defect.

11. Multiply each assumed number into the error which belongs to the other; and, if the errors are both plus, or both minus, divide the difference of the products by the difference of the errore. But, if one is a plus, and the other is a minus error, divide the sum of the products by the sum of the errors. In either case, the result will be the number sought, or an approximation to it.

Example 1.—There is a fish whose head in 8 feet long, his tail is as long as his head and half his body, and his body is as long as his head and tail; what is the whole length of the fish?

### OPRRATION.

Assume 24 ft. as the length of body.

Then tail = 8+4 of 24 = 8+12 = 20

Body = head + tail = 8+20 = 28

Assumed length of body = 24

Assumed length of body = 24

Then 64-9 = 82 = length of body 8+1 of 89 = 8+16 × 24 = " tail head

# 64 = length of fish.

EXAMPLE 2.—A laborer contracted to work 80 days for 75 cents per day, and to forfeit 50 cents for every day he should be idle during that time. He received \$25; now how many days did he work, and how many days was he idle?

them lays. mself ys. Il the

ened, ened, il the ours. ther,

hour,
th 3,
d 500
ours,
ours;
an be
od, if

od or nown either blem, f the y the n apwords e pro-

plier, given

lays, in and O the do the

# OPERATION. TELEVISION.

Suppose he worked 50 days; then he was idle 30 days.

Sum earned $= 50 \times 75 = $37.50$	True result = \$25.00
Sum forfeited $= 30 \times 50 = 15.00$	Result obtained = 22.50
Sum received = 22:50	Brror = 2'50

Again: suppose he worked 40 days; then he lost 40 days.

750 100

Difference of errors = 121. Difference of products = 650.

Therefore result required =  $650 \div 12 = 52$  days.

Number of idle days = 80-52 = 28. Ans.

PROOF.—Sum earned = 52 × 75 = \$39.00 Sum forfeited = 28 × 50 = 14.00

Sum received = \$25.00.

EXAMPLE 3.—What number is that which, being multiplied by 3, the product increased by 4, and that sum divided by 8, the quotient shall be 82?

### OPERATION.

Assume 40 to be the number.

Then  $40 \times 3 = 120 + 4 = 124 \div 8 = 151 = result obtained.$  82 = result required.

Error = -16

Again: assume 100 to be the number.

Then  $100 \times 3 = 300 + 4 = 304 \div 8 = 38 =$  result obtained. 32 = result required.

Sum of error = 221 Sum of products == 1890

> = 84. Ans. Required number ==

Proof.-84 × 8=252+4=256+8=32.

Norm.—In this example we take the sum of the errors for a divisor and sum of the products for a dividend, because the errors are not sothering to the means. Trun": 658 faringer dir.

है अस्ति व र १७% इंस्की रेसर व राज है। र १ ।

Ex

ART.

squar

It is multip upon w mation obtain

Fine they ar ent kin one of which

NOTE the assi ences o in the r proxim

EXA added

Assun

EXAMPLE.—What is that number which is equal to 4 times its square root + 21?

OPERA!	

ume 64		Assum	e 81
√64	= 8 4.	-	<b>√81 = 9</b> .
**	82 ·	dh e d	36 · · · · · · · · · · · · · · · · · · ·
	53, result obtained. 64, result required.	11,5	57, result obtained. 81, result required.
111	-11, difference,		-24, difference.
	891	**	1536 891

The first approximation is 49 6154

It is evident that 11 and 24 are not the errors in the assumed numbers multiplied or divided by the same quantity, and, therefore, as the reason upon which the rule is founded, does not apply, we obtain only an approximation. Substituting this, however, for one of the assumed numbers, we obtain a still nearer approximation.

### SECOND RULE.

Find the errors by the last rule; then divide their difference (if they are both of the same kind), or their sum (if they are of different kinds), into the product of the difference of the numbers and one of the errors. The quotient will be the correction of that error which has been used as multiplier.

Note.—This rule depends upon the principle that the difference between the assumed numbers and the true numbers are proportional to the difference between ences of the results obtained using the assumed numbers and that given in the problem. As in the last rule, when the question could not by algebra be resolved by an equation of the first degree, the rule gives only an approximation to the correct result.

EXAMPLE.—If to four times the price of my horse 2.0 be added the result will be £100. What is the price of my horse?

### OPHRATION.

	Assume	£10	and	second	ly £25 i	is the r	price of	the l	10186	1 Y' 5	1 12	
	4	The	n 19	1, 8 M 2.	Ye ji		1.7.1	25		Ø 14.	,	
	1	1 18	-	N 5.5" 15	1 1 1 1 1	· 7 , 2 00	7 1. 24	17.1	1		^	
ø	30 12	# 9 c	78	. treat	" 1 ,0	7 16 51 Fr 3.0	g. 74	100	1110	13. + + Fr	4	
				1				1	€ Ĭ	, , , , , , , , , , , , , , , , , , , ,	*	
٠,	2.21	88:	- 86,	the res	ult obt	sined.		1110	the	result of	btaine	d.
	7-11-3	1 1	100,	the res	ult req	uired.	17.17	100	, the	result r	Bouire	<u>d.</u> ,
	4,0	437	-14	is an em	ror of a	lefect.		-		error o		

ed by

24)84

numl

The errors are of different kinds: and their sum is 14+10=24; and the difference of the assumed numbers is 25-19=6. Therefore

is multiplied by 6, the difference of the numbers. Then divide by

and 3'5 is the correction for 19, the number which gave an

19+(the error being one of defect, the correction is to be added) 3.5=22.5 = 2.22 10s. is the required quantity.

# EXERCISE 161.

- 1. A son asked his father how old he was, and received the following answer: Your age is now 1 of mine, but 5 years ago it was only 5. What are their ages?

  Ans. 80 and 20.
- 2. Required what number it is from which if 34 be taken, 3 times the remainder will exceed it by 1 of itself?

  Ans. 582.
- 3. A and B go out of a town by the same road. A goes 2 miles each day; B goes 1 mile the first day, 2 the second, 3 the third, &c. When will B overtake A?

Suppose	A. B. 5 1 8 2 8	Suppose 7	1 2
943 12	40 4 5	56 28	5 8
w	)25 15 -5	7)28	28
of the s	35 20	20	inget ;

We divide the entire error by the number of days in each case, which gives the error in one day.

- 4. What are those numbers which, when added, make 25; but when one is halved and the other doubled, give equal results?

  Ans. 20 and 5.
- 5. Two contractors, A and B, are each to build a wall of equal dimensions; A employs as many men as finish 22½ perches in a day; B employs the first day as many as finish 6 per., the second as many as finish 9, the third as many as finish 12, &c. In what time will they have built an equal number of perches?

  Ans. 12 days.
- 6. What is the number whose 1, 1, and 7 multiplied together, make 24?

d the 17 1,

т. Х.

ve an

=22.2

d the out 5

d 20. times

 $.58_{7}^{2}$ 

miles 3 the

which

i; but equal and 5. equal erches 6 per., finish numdays.

gether,

Suppose 13	S	uppose 4
″ <u>‡</u> ≣\$		
Product = 18	Pro	
81, resu 24, resu	it obtained. it required.	8, result obtained 24, result required.
+57, erro 64, the	oube of 4.	—91, error, 1728, the cube of 12,
3648, pro	luct.	36288 to this product 3648 is added.
, 57 <b>+21=</b> 7	8	78)39234 is the sum,

And 512 the quotient. 2/512 = 8, is the required number.

We multiply the alternate error by the cube of the supposed number, because the error belongs to 3 part of the cube of the assumed numbers and not to the numbers themselves; for in reality it is the cube of some number that is required—since 8 being assumed, according to the question we have  $\frac{8}{2} \times \frac{8}{4} \times \frac{3 \times 8}{8} = 24$ ; or  $\frac{3}{64} \times 8^2 = 24$ .

- 7. What number is it whose 1, 1, 2, and 6, multiplied together, will produce 6998%? Ans. 36.
- 8. A said to B, give me one of your shillings and I shall have twice as many as you will have left. B answered, if you give me one shilling I shall have as many as you. How Ans. A 7, and B 5. many had each?
- 9. There are two numbers which, when added together, make 30; but the 1, 1, and 1 of the greater are equal to 1, 1, 1 of the lesser. What are they? Ans. 12 and 18.
- 10. A gentleman has 2 horses, and a saddle worth £50. The saddle, if set on the back of the first horse, will make his value double that of the second; but if set on the back of the second horse, will make his value treb!a that of the first. What is the value of each horse? Ans. - and £40.
- 11. A gentleman finding several beggars at his door, gave to each 4d. and had 6d. left, but if he had given 6d. to each. he would have 12d. too little. How many beggars were there?

16 12 1

gent . 24 hps light.

to

tra pay

wil be .

the

men

one

trac 100

# COMPOUND INTEREST.

35. Let P = the principal, I = the interest, A = the amount, t = the number of payments, and r = the rate per unit for one payment.

Then since r is the interest of \$1 for one payment, the amount of \$1 for one payment is 1 + r, and since the principal is always proportional to the

1: 
$$1+r: P: P: P: (1+r) =$$
 Amount of P at end of 1st period.  
1:  $1+r: P: P: (1+r): P: (1+r)^2 =$  Amount of P at end of 2nd period.

1: 
$$1+r: P(1-r)^2: P(1-r)^2: Amount of P at end of 3rd period.$$

1: 
$$1+r:: P(1+r)^3: P(1+r)^4 = Amount of P at end of 4th period.$$

And so on; hence at the end of the  $t^{th}$  period A = P(1+r); which is

$$\mathbf{A} = \mathbf{P} (\mathbf{1} + \mathbf{r}) \epsilon (\mathbf{I})$$

$$\mathbf{P} = \frac{\mathbf{A}}{(\mathbf{1} + \mathbf{r})^{4}} (\mathbf{II})$$

$$r=\sqrt{\frac{A}{P}}-1$$
 (III)

Obtaining as before 
$$(1+r)^i = \frac{\Lambda}{P}$$
 and applying the principle of logarithms we get log.  $(1+r)$   $\times t = \log \Lambda - \log P$ , and dividing each side

by log. 
$$(1+r)$$
 we get  $t = \frac{\log A - \log P}{\log (1+r)}$  which is (IV) of the margin.

$$t = \frac{\log n}{\log (1+r)}$$
 (V)

Lastly to find the time in which any sum of money will amount to 
$$n$$
 times itself at a given rate per cent. compound interest, we substitute  $nP$  for A in formula (1), which gives us  $nP = P (1+r)^s$  and dividing each of these by P we get  $n = (1+r)^s$  whence  $\log n = \log n$  (1+r) × t; or  $t = \frac{\log n}{\log (1+r)}$  which is formula (V).

# APPLICATIONS.

When the principal, rate per cent.. and time are given to find the amount:-

$$A = P (1+r)^t$$
 or  $\log A = \log P + \log (1+r) \times t$ . (I)

INTERPRETATION .- Multiply the logarithm of the amount of \$1 for one payment by the number of payments, and to the product add the logarithm of the principal; the result will be the isgarithm of the amount.

II. At the natural number corresponding to this logarithm and the read will be the answer.

PLE.—To what sum will \$750 amount in 3 years, at 2 cent., quarterly compound interest?

OPERATION. Then A = P (1 + r)' or  $\log A = \log P + \log (1 + r) \times t = 2.875061 + 0.00860 \times (2 = 2.978261 = \log Of Answer. Hence amount = $951.17.$  36. When the amount, rate, and time are given to find the principal:—

RULE.

$$P = \frac{A}{(1+r)^{r}}$$
; or log.  $P = \log A - \log (1+r) \times t$ . (II.)

INTERPRETATION.—Take the number expressing the amount of \$1 for one payment, and raise it to the power indicated by the number of payments.

II. Divide the given amount by the number thus obtained and the quotient will be the required principal.

# BY LOGARITHMS.

Take the logarithm of the amount of \$1 for one payment, and multiply it by the number of payments.

Subtract the logarithm thus obtained from the logarithm of the given amount; the remainder will be the logarithm of the required principal.

EXAMPLE.—What principal put out at compound interest, at the rate of 31 per cent. half yearly, will amount to \$8764.00 in 11 years?

Here A = 8764, r = 035 and t = 22.

Then  $P = \frac{\Lambda}{(1+r)^t}$  or log.  $P = \log \Lambda - \log (1+r) \times t$ .

 $\log_{100} P = 13.942702 - 0.014940 \times 22 = 3.942702 - 0.328680 = 3.614022$ . Hence P = 4411170. Ans.

37. When the amount, principal, and time are given to find the rate per cent:—

RULE.

$$r = t | \overline{\left(\frac{A}{P}\right)} - 1;$$
 or  $\log. (1+r) = \frac{\log. A - \log. P}{t}$  (III.)

INTERPRETATION.—Divide the amount by the principal, and extract that root of the quotient which is indicated by the number of payments.

II. Subtract 1 from the root thus obtained and the remainder will be the rate per unit, multiply this by 100 and the result will be the rate per cent.

### BY LOGARITHMS.

Subtract the logarithm of the principal from the logarithm of the given amount, and divide the difference by the number of payments; the result will be the logarithm of the amount of \$1 for one payment.

Find the natural number corresponding to this, and from it subtract 1, the result will be the rate per unit, and this multiplied by 100 gives the rate per cent.

= the

\$1 for to the eriod.

eriod. eriod. period.

nich is

(1+r)

pplying

ch side

sum of a given betitute us nP hese by = log.

given

(I) t of \$1 product garithm

hm and

rs, at 2

3 years. 375061 + 1'17. Example.—At what rate per cent. compound interest, payable half-yearly, will \$278 amount to \$6742 in 27 years?

Here A = 6742, P = 278 and t = 54.

Then log. 
$$(1+r) = \frac{\log A - \log P}{t} = \frac{3.828789 - 2.444045}{54} = \frac{1.384744}{54}$$

= 0255434. Hence 1+r=1.06, r=0.06, and rate per cent. = 6. Ans.

38. When the amount, principal, and rate are given to find the time:—

$$t = \frac{\log A - \log P}{\log (1+r)}$$
 (IV.)

INTERPRETATION.—Subtract the logarithm of the principal from the logarithm of the given amount, and divide the remainder by the logarithm of the amount of \$1 for one payment; the quotient will be the number of the payments.

EXAMPLE.—In what time will \$729 amount to \$7148 at 21 per cent. compound interest, quarterly?

Here 
$$A = 7148$$
,  $P = 729$  and  $r = 025$ .

Then 
$$t = \frac{\log A - \log P}{\log (1+r)} = \frac{3.853881 - 2.862728}{0.010724} = \frac{6.991153}{0.010724} = 92.42 \text{ payments} = 23.105 \text{ years} = 23 \text{ years 1 month 7.8 days.}$$

39. To find in what time any sum of money will amount to n times itself at any given rate per cent. compound interest:—

#### DTTT 'B

$$t = \frac{\log \cdot n}{\log \cdot (1+r)} \quad (V.)$$

INTELEPRETATION.—Find the logarithm of the number expressing to how many times itself the given sum is to amount, and divide it by the logarithm of the amount of \$1 for one payment; the result will be the required time.

EXAMPLE 1.—In what time will any sum of meney amount to five times itself at 5 per cent. per annum, compound interest?

Here 
$$n=5$$
 and  $r=05$ .

Then 
$$t = \frac{\log n}{\log (1+r)} = \frac{0.698970}{0.021169} = 32.987 \text{ yrs.} = 32.987 \text{ yrs$$

EXAMPLE 2.—In what time will any sum of money amount to nine times itself at 31 per cent. quarterly, compound interest?

1

•

3.

--

ß.

7.

9.

equ

upo 4 one

spec

fixed

### OPPRATION.

Here n = 9 and r = 035. Then  $t = \frac{\log n}{\log (1+r)} = \frac{0.054848}{0.014940} = 68.8716$  payments = 15.9679 years = 15 years 11 months 18 days. Ans.

# Exercise 162.

1. What is the amount and compound interest of \$713.29 for 7 years at 41 per cent. half yearly?

Ans. Amount = \$1320.96. Compound interest = \$ 607.67.

2. In what time will any sum of money amount to seven times itself at 1½ per cent. quarterly, compound interest?

Ans. 32 years 8 months 2 days.

3. In what time will \$111.11 amount to \$1111.11 at 8 per cent.

per annum, compound interest?

Ans. 29 years 11 mos.

- 4. At what rate per cent quarterly will \$222-22 amount to \$3333-33 in 30 years, compound interest being allowed?
- 5. In what time will any sum of money double itself at 7 per cent. per annum, compound interest?

  Ans. 10 years 2 months 28 days.
- 6. What principal put out at compound interest at the rate of 2½ per cent. quarterly will amount to \$100 in 7 years?
- 7. To what sum will \$2468.13 amount in 13 years at compound interest 32 per cent. half yearly?

  Ans. \$6427.705.
- 8. What principal will amount to \$7137.40 in 11 years, compound interest at the rate of 41 per cent. half yearly being allowed?

  Ans: \$2856.728.
- 9. In what time will any sum of money amount to 19 times itself at 51 per cent. half yearly, compound terest?

  Ans. 28 years 9 months 8 days.

# ANNUITIES:

40. An Annuity is any periodical income payable at equal intervals a yearly, half yearly, quarterly, &c.

41. An Annuity in possession is one that is entered

upon already.

42. An Annuity in reversion or a deferred annuity is one whose first payment is not to be made until after the expiration of a given time or until the occurrence of a specified event.

43. An Annuity certain is one that is to continue for a

fixed number of years.

able

ns. n to

from er by otient

per per

2 pay-

will com-

essing pide it result

unt to

nths 25

unt to

- 44. An Annuity contingent or a life annuity is one that is to continue to be paid only so long as one or more individuals shall live.
- 45. A Perpetuity is an annuity that is to continue for ever.
- 46. An Annuity is in arrears when one or more payments are retained after they have become due.
- 47. The amount of an annuity is the sum of the payments forborne (i.e. in arrears) and the whole interest due and them.
- 48. The present worth of an annuity is that sum which, being put out at interest until the annuity ceases, would produce a sum equal to what would have been accumulated had the annuity been left unpaid until that time.
- 49. Annuities are calculated at both simple and compound interest.

# ANNUITIES AT SIMPLE INTEREST.

**50.** Let a = a single payment of the annuity, t = number of payments r = rate per unit for one period, and A = amount of the annuity.

Then when the annuity is forborne a. number of payments, the last payment being made at the time it falls due, is equal to a; last payment but one = a + interest on a for one period = a + ar; last but two = a + interest on a for two payments = a + 2ar; last but three = a + 3ar; last but four = a + 4ar, a.c.; and hence the first payment = a + interest on a for one less than the number of payments = a + (t-1)ar.

Hence the payments forborne, with their interest, constitute a series in arithmetical progression where the first term is a, the last term a+(t-1) ar, the common difference ar, the sum of the series A, and the number of terms t.

Then (Art. 5.) A = a+(a+ar)+(a+2ar)+(a+3ar), &c.  $+\left\{a+(t-1)ar\right\}$ Whence (Art. 6.)  $A = \left\{a+(t-1)ar\right\} \frac{t}{2} = (1+\frac{(t-1)r}{2})$  to which is formula I in the margin.

$$\Delta = at \left(1 + \frac{(t-1)r}{2}\right) \quad \text{(I.)}$$

$$a = \frac{2\Delta}{t\left(2 + (t-1)\right)r} \quad \text{(II.)}$$

$$r = \frac{2(\Delta - at)}{at(t-1)} \quad \text{(III.)}$$

$$t = \sqrt{\left\{\frac{8r\Delta}{a} + (2-r)^2\right\} - (2-r)} \quad \text{(IV.)}$$

Formulas II., III., and IV., are derived from formula I, by transposition, &c.

one nore

for

pay-

paydue

iich.

ould

ated

com-

nents

e last

ment

= a +; last

ona

ies in

(t-1)ber of

)ar {

which

trans-

No general formula has yet been discovered for the summation of a series for finding the present value of an annuity at simple interest. The rule generally adopted for finding the present value of an annuity at simple interest is the following:—

Find the present worth of each payment by itself, discounting from the time it falls due—the sum of the present worth of all the payments will be the present worth of the annuity.

Note.—The absolute absurdity of purchasing annuities by simple interest is evident from the fact that the interest of the sum required to purchase an annuity, discounting at 5 per cent. simple interest, actually exceeds the annuity; i. e., to purchase an annuity to continue only a limited number of years, requires a sum which will yield a larger yearly interest for ever. Hence the various rules given for finding the present value of annuities at simple interest are, in effect, valueless.

# APPLICATIONS.

51. When the annuity, number of payments forborne, and the rate per cent. of interest are given, to find the amount:-

$$A = at \left\{ (1 + \frac{(t-1)r}{2} \right\}$$
 (1.)

INTERPRETATION.—Multiply the rate per unit by one less than the number of payments and to half the result add 1.

Multiply the number thus obtained by the product of the annuity by the number of payments and the result will be the required amount.

EXAMPLE.—If a pension of \$600 per annum be forborne 5 years, to what sum will it amount at 4 per cent. simple interest?

Here a = 600, t = 5, r = 04Then  $A = at \left\{ 1 + \frac{(t-1)r}{2} \right\} = 600 \times 5 \left\{ 1 + \frac{(5-1) \times 04}{2} \right\} = 8000 \times (1 + 1)$  $(08) = 3000 \times 1.08 = $3240$ . Ans.

52. When the amount of the annuity forborne, the number of payments forborne, and the rate per cent. of interest allowed, are given, to find the annuity: -

$$a = \frac{2.A}{t\{2 + (t-1)r\}}$$
 (II.)

INTERPRETATION.—Multiply the rate per unit by one less than

the number of payments and to the product add 2.

Multiply this sum by the number of payments, and divide twice the given amount of the annuity by the product thus obtained: the result will be the arnuity required.

For The after repre 55. riven.

intere

EXAMPLE.—What annuity payable quarterly, will amount to \$3225.25 in 7 years, at 41 per cent. per annum, simple interest?

### OPERATION.

Here since the rate is 41 per cent. per annum or '065 per unit per annum, the rate per quarter = '066 ÷ 4 = '01125.

Then 
$$t = 28$$
,  $A = $8225 \cdot 25$  and  $r = \cdot 01125$ .
$$a = \frac{2A}{t} = \frac{3225 \cdot 25 \times 2}{28 \cdot (28 - 1) \times \cdot 01125} = \frac{(6450 \cdot 50)}{28 \times (28 + \cdot 30875)}$$

6450'50 = 6450'50 = \$100 = quarterly payment, and hence annual annuity = \$400. Ans.

53. The application and interpretation of the remaining formulæ will be readily understood from the foregoing examples.

### Exercise 163.

1. In what time will an annuity of \$1000 per annum, payable half-yearly, amount to \$8365, allowing simple interest, at the rate of 6 per cent. per annum? Ans. 14 payments, or 7 years.

Note.—In this question we use formula IV, r being equal to '03 and a

2. If a rent of \$450 per annum, payable quarterly, be forborne for 11 years, to what does it amount, allowing 6 per cent. per annum simple interest?

NOTE.—Take a = \$112.50, r = 0.015 and t = 44.

3. At what rate per cent. per annum, simple interest, will an annuity of \$300, payable yearly, amount to \$1680 in 5 Ans. 6 per cent.

4. The rent of a farm is forborne for 8 years, and then amounts to \$2080. Now assuming the rent to be paid half-yearly, and simple interest at the rate of 8 per cent. per annum allowed, what was the rent of the farm? Ans. \$200.

# ANNUITIES AT COMPOUND INTEREST.

54. Let A, a, r, t = same quantities as in last articles and also let v = present value of the annuity.

Then, as before, the last payment of a forborne annuity being paid when due, = a; last payment but one, = a + interest of a for one payment = a + ar = a (1 + r); so also last payment but two, = a (1 + r); last but three = a (1 + r); and first payment = a (1 + r);

Hence A, the amount of the annuity  $= a + a (1 + r) + a (1 + r)^2 + a (1+r)^3 + a (1+r)^{-1}$  which is a geometrical series and is equal (Art. 18.)

int ple

ım,

76)

ual

ing ex-

ble est. its,

da

rne

nt.

71.

an.

5

nt.

nts

ly, ım

DO.

en e a out

8.)

$$A = \frac{a\{(1+r)-1\}}{r} (1)$$

$$a = \frac{Ar}{(1+r)-1}$$
 (II)

$$r = \sqrt[4]{\frac{dr+a}{a}} - 1 \text{ (III)}$$

$$t = \frac{\log. (Ar + a) - \log. a}{\log. (1+r)}$$
(IV)

$$v = \frac{a}{r} \left\{ 1 - \frac{1}{(1+r)^i} \right\}$$
 (V)

$$a = \frac{vr(1+r)^{t}}{(1+r)^{t}-1}$$
 (VI)

$$t = \frac{\log a - \log (a - vr)}{\log (1 + r)} (VII)$$

$$v = \frac{a}{r} \left\{ \frac{1}{(1+r)^{1}} \frac{1}{(1+r)^{r+1}} \right\} (rm)$$

$$v = \frac{a}{r} \text{ (IX)}$$

$$a = vr(X)$$

$$r = \frac{a}{v} (XI)$$

$$v = \frac{a}{r(1+r)!} \, (XII)$$

Since the present value of an annuity at compound interest is that principal which put out at compound interest for the given time, would produce the amount of the annuity we have from Art. 35, formula I, 
$$v$$
,  $(1+r)' = A = a \{(1+r)'-1\}$  whence by dividing by  $(1+r)'$ , we get formula V in the margin.

To find the present value of an annuity which is to commence after 
$$t$$
 years and then continue for  $s$  years, we have from formula  $V$ ,  $v$  for  $s+t$  years,  $=$   $\frac{a}{r}\left\{\frac{(1+r)^{s+t}-1}{(1+r)^{s+t}}\right\}$  and for  $t$  years

$$\frac{a}{r}\left\{\frac{(1+r)^{r+1}-1}{(1+r)^{r+1}}\right\} \text{ and for } t \text{ years}$$

alone, 
$$v = \frac{a}{r} \left\{ \frac{(1+r)i-1}{(1+r)^i} \right\}$$

$$\frac{a}{r} \left\{ \frac{(1+r)^{r+1}-1}{(1+r)^{r+1}} - \frac{(1+r)^{t}-1}{(1+r)^{t}} \right\}$$
or  $v = \frac{a}{r} \left\{ \frac{1}{r} - \frac{1}{r} \right\}$ 

or 
$$v = \frac{\sigma}{r} \left\{ \frac{1+r}{(1+r)^{r}} - \frac{1+r}{(1+r)^{r+r}} \right\}$$
 which is formula VIII in the margin.

When an annuity lasts for ever as in the case of landed property, 
$$(1+r)^{i}$$
 in formula V becomes infinitely great, and therefore

$$\frac{1}{(1+r)^c} = \frac{1}{cc} = 0$$
 and the formula for finding the present value of a perpetuity is reduced to the form given in IX.

Formulas X and XI are derived from IX.

The present value of a freehold estate to a person to whom it will revert after s years and then continue for ever, is found from formula VIII and is represented by formula XII in the margin.

<sup>55.</sup> To facilitate the calculation of annuities the following tables are given, the first showing the amount of an annuity of \$1 at compound interest, and the second, the present value of an annuity of \$1 at compound interest.

# TABLE OF THE AMOUNTS OF AN ANNUITY OF \$1 OR £1.

No. of Pay- ments.	8 per cent.	4 per cent.	5 per cent.	6 per cent
1 '	1.00000	1.00000	1.00000	1.00000
	2.08000	2.04000	2.05000	2.06000
2 3	8.09090	3.12160	8.15250	8.18860
4	4.18868	4.24646	4.81012	4.87462
5	5.80918	5.41632	5·52F38	5.68706
. 6	6.46841	6.63297	6.80.91	6.97582
6 7	7.66246	7.89829	8.14.401	8.39384
. 8	8.89234	9.21423	9.54.711	9.89747
9	10.15911	10.58279	11.02656	11.49181
10	11.46888	12.00611	12.57789	18.18079
11	12.80779	13.48635	14.20679	14.97164
12	14.19203	15.02580	15.91718	16.86994
18	15.61779	16.62684	17.71298	18.88214
14 15	17:08632	18-29191	19.59863	21.01506
16	18.59891	20.02359	21.57856	23.27598
17	20·15688 21·76159	21.82458	23.65749	25·67253 28·21288
18	28 41448	23·69751 25·64541	25·84037 28·13238	30.90565
19	25.11687	27.67123	80.53900	33.75999
20	26.27087	29.77808	33.06595	86.78559
21	28.67648	31.96920	35.71925	39.99273
22	30-53678	84-24797	38.50521	43.89229
28	82.45288	86-61789	41.43047	46.99588
24	34.42647	39.08260	44.50200	50.81558
25	36.45926	41.64591	47.72710	54.86451
26	88-55304	44.81174	51.11345	59.15639
27	40.70963	47.08431	54.66931	63.70576
28 29	42 93092	49.96758	58.40258	68-52811
29	45.21885	52.96629	62.82271	73.63980
80	47.57541	56.08494	66.43885	79.05819
81	50.00268	59.32833	70.76079	84.80168
82	52.50276	62.70147	75.29829	90.88978
83	55.07784	66.20958	80.06377	97.84816
34	57.78018	69.85791	85.06696	104.18375
.: 85	60.46208	78.65222	90.32031	111.48478
36	63.27594	77.59831	95.83623	119.12087
37 38	66.17422	81.70225	101·62814 107·70954	127·26812 185·90420
39	69·15945 72·23423	85·97034 90·40915	114.09502	145.05846
40	75.40126	95.02551	120.79977	154.76196
41	78-66330	99.82654	127.88976	165.04768
42	82.02320	104-81960	135.23175	175.95054
48	85.48389	110.01288	142-99334	187-50758
44	89.04841	115.41288	151-14300	199.75803
45	92.71986	121.02939	159.70015	212.74851
46	96-50416	126-87957	168-68516	226.50812
47	100-89650	132-94589	178-11924	241.09861
48	104 40839	189 26321	188-02539	256-56458
: 4 49 a	108-54065	145.88378	198-42666	272-95840
60	172-79-687	162 66708	209-84799	290.83590

# TABLE OF PRESENT VALUES OF AN ANNUITY OF \$1 OR £1.

No. of Pay- ments.	8 per cent.	4 per cent.	5 per cent.	6 per cent.
1	0.97097	0.96154	0.95238	0.94840
2	1.91847	1.88619	1.86941	1.83339
3	2.82861	2.77519	2.87519	2.67801
4 .	8 71710	3.62999	8.54595	8.46510
5	4.57971	4.45182	4.82948	4.21236
6	5.41719	5.24214	5.07569	4.91782
6 7 8 .9	6.28028	6.00205	5.78637	5.58238
. 8	7.01969	6.73274	6.46821	6.20979
. 9	7.78611	7.48538	7.10782	6.80169
10	8.58920	8.11089	7.72178	7.86009
11	9.25262	8.76068	8.80641	7.88687
12	9.95400	9.38507	8.86325	8.88884
18	10.68496	9.98565	9.89857	8.85268
14	11-29607	10·56312 11·11849	9.89864	9-29498
15	11.93794 12.56110	11.65239	10.87965	9·71225 10·10589
16 17	18-16612	12.16567	10.88777	10.10589
	18.75851	12.65940	11.68958	10.82760
18 19	14.32380	18.13394	12.08532	11.15811
20	14.87748	18.59032	12.46221	11.46992
20 21	15.41502	14.02916	12-82115	11.76407
22	15.93692	14.45111	18.16300	12.04158
23	16.44361	14.85648	18.48857	12.80888
24	16.93554	15.24696	13.79864	12.55036
25	17.41315	15.62208	14.09394	12.78835
25 26	17.87684	15.98277	14.87518	18.09316
27	18.32703	16.32958	14.64303	13.21058
28 29	18.76411	16.66306	14.89812	13.40616
29	19.18846	16.98371	15.14107	18.59072
30	19.60044	17.29203	15.87245	18.76483
31	20.00043	17.58849	15.59281	18-92908
32	20.38877	17.87855.	15.80267	14.08404
33	20.76579	18.14764	16.00255	14.23028
34	21.13184	18.41119	16.19290	14.36814
85	21.48722	18.66481	16.37419	14.49824
36	21·83225 22·16724	18·90828 19·14258	16.64685 16.71128	14·62099 14·73678
87	22.49246	19:36786		14.84602
38 39	22.80822	19.58448	16·86789 17·01704	14.94907
40	28.11477	19.79277	17.15908	15.94630
41	23.41240	19.99305	17.29436	15.18801
42	23.70136	20.18562	17.42320	15.22454
43	23.98190	20.37079	17.54591	15.80617
44	24.25428	20.54844	17.66277	15.88318
45	24.51871	20.72004	17.77407	15.45588
46	24.77545	20.88465	17.88006	15.52487
47	25.02471	21.04298	17.98101	15.58908
48	25.26677	21.19518	18.07714	15.6500
49	25.59166	21.50166	18.16872	15.7075
50	25.72977	21.72977	18-25592	15.7618

the

\$75

his

H

T

pay elaj and

pre

per

alld

H

# APPLICATIONS.

56. To find the amount of an annuity forborne for any number of years at compound interest:

RULI

$$A = \frac{a\{(1+r)^{i}-1\}}{r}$$
 (1.)

INTERPRETATION.—From the amount raised to the power indicated by the number of payments subtract 1 and multiply the remainder by the annuity. Lastly: divide the sum thus obtained by the rate per unit and the quotient will be the required amount.

By THE TABLE.—Find from the table the amount of \$1 for the given number of payments and at the given rate; multiply it by the given annuity and the quotient will be the amount.

EXAMPLE—If a yearly rent of \$400 be forborne for 23 years, to what sum will it amount at 5 per cent. compound interest?

### OPERATION.

Here 
$$a = 400$$
,  $t = 23$ ,  $r = .05$ .

Then  $A = \frac{a \left\{ (1+r)^4 - 1 \right\}}{r} = \frac{400 \left\{ (1.05)^2 \cdot 3 - 1 \right\}}{.05} = \frac{400 \times 2.071475}{.05} = \frac{828.590}{.05}$ 
= \$16571.80. Ans.

BY THE TABLE.—Amount of \$, at the given rate and time =  $\$41^{\circ}43047$ . Then  $\$41^{\circ}43047 \times 400 = \$16572^{\circ}188$ .

NOTE.—These two methods give results slightly different. This arises from the fact that the table shows only an approximation to the correct amount of the simulity for \$1; all the figures except the first five of its decimal being rejected.

57. To find the present value of an annuity at compound interest:—

$$v = \frac{a}{r} \left\{ 1 - \frac{1}{(1+r)^{t}} \right\} (v.)$$

INTERPRETATION.—Divide 1 by that power of the amount of \$1 which is indicated by the number of payments and subtract the result from 1.

Multiply the remainder by the quotient arising from the division of the given annuity by the rate per unit and the result will be the required present value.

BY THE TABLE.—Find the present value of an annuity of \$1 for the given number of payments and at the given rate, and multiply this by the given annuity.

U.

ny

rdithe

ned it. the

the

ars,

8.290

3047.

rises

und

f \$1 t the

vision

e the

1 for

ltiply

t?

Example.—What is the present value of an annuity of \$40, to continue 5 years, allowing 5 per cent. compound interest?

OPERATION.

Here a = 40, t = 5, and r = 05.

Then 
$$v = \frac{a}{r} \left\{ 1 - \frac{1}{(1 + r)^2} \right\} = \frac{40}{.05} \times \left\{ 1 - \frac{1}{(1 \cdot 05)^6} \right\} = \frac{4000}{5} \times (1 - .7835)$$

OR BY THE TABLE.—Present value of an annuity of \$1 for given rate and time = \$4\*32948 and \$4\*32948 × 40 = \$173\*179. Ans.

58. To find the present worth of a perpetuity:—

RULE

$$V=\frac{a}{r}$$
. (ix.)

INTERPRETATION.—Divide the annuity by the rate per unit and the quotient will be the value of the perpetuity.

EXAMPLE.—What is the present value of a freehold estate of \$75—allowing the purchaser 6 per cent. compound interest for his money?

OPERATION.

Here a = 75, and r = 06.

Then 
$$V = \frac{a}{r} = \frac{75}{06} = \frac{7500}{6} = $1250$$
. Ans.

59: To find the present worth of a perpetuity in reversion:—

RULE.

$$V = \frac{a}{r(1+r)}. \quad (x...)$$

INTERPETATION.—Find that power of the amount of \$1 for one payment that is indicated by the number of payments that have to elapse before the annuity reverts, multiply this by the rate per unit and divide the given annuity by the product—the result will be the present value.

EXAMPLE.—What is the present value of the reversion of a perpetuity of \$79.20 per annum, to commence 7 years hence—allowing the buyer 41 per cent. for his money?

OPERATION.

Here 
$$a = 79^{\circ}20$$
,  $s = 7$ , and  $r = 045$ .  
Then  $V = \frac{a}{r(1+r)^{\circ}} = \frac{79^{\circ}20}{045 \times (1+045)^{7}} = \frac{79^{\circ}20}{045 \times 1.360862} = \frac{79^{\circ}20}{06123879} = \frac{79^{\circ}20}{06123879}$ 

16 17 18

19 20, 21, 22

23 24, 25

1.

2.

60. With due attention to the foregoing interpretations and examples, the pupil will not experience any difficulty in applying the remaining formulæ.

# EXERCISE 164.

- 1. What is the annual rental of a freehold estate, purchased for \$3000 when the rate of interest is at 4 per cent.? Ans. \$120.
- 2. If a perpetuity of \$563 can be purchased for \$11260 ready money, what is the rate of interest allowed? Ans. 5 per cent.
- 3. A freehold estate producing \$75 per annum is mortgaged for the period of 14 years; what is its present value, reckoning compound interest at 5 per cent. per annum? Ans. \$757.608.
- 4. Required the present value of a deferred annuity of \$90, to be entered upon at the expiration of 12 years, and then to be continued for 7 years at 4 per cent. compound Ans. \$337.39. interest?
- 5. What is the present value of an estate whose rental is \$1500, allowing 5 per cent. compound interest? Ans. \$30000, or 20 years' purchase.
- 6. For how many years may an annuity of £22 be purchased for £308 12s. 10d., allowing compound interest at 4 per Ans. 21 years.
- 7. What is the present value of an annuity of \$154 for 19 years at 5 per cent. compound interest? Ans. \$1861.13.
- 8. What annuity, accumulating at 32 per cent. compound interest, will amount to £600 in 40 years? Ans. £6 13s. 11d.
- 9. In how many years will an annuity of \$8 per annum amount to \$187.315625 at 3 per cent. compound interest? Ans. 18 years.
- 19. What will an annuity of \$74 amount to in 30 years at 4 per cent. compound interest? Ans. 34150.28.

# QUESTIONS TO BE ANSWERED BY THE PUPIL.

NOTE.—The numbers after the questions refer to the numbered articles of the section.

- When are quantities said to be in arithmetical progression? (1)
   What are the extremes? What the means? (2)
   What five quantities are to be considered in arithmetical progression? (3)
   How are these related to each other? (3)
   How many cases arise from these combinations? (3)

culty

ed for

\$120. ready

cent. gaged value, num? 7.608. 90, to

1 then pound

37.39. \$1500, chase. chased t 4 per

years. 9 years

61.13. pound

s. 11d.

mount

years.

t 4 per

150.28.

articles

sion ?(3)

L.

- tions

  - 6. Deduce the fundamental formulæ for arithmetical progression. (4-7)
    7. When are quantities said to be in geometrical progression? (15)
    8. What five quantities are to be considered in geometrical progression? (16)
    9. How are these related and how many cases arise from their combinations? (16)
  - 10. Deduce the fundamental formulæ for geometrical progression. (17-19)

  - 11. What rule do you use when finding the sum of any infinite series when the ratio is less than 1? (25)

    12. Prove this rule. (25)

    13. How do we insert any number of arithmetical means between two given extremes ? (26)
  - 14. How do we insert any number of geometrical means between two extremes? (26)
    15. What is position? (27)

  - 16. Into what rules is position divided? (28)

  - 16. Into what rules is position divided? (29)

    17. When is a single position used? (29)

    18. What class of questions require the use of double position? (30)

    19. Give and prove the common rule for single position. (32)

    20. Give and prove the common rule for double position. (34)

    21. Deduce algebraically a complete set of rules for compound interest. (35)

    22. What is an annuity? (40)

    23. When is an annuity said to be in possession? (41)

    24. What is a deferred annuity or an annuity in reversion? (42)

    25. What is a contingent annuity? (41)

    26. What is a perpetuity? (45)

  - 26. What is a perpetuity? (45)
    27. When is an annuity said to be in arrears? (46)
    28. What is the amount of an annuity? (47)

  - 29. What is the amount of an annuity? (48)
    30. Deduce a set of rules for computing annuities at simple interest.
    31. Illustrate the absurdity and injustice of computing the present value of annuities at simple interest. (50)
    32. Deduce a set of rules for annuities at compound interest. (54)

### EXENCISE 165.

# EXAMINATION PROBLEMS.

## FIRST SERIES.

- 1. Write down as one number seven trillions and ninety millions, and nineteen and four million two hundred thousand and six hundredths of trillionths.
- 2. Deduct 19 per cent. from \$7580 and divide the remainder among A, B, C, and D, so that A may have \$111.11 more than B; B \$90.90 more than C, and D one third as much as A, B and C together.
- 3. A and B can perform a piece of work in 8 days, when the days are 12 hours long; A, by himself, can do it in 12 days, of 16 hours each. In how many days of 14 hours long will E do it?
- 4. Reduce £179 14s. 82d. to dollars and cents, and divide the result by '00000048.
- 5. What is the l. c. m. of 44, 18, 30, 77, 56 and 27?

6. In what time will any sum of money amount to 20 times itself at 5½ per cent. simple interest?

7. Divide 7342163 octenary by 61351 nonary, and give the answer in the duodenary scale true to two places to the right of the separating point.

8. Multiply 43 lbs. 3 oz. 17 dwt. 11 grs. by 7831.

9. Find the sum of the series 1+1+1+1, ad infinitum.

	1 06	2 06	100	3
10. Divide	# Ot	3. 01	194	43
				2

11. Extract the 17th root of 129140163.

12. There is a number consisting of two places of figures, which is equal to four times the sum of its digits, and if 18 be added to it, its digits will be inverted. What is the number?

# SECOND SERIES.

- 13. Divide \$897.43 among A, B and C, so that B may have \$93.40 less than A, and \$69.18 more than C.
- 14. If 7 lbs. of wheat contain as much nutritive matter as 9 lbs. of rye, and 5 lbs. of rye as much as 8 lbs. of oats, and 13 lbs. of oats as much as 21 lbs. of buckwheat, and 27 lbs. of buckwheat as much as 20 lbs. of barley, and 24 lbs. of barley as much as 26 lbs. of yeas, and 11 lbs. of peas as much as 35 lbs. of potatoes; how many pounds of potatoes contain as much nourishment as 16 lbs. of wheat?
- 15. Reduce  $\frac{2}{3}$  of  $4\frac{1}{4}$  of  $7\frac{1}{5}$  of  $\frac{9}{19\frac{1}{4}}$  of  $\frac{3}{6}$  of  $\frac{3}{2}$  of  $\frac{3}{13}$  of  $6\frac{1}{4}$  times 7 lbs. 3oz., Apothecaries Weight.
- 16. From 623.42793 take 93.4267192; mark distinctly the resulting repetend.
- 17. If I own a vessel valued at \$7493 and wish to insure it at a premium of 43 per cent. so as to recover, in case of the destruction of the vessel, both the premium paid and the value of the vessel, for what sum must I insure?
- 18. If 18 men in 20 weeks of 5 working days each, working 11 hours a day, dig 11 cellars, each 20 feet long, 16 feet wide

20. 21.

22.

23.

24.

25. 26.

27.

28.

40.

and 5 feet deep; how many men will be required to dig 24 cellars, each 22 feet square and 4 feet deep, in 36 weeks

of 6 days each, working 9 hours per day?

19. A certain number is divided by 9 and the quotient multiplied by 17; the product is then divided by 300 and 33 is added to the quotient; the result is next divided by 3, and from this quotient 31 is subtracted, and the resulting difference divided by 121. Now 1 of 3 of 4 of this last quotient is 2. Required the original number.

20. What is the 1. c. m. of 480, 768, 348, and 1176? 21. What is the G. C. M. of 17598, 46090, and 171347?

22. In a certain adventure A put in \$12000 for 4 months, then adding \$8000, he continued the whole 2 months longer; B put in \$25000, and after 3 months took out \$10000, and continued the rest for 3 months longer; C put in \$35000 for 2 months, then withdrawing 4 of his stock, continued the remainder for 4 months longer; they gained \$15000; what was the share of each?

23. Three merchants traffic in company, and their stock is £400; the money of A continued in trade 5 months, that of B 6 months, and that of C 9 months; and they gained £375, which they divide equally. What stock did each

put in?

24. A fountain has 4 pipes, A, B, C, and D, and under it stands a cistern, which can be filled by A in 6, by B in 8, by C in 10, and by D in 12 hours; the cistern has 4 pipes, E, F, G, and H; and can be emptied by E in 6, by F in 5, by G in 4, and by H in 3 hours. Suppose the cistern is full of water, and that 8 pipes are all open, in what time will it be emptied?

### THIRD SERIES.

25, Express 74938 and 17498679 in Roman Numerals.

26. 2310 loaves of bread are divided among charitable institutions in the following manner: as often as the first receives 4 the second receives 3, and as often as the first receives 6 the third gets 7; how many will each have?

27. How much sugar at 4, 5, and 9 cents a pound, must be mixed with 72 pounds at 12 cents a pound, so that the mixture

may be worth 8 cents a pound?

28. What principal put out at simple interest will amount to \$4444.44 in 4 years, 4 months 4 days at 4.44 per cent.?

29. For what sum must a ship valued at \$23470 be insured so as, in case of its destruction, to recover both the value of the vessel and the premium of 24 per cent.?

hat is

gures,

and if

the

the

have

g 9 lbs. and 13 27 lbs. lbs. of peas as of poieat?

scr. 5 times

the re-

e it at a of the and the

king 11 et wide

- 30. What principal will amount to \$7493.47 in 8 years, allowing simple interest at 7 per cent.?
- 31. I send to my agent in Manchester \$17460 and instruct him to deduct his commission at 3½ per cent., and invest the balance in broadcloths at \$2.95 per yard. When I receive the goods I have to pay in addition \$1347.90 for carriage, \$479.40 for insurance, \$169.83 for storage, wharfage, and harbour dues, and an ad valorem duty at 2½ per cent. on the invoice of goods. Required how many yards of cloth my agent ships to me and what I gain or lose per cent. on the whole transaction if I sell the goods for \$25000.
- 32. Transpose 134234 quinary into the ternary, octenary, and duodenary scales, and prove the results by reducing all four numbers to the denary scale.
- 33. What is the difference between  $\frac{2}{3}$  of  $\frac{9\frac{3}{2}}{\frac{1}{3}}$  of  $\frac{1}{16}$  of  $\frac{7}{3}$  of

£43 18s. 111d., and 3 of 171 of 56 of 1.75 of 61 times \$97.18?

34. Given the logarithm of 2 = 0.301030 3 = 0.47712113 = 1.113943

Find the logarithms of  $\frac{1}{13}$ , 19.5, 1125, 28.16, 65000, .0005, 152.1, and 8.112.

35. Extract the cube root of 871tet 72 duodenary true to two places to the right of the separating point.

36. A person passed & of his age in childhood, \(\frac{1}{13}\) of it in youth, \(\frac{1}{2}\) of it \(\frac{1}{2}\) 5 years in matrimony; he had then a son whom he survived 4 years, and who reached only \(\frac{1}{2}\) the age of his father. At what age did this person die?

### FOURTH SERIES.

37. Divide 63 miles 3 fur. 7 per. 3 yds. 2 ft. 7 in. by 7 fur. 23 per. 3½ yds.

55

56

58.

- 38. Divide 6.3 by .000000274.
- 39. If } yards of cloth cost \$19, how much will 613 yards cost?
- 40. Find the interest on \$4237.71 at 64 per cent. for 1.67 years.
- 41. In what time will \$674.30 amount to \$1000 at 81 per cent.?
- 42. What are the amount and compound interest of \$813.71 for 7 years at 4 per cent. half-yearly?
- 43. A owes B \$4300 to be paid as follows, viz.: \$300 down, \$700 at the end of 4 months, \$750 at the end of 7 months, \$850 at the end of 9 months, \$400 at the end of 13 months, and the balance at the end of 19 months. Required the equated time for the whole debt.

44. Deduct 23 per cent. from \$4200 and divide the remainder between A, B, C, D, and E, so that A may have \$17-10 more than B, C \$19.23 less than B, D \$42.11 less than C, and E half as much as A, B, C, and D together.

45. What principal put out at simple interest at 16 per cent. will

amount to \$3786.80 in 11 years?

46. Find the value of

$$\frac{\left(3\frac{3}{7}-2\frac{7}{6}\right)\times \cdot 46\div \frac{3}{7} \text{ of } \cdot 14285^{\frac{1}{7}}\right)\div 8\frac{1}{7} \text{ times } \left(\frac{1}{7}+\frac{1}{7}+\frac{3}{7}+\frac$$

47. Add together 312312302 and 2312132 quaternary; multiply the sum by twenty-three thousand and eleven times 4234 quinary; from the product subtract 555+444+333+222 +111 senary; divide the remainder by 6542 septenary. and give the answer in the octenary scale.

48. What is the square of 1 and also of 1?

### FIFTH SERIES.

49. Read the following numbers: 1000300500600.00070080009. 7600290034007.000000067400209.

50. Find the l. c. m. of 2, 9, 16, 27, 48, and 81.

51. In what time will any sum of money amount to 7 times itself at 6 per cent. per annum compound interest?

52. How often will a coach wheel turn in going from Toronto to Brampton, a distance of 20 miles; the wheel being 14 ft. 10 in. in circumference?

53. How many divisors has the number 1749600?

1 of 7 54. Divide  $\frac{2}{3}$  of  $\frac{1}{5}$  by

55. A can do a piece of work in 12 days, and A and B together can do it in 5 days; in what time can B alone do it?

56. What principal will amount to \$8899.77 in 11 years at 6

per cent. half yearly, compound interest?

57. Divide the number 10 into three such parts, that if the first be multiplied by 2, the second by 3, and the third by 4,

the three products will be equal.

58. There are three fishermen, A, B, and C, who have each caught a certain number of fish; when A's fish and B's are put together, they make 110; when B's and C's are put together they make 130; and when A's and C's are put together they make 120. If the fish be divided equally among them, what will be each man's share; and how many fish did each of them catch?

and all

im

he

Te

¿e,

ind

oth

on.

1.18?

3 of

0005, two

routh, whom age of

fur. 23

cost? years. cent.? .71 for

down, nonths, nonths, red the

- 59. What is the forty-seventh term and also the sum of the first 93 terms of the series 7, 11, 15, 19, &c.?
- 60. In what time will any sum of money amount to 21 times itself at 7 per cent. compound interest?

### SIXTH SERIES.

- 61. Divide \$3700 among three persons, A, B, and C, so that B may have \$387 less than A and \$196.87 more than C.
- 62. What are all the divisors of 5716?
- 63. What is the value of

$$\left\{ \frac{(17.7 - 108.9) - (\cdot 4 + \frac{1}{2} + \cdot 9 - \frac{1}{2})}{\cdot 6322632 \times \frac{1}{2} \text{ of } 9\frac{1}{2} \div (\frac{1}{2} \text{ of } 4\frac{1}{2} \text{ of } 85\frac{1}{2} + \frac{1}{2} \text{ 101})} \right\}$$

- 64. Divide \$7200 among 3 men, 4 women, and 17 children, giving each man twice as much as a woman, and each woman three times as much as a child. What is the share of each?
- 65. How many divisors has the number 25400?
- 66. What is the difference between  $\frac{9}{3}$  of  $4\frac{1}{2}$  of  $\frac{9\frac{3}{7}}{14}$  of  $\frac{1}{6}$  of £3 16s.

11\frac{1}{4}d. and 
$$\frac{3}{11}$$
 of  $\frac{43}{3}$  of  $\frac{19\frac{1}{4}}{13}$  of  $\frac{95}{117}$  of  $\frac{11}{23}$  of  $\cdot 85$  of  $\frac{1}{42\frac{1}{2}}$  of \$1783?

- 67. Compare together the ratios 7:13, 9:16, 8:15 and 10:19 and point out which is the greatest, which the least, and what the ratio compounded of these given ratios.
- 68. Divide 67.432 by 7.9036.
- 69. Reduce 9 per. 9 yds. 7 ft. 120 in. to the decimal of 1 of 3 of 3 of 35 acres 2 roods.
- 70. Add together 17.0342, 27.06357, 98.123456, 829.6423, 986.1234298, 9.876342, and 813.9864234567.
- 71. In the ruins of Persepolis there are two columns left standing upright. The one is 64 feet above the plain and the other 50. In a straight line between these stands a small statue, the head of which is 97 feet from the top of the higher column and 86 feet from the top of the lower, the base of which is 76 feet from the base of the statue. Required the distance between the tops of the columns.
- 72. In a mixture of spirits and water, \( \frac{1}{2} \) of the whole plus 25 gallons was spirits, but \( \frac{1}{2} \) of the whole minus 5 gallons was water. How many gallons were there of each?

### SEVENTH SERIES.

- 73. Extract the square root of 401241.3424 in the quinary weale.
- 74. A father being asked by his son how old he was, replied, your age is now tof mine; but 4 years ago it was only of what mine is now: what is the age of each?

75. Divide .72347 by .0032.

76. Extract the 11th root of 97294764.372.

- 77. Find two numbers, the difference of which is 30, and the relation between them as 7 18 to 31.
- 78. What is the l. c. m. of 35, 16, 18, 38, 62, 63 and 40?

9. Sum the series 1+7+13+19+&c., to 101 terms.

- 80. What is the ratio compounded of 19:7, 11:56, 35:121, 113:29, 8:42 and 44:3.
- 81. Find two numbers whose sum and product are equal, neither of them being 2.

Note.—In this question take any number for the first of the two, as for example 7. Then 7+some other number—7×that other number.

Assume for this second number any other, as 3.

Then  $7+3=10=7\times3$ , gives an error of—11. Assume some other for the second as 5.

Then  $7+5=12=7\times5$  gives an error of—23.

Then  $23\times3=69$   $11\times5=55$  Whence second number  $=\frac{14}{12}=1\frac{1}{6}$ .

82. Find the value of

$$\frac{\left(\left\{\left(9\frac{1}{3}+4\frac{1}{3}\right\}+3\frac{1}{7}-16\frac{3}{3}\frac{1}{6}\right)\times \cdot 54\right\}\div 14\right)\times 35 \text{ times } \cdot 142857.}{\left\{\cdot 97\times \cdot 24378\times \left(1\frac{1}{44}\times 4\frac{1}{467}\right)\right\}\times \left(4\frac{3}{1}-2\frac{1}{17}\right).}$$

83. The hour and minute hands of a watch are together at 12; when will they be together again?

84. Given the logarithm of 2 = 0.301030logarithm of 7 = 0.845098logarithm of 11 = 1.041393

Find the logarithms of 3850000, 3181.81, 0000154, 77,

# 1.571428 and 93.17.

#### EIGHTH SERIES.

85. Find the difference between the simple and compound interest of \$700 in 3 years at 41 per cent. per annum.

86. X, Y, and Z, form a company. X's stock is in trade 3 months, and he claims 12 of the gain; Y's stock is 9 months in trade; and Z advanced \$3024 for 4 months, and claims half the profit. How much did X and Y con-

at les

t B

31) 01)

cen, ach the

168.

183?


and


3 of

6423,


tandd the ds a op of ower, atue. ng.

us 25 llons






# IMAGE EVALUATION TEST TARGET (MT-3)



Photographic Sciences Corporation

23 WEST MAIN STREET WEBSTER, N.Y. 14580 (716) 872-4503

STATE OF THE STATE



87. There is a fraction which multiplied by the cube of 11 and divided by the square root of 17, produces 1; find it.

88. Find the cube root of 80677568161.

- 89. How much sugar, at 4d., 6d., and 8d. per lb. must there be in 112 lbs. of a mixture worth 7d. per lb.
- 90. Find three such numbers as that the first and 1 the sum of the other two, the second and i the sum of the other two, the third and 1 the sum of the other two, will make

NOTE.—Assume 40 as the sum of the three numbers.

Similarly assume some other number and apply the rule, and the true sum 58 will be found, from which the numbers may be easily obtained.

91. Insert 4 arithmetical means between 1 and 40.

- 92. The sum of all the terms of a geometrical progression is 1860040, the last term is 1240029, and the ratio is 3. What is the first term?
- 93. If 6 apples and 7 pears cost 33 pence, and 10 apples and 8 pears 44 pence, what is the price of one apple and one

94. Multiply 1 of 2 of 5 of  $\frac{281}{6}$  by 3 of 5 of 2.

- 95. From a sum of money, \$50 more than the half of it is first taken away; from the remainder, \$30 more than its fifth part; and again from the second remainder, \$20 more than its fourth part. At last there remained only \$10. What was the original sum?
- 96. A gentleman hires a servant, and promises him, for the first year, only \$60 in wages, but for each following year \$4 more than the preceding. How much will the servant receive for the 17th year of his engagement, and how much for all 17 years together?

### NINTH SERIES.

- 97. Write down as one number eleven trillions and eleven, and eleven tenths of billionths.
- 98. Reduce £749 16s. 53d. sterling to dollars and cents.

99. What are the prime factors of 177408?

100. At what rate per cent. per annum will \$704 amount to \$11111.11 in 11 years at compound interest?

101. How many scholars are there in a school to which if 9 be added the number will be augmented by one-thirteenth?

102. Three different kinds of wine were mixed together in such a way that for every 3 gallons of one kind there were 4 of another, and 7 of a third: what quantity of each kind was there in a mixture of 292 gallons?

103. Divide £500 among four persons, so that when A has £1,

B shall have £1, C 1, and D 1.

104. What is the present worth of an annuity of \$100 to continue 23 years, at 6 per cent. compound interest?

105. Twenty-five workmen have agreed to labor 12 hours a day for 24 days, to pay an advance made to them of \$900; but having each lost an hour per day, five of them engage to fulfil the agreement by working 12 days; how many hours per day must these labor?

106. A man has several sons, whose ages are in arithmetical progression; the age of the youngest is 5 years, the common difference of their ages is 6 years, and the sum of all

their ages is 161. What is the age of the eldest?

107. If a man dig a small square cellar, which will measure 6 feet each way, in one day, how long will it take him to dig a similar one that shall measure 10 feet each way?

108. A servant agreed to live with his master for £8 a year, and a suit of clothes. But being turned out at the end of 7 months, he received only £2 13s. 4d. and the suit of clothes: what was its value?

### TENTH SERIES.

- 109. What number is that of which \(\frac{1}{2}\), \(\frac{1}{2}\), and \(\frac{1}{2}\) added together, will make 48?
- 110. If an ox, whose girth is 6 feet, weighs 600 lbs., what is theweight of an ox whose girth is 8 feet?
- 111. Four women own a ball or butter, 5 inches in diameter. It is agreed that each shall take her share separately from the surface of the ball. How many inches of its diameter shall each take?
- 112. Divide 71213 43 by 12 342 in the nonary scale and extract the square root of the quotient true to three places to the right of the separating point.
- 113. Five merchants were in partnership for four years; the first put in \$60, then, 5 months after, \$800, and at length \$1500, four months before the end of the partnership; the second put in at first \$600, and six months after \$1800; the third put in \$400, and every six months after he added

ind

be of

her ake

3rd)

-2nd

true ned.

n is s 3.

970 8

first fifth nore \$10.

first r 34

rant how

and

t tg

\$500; the fourth did not contribute till 8 months after the commencement of the partnership; he then put in \$900, and repeated this sum every six months; the fifth put in no capital, but kept the accounts, for which the others agreed to pay him \$1.25 a day. What is each one's share of the gain, which was \$20,000?

114. In what time will any sum of money amount to 16 times itself at five per cent. per annum. 1st. at simple interest?

2nd. at compound interest?

115. Three persons purchased a house for \$9202; the first gave a certain sum; the second three times as much; and the third one and a half times as much as the two others to-

gether: what did each pay?

of workmen; the first numbered 25 men and the second 22; how many acres did each company clear, and what did the clearing cost per acre, knowing that the first company received \$86 more than the second?

117. The greatest of two numbers is 15 and the sum of their

squares is 346: what are the two numbers?

118. To what sum will \$1200 amount in 10 years at 64 per

6. cent. simple interest?

119. If 496 men, in 5½ days of 11 hours each, dig a trench of 7 degrees of hardness, 465 feet long, 3¾ wide, 2½ deep, in how many days of 9 hours long will 24 men dig a trench of 4 degrees of hardness, 337½ feet long, 5¾ wide, and 3½ deep?

120. Four men, A, B, C, and D, took a prize of \$6213, which they are to divide in proportion to a following fractions: if possible, A, B, and C, are to be \$7; B, C, and D, \$7; A, C, and D, \$70; and A, B, and D, \$10f the prize. What

does each receive?

# ELEVENTE SERIES.

121. Reduce : 7, :83, :727, :91325 and 8:671347 to their equivalent vulgar fractions.

122. Reduce 713301 undenary, and 12123100000 quaternary to

equivalent expressions in the denary scale.

123. Add together 33 of 23 of 720 of a £, 93 of 33 of a shilling, and 82 of 42 of a penny, and divide the sum by 12 of 574

of 1 of 31d.

124. If 24 pioneers, in 2½ days of 12½ hours long, can dig a trench 139.75 yds. long, 4½ yds. wide, and 2½ yds. deep, what length of trench will 90 pioneers dig in 4½ days of 9½ hours long, the trench being 4½ yds. wide, and 3½ yds, deep?

125. A person, by disposing of goods for \$182, loses at the rate of 9 per cent.; what ought they to have been sold for to realize a profit of 7 per cent.?

126. In what time will any sum of money amount to 11½ times itself at 6 per cent. per annum.

1st At simple interest?
2sd At compound interest?

- 127. It is desired to cut off an acre of land from a field 151 perches in breadth; what length must be taken?
- 128. Express a degree (69 miles) in metres, when 32 metres are equal to 35 yds.
- 129. Find 7 geometrical means between 3 and 19685.

130. Sum the infinite series  $7+1\frac{3}{4}+\frac{7}{16}$ , &c.

131. Four men bought a grindstone of 60 inches diameter.

Now, how much of the diameter must be ground off by
each man, one grinding his part first, then another, and
so on, that each may have an equal share of the stone, no
allowance being made for the axle?

132. Divide 100 guineas into an equal number of guineas, half-guineas, crowns, half-crowns, shillings, and sixpences, and reduce the remainder to a fraction of a pound.

# TWELFTH SERIES.

- 133. The owner of  $\frac{1}{1}$  of a ship sold  $\frac{1}{1}$  of  $\frac{3}{3}$  of his share for \$12\frac{3}{3}; what would  $\frac{2\frac{1}{4}}{4}$  of  $\frac{3}{4}$  of the ship cost at the same
- 134. At what rate per cent. per annum will \$700-90 amount to \$1679-40 in 5 years, compound interest being allowed?
- 135. A person paid a tax of 10 per cent. on his income; what must his income have been, when, after he had paid the tax, there was \$1250 remaining?
- 136. The sum of £3 13s. 6d. is to be divided among 21 men, 21 women, and 21 children, so that a woman may have as much as two children, and a man as much as a woman and a child: what will each man, woman, and child receive?
- ceive as much as A; C as much as A and B together, and D as much as A, B, and C together.
- 138. Find the difference between √3 and ∜3.
- 139. Reduce  $\frac{3273}{52737}$ ,  $17\frac{5}{12} + \frac{1}{15} + 144\frac{1}{21}$ ,  $2\frac{1}{23} \frac{1}{27}$ ,  $\frac{3}{2}$  of  $\frac{4}{7} \times \frac{1}{15}$  of  $\frac{4}{15}$ , and  $6347 \div 2\frac{3}{2}$ , to their simplest forms.
- 140. Find the cube root of 884736, and the fourth root of 95951

mes ? est?

the

900,

t in

hers

nare

the to-

nies cond what com-

per

their

h of p, in ench d 31

hich ions: , %; What.

uiva-

lling,

dig a deep, ys of yds,

- 141. A general levied a contribution of \$520 on four villages, containing 250, 300, 400, and 500 inhabitants respectively: what must they each pay?
- 142. A person had a salary of \$520 a year, and let it remain unpaid for 17 years. How much had he to receive at the end of that time, allowing 6 per cent. per annum co pound interest, payable half-yearly?
- 143. Insert four arithmetical means between 2 and 79; also find the 9th term and the sum of the first 207 terms of the series 3, 7, 11, 15, &c.
- 144. A. B. and C. start at the same time, from the same point, and in the same direction, round an island 73 miles in circumference; A goes at the rate of 6, B at the rate of 10, and C at the rate of 16 miles per day. In what time will they be all together again?

# ARITHMETICAL RECREATIONS.

- 1. If the third of 6 be 8 what must the fourth of 20 be?

britisms with takes note

- If the half of 5 he 7 what part of 9 will be 11?
  Place four states so that their sum shall be 100.
  What part of 5 pence is the third of two pence?
- 5. If a herring and a half cost 1 d, how much will 11 herrings cost ?
- 6. If 13 apples are worth 21 pears, and 8 pears cost a cent, what will be the e of 100 apples ? 7. Find a number such that 5 shall be the three-sevenths of it.
- undred hurdles are so placed as to inclose 200 sheep, and with two urdles more the field may be made to held 100 sheep, and with two hurdles more the field may be made to hold 400; how is this to be
- 9. A gentleman who owned four hundred acres of land an the form of a square, desired to keep 100 acres also in the form of a square in one corner, and divide the remainder, a b c d e f, equally among his four sons, so that each son should have his tot of the same shape as his brother's. How may this be done?



- 10. Place four threes so as to make 84.
- 11. Write down 18 in such a way that rubbing helf of it out 3 shall remain.
- 18. Two thirsty persons cast away on a desert inlead, find an 8 stallon cask of water. They wish to divide it equally between them, but have no other measures than the 8 gallon cask of five gallon cask and a three gallon cask. How can they divide it?
- 13. How must a board 16 inches long and 9 inches wide he out into two such parts, that when they are joined together they may form a square?
- 14. Place the 9 digits in the socompanying figure, one digit to each division, in such a way that when added vertically horisontally or diagonally, the sum shall glways be the



tive-

main co · -

o find eries

point. a cirf 10. will

be the

th two s to be

m cask and a

to two

15. Three persons bought a quantity of sugar weighing 51 lbs., and wish to part it equally between them. They have no weights but a 41b. weight and a 7 lb. weight. How can they divide it?

16. Suppose 26 hurdles can be placed in a rectangular form so as to inclose 40 square yards of ground; how can they be placed when two of them are taken away, so as to inclose 150 square yards?

17. A person has a fox, a goose and a peck of case to carry over a river, but on account of the smallness of the boat he can only carry over one at a time. How can this be done so as not to leave the fox with the goose, nor the goose with the cate?

18. In a distant and sparsely settled village of Canada, there was stationed a small detachment of treams application of a particular and 24 men.

goose, nor the goose with the oats?

In a distant and sparsely settled village of Canada, there was stationed a small detachment of troops consisting of a sergeant and 28 men. Having constructed temporary barracks, the sergeant and 28 men. Having constructed temporary barracks, the sergeant and 28 men. Having compartments, allotting the centre one to himself, and the rest to his men. One evening the sergeant wishing to ascertain if all were in, visited each compartment, and finding 3 men in each making 9 in each row, retired. Four men, however, went out, and the sergeant feeling shortly afterwards uneasy, returned to count his men, but still finding 9 in each row, retired again; the 6 men then came back, bringing each another man with him, and the sergeant upon going his round once more, counted as before, and retired perfectly satisfied. After he left, four more men were introduced, and once more the sergeant once men the sergeant imposed that all was not right, counted, but finding the number still the same lineach row, he left. No sooner had he left, then four more men come in, making 11 strangers; and once more the sergeant imposed the dominar monta to his satisfaction. Finally the 13 strangers left, taking with them 6 of the schliers, and the sergeant counting once more retired to year perfectly the montant of the sorders and the sergeant counting once more friend to year perfectly that no one had gone out or come in, and that his unspectors were unfounded. How was this possible?

Write down 15 so that by rubbing out one half 7 shall remain.

19. Write down 15 so that by rubbing out one half 7 shall remain.

20. Place the first 25 numbers 1, 2, 3, 4, 5, &c., in the divisions of the accompanying figure, so that the columns added in any order, i. e., upwards, horisontally, or disconally, may amount to the same sum.



11. What is the difference between half-a-dozen dozen and six dozen

22. If a cross be made of 13 counters as in the margin, wise
may be reckeded in three ways, i. s., by counting from
the bottom up to the top of the perpendicular line;
from the bottom up to the cross and then to the right;
or from the bottom up to the cross and then to the right;
or from the bottom up to the cross and then to the left. Now take away two of the counters and with
the others form a cross which shall possess the same
property of counting wise when thus reckoned.



property of counting with when thus reckoned.


28. Seven out of \$1 bottles being full of wine, 7 half full and 7 cuspts it is required to distribute them among 5 persons, so thus each may have the wine quantity of wine and the tame number of heetles.

24. Two travellers, one of whom had with him 5 bottles of wine and the other 3, were joined by a third person, who, after the wine was drunk, left 3 shillings for his just share of it; how is this to be divided between the other two!

25. A person having by accident broken a basket of eggs, offered to pay for them on the spot if the owner could tell how many he had; to which he counted them by 5's and 5's at a time none remained; but what he counted them by 5's and 5's at a time none remained; but what he counted them by 5's and 5's at a time none remained; but what he counted them by 5's and 5's at a time none remained; but what he counted them by 5 at a time there were 5 remaining; how many eggs had he?

- 26. It is required to find 4 such weights that they weigh any number of pounds from 1 to 40.
- pounds from 1 to 40.

  27. In the accompanying figure it is required to fill seven out of the eight points with counters in the following manner, i. e., the counter has to start from an encoupted point, pass along the line and be deposited at the other extremity. Thus, in commencing, the counter may start from any point, since all are unoccupied, starting from 1 the counter may be carried either to 6 or to 4 and there deposited, suppose it be deposited at 6, then the next counter may start from any point exmay start from any point ex-cept 6, and so on.



- 28. A brasen lion, placed in the middle of a reservoir, throws out water from its mouth, its eyes and its right foot. When the water flows from its mouth alone, it fills the reservoir in 6 hours; from the right eye it fills it in 2 days; from the left eye in 3 days, and from the foot in 4 days. In what time will the basin be filled by the water flowing from all these apertures at once?
- 20. Degree a person to think of any three numbers, each less than 10, and then tell him the numbers thought of.

  30. Three men, Jones, Brown, and Smith, with their sons Harry, Tom and Ned, had each a piece of land in the form of a square. Jones piece was 13 rods longer on each side than Tom's, and Brown's piece was 11 rods longer on each side than Harry's. Each man possessed 63 square rods of land more than his son. Which of the persons were father and son respectively?
- 51. A sea-captain, on a voyage, had a crew of 30 men, half of whom were blacks. Being becamed on the passage for a long time, their provisions began to fail, and the captain became satisfied that, unless the number of men were greatly diminished, all would periah of hunger before they could reach any friendly port. He therefore proposed to the sailors that they should stand in a row on deck, and that every minth man should be thrown over-board, until one-half of the crew were thus destroyed. To this they all agreed. How should they stand so as to save the whitee?
- s. Direct a person to multiply together two numbers, one of which you select, and, unseen by you, to rub out one of the digits of the product—it is required to tell, upon his reading the remaining digits of the product, what figure was rubbed out.
- 11 is required to write down beforehand the answer to a question in addition of a given number of lines, you writing the second, fourth, statis, &c., addends, and some other person the intermediate ones. er fin calific to field fit must filter had arrite to see a tiltering all his filtering and his call the second part of the sec

water r flows te right the foot flowing

10, and

om and si piece lece was essed 63 ns were

om were neir pro-i, unless erish of fore pro-eck, and e-half of

should

hich you the pro-digits of

fourth,

# MATHEMATICAL TABLES.

LOGABITHMS OF NUMBERS FROM 1 TO 10,000, WITH DIFFERENCES AND PROPORTIONAL PARTS.

2 700 2 700	e le,	, 4	2 1 1-3·	,01	from 1 to	Albert 1	holy as with a	(5,3)	allywell is closed the
۱۰.	Lug.	No.	Log.	No.	Log.	No.	Les.	No.	Log.
< -	0.000000	21	1-322219	41	1-612784		St.	h	Par. 7 " 3. 4
1	0-301030	22	1-342423	42	1-620040	61	1-796800	81	1-90048
3	0-477121	23	1.361728	100	1-633468	63	1-796541	ž 85	1-918814
_	0.602060		1.880211	4	1-643453	64	1-806180	84	1-43447
4	0-602000	1000	1.397940	45	1-653218	65	1-812018	85	1-600/1
9	\$ ?	-	1.09(1950	1 - 1	4 40210	Farty.			
100	1 ,17		34	. 3	6.6 6"		- I will	1	-125
6	0-778151	26	1.414978	46	1-662758	66	1.819544	: 86	1-00469
7	0-845098		1.431364	. 47	1.672008	67	1-826075	87	1-99061
8	0-903090	4	1-447158	48	1-681341	68	1-852000	85	1-94440
	0-954243	-	1.462398	40	1-690196	10	1.838849	80	1-94000
10	1 000000	30	1.477121	50	1-608076	3)	1.845098	1 90	1-16484
	1 10	300	1000	1	ake.		Asse 15		19/15
11	1.041303	31	1-491362	51	1-707570	71	1.851258	91	1-96004
12	1-079181	- 32	1.505150		1.716003	72	1.857382	§ 92	1-96878
13	1-113945		1.518514	, 58	1-724276	78	1.863323	98	1-96848
14	1-146128		1.531479		1.732394	.74	1-869232	1 96	1-97319
15	1.176091	35	3200008	55	1.740363	75	1.875061	95	1.87773
	Ade	100	4 4	-	1 -		16 1 565		E pillica e
16	1-204120	36	1-556303	56	1-748188	76	1.880614	96	1-96937
17	1-230440		1-566202		1.755875	77	1.886491	97	1.98677
18	1-258273	38	1.579784		1.763428	.78	1-802095	98	1-99120
	-	100			The second second		59	1 5 .	200

PP	N.	•	1	9	2	4	5		9			D.
_		A										
4	100	4951	000434 4751	000008 5181	001301 5600	. 6085		002508	7381	774	008891 8174	を
12		01.2837 7081	01202	9451 013600 7445	914100 834	010300 4651	010734	011147	011570	011998	012415 0616 080778	400
307 348	8	091186 5806	021 dos	6125	022435 6583	023541 023541 0042	7360	(1866) 7757	02 1076 8164	8571	4896 1 8978	
200 201 201 201	I	9884 988434 7486	9789 093836 7825	080196 4237 8238	080600 4628 8680	031004 . 5039 9017	031408 5430 9414	031818 5680 9611	082216 6230 040307	033619	009081 7088 04088	11864
h	110	041398	041787	049199	042576		045303	043755	044146	044540	044003	200
76 113	7 3 7 8	, 6548 9318	8014 9004 963463	9995	050380	-0559	7375	051538	061884 5068	0445	6600 052694	350 366
调		063078 0005 000698	7286 061075	7606	9046	8426 067306	8805 062583	9185	9568 068388	6142 9043 063709	0034 000390 4083	233 573 574
227 365	7	4458 8186	4832 8557	5206 8028	5590	5963	6326 070038	062958 6699 070407	7071	7448	7815	376 378 370
305 340	8	071888 5547	073250 5912	072617 6276	9298 072988 6640	9668 073362 7004	3718 7368	4085 7731	070776 4451 8004	4816 8457	5182 8619	366 363
	130	079181	079548 038144	079004	080266	060626	000007	081347 4934 8400	061707	082067	000406	100
104	į	000	6716 Caluada	7071	7400 7400 00000	7781	8196	8400	00250	9198 002721	089496 9054 9658 069071	25. 25. 28.
湿	- 4		腦	4122 7604 101080		4820	-	- 5518	9335	9681	100026	346
	7		4716	4487	401418	101747 5100	100001 5510	103434 5351	6191	108119 6531 9916	8463 6871 110863	343 341
388		110200	110008		111500	111984	112270	3 - 5 12 - 4	112040	113275	3600	335
	130	113949	11457 7018	214611 7944	1144	118278 8506 121888	115611 8936	118045	118276 9586	110008	11 <b>00</b> 40 120 <b>9</b> 45	333 330 336
1.2	8	120074 8688 7106	7439	4504 7788	121500 430	· 6790	··· 9491	122544 5806 9045	122871 6131 9364	6456	3626 6781 130012	1 330
198		3 85 90	180655	130077	131306	131619	131960 5136 8366		153589	152900	8319	311
35	1	6721 9879		7364 140608	7671	7987 141136	14140	5451 8618 141768 4885	5700 8034 142076	9249 142379	9564 143702	316 814
	140	148015	2537 146456	3680	1 4000	-	1574		-	5507	4818	311
8	1	9819 154968	9527	9835 152000	3805	147367 150449 3510 6540	147676 150746 3815	4120	4424	101076	148911 151982 5032	309 307 305 303 301
120	8	8068	182504 5640 8664	- 5943 9946	9246 9266	SOU!	3000	7154 100168	10040	100700	161068	303 301
180	6	161566 4858 7817	161007 4650 7613	161967 4947, 7908	5244	162064 5541 8497	5838	3161 6134 9086	9430 9380	5768 5726	4066 7082	200 207
240) 270	8	170003 51.86	170855 3478		171141	171434	171728	173019		172006 5512	172505	293 201
場と	150	176191	176381	176670	178969	177248 180126	177536	1778 <b>2</b> 5 180699	179119	178401	178600 181658	280
24	2	181844	9264 183139 4075	9552 182415 5250	182700 5543	180126 2985 5825	180413 3270 6108	180699 8565 6391	100900	181272	4807	267 265
113	4 6	100333		190892	191171	8647	191730	192010	9490 192280	192507	100051	370
100	7	3195 5800	3408	6453	5 300 0729	4257 7006	7281	4790	7833	807	2423 2429	278 274 274
202	9	201307	10003	9200 201943	202216	202488	2761	200908 3033	300677 3306	200000 3077	3245 201134	372

		;		medy nym		2					ì	
27002	N.	VI CHE.	1	2	3 *	4	5	.6	7	84	10 PB	Di.
.811	100	NA VAN	-	-	25 455A	200004	206475 8178	200745	200014	200000	31	31
75	3	1.13% 1.13%	313 MG	31 (F) 3 (F)	21 (MA)	210006 2400	210000	21[[2]	21 165	21 1414	nilli,	
躩	. 8	78	77.57	9473 9516		9465 221165	9768 221614	231676	- 991 966 961 961	995 995 20195	4	
號	7	- #16 - #16 - #16	SATE SATE SATE	4654 4664 8400	3.66	8765 6348 8913	4015 6600	4974	111	#	4051 (190	
257		7007	200704	2000	201216	201470	9170 531794	201979	20204	1000 1000 1000 1000 1000 1000 1000 100	307193 867/40	24
3			8791 8791	4554 0105	1 20	4011	484 9700	4617 7041	910 110 110 110 110	7,644	4376 1750	
			3400	341048 8884	94167 908	241146	341706	343)44 4625	24338 1 4772	2441 2619	3700	
竖	7	#13 773	919	3 days	6363 8709 261161	0350 0054	97.6 97.6 97.6	9449	7 107	740	2017	
		2003	2000	8883	90.00		4064	1300	4546	470	1661	
2			246514 7249	3100 8100			一新	246718 2116 211611		207196		器
71	8	3451 3451 4618	aut i		1			-	4100	4946 9702	- MIN	
遥	L.	9613 271842	7 404 97 46 27207 4	979906	7875 270913 2678		-	2 Miles	9014 27144	37	27 E	
器	8	4168 6463	4300	4480	4860 7151	機	機	188	8772 8007	300		
22	览	\$18764 \$11038	数数	<b>拉拉</b>	47-169 2017 L	Market Miles	270 de 200 de 20	*****	100 to 1	10 to	TO SERVICE	
45	2	5301 557 7503	2005 2005	8748 6007	ST.		鑑	4456	總	180	100	
误	5	201164 246	200257 247B	200	2007 (S)	8141	20 [G 20 8	10 150 10 150 10 150	00 100 1 3504	201818 484		
24 45 25 25 25 25 25 25 25 25 25 25 25 25 25	8	0005 0005	4607 668.4 807.1	4407 7104	5157	7545 7745	7761	57.87 13/10 10(1) 61	6107 6100 60078	9416 7416		
-	100	1000	201247	301464	001081	301.808	202114	diates	1000	Spirited Spirited		317
21 43 64 85 108 127 148 170 191	į	5351 742	8412 5686 7710	781 784	5006 8137	85	4275 6485 8464	481	6706 6684 ::8981	700	127	
86 108	40	. <b>30</b> 57.73	9545 311966	2177		2000	\$1000g \$112 4020	8776 8100 65 8100 65 8128	\$11118 \$554	811860 8445	9) [66 60 60 60 60 60 60 60 60 60 60 60 60 6	盟
操	77.8	570	5100 8573	6390 8481	6460 8880	0560 8566	7018	25%	7486	773	164	
	9	30146 30146	330354 833436		330769	320077	321184	221.501	321.006	321805	824012	307
888859	210	4483	4488	4494 6745	4500	5106 7155	4310 7340	5514 7568	9121 7767	5050 1972	誤	
61 81		830) 30414	8583 830617	8787 80819	8091 331022	OHEAL	9398 331427	9001 331630	9005	20006 2004	89(2) ] 1206	
1101	3	4454	4655	4656	5057 7000	疆	8447 6456 7450	7600	5800 7858	1000	100 107	
183	3	340446	340645	14000	9054 341089	1415	341435	341633	9640 3:11890	2400		
7.25	, ,		*:	w. c. of the same	a to the second	_	200			-	-	-

22	N.	•	1	9	8	4 5	5	•	7	8,	9,	D.
29.80 T M 13.64	200000000000000000000000000000000000000	848-483 4868 8905 8508-46 21.83 4106 6026 7935	34869) 4540 8540 8600 3576 2576 2517 8117 8120	349817 4786 6744 3694 350696 4408 6406 8316	348014 4061 9000 8000 250500 2761 4066 6500 8506	843918 5178 7136 9058 851098 2064 4676 6790 8006	845400 5874 7830 9778 851216 8147 5066 6061 8066	\$4000 5570 788 \$473 \$61410 \$890 5000 7172 \$076	845000 5700 7730 9000 801400 8418 5418 7818 1805		4 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	
174 19 88 874 111 119	2012245074	9636 861726 8612 5466 7864 9816 871068 2013 4748	361917 8800 8676 7842 9401 871863 8006 4963 6780	362106 362106 3668 5663 7730 9667 371437 3250 6115	360404 362304 4176 6040 7915 9772 871622 8464 6296 7134	360403 362463 4363 6396 8101 9058 371806 3647 5481 7306 9124	800788 800671 4551 4423 8267 370148 1901 8881 5664 7488	800073 6010 6010 8478 870538 2175 4018 5846 7677	261161 260048 4008 6798 8409 870513 2509 4198 6020	861300 846236 5114 6003 6003 6003 6004 4006 6013	261290 260 94 37 37 37 37 37 37 37 37 37 37 37 37 37 3	
167 18 35 88 71 88 106	S-30-00-00-00-00-00-00-00-00-00-00-00-00-	850811 2017 8615 8616 7800 9166 800056	8580 280802 2197 3005 5785 7568 9043 301112 2873	8761 880673 9877 4174 5004 7746 9090 801986	8043 8667 4868 6148 7088 801464	2787 4588 6821 8101 9875	200   43	9487 881296 8007 4901 6077 8456 80228 1908 8751 6501	7858 9666 881476 8877 889405 899405 8169	361666 3406 4940 7084 8611 390683 2546 4101	24 CENT CENT CENT CENT CENT CENT CENT CENT	355556 B
148	200 1 22 2 4 5 4	4458 6190 99740 9974 401401 8131 4894 6540	4657 6674 896114 9847 401678 8602 8005 67710 8410 410108	#902 6648 70020 1746 8464 5176 6081	896461 400192 1917 8636 7051	808634 400365 2080 8907	5886 7071 80660 400586 2861 8078 5686 7601 7601 1601	7945 306061 400711 9438 4140	8026 8676 7419 800154 400668 2606 4890 6020 7731	8850 7802 800828 401056 8777 4403 6190 7901	800601 401228 2040 4663 6670 9764 411451	の記記の
119	1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	9063 411620 8300 414073 6041 8301 9066 421604	410103 1788 8467 415140 6907 8467 420121 1768 8410	6081 8679 410271 1956 3835 416307 6078 8683 420286 1988	8740 410440 2124 8803 415474 7139 8798 420451 2007	7221 8018 410600 2298 8070 415641 7806 8064 420616 2261	2461 4137 415808 7473 9130 420781	7561 9287 410946 2629 4305 415974 7638 9295 420945 2500 4228	9426 411114 2796 4472 416141 7804 9460 421110 2754 4392	411283 9064 4630 416308 7970 9625 421275 2018	8132 4906 416474 8135 9791 421430	160 168 167 167 166 165 165
88 98 115 181 146	567 89 271 23	8246 4862 6511 8135 9753 481364 2000 4500 6163	8410 8045 6674 8297 9014 481525 8180 4729 6322	8074 5208 6896 8430 430075 431685 5290 4388	8737 5871 6009 8621 430236 451846 8430 5048 6640	3901 5554 7161 8788 430308 432007 3610 5207 6799	4065 5607 7324 8044 480550 482167 8770 8067	4328 8980 7486 9106 450730 432328 3090 5536 7116	4502 6023 7648 9268 450861 452488 4090 5665 7275	4555 6186 7811 9439 431042 432649 4349 5844 7438	4718 6340 7973 9001 431308 45000 4460 6004 7792	164 163 163 161 161 160 150 150
80 70 95 III 196 III	4 6 6 7 8 9	7751 9333 440000 2490 4045 5604	7900 9401 441066 2637 4901 8700	6481 8067 9648 441224 2798 4367 5015	8236 9806 441881 2950 4513 6071	8384 9964 441538 8106 4669 6226	8542 440128 1605 8263 4825 6362	8701 440279 1853 8419 4061 6687	8800 440437 2000 8878 5137 6603	0017	7802 9175 440752 2823 8880 5440 7008	1239年33

***	, 5	97.			100	3	1 -	) i	e 2	1 1	
127	e e . O . Est	10	9,1	₹ <b>8</b> 2	1,42	1 6	6.,	172	8,	10 101	<b>A:</b>
70 1 20 2 31 3		學	4948	(語)	47113	4770 4770 41018	4977	44704	1	细	
1 2 1	亞	湿	4987	49711 947 9177	345) 340)	4063 7463 7019			200 200	题	語
107 7	645 6066 7166	9013	8180 8184	- 45	645.4 607.8 1477	123	明時	7(40)		問問	띩
强。	400000	9643 461048	461 195	41348	40140	1649	40300	4947	40007	473	图
15 200	4980	4048	433	40047	400000	401.00	48354	4949	4000		漫
141		1000	77	7319	760	100 100 100 100 100 100 100 100 100 100	772			対	
74 6	4震	45年	40116	\$ 500 PM	4000	911	40704 3121	4780	100	THE STATE OF	
選	3000	- 245 416	400	4450 6197	聖		100	112 112 112 112 112 112 112 112 112 112		器	
41.09	<b>1133</b>		47(1)	<b>阿</b> 豐	41100	9784	The same	470130	1973	47.00E	础
1,3,8			486 486 486 486 486 486 486 486 486 486	4933	401100 2015 3440 4000	1300		411013	1,200		器
2573 2573 2573	100 100 100 100 100 100 100 100 100 100	4442 7 8053	0096	额	600	960	51 da 51 da 9472				
100 7 114 8 136 9	號	7380 8802 400080	40000 40000	7660 6974 667860	40000	7845 490061	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	9007 401941	40000	I	
810	401363	401502			401983 8819	492063	(0000)1 (J07	40041 6767	49040)		
14 1 26 2 41 3	## ##		9 15 15 15 15 15 15 15 15 15 15 15 15 15	4917F9 4977 4973 4973	4711 998 748	600 635	6576	918 0018		温	
E-9888	3311	1 246	110000	7344 8724 8000	800236	7621 8600 500374 1744	9789 9187 800611	9278 400648	9418 800786 3164	40.0000 10.0000 10.00000	題
97 7 110 8 124 9	501050 3427 3791	501196 2064 3027	100 100 100 100 100 100 100 100 100 100	1470 2837 4190	1007 2073 4335	1744 8109 4471	1880 3246 4007	2017 2022 4743	3164 8018 4978	\$291 \$885 \$617	題
-	506150	-	A08491	_	404400	ANARSE		506000 7451	M04284	406579	圖
13 13	7866 9203 610545	7991 9587	9776 8126 9471 510618 2151 8484 4613	5058677 6911 8260 9606	7046 8595 9740	7181 3630 9674	805064 7915 8664 510000	8799 510143	7098 8084 810877	40011 40011 1759	當
8 3 8 5 8 6	43	9587 510679 2017	510618 9151	\$606 510947 2284 3617 4946 6271	9740 511081 2418	511215 2551 8883	8064 510000 1340 2084 4016 5344 6008	510148 1482 2618 4140	810977 1416 2961	17.00 17.00	
107 B	8218 4545 5874 7196	E UVVV	4613 6139 7460	4946 6271	8780 8079 6403	8811 6686 7866	5344	5476 6800	460 460 660 660 660 660 660 660 660 660	豱	題
890	518514	518646		518900	7724 519040	519171	7987 519908	8119 519484	519666	\$1960r	温
11 1 26 2 30 3	881138	#11260 #11260		590221 1530 2885	690663 1661 2006 4266	520484 1792	520615 1922	2083 2083 2086	840676 3188 8466	231007 2314 3616	墨
54 5 56 5	2144 3746 8045	8876 5174	1400 2706 4006 5304 6886	8484	1563	4396	4526	4656 5951 7943	6061	4010	
104 8	(000 (000) (001)	23898 25888 25888	7888 9174	8016 9302	6856 8145 9430	8974 9559	7114 8402 9887 550068	9815	7372 8600 9943	7501 8788 89078	盟
H. P.	asharo	<b>500038</b>	500456	9902 550564	9450 530712	880840	SEPOCE SEPOCE	531006	5943 581273	21241	

3.5	was stall	d figur	althor the	to the King of	ru -	100	91 0	n 1	100	1		c. or 16.	ma 10
and the same	PP	N.	open n.	1 1 NA	2	3	4"	5	63	i y"	8	•	D
1 1 1 1 1 1	13	340	581479 2754 4026	531607 2882 4153	581784 2009 4280	831862 8136 4407	#31990 2264 4534	862117 8391 4661	252345 - 3518 - 4787	882373 8645 4014	552500 8772 5041	863687 8860 5167	路路
の一時の一人	13 38 5 5 7 8 10 11 11 11 11 11 11 11 11 11 11 11 11	5 6	5394 6558 7819 9076	5421 6685 7945 9202	5547 6811 8071 9827	6037 8197 9452	5800 7063 8822 9578	7180	7315 8074 9829	6180 7441 8600 9954	8772 5041 6306 7567 8825 540079	6438 7698 8951 840204	126 126 126 126
Marie of the same	101 118	789	540329 1579 2825	540455 1704 2050	540580 1829 8074	540705 1963 8190	840830 2078 3323	9708 540055 2203 3447	841080 2327 3571	541205 2452 3096	540079 1330 2676 8820	1454 2701 3044	126 126 124
1	278	350	544068 5807 6543 7775	544192 5431 6066 7898	544316 5555 6780 8081	544440 5678 6913 8144	544564 5802 7036 8267	544688 5026 7180	8512 8512	544086 6172 7405 8635	545060 6906 7820 8758	545183 6419 7652 8880	
and the same	3070	4.50	9008 550228 1450	9126 860351 1572	9249 550473 1694 2911 4126	9371 850595 1816	9494 800717 1988	8889 9616 480840 2089 3276	9739 590963	9861 851084 2308 3519	9064 801216 2428 3640	\$00106 1328 2647 2768	1988
1	50	789	8868 5004	2790 4004 8318	4126 5336	3083 4247 5467	4548 5578	8276 4489 8600	2398 4610 5820	4731 8940	0061	0182	100
di Conju	12	8	7507 8709 9007	7927 8820 860026	77.48 89.48 8601.46 13.40	7868 9068 801266	7988 9188 9188 1578	853006 8108 9308 850604 1668 2687	9428 9428	8849 8849 9648 590748	8450 9450 90007 900968 2056	8010 (E 8010 (E 317 (	
## C + # 4 5 h 1 h	35 40 T 55	5 6 7	561101 2393 8481 4666	2012 3600 4784	2531 3718 4008	2680 8837 8021	27/00 2068 5130 6320	2007 4074 10057	1817 8006 4192 8876	8136 4311 5404 6678 7649	24. 25. 26. 26. 26. 26. 26. 26. 26. 26. 26. 26	4548 5730	119 119 118 118
And the Principle of the Paris	95 107	8 870	7026 568902	8066 7144 668819	7803	7879	7407	7614 906788 9050	7735 50000	7849 860028 870198	7967 7967 #60140	8909 8064 560267 870426	118
The species of the	12286803	384	9574 570548 1709 2873 4051 5188	9401 870660 1835 2988	9608 870776 1942 8104	973. 870808 2068 8220	9643 971010 974 9836	<b>初1136</b> 2391 2483	570076 1248 2407 2568 4726 5880	1860	570000 1476 2600 2600 4677 6111	1592 2755 3915	117 116 116
design of the same		5678	4051 5188 6541 7408	4147 5308 6457 7607 8764	5419 6572	4570 8534 6687 7896 8088	4494 8800 8803 7951 9007	4510 87:55 6017 8065	· 8181	494 494 496 7147 896 7147	8410	8072 6236 7377 8535	
White diagnal and a con-	104	9 880	579784 580936	8764 8779308 581,089	77723 8868 680013	MECTION .	580341 1381	9818 880868 1498 2681	9626 050469 1608 2745	190000	9655 550697 1836	0600 080811 1950	114 114 114
1000	BACKE		2063 8196 4331	2177 8812 4444 5574	2201 3426	1267 2404 2539 4570 8780	2618 3633 4788 8012 7037	2651 8765 4506 9044 7140	3745 3579 3000 3137	1742 2008 2009 2009 2009	2772 4106 5035 6062	8085 4218 8346 6475	114 118 118 118
the same of the same	828	5780	6587 7711 8833	6700 7828 8944 890061	5686 6812 7988 9056 590178	8047 9167 800284	7087 8160 9279 600896	7140 8373 9391 89007	7263 8884 9508 800619	7374 8466 9515 890730	7486 8008 9736 690843	7800 8720 9836 890068	113 113 113 113 113
A. Comments	11	<b>390</b>	501065	801176	801587	601.800 251.0 861.8	301510 2621	401621	101733 2043	391848 2954 4061	3064 4171	802066 8176	H
100	1224281	34	8986 4398 5496 6607	2288 8307 4506 5006 6707	8608 4614 8717 6817	4734 5827 6037 8034	8730 4834 5937 7067 8184	2782 2840 4945 7146	#160 #165 6157 7266	5165 6267 7366	95/7 7476	4363 5386 6487 7586 8681	110 110 110 110
	8(878	740	7606 8791 9888 600678	7805 8900 9992 601082	7914 9009 600101 1191	9119 600010 1300	9228 600819 1408	9057 600428 1517	9446 600887 - 1665	9556 600646 1784	9985 9985 99755 1848	9774 600664 1951	109 109 109
		20			66 44		0 6		10				

143-1

1	PP	N.	•	1	9	3	4	5	6	7.	8		D.
一年 一年 一日 日本	111 211 823 43 54 64 75 86 96	400	602060 3144 4226 5306 6381 7455	602169 3253 4384 5413 6489 7562	602277 - 3361 4442 - 5521 6596 - 7669	602386 5469 4560 5628 6704 7777	902494 8577 4658 5736 6811 7884	602603 3686 4766 5844 6019 7991	602711 - 8794 - 4874 - 5951 - 7026 - 8098	602819 8902 4982 6059 7183 8206	602928 4010 5089 6166 7241 8312	608086 4118 5197 6274 7348 8419	108 108 108 108 107
The state of the s	64 75 86 96	6789	8526 9594 610660 - 1723	8633 9701 610767 1826	8740 9808 610873 1936	8847 9914 610979 2042	8954 610021 1086 2148	9061 610128 1192 - 2264	9167 610234 1298 2360	9274 610841 1405 2466	9881 610447 1511 - 2572	9488 610664 1617 2678	107 107 108 108
THE RESERVE THE PARTY OF THE PA	11 21 22 42 53 63 74 86	410 1 2 8 4 5 6 7 8 9	612784 3842 4807 5950 7000 8048 9003 620136 1176 2214	512890 3947 5003 6055 7105 8153 9198 620240 1280 2318	512096 4053 5106 6160 7210 8257 9302 620844 1384 2421	613102 4159 8213 6265 7315 8362 9406 620448 1488 2525	615207 4264 5319 6370 7420 8466 9511 620652 1592 2626	613313 4570 5424 6476 7525 8571 9615 620656 1606 2732	613419 4475 5529 6581 7629 8676 9719 620760 1799 2835	613525 4581 5634 6686 7734 8780 9624 690864 1903	615630 4686 5740 6790 7859 8863 9028 620068 2007 8043	613786 4792 5646 6865 7943 8969 620032 1072 2110 8146	106 106 105 105 106 106 104 104 104
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	10 20 81 41 51 61 71 82 92	420 1 2 3 4 5 6 7 8 9	628° 02 4 138 6340 7366 8389 9410 630428 1444 2457	625353 4385 541,5 6443 7468 8491 9512 630530 1545 2559	623456 4488 5618 6546 7571 8593 9613 630631 1647 2660	628550 4501 5621 6648 7673 8606 9715 630733 1748	623663 4696 5724 6751 7775 8797 9617 630635 1849 2862	623766 4796 5827 6853 7878 8000 9019 630936 1951 2063	623369 4901 5929 6966 7980 9002 630021 1038 2052 3064	625973 6004 6062 7058 8062 9104 630123 1139 2153 8165	624076 5107 6135 7161 8186 9206 630224 1241 2255 3266	634179 8210 6238 7263 8287 9806 630326 1348 2356 8307	108 108
Section 19 and 1	10 20 30 40 50 60 70 80	430123456789	633468 4477 5484 6488 7490 8489 9486 640481 1474 2465	633509 4578 5584 6588 7590 8589 9586 640581 1573 2563	633670 4679 5685 6688 7680 8689 9686 640680 1672 2662	635771 4779 5785 6789 7790 8789 9785 640779 1771 2761	633872 4880 5886 6889 7890 8888 9885 640679 1871 2860	633973 4981 5986 6989 7990 8988 9984 640978 1970 2959	634074 5081 6087 7089 8090 9088 640084 1077 2009 3068	684176 \$183 6187 7189 6190 9138 640183 1177 2138 3156	634976 5283 6867 7390 8390 9267 640285 1376 2267 8265	634376 8583 9385 7800 8080 9367 640363 1376 2366 8564	101 100 100 100 100 100 99
	10 20 29 39 49 59 69 78 88	440 11 22 3 4 5 6 7 8 9	643453 4439 5422 6404 7383 8560 9335 650308 1278 2246	643561 4537 5521 6502 7481 8458 9432 650405 1575 2343	643650 4636 5619 6600 7579 P555 9630 650502 1472 2440	643749 4734 5717 6698 7676 8653 9627 650699 1539 2536	643847 4832 5815 6796 7774 8750 9724 650696 1666 2633	643946 4951 5013 6894 7872 8848 9821 650793 1762 2730	644044 - 5029 - 6011 6092 7969 8945 659919 - 0890 1869 2826	644143 5127 6110 7069 8067 9043 650016 0987 1966 2923	644242 6226 6208 7187 8166 9140 650113 1064 2063 8019	644540 5324 6506 7285 8268 9237 650210 1161 2150 3116	96 96 96 96 96 96 96 96 96 96 96 96 96 9
The second second	10 10 30 38 48 58 67 77 86	450 1 2 3 3 4 5 6 7 8 9	653213 4177 5138 6098 7056 8011 8065 9016 600865 1813	9060 9060	653405 4369 5381 6290 7947 8202 9155 660106 2002	653602 4465 5427 6386 7343 8398 9250 660201 1150 2096	653596 4562 5523 6482 7436 8393 9546 660296 1245 2191	653696 4658 5619 6577 7584 8488 9441 600391 1339 2386	655791 4754 5715 6678 7629 8684 9636 660486 1484 2360	653888 4856 5810 6709 77725 2679 9631 600581 1829 2475	653084 4946 5006 6864 7820 8774 9726 660676 1623	654060 5042 6002 6960 7916 8676 9631 660771 1718 2043	96 96 96 96 96 98 98

1	1		. 5.	8	910	n 2	1	1	17 1		<i>1</i> 5-	F	100
4	PP	N.		1	· <b>2</b>	8.4	4 .	5.	•	17		, 3 <b>9</b> , 8	D.
	S. S.S.	400	000708	000002	063047	008041	068135	663230	003324	063418	008012	000007 4548 5487	94
-	19	\$1 \$62 .88	8701 4642 5581	4796 4796 4675	4880 4880 5760	8983 - 4984 5963	4078 5018 5066	5112 6050	4266 6206 6143	4300 8309 6237	4454 5893 6331	6494 6494	33333
ı	28 47	4	6518 7453	6612	7640	6799 7733	7826	7990 8852	7079	7173 8106	7266 8180	7300 8300 9324	94 93
1	66 66 70	126 117	8386 9817 670246	9410 670389	8572 9508 670481	8665 9596 670524	9689 670617	9782 670710	8945 9875 670802	9088 9967 670895	9181 070000 0988	670188 1080	93 93 93 98 98
-	85	9	11173	1265	1358	* 1451	1543	1636	1728	1821	0988 1913	2005	
1	9	470 11 2	672098 3021 3942 4861	672190 3113	8205 4126	672375 . 8297 . 4218	679467 . 8390 . 4310	672560 3482 4402 5820 6236	672652 - 8574 - 4494 - 5412	8066 4586	872836 8758 4677	872020 8850 4709	92 92 92 92 92 91 91
	18 28 37 46 55 64 74	1.8	5778	4034 4953 5870	5045 5962	5137 6063	5328 - 6145	5820 6236	0328	6419	# 6595 6511		92
	46 55	5 5	7604 7607 8618	7698 8609	6876 7789 8700	7881 8791	7059 7972 8883	7151 8063 8073	7242 8154 9064	7333 8345 9146	7424 8336	7516 8427 9387	91 91 91
2 4 1	74	89	9428 680836	9519 680426	9610 680517	9700 080007	9791 080098	9882 680789	9973 680879	680063 - 0970	9346 680154 1080	000245 1151	91 91
S. 150 S.	200	490	681241	681332	681422	681513 2416	681603	681693	681784	681874	681964 2867	682065	90
7.5	48 97	223	2145 3047 3947 4845	3137 4037	2326 3227 4117	3317 4217	681603 2806 3407 4307 5204 6180	2595 3497 4396	681784 2686 8587 4496 5383	2177 3677 4576	8767 4066	3887 4746	90
	ならいの	4	8 676 2	4037 4986 5881	5025 5021 6815	5114 6010	5204 6190	4396 5394 6189	004.5	6368	8767 4066 5663 6458	5652 6547 7440	90 80
. 4.	63 72	7	7529 8420	7618 8599 9398	7707 8598	7796 8687	7896 8776	7083 7975 8865	7173 8064 8063 9841	7261 8153 9042	7351 8242 9181	8331 9220	<u>998888888</u>
-	81	#9	/ 9500		9486	9675	9664	9753	-	. 9930	000019	690107	
- 0.0	s 9 18	490	#1081 1965	690285 1170 2053	690373 1258 2142	090462 1347 2230	690550 - 1435 - 2318	690639 1524 2406	690728 1612 2494	090816 1700 2583 3463	090005 1789 2671	1877 2750	89 88 88 88 88 87 87 87 87
100	1385488F	#3 #4	2847 8727 4805	2935 3815	3023 3903	7 8111	3199 4078	3287 4166	2494 3375 4254 5131	8463 4842 5219	2671 3551 4430 5307	4517	88 88
A	168	5 6 7	5482 6356	4003 5509 6444	4781 5657 6531	3991 4868 5744 6618	4966 5832 6706	5044 5919 6793	6007 6880	6094 6968 7839	2085	5394 6269 7142	87 87
1. 100	7078	8	7229 8101	7317 8188	7404 8275	7491 8362	7578 8449	7665 8535	7752 8622	7839 8709	7926 8796	8014 8883	87 87
47.0	4.0	200 21	698970 9838	699057	699144 7000H	609281 700098	699317 700184	699404 700271	609491 700358	699578	699664 700531	690751 700617	87
-		3	700704	700790 1664 2517	1741	0963 1827 2689 3549	1050	- 1104	# 1000	700444 1309 2172	1305	1482	86 86
1	48	.55	2431 3291 4151	2517 3877 4286	2003 8463 4322		2775 3635 4494	1999 2861 3721	2086 2947 3807 4665 5522	2172 5033 3893 4751	3119 3979 4837	3205 4065 4922	86 86
	1888	197	5008 5064	5094	6035	5265 6120	5350 6206	4579 5436 6291	6376	6462	5008 6547	5778 6632	87 86 86 86 86 86 86 86 85
2	/77	510	707570	6803 707655	6888	707826	7059	707996	708081	708166	708261	7485	
	8 17 25	1 2	8491 9270	8506 9855	707740 18591 9440	8676 9824	8761 9609 710456	8846	0091	-0015	9100	: 010K	**************************************
the spirit	25 34 49	345	710117	710202 1048 1802	710287	710371 1917 2000 2902	. 1301	710640 # 1385	9779 710625 1470 2313	710710 1564 2307	710794 1659 2461	0879 1798	85 84
2 1	1886	6	1897 9680 8491	2734	1976 2818 8669	2902 8742 4551	2144 2986 3826	8070 8010 4749 5586	3304	2507 2008 4078	4167	1246	84 84
	75	8	4830 6167	8675 6414 5261	44.97 6335	4461 5418	4665 5502	4749 5596	3/8994 6.4883 6.5609	4016	0000 0000	300€ 3000	84 84
	- 3	7 3		1		4		2				1	1

8 15 23 50 38 46 53	8 15 23 31 39 46 54 63 69	8 16 23 31 39 47 55 62 70	8 16 X 53 4 8 56 4 72	8 16 24 41 49 57 63 73	8 17 25 83 41 50 58 66 75	P
570 1	560 1 2 3 4 5 6 7 8	550 1 2 3 4 5 6 7 8 9	540138456789	530 123 34 56 789	530123456789	N.
755875 6636 7396 8155	748188 8963 9736 750508 1279 2048 2816 3583 4348 5112	740363 1152 1939 2725 3510 4293 5075 5855 6634 7412	782394 3197 3999 48900 5699 6397 7198 7987 8781 9572	724276 5095 5912 6727 7541 8354 9165 9974 730782 1589	716008 6886 7671 8502 9531 720159 0986 1811 2634 8466	1 0
755951 6712 7472 8230 8988 9743 760498 1251		740442 1230 2018 2804 3588 4371 5153 5933 6712 7489	732474 8278 4079 4880 5679 6476 7272 8067 8860 9651	724358 5176 5096 6809 7623 8435 9246 730055 0863 1669	716067 0021 7754 8585 9414 720242 1068 1803 2716 3538	11
7548	748343 9118 9891 750663 1433 2202 2970 3736 4501 5265	740521 1309 2096 2882 3667 4449 5231 6011 6790 7567	732555 3358 4160 4960 5750 6566 7352 8146 8939 9731	724440 5258 6075 6890 7704 8516 9327 780136 0944 1750	716170 7004 7887 8666 9497 720526 1151 1975 2798 3620	72
756103 6964 7624 8382	748421 9196 9968 750740 1510 2279 3047 3813 4578 5341	740600 1388 2175 2961 8745 4528 5309 6089 6868 7645	732638 8438 4240 5040 5888 6635 7431 8225 9018 9810	724522 5340 6156 6972 7785 8597 9408 730217 1024 1830	716254 7088 7920 8751 9580 730407 1233 2068 2881 8702	03/
756180 6940 7700 8458 9214	748498 9272 750045 0817 1587 2356 3123 3889 4654 5417	740678 1467 2254 3039 3823 4606 5387 6167 6945 7722	732715 3518 4320 5120 5918 6715 7511 8305 9097 9889	724604 5422 6238 7053 7866 8678 9489 730298 1105 1911	716357 7171 8003 8654 9665 720490 1316 2140 2963 8784	4:
756256 7016 7775 8533 9290 760045	748576 9350 750123 0894 1664 2433 3200 3966 4730 5494	740757 1546 2382 3118 3902 4684 5465 6245 7023 7800	782796 8598 4400 5200 5998 6795 7590 8384 9177 9968	724685 5503 6320 7134 7948 8759 9570 730378 1186 1991	716431 7254 8086 8017 9745 720573 1396 2222 3045 3866	5
756332 7092 7851 8609 9366 760121	748653 9427 750200 0971 1741 2509 3277 4042 4807 5670	740836 1624 2411 8196 3980 4762 5543 6823 7101 7878	782876 3679 4480 5279 6078 6874 7670 8463 9256 740047	724767 5585 6401 7216 8029 8841 9651 730450 1266 2072	716504 7388 8169 9000 9828 720655 1481 2306 3127 3946	18
756408 7168 7927 8685 9441 760196 0930		740015 1703 2459 3275 4058 4840 5621 6401 7179 7955	732956 3759 4560 5359 6187 6954 7749 8543 9336 740126	724840 5667 6488 7297 8110 8922 9732 730540 1347 2152	716588 7421 8255 9068 9911 720738 1563 2587 8209 4030	87°
756484 7244 8003 8761 9517	4960	740994 1788 2568 8358 4136 4019 5609 6479 7256 8053	733037 3839 4640 5439 6237 7034 7829 9622 9414 740205	724981 5748 6564 7379 8191 9003 9813 730621 1428 2233	710671 7504 8536 9105 9994 720821 1646 2460 8201 4112	8,1
756560 7320 8079 8836 9592	748685 9650 780431 1202 1972 2740 8806 4272 5036 5790	741073 1800 2647 8481 421.5 4907 5777 6556 7834 8110	788117 5919 4790 5619 6317 7113 7906 8701 9493 740284	728018 5890 6646 7460 8273 9084 9898 730702 1508 2813	716784 7567 8419 9846 720077 (968 1788 2552 8374 4194	
76 76 76 76	TETETETE TO	79 79 78 78 78 78 78 78	80 80 80 80 80 80 79 79 79 79	82 82 83 81 81 81 81 81	***************************************	D.
-	the name of Street, or other Designation of the last o	ı				-

ń		17	"			1			¥ .	1 . 3		No. 1		s
	PP	N.	€ • #	.1	2,	3	4	5	. 6	7	8	9	D.	
7.	7 15 22 80 87 44 54 56 67	480 1 2 3 4 5 0 7 8 9	763428 4176 4923 4609 6413 7156 7898 8638 9577 770115	763608 4251 4996 5743 6487 7230 7972 8712 9451 770189	763578 4326 5072 5818 6562 7304 8046 8786 9525 770263	763653 4400 5147 5892 6636 7379 8120 8800 9699 770336	763727 4475 5221 5966 6710 7453 8194 8934 9673 770410	763803 4550 5296 6041 6785 7527 8268 9008 9746 770484	763877 4624 5370 6115 6859 7001 8342 9082 9620 770657	763952 4699 5445 6190 6933 7675 8416 9156 9894 770631	764027 4774 5520 6264 7007 7749 8490 9230 9968 770705	764101 4848 5504 6338 7062 7823 8564 9303 770042 0778	75 75 76 74 74 74 74 74 74	* 2
	7 15 22 29 37 44 51 58 66	500 1234 56789	770862 1587 2322 3056 3786 4517 5246 6974 6701 7427	770926 1561 2395 3128 3860 4590 5319 6047 6774 7499	770999 1734 2468 8201 8933 4663 5392 6120 6846 7572	771073 1808 2542 8274 4006 4736 5465 6193 6919 7644	771146 1881 2615 3348 4079 4809 5538 6265 6992 7717	771220 1955 2688 3421 4152 4882 5610 6338 7064 7789	771293 2028 2762 3494 4225 4955 5683 6411 7137 7862	771367 2102 2835 3567 4298 5028 5756 6483 7209 7934	771440 2175 2908 3640 4371 5100 5829 6586 7282 8006	771514 2248 2961 3713 4444 5173 5902 6629 7854 8079	74 73 73 73 73 73 73 73 73 73	and the same of th
	7-14-22-29-86-43-50-58-65	60128466789	778151 8874 9596 780317 1037 1786 2473 5189 5904 4617	778224 8947 9669 780389 1109 1827 2544 5260 3975 4689	778296 9019 9741 780461 1181 1899 2616 3332 4046 4760	778368 9091 9613 780533 1263 1971 2688 3403 4118 4831	778441 9163 9885 780605 1324 2042 2759 3475 4189 4902	778513 9236 9967 780677 1396 2114 2831 3646 4261 4974	778585 9308 780029 0749 1468 2186 2902 3618 4332 5045	778658 9380 780101 0821 1540 2258 2974 3689 4403 5116	778730 9462 780173 0803 1612 2329 3046 3761 4475 5187	778802 9624 780245 0965 1684 2401 3117 3832 4546 5259	72 72 72 72 72 72 72 71 71 71	
45. 15.1	7 14 21 28 36 45 50 57 64	610 122 34 56 789	785330 6041 6751 7460 8168 8875 9581 790285 0988 1601	785401 6112 6822 7531 8239 8946 9651 790356 1059 1761	785472 6183 6893 7602 8310 9016 9722 790426 1129 1831	785543 6254 6964 7673 8381 9087 9792 790496 1190	785615 6325 7035 7744 8451 9157 9863 790567 1269 1971	785686 6396 7106 7815 8522 9228 9933 790637 1340 2041	786757 6467 7177 7885 8593 9299 790004 0707 1410 2111	785828 6558 7248 7956 8663 9369 790074 0778 1480 2181	785899 6609 7319 8027 8734 9440 790144 0848 1550 2252	785970 6680 7590 8098 8804 9510 790215 0918 1620 2522	กรากการกรร	
	7 14 21 28 36 42 49 56 63	620 1 2 3 4 5 6 7 8 9	792392 3092 8790 4488 5185 5880 6574 7268 7960 8651	792462 3162 3860 4558 5254 5949 6644 7337 8029 8720	792532 3231 3930 4627 5324 6019 6713 7406 8098 8789	792602 \$301 4000 4697 5393 6088 6782 7475 8167 8858	792672 8371 4070 4767 5463 6158 6852 7545 8236 8927	792742 3441 4159 4836 5532 6227 6921 7614 8305 8996	792612 5511 4209 4906 5602 6297 6990 7683 8374 9066	792882 3581 4279 4976 5672 6366 7060 7752 8443 9134	792952 3651 4349 5046 5741 6436 7129 7821 8513 9203	799022 3721 4418 5115 5811 6505 7198 7890 8582 9272	70 70 70 70 70 89 69 69 69	
1	7 14 21 28 35 41 48 55	630 1 2 3 4 5 6 7 8 9	799341 800029 0717 1404 2089 2774 8457 4139 4821 5501	799409 800098 0786 1472 2158 2842 3525 4208 4889 5569	799478 800167 0854 1541 2226 2910 3594 4276 4967 5637	799547 800236 0923 1609 2295 2979 3662 4344 5025 5705	799616 800305 0992 1678 2363 8047 3730 4412 5093 5778	799685 800373 1061 1747 2432 3116 3798 4480 5161 5841	799754 800442 1129 1815 2500 3184 3867 4548 5229 5008	799623 800511 1198 1884 2568 3252 3935 4616 5297 5976	799892 800580 1266 1952 2657 3321 4003 4645 5365 6044	799961 800648 1336 2021 2706 3389 4071 4753 5433 6112	69 69 69 69 69 68 68 68 68 68	

_		11						,,	Label .	10	1- 15	1.7
PP	N.	7	1	2	***** <b>3</b>	4	5	6 1 1 and	. <b>7</b>	. 8		S.A.
~32734448	9-9-9-8-8-8-8-8-8-8-8-8-8-8-8-8-8-8-8-8	806180 - 9858 - 7885 - 8211 - 8586 - 9560 - 810283 - 0904 - 1578 - 2246	806948 6925 7606 8979 8968 9637 810900 0971 1642 2813	806316 6004 7670 8846 9021 9004 \$10867 1700	806384 7061 77788 8414 9068 9762 810434 1106 1776 2446	806451 7129 7806 8481 9156 9829 810501 1178 1845 2513	806519 7197 7873 8640 9223 9896 810569 1240 1910	806887 7364 7941 8616 9380 9064 810686 1807 1977 2646	800655 7862 8006 8664 9686 810061 0708 1574 2044 2718	806728 7400 8076 8781 9428 81006 0770 1441 2111 2780	806790 7467 8143 8818 9462 810165 0857 1508 2178	***********
7 13 20 36 35 40 65 50 50 50 50 50 50 50 50 50 50 50 50 50	60188486789	813018 8581 4248 4013 5578 6241 6004 7565 8226 8886	812060 2648 4314 4060 5644 6306 6770 7631 8292 8061	813047 8714 4381 8046 8711 6374 7056 7696 8358 9017	818114 8781 4447 8113 8777 6440 7102 7764 8424 9068	818181 8848 4514 5179 8843 6506 7109 7850 8490 9149	813347 8014 4061 5346 5010 6873 7236 7896 8656 9215	813314 8061 4647 8812 8076 6689 7301 7962 8682 9861	813361 4048 4714 6378 6042 6705 7367 5008 8608 9546	813448 4114 4780 5445 6100 6771 7433 8004 8784 9413	813614 4181 4947 5611 6175 6088 7409 5160 9678	888888888
7 18 20 26 83 59 46 52 - 50	60123456789	819644 820201 0658 1514 3168 2822 3474 4126 4776 6426	819610 820267 0924 1579 2233 2887 8530 4191 4841 5401	819676 820533 0069 1645 2299 2052 8605 4256 4006 5656	819741 880599 1056 1710 2564 8018 8070 4521 4971 5681	819607 820464 1120 1776 2430 8063 8736 4366 8066	819873 890630 1186 1841 2496 3148 5800 4461 5101 6761	819999 890595 1261 1906 2560 8213 8965 4516 5166 8615	980004 0661 1817 1972 2636 8279 8950 4561 6231 6880	50 AS	800186 0792 1448 2108 2756 8409 4061 4711 5361 6010	66 66 65 65 65 65 65 65
6 13 19 26 32 38 45 51 58	670128466789	826075 6723 7369 8015 8660 9504 9947 830589 1230 1870	826140 6787 7434 8080 8724 9568 830011 0653 1294 1934	826204 6852 7499 8144 8789 9452 880075 0717 1558 1998	826269 6917 7568 8909 8853 9497 830159 0781 1422 2063	6263.74 6061 7628 8273 8918 9561 830204 0845 1486 2126	7046 7092 8338 8962 9635 830268 0909 1850 2189	826464 7111 7757 8402 9046 9690 830533 0973 1614 2253	836638 7176 7831 8467 9111 9754 830596 1037 1678 2317	896896 7240 7886 8631 9176 9618 850400 1102 1742 2381	826658 7305 7961 8696 9859 9862 830626 1166 1806 2445	65 65 65 65 65 65 65 65 65 65 65 65 65 6
13 19 25 52 53 54 50 57	680123456789	832509 \$147 5784 4421 5056 5691 6324 6957 7588 8219	832573 8211 8848 4484 5120 5754 6387 7020 7652 8282	832637 \$276 \$912 4548 5183 5817 6451 7088 7715 8345	832700 8338 3975 4611 5247 5881 6514 7146 7778 8408	832764 3403 4030 4676 5310 5844 6577 7810 7841 8471	832828 \$466 4108 4739 5373 6007 6641 7273 7904 8534	832892 8530 4166 4802 5457 6071 6704 7839 7997 8597	832956 8693 4230 4866 5500 6134 6767 7399 8030 8660		833063 8721 4367 4093 5627 6261 6894 7525 8156 8786	2222222
6 13 19 25 82 33 44 50 57	600 12 34 56 78 9	838849 9478 840106 0733 1359 1985 2609 8233 8855 4477	838912 9541 840169 0796 1422 2047 2672 3295 8918 4639	858975 9604 840232 0859 1485 2110 2754 3357 5980 4001	839038 9667 840294 0921 1547 2172 2796 3420 4042 4664	869101 9729 840357 0984 1610 2235 2859 3482 4104 4726	839164 9792 840420 1046 1572 2297 2921 8544 4106 4788	839227 9656 840482 1109 1735 2560 2963 3606 4229 4860	839289 9013 840545 1172 1797 2422 3046 3669 4291 4912	9981 840608	839415 840043 0671 1297 1922 2547 8170 5793 4415 5036	68 63 63 63 63 63 63 63 63 63 63 63 63 63

	500 mg	Sugar -	edter or	The .	1.7m is	- 0.5%	design	and O the	- de figure	-2 - 24	- Garin	the graphed	A STATE OF THE PARTY OF
1	1	×.	0 1	1 near au	<b>9</b>	8	27 1 1 4	. S	en f h S ugh de <sub>ga</sub> tro	7	<b>S</b>		5.
-	613	700	645(BB) 5718 9537	845160 5730 9600	845.202 5848 6461	840884 5004 6523	845346 5066 5665	845406 6026 6646	945470 9090 9705	845539 6151 6770	845004 6018 6006	845e56 6975 6844	8888
1	6428	1001	7578 8189 8806 9419	7684 6861 8868 9481	7606 8819 8088 9649 850156	7758 8374 8060 9004	7819 9435 9051 9665	7881 8407 9118 9786	1945 8580 9174 9788	8004 8020 9205 9840	8086 8239 9307 9011	88.36 87.48 99.48 99.48 99.72	61
	80 56	9	850083 0646	850095 0707	850156 0700	850917 0630	\$50979 , 0891	850340 .,0952	850401 1014	1075	1136	1197	22
1	61184	710	861258 1870 3480	251320 1981 2541 8150	851881 1992 2008 8211	851449 2053 2063	851506 2114 2794 8838	951564 2175 2785	\$51005 2298 2046 3445	851086 2397 2907 8516	2358 7968	951 990 9410 3480	61 61
-	17 28	Section in the section is	3608 4306	8780 4967	8820	8973 8881 4488 5095	8041	400a 4610 5316	4068	4194	200 4700	4945	61 61 61
	88690	57.800 9.000	5519 6134 6730	5580 6185 6780	5064 5640 6345 6850	8701 6306 6010	5186 5761 6366 6970	5888 6987 7871	5877 5889 6487 7091	4731 5887 5943 6548 7152	6008 6608 7212	6064 6068 7373	90 91 91 91
I	6	720	857732 7985	857308 7995	857458	857518   8116	857574 8176	857684 8236 8088	107604 8397	857765	867818 8417 9018	837675 ; 8477	20
21	出版	204	8587 9138 9780	8807 9198 9790	9258 9258 9880	9318 9318 9918	9379 9379 9978	9430	8998 9499 80008	9559 800148	9619 800218	9078 9079 900878	888
9	ななないのでは	5 67	800558 0987	800398 0998	300458 1056	360518 1116 1714	860578 1176 1773	0687 1886 1888	0007 1995 1898	0787 1855 1969	0817 1415 2013	0677 1475 9072	888888888
	48	89	2131 2728	2101 2787	2261 2847	2310 2006	2370	9430 8025	2480 8065	2540 3144	2006 3204	2063	80
	6	780	963943 8917 4511	863382 8077 4570 5163	86344 <u>9</u> 4096 4630	863501 4096 4689	200561 4155 4749	863690 : 4214	805380 4374 6967	863739 4383 4986 5519 6110	863790 4302	903008 4452 5045	888
I	1886	34	5104 5696 6287	5163 5755	5293 5814	2000	4748 5341 5988 6594	4808 5400 8008	4967 5450 6051	5519 6110	5578 6169 6760 7380	5637	888
	SACRBREGG	5678	6878 7467 8056	7698 8116	6406 6006 7585	5874 6468 7085 7044	7114	7178 7768 8380	789 781 889	7301 7880	73.60 79.80	7400 7400 7908	59 59 59 59 59
-	25	9	8644	8703	8174 8762	8651	8879	8988	8007	8468 9056	9114	9178	-
١	6	740	961233 9818 870404 0060	9677 870463	9085 870521	81.0(61.8) 800 % 91.0(61.8)	70058 70058 1458	800695 670111 0006	869684 870170 0785 1889	860649 870298 - 0818	870701 870267 0873	800700 870846 10000 1515	59 58 58 58 58 58 58 58 58
	1782 9	845	1573 1573 2156	1047 1031 2215	1106 1600 2278	1748 1748 2881	1886 1886 2880	1961 1865 2448 8080	1389 1923 2506	0618 1806 1981 2564 8146	2040	9,515 2008 2081	58 86 58
	ESPENDEN	5789	2739 8321 8902	3797 3379 3960	2855 3437 4018	2918 8495 4076	2072 3553 4134	8080 8611 4198	6660	8146 8797 4306	8904 8785	8368 8844 4434 5603	58 58
1	27	760	8909 4482	875119	4508 875177	975995	4714	4772	4830	4888 875466	4366 4045 875524	PRIKOD	.3
	13	4	575061 5640 6318 6795	5698 6376 6853	5756 6333 6010	5813	5871 6449 7026 7002	8029	5087 8564 7141	6045 6622 7190	6102 6650 7256	6180 6787 7314	56 58 58 58 57 57 57
4-	22 29	Series and Series	7371 7947	7429 8004	7487 8062	STIM	8177	6454	7717	7774 8849	7852	7000	56 57
-	111111111111111111111111111111111111111	07-8	9096 9096	8579 9153 9736	9211 9784	9268 9841	9525 9608	9565 9056	9440 940018	9497 880070	9665 880127	9616 860186	87 87
2	92	* 8	880843	880299	000000	880413	050471	680538	0585	0642	0600	0756	107

A 124	a stronger		A . A . A . A . A . A .	y B militage	1	en to make a	same or a	7	3088	Ma	40-517	E . 81
PP	N.	7 P	1 1 m	14 <b>2</b> 1200	3	A see	5 A	<b>6</b>	7	8,	O.	D.
11	750	1506 1908	880871 1442 2013	1400 2000	1866 1866 2136	1613 2188	881000 1670 2240	861186 1727 2307	961318 1784 2354	1841 1841 3411	061986 1806 9468	10 mm
2882	1999 C	1 CONT.	8180 8718	8775 4342	8364 8983	3001 3001 3000 4465		3434 4002 4560	8401 4060	35.66 411.6		<b>新教育者在18</b>
46 51	. 0	5361 5096	5418 5088		6096	8587		. 0000	5757 6321	6219	5470 6484	-
0 11 17	1200	7064 7617 8179	7111 7674 8236	7167	7233 7780 8348	7280 7842 8404	7836 7898 8460	7892 7955 8516	7440 8011 8078	7505 8007 8020	7001 8128 8665	56 56
28.34	10 T	890421	9358 9918 990477	0414	00000		9588 890141 0700	9638 890197 0756	9604	9780 90809 0088	9606 800365 0024	56 56 56 56 56 56 56 56 56
0.0	780	90900K	892150	802206	-	900917	1816 892578	1872		1968	2080 802505	-
11 17		8207 8762 4316	8617	8318 8878 4437	8378 8928 4489	3420 3084 4588	3464 4039 4598	8540 4094 4648	3040 3005 4180 4704	8651 4205 4759	8706 4261 4814	56 55 55
1884	5678	5428 5075 6526	6581	6636	5036 5588 6140 6692	5644 6195 6747	5146 5699 6261 6802	5754 5754 6306 6867	5257 5800 6361 6913	5313 5864 6416 6967		56 56 56 55 55 55 55 55
- F	790	807627	7138	4.00	-	7207	-	-	808012	808067	808192	-
1172	-	97% 9278 9821	9528 9528 9675			9493 900039 00596	9547 90004	60001 AD	900208	9711	900312	55 55 55
8844	\$7.80 \$400000000000000000000000000000000000	0918	1518	1028 1567 2112	1077 1623 2160	2221		1240 1786 2329	1395 1840 2384	1840 1804 2438	1404 1948 2492	56 55 55 55 55 55 55 54 54
. 7.	800	903000	908144	908199	910383 8799	908307		18.00	. 12	903524	908478	4
15831	1040	4716 5256 5796	4770 5610 5850	4894 5964 5904	4878 5418 5968	5472	4400	5040 5530 6119	5004 5634 6178	5148 5688	. 5742	54 54 54 54 54 54 54 54
844	0789	6874 7411 7949	0017 7465 8002	6981 7519 8056	7035 7573 8110	7069 7626 8168	7143 7680 8217	7196 7784 8270	7250 7787 8824	7804 7841 8378	7858 7895 8481	54 54 54
5 11	810 1	908485 9021 9556	908539 9074 9610	908592 9128 9663	906646 9181 9716	908609 9285 9770	906753 9289 9888	908807 9342 9877	908860 9896 9980	908914 9440 9984		54 54 53
16 21 27	848	910091 0624 1159	1211	: U/31	910251 0784 1317	910304 0838 1371 1908	910358 0891 1424	910411 0944 1477 2009	910464 0998 1580	910518 1051 1584 2116	0571 1104 1687	54 53 53 58 58 58 58
87 42 48	7 8 9	2222 2758 3284	2375 2806 8537	2328 2859 8590	2581 2013 3443	2486 2066 3406	2488 8019 8549	2541 8073 3602	2594 8125 8655	2647 8178 8708	2700 3431 3761,	153 155 155
	611177288 844 40 61117288 848 44 40 6111188 888 448 49 6111188 888 489 6111188 888 489 6111188 888 888 489 6111188 888 888 888 888 888 888 888 888	700 1117 22 45 50 770 1117 22	750 800514 11 1 1365 11 1 2 1365 11 2 1365 28 4 3005 29 6 5 3651 61 9 5006 6 1 7 675 6 1 770 885491 6 1 7 7054 11 2 8 8179 28 4 8 9853 39 7 800491 11 2 8397 17 3 8179 28 4 6 9853 39 7 800491 11 3 3397 17 3 8762 28 4 8 6525 44 8 6525 44 8 6525 44 8 6525 44 8 6525 44 8 6525 44 8 6525 44 8 6525 44 8 6525 44 8 6525 44 8 6525 44 8 6525 44 8 6525 44 8 6525 44 8 6525 44 8 6525 44 8 6525 5 1 8176 11 2 9777 770 877627 5 1 8176 11 2 9775 5 1 8176 11 2 9775 5 1 8176 11 2 9775 5 1 8176 11 2 9775 5 1 8176 11 3 9775 5 1 8176 11 3 9775 5 1 8176 11 3 9775 5 1 8176 11 3 9775 5 1 8176 11 3 9775 5 1 8176 11 3 9775 5 1 8176 11 3 9775 5 1 8176 11 3 9775 5 1 8176 11 3 9775 5 1 8176 11 3 9775 5 1 9775 5 1 9775 5 1 9775 5 1 9775 5 1 9775 5 1 9775 5 1 9775 5 1 9775 5 1 9775 5 1 9775 5 1 9775 5 1 9775 5 1 9775 5 1 9776 5 1 9776 5 1 9776 5 1 9976	750 S00614 S00671 1 1865 1448 11 1 1865 1448 28 1 1865 2011 17 3 1865 2011 17 3 1865 2011 17 3 1865 2011 17 3 1865 2011 28 4 3008 31800 29 6 3661 5718 40 7 4785 4812 44 6 4329 4326 45 8 5821 5418 51 9 6666 5088 770 806461 806647 11 7064 7111 11 7 7074 17 3 8179 8286 28 4 8741 77074 17 3 8179 8286 34 6 9663 9012 28 4 8741 77074 45 8 9802 9386 36 7 89042 80180 47 18 18 18 18 18 18 18 18 18 18 18 18 18	Top   Second   Second   Second   1   1366   1442   1460   1471	760   \$60.614   \$60.621   \$60.600   \$1.546   \$1.448   \$1.460   \$1.546   \$1.550   \$	780   SOUGLE   SOUG	780   800914   800871   80088   801088   851088   851088   851088   851088   851088   851088   851088   851088   851088   851088   851088   851088   851088   851088   851088   851088   85248   85248   85248   85248   85258   85188   85778   85283   85688   85188   85778   85283   85688   85188   85778   85283   85688   856	700   \$60.614   \$60.671   \$60.680   \$61.048   \$61.048   \$61.048   \$71.081   \$71.072   \$71.081   \$71.082	TRO	750   60-614   50-671   50-68-6   60-08-6   501-68   50-108-6   50-118-6	780   68051   68051   58058   68068   681048   681068   681186   681318

PP	r d.	. 62	1,	9	8	id.	. 6	6	7	8.	•	D.
5 11 16 21 22 25 45	80198488789	918614 4548 4672 6400 5007 6484 6060 7506 8080 8585	913067 4506 4026 5453 5060 6507 7063 7586 8063 8007	91,8000 44.40 4077 5506 6088 6560 7066 7611 8186 8680	913973 4593 8030 5036 6036 6612 7136 7663 8186 8712	914026 4555 5083 5611 6138 6664 7190 7716 8240 8764	914079 4608 51,84 6191 6717 7348 7766 8208 8616	914188 4600 5180 5716 6248 6770 7995 7890 8345 8869	914184 4713 5241 5760 6898 7348 7873 8307 8021	91 4987 47 68 8504 6549 6575 7 7025 8450 8978	914990 4619 5347 5576 6401 6007 7468 7978 8608 9026	*83 85 85 85 85 85 85 85 85
5 10 16 21 26 81 81	800	919078 9601 920123 0645 1106 2906 9726 8344 8762	919180 9688 920176 0607 1218 1738 2256 2777 3296 4 3814	919188 9706 920228 0749 1270 1790 2310 2620 3348 3866	919236 9766 920290 0801 1822 1842 2563 2881 8399	019287 9610 920332 0668 1874 1804 3414 2933 3451 3969	919340 9668 920394 0906 1426 1946 2466 2986 3508 4021	919392 9014 920436 0958 1478 1998 2518 8037 8666 4072	919444 9067 920489 1010 1530 2050 2570 8069 8007	919406 989019 0541 1062 1582 2102 2622 8140 8656 4176	919549 989071 0596 1114 1654 2154 2174 2074 2174 2172 2710	\$5.00 mm
5 10 16 20 20 81 26 41	840	924279 4796 5312 5828 6343 6867 7870 7888 8506 8506	924381 4848 5804 5879 6394 6908 7433 7936 8447 8959	924383 4809 5415 6445 6050 7478 7796 8408 9010	924434 4051 5467 5982 6497 7011 7524 8037 8549	924486 5003 5518 6054 6548 7062 7576 8088 8601 9113	924538 5064 5570 6086 6000 7114 7627 8140 8652 9163	924589 5106 5621 6157 6651 7166 7678 8191 8703	924641 5157 5678 6188 6702 7216 7780 8243 8754	924698 5809 5726 6240 6754 7781 8293 8805 9317	994744 6961 5776 6391 6806 7319 7832 8346 8867 9368	53 54 51 51 51 51 51 51
5 10 15 20 20 31 36 41 46	8	929419 9650 980440 9949 1458 1966 2474 2981 4 3487 2993	929470 9981 980401 1000 1809 9017 2524 3031 3538 4044	929621 950052 0643 1061 1800 2068 2576 5082 3589	929872 950063 0892 1103 1610 2118 2626 8133 8639	989623 980134 0648 1163 1661 2169 2677 3183 3690 4196	929674 980186 0694 1204 1712 2230 2727 8234 8740	929725 980236 0746 1254 1763 2271 9778 8286 3791 4296	929776 930287 0796 1305 1814 2322 2829 3335 8841 4347	929827 930338 0847 1356 1865 2372 2879 8386 8892 4397	990879 950589 0608 1407 1915 2423 2030 8487 5043	51 51 51 51 51 51 51 51 51 51
5 10 15 20 25 30 35 40 45	800 1 2 8 4 5 6 7 8	934498 5003 5507 6011 6514 7016 7518 8019 8520 9020	6564 7066 7568	934500 5104 5608 6111 6614 7117 7618 8119 8620 9120	934650 5154 5656 6162 6665 7167 7668 8169 8670 9170		934751 6255 5789 6262 6765 7267 7769 8269 8770 9270	934801 65306 5809 6313 6816 7317 7819 8320 8820 9320	934852 5306 5800 6363 6865 7367 7800 8870 9500	934902 5406 5910 6413 6916 7418 7919 8420 8020 9419	934958 5457 5000 6463 6066 7468 7969 8470 9469	50 50 50 50 50 50 50 50 50
8 10 15 20 25 80 85 40 45	870	939519 94001H 0516 1014 1511 2008 2504 3000 8495 8969	940068 0566 1064 1561 2058	939619 940118 0616 11114 1611 2107 2003 3090 3593 4088	940168 0666 1163 1660	939719 940218 0716 1213 1710 2207 2702 3198 3692 4186	939769 940267 0765 1263 1760 2256 2752 3247 3742 4236	939819 940317 0815 1315 1809 2306 2801 8207 8792 4285	939609 940367 0865 1362 1859 2355 2851 8346 8841 4336	939918 940417 0915 1412 1909 2405 2901 8396 3890 4384	930968 940467 0064 1462 1958 2455 2950 8446 5039 4433	50 50 50 50 50 50 50 49 49

は以中は響き

PP	N.	i. O in	1,	49		N.4	26	z G	7	8	20	D.
5 10 15 20 25 29 84 89 44		944468 4976 5469 8061 6452 6043 7434 7934 8418 8002	944888 5025 5518 6010 6501 6002 7488 7978 8463 8961	944561 8074 5067 6060 6861 7041 7583 8022 8511 8099	944681 5134 5616 6106 6600 7090 7581 8070 8560 9048	944680 5173 5666 6157 6640 7140 7630 8119 8600 9097	944739 5222 8716 6207 6698 7189 7679 8168 8657 9146	944779 8372 8764 9764 7728 7728 8217 8706 9195	944986 5321 5613 6805 6796 7787 7777 8306 8755 9244	944677 5370 5693 6364 6645 7836 7836 8315 8004 9202	944667 5419 6404 7866 7875 7875 8863 9841	*********
5 10 15 20 24 29 34 89 44	801984567-09	949390 9678 960366 0861 1336 1823 2306 2792 8276 3760	949439 9926 957414 0900 1886 1872 2856 2841 8326 8808	949488 9975 950463 0949 1455 1920 2406 2889 3373 U 3856	949536 950034 0511 0097 1488 1969 9453 2036 3421 3005	949585 950078 0560 1046 1532 9017 2502 2966 8470 8963	949634 960121 0606 1096 1580 2066 2500 8034 8618 4001	949083 960170 0667 1143 1629 2114 2590 8088 8666 4049	949781 960219 0706 1192 1677 2163 2647 8181 3615 4098	949780 960267 0754 1240 1796 2911 2006 3180 3063 4146	940039 950616 0805 1260 1775 2000 2744 8208 8711 4194	*****
5 10 14 19 34 29 34 38 43	90133455789	964243 4735 5207 5688 6168 6649 7138 7607 8086 8564	954291 4773 5255 5736 6216 6697 7176 7655 8134 8612	954339 4821 5303 5754 645 6745 77224 7708 8181 8659	954387 4869 5351 5632 6313 7793 7.272 7761 8229 8707	954435 4918 5399 5880 6361 6840 77320 7799 8277 8755	954484 4966 5447 6928 6409 6888 7368 7847 8325 8803	954532 5014 5495 6976 6457 6936 7416 7894 8373 8850	954580 5062 5543 6024 6505 6984 7464 7942 8421 8898	964648 5110 5892 6072 6563 7032 7512 7990 8468 8946	954677 (5156 5640 6120 6601 7560 8038 8516 8094	********
5 9 14 19 24 28 33 38 42	910128866780	959041 9518 9995 960471 0946 1421 1895 2309 2843 3316	1469 1943 2417 2890	969187 9614 960090 0566 1041 1516 1990 2464 2987 3410	959186 9661 960138 0613 1069 1563 2038 2511 2985 3457	969232 9709 960185 0661 1136 1611 2085 2559 3032 8504	959280 9757 960233 0709 1184 1658 2182 2606 5079 3552	959528 9804 960281 0756 1231 1706 2180 2653 5126	969675 9852 960328 0804 1279 1753 2227 2701 8174 3646	960423 9900 960376 0851 1806 1801 2275 2748 8221 8693	969471 9047 90423 0699 1374 1848 2322 2796 8266 8741	46 46 47 47 47 47 47 47
5 9 14 19 23 28 33 38 42	920 1 2 8 4 5 6 7 8	963788 4260 4731 6202 5672 6142 6611 7080 7548 8016	4307 4778 5249 5719 6189 6658 7127 7595	6236 6705	7220 7688	6329 6799 7267		964071 4542 5018 5484 5954 6423 6892 7361 7829 8296	964118 4590 5061 5631 6001 6470 6939 7408 7876 8343	964165 4637 5108 5578 6048 6517 6986 7454 7922 8390	964212 4684 5155 5625 6096 6564 7033 7501 7969 8436	क्षा क्षा क्षा क्षा क्षा क्षा क्षा क्षा
5 9 14 18 23 28 52 37 41	930 1 2 3 4 5 6 7 8	968483 8950 9416 9882 970347 0812 12740 1740 2203	8996 9468 9928 970398 0858 1322 1786	9043 9509 9975 970440 0904	9090 9556 970021	9602 970068 0533 0997 1461	9183 9649 970114 0579 1044 1508	9229	968810 9276 9742 970207 0672 1137 1601 2064 2527 2980	968856 9323 9789 970254 0719 1183 1647 2110 2573 3035	1220 1693 2157 2619	46

PP	N.	Fol	1	9	3 3	î <b>4</b>	5	1:6	17	8	69	D.
CHESSEST	20198456789	978126 8090 4061 4512 4973 6438 5691 6360 6966 7366	978174 8636 4097 4586 8018 5478 5957 6386 6354 7313	978230 8688 4143 4004 5064 5524 5088 6442 6000 7368	978866 8736 4189 4680 5110 5570 6039 6498 6046	5616	5000 5000	5707 6167 6025 7063	6213 6671 7139	8700	6304	**********
5914888841	900 1234 56789	977724 8181 8087 9088 9648 960008 0458 0913 1306	977769 8236 8665 9138 9604 960040 0603 0967 1411 1864	977815 8278 8728 9184 9630 980094 0540 1003 1456	977861 8817 8774 9230 9685 980140 0504 1048 1501	977906 8363 8819 9275 9730 980385 0640 1098 1547 2000	977952 8409 8865 9821 9776 980231 0685 1139 1592 2045	977998 8454 8911 9366 9831 980276 0730 1184 1637 2090	978043 • 8500 8966 9413 9867 980322 0776 1239 1683 2185	978080 8546 9002 9457 9913 980367 0821 1275 1728 2181	978186 8591 9047 9608 9984 980412 0867 1320 1773 2296	********
5914181878841 1181878841	960128456789	989871 2733 8176 8636 4077 4527 4977 6498 5676 6324	982316 2760 3520 8671 4122 4572 5022 5471 8020 6369	982362 2814 8266 8716 4167 4617 5067 5516 5965 6413	982407 2869 3310 5762 4212 4662 5113 5561 6010 6458	982452 2904 8356 8807 4257 4707 5187 5606 6055 6503	982497 2949 3401 3852 4753 5202 5661 6100 6548	982543 2994 3446 3897 4347 4797 5247 5696 6144 6593	982588 8040 8491 8942 4392 4842 5292 5741 6189 6637	982633 5066 8656 5967 4437 4887 5337 5786 6234 6682	989678 8180 8581 4083 4485 4082 5582 5880 6279 6777	****
5 9 14 18 27 28 64 1 27 28 64 1	97128456789	9867772 7219 7866 8113 8560 9006 9400 9806 990539 0788	986817 7264 7711 51.57 8004 9049 9494 9959 990888 0827	96661 7309 7756 8202 8648 9094 9559 9983 990428 0671	966906 7356 7800 8247 8695 9138 9585 90028 0472 0916	986951. 7398 7845 8291 8787 9183 9678 9676 0516 0900	986996 7443 7890 8336 8782 9227 9672 990117 0561 1004	987040 7488 7934 8381 8826 9272 9717 990161 0605 1049	987086 7532 7979 8426 8871 9516 9761 990206 0650 1093	987130 7577 8024 8470 8916 9561 9806 990250 0694 1187	967175 7692 8068 8514 8900 9405 9650 990294 0738 1182	***********
4 9 13 18 22 26 SI	960128456789	901226 1600 2111 2554 2905 8436 8477 4317 4757 6196	991270 1713 2156 2598 8039 8480 8921 4361 4801 5240	991315 1758 2200 2642 3085 3524 8965 4405 4845 5284	991369 1802 2244 2686 3127 3568 4009 4449 4889 5328	991408 1846 2288 2730 3172 3613 4053 4493 4933 5372	991448 1890 2333 2774 5216 8657 4097 4537 4977 5416	991498 1936 2577 2819 3260 3701 4141 4581 6021 5460	991536 1979 2421 2863 3304 3745 4185 4626 5065 5504	991580 2023 2465 2907 8548 8789 4229 4669 5108 5547	991625 2067 2500 2961 3392 3853 4273 4713 5152 5501	333333333
4 9 13 18 22 26 31 36 40	990123456789	996635 6074 6512 6040 7386 7888 8250 8695 9181 9566	995679 6117 6555 6993 7430 7867 8303 8739 9174 9609	996723 6161 6599 7037 7474 7910 8347 8347 8782 9218 9652	996767 6205 6643 7060 7517 7964 8390 8826 9261 9496	905811 6240 6687 7124 7561 7998 8434 8800 9605 9739	995854 6293 6731 7168 7605 8041 8477 8915 9348 9783	995898 6337 6774 7212 7648 8085 8521 8956 9392 9836	905942 6390 6818 7255 7692 8129 8664 9000 9446 9870	995986 6424 6862 7290 7736 8172 8608 9043 9479 9918	998030 6468 6906 7343 7779 8216 8662 9627 9622 9657	THE STATES

	lynäre.	"Other	Sq. Rook	CubaBoot	Ma.	Square.	d Opbe.	Sq: Root,	Onbollo
,		n ** 12	12-0000000	1.000000	4	2000	200144	9-0000000	40000
اؤ	44	M (13 8	1-4142136	1-200021	488	4006 4255 4256 4420	374035	9-0033077	4-0207
Ì	9	-8- 2 <b>64</b>	17890008 9-0000000		-	4366		9-19400b4	40413
H	16	-4- 1 <b>64</b>	3-0000000	1.587401	.07	4454	90748 900768 914488	2.2403113	40616
		125 216 348	3-440-4807	1.817121	. 20	4634 4781 4900	30000	2-2064200	4 1014
71	1 0	345	2-6457518	1.913961	70	8041	# 843000	8-8000000	41919
	. 4	513	3.5	2000000	171	8047	307911 378348	8-4061406	41400
	100	1000	8-1622777	9-154496	72	5184	978348 989017 406234	3-6440087	4 1790
	100	720 1000 1831	8-1622777 3-31-66248	3-223060 3-236428	74	8390 8476 8626	405224	8-6140067 8-6023268 8-6002540	41985
	144	1728	3-4641016 3-6065513	7-280428 7-351336	74 78	5626	421875	8-0002540	42171
	100	3197	3-7416574	3-410142	. 76	6776	456528		
	196	2744 8375	8-8720833	2-400213	78	5020 0084	474403	8-8317600	4272
6	256 250 224	4000	4-00000000	9-510049	70	0041	403080	8-8661944	42000
71	250	4018	41231066	2-671282 2-630741	80	6400 6661	812000	8-9443719	4/5000
	1 31 450	6883	4-1231056 4-2425407 4-3688080	2 660	81		881441	0.0533003	777
678901	351	8000	4-4731300	2 71 MIS 2 70 MIS 2 70 MIS	3	7724	5 × 571767	9-1104300	0.9020
i)	1) 484 1) 484	9261	4-4998787	2-758934	84	7056	. 5003704	9-1451514	43794
	11 484	9261 10648 13167	4-6004158 4-7958315	2-803060 2-843067 2-864400 2-924018	30	7	614128 686066	9-319648	4414
	1 576	18824	II A PROPERTY LAND	2-884400	87	7506 7500		9-2275791	4:4810
5		of a little to the	J 5-0000000	2-924018	. 92	7744	8 081472	編 2 . 1 . 5 1 1 6	4447
8	676	17576	5-0990195 5-1961634	3-963496	80	7921 8100	704969 729000	9-4839611	4:451
7	754 754 800	19068	6-201.5026	8-000000 8-006060 8-073817 8-107233 8-141681	, 80 91	8100	753571	0-10000000	
800	i dan	24300	5-8851648	8-072817	80	8281 8464 8640	763671 779688 804387	9-0916600	4514
ol	900	27000	THE RESERVE OF THE PERSON NAMED IN	8-107233	. 86	8640	10 804367	9-0406500	4:5800
w	961	20/01	5-5677644 5-6608642 5-7445686	3-141881 3-174802	94	0000	857875	The state of	
3	1084	4 - 3 TO 168	P.7445698	8-207534	95	9025 9216 9409	884736	9-79/7950	4578
	1156	20304	1.0.02077213	3-239612 3-271066	I 97	9400	1 913073	9-9466078	
6	1206	42878	5-9100798	8-301927	55   55	9604 9801	941198	-	4010
4 5 6 7		21.0000	6-0000000	0.201821	100	10000	970290 1000000	10-0000000	2011
al i	1300	84872 84872 80819	6-0HB7625 6-1644140 6-2449980	9-361978 8-301211	96 100 101	10201 10404	1080301	10-0408706	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
8	1681	10 80819	8-9449980	8-391211	108 108	10404	EVOLUCE	10-00-000	1001
OI.	1600	64000	0-3245553 0-4031242	3-419052 3-448817	104	10616	1124864	10-1405010	
1	1681 1764	74088	6-4807407	3-470007	106	11025	1107025	10-2400508	TO VALLE
3		74088 79607 85184	6-5074385	8-508306 8-530848	106	111256	91191016	10-2056501	4782
3	1986 2025	85184	6-6332496	8-556893	107 108	11440	1225048 1269712	10-3440814	4747
6	2116	91125	6-7082039 6-7823300 6-8656546	3 583048		11881	1296029	10-4409000	1 4776
žI.	2200 2304	97336 103823	6-8556546	3.606826	100 110	. 12100	1331000	10-4690000	4:7914
7 8	2304	Parameter (Contract	0.0000000		111	12321 12544	1367631 1404028	10-8956586 10-6830082	4.8066
21	2600	117640	7-0000000	3°659306 3°684031	112 113	12700	1404028 1442897 1461544	10-6830052	4-8066 4-8206 4-8346 4-8486
ĭ	* 960h	132651	7-1414984	3.708430	114	12996	1461544	10-6770785	4-848
3	2704	140000	7'2111026	8-732611	115	375.745	1620875	10-7235065	
3	2704 2809 2916	148877	7-2801009	3.756286	116 117	13456 13680	1601618	10-7703296	4-9000
2	3V16	187464 166375	7·8484692 7·4181985	3-779763 8-802963	1118	13924	1643032	10-8627806	4.004
š	8136	175616	7-4853148	8:802068 8:826802 3:848501	119	14161	1,000,000	10 10 10 10 10 10 10 10 10 10 10 10 10 1	4-0194
7	3240	185198	7-5408344	3.848501	130	14400	1728000	110:9544515	4952
01334567890	8304 8481	196112	7.6157731 7.6811457	8-870877	띮	14641	1771061 1810848	11-0000000 11-0453610	4-9460
6	9000	\$16000	7-7459667	5-892996 5-914867	133	15129	1800007	11-0905365	4-973
n	8781 8844	205370 210000 220061 238328	7-7459667 7-8102497	3-098A07	134	16376	1900024	11-1855287	A STATE
1223	3844	258328	7 8740079	3-967892	126	15626 15876	2000576	11-1808399	E-0000
0	g. 8000	250047	7-9372539	0.313001	100	. TOO! 0			1

SECESTE SECESTES SECESTES SECESTES SECESTES SECESTES SECUSION OF SECESTES SECUSION OF SECU 

							مستسمع		
	Square	Out.	Sq. Book	CubeRest	Mei	Severe.	Oube	Sa. Book	Cabelloot
		985	7				, ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,		
127	16120	2046268	11-2014077	#-0260E6	190	86100	4040000	18-7840488	874007
139	16364	2007188	110 11 14 1	5-059664 5-062774	101	86461 M6844	: 6067671	18-77-40-488 13-6908-780 13-865-4085	8-7800e6
	16900	2197000	11-8578167 11-4017548	8-063774	198 198	87340	7160007	18-8934140	8-778900
iği	17161	2348091	1114455251	8-078758	194	87696 86026	7801844	140000	8-780000
	17494 17680	2200000	11-4801258	8-091648 8-104469	196	86026 86416	7414678	14-0000000	5-908786
194	17966	3406104	11.5758369	6-117330	197 198	80800	7040878	14-0000000	I COLOR 4
185 186	18225 18496	3460975 3515456	11-6189500	5-120028 5-142563	198	80204 80204 80601	7762863	14-0713473 14-1067860	8-80064R
11.77	18700	2071268	11-6619088	5-166187	199	40000	8000000	14-1491356	0 00075 V
隐	19044	2628072	11-7473444	5-167640	201	40401	8190001 8842408	14:17744	1-8577 90
	19891 19800 19861	2685619	11-7894361 11-8321596	5-180101 5-193494	뾃	40804	8005497	14:3136704 14:3478068 14:3838660	
180		2744000 2003221	11-6743431	5-102494 5-204828	3488	41616	3 8400064	14-2000500	5-8777.10 5-186741 5-186865 5-1868941
143	20164	2003208	11-9168758 11-9682607	5-217103 5-220821	205 916	42025	8615125 8741816	14·8178271 14·3527001	5-805041
144	207736 21025	". SORFORA	12-0000000	5-241488		ADDAM	50 // 51	14.577 (346	E 5 916 (65
9 44		8048698 8112136	12-0415046	5-25356% 5-260657	206 200	43264 43681	800801	14-4222051 14-45633553	8-9240-1 8-984478
17	21,600	8176523 8841798	12-0680460 13-1243557	5-277632	210	44100	9261000	14-52-58800	
	21904	8941793	12-1655261 12-2065556	5-280672 5-301459 5-318293	211	44521	9261000 9393081 9626128	14-5258800	5-953841 5-963/81
140	22500	8307940	113 3474487	5-318293	218	44944		14-5002196 14-5945196	
頭	20001	8443081	13-2882056 13-8288280	5-825074 5-836808	214	45796	9800844	14-6267386 14-6626783 14-6060385	8-981426 5-990727
H	28104	8511896 8061.577	10.0000100	A-348481	315 316	46236	10077696	14-00000785	1 6 000000
H	28716	· QUICESUS	19-4004794	5-300108	217	47080	10218313	14-7309199	6-009244
166	24025	8723878 8796416	13-440-906	5-371685 5-383213	318 219	47524 47961	10300232	14·7643981 14·7986456	6-027660
1157	24640	3940A08	13-4400006 13-480006) 13-5200641	5-304601	230	48400	10849000	1 ( 1355 771)	6-030611 6-045043
냺	24064 26281	8044312 4019679	12-5609051	5-406120 5-417501	231	49841	10793861		6-045943 6-055048
100	25600	4000000	12-6491106	8-428835	223	49720	11089667 11239424	14:8996644 14:9931845 14:9666995	6-064124
161	25021 26244	4173231 4251526	120000770	5·440133	224	50176	11230424	14-9666295	6.073178
163 163		4330747	13-7379281 13-7671468	8-451862 8-462556	236	50625 51076	11390625 11543176	15-0000000 15-0832964	6-082201
144	26896 27225	4410944	72-8062486	5-478704 5-484806	237	K1 590	11697068 11862362	15-0665192	6-100170
185 186	27556	4574206	13-8640007	5-495965	新報報報	51984 53441	13000980	15-0996689 15-1327460	6-109115 6-118033
107	27889	M. 4607463	12-9228480	5·495865 5·506879	230	52900	12167000	15-1657500	0.130830
168 160	28224 28561	4826800	12-9614814 12-0000000	5-517848 5-528778	251 288	53361	12326391 12487168	15-1986842 15-2315462	6-135792
170	28900	4913000	18-0884048	5-528775 5-539658	253	54290	7004000	4	6.159440
汉	20241	8000211 8088448	13.0766968	5-550499 5-561298	25A 255	54756 55225	12812904	15-2970585 15-3970585 15-3297097	6-162239
说	20584 20020	6177717	18-1148770 13-1529464	8-572065	236	55696	13144266	15-3522015	6.179747
11/21	30276	5268024	13-1909060	5.582770	237	56169	13312063	15-3948043	6·179747 6·188463
175 176	30625 30976	5350375 5451776	13-2864992	5-593445 5-604079	238 230	56644 - 57121	13481273	15-4672486	6·197154 6·205821
Ш	81329	5545233	13-8041347	5-614673	240	57000	13824000	15-4919334	6-214464
出	31684 32041	5735389	18-3416641 18-3790882	5-625226 5-635741	2/17	88781	14172488	15-5241747 15-5548492	6·223084 3·231679
179 180	39400	5832000	18-4164079	5-646216	19	50 %	14172488	15:58:4573	6.240251
181 182	83761	5929741 6028568	13-4536240 13-4907376	5-656651 5-667051	3/4	60025	14526789	15-6204994 15-6524758	6-248800
188	33489 33856	6128487	13-5277493	5-667051 5-677411	246 246	60516	14706126 14886936 15069223 15262992	15-6843871	6·257324 6·265826
184	33856	6128487 6229504	13:5646600	5-687734	247 248	61000	15069223	15-7162336	6-274305
185 186	84225 - 84596	6331625 6434866	13-6014705 13-6381817	5:708267	348	61504	15438249	15·7480157 15·7797338	6·282760 6·201194
187	34980	6539203	18-6747943	8-718479	250	62500	15625000	15-8113883	6.299604
IK.	35344 35721	6644672	13-7113092 13-7477271	5·728654 5·738794	251 252	63001		15-8429795 15-8745079	6-307993 6-816359
.07	COLOR	AL ATTEN	(E(1-1)	-100192		WW.	TANADA	TO OI MANIE	A.OTOOOR

•	Square.	Ouba.	Nq. Root.	Oubsileet	Ma.	Équare,	Oube.	Sq. Root.	Cubell
	أخمية	16194277	15-9000787	4-304704	316	-	NI STATES	17-7700000	0411
M H	. 04516	16387064 16381878 16777316	16-987378 16-9887194 16-9987194 16-9999000 16-9812196 16-9823784	6:364704 9:339726 6:341236	317	100486 101134	81856018	17-001-000	200 S
듺	94516 94005	16381875	15-9087194	6-341836	317 318	101194	38157488	17-0205545	0.00
H		16974566	14-0012195	A 347041	319	101761	2565 1789 2676 2000	17(0000711	dia.
	- 00004	160745 <b>48</b> 47173613	16-0023784	6-366098	100	708041	2076161	17-9104720	0.50
4	, 67091		16-093-4760	4 9 ( 49) 1	1883	10000001	38396948	17 5443150	500
1	6121	17876060 177770061	16-12-15188	0.500576		104976	7/11(0)(0)	7 2 7 2 3 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	A 200
3	00044		14.1044141			104076	BASSES	15 10 2	000
$\overline{\gamma}$	7 (0)	101607	16-24/27/47 16-24/20/768 16-37/20/20 16-38/20/24	8-406958 8-418068	99.	10es76	34646976	18-068-4701 18-0681-413	
+11	70005	1	162 77 100	6-428186		107504	Carponal Gra	18 1107708	
	70786	10001000	16-3000044	1412	330	107584 108241	35257468 30611200		8-900
Ž	7199	19094168	16-500-03-4 16-30013-4 16-3707056 16-4012196 16-4010776 16-40-0776 16-40-0776 16-40-0776 16-40-0776 16-40-0776 16-40-0776 16-40-0776	9-447866 9-447866 9-445815 9-445804	330	100000 100561	36594601 36594868	18-1059021 18-1054084	9910
4	(Alter)	10445100	18-4012195	0.447,835		110034	36504301 36504348	10-10-01	0-917
1	* ******	19000000	18-4110707	6-443004	353	110000	30020087		6 991
4	78461	19000000 19000511 2012046	16-4400776	647187		111556	. 87200704		8-945
뷔	74000	-	110	0-470234 0-407154 0-40006	355 534	112226	37506876 87903344	18-3030052 18-3303038	
	70076	. 3377 TESA	10-49-0454	6-400065	藝	113560	37500476 37933006 36373783	18-307-000	8-965
Q.	10 7 7 7 7 7 7 7	20790078 21004578	10 0001340	6-400065 6-802066 6-810690	335	114360	30514473	111	6.966
9	79720	31100076	16-6139477	0-51055U		114921	20204000	711	GALF
ωII	17284	21464000	A COMPANY OF STREET	6-518694 6-534396	W	116281	20651221	18-4661863	0.000
Ž	77364 77841 78400	\$1717600	16-7000081	6-534395	349	116281 116964 117640	40001688	18-4000430	8-9-8-9
d	78400	21962000 20188041	16-7000001 16-7802006 16-7600566 16-760566 16-8220068 16-8522008	6-534536 6-542123 6-540012 6-565415 6-577126	345	117640	40500507 40707864	18-4590666 18-4661855 18-4661855 18-466260 18-5673670 18-5741756	7,000
HП		29420708	10-7900556	0-657672	3434	119005		13.77	7:00
Ş	79584 80089 80656 61235	20426768 22665187	16-8226068	0-565415	346	119716	41431736		A
41	81995	22140125	16-8622006 16-8619430	6-573130	847 848	190400	41781925	18-6279800	
	81796	23303686 20430008		6-573130 6-540644 6-806532 6-506002	340	181801	42144192 42508540	18: (18-7765) 18: 08:18417 18: 70:00:00 18: 73:400:00 18: 76:16:00	7-040
700	82060		16-9410748 16-9705827	6-506808	350	1325001	42878000	18-7062800	7-047
4	83521	24187560	17-0000000	6-605664 6-611480 6-819106	151 181 181	123901 123904	43249651 43614206	Light	7
ŏ	84100	24389000	17-0293864	4419196	242	124400	43004077	11	200
	84681	24642171	1N.AROTEGE	6-636705 6-634287 6-641852 6-640800		100	44361864	16 71 COV	7-007
4	85264	24897088 26153757 26412184	17-068/0278 17-1172428 17-1464283 17-1755640 17-2046506 17-2636679	6-634387	355	19036 19036 19736 19746 198164	44738878 45118016 45499208	19.981442\	البياكا
000000000000000000000000000000000000000	94494	25412184	17-1464283	6-640300	367	137440	45499208	18-8679635 18-8944456	
5	87025	25672378 25084336	17-1755640	6-656980	33588	128164	46882712 46268279	B F. I'r   1 1 1741	7·008
엙	87616	26198073	17-2046608	6-640000 6-656000 6-664444 6-671940	P 19 13	128881 139600	46656000	16 9473969	7·107 7·118
ŝ	88909 88804	26463592	17-2626762 17-2916165	6-679420	860 861	180821	47045881		7:120
9	89401	26463592 26730899 27000000	17-2016165	6-686862	363 863	181044	47457926	TO COLUMN !	7·130 7·133
1	90000	27000000	17·3205081 17·3493516	0.701750	365	181760 132496	47832147 48228544	19-0526680	7:140
al	91204	27543608	17.8781472	6-709173	365	10220	486371.25	19-1040785	I BOT'
8	91809	27818127	17-4068952	6·709173 6·716570	364 365 366 367	133956	49027896	19-1311265	
	92416	28094464	17:4355958 17:4642492	6·723951 6·731816	30/	134680	49470863 49836032	19-1579461 19-1833261	7.144
5	93636	28372625 28652616	17-4928557	6.738665	388 369	200200			7-172
7	94249	28934443 29218112	17.6014186	6-738665 6-745997 6-753313	870	186000	50653000 51064811	19-2353841	7·172 7·179 7·185
780	94864	20218112 20503620	17·5499288 17·5783958	6·763313 6·760614	371 372	187641 188384	51064811 51478848	19-2613008	7:185
o	96100	29791000		6-767899	373 878	139120	51895117	19-287301 <i>5</i> 19-3132079	L. TAT
П	96721	\$0080231	17-6351921	6.775169	374	139129 189876	52313624	19-3390796	7.204
3	973441	30371328	17·6635217 17·6918060	6-782423	375	140625 141376	52734375 53157376		
	97969 98596		17-7200451	6-789661 6-796884	376 377	142129	53157376 53582633	19-4164878	7-224
	99225		17-7482303	6-804092	378	142884	54010152	19-4422321	7-230

Ko.	Square	g Cube.	Sq. Root.	CubeRoo	Ne	Square.	Oube.	Sq. Root.	CubeRo
79	143641	54430939	19-4679223	7·286797 7·245156	443	195864	86350888	21-0237900	7-61741
	144400 145161	54872000	19-4935867	7-243156	443	196240	80038807	21-0475062	7-02534
1	145924	55306341 55742968	19-5192213 19-5448203	7·249504 7·255841	445	197136 198025	87528384 88121125	21-0713075 21-0950231	7-02000
	146689	56181887	19-5703858	7-252167	446	198916	88716536	21 1187721	7-6403
	147456	56623104	19-5959179	7:268482	447	199800	80314623	21-1423745	7-64600
35	148925	57066625	19-6214169	7-274786	448	200704	80915302 90518849	21-1660105	7-65171
記録を発	148996	57512456 57960603	19-6468827 19-6723156	7·281079 7·287362 7·293633	449	201601	91126000	21-1806201 21-2132034	7:65743 7:64306 7:66876
88	149700	58411072	19-6977156	7-293633	451	203401	91126000 91733851	21-2367606	7-0687
180 180	151821	÷ 58863800	19-7230829	7:299894	452	204304			7.0744
000	152100	59319000	19-7484177	7.306143	453	205209	92959077	21-2637967	7.68000
91 92	152881	60236288	19-7737199 19-7980899	7:312383 7:318611	454	206116	93576664 94196376	ZI.20/2(00	7-68573
103	154449	60698457	19.8242276	7.324829	45	207936	94818816	21-3641565	7-69700
104	155236	61162084	19-8494332	7:331087	457	207936 208849	95443993	21-3775583	7.70263
25	156025	61629875	19-8746069	7:337234	458	209764	96071912	21-4009346	7071832
96	156816	62099136	19-8997487 19-9248588	7·343420 7·349597	38	210681	96702579	21·4242853 21·4476106	7-7138
de.	157009	63044792	19-9499373	7.356762	161	211600 212521	97330000 97972181		7.7250
98 98	159201		19-9749844	7-361918	462	213444	98611128	21-4941853	7.7306
100	160000	64000000	20-0000000	7:368063	463	214369	99252847		7.73618
01 02	160801	64481201	20-0249844 20-0499377	7:374198 7:380322	464	215296 216225	99897344 100544625	21-6406892 21-6638687	7.74171
03	162409	65450827	20-0459577	7-386437	466	217156	· 701104404	21-5670331	7·74731 7·76280
04	163216	65939264	20-0997512	7:392542	467	218089	T0124/003	21-6101828	7.75840
05	164025	66430125	20-1246118	7:398636	408	219024	102503232	21-6333077	1.10000
60	164836		20-1494417	7:404720	409	219961	103161709	21-6564078	7.76040
07 08	166649	67419143	20·1742410 20·1990099	7:410795 7:416869	470 471	220900 221841		21-6794834 21-7025344	7-77496
00	167281	68417929	20-2237484	7:422014	472	222784	105154040	01-70KKE10	7-78899
10	168100	68921000	20-2484567	7-428959	473	25 720	105823817	21.7485632	7-79148
11	168921		20-2731349	7:434994	474	224676	106496424	21-7715411	7-79097
12	170669	69934528 70444997	20-2977831 20-3224014	7·441019 7·447034	476 476	225625 226576		21 <i>-</i> 7944947 21-8174242	7-80792
ii.	171396	70957944	20-5469899	7-453040	477	227529		21-8403297	7.81338
15	179998	71473375	20-3715488	7:459036	478	228484		21-8632111	7.81884
16	173056	71991296	20-3960781	7:465022	479	229441 230400		21-8860686	7·82420 7·82973
芸	173889 174724	72511713 73034632	20·4205779 20·4450483	7:470999	480 481	231361	111284841	21-9089023 21-9317122	7.83516
10	175561	73560059	20-4694896	7·470966 7·482924	482	232324	111284641 111980168	21-9544984	7-84059
20	176400	74088000	20-4939015	7.488872	483	233289	112678587	21.9772610	7.84601
21	177241		20-5182845	7:494811	484	234256	113379904	22.0000000	7.85142
22 23	178084		20·5426386 20·5669638	7:500741 7:506661	485 486	235225 236196		22-0227156 22-0454077	7.85682
24	179776	76225024	20.5912603	7:012571	487	237169	115501303	22-0680765	7-86761
25	180625	76765625	20-6155281	7-619479	488	238144	116214272	22.0907220	P.OTROC
26	181476	77308776	20-6397674	7.524365 7.530248	489	239121	116930169	22-1133444	7.87830
27 28	182329		20·6639783 20·6881609	7.536121	490 491	240100 241081	117649000 118370771	22·1359436 22·1585198	7·88373 7·88909
20	184041		20.7123152	7.541986	492	242064		22-1810730	7.89444
30	184900	79507000	20.7364414	7.547842	493	243049	110923157	22-2036033	7-89979
31	185761		20.7605395	7.553688	494	244036		22-2261108	7-90512
32 33	186624	80621568 81182737	20.7846097	7·559526 7·565355	495 496	245025 246016	121287375 122023936	22-2485955	7·91046 7·91578
33 34	188356	81746504	20-8086520 20-8326667	7.571174	497	247009	122763473	22-2710575 22-2934968	7-92110
35	189225	82312875	20.8566536	7.576985	498	248004	123505992	22:3159136	7.92640
36	190096		20.8806130	7.582786	499	249001		22-3383079	7-93171
37	190969		20-9045450 20-9284495	7·588579 7·594363	500	250000 251001		22-3606798	7·93700 7·94229
38 39	191844	84027672 84604619	20-9523268	7.600138	501 502	252004	125751501 126506008	22-3830293 22-4053565	7-94757
40	193600	85184000	20.9761770	7.605905	503	253009	127263527	22-4276615	7-95284
41	194481		21.0000000	7.611662	504	254016		22-4199443	7-95811

-

79		P		
	C 1914	127) marke		
	900			
	900		뛻	
	7.6	$\frac{\pi}{2}$		
	7.6			
	7.0			
	7·6 7·7	97( 92( 98)	125 126 100	4.1
3	7.7	13 19	35.52	
300	7.7	30 36	514 188	
	7   1   7	41 47 62		
B 7 R	7·7 7·7 7·7	88	402 030 462	
44	7.7	74	980 490	
2	7	91	457 97	
227	7.8	07 313	92i 38	
6	7.8	21 22 23	844 294 734	
24.5	7.1	335 340 846	169 696	
000	7.	851 856	12 12 12 13 13 13 13 13 13 13 13 13 13 13 13 13	
500	7.	307 374	61. 99	
468	7.7.7	871 881 880	73 73 109	5
999	7.	894 899 904	779	2
50.00	7.	910 910 92	046 578	03
666	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	92	640 171	8
25.50	7	94	229 757	34
335555555555557 L87544021727163240057504680385586985553	7	95	811	4
	_		-	

fo.	Square.	Cube.	Sq. Root.	CubeRoot	No.	Square.	Cube.	Sq. Root.	CubeRoo
05	255025	128787625	22-4722051	7-963374	568	322624	1882/0432	23-8327506	8-21863
00	256036	129554216	22-4944438	7-968627	569	823761	184220009	23-8537209	8-28649
07 08	257049 258064	130323843 131096512	22-5166605 22-5388553	7·973873 7·979112	570 571	324900 326041	185193000 186169411	23·8746728 23·8956063	8·2913 8·2961
09	259081	131872229	22-5610283	7-884944	572	327184	187149248	23-9165215	8-3010
10	260100	132651000	22-5831796	7.989570	573	328329	188132517	23-9374184	8-3068
П	261121	133432831	22-6053091	7:994788 8:000000	574	329476	189119224	23.9582971	8.3106
	262144 263169	134217728 135005697	22-6274170 22-6495033	8-000000	578 576	330625 331776	190109375 191102976	23·9791576 24·0000000	8·3155 3·3203
ĭ	264196	135796744	22-6715681	8-010403	577	232920	192100033	24-0208243	8-3251
15	265225	136590875	22.6936114	8.015695	578	334084	193100552	24.0416306	8-3299
16	266256	137388096	22.7156334	8-020779	579	335241	194104539	24.0624188	8:3347
17 18	267289 268324	138188413 138991832	22-7376340 22-7596134	8-025957 8-031129	580 581	836400	196112000 196122941	24·0831892 24·1039416	8.3395
Š	269361	139798359	22.7815715	8-036293	582	837561 838724	197137368	24-1246762	
20	270400	140608000	22-8035085	8-041451	583	334724 33988	198155287	24-1453929	8:3539
21	271441	141420761	22.8254244	8-041-61 8-046608	584	341056	199176704	24-1660919	8:3586
22	272484	142236648	22-8473198	8-051748	585	342225	200201625	24-1867732	
Ž	273529 274576	143055667 143877824	22·8691933 22·8910463	8.056886 8.062018	586 587	343396 344569	201230056 202262003	24-2074769	8:3682
26	275625	144703125	22-9128785	8-067143	588	845744	203297472	24-2487113	8:3777
26	276676	145531576	22-9346899	8·072262 8·077874	589	346921	204336460	24-2693222	8:3824
27	277729	146363183	22-9564806	8.077374	590	848100	205379000 206425071	24.2899156	8.3872
8	278784 279841	147197952	22-9782500	8.083480	591 592	349281	206425071	24-3104916	8:3919 8:3966
ő	280900	148035889 148877000	23:0000000 28:021723	8·087579 8·092672	593	350464 351649	207474688 208527857	24·3310501 24·8515913	8.4013
ñ	281961	149721291	23-0434379	8-097759	594	352836	209584584	24 3721152	
12	283024	150568768	23.0651252	8.102839	595	354025	210644875	24-3926218	8:4108
3	284089	151419437	23 0867928	8-107913	596	355216	211708736	24-4181112	
4	285156 286225	152273304 153130375	23·1084400 23·1300670	8-112980	597	356409 357604	212776173 213847192 214921799	24·4335834 24·4540385	8-4202
55 56	287296	153990656	23 1516788	8·118041 8·123096	598 599 600	358801	214021790	24 4744765	
7	287296 288369	154854158	23.1732605	8-128145	600	860000	216000000	24-4948974	8-4343
18	289444	155720872	23.1948270	8-153187	601	361201	217081801	24-5153013	8:4390
Ď	290521	156590819	23-2103785	8-138223	602	362404	218167208	24-5356883	8-4436
10	291600 292681	157464000 158340421	23·2379001 23·2594007	8·143253 8·148276	603 604	364816	219256227 220348964	24·5560583 24·5764115	8·4483 8·4530
2	293764	159220088	23:1008935	8.153294	605	366026	221445125	24-5967478	8-4576
2234	294849	160103007	23:3023604	8.158305	606	366025 367236	222545016	24.6170673	8.4623
И	295936	160989184	23-3238076	8.163310	607	368449 -369664	223648543	24.6373700	8.4670
10	297025	161878625	23-3452351	8.168309	608		224755712	24-6576560	8.4716
6	298116 299209	162771336 163667323	23·3666429 23·3880311	8·173302 8·178289	609 610	370881 372100	225866529 226981000	24·6779254 24·6981781	8·4762 8·4809
8	300304	164566592	23.4093998	8-183269	611	373321	228099131	24.7184142	
9	301401	165469149	23.4307490	8-188244	612	374544	229220928	24-7386338	8.4901
0	302500 303601	166375000	23-4520788	8-193213	613	875769	230346397	24.7588368	8·4948 8·4994
1 2	303001	167284151 168196608	23·4783892 23·4946802	8·198175 8·203132	614 615	376996 378225	231475544 232608375	24·7790234 24·7991935	8.5040
3	304704 305809	169112377	23.5159520	8.208082	616	379456	233744896	24.8193473	8.5086
ă	306916	170031464	23.5372046	8-213027	617	380689	234885113	24-8394847	8.5132
55	308025	170953875	23.5584380	8.217966	618	881924	236029032	24.8596058	
6	309136	171879616 172808693	23.6796522 23.6008474	8-222898	619 620	383161 384400	237176659 238328000	24·8797106 24·8997992	8-5224
7	311364	178741112	23.6320236	8:227825 8:232746	621	385641	238328000	24 9198716	8-5316
59	312481	174676879	23.6431808	8:237661	622	886884	240641848	24.9399278	8-5361
30	313600	175616000	23.0643191	8-242571	623	388129	241804367	24.9599679	8.5407
1	314721	176558481	23-6854386	8-247474	624	889376	242970624	24.9799920	8.5453
12 18	315844	177504328 178453547	23·7065392 23·7276210	8-252371	625 626	390625	244140625	25·0000000 25·0199920	8.55498
Н	318096	170406144	28.7486842	8·257263 8·262149	627	891876	245314376 246491883	25.0399681	8-5589
80	319225	179406144 180362125 181321496	28.7697286	8-267029	628	593129 594384	247673152	25.0599282	8.5635
86 80	320356	181921496	28-7907545	8-271904	ROD	895641	248858139	25-0798724	8.5680
57	321489	174384268	23-8117618	8-286778	6.40	396900	250047000	25,0998008	8.5726

No.	Square.	Cube.	Sq. Root.	CubeRoot	No.	Square.	Cube.	Sq. Ro	ot.	CubeRoo
131	398161	251239591	25-1197134	8-577152	694	481636	334255384	26:3438	797	8-85366
132 133	399424	252435968	25-1396102	8.581681	695	483025	835702375	26-3628	527	8.85784
33	400689	253636137	25-1534913	8-586205	696	484416	837153536	26.3818		8-86200
334 335 336	401956	254840104	25-1793566	8·590724 8·595238	697	485809	838608873	26-4007		8-8663
150	403225	250047875 257259456	25·1992063 25·2190404	8.599747	608 699	487204 488601	340068392 341532099	26:4196 26:4386		8·8705 8·8748
37	405769	258474853	25-2388585	8.604252	700	490000	843000000	26-4575		8.8790
38	407044	259694072	25-2586619	8-608753	701	491401	344472101	26-4764		8-8832
39	408321	260917119	25-2784493	8-613248	702	492804	345948408	26-4952		8-8874
140	409600	262144000	26-2982213	8-617739	703	494209	347428927	26.5141	472	8-8917
#	410881	263374721 264609288	25:3179778	8-622225	704 704	495616 497025	348913664 350402625	26·5329 26·5518	963 961	8-8959 8-9001
7	412164 413449	265847707	25·8377189 25·3574447	8·626706 8·631183	705 706	498436	351995816	26.5706	80¥ 201	8-9043
101121214 11112121414	414736	267089984	25-3771551	8-635655	707	499849	351895816 353393243	26-5894		8-9085
145	416025	268336125	25-3968502	8-640123	708	501264	354894912	26-6082		8-9127
346	417316	269586136	25.4165301	8-644585	709	502681	356400829	26.6270		8-9169
47	418609	270840023	25.4361947	8.649044	710	504100	357911000	26-6458		8-9211
347 348 349	419904 421201	272097792	25·4558441 25·4754784	8-653497 8-657946	711 712	505521 506944	359425431 360944128	26.6645 26.6833		8-9253
150	422500	273359449 274625000	25-4950976	8.662391	713	508369	362467097	26·7080		8-9336
151	423801	275894451	25.5147016	8-666831	714	509796	363994344	26.7207		8-9378
52	425104	277167808	25.5342907	8.671266	715	511225	365525875	26.7394	839	8-9420
53	426409	278445077	25.5538647	8.675697	716	512656	367061696	26.7581	763	8-9461
54	427716	279726264	25.5734237	8.680124	717	514089	368601813	26.7768		8-9503
55 56	429025 430336	281011375 282300416	25-5929678 25-6124969	8·684546 8·688963	718	515524 516961	370146232 371694959	26·7955 26·8141		8-9545
57		283593393	25.6320112	8.693376	719 720	518400	979249000	26.8328		8.9628
57 58	431649 432964	284890312	25-6515107	8.697784	721	518400 519841	374805361	26.8514		8.9669
59	434281	286191179	25-6709953	8.702188	722	521284	376367048	26.8700		8-97110
60	435600	287496000	25-6904652	8.706587	723	522729	377933067	26.8886		8-9752
61	436921	288804781	25.7099203	8.710983	724	524176	379503424	26-9072		8-9793
62	438244	290117528 291434247	25·7203607 25·7487864	8·715373 8·719759	725 726	525625 527076	381078125 382657176	26-9258 26-9443		8-98350 8-98763
63 64 65	440896	292754994	25.7681975	8-794141	727	528529	334240583	26-9629	0/ A 97K	8.9917
8K	442225	294079625	25.7875939	8·724141 8·728518	728	529984	385828362	26.9814		8-9968
66	443556	295408296	25.8069758	8.732892	729	531441	387420489	27.0000		9-0000
66 67	444889	296740963	25.8263431	8.737260	730	532900	889017000	27-0185	122	9-0041
68 69	446224	298077632	25.8456960	8.741624	731	534361	390617891	27-0370		9.0082
60	447561		25·8650343 25·8843582	8.745985	732	535824	392223168 393832837	27-0554		9-0123
70	448900	300763000 302111711	25 9036677	8-754691	733 734	537289 538756	395446904	27·0739 27·0924		9-0164
71	451584	303464448	25-9229628	8.759038	735	540225	397065375	27.1108	224	9.0246
73	452929	304821217	25-9422435	8.763381	736	541696	398688256	27-1293	199	9-0287
74	454276	306182024	25-9615100	8.767719	737	543160	400315553	27-1477	439	9.0328
75	455625		25.9807621	8.772053	738	544644	401947272	27.1661		9-0368
<u>76</u>	458976		26-0000000	8-776383	739	546121	403583419	27-1845	110	9-04090
77 78	700		26·0192237 26·0384331	8·780708 8·785029	740 741	547600 549081	406869021	27·2029 27·2213	FIU	9-0491
79	461041		26.0576284	8.789346	742	550564	408518488	27-2396	780	9-05318
79 80	462400	314432000	26.0768096	8.793659	743	552049	410172407	27-2580	263	9.0572
81	463761	315821241	26.0959767	8.797968	744	553536	411830784	27.2763		9-0613
82	465124		26.1151297	8.802272	745	555025	413493625	27.2946		9.06536
83 84	466489	318611987	26.1342687		746	556516	415160936	27.3130		9-06942
85 85	467856 469225	320013504 321419125	26·1533937 26·1725047		747 748	559009 559504	416832723 418508992	27:3313 27:3495		9-07347
NA.	470596	322828856	26-17250-17	8.819447	749	561001	420189749	27·3678		9-07752 9-08156
86 87	471969	324242703	26.2106848	8.823781	750	562500	421875000	27.3861		9.08560
88	473344	325660672	26-2297541	8.828009	751	564001	423564751	27-4043		9-08963
89	474721	327082769	26-2488095	8-832285	752	565504	425259008	27-4226	184	9.09367
90	476100		26.2678511		753	567009		27.4408		9-09770
91	477481	329939371	26-2868789		754	568516	428661064	27:45900		9.10172
92 93	478864 480249	331373888 332812557	26·3058929 26·3248932	8-845085 8-849344	755 756	570025 571536		27·47720 27·4954		9·10574
46.0	- WEAUGH	OUECIEUD/ I	AU JASOVJA	0.02015	100	0/1000		ALCENIA	2020	# 11 <i>75/</i> / 1

7	- 13	, 1 2	1					\$ e t	- /	
CubeRoot	No.	Square.	Cube.	Sq. Root.	CubeRoot	No.	Square.	Cube.	Sq. Root.	Oi
8-853598 8-857840 8-862095	757 758	573049 574564	433798093 436519512	27·5136330 27·5317998	9-113781 9-117793	820 821	672400 674041	551368000 553387661	28·6356421 28·6630976	1 9
8-862096	759	576081	437245479	27-5499546	9-121801	822	675684	555412248	28-6705424	1
8-870676	760 761	577800 579121	438976000 440711081	27·5680975 27·5862284	9·125805 9·129806	823 824	677329 678976	850476224	28·6879766 28·7054002	1 3
8-874810	762 763	580644	442450728 444194947 445943744 447697125 449455096	27·6043475 27·6224546	9-133803	825	680625	559476224 561515625	28-7228182	1
8·879040 8·883266	763	582169	444194947	27-6224546	9-137797	825 826	682276	563559976 565609283 567663552	28-7402157	1
8-887488	764 765 766	583696 585225	447697125	27·6405499 27·6586334	9·141788 9·145774	827 828	683929 685584	567663562	28-7576077 28-7749891	13
8·891706 8 8·895920	766	586756	449455096	27.6767050	9-149757	820	687241	56972278V	28 7923601	
8-900130 8-904336 8-904538	767 768 769	588289 589824	451217663 452984832	27·6947648 27·7128129	9-153737 9-157714	830 831	688900 690561	571787000 573856191	28·8097206 28·8270706	
8-904336	769	591361	454756609	27.7308492	9-161686	832	692224	575930368	28-8444102	
8-908538	770	592900	456533000	27.7488739	9-165656	833	693889	578009537	28-8617394	
8-916931	771	594441 595984	458314011 460099648	27·7668868 27·7848880	9·169622 9·173585	834 835	695556	580093704 582182875	28-8790582	ı
8-921121	773	597529	461889917	27-8028775	9-177544	835 836	698896	584277056	28-8963666 28-9236646	
8-925308 8-929490	774	599076 600625	463684824	27·8208555 27·8388218	9-181500 9-185443	837	700569 702244	586376253 588480472	28-9309523 28-9482297	ı
8-933668	775	602176	465484375 467288576	27-8567766	9.189402	838 839	703921	590589719		ı
8-937843 8-942014	777	603729	469097433	27.8747197	9-193347	840	705600	592704000	28-9827535	
8-942014 8-946181	778 779	605284	470910952 472729139		9-197289 9-201229	841 842	707281 708964	594823321 596947688	29-0000000 29-0172363	
8-950344	.  780	608400	474552000	27.9284801	9.205164	843	710649	599077107	29-0344623	
8-954503 4 8-958658 7 8-96809	781	609961 611524	476379541 478211768	27-9463772	9-209096 9-213025	844	712336 714025	601211584	29.0516781	
8.962809	783	613089	480048687	27-9642629 27-9321372	9-215025	846	715716	603351125	29-0688837 29-06607791	12
8-966957	782 783 784 785	614656 616225	480048687 481890304 483736625	28-0000000	9-220873	847	717409 719104	605495736 607645423	29·0860791 29·1032644	
8-971101 8-975240	785	616225	483736625 485587656	28-0178515 28-0356915	9-224791 9-228707	848 849	719104 720801	609800192	22.12042AC	34 3
8-979376	787	619369	487443403		9-232619	850	722500	614125000	29-1547595	
8-983509	788	620944	489303872	28-0713377	9-237528		724201	616295051		
8·987637 8·991762	789 790	622521	491169069 493039000	28-0891438 28-1069386			725904 727609	618470208	29-2061637	7
8-996883	791	625681	494913671	28-1247222	9.248234	854	729316	622835864	29-223278	
9-000000 9-004113	792 793	627264 628849	496793088 498677257	28·1424946 28·1602557	9-252130 9-256022	855 856	731025	625026375 627222016	29-2403830 29-2574777	
0.008223	1/32	630436	500566184	28-1780056	9.259911	857	734449	629422793	29-274562	3
9-0123254	1790	632025	502459875	28-1957444	9-263797		736164	631628712	29-29162	
9·016431 9·020529	796 797	633616 635209	504358336 506261573		9-267690 9-271559			633839779		
9-024624	17UH	RESERVA	508169592	28-2488938	9-275435	861	741321	638277381	29-342801	5
9-028715 9-032802	799 800	638401 640000	510032399 512000000	28·2665881 28·2842712	9·279308 9·283178	862 863			29-3598360 29-376861	
9-036886	801	641601	513922401	28-3019434	9 237044	1884	746496	644972544	29-393876	9
9-040960	801 802 803	643204	513922401 515849608	28-3196045	9-290907	865 866	748225 749956	644972544 64721462	29-4108823	3
9-045041 9-049114	804	644809 646416	517781627 519718464	28·3372546 28·3548938	9·294767 9·298624	867	751689	649461896 651714363		-
9-063183	80/5	648025	521660125	28.3725219	9.302477	868	753424	653972032	29-4618397	7
9-049114 9-053183 9-057248	806	649636	523606616		9·306328 9·310175			656234909 658503000		
9-065367	808		525557943 527514112	28-4253408			758641	66077 6311	29.512709	
06 9-069422	809	654481	529475129	28-4429253	9-317860	872	760384	663054848	29.529646	L
07 9·073473 9·077520	810  811		531441000 533411731			873 874		665338617	29·5465734 29·5634910	
9-077520 9-081563 79 9-085603 92 9-089639	812	659344	535387328	28-4956137	9.329363	875	765625	669921875	29.5803989	)
9.085603	813   814		537367797	28.5131549	9.333192	876	767376	672221376	3 29 5972972	3
9·089639 9·093672	815		539353144 541343375			877 878	769129 770884			
9-097701	816	665856	543338496	28-5657137	9.344657	1879	772641	679151439	29-6479325	5
9-101726 9-105748	817	667489 669124	545338513 547343432	28·5832119 28·6006993	9-348478 9-352286	880 881	774400 776161	681472000 683797841	29-6647939 29-6816442	
	818 819	670761	549353259	28-6181760	9-356095	882	777924	686128968	29-6984848	
542 9-109766		1,					1			1

19/	Square.	Cube.	Sq. Root.	CubeRoot	No.	Square.	Cube.	Sq. Root.	CubeRoo
	Edward.	, Oupo.	By. Moos	Cubertoor	10.	Didage	. 44	F F	0.00
888	779680	688465387	29.7153159	9-593716	942	887364	835896888	30-6920185	9-802804
884	781456	690807104	29-7321375	9-597337	943	889249	838561807	30-7083051	9.80627
995	783225	693154125	29-7489496	9-600955	944	891136	841232384	30-7245830	9.80973
	784996	695506456	29-7657521	9-604570 9-608182	945 946	893025	843908625 846590536	30·7408523 30·7571130	9.813199 9.816659
	786769 788544	697864103 700227072	29·7825452 29·7993289	9.611791	947	894916 896809	849278123	30 77 33651	9.82011
999	790321	702595369	29.8161030	9-615398	948	898704	851971892	30-7806086	9 823572
800	792100	704969000	29.8328678	9-619002	949	900601	854670349	30.8058436	9.82702
801 802 803 804	793881	707347971	29.8496231	9.622603	950	902500	857375090	30·8220700 30·8382879	9.83047
-22	795664	709732288	29.8663690	9-626201	951	904401	860085351	80.8382879	9.83392
	797449	712121957 714516984	29·8831056 29·8998328	9·629797 9·633390	952 953	906304 908209	862801408 865523177	30·8544972 30·8706981	9.83736
905 906	801025	716917375	29-9165506	9.636981	954	910116	868250664	30.8868904	9.84425
106	802816	719323136	29-9332591	9-640569	955	912025	870983875	30'9030743	9.84769
- 74	804609	721734273	29-9499583	9.644154	956	913936	873722816	30-9192497	9.85112
806 800	806404	724150792	29-9666481	9-647737	957	915849	876467493	30.9354166	9.85456
900	808201	726572699	29-9833287	9-651317	958	917764	879217912	30'9515751	9.85799 9.86142
901	810000 811801	729000000 731431701	30.00000000	9-654894 9-658468	959 960	919681 921600	881974079 884736000	30.9677251 30.9838668	9.86484
002	813604	733870808	30.0533148	9-662040	961	923521	887503681	31.0000000	9.86827
	815409	736314327	30-0499584	9.665609	962	925444	890277128	31 0161248	9.87169
003 004	817216	738763264	30.0665928	9.669176	963 964	927369	893056347	31.0322413	9.87511
105	819025	741217625	30.0832179	9.672740	964	929296	895841344	31.0483494	9.87853
06	820836	743677416	30.0998339	9.676302	965	931225	898632125	31.0644491	9.88194
07	822649	746142643	30-1164407	9.679860	966	933156	901428696	31.0805405	9.88535
006 009	824464 826281	748613312 751389429	30·1330383 30·1496269	9-683416 9-686970	967 968	935089	904231063 907039232	31.0966236 31.1126984	9.88876
10	828100	753571000	30 1450200	9.690521	969	937024 938961	909853209	31 1287648	9.89558
ni	829921	756058031	30-1827768	9-694069	970	940900	912673000	31 1448230	9-89898
12	831744	758550528	30-1993377	9-697615	971	942841	915498611	31-1608729	9-90238
13	831744 833569	761048497	30-2158899	9-701158	972	944784	918330048	31-1769145	9-90578
014	835396	763551944	30-2324329	9-704699	973	946729	921167317	31.1929479	9-90917
215	837225 839056	766060375	30-2489669	9.708237	974	948676	924010424	31-2089731	9.91257
)16 )17	840889	768 <b>57</b> 5296 771095213	30-2654919 30-2820079	9-711772 9-715305	975 976	950625 952576	926859375 929714176	31·2249900 31·2409987	9-91596
018	842724	773620632	30-2985148	9-718835	977	954529	932574833	31.2569992	9.91930
110	844561	776151559	30-3150128	9-722363	978	956484	935441352	31.2729915	9.92612
920 931	846400 848241	778688000	30-3315018	9-725888	979	958441	938313739	31-2889757	9-92950
931		781229961	30-3479818	9.729411	979 980	960400	941192000	81.3049517	9.93288
74	850084	783777448	30.3644529	9.732931	981	962361	944076141	31-3209195	9.93626
23	851929	786330467 788889024	30-3809151	9-736448	982	964324	946966168	31-3368792	9.93963
DK.	853776 855625	791453125	30·3973683 30·4138127	9.739963 9.743476	983 984	966289 968256	949862087 952763904	31-3528308	9·94300 9·94638
234 225, 226	857476	794022776	30-4302481	9.746986	985	970225	955671625	31·3687743 31·3847097	9-9-974
711	859329	796597983	30.4466747	9.750493	986	972196	958585256	31.4006369	9-95311
28	861184	799178752	30.4630924	9.753998	987	974169	961504803	31-4165561	9-95647
)28 )29 )30	863041 864900	801785089	30-4795013	9.757500	988	976144	964430272	31-4324673	9.95983
30	864900	804357000	30-4959014	9.751000	989	978121	967361669	31-4483704	9.96319
31	866761 868624	806954491 809557568	30·5122926 30·5286750	9·764497 9·767992	990 991	980100	970299000 973242271	31-4642654	9.96655
33	870189	812166237	30.5450487	9.771484	991	982081 984064	976191488	31·4801525 31·4960315	9·96990 9·97326
34	872356	814780504	30.5614136	9.774974	993	986049	979146657	31.5119025	9-97661
35	874225	817400375	30.5777697	9.778462	994	988036	982107784	31.5277655	0.07006
36	876096	820025856	30-5941171	9.782946	995	990025	985074875	31.5436206	9-98330
37	877969 879844	822656953	30.6104557	9.785429	996	992016	988047936	31.5594677	9.98664
38		825293672	30.6267857	9.788909	997	994009	991026973	31-5753068	9.98999
39	881721 883600	827936019	30-6431069	9.792386	998	996004	994011992	31.5911380	9-99332
49	885481	830584000 833237621	30.6594194 30.6757233	9-795861 9-799334	999	998001	997002999	31.6069613	9.99666
-	200,01	000401041	ON OLO 1700	9.188324	1000	1000000	1000000000	31.6227766	10.00000

# ANSWERS TO MISCELLANEOUS EXERCISES.

#### EXERCISE 8.

2. Sixty-seven trillions eight hundred and forty-five billions three hundred and ninety-eight millions six hundred and seventy-eight thousand nine hundred and four.

Five quadrillions nine hundred trillions seven hundred and four billions sixty millions forty thousands, and sixty thousand six hundred and four hundredths of millionths.

- 3. MVDCCLXIX.
- 4. 429860000.

- 5. \$67.31\frac{1}{2}.
  6. 77991.
  7. 605000070016-000009.
  8. 46978900.
  10. 69.800463.
  11. .8439.
  12. 678900000.
  13. 6043298600000000.
  14. 1000001000001001.00000000000001.
- **15**. ·0007609.

16. Ninety trillions eight hundred and seven billions sixty millions five hundred and four thousand and thirty.

Four quintillions four quadrillions forty trillions four hundred billions sixty thousand four hundred and thirty-two. and one trillion ten billion two hundred and three million forty thousand five hundred and six hundredths of trillionths.

- 18. 771 cords.
- 19. 717 cords 91 cubic feet.
- 20. DOCKVIII, DOXIV, CDXCIX, CMXCIX, VMMMDCXLIII, XOVMOXLIX, OLXMMMOMLXXXVI, ODXLMVODXLIV.
- 21. 333, 1989, and 1000001.
- 25. \$3.75 52, \$24.581, 713, and \$757.4711.

#### EXERCISE 17.

- 1. \$18029304.
- 2. \$13999999.73.
- 3. 36497318.
- 4. 35857536.
- 5. 27424500.
- 6. 271633.
- 7. 9504000. 8. 327040000.

- 9. 92438 lbs. 8 oz. 2 dr. 1 scr. 13 grs.
- 10. 1698728602536.
- 11. 78990 bushels.
- 12. \$64.97. 13. 9032 yds. 3 qrs. 2 ns.
- 14. 1037957601.5.
- 15. \$16444.9602.

#### Exercise 22.

42.000	AND WAS
1. \$34736.8421.	10. ·578 oz.
2. \$30634.9206.	11. 503.
3. 3308 dys. or 9 yrs. 201 dys	. 12. 250 lbs.
4. \$32.	13. 10.157.
<b>5. \$137.</b>	14. 2 bush. 1 pk. 1 gal. 2 qt
<b>6.</b> \$108.	
7. \$9.	15. 1898 385.
8. \$29.	16. 267 days 748295 hours.

17 pts.
15. 1898 <del>189</del> .
16. 267 days 718334 hours.
, , , , , , , , , , , , , , , , , , , ,
18m 23.
14. 0331632.
15. 475347 hhds.
16. \$6750.
17. 11 <del>41</del> .
18. 58 acres.
19. \$0.501.
20. \$37.
21. 3 lbs. 0.oz. 14 dwt. 13} grs.
22. 29 acres 0 roods 21 per.
23. 14 yds.
24. 15 lbs. 4 oz. 1 dwt. 14 grs.
25. \$3890·38 <del>2</del> .
26. 1032694.
0
<b>27.</b> 16800. <b>28. \$360·15.</b>
29. \$247.95.
30. \$132082.
<b>31</b> . 169·49.
<b>32.</b> \$79.99\frac{7}{2}.
33. \$59.85.
<b>34.</b> \$532·12\frac{1}{2}.
<b>==</b>
35. COCCCCCCCIX.
<b>36.</b> ·56218 <del> </del> .
**
<b>37.</b> 1869696969.69.
<b>89.</b> \$21·1433.
40. 236 <del>1</del> 9.

#### EXERCISE 40.

	CXERCISE 4	ŁU.	12	à-
1. \$4688.16.7g. 2. 27536 miles 1 fur. 21 0 yds. 1 ft. 6 in. 8. 96.	per.	202221	octenary 33 quinar 1·98275,	and y.

6. LXXMXCDXXIII and CCXXXMVDLXVII.

7. 277200.

8. See XLVIII Recapitulation. 13. 27. Sec. I., page 57.

9. 642762977065601.1.

15. 742000000905000078014.0000087200011.

16. Seventy-one trillions three [18.  $2^5 \times 5^2 \times 3 \times 23$ . dred millions two hundred thousand four hundred and |20. 011436. one, and seventy thousand 21. 16383. four hundred and two tril- 22. 4096. lionths.

One hundred and thirty-four quadrillions nine hundred 24. 336960. one billions one hundred thousand and one hundred, and two hundred million 26, 1023 and 512. twenty thousand and two 27. 99472 trillionths.

Four quadrillions seven hun- 29. 722487-0873859. dred trillions twenty thou- 80. 65 lbs. 7 oz. 0 drs. 1 scr. hundred and seventy-eight hundredths of trillionths.

17. £2272 0s. 31d.

11. See Table, page 125.

12. \$2689.513.

14. See Recapitulation XLVIII page 57.

hundred billions one hun- 19. 87 ft. 1' 1" 3" 0" 10"" 8""" 10""" 10""""

23. 11 acres 8 rds. 7 per. 19 yds. 0 ft. 130 in.

trillions one hundred and 25. Child's share, \$179.41 ; woman's, \$358.82 4; man's \$1794·12-4.

28. 48350 8979694.

sand and seven, and two 31. 1, 2, 4, 7, 8, 14, 19, 28, 38, 56, 76, 133, 152, 266, 532, 1064.

32. 82 40 yards.

#### EXERCISE 63.

\$\frac{2}{5}, \frac{1}{100}, \frac{1}{20}, \frac{1}{25}, \text{ and } \frac{4}{600}.
 \$\frac{5}{2}, \frac{1}{25}.

4. 136.

5. Gave away 20 and kept 11. 14. 1 and 1359.

6. 147.

8. Longer part 72 feet and 17. 488. shorter part 64 feet.

10. 14 80 and 8%. 11. \$134.15\$. 12. \$28387·061.

13. 311% bushels.

15. 214 bushels.

16.

18, 5% and 23%.

9. 105843 acres; \$13219.683. 19. \$1333.33 or 30 of the whole.

#### EXERCISE 77.

1. .8.

2. 1.4445566778.

3. 4 days 17 hours 55 min, 30 sec.

4. 18488,

5. 156.85931270094.

6. '739157196 of a mile.

7. 16 sq. ft. 10483 inches.

8. 1 acre 3 roods 13 per.22 yds,

2 qts.

131 grs.

14 grs.

l per.

ITS.

and

,	1		μ
9.	1118 and 170.		1- 13 - 18 18 - 18 1 1 1 1 1 1 1 1 1 1 1 1 1
	and the state of t	14.	13.6169633.
10.	26-7837428671.	15.	3, 3, 1, 4, 1, and 9.
-		12	
	71.86193.		476-65028119.
	11.546 oz.	17.	9. , w <sup>2</sup> . 2
19.	75% yards.	- 35	- m by g
-1	Exec	IST	<b>78.</b>
•	702000007030017-000000000	400	00 7 e
	w	<b>&amp;</b> UU	Out of the state of
8.	1017116666.6.		20790, 6 HR SHIBSON
	23.		13751·12 and 2049151.
	10,38.47		66. Let the state the the
6.	5044 bricks.	13.	1 day 23 hours 24 min. 3414
7.	111 sq. ft. 0' 9" 4" 4" 8""	3	seconds.
	Bunn.		19860 lbs. 2 os. 91 drs.
			<b>\$168.75.</b>
9.	12225 bush 2pks 0 gal 2 qts.	16.	9, 58, \$298, and #3180.
17.		000	00, 00000000704, 0000704,
	7.04. WES des 124.	A se	de the forest of the second
			. 13450 <del>  78</del> .
19.	Man's share=266 0s. 41d.,		1340621 lbs. or 134061 gals.
١.	woman's =£35 0s. 21d.,	26.	8295.59
474	child's = £11 05.04d.	27.	247345 W7 Edg. 11 1 . 2. 2
BO.	1904080		644. cores
MI.	1, 2, 3, 4, 5, 6, 9, 10, 11, 15,		decimal for hear of the state of
- 1	18, 20, 25, 27, 30, 36, 45,		29×3×5.
20	50, 54, 60, 75, 81, 90, 100,		\$45.59.
48.	108, 135, 150, 162, 180,	40	\$90.96 <del>11</del> .
1 12	225, 270, 300, 324, 405, 450, 540, 675, 810, 900,		. 3·185988.
	1350, 1620, 2025, 2700,		215947.
11 11	4050, 8100.		. \$21588·90.
92	4419	OF	\$142.8248.
22	Lunar month=29 days 12	88	293.
#U.	hours 44 min. 3 seconds.	89	1470, 1818, 1978, 4600,
		100	. maio, maio, majo) 3310)
	Solar year=365 days 5		1818, 1818, 1218.

30. 31. 32. 33. 34. 35.

1.

2.

12.

13. 14.

#### EXERCISE 89.

1.	2:3.	 4. Greatest 21:27;	least 9:13
2.	2:3. \$479.30§.	· · · · · · · · · · · · · · · · · · ·	
•	127.00	5 57-10055566187	2492

6. 53ee3 7787 dwodenary, 12014313 410042 quinary, and 76010 12574 undenary.

7. 5.57052 oz.

8. 3 yds. 3 grs. 0 ns. 011 in.

9. \$2962.70.

10. 1 bush. 2 pk. 0 gal. 1 qt.

23:11; 6:7 and 88:176; 19. 5034. 1173:616.

12. 39 per cent.

13. 2436.

23. 764876837 nonary; 10011110101000011001111010000 binary; 11146453021 septenary.

24. 188100.

25. 80199.

26. 48.

27. 415-471137804. 28. \$53-5966.

14. 10 13. - £2 1 ?id. nearly.

16. 3 & a. s.

20. 026856599989+.

21. .0778.

22. 4.32958 miles.

29. 1, 2, 3, 4, 5, 6, 7, 9, 10, 12, 14, 15, 18, 20, 21, 25, 28, 30, 35, 36, 42, 45, 50, 60, 63,

70, 75, 84, 90, 100, 105, 126, 140, 150, 175, 180, 210, 225, 252, 300, 315, 350, 420, 450, 525, 630, 700, 900, 1050, 1260, 1575,

2100, 3150, 3600.

30. \$5.04.

31. Each man's share, \$325.99123; each woman's, \$88.90144; each child's, \$25.40128.

**32.** 126, 5<sub>1</sub>87, 2<sub>16</sub>.

33. 3 yds. 2 ft. 83 in.

34. 104:5.

85. 71 miles 5 fur. 34 per. 3 89. 200. yards.

37. 2.65g. 38. 70 goats.

#### EXERCISE 92.

1. 7020400000, 7.0204, 70.204, 5. 5: 7; 9:13; 54:221. :0000070204, 7020.4, and 6. \$2070.3593. ·00000070204.

2. 6704866.561.

8. £399 19s. 5}\$\$\$d.

a to to be selected to be a selected to 4. 846.372095763.

.0007449164; 744916.4.

14. Binary 63 and 32, Quaternary 4095 and 1024, Senary 46655 and 7776, Octenary 262143 and 32768, Duodenary 2985983 and 248832.

7. They have none.

8. \$27431·31\frac{1}{2}.

9. 18, 799988, 38, and 17.

10. 2361 11. 126 days.

12. 744916400000; 7.449164; .00000000007449164; 7449.164;

15. 1, 2, 3, 4, 6, 8, 9, 12, 16, 18, 24, 27, 32, 36, 48, 54, 64, 72, 96, 108, 144, 192, 216, 288, 432, 576, 864, 1728.

16. 720720.

17. 79.789966677748855.

18. \$127.98.

19. 21.19117.

gals.

#### Exercise 165.

60

62 63 64

65 66 67.

84.

85. 86. 87. 88.

89.

90. 97. 98. 99.

103

1. 70\00000000010-0000004200006. 2. A, \$1639·32\frac{1}{2}; B,\$1528·21\frac{1}{2}; O,\$1437.31\frac{1}{2}; D,\$1584·95. 3. 12\frac{1}{2}. 4. \$1497803819·4444. 5. 83160. 6. 361 y'rs. 10 m'ths. 25 days. 7. 40·38. 8. 33943 lbs.4 oz.8dwt.14\frac{1}{2}grs. 9. 2. 10. 129\frac{1}{2}. 11. 3. 12. 24.	20. 5456640. 21. They have none. 22. A, \$3492.06; B, \$4761.91; C, \$6746.03. 28. A, £16714; B, £18924; C, £934s. 24. 275 hours. 25. LXXMVCMXXXVIII and XVMMCDXCVMMMDCLXXIX. 26. 1st gets 792 loaves; 2nd, 594; 3rd, 924. 27. 72, 18 and 54 lbs., or 24, 96,
13. A, \$384.47; B, \$291.07; O, \$221.89. 14. 13514 lbs. 15165229. 16. 530.00121864500. 17. \$7854.29. 18. 268. 19. 81000.	27. 72, 18 and 54 los., or 24, 56, and 96 lbs. respectively. 28. \$3725.764. 29. 24010.23. 30. \$4803.5004. 31. 5739.29 yds. Gain 253 per cent. 32
34. 2·886057; 1·290035; 3·0 4·698970; 2·182129; 0·903 35. t8·t2. 36. 84 years. 37. 66·80578 times. 38. 22992700·72992700. 39. \$5·482.	40. \$460.0034. 41. 5 yrs. 8 mos. 5 days. 42. Amount \$1409.07. Compound Int. \$595.36. 43. 10 months 18 days.
44. A, \$571.9676; B, \$554.8676 and E, \$1078. 45. \$1372.02898. 46. 1. 47. 117042723742437 octenary. 48. 01 and 012345679. 49. One quadrillion three hundred billions fifty million and six thousand, and seven hundred million eighty thousand and nine trillionths.	Seven trillions six hundred billions two hundred and ninety millions thirty-four thousand and seven, and sixty-seven millions four hundred thousand two hundred and nine quadrillionths.

91;

23 ;

and

CIX.

2nd,

, 96,

251

913 ;

Com-

5275;

ndred

d and

hirty-

seven.

llions

usand

nine

per acre.

```
52. 7119 A.
                                 68. 8.5318452.
58. 144.
                                  69. ·019156118.
54. 3517.
                                 70. 2781.848813156689829957.
55. 8# days.
56. $2469.71.
                                 71. 157.036 feet.
57. 47, 37, and 274.
                                 72. 85 spirits, 35 water.
58. Each man had 60; A caught | 73. 422.32.
       50, B 60, C 70.
                                 74. 70 and 14.
59. 191 and 17763.
60. 44.997 years.
                                 75. 223.82460585.
61. A,$1556.953; B,$1160.953;
                                 76. 5.32341.
       O, $973.083.
                                 77. 58 and 28.
62. 1, 2, 4, 1429, 2858, 5716.
                                 78. 156240.
68. 236.
                                 79. 30401.
64. Man's share = $919.144$,
                                 80. 2281: 1617.
       Woman's = $459.572+,
                                 81. 3 and 11, or 4 and 11, or 5
       and child's= $153.197.
                                        and 11, &c.
65. 24.
                                 82. 147.
66. $21.03.
                                 88. 5 fr minutes past 1 o'clock.
67. Greatest 9:18; least 10:19;
       comp. ratio 21: 247.
84. 6.585461; 3.502675; 5.187521; 2.113509; 0.196295;
       1.969276.
85. $4·314.
                                 91, 1, 84, 167, 244, 321, 40.
86. X $672 and Y $1120.
                                 92. 7.
                                 93. Apple 2d., pear 3d.
87. 17.
88. 4321.
89. 183 lbs. at 4d.; 183 lbs. at 95. $275.
       6d.; and 74% lbs. at 8d. 96. $124 and $1564.
90. 10, 22, 26.
97. 11000000000011.0000000011
98. $3649.3932.
                                 101. 117.
99. 2^8 \times 3^2 \times 7 \times 11.
                                 102. 624 gal., 833 gal., and 146
100. 281.
103. A, £194 16s. 1\flackfd.; B, £129 17s. 4\flackfd.; C, £97 8s. 0\flackfd.;
       D. £77 188. 537d.
104. $1230.338.
                                 111. 1st, '46 inches; 2nd, '57
105. 10 hours.
                                        in.; 3rd, 82 in.; 4th,
                                        3.149 in.
106. 41 years.
                                 112. 71.117.
107. 4.629 days.
108. £4 16s.
                                 113. $2019.651 ; $4871.803
109. 4413.
                                        $4815.805; $6467.739
110. 1422·2 lbs.
                                        $1825.
                                 114. 1st 30.0 yrs; 2nd 56.827 yrs.
115. 1st, $920.20; 2nd. $2760.60; 3rd, $5521.20.
116. Paid each workman $28.663; 1st company cleared 8725
       acres; 2d company, 771 acres; cost of clearing, $8-46
```

118. \$2340·00. 119, 132 days. 120. A, \$2180; B, \$1635; C, tient 32414.56. 124. 491 Tan yds. 125. \$214. 126. 1st 175 yrs.; 2nd 41.914 yrs. 127. 1014 perches. 128. 111104. 129. 9, 27, 81, 243, 729, 2187, 130. 91. W ...

Day of the state

ALCO TON OUT A MET TO ME

117. 15 and 11. | 181. 8.04 in. 9.534 in. 12.426 in. and 30 inches. 132. 51 of each, rem. £11%. 133. \$200. 134. 19 per cent. 135. \$1388.888. 136. 1s. 9d., 1s. 2d., and 7d. 137. A, \$25; B, \$25; C, \$50; Ď, \$160. 138. ·057. 139. 767; 162770; 1191; 454; 140. 96; 178. 141. \$8918; \$10717; \$14318; and \$179.9. 143. 173, 324, 481, and 631; 35 and 85905.

144. 361 days.

Application of the second

111

Au

ad

88 one wit

abs inte

me the

tion Wor by 1 to a San Edu orig sary

the

as a befo Trea tial yet

the: atlo the

it to ject

areg Cour

mato It co mak ledge

THE END. 

May Conference and the second of the second

#### Opinions of the Press on the National Arithmetic.

This is one of Lovell's series of School books, a series which we hope some This is one of Lovell's series of Senool books, a series which we nope some day to see introduced into all our Canadian Schools. It has been prepared expressly for these schools by the Mathematical Master of the Upper Canada Normal School. From the brief examination we have been enabled to give it, we are inclined to think it will give a more thorough knowledge of the science of numbers than any other Arithmetic we remember, and we hope Canadian teachers will give it a trial.—Montreal Gazette.

It is the production of one of our most useful and energetic teachers, and it shows a thorough knowledge of the subject, and adaptation to the antis of the country. Mr. Sangster, by this volume, has supplied a want

wants of the country. Mr. Sangster, by this volume, has supplied a want long felt; and it augurs well for the future teachers of our children that the Author of such a work as this, is Mathematical Master in our National Normal School.—Ottawa Citizen.

: 16!

Normal School.—Ottawa Citizen.

We think it is admirably adapted for, and should be speedily introduced into, all our Canadian schools.—Carleton Place, C. W., Herald.

This Arithmetic is not only infinitely better adapted to the wants of this country than any other in use, but the simplicity of its rules, the practical illustrations of the theory and practice of arithmetic in the many original problems, give it a stamp of nationality highly creditable to the Author. It is divested of all the useless, lumbering material to be found to a greater or less extent in those heretofore in use in this country. The problems possess an eminently practical character—and by that very adaptation to our wants, they are the more interesting. The learner, instead of covering his slate with figures, with a vague and confused notion as to what they all mean, works a problem in this arithmetic feeling at once that he is doing a useful and interesting work, and watches the result with a degree of interest that must help to make his schoolboy days cheerful and pleasant.—Markham, C. W., Economist.

We hail with much satisfaction the appearance of this work, rendered absolutely necessary by the recent introduction of the Decimal Currency into Canada. For a long time the want of a Canadian Treatise on Arithmetic, combining the above mentioned system with the application of the Modern Scientific methods of analysis and formulæ, to the elucidation of the various rules, was felt. Dr. Ryerson, conscious that such a work was needed, requested the Author to adapt the Arithmetic published by the Irish Board of Education, to the Decimal Currency of Canada, and to abbreviate some of the tedious reasons for the rules there given. Mr. Sanester in complying with the request of the Chief Superintendent of

to abbreviate some of the tedious reasons for the rules there given. Mr. Sangster in complying with the request of the Chief Superintendent of Education, transcribed ten or fifteen pages from the commencement of the original work, but finding so many "alterations and improvements" necessary, "abandoned "the design and determined to write a new Treatise on the subject. The admirable volume which now lies before us is the result of that determination.—From what fame says of Mr. Sangster's capabilities as an excellent Teacher, and an accomplished Mathematician, the volume as an excellent Teacher, and an accomplished Mathematician, the volume before us has not exceeded our expectations, though it surpasses every Treatise on the subject which has yet come into our hands in three essential requisites, namely: Methodical arrangement of matter; conciseness yet comprehensiveness in the demonstration of the various rules; and the immense practical utility which it possesses by the number of examination questions given at the end of each section to test the knowledge of the student as he progresses. These advantages must inevitably cause it to supersede in a very short time those spurious Treatises on the subject at present in existence throughout the Province: for this reason we are glad the work is entitled the "NATIONAL ARITHMETIC."—Brant, C.W., County Herald. County Herald.

Mr. Sangster's Book is the best going—has no competitor—cannot be matched—positively overflowing with matter. We highly recommend it. It combines beautiful printing, stout binding, with all that is wanted to make a young person have a complete storehouse of mathematical know-ledge at his fingers ends. No book we have yet seen, on this indispensable branch of knowledge, can compare with it.—Cayuga, C. W., Sentinel.

The undersigned, having long felt it would be highly desirable to have a SERIES OF EDUCATIONAL WORKS, PREPARED AND WRITTEN IN CANADA, and adapted for the purposes of Canadian Education, begs to call attention to the Text Books with which he has already commenced this series. These works have met with a very general welcome throughout the Province; and the Publisher feels confident that the eulogiums bestowed upon them are fully merited, as considerable talent and care have been enlisted in their preparation.

#### The following Text Books have already been published:

1. LOVELL'S GENERAL GEOGRAPHY, with 51 Colored Maps, 113 Beautiful Engravings, and a Table of Clocks of the World. By J. George Hodgins, LL.B., F.R.G.S.

[This Book is especially adapted for, and worthy of introduction into every College, Academy, and School in the British Provinces.

Parents should see that it is in their children's

2

R

2. NATIONAL ARITHMETIC, in Theory and Practice, adapted to Decimal Currency. By J. H. SANGSTER, Esq., M.A.

3. KEY to ditto. By the same.

4. ELEMENTARY ARITHMETIC, in Decimal Currency. By the same.

NATURAL PHILOSOPHY, PART I., including Statics, Hydrostatics, &c., &c. By the same.

6. GENERAL PRINCIPLES OF LANGUAGE; or the Philosophy of Grammar. By Thomas JAFFREY ROBERTSON, Esq., M. A.

7. AN EASY MODE OF TEACHING THE RUDIMENTS OF LATIN GRAMMAR TO BEGINNERS. By the same.

8. CLASSICAL ENGLISH SPELLING BOOK. By G. G. VASEY.

9. ENGLISH GRAMMAR MADE EASY. By the same.

10. THE A-B-C SIMPLIFIED, and Reading Made Easy to the capacity of Little Children. By the same.

11. STUDENT'S GUIDE TO ENGLISH GRAMMAR, or the Way to speak and Write Grammatically. By the Rev. J. G. Armstrong, M. A.

12. ELEMENTS OF ELOCUTION. By J. BARBER, M.R.C.S.

13. OUTLINES OF CHRONOLOGY. By Mrs. Gordon.

14. BRITISH AMERICAN READER. By J. D. Borthwick, Esq.

#### The following are in Press:

- 15. ELEMENTARY TREATISE ON ALGEBRA. By J. H. SANGSTER, Esq., M. A.
- 16. NATURAL PHILOSOPHY, PART II. By the same 17. KEY to the Elementary Arithmetic. By the same.

18. EASY LESSONS IN GENERAL GEOGRAPHY. By J. GEORGE Hodgins, LL.B., F.R.G.S.

19. BOOKKEEPING by Single and Double Entry. By John G. Dinning, Esq.

# The following Text Books, printed from New Stereotype plates and in good Bindings, have also been published:

20. The FIRST NATIONAL BOOK OF LESSONS.

21. The SECOND ditto ditto
22. The THIRD ditto ditto
23. The FOURTH ditto

24. The FIFTH ditto ditto
25. PINNOCK'S IMPROYED EDITION OF GOLDSMITH'S
HISTORY OF ENGLAND. By W. C. TAYLOR, LL.D.,

T.C.D.

26. LENNIE'S ENGLISH GRAMMAR.

27. KIRKHAM'S ENGLISH GRAMMAR.

28. FRENCH WITHOUT A MASTER.

29. FRENCH GENDERS taught in Six Fables.

In preparation:—A SCHOOL HISTORY OF CANADA, AND OF THE OTHER BRITISH PROVINCES, with Illustrations. By J. Grouge Hodgins, LL.B., F.R.G.S.

Messrs. R. & A. MILLER, Montreal and Toronto, are the General Agents in Canada for the sale of Lovell's Series of School Books.

## Extracts from Opinions on Lovell's General Geography.

Your work on Geography supplies a want which teachers have long felt and complained of.—Rev. Dr. Leach.

most excellent and in all respects suitable school book.—
Rev. Dr. Irvine.

Such portions as I have paid particular attention to, appear to me to be very accurate.—Rev. W. Snodgrass.

It will become a necessity in our Seminaries of Education.—
Rev. Dr. Bancroft.

The work under review seems to merit the highest commendation.—Rev. J. Ellegood.

The plan and manner of execution, are both admirable.—Rev. W. Scott.

The fairness and impartiality with which the different countries are described will commend it to general use.—Rev. E. J. Rodgets.

It seems to me just what is needed, suited to the requirements of the country in its matter, form, and price.—Rev. J. B. Bonar.

olored ocks of

sirable

ARED

oses of

Books works

vince :

stowed

d care

of introchool in ildren's

adapt-

y. By

cs, Hyne Phiertson,

TS OF me. G. G.

to the

he Way

.c.s.

r, Esq.

J. H.

I think the publication of the work ought to be regarded as a matter of sincere congratulation to the country at large.

—Rev. W. S. Darling.

Where all is excellent it is difficult to particularize, but I may state that I consider the introductory part deserving of especial commendation.—Rev. Dr. Brooke.

The classification appears to be faultless, the definitions concise and lucid, and the information given in regard to the derivation and pronunciation of proper names is very valuable. It is indeed multum in parvo, and will doubtless become the standard Geography of our Schools.—Rev. I. B. Howard.

I am really delighted that at last a School Geography, almost perfect, is provided for the youth of the British North American Provinces.—Rev. John Carry.

Your Geography is all that can be desired; and, after a thorough examination, I am convinced, from its merits, it will at once be adopted in all our schools. It is a marvel of cheapness, admirable in plan, and a fine specimen of what can be done by an enterprising and liberal publisher. We shall at once introduce it into our School, as its want has long been felt.—Rev. H. J. Borthwick, Principal of the County of Carleton Senior Grummar School.

I hail it as the best Geography extant for our Canadian Schools. I can give no better proof of my appreciation of its merits than by introducing it immediately as the standard textbook in our Academy.—J. Douglass Borthwick, Principal of Huntingdon Academy.

It is my intention to adopt at once this Geography as a textbook in the Grammar School department of this Institution.— Rev. S. S. Nelles, President of Victoria College.

I shall be most happy to recommend it to the schools in my superintendency, as well as to the heads of families, and hope it will be patronized as extensively as it deserves.—Rev. J. Gilbert Armstrong.

When it comes to be known by the public, I should think it must command a very extensive, if not universal, circulation in the Schools of British North America.—Rev. John Cordner.

The plan of your School Geography is excellent.—Rev. Dr. Leitch, Principal of Queen's College.

So far as I can judge, Lovell's General Geography is well adapted to our Canadian Schools.—Rev. A. J. Parker.

No existing work can be held to excel it .- Rev. A. de Sola.

Fight of the section of the section of the

OKS.

garded as at large.

but I may f especial

the deriable. It he stand-

y, almost American

it will at neapness, done by nee intro-Rev. H. Grummar

Janadian on of its ard text-ncipal of

s a texttution.—

ols in my d hope it . Gilbert

ld think culation rdner.

Rev. Dr.

is well

Sola.

