


**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1981

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

- Coloured covers/
Couverture de couleur
- Covers damaged/
Couverture endommagée
- Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée
- Cover title missing/
Le titre de couverture manque
- Coloured maps/
Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur
- Bound with other material/
Relié avec d'autres documents
- Tight binding may cause shadows or distortion
along interior margin/
La reliure serrée peut causer de l'ombre ou de la
distortion le long de la marge intérieure
- Blank leaves added during restoration may
appear within the text. Whenever possible, these
have been omitted from filming/
Il se peut que certaines pages blanches ajoutées
lors d'une restauration apparaissent dans le texte,
mais, lorsque cela était possible, ces pages n'ont
pas été filmées.
- Additional comments:/
Commentaires supplémentaires:

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured pages/
Pages de couleur
- Pages damaged/
Pages endommagées
- Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- Pages detached/
Pages détachées
- Showthrough/
Transparence
- Quality of print varies/
Qualité inégale de l'impression
- Includes supplementary material/
Comprend du matériel supplémentaire
- Only edition available/
Seule édition disponible
- Pages wholly or partially obscured by errata
slips, tissues, etc., have been refilmed to
ensure the best possible image/
Les pages totalement ou partiellement
obscures par un feuillet d'errata, une pelure,
etc., ont été filmées à nouveau de façon à
obtenir la meilleure image possible.

**This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.**

10X	14X	18X	22X	26X	30X	
						<input checked="" type="checkbox"/>

12X 16X 20X 24X 28X 32X

med here has been reproduced thanks
osity of:

National Library of Canada

appearing here are the best quality
considering the condition and legibility
al copy and in keeping with the
tract specifications.

ies in printed paper covers are filmed
with the front cover and ending on
e with a printed or illustrated impres-
back cover when appropriate. All
al copies are filmed beginning on the
ith a printed or illustrated impres-
nding on the last page with a printed
d impression.

orded frame on each microfiche
n the symbol → (meaning "CON-
or the symbol ▽ (meaning "END"),
pplies.

s, charts, etc., may be filmed at
duction ratios. Those too large to be
uded in one exposure are filmed
n the upper left hand corner, left to
p to bottom, as many frames as
he following diagrams illustrate the

L'exemplaire filmé fut reproduit grâce à la
générosité de:



Bibliothèque nationale du Canada

Les images suivantes ont été reproduites avec le
plus grand soin, compte tenu de la condition et
de la netteté de l'exemplaire filmé, et en
conformité avec les conditions du contrat de
filmage.

Les exemplaires originaux dont la couverture en
papier est imprimée sont filmés en commençant
par le premier plat et en terminant soit par la
dernière page qui comporte une empreinte
d'impression ou d'illustration, soit par le second
plat, selon le cas. Tous les autres exemplaires
originaux sont filmés en commençant par la
première page qui comporte une empreinte
d'impression ou d'illustration et en terminant par
la dernière page qui comporte une telle
empreinte.

Un des symboles suivants apparaîtra sur la
dernière image de chaque microfiche, selon le
cas: le symbole → signifie "A SUIVRE", le
symbole ▽ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être
filmés à des taux de réduction différents.
Lorsque le document est trop grand pour être
reproduit en un seul cliché, il est filmé à partir
de l'angle supérieur gauche, de gauche à droite,
et de haut en bas, en prenant le nombre
d'images nécessaire. Les diagrammes suivants
illustrent la méthode.





OUTLINES
OF
CLASSIFICATION OF PLANTS,

BY

D. P. PENHALLOW, B.Sc., F.R.S.C.,

Professor of Botany, McGill University,



MONTRÉAL:
E. M. RENOUEF, Publisher.
1895.

Entered according to Act of Parliament of Canada in the Year one thousand
eight hundred and ninety five by D. P. PENHALLOW, in the Office of the Minister
of Agriculture.

OUTLINES OF CLASSIFICATION.

Synopsis.

RANCH I.—Protophyta.

CLASS 1.—MYXOGASTRES, (Slime Moulds.)

- Orders*—Peritrichae, Columellifera, Lithodermæa, Calotrichæa.

CLASS 2.—SCHIZOPHYCE, (Fission Fungi)

- Orders*—Myxobacterae, Cystiphore, Nematogene.

From the nature of the organisms included in the protophyta, it is improbable that many of them will ever be found in the fossil state, although Renault claims to have recognized several species of bacteria in the tissues of carboniferous plants.

RANCH II.—Thallophyta.

CLASS 3.—ASCOMYCETES, (Sac Fungi)

- Orders*—Perisporiaceæ, Tuberaceæ, Pyrenomycetæ, Discomycetæ, Uredinaceæ, Ustilaginæ, Sphaeropsidæ, Melanconicæ, Hyphomycetæ.

Fossil representatives occur in the carboniferous and later formations, chiefly upon leaves and in lignite.

CLASS 4.—BASIDIOMYCETES, (Higher Fungi.)

- Orders*—Gasteromycetæ, Hymenomycetæ.

Fossil representatives occur in the carboniferous, *Grillidea carofermii*, *Gastromyces farinosus*, etc.

CLASS 5.—CHLOROPHYCE, (Green Alge.)

- Orders*—Protococcoideæ, Conguatae, Siphoniacæ, Conervoideæ.

Fossil representatives, in the form of siliceous diatom valves, are especially abundant in the Tertiary and Quaternary, often forming extensive deposits known under the name of infusorial earth.

CLASS 6.—PHILOPHYCE, (Brown Alge.)

- Orders*—Phaeosporen, Dictyotæ, Fucoideæ.

The earliest fossil representatives of the brown algae occur in the Upper Silurian whence they pass into the Lower Devonian. They appear in these early formations as plants of great size (*Arenophyton*), indicating that they must also have flourished at much earlier periods, but the remains of the vegetation which existed prior to the Upper Silurian are now represented only by a carbonaceous residue in the form of graphite. Philophyceæ are found more or less abundantly in all the later formations, and the genus *Fucus* is well defined in the Cretaceous.

CLASS 7.—COLEOCHILIFÆL.

- Order*—Coleochiliacæ.

No fossil representatives known.

CLASS 8.—RHODOPHYCE, (Red Alge.)

- Order*—Florideæ.

Fossil representatives from the Silurian upward, and especially abundant in the Cretaceous.

CLASS 9.—CHAROPHYCE, (Green Alge.)

- Order*—Characeæ.

Represented by many species in the lower Cretaceous, in the Tertiary and Quaternary.



Synopsis.—(Continued.)

NOTES.

Branch III.—Bryophyta.

CLASS 10.—HEPATICÆ.

- Orders—Jungerniaceæ,
Ricciaeæ,
Anthocerotææ,
Marchantiææ,

Found only in recent formations—Tertiary and Quaternary.
Tarachnites.

CLASS 11.—MUSCIÆ.

- Orders—Sphagnacæ,
Andreacæ,
Phæsacæ,
Bryacæ.

Fossils found only in the Tertiary and Quaternary. *Gymnomium*, *Sphagnum*, *Hypnea*.

Branch IV.—Pteridophyta.

CLASS 12.—FILICINÆ.

- Orders—Filicæ,
Salviniaeæ,
Marsiliææ,
Ophioglossacæ,
Marattiææ.

Fossil representatives numerous from the Devonian (*Parkeriostrophia*, *Nuropteris*, *Sphenopteris*, etc.) through the carboniferous to recent formations.

CLASS 13.—EQUSETINÆ.

- Orders—Equisetacæ,
Calamiteæ,
Annulariææ,
Astrophylliteæ.

Fossil representatives numerous from the Devonian (*Calatites*, *Astrophyllites*, etc.) and throughout the carboniferous.

CLASS 14.—SPHENOPHYLLEÆ.

- Order—Sphenophyllacæ.

Fossil representatives in the Silurian-Cambrian and upward.

CLASS 15.—LYCOPODINÆ.

- Orders—Iycopodiacæ,
Psilotacæ,
Selaginellacæ,
Isoetacæ.

Fossil representatives numerous and often very large, from the Upper Silurian (*Ptychophyton*, etc.) through the Devonian (*Leptodendron*) and Carboniferous.

Branch V.—Spermaphyta.

CLASS 16.—GYMNOSPERMÆ.

- Orders—Cycadacæ,
Conifereæ,
Taxaceæ,
Gnetacæ.

Fossil representatives very numerous from the Devonian (*Dadoxylon*, *Cordaites*, etc.) and through the Carboniferous and more recent formations.

CLASS 17.—ANGIOSPERMÆ.

- Sub-Class 1.—Monocotyledons.

- Sub-Class 2.—Dicotyledons.

Orders—numerous.

Fossil representatives are very numerous as leaves, lignite, &c., from the Permian upward, becoming more numerous in recent formations.



BRANCH I.—*Protophyta.*

NOTES.

Unicellular Plants often forming more or less extensive gelatinous plasmodia by coalescence.

CLASS I.—*MYXOGASTRES.* (Slime Moulds.)

Orders.

1. Peritrichiae.
2. Columellifera.
3. Lathodermiae.
4. Calotrichiae.

Asexual reproduction predominant.

CLASS II.—*SCHIZOPHYCEAE.* (Fission Fungi.)

Orders.

1. Myxobacteriaceæ.
2. Cystiphoreæ.
3. Nematogeneæ.

Asexual reproduction predominant.

Saprophytic plants devoid of a cell wall except in the spores. Reproduction through amoeboid swarm spores, asexual by simple fission, or sexual (?) (Goebel) by conjugation and the formation of plasmodia, from which asexual spores arise.

Parasitic or saprophytic plants sometimes forming gelatinous plasmodia. Reproduction wholly asexual by simple division of a mother cell, more rarely by endogenous spore formation.



BRANCH II.—*Thallophyta*.

Unicellular or multicellular plants, the latter often with a stem-like habit of growth.
Sexual generation predominant.

III. ASCOMYCETES. (Sac Fungi.)	IV. BASIDIOMYCETES. (Higher Fungi.)	V. CHLOROPHYCEAE (Green Algae.)	VI. PHYLLOPHYCEAE (Brown Algae.)	VII. COCCOCYCEAE
<i>Orders.</i>	<i>Orders.</i>	<i>Orders.</i>	<i>Orders.</i>	<i>Orders.</i>
1. Perisporaceae. 2. Tuberidae. 3. Pyrenomycetaceae. 4. Discoscytaceae. 5. Uredinae. 6. Ustilagineae. 7. Sphaeropsidaceae. 8. Melanconiae. 9. Hyphomycetaceae.	1. Gasteromycetaceae. 2. Hymenomycetaceae.	1. Protococcoideae. 2. Conglutinaceae. 3. Siphonine. 4. Confervaceae.	1. Phaeophoraceae. 2. Dictyotaceae. 3. Fucoidea.	1. Coleochaetaceae.
Asexual reproduction commonly predominant.	Asexual reproduction predominant.	Sexual reproduction often by conjugation with the formation of Zygospores. Asexual reproduction predominant.	Reproduction wholly sexual in Fucales; asexual in Phaeophora and Dictyotaceae.	Asexual reproduction by swarm spores often predominant.
I. Perfect plant ♂. 2. Antheridia ♀.	I. Perfect plant ♂ 2. Antheridia ♀. 3. Spermatozoids.	I. Perfect plant ♂ or ♀ 2. Antheridia ♀. 3. Spermatozoids.	I. Perfect plant ♂ 2. Antheridia ♀. 3. Spermatozoids.	I. Perfect plant ♂ 2. Antheridia ♀. 3. Spermatozoids.
I. Perfect plant ♂. II. Archicarps ♀.	I. Perfect plant ♂ II. Oogonium.	I. Perfect plant ♂ or ♀ II. Oogonia.	I. Perfect plant ♂ or ♀ II. Oogonia.	I. Perfect plant ♂ II. Oogonium III. Oosphere (Ovum).
No clearly defined sexual reproduction known.		III. Oosphere (Ovum).	III. Oospheres (Ova).	III. Oosphere (Ovum).
3 x III. Ascus. 4 x IV. Spores.	IV. IV. Oospore.	IV. IV. Oospore.	4 x IV. Oospore.	5 x V. Swarm spores.
	5 x V. Swarm spores.			



BRANCH III.—Bryophyta.

NOTES.

Thallophyta.—(Continued.)

II. RHODOPHYCEAE.
(Red Seaweeds.)

Order.
Florideae.

IX. CHAROPHYCEAE.

Order.
Characeae.

Asexual reproduction often predominantly by means of tetrasporidia.

Asexual reproduction often predominant by means of bulblets, or tubers, or special vegetative branches.

Perfect plant ♂ or ♀ or Spermogonium.

1. Perfect plant ♂
2. Antheridium ♀

Spermatia ♂
Perfect plant ♂ or ♀ or Procarp
Trichogyne.

3. Spermatozooids motile
1. Perfect plant ♂
II. Oogonium .

x III. Carpospores.
IV. Protomema.

I x IV. Oospore.
5 x V. Proembryo.

Cellular plants with roots, epidermis and stomata,
often with leafy stems.
Sexual generation predominant.

X. HEPATICAE.
(Liverworts.)

Orders.

1. Jungermanniaceae.
2. Ricciaceae.
3. Anthocerotaceae.
4. Marchantiaceae.

XI. MUSCIACE.
(Mosses.)

Orders.

1. Sphagnaceae.
2. Andreaeae.
3. Phascaceae.
4. Bryaceae.

Asexual reproduction often predominantly by means of tetrasporidia.

Asexual reproduction often predominant by means of simple separation of the vegetative axis by adventitious shoots or by gemmae.

Asexual reproduction predominant by the development of a protonema, more rarely through gemmae.

Perfect plant ♂ or ♀ or Spermogonium.

1. Perfect plant ♂ or ♀
2. Antheridium ♀

Spermatia ♂
Perfect plant ♂ or ♀ or Procarp
Trichogyne.

3. Spermatozooids motile
1. Perfect plant ♂ or ♀ .
II. Archegonium.

x III. Carpospores.
IV. Protomema.

II. Archegonium.
III. Oosphere (Ovum).
—
4 x IV. Oospore.

1. Perfect plant ♂ or ♀
2. Antheridium ♀

Asexual reproduction often predominantly by means of tetrasporidia.

Oospore Generation.
(Asexual Phase.)

Alternation of generations well defined.

Sporophyte Generation.
(Asexual Phase.)

Alternation of generations well defined.

5 x VI. Embryo rudimentary.
6 x VI. Sporogonium parasitic upon the sexual generation.

7 x VII. Spores with elaters.
8 x VIII. Protonema.

1. Perfect plant ♂ or ♀
2. Antheridium ♀

3. Spermatozooids motile
1. Perfect plant ♂ or ♀ .
II. Archegonium.

III. Oosphere (Ovum).

1. Perfect plant ♂ or ♀
II. Archegonium.

III. Oosphere (Ovum).

4 x IV. Oospore.
5 x V. Embryo rudimentary.

6 x VI. Sporogonium parasitic upon the sexual generation.

7 x VII. Spores.
8 x VIII. Protonema.

		XII.—FILICINA (Ferns.)	XIII.—EQUISETINA (Horse-tails)	
		A. Leptosporangiate.	B. Eusporangiate.	
		1. Homosporous.	2. Heterosporous.	Homosporous.
<i>Oospore</i> Generation. (Sexual Phase.) <i>Sporophyte</i> Generation (Asexual Phase.) Alternation of generations often well defined.		Orders. 1. Filices. Asexual reproduction wholly subordinate; when present, chiefly by extension and division of the main axis, more rarely by bulbs or by stolomiferous rhizoids, or even by apogamy.	Orders. 1. Salviniales. 2. Marsiliales. Asexual reproduction wholly subordinate.	Orders. 1. Ophioglossaceae. 2. Marattiaceae. Asexual reproduction subordinate, by means of adventitious buds on the roots.
I. Prothallus ♀ well formed. 2. Antheridium ♂. 3. Spermatozoids motile. 4. Prothallus ♀ well formed. II. Archegonia.		1. Prothallus ♂ well formed. 2. Antheridium. 3. Spermatozoids motile. 4. Prothallus ♂ well formed. II. Archegonia.	1. Prothallus ♂ well formed. 2. Antheridium. 3. Spermatozoids motile. 4. Prothallus ♂ well formed. II. Archegonia.	1. Prothallus ♂ well developed. 2. Antheridium ♂. 3. Spermatozoids motile. 4. Prothallus ♂ well developed. II. Archegonium ♂.
III. Oosphere (Ovum).		III. Oosphere (Ovum)	III. Oosphere (Ovum)	III. Oosphere (Ovum)
4 x IV. Oospore. 5 x V. Embryo rudimentary. 6 x VI. Normal plant. 7 x VII. Sporophyll. 8 x VIII. Sporangia. 9 x IX. Spores.		4 x IV. Oospore. 5 x V. Embryo rudimentary. 6 x VI. Normal plant. 7 x VII. Sporecarp. 8 x VIII. Sporangia and 9 x IX. Spores.	4 x IV. Oospore. 5 x V. Embryo rudimentary. 6 x VI. Normal plant. 7 x VII. Sporophyll. 8 x VIII. Sporangia and 9 x IX. Spores.	4 x IV. Oospore. 5 x V. Embryo rudimentary. 6 x VI. Normal plant. 7 x VII. Scales of sterile spike. 8 x VIII. Sporangia. 9 x IX. Spores with elaters.

ophyta.

Inet Vascular System.
(ptograms.)
ordinate.

NOTES.

XIV. SPHENOPHYLLEAE.

Heterosporous.

Order.

Sphenophyllaceæ.

Homosporous or Heterosporous,
the latter fossil only.

Order.

I. Lycopodiaceæ

Asexual reproduction sub-
ordinate, by means of axillary
bulbils, or by lateral budding
of underground tubers.

1. Prothallus ♀ or ♂ + rudimen-
tary.

2. Antheridium ♂

3. Spermatozoids motile.

4. Prothallus ♀ or ♂ + rudimen-
tary.

II. Archegonium ♀

III. Oosphere (Ovum).

—

4 x IV. Oospore.

5 x V. Suspensor.

6 x VI. Embryo rudimentary.

7 x VII. Normal plant.

8 x VIII. Sporangiferous leaf.

9 x IX. Sporangium.

10 x X. Spores.

XV. LYCOPODINÆ.

(Club Mosses.)

Homosporous.

Order.

I. Psilotaceæ.

Sexual reproduction pre-
dominant.

1. Prothallus ♀ + rudimen-
tary.

2. Antheridium ♂

3. Spermatozoids motile.

1. Prothallus ♀ + rudimentary.

II. Archegonium ♀

III. Oosphere (Ovum).

4 x IV. Oospore.

5 x V. Suspensor.

6 x VI. Embryo rudimentary.

7 x VII. Normal plant.

8 x VIII. Sporangiferous leaf.

9 x IX. Sporangium.

10 x X. Spores.

Heterosporous.

Orders.

I. Selaginellaceæ.

2. Isoetaceæ.

Asexual propagation sub-
ordinate, by division of the
main axis, rarely by apogamy.

1. Prothallus ♀ + rudimen-
tary.

2. Antheridium ♂

3. Spermatozoids motile.

1. Prothallus ♀ + rudimentary.

II. Archegonium ♀

III. Oosphere (Ovum).

4. Oospore.

5. Suspensor.

6. Embryo rudimentary.

7. Normal plant.

8. Sporangiferous leaf.

9. Microsporangia.

10. Macrospores.

IV. Oospore.

V. Suspensor.

VI. Embryo rudimentary.

VII. Sporangiferous leaf.

VIII. Normal plant.

IX. Macrosporangia.

X. Macrospores.



BRANCH V.—Spermaphyta.

True seed plants.

Sexual generation very subordinate, in the Dicotyledons becoming almost completely suppressed with respect to the general structure.

XVI.—GYMNOSPERMIA.

Carpels open, seeds naked.

XVII.—ANGIOSPERMIA.

Carpels forming closed seed vessels.

Heterosporous; prothalli two ⁺.

Orders.

1. Cycadaceae. 2. Coniferae.

3. Taxaceae. 4. Gnetaeæ.

Asexual propagation subordinate;
sometimes by tubers.

1. Prothallus rudimentary. ⁺
2. Antheridium rudimentary. (Pollen tube.)
3. Spermatozoids none. (Protoplasm of the antheridium.)

4. Prothallus (endosperm), formed before impregnation; parasitic upon the asexual generation. (Seeds albuminous.)
5. Archegonium well formed.
6. Oosphere (Ovum.)

7. Oospore.
8. Suspensor.
9. Embryo with 2 to several cotyledons.
10. Normal plant.
11. Anthophylla.
12. Microsporangia (Anthers.)
13. Microspores (Pollen.)

14. Oospore.
15. Suspensor.
16. Embryo with 2 to several cotyledons.
17. Normal plant.
18. Carpophyllum.
19. Macrosporangia (Ovules.)
20. Macrospore (Embryo sac.)

Heterosporous, the rudimentary prothalli often wanting.

1. Monocotyledons.

Orders numerous.

2. Dicotyledons.

Orders numerous.

Asexual propagation subordinate, rarely wholly replacing the sexual; varied, by means of runners, stolons, offsets, bulbs, suckers, tubers, and rarely by parthenogenesis.

1. Prothallus not developed.
2. Antheridium rudimentary. (Pollen tube.)
3. Spermatozoids none. (Protoplasm of the antheridium.)

4. Prothallus (endosperm) formed only after impregnation of the ovum; parasitic upon the asexual generation. (Seeds albuminous.)
5. Archegonium not formed.
6. Oosphere (Ovum. Germ cell.)

7. Oospore.
8. Suspensor.
9. Embryo with one cotyledon.
10. Normal plant.
11. Stamens.
12. Microsporangia (Anthers.)
13. Microspores. (Pollen.)

14. Oospore.
15. Suspensor.
16. Embryo with one cotyledon.
17. Normal plant.
18. Pistil.
19. Macrosorangiæ (Ovules.)
20. Macrospore (Embryo sac.)

