




**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**Photographic  
Sciences  
Corporation**

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503

15 28  
32 25  
36 22  
40  
18

**CIHM/ICMH  
Microfiche  
Series.**

**CIHM/ICMH  
Collection de  
microfiches.**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

10

**© 1987**

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- |  |  |
|--|--|
| <input type="checkbox"/> Coloured covers/<br>Couverture de couleur   | <input type="checkbox"/> Coloured pages/<br>Pages de couleur   |
| <input type="checkbox"/> Covers damaged/<br>Couverture endommagée  | <input type="checkbox"/> Pages damaged/<br>Pages endommagées   |
| <input type="checkbox"/> Covers restored and/or laminated/<br>Couverture restaurée et/ou pelliculée  | <input type="checkbox"/> Pages restored and/or laminated/<br>Pages restaurées et/ou pelliculées  |
| <input type="checkbox"/> Cover title missing/<br>Le titre de couverture manque   | <input checked="" type="checkbox"/> Pages discoloured, stained or foxed/<br>Pages décolorées, tachetées ou piquées   |
| <input type="checkbox"/> Coloured maps/<br>Cartes géographiques en couleur   | <input type="checkbox"/> Pages detached/<br>Pages détachées  |
| <input type="checkbox"/> Coloured ink (i.e. other than blue or black)/<br>Encre de couleur (i.e. autre que bleue ou noire)   | <input checked="" type="checkbox"/> Showthrough/<br>Transparence   |
| <input type="checkbox"/> Coloured plates and/or illustrations/<br>Planches et/ou illustrations en couleur  | <input type="checkbox"/> Quality of print varies/<br>Qualité inégale de l'impression   |
| <input type="checkbox"/> Bound with other material/<br>Relié avec d'autres documents   | <input type="checkbox"/> Includes supplementary material/<br>Comprend du matériel supplémentaire   |
| <input type="checkbox"/> Tight binding may cause shadows or distortion<br>along interior margin/<br>La reliure serrée peut causer de l'ombre ou de la<br>distorsion le long de la marge intérieure   | <input type="checkbox"/> Only edition available/<br>Seule édition disponible   |
| <input type="checkbox"/> Blank leaves added during restoration may<br>appear within the text. Whenever possible, these<br>have been omitted from filming/<br>Il se peut que certaines pages blanches ajoutées<br>lors d'une restauration apparaissent dans le texte,<br>mais, lorsque cela était possible, ces pages n'ont<br>pas été filmées. | <input type="checkbox"/> Pages wholly or partially obscured by errata<br>slips, tissues, etc., have been refilmed to<br>ensure the best possible image/<br>Les pages totalement ou partiellement<br>obscurcies par un feuillet d'errata, une pelure,<br>etc., ont été filmées à nouveau de façon à<br>obtenir la meilleure image possible. |
| <input type="checkbox"/> Additional comments:/<br>Commentaires supplémentaires:  |  |

This item is filmed at the reduction ratio checked below/  
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
12X	16X	20X	24X	28X	32X

ails  
du  
odifier  
une  
image

The copy filmed here has been reproduced thanks to the generosity of:




Library of Parliament and the  
National Library of Canada.

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol  $\rightarrow$  (meaning "CONTINUED"), or the symbol  $\nabla$  (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

La Bibliothèque du Parlement et la  
Bibliothèque nationale du Canada.

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole  $\rightarrow$  signifie "A SUIVRE", le symbole  $\nabla$  signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

rata  
D

elure,  
à

12X

NOVA SCOTIA  
LEGISLATIVE LIBRARY



PROVINCE HOUSE

3475

345

ON THE GEOLOGICAL RELATIONS OF THE IRON  
ORES OF NOVA SCOTIA.

*(Read before the American Association for the Advancement  
of Science.)*

By J. W. Dawson, LL.D., F.R.S.

The Iron Ores of Nova Scotia, long neglected, have recently begun to attract the attention of capitalists to an extent in some degree commensurate with their importance. The magnitude and variety of the deposits, the great richness of the ores, their proximity to the Atlantic and to great deposits of coal, are all features which give them very great economic value, and must eventually cause them to take no small part in contributing to the iron supply of the world. My purpose in the present paper is, with the aid of recent researches in which I have been occupied, to give a concise summary of the geological position and mode of occurrence of the principal deposits, and more especially of those facts which have been developed since the publication of my "Acadian Geology."

If we arrange these deposits in the first place under the two heads of *Beds* conformable to the stratification and *Veins*, we shall find that the former occupy three distinct geological horizons—that of the Lower Helderberg or Ludlow in the upper part of the Silurian, that of the Oriskany at the base of the Devonian, and that of the Lower and Middle Carboniferous. The latter occur in altered rocks which may be assumed to be of

NS  
5033

J  
TR

Silurian age, in the Lower Carboniferous, and at the junction of these two groups of rocks. We may shortly consider the deposits of these several kinds and ages in their order.

1. BEDDED ORES.

(1) *Great Hematite Bed of the Lower Helderberg Series.*

This, in so far as at present known, is most extensively developed in the vicinity of the East Branch of the East River of Pictou, and on the upper part of Sutherland's River. Here the rocks which rise unconformably from beneath the Carboniferous beds of the Pictou coal-field, consist in great part of gray and olive slates, usually coarse and unevenly bedded, and with occasional calcareous bands, holding the characteristic fossils of the "Arisaig group," a series in Nova Scotia equivalent to the Lower Helderberg of American geologists, though in its specific forms more nearly allied to the English Ludlow than to groups of this age on the great inland plateau of America. These beds are affected with slaty cleavages, highly inclined, much faulted, and folded in abrupt anticlinals, so that their detailed arrangement has not yet been satisfactorily traced. The great ore-band which forms one of the most conspicuous marks for unravelling their complexities, has been traced mainly along two distinct lines of outcrop, both somewhat curved and broken, and which seem to lie on the opposite sides of an anticlinal axis. It has also been recognized in two other localities where it must come up on distinct lines of outcrop, the precise relation of which to the others has not yet been ascertained.

The ore bed is accompanied by a thick band of olivaceous slates, and beneath this there appears hard ferruginous quartzite which Dr. Honeyman compares to the Medina sandstone. Lower than this and possibly unconformable to it are black and greenish slates with bands of quartzite and soft chloritic and nacreous schists which as yet have afforded no fossils. They are associated with hard beds or masses of rock rising into some of the highest eminences, and which have usually been described as trap, but which seem to consist for the most part of an indurated slaty breccia or conglomerate, corresponding very nearly in character to the typical graywacke of the older German geologists. These rocks may be of middle Silurian age, though possibly in part older, and we shall meet with them again in connection with the great vein of specular iron.

4355

The ore-bed where most largely developed attains a thickness of about thirty feet, and in places where it has been opened up by exploratory works, it has been found to afford from ten to twenty feet in thickness of good ore. This ore is a red hematite, sometimes compact and laminated, but more frequently of an oolitic character occasioned by the arrangement of the peroxide of iron in minute concretions enveloping grains of sand. By the increase of these silicious grains it passes in the poorer portions into a sort of ferruginous sandstone. Similar beds of fossiliferous ore are well known to occur in the Clinton group of New York and Pennsylvania, and Prof. Hall informs me that they are found also in the Lower Helderberg series of New York.

Along the different lines of outcrop above referred to, this bed has been traced for several miles, and being of a hard and resisting character, it rises into some of the higher elevations of the country. Though not one of the richest ores of the district, its great quantity and accessibility render it highly important for practical purposes. The analyses made of it show a percentage of metal varying from 43 to 54 per cent. The foreign matter is principally Silica, and the proportions of Phosphorus and Sulphur are small—one of the specimens analyzed affording none whatever, another .22 Phosphoric Acid and .29 Sulphur. These analyses were made at the instance of Mr. E. A. Prentice, now organizing a company to work this and other deposits in the district. The principal exposures of this bed are distant only twelve miles from the great collieries of the East River of Pictou, and less than ten miles from the Pictou and Halifax Railway. This deposit was first described by Mr. R. Brown in Haliburton's History of Nova Scotia, 1829, and subsequently by the writer in Acadian Geology. More recently exploratory works have been carried on and a practical report made by Mr. G. M. Dawson, Associate of the School of Mines, London; and the bed has been traced and collections of its fossils made by Mr. D. Fraser of Springville.

(2) *Hematite and Magnetic Iron of Nictaux and Moose River.*

This deposit takes us to the other extremity of Nova Scotia, and brings us a stage higher in geological time, or to the period of the Oriskany Sandstone. It would indeed appear that the conditions of ore-deposit so marked in Eastern Nova Scotia in the upper Silurian, were continued in the western part of the Province into the Devonian. In many specimens of the Nictaux

ore the chief apparent difference as compared with that of Pictou is in the contained species of fossils.

Where I have examined this bed, it appears to be six feet thick and enclosed in slaty rocks not dissimilar from those associated with the Silurian ore of Pictou. Recent explorations at Nictaux are said to have developed extensions of this deposit; but I have no details of these. As rocks of the Arisaig group are known to underlie the Nictaux beds, it is not impossible that additional beds of ore may be found in these. The normal condition of the iron of the Nictaux bed is that of peroxide; but locally it has lost a portion of its oxygen and has become magnetic. This I believe to be a consequence of local metamorphism connected with the immense granite dikes which traverse the Devonian rocks of this region.

The Nictaux ore is more highly fossiliferous than that of Pictou, and contains a larger proportion of Phosphate of Lime. In the attempts hitherto made to work this ore, the distance from coal has been a main disadvantage, but the construction of the Windsor and Annapolis railway has diminished this. The Devonian beds holding this bed are described in "Acadian Geology." An analysis of a specimen made many years ago gave 55 per cent of iron.

### (3) *Bedded Ores of the Carboniferous System.*

The most remarkable of these is a bed of crystalline *Spathic iron* or Siderite, occurring in the Lower Carboniferous series, near Sutherland's River in the County of Pictou. As described by Mr. G. M. Dawson, who prosecuted works of exploration in it last year, it is a conformable bed, occurring in the Lower Carboniferous red sandstones, and varying from six feet six inches to ten feet six inches in thickness. It is accompanied with smaller bands of the same mineral, and at no great vertical distance from it is a bed of gypsum. Its mode of occurrence is on the whole not dissimilar from that of the non fossiliferous sub-crystalline limestones which occur in some parts of the Lower Carboniferous series associated with the gypsum. This ore is a true Spathic Iron, granular and crystalline in texture, and when unweathered of a light gray colour. It affords from 42 to 43 per cent. of iron and contains from 2 to 8 per cent. of manganese. This bed is only four miles distant from the "Vale" colliery, and is intended to be worked in association with the Hematite already

described, and with the other ores on the East River of Pictou possessed by the same proprietors. From the Report of Mr. Andrews on the second geological district of Ohio, it would appear that similar beds, though on a smaller scale, occur in the Lower Carboniferous series of that State. In Nova Scotia this bed is at present altogether unique.

*Clay Ironstones* occur in many parts of the Nova Scotia coal-field. In the workings of the main seam of the Albion Mines, Pictou, considerable quantities of nodular black ironstone are extracted, and will, no doubt, be utilized. In the beds under the main seam there are also clays rich in ironstone concretions. Beds with ironstone balls also occur in the measures north of the New Glasgow conglomerate, and one of these is remarkable for the fact that the nodules were found by Dr. Harrington to contain nuclei of Blende, a mineral otherwise unknown in the carboniferous of Nova Scotia. No attention has yet been given to these ores as sources of iron, but it may be anticipated that a demand for them will arise in connection with the richer ores in the older formations.

## II. VEINS OF IRON ORE.

### (1) *Great Specular Iron Veins of the Silurian States and Quartzites.*

In a paper on the metamorphic and metalliferous rocks of Eastern Nova Scotia in 1848,\* I mentioned the fact that the inland series of metamorphic rocks (bounding the coast series now known as the gold-bearing series) and believed to be of Upper or Middle Silurian age, abound in veins of specular iron, associated with spathic iron and ferruginous dolomite, and occasionally with metallic sulphides, and I described some of these deposits. In the country eastward of Lochaber Lake, where this same formation occurs, not only are numerous small veins of specular iron and carbonate of iron found in it, but a rich vein of Copper Pyrites, noticed in "Acadian Geology," has recently been opened up and found to be very valuable.

In most parts of the region these iron veins, though very numerous, are of trifling thickness; but in two localities they are known to attain to gigantic dimensions, rendering them of great economic importance.

---

\* Journal of Geological Society of London.

The earliest known of these was the great vein of the Acadia mine in the Cobequid mountains, discovered by the late Mr. G. Duncan, and on which I reported in 1845. These hills consist on their southern side of parallel bands of olive and black slate with beds of quartzite, all very highly inclined. The iron vein is a great irregular fissure, extending for many miles parallel to the bedding, and apparently accompanying a band of quartzite. It contains in addition to crystalline and often micaceous Specular iron and Magnetic iron, large quantities of a rich earthy red ore, which from the crystalline planes which it presents, would seem to have been a Carbonate of Iron decomposed and oxidised. These iron ores are associated with large quantities of a crystalline ferruginous Dolomite, allied in composition to Ankerite. This may be regarded as the veinstone to which the iron ores are subordinate, and which in the thinner parts of the vein occupies nearly its whole breadth. At the outcrop of the vein it is in some places weathered to a great depth into a soft and very pure yellow ochre. Small quantities of sulphides of iron and copper and of sulphate of barium are occasionally present. In addition to the above, which may be regarded as the primary contents of the vein, there occur in some parts of it secondary deposits of concretionary Limonite, which have of late years afforded a very large part of the ore smelted by the Acadia Company.

In some places the thickness of this vein has been found to be 150 feet, with intercalated masses of rock, but it is very irregular, diminishing occasionally to mere strings of ankerite. It is remarkable that in the Cobequid mountains, which are cut by transverse ravines to the depth of about 300 feet, the vein does not appear to be well developed in the bottom of the ravines, but only in the intervening heights. At first I was disposed to account for this by supposing that the deposit is wedge-shaped, diminishing downward; but I have more recently been inclined to believe that the large development of the vein is dependent on differences in the containing rocks which have rendered them harder and more resisting at the points of such greater developments.

With respect to the age of these beds, they must be older than the Lower Helderberg rocks, which both at the eastern end of the Cobequids and at the East River of Pictou, rest upon them. They are on the other hand probably newer than the auriferous pri-

mordial rocks of the Atlantic coast. As they have afforded no fossils their age does not at present seem capable of more precise definition. With regard to the filling of the vein fissures, this, if coeval with the metamorphism of the containing beds or immediately subsequent thereto, would fall between the period of the lower Devonian and that of the lower Carboniferous, or within the Devonian age. The denudation connected with the Lower Carboniferous conglomerates and the fragments contained in these conglomerates, seem to imply that the ore-bearing slates were then in the same condition as at present. On the other hand the Lower Carboniferous sandstones themselves contain in places narrow veins of specular iron, which also occurs, as well as magnetic iron, in the fissures of the Triassic trap.


On the west side of the East River of Pietou, there occur rocks precisely similar to those of the Cobequid range, of which indeed they may be regarded as an Eastern continuation, and including an iron vein which must be regarded as the equivalent of that of the Acadia Mine, which it resembles perfectly in mineral character and mode of occurrence, differing only in the greater proportionate prevalence of the specular ore.\*

In New Lairg, a few miles from Glengarry Station, the most western portion of this vein known to me, contains much Ankerite, with strings of Specular iron; and in large loose pieces there are indications also of red ore which is not visible in place. Farther to the eastward on the West Branch of the East River of Pietou, there appears a band of quartzite thirty feet thick filled with veins of Limonite; but specular ore is not found at this place. Still farther to the eastward and near the east branch of the East River the specular vein attains a very large development, shewing in some places a thickness of twenty feet of pure ore. Its course is S. 60° to 70° E. or nearly coincident with that of the containing beds; and as on the Cobequids, its attitude is nearly vertical and it appears to be thickest and richest in the rising grounds. In one very deep ravine the bed of quartzite usually associated with the ore seemed to be wanting, and the vein was represented by innumerable strings of Ankerite, forming a network in the slate. As in the Cobequid vein, masses of Magnetic ore are occasionally mixed with the Specular. To complete

---

\* This vein was first described by the late Mr. Hartley in the Report of the Geological Survey of Canada, 1870.

the resemblance, loose masses of Limonite are found in the vicinity of the vein, giving rise to the expectation that a vein or veins of this mineral may be found to be associated with the specular ore. The ores of this vein in Pictou County are nearly pure peroxide of iron, containing from 64 to 69 per cent. of metal, and can be obtained in great quantity from the outcrop of the vein where it appears on the rising grounds.



*Ideal Section, showing the general relations of the Iron Ores of the East River of Pictou.*

1. Great bed of Red Hematite.
  2. Vein of Specular Iron.
  3. Vein of Limonite.
- (a) Older Slate and Quartzite series, with Trap, &c.  
 (b) Lower Helderberg formation and other Upper Silurian rocks.  
 (c) Lower Carboniferous of the East Branch of East River.

(2) *Limonite veins of the East River of Pictou.*

The valley of the East River of Pictou above Springville is occupied by a narrow tongue of Lower Carboniferous rocks, having at one side the slates containing the ore last mentioned, and on the other a more disturbed country already referred to as containing the great Lower Helderberg bed of Hematite. It is highly probable that the river valley follows the line of an old pre-carboniferous line of fracture, denuded and partially filled with the Lower Carboniferous beds, including large deposits of limestone and gypsum. At the line of junction of the Carboniferous and older rocks on the east side of the river, occurs the great Limonite vein of the district, forming a vein of contact of exceeding richness and value. It follows the sinuosities of the margin of the older rocks, and varies in thickness and quality in different places; being apparently richest opposite the softer slates and where these are in contact with a black manganese limestone, which here, as in many other parts of Nova Scotia, forms one of the lowest members of the Carboniferous series. The ore is sometimes massive but more frequently in fibrous concretionary balls of large size, associated with quantities of

smaller concretionary or "gravel" ore. In some places the ore of iron is associated with concretions or crystalline masses of Pyrolusite and Manganite.

Denuding agencies in the Post-pliocene period have removed portions of the vein and its wells, and have deeply covered the surface in many places with debris. Hence the outcrop of the vein was originally marked by a line of masses of the ore too heavy to be removed by water. From the analogy of the other veins to be mentioned in the sequel, I was led to believe that the source of these masses would be found in the Lower Carboniferous rocks, and so stated the matter in the first edition of *Acadian Geology* (1855). Subsequently, however, the vein having been exposed in situ, and one wall proving to consist of metamorphic slate, it was described by Dr. Honeyman and by Mr. Hartley of the Geological Survey as a vein in the Silurian rocks. Still more recently exploratory works conducted by Mr. G. M. Dawson, with the aid of Mr. D. Fraser, have clearly proved that the vein follows the junction of the two formations. The ore of this vein is of the finest quality, affording from 62 to 65 per cent. of metallic iron. The more productive portions of this vein, as well as of the specular vein in its vicinity, are in the hands of the parties already referred to, in connection with the Hematite bed.

(3) *Limonite of Shubenacadie, Old Barns and Brookfield.*

At the mouth of the Shubenacadie River, the lowest Carboniferous bed seen is a dark-coloured laminated limestone, in all probability the equivalent of the Manganesian limestone already referred to, as well as of the Manganiferous limestone of Walton, the Plumbiferous limestone of the Stewiacke, and the lower black limestone of Plaister Cove, Cape Breton.\* This limestone and the sandstones and marls overlying it, are traversed by large fissure veins, holding a confused aggregation of iron ores and other minerals, as Limonite, Hematite, Gothite, Sulphate of Barium, Calcite, &c., some of which appear sufficiently large and rich for profitable exploration. In the same formations, further to the eastward, at Old Barns, similar veins are found to be largely developed, and at Brookfield, fifty miles east of the Shubenacadie, and apparently near the junction of the Lower Carboniferous with older rocks, large surface masses of Limonite

\* See *Acadian Geology*.

appear to indicate an extensive deposit of similar nature, but which has not, I believe, been yet so far opened up as to establish its practical importance.

(4) *Iron Veins of the Triassic Trap.*

Veins of Magnetite and Specular Iron occur in several localities in the great beds of trap associated with the Triassic red sandstones of the Bay of Fundy, but so far as known these ores are insignificant in quantity.

It will be observed from the above notes, that while the iron vein of the Cobequid hills is at no great distance from the coal-field of Cumberland, with which it has now railway connection, the still larger and more important deposits of Pictou are very near to the extensive collieries of that district, and to railway and water communication, so that every facility appears to exist for their profitable exploration, and it may be anticipated that they will soon be rendered available for the supply of iron of superior quality, more especially to meet the large and increasing demand of the Dominion of Canada.

ure, but  
o estab-

l locali-  
ssic red  
se ores

he iron  
e coal-  
ection,  
re very  
ay and  
ist for  
t they  
uperior  
emand

