

**CIHM  
Microfiche  
Series  
(Monographs)**

**ICMH  
Collection de  
microfiches  
(monographies)**



**Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques**

**© 1998**



The copy filmed here has been reproduced thanks to the generosity of:



National Library of Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol  $\rightarrow$  (meaning "CONTINUED"), or the symbol  $\nabla$  (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

Bibliothèque nationale du Canada




Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole  $\rightarrow$  signifie "A SUIVRE", le symbole  $\nabla$  signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents.

Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.



MICROCOPY RESOLUTION TEST CHART

ANSI and ISO TEST CHART No. 2



APPLIED IMAGE, Inc.

1000 W. 10th Street  
Ann Arbor, MI 48106  
313/769-1000  
313/769-1001



PROVINCE OF BRITISH COLUMBIA.

DEPARTMENT OF AGRICULTURE  
(HORTICULTURAL BRANCH).

THE ONION-THRIPS

(*Thrips tabaci* Lind.)

THE onion-thrips is an annual source of loss, particularly to growers of onions in the drier and more arid sections of the Province. This loss is for the most part needless, inasmuch as effective sprays can be applied by which the attack by the thrips may be reduced below the margin of serious import.



The onion thrips (*Thrips tabaci*): Adult. Highly magnified. (U.S. Dept. Agric. Bull. 856. Orton and Cliftenden.)

The only question that need arise in the minds of the growers is, "Does it pay to spray?" A. L. Safro, writing in the *Journal of Economic Entomology*,<sup>6</sup> propounds the rule that "the cost of spraying truck crops for pests that threaten to destroy all or a large part of the crop does not become prohibitive until the immediate application in view, together with such following farm operations as can be definitely foreseen, have a total cost in excess of the reasonable expectation of gross returns from the crop in question." He further remarks that "when from \$150 to \$175 per acre has been expended in growing onions and a severe epidemic of onion-thrips threatens, it is certainly good business to spend even as much as \$50 per acre, if necessary, in spraying to save even as little as \$100 of the original investment." To these statements may be added the remark that efficient control in one year undoubtedly reacts upon the degree of



Onions attacked by thrips. Note the variation in size and the characteristic appearance of the leaves. (Author's illustration.)

infestation in the next year. The advisability of controlling the onion-thrips, therefore, is a matter which can only be settled by the individual grower himself or by a community of growers.

The onion-thrips passes through four distinct stages in its development—the egg, larva, pupa or nymph, and adult. The winter is passed

<sup>6</sup> Safro, J. E. E., Vol. 10, No. 6, 1917.

00924371

in the soil in the larval stage for the most part, development to the adult taking place in early spring. Eggs are laid in slits on the leaves of the plant and the young larvae issue to commence feeding on the leaf tissue. The illustration given below shows the characteristic work of another species of thrips upon oat leaves. The damage to onion-leaves is similar, and results in a sapping of the vitality of the plant by a process of sucking, causing a "silvery" appearance and a diminution in the growth of the onion. The life-history of an individual thrips, from the egg to the adult, is rapid; hence several generations may be produced in a season. Warm, dry weather fosters their development, but rain and cold weather undoubtedly checks them.



Work of thrips on leaves of oats, enlarged

#### Control.

*Water Spray.* In a small way, frequent and well-directed sprays of ordinary water will go far in reducing injury.

*Soap Spray.* Any spray containing either whale-oil soap, soft soap, or any of the non-caustic neutral soaps may be used. If applied without adding nicotine-sulphate or any other insecticide, these soaps, each individually, may be used at the ratio of 1 lb. to 8 imperial gallons of water. The soap is first dissolved in a small quantity of warm water and then added to the full amount.

*Nicotine-Sulphate-Soap Spray.* If standard 40 per cent. nicotine sulphate is used, soap may be added in lessened quantity to that required for a straight soap spray, and is included to serve as a "spreader."

The following formula is recommended to obtain the most satisfactory results in commercial plantations: 40 per cent. nicotine sulphate,  $\frac{1}{2}$  to  $\frac{3}{4}$  pint; whale-oil soap, 4 to 5 lb.; water, 80 (imperial) gallons. Mix in barrels of 40 gallons each.





*Tobacco Decoctions*—These may be made by steeping waste tobacco leaves and stems in cold water for a few days; 1 lb. of tobacco waste to 2 gallons of water.

The time of year at which these sprays are applied depends upon the powers of observance of the grower. A close watch should be kept for the presence of the small, active, greenish-yellow larvæ which may be found in the axils of the leaves close to the bulb. Spray should be applied as soon as they are readily apparent and as often as deemed necessary. The first spray should be applied about June, and particular care should be taken to direct it, under considerable pressure, into the axils of the leaves. On the degree of efficiency at which this first spray is applied will depend the necessary number of later applications.

Commercial growers of onions should provide themselves with suitable spraying-machines which will permit of the work being done rapidly and economically.

*Victoria, B.C., issued September, 1919.*

This circular has been prepared by R. C. Treherne, Field Officer for British Columbia Entomological Branch, Dominion Department of Agriculture, Vernon, B.C. at the request of the Horticultural Branch.

Copies of this circular may be obtained free of charge on application to the Horticultural Branch, Department of Agriculture, Victoria, B.C., or from local branch offices of the Department.

VICTORIA, B.C.

Printed by WILLIAM H. CLUGES, Printer to the King's Most Excellent Majesty,  
1919.



