CIHM Microfiche Series (Monographs) ICMH
Collection de
microfiches
(monographies)



Canadian Institute for Historical Microreproductions / Institut canadian de microreproductions historiques

(C) 1997

Technicai and Bibiiographic Notes / Notes techniques et bibiiographiques

	12x	16x		20x		24x		28x		32x
									TT	
	em is filmed at the redu cument est filmé au tau 14:	x de réduction in		sous.	22x		26x		30x	
	Additional commentaires sup									
	Blank leaves added within the text. Whe omitted from filming blanches ajouté apparaissent dans possible, ces pages	enever possible	e, these have le certaines ne restau lorsque cel	e been pages ration		possible ima colorations filmées deu possible.	variables	ou des d	iécoloration	ns son
	Tight binding may control interior margin / La l'ombre ou de la contérieure.	a reliure serré	e peut cau	ser de		Opposing discolouration	eilleure in pages w ons are fil	hage possibilith varying med twice	ble. ig colours to ensure	ation o
	Only edition availal Seule édition dispo					possible is partiellement pelure, etc.,	mage / I	Les page les par un f	s totaler euillet d'er	nent or
Ш	Relié avec d'autres	documents				Pages who				
	Bound with other m		diedi			Includes su Comprend				
	Coloured plates an Planches et/ou illus	d/or Illustration	ns /			Qualité Inég			st /	
<u></u>	Coloured Ink (I.e. o Encre de couleur (I	iner than blue i.e. autre que b	or black) / bleue ou no	ire)		Quality of p				
	Coloured maps / C			couleur	Image: Control of the	Showthroug			7.7	
	Cover title missing					Pages deta				
	Covers restored ar Couverture restaur					Pages disc	oloured, s	tained or fo	oxed /	
	Couverture endom	magée				Pages resta				
	Covers damaged /					Pages dam	aged / Pa	ges endon	nmagées	
7	Coloured covers /					Coloured p				,
copy may the signi	Institute has attempted available for filming be bibliographically images in the refleantly change the ked below.	g. Features of unique, which eproduction,	of this copy n may alter or which	which any of may	été plaire ogra ou q	titut a micro possible de : e qui sont p phique, qui p ui peuvent e ormale de fili	se procure eut-être u beuvent m exiger une	er. Les de niques du nodifier une modificat	étails de d point de e image re lon dans l	et exert vue bib produit a méth

The copy filmed here has been reproduced thanks to the generosity of:

National Library of Canada

The images eppearing here ere the best quelity possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or liustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol — (meaning "CONTINUED"), or the symbol V (meaning "END"), whichever applies.

Maps, plates, charts, stc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as meny frames as required. The following diagrams illustrats the method:

L'exempleire filmé fut reproduit grâce à le générosité de:

Bibliothèque nationale du Canada

Les images suivantes ont êté reproduites avec le plus grand soin, compte tanu de la condition at da la netteté de l'axamplaira filmé, at an conformité avec les conditions du contrat da fiimage.

Les exemplaires originaux dont la couverturs an papier est imprimée sont filmés an commançant par la premier piet et an terminent soit per la dernière page qui comports une emprainta d'impression ou d'iliustration, soit par la sacond plat, salon le cas. Tous les autres examplaires originaux sont filmés an commençent per la pramière page qui comports une ampreinte d'impression ou d'iliustration et an terminant par la dernière page qui comporte une taile empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaqua microficha, salon la cas: le symbole → signifia "A SUIVRE", la symbole ▼ signifie "FIN".

Les cartes, pianches, tableeux, etc., pauvant être filmés à des taux de réduction différents.

Lorsque le document est trop grend pour être raprodult en un seul cliché, il est filmé à partir de l'angle supérleur gauchs, de gauchs à droits, et de haut en bes, en prenent le nombre d'Images nécesseirs. Les diagrammes suivents illustrent le méthode.


1	2	3

1	
2	
3	

1	2	3
4	5	6

MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)






APPLIED IMAGE Inc

1653 East Main Street Rochester, New York 14609 USA (716) 482 - 0300 - Phone (716) 288 - 5989 - Fax





2 EN.SJ. 2/ Hilpin, Edwin. C2
3/ The minerals of N.S.
190






NO Miles





VINCE OF NOVA SCOTIA.





BY

EDWIN GILPIN, JR., A.M., LL.D., Fellow Royal Society of Canada, Etc., Etc.,


CHIEF INSPECTOR OF MINES, AND

DEPUTY COMMISSIONER OF PUBLIC WORKS AND MINES.




HALIFAX, N. S.: Commissioner of Public Works and Mines, King's Printer, 1901.

WM. MACNAB, PRINTER, 3 PRINCE ST., HALIFAX, N. S.



INDEX.

Geol	ogy of Nova	Scotia	9
Coal	-Cape Breto	A 60	10
**	Richmond		14
66	Pictou		15
64	Comberlan		_
66	Cannel, Sh	alan ata	16
Grap	hite. Petrole	A 60 Pro a	17
Gold	-General Re	n	18
44	History of	Minim	19
44	Districts.	Wannalas	23
44	11	Montage Oldham	24
44	46	Montague, Oldham	25
44	46	Mt. Uniacke, Renfrew	26
44	44	Caribou	27
44	**	Sherbrook	28
64	**	Wine Harbor, Salmon River	29
44	.,	Tangier, Fifteen Mile	30
••	.,	Lake Catcha	3)
"	"	Stormont	32
46		Miscellaneous	15
"	Mining—Co	ets, etc	36
	Concentrate	8	10
. "	Production .		14
Lead	Ores		15
Copp	er Ores		17
Iron (Ores	4	19
Misce	llaneous Mir	nerals:	
	Gypsum		5
	Manganese		7
	Antimony .	B	0
	Celestine, 8	Induction I foliability of the same and the	0
	Phosphate,	Salt, Building Stone 6	
	Limestone .		H
	Clay, Comn	Man.	5
	Paints		_
	Barytes		R
	Fire Clay		_
	Refractory	Stone Decision Stone	0
Tenur	e of Mineral	Tanda	-
Table	-Coal-Sal	aa 1000	2
	Cler	novel Durchastion	-
	Pre	education by Calling	
	Sale	no 1708 A. 1000	_
	Ext	mandad du Ilmidad 674.4	-
	Gold-Ger	neral Statement 77	•



HALIFAX, NOVA SCOTIA, May 1st, 1901.

The Honorable George H. Murray, Provincial Secretary:

SIR,-

I beg leave to submit the following report on the mines and mineral lands of Nova Scotia. I have endeavored to give in a brief form information intended to convey in a general manner the extent and value of the mineral resources of the Province.

I have the honor to remain,

Your obedient servant,

EDWIN GILPIN, JR.,

Deputy Commissioner and Chief Inspector of Mines.




INTRODUCTION,

In the following pages I have tried to describe briefly the mineral resources of the Province.

For its extent Nova Scotia compares more than favorably with any of the Provinces of the Dominion. Its wealth is not confined to minerals. Large tracts supply in abundance the vegetable products of the northern temperate zone. Its shores are surrounded by extensive and valuable fisheries. Its forests furnish an unlimited lumber supply, and abundant water powers are being brought into increased use.

The peculiarly diversified nature of the Nova Scotia minerals may be judged of by the fact that it has in close relationship the three most valuable, iron, coal, and gold. The initiation of extensive iron and steel works is directing attention to our large deposits of ore, and of fuel yielding coke claimed to be the best in the world. Among other minerals more or less worked may be mentioned lead, zinc, silver, copper, manganese, gypsum, barytes, etc.



GEOLOGY OF NOVA SCOTIA.

I give the following outline of the geology of Nova Scotia, as serving to indicate in a rough manner the extent and ages of the strata, which at one point present to the farmer soils well adapted to recompense his labor, and at another hold the treasures which excite the cupidity of the miner.

I am indebted to Mr. Ami, of the Canadian Geological Survey, for information on this subject, but it may be remarked that on several points

definite conclusions cannot yet be drawn.

The Laurentian system is well developed in Cape Breton, occupying the more elevated portions of the island. There are apparently two series—the lower series carrying an extensive calcarcous development in addition to the granites, syenites, felsites, slates, etc., common to both. These measures carry gold, silver, lead, zinc, graphite and iron ores, and doubtless will yield the other minerals characterizing the system in Quebec and Ontario.

The Huronian system is not yet recorded as occurring in Nova Scotia.

The upper and lower divisions of the Cambrian are represented here. To the former the gold-bearing series, consisting of an upper slate formation and a lower quartzite formation, has been referred. The foldings of these strata have been broken through by masses of granite at a date slightly preceding the Carboniferous cra.

Below McAdanis' Brook, Escasonic River, near McPhec's Point, and along the Mira River in Cape Breton, are beds referred to the upper Cam-

These strata frequently carry ores of iron.

It is interesting to note that the iron-ore deposits of Bell Island, in Newfoundland, are referred to the Ordovician system, and that large areas in the eastern counties are provisionally referred to this system on stratigraphical This determination, however, requires paleontological corroboraevidence. tion.

The upper division of Sir R. Murchison's Silurian is extensively developed in Nova Scotia. It is typically developed at Arisaig and extends into Pictou County. It also occurs from Nictaux to Wolfville, and at several points along the northern side of the Cobequid Mountains. This formation is noted at several points for large deposits of iron ore.

The Devonian is represented at Nictaux and Arisaig, at both points containing deposits of iron ore. Large areas in Cape Breton, Richmond and Guysboro counties are referred to this age, and contain iron and copper ores.

The Carboniferous system is typically developed in Nova Scotia, in the northern and eastern counties. The productive horizon is met in the counties of Cape Breton, Richmond, Inverness, Pictou and Cumberland. This is underlaid by milletone grit, carboniferous ilmestones, and lower or basal conglomerates. Along the shore of the Straits of Northumberland and in the interior of Cumberland county are still higher divisions of the Carboniferous apparently passing into the Permian, and they have been termed Permocarboniferous.

The Triassic system is represented on the Bay of Fundy and the Basin of M as by beds or bright red sandstone, associated with amygdaloidal traps and ash beds. Of the succeeding systems, the Cretaceous system is reported present in a few small areas resting on the Triassic eruptions, and the Quarternary system records the glacial action which this province has been subjected to in common with the rest of the Acadian region.

COAL, SHALES, PETROLEUM, ETC.

THE COAL FIELDS OF NOVA SCOTIA.

The coal mining industry began many years ago in Nova Scotia, but it has only assumed important proportions during the past few years. Mr. Richard Brown, in his books on Cape Breton coal, has given in detail the difficulties of the trade in its Infancy.

The coal found in Nova Scotia is bituminous, and is in nearly all cases, coking. It is a true coal from the middle or productive measures of the Carboniferous division, and resembles in appearance and quality the coals of Durham and Northumberland in England.

It may be mentioned here that all the Nova Scotia coal fields are on tide water, an advantage enjoyed by no other district on the Atlantic seaboard of America.

CAPE BRETON.

The island of Cape Breton, at the eastern end of the province, holds several important coal deposits. The Sydney coal field, in Cape Breton County, extends for 35 miles along the shore, and extends inland several miles. Its available area may be estimated at over 300 square miles, 2 though recent discoveries of coal may add materially to this estimate. The seams are presented in regular beds, dipping at easy angles to the eastward, under the Atlantic. The district is bisected by Sydney Harbor, one of the firm on the Atlantic coast, and has Louisburg Harbor at one end and the I als d'Or at the other end.

known in 1874.

in the in the rland. wer or and and of the been

Basin aloidal stem is as, and ace has

but it s. Mr.

ll cases, of the coals of

on tide

Breton several niles, s te. The astward, te of the

he seams

and the

			O	P NO	VA HC	OTIA.				
BOULARDARIE.	Seam.	Point Aconi.	Bonar.	Stubbert.	Seam C.	Mill Pond.	Black Rock.	Seam F.	Seem G.	
STDNET MINES.	S.am.		Lloyd's Cove.	Seam B	Sydney Main	Bryant	Edward	Seam F	Collins	
	Strate and Coal. ft. in.	3.0	6.5	12.1	8.0	78.0	4.0	95.0		6
LINGAN.	Seam.	Seam A	Carr	Barrasois.	Victoria	Seam D	North Head	Lingan Main	Seam G.	Seam H
GLACE BAY.	Seam,			Hub	Harbor	Boutillier	Back Pit	Phelan	Roes Emery.	Lorway
Cow Bar.	Seam.				Block House	Seam D	Seam E		South Head	Long Beach

Tracey, Mullins, and other Scams,

Explorations have since shown that lower down in the Carboniferous there are workable beds of good quality underlying these given in the section.

At present there are five companies working in the coal field. The largest, the Dominion Coal Company, has control of the Glace Bay section, where it works the Phalen and Emery and International seams on a large scale. The following collieries are in operation, the Caledonia, International, Reserve, and Dominion Nos. 1, 2, 3, 4. The last three are in process of development. These collieries are up to date in every respect, and last year produced 1,930,425 tons of coal, an amount greatly to be increased in 1901, as the newer mines come into full operation. The pits are filled with all modern improvements of endless haulage, compressed air for pumps and mechanical coal cutters, fens, picking tables, etc.

A railway connects the operations of this company with the Intercolonial (Government) railway, with two piers in Sydney harbor, and one at Louisburg, and with the plant of the Dominion Iron and Steel Company. There are a ut 40 miles of road, not including sidings, laid with 80 lb. rails, and

fully equipped.

The piers can take the largest steamers affoat and load them through shoots at any tide, and are provided with appliances for loading coal without breakage, etc. The Louisburg pier has also a system of conveying belts from large pockets capable of loading in addition to the regular pier work over 600 tons per hour. These three piers can readily load 18,000 tons per day.

In the Cow Bay district the Gowrie and Biockhouse Collieries are developing an extensive submarine area in the Gowrie seam, long known and worked on the land. Their works are on the shore, and the coal passes direct

to the vessels without the intervention of railway haulage.

On the north side of Sydney harbor the Nova Scotia Steel Company are working the Sydney main seam. The workings on this seam, begun over 100 years ago, are at this point entirely submarine. The coal enjoys a a reputation for house purposes, and as it makes an excellent coke will be largely used in the steel works the company contemplate building at this point. The output last year was 249,910 tons.

The Sydney Coal Company on an adjoining area are working a lower seam, I at have not yet effected a development permitting of a large annual output.

At the extreme northern end of the field, at New Campbelton, the Messrs. Burchell Bros. are working seams, considered the extension of those of the Nova Scotia Steel and Sydney Coal companies. They effect an annual output of 12,426 tons, and control a large and valuable field.

The total production of this coal field last year was 2,199,983 tous. This district ships largely to Newfoundland and Prince Edward island, sends about 1,000,000 tons up the St. Lawrence, and after supplying about 600,000 tons to the United States, distributes the remainder at various points in

Nova Scotia.

The complation of the great plant of the Dominion Iron and Steel Company will this year call for a much increased output, to which possibly in another year will be added the demands of the furnaces the Nova Scotia Figal Company contemplate erecting. So far as can be judged at present the colliaries will not be able to meet the demand. In addition to the large territories controlled by the companies named there are valuable tracts of coal lands yet untouched, and probable extensions in addition.

Quality of the Coals.

The following analyses will serve to show the character of the seams in this district:

Emery.	Gowrie.	Phalen,	International,	Bydney.
Moisture	.50	.52	.80	1.536
Volatile combustible matter 51.10	28.13	34.21	27.55	36.362
Flxed carbon 63.10	66.01	59.73	65.90	57.008
Sulphur 1.51	1.75	.81*	1.45	1.894
Ash 3,65	5.36	3.92*	4.30	5,084

^{*}Norz. -Ash and sulphur average of nearly two hundred assays.

In volume X. of the Transactions of the Nova Scotia Institute of Science, I have given summaries of sets of tests of Phalen and Harbor seam coals in closed coke ovens for coke and by-products. The following results were obtained from the Phalen seam. It yielded per long ton 11,012 cubic ... is of gas, having an average calorific power of 571.85 B. T. U., and an average illuminating power of 9.9 candles, and with a Welsbach burner, 54.34 candles. The gas contained:—

Carbon	di	οx	id	e									,				2.	7
Illumin	ante	8 .															2.	9
Oxygen																		13
Carbon	Mo	a	xc	id	le												5.	8
Marsh	gas															3	2.	3
Hydrog	en									٠						5	1.	1
Nitroger	n.			, .													5.	07

The total yield of coke was 74.68 per cent. The ammonium sulphate per ton amounted to 32.91 lbs., the tar amounted to 12.89 gals., and the benzole amounted to .103 gals.

I pointed out a number of years ago that this coal field contained beds capable of yielding a coke equal to that from Connelsville, in Pennsylvania, admittedly the best in America, and dwelt upon the importance of this factor in iron making. The coke made at the works of the Dominion Steel Company from unwashed Glace Bay coal has answered every expectation, and by washing the amount of sulphur and ash will be decreased to a point seldom attained in commercial coke.

The coals of this district are admirably adapted for gas, coke, domestic

tons

are

irect

oots

rous

the

The

on.

arge mal,

of of

year 901.

all

and

onial ouishere and

y are over

ll be

seam, atput.
, the those ct an

This sends 0,000 nts in

use, and for marine and land boilers. They ignite readily, burn with a long flame, and when properly burned do not make an excessive smoke.

In the district extending from Cow Bay to the Mira and Salmon Rivers, a number of small coal seams have been noticed, and explorations may

prove the presence of larger deposits.

At River Inhabitants, Riehmond County, isolated erops of workable seams occur over a large tract of country, but as yet have received little attention. In Inverness county, on the western side of Cape Breton, there are several coal fields. The most southerly is presented at Port Hood. Here the crops are known of two seams, each about six feet in thickness, and of another smaller one. These seams crop on the shore, and as the coal measures extend inland and along the shore for several miles, explorations will probably increase the area. As the Port Hood islands lie a short distance from the shore on the dip of the beds it is anticipated that a very large submarine area will be available. On the lower of the larger seams a slope is being sunk, and arrangements made for a large output, to be shipped within a few yards of the bankhead.

At Mabou there are similar outcrops, comprising a four feet bed, a thirteen feet bed, a seven feet bed, a five feet bed, and others reported. Operations are now under way to open submarine workings at this point.

At Broad Cove, explorations have shown a considerable land and sea area containing, it is said, the following section:—

First 8	Seam												2	feet.
Second													2	
Third	66	Ĭ											3	66
Fourth													4	
Fifth													2	**
Sixth													7	66
Seventl		-	-	-		- 6	- 6						4	66
Eighth	_												3	

The control of this district has passed into the hands of the Inverness and Riehmond Collieries and Railway Company, and they have commenced a modern colliery, capable of working a large land area, and of indefinite extension on the seams under the sea.

. Still further north comes the Chimney Corner coal field, which has a land area 15 miles long and 3 miles wide, with an equally large submarine area. The following is given as the section of the seams at the northern end of the district where a little mining was carried on a number of years ago. There is nothing known of the seams of the lower group. The upper contains a three feet bed, a three feet vix inchea and a five feet bed. The extension of these seams has not been traced any distance from the ahore, but about five miles away the crops of two six feet seams are known.

At Glendale, on the upper waters of the River Inhabitants, there is a

small coal basin holding two three feet seams of coal. No work has been done here to test the value of the deposits, which are presumably an outlier of the Richmond coal field already referred to.

Passing to the mainland there are outcrops of small seams at Pomquet, in Antigonish county. At Hollowell Grant, in the same county, are beds of shale so charged with bituminous matter as to possess some of the properties of coal, but they are probably more valuable as a source of oil, etc. Small seams of coal are found at Nine Mile River, Hants county, and along the southern flanks of the Cobequid Mountains in the counties of Colchester and Cumberland.

PICTOU.

In Picton county, coal has long been worked in a field the dimensions of which are fairly well ascertained. The district forms a main east and west synclinal, covering about 35 square miles. There are about thirteen seams of coal from 3 feet 6 inches to 30 feet in thickness. The greater part of the district is controlled by two companies, the Intercolonial and the Acadia. There are still unworked a number of valuable areas belonging to the Nova Scotia Steel Company, Messrs Gray, Barton, Richcy and others.

The Intercolonial company works the main and underlying cams at Westville, producing last year 244,000 tons of coal. It is connected with the Canadian Government (Intercolonial) Railway, which intersects the district, and with a private shipping pier at Granton, on the Middle River, distant about five miles. The Acadia company works the same seam at Westville, and has two collieries on the Albion seams at Stellarton, and another at Thorburn, in the eastern end of the district working the six feet seam. These collieries had an output last year of 294,884 tons, and are all connected with the Government railway and with a shipping pier in Pictou harbor, distant eight miles from New Glasgow, the centre of the district.

The coals of the Pictou district are less bituminous than those of Cape Breton. Some are possessed of good coking properties, while others are not suited for this purpose. They are however good steam coals, and well adapted for use in iron works, etc., and are largely used for domestic fuels.

The following analyses, though not recent, will serve to convey a general idea "their composition:—

	Acadia,	Intercolonial,	Deep.	Montreal and Pictou.
Moisture	2.10	1.52	.75	1.76
Volatile combustible matter		29.46	20.34	22.43
Fixed carbon	61.15	60.19	68.50	67.73
Ash		9.10	10.41	7.12
Sulphur	1.48	1.62	.94	.96
Theo. evaporative power	8.50	8.24	9.39	,

thirrted...

area

long

vers,

may

able

ittle

here

ood.

ness,

the lora-

short

very

ms a

rness enced finite

has a parine n end s ago. con-

e is a

e, but

Outcrops of small coal seams are known in the district between New Glasgow and Pictou, but geologists consider they are not indicative of large

deposits.

The coal of the district is used locally for the works of the Nova Scotia Steel Company, foundries, and other manufacturing works, on the railway, at Halifax for bunkering steamers, as well as in Prince Edward Island and Quebec.

CUMBERLAND COAL. FIELD.

The northern edge of this coal field begins at the Joggins, on Chignecto Bay, and runs casterly in an almost straight line for about 19 miles. From this point the continuation of the scams has not yet been traced to the southern edge at Springhill. At Springhill the reverse side of the field is presented in an outcrop several miles long, and folded in irregular curves. The country lying between Springhill and Apple River on Chignecto Bay has not yet been explored, but it may be anticipated that the continuation of the southern outcrop of the field will be found along this line. The proved extent of the coal field has been materially increased during the past two years in the Springhill district, but as yet it can only be said that it probably extends over 350 square miles.

· The principal seam worked on the northern side is known as the Joggins main seam, of the Canada Coals and Railway Company. This seam is about six feet thick and yields annually from 60,000 to 80,600 tons. Coal is shipped by water at the mine to various points on the Bay of Fundy, and by a branch railway ten miles long to the main line of the Intercolonial Railway. A number of smaller seams are worked along the northern side of the field on a limited scale to supply local winter markets. The coals of this section

are best known as steam and domestic fuels.

At Springhill is the large colliery of the Cumberland Railway and Coal Company. Here three seam from 6 to 14 feet in thickness are extensively worked, the output last year amounting to 428,904 tons. Other underlying

seams of workable size are known here and await development.

The Springhill coals are shipped over the company's railway 4 miles to the Government Railway, or about 30 miles to the company's piers at Parrsboro, on the Bay of Fundy. From this point shipments are maintained all winter by barges to Portland, St. John, etc. The coal is found to be specially adapted for steam purposes, and is largely used by the Government, Grand Trunk and Canadian Pacific roads. In quality it is less bituminous than that from Cape Breton, and has the following general analysis:-

Vol. Comb. Matter									_		_					28.55
Fixed Carbon	•	 •	•	•	•	•	•	•								62.78
Moisture	•	•	•	•	i											3.66
Ash	•		·	·	Ĭ											4.32
Sulphur				4									٠	٠	٠	1,20
Theo. Evap. Power																8,69

To the east of Springhill, there are seams of coal apparently belonging to a horizon lower? an that occupied by the seams just referred to. So far as any work has been done the seams are smaller, not exceeding 4 feet in thickness. They have been found to a limited extent at Salt Springs and at Oxford, and crops of similar beds are reported in the Thompson district. Further work may place this section in the list of those available for economic mining. It may be said that in this county the opportunition or the discovery of coal are greater than at the longer worked districts of Pictou and Cape Breton Counties, and that the facilities for marketing the coals are better, as it lies nearer New Brunswick and the United States.

CANNEL COAL, SHALES, ETC.

to

m he

is es.

ay

on

he

he

tit

ins

is
by
ay.
eld
ion

oal

rely

ing

the

ого,

all

eci-

ent,

nous

Beds of cannel coal are met at several places in the Sydney coal field. The deposits have, I believe, not been examined in any detail, and are reported to be limited in extent.

On the North side of East Bay the manufacture of oil, etc., has been started on be is of bituminous shale in lower carboniferous conglomerate rocks. There are a number of beds ranging for several miles, and the supply is said to be unlimited. The following results are said to have been attained on working tests:

Kerosene Oil	1.25
White Spindle Oil	1.25
Heavy Lubricating Oil	2.50
Pitch	1.25
Water	5.75
Coke	
Loss, gas, etc	.50

The products, which are now being introduced on the market, are said to be well received. Bituminous shales are not uncommon in the province in the lower divisions of the carboniferous. They occur at Port Hastings, Inverness county, and at Hollowell Grant, Antigonish county. At the latter place tests were made a number of years ago on several beds from 5 to 15 feet thick, with satisfactory results.

In Pictou county, on the property of the Acadia Company, among the lower bods of the coal field, there is an interesting cannel coal, known as Stellarite. It is associated with a layer of oil shale and of bituminous coal. The cannel and shale were worked to some extent a number of years ago, and yielded 60 gals. of crude and 35 of refined oil. Its ultimate analysis gave:—

Carbon	80.96 10.15
Nitrogen Sulphur Oxygen	.68
Oxygen)	8.21

It is said to be present on the northern dip of the coal field, in which case it would underlie a considerable territory. Other shales, said to be rich in oil, are reported from the eastern part of the district, and many portions of the 1,200 feet of shale overlying the main scam are stated to yield considerable amounts of oil. Shales, apparently highly bituminous, occur at Cheverie, Horton and Kennetcook. Should the attempts now being made to work the East Bay shales prove remunerative, there should be ample room for extensive works of a similar character in the province. The oil shale industry of Scotland has been established for many years, and has flourished in spite of the competition of the United States.

Graphite.—In the presilurian strata graphitic slates are frequently observed. They are often met as a distinctive rock in the upper division of the gold measure strata. In Cape Breton at Glendale, West Bay, Grand Narrows, East Bay, Hunter's Mountain, etc., deposits have been found, yielding good commercial results. While the abundant deposits met in the United States may prevent the importation of the raw material, it might be found that its manufacture could be carried on here.

Other localities are Parrsboro, Musquodoboit, Salmon River, Hammond Plains, etc.

Oil. Indications of petroleum have 1. A observed around Lake Ainslie, Inverness County, and for a number a years attempts have been made to find its source. A number of occeholes have been put down without positive results. It may be found that the district in the headwaters of the Mabou, Denys, and Inhabitants Rivers presents a more favorable field for prospecting. Oil is also found at Cheverie, Hants County, but I am not aware of any attempt to prove its presence from an economic point of view.

NOVA SCOTIA GOLD FIELDS.

The gold fields of Nova Scotia became known about the year 1860, and were soon proven to occupy the Atlantic coast of the province.

Investigation showed that this district was occupied by two divisions of rock, granite, and slates and quartzites. Geologists referred the latter to the lower Cambrian age, a series of rocks known to be auriferous in other parts of the world.

In Nova Scotia they are divided into the lower or quartzite group, and the upper or ferruginous and graphitic slate group. The lower group, to which a thickness of over 11,000 feet is assigned by Mr. Faribault, consists principally of quartzite, interstratified with numerous beds of slate varying in colour and texture, and with a few beds of compact conglomeritic rock. The upper group, which is over 4,000 feet thick, is almost wholly composed of bluish black soft slates.

By the labors of Mr. Faribault, the pioneer of the Geological Survey in the mapping of the gold fields, much detailed information has been collected, and I am indebted to him for his clear presentation of the most interesting facts relating to the stratigraphy of the rocks under consideration.

These measures, originally horizontal, have been moved by a powerful and uniform pressure from the south, exerted in a line roughly parallel to that of the coast, which has folded them into a series of sharp parallel undulations or folds. By denudation these folds have been so worn down that in a generally level country they have been extensively exposed in horizontal sections, showing within a few hundred yards the reverse dips, to the north and to the south. The rocks generally dip at an angle of 75° to 90°, seldom lower than 45°, and overturns are frequently noted.

Estimates allow that about one-half of the auriferous district is occupied by granite, making the extent of the gold fields about 3,000 square miles. The granite is presented in masses and dykes of varying shape and size, and appears, broadly speaking, to have melted through the slates and quartzites. The granite is not generally considered to be auriferous, although gold has been found in quartz veins in it at points far distant from the slates and quartzites.

The upper or slate group is not recognized by miners as a different, although quartz veins occur in it, and sometimes show gold. It is y be found, however, on further examination to hold extensive deposits of low grade ores.

at de ple oil

has

eh be

ny

of and, the

elie,
to
the
for

iew.

ond

Attention has therefore been confined principally to the lower or quartitic group. From Mr. Farihault's observations it appears that the gold belts occur at a distance below the base of the upper or slate group, varying from 2,800 to 8,000 feet, giving a thickness of about 5,000 feet of auriferous strata. Veins occur at other points in the lower group of rocks, but have

not yet been found to be pronouncedly rich in gold.

It will therefore be seen that wherever this section of the lower group has been folded, denuded and exposed, there the best fields are opened for exploration and work. Eleven of these auriferous anticlinals have been defined between Sheet Harbor and Caledonia. There are many others succeeding these, starting from the shore and passing obliquely away from it. The district to the eastward of Halifax has been carefully mapped by the Geological Survey, and the courses of these auriferous anticlinals laid . down approximately between the points where they are known exactly by mining operations. These maps are on a scale of one mile to the inch, and give minute topographical details, so that the localities where prospecting can be most advantageously carried on are readily observed. In order that information may it available as to the district lying west of Halifax, Professor Bailey has examined, in less detail, the counties of Lunenburg, Queens, Shelburne and Yarmonth. In these districts there are larger areas of granite, but the same general structure is preserved. The maps and reports relating to the gold fields are invaluable to gold miners, and can be obtained by payment of nominal prices from the office of the Canadian Geological Survey at Ottawa.

The auriferous quartz veins of Nova Scotia occur in these anticlinals, intercalated between the layers of quartzite and slate as beds running parallel with the strata. They are seen to turn on their course where the anticlinal axis has become depressed, and underground operations have shown them rising up on one side and dipping down on the reverse side without a break in their continuity, or giving any surface indication of their existence. They extend in many cases for thousands of feet, and have been followed to depths of seven hundred feet in their vertical extension. In size they vary from an inch up to twelve or fifteen feet; many of the most productive are from six to fifteen inches in thickness. They present in spite of their bedded position many of the characteristics of typical veins. Their essential ingredient is quartz, varying in texture and color; in many of the more productive veins presenting a smooth surface and bluish shade. There is always present a varying percentage of iron, copper, lead and zinc sulphides and traces of other minerals. Cross or fissure veins also occur at Rawdon, Caribou, Blockhouse, Oldham, Brookfield, etc., and are frequently productive.

The gold is present characteristically in the free state as irregular masses, varying from microscopic particles to irregular patches, often several ounces

in weight. It is also present in films covered by the metallic accompaniments, and as invisible particles. There are also quartz veins almost free from sulphides, etc., yielding gold in workable amounts, although it is not visible. But little attention has yet been paid to this class of veins, unattackable by the ordinary quartz mills, although adapted for chemical treatment by the more modern systems. In the veins the gold is sometimes distributed with comparative uniformity over considerable areas; usually however, it is more or less concentrated within cert...n defined limits, leaving spaces on each side comparatively barren. These enriched zones are known as pay streaks, are repeated in some veins, and follow certain laws not yet clearly understood. They have hitherto been the principal source of the gold production.

Gold is also found sparingly in the quartzite beds and more abundantly in the slates. The latter when in contact with the quartz frequently show it in thin platings, and the small veinlets of quartz seaming the slates often carry gold. When one or more quartz veins occur in connection with a slate bed, the whole bed is frequently rich enough to be worked as a low grade ore. As will be shown further on, practical experience has taught the miner that profitable low grade ore means material yielding in an ordinary stamp mill from two dollars to the ton and upwards.

The early prospecting developed promising alluvial doposits at several points. At Tangier it was shown that a large extent of surface carried gold, and several hundred ounces were washed out. The limited extent of the areas, however, did not allow of systematic work, and attention was diverted to the rich vein outcrops. The most interesting alluvial development was at the Ovens, near Lunenburg. Here the sea has worn a number of large cavities in the cliffs, which are composed of soft slates with small quartz veins often showing gold. The action of the waves concentrated the gold washed from the slates along the beach near these openings, and during the years 1861 and 1862 about 2,500 ounces were returned as taken out. It is believed, however, that only a small part of the gold extracted was accounted for. Since that date small amounts have been secured by persons living in the vicinity. The richness of the alluvium directed attention to the neighboring quartz veins, but they proved uncertain and unprofitable with the crude methods of that day. Trial crushings of the quartz and slate from several points in this district would indicate the feasibility of dealing with them on a low grade basis.

At Gay's River explorations have shown that a conglomerate of lower carboniferous age, lying on the gold bearing slates, carries at its junction with them considerable amounts of gold. At one time profitable mining was carried on here on a small scale along the edge of the outcrop of the conglomerate; but the miners turned their attention to richer districts. As the conglomerates at many points in this district show gold it is to be

masses,

tzite

belts

from

erous

have

group

d for

been

thers

from

d by

ly by

i, and

ecting

r that

, Pro-

nburg,

areas

s and

can be

nadian

, inter-

el with

al axis

rising

reak in

They

wed to

v vary

ive are

f their

ssential

e more

here is

lphides

lawdon,

quently

s laid .

regretted that more attention has not been paid to them, as it a certain degree a parallel can be traced between them and the South African gold

Promising alluvial ground exists on the Nine Mile and Meander Rivers, at Renfrew, in the southern part of Waverley, and in the vicinity of other gold fields. In fact at any favorable point south of the passage of an auriferous anticlinal sights of gold can be obtained by panning, which would in California or other countries where alluvial mining is a recognized practice, be given immediate attention by enterprising miners.

However, little attention has been paid to this source of gold for many years, although it has been well within the knowledge of miners that in overy district the immediate surface cover contains much gold in boulders and free in the earth. At Moose River, Mr. Touquoy, a miner who acquired his experience in Australia, for a number of years systematically passed the surface earth of his areas through his crusher with profitable returns. There can be no doubt that at many points this system could be advantageously united, as in his case, with the crushing of underground quartz.

An Act has been passed to allow of large tracts of ground being taken on almost nominal terms for a period of three mouths, to permit of preliminary investigation for the purpose of alluvial mining. This arrangement allows capital to secure cheaply unbroken stretches of ground which, if found valuable, can afterwards be brought under the regular conditions of the Mines Act.

In this connection it is interesting to rote that indications of gold deposite are found in alluvials outside of the recognised or working gold fields of the Province. Gold has been found in this manuer in Digby County, along the Cobequid Mountains, in the northern part of Guysboro County, and it almost every river flowing from the Precambrian table land of northern Cape Breton. It is not known yet if this gold is derived from workable veins carrying free gold, from the schists which sometimes show gold, or from the oxidation of sulphurets which often form a considerable percent age of certain beds. At Middle River and Whycocomagh, in Cape Breton free gold occurs in the quartzites and in quartz veins therein. At the latter place it also occurs in the sulphides with which many of the felsites, etc.

There appear to be two sources of alluvial gold in the province, first the free gold-bearing veins of the Atlantic coast, and the auriferous mineralise deposits, also in places containing free gold, and presented over a large section of the province, hitherto generally considered valueless for gold mining.

certain an gold

Rivers, of other e of an would

that in boulders acquired assed the There tageously

r many

taken on eliminary it allows und valuhe Mines

d deposits
lds of the
along the
y, and in
northern
workable
w gold, or
percentpe Breton,
the latter

, first the mineralised er a large es for gold

sites, etc.,

HISTORY OF GOLD MINING.

The discoveries of the gold deposits of California and Australia had prepared the minds of Nova Scotians, and a tremendous excitement arose when the presence of gold in the province was established. It is true that the first reports were listened to with incredulity by those in power, but public opinion, inflamed by the sight of the beautiful specimens from Tangier, compelled a legal recognition of the various districts.

The first claims were limited in size, and operators soon gave them up. Gradually companies acquired control of areas, put up mills, and started with high hopes. In many cases large profits were realized for a time in spite of every form of extravagance, of poor mills, and of unskilled min; This continued for a few years, when the laws of finance and mining resumed their sway. The veins opened on the richest points of the outcrop of their pay streaks gradually became too poor to pay; the treasuries of the companies, after lavishing dividends from the proceeds of the richest ore, had nothing left for a rainy day.

In 1872 this state of affairs had very materially changed in one respect, the system of working the mines. The tribute system had come into practice, and company mining had been largely suspended. It became evident that many leads that were considered unworkable by the directors of a company yielded a profit to a party of tributors. The chief objection to the tributors was the desultory nature of their operations, which in many cases led to the flooding and the crushing of the outcrops of good veins.

However, the tribute system marked the commencement of a new era. The enterprising men among the tributors, left to their own scanty resources, found that economy was the mainstay of mining. So by degrees high explosives, improved pumping and hoisting machinery, air drills, better systems of working, and last but not least great improvements in mills and milling, were introduced, until the practice in Nova Scotia is fairly abreast of the times. As the success of many of the tributors became known, capital was gradually again available, and at the present time the greater number of the companies working in the province are successful whenever they have been started on a business basis and properly managed.

DISTRICTS.

It would occupy too much space to give a detailed notice of all the gold districts. I therefore give a very brief reference to the principal ones.

The best known districts are Wine Harbor, Stormont, Sherbrooke, Tangier, Montagu, Waverley, Renfrew, Oldham, Uniacke, Caribou, Lake Catcha, Brookfield, Whiteburn, Malaga, Fifteen Mile Stream, and Salmon River. Among others that have received more or less attention may be named Gegoggin, Ovens, Indian Path, Millipsigate, Gold River, Kemptville, Gay'a River, South Stewnscke, Gold Lake, Killag, Beaver Dam, Ecum Secum, Harrigan Cove, Cow Bay, Lawrencetown, Rawdon, etc.

More or less detailed notices of all these localities can be found in the reports of the Mines Department. In the case of the minor districts, as may be gathered from preceding remarks, their development has been retarded by the persistence of the prospector in neglecting for years the problem of large supplies of low grade ore in favor of isolated rich velns.

WAVERLEY.

Waverley was one of the earliest proclaimed districts. Rich boulders were found on "American Hill" in the fall of 1861, and shortly afterwards on Laidlaw's Hill, on the east side of the Waverley lakes. The latter place carried a valuable vein in a shape then new to miners, and attracted much attention. The vein was presented in a horse-shoe shape, with its apex pointing to the westward, and corrugated, like logs of wood lying side by During the next three years about 5,000 tons of quartz were extracted by open work along the crop, and good average returns secured, with some extraordinary yields, in one case of 200 ounces from two tons. Finally the surface water proved too much for the individual operators, and the workings were abandoned for many years. At present these properties and others have been consolidated, and a tunnel driven in to cut the lode at the level of the lake, and an extensive mining plant installed. As at this level and for some distance lower it is auriferous, it has again proved a There are probably other unseen veins below this one valuable property. which would give large backs above the level of the lake. In West Waverley the Burkner property was for some years a large producer, having yielded nearly a quarter of a million of dollars, as was also the DeWolf property, yielding about 9,000 ounces. Several companies on the American Hill were amalgamated in 1864, and yielded during that year a profit of \$90,000. Other small operators in some cases did fairly well. These results were all obtained from shallow workings, in no case exceeding 300 feet. Up to 1867 there had been a yield of about 40,000 ounces of gold from 56,758 tone of quartz, from a narrow strip running through the district. The total yield up to date is 61,711 ounces, from 123,383 tons of quartz, or about \$1,200,000.00.

MONTAGU.

The district of Montagu sprang into notice on the discovery of a boulder of quartz yielding \$1,600.00 of gold. This led to the openin, of the Lawson lode which, after a career unprofitable on an average yield of over two ounces to the ton, fell into the hands of the Lawson Brothers. They worked it for about five years and extracted 10,000 ounces. The Rose lead was discovered by Mr. G. W. Stuart in 1879, and yielded large returns for some time, notably on one occasion 800 ounces from 80 tons of quartz. Finally it was lost on a fault.

The DeWolf property, after an uneventful career for some years, became known as the Annand Mine, and gave large returns for some time, when, owing to the death of the proprietor, little work was done until it came into the hands of the present owners.

The Symonds-Kaye property has been intermittently worked. In this district operations have been confined to the south dip of the anticlinal, and there is a large unworked but promising field on the reverse or northerly dip. The yield of the district to date is 40,606 ounces, valued at \$861,514.00, from 26,140 tons of quartz.

OLDHAM.

Work was begun in this district in 1862, and the following year saw eight crushers in operation. Up to 1884 an annual production of about 1,200 ounces was maintained, the yield in 1870 being 2,052 ounces from 2,644 tons of quartz. Among the leads worked during this period may be named the Britannia, Ohio, Sterling, Hall, Bonanza, Mayflower and Frankfort. At times very rich returns were made from small lots of quartz. Veins which passed obliquely from bed to bed were at one time quite extensively worked, and although small often gave rich returns, notably in one case 160 ounces from 10 tons of rock. The work in this district having been, during this period, largely in the hands of individual operators on small areas, the surface is greatly cut up, and the veins practically worked only along the outcrop. Some work was done on a cross lead in the eastern end of the district, and in 1893 the Dunbrack lode began to show valuable quartz. Operations were carried on briskly up to 1893, and the following returns were made :-

e gold

rooke, Lake Salmon nay be ptville, Ecum

in the icts, as been are the ins.

ards on er place d much ts apex side by tz were secured, wo tons. tors, and roperties e lode at s at this proved a this one In West

In Wester, having
I)eWolf
American
profit of
I. These

ding 300

YEAR. TONA.	Ounces.
1885	2.360
1885	9 100
18861,026	0.500
1407	
1000	
1889 1,391	2,709
1 100	2.774
1890	9 900
1901 2.019	
1900	
1893 (9 mos.) 2,389	3,171

The operations of Mr. Hardman were profitable, and conducted with skill a l economy; and this pleasing page in the history of this district is due to his enterprise and industry.

Operations in the leading mines were discontinued from causes other than the impoverishment of the properties, and little has been done since.

MOUNT UNIACKE.

Mining began here in 1867, and during the next few years several companies were working; however, the returns, which rose in 1868 to 3,247 ounces, were maintained for twenty years at figures varying from 100 to 1,700 ounces. During part of this time considerable attention was paid to the slate belts which were found to carry regular values over considerable areas. It may be found that in this district the best returns will be secured from these belts which often carry veins, anriferous, but not constant in their dimensions or values

About this time an important discovery on the Withrow areas at South Uniacke diverted attention to the new district. A vein which yielded excellent returns was worked by the Withrow Company, and its extension was mined first by Thompson and Quirk, and afterwards by the Golden Lode Company. All these parties realized profitable returns. Interesting information on the system of mining, etc., carried on by Mr. A. A. Hayward at the Golden Lode mine, has been contributed by that gentleman to the proceedings of the Nova Scotia Mining Society.

RENFREW.

Work was begun here in 1862, but little was done until 1866, when the returns showed £,423 ounces from 6,003 tons of quartz. The following year showed an increase of 1,500 ounces, succeeded by a decrease to ten ounces in 1874. Since that date the yield has fluctuated up to 1,679 ounces. Among the earlier companies may be named the Hartford Colonial, Ophir, Renfrew and McClure. In 1883 the Empress Company was started, and worked with much energy and ability. A suspension of activity followed

for some years; but in the autumn of 1899 returns from the Big Five and Jublice companies showed an extremely high grade of ore, one return giving from about 15 tons of quarts nearly \$8,000.00.

A very rich strike was made on the property of the Mesers. Thompson in the spring of 1900, and up to date 272 tons have yielded 5,071 ounces, or about \$100,000.

CARIBOL'.

This district includes the two districts generally known as Moose River, and Jennings, or Caribou. It was proclaimed in 1870. The returns, although not large for a number of years, showed good averages, and in many cases proved satisfactory. The North, South, Hyde, Comstock and other leads were worked, often with good returns.

In 1881 the Moose River Gold Company carried on extensive workings, but they were abandoned, as the value of the leads diminlahed. Before closing, however, this company demonstrated that it had extensive deposits of low grade ore on its property. Since their large amounts of this low grade ore have been passed through their ten stamp mill by Mr. McGregor, and show an average of about six pennyweights to the ton. As these operations were profitable on a small scale there should be an opportunity for systematic working. On the Tongnoy and adjoining properties there appear to be similar supplies of low grade ore in addition to workable veins, which are being extensively developed by the Touquoy Gold Mining Company, the Colonial Mining Company, and others. The slate and surface earth of the Moose River district has received much attention so far, and with satisfactory results, even on a very small scale of working, and in this connection the following figures are interesting. There were crushed in the Moose River mill from 1881 to date 38,054 tons of slate and low grade ore, with some lots of richer quartz, yielding 11,154 oz., 19 dwt., 13 gr.

The returns made by Mr. Touquoy, from his ten stamp mill, from 1888 to 1898, including some lots of fairly rich quartz, show that 60,943 tons of slate and surface ground yielded 8,640 ounces, an average of about 2 dwts, 20 grains. This work yielded a uniform profit, and lower averages per ton will be shown further on to have also yielded a profit.

In Caribou in 1884 the Lake Lode mine was opened by Mr. Stuart, and worked by Mr. Wadsworth and by Mr. Saunders, and is now being extensively developed by the Guffey-Jennings Gold Mining Company. Large amounts of gold have been taken from this property, and a bright future is anticipated for it. In the same locality the Dixon, Truro, Hetherington, McDonald, Elk and other mines have been opened and worked successfully. Up to date about 124,500 tons of quartz have yielded 41,814 onnce walued at \$794,472.42.

h skill is due

s other s since.

eal comto 3,247 100 to paid to iderable will be not con-

at South
yielded
extension
Golden
teresting
A. Hayentleman

when the wing year to ten 79 ounces, ial, Ophir, and y followed

SHERBROOKE.

Among the veins that have been worked may be mentioned the Cumminger, Hewitt, Hayden, Blue, Palmerston and Dewar. The following table shows the returns year by year from the district:—

anowa site to the state of	• •	
YEAR.	Tons.	Ounces.
1862	663	2,023
1863	•	3,304
1864	2.673	3,419
1865	2.511	3,424
1866	2.853	5,829
1867	7.378	9,463
1868	. 9.880	7,070
1960	. 11.500	5,546
1970	. 11.428	
1871	13.882	
1872	5.243	4,100
1873	7,187	5,026
1874		4,037
1875		5,818
1876		5,176
1877		8,237
1878		6,843
1879		7,389
1880		2,580
1881		2,542
1882		3,356
1883		2,668
1884 1885		1,238
1886		1,341
1887		585
1888		
1889	1.618	243
1890		
1891	464	119
1892	893	179
1893		
1894	708	
1895	3,397	
1896	7.177	3,287
1897	12,659	5 001
1000	16.891	
1899	18,437	4 763
1900	17,711	

representing a total value of about \$2,629,729.00.

The quartz mined in this locality has generally been of a high grade, and there can be no doubt that there are many more rich pieces of ground. The

district moreover presents several localities which should be profitably worked on a low grade basis.

WINE HARBOR.

This district for a few years during its early history yielded some good returns, but for a number of years past attempts at development have not been markedly successful. No resson can be given for this, as the geological conditions are apparently the same as elsewhere, and so ne of the veins were profitably productive. The total eterns to date are 26,872 ounces from 49,070 tons of quartz.

SALMON RIVER.

This district was opened in the year 1880, and work commenced in 1881 with a twenty stamp mill driven by water power. The yield for that year was 1,758 ounces. The following year the Hattie mine was opened on what was believed to be the eastern extension of the Dufferin vein, but the operations were not long continued. In 1885 operations had reached a depth of 150 feet; the vein had been followed for nearly a thousand feet, and the yield that year was 4,924 onnecs from 10,880 tons of quartz. Increased power having been secured, the mill, which had for some years contained thirty stamps, had its capacity increased by ten.

This vein, which in structure is very similar to the saddle back veins met in the Bendigo gold district, Australia, has been one of much interest to miners. It showed at the surface, and on being followed down it divided, going down on each side of the anticlinal, and increased in thinness, giving in places over fifteen feet of crushing material.

The working of the mine has been discontinued more or less during the past few years owing to change of ownership and litigation, but at the date of writing it has been reopened and provided with a thoroughly good equipment, and will, it is hoped, resume its leading position among our gold mines.

The following are the returns received, almost exclusively from the-Dufferin mine:-

YEAR.	Tons,	Ounces.
1881	1.640	1.785
1882	3,460	4.315
1883	7,602	3.885
1884	9.799	3,397
1885	10.880	4.924
1886	11.628	6.509
1887	10.602	3.258
1888	9.925	3.354
1889	7,633	2,032

grade, and

ed the

follow-

••	Tons.	Ounces.
) EAR.	0.435	2.070
1890	6,415	1.406
* 0.00	3 7 211	
1895	1,467	
1000		
1897		

TANGIER.

For the first few years operations were much hindered by the numerous small areas held by individuals. The greatest production was during the years 1869-70-71, when 6,988 tons of quartz gave 5,099 ounces. During all other years the yield has been under one thousand ounces. Recent developments in the northern part of the district, and a short distance east of the Tangier River, promise a development on a larger and more permanent scale. The total returns to date are 22,831 ounces, valued at \$433,802.90, from 46,938 tons of quartz. In this are included the returns from the Mooseland mines in the northern part of the district, which has not yet received the attention it deserves.

FIFTEEN MILE STREAM.

This district, situated at the head waters of the Sheet Harbor River system, has always been largely an object of interest to the people of Pictou County. Its comparatively inaccessible position retarded its development for a number of years, but roads are now built into it, and during the past few years its undoubted value has been shown by the enterprise of New Glasgow capitalists. The following sketch of the principal mine operating in the district will be of interest:—

In 1886 the Egerton Gold Mining Company succeeded the Hall-Anderson Company at Fifteen Mile stream. They worked with a 10 stamp mill for four years and were fairly successful. In 1890 the "Egerton" sold out to the present proprietors, who organized the New Egerton company, and placed Mr. James A. Fraser in charge as manager. A new 15 stamp mill was at once erected, and the new company was soon on a good paying basis. During the first two years (1890 and 1891), nearly 4,500 ounces of gold were secured. Though operations were confined during the next two or three years mainly to developing the property, the yield of gold continued sufficient to meet all running expenses and pay for extensive improvements in machinery, etc. In 1893 an amalgamation was effected with the Stanley Gold Mining Company, which had been operating for some time in the district, and had erected a large dam and a 10 stamp water mill. During

the years 1825-96, and the early part of '97, the company was particularly successful, the yield of gold for 2½ years averaging 225 ounces per month. In the summer of 1896 a new 30 stamp mill had been built, thoroughly equipped with new air compressor, etc. The first half of 1897 saw great attention paid to development work under the guidance of an experienced mining engineer. Notwithstanding this the mine continued yielding its regular returns till a severe crush in October caused a temporary abandonment of work. In May of the following year operations were resumed on an open cut with a belt of over 120 feet.

A Ledgerwood overhead cable-way has been put in, succeeding the old fashioned tramways as a carrier of rock to the mill. Ten additional stamps have been placed in the Stanley mill, which is now being operated with 20 stamps. This, with the 30 in the "Egerton" mill, makes a total of 50 stamps.

The property of the company embraces 300 mining areas at Fifteen Mile Stream, together with 4,000 acres of woodland from which fuel is obtained.

In 1896 the company was incorporated under the laws of Nova Scotia as the New Egerton Gold Mining Company, (Limited).

The returns of this company show that since their purchase of the Hall-Anderson Company property, 35,698 tons of quartz have yielded 17,617 ounces of gold.

LAKE CATCHA, (CHEZZETCOOK.)

The first mention of this district appears in the Mines Report for the year 1882, when the Oxford Gold Mining Company erected a complete milling and mining plant, with the necessary stores, dwellings, etc., and commenced mining operations. Other parties did considerable work during the same year. In the following year the same any produced nearly 2500 oz. of gold from 1475 tons of quartz.

In 1884 the Oxford Co. had proceed in their development work to such an extent that they were spoken of as owning one of the best mining plants in the Province.

A 10 stamp :...'I was built during the year 1886 by the Cambridge Company. Work was continued by the Oxford people, who in 1886 discovered a new and very rich lead. Up to the year 1888 over 40 leads had been cut on the Oxford property, and by 1891 the Company had discovered over 100 leads and done a large amount of development work. More or less work has also been done in the district by Jno. H. Anderson, Dr. Cogswell and others. During the present considerable development work has been done in a new part of the district and some of the old properties have been re-opened. The prospect for the district appears to be promising.

The returns from this district have not appeared in separate form in the

erous g the uring lecent e east more led at

h has

River Pietou opment he past f New erating

nderson
mill for
l out to
my, and
mp mill
ng basis.
of gold
two or
outinued
ments in
Stanley
in the
During

Mines Report for those years during which it has been worked only to a limited extent. The total returns show that 23,023 tons of quartz have yielded in round numbers 24,340 oz., valued at about \$462,467.

STORMONT.

The returns to date from this district show 66,545 ounces from 176,170 tons of quartz, etc. The history of this district for many years was similar to that given of the others. Some rich leads were worked profitably, but their abandonment left the district non-productive for a time. Attention has however been always directed to it as irequent rich returns were made from new veins, and numerous indications of gold bearing veins were found over a tract several square miles in extent. Finally a new anticlinal system, known as the Richardson, was found to the north of the Mulgrave or Isaac's Harbor anticlinal. A number of mines have been opened on these and on the Country Harbor Narrows and Forrest Hill anticlinals. The Doliver's Mountain and Richardson section present large and strong veins and belts, holding good gold values and highly mineralized. There is urdoubtedly room here for several mines like the Richardson, worked, however, with more attention to the tailings and concentrates. The workings of the Richardson mine, of which I give a few details further on, are of particular interest, as they have materially added to the proof that we can mine and work profitably deposits which a few years ago were considered beneath the miner's notice.

The output of the Richardson mine to date has been 21,623 ounces from 240,356 tons of quartz—and of the district. 66,545 ounces of gold, valued at \$1,264,370.90.

The followare remarks from the paper by Mr. Andrews, referred to elsc-where in this report, are interesting:—

"The Richardson belt is very heavily mineralized; and there is great cause for regret that only the free gold is saved. The results of a careful and elaborate series of tests of the tailings from this mine made by Mr. F. H. Mason, are somewhat surprising to many who do not consider the auriferous ores of Nova Scotia worth concentration. But facts speak loudly for themselves, and, much as we would like to have all our ore free milling,

the desire does not alter the refractory nature of some of it.

"According to these tests, made when the ore being crushed was of an unusually low grade, the average loss was 1 dwt. 18 gr. per ton, A sample of tailings from which all the concentrates were not extracted gave 1.3% of concentrates, which had an assay value of 1 oz. 10 dwt. 1 gr. per ton, and still left a value of 1 dwt. per ton in the tailings. Another sample of tailings gave 6.3% of sandy concentrates, having an assay value of 1 oz. 1 dwt. 13 grs. per ton of concentrates. In neither sample was free gold or amalgam detected. The majority of the arsenical prophyrites is contained in the slate; some samples of this slate assay very high. A chance sample gave the surprising result of 28 oz. 8 dwt. of gold to the ton of concentrates,

to a

6,170 similar y, but ention made found system,

n theso . Tho

is unowever, of the rticular ine and ath the

es from , valued

to elsc-

is great a careful Mr. F. sider the k loudly milling,

as of an A sample 1.3% of ton, and e of tailz. 1 dwt. or amaltained in e sample centrates, and yielded 36% of its total weight in concentrates. Two more assays of this slate gave concentrates valued respectively s* 4 oz. 2 dwt. 8 grs. and 5 oz. 6 dwt. 12 grs. per standard ton of concentrates. As this slate contains so little free gold but a small portion of it is crushed.

"Near the surface the slate is soft and partly decomposed. In this condition it yields considerable free gold when milled. As the depth increases the slate becomes harder, increasing perceptibly in bulk, and in the quantity and quality of its concentrates. Below a depth of about 100 feet it contains so little tree gold that it is unprofitable as a free milling ore.

"An analysis of clean concentrates taken from the sluices of the mill gives

the following composition :--

Siliea	2.65
Iron	
Sulphur16	
Arsenic	
Copper	ace "
Zincni	

An assay of these, concentrates gave 2 oz. 14 dwt. 21 gr. of gold per ton. A chlorination test of these concentrates obtained an extraction of 97% of gold contained.

"The cost of an eight or ten ton chlorination plant erected in Nova Scotia

will be about \$4,000, exclusive of building."

"Up to the present time this property has produced 43,000 tons of oro, which goes to show that the mining of low grade ores in Nova Scotia at a reasonable cost per ton, has got beyond the experimental stage and is a reality. The handling of the refractory ores has yet to be experimented with, and from the appearance of nearly all the ore that I have seen al ng the Gold group anticlinal, I am of the opinion that material for the experiment is not lacking."

I am able to give the following interesting figures :--

HALIFAX, May 30th, 1898.

Dr. E. Gilpin, Inspector of Mines:

DEAR SIR,-

As promised, I beg to furnish you with an exhibit of the operations of above company for the year just closed, as follows:

FORTY STAMP MILL.

Total tons crushed	tons.
Total gold won	
Product per ton 2 dwt.	6% gr.
Cost per ton, labor	1.151
" all charges, including renewals to	
	1.831

For the past year, after providing and paying for every dollar of expenditure in connection with the running and thorough maintenance of the property in the highest state of efficiency, and leaving a balance of several hundred dollars on hand, the original shareholders have received dividends equal to 291% on their outlay.

Yours truly,

J. W. CRICHTON, Secretary.

Cost of working and all expenses in connection with the "Richardson Mine," Isaac's Harbor, Nova Scotia, for month of May, 1898, embracing everything except insurance and wear and tear of machinery.

WAGES AS FOLLOWS:

WAGE.			
4 Drill men	475 25		
14 Drill men	615 90		
	102 00		
5 Quarry men	163 40		
2 Engine drivers at pit	83 70		
2 Timber men	66 70	•	
	61 80		
2 Trolley men " "	59 85		
1 17	48 00		
1 Might foremen	89 00		
at Tile alcomithe	96 20		
() (I) - a month own	78 15		
1 Cit of Ungineer	55 00		
4 4 1 mm m from	51 00		
o Ti deigrope of Dilli	78 50		
2 ()	68 40		
0 9 8 111 Lando	219 90 100 00		
Management	100 00	\$2,462 7	5
		Ψ2, 102 .	
OTHER EXPENSES.			
Olnen La. L.	2189 00		
140 Tons coal	171 00		
11 18 : aniaballuar (3/8/109//			
THE of choose and dies			
a and tannetts	- 00		
157 -4-			
Timber handles	15 00		
Shovels, picks, drills and handles			
The state of the control of the cont			
Horse feed			
Royalty	25 00		
Miscellaneous		\$ 811	99
		2000	7.4
Total		\$3,274	14
lotal			

A. B. COX.

Isaac's Harbor, June 6th, 1898.

Manager.

In the Western part of the province Malaga, Brookfield and Whiteburn have been the largest producers. These districts are comparatively new, and at first some very promising veins were opened and extensively worked, but little is now being done at Whiteburn and Malaga. Brookfield is interesting as it has the first chlorination plant built in the province. This was put up by Mr. W. L. Libbey and associates, and is run in connection with a valuable property owned by them. This plant has been in successful operation for some time. Mr. Libbey is prepared to treat, on reasonable terms, any parcels of concentrates that may be forwarded to him, guaranteeing an extraction of 90 per. cent of the assay value. He is also prepared to make working tests of small quantities of ore as to their adaptability for chlorination, etc. This should supply a favorable opportunity to our gold miners to get information as to the value of their concentrates. Mining has also been carried on at Leipsigate, Gold River, Block House, and a number of other places in Lunenburg and Queens Counties, but they have not yet been worked on a scale commensurate with their promising indications. There is no doubt that all these districts would repay proper examination and development, as they differ in no respect, that I am aware of, from those opened to the cast of Halifax.

Among the other districts that of Central Rawdon is interesting. Here the veins run across the measures, and two of them are found to unite in depth, thus reversing the method in which they are often presented in the other districts. Since 1880, 4,814 tone of quartz yielded 6,648 ounces of gold.

pendi-

e pro-

everal

idends

ary.

ardson

bracing

99

MINING.

At many of the mines shafts are sunk at the proper places, and ground systematically blocked out, and the operations underground are carried on with economy and system. Properly designed machinery is largely used for pumping, hoisting and crushing, and the mills are carefully built, and run with proper attention to the details essential to this metallurgical process, which is really much more complicated than would be believed from a casual glance at a mill. Interesting details on the best system of milling the quartz of this province are given by Mr. Hardman in a paper published in the Transactions of the Nova Scotia Institute of Mining Engineers, for the year 1892-93. These Transactions give many other interesting papers touching on gold mining practice in our gold fields. It can be said for the benefit of investors, that our gold mine managers can stand comparison with those of any other country in their ability to mine systematically, and to treat economically the extracted material.

To show this, I quote from a paper read by Mr. A. A. Hayward at a late meeting of the Nova Scotia Mining Society, in which he gave his experience in shaft sinking at South Uniacke and Montagu. At South Uniacke it was decided to sink a five feet six inch by twelve feet shaft 404 feet through hard quartzite. The work was completed in 140 days, of which 16 were occupied in timbering. The average rate sunk was 3.02 feet per day; the total cost of explosives per foot sunk was \$1.22; and the average monthly rate of sinking was 71 feet 6 inches, the highest rate being 85 feet in one month. The total cost, including management, fuel, and everything chargeable to the shaft, was \$4,647, or \$11.50 per foot, the shaft being left completed, timbered, and ready for occupation.

At Montagu, during 1897 the shaft of the Golden Group Company was sunk 100 feet below the 200 feet level. The work would have been finished in 25 days, had several days not been lost in taking down and saving a small rich lode on the foot wall. The cost of this sinking per foot was a little in excess of that at Uniackc. These two instances show that the ordinary cost of sinking can be most materially reduced when the work is fully planned out, and carried on systematically.

A paper read by Mr. W. L. Libbey at the same meeting, gives some interesting figures as to his experience in mining and milling at Brookfield, Queen's County. He compares hand and machine drilling as follows:—

In making comparisons with the cost of landing ore at the rock-breaker on other mines, it must be remembered that the fissure vein at Brookfield averages not over 24" in width of crushing material. The extreme depth perpendicularly of the working of the mine is 450 fcct. The figures given

include the cost of sinking and drifting, and also in the first table is covered the cost of excavating a large chamber to hold a double plunger Northey pump, and a cistern capable of holding mine water for twelve hours.

During six months, from May 1st, 1897, to November 1st, 1897, 5,606 tons of ore were sent to the mill at an average cost of \$2.54 per ton, as

follows, for ore landed at the rock-breaker :-

ound do on ed for drun occess, som a ng the hed in or the touch-benefit those or treat

a late erience it was

hrough

6 were

y; the

onthly

in one charge-

ft com-

finished saving a at was a hat the work is

some inrookfield, ws: reaker on brookfield ne depth res given

Labor (which includes blacksmiths and deck men) \$11,173	00
Timber and poles	
	20
	60
	96
	00
	00
	00
Candles	
	82
Fuel at pumping station and mill. 1,046	
Explosives	
Iron, including rails for tracks	
Miscellaneous expenses	
15 Lumber 45	UU
'total	91

Following are the figures for three months of work with the air plant. It should be borne in mind that not only have green men been broken in, but the method of stoping is being gradually changed from breast stoping to back stoping. The results are especially gratifying to the Brookfield Mining Co., as many old timers have flatly stated that a small lead could not be worked as cheaply by an air plant as by hand labor, and in one instance recently an air plant has been discontinued and a return made to hand drilling. The months taken are January, February and March, 1898, during which time 2,840 tons were sent to the mill at an average cost of \$2.44 per ton, as follows, for ore landed at the rock-breaker:—

Labor (which includes blacksmiths and deck men) \$5,078	95
11mber and poles	
Shovels	00
Picks o	00
Blacksmith's coal, 44 tons at \$10.66	
Charcoal, 150 bushels at 15c	
Hoisting ropes	
	-
Explosives 717	_
Explosives	-
Iron, including rails for tracks	45
Miscellaneous expenses	00
Lumber	00
Total	42

The result thus far is apparently to place our ore at the deck head ten cents per ton cheaper with an air plant than by hand work. We are, however, doing more than twenty-five per cent. more of sinking and drifting with the air plant than formerly by hand. In fact, it would be impossible to place men enough in the mine to equal by hand the work now done by

Following is a table showing the expense of running the 20 stamp mill for six months, commencing September 1st, 1897, and ending February 1st, 1898. During this time 5,910 tons of ore were milled and concentrated at

an average cost of 60 cents:

warana C	cost of 60 cents:		
Agruse c		\$876	00
Fuel	08 (0-ma)	860	00
Labor-	-2 firemen	900	w
44		252	00
66	a managem hove	242	
64		300	
		75	00
Miscel		397	53
Cost t	cotal for shoes and dies	47	55
			00
Screet	n wire. 192 it, at out.		
.,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	Total	\$3,546	73
	Total		

At the Richardson mine, Stormont, the lead has been found to be from 6 to 18 feet wide. In 1894 the mill contained twenty stamps, and the cost of mining and miling was \$2.90 per ton, inclusive of taxes, insurance, depreciation, and all charges. The beit was composed for the most part of one large iode from 1 & 4 feet in thickness, and a varying number of small lodes intermixed with slate. In places nearly the entire belt is quartz. Shortly after that date, the The slate is auriferous as well as the quartz. mine was remodelled in every respect, so that nine hand drills could produce regularly 2,000 tons of ore per month. The mill was enlarged to 40 stamps of 850 lbs. weight each, dropping 99 times per minute, and crushing very finely. With this plant, lighted by electricity, and driven by a central engine, the cost of mining and milling, including all charges, was \$1.65 per ton. The belt is very heavily mineralized, and the concentrates show gold up to 5 oz. per ton.

The paper from which these brief extracts are given was read by Mr. C. F. Andrews, at the Montreal Meeting of the Canadian Mining Society in 1897. I may add that arrangements are now being made to save and treat the concentrates, which should yield a considerable additional prof.

ten nowting

e by

mill 1st.

ed at

003

from

d the

urance.

part of

f small

quartz.

te, the

produce

stamps

ng very

central

1.65 per

ow gold

y Mr. C.

ociety in

and treat

of .

GENERAL REMARKS.

The climate presents no obstacles to mining, and operations are carried on throughout the year. There are no heavy falls of snow, and no heavy freshets, or rainy seasons. The mines themselves are with very few exceptions dry, and pumping charges are not heavy, when proper provision is made for diverting the surface water. All the districts are accessible by roads passable for heavy loads; some are close to railways; and many are on tide water, or only a few miles removed from it. Few, if any, gold mining districts are more favorably situated for procuring supplies. Mills, boilers, engines, and all the innumerable mechanical alds to mining are manufactured in the country, and within short distances of any district. Labor is plentiful. The men are quiet and orderly, and no communities are more law-abiding than our mining villages, which are always provided with schools, and frequently with churches. The men are intelligent and good miners, and in the mining districts of Western America they are preferred to all others.

It will be observed from the preceding notes that there are three additional openings for extending the operations of the gold mines in this province: treatment of the tailing and concentrates, alluvial mining, and the development of what might be termed the gold in the mineralized deposits, in part only free milling.

As to the first of these openings, the treatment of the tailings and their concentrates, it may be remarked that its feasibility has been demonstrated at the Brookfield mine, and the reports of the assayers show an equally good opening for the treatment of this material by chlorination, or an allied process, in every district.

It is true that the expense of these plants may preclude their adoption by many of the smaller mines and districts. As however, the greater number of the districts are readily accessible by water or railroad, it would appear that, if even the concentrates were saved and shipped to a chlorination plant, located at a convenient point, the Government would receive an increased royalty, and many mines would have an important additional source of revenue secured to them. It is the generally expressed opinion of all who have looked into the matter that such a plant, for the treatment of concentrates, located at Halifax, would greatly benefit the mining community.

The subject of the third opening has as yet received comparatively little attention. It is known that at a number of places in the province, veins and beds, carrying as essential ingredients lead or copper ores, or both, con-

tain gold and silver in amounts commercially valuable. It is also known that there are other localities in which there are deposits, sometimes showing free gold, which carry gold and silver in the sulphurets in amounts sufficient to warrant their reduction where the percentage of the concentrates is high enough. Localities coming under this latter classification occur in the Cobequid Mountains, in the counties of Pictou and Guysboro, and in northern Cape Dreton. In the first and last named districts the streams show sights of gold, and a certain amount of free gold occurs. In the Whycocomagh, Middle River, and Cheticamp districts, there are extensive mineralised zones in the Precambrian rocks, which present a good field for investigation. Free gold occurs in the alluvium in promising amounts; fine free gold is seen in the silicious beds, and the sulphurets carry gold. Given these data it would appear reasonable to expect that no great amount of prospecting should disclose localities for profitable mining. Examination will probably reveal points where a more particular enrichment has been caused by dykes, etc., and the mining developments of Western America will find a parallel in Nova Scatia.

CONCENTRATES.

Mr. F. H. Mason, Metallurgist, of Halifax, has kindly contributed to this report a memo. on the tailings and concentrates of our Nova Scotia gold ores, as follows:—

Until the beginning of the year 1897 it may be said that no systematic attempt had been made to obtain anything but free milling gold from the ores of Nova Scotia. The class of machines used was the "Golden Gate," a machine requiring constant attention, and even their giving results far from being satisfactory. Notwithstanding the disadvantage resulting from the class of machine employed, the concentrates obtained in a custom mill at Olcham were of such a high value that it is surprising the attention was not given to the concentrating of tailings in other part of the province at a much earlier date. The average value of the concentrates obtained at Oldham was from sixty to seventy dollars per to concentrates obtained at Oldham was from sixty to seventy dollars per to 2,000 lbs. I am given to understand, however, that no systemate record was kept of the percentage of concentrates saved, nor of the gold lose in the tailings after they had been over the concentrators.

At the Lake View Mines at Waverley, eight Frue Vanners were erect to treat the tailings from 30 stamps, or two-thirds the number required good practice. So that here, although the class of machines was excelled the number of machines was insufficient to handle the quantity of tailing going over them, and good results were consequently not obtained.

value of the concentrates, which at thus contained a considerable propertion of sand, ranged from eight to thirty deliars per ton.

About 1881 W. A. Mirral of Colorado, started a chlorination plant at Waverley. The plant consisted of a very small hand roasting furnace, and a home made Plattner Chlorination plant of the crudest description. The whole thing proved a complete failure. Two years later B. A. Temple of Waverley joined Mirral, and put in a new furnace and new leaching and precipitating vats, &c., to start a Plattner plant. I have examined this plant and have never seen or read of anything guite like it.

The early attempts at cyanking met with a fate similar to that of chlorination. Dr. Kendall and his assistants came from he States and erected a plant at North Briokfield, with a view to treating an old tailing dump at that time (about 1893) owned by the Brookfield Gold Mining Associates. Dr. Kendall experimented for some time without obtaining any results, and the Brookfield Associates who had employed him, gave him notice to quit, and continued the researches themselves for some time with no better results, and the attempt was abandoned.

These early failures naturally had a bad effect upon this branch of the gold mining industry, and gave the general public the false impression that even if the concentrates of Nova Scotia contained gold, it would cost more to extract it than the gold was worth, and the concentrates were allowed to go away with the tailings down streams and amongst brush wood, never to be recovered. At the time of my arrival in the province it had become generally recognized that the ores of Nova Scotia were free milling ores, and that gold which could not be saved in the mortars and on the plates of the stamp battery was not worth saving.

l at once turned my attention to studying the remarkable phenomenon of a country with a large area of auriferous formation, in which all the gold was said to be of a free milling nature, with a view to establish either the truth or fallacy of this then generally recognized opinion. By obtaining samples of a number of old tailing dumps and making a number of careful assays of them, I soon discovered the complete error into which the gold mining fraternity of Nova Scotia had fallen. Assays of these samples gave from two dollars to seventeen dollars per ton. Of course in many cases a natural concentration had taken place, and these samples did not represent actual losses in the mill, but it did clearly prove to me that considerable quantities of gold were being lost, a large percentage of which might have been saved by the employment of proper machinery.

During the last five years I have made a large number of assays of tailings taken both directly from the end of the plates and also from dumps. I have also assayed a number of samples of concentrates.

The following table of maximum and minimum results from different districts will be of interest:

ed to this

known show-

mounts

entrates

ceur in

streams

In the

xtensive field for

te; fine

Given

nount of mination

has been

America

that no lling gold was the even then sadvantage obtained in orising that ther parts ue of the ars per ton systematic he gold lost

ere erected equired by s excellent, of tailings ined. The

DISTRICT. Maximum. Minimum. Maximum. Minimum.		TAILINGS FROM PLATES.	PLATES.	TAILINGS FROM DUMPS.	Drups.	CONCENTRATES	RATES.
1 dwt. 18 grs. 21 grs. 6 dwt. 10 grs. 3 dwt. 6 grs. 2 dwt. 22 grs. 14 grs. 13 dwt. 2 grs. 2 dwt. 23 grs. 8 dwt. 4 grs. 1 dwt. 18 grs. 16 dwt. 22 grs. 3 dwt. 6 grs. 15 dwt. 8 grs. 1 dwt. 1 grs. 14 grs. 14 grs. 14 grs. 14 grs. 14 dwt. 17 grs. 2 dwt. 12 grs. 1 dwt. 20 grs. 1 dwt. 4 grs. 2 grs. 1 dwt. 20 grs. 7 dwt. 7 grs. 7 dwt. 7 grs. 7 dwt. 20 grs. 7 dwt. 7 d	DISTRICT.	Maximum.	Minimum.	Maximum.	Ninimum.	Maximum.	Minimum.
3 dwt. 12 grs. 21 grs. 6 dwt. 10 grs. 3 dwt. 28 grs. 1 4 grs. 13 dwt. 2 grs. 2 dwt. 23 grs. 1 5 dwt. 8 grs. 1 dwt. 18 grs. 13 dwt. 1 gr. 14 grs. 1 dwt. 1 grs. 1 dwt. 12 grs. 1 dwt. 1 grs. 1 dwt. 20 grs. 7 dwt. 1 grs. 7 dwt. 1 dwt. 10 grs. 7 dwt. 2 dwt. 1 grs. 1 dwt. 20 grs. 7 dwt. 2 dwt. 1 grs. 1 dwt. 2	Cow Bay	1 dwt. 18 grs.	21 grs.				1 oz. 8 dwt.
2 dwt. 22 grs. 14 grs. 13 dwt. 2 grs. 2 dwt. 35 grs. 15 dwt. 4 grs. 1 dwt. 18 grs. 16 dwt. 22 grs. 3 dwt. 6 grs. 1 dwt. 15 grs. 2 dwt. 12 grs. 6 dwt. 12 grs. 14 dwt. 16 grs. 1 dwt. 22 grs. 1 dwt. 17 grs. 6 dwt. 17 grs. 3 dwt. 12 grs. 1 dwt. 22 grs. 1 dwt. 22 grs. 1 dwt. 21 grs. 6 dwt. 17 grs. 3 dwt. 12 grs. 1 dwt. 22 grs. 7 dwt. 12 grs. 1 dwt. 20 grs. 7 dwt. 12 grs. 7 dwt. 10 grs. 7 dwt. 20 grs. 7 dwt. 20 grs. 7 dwt. 20 grs. 7 dwt. 2 dwt. 20 grs. 7 dwt. 2 dwt.	Sherbrook	3 dwt. 12 grs.	21 grs.	6 dwt. 10 grs.	3 dwt. o grs.	ers.	
8 dwt. 4 grs. 1 dwt. 18 grs. 16 dwt. 22 grs. 3 dwt. 6 grs. 15 dwt. 8 grs. 1 dwt. 18 grs. 13 dwt. 1 gr. 14 grs. 2 dwt. 22 grs. 21 grs. 6 dwt. 12 grs. 4 dwt. 16 grs. or 3 dwt. 12 grs. 1 dwt. 21 grs. 6 dwt. 17 grs. 3 dwt. 12 grs. 1 dwt. 20 grs. 7 dwt. 10 grs. 7 grs. 19 dwt. 20 grs. 7 dwt. County.	Carillou	2 dwt. 22 grs.	14 878.	13 dwt. 2 grs.	2 dwt. 23 grs.	13 oz 3 dwt. 3 grs.	l oz. 17 dwt. 21 grs.
15 dwt. 8 grs. 1 dwt. 18 grs. 15 dwt. 2 grs. 14 grs. 14 dwt. 15 grs. 21 grs. 6 dwt. 12 grs. 4 dwt. 16 grs. 1 dwt. 22 grs. 1 dwt. 17 grs. 3 dwt. 12 grs 21 grs. 1 dwt. 17 grs. 3 dwt. 12 grs 21 grs. 1 dwt. 20 grs. 7 dwt. 1 dwt. 10 grs. 7 dwt. 7 grs. 19 dwt. 20 grs. 7 dwt. 7 dwt	Uniacke	8 dwt. 4 grs.	or brain	:	3 dat. 6 grs.	136oz. 1 dwt. 19grs	1 oz. 7 dwt.
2 dwt. 15 grs. 6 dwt. 12 grs. 4 dwt. 16 grs. 4 dwt. 16 grs. 14 dwt. 17 grs. 4 dwt. 16 grs. 1 dwt. 12 grs. 1 dwt. 4 grs. 21 grs. 19 dwt. 20 grs. 7 dwt. 7 grs. 7 dwt. 7 grs. 7 dwt. 7 dwt. 20 grs. 7 dwt. 7 dw	Stormont	15 dwt. 8 grs.	1 dwt. 18 grs ‡		14 grs.	12 oz. 17 dwt.	1 oz. 3 dwt.
2 dwt. 15 grs. traces. 14 dwt. 17 grs. 4 dwt. 16 grs. 3 dwt. 12 grs. 1 dwt. 2 grs. 21 grs. 21 grs. 7 dwt. 20 grs. 7 dwt. 7 grs. 7 dwt. 7 dwt. 7 dwt. 7 dwt. 20 grs. 7 dwt. 7 dwt. 7 dwt. 7 dwt. 20 grs. 7 dwt. 7 dwt	Brookfield		ē	6 det. 12 grs.		4 oz. 14 dwt. 1 gr	1 oz. 8 dwt. 7 grs.
2 dwt. 22 grs. traces. 14 dwt. 17 grs. 3 dwt. 12 grs. 1 dwt. 21 grs. 21 grs. 19 dwt. 20 grs. 7 dwt. 7 grs. 7 dwt. 7 grs. 7 dwt.	Waverley	2 dwt. I5 grs.	-118 Iz		4 Just 16 grs.		1 oz. 2dwt. 4grs.
3 dwt. 12 grs. 1 dwt. 21 grs. 21 grs. 7 dwt. 20 grs. 7 dwt. 19 dwt. 20 grs. 7 dwt. 19 dwt. 20 grs. 7 dwt. 19 dwt. 20 grs. 7 dwt. 10 grs. 10 dwt. 20 grs. 10 dwt. 2	Central [Rawdon	2 dwt. 22 grs.	traces.	14 dwt. 17 grs.	3 dwt 12 grs		
1 dwt. 4 grs. 7 dwt. 19 dwt. 20 grs. 7 dwt. ty. 7 oz. 12 dwt. 20 grs. 7 dwt. ty.	Wine Harbor	3 dwt. 12 grs.	1 dwt.	6 dwt. 17 grs.			
13 dwt. 10 grs. 7 grs. 19 dwt. 20 grs. 7 oz. 12 dwt. 20 grs. 7 dwt.	Tangier	. 1 dwt. 4 grs.	21 6 TS.				
7 oz. 12 dwt. 20 grs.*	Fifteen Mile Stream		1-	19 dwt. 20 grs.		3 oz. 8 dwt.	1 oz. 8 dwt. 4 grs.
unty	Oldham			7 oz 12 dwt. 20 grs.			
	Gold River					99 oz. 16dwt. 18gr	8 21 oz. 1 dwt. 8 gre
	Lunenburg County.					7 oz. 11 dwt. 16g	rs 5 oz 9 dwt. 16 grs.

8 oz. 2 dwt. 8 grs. * | 1 dwt. 4 grs.

8 07. 2 dwt. 8 grs. 1

The results given in the foregoing table clearly show that the stamp mill only saves a part of the gold present in our Nova Scotia ores. It is true in many cases a greater percentage of gold might have been saved by more skillful crushing and amalgamation, but on the other hand it clearly shows the large amount of gold which is going to waste for the want of proper appliances to save it.

Concentrators have now been running for some time at North Brookfield. At Central Rawdon the orc gives 4% of concentrates, having an average assay value of \$30 in gold and \$1 in silver. I am unable to give any values obtained at Brookfield, as the B afield Gold Mining Co., Ltd.,

refuse to give any information on the subject.

Another considerable source of waste in Nova Scotia gold mining is to be found in the waste rock dumps. At many times it has been the custom to dump and stow away on the scaffolds nearly the whole of the slate rock mined, much of which is often auriferous. In a paper read by Mr. C. F. Andrews he gave assays made by himself of rock taken from the waste dump at the Richardson mine. These samples were not average samples of the dump, but were taken on account of the large proportion of mineral (mispickle) contained in them.

The following are the assays and per centage of concentrates:

Contained 30% of concentrates, having an assay value of 28 oz. 8 dwts. of gold per ton.

Assayed 11 dwt. 2 grains per ton, contained 12.05% of concentrates, having an assay value of 4 oz. 2 dwt. per ton.

3. Assayed 17 dwt. per ton, contained 15% of concentrates, having an assay value of 5 oz. 6 dwt. per ton.

Mr. Dimcck milled a trial lot of slate rock taken from his scaffolds at the Central Rawdon mine which he had previously been in the habit of dumping and stowing. The result was a saving of 60 cents per ton in the mill, while the tailings contained \$2.60 per ton.

Undoubtedly many other mines might obtain similar results from material which at the present time is being piled on the waste rock dump or stored on the scaffolds under ground.

I have made a large number of experiments in my laboratory upon concentrates from all parts of the province, and up to now I have not come across a single instance in which any difficulty has been found in obtaining from 90 to 96 per cent. of the assay value by chlorination. Cyanide, on the other hand, has not proved so satisfactory. In one case I obtained a 94% extraction. With this exception 78% is the best extraction I have obtained, while in many cases the extraction has fallen below 30%, even with prolonged treatment lasting over 30 days.

There are other sources of gold in Nova Scotia which up to now have received no attention. The stibnites of East Gore are highly auriferous,

running at times over 50 ounces of gold per ton and showing no free gold by panning. In several samples of copper pyrites I have found over half an ounce of gold per ton. The fahl ores of Cheticamp have assayed as high as \$15 in gold, while samples of rock carrying pyrrhotite have at times run over half an ounce. All of these, especially the first named, will pay for further investigation.

The foregoing remarks of Mr. Mason are interesting and important, and should be carefully considered by our gold miners.

The following table shows the production of the Nova Scotia mines for a number of years. The annual production has varied from 7,275 ounces in the year 1872, the first year during which official returns were made, to the year, 1867, which showed an output of 29,314 ounces. For some years past the production has been in the vicinity of the last named amount, although derived from a much lower average yield per ton of quartz. Thus in 1867, there were 31,386 tons of stuff crushed for 27,314 ounces, while in 1897 there was a return of 26,579 ounces from 76,559 tons of stuff.

NOVA SCOTIA GOLD.

PRODUCTION BY DISTRICTS FROM 1862 TO SEPT. 30, 1900.

	Tops	Total Yield of Gold.		Total Yield of Gold. Average Y per ton		rage Yi er ton.	eld	Value
DISTRICT.	Crushed	Oz.	Dwts.	(irs.	Oz.	Dwts.	Grs.	@ \$19 per Oz.
Caribou & Moose River Montague Oldham Renfrew Sherbrooke Stormont Tangier Uniacke Waverley Salmon River Brookfield Whiteburn Lake Catcha Rawdon Wine Harbor Fifteen Mile Stream Malaga Other Districts	26140 48243 48601 226169 176170 46938 54362 123383 117175 6394 23023 12178 49070 32893 20299	21098 9554 24340 9594 26872 3 16403 2 18563	15 5 17 17 18 18	5 20 22 18 5 10 2	1	18	21 3 18 22 22 23	400,879.08 181,542.87 462,467.80 182,300.65 510,575.68 311,665.75 852,699.22
Total		717881	10	10		1	1	13,639,748.97

NOVA SCOTIA LEAD ORES.

As yet little has been effected in the development of the ores of this metal in Nova Scotia. They occur principally in two geological horizons, the Lower Carboniferous or mountain limestone, and the Precambrian or Laurentian. As the former horizon in other countries has yielded productive deposits of galena, some attention has been paid to it here in this respect. Observation has shown that at many localities lead ore is more or less distributed through : . limestones. Among the places showing it may be mentioned the East and West Rivera of Pictou county; Gay's River and Musquodoboit, Halifax county; Smithfield and Pembroke, Colchester county; Milford Haven, Caledonia and Salmon River, in Guysboro county; and numerous points in Cape Breton Island. At Gay's River it occurs over a large tract of country, in limestone, disseminated in small crystals, in small nodules, and occasionally in thin veins. The average percentage at any place is small, and carries, I believe, up to 15 ounces of silver to the ton of lead. At Smithfield and Pembroke it occurs in greater quantity, and a considerable amount of exploratory work has been done. Here it is four in limestone also, apparently replacing it in part, and as a residual concentration from the removal of the rock. A very considerable amount of lead ore has been shown here at one point, but as the silver contents of the ore were not high the low lead prices discouraged prospecting. It is, however, evident that in this district there are widespread and promising galena ores, which merit much more attention than has yet been paid to them. In Musquodoboit, near the Crawford Settlement, pockets of ore in the granite have yielded lead carrying large amounts of silver.

At Caledonia, small veins of lead ore have received occasional attention. The ore is essentially galena, carrying from 15 to 18 ounces of silver to the ton.

Galena also occurs in considerable amount near Arichat, Richmond county, and on the head waters of the Gold and LaHave Rivers to the west of Halifax. The latter ores carried in some cases as much as 100 ounces of silver to the ton of lead, but, as far as I am aware, no attention has been directed to them. These ores are presumably in the Precarboniferous rocks. On the Salmon River, and near Sydney, in the county of Cape Breton, galena also occura in limestone, at the former place in considerable amount. Near Burnt and Boulacet Harbours, on the Bras d'Or Lake, are quartz veins carrying galena, copper pyrites, etc. Samples have shown 18½ ounces of gold and 97 ounces of silver to the ton. The adjoining rocks are in places heavily mineralised. On the North and Barrasois Rivers of St. Ann's, in

and

balf half high

run for

for a es in the years ount, Thus while

due per Oz.

,472.42 ,524.06 ,5258.48 ,083.80 ,729.56 ,370.82 3,802.90 ,685.53 2,523.45 8,258.55 0,879.03 1,542.87 2,467.80 2,300.65

0,575.68 1,665.75 2,699.22 7,908.90

9,748.97

the county of Victoria, and several other places in that locality, in Precambrian rocks, are veins of quartz and zones of rock showing galena with copper pyrites, blende, etc. A sample of 900 lbs. of ore from McDonald's farm, one mile north of the bridge over the North River, yielded 155 lbs. of lead and 3 oz. of silver.

The most promising deposit of silver lead ore yet found is being developed at L'Abime Brook, Cheticamp, Inverness county, by S. M. Brookfield of Halifax and his associates. The ore occurs as a vein in the Precambrian felsites, etc., and has been traced for several hundred feet, and where opened shows twenty feet of very high grade galena carrying a little copper pyrites.

Mr. Mason, of Halifax, states that from assays made by him and by others, the ore carries an average of one ounce of silver to each unit of lead. Gold also occurs in some assays as high as 14 dwts. per ton, but it does not

appear to be a regular constituent.

From his examination of the openings he considers that 50 tons of ore contains 5 tons of galena, and 1,500 lbs. of copper pyrites. This would yield, allowing for loss in dressing and smelting, 3 tons 1,200 lbs. of lead, and 360 ounces of silver, in addition to the copper. The figures, coupled with the accessibility of the mine, etc., etc., and the duty on lead, would indicate very promising results. Other deposits of galena are reported in the same district, but as yet little work has been done on them. Galena ores occurnortherly from the Claticamp River to beyond the McKenzie River, and native silver and carbonate of silver occur in the valley of the latter river and its tributaries. It is probable that the opening of the Cheticamp mine will lead to a careful prospecting of this district, and it is, I feel sure, from information that I have gathered for some years, one of the most promising localities in the province.

From these remarks it will be seen that the lead ores of the province are widely scattered, and have hitherto received little attention. It is very probable that the Lower Carboniferous limestones, which cover a great extent of country, may reasonably be expected to afford workable deposits at favorable points; for instance, in proximity to intrusive rocks of later dute, or wherever they appear to have been in a position favorable to the concentration, aggregation, or deposition of such ores by aqueous agency.

Prewith nald's bs. of

eloped eld of brian pened yrites, nd by f lead, es not

of ore lyield, d, and d with indicate e same s occur er, and er river p mine e, from omising

ince are
is very
t extent
it favordute, or
concen-

NOVA SCOTIA COPPER ORES.

The remarks I have made about the lead ores of this province apply in a general manner to those of copper. The metal occurs frequently, in the metallic form, in voins and joints in the Triassic trap range running along the south shore of the Bay of Fundy, and in the isolated masses of the same mineral on the opposite side of the Bay. It also occurs in a similar manner in the consolidated ash accompanying the trap, and has been observed in the sandstones of the district, which are referred to the same geological horizon. As yet none of the deposits have, from the superficial tests hitherto made, appeared to be of permanent value. In some localities it is observed disseminated in the rock, and all such occurrences should be carefully tested. It is well known that in the Lake Superior district very small percentages of copper have yielded rich returns to systematic and economical systems of mining and milling. It is true that the two districts differ widely from a geological point of view, but as both yield metallic copper in masses, etc., it does not appear unreasonable to expect that here, as at Lake Superior, it may be found in some localities, scattered in fine grains in the trap, in quantity sufficient to allow of its profitable extraction. Copper also occurs native in the Lochaber district, in Antigouish County, and at Cheticamp, Inverness County.

The carboniferous strata of Cumberland, Pictou, Colchester, Autigonish, and other counties, frequently show outcrops of nests and layers of vitreous sulphuret and green carbonate of copper, sometimes associated with coaly matter. Prospecting has shown these outcrops at Maccan, Wallace River, Pugwash, Tatamagouche, Athol, Oxford, River John, Salmon River, Caribon, Durham and Waugh's River, in the counties of Cumberland, Colchester and Pictou. A few attempts have been made to work these deposits, but the c. Ithough rich is variable and irregular in its occurrence, and work .doned after the extraction of a few tons. A good deal of work nable a lone during the past two years on some defined beds of sandstone and shale, more or less uniformly impregnated with these and other ores of copper. Among these localities so tested may be named Doherty Creek, River Philip, Malagash Point, Fox Harbor, Gulf Shore, New Annan, Wentworth, and Henderson's Settlement in Cumberland County. The ores are also present at some points in beds of clay resting on cupriferous sandstones, etc. Working samples are reported to have yielded as high as 18 per cent. of copper, with in some cases small amounts of gold and silver. It is understood that the work of testing these deposits will be continued, as the parties interested have so f r been much encouraged by the results.

The numerous outcrops of the ores over so wide a tract of country warrant much more attention than has hitherto been paid to them.

At East Dalhousie, in Kings County, there are numerous shows of copper ore, which have hitherto been almost unnoticed. Copper pyrites occurs at Blandford, Lunenburg County, and at several points in Queens and Yarmouth Counties, and is commonly present, in small amount, in the auriferous quartz veins of our gold fields.

The district extending from the head waters of the East River of Pictou, along the county line, to Polson's Lake, in Antigonish County, has yielded numerous indications of the presence of copper ores.

At the head waters of the East River specimens of copper ore occur with a gangue of carbonate of iron, but no attempt has been made to search for their source. In the vicinity of the Garden of Eden, there are veins of spathic iron ore, up to several feet in thickness, holding crystals of copper pyrites.

At Lochaber there are a number of voins carrying copper pyrites, and associated with diorite dykes. These veins have been prospected a little, and are apparently valuable, as the copper contents in the case of the largest vein, about six feet thick, were returned from large average samples at 19 per cent. From this point the cupriferous belt has been traced about four miles to Polson's Lake. Here the presence of large rich boulders instigated desultory prospecting for a number of years. Finally a vein of spathic ore, holding copper pyrites, was found in the year 1875, and traced several hundred feet. Its width, as proved by several shallow pits, varied from 6 to 11 feet. Large average samples yielded from 5.6 to 11.7 per cent. of copper. The distance of this district from a railroad and a shipping point, has apparently postponed its development, although it is probably well worth the cost of a railroad. On the Salmon River, in Guysboro County, some veins were tested a number of years ago, containing copper pyrites and erubescite, and yielding as high as 39 per cent. of copper. The age of the rocks holding these deposits in Pictou, Antigonish and Guysboro counties, is I believe, determined as Devonian, and is in favor of their richness and permanence.

At Ohio, St. Joseph, Brierly Brook, and other points in Antigonish County, deposits of copper ore occur at the junction of carboniferous with older rocks. A little prospecting has shown small veins of rich ore, but as yet enough work has not been done to enable a correct idea to be formed of their extent.

The reports of Mr. Fletcher, of the Canadian Geological Survey, on the Island of Cape Breton, contain frequent references to copper ores. The following places may be mentioned: Benacadie, White Granite Hills, Gillis Brock, Spruce Brook, Irish Cove, East Bay, Washaback, Middle and North Rivers, French Road, Gillis Lake Road, Mira, Cape North, etc.

At Washaback, the cres yielded in addition to the copper, gold up to about one ounce to the ton. At Eagle Head, in Gabarus Bay, Cape Breton County, several shafts have been sunk showing quite large beds, carrying mixtures of copper pyrites with other ores. Those acquainted with the operations consider that the deposits are well worth further examination.

At Cheticamp, Inverness County, there are numerous signs of copper deposits, and occasionally attempts have been made on a small scale to determine their value. This district shows copper ores over so large an extent of ground, that it is probable that this metal will be found present in workable amounts, as well as the lead already referred to. At George's River, near North Sydney, a little prospecting has been done on a very promising outcrop. On the Coxheath Mountains, a few miles to the south, a quite extensive development has shown the presence of several leads carrying copper pyrites, with some silver, from three to twelve feet thick. The copper contents of these beds vary from 3 to 10 per cent, from extraction of large lots. The shafts and levels have yielded several thousands of tons of good ore. Several causes, including financial ones as well as the low prices of copper, have retarded the development of this property; but as the outlook is now encouraging, arrangements are being made for working it.

In the Island of Cape Breton the copper ores occur in the Precambrian felsites, etc., at Gabarus, Coxheath, George's River, Cheticamp, etc., and are found also in Carboniferous strata.

These brief remarks on the occurrence of copper ores in Nova Scotia will show that they are very wide-spread. They have as yet received almost no systematic examination, and local capital has preferred to venture in better known channels. It may, however be justly claimed that the province presents, in several districts, unusually promising deposits, and that in better known and richer communities, they would before this have supported at least several profitable mines.

IRON ORES.

The iron ores of Nova Scotia are found in almost all its geological horizons. They are of every variety, and in a number of dist. its are present in large amounts. Practical work has shown their adaptability for the manufacture of iron and steel. At present the extraction of iron one is confined to the mines of the Nova Scotia Steel Company at Bridgeville, in Pictou county. In view of the presence of large iron and steel works at Sydney, in Cape Breton, the reason for this small output may be presented to explain any surmise as to the non adaptibility of our ores for economic purposes. On Bell island, near St. John's, Newfoundland, there are large

pper

rant

Yarrifer-

ctou, elded

with h for ns of opper

, and little, f the mples about ulders ein of

traced varied r cent. ipping obably ysboro

The ysboro f their

gonish
s with
but as
med of

on the . The ., Gillis .

· deposits of red hematite, most conveniently situated for working and shipplng. Owing to a short sea transport, it is claimed that this ore can be laid down at Sydney for less than \$1.25 a ton. This deposit is divided between the Nova Scotia Steel Company and the Dominion Iron and Steel Company, and it is found to be cheaper than the local ores, so far acquired by these companies. It will, however, be pointed out that there are localities in the province which will probably on further examination prove capable of competing with the Newfoundland ore.

: In the island of Cape Breton Iron ores are known to occur at a number of places, but hitherto they have received little attention. At Big Pond, Ben Eoin, Red Island, Loran, Lake Ainslie, Mabou, Cheticamp, Craignish, Loch Lomond and Grand Anse, red hematites are reported. In some cases these deposits are at the junction of Lower Carboniferous strata with older measures. In some cases openings have been made and samples of good

-quality extracted.

At George's River and Barrasois, on the Little Bras d'Or, in rocks of Lower Silurian and Cambrian age, are red hematites and magnetites. These ores run diagonally across to Escasoni, on West Bay. At Gillis Lake a red hematite from 6 to 12 feet wide was traced by Mr. Moseley for several miles. The magnetites and hematites at Barrasois are reported to be in large quantity, but have as yet not been thoroughly tested as to their quality. Red hematite also occurs at Smith's Brook, East Bay, on a bed said to be 15 feet wide.

On Boularderie Island, at Long Point, spathic ore occurs in a bed several feet wide. In the district lying east of St. Peter's there are several deposits of specular ore up to five feet in thickness, reported to be of fair quality. In the Mira district red hematites from 2 to 6 feet in thickness, have been

opened at a number of points, and are of good quality.

The locality which has so far shown the best promise is that extending westwardly from Whycocomagh. Here a number of outcrops of beds from 3 to 12 feet in thickness have been opened. The ore, which is red hematite and magnetite, is of good quality, but the beds have been traced for a limited distance only. As traces of ore are abundant here over a territory 12 by 3 miles, there is every reason to believe that very large deposits are present. This locality permits of cheap mining, is on deep water, and from 3 to 5 miles from the Intercolonial Railway.

The following analyses will serve to show the quality of some of these

. ores :--

Comquired calities apuble umber Pond, iignish, o cases older f good

shiplaid tween

These se a red several o be in quality d to be

several deposits quality.

ctending eds from a is red in traced r a terrideposits ater, and

of these

	Loch Lonowl.		East Bay. Whycocomagh	Loren.	Big Pond.	George's River.
Metallic Iron	64.49	57.92	60.90	63.09	61.39	62.50
Silica	7.76	12.80	10.80	5.45	9.04	 8
Phosphorus	.03	į	trace.	:	trace.	80.
Sulphur	70.	trace.	trace.	01.	trace.	trace.
Phosphoric Acid	:	91.	:	11.	:	:
Alumina	:	1.55	1.40	:	:	•
Magnesia	:	09.	1.64			
Lime	:	1.30	1.85	05.4 —		æ.
Water	:	1.30	:	:	1.53	19.
Manganese	28.6	.26	:	:	:	1.10

Passing to Nova Scotla proper, the first deposits met are those of Guysboro County. A large tract of Devonian strata between Milford Haven and Salmon River contains specular ore at numerous points. At Erinville two large bodies of ore were tested some years sgo. They were 12 and 65 feet wide. No doubt other large deposits will be found by systematic prospecting. These ores occur on tide water, and at distances therefrom up to 12 miles. These ores are fairly compact, mining from 55 to 62 per cent. metallic iron, and are low in sulphur and phosphorus. The Nova Scotla Steel, and the Dominion Irou and Steel Companies have acquired large tracts in this district. Similar ores, of good quality, occur in Boyleston and Manchester townships, but the velns hitherto opened are small.

At Arisaig and Doctor's Brook, in Antigonish County, there are numerous bedded deposits of red hematites in Silurian strata. These beds, which are receiving some attention, are from 3 to 20 feet in thickness and are reported in many cases to run 60 per cent. in iron, and to be low in phosphorus. There is no doubt that there are large quantities of ore in this district, and it is anticipated that exploratory work will prove the deposito to be uniform in quality and unbroken in extent. They are close to water and about 8 miles from the Intercolonial railway.

Signs of the passage of these ores are found until at a distance of 25 miles a point is reached on the East River of Pictou where explorations have shown the presence of large deposits. Here, in a district 8 by 12 miles, 4 beds from 5 to 30 feet in thickness have been proved to underlle large tracts of country. At the junction of these measures with Lower Carboniferous strata, from Springville to Sunnybrae, there are numerous deposits of brown hematite. They have been worked for a number of years by the Nova Scotia Steel Company for mixture with their Newfoundland and Torbrook red hematites. The production last year for this purpose was about 16,000 tons. This company has an 80 ton furnace and washing and retort or an plant on the East River, near Hopewell.

On the West side of the East River (East Branch) there are extensive signs of the presence of brown hematite deposits, and a vein of specusion or has been traced several miles. In places it yields 15 feet of ore. Still further west on a tributary of the West Branch the presence of brown hematite ore is marked by numerous deposits of drift ore. Spathic ore occurs on McLellan's Brook and Sutherland's River, and at the latter place some exploratory work showed a bed 6 to 10 fect thick. Still further east on French River there are a number of beds of a nodular red hematite varying in thickness up to 5 feet. This district contains very large amounts of ore, a prably situated for working, and near the Pictou coal field with which the is connected by a branch railway running from Sunny Brae to the Intercolonial Railway at Ferrona.

The following analyses will selve to show the quality of some of these ores:

Manganese oxide	ilar. Brown hematite.	Red hemalite.
Alumina	01	
Line each water	21	
Lime carbonate 1.		3.03
Magnesia "	43	
Phosphoric acid	08 .15	.20
Sulphur	16 trace	
Silica 3.0	68 4.26	23.82
Metallic iron 64.4	11 56,83	45.47
BPATINC (one.	
Sesquioxide of iron	20.52	
Carbonate of iron	57 10	82.11
" manganese		4.70
" line	4.02	2.37
" magnesia	5.66	9.06
Silica	2.38	
Moietune	2.00	1.69
Moisture	1.43	
Sulphur	· · · · · · · · · none.	.10
Phosphorus	none.	none.
Metallic iron	42.07	39.64

Small beds of clay ironstone occur in the Pictou coal field, and in the district north of New Glasgow.

Deposits of iron ore are known on the upper waters of the Stewiacke River, at Newton Mills, Grand Lake, Goshen, Clifton, Selma, Brookfield, Mount Tom, Kemptville and North River. Those at Brookfield and Selma have been worked to a small extent.

In the Londonderry district, in Colchester county, there is a band of slates carrying bodies of brown hematite included in a stratum vein of carbonates of iron, lime and magnesia. These were worked for a number of years to supply a charcoal furnace, and two coke furnaces. During the last depression in the iron trade the furnaces were closed, and have since remained idle. The deposits extend for a number of miles, and are available for large supplies of a high grade brown hematite. Near Pugwash, in Cumberland County, similar ores are said to occur in quantity. The traprocks, of Triassic age, extending along the south side of the Bay of Fundy, contain numerous veins of magnetite and red hematite, not, however, yet observed of workable size.

The Devonian or Upper Silurian measures of the Annapolis Valley extend in a narrow band from Digby to a point a few miles west of Windsor, with several interruptions due to granitic masses. At Clementsport two beds of ore, from 2 to 6 feet thick, have been noticed. In the Nictaux and Torbrook

,

e. Still f brown athic ore er place further

Guys-

en and le two 55 feet rospecto 12 r cent. Scotla d large on and

merous ich are eported phorus. ct, and iniform

of 25 orations

by 12

lie large

Carboni-

osits of

he Nova

orbruok

16,000

rt o on

xto...ve

u. ore

er place further

ins very
e Pictou
ng from

districts a large number of beds of red hematite and magnetite have been observed over a district 10 by 4 miles, and are from 2 to 20 feet in thickness. As yet enough work has not been done to permit of their correllation, but the general structure of the district appears to be that of two synclinsl folds repeating the beds. These deposits have been traced in a continuous line from Black River to Nictaux, a distance of six miles, and have been proved by shafts and boreholes that in their quality and size to a depth of 400 feet. Their content was from 0 to 62 per cent of iron, are low in sulphur, and vary in phospherical from 0 to 130. There are in this district enormous amounts of or a with extracted, intersected by railways, and about 32 miles from shipping a Annapolis via the Dominion Atlantic railway. The Torbrook Iron Company worked for several years, with an annual output of 30,000 tons content to the earlier bed 6 to 10 feet thick, until the Londonderry furnaces were seed, and the North Scotia Steel Company drew its supplies from Newformaland.

From these notes it may be interest to the street of Nova Scotia are very considerable, and company favorably with any deposite, except a few of international calabrity. The Corbrook, Nictaux, London-derry, Pictou, Antigonish, Guysboro, Barrasois and Whyhogomah districts undoubtedly contain large amounts of ore, and the indications in other districts warrant attention.

There can be no doubt that as the expense attending the working of the deposits at present supplying high grade ores to the manufacturing districts of Europe increases, the ores of this province will receive greater attention. It may be said that at this date they could, in conjunction with our excellent fuels, maintain several establishments capable of producing pig iron at rates cheaper than those now obtaining in Europe.

e been thickllation, nclinal tinuous

o been

a depth

low ln

his disays, and

tic railwith an ck, until ompany

of Nova

deposite,

Londondistricts

In other

ig of the

districts

ttention.

ur excel-

ig iron at

MISCELLANEOUS MINERALS.

Gypsum.—This nilneral may be considered specially characteristic of Nova Scotia, as deposits of equal extent and variety are met at no other point known to the writer. It occurs as soft or hydrated gypsum, having the following typical composition:—

Lline												48 R1
											•	100,00

and as hard or anhydric gypsum, containing 41 per cent. of lime and 55 per cent. of acid. It is usually massive or granular, but occurs as selenite, in tabular crystals, and in segregated crystals, and as fibrous gypsum.

The exposures show beds from a few inches up to 100 feet in thickness, and it is often found in grains, veins, and crystals in the associated shales, marls, and limestones. In this province it occurs in the lower carboniferous limestone formation. As these measures occur in Halifax, Hants, Kings, Cumberland, Colchester, Pictou, and Antigonish counties, and in all the counties of Cape Breton, it would be impossible to give all the localities. The amount available in the province is inexhaustible. As might be expected it is not everywhere of the best grade, but every district contains deposits available for agricultural purposes, and in many localities there are abundant deposits of the best quality. The chief localities where it is worked are Windsor, Cheverie, Maitland, Walton, Hantsport, Wallace, Antigonish, Mabou, Judique, Port Hood, Port Hawkesbury, Lennox Passage, St. Ann's, and Big Harbor, near Baddeck.

Among the minerals found in the gypsum may be mentioned: glauber and common salt, calcapar, magnesite, arragonite, carbonate and oxide of iron, silica, and free sulphur. In the Windsor district, considerable amounts of boro silicate lime compounds have been noticed. Among them is ulexite, a very pure form of the Peruvian "tiza." These minerals can be utilized in glazing operations, and would be of value if found in quantity. I give an analysis of natroboro calcite (ulexite), as found at Windsor:—

Water													34.29
Lime													14.20
Boracic acid	•	•	•	•	•	٠	•	•	٠				44.10
Soda													14.20

The hard gypsum frequently occurs in lenticular masses imbedded in the soft gypsum, and as beds underlying, and alternating with it. The attention of the quarrymen however is directed to the softer variety. Gypsum has been largely exported to the United States from the Windsor district, and smaller shipments have been made from Cape Breten It is shipped as rock, and not prepared in any way, owing to the duty being virtually prohibitive on any manufactured form. It was at one time used as a top dressing for cotton crops, but now finds its principal use, in the States, for architectural purposes, and in the manufacture of fertilizers.

The transparent c.ystalline selenite commands a higher price than the ordinary gypsum, and is utilized as a filler for paper. Small lots are occasionally shipped, but as yet it has not been observed in amounts permitting of regular production.

SHIPMENTS OF GYPSUM FROM NOVA SCOTIA FROM 1890 TO 1901.

	Tons.
Year. 1890	146,003
1890	161.934
1891	162,285
1891	98.247*
1893	106,171
1894	133,300
1895 1896	130,489
1896	125,000
1896	131,000
1898	140,000
189	122.281
1900	,_

^{*}Nine months only.

The gypsum marketed in this province is in the manufactured form of "Land Plaster" and Plaster of Paris. The crude gypsum is exported to the United States, being quarried and delivered on board ship for about \$1.00 or a little less per ton.

Inferior grades of gypsum are ground and sold as a fertilizer, being called "Land Plaster." It is used in the province as a fertilizer, either alone of in combination with other ingredients. The superabundance of commercia fertilizers now in the market, however, cannot but limit the sale of any on kind. The ease with which crude gypsum can be obtained in some of ou agricultural districts also perhaps makes the necessity of a manufacture article less felt. This ground gypsum, or "Land Plaster," is also being used as an absorbent in stables and poultry houses, and is beginning be employed for the same purpose in slaughter houses. The Winds Plaster Co's. "Land Plaster" sells at present for about 90 cents per bl in carload lots at Halifax.

Plaster of Paris, or Calcined Plaster, is also extensively manufactured at n the Windsor by the Windsor Plaster Co., and is sold throughout the province tion of to a large extent, bringing at present about \$1.25 per bbl. in carload lots at s been Halifax. It is largely used in giving a "hard finish" to walls and ceilings, maller making cornices, etc. By more careful manufacture and finer bolting, Plaster k, and of Paris of the very finest grade could be manufactured in the province, on any whereas at present the most superior grade is usually brought from the cotton United States, where it has been largely manufactured from our own ectural gypsum. In the local market the Windsor Plaster of Paris enters largely into competition with that of Hillsboro, N. B. nan the

What is probably a coarser grade of Plaster of Paris, with possibly other ingredients, is gaining a market here under the trade name of "English Selenite Cement," and is used in plastering walls as "hard finish," the advantage over ordinary plaster being that it sets very quickly and is very hard. Its wholesale price at present at Halliax is about \$1.30 per bbl.

Manganese.—The exceptional purity of some of the manganese ores found in Nova Scotia makes them interesting to the mineralogist, and valuable in certain operations of the manufacturer. The attention paid to these ores is by no means proportionate to their value, and to the great extent of the geological formation to which they appear to be chiefly confined.

The least valuable, but the most common of these ores is wad or bog manganese. It is found as a superficial deposit in connection with every geological formation known in the province. Among the localities yielding it may be mentioned Jeddore, Ship Harbor, St. Msrgaret'a Bay, Shelburne, La Have, Chester, Parrsboro, Springhill, and several localities in the counties of Pictou and Antigonish. On Boularderie Island, Cape Breton County, it occurs as a bed several feet thick, containing from 25 to 44 per cent. of manganese per oxide, from 12 to 35 per cent. of sesquioxide of iron, and from 10 to 33 per cent. of insoluble matter. At Londonderry, in the iron mines, in precarboniferous strata, the ores have, in places, been enriched by manganese oxide to the extent of 14 per cent.

Pyrolusite occurs near Mount Uniacke in pockets and veins penetrating granite, and in the quartzites of the auriferous rocks. It also occurs as veinlets in granite at Musquodoboit and Ship Harbor. In the slates of Devonian age, south of Wolfville, it is found in small masses and stringers.

Pyrolusite and its associated ores occur however most conspicuously in the lower horizons of the Carboniferous and in some of the upper divisions of the Devonian. In the northern part of Hants County a manganiferous limestone belt appears to pass from the mouth of the Shubenacadie River to Tenny Cape, and to Walton and Cheverie. This appears to underlie the gypsiferous strata. The measures carrying manganese ores reappear again south of Windsor, and at Douglas, fifteen miles south of Tenny Cape, and

ots are

ats per-

01.

d form of ted to the pout \$1.00

eing called or alone or commercial of any one one of our unufactured also being eginning to e Windsornts per bbl.

near the line of junction with the gold bearing rocks. In this range of measures the manganese of Tenny Cape appears to be principally connected with a red and gray limestone, dolomitic in composition. At the western end of the district it occurs as veins in conglomerates and sandstones, and also in limestones in places magnesian.

The Tenny Cape ore occurs in irregular nests and in seams erroded on the bedding planes and cross fractures. The ore is chiefly a fibrous pyrolusite, with splendent lustre, based on a compact or granular ore consisting of pyrolusite, of psilomelane, and of manganite, the latter not present in large quantity. The quality of these ores is very high, judging from the fact that many tons have been shipped yielding from 88 to 95 per cent. of available oxide.

The following analyses show the general character of these ores:-

	Douglas.	Cheverie.
Moisture	- 000	2.05
Water of composition	. 3.630	
Iron peroxide		2.55
Oxygen		
Baryta		1.12
Insoluble		2.80
Phosphoric acid		1.020
Manganese oxides	. 84.620	
Peroxide of manganese		90.15
Lime		trace.
	100.000	99.699

At Walton and Cheverie manganite is more common; its composition may be shown by the following analyses:—

	Tenny Cape.	Cheverie.
Manganese oxides	85 54	86.81
Iron peroxide	1.18	2.05
Baryta Insoluble matter	1.27	1.14
Phosphoric acid		10.00
Available oxygen	W 4 F 4	17.73

Calcite, red and brown hematite, barite, and selenite occur with the ores of this district.

Minudie, in Cumberland County, has yielded small quantities of soft fine-grained pyrolusite giving 97.04 per cent. of binoxide of manganese. Similar ores are found at Onslow, and on the Salmon River, near Truro. Boulders containing a mixture of psilomelane and manganite are found associated with brown hematites at the Bridgeville iron mines, and in places the ore is heavily charged with manganese.

In Antigonish County similar manganiferous iron ores are found in the drift at several places.

In Cape Breton County, at Salmon River, several shipments of manganese ore have been made from the Moseley mines. The deposits occur as beds and irregular layers and nodules in soft arenaceous shale, and associated with a dark manganiferous limestone, both of lower carboniferous age. The ore is generally a soft fine-grained or subcrystalline pyrolusite, sometimes containing manganite. The ores run as high as 88.9 per cent. of binoxide, and are very low in iron.

During the past two years a few shipments of ore have been made from the college grant, New Ross, Lunenburg county. The ores extend over a large tract of country and are said to occur as veins sometimes three feet in thickness. The distance of the district from shipping has proved a drawback to mining. The ore is apparently a mixture of psilomelane and manganite, and is said to carry up to 50 per cent. of manganese with traces of phosphorus, and to be suitable for ferromanganese.

The shipments of these eres have never risen to a large figure, and for many years have ranged from 50 to 500 tens, principally from Tenny Cape, with small lots from Walton, Cheverie, Truro, New Ross, and Cape Breton County.

There are no doubt localities in which these widely spread ores will be found in considerable amounts, but as yet the indications have received little attention. The demand for the rich Tenny Cape ores seems to have fallen off, presumably owing to the introduction of methods permitting the utilization of cheaper sources of the oxide. There is, however, a large demand for manganese ore and manganiferous iron ore, suitable for steel-making processes, and a search would probably, owing to the presence of the purer deposits, reveal the existence of the less highly manganiferous bodies adapted for the use of the iron and steel maker.

Antimony.—The sulphide of this ore occurs at Rawdon, in Hants county. A mine was worked here for some years and about 3,000 tons shipped. Owing to the distance the ore had to be hauled to reach the railway, it was found profitable to ship only the richest, and a fall in prices caused the closing of the mine. A few years ago properties were acquired here by Mr. A. McNcil and others, and a number of veins were found in the vicinity of the one originally worked. The ore carries considerable quantities of gold, and the opening of the Truro-Windsor railway will assist in the development of this district, which is undoubtedly of importance.

The following assays were made for the Department by Mason and Asquith of samples from the veins recently opened:—

osition

ge of

ected

stern

, and

n the

usite,

ng of

large

fact avail-

he ores

of soft iganese. Truro. e found in places

	No. 1.	No. 2.
Antimony* *Gold	45.75% 2.48 oz. per ton.	18.21 .23 oz. per ton-
*Silver	10 " "	.13 " "

*Ton of 2,000 lbs.
N. B.—No. I samples were taken from the pay streak, and were a fair average.
No. 2 were from the large bodies of low grade ore.

Celestine.—This mineral, the Sulphate of Strontium, is reported by Mr. H. Fletcher as occurring on the right bank of the Sydney River, about a mile and a half above the bridge, in a bed about a foot in thickness.

Sulphur.—Common and magnetic iron pyrites are found in many places in the province. It is reported as occurring in Queeus and Shelburne-Counties in masses and aggregations of crystals. A considerable body of rather low grade pyrites is known at the head water of the George's River, Cape Breton County. It is found near Salt Springs, Pictou County, cementing a coarse sandstone, but has not yet been found in the province in quantity permitting of its utilisation for acid making. It is reported as occurring in gypsum, near Brookfield.

Molybdenum.—The sulphide of this metal is found at several points, as at Gabarus and Louisburg in Cape Breton, where it occurs in small quantities over a large district. Hammonds Plains, Bedford, Lower Musquodoboit, New Germany, and New Ross have yielded specimens, those from the latter locality being frequently in large and handsome crystals.

Cobalt and Nickel.—These metals are not uncommon in the province, but hitherto have been observed only in traces. It occurs in mispickel at Montagu and Cochran's Hill, in copper pyrites at Caribou, Pictou Co., and at Londonderry.

It has been reported in magnesia alum from Newport, and in magnetic iron pyrites from Nictaux and Geyser's Hill. Cobalt is noticed also in bog manganese.

Zinc.—The sulphite, zinc blende, is frequently found in auriferous quartz veins. It also occurs with galena at Cheticamp, Inverness Co., where development work is being done on deposits of considerable size, which carry valuable amounts of silver and lead. Samples of similar ores are reported from the district extending along that shore for fifteen miles north of Cheticamp.

Tin.—Tinstone is reported as occurring at Tangier, Shelburne and Country Harbor, associated with granitic debris.

Arsenic.—Arsenical iron pyrites is abundant in the auriferous quartz veins.

Phosphates.—Mr. Fletcher (Geological Survey Report, 1876-77, p. 453,). mentions the occurrence in the Lower Silurian limestones of MacIntosh Brook, Cape Breton, of layers of phosphatic nodules. Similar deposits have been observed at Arisaig, and in Colchester county samples are said to have—yielded as high as 40% of phosphate of lime. The blue phosphate has been observed near the town of Antigonish. It is doubtful if any of these deposits are of economic value. Discoveries of apatite similar to that extensively worked some years ago in the province of Quebec, may be anticipated in the Precambrian limestones, etc., of several parts of Cape Breton.

Salt.—This mineral is found in crystals in many of the gypsum quarries, and occurs abundantly as brine at numerous points in the lower carboniferous measures. The apparent similarity of the conditions attending the accumution of the gypseous series in parts of Nova Scotia and in Europe leads to ahope that heds of salt may be found here by means of borcholes. This subject is one of considerable importance, but has not yet received serious attention in Nova Scotia.

In this province salt has been made at various times from brine springs, and in some cases a superior article was produced. At present there is no attempt being made to utilize them. Among the various springs may be mentioned those of Renfrew, Walton, Salt Springs, Irish Mountain, Sutherland's River, Antigonish, Ohio, Baddeck, Whylogomah, River Philip and Black Brook.

The following analyses will show the character of some of these-brines:-

	Walton.	West River.	Little Narrows.
Lime carbonatc	14.73	3,775	
Magnesia "	trace.	2.932	• • • •
Iron "			
Phosphoric acid			
Magnesium chloride	4.48	27.330	.1593
Lime sulphate		154.730	5.6810
Common salt		4133.500	50.6881
Silica		.560	trace.
Calcium chloride		51.910	.1942
Total	967.48	4374.737	56.7226
Salt per cent	1.1	5.9	

The analyses of Walton and West River brines give the contents ingrains to the imperial gallon; that of Little Narrows gives the solid contents in 1000.00 parts.

Building Stones.—The stone used for structural purposes in this province is either sandstone or granite. There are several varieties of the former, passing from a very light fawn to a purplish red. The granite used is of a.

many burnedy of River, ounty, ovinceted as

ton-

erage.

Mr.

out a.

nts, asquantidoboit,...

ckel at

agnetic in hog

s quartz,
where
ch carry
reported
of Cheti-

Country

s quartz

standard gray colour. As the sandstones and freestones fit for the quarryman's work occur chiefly in the Upper Coal measures and the Permocarboniferous, we find the principal quarries extending along the straits of Northumberland from Merigomish, in Pictou County, to the Joggins, in Cumberland County. The quarries are in most cases conveniently situated for either railway or water transport, and for effective quarry work.

Samples of these stones were forwarded to the Paris and Glasgow exhibitions, and the following notes from a paper on the Nova Scotia minerals for the Paris exhibitions, read before the Nova Scotia Institute,

session 1899-1900, will serve to show their position and extent.

QUAPRIES.

These notes refer to the quarries in the northern part of Cumberland County, which furnished samples of their products.

Quarries at River John.-No. 1. This is the only quarry at present working in this district. It is situated at River John and about a quarter of a mile from I. C. Railway, and is connected therewith by a good road.

It contains a reddish sandstone of fine grain, and has been opened for about 350 feet in length, exposing a face so far of about 14 feet. Stones are cut here up to about 33 cubic feet, though almost any size could be obtained with larger machinery. The seams are very regular in formation and lie nearly horizontal. Worked for nearly a year.

No. 2 adjoins the first quarry, and resembles it in general character-

istics, though the stone is of a lighter colour.

No. 3, shout a quarter of a mile up the river from No. 1, was worked for six years intermittently. Many grindstones were cut in this quarry,

which yields a firm grey sandstone.

Wallace Harbor.-The Wallace Gray Stone Co., Wallace Harbor, John Stevenson, manager. This quarry is situated at Wallace, and a great part of the stone is shipped by water, though it is connected with the I. C. Rail-

way hy a good waggon road about two miles long.

Though the stone is carried to the wharf by horses, a tramway (gravity) could easily be operated, the quarry being situated on a hill. The distance is about a quarter of a mile. This quarry has been worked for a period of nearly thirty years off and on, and is still only partially developed. It produces an average of about 1,500 tons ("quarry") a year, of fine grained sandstone in two colours-" olive" and "bluish."

Blocks up to ten tons in weight and measuring fourteen feet are cut, and the greater part of the stone is shipped to the Boston and New York

markets. The poorer stone is sold locally.

Wallace Harbor.-The G. P. Sherwood Co., T. C. Dobson, manager. This quarry adjoins the quarry of the Wallace Gray Stone Co., and the same remarks apply to it.

At Wallace Bridge the famous Battye Quarry is being operated by George Battye. Stone has been taken from this quarry since the year 1809, and there is still much in sight. It is situated on the I. C. Railway and the Wallace River. Chief market New York and Eastern States cities. Blocks up to ten tons are cut. At present 25 feet of rock is shown in the face, with seams measuring from 2 to 6 feet in thickness. This is composed of a very uniform and beautiful sandstone, suitable for monuments as well as construction work.

On the River Philip, about five miles from Pugwash, is situated the quarry of McLeod & Embree. It produces a handsome red sandstone contained in scams from 2 to 7 feet, and shows altogether 20 feet in the face. Blocks cut to 8 tons. Has been operated for upwards of 30 years, and usually ships to the States. This year all the stone quarried is being supplied to Toronto.

The Atlantic Stone Co., Limited, R. S. Hibbard, Manager.—The quarry of this company is situated on Cumberland Basin, 3½ miles from Joggins Station, on the Canada Coal Company's Railway, and 16 miles from I. C. Railway. The stone is shipped chiefly by water, in vessels up to about 300 tons. The market is mainly in the New England States, though the stones are occasionally sent much further west. 2,000 tons shipped per year. This quarry produces a very superior form of grindstone. Stones from half an inch to 14 inches thick, and up to 7 feet in diameter are cut, though almost any size that could be handled is procurable.

At Lime Rock, West River, Pietou Co., are sandstone quarries yielding good building stone. Samples are shown by Mr. J. H. Fraser. In the Merrigomish district the strata lying above the Productive measures yield grindstones and fair qualities of freestone.

Other localities are Pieton, and Glenfallock. On the Basin of Minas, Cornwallis, Johnston Brook, Ilorton, Falmouth, Kenneteook, Nine Milo River, and Old Barns, have yielded freestones in some eases of delicate shades and good texture. In Cape Breton sandstones from the Millstone grit and the Coal measures have been used to a limited extent for building purposes.

Syenites, Porphyries and Granites.—Granite is very abundant among the older rocks of the Province. Among localities which have furnished it for building purposes, may be mentioned Shelburne, Queens and Lunenburg Counties. It also occurs at Aspotogan and various points thence to Halifax. As already mentioned, in describing the Gold fields, it runs continuously from Halifax to Windsor, and thence westward. It occurs again at Waverley, and runs through Musquodoboit, Jeddore, Ship Harbor, Sherbrooke, and Country Harbor to Canso. It occurs inland at the head waters of many of the Eastern rivers, and is estimated to cover a large area of the Atlantic coast

nd

y.

ni-

th.

er-

for

ow

tia

te,

ent rter .

for nes

ter-

ohn

part Railvity)

anee d of pro-

and York

iined

ager.

district. It has however, been quarried only at points accessible to shipping At Halifax, it has been used a good deal about the fortifications, and a number of houses have been constructed of it, its cost, rough, being from \$2.25 to \$4.00 a ton.

In the Cobequids, there are masses of ficeli and red colored syenite, which have afforded very handsome polished samples; but as yet have not been worked for construction. Porphyries and syenites occur in various parts of Cape Breton, but their economic value has not been tested. The following localities may be mentioned,—St. Ann's, Boisdale and Coxheath. The crystalline diorites of Louisburg were used by the French in huilding their fortifications.

Limestones.—This material has not been used to any extent in Nova Scotia, for building purposes, although it is frequently found to stand exposure well, and to be readily quarried. Among localities yielding it, may be mentioned the Shubenacadie River, Kennetcook, Lower Horton, Thompson Station; Glengarry and Springville, Pictou County. Stones from a quarry here, retain, after an exposure of sixty years, every trace of the chisel or pick. The marbles will be noticed further on.

As before mentioned the anhydric or hard gypsum has been used to a small extent for walls, foundations, etc. A flaggy arenaceous schist, known as iron stone, was extensively used some years ago, for warehouses and walls in Halifax. Many of the metamorphic sandstones of the Atlantic coast would furnish a most pleasing and durable building material.

At present the supply of wood for building purposes is so plentiful, that brick or atone houses are the exception. Even public buildings, churches, halls, etc, are almost always of wood; but as this material becomes more expensive, the labor of the quarrymen will succeed that of the lumbermen, and our towns become something better than wooded shells blackened by smoke.

Flags and States.—A small amount of flagstone has been quarried on the North-West Arm of Halifax Harbor, and at Beaver Bank. Slates were quarried to a small extent at Rawdon, and various places in Hants County, and the quality and quantity are equal to any demand. Dalhousie Mountain and West River, Pictou, are said to have good slate beds, and it is also reported from the South Mountain, in Digby and Yarmouth Counties.

At the Provincial Exhibition of 1879, slates were shown from Sackville, River John and Upper Stewiacke, which, although in the rough, were of good material.

The demand for roofing states will become general in the province in a few years, as their superior y over the shingles in ordinary use becomes apparent.

Brick Clay.—These clays are found in many places, presenting an unlimited supply of raw material for the brick maker. As yet, brick is but little used in the province, although when well made and used according to proper designs, it makes the best building material for this climate. The best known clay fields are those of Shubenacadie, along the line of the Intercolonial Railway, and many points in the carboniferous districts of Colchester, Pictou, Cumberland and Antigonish.

Limestone.—In Nova Scotia the limestones are confined practically to the Lower Carboniferous, and are generally associated with the gypsums. There are also beds of this material, sometimes metamorphosed into marble, in the Laurentian, etc., of Cape Breton, and in the Cambrian and Silurian measures, but they do not usually form deposits of economic value in the latter measures. The carboniferous limestones are strongly developed in Cumberland, Colchester, Hants, Kings, Pictou and Antigonish Counties, and at many points in Cape Breton. They occur in beds varying in thickness from a few inches to 50 feet, and in some localities their aggregate dimensions will exceed 400 feet. Their quality varies from calcareous sandstones and clays to the crystalline pure mineral.

At Windsor, Brookfield and many other localities beds are found composed entirely of fossils characterising the Marine Limestone formation, and give the following component parts on analysis by Dr. How:—

Carbonate of lime	97.64
" niagnesia	1.10
Oxide of iron	.07
Phosphoric acid	tracc.
Insoluble residue	.68

A limestone similar to the above was extensively quarried at Brookfield as a flux for the Londonderry iron ores.

The limestones of Pietou County are also well adapted for fluxes. The following analyses of a limestone from Lime Brook, Springville, were made for the Haifax Company at the Durham College of Physical Science —

	1.	11.
Lime earbonate	93.90	96.26
Magnesia "	2.45	2,33
Iron peroxide	.59	.57
Manganese peroxide		.55
Alumina	.12	.10
Sulphur	.08	.02
Phosphoric acid	.03	.03
Silica	2.10	1.99
Moisture	.18	.17

These results are confirmed by an extensive series of analyses made some years ago by the writer, embracing all the more important exposures of that

and g it, ton, ones

e of

ng

m-.25

ite.

not

Otta

Che

th.

ing

ova

to P own valls

coast

that ches, more men,

d by

were unty, foun-

ville, ere of

in a

mineral in the vicinity of the Pictou coal and iron deposits, but the space at my disposal would forbid its insertion. The position of the East River limestones forms an important item in their adeptability for fluxing purposes. They occur as a band everywhere between the coal and iron, so that their transport becomes a metter of comparatively low cost, and large quentities are available by simple quarry work. A quarry at Black Rock, above Bridgeville on the East Branch, has furnished flux for many years to the Ferrona furnace.

Some limestones at Horton, Onslow, the Joggins, Pugwash and other places are hituminous, and contain notable quantities of phosphoric ecid. As already mentioned, some of the East River limestones contain important percentages of carbonate of iron, and may, at some localities, afford an ore of

this metal, in addition to the spathic ores referred to before.

In Cape Breton, several places are known which afford marble believed to be well adapted for building and decorative purposes. The finest deposit of workable limestone yet discovered is on West Bay, Bras d'Or Lake. In variety of color and tint this rock resembles the limestones of the George River series, of which it forms a part, but it contains little or no admixture of foreign materials, and is uniform to texture and in unequalled abundance. The following varieties have been recognized:—

- 1. Fine white statuary marble.
- 2. Fine white building "
- 3. Coarse white building "
- 4. Blue and white clouded marble.
- 5. Brocatello marble, mixed with six varieties of colored marbles.
- 6. Fine flesh-colored marbles, often striped and variegated.

The locality offers every facility for quarrying and shipping, and blocks of any required dimensions can be shipped cheaply to the United States, equal in quality to those already admired as samples. These marbles are in many cases magnesian. At present large quantities of lime are burned here, and the Dominion Steel Company have extensive openings furnishing the flux for their furnaces at Sydney. The same company has also drawn largely upon the marble of George's River, in Cape Breton County, as a flux and for use in the concrete foundations of its furnaces, ovens, etc.

At St. Ann's Mountain, Cape Dauphin, Salmon Creek, Whyhogomah, River Dennis, George's River, French Valley and Escasoni, marbles are also found. At the latter place they are usually too much broken and mixed

with other rock to be available for artistic purposes.

A magnesian limestone on the property of the Messrs. Burchell, at Kelly's Cove, Victoria County, has been extensively quarried by the Nova Scotia Steel Co. for linings for convertors, etc.

At Five Islands, Colchester County, promising marble deposits are known.

The carboniferous and other limestones are quartied at all points for lime for building and agricultural purposes. That from East Bay has been extensively burned for lime. The Nova Scotia lime is frequently brown, arising probably from carelessness in selecting and burning; its price per barrel may be averaged at 95 cents. At numerous points in the province the limestones contain foreign ingredients, indicating the presence of a certain amount of hydraulicity, such as alumina, carbonate of iron, magnesia, silica, etc.

er

T-

at

to

d.

nt

of

ed

sit

In

ge

ire

ce.

cks

tes,

are

ned

ish-

also

ity,

ah,

also

xed

lly's

otia

wn.

c.

The well-known Portland cement was invented in accordance with the desire for a material which would set rapidly, and remain indurated in water. Lime, itself, is useless for this purpose, unless mixed with the proper proportions of an argillaceous compound. Many good natural cements have been used at various times, but the want of uniformity in the composition of the calcares argillaceous rocks they were made from soon rendered their action even dangerous.

In this province, although many limestones possess hydraulic qualities, their uniformity cannot be relied on. Hence, our cements, which are natural ones, have never proved permanently satisfactory. The limestones and clays of the province will answer for the manufacture of cement; but the absence of chalk requires more care in the manipulation of the limestone, and a consequently increased cost.

At present I am not aware of any Nova Scotia cement in the market, the Portland cement being generally called for in contracts and specifications. Among Nova Scotia limestones possessing hydraulicity may be mentioned those from Horton, Windsor, St. Peter's, Chester, Onslow, Shubenacadie, Springhill, Whyhogomah, etc. In the lockwork of St. Peter's Canal, Portland cement alone was used.

The quantity burnt for agricultural purposes must be very considerable; but no returns are made of the amount thus used, and in many parts of the province nature has supplied this fertilizer in abundance. The Mines Department gives the amount of limestones and ankerite quarried during the year 1900 at 50,000 tons, but this refers only to that used for smelting purposes, and does not include the amount extracted by individuals throughout the province.

Paints.—Mineral paints as found in Nova Scotia may be divided into two classes. The first consists essentially of iron ores, composed of hydrated peroxide of iron, with manganese, silica, ulumina, and water of hydration. They are frequently, as far as their composition goes, suitable for smelting, having been originally carbonates of iron, lime, and magnesia, and having had the lime, phosphates, sulphur compounds, etc., leached out. But their friability and excess of moisture would require their manufacture into briquettes before being smelted. Good examples of these ores are met at

the Londonderry mines, where they appear to have been formed from the ankerite and sideroplesite forming the vein stone.

The following analyses show the original and present composition of these

minerals :-

	OCHRE.		
oxide. 74.52 70 xide. 40 5 xide 6.20 14.80	Red. 0.20 5.50 2.80 .45 4.40 6.80		
	1,00		

The red colours produced by burning the ochres are modified to purples by manganese, and to browns and grays by mixtures of white aluminous clays and other impurities. Similar others occur in Antigonish and Picton Counties, in connection with the specular ore veins, and at Chester and numerous other points. At Chester the others are produced by the oxidation of Lower Carboniferous limestones containing manganese, affording purplish paints which were formerly manufactured to some extent.

The other class of ochres appear to have been formed by water depositing in beds of earth, marl, etc., the oxides of iron, manganese, etc. Such deposits occur at Folly Mountain, Onslow, Kentville, Cornwallis, Polson's Lake, Chezetcook, Lunenburg, and in Cape Breton. Small quantities of these ores have been treated at several points for local use. These deposits pass into bog iron ore or wad manganese, as the conditions of disposition permitted.

These mineral paints occur abundantly in Europe, and their price is too low to permit of the utilization of the local deposits for the limited home market or for export. The price of the continental ochres and umbers depends not only on the original ore, but also on the care with which they are cleaned and manufactured.

Barytes.—This mineral is frequently met in the province, and a small annual production has been maintained for a number of years. It is confined to no particular geological horizon, but is met in the Carboniferous and all the older formations. It is principally used in the manufacture of paints and enamels. Its value depends largely on its freedom from iron, copper and other ores. It is found in irregular pockets in Lower Carboniferous at Five Islands, Colchester County, usually in characteristic tabular crystals, with calcapar and copper pyrites. It has been mined at River John, and at Brookfield and Stewiacke. It occurs with limonite at the mouth of the Shubenacadie River, at Clifton. Among other localities may be mentioned the Lower Carboniferous of the Avon, Musquodoboit and

Wallace Rivers, and at Greenville, Cumberland County. It is frequently found with the manganese ores of Hants and Colchester counties. In Cape Breton it was mined for several years at Lake Ainslie, and at present an annual output of 783 tons is secured from Cape Rouge, Inverness County. Other localities reported in Cape Breton are Whyhogomah, Cheticamp, Loch Lomond, St. Annes and East Bay.

Fire Clays.—Clays are essentially hydrated silicates of alumina, also holding moisture. Their plasticity when mixed with water depends on the presence of the water of composition, and when heated to redress, it is driven off, and they lose this property. When these clays stand a high temperature without meiting or softening, they are termed refractory or fire-clays. They occur in various geological formations, those of the best quality being generally obtained from the coal measures, where they usually form the beds underlying the coal seams.

The mechanical mixtures accompanying clays are silica, carbonates of lime and magnesia, pyrites, etc. They affect the hested clay in various ways, by causing unequal expansion and cracking, and by forming fusible slags. These effects depend not only on the amounts of the impurities, but also on their relative proportions, as in the case of furnace slags. Among the purer clays are those holding large proportions of pure silicious sand, such as the celebrated Dinas brick clay.

Fire bricks are required to stand in some cases, high and prolonged heat, alternations of temperature, not to soften when heated, or run when in the presence of heated metallic oxides. Any one clay will seldom meet all of the above requirements, and various materials are added to counteract the failing, such as sand, powdered millstone grit, clay previously burned and ground, etc.

The method of making fire brick resembles that followed in the manufacture of common brick. The clay, after weathering, is carefully ground, etc., and burned for five or six days in kilns of 15 to 20 thousand.

The following analyses will show the composition of several Nova Scotia fire clays:---

nre ciaya :				
Lingan,	Deep Seam,	Opringhill.	Stewlacks.	Coxheath, C. B.
Silica	55.15	50.47	45.611	76.260
Alumina 32.10	22.57	32.69	14.000	
Iron oxide 1,87	6.48	8.01		19.152
Titannia anid			ช.100	trace.
Titannic acid	1.75	1.88		
Lime	1.27	1.85	trace.	.555
Lime	1.58	2.11	1.096	
Alkalies	2.66	1.96		.170
Comb'd suntan 8 20				.259
Comb'd water 5.30	4.23	1.03	6.550	4,300
Moisture	2.88		1.000	
Carbonic acid	5.04			• • • •
Organic matter	1.16	• • • •		• • • •
Samil	1.10			
Sand			25 .689	
5				• • • •

home nbers vhich

smali
It is
nifernufacfrom

the

686

ples

nous

cton

and

the

ding

iting

Such

son's

e of

osita

ition

too

Lower eristic sed at ite at

alities oit and

The fire clay from Coxheath Hills, Cape Breton, is an altered felsite, compact, lustre pearly to dull, color pearl grey, greasy to the touch. The analyst, Mr. Hoffman, G. S. R., 1875-76, gives the results of numerous experiments, from which it appeared that the rock alone would not make a brick, as after burning the mass remained friable. When from half to one per cent of lime was added an excellent refractory brick was produced. Large quantities of this rock, which is well worth a practical test, are reported by Mr. Fletcher to be exposed at Watson's Brook and several localities in the Coxheath and East Bay Hills, at Big Pond, Gillis and Forks Lake Brooks.

During 1879 a few tons of fire clay were burnt into good brick by Mr. Cameron, of Stellarton, and used in the construction of coke ovens by the Halifax Company. The clay is from a bed overlying the McGregor seam, and is four feet thick where opened on the land of the Halifax Company.

Doubtless as the search is extended other clays will be found suitable for refractory bricks, linings, etc., and in this connection the large deposits of infusorial or silicious earth found in many of our lakes, etc., may prove of value.

Refractory Stones.—There has been so little demand for such materials that scarcely any information can be given. Some of the metamorphosed Upper Silurian clay slates of Pictou County afford stones which have been successfully used for lining cupolas, boiler furnaces, etc. I have tested the fire resisting qualities of several sandstones from the Pictou coal field, but did not find them valuable. The upper coal measures may yield sandstones adapted for furnace hearths, etc.

Scapetone has been found, I am informed, in Cape Breton, near Louisburg, and at other points; but I am in possession of no details as to its

quality, etc.

At late exhibitions very handsome samples of pottery, vases, jars, ornaments, besides tiles, drain pipes, dishes, etc., have been shown. From these the adaptability of the various white and red clays to the potter's art was satisfactorily shown. Some years ago pottery was made near New Glasgow from a coal measure clay, but work has been discontinued.

Kaolin has been found in small beds at several places on the Atlantic coast, resulting probably from the decomposition of feldspar. As yet, however, it has not received any attention. In Cornwall a large business is carried on in this material, some of which is worked into porcelain, etc., on the spot, and the rest taken to the Midland counties.

Precious Stones .- Our list of precious stones for the purpose of ornament, etc., is a very limited one. A specimen of topaz was shown at the London exhibition of 1872, said to have come from Cape Breton. This may have been a yellow corundum. Garnets are very common in the more highly

metamorphosed slates of the gold districts. They vary in size from a pin's head to crystals one-half an inch in diameter, but, I believe, are generally found to be too britle for the purpose of the jeweller. Crystals of quarts, commonly known as rock crystals, are very common in the older rocks, and are met in great beauty in some of the gold leads. The mineral, however, is so abundant that they would be valuable only if there were a home demand to admit of their being worked here.

Amethyst, purple quartz, is a very common mineral in the trap districts, where it occurs as crystals and incrustations often of considerable size. Its value depends entirely on its depth and uniformity of color. Specimens from Nova Scotia have been much admired in England and on the continent. The early French settlers sent considerable quantities home, and one very handsome specimen was divided in two and placed in the crown of one of the French kings. Among localities yielding it may be mentioned Parraboro, Digby Neck, Cape Sharp, Blomidon, Cornwallis, Sandy Cove, etc. Smoky quartz, cairn gorm stone, is very abundant at Paradise, Annapolis County, and many of our auriferous veins present varieties of it. Heliotrope, Jasper, Chalcedony, Agate and Common Opal are also frequently met in the trap districts. The Heliotrope or blocd-stone finds a ready sale when in good specimens.

The trap of the Bay of Fundy has acquired an almost world-wide reputation for the variety and beauty of its minerals, which have been formed by the dissolving and concentrating agency of water acting through long ages.

Dr. Gesner, the late Dr. How and others, devoted much time to the determination and collection of these minerals, and very beautiful cabinets have been exhibited at various times.

It would be foreign to the aims of this report to describe, even in the briefest terms, the various minerals and their localities, as they possess scarcely any economic value at present. The following list embraces those best known:—Albin, Analcime, Anatase, Apatite, Apophyllite, Calcspar, Centrallasite, Chabazite, Chlorophœite, Chlorite, Cyanolite, Farœlite, Green Earth, Green Calcspar, Gyrolite, Heulandite, Laumonite, Lederite, Mesolite, Mordenite, Moss Agate, Natrolite, Obsidian, Phrenite, Poonah Farth, Scholesite, Sinter, Stilbite, Tremolite, Thomsonite, and Wichtesite. Those who desire further information on the subject of these trap minerals, the identification, physical chalacteristics, etc., of their varieties, will find Dana's Mineralogy the best text book.

The nerous take a to one duced. set, are several

Forks

by Mr.
by the
seam,
cany.
cuitable
leposits
y prove

rphosed re been ted the eld, but adatonea

r Louis-

es, jars,
From
ter's art

Atlantic ret, howsiness is , etc., on

Indonia have highly

TENURE OF MINERAL LANDS.

The grants of land to the early settlers in this province contained no regular reservation of minerals; in some instances gold, silver and precious stones alone were reserved, in other cases, the gold, silver, coal, iron, copper, lead, etc., were retained for a source of revenue to the Crown. After the year 1868 the reservations were gold, silver, lead, copper, tin, coal, iron, and precious stones. In 1827 all ungranted minerals became the property of the Duke of York and were transferred by him to the General Mining Association. After the agreement with the General Mining Association terminating their monopoly, the government passed an Act in 1858, by which they retained in previous grants, the gold, silver, copper, iron, lead, tin, coal and preciou astones whenever mentioned, and made the above reservations in all future grants for the purpose of revenue. This Act releases to the owner of the land, all minerals not mentioned above, such as limestones, gypsum, fire clay, barytes, manganese, etc., and any of the above reservations whenever they are not specified in the grant. There is no complete list published of the various grants, but full information as to the position, date and reservation of every grant, can be obtained from the Crown Land Office.

In 1892 the government reserved all minerals excepting gypsum, lime-

stone and building material.

The Department of Public Works and Mines is entrusted with the management of the Crown minerals. Gold areas are obtainable under either prospecting licenses or leases. The unit area is a piece of ground, with vertical boundaries, 150 by 250 feet. Prospecting licenses are granted in blocks containing not less than six, nor more than 100 areas, and not in length exceeding twice the width. These cost at the rate of 50 cents an area, run for 12 months, and can be converted into leased areas. Areas can be applied for directly under lease, or selected from prospecting licenses, at the rate of \$2.00 per area. The leases run for 40. years, and when not working pay an annual rental of 50 cents an area. Mill licenses are issued, and the royalty on gold is at the rate of 2 per cent. on unsmelted gold, valued at \$18.50 per ounce, Troy, and at \$19.00 an ounce on smelted gold.

For minerals other than gold, licenses to search are issued for 18 months, covering an area 2 by 21 miles. During the term a lesse can be selected covering one square mile for coal or iron, one half square mile for lead, copper, etc. These leases cost \$50.00, and when work is not performed pay an annual rental of \$30.00. The leases are for four renewable terms of 20

years each.

The royalty charged on the coal sold by the Dominion Coal Company is at the rate of 12½ cents per ton of 2,240 lbs., on account of their having a 99 year lease. On the coal sold by the other companies the royalty is ten cents. Coal used for colliery engines and by workmen is not liable to royalty. Proportionate royalties are charged on other minerals.

The Department is constituted a court of record and justice for all mining titles, and is administered under a law framed to facilitate the acquisition and working of mines, with a proper regard to the royalties, which form an

important source of revenue.

A mines regulation Act is enforced for the safe working of the properties. Coal mining officials are required to undergo examination, and all colliers are examined as to their experience and ability before taking charge of working faces. Similar regulations govern the employment of engine-drivers at man-hoisting shafts and slopes.

Diamond and Calyx drills are obtainable from the government for exploring work.

iron, above is Act , such of the here is n as to

ed no

ecious

iron,

rown.

r, tin,

ne the

eneral

A880-

1858,

, lime-

om the

th the under ground, granted not in tents an eas can nees, at hen not e issued, ed gold, months,

selected for lead, med pay us of 20

COAL—SALES.
YEAR ENDED SEPTEMBER 30, 1900.

Names.	lst Quarter.	2nd Quarter.	3rd Quarter.	4th Quarter.	Year 1900.	Year 1899.
Nova Scotia: Land Sales . Sea Borne		98,648 94,815		90,457 145,663	379,205 184,695	336,807 392,670
Total N. S. New Brunewick	92,175	193,463 109,137	109,421	95,786		327,504
Newfoundland . P. E. Island		46	20,756	21.567	68,103	
Quebec			314,399	391,833 138,300		6,04
United States Oth'r Countries		100,010	140,000	1,215		,
Total	743,699	549,353	806,443	898,051	2,997,546	2,419,13

COAL—GENERAL STATEMENT.
YEAR ENDED SEPTEMBER 30, 1900.

Barrier de compression de la la compression de la compression della compression dell	1900.	Production.	Colliery Consumption.	Sales.
lst Q 2nd 3rd 4th	Quarter	756,983 834,945	58,606 45,749	743,699 549,353 806,443 898,051
	Total	3,238,245	209,082	2,997,546

COAL PRODUCE OF NOVA SCOTIA FOR YEAR ENDED SEPTEMBER 30, 1900.

331-6

			COLLIERY CONSUMPTION.	NSUMPTION.
COLLIERY.	Produce.	Sales.	Engines.	Workmen.
Joggins	67,290	55,456	7.131	1.490
ubilee	80.5	58	27	
Scotia	525	417		286
Springhill	458,904	403,170	18,448	7.27
Acadia	294,884	258,378	32,290	5.291
Intercolonial	244,000	226,535	14,158	3.826
Dominion Coal Co	1,930,425	1,808,694	65,476	22.566
ova Scotia Steel Co.	249,910	225,932	16,277	11.756
North Sydney	7,222	7,180	115	345
ew Campbellton.	12,426	9,891	1,743	792
Port Hood	2,323	1,770	443	22
Broad Cove.	195	65		
	3.238.245	2 997 546	158 108	89.074

Nora Scotia Coal Sales, 1785 to 19(4) (inclusive.)

Your.	Sales,	Total.	Year.	Sales.	Total.
1786 1786 1787, 1788 1780 1790	1,008 2,000 10,001	14,249	1841 1842 1843 1844 1845 1846 1847	148,296 139,704 105,151 106,482 180,674 147,504 201,650	Forw'd 1,916,150
1791 1792 1796 1794	3,670 2 143 1,996 4,405		1848 1849 1850	187,643 174,6-2 180,064	1,832,794
1796 1796 1797 1798 1799 1800	5,320 8,249 6,039 ,5,948 8,947 8,401	51,048	1851 1862 1863 1854 1855 1856 1857	188 499 188 076 217,448 284,812 288,815 238,492 294,195	
1801 1802 1803 1804	\$,755 7,769 6,601 \$,976		1856 1850 1860	226,726 270,293 322,693	2,309,319
1805 1806 1907 1806 4809 1810	10,130 4,938 6,119 6,616 8,918 8,908		1861 1862 1863 1864 1866 1866 1867	326.429 306,637 429,351 576,965 636,966 568,520 471,185	
1811 1612 1613 1614	8,816 9,870 9,744 9,864	•	1868 1849 1870	453,634 811,795 868,277	4,927,330
1815 1818 1817 1816 1819 1820	9,336 8,619 9,294 7,920 8,692 0,960	91.527	1871 1872 1873 1874 1875 1876 1877	596,418 785,914 811,106 749,127 706,795 634,307 697,668	
1921 1822 1823	11,816 7,512	VI,821	1878 1579 1880	693,511 688,628 164,659	7 217,430
1824 1825 2936 1827 1826 1829 1830	27,000 12,000 12,149 20,967 21,935 27,209	140,620	1881 1882 1883 1884 1886 1886 1887	1,086,014 1,250,179 1,297,823 1,201,65: 1,254,510 1,373,66: 1,519,684	
1831 1832 1833 1834	87,170 50;869 64,743 50,833		1888 1589 1390	1,576,69/2 1,785,107 1,786,111	18,910,130
1836 1834 1867 1838 1839 1849	86,4 44 107,593 118,942 116,780 145,962 101,1116	839,964	1891 1892 *1893 *1994 1895 1896 1897 1899 1899	1,840,945 1,752,934 1,455,904 2,019,742 1,831,357 2,047,133 2,013,421 2 116,197 2,419,137 2,997,546	90,552,53

SUMMARY,

1786 to 1790	
1901 4- 1900 51 048 1851 to 1960	
1811 to 1820. 91,527 1871 to 1880	
1891 to 1890	į.
1981 to 1840	

^{*}Nine months only. Flecal year begins Oct. 1 and ends Sep. 30. (Cap. 4, Acts 1843.)

COAL.

NOVA SCOTIA EXPORTED TO THE UNITED STATES.

Yeurs.	Tons.	Duty.	Years.	Tons.	Duty
1850	118,173	24 nd.	1876	71,634	75
1851	116,274	"	1877	118,216	66
1852	87,542	66	1878	88,495	44
1853	120,764	66	1879	51,641	46
1854	139,125	Free.	1880	123,423	44
1855	103,222	66	1881	113,728	. 44
1856	126 152	44	1882	99,302	44
1857	123,335	66	1883	102,755	66
1858	186,743	44	1884	64,515	44
1859	122,720	44	1885	34,484	- "
1860	149,289	44	1886	66,003	- 44
1861	204,457	44	1887	73,892	"
1862	192,612	44	1888	30,198	
1863	282,775	••	1889	29,986	66
1864	347,594	44	1890	50,854	66
1865	465, 194	44	1891	25,431	66
1866	404,252	.4	1892	13,883	66
1867	338,492	\$1.25	1893	16,099	44
1868	228,132	44	*1894	79,837	40
1869	257,485	• •	+1895	73,097	- 46
1870	168,180	64	11896	174,919	66
1871	165,431	" "	1897	106,279	44
1872	154,092	75	1898	98,027	- "
1873	254,760	4.	1899	153,188	16
1874	138,336	64	1900	624,273	46
1875	89,746	66		,	

NOTE.—The quantities given for the years 1852 to 1872 are on the authority of the Board of Trade, Philadelphia, and are probably under-estimated.

*Nine months only.

†Note.—After August 1st, 1891, duty on Round Coal 40 cents, on Culm or Slack, 15 cents.

Fiscal year begins October 1st, and ends September 30th.

GENERAL GOLD STATEMENNT.

Year ended Sept. 30th, 1900.

DISTRICT.	Days'	Tone Crushed.				Total yield of Gold.		
	Labor.		Oz.	Dwta.	Gre.	Oz.	Dwts.	Gro.
Oldham Sherbrooke Wine Harbor Lake Catcha Montague Harrigan Cove. Brookfield Renfrew Caribou & Moose River. Stormont Salmon River Other Districts	34632 18975 6414 436 13333 8348 33467 17959	9291 459 8348 26338 8825	9	11 13 15 18 5 13 3 5	2 9 7 10 3 19 20 22 21 11 7	1379 4763 2088 507 481 1705 2726 4450 1633 7745 1027 1888	16 12 17 3 3 9 16 15 6 18 14	8 9 0 66 77 12 13 222 22 10
		. 65744				30399	14	1 10

