

**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1997

The copy filmed here has been reproduced thanks to the generosity of:



Stauffer Library
Queen's University

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shell contain the symbol \rightarrow (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemple filmé fut reproduit grâce à la générosité de:



Stauffer Library
Queen's University

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemple filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.


Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole \rightarrow signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.



MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)



APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482-0300 - Phone
(716) 288-5989 - Fax

REPORT
OF THE
CANADIAN ARCTIC EXPEDITION
1913-18

VOLUME VIII:
MOLLUSKS, ECHINODERMS, COELENTERATES, Etc.

PART B:

CEPHALOPODA - - - By S. Stillman Berry
PTEROPODA - - - By William Healey Dall



G
670
1913 C2t
V 8 : 1 B

OTTAWA
F. A. ACLAND
PRINTER TO THE KING'S MOST EXCELLENT MAJESTY
1925

Issued Aug. 6, 1925

6
670
1913
C2t
v.008
pt.B
c.3

Report of the Canadian Arctic Expedition, 1913-18

VOLUME I: GENERAL INTRODUCTION, NARRATIVE, ETC.

- Part A: NORTHERN PARTY, 1913-18..... (To be prepared).
Part B: SOUTHERN PARTY, 1913-16. By Rudolph Martin Anderson..... (In preparation).

VOLUME II: MAMMALS AND BIRDS

- Part A: MAMMALS OF WESTERN ARCTIC AMERICA. By Rudolph Martin Anderson..... (In preparation).
Part B: BIRDS OF WESTERN ARCTIC AMERICA. By R. M. Anderson and P. A. Taverner..... (In preparation).

VOLUME III: INSECTS

- INTRODUCTION. By C. Gordon Hewitt..... (Issued December 10, 1920).
Part A: COLLEMBOLA. By Justus W. Folsom..... (Issued July 10, 1919).
Part B: NEUROPTEROID INSECTS. By Nathan Banks..... (Issued July 11, 1919).
Part C: DIPTERA.
Crane-flies. By Charles P. Alexander.
Mosquitoes. By Harrison G. Dyar.
Diptera (excluding Tipulidae and Culicidae). By J. R. Malloch..... (Issued July 14, 1919).
Part D: MALLOPHAGA AND ANOPLURA.
Mallophaga. By A. W. Baker.
Anoplura. By G. F. Ferris and G. H. F. Nuttall..... (Issued September 12, 1919).
Part E: COLEOPTERA. By
Forest Insects, including Ipidæ, Cerambycidae and Buprestidæ. By J. M. Swaine.
Carabidæ and Silphidæ. By H. C. Fall.
Coccinellidæ, Elateridæ, Chrysomelidæ and Rhynchophora (excluding Ipidæ). By C. W. Leng.
Dytiscidæ. By J. D. Sherman, Jr..... (Issued December 12, 1919).
Part F: HEMIPTERA. By Edward P. Van Duzee..... (Issued July 11, 1919).
Part G: HYMENOPTERA AND PLANT GALLS.
Sawflies. (Tenthredinoidea). By Alex. D. MacGillivray.
Parasitic Hymenoptera. By Charles T. Brues.
Wasps and Bees. By F. W. L. Sladen.
Plant Galls. By E. Porter Felt..... (Issued November 3, 1919).
Part H: SPIDERS, MITES AND MYRIAPODS.
Spiders. By J. H. Emerton.
Mites. By Nathan Banks.
Myriapods. By Ralph V. Chamberlin..... (Issued July 14, 1919).
Part I: LEPIDOPTERA. By Arthur Gibson..... (Issued January 10, 1920).
Part J: ORTHOPTERA. By E. M. Walker..... (Issued September 4, 1920).
Part K: INSECT LIFE ON THE WESTERN ARCTIC COAST OF AMERICA. By Frits
Johansen..... (Issued November 7, 1921).
PART L GENERAL INDEX..... (Issued December, 1922).

VOLUME IV: BOTANY

- Part A: FRESHWATER ALGAE AND FRESHWATER DIATOMS. By Charles W. Lowe..... (Issued February 20, 1923).
Part B: MARINE ALGAE. By F. S. Collins and M. A. Howe..... (In preparation).
Part C: FUNGI. By John Dearness..... (Issued June 1, 1923).
Part D: LICHENS. By G. K. Merrill..... (Issued July 16, 1924).
Part E: MOSSES. By R. S. Williams..... (Issued February 8, 1921).
Part F: MARINE DIATOMS. By Albert Mann..... (In press).

VOLUME V: BOTANY

- Part A: VASCULAR PLANTS. By James M. Macoun and Theo. Holm..... (Issued October 14, 1921).
Part B: CONTRIBUTIONS TO MORPHOLOGY, SYNONYMY AND GEOGRAPHICAL DISTRIBUTION OF ARCTIC PLANTS. By Theo. Holm..... (Issued February 10, 1922).
Part C: GENERAL NOTES ON ARCTIC VEGETATION. By Frits Johansen..... (Issued October 7, 1924).

VOLUME VI: FISHES, TUNICATES, ETC.

- Part A: FISHES. By F. Johansen..... (In preparation).
Part B: ASCIDIANS, ETC. By A. G. Huntsman..... (Issued November 29, 1922).

VOLUME VII: CRUSTACEA

- Part A: DECAPOD CRUSTACEANS. By Mary J. Rathbun..... (Issued August 18, 1919).
Part B: SCHIZOPOD CRUSTACEANS. By Waldo L. Schmitt..... (Issued September 22, 1919).
Part C: CUMACEA. By W. T. Gilman..... (Issued October 15, 1920).
Part D: ISOPODA. By P. L. Boone..... (Issued November 10, 1920).
Part E: AMPHIPODA. By Clarence R. Shoemaker..... (Issued September 7, 1920).
Part F: PYCNOGONIDA. By Leon J. Cole..... (Issued January 3, 1921).
Part G: EUPHYLLOPODA. By F. Johansen..... (Issued May 10, 1922).
Part H: CLADOCERA. By Chaney Juday..... (Issued June 23, 1920).
Part I: OSTRACODA. By G. O. Sars..... (In preparation).
Part J: FRESHWATER COPEPODA. By C. Dwight Marsh..... (Issued April 21, 1920).
Part K: MARINE COPEPODA. By A. Willey..... (Issued June 3, 1920).
Part L: PARASITIC COPEPODA. By Charles B. Wilson..... (Issued August 1, 1920).
Part M: CIRRIPEdia. By H. A. Pilsbry..... (In preparation).
Part N: THE CRUSTACEAN LIFE OF SOME ARCTIC LAGOONS, LAKES AND PONDS.
By F. Johansen..... (Issued December 30, 1922).

REPORT
OF THE
CANADIAN ARCTIC EXPEDITION
1913-18

VOLUME VIII:
MOLLUSKS, ECHINODERMS, COELENTERATES, Etc.

PART B:

CEPHALOPODA By S. Stillman Berry
PTEROFOIDA By William Healey Dall



OTTAWA
F. A. ACLAND
PRINTER TO THE KING'S MOST EXCELLENT MAJESTY
1925

Issued Aug. 6 1925

2

The Cephalopoda Collected by the Canadian Arctic Expedition, 1913-18

By S. STEILMAN BERRY,

Rollands, California.

The cephalopods collected by the Canadian Arctic Expedition during the years 1913 to 1916 comprise only a few fragments, principally mandibles, taken from the stomachs of two species of seals, the Bearded Seal, *Erymnathus barbatus* (Erxleben), and the Common Rough Seal, *Phoca hispida* Schreber. In the light of present knowledge alone, proper identification of such material at the very least, is hazardous. But from the notes, and more especially from the figures given, it is just possible that a way will be left open for throwing a more illuminating light upon the material at some time in the future.

It is hoped that the bibliographical portion of the present paper may prove of more immediate utility. It is unfortunate that knowledge of the cephalopods of the American Arctic is so incomplete, scattering, and fragmentary, as the study of this list shows.

MATERIAL COLLECTED

For convenience in reference, the material collected is listed in order according to the numbers under which it is entered in the author's card register.

[726]

Numerous fragments taken from stomach of *Phoca hispida* Schreber, station 29f, ca. 30 fathoms, lat. 70° 13' N., long. 140° 50' W., a little east of Alaska-Yukon boundary, April 4, 1914.

The material here entered consists of 3 fragmentary and semi-digested buccal masses of cephalopods, together with a few odd shreds of tissue and some 58 fragments of beaks, the latter consisting of half mandibles only. None even those persisting in the buccal tissues are entire. Each is split mesially. Evidently the median region of the mandibles is not only narrow in this species but more than ordinarily weak. In fact, one is reminded very much of the appearance of the valves of the chiton *Schizoplax* after the dissolution of the median uncalcified wedge. There are other peculiarities of shape and proportion (Fig. 1). On the whole the mandibles are fairly uniform in size. None




Fig. 1. - The four mandibular fragments from the buccal mass of a cephalopod taken at Station 29f [726]; camera outline; the two outermost figures are internal views.

are very large, particularly in comparison with the relatively enormous size of the buccal mass. Such structures should surely be identifiable with the passage of time—consequently it seems worth while to add figures of the four fragments dissected from the best preserved of the buccal masses. It is not, however, a glad commentary on the sufficiency of our knowledge regarding animals possessing such importance in the economy of the sea as these, that without more complete specimens or a better base of attack than is afforded by the literature, one dares not risk a guess at even the genus or family of cephalopods represented by the fragments.

[727]

1 mandible from stomach of *Phoca hispida* Schreber, station 29f, about 30 fathoms, lat. $70^{\circ} 13' N.$, long. $140^{\circ} 50' W.$, a little east of Alaska-Yukon boundary, April 4, 1914.

A single dorsal mandible from the same stomach as the specimens entered as No. 726 is larger than the others, altogether differently shaped, and entire. Its cutting edge is sharp and strongly curved, as in many decapods (Fig. 2). It clearly represents a quite different species.




Fig. 2. Mandible of cephalopod from Station 29f [727], camera outline.

[728]

1 mandible from stomach of *Eriquathus barbatus* (Erleben), Station 42a, Bernard harbour, Dolphin and Union strait, Northwest Territories, October 22, 1915.

This is a single dorsal mandible somewhat resembling No. 727, but with the cutting edge appreciably shorter and stouter (Fig. 3). This specimen, which is preserved dry, is very likely representative of a third species.




Fig. 3. Mandible of cephalopod from Station 42a [728], camera outline, same scale as Fig. 2.

In the diary of Captain Robert A. Bartlett, Report of the Department of the Naval Service for the Fiscal Year ending March 31, 1915, Ottawa, p. 35, there is an entry under Nov. 21st, (1913): "10 a.m. Sounding, 36 fathoms. Dredge to-day secured an octopus Lat. $72^{\circ} 56'$ — Long. $163^{\circ} 54' W.$ at 5 p.m." All the specimens secured by Mr. James Murray, marine biologist and the scientific staff of the *Karluk* during the drift northwest of Alaska in 1913 were lost with the ship north of Herald island, on January 11, 1914.

BIBLIOGRAPHY

It has been suggested that a bibliography for the Cephalopoda of the American Arctic would be found useful in connection with other bibliographies appearing in this series of papers. Defining the American quadrant as those Arctic waters lying between Davis and Bering straits, with possibly the 60th parallel as an arbitrary southern boundary, the following list of titles has been compiled as a contribution to this end. It is thought to be complete, at least to the point of serviceability, if unfortunately not to that of perfection. No attempt is here made to correct the synonymy quoted.

A few titles incuded have not been available for consultation and are consequently quoted from the citations of other authors. Such are enclosed in brackets.

DALL, W. H.

1881. New or specialty interesting shells of the Point Barrow Expedition. Proceedings United States National Museum, v. 7, pp. 523-526, pl. 2. f. 8. 1881.

Records *Octopus gröenlandicus* from Point Barrow, Alaska.

DEWHURST, H. W.

1831. The natural history of the order Cetacea, and the oceanic inhabitants of the Arctic regions. p. i - xx, 1 - 291 (331), with numerous pls. and figs., 8 vo., London, 1831.

Contains note on *Sepia gröenlandica*, n. sp. (*nomen nudum*, p. 263), from "Greenland Seas," and various references to cephalopods as food of Arctic whales.

ESCHRICHT, D. F.

1836. Cirroteuthis Mülleri, eine neue Gattung der Cephalopoden bildend. Acta Academiae Caesaris Leopoldino Carolina Naturae Curiosorum, v. 18, p. 627-634, pl. 46-48, 1836.

Cirroteuthis Mülleri, n. gen. and sp. described from Jacobshavn, Greenland.

FABRICIUS, O.

- [1780. Fauna Groenlandica. 8vo, Hafniae et Lipsiae, 1780, pp. 358-60.]
Records *Sepia loligo* and *S. octopodia* from Greenland.

GRIEG, J. A.

1896. Bidrag til kundskaben om Vestlandets mollusker. Bergens Museums Aarbog, no. 10, pp. 1-33, pl. 1, 1896.

Records *Rossia glaucopsis* from 100-150 fath., off Kraakøskallen; 120 fath., off Baardshaug, Selbjørnefjorden; and 60-70 fath., Hjeltefjorden. *Scpiola rondletii* from Kraakøskallen and Godø Sund, Greenland.

1909. Brachiopods and mollusks with a supplement to the echinoderms. Report 2nd Norwegian Arctic Expedition in the *Fram*, 1898-1902, No. 20, pp. 1-15, pl. 1, October, 1909.

Records *Rossia palpibrosa* from 8 m., entrance to Stordalen, Havnefjord, Ellesmere island.

HOYLE, W. E.

1886. Report on the Cephalopoda collected by H.M.S. *Challenger* during the years 1873-76. Voyage of the *Challenger*, v. 16, pt. 44, pp. i-vi, 1-246, text f. 1-10, map, pl. 1-33, 1886.

On p. 222 gives summary of the then known cephalopod fauna of the Arctic faunal region, listing 9 species, referable to 6 genera.

MÖLLER, H. P. C.

1842. Index Molluscorum Groenlandiæ. Naturhistorisk Tidsskrift, v. 1, p. 76-97, 1842.

Lists following cephalopods from Greenland:

Onychoteuthis Fabricii
Onychoteuthis? amoena n. sp.
Rossia palpebrosa
Cirrotheuthis Mülleri
Octopus granulatus

MÖREN, O. A. L.

[1857. Fortegnelse over Groenlands Bløddyr (Mollusca Groenlandica). Naturh. Tillaeg til en Faunistisk og Statistisk beskrivelse af Groenland (H. Rink), pp. 75-100, Kjøbenhavn, 1857.]

Following Greenland cephalopods listed on pp. 88-89 (*teste* Johannsen).

Octopus groenlandicus
Cirrotheuthis Mülleri
 Jakobshavn
Rossia palpebrosa
Rossia Mølleri
Leachia hyperborca
Gonatus Fabricii
 Amikok
 ? *Sepioida atlantica*

OWEN, R.

1834. Mollusca—Cephalopoda (with preliminary note by J. C. Ross). Narrative Second Voyage in Search of a North-West Passage, etc., by Sir John Ross, C.B., etc., Appendix, Natural History, pp. xcii-xcix, pl. B, f. 1, pl. C, June, 1834. (Complete work dated 1835).

Records *Rossia palpebrosa* n. gen. and sp. from Etwin bay, Prince Regent's inlet. [Somerset island, Regent inlet.]

PFEFFER, G.

1908. Cephalopoda. Nordisches Plankton, 9 Lief., pt. 4, pp. 1-116, text f. 1-120, 1908.

1912. Die Cephalopoden der Plankton-Expedition. Zugleich eine monographische Übersicht der oegopsiden Cephalopoden. Ergebnisse Plankton-Expedition der Humboldt-Stiftung, Bd. 2, pp. i-xxi, 1-815, atlas of 48 pls., 1912.

On p. 241 records *Gonatus fabricii* from Umanak and Disco, Greenland. Brief discussion of Arctic fauna on pp. 793-797.

POSSELT, H. J.

1898. Grønlands Brachiopoder og Bløddyr. Meddelelser om Grønland 23, pp. i-xix, 1-298, pl. 1-2, map, 1898.

Contains the following West Greenland locality records (pp. 269-283):

Cirrotheuthis Mülleri
 Jakobshavn
 North Greenland.
Octopus arcticus
 Julianehaab
 Ivigtut
 Lat. 65° 36' N., long. 56° 24' W.
 Holstensborg.

Rossia palpebrosa

West Greenland.

Rossia Mølleri

West Greenland.

Rossia subleris

289 fath., lat. 65° 30' N., long. 55° 26' W.

349 fath., lat. 65° 36' N., long. 56° 24' W.

235 fath., lat. 66° 19' N., long. 56° 28' W.

Rossia glaucopsis

West Greenland.

Rossia Hyalli

25-35 fath., Umanak

40 fath., Prøven.

Rossia megaphra

349 fath., lat. 65° 36' N., long. 56° 24' W.

Spirula

Waigat, lat. 69° 41' N.

Illex illecebrosus

Frederikshaab.

Architeuthis sp.

Holstensborg.

Gonatus Fabricii

Julianehaab, Ivigtut, Godthaab, Sukkertoppen, Egedesminde, Christianshaab, Jakobshavn, Godhavn, Ritenbenk, and Umanak.

Taonius hyperboreus

North Greenland.

PROSCH, V.

1847. Nogle nye Cephalopoder, beskrevne og anatomisk undersøgte. Kongelige Danske Videnskabernes Selskabs Skrifter (5), naturvidenskabelig og matematisk Afdeling, v. 1, pp. 53-72 [1-20], pl., 1847.

Octopus arcticus n. sp., described from Greenland.

REINHARDT, J. T., AND PROSCH, V.

1846. On *Sciadephorus Mülleri* (Eschr.). Kongelige Danske Videnskabernes Selskabs Skrifter, v. 12, pp. 165-221 [1-40], pl. 1-5, 1846.

STEENSTRUP, J. J. S.

1856. Hectocotylidannelsen hos Octopodslægterne Argonauta og Tremoctopus, oplyst ved Iagttagelse af lignende Dannelse hos Blæksprutterne i Almindelighed. Videnskabernes Selskabs Skrifter (5), naturvidenskabelig og matematisk Afdeling, v. 1, pp. 185-216, pl. 1-2, 1856.

On p. 200 describes *Leachia hyperborea* n. sp. from North Greenland.

1857. Hectocotylus-formation in Argonauta and Tremoctopus explained by observations on similar formations in the Cephalopoda in general. Annals and Magazine Natural History (2), v. 20, pp. 81-114, pl. 2-3, August, 1857.

An English translation of preceding paper, the reference to *Leachia hyperborea* being found in the footnote on p. 96.

1861. Overblik over de i Kjöbenhavns Museer opbevarede Blæksprutter fra det aabne Hav (1860-61). Oversigt over det Kongelige Danske Videnskabernes Selskabs Forhandling 1861, pp. 69-86 [1-18], 1861.

1880. Orientering de Ommatostrephagtige Blæksprutters indbyrdes Forhold. Oversigt over det Kongelige Danske Videnskabernes Selskabs Forhandlinger 1880, pp. 73-110 [1-10]. 13 f. in text, pl. 3, 1880.

Note on distribution of *Todarodes* and *Ommatostrephes* on p. 107. *Ommatostrephes Bartramii* recorded from stomachs of fish taken in Davis strait, etc.

1881. Professor A. E. Verrils [Verrill] to nye Cephalopod-lægter: *Sthenoteuthis* og *Lestoteuthis*. Oversigt over det Kongelige Danske Videnskabernes Selskabs Forhandlinger 1881, pp. [1-27], text f. A-B, pl. 1, 1881.

VAN HOFFEN, E.

[1897. Die Fauna und Flora Grönlands. In Grönland-Expedition d. Gesellsch. f. Erdkunde zu Berlin 1891-1893, 1897.]

NOTE. For courteous help in the compilation of the foregoing bibliography, as well as securing certain of the works therein noted, thanks are due to Mr. Frits Johansen, formerly of the Canadian Arctic Expedition, 1913-18, and to Dr. Theodor Mortensen of the Copenhagen Museum.

For comparison of the Arctic cephalopod fauna with that of the Antarctic see

BERRY, S. S.

1917. Cephalopoda. Australasian Antarctic Expedition, 1911-1914, Scientific Reports Series C—Zoology and Botany, v. 4, pt. 2, pp. 1-39, text f. 1-30, pl. 10-14, March, 1917.

And the papers cited in the further bibliography there appended.

The Pteropoda Collected by the Canadian Arctic Expedition 1913-18, with description of a New Species from the North Pacific

BY WILLIAM HEALEY DALL¹

Honorary Curator of Mollusks, United States National Museum.

A small collection of Pteropoda was referred to me by the Arctic Publications Committee for a report upon them. They consisted mainly of material collected by Mr. Frits Johansen, biologist with the Canadian Arctic Expedition, with a few Opisthobranchs and other specimens added from other northern Canadian localities.

A peculiarity of the expedition series is the absence in great measure of adult specimens, nearly all the specimens of *Clione* and all the specimens of *Spiratella* (*Limacina* of authors), were young, especially the latter; although Mr. Johansen frequently mentions its presence, not a single adult specimen is contained in the series. As most of the specimens were obtained near shore, it may indicate that it is the habit of the young to avoid the open sea until mature.

Clione borealis Phipps.

Station 6 *a, b*.² Latitude 56° 26' N., longitude 133° W., June 21, 1913. — One *Spiratella* and one young *Clione* obtained at the surface and near the surface (F.J.).

Station 9*a*. Latitude 55° 2' N., longitude 141° W., June 27, 1913. Water temperature at 1 p.m. 51° to 51°.6 F. Several young *Clione* and many *Spiratella*. (F.J.)

Station 12*a, f*. Latitude 54° 38' N., longitude 157° 45' W., June 30, 1913. Water temperature 50°.7 to 51°.3 F. Many young *Clione* and one *Spiratella* from, at or near the surface. (F.J.)

Station 14. Latitude 54° 23' N., longitude 164° 45' W., July 2, 1913. One young *Clione* obtained at the surface. (F.J.)

Station 57*a*. Cape Smyth, near Point Barrow, Alaska, August 5, 1916. Several *Clione* were obtained at or near the surface in a strong northerly current. (F.J.)

Station 25*b, c*. Ten miles east of Point Barrow, Alaska, off the Plover islands, August 27-28, 1913. Water temperature 30°.2 to 30°.3 F. Depth about two fathoms. One *Clione* obtained. (F.J.)

Station 27*c*. Camden bay, inside of Collinson point, Alaska, September 15, 1913. Several *Clione* captured. (F.J.)

Station 27*n*. Off Collinson point, Alaska, Oct. 5, 1913. One *Clione* from under ten inches of ice in about six feet of water at a temperature of 30°.5 F. (F.J.)

¹ By permission of the Director of the United States Geological Survey.

² The collection stations are arranged in order from west to east.

Station 28k. Jan. 4, 1914. Camden bay, inside Collinson point, shallow water under four feet of ice; water temperature 28°.5 F. (F.J.)

The following data are derived from Mr. Johansen's notes:

One adult *Clione* was 30 mm. long, the spread of the flippers 20 mm. when alive. The muscles of the tail were clearly visible through the transparent body. The purplish tint of the tail extended with decreasing intensity forward for about 5 mm. The oral tentacles were deep rose purple; the oesophagus rose colour, as also the edge of the body around the base of the parapodia; the intestinal mass was blackish brown. The rest of the body was pale and translucent. The protruded oral tentacles were about 3 mm. long, the hook bearing captacula when fully extended, appeared longer.

When swimming, the parapodia were bent toward the ventral side, returning to a nearly horizontal position, and repeating.

The animal came up with the water in a hole cut through the ice and was caught swimming just below the surface film of new ice.

Placed in a bottle of sea water which began to freeze, the animal became immobile among the ice particles. Only after arrival at camp when the ice began melting did it begin to move again. The tail is much used by the animal as a kind of rudder when changing direction of movement. When moving up or down the tail is recurved dorsally or sometimes ventrally. At times the animal would revolve in one place with the tail curved against the side of the body and the fins continually moving. When swimming horizontally it always tried to keep the ventral side downward. When at rest it remained in a vertical position, head downward, or sank upon its ventral side to the bottom of the container. The resting periods were always shorter than those of activity. The fins if brought into contact with ice or the glass of the container, or when the animal is moribund, are folded up and held close to the body. They are subject to extreme contraction when placed in a preservative.

In the warmth of the laboratory the movements decreased and the animal became sluggish, but if the container was placed in a colder situation activity revived. From time to time the animal exuded mucus which seemed to embarrass its movements. On January 5, the container was put in a cool place and the water congealed during the night. When thawed the Pteropod was still alive, though less active and this was continued for a week. January 12, it died, having been kept for eight days without renewal of water or food.

Station 37r. Inner harbour at Bernard harbour, Dolphin and Union strait, Oct. 16, 1914. Under four inches of ice in about seven feet of water at a temperature of 30°.1 F. One adult *Clione*. (F.J.)

Station 43a. Off Cockburn point, Dolphin and Union strait, September 13, 1915. Water temperature 32° F. about fifty fathoms deep. One *Clione* obtained. (F.J.)

In view of the fact that so many good dissections have been published of this species, and that nearly all the specimens were immature and not in the best condition, it is not thought worth while to attempt anatomical work upon them.

It may be noted that young are more deeply tinted than the adults and one specimen of which Mr. Johansen has made a rough coloured sketch is quite noticeably red, while others are less so.

Spiratella pacifica Dall?

This genus is more generally known by the name of *Limacina*. The minute young specimens in the collection, perhaps from being preserved with formalin, have lost their shelly matter, and it is impossible to say whether they belong to the Pacific type or to the better known and much more common Atlantic species *S. helicina* Phipps.

Station 13g, b. Latitude 54° 30' N., longitude about 157° W., June, 1913. In marine plankton, numerous young mixed with young *Clione*. (F.J.)

Station 6a, b. Latitude 56° 26' N., longitude 133° W., June 24, 1913. One *Spiratella* from the surface. (F.J.)

Station 9a. Latitude 55° 02' N., longitude 144° W., June 27, 1913. With marine Plankton, numerous larval or very young specimens mixed with young *Clione*. (F.J.)

Station 12a, f. Latitude 51° 38' N., longitude 157° 45' W., June 30, 1913. Water temperature 50°.7 to 51°.3 F. One *Spiratella* with many young *Clione*, from at or near the surface. (F.J.)

Station 13a, f. Latitude 54° 30' N., longitude 159° 42' W., July 1, 1913. Water temperature 47°.5 to 48°.1 F. Many *Spiratella* at or near the surface. (F.J.)

Station 15a, f. Latitude 55° 22' N., longitude 165° 45' W., July 3, 1913. Noon to 3 p.m. Many *Spiratella*. (F.J.)

Station 20a. Grantley harbour, Port Clarence, Alaska, July 30, 1913. Several *Spiratella* from the surface. (F.J.)

Station 40d. Off Bernard harbour, Dolphin and Union strait, June 8, 1915. Depth of water nine fathoms, temperature 32° F. Six vertical hauls through a hole in the ice to five fathoms depth, produced several veliger larvae of *Spiratella*. (F.J.)

Station 46k. Off west end of Chantry island, in Dolphin and Union strait, June 10, 1916. Depth of water about eight fathoms, the temperature 32° F. A vertical haul through four feet of ice to depth of five fathoms produced several veliger larvae of *Spiratella*. (F.J.)

SPECIMENS FROM OTHER SOURCES

Clione borealis Phipps.

1. Half digested young from the stomach of *Myotæcephalus grönlandicus*, Nain, Labrador; collected by Owen Bryant, 1908.

2. Port Burwell and Wakeham Bay Ungava, *Neptune* Expedition; collected by A. Halkett, 1904.

3. Black Tickle, Labrador; collected by A. Halkett, 1903.

Larval Gastropod

Station 13a, c. Latitude 54° 30' N., longitude 159° 42' W. Specimen in marine plankton, not identifiable.

Gastroteron (pacificum Bergh var.?) cinereum Dall, n. sp.

1. Skidegate inlet, Queen Charlotte islands, British Columbia, July, 1910, W. Spreadborough, collector. Victoria Memorial Museum, Mollusks, No. 3177; cotype, presented to United States National Museum.

2. West side of Vancouver island, Ucluelet, British Columbia at low tide, C. H. Young and W. Spreadborough, May, 1919. Cotype, Victoria Memorial Museum, Ottawa, Cat. Mollusks, No. 3176. (Fig. 4).¹

¹Drawn from alcoholic specimen, by Claude E. Johnson, artist of Biological Division, Victoria Memorial Museum. The lack of lateral symmetry is probably due to contraction in the preservative medium.

Both the North Atlantic, Asiatic, and North Pacific species of *Gastropteron* are of a reddish colour sprinkled densely with darker red dots. The present form is of a uniform dusky slate colour and of a smaller size than the Pacific species described by Bergh. It is an interesting addition to the sparse list of known species. The spread of the parapodia totals 10 to 15 mm., the length of the body 8 to 11 mm. (in spirits), in the living state it was probably larger. The anterior shield is about one-third the whole length of the body. Like *G. pacificum*, it lacks a posterior flagellum on the mantle.




Fig. 1. *Gastropteron pacificum* Bergh var. *G. canicum* Dill, new species. Left, dorsal aspect; right, ventral. X 4. Upper outline figure, natural size.

Eggs of fish?

Station 7a, b. Latitude 55° 42' N., longitude 136° 20' W., June 25, 1918. Eggs, probably of fish, were found floating on the surface. (F.J.)

Young of Sipunculoid worm?

Station 57. 1 fathom in marine plankton, off Cape Smyth, near Point Barrow, Alaska.

This puzzled our experts, but is left without positive determination, being badly contracted by the preservative.



VOLUME VIII: MOLLUSKS, ECHINODERMS, COELENTERATES, ETC.

- Part A: MOIUSES, RECENT AND PLEISTOCENE.** By William H. Dall. (Issued September 24, 1919).
- MOLLUSKS (Supplementary).** By W. H. Dall. (Issued November 27, 1924).
- Part B: CEPHALOPODA AND PTEROPODA.**
 Cephalopoda. By S. S. Berry.
 Pteropoda. By W. H. Dall. (In press).
- Part C: ECHINODERMS.** By Austin L. Clark. (Issued April 6, 1920).
- Part D: BRYOZOA.** By R. C. Osburn. (Issued February 20, 1923).
- Part E: ROTATORIA.** By H. K. Harring. (Issued December 31, 1921).
- Part F: CHAETOGNATHA.** By A. G. Huntsman. (In preparation).
- Part G: ALGYONARIA AND ACTINARIA.** By A. E. Verrill. (Issued April 23, 1922).
- Part H: MEDUSAE AND CTENOPHORA.** By H. B. Bigelow. (Issued June 30, 1920).
- Part I: HYDROIDS.** By C. McLean Fraser. (Issued August 21, 1923).
- Part J: PORIFERA.** By A. Dendy and L. M. Frederick. (Issued July 5, 1924).

VOLUME IX: ANNELIDS, PARASITIC WORMS, PROTOZOANS, ETC.

- Part A: OLIGOCHAETA.**
 Lumbricidae. By Frank Smith.
 Enchytraeidae. By Paul S. Welch. (Issued September 29, 1919).
- Part B: POLYCHAETA.** By Ralph V. Chamberlin. (Issued November 16, 1920).
- Part C: TIRIDINEA.** By J. P. Moore. (Issued February 4, 1921).
- Part D: GEPHYREA.** By Ralph V. Chamberlin. (Issued June 10, 1920).
- Part E: ACANTHOCEPHALA.** By H. J. Van Cleave. (Issued April 7, 1920).
- Part F: NEMATODA.** By N. A. Cobb. (In preparation).
- Part G-H: TREMATODA AND CESTODA.** By A. R. Cooper. (Issued February 4, 1921).
- Part I: TURBELLARIA.** By A. Harsell. (In preparation).
- Part J: POLYCHAETA (Supplementary).** By J. H. Ashworth. (Issued September 29, 1924).
- Part K: NEMERTINI.** By Ralph V. Chamberlin. (In preparation).
- Part L: SPOROZOA.** By J. W. Mavor. (In preparation).
- Part M: FORAMINIFERA.** By J. A. Cushman. (Issued February 6, 1920).

VOLUME X: HYDROGRAPHY, TIDES, ETC.

- Part C: TIDAL OBSERVATIONS AND RESULTS.** By W. Bell Dawson. (Issued October 1, 1920).
- Part D: HYDROGRAPHY.** (In preparation).

VOLUME XI: GEOLOGY AND GEOGRAPHY

- Part A: THE GEOLOGY OF THE ARCTIC COAST OF CANADA, WEST OF THE KENT PENINSULA.** By J. J. O'Neill. (Issued July 8, 1924).
- Part B: MAPS AND GEOGRAPHICAL NOTES.** By Kenneth C. Chipman and John R. Cox. (Issued July 8, 1924).

VOLUME XII: THE COPPER ESKIMOS

- Part A: THE LIFE OF THE COPPER ESKIMOS.** By D. Jenness. (Issued January 14, 1922).
- Part B: THE PHYSICAL CHARACTERISTICS OF THE WESTERN AND COPPER ESKIMOS.** By D. Jenness. (Issued May 23, 1923).
- Part C: THE OSTEOLOGY OF THE WESTERN AND CENTRAL ESKIMOS.** By John Cameron. (Issued June 23, 1923).

VOLUME XIII: ESKIMO FOLK-LORE

- Part A: ESKIMO MYTHS AND TRADITIONS FROM ALASKA, THE MACKENZIE DELTA AND CORONATION GULF.** By D. Jenness. (Issued November 15, 1924).
- Part B: STRING FIGURES OF THE ESKIMOS.** By D. Jenness. (Issued August 8, 1924).

VOLUME XIV: ESKIMO SONGS

- SONGS OF THE COPPER ESKIMOS.** By Helen H. Roberts and D. Jenness. (In press).

VOLUME XV: ESKIMO LANGUAGE AND TECHNOLOGY

- Part A: COMPARATIVE GRAMMAR AND VOCABULARY OF THE ESKIMO DIALECTS OF POINT BARROW, THE MACKENZIE DELTA AND CORONATION GULF.** By D. Jenness. (In preparation).
- Part B: TECHNOLOGY OF THE COPPER ESKIMOS.** (To be prepared).

VOLUME XVI: ARCHAEOLOGY

- CONTRIBUTIONS TO THE ARCHAEOLOGY OF WESTERN ARCTIC AMERICA.** (To be prepared).

(Index issued with last number of each Volume.)



