

**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**




Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 2000

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- ☐ Coloured pages / Pages de couleur
- ☒ Pages damaged / Pages endommagées
- ☐ Pages restored and/or laminated / Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed / Pages décolorées, tachetées ou piquées
- ☐ Pages detached / Pages détachées
- ☐ Showthrough / Transparence
- ☐ Quality of print varies / Qualité inégale de l'impression
- ☐ Includes supplementary material / Comprend du matériel supplémentaire
- ☐ Pages wholly or partially obscured by errata slips, tissues, etc., have been refiled to ensure the best possible image / Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.
- ☐ Opposing pages with varying colouration or discolourations are filmed twice to ensure the best possible image / Les pages s'opposant ayant des colorations variables ou des décolorations sont filmées deux fois afin d'obtenir la meilleure image possible.

**This item is filmed at the reduction ratio checked below /
Ce document est filmé au taux de réduction indiqué ci-dessous.**



The copy filmed here has been reproduced thanks to the generosity of:


Gerstein Science Information Centre
University of Toronto

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol ➡ (meaning "CONTINUED"), or the symbol ▼ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:


Gerstein Science Information Centre
University of Toronto

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.


Un des symboles suivants apparaît sur la dernière image de chaque microfiche, selon le cas: le symbole ➡ signifie "A SUIVRE", le symbole ▼ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.



MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)



APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482 - 0300 - Phone
(716) 288 - 5989 - Fax

ack.
25/9/10
Pamph.
Astroph.
12

With Compliments of the Author.

Reprinted from the ASTROPHYSICAL JOURNAL, Vol. XXVIII, No. 4, November 1908

THE ORBIT OF γ ORIONIS

3 1761 04304 7199

J. S. PLASKETT

PRINTED AT THE UNIVERSITY OF CHICAGO PRESS



THE ORBIT OF ι ORIONIS

By J. S. PLASKETT

Acting on a suggestion kindly made to me by Dr. Schlesinger, a least-squares solution has been applied to the elements of the orbit of ι *Orionis* recently published in this *Journal*.¹ As preliminary elements those determined in the above paper were taken, and the 113 plates were grouped into 26 normal places, the same as before except for some minor changes introduced by a consistent system of weighting. An ephemeris having been computed from the preliminary elements, the coefficients of the unknowns in the observation equations were calculated from the formulae of Lehmann-Filh  s² for each of these places. The period was considered as very closely determined by the method previously used and a correction considered unnecessary.

From the observation equations were formed the normal equations below, the following factors being introduced for homogeneity:

$$\begin{aligned}x &= \delta K \\y &= K \delta e = 112 \delta e \\z &= K \delta \omega = 112 \delta \omega \\u &= \frac{K \mu}{(1-e^2)^{3/2}} \delta T = 83.46 \delta T \\\tau &= \delta \gamma\end{aligned}$$

NORMAL EQUATIONS

$$\begin{aligned}+44.968x &- 29.522y + 3.322z &+ 0.327u &- 14.201\tau &+ 98.606 = 0 \\+228.507y &- 41.899z &+ 65.630u &+ 39.089\tau &- 244.074 = 0 \\+51.724z &- 5.5431u &- 33.920\tau &- 33.442 = 0 \\&&+ 76.940u &+ 25.424\tau &- 56.373 = 0 \\&&&+ 95.000\tau &- 15.870 = 0\end{aligned}$$

The solution of these equations gave the following corrected elements:

¹ *Astrophysical Journal*, **27**, 272, 1908.

² *Astronomische Nachrichten*, **136**, 17, 1894.

	Preliminary	Corrected
K	112.0	109.92
e	0.75	0.7552
ω	110°	113.31
T	1.94 days	1.991 days
γ	+20.7 km	+21.34 km
Period	29.136 days	29.136 days
$a \sin i$	29,680,000 km	28,867,000 km


A comparison of the residuals obtained from an ephemeris and from substitution in the observation equations showed some differences of over a kilometer and a second solution was necessary. For preliminary elements those obtained in the first solution, except T , which was increased to 2.01 days, were used. The change in T was due to a better agreement thereby produced in the residuals. Using the same substitutions for homogeneity there result the normal equations

$$\begin{aligned}
 +45.058x - 28.047y + 2.576z + 0.726u - 14.660v - 0.771w &= 0 \\
 +266.994y - 45.686z + 65.208u + 43.697v + 117.212w &= 0 \\
 +50.372z - 48.247u - 33.053v - 70.762w &= 0 \\
 +64.085u + 22.897v + 95.820w &= 0 \\
 +95.000v + 34.800w &= 0
 \end{aligned}$$

Their solution gives for final elements the following:

K	109.90 ± 1.100 km
e	0.7543 ± 0.0046
ω	113.28 ± 1.083
T	1.993 ± 0.022 days = Julian Day 2,417,587.993
γ	+21.34 ± 0.856 km
Period	29.136 days
$a \sin i$	28,907,000 km

An ephemeris computed from these elements shows that $\Sigma p\eta$ has been reduced from 2094 to 2181, the probable error of an observation of unit weight becoming ± 6.88 km, while the probable errors of the elements become those given above. The changes from the first solution are very small but the agreement between the residuals is now satisfactory. The velocity-curve corresponding to the final elements is given in the accompanying figure, with the positions of the normal places as small circles. A comparison of



this with the previous figure shows that the general trend of the observations is more closely followed. The probability of a secondary disturbance seems somewhat less than with the original elements, and this is further lessened by a knowledge of the fact that the normal places with the highest residuals contain always one or more observations with the universal spectroscope where the temperature control was poor and the spectra contained only two measurable lines.

The result of this computation seems to justify Dr. Schlesinger's contention that, even in the case of spectra in which accurate measurement is impossible, the least-squares solution will give the best determination of the elements of a binary orbit.

DOMINION OBSERVATORY, OTTAWA

July 1908

