







# IMAGE EVALUATION TEST TARGET (MT-3)



APPLIED IMAGE, Inc  
1653 East Main Street  
Rochester, NY 14609 USA  
Phone: 716/482-0300  
Fax: 716/288-5989

© 1993, Applied Image, Inc., All Rights Reserved



**CIHM  
Microfiche  
Series  
(Monographs)**

**ICMH  
Collection de  
microfiches  
(monographies)**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

**© 1993**

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

☒ Coloured covers/  
Couverture de couleur

☐ Covers damaged/  
Couverture endommagée

☐ Covers restored and/or laminated/  
Couverture restaurée et/ou pelliculée

☐ Cover title missing/  
Le titre de couverture manque

☐ Coloured maps/  
Cartes géographiques en couleur

☐ Coloured ink (i.e. other than blue or black)/  
Encre de couleur (i.e. autre que bleue ou noire)

☐ Coloured plates and/or illustrations/  
Planches et/ou illustrations en couleur

☒ Bound with other material/  
Relié avec d'autres documents

☒ Tight binding may cause shadows or distortion  
along interior margin/  
La reliure serrée peut causer de l'ombre ou de la  
distorsion le long de la marge intérieure

☐ Blank leaves added during restoration may appear  
within the text. Whenever possible, these have  
been omitted from filming/  
Il se peut que certaines pages blanches ajoutées  
lors d'une restauration apparaissent dans le texte,  
mais, lorsque cela était possible, ces pages n'ont  
pas été filmées.

☒ Additional comments: /  
Commentaires supplémentaires:

Pagination is as follows: p. [99]-130.

This item is filmed at the reduction ratio checked below/  
Ce document est filmé au taux de réduction indiqué ci-dessous.

|                          |                          |                          |                          |                                     |                          |
|--------------------------|--------------------------|--------------------------|--------------------------|-------------------------------------|--------------------------|
| 10X                      | 14X                      | 18X                      | 22X                      | 26X                                 | 30X                      |
| <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input checked="" type="checkbox"/> | <input type="checkbox"/> |
| 12X                      | 16X                      | 20X                      | 24X                      | 28X                                 | 32X                      |

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

☐ Coloured pages/  
Pages de couleur

☐ Pages damaged/  
Pages endommagées

☐ Pages restored and/or laminated/  
Pages restaurées et/ou pelliculées

☒ Pages discoloured, stained or foxed/  
Pages décolorées, tachetées ou piquées

☐ Pages detached/  
Pages détachées

☒ Showthrough/  
Transparence

☐ Quality of print varies/  
Qualité inégale de l'impression

☐ Continuous pagination/  
Pagination continue

☐ Includes index(es)/  
Comprend un (des) index

Title on header taken from: /  
Le titre de l'en-tête provient:

☐ Title page of issue/  
Page de titre de la livraison

☐ Caption of issue/  
Titre de départ de la livraison

☐ Masthead/  
Générique (périodiques) de la livraison

The copy filmed here has been reproduced thanks to the generosity of:



National Library of Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol → (meaning "CONTINUED"), or the symbol ▼ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:


Bibliothèque nationale du Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole → signifie "A SUIVRE", le symbole ▼ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.





512.3

Y8

SOLVABLE QUINTIC EQUATIONS WITH COMMENSURABLE COEFFICIENTS.

By GEORGE PAXTON YOUNG, UNIVERSITY COLLEGE, TORONTO, CANADA.

*Reprinted from American Journal of Mathematics, Vol. X, No. 2.*

512.3

Y8

QA404

.5

Y68

1887

fol.

31843

## ***Solvable Quintic Equations with Commensurable Coefficients.***

BY GEORGE PAXTON YOUNG, *University College, Toronto, Canada.*

### OBJECT OF THE PAPER.

§1. Some time ago, in the *American Journal of Mathematics* (Vol. VI, page 103), the present writer sketched a general method for finding the roots of solvable irreducible equations of the fifth degree. The method was partially developed, and its application to certain forms of quintic equations was shown. It is now proposed to give the method the farther development necessary to make it applicable, by a definite and certain process, and without any difficulty beyond the labor of operation, to all solvable irreducible quintics having commensurable coefficients. The following equations will be solved as examples of the application of the theory:

1.  $x^5 + 3x^2 + 2x - 1 = 0.$
2.  $x^5 - 10x^3 - 20x^2 - 1505x - 7412 = 0.$
3.  $x^5 + \frac{625}{4}x + 3750 = 0.$
4.  $x^5 - \frac{22}{5}x^3 - \frac{11}{25}x^2 + \frac{11 \times 42}{125}x + \frac{11 \times 89}{3125} = 0.$
5.  $x^5 + 20x^3 + 20x^2 + 30x + 10 = 0.$
6.  $x^5 + 320x^2 - 1000x + 4238 = 0.$
7.  $\left(\frac{x}{10}\right)^5 + 40\left(\frac{x}{10}\right)^2 - 69\left(\frac{x}{10}\right) + 108 = 0.$
8.  $x^5 - 20x^3 + 250x - 400 = 0.$
9.  $x^5 - 5x^3 + \frac{85}{8}x - \frac{13}{2} = 0.$
10.  $x^5 + \frac{20x}{17} + \frac{21}{17} = 0.$



11.  $x^5 - \frac{4x}{13} + \frac{29}{65} = 0.$
12.  $x^5 + \frac{10x}{13} + \frac{3}{13} = 0.$
13.  $x^5 + 110(5x^3 + 60x^2 + 800x + 8320) = 0.$
14.  $x^5 - 20x^3 - 80x^2 - 150x - 656 = 0.$
15.  $x^5 - 40x^3 + 160x^2 + 1000x - 5888 = 0.$
16.  $\left(\frac{x}{2}\right)^5 - 50\left(\frac{x}{2}\right)^3 - 600\left(\frac{x}{2}\right)^2 - 2000\left(\frac{x}{2}\right) - 11200 = 0.$
17.  $x^5 + 110(5x^3 + 20x^2 - 360x + 800) = 0.$
18.  $x^5 - 20x^3 + 320x^2 + 540x + 6368 = 0.$
19.  $x^5 - 20x^3 - 160x^2 - 420x - 8928 = 0.$
20.  $x^5 - 20x^3 + 170x + 208 = 0.$

The first equation in this group was brought under the notice of the writer by a mathematical correspondent; the fourth has been treated by Lagrange; the others were formed by the writer with a view to the full illustration of his theory.

#### THE METHOD.

§2. In the article of the *Journal* above referred to, certain principles were assumed, as having been previously established, or as being known to mathematicians. It was taken for granted that the root  $x$  of the solvable irreducible quintic

$$x^5 + p_1x^4 + p_2x^3 + p_3x^2 + p_4x + p_5 = 0 \quad (1)$$

is of the form

$$\frac{1}{5}(\Delta_1^{\frac{1}{5}} + \Delta_2^{\frac{1}{5}} + \Delta_3^{\frac{1}{5}} + \Delta_4^{\frac{1}{5}}),$$

or, putting  $u_1$  for  $\frac{1}{5} \Delta_1^{\frac{1}{5}}$ ,  $u_2$  for  $\frac{1}{5} \Delta_2^{\frac{1}{5}}$ ,  $u_3$  for  $\frac{1}{5} \Delta_3^{\frac{1}{5}}$ , and  $u_4$  for  $\frac{1}{5} \Delta_4^{\frac{1}{5}}$ ,

$$u_1 + u_2 + u_3 + u_4,$$

where  $u_1, u_2, u_3$  and  $u_4$  are the roots of a quartic equation, which, when irreducible, as it is in the most general case, that includes all the others, is a uni-serial Abelian. The expressions  $u_1, u_2, u_3$  and  $u_4$  are such that

$$\left. \begin{aligned} u_1u_4 &= g + a\sqrt{z} \\ u_2u_3 &= g - a\sqrt{z} \end{aligned} \right\}; \quad (2)$$

and

$$\left. \begin{aligned} u_1^2u_3 &= k + c\sqrt{z} + (\theta + \phi\sqrt{z})\sqrt{(hz + h\sqrt{z})} \\ u_2^2u_4 &= k + c\sqrt{z} - (\theta + \phi\sqrt{z})\sqrt{(hz + h\sqrt{z})} \\ u_3^2u_1 &= k - c\sqrt{z} + (\theta - \phi\sqrt{z})\sqrt{(hz - h\sqrt{z})} \\ u_4^2u_2 &= k - c\sqrt{z} - (\theta - \phi\sqrt{z})\sqrt{(hz - h\sqrt{z})} \end{aligned} \right\} \quad (3)$$

where  $g, k, a, c, h, \theta$  and  $\phi$  are rational; and  $z = e^2 + 1$ ,  $e$  being rational. It is readily seen that

$$g = -\frac{p_2}{10}, \text{ and } k = -\frac{p_3}{20}.$$

Because  $u_1^5 = \frac{(u_1^2 u_3)^2 (u_2^3 u_4)}{(u_3 u_5)^2}$ , it follows from (2) and (3) that

$$\left. \begin{aligned} u_1^5 &= B + B'\sqrt{z} + (B'' + B'''\sqrt{z})\sqrt{(hz + h\sqrt{z})} \\ u_1^5 &= B + B'\sqrt{z} - (B'' + B'''\sqrt{z})\sqrt{(hz + h\sqrt{z})} \\ u_2^5 &= B - B'\sqrt{z} + (B'' - B'''\sqrt{z})\sqrt{(hz - h\sqrt{z})} \\ u_2^5 &= B - B'\sqrt{z} - (B'' - B'''\sqrt{z})\sqrt{(hz - h\sqrt{z})} \end{aligned} \right\} \quad (4)$$

where  $B, B', B''$  and  $B'''$  are rational functions of  $a, c, e, h, \theta$  and  $\phi$ . In like manner, because  $u_1^3 u_3 = \frac{(u_1^2 u_3)(u_2^3 u_4)}{u_2 u_3}$ , we have from (2) and (3)

$$u_1^3 u_2 = A + A'\sqrt{z} + (A'' + A'''\sqrt{z})\sqrt{(hz + h\sqrt{z})},$$

where  $A, A', A''$  and  $A'''$  are rational. The value of  $A$  is

$$A = \frac{1}{g^2 - a^2 z} \{ g(k^2 - c^2 z) + azhe(\theta^2 - \phi^2 z) \}. \quad (5)$$

From these data, the six equations, involving the six unknown quantities  $a, c, e, h, \theta$  and  $\phi$ , are (see *Journal of Mathematics* as above) obtained:

$$\left. \begin{aligned} p_4 &= -20A + 5g^2 + 15a^2 z \\ p_5 &= -4B + 40acz \\ B'' &= 1 \\ B''' &= 0 \\ hz(\theta^2 + \phi^2 z + 20\phi) &= k^2 + c^2 z - g(g^2 - a^2 z) \\ h(\theta^2 + \phi^2 z + 20\phi z) &= 2ke - a(g^2 - a^2 z) \end{aligned} \right\} \quad (6)$$

Our business is to obtain  $u_1^5, u_2^5, u_3^5$  and  $u_4^5$  from these equations.

§3. It will be found that  $a^2 z$  is the root of an equation  $F(y) = 0$ , whose coefficients are rational functions of  $p_2, p_3, p_4$  and  $p_5$ , and which, when  $p_2$  is zero, is of the sixth degree. Since  $a^2 z$  is rational, it follows that, when the coefficients of the given quintic are commensurable, the equation  $F(y) = 0$  has a commensurable root. Let this be found. Then  $a^2 z$  is known. The formulæ from which the equation  $F(y) = 0$  is obtained give us, along with  $a^2 z$ , the value of  $\frac{c}{a}$ . The remaining elements necessary for the determination of  $u_1^5, u_2^5, u_3^5$  and  $u_4^5$  may then be obtained from linear equations, without finding

$a, c, e, h, \theta$  and  $\phi$  separately. Thus a solution of the given quintic is effected. It will be pointed out how  $a, c, e, h, \theta$  and  $\phi$  can be found separately, should we desire to obtain their values.

## THE PROOF.

Case in which  $p_2$  is zero.

§4. When  $p_2 = 0$ , the investigation is much simplified. By beginning with this case, and presenting a full description of it, we shall be prepared for giving an exposition, less detailed, but still sufficiently minute to make the theory intelligible, of the case in which  $p_2$  is not assumed to be zero. When  $p_2$  is zero, equations (2) and (5) become

$$\left. \begin{aligned} u_1 u_4 &= a\sqrt{z} \\ u_2 u_3 &= -a\sqrt{z} \\ A &= -\frac{he(\theta^2 - \phi^2 z)}{a} \end{aligned} \right\} \quad (7)$$

and

Since  $B$  is the coefficient of the rational part,  $B'$  the coefficient of  $\sqrt{z}$ ,  $B''$  the coefficient of  $\sqrt{hz + h\sqrt{z}}$ , and  $B'''$  the coefficient of  $\sqrt{z}\sqrt{hz + h\sqrt{z}}$ , in the expansion of  $u_1^2$ , their values are given by the equations

$$\left. \begin{aligned} a^2 z B &= 2k(k^2 - c^2 z) - a^2 c z^2 + 2c z h e(\theta^2 - \phi^2 z) \\ a^2 z B' &= 2c(k^2 - c^2 z) + a^2 k z + 2k h e(\theta^2 - \phi^2 z) \\ a^2 z B'' &= 2(k^2 - c^2 z)\theta - \frac{2(k^2 + c^2 z)(\theta + \phi z)}{e} + \frac{z(\theta + \phi)(4kc + a^2 z)}{e} \\ a^2 z B''' &= 2(k^2 - c^2 z)\phi + \frac{2(k^2 + c^2 z)(\theta + \phi)}{e} - \frac{(\theta + \phi z)(4kc + a^2 z)}{e} \end{aligned} \right\} \quad (8)$$

The equations (6), when  $p_2$  is zero, become

$$\left. \begin{aligned} p_4 &= -20A + 15a^2 z \\ p_5 &= -4B + 40acz \\ B'' &= 1 \\ B''' &= 0 \\ h z(\theta^2 + \phi^2 z + 2\theta\phi) &= k^2 + c^2 z \\ h(\theta^2 + \phi^2 z + 2\theta\phi z) &= 2kc + a^2 z \end{aligned} \right\} \quad (9)$$

Let

$$\left. \begin{aligned} y &= a^2 z \\ t &= \frac{c}{a} \end{aligned} \right\}$$

and

(10)

Substituting in the first of equations (8) the value of  $B$  obtained from the second of equations (9), and the value of  $he(\theta^3 - \phi^3 z)$  obtained from (7), and making use of (10),

$$p_5 y = -8k(k^3 - t^2 y) + 44ty^2 + 8tyA.$$

Therefore, by the first of equations (9),

$$8kyt^2 + \left(50y^3 - \frac{2}{5} p_4 y\right)t = 8k^3 + p_5 y. \quad (11)$$

§5. Again, from the last two of equations (9), because  $z = t^2 + 1$ ,

$$he^3(\theta^3 + \phi^3 z) = (k^3 + t^2 y) - (2kt + y).$$

Therefore, by (10),

$$he^3(\theta^3 + \phi^3 z) = (k^3 + t^2 y) - a(2kt + y). \quad (12)$$

Similarly, from the last two of equations (9),

$$2hze^2\theta\phi = az(2kt + y) - (k^3 + t^2 y). \quad (13)$$

And

$$(\theta^3 - \phi^3 z)^3 = (\theta^3 + \phi^3 z)^3 - 4z(\theta^3 \phi^3).$$

Therefore, from (12) and (13),

$$\begin{aligned} (he^3)^3(\theta^3 - \phi^3 z)^3 &= \{(k^3 + t^2 y) - a(2kt + y)\}^3 - \frac{1}{z} \{az(2kt + y) - (k^3 + t^2 y)\}^3 \\ \therefore z h^3 e^3(\theta^3 - \phi^3 z)^3 &= (k^3 + t^2 y)^3 - y(2kt + y)^3. \end{aligned}$$

Hence from the value of  $he(\theta^3 - \phi^3 z)$  in (7),

$$(k^3 + t^2 y)^3 - y(2kt + y)^3 = za^3 A^3 = yA^3.$$

But, by the first of equations (9),

$$A = \frac{3y}{4} - \frac{p_4}{20}. \quad (14)$$

Therefore,

$$(k^3 + t^2 y)^3 - y(2kt + y)^3 = y\left(\frac{3y}{4} - \frac{p_4}{20}\right)^3. \quad (15)$$

And, from (11),

$$8k(k^3 + t^2 y) = (16k^3 + p_5 y) - t\left(50y^3 - \frac{2}{5} p_4 y\right).$$

Substitute in (15) the value of  $k^3 + t^2 y$  here given. The result is

$$\begin{aligned} t^3 \left\{ \left(50y^3 - \frac{2}{5} p_4 y\right)^3 - 256k^4 y \right\} + t \left\{ -2(16k^3 + p_5 y)\left(50y^3 - \frac{2}{5} p_4 y\right) - 256k^3 y^3 \right\} \\ = 64k^2 y \left(\frac{3y}{4} - \frac{p_4}{20}\right)^3 + 64k^2 y^3 - (16k^3 + p_5 y)^3. \end{aligned} \quad (16)$$

§6. In (11) and (16) we have two equations, with the two unknown quantities  $y$  and  $t$ . The equations may be written

$$\begin{cases} 8kyt^2 + myt = n \\ vyt^2 + qyt = r \end{cases} \quad (17)$$

and

where

$$m = 50y - \frac{2}{5}p_1,$$

$$n = 8k^3 + p_5y,$$

$$v = y\left(50y - \frac{2}{5}p_1\right)^2 - 256k^4$$

$$= 2500y^3 - 40p_1y^2 + \frac{4}{25}p_1^2y - 256k^4,$$

$$q = -2(16k^3 + p_5y)\left(50y - \frac{2}{5}p_1\right) - 256k^3y$$

$$= -100p_5y^2 + \left(\frac{4}{5}p_1p_5 - 1856k^3\right)y + \frac{64}{5}k^3p_1,$$

$$r = 100k^3y^3 - \left(\frac{24p_1k^3}{5} + p_5^2\right)y^2 + \left(\frac{4}{25}p_1^2k^3 - 32k^3p_5\right)y - 256k^4.$$

§7. The elimination of  $t$  from the equations (17) gives us

$$(vn - 8kr)^2 + y(qn - mr)(vm - 8kq) = 0.$$

From the values of  $m$ ,  $n$ ,  $v$ ,  $q$  and  $r$  in §6,

$$vn = 2500p_5y^4 + (20000k^3 - 40p_1p_5)y^3 + \left(\frac{4}{25}p_1^2p_5 - 320p_1k^3\right)y^2 \quad (18)$$

$$+ \left(\frac{32}{25}p_1^2k^3 - 256k^4p_5\right)y - 2048k^4;$$

$$8kr = 800k^3y^3 - \left(\frac{192}{5}p_1k^3 + 8p_5^2k\right)y^2 + \left(\frac{32}{25}p_1^2k^3 - 256k^4p_5\right)y - 2048k^4;$$

$$qn = -100p_5^2y^3 + \left(\frac{4}{5}p_1p_5^2 - 2656k^3p_5\right)y^2 + \left(\frac{96}{5}p_5k^3p_1 - 14848k^3\right)y + 512k^3p_1;$$

$$mr = 5000k^3y^4 - (280p_1k^3 + 50p_5^2)y^3 + \left(\frac{248}{25}p_1^2k^3 - 1600k^3p_5 + \frac{2}{5}p_1p_5^2\right)y^2$$

$$- \left(12800k^4 + \frac{8}{125}p_1^3k^3 - \frac{64}{5}k^3p_1p_5\right)y + \frac{512}{5}k^4p_1;$$

$$vm = 125000y^4 - 3000p_1y^3 + 24p_1^2y^2 - \left(12800k^4 + \frac{8}{125}p_1^3\right)y + \frac{512}{5}p_1k^4;$$

$$8kq = -800p_5ky^3 + \left(\frac{32}{5}kp_1p_5 - 14848k^4\right)y + \frac{512}{5}p_1k^4.$$

$$\therefore vn - 8kr = y^2 \left\{ 2500p_5y^2 + (19200k^3 - 40p_1p_5)y + \left(\frac{4}{25}p_1^2p_5 - \frac{1048}{5}p_1k^3 + 8p_5^2k\right) \right\},$$

$$\begin{aligned}
 (17) \quad qn - mr &= y \left\{ -5000k^3y^3 + (-50p_5^3 + 280p_4k^3)y^3 \right. \\
 &\quad + \left( -\frac{2}{5}p_4p_3^3 - 1056k^3p_5 - \frac{248}{25}p_4^3k^3 \right) y \\
 &\quad \left. + k^3 \left( -2048k^4 + \frac{8}{125}p_4^3 + \frac{32}{5}kp_4p_5 \right) \right\}; \\
 rm - 8kq &= y \left\{ 125000y^3 - 3000p_4y^3 + (24p_4^3 + 800kp_5)y \right. \\
 &\quad \left. - \left( -2048k^4 + \frac{8}{125}p_4^3 + \frac{32}{5}kp_4p_5 \right) \right\}.
 \end{aligned}$$

By substituting in (18) these values of  $rn - 8kr$ ,  $qn - mr$ ,  $rm - 8kq$ , we get

$$F(y) = q_0y^6 + q_1y^5 + \dots + q_5y + q_6 = 0, \quad (19)$$

where

$$\begin{aligned}
 (18) \quad q_0 &= -625000000, \\
 q_1 &= 50000000p_4, \\
 q_2 &= -200000(200kp_5 + 11p_4^3), \\
 q_3 &= 102400000k^4 + 1280000kp_4p_5 + 44800p_4^3, \\
 q_4 &= -(25600kp_4^3p_5 + 4096000k^4p_4 + 640000p_5^3k^3 + 448p_4^4), \\
 q_5 &= 8192k^3p_4p_5^3 - 2048 \times 1856k^5p_5 + 64k^3 + \frac{49152}{5}k^4p_4^3 + \frac{6144}{25}kp_4^3p_5 + \frac{6784}{3125}p_4^5, \\
 q_6 &= -\left( \frac{8}{125}p_4^3 - 2048k^4 + \frac{32}{5}kp_4p_5 \right)^2.
 \end{aligned}$$

§8. Assuming now that the coefficients  $p_3$ ,  $p_4$ ,  $p_5$  are commensurable quantities, let the commensurable root  $y$  of equation (19) be found. Then  $a^2z$  is known. Then, from (11) and (16),  $t$  or  $\frac{c}{a}$  is found.

§9. At this stage, as was indicated in §3, two courses are open to us. One is to proceed to find  $u_1^5$ ,  $u_2^5$ ,  $u_3^5$ ,  $u_4^5$  without troubling ourselves to inquire what  $a$ ,  $c$ ,  $\theta$ ,  $\phi$  and  $h$  are separately. This, the natural and the shortest course, we will now follow. Since  $a^2z$  and  $\frac{c}{a}$  are known, their product  $acz$  is known. And, by (14),  $A$  is known. Therefore  $czhe(\theta^2 - \phi^2z)$ , which, by the last of equations (7), is equal to  $-aczA$ , is known. Hence by the first of equations (8),  $B$  is known. We might even more simply,  $acz$  being known, find  $B$  from the second of equations (9). The second of equations (8) gives us

$$y(B\sqrt{z}) = 2c\sqrt{z}(k^2 - c^2z) + a^2zk(a\sqrt{z}) - 2kAa\sqrt{z}. \quad (20)$$

Now  $acz$  is known, and it is the same as  $(a\sqrt{z})(c\sqrt{z})$ ; consequently, the signs

with which  $c\sqrt{z}$  and  $a\sqrt{z}$  must be taken relatively to one another are known. Therefore  $B'\sqrt{z}$  is known. Therefore  $B + B'\sqrt{z}$  is known. But, by the manner in which  $B, B', B''$  and  $B'''$  were taken, keeping in view the values of  $B'$  and  $B'''$  in (6),

$$u_1^5 = B + B'\sqrt{z} + \sqrt{(hz + h\sqrt{z})},$$

and

$$u_4^5 = B + B'\sqrt{z} - \sqrt{(hz + h\sqrt{z})}.$$

$$\therefore (u_1 u_4)^5 = (B + B'\sqrt{z})^2 - (hz + h\sqrt{z}).$$

And, by (7),  $u_1 u_4 = a\sqrt{z}$ . Therefore  $hz + h\sqrt{z} = (B + B'\sqrt{z})^2 - (a\sqrt{z})^5$ . This gives us

$$u_1^5 = B + B'\sqrt{z} + \sqrt{\{(B + B'\sqrt{z})^2 - (a\sqrt{z})^5\}},$$

$$u_4^5 = B + B'\sqrt{z} - \sqrt{\{(B + B'\sqrt{z})^2 - (a\sqrt{z})^5\}},$$

$$u_2^5 = B - B'\sqrt{z} + \sqrt{\{(B - B'\sqrt{z})^2 + (a\sqrt{z})^5\}},$$

$$u_3^5 = B - B'\sqrt{z} - \sqrt{\{(B - B'\sqrt{z})^2 + (a\sqrt{z})^5\}}.$$

Hence  $u_1 + u_4 + u_2 + u_3$ , the root of the given quintic, is known.

To find  $a, c, e, \theta, \phi$  and  $h$  separately.

§10. If we desire to obtain the values of  $a, c, e, \theta, \phi$  and  $h$  separately, we may first find  $a$  by means of a quadratic equation. By (12) and the third of equations (7),

$$hc^2(\theta^2 + \phi^2 z) = (k^3 + l^2 y) - a(2kt + y)$$

and

$$hc^2(\theta^2 - \phi^2 z) = -aeA.$$

Therefore,

$$2hc^2\theta^2 = (k^3 + l^2 y) - a(2kt + y) - aeA.$$

Also, by (13),

$$2hze^2\theta\phi = az(2kt + y) - (k^3 + l^2 y).$$

Therefore,

$$\frac{\theta}{\phi z} = \frac{(k^3 + l^2 y) - a(2kt + y) - aeA}{az(2kt + y) - (k^3 + l^2 y)}. \quad (21)$$

But, by (9),  $B''' = 0$ . Therefore, from the last of equations (8),

$$\begin{aligned} \frac{\theta}{\phi z} &= \frac{2e(k^3 - c^2 z) + 2(k^3 + c^2 z) - z(4kc + a^2 z)}{z\{(4kc + a^2 z) - 2(k^3 + c^2 z)\}} \\ &= \frac{2e(k^3 - l^2 y) + 2(k^3 + l^2 y) - az(4kt + y)}{az(4kt + y) - 2z(k^3 + l^2 y)}. \end{aligned} \quad (22)$$

From (21) and (22),

$$\frac{(k^3 + l^2 y) - a(2kt + y) - aeA}{az(2kt + y) - (k^3 + l^2 y)} = \frac{2e(k^3 - l^2 y) + 2(k^3 + l^2 y) - az(4kt + y)}{az(4kt + y) - 2z(k^3 + l^2 y)},$$

$$\therefore \frac{(k^3 + l^2 y) - a(2kt + y) - aeA}{a^2 z(2kt + y) - a(k^3 + l^2 y)} = \frac{2ae(k^3 - l^2 y) + 2a(k^3 + l^2 y) - a^2 z(4kt + y)}{a(a^2 z(4kt + y) - 2a^2 z(k^3 + l^2 y))},$$

$$\text{or, } \frac{(k^3 + l^2 y) - a(2kt + y) - aeA}{y(2kt + y) - a(k^3 + l^2 y)} = \frac{2ae(k^3 - l^2 y) + 2a(k^3 + l^2 y) - y(4kt + y)}{ay(4kt + y) - 2y(k^3 + l^2 y)}.$$

Put

$$\begin{aligned}\beta &= (k^2 + l^2 y) - a(2kt + y), \\ \gamma &= y(2kt + y) - a(k^2 + l^2 y), \\ \delta &= ay(4kt + y) - 2y(k^2 + l^2 y), \\ \sigma &= 2a(k^2 + l^2 y) - y(4kt + y), \\ \tau &= k^2 + l^2 y.\end{aligned}$$

Then

$$\frac{\beta - acA}{\tau} = \frac{2a\tau + \sigma}{\delta},$$

$$\therefore ac(\delta A + 2\gamma\tau) = \beta\delta - \gamma\sigma. \quad (23)$$

But  $\beta\delta = -a^2 y(2kt + y)(4kt + y) + ay(k^2 + l^2 y)(8kt + 3y) - 2y(k^2 + l^2 y)^2$   
 and  $\gamma\sigma = -y^2(2kt + y)(4kt + y) + ay(k^2 + l^2 y)(8kt + 3y) - 2a^2(k^2 + l^2 y)^2$ ,  
 $\therefore \beta\delta - \gamma\sigma = (y - a^2)\{y(2kt + y)(4kt + y) - 2(k^2 + l^2 y)^2\}.$

And  $y - a^2 = a^2 z - a^2 = a^2(z - 1) = a^2 e^2$ . Therefore

$$\beta\delta - \gamma\sigma = a^2 e^2 \{y(2kt + y)(4kt + y) - 2(k^2 + l^2 y)^2\}.$$

Therefore, from (23),

$$\delta A + 2\gamma\tau = ae \{y(2kt + y)(4kt + y) - 2(k^2 + l^2 y)^2\}, \quad (24)$$

$$\begin{aligned}\therefore (\delta A + 2\gamma\tau)^2 &= a^2 e^2 \{y(2kt + y)(4kt + y) - 2(k^2 + l^2 y)^2\}^2 \\ &= (y - a^2)^2 \{y(2kt + y)(4kt + y) - 2(k^2 + l^2 y)^2\}^2.\end{aligned} \quad (25)$$

Now  $k$ ,  $y$  and  $l$  are known. And, by (14),  $A$  is known. Therefore (25) is a quadratic equation from which  $a$  can be found. The quadratic has its roots commensurable, and care must be taken in each case to select that one which satisfies all the conditions of the problem. When  $a$  has been found, since  $a^2 z$  and  $\frac{c}{a}$  are known,  $z$  and  $c$  are known; and, because  $z = e^2 + 1$ , the absolute value of  $e$  is known. The sign with which  $e$  is to be taken is determined by (24). Next, to find  $\theta$ ,  $\phi$  and  $h$ , the third and fourth of equations (9) are  $B'' = 1$ ,  $B''' = 0$ . Hence, taking the values of  $B''$  and  $B'''$  in (8),

$$\begin{aligned}a^2 zc &= 2e(k^2 - c^2 z)\theta - 2(k^2 + c^2 z)(\theta + \phi z) + 2(\theta + \phi)(4kc + a^2 z) \\ 0 &= 2e(k^2 - c^2 z)\phi + 2(k^2 + c^2 z)(\theta + \phi) - (\theta + \phi z)(4kc + a^2 z)\end{aligned} \quad (26)$$

But  $e$ ,  $z$ ,  $a$ ,  $c$ ,  $k$  are known. Therefore, from the simultaneous equations (26),  $\theta$  and  $\phi$  are known. Therefore, from (7),  $h$  is known.



Another way of finding  $a, c, e, \theta, \phi$  and  $h$ .

§11. The values of  $a, c, e, \theta, \phi$  and  $h$  may be arrived at in another way. Let  $B$  and  $B'\sqrt{z}$  be found as in §9. Then, because (see §9)

$$hz + h\sqrt{z} = (B + B'\sqrt{z})^2 - (a\sqrt{z})^2,$$

we have

$$hz = B^2 + (B'\sqrt{z})^2,$$

and

$$h\sqrt{z} = 2B(B'\sqrt{z}) - (a\sqrt{z})^2.$$

Here the quantities to which  $hz$  and  $h\sqrt{z}$  are equated are known. Therefore  $h$  and  $z$  are known. When  $z$  is known, because  $a^2z$  and  $\frac{e}{a}$  are known,  $a$  and  $c$  are known. Finally,  $\theta$  and  $\phi$  are obtained, as in §10, from the equations (26).

§12. *First Example.*—To exemplify the theory, let us take the equation

$$x^5 + 3x^2 + 2x - 1 = 0.$$

Because  $k = -\frac{p_3}{20} = -\frac{3}{20}$ ,  $p_4 = 2$ ,  $p_5 = -1$ , the equations (11) and (16)

become

$$\left. \begin{aligned} \frac{6t^2y}{5} + t\left(\frac{4y}{5} - 50y^2\right) &= y + \frac{27}{1000}, \\ t^2\left(\frac{125^2}{9}y^4 - \frac{500y^3}{9} + \frac{4y^2}{9} - \frac{9y}{100}\right) + t\left(\frac{625y^3}{9} + \frac{583}{180}y^2 - \frac{3y}{50}\right) \\ &= \frac{25y^3}{16} - \frac{38y^2}{45} - \frac{13y}{200} - \frac{81}{40000} \end{aligned} \right\} \quad (28)$$

The equation (19), obtained by eliminating  $t$  from the equations (28), is

$$F(y) = \left(\frac{25 \times 125^4}{36}\right)y^6 - \left(\frac{125^4}{9}\right)y^5 + \left(\frac{37 \times 125^3}{18}\right)y^4 - \left(\frac{1241 \times 125^3}{90}\right)y^3 \\ + \left(\frac{3209 \times 125}{36}\right)y^2 - \left(\frac{5159}{136}\right)y + \frac{109^2}{22500} = 0,$$

and this has the commensurable root  $\frac{1}{125}$ . Therefore

$$y = a^2z = \frac{1}{125}.$$

Hence, from the two equations (28),

$$t = \frac{e}{a} = \frac{7}{4}.$$

In subsequent examples, when  $y$  and  $t$  have been found, we shall proceed at once to find  $B$  and  $B'\sqrt{z}$ , as in §9, without inquiring what  $a, c, e, \theta, \phi$  and  $h$  are separately. But we desire to illustrate, in one instance, the method of

obtaining the values of these quantities which was described in §10, and we will use the present instance for that purpose. To find  $a$  we take the equation (25),

$$(\delta A + 2\gamma\tau)^2 = (y - a^2)\{y(2kt + y)(4kt + y) - 2(k^2 + t^2y)^2\}^2. \quad (29)$$

By (14),

$$A = \frac{3y}{4} - \frac{p_4}{20} = -\frac{47}{500}.$$

Also, from the values of  $k$ ,  $y$  and  $t$  above given,

$$2kt + y = -\frac{517}{1000},$$

$$4kt + y = -\frac{521}{1000},$$

$$k^2 + t^2y = \frac{47}{1000}.$$

Therefore

$$\delta = ay(4kt + y) - 2y(k^2 + t^2y) = -\frac{521a + 47}{62500},$$

$$\gamma = y(2kt + y) - a(k^2 + t^2y) = \frac{-47(125a + 11)}{125000},$$

$$\tau = k^2 - t^2y = -\frac{1}{500}.$$

Hence (29) becomes

$$125(323a + 29)^2 = 36^2(1 - 125a^2). \quad (30)$$

One root of this equation is  $-\frac{11}{125}$ . But this root proves on examination to be inadmissible. We must therefore take the other root, which is  $-\frac{9439}{25^2 \times 13^2}$ .

Then, since  $c = \frac{7a}{4}$ , and  $a^2z = \frac{1}{125}$ , we have

$$a = -\frac{9439}{25^2 \times 13^2} = -\frac{9439}{25 \times 4225},$$

$$c = -\frac{7 \times 9439}{422500},$$

$$z = \frac{5 \times 4225^2}{9439^2},$$

$$e = \frac{398}{9439}.$$

The sign of  $e$  is determined in the way pointed out in §10. By means of the values of  $e$ ,  $z$ ,  $a$ ,  $c$  that have been obtained, we get, from the two equations (26),

$$\theta = \frac{125}{199}, \phi = -\frac{18 \times 9439}{199 \times 845} \therefore \theta^2 - \phi^2z = -\frac{125}{199}.$$

Therefore, from (7), keeping in view that  $A = -\frac{47}{500}$ ,

$$h = \frac{47}{8} \left( \frac{9439}{13 \times 25 \times 125} \right)^2.$$

Also, from the first two of equations (8),  $B = \frac{39}{100}$ , and  $B' = \frac{91 \times 9439}{500 \times 4225}$

$$\therefore B'\sqrt{z} = \frac{91\sqrt{5}}{500},$$

and

$$hz + h\sqrt{z} = \frac{1}{625} \left\{ \frac{47}{8} (21125 + 9439\sqrt{5}) \right\}$$

Therefore,

$$\left. \begin{aligned} u_1^5 &= \frac{13}{100} \left( 3 + \frac{7\sqrt{5}}{5} \right) + \frac{1}{625} \sqrt{\left\{ \frac{47}{8} (21125 + 9439\sqrt{5}) \right\}} \\ u_4^5 &= \frac{13}{100} \left( 3 + \frac{7\sqrt{5}}{5} \right) - \frac{1}{625} \sqrt{\left\{ \frac{47}{8} (21125 + 9439\sqrt{5}) \right\}} \\ v_2^5 &= \frac{13}{100} \left( 3 - \frac{7\sqrt{5}}{5} \right) + \frac{1}{625} \sqrt{\left\{ \frac{47}{8} (21125 - 9439\sqrt{5}) \right\}} \\ u_3^5 &= \frac{13}{100} \left( 3 - \frac{7\sqrt{5}}{5} \right) - \frac{1}{625} \sqrt{\left\{ \frac{47}{8} (21125 - 9439\sqrt{5}) \right\}} \end{aligned} \right\} \quad (31)$$

Therefore the root of the given quintic is known.

§13. To verify this result, we have  $9439\sqrt{5} = 21106.2456$ . Therefore

$$\frac{1}{625} \sqrt{\left\{ \frac{47}{8} (21225 + 9439\sqrt{5}) \right\}} = .79696796,$$

$$\frac{1}{625} \sqrt{\left\{ \frac{47}{8} (21225 - 9439\sqrt{5}) \right\}} = .01679462.$$

Also,

$$\frac{13}{100} \left( 3 + \frac{7\sqrt{5}}{5} \right) = .79696437$$

and

$$-\frac{13}{100} \left( 3 - \frac{7\sqrt{5}}{5} \right) = .01696437.$$

Therefore

$$\begin{aligned} u_1^5 &= 1.593932, & u_1 &= 1.09773 \\ -u_4^5 &= .00000359, & -u_4 &= .08147 \\ -u_2^5 &= .00016975, & -u_2 &= .17618 \\ -u_3^5 &= .03375899, & -u_3 &= .50778. \end{aligned}$$

Therefore  $u_1 + u_2 + u_3 + u_4 = .3323$ . Therefore

$$x^5 = .004052$$

$$3x^3 = .331248$$

$$2x = .66448$$

$$.99978$$

§14. If we wish to exhibit the root as in §9, we find from (8) that

$$B + B'\sqrt{z} = \frac{13}{100} \left( 3 + \frac{7\sqrt{5}}{5} \right).$$

Therefore, since  $a^5 z^2 \sqrt{z} = -\frac{\sqrt{5}}{625}$ ,

$$u_1^5 = \frac{13}{100} \left( 3 + \frac{7\sqrt{5}}{5} \right) + \sqrt{\left\{ \left( \frac{13}{100} \right)^2 \left( 3 + \frac{7\sqrt{5}}{5} \right)^2 + \frac{\sqrt{5}}{625^2} \right\}},$$

with corresponding expressions for  $u_2^5, u_3^5, u_4^5$ . As a matter of fact,

$$\left( \frac{13}{100} \right)^2 \left( 3 + \frac{7\sqrt{5}}{5} \right) + \frac{\sqrt{5}}{625^2} = \frac{1}{625^2} \left\{ \frac{47}{8} (21125 + 9439\sqrt{5}) \right\}.$$

§15. It is interesting to observe the application of the theory to the equation

$$x^5 - 3x^2 + 2x + 1 = 0, \quad (32)$$

whose roots, with the signs changed, are the same as the roots of the equation (27). By reference to §7, keeping in view that  $k = -\frac{p_3}{20}$ , it will be seen that, wherever an odd power of  $p_3$  occurs as a factor in a term of any one of the coefficients of the equation  $F(y) = 0$ , an odd power of  $p_5$  occurs as a factor of the same term. It follows that, by changing the signs of both  $p_3$  and  $p_5$  in the equation (27), in other words, by passing from the equation (27) to the equation (32),  $F(y)$  remains unchanged. Therefore the commensurable root of this equation, which we have seen to be  $\frac{1}{125}$ , is the value of  $a^2 z$  for the equation (32)

as well as for the equation (27). To find  $t$  or  $\frac{c}{a}$ , the equations (17) give us

$$yt = \frac{vn - 8kr}{vm - 8kq}.$$

The values of  $vn - 8kr$  and  $vm - 8kq$  given in §7 show that, in passing from (27) to (32),  $vn - kr$  simply changes its sign, while  $vm - kq$  remains unaltered.

Hence  $t$  or  $\frac{c}{a}$  has the same absolute value for the equation (32) as for the equation (27), the signs, however, being different in the two cases. Consequently, for the equation (27),  $\frac{c}{a} = -\frac{7}{4}$ . Thus we get

$$a = -\frac{9439}{25 \times 4225}, \quad c = \frac{7 \times 9439}{422500}, \quad z = \frac{5 \times 4225^2}{9439^2}, \quad e = -\frac{398}{9439}.$$

Hence, by the first two of equations (8),

$$B + B'\sqrt{z} = -\frac{13}{100}\left(3 + \frac{7\sqrt{5}}{5}\right).$$

Therefore, for the equation (32), the values of  $u_1^5$ ,  $u_4^5$ ,  $u_2^5$  and  $u_3^5$  are

$$\begin{aligned} u_1^5 &= -\frac{13}{100}\left(3 + \frac{7\sqrt{5}}{5}\right) - \sqrt{\left\{\frac{13^3}{100^3}\left(3 + \frac{7\sqrt{5}}{5}\right)^2 + \frac{\sqrt{5}}{625^2}\right\}}, \\ u_4^5 &= -\frac{13}{100}\left(3 + \frac{7\sqrt{5}}{5}\right) + \sqrt{\left\{\frac{13^3}{100^3}\left(3 + \frac{7\sqrt{5}}{5}\right)^2 + \frac{\sqrt{5}}{625^2}\right\}}, \\ u_2^5 &= -\frac{13}{100}\left(3 - \frac{7\sqrt{5}}{5}\right) - \sqrt{\left\{\frac{13^3}{100^3}\left(3 - \frac{7\sqrt{5}}{5}\right)^2 - \frac{\sqrt{5}}{625^2}\right\}}, \\ u_3^5 &= -\frac{13}{100}\left(3 - \frac{7\sqrt{5}}{5}\right) + \sqrt{\left\{\frac{13^3}{100^3}\left(3 - \frac{7\sqrt{5}}{5}\right)^2 - \frac{\sqrt{5}}{625^2}\right\}}. \end{aligned}$$

$p_2$  not assumed to be zero.

§17. Let us now consider the more general case in which  $p_2$  is not assumed to be zero. The method that has been illustrated above is still applicable, though the labor of operation, in dealing with particular instances, is increased.

Putting, as before,  $y$  for  $\alpha^2 z$ , and  $t$  for  $\frac{c}{a}$ , we form equations corresponding to (11) and (16); from these we obtain the values of  $y$  and  $t$ ; then we find  $B$  and  $B'\sqrt{z}$  from equations corresponding to the first two of the group (8); or,  $B$  can be more readily found from the second of equations (6), and, on the principles of §9, when  $B + B'\sqrt{z}$  is known,  $u_1 u_4$  or  $g^2 - y$  being also known, the root of the given quintic is known.

§18. The values of  $B$  and  $B'\sqrt{z}$  which correspond, when  $g$  or  $-\frac{p_2}{10}$  is not zero, to those given in (8) for the case in which  $g$  is zero, are obtained from the equations (2) and (3) by keeping in view that, according to the first of equations (4),  $B$  is the rational part and  $B'$  the coefficient of  $\sqrt{z}$  in the expansion of  $u_1^5$ .

Put

$$\begin{aligned} A' &= ch(\theta^2 - \phi^2 z) \\ P &= 2k(k^2 - t^2 y) - (g^2 - y)(gk - ty) \\ Q &= 2t(k^2 - t^2 y) - (g^2 - y)(k - gt) \end{aligned} \quad (33)$$

Then

$$(g^2 - y)^2 B = (g^2 + y)P + 2gyQ + 2azA'\{t(g^2 + y) + 2gk\}$$

and

$$(g^2 - y)^2 B' = a\{2gP + (g^2 + y)Q\} + 2A'\{k(g^2 + y) + 2gt y\} \quad (34)$$

By (5) and (6)

$$A(g^2 - y) = g(k^2 - t^2y) + azA',$$

and

$$p_4 = -20A + 5g^2 + 15y.$$

Therefore

$$azA' = \frac{g^2 - y}{20} (5g^2 + 15y - p_4) - g(k^2 - t^2y). \quad (35)$$

§19. From the second of equations (6),

$$p_5(g^2 - y)^2 + 4B(g^2 - y)^2 - 40ty(g^2 - y)^2 = 0.$$

Therefore, from the first of the two equations (34),

$$(p_5 - 40ty)(g^2 - y)^2 + 4(g^2 + y)P + 8gyQ + 8azA'\{t(g^2 + y) + 2gk\} = 0.$$

Putting for  $azA'$  its value in (35),

$$\begin{aligned} (g^2 - y) \left[ (p_5 - 40ty)(g^2 - y) + 2 \left( g^2 + 3y - \frac{1}{5}p_4 \right) \{t(g^2 + y) + 2gk\} \right] \\ + [4(g^2 + y)P + 8gazQ - 8g(k^2 - t^2y)\{t(g^2 + y) + 2gk\}] = 0. \end{aligned} \quad (36)$$

But, from the values of  $P$  and  $Q$ ,

$$\begin{aligned} 4(g^2 + y)P + 8gazQ - 8g(k^2 - t^2y)\{t(g^2 + y) + 2gk\} \\ = \{ -4(g^2 + y)(gk - ty) + 8gy(k - gt) - 8(k + gt)(k^2 - t^2y)\}(g^2 - y). \end{aligned}$$

Therefore, rejecting the common factor  $g^2 - y$ , (36) becomes

$$\begin{aligned} (p_5 - 40ty)(g^2 - y) + 2 \left( g^2 + 3y - \frac{1}{5}p_4 \right) \{t(g^2 + y) + 2gk\} \\ - 4(g^2 + y)(gk - ty) + 8gy(k - gt) - 8(k + gt)(k^2 - t^2y) = 0. \end{aligned}$$

Arranging according to the powers of  $y$ ,

$$\left. \begin{aligned} 50gt^2t + y \left\{ 8gt^2 + 8kt^2 - t \left( 20g^2 + \frac{2}{5}p_4 \right) - p_5 \right\} \\ + t \left\{ 2g^3 \left( g^2 - \frac{p_4}{5} \right) - 8gk^2 \right\} + g^2p_5 - 8k^3 - \frac{4}{5}gkp_4 = 0. \end{aligned} \right\} \quad (37)$$

This is one equation between the unknown quantities  $y$  and  $t$ .

§20. From the last two of the equations (6),

$$he^2(\theta^2 + \phi^2z) = (k^2 + t^2y) - 2kat + (g^2 - y)(a - g)$$

and

$$2kze^2(\theta\phi) = 2kzat - (k^2 + y^2t) - (g^2 - y)(az - g).$$

But

$$(\theta^2 - \phi^2z)^2 = (\theta^2 + \phi^2z)^2 - 4z\theta^2\phi^2.$$

Therefore

$$\begin{aligned} zhe^2e^4(\theta^2 - \phi^2z)^2 = z \{ (k^2 + t^2y) - 2kat + (g^2 - y)(a - g)^2 \} \\ - \{ 2kzat - (k^2 + t^2y) - (g^2 - y)(az - g)^2 \}. \end{aligned}$$

Therefore

$$z \{ h e (\theta^2 - \phi^2 z) \}^2 = (k^2 - t^2 y)^2 + (g^2 - y)^2 + (g^2 - y) \{ 4kty - 2g(k^2 + t^2 y) \}.$$

But  $A' = h e (\theta^2 - \phi^2 z)$ . Therefore, by (36),

$$\frac{1}{y} \left\{ \frac{g^2 - y}{20} (5g^2 + 15y - p_4) - g(k^2 - t^2 y) \right\}^2 = (k^2 - t^2 y)^2 + (g^2 - y)^2 + (g^2 - y) \{ 4kty - 2g(k^2 + t^2 y) \};$$

or, arranging according to the powers of  $y$ ,

$$\begin{aligned} & 25y^3 - y^2 \left( 16t^4 + 56gt^2 - 64kt + \frac{6}{5} p_4 + 35g^2 \right) \\ & - y \left\{ 8gt^2 \left( g^2 - \frac{1}{5} p_4 \right) - 32k^2 t^2 + \left( g^2 - \frac{1}{5} p_4 \right) \left( 5g^2 + \frac{1}{5} p_4 \right) + 8gk^2 - 16gt^4 \right\} \\ & - g^2 \left( g^2 - \frac{1}{5} p_4 \right)^2 + 8gk^2 \left( g^2 - \frac{1}{5} p_4 \right) - 16k^4 = 0. \end{aligned} \quad (38)$$

This is the second equation between the unknown quantities  $y$  and  $t$ .

§21. We may now either eliminate  $t$  from the two equations (37) and (38) so as to obtain an equation

$$F(y) = 0$$

whose coefficients are rational functions of the coefficients of the quintic to be solved, or we may eliminate  $y$  so as to obtain an equation

$$\psi(t) = 0$$

whose coefficients are rational functions of those of the quintic to be solved. In the former case, let the commensurable root  $y$  of the equation  $F(y) = 0$  be found. Then, by (37) and (38),  $t$  is known. In the latter case, let the commensurable root  $t$  of the equation  $\psi(t) = 0$  be found. Then, by (37) and (38),  $y$  is known. When  $y$  and  $t$  have thus been found, we find  $B$  and  $B'\sqrt{z}$ , exactly as in §12, from the equations (34), or  $B$  can more readily be found from the second of equations (6). Then

$$\begin{aligned} u_1^5 &= B + B'\sqrt{z} + \sqrt{\{(B + B'\sqrt{z})^2 - (u_1 u_4)\}} \\ &= B + B'\sqrt{z} + \sqrt{\{(B + B'\sqrt{z})^2 - (g + a\sqrt{z})^5\}}. \end{aligned}$$

Therefore  $x = u_1 + u_4 + u_5 + u_3$

$$\begin{aligned} &= [B + B'\sqrt{z} + \sqrt{\{(B + B'\sqrt{z})^2 - (g + a\sqrt{z})^5\}}]^{1/5} \\ &+ [B + B'\sqrt{z} - \sqrt{\{(B + B'\sqrt{z})^2 - (g + a\sqrt{z})^5\}}]^{1/5} \\ &+ [B - B'\sqrt{z} + \sqrt{\{(B - B'\sqrt{z})^2 - (g - a\sqrt{z})^5\}}]^{1/5} \\ &+ [B - B'\sqrt{z} - \sqrt{\{(B - B'\sqrt{z})^2 - (g - a\sqrt{z})^5\}}]^{1/5}. \end{aligned}$$

It need scarcely be pointed out that since  $y = a^2 z$ ,  $a\sqrt{z}$  is known.

§22. *Second Example.*—As an illustrative example, let

$$x^5 - 10x^3 - 20x^2 - 1505x - 7412 = 0.$$

Here  $g = -\frac{p_2}{10} = 1$ , and  $k = -\frac{p_3}{20} = 1$ . Therefore the equations (37) and (38) become

$$50y^2t + y(8t^3 + 8t^2 + 528t + 7412) + 596t - 6216 = 0,$$

and

$$25y^3 - (16t^4 + 56t^3 - 64t - 1771)y^2 - (2384t^3 - 89384)y - 88804 = 0.$$

The commensurable values of  $y$  and  $t$  which satisfy these equations are  $y = 2$ ,  $t = -4$ . Then, we can get the values of  $B$  and  $B'\sqrt{z}$  from the equations (34), keeping in view that  $azA'$  is known by (35). As far as  $B$  is concerned, it is simpler to make use of the second of equations (6), keeping in view that  $acz = ty = -8$ . Therefore

$$4B = 7412 - 320,$$

$$\therefore B = 1773.$$

In order to obtain the value of  $B'\sqrt{z}$ , we must find  $P$ ,  $Q$  and  $azA'$ . By (33) and (35),

$$P = -62 + 9 = -53,$$

$$Q = 248 + 5 = 253,$$

and

$$azA' = -77 + 31 = -46.$$

Therefore

$$B'\sqrt{z} = 653a\sqrt{z} + 2A'\sqrt{z}$$

$$= 653a\sqrt{z} - \frac{26(azA')a\sqrt{z}}{a^2z}$$

$$= a\sqrt{z}(653 + 598) = 1251\sqrt{z},$$

$$\therefore B + B'\sqrt{z} = 9(197 + 139\sqrt{2}).$$

Hence, since  $u_1u_4 = g + a\sqrt{z} = 1 + \sqrt{2}$ ,

$$\begin{aligned} x = & 9^{\frac{1}{4}} \left[ (197 + 139\sqrt{2}) + \sqrt{\{(197 + 139\sqrt{2})^2 - \frac{1}{81}(1 + \sqrt{2})^5\}} \right]^{\frac{1}{4}} \\ & + 9^{\frac{1}{4}} \left[ (197 + 139\sqrt{2}) - \sqrt{\{(197 + 139\sqrt{2})^2 - \frac{1}{81}(1 + \sqrt{2})^5\}} \right]^{\frac{1}{4}} \\ & + 9^{\frac{1}{4}} \left[ (197 - 139\sqrt{2}) + \sqrt{\{(197 - 139\sqrt{2})^2 - \frac{1}{81}(1 - \sqrt{2})^5\}} \right]^{\frac{1}{4}} \\ & + 9^{\frac{1}{4}} \left[ (197 - 139\sqrt{2}) - \sqrt{\{(197 - 139\sqrt{2})^2 - \frac{1}{81}(1 - \sqrt{2})^5\}} \right]^{\frac{1}{4}}. \end{aligned}$$



To verify this result,

$$\begin{aligned}\frac{1}{81}(1 + \sqrt{2})^5 &= 1.012496, \\ \frac{1}{81}(1 - \sqrt{2})^5 &= -.000150535, \\ 197 + 139\sqrt{2} &= 393.57568, \\ (197 + 139\sqrt{2})^3 &= 154901.8, \\ 197 - 139\sqrt{2} &= .424315, \\ (197 - 139\sqrt{2})^3 &= .1800434, \\ \sqrt{\left\{(197 + 139\sqrt{2})^3 - \frac{1}{81}(1 + \sqrt{2})^5\right\}} &= 393.57436, \\ \sqrt{\left\{(197 - 139\sqrt{2})^3 - \frac{1}{81}(1 - \sqrt{2})^5\right\}} &= .180194.\end{aligned}$$

Therefore

$$\begin{aligned}u_1 &= 5.888, \\ u_4 &= .412, \\ u_2 &= 1.502, \\ u_3 &= -.276, \\ \therefore x &= 7.526.\end{aligned}$$

#### MODIFICATION OF THE METHOD TO MEET SPECIAL CASES.

*First special case: When  $p_2$  and  $p_3$  are both zero.*

§23. When  $p_2$  and  $p_3$  are both zero, a modification of the general method is rendered necessary by the circumstance that the equations (11) and (16), from which  $y$  and  $t$  are to be found, are then virtually one, and so are insufficient to give us the values of  $y$  and  $t$ . In fact, they become

$$\begin{aligned}& \left. \begin{aligned}t\left(50y^3 - \frac{2}{5}p_4y\right) - p_5y &= 0 \\ \text{and } \left\{t\left(50y^3 - \frac{2}{5}p_4y\right) - p_5y\right\}^2 &= 0\end{aligned} \right\} \quad (39)\end{aligned}$$

§24. In an article which appeared in No. 2, Vol. VII of this *Journal*, the present writer showed that, when  $p_2$  and  $p_3$  are both zero,  $p_4$  and  $p_5$  have the forms

$$\begin{aligned}p_4 &= \frac{5n^4(3-m)}{16+m^2} \\ p_5 &= \frac{n^5(22+m)}{16+m^2}\end{aligned} \quad (40)$$

These expressions for  $p_4$  and  $p_5$  furnish the criterion of solvability for the quintic

$$x^5 + p_4x + p_5 = 0. \quad (41)$$

The root of the equation is

$$x = \theta^{\frac{1}{5}} + \alpha\theta^{\frac{2}{5}} + \lambda\alpha^2\theta^{\frac{3}{5}} - \lambda\alpha^3\theta^{\frac{4}{5}},$$

where  $\lambda$  is a root of the quartic equation

$$\lambda^4 - m\lambda^3 - 6\lambda^2 + m\lambda + 1 = 0,$$

and

$$\alpha = -\frac{\lambda^2 + 1}{n\lambda(\lambda - 1)},$$

and

$$\theta = -\frac{n^3\lambda(\lambda - 1)^3}{(16 + m^2)(\lambda + 1)(\lambda^2 + 1)}.$$

In the same issue of the *Journal* in which these results were established, Mr. J. C. Glashan of Ottawa, in "Notes on the Quintic," gave the relations between the coefficients of the solvable quintic

$$x^5 + p_5x^4 + p_4x^3 + p_3x^2 + p_2x + p_1 = 0,$$

and, in his wider formulæ, the forms of  $p_4$  and  $p_5$  in (40) are included. They were subsequently announced by Mr. Emory McClintock, who had discovered them independently. It is to be regretted that Mr. Glashan has not made public the method by which his conclusions were reached.

§25. From our present position the criterion of solvability of the quintic (41) can be at once deduced, and the solution of the equation effected more readily than by the process employed in the article of the *Journal* just referred to. For, put

$$m = \frac{4A}{y} \text{ and } n = 2t;$$

then the first of the equations (9) may be written

$$p_4 = 5y(3 - m). \quad (42)$$

Also, by the second of equations (9),

$$p_5 = -4B + 40ty. \quad (43)$$

But, from the first of equations (8),  $B = -t(y + 2A)$ . Therefore

$$p_5 = 44ty + 8At = 2ty(22 + m). \quad (44)$$

And, by (15), in connection with (14),

$$t^4y - y^3 = A^2 = \frac{m^2y^2}{16},$$

$$\therefore y = \frac{16t^4}{16 + m^2} = \frac{n^4}{16 + m^2}.$$

Therefore also

$$2ty = \frac{2tn^4}{16+m^2} = \frac{n^4}{16+m^2}.$$

By the substitution of these values of  $y$  and  $2ty$  in (42) and (43), the formulæ (40) are obtained,  $n$  in (40) being what we have called  $2t$ . To find now the root of the equation (41), eliminate  $m$  from the equations (40). The result is a sextic equation,

$$\psi(n) = 0.$$

When the coefficients of the quintic (41) are commensurable, the sextic  $\psi(n) = 0$  has a commensurable root. Let this be found. Then  $n$  is known. Consequently, since  $n = 2t$ ,  $t$  is known. Then  $y$  is known from (39). Then  $B$  is obtained from the second of equations (9), and  $B'\sqrt{z}$  from (20). Also

$$u_1 u_4 = g + a\sqrt{z} = a\sqrt{z} = \sqrt{y}.$$

Therefore  $u_1 u_4$  is known. Therefore, as in §9, the root of the given quintic is found.

§26. *Third Example.*—As an illustrative example, let

$$x^5 + \frac{625}{4}x + 3750 = 0.$$

Here the equations furnishing the criterion of solvability are

$$\frac{625}{4} = \frac{5n^4(3-m)}{16+m^2},$$

$$3750 = \frac{n^5(22+m)}{16+m^2}.$$

These are satisfied by the values  $m = 2$ ,  $n = 5$ . Therefore

$$t = \frac{5}{2}.$$

Therefore, by (39),

$$y = \frac{125}{4}.$$

Therefore, by the second of equations (9),

$$B = -\frac{625}{4}.$$

And, by (20),

$$yB'\sqrt{z} = -2(c^2z)(c\sqrt{z}) = -2(t^2y)(t\sqrt{y}),$$

$$\therefore B'\sqrt{z} = -2t^3\sqrt{y} = -\frac{625}{8}\sqrt{5}.$$

$$\therefore B + B'\sqrt{z} = -\frac{625}{4}\left(1 + \frac{\sqrt{5}}{2}\right).$$

And  $u_1 u_4 = a\sqrt{z} = \frac{5\sqrt{5}}{2}$ . Therefore

$$\begin{aligned} x = & \left[ -\frac{625}{4} \left( 1 + \frac{\sqrt{5}}{2} \right) + \sqrt{\left\{ \left( \frac{625}{4} \right)^2 \left( 1 + \frac{\sqrt{5}}{2} \right)^2 - \left( \frac{5\sqrt{5}}{2} \right)^2 \right\}} \right]^{\frac{1}{2}}, \\ & + \left[ -\frac{625}{4} \left( 1 + \frac{\sqrt{5}}{2} \right) - \sqrt{\left\{ \left( \frac{625}{4} \right)^2 \left( 1 + \frac{\sqrt{5}}{2} \right)^2 - \left( \frac{5\sqrt{5}}{2} \right)^2 \right\}} \right]^{\frac{1}{2}}, \\ & + \left[ -\frac{625}{4} \left( 1 - \frac{\sqrt{5}}{2} \right) + \sqrt{\left\{ \left( \frac{625}{4} \right)^2 \left( 1 - \frac{\sqrt{5}}{2} \right)^2 + \left( \frac{5\sqrt{5}}{2} \right)^2 \right\}} \right]^{\frac{1}{2}}, \\ & + \left[ -\frac{625}{4} \left( 1 - \frac{\sqrt{5}}{2} \right) - \sqrt{\left\{ \left( \frac{625}{4} \right)^2 \left( 1 - \frac{\sqrt{5}}{2} \right)^2 + \left( \frac{5\sqrt{5}}{2} \right)^2 \right\}} \right]^{\frac{1}{2}}. \end{aligned}$$

Second special case: When  $u_1 u_4 = u_2 u_3$ .

§27. In the case in which  $u_1 u_4 = u_2 u_3$ ,  $a = 0$ . Consequently, if  $c$  is distinct from zero,  $t$  or  $\frac{c}{a}$  is infinite; while, if  $c$  is zero,  $t$  assumes the form  $\frac{0}{0}$ . As we cannot here proceed by finding  $t$ , the general method has to be modified.

§28. Because  $a = 0$ ,  $y = a^2 z = 0$ , and  $ty = \left( \frac{c}{a} \right) (a^2 z) = 0$ . Also  $t^2 y = c^2 z$ . Equation (5) becomes

$$gA = k^3 - c^2 z.$$

Therefore, from the first of equations (6),

$$k^3 - c^2 z = \frac{g}{20} (5g^3 - p_1), \quad (45)$$

and, from the second of equations (6),

$$p_2 = -4B. \quad (46)$$

From (33) and (35),

$$\begin{aligned} P &= 2k(k^3 - c^2 z) - (g^3 - a^2 z)(gk - acz) \\ &= 2k(k^3 - c^2 z) - g^3 k, \end{aligned}$$

and

$$aQ = 2at(k^3 - c^2 z) - (g^3 - a^2 z)(ak - gc) = g^3 c.$$

Therefore, from the first of equations (34), taken in connection with (46),

$$\frac{1}{4} g^3 p_2 + k \{ 2(k^3 - c^2 z) - g^3 \} + 2czA' = 0. \quad (47)$$

But,  $A'$  having been put for  $he(\theta^2 - \phi^2 z)$ , we have, from the value of  $z \{ he(\theta^2 - \phi^2 z) \}$ , obtained in §20,

$$z(A')^2 = (k^3 - c^2 z)^2 + g^4 - 2g^3(k^3 + c^2 z). \quad (48)$$

Therefore, from (47), by the elimination of  $A'$ ,

$$4c^2z = \frac{\left[ \frac{1}{4} g^2 p_5 + k \{ 2(k^2 - c^2z) - g^2 \} \right]^2}{(k^2 - c^2z)^2 + g^2 - 2g^2(k^2 + c^2z)}. \quad (49)$$

But, by (45),  $c^2z$  is known in terms of the coefficients of the given quintic. Therefore the equation (49) gives a relation necessarily subsisting between the coefficients  $p_3, p_4, p_5$  and  $p_6$  of the solvable quintic in which  $u_1u_4 = u_2u_3$ , in order that  $u_1u_4$  may be equal to  $u_2u_3$ . To find now the root of the quintic,  $c^2z$  is known by (45), and  $B$  is known by (46). To find  $B'\sqrt{z}$ , making use of the values of  $P$  and  $aQ$  and  $(A'\sqrt{z})^2$  obtained above, we have, from the second of equations (34),

$$g^2(B'\sqrt{z}) = c\sqrt{z} \{ 2(k^2 - c^2z) + g^2 \} + 2k(A'\sqrt{z}). \quad (50)$$

Now, by the first of equations (34), keeping in view the value of  $azA'$  in (35),

$$\begin{aligned} g^2B &= g^2 \{ 2k(k^2 - c^2z) - g^2k \} + 2azA' \left( \frac{g^2c}{a} \right) \\ &= g^2 \{ 2k(k^2 - c^2z) - g^2k \} + 2g^2czA'. \end{aligned}$$

As  $B$  and  $c^2z$  are known, this equation determines the sign of  $czA'$  or  $(c\sqrt{z})(A'\sqrt{z})$ , and therefore determines the signs with which  $c\sqrt{z}$  and  $A'\sqrt{z}$  are to be taken relatively to one another. Hence  $z(A')^2$  being given by (48),  $B'\sqrt{z}$  is given by (50). Therefore  $B + B'\sqrt{z}$  is known. And  $u_1u_4 = g$ . Therefore, because

$$u_1^5 = B + B'\sqrt{z} + \sqrt{\{ (B + B'\sqrt{z}) - g^5 \}},$$

the root of the quintic is known.

§ 29. *Fourth Example.*—As an illustrative example, let

$$x^5 - \frac{22}{5}x^3 - \frac{11}{25}x^2 + \frac{11 \times 42}{125}x + \frac{11 \times 89}{3125} = 0.$$

Finding the value of  $c^2z$  from (45), and substituting in (49), we find that (49) is satisfied. Then, as in the preceding section,

$$p_5 = -4B \therefore B = -\frac{11 \times 89}{4 \times 5^5}.$$

The value of  $c^2z$  is  $\frac{5}{16} \left( \frac{11}{25} \right)^2$ . Therefore

$$k^2 - c^2z = \left( \frac{11}{25} \right)^2 \left( \frac{1}{400} - \frac{5}{16} \right) = -\frac{31}{100} \left( \frac{11}{25} \right)^2,$$

and

$$k^2 + c^2z = \left( \frac{11}{25} \right)^2 \left( \frac{1}{400} + \frac{5}{16} \right) = \frac{63}{200} \left( \frac{11}{25} \right)^2.$$

From (48),  $(A'\sqrt{z})^2 = \left(\frac{11^2 \times 5}{4 \times 5^3}\right)^2$ .

The value of  $czA'$ , obtained from (47), is negative. Therefore  $c\sqrt{z}$  and  $A'\sqrt{z}$  must have different signs. Put

$$c\sqrt{z} = \frac{11}{25} \frac{\sqrt{5}}{4}$$

and

$$A'\sqrt{z} = -\frac{1}{5} \left(\frac{11}{25}\right)^2 \frac{\sqrt{5}}{4}.$$

Then, from (50), dividing by  $g^2$ ,

$$B'\sqrt{z} = -\frac{11\sqrt{5}}{4 \times 5^3}.$$

And  $u_1 u_4 = g$ . Therefore

$$u_1^5 = \frac{-11}{4 \times 5^3} (89 + 25\sqrt{5}) + \sqrt{\left\{ \frac{121}{16 \times 25^3} (89 + 25\sqrt{5})^2 - \left(\frac{11}{25}\right)^5 \right\}}.$$

This is, in a different form, the value of  $u_1^5$  given by Lagrange.

§30. *Fifth Example.*—In the instance just considered,  $c$  is distinct from zero. It may be well to give an example in which  $c$  is zero, as the mode adopted above of determining the sign of  $A'\sqrt{z}$  does not apply in that case. Let

$$x^5 + 20x^3 + 20x^2 + 30x + 10 = 0.$$

Here  $g = -2$ , and  $k = -1$ . Therefore (45) becomes

$$20(1 - c^2z) = -2(20 - 30).$$

Therefore  $c^2z = 0$ . This value of  $c^2z$  satisfies (49). Then, by (46),  $B = -\frac{5}{2}$ .

From 48,  $(A'\sqrt{z})^2 = 1 + 64 + 16 = 81 \therefore A'\sqrt{z} = 9$ .

Because  $c$  is zero, we cannot determine the sign of  $A'\sqrt{z}$  relatively to that of  $c\sqrt{z}$ . But the root is the same, whatever sign be taken. Having found  $A'\sqrt{z}$ ,

we have, from (50),  $4B'\sqrt{z} = -2 \times 9 \therefore B'\sqrt{z} = -\frac{9}{2}$ . Therefore

$$B + B'\sqrt{z} = -\frac{5}{2} - \frac{9}{2} = -7, \quad B - B'\sqrt{z} = -\frac{5}{2} + \frac{9}{2} = 2.$$

And  $u_1 u_4 = g = -2$ . Therefore

$$u_1^5 = -7 + \sqrt{\{49 - (-2)^5\}} = 2,$$

$$u_4^5 = -7 - \sqrt{\{49 - (-2)^5\}} = -16,$$

$$u_2^5 = 2 + \sqrt{\{4 + 2^5\}} = 8,$$

$$u_3^5 = 2 - \sqrt{\{4 + 2^5\}} = -4.$$

Therefore  $x = 2^{\frac{1}{5}} - 2^{\frac{3}{5}} + 2^{\frac{2}{5}} - 2^{\frac{4}{5}}$ .

§31. In the article of the *American Journal of Mathematics* (Vol. VI, page 103) referred to in the opening paragraph of this paper, the solvable irreducible quintic in which  $u_1u_4$  is equal to  $u_2u_3$  was discussed, and the roots of the equation were shown to be determinable in terms of the coefficients  $p_2, p_3$ , etc., even while these coefficients have no definite numerical values assigned to them, but remain symbolical. The solution that has now been given is much simpler than the former; equally with the former, it is applicable to equations with symbolical coefficients, the assumption being of course made that the coefficients are related as in (49); and it possesses the advantage of being part of a general theory.

## ADDITIONAL EXAMPLES.

§32. *Sixth Example.*—Let

$$x^5 + 320x^3 - 1000x + 4288 = 0.$$

Here  $g = 0$ ,  $k = -16$ . Because  $g = 0$ , we use the formulæ (11) and (16). The commensurable values of  $y$  and  $t$  which satisfy (11) and (16) are

$$y = 8, \quad t = 6.$$

Also, by (14),

$$A = 56.$$

Therefore, from the second of equations (9) and from (20),

$$\begin{aligned} B &= -16 \times 37, \quad B'\sqrt{z} = -16 \times 20, \\ \therefore B + B'\sqrt{z} &= -16(37 + 20\sqrt{2}). \end{aligned}$$

And  $u_1u_4 = -2\sqrt{2}$ . Therefore

$$\begin{aligned} x &= u_1 + u_4 + u_2 + u_3 \\ &= [-16(37 + 20\sqrt{2}) + \sqrt{\{256(37 + 20\sqrt{2})^3 - (-2\sqrt{2})^5\}}]^{\frac{1}{4}} \\ &\quad + [-16(37 + 20\sqrt{2}) - \sqrt{\{256(37 + 20\sqrt{2})^3 - (-2\sqrt{2})^5\}}]^{\frac{1}{4}} \\ &\quad + [-16(37 - 20\sqrt{2}) + \sqrt{\{256(37 - 20\sqrt{2})^3 - (2\sqrt{2})^5\}}]^{\frac{1}{4}} \\ &\quad + [-16(37 - 20\sqrt{2}) - \sqrt{\{256(37 - 20\sqrt{2})^3 - (2\sqrt{2})^5\}}]^{\frac{1}{4}}. \end{aligned}$$

§33. *Seventh Example.*—Let

$$\left(\frac{x}{10}\right)^5 + 40\left(\frac{x}{10}\right)^3 - 69\left(\frac{x}{10}\right) + 108 = 0,$$

or,

$$x^5 + 40000x^3 - 690000x + 10800000 = 0.$$

Here  $g = 0$ ,  $k = -2000$ . Because  $g = 0$ , we use the formulæ (11) and (16).

The commensurable values of  $y$  and  $t$  which satisfy (11) and (16) are,

$$y = 2000, \quad t = 50.$$

Also, by (14),

$$A = 36000.$$

Therefore, from the second of equations (9) and from (20),

$$B = -1700000, \quad B'\sqrt{z} = -400000\sqrt{5}.$$

And  $u_1u_4 = -20\sqrt{5}$ . Therefore

$$\begin{aligned} x &= u_1 + u_4 + u_2 + u_3 \\ &= [-100000(17 + 4\sqrt{5}) + \sqrt{\{100000^2(17 + 4\sqrt{5})^2 + (20\sqrt{5})^2\}}]^{1/2} \\ &\quad + [-100000(17 + 4\sqrt{5}) - \sqrt{\{100000^2(17 + 4\sqrt{5})^2 + (20\sqrt{5})^2\}}]^{1/2} \\ &\quad + [-100000(17 - 4\sqrt{5}) + \sqrt{\{100000^2(17 - 4\sqrt{5})^2 - (20\sqrt{5})^2\}}]^{1/2} \\ &\quad + [-100000(17 - 4\sqrt{5}) - \sqrt{\{100000^2(17 - 4\sqrt{5})^2 - (20\sqrt{5})^2\}}]^{1/2}. \end{aligned}$$

§34. *Eighth Example.*—Let

$$x^5 - 20x^3 + 250x - 400 = 0.$$

Here  $g = 2$ ,  $k = 0$ . Because  $g$  is distinct from zero, we use not the formulæ (11) and (16) as in the two preceding examples, but (37) and (38). The commensurable values of  $y$  and  $t$  which satisfy (37) and (38) are

$$y = 2, \quad t = -2.$$

Therefore, from the second of equations (6),  $B = 60$ . The value of  $azA'$  is obtained from (35). Thus, since  $(A'\sqrt{z})(a\sqrt{z}) = azA'$ , and since  $a\sqrt{z} = \sqrt{y} = \sqrt{2}$ , the value of  $A'\sqrt{z}$  is known. Then, by the second of equations (34),  $B'\sqrt{z} = 44\sqrt{2}$ .

Therefore

$$B + B'\sqrt{z} = 4(15 + 11\sqrt{2}).$$

And  $u_1u_4 = g + a\sqrt{z} = 2 + \sqrt{2}$ . Therefore

$$\begin{aligned} x &= u_1 + u_4 + u_2 + u_3 \\ &= [4(15 + 11\sqrt{2}) + \sqrt{\{16(15 + 11\sqrt{2})^2 - (2 + \sqrt{2})^2\}}]^{1/2} \\ &\quad + [4(15 + 11\sqrt{2}) - \sqrt{\{16(15 + 11\sqrt{2})^2 - (2 + \sqrt{2})^2\}}]^{1/2} \\ &\quad + [4(15 - 11\sqrt{2}) + \sqrt{\{16(15 - 11\sqrt{2})^2 - (2 - \sqrt{2})^2\}}]^{1/2} \\ &\quad + [4(15 - 11\sqrt{2}) - \sqrt{\{16(15 - 11\sqrt{2})^2 - (2 - \sqrt{2})^2\}}]^{1/2}. \end{aligned}$$

§35. *Ninth Example.*—Let

$$x^5 - 5x^3 + \frac{85}{8}x - \frac{13}{2} = 0.$$

Here  $g = \frac{1}{2}$ ,  $k = 0$ . Because  $g$  is distinct from zero, we use the formulæ (37)



and (38). The commensurable values of  $y$  and  $t$  which satisfy (37) and (38) are

$$y = \frac{1}{8}; t = -1.$$

Therefore, from the second of equations (6),  $B = \frac{3}{8}$ . Finding  $A'\sqrt{z}$  as in the immediately preceding example, we have, from the second of equations (34),  $B'\sqrt{z} = \frac{3}{8}\sqrt{2}$ . Therefore

$$B + B'\sqrt{z} = \frac{3}{8}(1 + \sqrt{2}).$$

And  $u_1 u_4 = \frac{1}{4}(2 + \sqrt{2})$ . Therefore

$$\begin{aligned} x &= u_1 + u_4 + u_2 + u_3 \\ &= \left[ \frac{3}{8}(1 + \sqrt{2}) + \sqrt{\left\{ \frac{9}{64}(1 + \sqrt{2})^2 - \frac{1}{4^5}(2 + \sqrt{2})^5 \right\}} \right]^{\frac{1}{2}} \\ &\quad + \left[ \frac{3}{8}(1 + \sqrt{2}) - \sqrt{\left\{ \frac{9}{64}(1 + \sqrt{2})^2 - \frac{1}{4^5}(2 + \sqrt{2})^5 \right\}} \right]^{\frac{1}{2}} \\ &\quad + \left[ \frac{3}{8}(1 - \sqrt{2}) + \sqrt{\left\{ \frac{9}{64}(1 - \sqrt{2})^2 - \frac{1}{4^5}(2 - \sqrt{2})^5 \right\}} \right]^{\frac{1}{2}} \\ &\quad + \left[ \frac{3}{8}(1 - \sqrt{2}) - \sqrt{\left\{ \frac{9}{64}(1 - \sqrt{2})^2 - \frac{1}{4^5}(2 - \sqrt{2})^5 \right\}} \right]^{\frac{1}{2}}. \end{aligned}$$

§36. *Tenth Example.*—Let

$$x^5 + \frac{20x}{17} + \frac{21}{17} = 0.$$

This and the next two examples are intended as additional illustrations of the method to be followed when  $p_3$  and  $p_5$  are both zero. The equations furnishing the criterion of solvability given in §24 are

$$\frac{20}{17} = \frac{5n^4(3-m)}{16+m^2}$$

and

$$\frac{21}{17} = \frac{n^5(22+m)}{16+m^2},$$

and they are satisfied by the commensurable values  $m = -1$ ,  $n = 1$ . But, by §25,  $n = 2t$ . Therefore  $t = \frac{1}{2}$ . Putting  $k = 0$  and  $t = \frac{1}{2}$ , equation (11)

becomes

$$\frac{1}{2} \left( 50y - \frac{2}{5} p_4 \right) = p_5.$$

$$\therefore y = \frac{1}{17} \therefore a\sqrt{z} = \sqrt{y} = \frac{\sqrt{17}}{17}.$$

Therefore also  $c\sqrt{z} = t(a\sqrt{z}) = \frac{\sqrt{17}}{34}$ . Therefore, by (20),

$$yB'\sqrt{z} = -2(c^2z)(c\sqrt{z}) \therefore B'\sqrt{z} = -\frac{\sqrt{17}}{68}.$$

And, by the second of equations (9),  $B = -\frac{1}{68}$ . Therefore

$$B + B'\sqrt{z} = -\frac{1}{68} (1 + \sqrt{17}).$$

And  $u_1 u_4 = \frac{\sqrt{17}}{17}$ . Therefore

$$\begin{aligned} x = & \left[ -\frac{1}{68} (1 + \sqrt{17}) + \sqrt{\left\{ \left( \frac{1 + \sqrt{17}}{68} \right)^2 - \left( \frac{\sqrt{17}}{17} \right)^5 \right\}} \right]^{\frac{1}{4}} \\ & + \left[ -\frac{1}{68} (1 + \sqrt{17}) - \sqrt{\left\{ \left( \frac{1 + \sqrt{17}}{68} \right)^2 - \left( \frac{\sqrt{17}}{17} \right)^5 \right\}} \right]^{\frac{1}{4}} \\ & + \left[ -\frac{1}{68} (1 - \sqrt{17}) + \sqrt{\left\{ \left( \frac{1 - \sqrt{17}}{68} \right)^2 + \left( \frac{\sqrt{17}}{17} \right)^5 \right\}} \right]^{\frac{1}{4}} \\ & + \left[ -\frac{1}{68} (1 - \sqrt{17}) - \sqrt{\left\{ \left( \frac{1 - \sqrt{17}}{68} \right)^2 + \left( \frac{\sqrt{17}}{17} \right)^5 \right\}} \right]^{\frac{1}{4}}. \end{aligned}$$

§37. *Eleventh Example.*—Let

$$x^5 - \frac{4x}{13} + \frac{29}{65} = 0.$$

The equations furnishing the criterion of solvability are

$$\begin{aligned} -\frac{4}{13} &= \frac{5n^4(3-m)}{16+m^2}, \\ \frac{29}{65} &= \frac{n^5(22+m)}{16+m^2}, \end{aligned}$$

and they are satisfied by the values  $m = 7, n = 1$ . By §25,  $n = 2t$ . Therefore  $t = \frac{1}{2}$ . Therefore, from (11),  $y = \frac{1}{65}$ . Therefore

$$a\sqrt{z} = \frac{\sqrt{65}}{65} \therefore c\sqrt{z} = t(a\sqrt{z}) = \frac{\sqrt{65}}{130}.$$

Therefore, by (20),  $yB'\sqrt{z} = -2(c^2z)(c\sqrt{z})$ . Therefore  $B'\sqrt{z} = -\frac{\sqrt{65}}{260}$ . And,

by the second of equations (9),  $B = -\frac{9}{260}$ . Therefore

$$B + B'\sqrt{z} = -\frac{9 + \sqrt{65}}{260}.$$

And  $u_1 u_4 = \frac{\sqrt{65}}{65}$ . Therefore

$$\begin{aligned} x = & \left[ -\frac{9+\sqrt{65}}{260} + \sqrt{\left\{ \left( \frac{9+\sqrt{65}}{260} \right)^3 - \left( \frac{\sqrt{65}}{65} \right)^5 \right\}} \right]^{\frac{1}{3}} \\ & + \left[ -\frac{9+\sqrt{65}}{260} - \sqrt{\left\{ \left( \frac{9+\sqrt{65}}{260} \right)^3 - \left( \frac{\sqrt{65}}{65} \right)^5 \right\}} \right]^{\frac{1}{3}} \\ & + \left[ -\frac{9-\sqrt{65}}{260} + \sqrt{\left\{ \left( \frac{9-\sqrt{65}}{260} \right)^3 + \left( \frac{\sqrt{65}}{65} \right)^5 \right\}} \right]^{\frac{1}{3}} \\ & + \left[ -\frac{9-\sqrt{65}}{260} - \sqrt{\left\{ \left( \frac{9-\sqrt{65}}{260} \right)^3 + \left( \frac{\sqrt{65}}{65} \right)^5 \right\}} \right]^{\frac{1}{3}}. \end{aligned}$$

§38. *Twelfth Example.*—Let

$$x^5 + \frac{10x}{13} + \frac{3}{13} = 0.$$

The equations furnishing the criterion of solvability are

$$\begin{aligned} \frac{10}{13} &= \frac{5n^4(3-m)}{16+m^3}, \\ \frac{3}{13} &= \frac{n^5(22+m)}{16+m^2}, \end{aligned}$$

and they are satisfied by the values  $m = -7$ ,  $n = 1$ . By §25,  $n = 2t$ . Therefore  $t = \frac{1}{2}$ . Therefore, from (11),  $y = \frac{1}{65}$ . Therefore

$$c\sqrt{z} = i\sqrt{y} = \frac{\sqrt{65}}{130}.$$

And, by (20),  $yB'\sqrt{z} = -2(c^2z)(c\sqrt{z})$ . Therefore  $B'\sqrt{z} = -\frac{\sqrt{65}}{260}$ . And, by the second of equations (9),  $B = \frac{1}{52}$ . Therefore

$$B + B'\sqrt{z} = \frac{5-\sqrt{65}}{260}.$$

And  $u_1 u_4 = \frac{\sqrt{65}}{65}$ . Therefore

$$\begin{aligned} x_1 = & \left[ \frac{5-\sqrt{65}}{260} + \sqrt{\left\{ \left( \frac{5-\sqrt{65}}{260} \right)^2 - \left( \frac{\sqrt{65}}{65} \right)^5 \right\}} \right]^{\frac{1}{3}} \\ & + \left[ \frac{5-\sqrt{65}}{260} - \sqrt{\left\{ \left( \frac{5-\sqrt{65}}{260} \right)^2 - \left( \frac{\sqrt{65}}{65} \right)^5 \right\}} \right]^{\frac{1}{3}} \\ & + \left[ \frac{5+\sqrt{65}}{260} + \sqrt{\left\{ \left( \frac{5+\sqrt{65}}{260} \right)^2 + \left( \frac{\sqrt{65}}{65} \right)^5 \right\}} \right]^{\frac{1}{3}} \\ & + \left[ \frac{5+\sqrt{65}}{260} - \sqrt{\left\{ \left( \frac{5+\sqrt{65}}{260} \right)^2 + \left( \frac{\sqrt{65}}{65} \right)^5 \right\}} \right]^{\frac{1}{3}}. \end{aligned}$$

§39. *Thirteenth Example.*—Let

$$x^5 + 110(5x^3 + 60x^2 + 800x + 8320) = 0.$$

Here  $g = -55$ ,  $k = -330$ , and the commensurable values of  $y$  and  $t$  which satisfy (37) and (38) are

$$y = 5 \times 11^2, \quad t = 10.$$

Therefore, from the second of equations (6),  $B = -220 \times 765$ . Finding  $A'\sqrt{z}$  as in the second example, we have, from the second of equations (34),  $B'\sqrt{z} = -220 \times 337\sqrt{5}$ . Therefore

$$B + B'\sqrt{z} = -220(765 + 337\sqrt{5}).$$

And  $u_1u_4 = -11(5 + \sqrt{5})$ . Therefore

$$\begin{aligned} x = & [-220(765 + 337\sqrt{5}) + \sqrt{\{220^3(765 + 337\sqrt{5})^3 - (-55 - 11\sqrt{5})^5\}}]^{\frac{1}{5}} \\ & + [-220(765 + 337\sqrt{5}) - \sqrt{\{220^3(765 + 337\sqrt{5})^3 - (-55 - 11\sqrt{5})^5\}}]^{\frac{1}{5}} \\ & + [-220(765 - 337\sqrt{5}) + \sqrt{\{220^3(765 - 337\sqrt{5})^3 - (-55 + 11\sqrt{5})^5\}}]^{\frac{1}{5}} \\ & + [-220(765 - 337\sqrt{5}) - \sqrt{\{220^3(765 - 337\sqrt{5})^3 - (-55 + 11\sqrt{5})^5\}}]^{\frac{1}{5}}. \end{aligned}$$

§40. *Fourteenth Example.*—Let

$$x^5 - 20x^3 - 80x^2 - 150x - 656 = 0.$$

Here  $g = 2$ ,  $k = 4$ , and the commensurable values of  $y$  and  $t$  which satisfy (37) and (38) are

$$y = 2, \quad t = 2.$$

Therefore, from the second of equations (6),  $B = 204$ . Finding  $A'\sqrt{z}$  as in the immediately preceding example, we have, from the second of equations (34),  $B'\sqrt{z} = 144\sqrt{2}$ . Therefore

$$\begin{aligned} B &= 12 \times 17, \text{ and } B'\sqrt{z} = 144\sqrt{2}, \\ \therefore B + B'\sqrt{z} &= 12(17 + 12\sqrt{2}). \end{aligned}$$

And  $u_1u_4 = 2 + \sqrt{2}$ . Therefore

$$\begin{aligned} x_1 = & u_1 + u_4 + u_3 + u_3 \\ & = [12(17 + 12\sqrt{2}) + \sqrt{\{144(17 + 12\sqrt{2})^3 - (2 + \sqrt{2})^5\}}]^{\frac{1}{5}} \\ & + [12(17 + 12\sqrt{2}) - \sqrt{\{144(17 + 12\sqrt{2})^3 - (2 + \sqrt{2})^5\}}]^{\frac{1}{5}} \\ & + [12(17 - 12\sqrt{2}) + \sqrt{\{144(17 - 12\sqrt{2})^3 - (2 - \sqrt{2})^5\}}]^{\frac{1}{5}} \\ & + [12(17 - 12\sqrt{2}) - \sqrt{\{144(17 - 12\sqrt{2})^3 - (2 - \sqrt{2})^5\}}]^{\frac{1}{5}}. \end{aligned}$$

§41. *Fifteenth Example.*—Let

$$x^5 - 40x^3 + 160x^2 + 1900x - 5888 = 0.$$

. There-

And, by

Here  $g = 4$ ,  $k = -8$ , and the commensurable values of  $y$  and  $t$  which satisfy (37) and (38) are

$$y = 8, t = -4.$$

Therefore, from the second of equations (6),  $B = 1152$ . Finding  $A'\sqrt{z}$  as in the two preceding examples, we have, from the second of equations (34),  $B'\sqrt{z} = 816\sqrt{2}$ . Therefore

$$B + B'\sqrt{z} = 48(24 + 17\sqrt{2}).$$

And  $u_1u_4 = 4 + 2\sqrt{2}$ . Therefore

$$\begin{aligned} x = & [48(24 + 17\sqrt{2}) + \sqrt{\{48^2(24 + 17\sqrt{2})^2 - (4 + 2\sqrt{2})^5\}}]^{\frac{1}{2}} \\ & + [48(24 + 17\sqrt{2}) - \sqrt{\{48^2(24 + 17\sqrt{2})^2 - (4 + 2\sqrt{2})^5\}}]^{\frac{1}{2}} \\ & + [48(24 - 17\sqrt{2}) + \sqrt{\{48^2(24 - 17\sqrt{2})^2 - (4 - 2\sqrt{2})^5\}}]^{\frac{1}{2}} \\ & + [48(24 - 17\sqrt{2}) - \sqrt{\{48^2(24 - 17\sqrt{2})^2 - (4 - 2\sqrt{2})^5\}}]^{\frac{1}{2}}. \end{aligned}$$

§42. *Sixteenth Example.*—Let

$$\left(\frac{x}{2}\right)^5 - 50\left(\frac{x}{2}\right)^3 - 600\left(\frac{x}{2}\right)^2 - 2000\left(\frac{x}{2}\right) - 11200 = 0,$$

or

$$x^5 - 200x^3 - 4800x^2 - 32000x - 3200 \times 112 = 0.$$

Here  $g = 20$ ,  $k = 240$ , and the commensurable values of  $y$  and  $t$  which satisfy (37) and (38) are

$$y = 80, t = 20.$$

Therefore, from the second of equations (6),  $B = 640 \times 165$ . Finding  $A'\sqrt{z}$  as in the preceding examples of the same type, we have, from the second of equations (34),  $B'\sqrt{z} = 640 \times 73\sqrt{5}$ . Therefore

$$B + B'\sqrt{z} = 640(165 + 73\sqrt{5}).$$

And  $u_1u_4 = 4(5 + \sqrt{5})$ . Therefore

$$\begin{aligned} x = & [640(165 + 73\sqrt{5}) + \sqrt{\{640^2(165 + 73\sqrt{5})^2 - (20 + 4\sqrt{5})^5\}}]^{\frac{1}{2}} \\ & + [640(165 + 73\sqrt{5}) - \sqrt{\{640^2(165 + 73\sqrt{5})^2 - (20 + 4\sqrt{5})^5\}}]^{\frac{1}{2}} \\ & + [640(165 - 73\sqrt{5}) + \sqrt{\{640^2(165 - 73\sqrt{5})^2 - (20 - 4\sqrt{5})^5\}}]^{\frac{1}{2}} \\ & + [640(165 - 73\sqrt{5}) - \sqrt{\{640^2(165 - 73\sqrt{5})^2 - (20 - 4\sqrt{5})^5\}}]^{\frac{1}{2}}. \end{aligned}$$

§43. *Seventeenth Example.*—Let

$$x^5 + 110(5x^3 + 20x^2 - 360x + 800) = 0.$$

Here  $g = -55$ ,  $k = -110$ , and the commensurable values of  $y$  and  $t$  which satisfy (37) and (38) are

$$y = 5 \times 11^2, t = -10.$$

Therefore, from the second of equations (6),  $B = -11 \times 7500$ . Finding  $A'\sqrt{z}$  as in preceding examples of the same type, we have, from the second of equations (34),  $B'\sqrt{z} = -11 \times 2700\sqrt{5}$ . Therefore

$$B + B'\sqrt{z} = -3300(25 + 9\sqrt{5}).$$

And  $u_1u_4 = -11(5 + \sqrt{5})$ . Therefore

$$\begin{aligned} x = & [-3300(25 + 9\sqrt{5}) + \sqrt{\{3300^2(25 + 9\sqrt{5})^2 + 11^5(5 + \sqrt{5})^5\}}]^{1/2} \\ & + [-3300(25 + 9\sqrt{5}) - \sqrt{\{3300^2(25 + 9\sqrt{5})^2 + 11^5(5 + \sqrt{5})^5\}}]^{1/2} \\ & + [-3300(25 - 9\sqrt{5}) + \sqrt{\{3300^2(25 - 9\sqrt{5})^2 + 11^5(5 - \sqrt{5})^5\}}]^{1/2} \\ & + [-3300(25 - 9\sqrt{5}) - \sqrt{\{3300^2(25 - 9\sqrt{5})^2 + 11^5(5 - \sqrt{5})^5\}}]^{1/2}. \end{aligned}$$

§44. Eighteenth Example.—Let

$$x^5 - 20x^3 + 320x^2 + 540x + 638 = 0.$$

Here  $g = 2$ ,  $k = -16$ , and the commensurable values of  $y$  and  $t$  that satisfy (37) and (38) are

$$y = 8, \quad t = -5.$$

Therefore, from the second of equations (6), and the second of equations (34),

$$\begin{aligned} B &= -12 \times 166, \quad B'\sqrt{z} = -12 \times 117\sqrt{2}, \\ \therefore B + B'\sqrt{z} &= -12(166 + 117\sqrt{2}). \end{aligned}$$

And  $u_1u_4 = 2(1 + \sqrt{2})$ . Therefore

$$\begin{aligned} x = & [-12(166 + 117\sqrt{2}) + \sqrt{\{144(166 + 117\sqrt{2})^2 - 32(1 + \sqrt{2})^5\}}]^{1/2} \\ & + [-12(166 + 117\sqrt{2}) - \sqrt{\{144(166 + 117\sqrt{2})^2 - 32(1 + \sqrt{2})^5\}}]^{1/2} \\ & + [-12(166 - 117\sqrt{2}) + \sqrt{\{144(166 - 117\sqrt{2})^2 - 32(1 - \sqrt{2})^5\}}]^{1/2} \\ & + [-12(166 - 117\sqrt{2}) - \sqrt{\{144(166 - 117\sqrt{2})^2 - 32(1 - \sqrt{2})^5\}}]^{1/2}. \end{aligned}$$

§45. Nineteenth Example.—Let

$$x^5 - 20x^3 - 160x^2 - 420x - 8928 = 0.$$

Here  $g = 2$ ,  $k = 8$ , and the commensurable values of  $y$  and  $t$  which satisfy (37) and (38) are

$$y = 72, \quad t = -\frac{7}{3}.$$

Therefore, from the second of equations (6),  $B = 552$ . Finding  $A'\sqrt{z}$  as in preceding examples of the same type, we have, from the second of equations (34),  $B'\sqrt{z} = -284\sqrt{2}$ . Therefore

$$B + B'\sqrt{z} = 4(138 - 71\sqrt{2}).$$

And  $u_1 u_4 = 2 - 6\sqrt{2}$ . Therefore

$$\begin{aligned} x = & [4(138 - 71\sqrt{2}) + \sqrt{\{16(138 - 71\sqrt{2})^2 - (2 - 6\sqrt{2})^2\}}]^{\frac{1}{2}} \\ & + [4(138 - 71\sqrt{2}) - \sqrt{\{16(138 - 71\sqrt{2})^2 - (2 - 6\sqrt{2})^2\}}]^{\frac{1}{2}} \\ & + [4(138 + 71\sqrt{2}) + \sqrt{\{16(138 + 71\sqrt{2})^2 - (2 + 6\sqrt{2})^2\}}]^{\frac{1}{2}} \\ & + [4(138 + 71\sqrt{2}) - \sqrt{\{16(138 + 71\sqrt{2})^2 - (2 + 6\sqrt{2})^2\}}]^{\frac{1}{2}}. \end{aligned}$$

§46. *Twentieth Example.*—Let

$$x^5 - 20x^3 + 170x + 208 = 0.$$

Here  $g = 2$ ,  $k = 0$ , and the commensurable values of  $y$  and  $t$  which satisfy (37) and (38) are

$$y = 2, \quad t = 2.$$

Then in the usual way we get

$$B + B'\sqrt{z} = -12(1 - \sqrt{2}).$$

And  $u_1 u_4 = 2 + \sqrt{2}$ . Therefore

$$\begin{aligned} x = & [-12(1 - \sqrt{2}) + \sqrt{\{144(1 - \sqrt{2})^2 - (2 + \sqrt{2})^2\}}]^{\frac{1}{2}} \\ & + [-12(1 - \sqrt{2}) - \sqrt{\{144(1 - \sqrt{2})^2 - (2 + \sqrt{2})^2\}}]^{\frac{1}{2}} \\ & + [-12(1 + \sqrt{2}) + \sqrt{\{144(1 + \sqrt{2})^2 - (2 - \sqrt{2})^2\}}]^{\frac{1}{2}} \\ & + [-12(1 + \sqrt{2}) - \sqrt{\{144(1 + \sqrt{2})^2 - (2 - \sqrt{2})^2\}}]^{\frac{1}{2}}. \end{aligned}$$

ents.

isfy (37)



