





**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

24 28 25
26 32 22
36 20
8

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

10

© 1982

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured covers/
Couverture de couleur
- Covers damaged/
Couverture endommagée
- Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée
- Cover title missing/
Le titre de couverture manque
- Coloured maps/
Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur
- Bound with other material/
Relié avec d'autres documents
- Tight binding may cause shadows or distortion
along interior margin/
La reliure serrée peut causer de l'ombre ou de la
distortion le long de la marge intérieure
- Blank leaves added during restoration may
appear within the text. Whenever possible, these
have been omitted from filming/
Il se peut que certaines pages blanches ajoutées
lors d'une restauration apparaissent dans le texte,
mais, lorsque cela était possible, ces pages n'ont
pas été filmées.
- Additional comments:/
Commentaires supplémentaires:

- Coloured pages/
Pages de couleur
- Pages damaged/
Pages endommagées
- Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- Pages detached/
Pages détachées
- Showthrough/
Transparence
- Quality of print varies/
Qualité inégale de l'impression
- Includes supplementary material/
Comprend du matériel supplémentaire
- Only edition available/
Seule édition disponible
- Pages wholly or partially obscured by errata
slips, tissues, etc., have been refilmed to
ensure the best possible image/
Les pages totalement ou partiellement
obscurcies par un feuillet d'errata, une pelure,
etc., ont été filmées à nouveau de façon à
obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	12X	14X	16X	18X	20X	22X	24X	26X	28X	30X	32X
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

The copy filmed here has been reproduced thanks to the generosity of:




Library,
Geological Survey of Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol → (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

Bibliothèque,
Commission Géologique du Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole → signifie "A SUIVRE", le symbole ∇ signifie "FIN".


Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

ails
du
modifier
une
page

rrata
o

pelure,
n à

32X



SMITHSONIAN MISCELLANEOUS COLLECTIONS.

256

MONOGRAPHS

OF THE

D I P T E R A

OF

NORTH AMERICA.

PART III.

PREPARED FOR THE SMITHSONIAN INSTITUTION

BY

H. LOEW.



WASHINGTON:
SMITHSONIAN INSTITUTION.

DECEMBER, 1873

ADVERTISEMENT.

THE present publication is Part III of a work upon the Diptera of North America, prepared at the request of the Smithsonian Institution, by Dr. H. Loew, of Guben, Prussia, well known as one of the most eminent cultivators of this branch of entomology.

The first part of this series of monographs was published in 1862, and included the families of *Trypetidæ*, *Sciomyzidæ*, *Ephydriidæ*, and *Cecidomyidæ*. The second part appeared in 1864, and consisted principally of a monograph of the *Dolichopodidæ*. The fourth part was issued in 1869, embracing a monograph of part of the *Tipulidæ*.

They were not published in sequence, but in the order in which materials could be collected for their preparation. The original manuscript of Dr. Loew was written in German, and the Institution is indebted to Baron R. Osten-Sacken for translating it into English; and to Mr. R. A. Witthaus, Jr., of New York, for revising and correcting the proof-sheets, in the absence of Baron Osten-Sacken.

JOSEPH HENRY,

Secretary S. I.

WASHINGTON, Dec. 1873.

TABLE OF CONTENTS.

THE FAMILY ORTALIDÆ.

PART FIRST.—INTRODUCTION.

	PAGE
THE EUROPEAN ORTALIDÆ	2
Meigen's genus <i>Ortalis</i>	2
1. Characters which Meigen's species of <i>Ortalis</i> have in common	2
2. Species erroneously placed in the genus <i>Ortalis</i>	3
The other European <i>Ortaliidæ</i>	4
1. Forms resembling <i>Ortalis lamed</i>	5
2. Forms resembling <i>Ortalis syngenesic</i>	6
3. Forms resembling <i>Ortalis paludum</i>	7
4. Forms not resembling any of Meigen's species	8
Aciphorea not belonging to the <i>Ortaliidæ</i>	8
Summary of the European <i>Ortaliidæ</i>	10
THE ORTALIDÆ OTHER THAN EUROPEAN	11
<i>a.</i> In Wiedemann	11
<i>b.</i> In Rob. Desvoidy	15
<i>c.</i> In Macquart	19
<i>d.</i> In Walker	26
<i>e.</i> In Bigot, Gerstæcker, Doleschall, and Saunders	27
NATURAL CHARACTERS OF THE FAMILY ORTALIDÆ	28
RELATIONSHIP OF THE ORTALIDÆ	30
DIAGNOSTIC OR ARTIFICIAL DEFINITION OF THE ORTALIDÆ	31

PART SECOND.—SYSTEMATIC DISTRIBUTION OF
THE ORTALIDÆ.

	PAGE
<i>FIRST DIVISION: ORTALIDÆ HAVING THE FIRST LONGI- TUDINAL VEIN BRISTLY OR HAIRY</i>	32
	PAGE
FIRST SECTION: PYRGOTINA	33
THIRD SECTION: CEPHALINA	48
SECOND SECTION: PLATYSTO- MINA	36
FOURTH SECTION: ORTALINA	52
FIFTH SECTION: PTEROCAL- LINA	58
<i>SECOND DIVISION: ORTALIDÆ HAVING THE FIRST LONGI- TUDINAL VEIN BARE</i>	63
FIRST SECTION: ULIDINA	64
SECOND SECTION: RICHARDINA	66

PART THIRD.—THE NORTH AMERICAN
ORTALIDÆ.

<i>FIRST DIVISION: ORTALIDÆ HAVING THE FIRST LONGI- TUDINAL VEIN BRISTLY OR HAIRY</i>	72
FIRST SECTION: PYRGOTINA	72
4. variabilis n. sp. ♂	91
Gen. I. PYRGOTA Wied.	72
5. flavimana n. sp. ♂. ♀	92
1. millepunctata Lw. ♀	73
6. micans n. sp. ♀	94
2. nudata Wied. ♂. ♀	77
7. pallida n. sp. ♂. ♀	95
3. vespertilio Gerst. ♂	79
4. pterophorina Gerst. ♀	81
Gen. IV. STENOPTERINA Macq.	96
1. cærulescens n. sp. ♂	97
Gen. V. MISCHOGASTER Macq.	98
SECOND SECTION: PLATYSTO- MINA	83
Gen. VI. MYRMECOMYIA R. Descr.	99
Gen. I. AMPHICNEPHES NOV. gen.	83
1. myrmecoides Lw. ♂. ♀	100
1. pertusus nov. sp. ♂. ♀	84
Gen. II. HIMERÖESSA n. g.	85
1. pretiosa n. sp. ♂	85
Gen. III. RIVELLIA R. Descr.	87
1. conjuncta n. sp. ♀	88
2. viridulans R. Descr. ♂. ♀	88
3. quadrifasciata Macq. ♂	90
THIRD SECTION: CEPHALINA	102
Gen. I. TRITONA n. g.	102
1. flexa Wied. ♂. ♀	102
2. inenrva n. sp. ♂. ♀	104
3. cuneata n. sp. ♂. ♀	107
Gen. II. CAMPTONEURA Macq.	108
1. picta Macq. ♂. ♀	109

TABLE OF CONTENTS.

v

	PAGE		PAGE
Gen. III. DIACRITA <i>Gerst.</i>	111	Gen. V. ANACAMPTA <i>Lw.</i>	129
1. costalis <i>Gerst.</i> ♂	111	1. latiuscula <i>Lw.</i> ♂. ♀	130
2. remula n. sp. ♀	114		
Gen. IV. IDANA n. g.	115	Gen. VI. APOSPASMICA n. g.	131
1. marginata <i>Say</i> , ♀	115		
FOURTH SECTION: ORTALINA	118	FIFTH SECTION: PTEROCAL-	
Gen. I. AUTOMOLA n. g.	118	LINA	132
		Gen. I. PTEROCALLA <i>Rondl.</i>	132
Gen. II. TETANOPS <i>Fall.</i>	119	1. strigula n. sp. ♀	133
1. luridipennis n. sp. ♂. ♀	119		
2. integra n. sp. ♂	121	Gen. II. STICTOCEPHALA	134
Gen. III. TEPHRONOTA <i>Lw.</i>	122	1. cribellum n. sp. ♂. ♀	134
1. humilis <i>Lw.</i> ♂. ♀	123	2. cribrum n. sp. ♀	135
		3. corticalis <i>Fitch</i> , in litt. ♂. ♀	136
Gen. IV. CEROXYS <i>Macq.</i>	125	4. van <i>Say</i> , ♀	138
1. obscuricornis n. sp. ♂. ♀	126		
2. ochreicornis n. sp. ♀	126	Gen. III. CALLOPISTRIA n. g.	140
3. similis n. sp. ♀	127	1. annulipes <i>Macq.</i> ♂. ♀	141
4. canus <i>Lw.</i> ♂. ♀	128		
		Gen. IV. MYENNIS <i>R. Descr.</i>	142
		1. scutellaris <i>Wied.</i> ♂. ♀	143
SECOND DIVISION: ORTALIDÆ HAVING THE FIRST LONGITUDINAL VEIN BARE			
FIRST SECTION: ULIDINA	145	4. nitidiventris n. sp. ♀	157
Gen. I. DASYMETOPIA <i>Lw.</i>	145	5. costalis <i>Fbr.</i> ♀	158
Gen. II. OEDOPA <i>Lw.</i>	146	6. quaternaria <i>Lw.</i> ♀	159
1. capito <i>Lw.</i> ♂. ♀	146	7. binotata <i>Lw.</i> ♂	160
		8. annonæ <i>Fbr.</i> ♂. ♀	162
Gen. III. NOTOGRAMMA <i>Lw.</i>	148	9. Thomæ <i>Lw.</i> ♂. ♀	163
1. stigma <i>Fbr.</i>	148	10. abdominalis <i>Lw.</i> ♂. ♀	163
		11. alternans <i>Lw.</i> ♂	165
Gen. IV. EUPHARA <i>Lw.</i>	150	12. stigmatias <i>Lw.</i> ♂. ♀	166
Gen. V. ACROSTICTA <i>Lw.</i>	151	13. elata <i>Lw.</i> ♂. ♀	168
Gen. VI. SEOPTERA <i>Kirby</i>	151		
1. colon <i>Lw.</i> ♂. ♀	152	Gen. VIII. CHÆTOPSIS <i>Lw.</i>	169
		1. aenea <i>Wied.</i> ♂. ♀	170
Gen. VII. EUCESTA <i>Lw.</i>	153	2. debilis <i>Lw.</i> ♀	172
1. spoliata <i>Lw.</i> ♂	154		
2. pusio <i>Lw.</i> ♀	155	Gen. IX. Hypoëta <i>Lw.</i>	173
3. notata <i>Wied.</i> ♂. ♀	156	Gen. X. Stenomyia <i>Lw.</i>	173
		1. tenuis <i>Lw.</i> ♂	174

REVIEW OF THE NORTH AMERICAN TRYPETINA.

	PAGE
Introduction	211
List of the described species	214
Description of the species	216
Analytical table of the smaller genera, adopted for the North American Trypetæ	327
Distribution of the North American Trypetæ among the adopted smaller genera	329
Comparison between the European and the American fauna of Trypetina	331

APPENDIX I.

Critical enumeration of all the North American Trypetina described by other Authors	335
---	-----

APPENDIX II.

Description of the five North American <i>Trypetidæ</i> contained in "Eugenies Rusa, etc.," Zoologi, Part VI	340
--	-----

	PAGE		PAGE
Trypeta hogaster Thoms.	340	Trypeta aurifera Thoms.	343
femoralis Thoms.	342	genalis Thoms.	344
acutangula Thoms.	342		
INDEX			345
EXPLANATION OF THE PLATES			349

NOTE FROM THE TRANSLATOR.

Readers of these "Monographs" may notice discrepancies in some minor points of the terminology used in the first volume from that of the following ones. A few words of explanation are therefore necessary. The first volume was translated from Mr. Loew's German manuscript into English by a German friend of his (see Vol. I, p. v). The second and third volumes were translated either by me, or under my supervision. Although in all essentials the terminology adopted in Vol. I was followed, some changes, which I thought would be improvements, were introduced. Thus, feet was used for *legs*; ovipositor, for *borer*; crossvein, for *transverse vein*; arista, for *antennal bristle*; thus *transverse shoulder vein* came to be humeral crossvein, and *hinder transverse vein*, posterior crossvein, etc. None of these changes can give rise to any error or uncertainty.—O. S.

o
o
l
l
g

d
fi
it
w
b
g

se
th
w
m

DIPTERA
OF
NORTH AMERICA.

PART III.

THE FAMILY ORTALIDÆ,

I.

INTRODUCTION.

IN the variety of forms the *Ortalidæ* are hardly surpassed by any other family of diptera; at the same time, they are hardly equalled by any in the importance of the structural differences occurring among them; hence, they may be considered as one of the most interesting families of the order. Nevertheless, but little has been done as yet for the exact definition of the limits of this family, as well as for its subdivision into smaller groups.

It would be impossible, therefore, to attempt a satisfactory description of the North American species of *Ortalidæ*, without first settling the questions of the true limits of the family, of its relationship to other families, and of the characters upon which it is established. It would also be indispensable to break the family up in subordinate groups and these groups in genera.

Of all these requirements, only one has been, as far as I can see, fulfilled, and that is, the definition of the limits between the *Trypetidæ* and the *Ortalidæ*, which I have tried to draw as well as I could, in the first volume of these monographs and in my earlier work on European *Trypetidæ*.

Through the successive, although disconnected, publications of several authors the systematic arrangement of the *Ortalida*, like that of some other families of diptera, has gradually reached a state of confusion which it is impossible to unravel without a detailed and somewhat lengthy discussion. I may be excused, therefore, if I preface the description of a comparatively small number of American species by an unusually long introductory chapter.

In order to point out the relationship of the *Ortalida* with other families, as well as the differences which distinguish them, it is necessary, first of all, to discover those characters which serve to define the family.

THE EUROPEAN ORTALIDÆ.

The examination of the works of Meigen, the founder of systematic dipterology, will afford a solid basis to proceed from, and I will begin with the European species which are the longest known and the best investigated.

Meigen's Genus Ortalis.

The nucleus of the European *Ortalida* is formed of those species which Meigen has brought together in the genus *Ortalis*, a genus which, in his acceptation, far exceeded the limits which we are accustomed to give to genera at present.

1. Characters which Meigen's species of *Ortalis* have in common.

I discovered a number of characters which the species of this genus have in common, and which also occur in many species added to the genus by subsequent authors. These common characters, to the exclusion of those which also belong to most of the neighboring families, are the following:—

Front broad in both sexes; a more or less distinctly developed small callosity runs from each corner of the vertex down the front; it bears in front of the lateral bristle of the vertex one or two distinct additional bristles; beyond this, the front is beset with only short hairs, or else quite bare; it never has the second row of bristles, nearer the orbit, which distinguishes all the genera of *Trypetida*.

Eyes bare, even under a strong lens; the compound microscope shows only some sparse, very short hairs.

The frontal fissure is distinct, but the frontal lunule is not pushed up as far as the front, so as to seem to form a part of it; on the contrary, it appears as the upper part of the face.

Vibrissæ, such as they appear in all the genera of *Helomyzidae*, are not extant.

The clypeus is always distinct. The palpi are rather broad; the proboscis more or less stout.

The metathorax is larger than usual, very much projecting inferiorly and posteriorly.

The feet short and strong, rather than long and slender; middle tibiæ distinctly spurred; front and hind tibiæ spurless; the erect bristle extant in several families of the *Diptera acalyp-tera* on the upper side of the tibiæ, near their end, for instance in the *Supromyzidae* and *Sciomyzidae*, does not occur here in any species.

The abdomen of the male has four segments, the first of which, like the first segment of the female abdomen, is formed of two coalescent segments; the diminutive fifth segment of the male abdomen forms the small, more or less imbedded hypopygium; the sometimes more filiform, in other instances tape-like, rolled up penis is of an extraordinary length.

The abdomen of the female consists of five segments; the sixth, seventh, and eighth segments are converted into a flattened, extensile ovipositor, the first joint of which surpasses the two following in breadth very much, and is often colored like the rest of the abdomen; the slender last joint of the ovipositor ends in a simple point.

The wings show the complete venation of the *Diptera acalyp-tera*; the auxiliary vein is entirely separated from the first longitudinal vein, although it is sometimes rather approximated to it; it ends at a more or less acute angle in the costal vein, without becoming less distinct at its end; the first longitudinal vein is provided with bristles, at least at its end; the two posterior basal cells are comparatively large.

2. Species erroneously placed in the genus *Ortalis*.

The agreement of all the species placed by Meigen in the genus *Ortalis* would have been complete if *O. parviflora* (*fulminans* M.), *connera*, *vibrans*, and *syngenesia* did not show differences, which require a special mention.

O. parviflora and *connera* differ from the other species in

the presence of a row of bristles on each side near the orbit, besides the bristles common to all the *Ortalidæ*. These bristles are a character so exclusively peculiar to the *Trypetidæ* that we cannot but consider those two species as belonging to that family (as I have already proved it elsewhere). They cannot, therefore, be further considered here.

Ortalis vibrans, the female of which has only four segments on the abdomen, approaches, in the absence of bristles upon the first longitudinal vein, *Palloptera* and the related genera so much, that one might be tempted to place it among the *Pallopteridæ*; but there are other genera having the first longitudinal vein bare, to which *Ortalis vibrans* is still more closely allied, and which, as I will have occasion to show hereafter, cannot possibly be separated from the *Ortalidæ*. Such being the case, *O. vibrans* has to remain in this family, and this is also justified by the large size of the two posterior cells in this species, which is a proof of its relationship to the other *Ortalidæ*. There is no doubt, at the same time, that this species is a stranger in Meigen's genus *Ortalis*.

Ortalis syngenesia is also distinguished from the other species of *Ortalis* by its abdomen, which has only four segments in the female; in other respects it is more related to them than *O. vibrans*; it is more closely allied to the species of the genus *Platystoma* than to the other species of Meigen's *Ortalis*.

The other European Ortalidæ.

After having thus disposed of those species placed by Meigen in the genus *Ortalis*, which have either to be entirely withdrawn from the family of *Ortalidæ*, or which can only conditionally be received in it, the next step to be taken, in order to chalk out the whole extent of the family *Ortalidæ*, is to discover such other genera as may likewise possess the characters common to the species of *Ortalis*. After this, we will have to point out such genera as possess not all, but most of those characters only, and especially the principal ones; and thus we will reach a limit beyond which only such genera will be found, as, on account of important differences from the species of *Ortalis*, cannot any more be united in one family with them. This research has also to show us which among the characters common to the above enumerated species of *Ortalis* have to be

stricken out, or at least to be modified, in order to leave, as a residue, the true characters of the family *Ortalidæ*.

Here also I begin with the European fauna, as the genera and species composing it are by far the best known.

The variously organized groups of species, within the genus *Ortalis*, each have, outside of this genus, a circle of relationship of their own.

1. Forms reminding of *Ortalis lamed*.

If we begin with *Ortalis lamed* (*pulchella* Meig.), we are led at once towards *Sciomyza bucephala* Meig., which Macquart has united, with several other heterogeneous species, in the genus *Orites*, and for which I have later established the genus *Cormocaris*.

Cormocaris brings us to *Tetanops*, which agrees in its principal characters with *Cormocaris bucephala*, quite erroneously placed in the genus *Sciomyza* by Meigen. In this species, as well as in all the European species of *Tetanops* which I know of, none of the characters are wanting the presence of which distinguishes the genus *Ortalis*.

The genus *Tetanops* again leads us towards *Dorycera*; the remarkable elongation of the second antennal joint is a peculiar character of most species of this genus, a character not to be found in the species of *Ortalis*. However, the difference in the length of this joint in different species of *Dorycera* sufficiently shows that too much systematic stress ought not to be laid upon this character; all the other characters agreeing with those of the *Ortalidæ*, *Dorycera* must necessarily be placed in this family.

Next to *Dorycera* I find the genus *Adapsilia*, founded by Waga, which, like most *Doryceræ*, has an elongated second antennal joint. It is distinguished by a very projecting front, very approximated antennæ, and the want of ocelli; with the species of *Ortalis* it agrees in the characters already specified, except that the first joint of the ovipositor of the female is not flattened, as in all the species of *Ortalis*, but capsule-shaped, swollen; as, however, in other respects the structure of this ovipositor resembles that of *Ortalis*, *Adapsilia* must also be added to the *Ortalidæ*. At the same time, the statement concerning the shape of the ovipositor of this family must be somewhat modified to be applicable to *Adapsilia*.

I know of no other European genus which, although still more distant from *Ortalis* in the direction of *Adapsilia*, would nevertheless be admissible into the family of *Ortalidæ*.

2. Forms reminding of *Ortalis syngenesiæ*.

If, in our search for forms related to *Ortalis*, we start from *Ortalis syngenesiæ*, distinguished by its four-jointed female abdomen, the genera *Ulidia*, *Timia* and *Platystoma* at once claim our attention.

Ulidia, in Meigen's sense, is not a homogeneous genus. *Ulidia demandata* is too aberrant to remain in it. Together with several exotic species allied to it, it has to form a separate genus for which *Chrysomyza*, a name already used by Fallen for *Ulidia demandata*, may be applied.

Timia apicalis, described by Meigen, is nothing but an *Ulidia*, and must be referred to this genus; the differences which appear in Meigen's statements and his figures do not exist in nature.

Timia erythrocephala, upon which Wiedemann, in the *Analecta*, has founded the genus *Timia*, differs from *Ulidia* only in its extreme glabrousness, its swollen head, much more projecting beyond the eyes in profile, and perhaps also the somewhat less developed clypeus; in all the other important characters both genera agree.

In all the species hitherto placed in the genera *Timia* and *Ulidia*, and consequently also in the species of *Chrysomyza*, the first longitudinal vein is bare. In all other respects, these species share all the characters common to the species of *Ortalis*, so that, in my opinion, their position among the *Ortalidæ* cannot well be disputed, unless we separate from this family all the species the first longitudinal vein of which is bare. Nevertheless, the relationship between the species of *Ulidia*, *Timia*, and *Chrysomyza* to *Ortalis syngenesiæ* cannot be considered as unusually close, because they differ from it, not only in the bareness of the first longitudinal vein, but also in the presence of a fifth, very much abbreviated, segment of the female abdomen.

A genus agreeing with them in the bareness of the first longitudinal vein, and most closely related to them, is the genus *Empylocera*, introduced by me.

The genus *Lonchæa* also seems related to *Ulidia*; I will, therefore, in the sequel, explain its systematic location.

The species of *Platystoma* differ somewhat from *Ortalis* in the

shape of the ovipositor: its first joint is smaller, narrower, and somewhat less flattened; generally also more withdrawn in the last abdominal segment. The hypopygium, formed by the upper half of the fifth abdominal segment, is unusually small; whether the penis has the shape of an unrolled tube or thread I cannot ascertain at present, as I have no fresh specimens at hand, but I have every reason to suppose that such is the case, as the female ovipositor, in its structure, is absolutely analogous to that of the species of *Ortalis*. The agreement of all other characters compels us to admit *Platystoma* among the *Ortalidæ*; and this genus really shows, in the four-jointed abdomen of the female, the absence of bristles on the pleuræ and an analogous structure of the mouth, a close relationship to *Ortalis syngenesiæ*.

I know of no other European genera which, in following the same direction of relationship, might be still more distant from *Ortalis* than the species of *Platystoma* are, and which, nevertheless, would show a sufficient agreement with the *Ortalidæ* to be placed among them. I, therefore, hold *Platystoma* to be one the more distant genera, placed on the extreme limit of the family.

3. Forms reminding of *Ortalis paludum*.

Species like *Ortalis paludum*, *luctuosa*, and others of the same group, remind of the genera *Psairoptera* and *Cephalia*.

The comparatively low head, the transversely oval eyes, and the small development of the clypeus give *Psairoptera* a very peculiar appearance; nevertheless in all the other important characters it agrees with the species of *Ortalis* so well, that its position among the *Ortalidæ* cannot be disputed, although its precise location within this family may not be very easy to determine. The relationship of *Psairoptera* with the above-named species of *Ortalis*, far from being a close one, can rather be called distant.

In *Cephalia* I cannot discover a single character which would justify its separation from the *Ortalidæ*. To place this genus among the *Sepsidæ* seems to me utterly impracticable, as the distinctive character of the latter family, the rudimentary structure of the palpi, must be maintained, unless we render the limits of the family altogether doubtful. Moreover, *Cephalia* does not show any vestige of vibrissæ which the *Sepsidæ* possess, and more than all, the structure of the ovipositor of *Cephalia* is like

that of the *Ortalidæ*, and not like that of the *Sepsidæ*. *Cephalia* is more closely related to the above-named species of *Ortalis* than *Psairoptera*.

4. Forms not resembling any of Meigen's species.

A fly which possesses all the essential characters of the species of *Ortalis*, and undoubtedly belongs to the *Ortalidæ*, is the *Scatophaga fasciata* of Fabricius, erroneously placed by Meigen in the genus *Trypeta*. The *Musca octopunctata* of Coquebert, Dec. III., Tab. XXIV., is probably identical with it. The circumstance that there is no other European *Ortalida* resembling this species probably caused Meigen to overlook its true relationship. Similar forms are more frequent in other parts of the world, especially in America. Among them I will name *Dictya ocellata* Fabr., *Ortalis van* Say, and *Platystoma annulipes* Macq., which, by the way, is no *Platystoma* at all.

Aciphorea not belonging to the ORTALIDÆ.

That group of genera which, on account of its peculiar, three-jointed, horny ovipositor, ending in a simple point, has been called, and not improperly, *Diptera aciphorea*, is represented in Europe, besides the *Trypetidæ* and those genera which, on the preceding pages, I have claimed for the *Ortalidæ*, only by *Lanchoæa*, with the genus *Earomyia*, separated from it by Zetterstedt, and by *Palloptera* and *Toxoneura*. As it would be useless to look for *Ortalidæ* outside of the *Diptera aciphorea*, it remains for us at present to define the position of those genera with regard to the *Ortalidæ*.

The number and position of the frontal bristles, the distinctness of the clypeus, the absence of vibrissæ, and the want of the characteristic bristle on the upper side of the tibia, before its end, which is distinctive of several families, the spurred middle tibia, the spurless front and hind tibiæ, as well as the completeness of the venation, clearly prove the close relationship of these genera with *Ortalis*. They are less closely allied to the *Trypetidæ*, from which they differ in the arrangement of the frontal bristles and in the course of the auxiliary vein. All four differ from all the genera, the location of which among the *Ortalidæ* I have proved in the preceding discussion, by a much smaller size of the two posterior, usually called small, basal cells

of the wings, and all four agree among themselves in the absence of bristles on the first longitudinal vein, and this character they share with some of the genera placed among the *Ortalidæ*.

The genera *Paloptera* and *Toxonevra* possess moreover another striking character, which occurs also among some few of the genera of *Ortalidæ*, but in none to that marked extent; they have, upon the middle of the otherwise short-pilose, thoracic dorsum, as far as its anterior part, a series of binary bristles, distinguished by their length, stoutness, and regular arrangement. The difference in the venation already spoken of, together with this marked peculiarity in the arrangement of the bristles of the thorax, seem to afford sufficient ground for excluding those two genera from the family *Ortalidæ*. I consider them as the nucleus of a separate family, which I call *Palopteridæ*.

The systematic position of *Lonchæa* is more difficult to decide upon than that of *Paloptera* and *Toxonevra*. While the venation of *Lonchæa* closely approaches these two genera, the position of the bristles on its thorax is more like that of many *Ortalidæ*, as there are no stronger bristles on the thoracic dorsum, anterior to the suture; this genus stands therefore nearer to the undoubted *Ortalidæ* than *Paloptera* or *Toxonevra*. Against its being united with the *Ortalidæ* may be urged (not to mention the smallness of the two posterior basal cells), not so much those characters which are common to all *Lonchææ*, as a number of peculiarities, which do not occur among the *Ortalidæ*, and which distinguish different species of *Lonchæa*, and are quite proper to form the basis of a subdivision of this widespread and rather numerous genus. As such characters I consider the long and strong hairs upon the whole body of some species, the long and dense pubescence of the eyes of others, the partial coalescence of the auxiliary vein with the first longitudinal in several, and finally the circumstance that in the females of some species the sixth abdominal segment does not take part in the formation of the ovipositor quite in the same manner as among the *Ortalidæ*. I am afraid that the *Ortalidæ*, as a family, would lose too many of their well-defined characters, if, in order to accommodate *Lonchæa* among them, we undertook to modify these characters in accordance with the above mentioned peculiarities of the latter genus. The nature of the

venation of the wings having proved, in so many cases, to be the most trustworthy character for the distinction of the families of diptera, we have to take care not to attach too little importance to the smallness of the posterior basal cells in *Lonchæa*, cells which, in the *Ortalidæ*, always are of a considerable size. These reasons induce me to exclude *Lonchæa* from among the *Ortalidæ*. Those entomologists who take the European fauna alone in consideration, will, I have no doubt, justify this course, as that fauna does not contain any intermediate forms between *Lonchæa* and the genera of *Ortalidæ*, but I am not quite as sure of the approbation of those who have a wide acquaintance with the diptera from all parts of the world, because, among the number, forms occur which seem to be intermediate between *Lonchæa* and the genera of *Ortalidæ* allied to *Ulidia*, and it is possible that the discovery of a large number of such forms may, at some future time, render the exclusion of *Lonchæa* from the *Ortalidæ* less plausible than it appears to me now. In the first volume of these monographs, I placed this genus in the family of the *Pallopteridæ* and considered it as the typical genus of a second group in this family. Whether this arrangement, which I for the present retain, is satisfactory, or whether it would not be better to take *Lonchæa* as the typical genus of a separate, small family, intermediate between the *Pallopteridæ* and the *Ortalidæ*, is beyond the scope of the present discussion, and may, therefore, be left for future investigation.

The genus *Earomyia* is so near *Lonchæa*, that, with regard to its systematic position, whatever I said of the latter may be applied to the former.

Summary of the European Ortalidæ.

From what precedes may be deduced the following list of genera and species of European diptera, which I place in the family of *Ortalidæ*: all the species of *Ortalis*, in Meigen's sense, with the exception of *O. pæciloptera* and *connexa*; *Sciomyza bucephala*; the genera *Adapsilia*, *Dorycera*, *Tetanops*, *Psairoptera*, *Cephalia*, *Plotystoma*, *Timia*, *Ulidia*, *Chrysomyza*, *Empyclocera*, and, finally, *Trypeta fasciata*.

THE ORTALIDÆ OTHER THAN EUROPEAN.

I will now try to find the genera and species from other parts of the world than Europe, which must be placed in the family *Ortalidæ*.

(a.) In Wiedemann.

I begin by the *Ortalidæ* contained in Wiedemann's writings. Besides his species of the genus *Ortalis*, against the location of two of which, however, I will have to raise some doubts, and besides his *Timia erythrocephala*, which occurs in the southeast of Europe and in the neighboring provinces of Asia, the species of *Cephalia* described by him undoubtedly belong to the *Ortalidæ*. They differ somewhat from the European *Cephalia rufipes*, and belong in the relationship of those species which Rob. Desvoidy distributed among his genera *Polistodes* and *Myrmecomyia*: Mr. Macquart has established for them the genus *Michogaster* (better *Mischogaster*), which coincides with the genus *Conopsidea*, introduced by him at a later time.

The two *Ortalis* of Wiedemann, the systematic position of which seems doubtful to me, are *Ortalis trifasciata* and *atomaria*, both from Brazil, both closely related to each other, and somewhat reminding, in their general appearance, of *Richardia* and *Rhopalomera*. Both have an erect bristle before the end of the tibiæ, which I cannot take for anything else but the præapical bristle, wanting in all the *Ortalidæ*. Considering the importance which the presence or absence of this bristle has in the classification of all the *Diptera acalyptera*, I would be very much inclined to exclude both of those species from the family *Ortalidæ*, if I could assign them a fitting place in some other family. The structure of the ovipositor clearly proves that they belong in the circle of the *Diptera aciphorea*, but even in this wider circle the existence of their, however weak, præapical bristle assigns them a rather isolated position. The venation and the position of the frontal bristles, in which they agree with the *Ortalidæ*, do not allow their introduction among the *Trypetidæ*. They have still less connection with the *Pallopteridæ*. Hence, nothing remains to be done, as it seems, but to tolerate them in the family *Ortalidæ*, however unwelcome they may be among them, as, in consequence of their appearance, the absence of a

principal bristle ceases to be an undoubted distinctive character of the *Ortalidæ*. That these two species, to which several undescribed South American forms have to be added, are to form the nucleus of a new genus is beyond question. I propose for it the name of *Autonola*.

Whether the two species of Fabricius, which Wiedemann quotes among the species of *Ulidia*, really belong to this genus cannot be decided without seeing the original specimens; but I have no doubt that they belong to the *Ortalidæ*. I would suppose that *Ulidia stigma* belonged to the genus *Notogramma*, and that *Ulidia ænea* is a *Chrysonomyza*.

Wiedemann's genus *Pyrgota*, with which *Orycephala*, Macq. is absolutely identical, shows all the characters of the *Ortalidæ*. It is closely allied to *Adapsilia*, the only difference being that the antennal foveæ are shorter, while in *Adapsilia* they are parallel, and run down to the edge of the mouth; but, as in different species of *Pyrgota* these foveæ vary in length, this difference has so little importance that *Adapsilia* might, without any inconvenience, be united with *Pyrgota*.

The genus *Dacus*, in Wiedemann's writings, is a mixture of many very different forms of diptera, most of which are *Ortalidæ* and two species are *Trypetidæ*. Two of the species of which Wiedemann formed the first section of the genus *Dacus*, form now, together with other species added since, the genus *Steopterina*, which Macquart established under the name of *Senopterina*, and which he placed quite correctly among the *Ortalidæ*. The *Dacus flavicornis*, placed by Wiedemann in the first division as a third species, has a certain general resemblance to the two former species; it differs, however, in the bareness of the first longitudinal vein and in several other characters, too much to be united in the same genus with them; nevertheless, this species, as well as the two others, belong to the *Ortalidæ*. Among the species of Wiedemann's second division of *Dacus* *D. succinctus* must be referred to the *Ortalidæ*; it belongs in the immediate relationship of *O. syngenesiæ*. *Dacus bicolor* likewise belongs to the *Ortalidæ*. The remaining *Dacus* of Wiedemann's second division are *Trypetidæ*; some of them belong to the genus *Trypeta*, if we take it in the wider sense of Meigen and Wiedemann; for instance, *Dacus parallelus*, fra-

terculus, serpentinus; the greatest part of the residue are species which may be left in the genus *Dacus*.

On the other hand, Wiedemann has placed in the genus *Trypeta* several species which do not belong to the *Trypetidæ* at all and have all the characters of the *Ortalidæ*. Such species are: *Trypeta ocellata*, which Macquart described again as a supposed new species, under the name of *Platystoma ocellata*, and upon which Rondani established later the genus *Pterocalla*: *Trypeta obscura*, which is very closely allied to the former, and which Macquart very improperly placed in the genus *Camptoneura*, while its place is in the genus *Pterocalla*, next to *P. ocellata*; moreover *Trypeta picta*, the typical species of the Ortalideous genus *Camptoneura*: *Trypeta flexa*, which may be placed in the genus *Mischogaster*; *Trypeta trimaculata*, re-described by Macquart as *Carlometopia ferruginea*; *Trypeta cyanogaster, basilaris, scutellaris*, and perhaps several others among Wiedemann's *Trypetæ*, which I have not had the occasion to compare.

Those species which Wiedemann placed in the genus *Platystoma*, with the exception of his *Platystoma decora*, really belong to that genus, and consequently to the *Ortalidæ*. *Platystoma decora*, which induced Macquart to establish the genus *Loxoneura*, is also to be placed among the *Ortalidæ*.

Tetanops sanguineiceps was described by Wiedemann from a specimen of the Berlin Museum; I have seen this species, unless my memory deceives me, not in the Berlin Museum, but in Wiedemann's collection. I found that in the structure of the head and in the venation it does not sufficiently agree with the European species of *Tetanops* to be left in the same genus with them, but, at the same time, that it undoubtedly belongs to the family of the *Ortalidæ*. I am sure that the *Dichromyia brasiliensis* of Rob. Desvoidy, described as the type of the new genus *Dichromyia*, is the same species.

The *Scatophaga bispinosa* Fab., placed by Wiedemann in the genus *Tetanocera*, differs from the other *Ortalidæ* in the venation as well as in the shape of the scutellum very much, but nevertheless, judging from Wiedemann's statements, and especially from his figure, it undoubtedly belongs in that family, where Macquart also places it in establishing for it the genus *Notacanthina*. Should we judge, however, from Macquart's

figure (in the *Diptères Exotiques*, II., 3, Tab. xxviii., fig. 8), we would not place it among the *Ortalidæ*, as it shows distinctly spurred front and hind tibiæ; these spurs, however, as well as many other things in Mæquart's figures, are probably productions of the draughtsman's fancy. In the most slovenly figure of the same species in Mæquart's *Suites à Buffon*, no such spurs are to be found.

That *Dacus podagricus* Fab., placed by Wiedemann in *Cordylura*, does not belong to that genus, nor to the *Cordyluridæ* in general, has been recognized long ago. For this species, as well as for similar ones, the genus *Richardia* has been established by Rob. Desvoidy in the family of the *Ortalidæ*.

The systematic location of *Dictya externa* Fab. cannot well be ascertained, owing to the insufficient statements of Fabricius as well as of Wiedemann; the latter are in an insoluble contradiction to Wiedemann's figure in what regards the shape of the head and the picture on the thorax; judging by the figure, it would seem that the fly does not belong to the *Diptera acalyptera* at all.

The genus *Rhopalomea*, Wied. seems to have been by all later authors unhesitatingly referred to the *Ortalidæ*. I look upon this decision as far from unobjectionable, but can very well conceive that a certain resemblance in the shape of the head between the species of *Rhopalomea* and *Platystoma* (with the genera allied to it), may easily have given rise to such an opinion. The species of *Rhopalomea* differ in a striking manner from all the *Ortalidæ* in the structure of the hypopygium of the male, while in this respect they show a most decided leaning towards the *Sapromyzidæ*, *Sciomyzidæ*, and the families immediately connected with them. The females are not provided with a borer-like ovipositor, composed of elongated, retractile joints; the metanotum is but very little developed, less than usual among the *Ortalidæ*; the front and middle tibiæ have, on the upper side, before their end, an erect bristle; upon the upper side of the hind tibiæ, this bristle, in most species, is not distinctly visible among the general pilosity of the tibia; nevertheless, it is easily recognizable in some species, for instance *Rhopalomea pleuropunctata* Wied. Such are the characters which, in my opinion, not only render the location of *Rhopalomea* among the *Ortalidæ* doubtful, but even impossible. If, among the

diptera I am acquainted with, I look for the immediate connections of *Rhopalomera*, I find them unmistakably among the South African species of the genus *Cestrotus*, erected by me. Before all, the striking structure of the head, reminding partly of some genera of *Ephydriidæ*, partly of the *Ortalidæ*, is very much alike in both genera; this resemblance extends to the mode of pilosity of the face, the shape of the antennæ, and the feathery pubescence of the arista. Moreover, the small development of the metanotum, the shape of the hypopygium, and the structure of the last segments of the female abdomen are very much alike. Now, as the genus *Cestrotus*, through the intermediate steps of *Prosoptomya* and *Physogenia*, approaches the family of the *Sapromyzidæ* very closely, I do not find any serious objection to placing *Rhopalomera* in the same family. That *Rhopalomera* is one of the extreme genera of the family cannot be doubtful; the size of the two posterior basal cells especially distinguishes it from all the other genera of *Sapromyzidæ* in a very striking manner, and connects it with the *Sciomyzidæ*; for this reason it would be also possible, by slightly modifying the definition of the boundary between those two families, to place *Rhopalomera* among the *Sciomyzidæ*. Those who will not share either of these two views, and prefer to place among the *Ortalidæ* a genus which is far apart from all the *Diptera aciphorea* in the structure of the ovipositor, may locate *Rhopalomera* in the vicinity of *Richardia*, on account of the bareness of the first longitudinal vein, the rounded end of the posterior basal cell, and the spines on the femora.

Thus, the following diptera, described in Wiedemann's works, belong to the *Ortalidæ*: his species of *Ortalis*; all the species which he brings in the genera *Timia*, *Ulidia*, *Cephalia*, *Platystoma*, *Tetanops*, and *Pyrgota*; in his genus *Dacus*, the three species in the first division, and *Dacus succinctus* and *bicolor* in the second; in the genus *Trypeta*, *Trypeta ocellata*, *obscura*, *picta*, *flava*, *trimaculata*, *basilaris*, *cyanogaster*, and *scutellaris*; in the genus *Tetanocera*, his *T. bispinosa*, and finally, his *Cordylura podagrica*.

(b.) *In Robineau Desvoidy.*

I turn now, not without reluctance, to the writings of R. Desvoidy. In his well-known *Essai sur les Myodaires* he united

the genera which we are considering under the general name of *Phytomydæ Myodinae*. This generalization may be considered as successful, as it contains but little which is foreign, that is which would be better placed among his *Aciphoreæ*, equivalent to the family *Trypetidæ*, and as at the same time it excludes but little of what really belongs to the *Ortalidæ*. The position also which Rob. Desvoidy assigns to the *Phytomydæ Myodinae*, next to the *Phytomydæ Thelidomydæ*, that is, the *Micropozidæ*, cannot but be sustained, as the latter are closely related to the *Ortalidæ*. After his *Phytomydæ Thelidomydæ* Rob. Desvoidy places his *Aciphoreæ*, that is, the *Trypetidæ*, while he would have done better in reversing this order of his two divisions, on account of the close relationship between the *Ortalidæ* and the *Trypetidæ*. The sovereign neglect of all previous publications, the wretched manner in which most of his genera are established, chiefly upon merely relative differences (for instance, a somewhat longer third antennal joint, a somewhat more pubescent arista, etc.), without regard to the most striking plastic characters, the very slovenly description of many species of unknown *habitat*, etc., have, long ago, put this author's writings in such bad repute that it would not be easy to add anything to it. It would be unjust, however, after this fully deserved blame, not to recognize that Rob. Desvoidy's judgment, with regard to questions of relationship, in this case, as in many others, was a very correct one.

The genera which he places among the *Phytomydæ Myodinae* are: *Dichromya*, *Palpomya*, *Hesyquillia*, *Heramya*, *Myoris*, *Oscinis*, *Blainvillia*, *Meckelia*, *Meliera*, *Myennis*, *Strauzia*, *Vidalia*, *Delphinia*, *Acidia*, *Myrmeconmya*, *Polystodes*, *Stylophora*, *Herina*, *Myodina*, *Richardia*, *Rivellia*, *Boisduvalia*, *Clidonia*, *Setellia*, *Chlorophora*. Concerning these genera and their names, I will offer the following remarks:—

The genus *Dichromya* (the name ought to be improved to *Dichromyia*) is adopted by Macquart in his *Diptères Exotiques*, and placed among his *Heteromyzides*. The *Dichromyia brasiliensis* of Rob. Desvoidy is the same as the *Platystoma microcera* of Macquart's *Suites à Buffon*, and was described still earlier as *Tetanops sanguineiceps* by Wiedemann. Not being a *Tetanops* this species must therefore be considered as the type of the genus *Dichromyia*. The position among the *Ortalidæ*,

assigned to it by R. Desvoidy, I hold to be correct; with *Platystoma* it has nothing to do.

The genus *Palpomya*, a hybrid name, being formed out of a Latin and a Greek word, and not rendered more valuable by its improvement in *Palpomyia*, is identical with *Platystoma*; the typical *Palpomyia Lalandi* is nothing else but the well-known *Platystoma asphaltina* Wied. The generic characters given by R. Desvoidy are entirely erroneous.

Under the name of *Hesyquillia* Rob. Desvoidy describes *Platystoma seminationis* Fab., and under that of *Hesyquillia lugubris* the *Platystoma umbrarum* Fab.; thus, the genus *Hesyquillia* likewise coincides with *Platystoma*.

The genus *Heramyia*, which ought at least to be called *Heramyia*, is based upon *Sciomyza bucephala*, which R. Desvoidy did not recognize, as well as upon another species which is very like it, if not identical. Macquart united this species with *Myoris* (a name which it is difficult to explain), a genus not distinguished by a single character of any value, and with *Blainvillia* (a preoccupied name), and thus formed his genus *Otilis* (a name which Latreille had already used in a broader sense); but he placed in it moreover some true *Sciomyzidæ*.

The genus *Oscinis*, as understood by R. Desvoidy, is identical with *Dorycera*; it has nothing in common with the genus of the same name to which Fallen reduced the much more comprehensive genus *Oscinis* of Latreille.

Meckelia (an already preoccupied name) and *Meliera* (probably also a dedication name), contain species belonging to Macquart's Ortalideous genus *Cerorhys*.

The genus *Myennis* (a badly formed name), is established for *Scatophaga fasciata* Fab., which Macquart, in the *Suiles à Buffon*, describes as *Ortalis fasciata*, after Rob. Desvoidy, and, for a second time, as *Tephritis fasciata*, after Meigen.

Strauzia (as the genus is dedicated to Strauss-Dürkheim, the name should be spelt *Straussia*) does not belong to the *Ortalidæ* at all, but to the *Trypetidæ*; the two species described by Rob. Desvoidy are nothing else but the male and female of *Trypeta longipennis* Wied., which Rob. Desvoidy did not recognize.

Vidalia seems likewise to belong to the *Trypetidæ*; not

having succeeded yet, however, in identifying the species, I am not positive about it.

The genus *Delphinia* is established for *Trypeta picta*, Fab., which Rob. Desvoidy did not recognize; the unbecoming generic name was afterwards replaced by *Camptoneura* Macq.

The genus *Acidia* belongs to the *Trypetidæ*.

Myrmecomya (more correctly *Myrmecomyia*) and *Polystodes* (better *Polistoides*) taken together nearly correspond to the genus *Michogaster* (better *Mischogaster*) of Macquart, placed by the latter among the *Sepsidæ*. The size of the palpi and the structure of the ovipositor do not justify this location, and the genus undoubtedly belongs to the *Ortalidæ*.

Of the position of the genus *Stylophora* in the system I cannot judge, not knowing the species upon which it is based.

Herina (the derivation of the name is not apparent) comprises species from the relationship of *Ortalis paludum*.

The genus *Myodina* (again a name of obscure derivation) is based upon *Ortalis vibrans*, which R. Desvoidy took for *Ortalis urticæ*. Macquart, in the *Suites à Buffon*, very erroneously united this genus with *Ortalis*, throwing together various very different species. Long before Rob. Desvoidy, Kirby had used for *Ortalis vibrans* the generic name of *Seioptera*.

Richardia is founded either upon *Dacus podagricus* Fab., not recognized by Rob. Desvoidy, or else on some closely allied species.

Rivellia (probably a dedication name) contains species related to *Ortalis syngenesisæ*, and among them this very species, as usual, not recognized by Rob. Desvoidy. Macquart in the *Suites à Buffon* unites *Rivellia* with *Herina*, while the species really belonging to it are put in the genus *Urophora*, or even in *Platystoma*; and upon one of them, in his later works, he even establishes a new genus, *Epidesma*.

Whether the genus *Boisduvalia* really differs from the preceding only in the length of the third antennal joint seems very doubtful; should this be the case, the separation of these two genera would not be justified.

Clidonia is considered by the author himself as belonging to quite a different family, in which we will not contradict him.

Setellia seems to contain *Ortalidæ* resembling *Micropezidæ* in their general appearance.

Chlorophora may also belong there, as Rob. Desvoidy especially mentions its relationship to *Setellia*.

The following among Rob. Desvoidy's genera belong therefore to the *Ortalidæ*: *Dichromyia*, *Palpomyia*, *Hesychyia*, *Heramyia*, *Myoris*, *Oscinis*, *Blainvillia*, *Meckelia*, *Melieria*, *Myennis*, *Delphinia*, *Myrmecomyia*, *Polistoides*, *Herina*, *Myodina*, *Richardia*, *Rivellia*, *Boisduvallia*. Very probably *Setellia* and *Chlorophora* have to be added to them. The systematic position of *Stylophora* is doubtful. Genera not belonging to the *Ortalidæ* are: *Straussia*, *Vidalia*, *Acidia*, *Clidonia*.

(c.) *In Macquart.*

During his long career as an entomological writer, Macquart has several times changed his views with regard to the classification of the *Diptera acalyptera*, as was to be expected from the great difficulty of the subject. His opinion, however, on the extent of the family *Ortalidæ* has, during that time, undergone but little change. As, strictly speaking, he is the only writer who has attempted to establish a general system of the diptera, embracing all parts of the world, I consider it as my duty to give a detailed account of his views, the more so as they differ from mine in a not unimportant manner. To attain this end I will enumerate all those of his families, with their genera, which, according to my opinion, contain genera belonging to the *Ortalidæ*, as well as to the families closely connected with them, for instance, *Palloptra*, *Toxoneura*, *Lonchæa*. In order to show the progress made by Macquart during his dipterological studies I will give this in a twofold manner, that is, first after the *Suites à Buffon* and next after the *Diptères Exotiques*. Those genera which I consider as undoubtedly *Ortalideous* I have marked with an exclamation; those doubtfully introduced into this family I have designated by an interrogation. The genera related to the *Ortalidæ*, which I have united in the family *Palloptra*, I have inclosed in brackets; the same I have done with the genus *Sapromyza*, because Macquart does not separate the species of *Palloptra* from the *Sapromyzæ*, although the typical *Sapromyzæ* have no relationship whatever with the *Ortalidæ*.

The review of the part of the system above alluded to, from the *Suites à Buffon*, is as follows:—

<i>Scatomyzidæ.</i>	! Amethysa,	<i>Thyreophoridae.</i>
Scatophaga,	! Notacanthina,	Thyreophora.
Dryomyza,	Rhopalomera,	
(Sapromyza,) ! Eurypalpus,		<i>Leptopoditæ.</i>
(Toxoneura,) ! Platystoma,		Tanypeza,
Sciomyza, ! Loxoneura.		Calobata,
Lucina,		Teniaptera,
Helomyza, <i>Tephritidæ.</i>		Micropeza,
Blephariptera,	Dacus,	Nerius,
Heteromyza.	Leptoxys,	Longina,
	Bactrocera,	! Setellia.
	! Scuopterina,	
<i>Psilomydæ.</i>	Petalophora,	<i>Ulidini.</i>
Orygma,	Urophora,	Actora,
Trigonometopus,	Terellia,	Cœlopa,
Eurina,	Tephritis,	Gymnopoda,
Psilomyia,	Acinia,	Lipara,
! Tetanops,	Ensiua.	! Ulidia.
! Pyrgota,		
! Otites,	<i>Sepsidæ.</i>	<i>Lauxanidæ.</i>
Platycephala,	Sepsis,	Lauxania,
! Dorycera.	Chelligaster,	Pachycerina,
	Nemopoda,	(Lonchæa,)
<i>Ortalidæ.</i>	! Cephalia,	(Teremyia,)
! Herina,	! Michogaster,	Pterodontia,
! Ortalis,	Diopsis.	Celyphus.
! Ceroxys,		
Clettamia,		

In the *Diptères Exotiques* the corresponding part of the system assumes the following shape, about which I have only to observe that in this work Macquart brings in only those genera in which he intended to describe, or at least to mention, exotic species; the genera *Toxoneura*, *Lucina*, *Tetanops*, *Otites*, *Platycephala*, etc., although not mentioned in this list, ought, in order to render it complete, to be transferred to it from the former.

<i>Scatomyzidæ.</i>	Sciomyza,	<i>Ortalidæ.</i>
Scatophaga.	Helomyza,	! Oxycephala,
	Curtonotum.	! Loxoneura,
<i>Sciomyzidæ.</i>		! Platystoma,
Dryomyza,	<i>Psilomydæ.</i>	! Camptoneura,
Tapeigaster,	! Eumetopia,	! Heterogaster,
(Sapromyza,) Ectecephala,		Rhopalomera,
Physegenua,	! Dorycera.	! Euripalpus,

! <i>Eniconeura</i> ,	<i>Cardiacera</i> ,	<i>Sepsis</i> .
<i>Cleitamia</i> ,	<i>Dacus</i> ,	
! <i>Richardia</i> ,	! <i>Meracantha</i> ,	<i>Diopsidæ</i> .
! <i>Senopterina</i> ,	<i>Bactrocera</i> ,	<i>Diopsis</i> .
! <i>Herina</i> ,	<i>Enicoptera</i> ,	
! <i>Epidesma</i> ,	<i>Ceratitis</i> ,	<i>Leptopoditæ</i> .
! <i>Ceroxys</i> ,	<i>Acanthoneura</i> ,	<i>Longina</i> ,
! <i>Ortalis</i> ,	<i>Urophora</i> ,	<i>Nerius</i> ,
! <i>Amethysa</i> ,	! <i>Toxura</i> ,	<i>Cardiacera</i> ,
! <i>Lamprogaster</i> ,	<i>Tephritis</i> ,	<i>Calobata</i> ,
! <i>Euprosopia</i> ,	<i>Terellia</i> ,	<i>Toxopoda</i> ,
! <i>Cœlometopia</i> ,	<i>Acinia</i> ,	<i>Tanypeza</i> ,
! <i>Notacanthina</i> ,	! <i>Epicerella</i> ,	! <i>Setellia</i> .
! <i>Cruphiocera</i> ,	<i>Ensina</i> .	
! <i>Plagiocéphala</i> ,		<i>Lauxanidæ</i> .
! <i>Campigaster</i> .	<i>Sepsidæ</i> .	(<i>Lonchæa</i> .)
	! <i>Cephalia</i> ,	<i>Lauxania</i> ,
<i>Tephritidæ</i> .	! <i>Omalocéphala</i> ,	! <i>Ulidia</i> ,
! <i>Odontomera</i> ,	! <i>Conopsida</i> ,	<i>Zygothrica</i> ,
<i>Leptoxys</i> ,	<i>Nemopoda</i> ,	<i>Celyphus</i> .

In the *Diptères Exotiques*, after the families I have enumerated the *Helomyzidæ* and *Geomyzidæ* follow, and after them the

Heteromyzidæ.

Heteromyza,
Actora,
 ! *Dichromyia*,
Cœlopa.

In examining the systematic distribution, introduced by Macquart in the *Suites à Buffon*, we soon find that as early as that work, he had, if not a definite knowledge, at least a correct instinct of the true characters of the *Ortalidæ*, less correct, however, than Rob. Desvoidy, who wrote before him.

Those genera which, in that work, he united in the family *Ortalidæ* really belong to it, with the exception of *Rhopalomera* and, very probably, of *Cleitamia*; the latter genus seems to be hardly distinct from *Henicoptera*, which belongs to the *Trypetidæ*.

A double error seems to be contained in the separation of the genera *Tetanops*, *Pyrgota*, *Otitis*, and *Dorycera* from the *Ortalidæ* and their combination with *Orygma*, *Trigonometopus*, *Eurina*, *Psilomyia*, and *Platycephala* into one family, the *Psi-*

lomydæ. Their relationship with the *Ortalidæ* is evident. Among the genera which Macquart places in one family with them, *Eurina* and *Platycephala* belong to the *Oscinidæ*, each of the others to some other dipterous family; none shows any close relationship to the *Ortalidæ*. In the *Diptères Exotiques* Macquart has in part corrected this error, as at least *Oxycephala*, of the identity of which with *Pyrgota* he was not aware, is put among the *Ortalidæ*.

A second error is that the ortalideous genus *Stenopterina* (Macquart incorrectly writes *Senopterina*) has been placed in his family *Tephritidæ*. In the *Diptères Exotiques* Macquart has amended this error.

A third mistake consists in Macquart having placed in his genus *Urophora* several species which do not at all belong to his family *Tephritidæ*; his *Urophora quadrivittata*, *fulvifrons*, and several others, are true *Ortalidæ*.

Fourthly, the position of the genera *Cephalia* and *Michogaster* (better *Mischogastra*, or at least *Mischogaster*) among the *Scpsidæ* cannot be sustained. As has been observed already, we agree with Rob. Desvoidy in considering both as true *Ortalidæ* on account of the large development of the palpi as well as of the structure of the ovipositor.

Neither can I, in the fifth place, agree with Macquart in putting *Setellia* among his *Leptopodidæ*; I refer it also to the *Ortalidæ*, and this once more in agreement with Rob. Desvoidy.

A sixth error is the great interval between *Ulidia* and the other *Ortalidæ*, as well as the whole composition of the family *Ulidini*. *Lipara*, with which Macquart's genus *Gymnopoda* is synonymous, belongs to the *Oscinidæ*; *Cælopa* and *Actora* do not belong to the same family, neither with *Lipara*, nor with *Ulidia*, nor together. In the *Diptères Exotiques* Macquart did rightly in dropping altogether the ill-conceived family of *Ulidinæ*.

I will not expatiate here on the incorrectness of the position of *Palloptera*, *Toxoneura*, *Lonchæa*, and *Teremyia* (established for *Lonchæa laticornis*), as this inquiry is of no especial importance to us.

It is easy to perceive that the system is improved in the *Diptères Exotiques*; but even here *Dorycera* is misplaced among the *Psilomydæ*, together with *Eumetopia* (which belongs to the *Ortalidæ*).

In his family *Tephritidæ* the genus *Odontomera* is established, which is closely related to *Cælometopia* on one side and *Setellia* on the other, and must therefore be transferred to the *Ortalidæ*.

The same may be said of the genus *Meracantha*, the true place of which is in the vicinity of *Odontomera*, *Setellia*, *Cælometopia*, *Richardia*, etc.

The genus *Toxura*, judging from the published figure, also belongs to the *Ortalidæ*, and indeed in the circle of relationship of *Pyrgota*; whether the examination of the insect itself would lead to the same result I do not pretend to affirm, as I have not seen it.

The figure of the head of *Epicerella* (*Dipt. Exot.*, Suppl. iv., Tab. xxvii.) might perhaps justify the supposition that the genus belongs to the *Ortalidæ*; nevertheless I think it more probable either that the frontal bristles, characteristic of the *Trypetidæ*, were broken off in Macquart's specimen, or that they have been omitted in the drawing. Thus I do not dare to express any opinion as to the correctness of the position assigned by Macquart to this genus.

Cephalia, in the *Diptères Exotiques*, is likewise put among the *Sepsidæ* instead of among the *Ortalidæ*.

Omalocephala (better *Homalocephala*, at all events, a preoccupied name) seems to belong in the vicinity of *Setellia*, *Cælometopia*, etc., that is, in the family *Ortalidæ*.

The genus *Conopsidea*, as Macquart informs us, is founded upon *Cephalia femoralis* Wied.; in the *Suites à Buffon*, this same and two more species gave him occasion to establish the genus *Michogaster*. If these two data be correct, as we have every reason to suppose, *Conopsidea* would be a synonym of *Michogaster*; the emendation of the incorrectly formed name *Conopsidea* thus becomes useless.

The erroneous location of *Setellia* at the end of the *Leptopoditæ* is preserved.

Ulidia is transferred to the family *Lauxanidæ*, where it is a perfect stranger.

About the systematic position of *Zygothrica* (not *Zygotricha*, as Gray, in the *Animal Kingdom*, spoils, in trying to improve it), a genus already proposed by Wiedemann in his essay on *Achias*, I can only form an opinion from the statements of Wiedemann and Macquart on the typical species, *Z. dispar*, as well as from

their figures. It seems to me that this species may belong to the *Drosophilidæ*. In the Berlin Museum there is a little fly which apparently belongs to this genus; I have not been able to ascertain whether this species is *Z. dispar*, but I have seen enough not to doubt in the least of its belonging to the *Drosophilidæ*.

Dichromyia is wrongly placed by Macquart among the *Heteromyzidæ*, between *Actora* and *Calopa*. I will maintain for the present its position among the *Ortalidæ*, although I cannot deny that a better place might perhaps be found for it; however, no such place has been pointed out yet. Besides the typical species, *Dichromyia sanguiniceps*, Macquart has another species from Africa, which, as I will show hereafter, cannot well belong to this genus.

About the genera which Macquart, in the *Diptères Exotiques*, places in the family *Ortalidæ*, I will make the following remarks:—

Oxycephala, as was mentioned before, is identical with *Pyrgota*.

Loxoneura is established for *Platystoma decora*.

Platystoma is misused for the location of a number of heterogeneous forms; whatever had broad wings, with a dark picture, among the rest a *Trypeta*, was taken by Macquart for a *Platystoma*.

Camptoneura is a true ortalideous genus, based upon *Trypeta picta* Wied., and, as observed above, identical with *Delphinia* Rob. Desv. Macquart has likewise used this genus for the introduction of species not belonging there at all, for instance, of *Trypeta obscura* Wied.

Heterogaster (a preoccupied name) is a well founded genus in the neighborhood of *Pyrgota*.

Euripalpus (a hybrid name), judging from Macquart's data, belongs to the *Ortalidæ*.

The genus *Eniconeura* (better *Heniconeura*) is said to be distinguished by its spurless middle tibiæ. If such were really the case the genus could not belong to the *Ortalidæ*, nor to any of the allied families. But in *Heniconeura fenestralis* Macq., I perceive at the end of the middle tibiæ a rather strong spur, which is closely applied to the tarsus when the latter is stretched

out. There cannot be any doubt, therefore, that the genus really belongs to the *Ortalidæ*.

Richardia, in the *Diptères Exotiques*, is with good reason entirely separated from *Herina*, with which it was united in the *Suites à Buffon*.

Senopterina (I have already corrected the name to *Stenopterina*) has been placed here where it really belongs, among the *Ortalidæ*.

Herina is a mixture of heterogeneous forms, which must be generically kept apart.

Epidisma is probably synonymous with *Rivellia* R. Desv.; moreover, Macquart has placed species of the latter genus under the head of *Herina*, of *Urophora*, and even of *Ceroxys*.

Ceroxys is a rather well founded genus, established at the expense of *Ortalis* Meig. But in the *Diptères Exotiques* Macquart adds species to it which do not at all share its characters; for instance *Ceroxys cærulea*, etc. It almost seems, in such instances, that he mistakes this genus for another.

The genus *Ortalis* is a mixture of heterogeneous species; how is it possible to crowd together in one and the same genus such species as *Ortalis ornata* Meig., *fasciata* Fab., *connexa* Fab., *frondescentiæ* Lin., *vibrans* Lin., and even the *Ortalis dentipes* Macq., said to be provided with spurs on the hind tibiæ? Either Macquart has not known these species or he has not closely examined them, otherwise he could not possibly have committed such a mistake; how very confused his ideas about the systematic position of these species was, appears from the fact that he described *Scatophaga fasciata* Fab. as an *Ortalis*, and for a second time as *Tephritis* and that *Dictya connexa* Fab. even appears three times in his writings, as *Cordylura*, as *Ortalis*, and as *Tephritis*! (*Dipt. Exot.*, Suppl. iv., p. 292, *Tephritis dorsalis*.)

The true characters of *Amethysa* are not to be gathered from Macquart's definition of this genus. As the name alludes to the color of the African species, upon which the genus is established, it should be improved to *Amethysta*.

Lamprogaster is a well founded genus; but the species belonging to it show considerable differences in their organization which would fully justify a subdivision in several genera. It belongs in the vicinity of *Platystoma*.

Cælonetopia seems to be founded on *Trypeta trimaculata* Fab., which Macquart did not identify; it is closely allied to *Odontomera* and *Setellia*.

Euprosopia undoubtedly belongs to the *Ortalidæ*.

Notacanthina is founded upon *Tetanocera bispinosa* Fab.

The figure of the head of *Crithiocera* (better *Cryphiocera*) seems to indicate that the species would be better placed in some other part of the system, as it has strong bristles on the forehead; the other characters, however, prove that its location among the *Ortalidæ* cannot well be called in doubt.

The position of *Plagioccephala* among the *Ortalidæ* likewise cannot be doubted; it seems closely related to *Richardia*, which also contains broad headed species.

Campigaster (a frightful compound) is undoubtedly well placed among the *Ortalidæ*, but the name cannot be preserved in its present shape.

(d.) *In Walker.*

Although Macquart's publications do not always define with sufficient precision the systematic position of the genera introduced by him, this position could, in most cases, be made out, and moreover, the attempt, on his part, of a systematic distribution is always apparent. Walker's publications on exotic diptera do not, unfortunately, deserve this praise. The systematic department, as well as everything else in them, is treated with the same superficial carelessness. In most cases it would be impossible to make out, from his statements, the real place in the system which the genera, introduced by him, must occupy, unless they were accompanied, as is often the case, by the excellent figures of Westwood. These usually furnish the necessary data concerning the relationship of the new genera; they would have done so in all cases if Westwood's attention had been directed to the sometimes very minute characters which are used in the classification of the diptera and especially of the *Diptera acalyptera*; the fact that in the majority of cases these characters are reproduced in the figures, give a most brilliant proof of the accuracy of Westwood's drawings, and of his keen perception.

Walker's publications in the *List of Dipt. Ins. of the Brit. Mus.*, and in the *Insecta Saundersiana*, do not raise our expectations very high, as the *Ortalidæ* and *Trypetidæ* are mingled together

generally; forms such as *Camptoneura picta* Fab., *Trypeta arcuata* Walk., *T. albovuria* Walk., *T. excepta* Walk., etc., are certainly no *Trypetidæ*! In Walker's later publications, the systematic confusion is still greater. As far as I can ascertain, among the genera published in the latter, *Adrana*, *Brea*, *Valonia*, are *Ortalidæ*; the two latter belong in the vicinity of *Platystoma*. The genera *Themara*, *Strumeta*, *Sophira*, and *Riora* belong to the *Trypetidæ*. The genus *Xangelina* is closely related to *Physogenia*, perhaps identical with it, and hence, has to be placed among the *Sapromyzidæ*. The position of the genus *Xiria* remains doubtful, even in the presence of Westwood's figure; it shows some characters which make one doubt that it belongs to the *Diptera aciphorea* at all. The genera *Duomyia* and *Chromatomyia*, which, taken together, seem to correspond to *Lamprogaster* Macq., and *Zona*, which is apparently identical with *Loxoneura*, are *Ortalidæ*; Walker, in the *List of Dipt. Ins.* etc., has erroneously placed them among the *Tachinidæ*, together with *Trigonostoma*, which likewise belongs to the *Ortalidæ* (however, he corrected this error in one of his later publications.)

(e.) In Bigot, Gerstæcker, Doleschall and Saunders.

In recent times it is to Bigot and Gerstæcker that the increase of our knowledge of exotic diptera is principally due.

The genera *Terastomyia*, *Maria*, *Agastrodes*, and *Pterogenia*, established by Bigot, belong to the *Ortalidæ*. *Elassogaster* likewise, although placed among the *Helomyzidæ* by Bigot, must be referred to the *Ortalidæ*. His genus *Elaphromyia*, on the contrary, if description and figure be correct, belongs to the *Trypetidæ*.

Gerstæcker has established the ortalideous genera, *Phytalmia*, *Gorgopis*, *Toxotrypana*, and *Diacrita*, and described several new species of *Richardia* and *Mischogaster*. *Phytalmia* has a synonym in Saunders's genus *Elaphomyia* (*Elaphomyia* Wallacei Saund. = *Phytalmia megalotis* Gerst.; *Elaphomyia cervicornis* Saund. = *Phytalmia cervicornis* Gerst.). The genus *Gorgopis* seems, as the author himself supposes, to be synonymous with *Zygenula paradoxa*, described somewhat earlier by Doleschall. If in the genus *Toxotrypana* the outer row of frontal bristles is really wanting, and it thus should really belong to the *Ortalidæ*, the not flattened ovipositor of this genus would place it in the

neighborhood of *Pyrgota*, with which it also agrees in the small development of the clypeus. However, the ocelli are fully developed, and the structure of the head is rather like that of the true species of *Dacus*, as *Dacus oleæ*, etc., so that it might perhaps be considered as a genus of this group, in which, in conformity to the striking shortness of all the hairs of the body, the lateral bristles of the front have disappeared. This supposition seems confirmed by the scutellum which has only two bristles at the tip; and the uncovered last abdominal segment of the female, which is generally wanting in the *Dacina*, or is altogether concealed under the preceding segment, is not a positive objection, as this segment is very much abbreviated and much less horny than the preceding ones, and thus can very easily be concealed in the living insect.

Among the scattered publications of various authors many forms may be found which belong to the *Ortaliidæ*. I purposely omit what I know of them, especially the gradually published species of the genera already discussed by me. It is not in my power to collect the residue, and I doubt whether such a work would materially alter the limits of the family *Ortaliidæ* as they have resulted from the preceding discussion.

NATURAL CHARACTERS OF THE FAMILY ORTALIIDÆ.

If we ask now what we have to erase or to modify in the characters of the original genus *Ortalis*, in Meigen's and Wiedemann's sense, in order to obtain the characters defining the whole family, the answer will be that it is very little indeed. In the first place, the mention of the pilosity of the front must be modified a little, as there are genera among the *Ortaliidæ* which have no other bristle before the bristles of the vertex. Next to that, the description of the structure of the feet has to be changed thus, that in most genera they are short and strong, but in some rather elongate. In the third place, the statement about the female abdomen must be modified by saying that it has generally five segments, but that the fifth is very often shortened and concealed under the fourth, and that, in some cases, it entirely disappears, and then the abdomen has only four segments. In the fourth place, the introduction of *Pyrgota* and of the related genera in the family, requires a modification in the statement about the structure of the ovipositor, which is not flattened here; the chief

stress in this statement should be laid upon the remainder of the structure, which is the same in all the genera. In the fifth place, the mention of the bristles on the first longitudinal vein should not be admitted in the definition of the family.

The definition of the *Ortalidæ* can therefore be put in the following manner: Front broad in both sexes; on both sides of the vertex a more or less developed swelling runs down the front, upon which, before the bristle of the vertex, one or two erect bristles are inserted, which, however, are wanting in some genera. Otherwise the front has only the ordinary pubescence, or is quite bare, but never provided with a second row of strong bristles along the orbit, even when the hairs on both sides of the *vitta frontalis* almost require, in some few genera, the character of bristles. Frontal fissure distinct; frontal lunule never pushed so far up as to appear to be a part of the front; even in those genera in which, on account of the great curvature of the frontal fissure, as in *Ædopa*, the lunule happens to lie higher than the antennæ, it always distinctly appears as a part of the face; in many genera it is not distinguishable from the face. The vibrissæ are always wanting. The eyes are bare. The clypeus is always distinct, of various size, usually well developed. Proboscis more or less stout. Palpi rather broad, often very broad, very seldom narrow. Metanotum larger than usual, strongly projecting posteriorly and inferiorly. Feet generally rather stout and short, in some genera, however, of a considerable, although not striking, length and slenderness. Middle tibiæ distinctly spurred; front and hind tibiæ spurless; no erect preapical bristle before the end of the upper side of the tibiæ. The abdomen of the male has four segments, however the first consists of two confluent segments, which is also the case in the females; the but little developed fifth segment represents a small, more or less imbedded hypopygium; the tape-like or thread-like penis is of an extraordinary length, rolled up in a spiral. The female abdomen consists of five segments, the fifth of which is often very much abbreviated, sometimes wanting, so that the abdomen of the female then seems to consist of only four segments; the sixth, seventh, and eighth segments of the abdomen are converted into the three telescope-like, extensile joints of the ovipositor, ending in a simple, hairless point; in most cases the ovipositor is flattened, and then its first joint often differs but little in its nature

and coloring from the preceding abdominal segments. The wings show the complete venation of the *Diptera acalyptera*; the auxiliary vein is entirely separated from the first longitudinal vein, although often very much approximated to it; it runs into the costa at a more or less acute angle, without becoming indistinct at its end; the two posterior, so-called small basal cells, are of a rather considerable size.

RELATIONSHIP OF THE ORTALIDÆ.

The great variety of forms occurring among the *Ortalidæ* accounts for the number of their near or distant connections among other families. A relationship of the first degree, which finds its most distinct expression in the similarity of the structure of the male hypopygium and of the female ovipositor, connects them with the *Trypetidæ* and the *Pallopteridæ*, as well as these two families with each other. All three form a very close circle of relationship, the members of which have very similar habits.

The *Ortalidæ* differ from the *Trypetidæ* in the absence of a second, external row of frontal bristles, and in the course of the auxiliary vein, which, in the *Trypetidæ*, is obliterated at the end and turns rather abruptly, at a more or less right angle, towards the costa.

From the *Pallopteridæ*, the *Ortalidæ* differ in the more considerable size of the two posterior basal cells.

A relationship of the second degree connects the *Ortalidæ* with the *Sepsidæ* and *Calobatidæ*, as well as these families with each other. Both differ from the *Ortalidæ* distinctly in the structure of the male hypopygium and the want of a horny, three-jointed ovipositor, ending in a simple, hairless point. The *Sepsidæ* differ moreover in their rudimentary palpi from the *Ortalidæ*, as well as from the *Calobatidæ*.

With those of the closely related families which, among their characters, have an erect preapical bristle before the tip of the tibiæ, and, at the same time, do not have any vibrissæ, that is, with the *Sapromyzidæ* and *Sciomyzidæ*, the *Ortalidæ* have only a very distant relationship. I would have left it unmentioned if the genus *Rhopalomera*, which I consider as belonging to the *Sapromyzidæ*, had not been placed among the *Ortalidæ*. The presence of an erect bristle before the end of the tibiæ, the different structure of the hypopygium in the male, the absence of an

ovipositor, similar to that of the *Ortalidæ*, sufficiently distinguish the *Sapromyzidæ* and *Sciomyzidæ*.

DIAGNOSTIC OR ARTIFICIAL DEFINITION OF THE ORTALIDÆ.

The statements about the relationship of the *Ortalidæ* prove that the following characters are sufficient to distinguish this family from all the others, in other words, to constitute its artificial definition.

Male with a rolled-up, long penis; female with a three-jointed, horny ovipositor, ending in a simple point. Front without a second lateral row of bristles. No vibrissæ. Complete venation of the *Diptera acalyptera*; auxiliary vein distinct to its very tip, ending in the costa at an acute angle; the two posterior basal cells large. The middle tibiæ alone are provided with spurs; all the tibiæ are without an erect bristle before the end of their upper side.

II.

SYSTEMATIC DISTRIBUTION OF THE ORTALIDÆ.

THE last, but not the easiest, task which it remains for me to fulfil is the systematic distribution of the family *Ortalidæ*. In attempting it, I will principally confine myself to those genera and species which I possess in my own collection. Only in exceptional instances, and with especial caution, will I allow myself to transgress the limit of what I have, or have had, before me, as the statements concerning the other genera and species which have been published are seldom complete enough to afford the necessary data for the discrimination of their position in the system.

In order to obtain a preliminary survey I first divide the *Ortalidæ* in two large divisions; to the first belong those which have the first longitudinal vein beset with bristles or hairs; to the second, those the first longitudinal vein of which is bare.

FIRST DIVISION.

ORTALIDÆ WITH A BRISTLY OR HAIRY FIRST LONGITUDINAL VEIN.

Among the European *Ortalidæ* of this division five diverging forms will easily be noticed: 1. *Adapsilia*; 2. *Ortalis* Meig., of course to the exclusion of *O. syngenesiæ* and *vibrans*; 3. *Platystoma*; 4. *Cephalia*; and 5. *Scatophaga fasciata* Fab.

All the other European genera with a bristly first longitudinal vein can be grouped around these five types, with the exception perhaps of the somewhat recalcitrant genus *Psairoptera*. The same may be said of all the exotic *Ortalidæ* of this division which I know of. Thus, the *Ortalidæ* of the first division may be naturally divided into five groups.

We will characterize these groups only after having made out the genera which belong to them, and we will proceed to the discovery of these genera by means of the principal characters which distinguish the above-mentioned five types.

Adapsilia shows a striking character, distinguishing it from all the others, in the absence of ocelli and the not flattened ovipositor.

Scatophaga fasciata with its broad and low head, the circular shape of its third antennal joint, and the considerable distance intervening between the end of the auxiliary vein and that of the first longitudinal, has a general appearance which differs from the four other types so much that for a long time the close relationship of this species with the others was, for this reason, misunderstood.

Ortalis, *Platystoma*, and *Cephalia* differ in a very marked way in the mode of insertion of the bristles upon the pleuræ.

Ortalis has a strong bristle immediately above the basis of the fore coxæ; this bristle is not extant in *Cephalia* and *Platystoma*.

Cephalia has above the middle coxæ, but below the longitudinal suture of the pleuræ, a strong bristle, which is also present in *Ortalis*, but entirely wanting in *Platystoma*. If, for the sake of brevity, I call the first *prothoracic*, the second *mesothoracic bristle*, the difference between these three genera will be as follows: *Ortalis* has a prothoracic and a mesothoracic bristle; *Cephalia* has the mesothoracic bristle only; in *Platystoma* both are wanting.

First Section: Pyrgotina.

I borrow the name of this group from the genus *Pyrgota* Wied., to which *Adapsilia* is most closely related. Both genera agree in the absence of ocelli, in the projecting front, the prolonged second antennal joint, the retreating face, the comparatively but little developed clypeus, the prolongation of the first abdominal segment in both sexes, and the contraction of the following segments in the female, as well as in the capsule-shaped structure of the first joint of the ovipositor, and in several other subordinate characters.

The principal difference between these genera consists in the structure of the antennal foveæ, which, in *Adapsilia*, run down in a parallel direction as far as the edge of the mouth, and are separated by a straight ridge, while in *Pyrgota* they end at some distance from the edge of the mouth, and are more or less coalescent.

The South-African genus, *Hypotyphla*, founded by me, agrees

with *Pyrgota* and *Adapsilia* in the want of ocelli, and resembles *Adapsilia* very much in the structure of the face; but it differs in the but inconsiderable elongation of the first abdominal segment, in the greater length of the other segments of the female abdomen, and especially in the long, elongated-conical, but not flattened ovipositor.

Judging by the figure which Macquart gives of his *Toxura maculipennis*, I must suppose with a considerable degree of probability, that it likewise belongs in this circle of relationship.

I have no doubt that the interesting genus *Toxotrypana* Gerst., if placed in the family *Ortaliidæ*, would find its location in the section *Pyrgolina*, on account of its not flattened ovipositor, its hairy first longitudinal vein, and the small development of its clypeus. The presence of ocelli, the enormous length of the ovipositor, and the elongation of the posterior angle of the anal cell into a very long lobe distinguish this genus from the other genera of the group in a most marked manner. I have already alluded to the fact that this genus shows some characters which would seem to justify its location not among the *Ortaliidæ* at all, but among the *Trypetidæ* of the group *Dacina*.

Mr. Macquart has established the genus *Heterogaster* for a South-African species. As the name he gave to this genus was preoccupied a long time ago, I replace it by the name of *Sphenoprosopa*. This genus is very like *Adapsilia* in the structure of the head; in the profile it projects considerably in front of the eyes: the middle of the face forms a high and straight ridge descending perpendicularly; alongside of it the antennal foveæ, which are further from the middle than usual, descend perpendicularly to the edge of the mouth. The cheeks are very broad. The oral opening is very small, the clypeus but little developed, and the proboscis not incrassated. *Sphenoprosopa* differs from *Adapsilia*, *Pyrgota*, and *Hypotyphla* by the presence of distinct ocelli, the great elongation of the third antennal joint, which nearly reaches the edge of the mouth, the enormous development of the last segment of the abdomen of the male, very approximated cross-veins, very parallel longitudinal veins, and a not acute posterior angle of the anal cell. The first and third longitudinal veins are distinctly bristly. I have no doubt that *Sphenoprosopa* belongs to the *Pyrgolina*, although, on the other hand, I must acknowledge that several of the above-

quoted characters seem to point towards a relationship with *Platystoma*. But I am prevented from laying much stress upon them by the small development of the clypeus and the not incrassated proboscis, characters which are not usual in the circle of relationship of *Platystoma*.

The typical species of the genus *Dichromyia*, proposed by Rob. Desvoidy, is Wiedemann's *Tetanops sanguiniceps* from Brazil. Macquart afterwards described a second species, *Dichromyia caffra*. I cannot approve of these two species being united in the same genus. The front of *D. caffra* is much shorter, and anteriorly it does not project as much in the profile as in *D. sanguiniceps*; moreover the ocelli are wanting here, while the other species has them, and the vertical diameter of their eyes is much longer than the horizontal, while in *D. sanguiniceps* the horizontal diameter exceeds the vertical; the scutellum is convex, and the tegulæ very large, while *D. sanguiniceps* has a flat scutellum and small tegulæ. Whether the longitudinal veins are beset with bristles in the same manner in both species or not, I cannot state positively; in *D. caffra* the first and third veins are very distinctly beset with hairs; in *D. sanguiniceps*, if I remember right, the first vein is beset with a hardly perceptible pubescence, but I cannot positively affirm that such is the case. But without insisting upon this difference, the others are sufficient to justify a generic separation. For this reason I have established for *D. caffra* Macq. a species generally found on an offensively smelling plant, the new genus *Bromophila*.

As to the final decision about the place of the American genus *Dichromyia*, I must leave it in abeyance until I have an opportunity to examine both sexes of *D. sanguiniceps*.

The ovipositor of the species of *Bromophila* is much more retracted than in the other genera of the present group; and although not flattened, it is not at all incrassated; unfortunately I have not been able to ascertain on any female specimen whether the ovipositor ends in a simple point, as it seems to me it does. Should this not be the case, the genus would not belong to the *Ortalidæ* at all. At present I cannot find a better place for it than in the neighborhood of *Pyrgota*.

I know of no other genera belonging to the *Pyrgotina*. At present, therefore, the section is composed as follows:—

1. PYRGOTA Wied. ; 2. ADAPSILIA Waga. ; 3. TOXURA Macq. ;

4. HYPOTYPHILA Lw.; 5. TOXOTRYPANA Gerst.; 6. SPHENOPROSOPA, Lw.; 7. BROMOPHILA, Lw.; and 8. DICHROMYIA, R. Desv. The final decision about the location of *Toxotrypana*, *Bromophila*, and *Dichromyia* is, of course, reserved.

The characters common to these eight genera are: oral opening small; proboscis not incrassated; clypeus but little developed; no bristle upon the broad cheeks, and no bristle immediately over the fore coxæ; the first longitudinal vein hairy; the costal vein soon attenuates beyond the end of the third longitudinal vein. The ovipositor is not flattened.

Second Section: Platystomina.

The name of this section is derived from *Platystoma* Meig., the oldest and best known genus in it.

PLATYSTOMA is represented in Europe by a number of closely allied species which must be considered as typical. We may entertain different views on the extent of the genus *Platystoma*, still we would not be justified in introducing in it, as has often been done, species which, in the majority of the most important characters, differ from the European *Platystomæ*. In fact, most of the exotic species, described by different authors as belonging to *Platystoma*, do not belong to it at all.

The *Dictya decora* Fabr., identical with *Tephritis violacea* Gray, and placed by Wiedemann among the *Platystomæ*, has the posterior angle of the anal cell drawn out into a long lobe; this character at once distinguishes this species, not only from *Platystoma* proper, but from all the genera closely related to it. Macquart was right in establishing the new genus LOXONEURA for it. Walker afterwards called it *Zona*. Judging from the figure of the head of *Loxoneura decora*, in profile, given by Macquart in the *Diptères Exotiques*, this genus must belong to the *Platystomina*; the absence of the pro- and mesothoracic bristles, and the only four-jointed abdomen of the female confirm the correctness of this location; the fore femora are spinous.

Whether the South-American *Platystoma stictica* Fab. really belongs to *Platystoma* is very doubtful.

Only a few of the species, placed by Macquart in the genus *Platystoma* really belong there, for instance, none of his American species. *Platystoma fascipennis* and *ocellata* are *Ortalidæ*, but belong to the *Pterocallina*, not to the *Platystomina*. *Platy-*

stoma lunulata belongs, unless the figure of the head is entirely incorrect, to the *Trypetidæ* and not to the *Ortalidæ*. The same may be said of *Platystoma latipennis*, of which Macquart does not give the *habitat*, but which is American.

In the same way as those species of Macquart, Walker's *Platystoma australis*, from Australia, does not belong to this genus. It seems even that not one of the Australian *Platystomæ* hitherto described is a real *Platystoma*, and that this genus is confined to the three old continents.

Should we even confine, as we must necessarily do, the genus *Platystoma* to those species only which agree with the European species in the formation of the head, in the venation, and in the peculiar picture of the wings, we will find species in it which show some, and not unimportant, plastic differences. To the European species, the arista of which has only a short pubescence, may be opposed African species, some of which have the arista perfectly bare, and the scutellum very much swollen, with only four bristles upon it (for instance, *Platystoma asphaltina* Wied.); others, on the contrary, with a feathery arista. The latter are again divided in species, in which, as in the European species, the scutellum has six bristles, and the femora are unarmed (for instance, *Platystoma nigronotata* Lw.); and in such the scutellum of which has four bristles, and the front femora of which, on the under side, towards the tip, are armed with a few little spines. The latter, and among them *P. pectoralis* Lw., differ moreover from the former in the usually more metallic coloring of the conspicuously broad abdomen, the upper half segments of which have a much harder consistency than in the other species; and besides, in such species, the two parts of the first abdominal segment, which represent the first two abdominal segments of other diptera, are not completely coalescent. It results from the foregoing that *Platystoma* may easily be subdivided in four smaller genera, which can be distinguished by the following characters:—

1. Arista bare; femora unarmed; scutellum swollen, with four bristles; type: *P. asphaltina* Wied.

2. Arista with a very short pubescence; femora unarmed; scutellum moderately convex, with six bristles; type: *P. umbrarum* Lw.

3. Arista feathery; femora unarmed; scutellum moderately convex, with six bristles; type; *P. nigronotata* Lw.

4. Arista feathery; front femora spinous; scutellum but little convex, with four bristles; type: *P. pectoralis* Lw.

As in the remaining parts of the organization there is a great deal of agreement among all the *Platystomæ*, and as at the same time the number of the described species is not large enough to require a further subdivision of the genus, we may leave it undivided for the present. Walker's genus *VALONIA* is closely allied to *Platystoma*. Unfortunately, I possess only a single male of *Valonia complicata* Walk., which, moreover, is not very well preserved. The structure of the head, the thorax, and the feet, as well as the venation, do not show anything which would justify a generic separation from *Platystoma*. The facts that the second longitudinal vein is a little shorter, and more curved forward, and that the small crossvein is a little nearer the end of the discal cell, are evidently not sufficient for such a course. The very much swollen and apparently only too bristly scutellum, as well as the moderate breadth and smooth surface of the upper abdominal segments, would furnish a better ground for a separation from *Platystoma*. At all events, thus much is evident, that *Valonia* does not show any distinctive characters more important than those of the four genera would be in which, as I have shown above, *Platystoma* might be subdivided.

Platystoma cincta, from Port Jackson, described by Guérin (*Voyage de la Coquille*), may be considered as the type of a separate genus, allied to *Platystoma*. Several Australian and African species are closely connected with it. If I remember right, such species are designated in the Berlin Museum by the new generic name of *Pachycephala*. But as a genus *Pachycephalus* exists already, I propose the name *SCHOLASTES*. Such species differ from *Platystoma* in the head being larger, the front much broader, the portion of the face between the foveæ much more excavated, and the much narrower clypeus not protruding; the oeciput likewise is much less swollen, so that the head is much more closely applied to the thorax, and appears entirely sessile in the profile; the thorax is much broader and flatter; the scutellum likewise, much larger and flatter, but with six bristles; the tegulæ are as much developed as in *Platystoma*. The structure of the abdomen and of the feet, as well as the venation,

do not show any important difference from *Platystoma*; still it is worthy of notice that the under side of the front femora is beset with a row of little black bristles, which in the larger species assume the shape of slender spines. The coloring of the body is generally ochre, or ferruginous-yellow, usually with black longitudinal stripes on the thorax; the picture of the wings consists of numerous black spots, which often coalesce into cross-bands. In *Scholastes cinctus* Guér., and the species from Australia allied to it, the first half of the arista is feathery, the second bare, and on the thoracic dorsum there are two rather distant rows of short, but strong bristles. The African *Scholastes*, as the type of which I consider *S. nepticula* Lw., from Guinea, have the whole arista bare and no trace of rows of bristles on the thorax. These characters may afford a ground for dividing *Scholastes* in two genera, in which case the present generic name would have to remain with the genus containing *S. cinctus* Guér.

Another genus, closely related to *Platystoma*, containing, as it seems, exclusively Australian species, is the genus *LAMPROGASTER* Macq., with which *Chromatomyia* Walk. is synonymous. The structure of the thorax, of the abdomen, and of the feet, as well as the venation, are very much in agreement with *Platystoma*. The tegulæ are large, larger than those of most *Platystomæ*. The structure and the arrangement of the bristles of the front are likewise similar to those of *Platystoma*; only the third antennal joint is much longer; not only are the antennal foveæ also longer, but deeper and more sharply defined, on their inside especially; the clypeus is of the same breadth as in *Platystoma*, but not projecting; the palpi usually towards their end are not as broad as in *Platystoma*, and the occiput is less swollen; the scutellum is strikingly swollen and provided with six bristles. The abdomen of all the species is of a brilliant metallic color, which the scutellum and the middle of the thorax often share with it; on the latter, however, the metallic color is generally concealed by the presence of pollinose longitudinal stripes, and of an appressed pubescence of a light color. Otherwise, the coloring of the thorax generally is brown or chestnut-red, which color, in many specimens, also extends over the scutellum; the coloring of the wings consists of a few black spots. All the known species of *Lamprogaster* have unarmed femora and a bare arista;

in general, their plastic characters are so much alike that I cannot point out any peculiarity, among the species I know of, which might give rise to a generic subdivision.

Next to *Lamprogaster* stands a genus of which *Senopterina decora* Macq., from Tasmania, may be considered as the type; I call it EUCHALCOTA. The front is of about the same breadth as in *Lamprogaster*, but is flatter and altogether furrowed-scribulate. The third antennal joint is still longer here; the sharply defined, deep antennal foveæ are once and a half the length of those of *Lamprogaster*, and reach almost altogether as far down as the front part of the lateral edge of the mouth; the arista is beset with a short pubescence near its basis, otherwise bare; the clypeus is perceptibly narrower; the occiput is less swollen, so that the head is more closely applied to the thorax. The latter is strongly built, but not as broad in the region of the wings, and hence, of a more equal breadth; scutellum convex, but not swollen, provided with six bristles. The venation is similar to that of the preceding genera, but differs in the fourth vein being gently curved forward before its end, and in the third vein being gently bent backward, so that the first posterior cell is distinctly attenuated towards its end. The coloring of thorax and abdomen is altogether metallic. In thus defining the characters of the genus, I have taken in consideration some species from Australia, which can very well be placed in the same genus with the above named typical species; nevertheless, they show the following differences: the wings are comparatively longer and without any picture, while in *Euchalcota decora*, the cross-veins have dark borders alongside of them; there are no other bristles in front of the row of bristles along the posterior part of the thoracic dorsum, while in *E. decora*, there are some few shorter and thinner bristles immediately in front of that posterior row. There is no necessity for a generic separation yet. I cannot identify any of my species from Australia in a satisfactory manner; it may be that *Chromatomyia laeta* Walk. belongs here.

It would be difficult to explain why Macquart places *Euchalcota decora* in his genus *Senopterina*. The structure of the face and the shape of the thorax are entirely different. The comparatively narrow abdomen of the male (I have not seen the other sex) is almost the only point of resemblance.

The genus *DUOMYIA*, of Walker, is probably closely allied to the above named two genera. Its definition is too insufficient to enable us to form a trustworthy opinion. The irregularly formed name cannot possibly be preserved.

Two species closely related to each other, belonging to the section *Platystomina*, Macquart (in the *Dipt. Exotiques*) has described as *Tephritis caerulea*, and *strigipennis*. With several other Australian species, very similar to them, they can be united in a genus which may be called *CELETOR*. The very striking characters of this genus are the following: The structure of the body is *Trypeta*-like, with the exception, however, of the head. Front of an equal breadth, very steep and long, so that the antennæ are situated much deeper than in any other genus of the present group; the front is evenly and rather densely pilose; the bristles of the vertex and the lateral bristles, closely approximated to them are rather stout; the ocelli are near the edge of the vertex and closely approximated to each other; the two bristles, which otherwise are near them, are wanting here. Antennæ short, hardly reaching beyond the middle of the face; their third joint of equal breadth, with an acute anterior angle; antennal arista slender, bare. Middle portion of the face concave; clypeus rather broad, projecting; proboscis stout; palpi rather broad towards their end. Eyes very high and narrow; cheeks broad; the lower part of the occiput strongly turgid. Thorax strongly developed, rather of an equal breadth; scutellum turgid, overhanging the perpendicular metathorax, with six bristles. Abdomen with four segments in both sexes, as in the preceding genera; the last segment of the female abdomen generally of a softer consistency. The first joint of the ovipositor flattened, always entirely protruding, suddenly attenuated near the basis, more gradually towards the tip, thus having an almost oval outline. Wings rather large, broad towards the basis; the auxiliary vein lies very near the first longitudinal, its end, however, diverges from it at an obtuse angle towards the costa, and preserves its distinctness and stoutness to the very tip. Otherwise, the venation is not unlike that of *Platystoma*, *Lamprogaster*, etc., only the small crossvein is beyond the last third of the discal cell, a position somewhat reminding of *Valonia* Walk. The coloring of the body is blackish-blue, seldom verging on greenish; the front red or reddish-brown; the lateral borders

with white pollen; wings hyaline; all the four species known to me have crossbands, connected near the anterior margin, thus forming an inverted Δ , and with a third black band, bordering the apex; moreover, near the basis of the wing there is a large spot in the shape of a band, or numerous black spots which form a kind of network, not unlike that of some species of *Petalophora*. This difference in the picture of the wings is accompanied by some plastic differences which, if the number of species were larger, could serve for a subdivision in two genera. Those species which have the large spot in the shape of a crossband near the basis of the wings have at the same time the lateral parts of the face very broad, while they are very narrow in the species which have the picture in the shape of a network; the former have the posterior angle of the anal cell smaller, the latter larger than a right angle, so that in the former, the angle is a large acute one, in the latter, a small obtuse one. One of the species from Australia in my collection, belonging to the second group, is distinguished by the very abnormal structure of the hind tibiae of the male. Among the species already published, besides the two described by Macquart, and mentioned above as typical, *Ortalis trifasciata* Doleschall, from Amboina, may likewise, perhaps, belong to the genus *Celetor*.

Macquart, in the *Diptères Exotiques*, describes as *Eniconeura violacea* a species distinguished by some peculiar characters, which undoubtedly is to be considered as the type of a distinct genus of *Platystomina*. The name *Eniconeura*, or more correctly *Heniconeura*, cannot be retained, as it has been already used by Macquart himself for a genus of *Bombylidæ*. The genus may be called *CLITODOCA*. According to that author it inhabits the East Indies; but this statement may perhaps be erroneous, as I have seen a fly said to be from Guinea, and in which I think I recognize Macquart's species; there is a slight difference in the picture of the wing, as represented on Macquart's figure, but the agreement of the description is perfect, and seems fully to justify my supposition. By all means the species is a *Clitodoca*. The characters of *Clitodoca* may be put down as follows: head large, almost square, with a very short longitudinal diameter; antennæ narrow, descending to the middle of the face; arista with a distinct pubescence; face concave, its lateral portions very narrow; oral opening very large, broader than long; clypeus not

disciform, but representing a swelling of the gula, and hence, reminding of a similar structure in *Loxoneura*, in which it fills the greater part of the oral opening. Proboscis but little swollen; palpi of a moderate breadth. Thorax very stout. Abdomen comparatively very short and narrow, consisting of four segments. Feet long; wings very large; the end of the auxiliary vein almost obliterate; the second longitudinal vein very strongly bisinuate; the third and fourth strongly convergent towards their end; the posterior crossvein very oblique; all the basal cells very long; the anal cell has an acute posterior angle.

Among the species which Wiedemann places in the genus *Ortalis*, there are three closely allied ones, which do neither belong in the genus *Ortalis*, nor in the group *Ortalina*. They have to form a separate genus in the group *Platystomina*, which I will call **ENGISTONEURA**. They are: *Ortalis moerens* Fab., *parallela* Wied., and *lugens* Fab.; *Trypeta albovaria* Walk., may be added as a fourth species, unless it is synonymous with *O. moerens* Fab., which may possibly be the case. The following characters distinguish the genus *Engistoneura*. They are large flies of yellowish coloring, with a very much developed thorax, especially broad between the roots of the wings; its convexity, however, is very small; the abdomen is comparatively small, of a metallic violet color. The structure of the head somewhat reminds of *Dacus*. The antennæ reach the middle or a little below the middle of the face; the long arista is distinctly feathery. The foveæ, which reach a little below the middle of the face, are very sharply defined. Clypeus distinct; proboscis of moderate stoutness, with a but little developed mentum; palpi rather broad. Scutellum large, but little convex, overhanging the metathorax more than in most of the other genera of the *Platystomina*; it has six bristles. Abdomen rather cylindrical. Feet of moderate length and not very strong; the front femora on the under side, in the vicinity of the tip, with a few bristle-like spines. Wings large, rather narrow towards the basis, broad towards the apex; auxiliary veins of moderate length, turning abruptly towards the costal margin, and becoming almost obliterate; the first longitudinal vein approaches closely to the margin beyond the end of the auxiliary vein, and runs alongside of it as far almost as the end of the second longitudinal vein; the third longitudinal vein is strongly bent back-

ward, the fourth vein slightly forward, so that the first posterior cell, very broad in the middle, is rather narrow at the end; the small crossvein is beyond the middle of the discal cell; the two posterior basal cells are of a rather considerable and equal length; the posterior angle of the anal cell is rounded. The extensive picture of the wings forms, in the vicinity of the apex, more or less regular crossbands.

The genus AMPHICNEPHES, which I have established for a North American species, will be characterized in the sequel. It is somewhat like *Platystoma*, but distinguished by the not swollen occiput, the flat scutellum, provided with only four bristles, the broad wings and the striking divergency of the longitudinal veins.

A pretty *Ortalida* from Cuba, which cannot conveniently be placed in any of the existing genera, gave occasion for the establishment of the genus HIMEROËSSA, which I will characterize below among the other North American genera. It is distinguished by the narrowness of the marginal and submarginal cells; moreover, the posterior crossvein is prolonged inside of the first posterior cell.

Ortalis syngenesiæ Linn. is the type of a very well justified genus, existing in Europe, Africa, Asia, and America, which Rob. Desvoidy called RIVELLIA. Although the name is not particularly well chosen, the objections against it are not serious enough for its rejection. Besides the species described by Rob. Desvoidy, the following belong to the genus *Rivellia*: *Trypeta basilaris* Wied., *Dacus succinctus* Wied., *Ceroxys quadrisciata* Macq., *Ortalis Ortoeda* Walk., *Tephritis melliginis* Fitch., and several others. Most of them agree quite well with the species placed in the genus *Rivellia* by Rob. Desvoidy; others, however, show a very gradual transition towards allied forms, which cannot very well be united in the same genus with the typical *Rivelliæ*. Thus Macquart has established for one of them the genus *Epidesma*. The transitions, however, are so gradual, that it is not very easy to decide upon the best boundary for the genus *Rivellia*. *R. viridulans* R. Desv., and all the North American species which I know of, agree in their generic characters with *Rivellia syngenesiæ* completely; the same is the case with *R. basilaris* Wied., and with several *Rivelliæ*, from the southeastern region of Asia, which I possess in my collection;

the only difference shown by the latter species is a somewhat smaller length of the third antennal joint. Next to these are some South African *Rivellia*, for instance, *Rivellia atra* Lw., which have the third antennal joint a little shorter still; all these species, however, cannot be separated from *Rivellia*, as the diminution of the third antennal joint is a very gradual one, not affording any distinct limit for a separation.

Macquart's *EPIDESMA fascipennis*, from the Cape, is likewise but very little different from the typical *Rivellia*. The occiput is somewhat more convex; the third antennal joint has a somewhat sharper anterior angle, the thorax is comparatively a little smaller, and the first section of the fourth longitudinal vein shows but very little of the sinuosity, so characteristic of the true *Rivellia*, and which renders the anterior part of the discal cell more narrow; at the end of the convex scutellum there are two strong bristles; whether the second pair of bristles, which exists in the other *Rivellia*, is wanting here, or whether they were accidentally broken off in the specimen I had before me, I am unable to decide; I rather incline to favor the former supposition. If I am right, *Epidesma* would deserve to be retained as a separate genus; in the opposite case, it would be better to place *Epidesma fascipennis* in the genus *Rivellia*, because then the whole difference between them would merely consist in comparative characters.

Among the species from the southeast of Asia, there are several which are closely allied to *Rivellia*, but differ from the typical species in the greater length and lesser breadth of the marginal cell, a more straight third longitudinal vein, and a hardly perceptible sinuosity of the first section of the fourth vein; moreover, the thorax is less strongly developed, so that their stature shows some, although only a distant, resemblance to the species of *Stenoptera*. They are easily distinguished by the picture of their wings, which is very different from that of the *Rivellia*; it consists in a conspicuous black border along the costal margin and the apex, not unlike that of *Diacrita* and *Molanotoma*, while the *Rivellia*, besides the apex, which is margined with black, also have black crossbands. I propose for this genus the name of *SCOTINOSOMA*.

Species having the first section of the fourth longitudinal vein straight, must, most decidedly, be eliminated from *Rivellia*.

Such is a group of closely related African species, which I unite in the genus ARDELIO. The lateral portions of their face are distinctly broader than in *Rivellia*, the eyes not so high, and the cheeks, for this reason, broader; the clypeus is narrower and the thorax more strongly developed; the convex scutellum has four bristles, like *Rivellia*. They almost show more affinity to *Platystoma* than to *Rivellia*; all the species known to me are black, with longitudinal lines of white dust on the thorax, and their wings have black crossbands, between which, along the costal margin, there are black spots or streaks. The single species show, in the length of the third antennal joint, still more considerable variations than the species of *Rivellia*, and it almost seems that, in this respect, they might be divided in two sections, one of which would be represented, as a type, by *Ardelio longipennis* Lw., the other by *A. brevicornis* Lw.

The genus EPICAUSTA, established by me for two African species, is less allied to *Rivellia* than to *Stenopterina*, which will be discussed below. These species are like *Stenopterina* in their stature, but are not so slender; the head is not unlike that of the species of *Dacus* proper; the antennæ are not quite as long as in *Stenopterina*; the fore coxæ are much shorter, and not so movable; the thorax, seen from the side, is not attenuated in front, as is the case with *Stenopterina*; the scutellum has four bristles, as in the latter genus; the wings are conspicuously shorter, and the last section of the fourth longitudinal vein is much more bent forward. The small crossvein is not oblique, as in all *Stenopterinae*, but perpendicular. The picture of the wings, in both of the species known to me, consists only in a large black spot at the tip.

STENOPTERINA, a genus proposed by Macquart in the *Suites à Buffon*, is well founded, as long as it is confined to the species of the immediate relationship of *Dacus brevicornis* Fab. and *æneus* Wied. Unfortunately the same author, in the *Diptères Exotiques*, has entirely left out of sight the characters of this genus, established by himself, and has introduced in it a number of heterogeneous forms, and, at the same time, placed in the genus *Herina* species which either belong to *Stenopterina* or are more closely related to it than to any other genus. His *Stenopterina femorata* and *immaculata*, both from Bourbon, seem to belong rather to *Epicausta* than to *Stenopterina*;

Stenopterina decora Macq. is, as has been observed above, the typical species of the genus *Euchalcota*; *S. gigas*, *scutellaris*, and *nigripes* of Macquart, all three from Tasmania, are certainly no *Stenopterina*. There would be more ground to place in that genus the *Ortalis violacea* of Macquart, which is probably correctly identified in the Berlin Museum with *Dacus macularis* Fab. *Herina mexicana* Macq. also belongs to *Stenopterina*, and *H. calcarata* Macq., although perhaps not a true *Stenopterina*, is closely related to that genus. The three species described by Walker (*List of Dipt. Ins.*), *bicolor*, of unknown origin, *trivittata*, from the Philippine Islands, and *basalis*, from Australia, do not seem to have anything in common with true *Stenopterina*. A true *Stenopterina* is *S. submetallica* Lw., from Mozambique; and *Herina chalybea* Doleschall, belongs probably to the same genus.

As I will have to characterize *Stenopterina* in detail among the North American genera of *Ortaliidæ*, it will suffice here to indicate the principal characters. Head resembling that of *Dacus* in structure; occiput convex, but not swollen. Front of a considerable and even breadth. Antennæ long and narrow, generally descending a little beyond the anterior edge of the mouth, which is somewhat drawn upwards; clypeus broad; proboscis stout. Thorax narrow; the pectus ascending obliquely in front, so that the thorax, seen from the side, is rather conspicuously attenuated anteriorly. Fore coxæ remarkably long, inserted unusually near the neck and very movable in this insertion. Scutellum with four bristles. Abdomen narrow; wings long and narrow; little crossvein oblique, placed beyond the middle of the long discal cell; the third and fourth longitudinal veins, in the majority of the species, are somewhat bent towards each other, so that the first posterior cell becomes narrower towards its end. In all the species I know of, the stigma, as well as a border between it and the apex, and the first basal cell, up to the small crossvein, are tinged with brown; in most species the posterior crossvein has likewise a dark border.

The next genus to be mentioned here is the genus MISCHOGASTER Macq., founded upon *Cephalia femoralis* Wied. *Mischogaster pernix* and *diffusus* Gerst., belong to it. It differs from *Cephalia* in the absence of a mesothoracic bristle, and in the face, which does not project inferiorly; from the following genus it is

distinguished by the first abdominal segment being beset with bristles. This character, as well as the somewhat advanced position of the anterior ocellus, remind of the *Richardina*. The face is rather short and somewhat excavated.

The last genus which I place among the *Platystomina* forms the transition from this group to the *Cephalina*, and shows a good deal of approximation to the genus *Cephalia*. As the typical species of this genus I consider *Cephalia myrmecoides* Loew. Besides the want of a mesothoracic bristle, this genus differs from the true *Cephalia* in the fact that the first abdominal segment is so coarctate in its middle that its anterior part forms a knot-shaped swelling; moreover, the shape of the body is still more slender; the wings still narrower and still more cuneiform towards the basis, so that the anal angle and the alula disappear entirely, whereas in *Cephalia*, there is at least a rudiment of them. The statements which Rob. Desvoidy makes about his genus *Myrmecomylia* render it probable that the above-mentioned species belongs to this genus. Certainty in this case is not possible, without the comparison of the species upon which Rob. Desvoidy established the genus. Not wishing to run the risk of introducing a useless generic name, I prefer to use the name of **MYRMECOMYLIA** for my species. The pleonastic name which the species thus obtains, *Myrmecomylia myrmecoides*, is not good, but may be tolerated in view of the fact that nothing is more like an ant than this dipteran.

A review of the genera which I placed among the *Platystomina* shows that, besides the bristles upon the first longitudinal vein, and the absence of prothoracic and mesothoracic bristles, which define this group, these genera have the following characters in common: The oral opening is very large; the clypeus generally very much developed, and the proboscis proportionally stout; the third antennal joint is elongate; the thoracic dorsum bristly upon its hind part only; the female abdomen has four segments, as the fifth is either altogether wanting, or only rudimentary and then completely hidden under the fourth segment.

Third Section: Cephalina.

I call this group after the genus which was first made known in it. It differs from the *Platystomina* in the presence of a

metathoracic bristle, from the *Ortalina*, in the absence of the prothoracic one. With the former it moreover agrees in the larger size of the oral opening, the greater development of the clypeus and the stouter proboscis; with the latter it has the more or less distinct development of the fifth segment of the female abdomen in common. While some of the genera show a very close affinity to the *Platystomina* in general appearance, others stand as near to the *Ortalina*, so that the *Cephalina* seem to form a transition from the first to the second of those sections.

The genus *CEPHALIA*, introduced by Meigen, shows some affinity to those genera of *Platystomina*, the species of which are distinguished by their slender shape, especially to the genera *Mischogaster* and *Myrmecomylia*. It necessarily must be confined to those species which, like the typical *Cephalia rufipes* Meig., have a mesothoracic bristle. The species added later to it, although in their general shape and their coloring they more or less resemble the true *Cephalia*, do not show the necessary agreement with them in those characters which are the most trustworthy in the establishment of the genera of *Ortalidæ*. They belong in the group *Platystomina* and principally in the genus *Mischogaster*, in part also in the genus *Myrmecomylia*. The genus *Cephalia*, in this narrower sense, does not contain as yet any American species. As, for this reason, I will have no occasion to refer to it again, I will characterize it here:—

Body slender, abdomen narrow at the basis, its first segment without any knot-shaped swelling; feet rather long and slender. Hairs on the body, extremely short; thorax with a few small bristles on the lateral and the posterior portions only; the bristles before the scutellum and its own lateral bristles are very short.

Antennæ long and slender; their second joint short. Face shield-like, convex, without antennal foveæ.

Palpi very broad; proboscis rather stout and mentum somewhat swollen.

Wings attenuated towards the basis in the shape of a wedge, with a very narrow alula; the second longitudinal vein hardly sinuose at all; the third and fourth longitudinal veins normal in their course; the anterior basal cell of equal breadth; the first longitudinal vein bristly towards its end only; the crossveins rather distant from each other; the picture of the wings usually

consists of an infuscation of the stigma and of a black spot on the apex.

Cephalia is immediately connected with a genus embracing *Trypeta flexa* Wied. and the genera related to the latter. As this genus does not coincide with any one of the hitherto adopted genera, it must receive a new name. I call it TRITOXÆ, the name alluding to the peculiar picture of the wings. The *Tritoxæ* differ from the *Cephalix* in the presence of a strong bristle before the end of the fore tibiæ, on their upper side, and in the presence of a weak indication of antennal foveæ, especially, however, in the fact that the third and fourth longitudinal veins have an irregular course, in consequence of which the anterior basal cell is expanded before its end; moreover also in the first longitudinal vein being, to a great extent, covered with bristles and in the approximation of both crossveins to each other. The wings have a dark coloring and the picture consists of three oblique, more or less arcuated, hyaline crossbands. The other characters the genus *Tritoxa* shares with the genus *Cephalia*.

After *Tritoxa* CAMPTONEURA naturally follows. The typical species is the well-known North American species, described by Fabricius as *Musca picta*, and afterwards erroneously placed by Wiedemann in the genus *Trypeta*. Rob. Desvoidy was the first to found a new genus for it, which he called *Delphinia*; Macquart established later for the same species the genus *Camptoneura*, which thus coincides with *Delphinia*. As the name *Delphinia* cannot be retained for reasons of priority, Macquart's name must be adopted. *Camptoneura* differs from *Tritoxa* in a striking manner in the structure of the wings; they are broad, and show, on the costal margin, near the end of the auxiliary vein, a shallow, but very striking excision; the third longitudinal vein is very remarkably sinuate, and the anal cell rounded at the end. The picture of the wings has a distant resemblance to that of the species of *Acitura*.

The other genera of *Cephalina* which I know of contain species of a less slender stature than the three genera which I have just examined.

Among them the genus PIARA, founded by me for an African species, is remarkable for its close relationship to the *Platystomina*. It may be characterized as follows:—

Body rather robust, the bristles upon the vertex, upon the posterior part of the thorax and upon the scutellum long.

Antennæ of medium length; the anterior corner of the third joint acute; arista feathery. Face excavated above, and with a projecting bump below.

Oral opening broader than long; proboscis very stout.

Wings rather broad; longitudinal veins diverging; the first, third, and fifth beset with bristles; posterior angle of the anal cell not acute; the picture of the wings is not unlike that prevailing in the genus *Aciura*.

Rather closely related to *Piara* is the genus *TRAPHERA*, which I propose to establish with *Ortalis chalybea* Wied. for its type. It also stands very near the *Platystomina* and may easily be considered as one of them, as the mesothoracic bristle is but very little conspicuous and the fifth segment of the female abdomen is also very much abbreviated. The principal differences between *Traphera* and *Piara* lie in the structure of the head and of the wings. The head of *Traphera* is not unlike that of *Platystoma*, but the lower part of the occiput is but very little turgid; the vertical diameter of the eyes is very long, the horizontal, on the contrary, very short; the first two antennal joints are short; the third jointed oval; the arista feathery; the face is descending obliquely, excavated under the antennæ, convex below; the clypeus is very much developed, its vertical diameter rather large, the horizontal one small; proboscis very much incrassated. Thorax stout and convex: its dorsum is provided with bristles only on the sides and posteriorly. Scutellum generally with eight bristles. Wings comparatively short and broad, with bristly hairs on their anterior margin; the whole of the first longitudinal vein is strongly bristly and shows, in the vicinity of the somewhat obliterate end of the auxiliary vein, a peculiar break; the basal half of the third longitudinal vein is beset with bristles; the posterior crossvein is oblique, so that the posterior angle of the discal cell is very acute; the anal cell is rounded at the end and its posterior angle withdrawn in a peculiar manner. The wings are of a dark color, marked with pale bands starting from the posterior margin and abbreviated in front.

While both of these genera are very near the *Platystomina*, the two which we have yet to mention approach the *Ortalina*. They are: *DIACHTA*, introduced by Gerstæcker, and a genus to be

adopted for *Ortalis marginata* Say, for which I propose the name of IDANA.

Diacrita is easily distinguished from *Idana* by the shape of the posterior angle of the anal cell, which is drawn out in a very long lobe, and by the picture of the wings, which consists only in a very broad dark border, extending to the very apex of the wing. The more extended picture on the wings of *Idana* is not unlike that of *Pteropœcila* and the posterior end of its anal cell forms only a short angle. As both genera contain North American species, I will have occasion to refer to them again more in detail.

Fourth Section: *Ortalina*.

The *Ortalina* have a prothoracic, as well as a mesothoracic bristle, while among the *Cephalina*, the former, among the *Platystomina*, both are wanting. The *Ortalina* are also distinguished from the two above-named groups by a smaller oral opening, a less developed clypeus, a less stout proboscis, a less turgid mentum and smaller palpi. In several genera, moreover, the thoracic dorsum is beset with bristles as far as its anterior portion. The abdomen of the female has five segments, which brings this group nearer to the *Cephalina* than to the *Platystomina*.

The geographical distribution of the *Ortalina* is, as far as known, confined exclusively to America and to Europe, with those parts of Asia which belong to the faunal province of the latter. Very striking is the great agreement between the European and North American forms of this group. As the knowledge of the latter is still very fragmentary, the generic distribution of the probably numerous species which may be discovered yet would offer great difficulties, or lead into error, unless based upon the knowledge of the European genera. I will give here, for this reason, a review of all the European genera adopted at present. Besides these, however, to the *Ortalina* must be reckoned the genus APOSPASMICA, which I propose to establish for the South American *Ortalis fasciata* Wied. and the genus AUTOMOLA, which I have adopted above for *Ortalis trifasciata* Wied. and *atomaria* Wied.

The European genera of *Ortalina* are the following:—

1. DORYCERA Meig.

Charact.—Eyes round; cheeks very broad; face very much projecting in profile; inferiorly it is very strongly retreating, carinate.

The hairs on the body have the ordinary length, or a little over the ordinary; thoracic dorsum bristly on its hind portion only.

Antennæ projecting, either of ordinary breadth and median length, with the third joint oval; or narrow and elongate, with an elongate third joint.

The first longitudinal vein bristly at its end only.

This genus contains gray species, their faces with dark spots, and with well-marked black stripes upon the thorax; the wings are either without any picture, or it consists of blackish-gray longitudinal lines, which are more confluent towards the apex, and even, in the male of one species, form a large, black spot.

The genus may be divided into two sections, which it will be necessary, when the number of species grows larger, to separate as genera.

Sect. 1. (*Dorycera*, sensu strict.) Antennæ narrow and very much prolonged; the pilosity of the body is of an ordinary length.

Typical species: *graminum* Fab.

Sect. 2. (*Percnomatia* Lw.) Antennæ of ordinary breadth and of medium length; pilosity of the body longer than usual.

Typical species: *inornata* Lw.

2. TETANOPS Fall.

Charact.—Eyes rounded-ovate; cheeks broad. Face in the profile very much projecting, more or less retreating inferiorly.

The hairs upon the whole body extremely short; the middle of the thoracic dorsum bristly on its hind part only; the prothoracic bristles are smaller than in all the other genera of *Ortalina*.

Antennæ short, often strikingly short; their third joint oval; somewhat longer than the second.

The first longitudinal vein is bristly at its end only.

This genus contains remarkably glabrous species; there are no thoracic stripes; the first segment of the ovipositor is comparatively large; there is no picture on the wings at all, or it consists only in narrow borders along the crossveins, or in more or less faded spots at the end of the longitudinal veins, thus resembling the picture of *Cerorhys*.

Typical species: *myopina* Fall.

3. CORMOCARIS LW.

Charact.—Eyes round; cheeks very broad; face in the profile strongly projecting, very much retreating inferiorly, not carinate.

Hairs on the body comparatively long; thoracic dorsum hairy and bristly as far as its anterior portion.

Antennæ short; the rounded oval third joint hardly as long as the second.

First longitudinal vein bristly at its end only.

Gray species, the abdomen and thorax of which are without any picture, and the wings dusky and somewhat spotted along the anterior margin.

Typical species: *bucephala* Meig.

4. PTEROPECILA LW.

Charact.—Eyes small, rounded oval; cheeks broad; front very much projecting.

Hairs on the body of the usual length; the middle of the thorax bristly on its hind portion only.

The rounded third joint of the antennæ short; the second likewise short.

The first longitudinal vein is hairy upon its whole length.

The coloring of the body is gray; the picture of the wings is not unlike that of *Idana marginata* Say.

Typical species: *lamed* Schrk.

5. PTILONOTA LW.

Charact.—Eyes elongated oval; front but little projecting.

Thorax bristly upon its middle, as far as its anterior portion.

The third antennal joint rounded oval; the second shorter.

The first longitudinal vein bristly at its end only.

Cinereous-gray species, the thorax of which is marked with four somewhat darker longitudinal lines; the picture of the wings consists of large blackish spots; in several species these spots are so much confluent that the picture of the wings can almost be called guttate.

Typical species: *centralis* Fab.

6. ORTALIS FALL.

Charact.—Eyes rather large, elongate oval; front only moderately projecting.

Hairs on the body of the usual length; the middle of the thoracic dorsum bristly on its hind portion only.

The rounded third antennal joint short, the second of the same length with it.

Both crossveins not more approximate than usual; the first longitudinal vein bristly at its end only.

The genus *Ortalis* contains species above the average size, some of them rather large; the abdomen is banded with gray; the thorax strongly pollinose, in most species with conspicuous black, in some, with gray longitudinal stripes, in a few, without any stripes. The wings are more or less spotted.

Typical species: *ruficeps* Fab.

7. SYSTATA LW.

Charact.—Eyes rather large, elongate oval; front only little projecting.

The hairs on the body as usual; the middle of the thoracic dorsum with bristles upon its hind portion only.

The rounded third antennal joint is short; the second of equal length with it.

The two crossveins are very closely approximated; the first longitudinal vein bristly at its end only.

The species of this genus differ from those of *Ortalis* in the very close proximity of the crossveins, but agree with them in the remainder of the organization. The picture of the wings consists in bands.

Typical species: *rivularis* Fab.

8. LOXODESMA LW.

Charact.—Eyes large, elongate; front but little projecting; face rather strongly carinate; cheeks narrower than in most other genera.

Hairs on the body as usual; thoracic dorsum with bristles upon its hind part only.

Third joint of the antennæ more or less prolonged, rounded at the tip; the second much shorter.

Both crossveins very much approximated; the first longitudinal vein bristly at its end only.

The species belonging here remind of the *Systatæ* in the striking proximity of the crossveins, differ however in other respects very much from them, and that in the same way as the species of *Pteropæctria* differ from *Ortalis*. The relation of *Loxodesma* to *Pteropæctria*, which is by far the most closely allied genus to

it, is exactly the same as that of *Systata* to *Ortalis*. The coloring and the picture of the wings resemble those of the first section of *Pteropæctria*, only the obscure borders of the crossveins coalesce more or less, on account of their proximity, into a single cross-band.

Typical species: *lacustris* Meig.

9. PTEROPÆCTRIA LW.

Charact.—Eyes large, elongate; front but very little projecting, face rather strongly carinate; cheeks narrower than in most other genera.

Hairs on the body of the usual length; the middle of the thoracic dorsum bristly on its posterior portion only.

Third antennal joint more or less elongate, rounded at the end; the second very much shorter.

The crossveins are at the usual distance from each other; the first longitudinal vein has bristles upon its end only.

This genus contains small, shining black species, the thorax of which shows only a faint trace of pollen. The picture of the wings generally consists in the dark color of the costal and subcostal cells, a more or less distinct black border of the crossveins and a black spot on the costa, lying a little before the apex, or upon it; in some species, however, this picture expands into four crossbands which are connected, two and two, near the costa.

The genus is divided into two sections, which may even be considered as separate genera. They are easily distinguished by the picture of the wings, which is in keeping with a corresponding difference in the rest of the organization.

Sect. 1. (*Pteropæctria*, sensu strict.) with spotted, or incompletely banded, wings.

Typical species: *palustris* Meig.

Sect. 2. (*Thryophila* Lw.); bands on the wings complete.

Typical species: *frondescentiæ* Lin.

10. TEPHRONOTA LW.

Charact.—Third antennal joint, although not excised on the upper side, still with a sharp anterior corner.

Thoracic dorsum, upon its middle, not bristly in front of the region of the suture.

First longitudinal vein bristly upon its end only; the fourth not bent forward; the posterior angle of the anal cell not prolonged in a lobe.

Tephronota begins the series of those genera, the third antennal joint of which is not rounded at the tip, but ends above in a sharp corner. It contains small species which, in the shape of their body, and especially in the structure of the head, remind of the *Pteropæctriæ* very much. But they can always be distinguished by their thorax, which is thickly covered with a gray pollen, even should the third antennal joint, in drying, have lost the sharpness of its upper corner. The picture of the wings consists either of complete crossbands, or of spots and half-bands, or even of spots only.

Typical species: *gyrans* Lw.

11. CEROXYS Macq.

Charact.—Third antennal joint distinctly excised on its upper side.

Thorax upon its middle beset with bristles as far as its anterior portion.

First longitudinal vein bristly upon its end only; fourth longitudinal vein not bent forward; the posterior angle of the anal cell not drawn out in a lobe.

Yellowish-gray or cinereous-gray species, with a thorax without stripes, and with wings having large dark spots; the arista is always distinctly pubescent.

Typical species: *crassipennis* Fab.

12. HYPOCHIRA Lw.

Charact.—Third antennal joint distinctly excised on its upper side.

Thorax, upon its middle, not bristly in front of the region of the suture.

First longitudinal vein with bristles upon its end only; fourth longitudinal vein not bent forward; posterior angle of the anal cell not drawn out in a lobe.

Small, grayish-white species, with a very limited picture of the wings, generally consisting of a very narrow border of the crossveins.

Typical species: *albipennis* Lw.

13. ANACAMPTA Lw.

Charact.—Third antennal joint distinctly excised upon its upper side.

Thorax, upon its middle, not bristly in front of the region of the suture.

First longitudinal vein bristly upon its end only; fourth longitudinal vein bent forward towards its end; posterior angle of the anal cell not drawn out in a lobe.

Rather robust species having the thorax pollinose with gray, without stripes or with weak ones, a black, shining abdomen, generally with gray bands, and wings which have black crossbands, or spots almost forming such crossbands.

Typical species: *urticæ* Lin.

14. HOLODASIA Lw.

Charact.—Third antennal joint distinctly excised on its upper side.

Thorax without bristles upon its middle, in front of the region of the suture.

First longitudinal vein bristly upon its whole length; fourth longitudinal vein curved forward at the end; posterior angle of the anal cell not drawn out in a point.

Holodasia differs from *Anacampta* (which it otherwise resembles very much) in the fact that the first longitudinal vein is bristly upon its whole extent, and not upon its end only. In this it agrees with *Pteropæcila*, from which it differs in the not projecting front, longer antennæ, the third joint of which is excised upon its upper side and pointed at the tip and in the fourth longitudinal vein being curved forward.

Typical species: *fraudulosa* Lw.

Fifth Section: Pterocallina.

At the beginning of the chapter on the Systematic Distribution of the Ortalidæ, I have pointed out *Scatophaga fasciata* as the species of this group known for the longest time and which may be considered as typical. It was described under that name by Fabricius in the *Systema Antliatorum*, was transferred by Meigen to the genus *Trypeta* and by Robineau Desvoidy to his new genus MYENNIS. It is very probably the same fly which was described by Coquebert in his *Iconographia*, Dec. III, under the name of *Musca octopunctata*, although it has nothing of the picture of the thorax shown in Coquebert's figure and which gave rise to the specific name. Although the publication of Coquebert's name is probably a little anterior to that of Fabricius, the choice of this name, based upon a non-existing character, as well as the nature of the entomological correspondence, which existed

between both authors, forbid us from giving Coquebert's name the priority over Fabricius's.

At the same place I have also observed how very distinct a species *Myennis fasciata* is, with its *Trypeta*-like stature, its low head and especially the very large distance between the ends of the auxiliary and of the first longitudinal vein; the latter character especially is quite peculiar among the *Ortalidæ* with a bristly first longitudinal vein.

Among the *Ortalidæ* hitherto described, the following species, as far as known to me, show a sufficient agreement, in their characters, with *Myennis fasciata* to be considered as belonging to the same circle of relationship: *Trypeta ocellata* Wied., from the environs of Bahia, Brazil; *Ortalis obscura* Wied., from Brazil, *Ortalis vau* Say, and *Platystoma annulipes* Macq., the two last from the United States. The numerous characters which all these species share with *Myennis fasciata*, besides the already mentioned peculiarities belonging to this species in particular, are: 1, the unmetallic coloring of the body; 2, the comparatively low, but rather broad head; 3, the broad front; 4, the rounded, more or less protruding eyes; 5, the round, or very short rounded-oval shape of the third antennal joint; 6, the shortness of the more or less concave face; 7, the small development of the clypeus; 8, the comparatively large development of the chest; 9, the prothoracic bristle, represented by a very small hair only; 10, the middle of the thorax, which is beset with bristles upon its hind part only; 11, the convex scutellum, provided with four bristles; 12, the very much abbreviated fifth segment of the female abdomen, which is very often quite withdrawn under the preceding segment; 13, the posterior angle of the anal cell, which is drawn out in a point, or even in a lobe.

Although the agreement in so many characters affords a distinct proof of the close relationship of these species, each of them shows at the same time plastic differences of such an importance, that one might be tempted to establish a separate genus for almost every one of them. These differences principally consist in the different shape of the wings, and in the different course of their veins, while the rest of the organization shows a remarkable agreement.

In the shape of the wings two remarkable modifications are worthy of notice, and may serve at some future time for a further subdivision of this group.

The wings of *Trypeta ocellata* and *obscura* differ from the usual shape of the wings of the *Ortaliidæ* by their narrowness, the parallelism of their anterior and posterior margins, their broad and rounded apex and their comparatively great length. Macquart placed the first of these species in the genus *Platystoma*, and the second, still more oddly, in the genus *Camptoneura*. Rondani has had a better eye for the plastic peculiarities of *Trypeta ocellata* and established the genus *PTEROCALLA* for it. I have derived the name of the present group from this well-founded genus of Rondani's, and not after Rob. Desvoidy's *Myennis*, established for *Scatophaga fuscata*, because the latter name, although much earlier in date, is a senseless malformation.

Trypeta obscura is, as Wiedemann has correctly observed in its description, a near relative of *Pterocalla ocellata*. As what occupies us now is the systematic location of only a small number of species, we can, without any hesitation, unite both of these species in the same genus, although the venation of *T. obscura* differs from that of *Pterocalla ocellata* in the second longitudinal vein being more arcuate than undulated, and in the fourth longitudinal vein being distinctly curved forward.

A small North American species, which will be described below, stands close enough to those two species in the shape of its wings and its venation to be placed in the same genus. It differs however in the second, third, and fourth longitudinal veins being quite straight, and neither wavy nor arcuate.

A most striking resemblance to this *Pterocalla strigula* is exhibited by *Trypeta ulula*, a South African species, described by me (*Berl. Entom. Zeitschr.*) after an incomplete specimen, without head. Already in describing this species, I drew attention to the fact that it differs from the ordinary venation of the *Trypetina* in the great distance intervening between the tips of the auxiliary and of the first longitudinal veins. I do not doubt now that this species is a *Pterocalla*, and that I would have recognized this earlier if I had had a complete specimen before me. Both species agree very well in all their plastic characters, especially in the shape of the wings and in the venation; the only difference which I notice in *P. ulula* is the position of the posterior crossvein, which is much steeper.

The genus *Pterocalla*, as I define it here, thus embraces all those *Pterocallina* which, in the outline of their wings, resemble

Pterocalla ocellata, so that this outline must be considered as the principal diagnostic character of this genus.

Among the numerous undescribed *Pterocallina*, which I have seen, I know of no one which may be placed in the genus *Pterocalla*, although several of them agree with the species of this genus in some one point pertaining to the venation. But none of those species has the wings of that peculiar shape which characterizes *Pterocalla*; on the contrary, the outline of the wings of all these species does not, in any marked degree, differ from that of the ordinary *Ortalidæ*. Like the species enumerated above, they have this peculiarity, that each species, although agreeing with the others in the characters belonging to the group, at the same time shows such important plastic differences, that the establishment of a series of new genera becomes indispensable. I regret not to be able to enter here into the detail of this subject, as, without plates, it is impossible to define those genera sufficiently. Thus much only will I mention, that among them there is a species which has the posterior angle of the anal cell rounded. The generic distribution of the North American species, which will be described below, does not, fortunately, require these South American forms to be taken into consideration.

Among the North American *Pterocallina*, *Ortalis vau* Say is the nearest to *Myennis fasciata* Fab. The venation, however, is different enough to prevent us from placing them in the same genus. The two crossveins in *O. vau* are less approximated, and the anterior end of the posterior one is further from the basis of the wing than the posterior end, while in *Myennis fasciata* the contrary is the case, so that the posterior crossvein of this species has a different position. Moreover, the first segment of the ovipositor of the female of *Ortalis vau* has not the conically attenuated shape which it has in *Myennis fasciata* and in many *Trypetina*; it is broader, somewhat attenuated from its middle only, like the ovipositor of the majority of the *Ortalidæ*. I consider, therefore, *Ortalis vau* as the type of a new genus, which I call *STICTOCEPHALA*.

To *Stictocephala vau* must be added a second North American species, which I received from Baron Osten Sacken, under the name of *Tephritis corticalis* Fitch in litt., and which will be described by me under the same name. The venation resembles

that of *S. vau* so closely that I have no hesitation in placing it in the same genus.

There are two other North American species which I take to be undescribed, and which also belong to *Stictocephala*. As their wings are not pictured like those of the two preceding species, but simply banded, the difference between them seems, at first glance, to be greater than it really is. A close examination does not disclose any plastic difference which would justify their generic separation from *Stictocephala*. I will describe them as *Stictocephala cribrum* and *cribellum*.

The North American species described by Macquart as *Platystoma annulipes* shows, in the detail of its structure, an almost complete agreement with the species of *Stictocephala*, but differs so much in the outline of the wings and still more in the venation, that it cannot be placed in that genus. The difference in the outline of the wings consists in the fact that the posterior margin is more convex, and hence, the wings are broader; the difference in the venation appears in the posterior angle of the anal cell being drawn out in a very long lobe, and in the position of the posterior crossvein, the anterior end of which is much nearer to the apex of the wing than the posterior end. As this species does not find a convenient place in any of the existing genera, I am compelled to establish a new one for it, which I call CALLOPISTRIA.

This would close the series of the few genera of *Pterocallina*, hitherto sufficiently defined, if we had not to advert to the genus *PSAIROPTERA* Wahlb., occurring in northern and central Europe, as well as in northern Asia, a genus for which it is not easy to find an appropriate place in the system. The species of this genus resemble the *Ulidina* in their general appearance, and I would not have hesitated to place them in that section, if their third longitudinal vein was not distinctly beset with hairs. I acknowledge that their location among the *Ulidina* is more natural than among the *Pterocallina*. Nevertheless, I place the genus among the latter and thus put a greater stress upon the artificial character, derived from the pilosity of the third vein, than upon more close and natural affinities, but which are more difficult to explain in words. If I do this, it is because I hold that a strict adherence to those characters, by means of which I have tried to introduce into the systematic chaos of the

Ortalidæ a satisfactory distribution in groups, is more apt to insure the recognition within these groups of available genera, than if we should attempt to avail ourselves of affinities, which, although visible to the eye, do not admit of exact definitions.

Psairoptera finds a fitting location at the end of the *Pterocallina*, so as to be immediately followed by the *Ulidina*. The principal differences from the above-mentioned genera of *Pterocallina* consist in the posterior angle of the anal cell, which is more or less a sharp right angle, and in the much smaller distance between the end of the auxiliary vein and that of the first longitudinal. The shape of the head likewise shows not unimportant differences from the other genera of the group, and some of the species of *Psairoptera* have, moreover, the last antennal joint of a more elliptical shape.

In enumerating the most characteristic distinctive marks of the *Pterocallina*, we cannot, for the above stated reasons, lay the same stress upon *Psairoptera* as upon the other genera of this group. These characters may be summed up as follows:—

Habitus *Trypeta*-like; coloring non-metallic; head rather broad, but low, with rather protuberant eyes; face short, perpendicular, excavated in the middle; clypeus but little developed; third antennal joint round or rounded ovate; thoracic dorsum bristly upon its posterior part only; third longitudinal vein hairy; and above all, as the most important character, the unusually large distance between the end of the first longitudinal and that of the auxiliary veins.

For the *Pterocallina* from North America, hitherto known, we can add to the above-enumerated characters the posterior angle of the anal cell, which is drawn out in a long lobe.

SECOND DIVISION.

ORTALIDÆ HAVING THE FIRST LONGITUDINAL VEIN BARE.

The European genera belonging here are: *SEOPTERA* Kirby, *TIMIA* Wied., *ULIDIA* Meig., *CHRYSOMYZA* Fall., with which *Chloria* Schin. is coincident, and *EMPYELOCERA* Lw. They are allied enough to each other to be united in the same group.

A type, very different from the preceding genera, appears in the genus *RICHARDIA* Rob. Desv., which seems to be rather

abundantly represented in America and likewise belongs to this division. A whole series of related genera, peculiar to America, may be classed with *Richardia*: like the latter, they are all distinguished by armed femora.

This is the reason why, in a former publication, I separated the whole second division of the *Ortalina* in two groups, the one with unarmed, the other with armed femora; the first I called *Ulidina*, the second *Richardina*; and in the *Berlin Entom. Zeitschrift*, Vol. XI, I described the American *Ulidina* which, at the time, were known to me. Now, however, that I have become acquainted with a larger number of forms belonging in this division, I incline to think that its separation in the groups *Ulidina* and *Richardina* becomes more natural, if, as a distinguishing character of these groups, we assume, not the armed or unarmed femora, but the shape of the anal cell. All the genera having the posterior angle of the anal cell more or less pointed belong to the *Ulidina*; those genera, on the contrary, where this is not the case are to be placed with the *Richardina*. This modification does not much alter the distribution of the genera among these two groups, as all the genera with armed femora, at present known, will, in the new distribution, be likewise referred to the *Richardina*. Among the genera which, in the above-quoted publication, I placed with the *Ulidina*, *Epipletca* alone will have to be transferred among the *Richardina*. Among the genera of *Richardina*, enumerated below, *Steneretma*, according to the former mode of subdivision, would have belonged to the *Ulidina*, and thus would not have been placed near *Idiotypa*, which is closely allied to it. With the former mode of distribution, the position of the new genus *Coniceps*, based upon a North American species, would have been a somewhat doubtful one, as the under side of its hind femora bears a few stronger hairs, but can hardly be called armed.

First Section: Ulidina.

The five genera of *Ulidina* represented in Europe, and enumerated in the preceding paragraph, are not confined to this part of the world. The European *SEOPTERA vibrans* also occurs in the adjoining provinces of Asia, and is represented in America by a species most closely resembling it. European species of

TIMIA, EMPYLOCERA, and ULIDIA occur in Asia together with other species, peculiar to that part of the world. CHRYSOMYZA *demandata* likewise ranges over a considerable part of Asia and Africa; both countries contain besides species of this genus peculiar to them.

The South American *Ulidia stigma* Wiedemann and the Brazilian *Ulidia bipunctata* Macq. are not *Ulidiæ* at all, although they probably belong to the group *Ulidina*, the first to the genus *Notogramma*, the last to *Euxesta*. *Ulidia metallica* Bigot, from Cuba, is perhaps a *Chrysomyza*; as to the *Ulidia fulvifrons* Bigot, from the same locality, it is impossible, from Bigot's description, to come to any conclusions about its place in the system.

America seems in general to be very rich in forms belonging to the *Ulidina*. For the species which came within my knowledge I have established the genera: DASYMETOPA, OEDOPA, NOTOGRAMMA, EUPIARA, ACROSTICTA, EUXESTA, CHÆTOPSIS, HYPOCTA and STENOMYIA.

The species described by former authors, which belong in the circle of the above-mentioned genera, are to be found in Wiedemann partly in the genus *Ortalis*, partly in *Ulidia*. In Macquart, as far as I can ascertain, they are scattered among the *Ulidia* or even in *Cerorhys* and *Urophora*, which shows, on that author's part, an utter neglect of their plastic characters. The genus EUMETOPA established by Macquart in his family *Psilomydæ*, does not belong to it, but to the *Ulidina*.

It is not doubtful at all that Asia and Africa, besides the genera which they have in common with Europe, harbor some genera of *Ulidina* which are peculiar to them. GORGOPIS Gerstæcker, described by Doleschall, some time previously, under the inappropriate name of *Zygenula*, probably belongs to this group. It differs, it is true, from all the known *Ulidina* very much; still the structure of the head in *Oedopa* may be indicative of an affinity.

Unfortunately I cannot give any further information concerning other exotic *Ulidina* of the old world, as I have none in my possession. The existing descriptions of a number of *Ortalidæ* which may possibly belong to the group *Ulidina*, are not accurate enough to admit of any positive conclusions.

I have not met with any *Ulidina* from Oceanica yet.

Second Section: Richardina.

Among all the genera of this group, RICHARDIA Rob. Desv., distinguished by its posterior femora, armed with spines, is the best and longest known. This circumstance induced me to derive the name of the section from it. It seems to be exclusively American; the *Richardia flavilarsis* Macq., from the Marquesas Islands, does not belong to this genus, and if the manner in which the auxiliary vein is represented upon Macquart's figure be only of average correctness, we may even infer that it does not belong to the *Richardina* at all. The other species which Macquart, Rondani and Gerstæcker have added to the genus *Richardia* are all natives of America. The two males of *Richardia* described by Gerstæcker are distinguished by the dilatation of their head, somewhat in the manner of *Achias*; their females are not known yet; still the analogy of *Achias* and of other genera, having a similar structure of the head, justifies us in supposing that their heads do not show any extraordinary dilatation.

The fly of unknown *habitat* which Macquart described as ODONTOMERA *ferruginea* undoubtedly belongs in the immediate affinity of *Richardia*. As I have never seen it, my knowledge of it is based exclusively upon Mr. Macquart's statements. These, however, are entirely sufficient to prove that the fly belongs in the family *Ortalidæ*, and not in the *Trypetidæ*, where Macquart places it. That it belongs to the *Richardina* I infer from the evidently very close relationship which exists between it and the *Sepsis Guérinii* Bigot from Cuba. The generic name must be changed, on account of the already existing *Odontomcrus* Gravenh.

This *Sepsis Guérinii* agrees in so many characters with *Odontomera ferruginea* Macq. that one might be tempted to place it in the genus *Odontomera*. Should Macquart's statements, however, be correct, this would not be admissible, as *Odontomera ferruginea* possesses not only much stouter femora and a much more projecting front, but also an auxiliary vein which is much less approximated to the first longitudinal than in *Sepsis Guérinii*. We are compelled, therefore, to consider *Sepsis Guérinii* as a separate genus of the *Richardina*, which we will call STENOMACRA.

We have, in the next place, to mention the genus SETELLIA.

It was founded by Rob. Desvoidy, and *Setellia atra* Rob. Desv. must be considered as its type. I have not seen this species, and, unfortunately, the statements of Rob. Desvoidy are not sufficient to enable me to decide whether *Setellia atra* belongs to the *Richardina* or to the *Cephalina*. In the same way, I am unable to decide whether the Brazilian species, subsequently described by Macquart as *Setellia apicalis* really belongs in the same genus with *Setellia atra*. As Rob. Desvoidy does not allude at all to the femora of his species being spinous, while Macquart's species is remarkable for all its femora being armed in a rather striking manner, it becomes exceedingly doubtful whether Macquart's species is a *Setellia* in the sense of Rob. Desvoidy's.

I do not know of any species more related to *Setellia apicalis* Macq. than that species from Colombia, South America, which Gerstæcker described under the name of *Michogaster egregius*. As its first longitudinal vein is bare and its femora are armed, it cannot possibly remain connected with the true species of *Michogaster*, but must be considered as the type of a separate genus of *Richardina* for which I propose the name of *EUOLENA*.

To place *Setellia apicalis* in the genus *Euolena* is not possible; it has no stump of a vein upon the second longitudinal vein inside of the submarginal cell, a character distinguishing *Euolena egregia*; its third and fourth longitudinal veins converge more distinctly towards their end, and the posterior angle of the anal cell is not rounded. *Setellia apicalis* will also have to be considered as the type of a separate genus, which may be called *SYNTACES*. In the supposition that the first longitudinal vein of *Syntaces apicalis*, like that of its relative *Euolena egregia*, is bare, I think that the best location for this genus is among the *Richardina*. It is true that the posterior angle of the anal cell, in Macquart's figure, is almost acute; in the generic diagnosis, however, he calls the anal cell: "terminée carrément," so that the shape of this cell cannot be an obstacle to the location of the genus among the *Richardina*; and this view is supported by the spinous femora, a character common to nearly all the genera of this group. Should, however, the first longitudinal vein of *Syntaces* be hairy or bristly, then the location of the genus among the *Richardina* would be impossible.

Next to *Euolena* is the genus *IDIOTYPA*, which I establish for a new species from Cuba. In its general habitus it is almost

like one of the more corpulent American species of *Baccha*; for instance, *Baccha capitata* Lw. The second longitudinal vein, which in *Euolena* forms a short stump inside of the submarginal cell, bears, in this genus, almost at the same place, similar stumps, not only in the submarginal, but also in the marginal cell. The most striking difference, however, lies in the structure of the feet, as *Euolena* has the four posterior femora remarkably long and slender, which is not in the least the case with *Idiotypa*.

The genus *STENERETMA*, which will be characterized in the third part, treating of the North American species, is related to *Idiotypa*.

The South American species described by Fabricius once as *Scatophaga trimaculata* and another time as *Dacus flavus*, and which Wiedemann placed in the genus *Trypeta*, does not belong in this genus at all, but in the present group of the *Ortaliidæ*. The description, which Macquart gave of his *CÆLOMETOPIA ferruginea*, contains so much which is entirely applicable to Fabricius's species, that I have no doubt that the latter species was the very same from which the description of *Cælometopia ferruginea* was drawn. When Macquart says that in *C. ferruginea* the middle femora alone are armed, this statement is probably based upon an insufficient observation; when he calls the last three tarsal joints white, this seems to be a *lapsus calami*, as the figure shows nothing of the kind, and as on two of the tarsi the first joints are even represented as much paler than the following ones; the latter probably being infuscated, as they are in Fabricius's species. Should even, contrary to my supposition, Macquart's species be different from that of Fabricius, they will at all events belong to the same genus.

The *Odontomera maculipennis* of Macquart from Colombia, South America, seems very closely allied to *Cælometopia*; Macquart's own statements show that it agrees in so many characters with *Cælometopia trimaculata*, that it may be transferred to the same genus with it; one would even be led to suppose that it is nothing but the female of *Cælometopia trimaculata*. With the above mentioned *Odontomera ferruginea* Macq. (not *Cælometopia ferruginea* Macq.) *Odontomera maculipennis* has too little in common to be considered as belonging to the same genus.

A pretty species from Cuba, which will be described in the

sequel, can also be placed in the genus *Carlometopia*, although the ocelli, which here, as well as in the latter genus, are rather much forward on the front and close to each other, are placed here upon a very gentle elevation, while in *Carlometopia* the projection which bears them is quite high.

Closely related to *Carlometopia* is the species described by Wiedemann as *Trypeta cyanogaster*. It is not a real *Carlometopia*, as its posterior ocelli are less remote from the vertex and the anterior one quite distant from them; moreover the third and fourth longitudinal veins are parallel here and the hind femora alone bear a few bristles, while in *Carlometopia* all the femora are beset with spines. For this reason *Trypeta cyanogaster* has to be considered as the type of a new genus, which may be called **MELANOLOMA**. A second species of this genus, from Brazil, has the same picture of the wings as *M. cyanogaster*, consisting in a black border of the costal margin and of the small crossvein.

Other Brazilian *Ortalidæ* resemble the genus *Melanoloma* in the fact that the third and fourth longitudinal veins are parallel; the agreement in the structure of the rest of the body, especially of the head, is very striking. These species differ, however, in all the femora being spinous, in the arista being distinctly pubescent, in the still greater distance between the anterior ocellus and the two posterior ones, in the close proximity of the two crossveins of the wings, and in the picture of the wings, which does not consist in a black border on the costa, but in large, crossband-like spots. I deem it useful to introduce for such species a new genus, which I will call **HEMIXANTHA**; a species of this genus, *H. spinipes*, will be described below.

That *Dacus flavicornis* Wied., from Brazil, belongs in the same circle of relationship is proved by the original specimen, preserved in the Berlin Museum.

Before having subjected that specimen to a second and more close examination, I would not venture to decide whether it can be placed in any of the genera, which I have just discussed. As far as I remember, its scutellum bears only two bristles; this would prevent its identification with any one of those genera, as it is very unlike just those among them which share that character with it. Otherwise it has the same *Dacus*-like structure of the face as most *Richardina*; the third antennal joint is elongated;

the slender arista is distinctly pubescent; the abdomen is of an equal breadth; the posterior angle of the anal cell is not neutre and the fourth longitudinal vein somewhat convergent with the third; all the femora are armed.

I have also to mention the genus *CONICEPS*, which I find necessary to establish for a North American species. On account of the retracted posterior angle of the anal cell it must likewise be placed among the *Richardina*, although in its general appearance it is more like certain *Ulidina*, especially *Eumetopia*.

The reason why I place *EPIPLATEA* among the *Richardina* has been alluded to above.

Thus I have reached the limit of the genera, the location of which among the *Richardina* appears to me beyond doubt. It is certain that the number of *Richardina* which may yet remain unrecognized among the existing descriptions is far from exhausted by me; but who would venture, upon the statements of most of these descriptions, to form an opinion on the systematic location of the species which they mean to represent!

It will hardly be necessary to mention here the East Indian genus *MERACANTHA*. Its spinose femora may suggest the supposition that it belongs to the *Richardina*. But as this character does not belong exclusively to this group, and as the very acute angle of the anal cell of *Meracantha* does not occur among the *Richardina* in the acceptation of that group as I understand it here, I cannot consider *Meracantha* as belonging to the *Richardina*.

Besides the bareness of the first longitudinal vein and the not acute posterior angle of the anal cell, which two characters constitute the diagnosis of the *Richardina*, the following characters are common to all the genera which I have had occasion to examine in detail: a break in the costal vein immediately before the end of the auxiliary vein; the great proximity between the auxiliary and first longitudinal veins and the very small distance between their ends; finally the thoracic dorsum being beset with bristles upon its posterior part only.

III.

THE NORTH AMERICAN ORTALIDÆ.

It is a long time since I intended to publish a monograph of the North American Ortalidæ. The hope and expectation, however, of increasing in a measure my very fragmentary knowledge of this family by the addition of more species, either new or not yet seen by me, induced me to postpone for some time the completion of my work. Unfortunately, this expectation has not been fulfilled. Within the last four years, only five species were added to those previously known by me, and it became evident that if I had to wait for a tolerable increase of my acquaintance with the Ortalidæ, my work would run the risk of remaining unpublished. I let it appear, therefore, in the best shape I could give it, with the scanty materials at my command. I have no doubt that North America contains a far larger number of genera than those which came within my knowledge. In order to define, with some approximation, the systematic position of the genera of which I have not had any representatives for comparison, I have included in this monograph all the South American genera of which I possess specimens; inasmuch as it is very probable that most of them occur at least in the southern portion of North America. The striking analogy between the North American and European *Ortalina* renders it very probable that the number of genera in this group, common to both continents, is larger than it appears at present. For this reason I have deemed it useful to include in the general characters of the *Ortalina* all the data necessary for the recognition of the more difficult and less well known among the European genera.

Synopsis of the Distribution of the Family.

DIVISION I.—First longitudinal vein bristly or distinctly hairy.

A. Ovipositor not flattened.

Section I. *Pyrgotina*.

B. Ovipositor flattened.

a. Third antennal joint not circular.

1. No prothoracic, no mesothoracic bristle.

Section II. *Platystomina*.

2. No prothoracic, but a mesothoracic bristle.

Section III. *Cephalina*.

3. A prothoracic and a mesothoracic bristle.

Section IV. *Ortalina*.

b. Third antennal joint circular.

Section V. *Pterocallina*.

DIVISION II.—First longitudinal vein bare.

A. Posterior angle of the anal cell drawn out in a point, or, at least, more or less acute.

Femora never armed.

Section I. *Ulidina*.

B. Posterior angle of the anal cell obtuse, rounded or retracted.

Femora armed in most of the genera.

Section II. *Richardina*.

FIRST DIVISION.

ORTALIDÆ WITH THE FIRST LONGITUDINAL VEIN BRISTLY OR DISTINCTLY HAIRY.

*First Section: PYRGOTINA.*GEN. I. **PYRGOTA** WIED.*Charact.*—Front of equal breadth, without ocelli, very much projecting in profile.*Antennæ* drooping, second joint rather long, third more or less ovate; arista pubescent.*Face* retreating, under the antennæ with deep foveæ, separated by a very low ridge; they reach as far as the middle of the face, or only a little below; lateral parts of the face very broad, still more approximated on the lower half of the face; *oral opening* comparatively small; *clypeus* but little developed; *proboscis* not stout.*Scutellum* with many bristles.

¹ It may not be useless to refer here to Vol. I, p. xxiv, of these *Monographs*, where (fig. 1) a wing of *Ortalis* is represented. The anal cell is marked *M* on the figure, and is the same as the third basal cell, or the posterior one of the small basal cells. Although this synonymy is not mentioned in the explanation of the figure (at the foot of the same page), it may be found in the same volume, p. xx, line 18 from the top.—O. S.

Abdomen: in the male with four segments, the first of which strikingly prolonged, the following ones considerably shorter; in the female with five segments, the first of which very remarkably prolonged, the following ones quite as remarkably shortened; *ovipositor* large, not flattened, almost capsule-shaped.

Spurs of the middle tibiae only bristle like; very weak in the species with less coarse hairs.

Wings large; posterior angle of the anal cell acute; small crossvein beyond the middle of the long discal cell; third longitudinal vein curved backwards towards its end; the last section of the fourth longitudinal vein arcuated, but little diverging from the third.

Maequart's genus *Oxycephala* is identical with *Pyrgota*. Harris, in his Catalogue of the Insects of Massachusetts, calls this genus *Sphecomyia*.

Real *Pyrgotæ* are known to occur with certainty in North America only. As in Europe and Africa genera occur, which are closely allied to *Pyrgota*, it is not impossible that Walker's *P. latipennis* (List of Dipt. p. 1087) from Sierra Leone is a real *Pyrgota*; however, his description is altogether silent concerning those characters which are indispensable for the recognition of the genus. Whether *P. pictipennis* Walker (List, etc. 1162) belongs to this genus is very doubtful; the author himself introduces it with a doubt, but remains silent as to the motives of this location as well as the cause of the doubt.

The North American *Pyrgotæ* at present known may be divided in two groups: in the first, the arista is only two-jointed, and, at the same time, the usual bristles on the vertex, as well as those bristles which in other genera protect the ocelli, are present; in the other group, the arista is distinctly three-jointed, and there are no conspicuous bristles either on the vertex, or round the spot where, in other genera, the ocelli are placed. *Pyrgota millepunctata* belongs to the first, all the other species to the last group. Were the number of the species larger, these characters would justify a subdivision in two genera; at present, with the small number of species, all easy to identify, this subdivision would be useless.

I. *P. millepunctata* Lw. ♀.—Fusco picea, seta antennali biarticulata; alæ infuscatæ, guttulis numerosis subpellucidis aspersæ.

Pitch-brown; arista two-jointed; wings infuscated, dotted with numerous pellucid spots. Long. corp. 0.38—0.43, cum terebra 0.51—0.55, long. al. 0.49—0.55.

SYN. *Pyrigota millepunctata* LOEW, Neue Beitr. II, 22, 50.

?*Orycephala maculipennis* MACQ. Dipt. Exot. Suppl. I, p. 210. Tab. xxviii, f. 2.

Sphecomylia valida HARRIS, Catal. Ins. Mass.

Prevailing color of the body pitch-brown, reddish-brown or even brownish-red in less intensely colored specimens, with a black pubescence, which is perceptibly coarser than in the following species. The occiput has, behind the vertex, a distinct black triangle, with its point directed downwards, which is connected with a black spot on the place where the ocelli should be; at some distance from this triangle there is, on each side, a large black spot, reaching from the posterior orbit of the eye almost to the point of attachment of the head; between these spots and the triangle the color is clay-yellow, almost wax-yellow; the sides of the occiput are generally of a similar yellow color, but become more infuscated towards the orbits and the cheeks, or are tinged with brownish as far as the black spots above. The front has a broad black stripe, which is divided longitudinally in two by a more or less complete and more or less narrow, sometimes more yellow, sometimes brownish, line; on both sides, near the orbits, the stripe is margined with yellow. The ordinary strong bristles on the vertex, the bristle placed in front of these, on each side, near the orbit, and those bristles which are inserted in the region of the ocelli (which here are wanting), are all present. The first antennal joint is generally rather dark-brown, except at the basis; the second is usually of a dirty brownish-yellow; the third agrees in its coloring sometimes with the first, sometimes more with the second joint; in some specimens, it is altogether ochre-yellow; the arista is distinctly two-jointed, the first joint short. The face is usually of a dark ferruginous-brownish coloring, often verging on ochre-yellow on the sides. The antennal foveae are somewhat less deep than in *P. undata*, but perceptibly longer and separated by a higher ridge. The sides of the face are approximated on the lower half, but not so much by far as in *P. undata*, so that the middle part of the face has about double the breadth of the other species. The oral opening is more horizontal than in *P. undata*. The but little developed clypeus is black, the palpi generally yellow; their shape is almost the same as in *P. undata*. The ground color of the thorax is clay-yellow or wax-yellow, but with very broad pitch-brown stripes,

which occupy everything but the humeri and the narrow intervals between the stripes, so that the prevailing color is the brown one; the middle stripe, which is of equal breadth, is longitudinally divided in two by a lighter longitudinal line; the stripe stops at the last quarter of the thorax, however, beyond it, at the posterior margin of the thorax, there is a brown spot; the very broad lateral stripes are strongly abbreviated anteriorly, attenuated and interrupted at the transverse suture; moreover, the lateral margin has a broad brown border. Scutellum blackish-brown, paler on the sides; the numerous bristles are more conspicuous in this species on account of their stoutness and their black coloring. Pleuræ pitch-brown, clay-yellowish on the sutures. Abdomen usually blackish-brown or dark pitch-brown, sometimes ferruginous-brown or yellowish-brown; the first segment is, about once and a half the length of the four following segments taken together. The capsule-shaped ovipositor is of the same color as the abdomen, or somewhat paler; its shape is nearly the same as in *P. undata*, but it is a little less pointed; on each side, not far from the basis, it has a large, impressed spot. The color of the feet is as variable as that of the remainder of the body; blackish-brown in more intensely colored specimens, otherwise ferruginous-brownish; the knees are always clay-yellow; paler colored specimens have the extreme tip of the tibiæ and the tarsi of a dirty ferruginous-yellow or ochre-yellow color. The shape of the wings is not unlike that in *P. undata*, but towards the apex they are broader. The chief differences in the venation are the following: the little stump of a vein on the second longitudinal vein existing in *P. undata*, is wanting here; the discal cell is much broader, especially towards its tip; the posterior transverse vein is nearer the margin of the wing, much longer and more straight; the last section of the fourth longitudinal vein is less strongly arcuated and the second posterior cell much smaller; the posterior angle of the anal cell is more drawn out in a point. The whole surface of the wings has a rather uniform dark-brownish coloring; this color is variegated by numerous transparent dots of a gray-yellowish tinge; the shape of these dots is rather irregular; they are often confluent, as often distinctly separated; round the root of the second longitudinal vein and round the small crossvein, the dark coloring is more continuous and less interrupted by dots; the brown

is also more intense along the costal margin, than upon the remaining surface.

Hab. Carolina (Zimmerman); Washington, D. C., New York, Illinois (Osten-Sacken); Massachusetts (Harris).

Observation 1.—Mr. Macquart (Dipt. Exot. Suppl. I, p. 210) describes as *Oxycephala maculipennis* from Texas (figured on Tab. XIX, f. 12), a species which either is a *Pyrgota* or is closely allied to this genus. In several respects this species shows a decided resemblance to *P. millepunctata*, and the question as to their diversity is a very doubtful one. The conformity is especially apparent in the picture of the wings and the venation, also in the coloring of the front and even in that of the thorax. But Macquart says that the thoracic stripes are interrupted near the suture (which is also rendered in his figure); moreover, according to the figure, the posterior angle of the anal cell is drawn out in a much longer point than is the case in *P. millepunctata*. These discrepancies alone, however, with Macquart's well-known inaccuracy in description and figure, would not be sufficient to neutralize the evident analogies. A more weighty ground for doubt is to be found in the representation of the abdomen; nothing like its remarkable breadth has been observed in any known *Pyrgota*; moreover, it shows, instead of five segments, only four, the first of which is abbreviated, and the second the longest; the ovipositor hardly exceeds one-third of the length of the abdomen, while in the other *Pyrgotæ* it equals the abdomen in length. If these statements were based on Macquart's figure alone, I would have been inclined to think that the abdomen, wanting in the specimen, had been supplied by the imagination of the draughtsman; but this supposition does not hold good in presence of the fact, that Macquart mentions expressly, that he had a female before him; and we know that the sex of a *Pyrgota* can only be recognized by the structure of the abdomen. Macquart also says that the ovipositor is flattened, which is not in the least the case with *P. millepunctata*. These grounds seem sufficient to justify the belief that Macquart's *Oxycephala maculipennis* is a different species from *P. millepunctata*, unless we assume that Macquart's specimen had the abdomen of a different species fastened to it. Should this not be the case, there is every reason to doubt whether the species is a *Pyrgota* at all. It is rather strange that in the list of the exotic species described in Macquart's

work, which is appended to his fourth supplement, *O. maculipennis* is omitted. The cause of this omission is not apparent. Should Macquart have discovered that it belonged to a different genus, he would have transferred it to that genus; but the species is altogether omitted in the list.

Observation 2.—*Sphecomyia valida* of Harris's Catalogue of the Insects of Massachusetts, is, according to a communication from Baron Osten-Sacken, nothing else but *Pyrgota millipunctata*. As a matter of course, Harris's name, being merely a catalogue name, has no claim of priority.

2. P. undata WIED. ♂ ♀.—Ex ochraceo ferruginea; antennarum articulus tertius secundo æqualis; seta antennalis triarticulata, articulis primis duobus subæqualibus; alarum vena longitudinalis secunda appendiculata.

Yellowish-ferruginous; the third antennal joint equal to the second in length; arista three-jointed; its first two joints of nearly equal length; the second longitudinal vein with a stump of a vein upon it. Long. corp. ♂ 0.4—0.43; ♀ cum terebrâ 0.5—0.53; long. al. 0.5—0.58.

Syn. *Pyrgota undata* WIED. Auss. Zweifl. II, p. 581. Tab. X. 6.

Pyrgota undata MACQ. Suites, etc., II, p. 423. Tab. XVIII, f. 23 (were mentioned after Wiedemann).

Myopa nigripennis, GRAY, Anim. Kingd. Tab. 125, f. 5.

Orycephala fuscipennis MACQ. Dipt. Exot. II, 3, p. 198. Tab. XXVI, 6.

Sphecomyia undata HARRIS, Cat. Ins. Mass.

Pyrgota undata GERST. Stett. Ent. Zeit. xxi, p. 188.

Yellowish-ferruginous or more ochre-brownish. Front rather broad, projecting almost in the shape of a tower, and with a short, rather inconspicuous pubescence; without stronger bristles in the region of the vertex or round the place where the ocelli usually are. Antennæ yellow; the first two joints with a yellowish pubescence; the third sometimes ochre-brown, of the same length as the second. Arista distinctly three-jointed; the first two joints almost of equal length. The face very much retreating when seen in profile; the very deep antennal foveæ reach only to its middle and are separated by a very low ridge, which is usually tinged with brown; below them, the middle portion of the face is remarkably narrow, groove-like and bordered on each side by a brownish-black ridge. A brown or brownish-black, somewhat curved stripe generally extends from the middle of the inner orbit of the eyes towards the region of the antennæ.

The oral opening is cut obliquely upwards; the but little developed clypeus is tinged with blackish; the rather broad palpi are usually tinged with yellowish-red towards the tip, sometimes they are altogether ferruginous. The thoracic dorsum has an extended ferruginous-brown spot upon it, formed by the almost complete coalescence of a broad intermediate stripe with two broad lateral stripes, which are abbreviated in front. The meta-thorax and the greater part of the pleuræ are often tinged with dark pitch-brown. The coloring of the abdomen on the first two segments, and also at the basis and along the middle of the following ones, often becomes pitch-brown or brownish-black, this is especially often the case in male specimens. The first abdominal segment is very much elongated in both sexes; in the male it is not quite as long as the three remaining segments taken together; in the female, the last four segments are so much shortened, that, taken together, they are much shorter than the first joint. The capsule-shaped ovipositor is conical, bent downward towards its end. The feet are ochre-yellowish, but the femora brown up to the tip; the tibiæ likewise are more or less infuscated, except the basis and the extreme tip. Wings large, the greater portion of them is uniformly tinged with brown, which color covers the costal, marginal, submarginal, the first posterior and the discal cells, also the basal cells, with the exception of a pale stripe in the anal cell, moreover, this color forms a broad border along the inner portion of the second posterior cell, and a narrower one along the anterior margin of the third posterior cell; within this brown coloring some specimens do not show any paler spots, the majority, however, show, in the submarginal cell, a little beyond the small crossvein, a rounded or oval, almost hyaline spot, which attains sometimes a considerable size; moreover, a great many specimens show some scattered, small, hyaline dots, not far from the end of the same cell, of the first posterior and of the discal cells; the posterior limit of the brown coloring has a whitish-hyaline border, which, following the course of that limit, forms a steep curve in the second posterior cell; in the third posterior cell it takes the shape of a gently arched longitudinal stripe; within this border, the surface of the wing has a uniform brownish coloring, which is perceptibly more intense only in the region of the axillary incision; in some cases, near the posterior side of the sixth longitudinal vein, a little

beyond the end of the anal cell, there is a small, almost hyaline spot; the alula is almost hyaline, or infuscated towards the posterior margin only. The second longitudinal vein, opposite the posterior crossvein, shows a small fold, the tip of which, directed backwards, emits a short stump of a vein; the last section of the fourth longitudinal vein is very strongly curved; the posterior angle of the anal cell forms a sharp, but not very acute angle.

Hab. United States; Carolina (Zimmerman), Massachusetts (Harris), etc.

Observation.—I am not able to compare the figure of *Myopa nigripennis* Gray, but I do not hesitate, on Gerstæcker's authority, to place this name among the synonyms of *P. undata*. The synonymy of *Sphecomya undata* Harris is based upon a statement of Mr. Walker, who seems to have received specimens from the author.

3. *P. vespertilio* GERST. ♂.—Antennarum articulo tercio præcedente plus dimidio brevior, rotundato ovato, fusco, aristæ articulo primo brevissimo, secundo elongato: fronte oculis duplo latiore, palpis filiformibus: alis venâ longitudinali secundâ nec fractâ, nec appendiculatâ, alulâ strigisque duabus marginis posterioris hyalinis.

Third antennal joint not half so long as the second, rounded oval, brown; the first joint of the arista very short, the second elongated; front double the breadth of the eyes; palpi linear; second longitudinal vein of the wings without fold or stump of a vein; the alula and two stripes near the posterior margin hyaline. Long. corp. 0.64; long. al. 0.56.

Syn. *Pyrgota vespertilio* GERST. Stett. Entom. Zeitschr. xxi, p. 189, Tab. II, f. 8.

Head comparatively stouter than in the preceding species; front, when viewed from above, and taken as far as the anterior border of the eyes, at least by one-half broader than long; the gibbosity projecting over the eyes is not of equal breadth, as in *P. undata*, but conically attenuated anteriorly; its tip is as broadly truncated as in the other species; viewed in profile, this projection is as high as in *P. undata*; its anterior side, however, does not ascend in a straight line, but shows a strong convexity, so that the tip itself is retreating. The cheeks are considerably broader and more sunken. The eyes are comparatively smaller, the excavated upper part of the face perceptibly shorter. The coloring on the front, especially on the inside of the eyes and

upon the gibbosity, is darker, more brown; upon the cheeks, with the exception of the ferruginous-yellow border of the eyes, chocolate-brown; the two black ridges, bordering the middle of the face, are present, as in *P. undata*, but even more distinctly marked and descending lower. The first two joints of the antennæ are pale ferruginous-yellow; the third joint dark-brown; the arista ferruginous-yellow at the basis, whitish towards the tip; the second antennal joint is not quite as long as in the preceding species, chiefly because it is but very little less drawn out forwards above than below; the last joint is at least by one half shorter than the second, rounded oval, ending in a blunt point; the arista is inserted on the middle of its length, on the outside, near the upper margin; of its two basal joints the second has four times the length of the very short first joint. The palpi are slender, filiform, tinged with brown, like the proboscis. The thoracic dorsum shows three deep black stripes; the middle one is very broad, begins at the anterior margin and ends some distance before the scutellum; the lateral stripes are abbreviated anteriorly and posteriorly; the portion of them behind the suture is larger than that in front of it. The greater part of the pleuræ, a spot on each side at the posterior margin of the scutellum, as well as the metathorax, dark-brown. On the abdomen, the anterior part and the middle line of the first segment are pitch-black and somewhat shining; on each of the following three segments is a triangular spot, of the same coloring, the basis of which is directed anteriorly, and which occupies the whole breadth of the segment. The upper part of the abdomen has delicate transverse grooves, the under side on the contrary is strongly grooved in a longitudinal direction, opaque velvet-black, with a narrow, ochre-yellow middle line; the projecting male organ of copulation is of a shining reddish-brown. The feet are light ferruginous, with yellow hairs; the femora, to the exclusion of the tip and tibiæ, with the exception of the basis and of the extreme tip, are chestnut-brown. The second longitudinal vein of the wings is hardly perceptibly broken and without any vestige of a stump; the wings in general are comparatively shorter than in *P. undata*, darker and more evenly earth-brown; a very delicate streak near the posterior border of the first longitudinal vein, not far from the origin of the second vein, the whole alula and two streaks near the posterior margin, the position of which

corresponds to the entirely discolored spots in *P. undata*, are hyaline. These two streaks have a very definite outline, and the space beyond them is as dark-brown as the remainder of the wing; the longer one is almost straight, the shorter one sickle-shaped. The halteres are pale ferruginous-yellow.

Hab. Carolina (Zimmerman).

Observation.—The above description is the reproduction of that prepared by Dr. Gerstæcker, l. c., from a single specimen in the Berlin Museum. I have had a passing view of the specimen; it is very like *P. undata*. The differences in coloring, noticed by this author, are in my opinion of but little importance, as most of them occur among the varieties of the very variable *P. undata*. More important are the plastic differences, mentioned by Dr. Gerstæcker. Although the shape of the head in different specimens of *P. undata* is variable (evidently, however, in consequence of different degrees of shrinkage in drying), although the size of the third antennal joint is subject to slight variations, and although the relative length of the first two joints of the arista is not altogether constant, it is hardly credible that all these discrepancies should reach the degree which Dr. Gerstæcker noticed in his *P. vespertilio*.

4. *P. pterophorina* GERST. ♀.—Antennarum articulo tertio præcedente paulo longiore, oblongo ovato, aristâ brevissimâ, crassâ: fronte oculis latiore, fortiter prolongatâ, palpis cochlearibus; alis latis, venâ longitudinali secundâ geniculatâ, nec appendiculatâ, fuscis; alulâ, maculis duabus, posticis magnis, semilunaribus, guttisque duabus hyalinis.

The third antennal joint is somewhat longer than the preceding one, elongated-oval, with a very short, stout arista; front broader than the eyes, very much prolonged; palpi spoon-shaped; wings broad, with a second longitudinal vein which is geniculate, but has no stump of a vein upon it; coloring on the wings brown; alula, two large crescent shaped spaces on the posterior margin and two dots hyaline. Long. corp. 0.4; long. al. 0.44.

Syn. *Pyrgota pterophorina* GERST., Stett. Entom. Zeit. xxi, p. 190, Tab. II, f. 6.

Body small, slender, pale-ferruginous, shining. Head, viewed from above, by one-third longer than broad; front broader than the eyes, but, taken as far as the anterior margin of the eyes, nevertheless longer than broad; the gibbosity only a little shorter

and very little attenuated anteriorly; viewed in profile, this gibbosity is less elevated than in the two preceding species; on the contrary, it is, to its very much protruding tip, almost on the same level with the remaining portion of the front; this causes the anterior margin, which, with a slight convexity, is strongly retreating, to lie almost entirely on the under side; cheeks likewise broader and descending lower than in *P. undata*. The coloring of the head is altogether pale-ferruginous, even the black lines, bordering the middle portion of the face, are wanting. The antennæ likewise are altogether ferruginous-yellow; the two apical joints are almost of equal length; the third appears a little longer, only when viewed from the outside, along the lower margin, because, at this point, this joint is less covered by the second than above and on the inside; the first two joints are beset with blackish bristles, as in the two preceding species; the third joint is elongated-oval; the arista is inserted in the middle of its length, near the upper margin; it is stout and very short, shorter than the third antennal joint; the second joint of the arista is one-half longer than the first; the styliform third joint is but little longer than the first two taken together. Palpi elongated, slightly curved, somewhat spoon-shaped at the tip, pale ferruginous-yellow, with black bristles; the proboscis brown. Thorax uniformly ferruginous-yellow; clothed, as the head, with delicate black bristles. Abdomen of a similar color, but more shining, beset with long black bristles, forming bunches, especially on the sides; the upper side of the first abdominal segment is infuscated beyond the middle. The horny capsule, which forms the end of the fifth segment of the abdomen of the female, has, in profile, the appearance of a sparrow's bill; it is convex above, concave below, obtuse at the tip and somewhat shorter than the last three abdominal segments taken together. Feet perceptibly longer and more slender than in the two preceding species, with dense and rather long hairs, light brown; the basal third of the tibiæ and the tarsi pale yellowish; the hind tibiæ are much more incrassated toward the tip than the middle ones. Wings remarkably broad, obtusely rounded at the apex; the second longitudinal vein strongly bent and then broken in the shape of an angle, but without stump of a vein; ground color of a saturate earth-brown; a trapezoidal spot, extending from the

costa to the third longitudinal vein and situated before the break in the second vein, a round spot between both crossveins, the alula and two large crescents on the posterior margin hyaline; the crescents show a pale shade of brownish towards the posterior margin. Halteres altogether pale yellow.

Hab. Carolina (Zimmerman).

Observation 1.—The above is a translation of Gerstæcker's description of the specimens in the Berlin Museum. The species is distinguished enough to render the discussion of its specific rights useless. I will only notice here, that when the author says that the fifth abdominal segment in the female gradually passes into the capsule-shaped ovipositor, this expression is not to be understood literally; in the two species which I have seen, such a transition is not visible. When the author calls the first two antennal joints of *P. pterophorina* "beset with blackish bristles, as in the preceding species (*P. undata* and *vespertilio*)," I would observe that in *P. undata* this pubescence is in reality yellow, and assumes a ferruginous or even blackish tinge only when seen against the light.

Observation 2.—A fifth American species is described by Macquart (Dipt. Exot. Suppl. IV, p. 281, Tab. XXVI, f. 1) as *Oxycephala fenestrata*. His data are not even sufficient to ascertain whether the species really is a *Pyrgota*. Moreover it is not distinctly stated whether this species belongs to North America.

Second Section: PLATYSTOMINA.

Gen. I. AMPHICNEPHES nov. gen.

Charact.—*Front* of medium breadth, not narrower anteriorly.


Antennæ reaching down to the edge of the mouth.



Face excavated, without distinct antennal foveæ; occiput but little turgid; eyes high; cheeks narrow.

Scutellum large, flat, with four bristles.





Wings very broad; the longitudinal veins straight and conspicuously diverging; anal cell shorter than the preceding basal cell; its posterior angle rounded.

Small, metallic-colored species, the wings of which show a picture not unlike that of the species of *Platystoma*, and the general habitus of which is less like the species of *Rivellia* than





**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

15
16
18
20
22
25

10
11

those of *Platystoma*. They are, however, easily distinguished from the latter by the narrower front, the much less turgid occiput, the larger and flatter scutellum and the much broader wings, with straight, very much diverging longitudinal veins.

1. *A. pertusus* n. sp. ♂ and ♀.—(Tab. VIII, f. 1) Viridis, nitidus alae nigrae, guttis et fasciâ subapicali hyalinis.

Green, shining; wings black with hyaline dots and a hyaline crossband before the tip. Long. corp. 0.13—0.14; long. al. 0.11—0.12.

Dark metallic-green, shining. Head black; the front blackish-brown, even, rather long, but only of a medium breadth, not narrowed anteriorly; the ocelli are closely approximated to each other near the edge of the vertex; the small ocellar triangle and the little stripes running down at the corners of the vertex are of a shining blackish-green. Bristles of the vertex rather long, directed backwards; the bristle which is in front of them on each side is short; the ocellar bristles are not distinct. Antennae reaching down to the edge of the mouth, brownish-yellow; their narrow third joint is blackish at the tip; often the greater part of its outer side is brownish. Face excavated; its lateral portions very narrow; antennal foveae indistinct. The shining black clypeus broad. Palpi broad, shining black, with a paler border on the under side and at the tip; proboscis of moderate thickness; mentum but little swollen. Eyes much higher than broad; cheeks narrow; occiput but little turgid. Thorax very delicately transversely aciculate. Scutellum large, flat, weakly rugose, with four bristles. Abdomen more distinctly rugose. Ovipositor black, considerably extensile. Feet black, brownish-black in less mature specimens; the first joint of the front and hind tarsi and the first three joints of the middle tarsi of a dirty ochre-yellow. Halteres black, tegulae but little developed. Wings rather broad, black, more grayish-black near the hind margin; immediately before their apex is a conspicuous, arcuated, hyaline crossband; before this band there is a moderate number of hyaline dots of regular shape, which become more sparse towards the anterior margin; five dots which are nearest to the crossband form a row, parallel to the latter; the blackish-gray coloring near the hind margin of the wings has no hyaline spots. The veins are much more straight than in *Platystoma* and very diverging;

the two posterior basal cells are rather striking for their large size; however, the anal cell, which has an obtuse posterior angle, is shorter than the basal cell lying in front of it; the small cross-vein is in the middle of the discal cell; the first half of this cell is by no means attenuated, as is the case in the species of *Rivellia*.

Hab. Carolina (Zimmerman); Washington, D. C., Connecticut (Osten-Sacken).

Gen. II. **HIMEROËSSA** nov. gen.

Charact.—*Front* of equal breadth, distinctly projecting in profile.

Antennæ reaching almost to the mouth, arista bare.

Face moderately excavated, somewhat retreating below; occiput moderately turgid, eyes high; cheeks narrow.

Scutellum convex; with six bristles.

Wings: marginal and submarginal cells very narrow; second section of the fourth longitudinal vein straight; posterior crossvein prolonged beyond the fourth vein; posterior angle of the anal cell rounded.

As I have seen only a single species of this genus, the one which is described below, the definition of the generic character can naturally be only a provisional one. Should the peculiar prolongation of the posterior crossvein, which distinguishes *H. pretiosa*, be wanting in some allied species, it would then be necessary to omit this character from the definition of the genus; the remaining characters are amply sufficient for the purpose.

1. *H. pretiosa* n. sp. ♂.—(Tab. VIII, f. 2.) Rufo testacea, abdomine violaceo, pedibus anticis totis, posteriorumque tibiis et tarsis nigris; alæ hyalinæ, inæquali costæ limbo et fasciâ tenui subinterrupta nigro-fuscis.

Yellowish-red, with a violet abdomen; the front feet altogether, the tibiæ and tarsi of the four posterior feet, black; wings hyaline with an irregular costal border and a narrow, somewhat interrupted crossband, blackish-brown. Long. corp. 0.38, long. al. 0.3.

Yellowish-red, shining. Front darker, opaque, of equal breadth, with very indistinct traces of flat pits and a very delicate border of white pollen along the orbits; distinctly projecting in profile; the little stripes, descending from the vertex along the sides of the front, and the ocellar triangle are distinct, and somewhat more shining; the latter is somewhat larger than

usual; ocelli very near the edge of the vertex, rather large, but little approximated; the four bristles on the edge of the vertex rather strong; the lateral, as well as the ocellar bristles replaced by shorter, bristle-like hairs. Antennæ of the coloring of the body, almost reaching to the anterior edge of the oral opening; arista bare. Face but moderately concave, somewhat retreating on the under side, pollinose with white, except in the vicinity of the oral opening; in the well-marked foveæ this pollen is thicker and more conspicuous; the lateral portions of the face, bordering on the eyes, are very narrow and likewise clothed with white pollen. Eyes much higher than broad; cheeks narrow. Clypeus of a moderate breadth, distinctly projecting over the edge of the mouth; palpi not very broad, almost ferruginous. Proboscis rather stout; occiput moderately turgid. The whole thorax and the convex scutellum shining, with a very faint trace of a reddish metallic reflection. Scutellum with six bristles. Abdomen of a metallic reddish-violet coloring, which, in a different light, assumes upon the first three segments a bronze-green tinge; this is not the case with the last segment. Front feet with the coxæ brownish-black; on the four posterior feet the tibiæ and tarsi alone have this coloring; the coxæ and femora have the color of the thorax. Halteres yellowish-red, with an infuscated knob. Wings hyaline, with brown veins, which are not in the least sinuous; their anterior margin has a conspicuous, but unequal brown border, which, near the apex, extends as far as the fourth vein; from the root of the wing to the small crossvein, which is still included in this border, it becomes gradually broader and reaches here almost to the fifth longitudinal vein; it contracts immediately beyond the small crossvein, to the second longitudinal vein; opposite the posterior crossvein it expands again towards the third longitudinal vein, and runs immediately behind this vein as far as the apex of the wing, where it suddenly turns towards the fourth longitudinal vein, which forms the limit of this dark border; the very steep posterior crossvein projects in an unusual way beyond the fourth longitudinal vein; it is bordered with brown; this border forms a narrow, perpendicular cross-band, which growing paler and more indistinct, extends to the dark border of the anterior margin, or quite near it; the costal cell is clay-yellow, except at the basis and at the tip, which are more brownish. The marginal and submarginal cells are re-

markably narrow; the small crossvein is in the middle of the discal cell; the posterior angle of the anal cell is rounded and the last section of the fourth longitudinal vein is parallel to the third.

Hab. Cuba (Gundlach).

Gen. III. **RIVELLIA** R. DEBY.

Charact.—*Front* of equal breadth, not projecting in profile.

Antennæ usually reaching down to the margin of the mouth; third joint long and narrow; arista with a very short pubescence.

Face rather excavated, its lower part projecting; the lateral portions very narrow; clypeus broad; occiput moderately turgid; eyes high; cheeks moderately broad.

Scutellum convex, with four bristles.

Wings: Marginal and submarginal cells comparatively broad; the second section of the fourth longitudinal vein remarkably sinuate, with the convexity encroaching upon the discal cell, so that the latter appears much narrower before the small crossvein than behind it; the last section of the fourth longitudinal vein is parallel to the third vein or very slightly diverging; posterior angle of the anal cell rounded.

A large number of closely resembling species belong to this genus; the picture or the wings of most of them is nearly the same, so that this picture alone helps to recognize the species belonging here; it consists of four brown or blackish-brown crossbands; the first starts from the root of the wing and is the most oblique of all and the shortest; the second, somewhat longer and less oblique, runs over the small crossvein; the third, which covers the posterior crossvein, is perpendicular and reaches from the anterior to the posterior margin of the wing; the fourth starts from the anterior margin, near the origin of the third band, and forms a border along the apex of the wing. The North American fauna seems to abound in these species. The apparently total absence of plastic differences between them and the, as it seems, not unimportant variation in the coloring of some of them, render their separation very difficult, especially when there are only single specimens for comparison. I hope not to have gone amiss in the definition of those which I know. Whether I was mistaken or not, those may judge who have the opportunity of observing these species in life.

Among the species described below, *Rivellia conjuncta* is the

only one which does not belong to the difficult group just characterized; it is distinguished from it not only by a different picture of the wings, but also by some easily tangible plastic differences.

1st Group. *Crossbands contiguous near the posterior margin.*

1. *R. conjuncta* n. sp. ♀.—(Tab. VIII, f. 3.) Nigro-viridis, pedibus præter tarsorum basin nigris, tribus primis alarum fasciis postice coherentibus.

Blackish-green; the feet, with the exception of the root of the tarsi, black; the first three crossbands of the wings contiguous posteriorly. Long. corp. 0.16; long. al. 0.14.

Blackish-green, shining. Front moderately broad, dusky ferruginous-brown, almost black, laterally with a rather broad border, pollinose with white. Antennæ reaching almost down to the edge of the mouth, brick-red, except the third joint which turns brownish or blackish towards its tip. Face and clypeus metallic-black. Feet black; the basis of the tarsi brick-red or dirty reddish-yellow to a considerable extent. Halteres black. Wings hyaline; the four crossbands much broader than in the following species, especially the first; the second coalesces with the first in the discal cell and the third unites with the first near the posterior margin of the wing; the band which forms a border along the end of the anterior margin and the apex is connected in the usual way with the third, at the anterior margin. The small crossvein is but little beyond the middle of the discal cell; the second section of the fourth longitudinal vein is strongly arcuated, and the posterior crossveins bisinuate.

Hab. Maryland (Osten-Sacken).

2d Group. *Crossbands separated near the posterior margin.*

2. *R. viridulans* R. DESV. ♂ ♀.—(Tab. VIII, f. 4.) Nigro-viridis, interdum chalybescens, pedibus præter tarsorum basin nigris, primis tribus alarum fasciis separatis.

Blackish-green, sometimes more steel-blue; feet, with the exception of the root of the tarsi, black; the first three crossbands of the wings isolated from each other. Long. corp. 0.18—0.21; long. al. 0.15—0.2.

SYN. Rivetia viridulans R. DESV. Myod. p. 729, 2.

Trypeta quadrifasciata HARRIS, Cat. Ins. Mass.

Ortalis Ortoeda WALK. List, IV, p. 992.

Ortalis quadrifasciata WALK. List, IV, p. 993.

Herina rufitarsis MACQ. Dipt. Exot. Suppl. V, p. 123, 7.

Tephritis melliginis FITCH, First Rep. 65.

Blackish-green, shining; the upper side of the thorax sometimes less so; recently excluded specimens acquire a somewhat steel-blue tinge after drying. Front reddish-brown, often very dark, of the usual breadth, with a very narrow border of white pollen on each side. Face and clypeus metallic black; the narrow lateral portions of the face, bordering on the eyes, brownish-red, more seldom dark-brown. Antennæ reaching to the edge of the mouth, brick-red or yellowish-red; the third joint gradually turning black towards the tip. Ovipositor and feet black; the tips of the four anterior tibiæ usually brownish brick-red; the first joint of the two front tarsi and the first two joints of the four posterior tarsi pale brick-red. Crossbands of the wings black, rather narrow; the first three, which are entirely separated from each other, reach from the anterior margin to the fifth longitudinal vein; the fourth band, bordering the end of the anterior margin and the apex, is often connected with the third only by a rather narrow black border of the anterior margin; the portion of the costal cell between the first and the second crossbands has a dingy, somewhat yellowish appearance. The small crossvein is far beyond the middle of the discal cell and the second section of the fourth longitudinal vein is very much arcuated. Halteres black.

Hab. New York; Georgia; Distr. Columbia (Osten-Sacken).

Observation 1.—The attentive reader of Walker's description of *Ortalis Ortoeda* will easily notice that, before the end of the fourth line, previous to the comma, several words have been accidentally omitted, so that the end of the sentence does not refer, as it should, to the second, but to the third crossband. What Mr. Walker meant to say results sufficiently from the next description, that of *O. quadrifasciata*, which reproduces again the present, apparently very common, species. The fact that the measurements of *O. Ortoeda* and *quadrifasciata* are different in Walker does not prevent me from considering them as one and the same species. Under the former name Walker describes a male; under the latter, a female; hence, the greater size of the latter has nothing surprising. Instead of the length of the single

wing, Walker gives the breadth of the wings from apex to apex, a datum which is to be obtained only by approximation. This breadth in *O. Ortoeda* is said to be three, in *O. quadrifasciata* four lines, a difference which is somewhat considerable, but, owing to its uncertain nature, not to be relied upon exclusively for separating the two species, as the female of *R. viridulans* really has longer wings than the male. The quotation from Harris's Catalogue has been introduced upon the authority of Walker, who seems to have had original specimens of this author; but as the species has never been described, the quotation might as well have been omitted. That *Herina rufitarsis* Macq. belongs here is not doubtful. I have been able to compare a typical specimen of *Tephritis melliginis* Fitch.

Observation 2.—The following species agree so much with *R. viridulans* in the breadth of the front, the shape and the length of the antennæ, and in the venation, that every statement about these points would be useless. In speaking of the picture of the wings, a statement about the points of difference will be more useful towards discriminating the species than a detailed description.

3. *R. quadrifasciata* Macq. ♂.—(Tab. VIII, f. 5.) Thorace viridi, capite præter occiput, abdomine, pedibus, halteribusque luteis.

Thorax green; the head, with the exception of the occiput, the abdomen, the feet, and the halteres, dark-yellow. Long. corp. 0.2; long. al. 0.19.

SYN. *Herina quadrifasciata* Macq. Suites, etc., II, p. 433, *l.*

Head dark-yellow, the occiput metallic dark-green. Front dusky red, with a narrow border of white pollen on each side. Antennæ dark yellowish-red; the third joint, with the exception of the root, brown; blackish towards the tip. Palpi dark-yellow. Thorax, including the scutellum, of a blackish-green, metallic coloring, shining. Abdomen dark-yellow, more brownish-yellow towards its end. Coxæ and feet dark-yellow; hind tibiæ yellowish-brown; the last four joints of the front tarsi, and the last three, more seldom the last four, joints of the middle and hind tarsi infuscated. Halteres dark-yellow. The crossbands on the wings as narrow and nearly in the same position as in *R. viridulans*, but less dark; the first band is narrower and crosses the fourth longitudinal vein but very little; the second reaches not

quite as far as the fifth longitudinal vein; the hyaline interval between them is broader and the intervening portion of the costal cell of a darker coloring than in *R. viridulans*; the costa itself, from the extreme basis as far as about the middle of the costal cell, is of a dirty-yellowish coloring.

Hab. Nebraska (?). [I possess a specimen from Washington, D. C., which agrees exactly with the above description. O. S.]

4. *R. variabilis* n. sp. ♂.—(Tab. VIII, f. 6.) Rufo-testacea, capite pectoreque piccis, abdomine nigro-picco, basin versus plerumque dilutius picco, pedibus luteis, tibiis posticis tarsorumque apice fuscis.

Brick-red; head and chest pitch-brown; abdomen pitch-black, towards the basis usually of a lighter pitch-brown; feet dark yellow; hind tibiæ and the tip of all the tarsi brown. Long. corp. 0.18—0.21; long. al. 0.15—0.2.

Brick-red. Head pitch-brown or reddish-brown. Front of an opaque dark-red coloring, on each side near the orbit with a very narrow border of white pollen. Antennæ reaching down to the mouth; the first two joints dark reddish-yellow; the third, with the exception of the basis, dark-brown, blackish towards the tip. Palpi dark-brown. The chest and the lower part of the pleuræ dark pitch-brown. Abdomen pitch-black, generally lighter pitch-brown near the basis. Coxæ and feet dark-yellow; the four anterior tibiæ but little infuscated; the hind tibiæ and the last three or four tarsal joints dark-brown. Halteres dark-brown. The picture of the wings almost entirely like that of *R. quadrifasciata* in coloring and design, only the first two crossbands are a trifle longer and the first a little broader; the brown coloring in the anterior basal cell is a little less extended.

Hab. District Columbia (Osten-Sacken).

Observation.—I have a female, from the same locality, which I think belongs to the present species. It differs from the male, described above, in having the antennæ of an altogether dark-yellow coloring, except the slightly infuscated tip of their third joint; the color in the middle of the thoracic dorsum almost verges on blackish; the first crossband on the wings is a little longer, the front and middle tibiæ do not show any distinct infuscation and the tip of the tarsi is but little infuscated.

5. *R. flavimana* n. sp. ♂ ♀.—(Tab. VIII, f. 7.) *Viridi-nigra*, vel nigro-chalybea, pedibus anticis luteis, posterioribus semper ex parte, plerumque maximâ ex parte, nigris vel fuscis.

Greenish-black, or more bluish-black; the front feet dark-yellow, the hind feet partly, and usually for the most part, black or bluish-brown. Long. corp. 0.16; long. al. 0.14.

SYN. ? *Herina metallica* v. d. WULF. Tijdschr. voor. Ent. x, p. 154. Tab. V, f. 10.

Very like *R. viridulans*, but easily distinguished by its smaller size and the paler, although very variable, coloring of the feet. Metallic blackish-green or almost blackish-blue. Head shining black; occiput of a metallic greenish-black; front dusky reddish-brown, often blackish-brown, on each side near the orbit with a very narrow border of white pollen. The first two antennal joints brownish-red, the third blackish-brown or black. The coloring of the abdomen towards the tip, in the male, verges more on bronze-black; the only female which I can compare has no trace of this color. Fore coxæ and tibiæ yellowish; the upper side of the femora and the basis of the tibiæ very seldom show a trace of infuscation. The four posterior feet have the coxæ, femora, and tibiæ black or brownish-black, the tarsi yellow. The above-mentioned female has the tip of the middle femora and the middle tibiæ, with the exception of the dark-brown basal third, of a brownish-yellow color; the tips of the tarsi in this specimen are hardly infuscated at all, while the male specimens have the three or four terminal joints of the front tarsi and the last three or four joints of the middle and hind feet somewhat dark-brownish. Halteres brownish-black. The picture of the wings recalls, in design and coloring, that of *R. viridulans*, only the crossbands are a little narrower; in general also the second, and especially the first, reach less near the fifth longitudinal vein; the black coloring, which is apparent on the root of the anterior basal cell of *R. viridulans*, is wanting in *R. flavimana*, and this affords a good character for distinguishing the latter species from those allied to it.

Hab. Nebraska (Dr. Hayden).

Observation 1.—I possess a male, the four posterior feet of which, with the exception of the hind tibiæ, are yellow; it is also distinguished by the color of the antennæ, which are reddish-yellow as far as beyond the middle of the third joint, and by the

somewhat narrower crossbands. Nevertheless, I consider it only as a variety of *R. flavimana*, which seems to be very variable in the coloring of the feet.

Observation 2.—*Rivellia Boscii* R. Desv. cannot very well be identified with *R. flavimana*, as it is described as considerably larger than *R. viridulans*, whereas *R. flavimana* is distinctly smaller. I did not succeed in identifying this species of Rob. Desvoidy; his data concerning the coloring do not agree with *R. quadrifasciata* and *variabilis*, and *R. pallida* is still less to be taken into account.

Observation 3.—At first, while in possession of insufficient materials, I took *R. viridulans*, *quadrifasciata*, *variabilis*, and *flavimana* for varieties of the same species, and it is only later that more abundant materials convinced me that they are actually different, although closely allied, species. It is in conformity with my former view that I have identified with *R. viridulans* the *Herina metallica* described and figured by v. d. Wulp in the Tijdschrift voor Entomologie, x, p. 154, Tab. V, f. 10. If my present separation of these species be correct, the only ones which can be taken into consideration in interpreting Mr. v. d. Wulp's species are *R. Boscii*, *flavimana*, and perhaps *R. micans*. *R. Boscii* is so inaccurately described by R. Desvoidy that its identification is very difficult anyhow; but as this species is 3 lines long, that of v. d. Wulp only $1\frac{2}{3}$, I consider their identity as not probable. The assumption that my *R. micans* is the *Herina metallica* of v. d. Wulp is contradicted by the very brilliant metallic-green coloring of the former. Moreover, v. d. Wulp's figure does not show, at the basis of the first basal cell, the dark coloring existing in *R. micans*, which coloring has the same extent, although not the same intensity, as in *R. viridulans*. If the correctness of the figure of the wing of *Herina metallica* could be implicitly relied upon, its specific diversity from *R. micans* would be a matter of certainty. But in this case I would have also to admit that *H. metallica* does not coincide with any of the species of *Rivellia* known to me, as the said figure differs from those species, especially in the broad interval between the first and second crossbands, which does not occur to that extent in any of them. We are forced to assume, therefore, that the figure of the wing is only of an average correctness, and to pay attention, in its interpretation, to the principal features

only. If the want of a dark coloring at the basis of the first basal cell be singled out as a characteristic feature, the supposition suggests itself that the species is identical with *R. flavimana*, which also partakes of this character; the shortening of the first two crossbands, as well as the data concerning the size and coloring of *H. metallica*, do not contradict such an assumption; even the statement about the coloring of the feet could be applied to unusually pale specimens of *R. flavimana*, although I have never met with specimens of this degree of paleness. Hence, it appears not improbable, although far from certain, that *Herina metallica* is identical with *R. flavimana*.

6. *R. micans* n. sp. ♀.—Speciebus precedentibus minor, læte æneoviridis, nitida, pedibus omnibus luteis, fasciis alarum fusco-nigris.

Smaller than the preceding species, metallic-green, shining; all the feet saturate-yellow; the crossbands on the wings brownish-black. Long. corp. 0.13—0.15; long. al. 0.13.

Not reaching the size of *R. variabilis* and perceptibly smaller than the other preceding species; of a metallic-green, bright and shining coloring. The front, the lateral stripes on the face and the lower part of the occiput of a reddish-yellow, seldom of a brownish-red coloring; antennæ, as far as the basal third or the middle of the third joint, reddish-yellow; beyond that, brown. The abdomen shows a diluted, half-pellucid, reddish crossband at the place where the first and second segments are soldered together; in some cases this band is wanting. Coxæ and feet saturate-yellow, the former sometimes more brownish-yellow; the tarsi, towards their tips, are strongly infuscated. The picture of the wings, in its design, is not unlike that of *R. viridulans*, but is rather brownish-black than deep black; the dark crossbands are a little narrower, especially the first and second, so that the hyaline interval between them is comparatively broader, almost equal in breadth to the interval between the second and third bands (in *R. viridulans* the first interval is considerably narrower than the second); the first and second crossbands stop about the middle of the interval between the fourth and fifth longitudinal veins; however, single specimens occur in which they are shorter; in other specimens they reach very near the fifth vein; the third band, towards its end, is perceptibly narrowed. The second section of the fourth longitu-

dinal vein is less arcuated towards the small crossvein than in *R. ciridulans*. The coxæ and feet are dark-yellow; the hind tibiæ, towards their end, grow gradually, but very slightly, more brownish-yellow; the tarsi, from about the basis of the third joint, dark brown.

Hab. Texas (Belfrage).

Observation.—The present species differs from the preceding ones by the more pure and brilliant metallic-green color. Varieties of *R. flavimana*, with very pale feet, are nearest to it; but such specimens have at least the hind tibiæ, with the exception of the extreme basis and the extreme tip, brown. Moreover, they differ from *R. micans* by the coloring of the first basal cell, which is hardly perceptibly tinged with gray at its extreme basis only, while in the latter species it is infuscated up to the last third of the second basal cell.

7. *R. pallida* n. sp. ♂ ♀.—(Tab. VIII, f. 8.) Flavo-testacea, Rivelliæ micanti æqualis, reliquis speciebus minor, fasciis alarum nigrofuscis.

Yellowish brick-red, of the size of *R. micans*, but smaller than the other species; the crossbands of the wings blackish-brown. Long. corp. 0.14—0.15; long. al. 0.13.

Yellowish brick-red. Head concolorous; front more ferruginous; on each side with a narrow border of white pollen. Antennæ of the same color with the remainder of the body, only the third joint a little blackish at the extreme tip. One of my specimens has the first two segments of the abdomen black at the basis; but this color seems to have originated after death, being produced by the contents of the abdomen. Ovipositor not darker, or but a little darker, than the rest of the abdomen. Feet dark-yellow; last two, at the utmost last three, joints of the tarsi brown. Knob of the halteres brown. The picture of the wings reminds of that of *R. flavimana*, but instead of black it is blackish-brown; the costal cell is tinged with brown at the spot only where the first crossband has its beginning, elsewhere it is of a dingy yellowish; the root of the first basal cell shows, as in *R. flavimana*, no dark coloring; the first and the second crossbands usually reach very near the fifth longitudinal vein. This species is easily recognized by its smaller size and lighter coloring.

Hab. Washington, D. C. (Osten-Sacken.)

Gen. IV. STENOPTERINA Macq.

Charact.—*Body* long and narrow.

Head almost like that of *Dacus*; *front* of a considerable and equal breadth, somewhat projecting in profile; *face* somewhat excavated in profile, perpendicular towards the somewhat upturned anterior edge of the mouth, or but little projecting; the shallow *antennal foræ* long and narrow, not distinctly separated from the convex middle portion of the face; the lateral portions of the face very narrow; *clypeus* very large; eyes large; cheeks not very broad; occiput only moderately turgid.

Antennæ: The first two joints short; the third narrow and very long, generally reaching a little below the anterior edge of the mouth; arista apparently bare, or with a pubescence which is so short as to be almost imperceptible.

Thorax long and narrow; the transverse suture runs across the whole dorsum in the shape of a shallow depression; viewed laterally, the thorax appears remarkably attenuated towards its anterior end, as the pectus is truncated obliquely in front; *scutellum* with four bristles.

Abdomen remarkably narrow; the first segment more or less prolonged in the male.

Feet slender; the fore coxæ very long, inserted remarkably near the collum, and unusually movable at the point of insertion

Wings rather narrow; stigma long and narrow; small crossvein oblique, inserted more or less beyond the middle of the long discal cell; second section of the fourth longitudinal vein straight; posterior angle of the anal cell rounded; the picture of the wings consists chiefly in a dark border of the costa, reaching from the basis of the stigma to the apex of the wing, and in the darker coloring of the entire anterior basal cell, to which, in most of the species, is added a brown cloud along the posterior crossvein.

The great uncertainty which seems to have hitherto prevailed concerning the characters of the genus *Stenopterina* has induced me to enter in more detail about them than about the other genera. If my limitation of this genus be correct, it will contain only species closely related in their plastic characters. Their venation alone shows some differences; some species have the third and fourth longitudinal veins convergent towards their ends, the second longitudinal vein perceptibly shorter, more distant from the costa, and meeting it at a less acute angle; other species show the opposite of all these characters. As far as I can judge at present, the species of the former group seem to belong principally to the old world.

S. ænea Wied. and *brevipes* F. may be considered as the types of the genus.

1. *S. cærulescens* n. sp. ♂.—Viridi-chalybea, humeris concoloribus, thoracis dorso magis violaceo, halteribus nigris, alarum hyalinarum limbo costali inde a venæ auxiliariæ apice usque ad venam quartam pertinente, cellulâ basali primâ et venæ transversalis posterioris limbo fusco-nigris.

Greenish-steelblue, with concolorous humeri and the thoracic dorsum more violet; halteres black; wings hyaline, a costal border, reaching from the end of the auxiliary to the end of the fourth longitudinal vein, the first basal cell and a border along the posterior crossvein brownish-black. Long. corp. 0.32—0.39; long. al. 0.26—0.31.

Of a greenish-steelblue coloring, which on the abdomen has a somewhat stronger admixture of green and verges on violet on the thoracic dorsum; the humeral callosities and the pleurae have the same greenish-blue color. Head dark-yellow, almost brownish-yellow; clypeus and palpi of the same color; front strongly infuscated anteriorly, this coloring having more or less extent; at the bottom of each of the foveæ a distinct brownish-black longitudinal streak; first and second antennal joints, as well as the root of the third, to a greater or lesser extent, dark-yellow; the third joint, towards its end, becomes more and more brown, even brownish-black. The last abdominal segment is only a little shorter than the penultimate. The hairs on thorax and abdomen are whitish, with the exception of the few and comparatively short bristles on the posterior end of the thoracic dorsum and of the four bristles of the scutellum. The coloring of the coxæ and feet is very variable, as that of the front and of the antennæ; the palest specimens in my possession have brownish-yellow coxæ, more yellowish feet, with a dark metallic streak, reflecting greenish-blue, upon the anterior side of the hind femora, and with tarsi which are dark-brown towards the tip; the darkest specimens in my collection have metallic-black coxæ, the femora almost black, with a bright metallic bluish-green lustre, excepting the tips of all the femora, which are brownish-red, and of the brownish-red basis of the middle ones; tibiæ and tarsi dark brownish-red; the latter, towards their end, colored with brownish-black to a considerable extent. Halteres black, only the basis of their stem a little paler. Wings hyaline;

their brownish-black picture consists of a narrow border along the anterior margin, which reaches from the end of the auxiliary vein to that of the fourth longitudinal vein, in the darker coloring of the first basal cell, which even crosses a little the small cross-vein and in a narrow border along the posterior crossvein.

Hab. Texas (Belfrage).

Observation 1.—The South American *S. brevipes* Fab. is distinguished from the present species by the ochre-yellow color of the humeri and the ferruginous-yellowish color of the halteres.

Observation 2.—*Herina metallica* Macq. (Dipt. Exot. II, 3, p. 208), from Mexico, is evidently no *Herina* at all, but a *Stenopterina*. It would seem possible, therefore, that *Stenopterina cærulescens* is that very species. Many of the statements in Macquart's description agree with *S. cærulescens*. It must be borne in mind, however, that these statements refer for the most part to characters which a whole series of *Stenopterinae* have in common. The statement that the wings are yellowish is not applicable to *S. cærulescens*, and none of the varieties of this species which are in my possession have the black feet mentioned in Macquart's description of *H. metallica*. Nevertheless, I would not have doubted this synonymy if I had nothing but Macquart's description to consult. The figure of the wing, however, which Macquart gives (l. c. Tab. XXIX, f. 2) sets this supposition entirely aside, by showing an unusually broad dark border along the anterior margin, by which Macquart's species differs conspicuously from *S. cærulescens* and similar species with the ordinary narrow border of the anterior margin.

Gen. V. MISCHOGASTER MACQ.

Charact.—*Front* of a considerable, rather equal, breadth; the anterior ocellus rather distant from the two others.

Face excavated in profile, hardly projecting below.

Antennæ rather long; arista with a distinct pubescence.

Wings narrowed towards the basis; auxiliary and first longitudinal veins closely approximated; posterior angle of the anal cell rounded.

Abdomen narrow, still more attenuated towards the basis; first segment beset with strong bristles; ovipositor rather conical.

The characters, as given here, are very incomplete, and require an entire revision. Unfortunately, I had no specimen at

hand for comparison, and was obliged to write from memory. The bristles on the first abdominal segment, the distance intervening between the anterior ocellus and the posterior ones, and even the shape of the ovipositor remind very much of some genera in the group of *Richardina*, from which, however, *Mischogaster* is easily distinguished by the distinct bristles on the first longitudinal vein and the unarmed femora.

The typical species of the genus is the *Cephalia femoralis* Wied. No species from North America are as yet known.

Gen. VI. **MYRMECOMYIA** R. Desv.

Charact.—*Body* slender, not unlike that of an ant.

Head comparatively large; occiput conspicuously stout behind the vertex.

Front of a uniform, considerable breadth, very long and steep, so that the antennæ are below the middle of the head; the very large lateral stripes of the front have wrinkle-shaped cross impressions.

Antennæ reaching a little below the anterior edge of the mouth; arista with a rather short pubescence.

Front convex, not excavated in profile, but descending in an inclined plane; clypeus of a moderate transverse diameter; cheeks rather broad.

Thorax somewhat narrowed anteriorly; *scutellum* small, with two bristles.

Abdomen very much attenuated at the basis; the narrow first segment without bristles; about its middle it is so coarctate that its anterior portion assumes the shape of a knot.

Feet very slender.

Tegulae wanting; *wings* narrow, running into a point towards the basis, so that the posterior angle of the wing and the alula are wanting; auxiliary and first longitudinal veins closely approximated; the two posterior basal cells small; the posterior angle of the anal cell rather sharp.

The very peculiar structure of the head, the approximated ocelli, the absence of bristles on the first abdominal segment and its peculiar coarctation, sufficiently distinguish this genus from *Mischogaster*. The species upon which it was founded by R. Desvoidy are unfortunately unknown to me, so that I cannot affirm with certainty whether the characters as based by me upon the species described below would in all particulars apply to them. Judging by his statements, however, it seems very probable that the discrepancies are not important.

Myrmecomyia is not only very like *Cephalia* in appearance, but closely allied to it in reality. However, they may be distinguished by the presence, in *Cephalia*, of a mesothoracic bristle, and by the absence of the coarctation of the first abdominal segment, peculiar to *Myrmecomyia*. The alulæ and tegulæ in *Cephalia*, although small, are not wanting; the posterior angle of the wing, although very shallow, is likewise apparent.

M. myrmecoides Loew. ♂ ♀.—(Tab. VIII, f. 9.) Nigra, alarum hyalinarum imâ basi et apice extremo nigris.

Black; wings hyaline, extreme root and apex black. Long. corp. 0.25—0.27; long. al. 0.21.

SYN. *Cephalia myrmecoides* Loew, Wien. Ent. Monatschr. IV, p. 83.

Black, glossy. Head shining black, face and cheeks usually brown. The very broad and long front, descending in a steep slope, has a very narrow middle stripe of velvet black, which does not reach much beyond the middle of the front, but is connected by a furrow with the frontal fissure; the latter is not in the shape of an arc, but of an angle. Ocelli approximated to each other. The vertex bears two strong bristles, and on both sides of them two shorter ones; moreover, far back of the ocelli there are two small erect bristlets, while there are none in the immediate vicinity of the ocelli. The conspicuously large lateral parts of the front have irregular, wrinkle-like, transverse impressions, and along the orbits a very narrow border of white pollen. Antennæ long and narrow, reaching to the anterior edge of the mouth; the first two joints brownish-red, the third black; arista with a very short pubescence. Face convex, descending obliquely in profile, but not excavated; the anterior edge of the mouth not drawn upwards; antennal fovæ indistinct; the very narrow lateral parts of the face with a thin white pollen. Eyes higher than broad. Cheeks rather broad. Clypeus projecting over the anterior edge of the mouth, however its longitudinal diameter does not equal its moderate transverse diameter; the rather broad palpi blackish-brown. Thorax rather long and narrow, broader in the region of the wings than before and behind. Scutellum very small, convex, with two bristles. The metathorax descends in an inclined plane, and is conspicuously long; the pectus rises obliquely from the middle coxæ towards the front

coxæ. Thoracic dorsum with a thin gray bloom, the impressions indicating the lateral beginnings of the transverse suture are more densely pollinose; the pleuræ, above the middle coxæ, are clothed with a very dense white pollen. The shining black abdomen is much narrower at its basis; its first segment is longer than each of the following ones; about its middle it is so attenuated that its smaller anterior portion is knot-shaped, the larger posterior portion funnel-shaped; the last abdominal segment is somewhat shorter than each of the two preceding ones. The comparatively large hypopygium is usually pitch-brownish, seldom blackish; the first segment of the black ovipositor is flat and rather broad. Feet very slender; anterior coxæ yellow; the four posterior coxæ yellowish-red or chestnut-brownish; all are clothed with white pollen. Front feet brownish-yellow, with pitch-brown femora; the tarsi, from the tip of the first joint, are blackish-brown; the four posterior feet are brownish-black; the knees, the extreme tip of the tibiæ and the root of the tarsi brownish brick-red; in very pale-colored specimens the light coloring of the tarsi is much more extensive. Halteres black. No tegulæ. Wings hyaline, with delicate black veins; the wings, towards the basis, are very much attenuated, without any posterior angle and without alula; auxiliary vein short, very much approximated to the first longitudinal vein; the latter rather stout, very gradually merging into the costa, so that the stigma is narrow, linear; second longitudinal vein very long and straight; the last section of the third longitudinal vein gently inflected backwards, so that it strongly diverges from the second longitudinal vein and ends in the extreme apex; small crossvein perpendicular, inserted but little beyond the middle of the long discal cell; the last section of the fourth longitudinal vein rather straight, only very little convergent towards the third; posterior crossvein straight; the two posterior basal cells comparatively small; the posterior angle of the anal cell rather acute, but not pointed; the sixth longitudinal vein rather short, but reaching distinctly to the margin. The picture of the wings consists in an obscuration of the extreme root and the extreme tip; the first extends in the costal cell a little beyond the humeral crossvein; behind the first longitudinal vein, however, it reaches as far as the posterior basal cells; the obscuration of the apex has its greatest breadth at the end of the first posterior cell; it

hardly crosses the fourth longitudinal vein posteriorly; anteriorly it extends as a rapidly contracting border along the costa as far as the end of the second longitudinal vein, so that it has rather the shape of an apical spot than of an apical border.

Hab. Washington, D. C. (Osten-Sacken.)

Third Section: CEPHALINA.

Gen. I. **TRITOXA** nov. gen.

Charact.—*Body* slender; abdomen narrow at the basis; feet rather long, front tibiæ before the end of the upper side with a stronger bristlet. Hairs and bristles rather short; thoracic dorsum with bristles along the sides and upon its posterior margin only.

Antennæ long and narrow; the second joint short, arista with short hairs. Face almost shield-shaped, with rather indistinct antennal foveæ.

Palpi very broad; proboscis rather stout, mentum but little inflated. *Wings* cuneiform towards the basis, with a very narrow alula; second longitudinal vein not conspicuously arcuated; third and fourth irregular in their course, which causes the anterior basal cell to expand before its end; first longitudinal vein beset with bristles upon the greater portion of its course; crossveins approximated to each other.

This genus contains reddish-brown and black species, with dark wings, marked with three hyaline, oblique, more or less arcuated crossbands.

1. *T. flexa* WIED. ♂ ♀.—(Tab. VIII, f. 10.) Nigra, capite thoraceque interdum fuscis; alæ nigræ, fasciis hyalinis valde angustis secundâ et tertiâ arcuatis, hac ab alæ apice late remotâ, venâ transversâ posteriore subnormali.

Black, head and thorax sometimes brown; the wings black, with three very narrow hyaline bands, the second and third of which are arcuated; the latter is rather remote from the apex of the wing; posterior crossvein almost perpendicular. Long. corp. 0.24—0.28; long. al. 0.21—0.23.

SYN. *Trypeta flexa* WIEDERMANN, *Auss. Zweifl.* II, p. 483, 11.

Trypeta arcuata WALKER, *Ins. Saunders*, p. 383. Tab. VIII, f. 3.

Fully colored specimens are altogether deep black; in very pale specimens, on the contrary, the whole head, the thorax, and the feet, the latter usually with the exception of the upper side of the femora, are often brown; vestiges of this color frequently

occur in a greater or lesser measure on specimens the prevailing color of which is black. Most specimens have the greater part of the front brown, some reddish-brown; the usual coloring of the antennæ, also, is more brown than black, especially towards the basis. The pubescence of the arista is short, but distinct. The front has on both sides a very narrow, the face a broader, border of white pollen; the face, also, is slightly hoary with white, which is not equally distinct in all specimens, nor from all points of view; it is most perceptible around the antennæ. The rather indistinct pollen on the thoracic dorsum forms two rather broad parallel lines. The first segment of the flattened ovipositor resembles in its nature the preceding abdominal segments, to which it is closely applied; it is clothed, like those segments, with short, black hairs. The wings are strongly cuneiform towards their basis, and towards their tip they are rounded in such a manner that the extreme apex is much nearer the posterior than the anterior margin; the second longitudinal vein is slightly wavy upon the first two-thirds of its course; its strongest curvature is just above the small crossvein; the latter is rather oblique; the posterior crossvein, on the contrary, is steep, almost perpendicular, slightly bisinuated in the shape of an S. The color of the wings is black; only very immature or faded specimens have it brownish-black; the three usual crossbands have an almost whitish tinge, and are very narrow; the first among them is so oblique that it almost assumes the appearance of a longitudinal stripe; it starts at the basis of the third posterior cell, diverges gently and moderately from the fifth longitudinal vein, becomes more and more attenuated and pointed, and ends already some distance from the posterior margin; the second pale crossband, which likewise has a very oblique position, begins at the tip of the costal cell, just before the end of the auxiliary vein, and runs to the posterior angle of the discal cell; it is perceptibly more arcuated on its anterior than on its posterior portion; the third crossband, running from the anterior to the posterior margin, likewise has a very oblique position, although less so than the second; between the posterior margin and the third longitudinal vein its course is straight; from there to the anterior margin it is more and more arcuated; the distance between the third crossband and the apex of the wing is very large, as it almost equals one-third of the length of the wing. In the imme-

diate vicinity of the small crossvein the coloring of the wing is more ferruginous-brown than black, which is especially perceptible by transmitted light; specimens also occur which have other pale streaks in one or the other of the cells.

Hab. Northern Wisconsin River (Kennicott); Illinois (H. Shimer).¹

Observation.—Wiedemann probably prepared his description of *Trypeta flexa* from a very imperfectly colored specimen. A drawing of the wing, which I prepared some twenty years ago after an original specimen in the Berlin Museum, proves conclusively that *Trypeta flexa* is distinct from *Tritoxa incurva* and *cuneata*. The former is proved by the dark coloring at the tip of the wing having a much greater extent than in *T. incurva*, and by the course of the third crossband in *T. flexa*, which is not arcuated towards its end, but almost straight; in *T. cuneata* the different shape of the wing and the entirely distinct delineation of the crossbands altogether exclude the possibility of its synonymy with *T. flexa*. The figure of the wing drawn by me and above alluded to agrees with the present species so well that I consider my opinion about the identification of this species as well founded. Should this not be the case, then *T. flexa* Wied. is a species which I do not possess. The statement of Wiedemann, that the ovipositor of the female is two jointed, rests upon an error, which is easily explained away by the resemblance of the first joint with the preceding abdominal segment. That Walker's *Trypeta arcuata* is synonymous with the present species is not in the least doubtful, although in the figure of the head the arista is made too short and its pubescence too long.

2. *T. incurva* n. sp. ♂ ♀.—(Tab. VIII, f. 12.) *Badia*, abdomine nigro; alæ fuscæ, fasciis hyalinis modicæ angustis, secundâ et tertiâ arcuatis, hæc ab alæ apice minus late quam in speciebus reliquis remotâ, venâ transversâ posteriore obliquâ.

Reddish chestnut-brown, with a black abdomen; the wings brown, with

¹ Mr. H. Shimer, from Mt. Carroll, Ill., informed me, in 1865, that this fly is very injurious to onion-plants, the larva occurring in the bulb. This fact has, since then, been mentioned in the *Practical Entomologist*, I, p. 4; II, p. 64 (with figures of larva and imago); *American Entomologist*, II, p. 110. Specimens of *Tritoxa incurva* were found by Mr. Shimer, together with *T. flexa*, and taken for a mere variety of that species.

only moderately narrow hyaline bands, the second and third of which are arcuated; the latter is less remote from the apex of the wing than in the other species; posterior crossvein oblique. Long. corp. 0.25—0.3; long. al. 0.22—0.26.

Reddish, chestnut-brown, with a black abdomen. Front opaque, with the exception of the edge of the vertex and of the small callosities descending from it, and bearing the bristles; along the orbits the front has a narrow border of white pollen, which also extends over the face, but is much broader here. The remainder of the face has a very thin, somewhat yellowish pollen upon it, which is most perceptible in the proximity of the antennæ. Antennæ reddish-brown; third joint darker brown towards its end; pubescence of the arista short, but distinct. The thoracic dorsum has a broad shining border upon its sides, otherwise it is opaque. Its thin whitish pollen is a little more perceptible than in *T. flexa*, and forms, as in that species, two broad, parallel longitudinal stripes, the position of which corresponds to that of the intervals between the ordinary thoracic stripes; upon the intermediate stripe between them the pollen has a somewhat yellowish tinge, and is much more dense upon the longitudinal line, which divides this stripe in two; well-preserved specimens show the white pollen on the sides of the thoracic dorsum also, while in less good specimens this is not visible, and often very little of the pollen is left on the whole surface. Scutellum, metanotum, and pleuræ are shining, the latter with a thin white bloom. Abdomen black or brownish-black, with a black pubescence, sometimes chestnut-brown on the sides of the first and second segments. The flattened first joint of the ovipositor is of the same nature as the preceding segments of the abdomen; it is very broadly truncated at the tip. The feet have the same coloring as the thorax, often, however, not only the upper side of the fore femora, the middle femora towards their basis, and the hind femora, with the exception of their last quarter, are more strongly infuscated, but also the fore tibiæ towards their tip, as well as the entire fore tarsi; the middle tarsi, with the exception of their basis and the entire hind tibiæ and hind tarsi, are dark brown. Halteres yellowish. Wings narrowed towards the basis, although not quite as cuneiform as in *T. flexa*, the portion lying beyond the sixth longitudinal vein not being quite as narrow as in that species; the end of the wing

is rounded in such a manner that the apex is equidistant from the anterior and the posterior margins; the second longitudinal vein, the course of which is rather wavy, has its strongest sinuosity only little beyond the small crossvein; the anterior end of the latter is nearer to the root of the wing than its posterior end, so that its position is entirely oblique; the posterior crossvein is oblique in the opposite direction, as its anterior end is nearer to the apex of the wing than the posterior. The coloring of the surface of the wing is a brown of unequal intensity; the design consists of the three hyaline bands usual in this genus; the portion of the surface of the wing beyond the third band is dark brown, with a large yellowish-brown spot, which leaves in the submarginal cell only a dark brown border along the margin of the wing, and, so far as it extends in this cell, also somewhat crosses the third longitudinal vein; the interval between the second and third bands, which has the shape of a crossband, is yellowish-brown, margined with dark brown on each side, and also dark brown at the end; the interval between the second and first crossbands is dark brown, with a large yellowish-brown spot, which fills up the basis of the submarginal cell, and, to a great extent, that of the first basal cell, so that in the former almost nothing is left of the dark brown color, in the latter only a border; the root of the wing is tinged with yellowish-brown as far as a little beyond the humeral crossvein; towards the place of insertion of the wing, however, the dark brown color appears again; the posterior angle of the wing, lying behind the first crossband, is only tinged with gray. The hyaline crossbands are distinctly broader than in *T. flexa*, and the last of them is much nearer the apex, so that the dark coloring of the latter assumes the shape of a broad crossband. The first hyaline crossband is so oblique that it almost assumes the appearance of a longitudinal stripe; it starts, as in *T. flexa*, from the basis of the third posterior cell, but is broader than in that species, and does not diverge from the fifth vein; gradually becoming more pointed, it ends some distance from the margin of the wing, and differs but little in intensity of coloring from the gray posterior angle of the wing; the second pale crossband, which has a very oblique position and is only gently curved, runs from the tip of the costal cell to the posterior corner of the discal cell; however, the tip of the costal cell itself is hyaline to a

very small extent only, so that the crossband appears somewhat abbreviated near the anterior margin of the wing; the third hyaline crossband, which is almost as oblique as the second, is more curved upon its posterior than upon its anterior portion.

Hab. Illinois (Dr. Schimer).¹

3. *T. cuneata* n. sp. ♂ ♀.—(Tab. VIII, f. 11.) Rufo-badia, abdomine nigro; alæ fuscæ, fasciarum hyalinarum secundâ obliquâ et levissime arcuatâ, tertiâ subnormali et rectâ.

Reddish chestnut-brown, with a black abdomen; wings brown, their second hyaline crossband oblique and only gently curved; the third almost perpendicular and straight. Long. corp. 0.23—0.25; long. al. 0.21—0.22.

Reddish chestnut-brown, with a black abdomen. Front opaque, however, with the exception of the edge of the vertex and of the two callosities, descending from it, and bearing the strong frontal bristles, of a rather reddish coloring; with a very narrow border of white pollen near the orbit; this border also extends over the face, but is not very perceptible here. The remainder of the face is covered with a very delicate whitish pollen, which is more perceptible near the antennæ only. The third antennal joint, with the exception of its basis, brown; arista with a very short, yet distinctly perceptible, pubescence. Thoracic dorsum upon its sides with a broad shining border, otherwise opaque; the rather whitish pollen which covers it is very distinct in well-preserved specimens, but even in such specimens it does not form any distinct longitudinal stripes. Scutellum, metathorax, and pleuræ shining, the latter with a white bloom. Abdomen black or brownish-black, with a black pubescence, usually reddish chestnut-brown upon the sides of the first and second segments. The feet are of the color of the thorax; the fore tarsi usually altogether dark brown; the middle and hind tarsi towards their end dark-brown to a great extent. Halteres yellowish-white. Wings comparatively narrower than in *T. incurva*, attenuated to a rather cuneiform shape towards their basis; second longitudinal vein only slightly wavy; the small crossvein very steep, almost perpendicular; the posterior crossvein oblique, its anterior end somewhat nearer the apex of the wing, so that the posterior angle of

¹ *Tritoxa incurva* occurs together with *T. flexa*, so that Dr. Schimer, who sent me specimens of both, took it for a mere variety of his *onion-fly*.—O. S.

the discal cell is a little larger than a rectangle. The coloring of the surface of the wing is an uneven brown; the design is formed of the usual three hyaline crossbands, the first of which, however, is but little apparent. The portion of the surface of the wing lying beyond the last hyaline crossband is rather dark-brown, more brownish-yellow towards the anterior more grayish-brown towards the posterior margin; the interval between the third and second bands is dark-brown below the fourth longitudinal vein, above it, yellowish-brown with dark-brown borders; the latter are broader, even sometimes coalescent, within the submarginal cell; the interval between the second and the first hyaline crossbands is dark-brown, its inner portion more yellowish-brown; the basis of the wing yellowish-brown; beyond the fifth longitudinal vein the brown coloring still continues, but soon verges on grayish. The first crossband has the same position as in the preceding species; only it is broader, less attenuated, and much shorter; its outline can be plainly visible only when the surface of the wing is viewed in an oblique direction; the second pale crossband, which is very oblique, begins below the tip of the costal cell, in the marginal cell, and reaches as far as the fifth longitudinal vein, which it touches already before the posterior corner of the discal cell; this band is but little curved; about its middle, it is more or less expanded in the shape of an angle, in consequence of its margin (the one nearest to the apex of the wing), between the third and fourth longitudinal veins, not running in the direction of the band itself, but being more or less perpendicular to the axis of the wing; the third hyaline band, running at some distance from the apex of the wing, is very steep, but by no means entirely perpendicular, and somewhat broader anteriorly than posteriorly; it begins at the anterior margin and completely or almost completely reaches the posterior one.

Hab. Nebraska (Dr. Hayden).

Gen. II. **CAMPTONEURA** Macq.

Charact.—*Body* slender, feet rather long; the hairs very short everywhere; the thorax with bristles on the lateral and posterior margins only.

Antennæ long and narrow; the second joint short. Face almost shield-shaped, convex, with rather indistinct foveæ.

Palpi very broad. Proboscis rather stout, with a but moderately turgid mesothum.

Wings broad, first longitudinal vein provided, to a great extent, with bristles; second longitudinal vein arcuated in a very striking manner; anal cell rounded at the tip; the anterior margin of the wings, at the end of the auxiliary vein, has a shallow, but very striking excision.

1. C. picta FABR. ♂ ♀.—(Tab. VIII, f. 13.) Badia, abdomine nigro; alæ nigro-fuscæ, maculis costalibus binis trigonis, binisque guttulis discoidalibus, marginis denique postici maculâ trigonâ et strigâ obliquâ hyalinis, angulo postico et alulâ omerascentibus.

Chestnut-brownish with a black abdomen; the wings blackish-brown; two triangular spots on the anterior margin, two dots on the middle of the wing, a triangular spot and an oblique streak beginning at the posterior margin, hyaline; posterior corner and alula grayish. Long. corp. 0.25; cum terebrâ 0.32—0.34; long. al. 0.22—0.25.

Stn. *Musca picta* FABRICIUS, Ent. Syst. IV, p. 355, 175.

Dictya picta FABRICIUS, Syst. Antl. p. 330, 18.

Tephritis conica FABRICIUS, Syst. Antl. p. 318, 10.

Trypeta picta WIED. Auss. Zweill. II, p. 489, 20.

Delphinia thoracica R. DESVOIDY, Myod. p. 720, 1.

Camptoneura picta MACQ. Dipt. Exot. II, 3, p. 201. Tab. XXVII, f. 4.

Trypeta picta WALK. List, IV, p. 1041.

Head and thorax chestnut-brownish or reddish chestnut-brown; thoracic dorsum sometimes darker brown; abdomen always black or brownish-black. Front opaque, usually more ferruginous-red than orange-red, sometimes darker, with a very narrow border of white pollen along the orbits; this border also extends over the face, but although broader here, it is less distinct, or at least more perceptible only a little distance below the antennæ. The remainder of the face is a little pollinose in the vicinity of the antennæ only. The third antennal joint is usually strongly infuscated, with the exception of its basis. Thoracic dorsum with a grayish-white pollen, which does not form any distinct stripes, while the ground color of the broad intermediate stripe is often darker than its surroundings, so that it becomes distinctly visible. Feet yellowish-brown, tarsi strongly infuscated towards their tip. Halteres whitish-yellow. Wings comparatively large and broad with a rather strongly projecting posterior angle, and a rather narrow alula; at the anterior margin there is an excision, which is very conspicuous, although it forms only an obtuse angle; it is

caused by considerable sinuous expansion of the costal cell; the second longitudinal vein is very conspicuously arcuated; the two crossveins are rather approximated and perpendicular, the posterior one somewhat curved; the posterior angle of the discal cell is acute. The coloring of the wings is blackish-brown, more yellowish-brown near the root, grayish in the posterior angle; on the anterior margin there are two triangular hyaline spots, which attain the third longitudinal vein more or less completely with their very sharp points; the first of these spots covers, near its anterior end, the tip of the costal cell and the basis of the stigma, while the second is immediately beyond the stigma; the dark crossband between these two spots is tinged with brownish-yellow inside of the marginal cell, with the exception of a brown border, which becomes narrower towards the first longitudinal vein. The stigma, towards its end, gradually assumes the same brownish-yellow coloring, so that the first hyaline spot has no well-defined limit within it. Upon the middle of the wing there are two hyaline drops, elongated in a direction perpendicular to the axis of the wing; the one is in the discal cell, somewhat this side of the small crossvein, the other in the first posterior cell, over the posterior crossvein. On the posterior margin of the wing, in the second posterior cell, there is a triangular spot, concave towards the apex of the wing, convex on the other side, which is near the posterior crossvein and separated by a narrow, brownish border from it. The sharp point of this spot is directed towards the dot in the first posterior cell, and is often connected with it, while, in other specimens, it does not even reach the fourth posterior vein. Near the basis of the wing there is a narrow, oblique, hyaline streak, beginning in the first basal cell, crossing the end of the second basal cell and entering the third posterior cell; here it runs along the sixth longitudinal vein and thus reaches the margin of the wing, where it becomes a little grayish.

Hab. United States, common.

Observation.—The description which Fabricius gives of his *Musca picta* in the *Entomologia Systematica* might suggest doubts as to its identity with the above described species, doubts, however, which I hold to be without foundation. First of all, it is certain that Wiedemann's *Trypeta picta* is identical with our species; his description, as well as the types of his collection,

proves it conclusively. Not less certain, according to my opinion, is the fact that Wiedemann's *Trypeta picta* and the *Tephritis conica* of Fabricius's *Systema Antliatorum* are synonyms. What Wiedemann says about the feet of his *Trypeta picta* clearly proves that he had examined the type in Fabricius's collection; moreover, Fabricius's description contains nothing to render this identification of *Tephritis conica* doubtful. In the preface to his first volume, Wiedemann gives a large number of synonymic and systematic emendations, the result of the examination of Fabricius's collection, undertaken by him; among them we find the statement that *Tephritis conica* and *Dictya picta* are the same species. But as *Dictya picta* of the *Systema Antliatorum* is nothing else but the *Musca picta* of the *Entomologia Systematica*, the synonymy of *Musca picta* F. with *Trypeta picta* Wied. and the above described *Camptoneura picta* seems to be sufficiently established. The correctness of this view seems confirmed by the fact, that *Musca picta* F. was described from a North American specimen, and that hitherto, besides *Camptoneura picta*, which has a wide range and is a common species, no other North American species is known which might come in conflict with it.

Gen. III. DIACRITA GERST.

Charact.—*Body* rather robust. Pubescence everywhere very short; thorax with some bristles upon the posterior and lateral margins only.

Antennæ of medium length; the oval third joint longer than the only moderately sized second joint. The face, retreating above between the rather short antennal foveæ, and obtusely carinate; below, it is again projecting and convex.

Palpi rather large, mentum swollen.

Wings narrow and long, the first longitudinal vein bristly at its end only; the third and fourth longitudinal veins converging towards the end; posterior angle of the anal cell drawn out in a very long point.

This genus contains brown or brownish-yellow species, rather opaque on account of the pollen which covers them; the thorax is usually spotted with black; the wings, on the anterior margin and the apex, have a broad black border.

I. D. costalis GERST. ♂.—(Tab. VIII, f. 14.) Fusca, polline cinereo aspersa. thoracis maculis nigris ante suturam sex, pone suturam duabus, binisque minutissimis utrinque adjectis.

Almost chocolate-brown, with a grayish pollen; thoracic dorsum with six black spots before the suture and with two beyond it, to which are added on each side two very small dots. Long. corp. 0.32; long. al. 0.37.

Syn. *Diacrita costalis* GERST. Stett. Ent. Zeitschr. xxi, p. 197. Tab. II.

Almost chocolate-brown, covered with a whitish-gray pollen and opaque. Head dark-yellow, the upper part of the occiput generally brownish-yellow; the broad front, in the vicinity of the ocelli and in front of these, more reddish-yellow; on both sides, near the orbit, there is a rather large, shallow impression, covered with white pollen; on the anterior end of the front there is a small triangular spot, covered with snow-white pollen. Immediately below each of these spots, upon the face, there is a velvet-black round spot, contiguous with the orbit, and immediately below the latter a spot covered with snow-white pollen. The upper part of the face, which is carinate and retreating, has, on each side, a transverse spot, clothed with white pollen. In the same way, the posterior orbit of the eyes has a pollinose white border, which also extends over the cheeks in the shape of a stripe. The antennæ are almost ochre-yellow, their third joint elongated-oval; the basal joint of the arista is so short as to be almost imperceptible; the second joint is comparatively long, both dark ochre-yellow; the third joint is blackish, with the exception of its extreme basis; in the vicinity of the basis, it is as stout as the first two joints, more attenuated afterwards, and clothed with an extremely short pubescence. The humeral callosities are brownish-yellow, and rather shining; thoracic dorsum marked with moderately large, rounded-oval, brownish-black spots; before the region of the transverse suture there are six of them, arranged in two regular transverse rows; beyond this region there are two approximated spots, the interval of which is equal to that between the spots of the first two rows; moreover, behind the region of the suture, on each side, may be noticed two very small, almost punctiform dots, placed one behind the other; of these, the anterior one is situated before, the posterior one at an equal distance behind the last two of the larger spots. The coloring of the convex scutellum, which is beset with four, not very long bristles, approaches the chestnut-red. The feet are concolorous with the remainder of the body; an admixture of yellow is perceptible on the first joint of the tarsi only. Halteres whitish-

yellow. Wings comparatively long and narrow, of a very equal breadth, in the middle only a little broader than at the basis and at the apex; stigma strikingly long; the third longitudinal vein gently curved backwards towards the tip, and hence, the submarginal cell very much expanded towards its end; the crossveins very distant from each other; the fourth longitudinal vein, towards its end, gently bent forward, and hence, the first posterior cell narrowed towards its end; the posterior angle of the anal cell is drawn out in a narrow lobe, which is considerably longer than the cell itself. The surface of the wing is bright, shining, hyaline, upon its posterior half only with a weak trace of a grayish-brown tinge. The design on the wing consists of a broad, black, or blackish-brown border of the costal margin and of the apex; the posterior limit of this border runs, at the basis of the wing, along the fifth longitudinal vein; at the basis of the discal cell, it suddenly turns towards the fourth longitudinal vein, and, after running alongside of it for a short distance, it turns suddenly towards the third longitudinal vein, alongside of which it runs as far as a little beyond the small crossvein, here, just opposite the end of the first longitudinal vein, it abruptly turns towards the second longitudinal vein, leaves open a small segment of a circle just above it, returns towards the second vein, follows it for some distance, and, abruptly turning again, crosses the submarginal and first posterior cells, turning towards the apex in the vicinity of the fourth vein, alongside of which it reaches the margin. This border is perceptibly broader at the tip than along the anterior margin, and can therefore also be described as a large spot, entirely confluent with the border along the anterior margin. Inside of the dark anterior border, there are three small, almost hyaline spots; the first lies at the end of the second basal cell, the second, almost euneiform, is in the marginal cell, before the origin of the third longitudinal vein, the third at the extreme tip of the costal cell; in the marginal cell, beyond the end of the first longitudinal vein, between the small hyaline spot in the shape of a segment of a circle and the costa, there is a spot, tinged with yellowish-brown; the broad black border along the apex is sometimes a little diluted in its middle.

Hab. Mexico (Germar).

Observation.—In the register of the second part of Wiedemann's *Aussereur. Zweifl. Insecten*, there is a *Platystoma costalis*,

which is not described in the work itself. Wiedemann's collection proves that this species is identical with the present one.

2. *D. annula* n. sp. ♀.—(Tab. VIII, f. 15.) Lutea, thoracis dorso fusco, maculis nigris ante suturam quatuor, pone suturam nullis.

Clay-yellow, dorsum of the thorax brown, with four brown spots before the transverse suture and none beyond it. Long. corp. 0.25; cum terebrâ 0.36; long. al. 0.31.

Very like the preceding in all plastic characters. Almost more ochre-yellow than clay-yellow, the thoracic dorsum alone strongly infuscated. The front, as in *D. costalis*, has on each side, near the orbit, a shallow impression, clothed with white pollen; below it is a round, velvet-black spot, and immediately below the latter again a spot of snow-white pollen, only the black spot is smaller than in the preceding species; also the two snow-white transverse spots on the upper part of the face are apparent, as in *D. costalis*. On the thoracic dorsum there are not six, but only four rounded oval velvet-black spots before the transverse suture, which correspond to the outward ones of the preceding species; there is no trace of black spots on the other side of the suture. The scutellum is convex and has four bristles; the large first segment of the flattened ovipositor is brownish-yellow, long, only moderately attenuated towards its end. Feet of the same coloring with the remainder of the body; the tarsi only moderately infuscated towards their end. Halteres whitish-yellow. Wings of the same shape as in *D. costalis*, only less long, especially their second half less elongate, so that the small crossvein is somewhat nearer the tip of the wing than in *D. costalis*, and that the last section of the longitudinal veins, ending in the apex of the wing, is shorter; otherwise the venation almost entirely agrees with that of *D. costalis*. The surface of the wing is hyaline; its posterior half strongly tinged with a smoky-brownish. The brownish-black design resembles that of the preceding species, differs, however, from it by the dark border along the apex being much narrower; the posterior limit of the border along the costa is also similar to that in the preceding species, but not quite identical; especially where, in *D. costalis*, this limit crosses the second longitudinal vein and leaves on the other side a hyaline segment of a circle; instead of the latter there is here only an indistinct paler dot and between this and the costa no spot of a paler coloring; the three

hyaline dots, contained within the black border of the costa, are much less clear in the present species, especially the first and the third among them.

Hab. California (Agassiz).

Gen. IV. **IDANA** nov. gen.

Charact.—*Body* robust. Hairs very short everywhere; thorax with bristles on its posterior and lateral borders only.

Antennæ of medium size; third joint oval, but little longer than the rather large second joint. Face obtusely carinate between the very long antennal foveæ.

Palpi of moderate size; the mentum moderately turgid.

Wings narrow and very long; first longitudinal vein towards its end provided with bristles to a considerable extent; third and fourth longitudinal veins converging towards their end; anal cell not drawn out in the shape of a lobe.

This genus contains conspicuous pollinose species; their thorax is marked with distinct black stripes and the abdomen banded with black, the design of the wings not unlike the genus *Pteropœcila*, while the general shape of the body reminds of the true species of *Ortalis*.

1. I. marginata Say. ♀.—(Tab. VIII, f. 16.) Alæ colore fusconigro et luteo pulchre variegatæ, inâ cellulæ marginalis basi, triente apicali cellulæ basalis primæ, cellulisque posterioribus duabus primis præter venarum limbos pure hyalinis, angulo postico et cellulâ posteriore tertiâ fere totâ cinereo-hyalinis.

The wings with a handsome brownish-black and brownish-yellow picture; the extreme basis of the marginal cell, the last third of the first basal cell, as well as the first two posterior cells, pure hyaline, with the exception of the borders of the veins, inclosing them; the posterior angle and the greater part of the third posterior cell grayish hyaline. Long. corp. 0.34; cum terebrâ 0.45; long. al. 0.46 lin.

Syn. *Ortalis marginata* SAY, Journ. Acad. Phila. VI, p. 183, 2.

Head reddish-yellow. Front orange-yellow, opaque, with the exception of the immediate proximity of the ocelli and of the two little callosities, descending from the vertex and bearing the frontal bristles; the sides more orange-red, usually infuscated above the antennæ; on each side a rather narrow border of yellowish pollen. Antennæ of medium length; the first two joints of the coloring of the head; the second rather large; the third almost orange-yellow, of an oval shape, and but little longer than

the second; arista of medium length, with a short, but distinct pubescence. The vertical diameter of the eyes more than twice the length of the horizontal one. Face with very deep and long antennal foveæ, which run down in a perpendicular direction; their bottom is tinged with brownish-black. The face, between the foveæ, is strongly, the lower part sharply carinate, and that in such a manner that in profile the face runs down perpendicularly and in a straight line. Cheeks broad; at the lower corner of the eye with an infuscated spot. Oral opening rather large, somewhat drawn up above, so that the strongly developed, although transversely narrow, elypeus, projects a great deal beyond the peristomium. The reddish-yellow palpi rather large, broader towards the end; the brown proboscis of medium stoutness and the reddish-yellow chin only moderately swollen. The whole occiput is strongly and evenly convex. Thorax comparatively stout, but not strongly convex, distinctly narrowed anteriorly. Thoracic dorsum with a very dense, almost ochre-yellow dust, and with well-defined black longitudinal stripes; lateral border, and usually also the anterior one, chestnut-brownish or more chestnut-red; the intermediate stripe, running at an equal breadth from the anterior to the posterior border, is divided in two halves by a stripe-shaped intermediate line, which is of the same breadth with both halves of the intermediate stripe itself; the lateral stripes, which are but very little abbreviated anteriorly and posteriorly, are crossed by the yellowish-pollinose transverse suture; their posterior part moreover has alongside of it a black longitudinal stripe, which is not distinctly separated from the anterior part of the lateral stripe. Pleuræ chestnut-brownish, about their middle with a broad longitudinal stripe, which is clothed with pale ochre-yellowish pollen and gradually disappears posteriorly. Scutellum brownish-yellow. Abdomen black, but little shining, more or less chestnut-reddish at the extreme basis and on the sides of the first two segments; the second and each of the following segments have a crossband, of a dingy ochre-yellow, very thickly laid dust, occupying almost the whole of their anterior half, and narrowed on each side. The first segment of the ovipositor is black, flat, broad, nevertheless strongly attenuated towards its end. Feet brownish-yellow, tarsi strongly, but gradually infuscated towards the end. Halteres yellowish. The wings strikingly elongated, of a comparatively small and rather

equal breadth; stigma rather long but not broad; the crossveins far distant from each other; the posterior crossvein rather oblique, its anterior end nearer the apex of the wing than the posterior end; fourth longitudinal vein strongly bent forward towards the end; the first posterior cell considerably narrowed in consequence towards the apex; posterior angle of the anal cell pointed, but not drawn out in the shape of a lobe. The picture of the wings consists, as to color, of brownish-black and brownish-yellow and some hyaline cells of a peculiar shape. The root of the wings is yellow, as far as the origin of the third longitudinal vein; the extreme basis, however, is strongly infuscated; there is a rather dark-brown crossband in the region of the humeral crossvein, and the basis of the marginal cell is hyaline. A dark-brown color follows next, the first portion of which forms a curved crossband, reaching backwards as far as the posterior basal crossvein; anteriorly it is prolonged in the marginal cell, as far as the end of the first longitudinal vein, where it stops short abruptly. After some interruption, the brownish-black color forms a broad border of the anterior margin, beginning somewhat above the posterior crossvein, which does not only occupy the whole breadth of the marginal cell, but also encroaches on the submarginal cell, follows the apex of the wing at the fourth longitudinal vein as far as the small crossvein and also covers the latter; posteriorly, it not only runs along the posterior crossvein and extends over the end of the discal cell, but follows also some distance along the end of the fifth longitudinal vein, upon its posterior side; the third longitudinal vein is bordered with brownish-black upon its whole length. The portions of the marginal, submarginal and discal cells, free from the brownish-black color, are tinged with brownish-yellow; the first basal cell, as well as the first two posterior cells, are hyaline. The alula, as well as the anal angle of the wing and the adjoining portion of the third posterior cell, is grayish-hyaline, with a tinge of yellow; the posterior side of the fifth longitudinal vein has a brownish-yellow border, the middle of the third posterior cell is rather pure hyaline, only more grayish towards the posterior margin of the wing.

Hab. Virginia, Pennsylvania (Osten-Sacken).

*Fourth Section: ORTALINA.*Gen. I. **AUTOMOLA** gen. nov.

Charact.—*Front* broad, very much narrowed anteriorly. Eyes rather large, slightly protruding, irregularly rounded. Face in profile somewhat concave, obtusely carinate between the distinct antennal foveæ. The anterior edge of the mouth very much drawn upwards, so that the rather strongly developed clypeus projects considerably beyond it. Cheeks broad.

Antennæ reaching beyond the middle of the face; the first two joints short; the narrow third joint more than twice as long as the first two taken together, rounded at the end; antennal arista thin, slightly stronger at the basis only, with a very short pubescence.

Thoracic dorsum not bristly on its middle, before the region of the transverse suture. The tibiæ, before the end of their upper side, with a præapical bristle.

The first longitudinal vein bristly before its end; the auxiliary vein very much approximated to it; the costa more or less incrassated beyond the end of the first longitudinal vein; the third and fourth longitudinal veins parallel towards their end; the crossveins not approximated; the second basal cell and the anal cell comparatively rather small, the latter rounded at the end; the sixth longitudinal vein complete, but remarkably short, and hence, the anal angle of the wing very small; alula comparatively large.

The genus *Automola* contains unmetallic species. The picture of their wings generally consists in black spots upon the root of the wings and three black crossbands, the first of which is only at a short distance from the basis, while the second runs over the posterior crossvein and the third lies between the second and the apex of the wing; these bands being more or less incomplete, or the second and third expanding or even coalescing into one large spot, give rise to different modifications of the design of the wings.

Ortalis atomaria Wied. and *trifasciata* Wied. from Brazil, may be considered as the types of the genus. North American species have not been discovered yet.

I have already had occasion to mention in the Introduction that *Automola*, on account of the præapical bristles on the tibiæ, which distinguish it from the other genera, is not very well placed in the family of *Ortalidæ*.

Gen. II. **TETANOPS** FALL.

Charact.—*Front* of a considerable and uniform breadth. *Eyes* rounded-oval, or oval. *Face* strongly projecting in profile, more or less retreating below. *Clypeus* small, but projecting beyond the edge of the mouth. *Oral opening* comparatively small; *proboscis* but little incrassated.

The hairs and bristles on the body remarkably short, especially the bristles of the prothorax much smaller than in any other genus among the *Ortalina*; *thoracic dorsum* upon its middle only posteriorly with a few bristles.

Antennæ short, sometimes strikingly short; third joint oval, longer than the second.

First longitudinal vein bristly towards its end only; the *crossveins* rather distant; the second and third longitudinal veins parallel towards their end, or only gently convergent; posterior angle of the anal cell pointed, but not prolonged in the shape of a lobe.

The North American species of *Tetanops* are distinguished from the European ones by the more distinct and sharper anterior edge of the mouth, while in the latter the anterior end of the oral opening hardly shows a distinct margin. As one of the American species, known to me, has, moreover, the vertical diameter of the eyes considerably larger than the European species, I was for some time in doubt, whether it would not be better to separate generically the North American from the European species. Nevertheless, they possess enough characters in common, to render such a separation, at least for the present, unnecessary. Besides the stout head, with the very broad front, the striking bareness of the whole body and the great shortness of the prothoracic bristle, the absence of any picture on the wings, except some very faint spots along the costa, easily distinguishes the species of *Tetanops*.

1. *T. luridipennis* n. sp. ♂ ♀.—(Tab. VIII, f. 17.) Frons præter vittam mediam punctata; alæ sordide lutescentes, ad costam obsolete-sime lurido-maculatæ.

Front, with the exception of a median stripe, punctate; *wings* of a dingy clay-yellow, with very indistinct brownish-clay-yellow spots along the costa. *Long. corp.* ♂, 0.21; ♀ cum terebrâ 0.28—0.32; *long. al.* 0.18.

Head reddish-yellow. The very broad front more red; it has a small median stripe, which is not pollinose, and has, on each side, a brown border; the latter sometimes becomes indistinct

above, and, on the anterior part of the front, is somewhat turned sideways, generally also more expanded and darker. The sides of the front, each of which is nearly double the breadth of the median stripe, are covered with white pollen, rendered cribose by a dense punctation of pollenless dots, so that of the pollinose surface, nothing but a network is left. The face, in profile, projects very much in front of the eyes, and retreats very considerably below; its intermediate portion is, as in all the species of *Tetanops*, comparatively narrow. The antennal foveae are deep and sharply defined, shining-black, except on their upper portion. Eyes rounded-oval; cheeks very broad. The upper part of the occiput is clothed with white pollen; in the vicinity of the orbits and of the edge of the vertex this pollen is likewise interrupted by punctiform pollenless dots. Antennae yellowish-red, the third joint, with the exception of the basal third, more or less infuscated. Although the ground color of the thorax is shining-black or brownish-black, it is, with the exception of the humeri, concealed by a thick grayish-white pollen, sometimes yellowish on the thoracic dorsum; numerous punctiform, pollenless dots interrupt this pollen and give it a cribose appearance; the region of the prothoracic spiracle alone is free from these dots. The pollen covering the scutellum is similar in coloring to that of the thorax, but it is, to a considerable extent, much less thick upon its sides. The abdomen has the same color and the same pollinose surface, interrupted by punctiform, pollenless dots, as the thorax, but the pollen is a little less thick and the punctiform dots a little larger, so that, here and there, they coalesce and the ground color becomes more apparent. The first segment of the flattened ovipositor is shining black, very broad, rather strongly attenuated, however, towards its end. Femora blackish-brown, the tip of the front ones yellowish-red to a small, the tip of the hindmost ones to a greater extent. Front tibiae blackish-brown, with a yellowish-red basis; middle tibiae usually entirely yellowish-red or but little infuscated towards their end; hind tibiae blackish brown, with a yellowish-red basis and generally also the extreme tip of the same color. Tarsi yellowish-red at the basis, the front ones from about the tip of the first joint, the posterior ones from about the tip of the second or third joint, blackish-brown. Wings of a dingy clay-yellow, almost brownish in fully colored specimens, without any distinct picture; however,

indistinct traces of three somewhat darker clouds are apparent; the first in the marginal cell, above the origin of the submarginal cell, the second at the end of the stigma, and the third, which sometimes is wanting, fills up the end of the marginal cell; all three are so little apparent that they can easily be overlooked.

Hab. Nebraska (Dr. Hayden).

2. *T. integra* n. sp. ♀.—(Tab. VIII, f. 18.) *Frons tota punctata; alæ cineræ, immaculatæ.*

The whole front is punctate; wings gray, without any picture. Long. corp. cum terebrâ 0.28—0.31; long. al. 0.17.

Head brownish-black, rather dusky brownish-red upon the greater part of the front, the cheeks, and near the anterior edge of the mouth. The front has no median stripe, and is altogether covered with grayish-white pollen, rendered cribrose by numerous small and very dense pollenless dots; a fine network, covering the whole front, is all that remains pollinose. The pollen extends, from the front over the very broad lateral portions of the face, as far as the cheeks; the pollenless dots, however, do not reach beyond the middle of the face. The face in profile is less projecting in front of the eyes, and less retreating below, than in *T. luridipennis*. The antennal forcæ, on their outside slope, are covered, to a considerable extent, by a white pollen; at the bottom they are shining black. The flattened ridge of the carina, separating them, has also a whitish pollen. The vertical diameter of the eyes is larger than in the preceding species or in any of the species of *Tetanops* to me known. The cheeks are very broad, although somewhat narrower than in *T. luridipennis*. The upper half of the occiput is clothed with a whitish pollen, extending upon the hind side of the cheeks as far as the edge of the mouth; in the vicinity of the posterior orbit and of the edge of the mouth, this pollen is interrupted by pollenless punctiform dots. Antennæ brownish-red, the third joint for the most part blackish-brown. The ground color of the thorax is glossy, almost shining-black, but altogether covered by a whitish-gray or more yellowish-gray pollen, interrupted by countless dots, which are, however, much smaller and less sharply defined than in the preceding species. Quite in front, the thoracic dorsum shows an indistinct beginning of a median stripe, in the shape of two dark longitudinal lines, which are rather distant from each other.

Upon the pleuræ the pollen is perceptibly less dense than upon the thoracic dorsum, so that they appear shining. Upon the sides of the scutellum the pollen is thick and not interrupted, while that upon its disk somewhat resembles the pollen on the surface of the thoracic dorsum, only it is a little thinner and has no distinct pollenless dots. The abdomen is shining black, covered, towards the basis, with a gradually increasing, uninterrupted, but not very thick ash-gray pollen. The first joint of the flattened ovipositor is shining black, very broad, but little narrowed towards its end, with somewhat convex sides and comparatively shorter than that of *T. luridipennis*. Feet black or brownish-black; the extreme tip of the femora, the basis and extreme tip of the tibiæ, as well as the tarsi, yellowish-red; however, the last three or four joints of the fore tarsi and the last two joints of the hind tarsi, brownish-black. Wings rather hyaline, gray, with a delicate tinge of brownish-chaly-yellow, without any picture.

Hab. Illinois (Osten-Sacken).

Gen. III. **TEPHRONOTA** Loew.

Charact.—*Head* high and short. Front of a moderate and equal breadth, comparatively long. Face rather sharply carinate, only little protruding in front of the eyes in profile; almost vertical. The vertical diameter of the eyes almost double the size of the horizontal one. Anterior edge of the mouth not drawn upwards. Cheeks very narrow.

Antennæ of a medium length; the first two joints short; the third ending at a sharp angle, although not excised above.

Thorax upon its middle with bristles on the hind part only; covered with a gray dust.

The first longitudinal vein with bristles upon its end only; the end of the fourth longitudinal vein not curved forward; the posterior angle of the anal cell, although sharp, is not extended in the shape of a lobe.

This genus contains only small-sized species, which, in their whole organization, approach the species of *Pteropaectria*; this is still more the case with the European species, than with the only American one which I know. The latter, however, agrees in so many characters with the European *Tephronotæ*, that it can be placed, without any hesitation, in that genus. Its antennæ are a little shorter and their third joint somewhat broader; the

pollen on the body is thicker and more extended than in the European species; the crossbands of its wings are incomplete.

I. T. humilis Low. ♂ ♀.—(Tab. VIII, f. 24.) Nigricans, cinereo-pollinosa, capite flavo, pedibus luteis, alæ albido-hyalinæ, fasciis tribus nigris, intermediâ integrâ, reliquis postice abbreviatis.

Rather black, covered with gray pollen; with a yellow head, and rather clay-yellow feet; wings whitish-hyaline, with three black crossbands, the median of which is entire; the two others are abbreviated. Long. corp. ♂ 0.12—0.14; cum terebrâ 0.11—0.16; long. al. 0.1—0.13.

Syn. *Herina ruficeps* v. d. WULF, Tijdschrift voor Entomol., Jaarg. IX, p. 156.

Head yellow. Front brighter yellow, almost orange-red upon its anterior end; on each side with a conspicuous border of white pollen, which, becoming broader, extends below over the face as far as the cheeks. The occiput becomes blackish above, but is rather evenly covered with a rather thick whitish pollen. The ground color of the thorax is rather black, more brownish on the humeri and upon the lateral border, as well as below the root of the wings; this color, in well-preserved specimens, is covered by a grayish-white pollen; upon the thoracic dorsum there are two longitudinal stripes, of a somewhat darker color, very little apparent and abbreviated posteriorly. The color of the scutellum, which is likewise covered with gray pollen, verges more on dingy brownish, and on clay-yellow along the edges; in less fully colored specimens the whole scutellum is clay-yellow. The color of the abdomen is likewise rather black, sometimes only brown at the basis. In the male, this color appears distinctly as black or brownish-black upon the last segment and on the hypopygium, both of which are pollenless, while on the preceding segments this color is concealed under a rather thick pollen, which on the anterior portion of the segment has a light whitish-gray, on the posterior half a brown coloring. The female has the last abdominal segment likewise pollinose, the pollen being generally light white-grayish, or verging on brownish about the middle of the abdomen only; the pollen on the preceding segments is the same as in the male. The first segment of the altogether flattened ovipositor is not very long, but very broad and very broadly truncate at its end; its pollen is very little perceptible, so that it is glossy-black, more brownish-black in

immature specimens. Feet of a dirty clay-yellow, femora in the middle and tarsi towards the tip, somewhat infuscated. Halteres whitish-yellow. Wings whitish-hyaline, with three broad, perpendicular, more grayish-black than black crossbands. The first of these bands covers, near the anterior margin, the latter half of the costal cell, and reaches, without becoming more narrow, the fourth or fifth longitudinal vein; in the first case it becomes perceptibly wider between the third and fourth, in the second case between the fourth and fifth longitudinal veins. The second band covers, near the anterior margin, the apical half of the stigma and reaches there, in most specimens, even a little beyond the end of the first longitudinal vein; without attenuating, it runs over the small crossveins as far as the fourth longitudinal vein, forms a very broad border along the section of the fourth vein lying between the two crossveins, and runs, afterwards, along the posterior crossvein towards the fifth longitudinal vein; its breadth is not the same in all specimens; when narrower, this crossband shows a distinct knee-shaped bend, depending upon its passage from the small to the posterior crossvein (this is the case with the specimen figured by Mr. v. d. Wulp); when broader, this crossband extends, in the shape of a blackish-gray shadow, as far as the third posterior cell, so that of the knee-shaped bend only a trace is left, which is due to a dilated spot upon the inner side of the crossband, near the posterior margin of the discal cell (as represented in my figure). The third band covers, on the anterior margin, the end of the marginal cell to a considerable extent, becomes gradually more narrow posteriorly and reaches more or less completely the fourth longitudinal vein, where it suddenly is interrupted. The root of the wing is tinged with blackish-gray as far as a little beyond the humeral crossvein. The second and third longitudinal veins are strongly divergent towards their end; the last section of the fourth longitudinal vein slightly converges towards the third vein and is not quite so straight as usual in the species of *Tephronota*, but, at the same time, not so much curved forward by far as in the case of the species of *Anacampta*, *Holodasia*, and *Apospasmica*. The crossveins are very much approximated, as the distance between them is not much larger than the length of the small crossvein, but smaller than the posterior crossvein. The posterior angle of the anal cell is short and sharp, and not prolonged in the shape of a lobe. The sixth

longitudinal vein is weak and indistinct soon after its middle, so that it appears interrupted a long distance before the margin of the wing.

Hab. New York (Osten-Sacken); Virginia; Texas (Belfrage).

Observation.—The description of *Herina ruficeps* by v. d. Wulp, contains only one datum which might render its identification with *T. humilis* doubtful. He says that the third antennal joint is four times as long as the second, while in all my specimens it hardly reaches three times its length. As, in other respects, the agreement of the very good description is perfect, I have not the slightest doubt that this difference arises from a different mode of viewing or measuring the antennæ. Unfortunately, the name given by Mr. v. d. Wulp cannot be preserved, as it has been preoccupied by Fabricius.

Gen. IV. CEROXYS MACQ.

Charact.—*Head* rather rounded. Front very broad, somewhat narrowed above, without stripe. The perpendicular diameter of the eyes is much larger than the horizontal one. Cheeks of medium breadth.

Third antennal joint upon its upper side distinctly excised, very much pointed at the tip. Arista distinctly pubescent.

Thorax, upon its middle, with bristles as far as its anterior portion.

First longitudinal vein with bristles upon its end only; the fourth longitudinal vein not curved forward. The posterior angle of the anal cell acute, but not prolonged in the shape of a lobe.

The genus *Ceroxys* contains species which are very much alike; the thorax and abdomen are thickly covered with yellowish or grayish dust; the head is yellow. The picture of the wings, consisting of comparatively large blackish-brown or black spots, is the same in all the species; it consists of seven spots, the first of which lies on the basis of the submarginal cell, the second upon the end of the stigmatal (third costal) cell; the third covers the small and the fourth the posterior crossvein; the last three spots lie on the ends of the second, third, and fourth longitudinal veins; the last two generally coalesce completely, while the one placed at the end of the second vein is generally less completely united with them.

The species are easily distinguished by the shape and color of the third antennal joint, by the presence or absence of a dark crossband on the posterior margin of the abdominal segments, by

the greater or smaller extent of the spots on the wings, especially by the relative position of the stigmatal spot to the one covering the small crossvein, and by the separation or coalescence of both.

1. *C. obscuricornis* n. sp. ♂ ♀.—(Tab. VIII, f. 20.) Polline ex cinereo lutescente vestitus, tertio antennarum articulo fusco-nigro, pedibus luteis, alarum maculâ stigmatali et limbo venæ transversalis mediæ fasciolam arcuatam efficientibus.

Covered with a grayish-clay-yellow pollen; third antennal joint brownish-black, feet clay-yellow; the spot at the end of the stigmatal cell and the one covering the small crossvein form a curved crossband. Long. corp. ♂ 0.21; ♀ cum terebrâ 0.25; long. al. 0.2—0.21.

The first two antennal joints brownish-ferruginous-yellow, or brownish-yellow; third joint brownish-black, of medium breadth; arista black. Scutellum upon its edge only indistinctly yellowish-brown. Abdomen without any trace of dark crossbands, except that the pollen, towards the posterior portion of the segments, becomes more brownish-gray in a hardly perceptible degree. The first joint of the flattened ovipositor is only moderately long, very broad; its truncate very broad also; the coloring and the pollen are the same as those of the abdomen. Feet clay-yellow; tarsi, with the exception of the basis, more or less strongly infuscated; the only male in my possession has the front femora very much infuscated upon the greater part of the posterior side; it is not probable, however, that this is a constant sexual character. The first spot on the wings extends from the first to a little beyond the fourth vein; the spot lying upon the end of the stigmatal cell is more or less completely coalescent with the one covering the small crossvein, and forms with it a rather oblique, distinctly arcuated crossband; the other spots have nothing peculiar about them.

Hab. Nebraska (Dr. Hayden).

2. *C. ochricornis* n. sp. ♀.—(Tab. VIII, f. 21.) Polline ex cinereo lutescente vestitus, segmentis abdominalibus postice anguste fuscolimbatis, anteunis ex-ferrugineo ochraceis, pedibus luteis, alarum maculâ stigmatali et limbo venæ transversalis mediæ in fasciolam rectam conjunctis.

Covered with a grayish-clay-colored pollen; the segments of the abdomen with narrow brown borders posteriorly; antennæ ochre-brownish, the

feet clay-yellow; the spot upon the end of the stigmatal cell and the one covering the small crossvein, in coalescing, form a straight crossband. Long. corp. cum terebrâ 0.25; long. al. 0.21.

Antennæ altogether ochre-brownish; third joint distinctly broader than in *C. obscuricornis*; arista brownish-black. Scutellum generally yellowish, with the exception of its middle. Abdominal segments, with the exception of the last one, with very narrow, but very sharply limited and conspicuous brown posterior margins. The first segment of the very flattened ovipositor is only moderately long, very broad, and very broadly truncate at the end; its coloring and its pollen are similar to those of the abdomen. Feet clay-yellow; tarsi strongly infuscated, generally paler towards the basis. The first spot upon the wings reaches from the first to the fourth longitudinal vein; the spot upon the end of the stigmatal cell is more or less completely connected with the spot covering the small crossvein, forming a straight, almost perpendicular half-crossband; the other spots have the ordinary appearance.

Hab. Northern Wisconsin River (Kennicott).

3. *C. similis* n. sp. ♀.—(Tab. VIII, f. 23.) Polline Intescente vestitus, segmentis abdominalibus postice nigro-limbatis, alarum maculâ subbasali in fasciam dilatata, maculâ stigmatali et limbo venæ transversalis mediæ in fasciolam conjunctis.

Covered with clay-yellow pollen; the abdominal segments margined with black posteriorly; the spot near the basis of the wing is extended in the shape of a crossband; the one at the end of the stigmatal cell forms a half-crossband with the spot covering the small crossvein. Long. corp. ♂ 0.22; ♀ cum terebrâ 0.27—0.28, long. al. 0.21—0.22.

First two antennal joints yellow; the third joint is unfortunately lost in all the three specimens which I have before me, but is probably of the same color. Scutellum yellow, or grayish upon its middle only. The segments of the abdomen have all, without exception, a brownish-black, narrow, well-defined border, upon their posterior side. The first segment of the flattened ovipositor is only moderately long, very broad, very broadly truncate at the end; its coloring and the pollen upon it, are of the same color as on the abdomen. Feet clay-yellow; tarsi, especially towards their tip, rather strongly infuscated. The first spot on the wings expands into a crossband, reaching anteriorly as far as the costa,

posteriorly it extends, although somewhat paler, along the sixth longitudinal vein, which it finally crosses, as far as the posterior margin of the wing, on the fifth longitudinal vein it forms an obtuse angle, at which place, on the sides of the fifth longitudinal vein, it is very faint, sometimes almost interrupted; the spot at the end of the stigmatal cell coalesces with the one covering the small crossvein, forming a steep, somewhat enrvd half-cross-band; the spot covering the posterior crossvein is rather large; the three other spots are of the usual shape.

Hab. Connecticut (Osten-Sacken).

Observation.—The name which I give to this species is intended to call to mind its extraordinary resemblance to *C. crassipennis* Fab., occurring in Europe. This resemblance is so great, that I would doubt the specific distinctness of the two species, if the femora of the American one were not altogether yellow, while those of *C. crassipennis* are blackish-brown from the basis as far as the middle. In order to overlook this difference and to maintain the identity of the two species, the proof of a perfect agreement in all, even the minutest, plastical characters would be required. The three specimens of *C. similis* in my possession are not well preserved enough to enable me to undertake such a comparison.

4. *C. canus* Loew. ♂ ♀.—(Tab. VIII, f. 22.) Polline ex lutescente cinereo vel albido-cinereo vestitus, tertio antennarum articulo pedibusque fuscis, alarum maculâ stigmatali et limbo venæ transversalis mediæ separatis.

Covered with a yellowish-gray or grayish-white pollen; third joint of the antennæ and the feet brown; the spot on the stigmatal cell entirely separated from the one which covers the small crossvein. Long. corp. ♂ 0.16; ♀ cum terebrâ 0.23; long. al. 0.16—0.18.

SYN. Ortalis cana Loew, Berl. Entom. Zeitschr. II, p. 374.

Smaller than the preceding species, with a grayish or whitish-gray pollen, verging less on yellow. The first two antennal joints brownish-yellow or yellowish-brown; the third joint of medium breadth and rather blackish-brown. Antennal arista black. The scutellum at most indistinctly yellowish-brown along the edges only. Abdominal segments without any trace of darker borders. The first joint of the flattened ovipositor distinctly longer than in the three previous species and somewhat less broadly truncate at

the end; its coloring and the pollen upon it are the same as those on the abdomen. Coxæ and feet blackish-brown; the second coxal joint, the tip of the femur, the basis of the tibiæ and the extreme tip of the middle tibiæ are yellowish-red. In the European specimens this yellowish-red coloring has often a much greater extent and also occurs at the basis of the tarsi; it is probable that the same is the case with some American specimens. The surface of the wings is much more whitish than in the other species; the first spot is small, although it reaches from the first to the fourth vein; the spot at the end of the stigmatal cell is also comparatively small, does not quite reach the second longitudinal vein, and remains quite separated from the spot covering the small crossvein; the spot covering the posterior crossvein is of a moderate breadth; the spots upon the ends of the longitudinal veins are of the ordinary size.

Hab. Yukon River, Alaska (Kennicott); Nebraska (Dr. Hayden).

Observation.—Of this species I possess only a male from Nebraska and a female from Hudson's Bay Territory. The most careful comparison with specimens of *Ceroxys canus* from the southern part of middle Europe and from southern Europe has not revealed any character indicative of a specific distinctness of the European and the American specimens.

Gen. V. ANACAMPTA Loew.

Charact.—*Head* hemispherical, rather than round; front broad, somewhat narrower above; the vertical diameter of the eye much larger than the horizontal one; cheeks broad.

Third antennal joint distinctly cut out upon its upper side; pointed at the end.

Thorax upon its middle provided with bristles near the posterior margin only.

First longitudinal vein with bristles upon its end only; the end of the fourth longitudinal vein curved forward in a striking manner; posterior angle of the anal cell sharp, but not prolonged in the shape of a lobe.

The genus *Anacampta* contains species of large size, which resemble *Ceroxys* in their general appearance, as well as in the picture of the wings. They differ, however, sufficiently in the black color of the body, in the thoracic dorsum not being provided with bristles as far as its anterior part and in the conspicuous

curvature of the end of the fourth longitudinal vein. The black coloring of the body they have in common with the species of *Holodasia* and *Apospasmica*, which they approach in the whole structure of their body. They differ from *Holodasia* in the fact that the first longitudinal vein is not provided with bristles upon its whole course, but at its end only. From *Apospasmica* they differ in the shape of the anal cell, the posterior angle being only acute here, while in *Apospasmica* it is drawn out in a long lobe; moreover, in the latter genus, the end of the fourth longitudinal vein is not curved forward; in *Anacampta* the picture of the wings consists rather of spots, or bands consisting of spots, while in *Apospasmica* there are complete crossbands. The structure of the third antennal joint of *Anacampta* affords a character for the distinction of it from all the other genera of *Ortalina*, which renders any further developments superfluous.

1. *A. latiuscula* n. sp. ♂ ♀.—(Tab. VIII, f. 19.) Nigra, thorace abdominisque fasciis duabus cinereo-pollinosi, capite ex rufo luteo, pedibus rufis, alis nigro-maculatis.

Black, thorax and two crossbands on the abdomen covered with gray pollen; head reddish-yellow; feet red; wings spotted with black. Long. corp. ♂ 0.31, ♀ cum terebrâ 0.33—0.34; long. al. 0.26.

One of the largest species of the genus, and broader in shape than most of them. Head reddish-yellow, opaque, covered with a very thin, and hence not easily perceptible greenish-white pollen; occiput more thickly pollinose with white. Front broad, somewhat narrower above; the not very distinct frontal stripe very much narrowed above, of a purer yellowish color and almost pollenless; the comparatively thick pubescence of the broad lateral portions of the front is inserted in very small, but distinct brownish dots. Antennæ ochreous-brown, the color of the first two joints more yellowish, that of the third joint more brownish. Ground color of the thorax, with the exception of the brick-red humeral callosities, black, but altogether covered with an ashy-gray pollen, which is not quite so thick on the pleuræ as on the thoracic dorsum. The hairs and bristles of the thoracic dorsum are inserted on small, but distinct black dots. Sentellum black, with a broad brick-red border, pollinose with ashy-gray. Abdomen shining-black, with black hairs and two broad crossbands of whitish-gray pollen, situate on the anterior portion of the

second and third segments; they gradually become indistinct on the sides and finally disappear near the lateral margin. The fifth segment of the female abdomen is very much shortened. The first joint of the ovipositor is shining-black, with black hairs, about as long as the penultimate segment of the abdomen, not very broad, and, towards its end, rather narrowed. Feet brick-red; tarsi infuscated towards their end, the front ones much more than the four posterior ones; the front tibiæ also show sometimes a browner coloring. Wings grayish-hyaline, quite gray towards the posterior border, not very transparent; more yellow towards the basis, especially in the costal cell; stigma ochre-yellow, with a somewhat infuscated end. The picture of the wings is brownish-black; it comprises: 1, a spot upon the humeral crossvein, reaching as far as the fourth longitudinal vein; 2, a perpendicular crossband, covering the end of the costal cell near the anterior border, and reaching posteriorly as far as the sixth longitudinal vein; between the fifth and the sixth longitudinal veins it is much paler and disappears gradually in the gray coloring of the surface of the wing; 3, a perpendicular half-crossband, beginning near the anterior margin, immediately beyond the end of the first longitudinal vein, running over the small crossvein and reaching a little beyond its posterior end; 4, a spot, broadly covering the posterior crossvein in the shape of a half-crossband; 5, a spot occupying the end of the marginal cell and, with the end nearer to the root of the wing, reaching into the submarginal almost in the shape of a hook, without touching the third vein; 6, a spot near the apex of the wing, the limit of which runs almost perpendicularly from the end of the second longitudinal vein to the fourth longitudinal, beyond which it occupies only a small space at the extreme end of the second posterior cell.

Hab. California (Alex. Agassiz).

Gen. VI. **AOSPASMICA** nov. gen.

Charact.—Front of equal breadth. Face rather strongly carinate; rather perpendicular and straight in profile; the vertical diameter of the eyes very much larger than the horizontal one.

Third antennal joint, on its upper side, gently but distinctly excised, very pointed at the end; arista very bare.

Thorax along the middle with bristles on its hind part only.

First longitudinal vein with bristles towards its end only; the end of the fourth longitudinal vein not curved forward; the posterior angle of the anal cell drawn out in a narrow, exceedingly long lobe.

Robust, black species, of the same general appearance as *Holodasia* and *Anacampta*; the structure of the head more like that of *Pteropaectria*; the thorax generally shows longitudinal lines of a paler-colored dust, answering to the intervals of the ordinary thoracic stripes. The wings have complete crossbands.

The typical species is the *Ortalis fasciata* of Wiedemann, from Chile, which is identical with the *Tephritis quinquefasciata* Macq. Dipt. Exot. Suppl. IV, 291.

The shape of the anal cell reminds very much of *Diacrita*; nevertheless, there are no other points of relationship between the two genera.

Observation.—Should an American species be found which does not well fit in any of the above-described six genera, the characters of the European genera should be compared; they have been given in the part treating of the systematic distribution of the *Ortalidæ* in general.

Fifth Section: PTEROCALLINA.

Gen. I. **PTEROCALLA** Rond.

Charact.—*General appearance:* Trypeta-like.

Wings very narrow, in comparison to their length, of a rather striking shape on account of their equal breadth, very broadly rounded at the root and at the tip; auxiliary vein much shorter than the first longitudinal vein, so that the distance between the ends of both is strikingly large; first basal and discal cells very long; posterior crossvein very oblique, its anterior end being much nearer the apex of the wing than its posterior end; the posterior angle of the anal cell drawn out in a moderately long lobe.

The peculiarities in the outline of the wings and in the venation of the species belonging to this genus are so striking, that no doubt can possibly arise about the location of any of them. In some other respects, these species differ considerably from each other, so that, should their number increase, it would be necessary to break up the genus *Pterocalla* into smaller genera. The name *Pterocalla* would, in this case, remain to the genus which contains *P. ocellata* Fab., as Mr. Rondani established the genus for this species.

1. *P. strigula* n. sp. ♂.—(Tab. VIII, f. 30.) Albido-pollinosa, punctis maculisque deformibus fusco-nigris aspersa; alæ fusco-nigræ, disco dilutius fusco, punctis maculisque fusco-nigris variegato, marginibus antico macularum hyalinarum serie, postico limbo latiusculo hyalino ornatis, venis longitudinalibus non undulatis.

Clothed with white pollen, marked with brownish-black dots and irregular spots; wings brownish-black, of a paler brown upon their middle, and with brownish-black spots and dots; the anterior margin with a row of hyaline spots and the posterior margin with a rather broad hyaline border; longitudinal veins not undulated. Long. corp. 0.12—0.13. Long. al. 0.17—0.18.

In the structure of the head and of its parts, the coloring and picture of the whole body, this species resembles *Myennis vau* very much, but it differs considerably in the narrow wings with almost parallel sides, with a different venation and a different picture. The ground color of the body is an opaque brownish-black, for the most part covered with a thick white dust; the latter's surface on the upper side is broken through by brownish-black dots and a number of rather regularly arranged, but very irregularly shaped, brownish-black spots; the face does not show any such broken through places; the upper, larger half of the pleuræ shows numerous brownish-black dots, which almost coalesce above into a stripe; a little below the middle of the pleuræ there is a brownish-black longitudinal stripe and immediately below it a narrower stripe, formed by a white pollen; the pectus is brownish-black. Femora and tibiæ brownish-black (the intermediate femora in the described specimen are paler perhaps in consequence of immaturity); all the femora have, upon their last third, a more or less complete ring of white pollen; their extreme tip, as well as the basis of the tibiæ, are tinged with yellowish-white; each tibia shows, upon its middle, a very conspicuous white ring and a very sharply limited white tip. The yellowish-white feet are somewhat infuscated towards the end. Wings strikingly long and narrow, of an unusually equal breadth; very obtuse at the end, like in other species of *Pterocalla*; the auxiliary vein is remarkably short, so that the distance between its end and the end of the first longitudinal vein is remarkably large; the second longitudinal vein is rather long; the third ends not far from the apex of the wing, and has, like the others, a very straight and not at all undulated course; the ends of the third and fourth veins hardly show a vestige of convergency; the crossveins are rather

closely approximated; the posterior crossvein, with its anterior end, is nearer to the apex than with its posterior end; the posterior angle of the anal cell is drawn out in a very long and pointed lobe (the figure makes it too short and heavy). The extended and entirely uninterrupted picture of the wings leaves near the anterior margin an irregular row of hyaline spots and on the posterior margin a broader hyaline border, with an irregularly undulated outline; the coloring of the picture is brownish-black; its inner part is paler brown, with numerous brownish-black dots and spots.

Hab. Georgia (Berlin Museum).

Gen. II. **STICTOCEPHALA** nov. gen.

Charact.—*General appearance:* Trypeta-like.

Front very broad, with punctures; cheeks comparatively broad; clypeus somewhat projecting over the edge of the mouth.

Wings of the usual shape; the ends of the auxiliary and of the first longitudinal veins are far distant from each other; posterior crossvein steep; posterior angle of the anal cell acute; the third and fourth longitudinal veins, towards their end, at least with a trace of a convergency.

All the species belonging here are opaque in their coloring; thorax and abdomen are punctate in all of them; moreover, they are generally marked with other pictures.

The species which I know of may be separated in two groups, on account of the different size of the hairs on the front. *Stictocephala cribrum* and *cribellum*, would belong to the first group, *S. corticalis* and *vau* to the second. In the two latter species, the two uppermost of the short hairs, inserted on the lateral border of the front, assume the appearance of bristles, so that in this respect these species are like the *Trypetina*, while this is not the case with the two preceding species.

1. S. cribellum n. sp. ♂ ♀.—(Tab. VIII, f. 26.) Cinerea, frontis parte antica, antennis, facie, genis, proboscide, palpis pedibusque luteis; alæ hyalinæ, fasciis quatuor, præter secundam, posticæ abbreviatis, maculâ apicali et venæ transversalis posterioris limbo fuscis.

Gray; the anterior part of the front, antennæ, face, cheeks, proboscis, palpi, and feet clay-yellow. Wings hyaline, with four bands, which are abbreviated posteriorly, except the second; a spot at the apex and a border along the posterior crossvein, brown. Long. corp. 0.13—0.15; long. al. 0.14—0.15.

Light gray, front somewhat yellowish towards its anterior margin, covered with rather coarse punctures; the uppermost hairs near the lateral margin of the front are not longer and stronger than usual. Antennæ clay-yellow, third joint rounded-ovate, sometimes rather brownish-yellow. Ground color of face and cheeks clay-yellowish, covered with a whitish pollen. Proboscis and palpi clay-yellowish. Thoracic dorsum with somewhat scattered blackish-brown dots, which sometimes coalesce in lines upon its posterior portion; moreover with four brownish-black spots in a row corresponding to the transverse suture. Scutellum with four bristles, turgid, pale-gray, with two conspicuously large shining-black spots at the end. Metathorax black, pruinose with whitish-gray. Pleuræ dotted with brownish-black above. Abdomen with similar dots, usually with a more clay-yellow ground color at the basis; this color is sometimes more extended and gives the abdomen a more yellowish-gray tinge, while the thorax is whitish-gray. Coxæ and feet clay-yellow; posterior coxæ at the basis and the tarsi towards their tip, somewhat infuscated. Wings hyaline with four perpendicular, not very dark, brown bands, a broad brown border on the posterior crossvein and a brown apex; the first band begins near the anterior margin immediately beyond the humeral crossvein, and is not distinctly perceptible beyond the sixth longitudinal vein; the anal cell is just filled out by it; the second band begins at the anterior margin quite near the end of the auxiliary vein, and ends upon the end of the sixth longitudinal vein; the third band begins immediately before the end of the first longitudinal vein and runs across the small crossvein, at the end of which it is interrupted; the fourth band generally reaches from the anterior margin not quite as far as the third longitudinal vein, or is continued a little beyond it in the shape of a faint shadow.

Hab. Nebraska (Dr. Hayden).

2. *S. cribrum* n. sp. ♀.—(Tab. VIII, f. 25.) Præcedenti simillima, sed major, alarum picturâ simili, sed saturatiore, fasciâ tertiâ et venâ transversalis posterioris limbo in fasciam integram confluentibus, tibiarum omnium apice, posticarumque annulo medio, apice denique tarsorum nigris.

Very like the preceding, but larger; the same picture of the wings, but darker; the third band and the infuscation along the posterior crossvein

coalesce into an incomplete crossband. The tip of all the tibiæ, a ring on the middle of the hind ones and the tip of all the tarsi, black. Long. corp. 0.21; long. al. 0.20.

Unfortunately, I possess only a single, badly preserved specimen of this insect. The resemblance to the preceding species is so great, that only the observation of the living insect or the comparison of a large number of specimens, will enable one ultimately to decide about their specific diversity. The considerably larger size, the darker coloring of the picture of the wings, the confluence of the third crossband of the wings with the infuscation on the posterior crossvein into a complete band, the difference in the coloring of the feet (in *S. cribellum* the tibiæ show only a weak trace of a darker coloring at the tips, and there is no trace whatever of a ring on the hind tibiæ, the tarsi are but slightly infuscated towards the end)—all these differences render a specific distinctness probable, although, on the other hand, the great resemblance of all the other characters tends to diminish this probability.

Hab. Middle States (Osten-Sacken).

Observation.—In case the specific identity of *S. cribellum* and *cribrum* is proved, the latter name should be retained for the species, as representing the more fully colored, and hence, normal specimens, while *S. cribellum* would then be regarded as a smaller and paler variety.

3. *S. corticalis* Fitch in litt. ♂ ♀.—(Tab. VIII, f. 28.) Fusco-nigra, polline albo-cinereo aspersa, punctis, maculisque fusco-nigris variegata; alæ albido-hyalinæ, venis omnibus, maculisque numerosis nigris.

Brownish-black, covered with a whitish-gray pollen and with brownish-black spots and dots; wings whitish-hyaline, with black veins and numerous black spots. Long. corp. ♂ 0.15; ♀, 0.19. Long. al. 0.17.

The ground color of the body is an opaque brownish-black. Head of the same coloring, only the front, towards its anterior margin, seems to have a more or less reddish-brown or brownish-red ground color; the pollen on the whole head is whitish-gray; on the extreme lateral margin of the front it is more dense and almost white; upon the middle of the front and at a considerable distance from its sides, there are two oval, oblique, opaque, brownish-black spots; a spot of the same coloring surrounds the ocelli, and has, upon each side a smaller spot, upon which the

inner vertical bristle is inserted. The two superior hairs upon the sides of the front are prolonged and incrassated to the size of distinct bristles; above the two spots upon its middle, the front has no hairs, besides these bristles; below the spots, however, the front is beset with erect black hairs, inserted upon hardly perceptible dark dots. Antennæ ferruginous-brown, more distinctly ferruginous on their inner side towards the basis; the third joint round, black towards the end. Arista slightly incrassated at the basis and blackish-brown upon the incrassation, then pale yellowish and again darker towards the end. Thorax covered with a white-grayish pollen and with a brownish-black punctation and picture; the latter consists of ten regularly arranged spots upon its disk, and of a longitudinal stripe on each side, which begins at the anterior end and reaches up to the root of the wings; the picture of the pleuræ consists of two irregular longitudinal stripes; the pectus is neither punctate nor pictured, and the pollen upon it is not equally distinct when viewed from different sides. The rather turgid scutellum has a brownish-black picture, the whitish-gray pollen remaining visible on the lateral corners and at the end only. The abdomen agrees with the thorax in its coloring and has, besides the punctation, a regular and elegant brownish-black picture, which is more fully developed in the female than in the male; it consists of two small, approximated longitudinal stripes in the middle of the abdomen, which begin at the posterior end of the second segment and end at the posterior end of the fourth segment; on both sides of these stripes, between them and the lateral margin, there is a row of conspicuous spots, placed near the anterior margin of the segments and not reaching the posterior one. The first segment of the ovipositor is very broad and broadly truncate at the end; brownish-black, like the rest of the body; its basis is marked with two very large brownish-black spots, reaching as far as the middle and which have only a narrow stripe between them; the latter, as well as the posterior half are covered with a thin, whitish-gray pollen, and punctate with brownish-black. Feet brownish-black; knees, a rather broad ring upon the middle of the tibiæ and basis of the tarsi yellowish. Halteres blackish-brown, the stem, with the exception of its basis, of a dirty whitish. Wings hyaline, viewed obliquely strikingly whitish; all the veins black upon their whole extent. The picture consists of rather numerous black spots,

which seem to be rather constant in their position, but less constant in their extent; the figure is drawn from a female specimen, which has them less extended; usually, the inside of these spots is distinctly paler, but this varies in different specimens; very characteristic is the part of the picture surrounding the small crossvein, which does not seem to be subjected to any important variation.

Hab. New York (A. Fitch).

Observation.—The described specimens, a male and a female, were obtained by Baron Osten-Sacken from Dr. Fitch under the name of *Trypeta corticalis*.

4. S. vau SAY. ♀.—(Tab. VIII, f. 29.) Fusco-nigra, polline albocinereo aspersa, punctis maculisque fusco-nigris variegata; alæ hyaline, maculis octo nigris, quatuor costalibus, unicâ apicali, unicâ margini postico contiguâ reliquis majore et venam transversalem posteriorem includente, duabus denique minoribus venæ longitudinali sextæ appositis; præterea macula ovata lutescens permagna, a muculâ costali secundâ usque ad secundam venæ sextæ maculam pertinens conspicitur et macula costæ tertiæ eodem colore luteo cum maculâ marginis postici conjungitur, ita ut fascia integra, in mediâ alâ multo dilutior, appareat.

Brownish-black, powdered with whitish-gray, marked with brownish-black dots and spots; wings hyaline, with eight black spots, four on the costa, one at the apex, one, larger than the others, near the posterior margin, covering the posterior crossvein, two smaller spots upon the sixth longitudinal vein; besides, there is a very large ovate, brownish-yellow spot, extending from the second spot on the anterior margin to the second of the two smaller spots on the sixth longitudinal vein; the third spot on the anterior margin is connected by the same coloring with the spot upon the posterior crossvein, thus forming a complete crossband, which is much paler in the middle of the wing. Long. corp. 0.16—0.19; cum terebrâ 0.18—0.24; long. al. 0.14—0.18.

SYN. *Ortalis vau* SAY, Journ. Acad. Phil. VI, 184, 4.

Ground color opaque brownish-black. Head of the same color, but the anterior portion of the front and the face of a reddish-brown or dirty brick-red coloring, which sometimes also extends to the middle line of the front. The two superior hairs upon the lateral margin of the front are bristle-like. The front, from the anterior margin nearly as far as the ocelli, is clothed with black hairs, inserted upon impressed punctures; there is no definite picture upon it. The thorax is covered with a white pollen and punctate with brownish-black. The picture on the thoracic

dorsum consists of ten small spots, the inner ones among which are sometimes dissolved into dots, and moreover, on each side, of a row of spots, almost coalescent into an irregular longitudinal stripe, closely approximated to the exterior margin; upon the pleuræ there are two irregular longitudinal stripes; the pectus itself is brownish-black without any paler pollen or paler picture. Abdomen with a whitish-gray pollen, with brownish-black dots and with four longitudinal rows of brownish-black spots, placed upon the anterior portion of the segments; between them, upon the middle of the third and fourth segments, there are two still smaller spots upon the posterior portion of these segments. The first segment of the ovipositor is brownish-black, opaque, without any whitish-gray pollen and without picture. Feet brownish-black, sometimes only dark-brown in not fully colored specimens; knees, a ring in the middle of the tibiæ and the tarsi clay-yellow; usually the last three joints of the front tarsi and the last two on the posterior tarsi, are more or less infuscated. Malteres yellowish-white. Wings hyaline, rather whitish when viewed obliquely, their picture consisting partly of a black, partly of a clay-yellow or brownish-yellow coloring; there are four deep-black spots upon the costa; the first is composed of the incrassated humeral crossvein, and a short line, immediately beyond it, between the costa and the auxiliary vein, so that it has the shape of a fork, or almost of a ring; below the humeral crossvein, as well as below the small areuate crossband, there are small black dots (one under each); the second deep-black spot on the anterior margin lies in the costal cell, but little beyond the end of the small basal cells; it is circular; between it and the third spot on the anterior margin, there is a small deep-black dot, placed at the end of the auxiliary vein; the third, likewise deep-black spot on the anterior margin, lies on the end of the subcostal cell and reaches the second longitudinal vein; the fourth spot on the anterior margin lies before the end of the marginal cell; inside of this cell it is deep-black, but turns beyond it into brown and further into yellowish-brown; it ends in the middle between the third and fourth longitudinal veins, thus assuming the shape of a perpendicular crossband, which is broader at its anterior end. At the apex of the wing there is another black spot, which begins immediately beyond the termination of the second longitudinal vein and extends but little beyond the end of the fourth longitudi-

dinal vein. The posterior crossvein is covered by a brownish-black spot, which is especially expanded near the posterior margin and the anterior end of which is connected by an ochre-yellowish or somewhat brownish-yellow coloring with the third spot of the anterior margin, thus forming a complete crossband, somewhat expanded posteriorly and tinged with yellowish in the middle. Sometimes, however, the brown spot upon the posterior crossvein is somewhat more isolated from the yellowish coloring and extends in the direction of the half-crossband, formed by the fourth spot on the anterior margin. This less common variety is the one described by Say, l. c.; the ordinary picture is represented on Tab. VIII, f. 29, of the present volume. On the anterior side of the sixth longitudinal vein there are two black spots of only moderate size and rounded shape. Of them, the second only crosses that vein, gradually to fade away. Between the second costal spot and the second of the two spots of the sixth vein, there is a very large oval ochre-yellowish or more brownish-ochreous spot; it reaches on one side as far as the posterior basal cell, and assumes within the marginal cell a rather dark-brown coloring. The veins of the wing are black or brownish-black inside of the picture, clay-yellow elsewhere. The third and fourth longitudinal veins converge towards their end a little more than in the preceding species.

Hab. United States.

Observation.—I possess six female specimens and no male, but have seen the latter in other collections. It does not show any perceptible difference from the female, except in the sexual marks.

Gen. III. **CALLOPISTRIA** nov. gen.

Charact.—*General appearance* almost Trypeta-like.

Front exceedingly broad, with impressed punctures; cheeks comparatively broad; clypeus somewhat projecting over the edge of the mouth, sometimes withdrawn inside of the oral opening.

Wings with an unusually convex posterior margin; posterior crossvein very oblique, its anterior end much more approximated to the apex of the wing, than the posterior end; the posterior angle of the anal cell is drawn out in a very long, acute lobe.

The species upon which this genus is based, cannot well be placed in the genus *Stictocephala* on account of the remarkable difference in the outline of the wings as well as in the venation.

In other respects this species agrees with the preceding genus in the structure of the body; with *S. corticalis* and *vau* it even agrees in the coloring and the picture of the body, as well as in the bristle-like nature of the upper hairs on the sides of the front.

1. *C. annulipes* Macq. ♂ ♀.—(Tab. VIII, f. 27.) Fusco-nigra, albido-pollinosa, et punctis maculisque fusco-nigris variegata, tibiis tarsisque pallide lutescentibus, illis nigro-triannulatis, his apicem versus infuscatis; alæ hyalinæ, maculis punctisque nigris confertim aspersæ. Brownish-black, with a whitish pollen, pictured with brownish-black spots and dots; tibiæ and tarsi pale-yellowish, the former with three black rings, the latter brown towards their end. Wings hyaline, densely covered with black spots and dots. Long. corp. ♂ 0.14—0.15; ♀ 0.17; long. al. 0.16—0.18.

SYN. *Platystoma annulipes* MACQUART, Dipt. Exot. Suppl. V, p. 121.

The ground color of the body is brownish-black and opaque; the pollen, covering it, is whitish-gray. Head of the same color, covered everywhere with brownish-black spots, moreover, dotted with brownish-black upon the front and the cheeks; upon the posterior orbit especially there is a conspicuous short row of brownish-black spots. Front very broad, perceptibly narrower anteriorly, where it is yellowish or yellowish-red. Eyes rather strongly projecting. Antennæ brown, the first two joints and the larger part of the inner side of the third joint yellowish-brown, sometimes much paler. Thoracic dorsum with brownish-black dots, which coalesce into ill-defined, although regularly arranged, spots. Scutellum somewhat swollen, with four bristles, two brownish-black longitudinal stripes and two blackish-brown dots, upon which the lateral bristles are inserted. Pleuræ likewise with brownish-black dots and spots; the latter form two irregular and incomplete longitudinal stripes. Pectus brownish-black, with a brown, but little perceptible, pollen. Abdomen with brownish-black dots and regularly arranged spots; the first segment of the ovipositor is for the most part covered with a whitish-gray pollen and punctate with brownish-black. Femora brownish-black, with a more or less distinct, broad, irregular ring, covered with gray pollen, and with black dots; the tip is pale-yellowish. Tibiæ pale-yellowish with three regular brownish-black rings; the first near the basis, the last before the apex; tarsi of the same color as the tibiæ, infuscated towards the tip. Halteres pale-yellowish.

Wings of an unusual shape, on account of the great convexity of the posterior margin, hyaline, with black veins and numerous, partly only punctiform, partly rather large black spots of an irregular shape; the punctiform dots prevail in the middle, while the borders of the wing are principally occupied by larger spots, among which those along the posterior margin do not entirely reach the latter. The peculiarities of the venation are indicated above, among the generic characters.

Hab. United States; very common.

Observation.—I do not entertain the slightest doubt that Macquart's *Platystoma annulipes* is the above-described species. His description agrees perfectly well, with the exception of the words: "face blanche, une petite tache ronde d'un noir luisant de chaque côté." All my specimens have, on the sides of the face, or rather on the cheeks, nothing but brownish-black, opaque, irregular spots.

Gen. IV. **MYENNIS** R. Desv.

Charact.—*General appearance:* Trypeta-like.

Third antennal joint oval; cheeks broad, clypeus small, projecting over the edge of the mouth.

Wings narrow in comparison to their length, a little more attenuated towards the apex; the first longitudinal vein beset with bristles upon the portion only, which forms the limit of the very long stigma; before this spot the first longitudinal vein appears almost bare, the pubescence being very short and delicate; the two posterior basal cells are comparatively large; the posterior angle of the anal cell is pointed; the posterior end of both crossveins is nearer the apex of the wing than their anterior end, so that their position is a very oblique one.

The genus *Myennis* was established by Rob. Desvoidy for *Scatophaga fasciata* Fab. As *Trypeta scutellaris* Wied. agrees with that species in the above-enumerated characters, we can, for the present, unhesitatingly refer it to *Myennis*. The peculiarity, however, of the *Pterocallina*, of showing considerable plastic differences almost from species to species, appears again in the two above-named species. In *P. scutellaris* Wied. the eyes are less round, the cheeks broader, the scutellum less swollen, the crossveins less approximated, the longitudinal veins, instead of straight, somewhat undulated, and the third and fourth longitudinal veins, towards their end, not distinctly convergent, but parallel.

1. M. scutellaris WIED. ♂ ♀.—Cinerea, antennis flavis, pedibus ex fusco testaceis, thoracis margine laterali atro-maculato, scutello tumido; alæ angustæ, hyalinæ, fasciolâ basali, fasciis duabus discoidalibus antice connatis, plagâque apicali ex nigro fuscis pictæ, præterea in cellulis marginali et submarginali maculis aliquot fuscis variegatæ.

Var. ♂ fasciâ discoidalî secundâ inter venas transversales late interruptâ.

Cinereous, with yellow antennæ and brownish-yellow feet; the lateral margin of the thorax with black spots; the scutellum swollen; the narrow wings are hyaline; a small crossband at the basis, two crossbands, connected anteriorly, upon the middle of the wing and a large spot upon the apex, brownish-black; moreover several brown spots in the marginal and submarginal cells.

Var. ♂ the second of the two bands upon the middle of the wing, is broadly interrupted in the middle.

Long. corp. 0.17—0.18; long. al. 0.17—0.18.

Syn. *Trypeta scutellaris* WIEDEMANN, Auss. Zweifl. II, p. 484.

Trypeta? scutellaris LOEW, Monogr. of N. A. Dipt. I, p. 92. Tab. II, f. 26, 27.

Very like a *Trypeta* in its general appearance. Head comparatively high. The under side of the occiput rather tumid. Front yellow, of a medium breadth, long, its anterior margin rather projecting. Face somewhat retreating, a little excavated, infuscated inferiorly, covered with a pale-colored dust; antennal foveæ hardly indicated. Eyes oval. Checks brown, very broad. Proboscis not perceptibly incassated. Palpi short, but broad, of a dusky reddish-yellow; clypeus small and narrow. Antennæ ochre-yellow; the third joint oval, altogether rounded at the end; arista rather long and bare; it is thin at its end, but gradually stouter towards the basis. The upper part of the thorax dark-gray from a thick dust; the ground color of the humeri more or less ferruginous-yellow. Upon the lateral border of the thorax there is an irregular row of, for the most part contiguous, black spots; the largest among them is near the posterior corner; one is higher upon the upper part of the thorax than the others and near the transverse suture. The hairs and bristles are also placed upon very small, and but little perceptible, black dots. Scutellum with four bristles, rather turgid, of a shining dark-brown, with a clay-yellow median stripe; sometimes the clay-yellow color is more extended. Pleuræ blackish-brown, the posterior part yellowish-brown. Abdomen of the male cinereous; the penultimate segment shining-black, more thickly dusted towards the posterior

margin, and hence gray and opaque; the last segment is similar to the penultimate, only the dust on the posterior margin is less extended. The female has a blackish-gray abdomen (its coloring, however, seems to have been unnaturally modified in the four specimens which I had for examination); at the basis of the last three segments a darker coloring is perceptible, but it is not shining. The flattened, broad, yellowish-brown ovipositor is but very little attenuated towards its end. Feet brownish-yellow, the front femora at the basis, the four posterior ones near the apex, brown. The more maturely colored male has the greater part of the femora dark-brown, the first half of the tibiæ and a faded ring upon the middle of their second half, yellowish-brown. Wings comparatively long and narrow; the first longitudinal vein reaches far beyond the middle of the anterior margin and is beset with bristles along the side of the very long stigma only; the longitudinal veins have a very irregular undulated course; both crossveins have their anterior end nearer to the root of the wing, than the posterior end; their position is consequently a distinctly oblique one and both are slightly bisinuated; the third longitudinal vein is not beset with bristles. Both small basal cells are rather large in size; the posterior angle of the anal cell is strongly pointed; the third and fourth longitudinal veins are parallel towards their end. The stigma is brownish-black; a brownish-black picture is contiguous to it, which has almost the shape of an inverted V; it is formed by two crossbands which are coalescent in front; the first is broader and runs from the basis of the stigma over the basis of the discal and of the third posterior cells rather perpendicularly, almost reaching the posterior margin of the wing, while the narrower second band takes an oblique course over both crossveins, as far as the posterior margin; a short, but rather broad brownish-black crossband runs from the humeral crossvein as far as the basis of the anal cell; upon the apex there is a very large blackish-brown spot, beginning at the end of the marginal cell and extending to the tip of the second posterior cell; in the submarginal cell, between this large spot and the preceding crossband, there is a brownish-black spot of a considerable size, which, however, is very variable in different specimens; the portion of the marginal cell situated between the stigma and the apical spot has blackish-brown, brownish and almost hyaline spots; a small spot of a much darker

tinge lies near the anterior side of the second longitudinal vein, below the point of the stigma. The picture of the wings seems to be rather variable, the end of the exterior costal cell being sometimes blackish-brown, sometimes hyaline; the other dark spots are sometimes faded upon their middle, sometimes also less extended. In a male in the Berlin Museum, the only specimen of that sex which I have seen, the crossband covering both crossveins is broadly interrupted between them (compare the figure in Monographs, etc., Vol. I, Tab. II, f. 26). At first, I supposed this difference to be a sexual one, but I doubt this now, since I have had an opportunity of ascertaining the great inconstancy of the picture of the wings of the female.

Hab. Mexico.

Observation.—The figures given in the Monographs, etc., Vol. I, Tab. II, f. 26, 27, are sufficiently correct as far as the picture of the wing is concerned, but the outline of the wing is not well rendered; they are represented as too broad in proportion to their length.

SECOND DIVISION.

ORTALIDÆ HAVING THE FIRST LONGITUDINAL VEIN BARE.

First Section: ULIDINA.

GEN. I. DASYMETOPA LOEW.

Charact.—*Front* broad, narrower anteriorly, abundantly hairy on the whole surface, the hairs on its sides not longer.

Antenne rather short, third joint elongated-oval, with a thin, bare arista.

Face not excavated, descending vertically; *clypeus* projecting over the border of the mouth; *opening of the mouth* not large; *proboscis* but little thickened.

Thorax bristly on its hind part only; *scutellum* with a rather even surface and with four bristles.

Wings broader than those of the related genera; stigma of a very conspicuous size; posterior crossvein oblique, its anterior end being much nearer the apex of the wing than the posterior; the last section of the fourth longitudinal vein is strongly bent forward; the posterior angle of the anal cell is drawn out in a point.

The general appearance of the species of this genus is very much like that of *Trypeta*; the coloring of the species at present

known is not metallic. The peculiar venation distinguishes these species from all the others of the present group. The typical species is *D. lutulenta* Loew (Berl. Entom. Zeitschr. XI, 285; Tab. II, fig. 1), from Surinam.

No *Dasymetopæ* from North America are as yet known.

Gen. II. **OEDOPA** Loew.

Charact.—*Head* conspicuously large; *front* unusually broad; ocelli on the edge of the vertex, very closely approximated.

Antennæ very short and very distant from each other; third joint rounded, with a thin, bare arista; frontal fissure running in an almost straight line from antenna to antenna; no frontal lunule.

Face broad, somewhat convex, with a small excavation under each antenna; its lateral portions conspicuously broad, distinctly separated from the middle portion.

Eyes rather round, but somewhat broader than high, comparatively small, hardly reaching the middle of the height of the head; hence, the cheeks unusually broad.

Clypeus not horseshoe-shaped and thus surrounding the proboscis, but lobiform, connate with the anterior edge of the comparatively small oral opening; *proboscis* small.

Thorax with bristles on its hind part only; *scutellum* flat, with four bristles.

Wings: the last section of the fourth longitudinal vein, towards its tip, is somewhat curved forward and thus convergent towards the third vein; posterior crossvein curved in the shape of an *S*; posterior angle of the anal cell drawn out in an elongated point.

The body appears very bare on account of the sparseness and shortness of the hairs, as well as of the shortness of the bristles. The structure of the head resembles somewhat that of some South Asiatic *Ortalidæ*, while similar American forms have, before now, not been known.

1. O. capito Loew. ♂ ♀.—(Tab. IX, f. 1-3.) Albicans, fasciâ frontis tenui, thoracisque vittis nigris, in supero faciei margine maculis atris tribus, lateralibus ovatis, mediâ didymâ.

Whitish; front with a black transverse band, thorax with black longitudinal stripes; the upper margin of the face with three deep black spots; the lateral ones oval, the middle one double. Long. corp. 0.18—0.25; long. al. 0.15—0.22.

Syn. *Oedopa capito* Loew, Berl. Ent. Zeitschr. XI, p. 287, Tab. II, f. 2.

Head yellowish-white, only the middle of the occiput somewhat blackish; the ocelli are placed upon a punctiform black dot; the

very broad front has, somewhat below its middle, a narrow, gently curved, blackish crossband, above which the single, rather sparse hairs are inserted in small, somewhat darker colored pits; this is not the case below the crossband; no stripes run from the vertex down along the orbits of the eyes. Antennæ yellow, the place of insertion of the arista infuscated or blackened; between the eye and the antenna there is, on each side, a transversely-oval, velvet-black spot; between the antennæ and next to the frontal fissure is another velvet-black transverse spot; which consists of two small semi-oval transverse dots. The face, including the clypeus and the very broad cheeks, is more whitish than the front; the cheeks with a very delicate, easily rubbed off, whitish down. Eyes during life with two narrow crossbands, which are sometimes perceptible even in dry specimens. Palpi yellow, with delicate, pale hairs. Thorax and scutellum whitish-yellow; the dorsum of the thorax with six parallel, blackish longitudinal stripes; the two intermediate ones extend also over the flat scutellum. Pleuræ with three blackish longitudinal stripes, the upper one of which occupies the border between the dorsum and the pleura; quite downwards, moreover, there is a stripe-shaped black spot, which, however, seems to be produced by the rubbing off of the dust on the upper part of the pectus. Abdomen flat and rather narrow, whitish in consequence of the very dense dust which covers it; the ground color, however, is blackish, except the posterior part of the last segment in the female; the short, black hairs are inserted on small black dots, which are so closely approximated in the vicinity of the lateral border that they appear confluent, as irregular longitudinal spots; the last segment of the abdomen of the male is very much elongated; the first segment of the female ovipositor is attenuated towards its end, otherwise it looks like the remainder of the abdomen; its punctation, however, is much closer and finer; its adaptation to the abdomen is so close, that it might easily be taken for the last abdominal segment, especially when, as often happens, the black second and the yellowish third joint of the ovipositor are altogether withdrawn into it. Feet yellowish with whitish dust; the posterior femora generally with a blackish spot, on the under side before the tip; all the tibiæ with two black rings, the upper one of which is narrower and usually interrupted on the upper side of the tibia; fore tarsi blackened beyond the

tip of the first joint; the other tarsi blackened to a smaller extent. Halteres yellowish-white. Wings rather hyaline, of a dirty whitish tinge; the second and the next following longitudinal veins, as well as the crossveins which connect them, are black; the other veins yellowish; stigma small, of the same coloring as the rest of the wing; the picture of the wing consists of five brown spots with somewhat paler nuclei; three of them are in the marginal cell, near the anterior margin: the first, which like the second is oblique, is placed at the tip of the first longitudinal vein; the last is at the end of the marginal cell; beyond this is the fourth, a transverse spot in the submarginal cell, immediately under the tip of the second longitudinal vein; and again under the latter is the last spot, which is rounded and placed in the first posterior cell. The last three spots have the appearance of a narrow, very much shortened transverse band, which appearance is more distinct in those specimens, in which these spots are somewhat larger than usual. The small crossvein is beyond the middle of the discal cell, but before the tip of the first longitudinal vein.

Hab. Nebraska (Dr. Hayden).

Gen. III. NOTOGRAMMA Loew.

Charact.—*Front* of an equal, rather considerable breadth, scrobiculate.

Antennæ rather long; third joint elongated, with a thin, bare arista.

Face very short, the anterior edge of the mouth very much drawn upwards; *clypeus* considerably projecting over it.

Thorax with bristles on its hind portion only; *scutellum* flat, with sharp edges.

Wings: posterior angle of the anal cell drawn out in a point; second half of the last section of the fourth longitudinal vein very much bent forward; posterior crossvein perpendicular; auxiliary vein unusually short, and hence, the narrow stigma very long.

1. *N. stigma* FAB. ♀.—(Tab. IX, f. 5.) Nigro-chalybea, thorace lineis alternantibus nigris et læte virescentibus variegato, alarum limbo costali maculisque parvis nigris.

Blackish-steelblue, thorax with lines, showing alternately a blackish and a pale-green reflection; wings with the anterior margin bordered with black, and with small black spots. Long. corp. 0.11—0.16; long. al. 0.1.

SYN. *Musca stigma* FABRICIUS, Ent. Syst. Suppl. p. 563, 72.

Musca stigma FABRICIUS, Syst. Antl. p. 303, 96.

Dacus obtusus FABRICIUS, Syst. Antl. p. 278, 30.

Midia stigma WIEDEMANN, Auss. Zweifl. II, p. 565, 1.

Notogramma cimiciformis LOEW, Berl. Entom. Zeitschr. XI, p. 289, Tab. II, fig. 3.

Head rather disciform. Front reddish-brown, scrobiculate, remarkably hairy; the rather conspicuous stripes, descending from the vertex along the orbits of the eyes, and the elongated ocellar triangle are steel-bluish, shining; the ocelli are placed near the edge of the vertex, and are approximated to each other. The first two antennal joints brownish-black; the elongated third joint brownish-brickred, brown towards the tip. Face and clypeus metallic blackish-green, but little tinged with blue. The dorsum of the thorax has numerous black longitudinal stripes, which are separated by finer lines, having a metallic, light-green reflection and traced as if with a trembling hand. Pleuræ metallic blackish-steelblue, strongly tinged with greenish; above the fore coxæ with a large spot, covered with white pollen; from this place to the suture which runs down from the root of the wings, the pleuræ are covered with deep-black, punctiform dots, upon which single hairs are inserted. Scutellum rather large, flat, sharp-edged, metallic greenish-black, but rather dusky. Abdomen shining, blackish-steelblue; the first segment of the flattened ovipositor is of the same color, and attenuated towards its end. Feet black; tarsi brick-red, the foremost ones from the tip of the first joint, the four posterior ones from the tip of the second joint, brownish-black; the hind tibiæ somewhat compressed. Halteres dirty-yellow. Wings comparatively short, rather hyaline, with conspicuous, black veins; the costal and marginal cells have an altogether black coloring, which forms a border along the apex of the wing, extending from the tip of the marginal cell across that of the submarginal and of the first posterior cells; it becomes less intense here; in the submarginal cell, above the small crossvein, there is a black dot and farther towards the apex a small, triangular black spot; between the two again a black longitudinal line, which extends as far as the triangular spot; the picture in the first posterior cell is a similar one, only the first black dot is wanting and the two other black spots are somewhat more approximated to the apex of the wing; in the discal cell there are also two black spots, the smaller one before, the larger one beyond its middle; the second posterior cell is marked in the

middle with a punctiform black dot;¹ finally, in the third cell, not far beyond the fifth longitudinal vein, there are two successive punctiform blackish spots; the small crossvein is in the middle of the discal cell; the posterior crossvein is straight.

Hab. Cuba (Gundlach).

Observation.—The accurate knowledge which Wiedemann had of Fabricius's collection enables us to admit his authority as to the synonymy of *Dacus obtusus* Fab. with *Musca stigma* Fab. Wiedemann had a large number of specimens of *Musca stigma* (which he placed in the genus *Ulidia*) for comparison, and it is upon the ground of this comparison that he affirmed that the presence or absence of a pale spot upon the black border of the costa does not constitute a specific character. We can therefore safely accept the synonymy of *Musca stigma* Fab. with *Noto-gramma cimiciformis* Loew, the latter being the variety in which the pale spot is wanting.

Gen. IV. EUPHARA LOEW.

Charact.—*Front* of an equal, moderate breadth; scrobiculate, coarsely hairy.

Antennae almost more than of medium length; third joint elongated, with a thin, bare arista.

Face excavated; *clypeus* projecting.

Thorax with bristles on its hind part only; *scutellum* convex, with four bristles.

Wings: Posterior angle of the anal cell drawn out in a point; the last section of the fourth longitudinal vein parallel to the third; the small crossvein rather approximate to the posterior crossvein; the latter perpendicular.

The principal characters which distinguish this genus from the following one, to which it stands nearest, are the shorter and not attenuated stigma and the parallelism of the third and fourth longitudinal veins. Moreover, all the species of this genus seem to have black crossbands on the wings, while in those of the next following genus only the costal cell, the stigma, and the apex of the wing are blackened. The typical species is *Ceroxys coerulea* Macq. (Dipt. Exot. Suppl. III, p. 62, Tab. VII, f. 6), from Brazil, again described by me as *Euphara coerulea* (Berl. Ent.

¹ It is inadvertently omitted in the figure; the spots in the next cell likewise are but very feebly marked.

Zeitschr. XI, p. 291, Tab. II, f. 4; the figure of the wing is reproduced in the present volume, Tab. IX, f. 4).

I have not seen any North American *Eupharæ* yet.

Gen. V. ACROSTICTA LOEW.

Charact.—*Front* of an equal, moderate breadth, scrobiculate, rather coarsely hairy.

Antennæ rather short; the third joint elongate-ovate, with a thin, bare arista.

Face excavated, *clypeus* projecting.

Thorax with bristles on its hind part only; *scutellum* convex, with four bristles.

Wings: posterior angle of the anal cell drawn out in a point; the last section of the fourth longitudinal vein converges towards the third longitudinal vein; posterior crossvein perpendicular; stigma narrow and very long.

The difference between this genus and the preceding has been mentioned under the head of the latter. The characters which distinguish *Acrosticta* from *Euxesta* are: the elongated shape of the third antennal joint, the front, which is marked with pits, the stouter proboscis and the very long, narrow stigma. The picture of the wings resembles that of the species of *Seoptera*, except that the somewhat turgid front of the latter shows no vestige of pits and the face is not transversely excavated, but carinate. As typical species may be considered either *A. scrobiculata* Loew (Berl. Ent. Zeitschr. XI, p. 293, Tab. II, f. 5) or *A. foveolata* Loew (ibid. p. 294), both from Brazil.

No North American species is as yet known.

Gen. VI. SEOPTERA KIRBY.

Charact.—*Front* of equal breadth, somewhat elevated, with very short hairs.

Antennæ rather long, the broad third joint elongate-oval, with a thin, bare arista.

Face carinate, *clypeus* projecting.

Thorax with bristles on its hind part only; *scutellum* convex, with four bristles.

Wings comparatively long; the posterior angle of the anal cell pointed; the very long last section of the fourth longitudinal vein converges towards the third vein.

Feet somewhat longer and more slender than those of the related genera.

Kirby called this genus *Scioptera*. Following the usual rule of latinization, I modified the name to *Scoptera*. Later, Rob. Desvoidy called this genus *Myodina*; this name, however, cannot supersede the older one of Kirby, which, moreover, characterizes very well the peculiar habit of the species belonging here.

1. *S. colon* Loew. ♂ ♀.—(Tab. IX, f. 6.) Nigra, nitida, fronte rufâ, antennis et facie ex rufo flavis, alarum maculâ apicali triangulâ et cellulæ costalis basi nigris, stigmatè subfusco.

Shining black, front red, antennæ and face reddish-yellow; a triangular spot on the apex of the wing and the basis of the costal cell black; stigma brownish. Long. corp. 0.19—0.21; long. al. 0.19—0.22.

Syn. *Scoptera colon* Loew, Berl. Ent. Zeitschr. XI, p. 296, Tab. II, f. 6.

Of a shining black, somewhat bluish-black color; the abdomen more glossy than shining. Front of a fiery red, opaque, along the orbit of the eyes with a delicate line, powdered with white pollen. Antennæ yellowish-red; the third, elongate-oval joint is rather broad. Face and clypeus brilliant reddish-yellow, the latter often, the former seldom, tinged with chestnut-brownish. On the dorsum of the thorax there are two narrow lines of whitish pollen, which extend beyond its middle; they are easily overlooked, although very distinct in well-preserved specimens. Feet black, the tips of the femora and tibiæ and the basis of the hind tarsi have a reddish-brown tinge, even in specimens of the darkest coloring; in lighter specimens this coloring is brownish-brickred, and extends not only over the greater part of the tibiæ and the hind tarsi, but is also perceptible at the root of the fore tarsi. Halteres pale-yellowish. Wings hyaline; costa, auxiliary vein, and first longitudinal vein black; the other veins much paler, generally yellowish when seen in a reflected light. The costal cell blackened as far as the humeral crossvein; the stigma, as well as the whole subcostal celi, at the end of which it is placed, brownish; at the apex of the wing there is a triangular black spot, which covers the extreme tip of the marginal cell as well as the tip of the submarginal cell, and crosses a little beyond the third longitudinal vein. The small crossvein is nearly under the middle of the stigma, but beyond the middle of the discal cell; the last section of the fourth longitudinal vein is particularly long, straight, gradually converging towards the third; the anal cell is

broad and has a sharp posterior angle, although it is hardly drawn out in a point.

Hab. New York (Osten-Sacken); Illinois (Kennicott).

Observation 1.—This species, as far as I know, is undescribed, although not absolutely new, because Wiedemann, as his collection shows, received it from Say under the name of *Ortalis colon*. Harris, in his Catalogue of the Insects of Massachusetts, also has *O. colon*, which is undoubtedly the same species. I preserved the name which Say gave it, although I do not find it described in his works.

Observation 2.—*Scoptera colon* is so exceedingly like the European *S. vibrans* Lin., that as long as I had only indifferently preserved specimens of it, I took it for the latter species. Although the differences are only slight, they are so constant that the specific distinctness of the two species cannot be called in doubt. The front of *S. colon* is somewhat broader than that of *S. vibrans*; the two whitish stripes of the thorax in *S. colon*, although but little apparent, can easily be traced beyond the middle of the dorsum, while in *S. vibrans* it is not without difficulty that their anterior end alone can be perceived. The abdomen of *S. colon* is always less shining, and its blackish color more bluish, while *S. vibrans* has it more blackish-green. The costal cell of *S. colon* is blackish as far and even a little beyond the humeral crossvein; in *S. vibrans* this cell is entirely hyaline as far as its extreme basis; the stigma of *S. colon* is brownish, that of *S. vibrans* black or brownish-black; finally the black spot at the tip of the wings is somewhat different in both species; that portion of it which crosses the third longitudinal vein is of more equal breadth in *S. colon*, whereas it becomes more narrow towards the margin of the wing in *S. vibrans*.

Gen. VII. EUXESTA Loew.

Charact.—*Front* of equal, medium breadth, even, rather coarsely hairy.

Antennæ short, the third joint almost round or rounded-oval, with a thin, bare arista.

Face more or less excavated, *clypeus* projecting.

Thorax with bristles on the hind part only; *scutellum* convex.

Wings: posterior angle of the anal cell drawn out in a point; the last section of the fourth longitudinal vein converges towards the third; posterior crossvein perpendicular.

The general appearance of the species belonging here is not unlike *Trypeta*. Legs short. The coloring is metallic; the black picture of the wings consists either of some large spots along the anterior margin or of crossbands. The plastic characters of the species do not afford any features for their satisfactory distribution into groups; for this reason the following three groups are merely based upon the picture of the wings.

1st Group. *Wings with spots along the anterior margin.*

1. *E. spoliata* LOEW.—(Tab. IX, f. 7.) Viridis, capite pedibusque flavis, extremo femorum apice fusco, tibiis anticis fere totis, reliquarum apice tarsisque inde ab articuli primi apice nigris, alarum stigmatate nigro, maculâ subapicali nigricante.

Green, head and feet yellow, the extreme tip of the femora brown, fore tibiæ almost entirely, the tips of the four posterior tibiæ and the tarsi, from the tip of the first joint, black; wings with a black stigma and with a blackish spot immediately before the tip. Long. corp. 0.12; long. al. 0.12—0.13.

SYN. *Euxesta spoliata* LOEW, Berl. Ent. Zeitschr. XI, p. 298, Tab. II, f. 7.

Metallic-green, shining; the color of the scutellum and of the anterior segments of the abdomen is somewhat more bluish-green. Head yellow; the upper part of the occiput is blackish-green; front ferruginous-yellow; the swellings descending from the vertex along the orbit of the eyes and the immediate vicinity of the ocelli is metallic greenish-blue. Antennæ of a dark ferruginous-yellow; third joint round. Face shorter than in most of the other species. Clypeus yellow, protruding considerably beyond the anterior border of the mouth, although projecting but little in profile. Feet yellow; all the femora distinctly infuscated at the extreme tip; fore tibiæ rather stout, brownish-black, before the middle with an incomplete yellow ring; the intermediate tibiæ are blackened at the extreme tip only, the hind tibiæ also at the tip, but to a greater extent; the first joint of the hind tarsi is yellow, except the tip; the following joints are black (the intermediate and hind tarsi are wanting in the described specimen) Halteres yellowish. Wings pure hyaline with pale clay-yellow veins; extreme root of wings pale yellowish; the last section of the fourth longitudinal vein is but very slightly arcuated, but converges in its whole length towards the third vein, its tip thus approaching very near this vein; stigma blackened;

immediately before the tip of the wing there is a blackish spot, which reaches from the anterior margin to the third longitudinal vein and covers the extreme end of the marginal cell; the extreme end of the submarginal cell is not covered by it. It may be that, in more fully colored individuals, this spot is darker.

Hab. Cuba (Riehl).

2. *E. pusio* Loew; ♀.—(Tab. IX, f. 8.) Viridis vel ex chalybeo viridis, thoracis dorso albido-pollinoso, pedibus piceo-nigris, genibus, tibiæ apice tarsisque totis luteis, alarum stigmatibus et maculâ subapicali nigris.

Green or bluish-green; dorsum of the thorax covered with a white pollen; feet piceous-black; knees, tips of the tibiæ and the whole of the tarsi of a dirty-yellow; wings with a black stigma and a black spot immediately before the apex. Long. corp. 0.12; long. al. 0.13.

Syn. *Euxesta pusio* Loew, Berl. Ent. Zeitschr. XI, p. 299, Tab. II, f. 8.

Metallic bluish-green; thorax and scutellum rather opaque, in consequence of a comparatively dense white pollen; abdomen shining; its first segment of a dirty-yellow towards its sides. The very broad first segment of the flattened ovipositor is almost as long as two-thirds of the abdomen. Head of a reddish-brick color; the sides of the front, the frontal lunule, the face, including the clypeus and the cheeks, are covered with a rather dense, white pollen. The black hairs on the front are not conspicuous. Antennæ brownish-ferruginous, or rusty-brown; third joint round. Face rather short, considerably excavated; clypeus but little projecting beyond the opening of the mouth. Occiput apparently altogether metallic black, but the ground color is very much concealed by a thick whitish pollen. Feet piceous black; the second joint of the coxæ, the knees, almost the whole latter half of the tibiæ and the whole tarsi dirty-yellow or brick-red. Halteres whitish-yellow. Wings somewhat whitish hyaline, the veins pale; stigma of a blackish color, which, on its first half, extends as far as the middle of the marginal cell; immediately before the apex of the wing there is a black spot, extending from the anterior border as far as a little beyond the third longitudinal vein, the tip of the marginal cell is also covered by it, that of the submarginal cell, however, is not; the last section of the fourth longitudinal vein in its whole course, converges towards the third and comes very near it at its tip; it is not perceptibly arcuate.

Hab. Cuba (Gundlach).

3. E. notata WIED. ♂ ♀.—(Tab. IX, fig. 9.) *Chalybeo-nigra*, abdomine feminae fasciâ apicali flavâ ornato, pedibus nigris, genibus, tarsorumque basi rufis, alarum maculis duabus nigris, alterâ costali minutâ, alterâ apicali trigonâ, cellulæ costalis basi et stigmatæ cinereis.

Bluish-black, abdomen of the female with a yellow crossband at the tip, feet black, knees and the root of all the tarsi red; wings with a small black dot in the middle of the costa and with a larger triangular spot at the tip; basis of the costal cell and stigma gray. Long. corp. 0.15—0.16; long. al. 0.15.

SYN. *Ortalis notata* WIED. Auss. Zweif. II, p. 462, 9.

Euzesta notata LOEW, Berl. Ent. Zeitschr. XI, p. 300, Tab. II, f. 9.

Of a blackish-steelblue, generally verging on green-blue, often with a violet hue on the middle of the abdomen; rather shining. Front of a saturate yellowish-red, sometimes almost yellowish-brown; with a whitish pollen along the lateral orbit of the eyes; the black hairs are scattered and not conspicuous; the swellings running from the vertex downwards, along the borders of the eyes, generally also the immediate vicinity of the ocelli are shining bluish-black or black. Antennæ brown, ferruginous-red at the basis, which color is more extended on the inner side; third joint rounded. The very considerably excavated face, together with the rather projecting clypeus are bluish-black, very shining; the upper portion rather densely pollinose, and hence opaque, the ground color not being distinctly visible; the lateral swellings of the face are tinged with brownish-red and thinly whitish pollinose. The female has the latter part of the last abdominal segment, as well as the basis of the ovipositor of a saturate yellow color; in the male, I have never observed any trace of this yellow coloring. The first segment of the very much flattened ovipositor is of a very moderate breadth, brownish-black, but with a more or less distinct coppery-red reflection. Feet black, femora in part metallic-black or bluish-black; knees and the root of all the tarsi brick-red, on the front tarsi this red generally reaches only as far as the middle of the first joint, on the hind tarsi as far as the tip, on the intermediate ones as far as the basis of the next joint. Knob of halteres yellowish; stem generally infuscated. Wings hyaline with rather dark veins; at the tip of the costal cell there is a small black dot, which extends posteriorly as far as the second longitudinal vein; at the apex of the wing there is a larger triangular black spot, occupying the end of the sub-

marginal cell and crossing to a small extent the third longitudinal vein, but being exactly limited by this vein near the apex of the wing; the costal cell is tinged with brownish-gray as far as a little beyond the humeral crossvein; the stigma has the same color, but this is sometimes more blackened, especially towards its end; the last section of the fourth longitudinal vein converges in its whole course towards the third vein and approaches it considerably towards its end; shows, however, hardly any perceptible curvature; the fifth longitudinal vein does not quite reach the margin of the wing.

Hab. District of Columbia, New York, Illinois, Connecticut (Osten-Sacken).¹

Observation.—Wiedemann gives a description of the male of this species which might easily lead to the conclusion that he had before him a species different from the one I have just described. According to his statement, the male has, on the posterior margin of the penultimate abdominal segment, a saturate yellow crossband. But as Wiedemann's collection contains as *Ortalis notata* the very species which I described under this name and as, among a considerable number of males which I have before me, not a single one is provided with such a crossband, I am compelled to come to the conclusion that Wiedemann mistook the sex of the specimen from which he drew his description; he may have had before him a female the ovipositor of which was bent under the abdomen.

4. *E. nitidiventris* n. sp. ♀.—Nigro-viridis, nitida, abdomine femine toto seneo-viridi et nitidissimo, pedibus gilvis, tibiis anticis totis posterioribusque adversus apicem infuscatis, tarsis adversus apicem fusco-nigris, alarum maculis duabus nigris, alterâ costali minutâ, alterâ apicali trigonâ, cellulâ costalis basi luteâ, stigmatè ex luteo cinereo.

Shining black-green, the entire abdomen of the female metallic-green, very shining. Feet saturate yellow, the entire fore tibiæ and the posterior ones towards their tip, infuscated; tarsi brownish-black towards the tip; wings with a small black dot on the middle of the costa and with a larger triangular spot at the apex of the wing; basis of the costal cell clay-yellow; stigma yellowish-gray. Long. corp. 0.14—0.15; long. al. 0.14—0.15.

¹ Mr. Riley gave me a male specimen of *E. notata* which he bred from the pulp of an osage-orange (*Maclura*).—O. S.

Dark-green, shining, the abdomen altogether of a vivid metallic green, very shining. The femora of a saturate dark-yellow; this coloring changes into brownish on the fore tibiæ from the very basis, on the posterior tibiæ farther down; the fore tarsi are saturate yellow at the basis as far as the tip of the first joint, the posterior tarsi nearly as far as the end of the second joint, beyond this the tarsi are brownish-black. The basis of the costal cell is clay-yellow, or pale ferruginous-yellow, as far as a little beyond the humeral crossvein; the stigma is yellowish-gray. In all other respects this species is so very like *E. notata*, that one would be inclined to take it for a mere variety of coloring, unless the much lighter coloring of the feet, combined with the darker coloring of the much more shining abdomen, proved the contrary.

Hab. Texas (Belfrage).

5. *E. costalis* FAB. ♀.—(Tab. IX, f. 10.) Nigro-chalybea, pedibus nigris, genibus tarsorumque basi rufis, alarum maculis duabus magnis, alterâ costali, alterâ apicali, nigris.

Blackish-steelblue, feet black, knees and roots of the tarsi red; wings with two large black spots, the first in the middle of the costa, the second at the apex of the wing. Long. corp. 0.15; long. al. 0.15.

SYN. *Musca costalis* FAB. Ent. Syst. IV, p. 360, 196.

Dacus costalis FAB. Ent. Syst. Antl. p. 278, 25.

Dacus aculeatus FAB. Syst. Antl. p. 275, 14.

Ortalis costalis WIED. Auss. Zweiff. II, p. 464, 13.

Euzesta costalis LOEW, Berl. Ent. Zeitschr. XI, p. 301, Tab. II, f. 10.

Very like both preceding species, but easily distinguished by the narrower front, the absence of a yellow crossband at the end of the abdomen of the female, the perceptibly larger size of the black spot on the middle of the anterior margin of the wings, the altogether black stigma and the course of the fifth longitudinal vein, which reaches the margin of the wing. Blackish-blue, shining; the head brick-red or of a rusty-red; front anteriorly of a more saturate coloring, narrow, somewhat whitish pollinose on the orbit of the eye; the hairs upon it are rather sparse and not at all conspicuous; the stripes, descending from the vertex along the orbits of the eyes and the immediate surroundings of the ocelli are steel-blue, shining. Occiput blackish, its lower portion and a spot back of the region of the ocelli, brick-red. Antennæ brick-red or more yellowish-red; third joint rounded-

oval, generally somewhat infuscated on the outer side, towards the tip. The larger portion of the rather excavated face shows a steel-blue, shining color, which is but little concealed by the whitish pollen; the projecting clypeus also has a steel-blue reflection. The first joint of the flattened, black ovipositor is of moderate breadth. Feet pitch-black, femora almost bluish-black, knees and basis of all the tarsi brick-red. Halteres of a dirty-whitish or yellowish color. Wings hyaline, almost grayish, with very dark veins; the root of the wings up to the humeral cross-vein and a little beyond, blackish; the whole stigma, as well as the tip of the costal cell and a spot connected with the latter and reaching as far as the fourth vein, are black; at the apex of the wing there is a large, triangular black spot, covering the tip of the marginal and the end of the submarginal cell, and, moreover, crossing to a considerable extent the third longitudinal vein, so that its posterior limit is not far from the fourth longitudinal vein and runs parallel to it. The last section of the fourth longitudinal vein is only very gently curved and converges towards the third in its whole course, approaching the latter considerably towards its end.

Hab. West Indies (coll. Wied.).

6. *E. quaternaria* Loew. ♀.—(Tab. IX, f. 11.) Nigro-violacea, dimidio apicali abdominis flavo, alarum maculis costalibus quatuor nigris.

Blackish-violet, second half of the abdomen yellow; wings with four spots along the anterior margin. Long. corp. 0.12—0.14; long. al. 0.13—0.14.

Syn. *Euxesta quaternaria* Loew, Berl. Ent. Zeitschr. XI, 302, Tab. II, f. 11.

Blackish-violet, the middle of the thoracic dorsum, a large portion of the pleuræ and the sides of the abdomen often more blackish-blue. Front rather narrow, ferruginous, along the orbits of the eyes with a very delicate border of white pollen and with coarse black hairs; the little stripes, running from the vertex down the sides of the front are blackish, but hardly shining. Antennæ ferruginous-brown, more reddish at the basis, sometimes of a lighter coloring; the third joint is rounded. Face, including the but little projecting clypeus and the cheeks brownish-red, less excavated than in most of the other species. Occiput for the most part black. Thoracic dorsum with a thin covering of whitish-gray pollen, and hence but little shining; more so on its

sides; the pleuræ likewise are rather shining. The anterior part of the abdomen of a metallic dark-violet hue; the apical half yellow, sometimes with a dark stripe in the middle. The first segment of the very much flattened ovipositor rather broad and long, black, usually with a bronze reflection. Feet black, only the extreme tip of the femora reddish-brown and the basis of all the tarsi brick-red. Halteres yellowish, the stem usually infuscated. Wings hyaline, with four black spots on the anterior margin; the first among these spots, placed on and immediately beyond the humeral crossvein, extends as far as the basis of the anal cell, so that the extreme root of the wing itself is hyaline; the second spot, covering the tip of the costal cell and the very short stigma, with the exception of its extreme end, runs perpendicularly and preserves the same breadth, as far down as the fourth longitudinal vein, beyond which it is still perceptible as a blackish-gray shadow; the third black spot lies opposite the posterior crossvein, is of an elongated triangular shape, and reaches with its tip as far as midway between the third and fourth longitudinal veins, the fourth spot has an irregularly rounded shape and lies quite near the apex of the wing; it covers the extreme end of the marginal cell and the end of the submarginal with the exception of its extreme tip; on its posterior side (that is the side which is nearer the basis of the wing) it crosses the third longitudinal vein; the last section of the fourth longitudinal vein, which is distinctly, although not strongly, curved, converges in its whole course towards the third longitudinal vein, without approaching it more, however, than in the several preceding species.

Hab. Cuba (Gundlach).

2d Group. *Wings with two, very much abbreviated, crossbands.*

7. E. binotata Loew. ♂.—(Tab. IX, f. 12.) Nigro-chalybea, capite, lateribus segmentorum abdominalium primi et secundi femoribusque luteis, tibiis tarsisque fusco-nigris, alarum fasciis duabus postice valde abbreviatis nigris.

Dark steel-blue, the head, the sides of the first two abdominal segments and the femora yellow; the tibiæ and tarsi brownish-black; wings with two very much abbreviated black bands. Long. corp. 0.12; long. al. 0.13.

Syn. *Euresta binotata* Loew, Berl. Ent. Zeitschr. XI, p. 304, Tab. II, f. 12.

Head dark-yellow; front rather broad, with a very narrow border of white pollen; the hairs upon the front are not conspicuous. The stripes descending from the vertex along the sides of the front and the immediate surroundings of the ocelli are steel-bluish, somewhat shining. Antennæ dark-yellow; their third joint rather round. Face rather excavated, with a white pollen which is less dense in the vicinity of the anterior border of the mouth, and from under which a faint steel-blue reflection is still visible. Clypeus but moderately projecting over the anterior edge of the mouth, generally of a dark-yellow color, seldom with a faint trace of a steel-blue reflection. The upper portion of the occiput, with the exception of a large spot behind the vertex, is steel-blue, with a whitish pollen. Thorax steel-bluish, with a rather whitish pollen and hence but moderately shining. Scutellum, metathorax and abdomen bright, shining, almost metallic black; the sides of the first and second segments of the abdomen have a yellow coloring, which, however, usually does not reach the posterior margin of these segments and sometimes is more expanded in the middle. Front coxæ and femora dark-yellow; tibiæ, with the exception of the extreme basis, and the tarsi brownish-black. Halteres whitish with a dirty-brownish stem. Wings hyaline; immediately beyond the humeral cross-vein there is a small black spot, which extends, in the shape of a crossband, as far as the root of the anal cell; the rather long stigma is black; from its basis a black crossband extends in a somewhat oblique direction as far as the middle of the discal cell; immediately before the apex of the wing, another black perpendicular crossband is situated; anteriorly it is somewhat widened, posteriorly it crosses the fourth longitudinal vein, the last section of the fourth longitudinal vein is moderately but distinctly curved, and converges with the third longitudinal more in its latter half than in its first. The intervals between the black crossbands of the wings of this species, as in most of the others, by transmitted light assume a rather indistinct white coloring, in a similar light, however, the apex of the wings of this species assumes a very striking whitish coloring.

Hab. Cuba (Gundlach).

3d Group. *Wings with four crossbands.*

S. E. annonæ FAB. ♂ ♀.—(Tab. IX, f. 13.) Nigro-chalybea, fasciis alarum nigris quatuor, secundâ postice abbreviatâ et reliquis paulo latiori.

Dark steel-blue; wings with four black bands, the second of which is abbreviated posteriorly and is somewhat broader than the others. Long. corp. 0.14—0.15; long. al. 0.14—0.15.

SYN. *Musca annonæ* FAB. Ent. Syst. 358, 189.

Tephritis annonæ FAB. Syst. Antl. IV, p. 320, 19.

Ortalis annonæ WIED. Auss. Zweifl. II, p. 463, 11.

Urophora quadri vittata MACQ. Suites, II, p. 456, 5.

Euxesta annonæ LOEW, Berl. Ent. Zeitschr. XI, p. 305, Tab. II, f. 13.

Head brick-red; the little stripes running down from the vertex and the region of the ocelli steel-blue, rather shining; the larger portion of the occiput blackish, with a grayish-white pollen. The front is of only moderate breadth; the hairs upon it are strikingly coarse, more dense upon the pollinose lateral borders, scarce upon the remainder of the surface. Antennæ brick-red, third joint rounded-oval, towards its end brownish and more so on its outer than on its inner side. The face is very moderately excavated; the clypeus moderately projecting, both with a steel-blue reflection and a white pollen. Thorax of a very dark color, verging sometimes on green, sometimes more on steel-blue or violet, and always covered with a rather dense whitish pollen. The scutellum is of a still more dark violet-black color, and less pollinose. The abdomen is of a metallic, but very dark bluish-black or violet-black color. The first segment of the flattened ovipositor is generally still darker. Fore coxæ, with the exception of their basis and the tip of the hind coxæ, brownish-brickred, the former with a white pollen. Femora black; the first pair, and generally also the last, more metallic-black; all are brownish-brickred at the tip; tibiæ blackish-brown; dark brick-red at the tip and often also at the extreme root; tarsi brick-red at the basis, blackish-brown towards the tip. Halteres clay-yellow. Wings hyaline, with four black crossbands. The first lies upon and a little beyond the humeral crossvein and reaches the basis of the anal cell; the second begins at the anterior margin with the but moderately long, black stigma and the blackened extreme tip of the costal cell; it is perpendicular and reaches beyond the fifth

longitudinal vein, without, however, reaching the margin of the wing; the small crossvein lies exactly upon its external limit; the internal one is always sinuate in the vicinity of the fifth longitudinal vein; the third and fourth bands are connected at the anterior margin in such a manner, that the hyaline space between them reaches either exactly as far as the second longitudinal vein, or goes very little beyond this vein; the third band, which is nearly straight and rather perpendicular, runs over the posterior crossvein and almost reaches the posterior margin of the wing; the fourth crossband is of considerable breadth, reaches as far as the fourth longitudinal vein and is continued even beyond it, in the shape of a gray shadow; the second half of the last section of the fourth longitudinal vein is very gently curved anteriorly, so that it converges towards the third longitudinal vein, without approaching it, however, to any considerable extent.

Hab. Cuba (Gundlach).

9. E. Thomæ LOEW. ♂ ♀.—(Tab. IX, f. 14.) *Læte chalybea, nitidissima, alarum fasciis nigris quatuor subintegris, ultimis tribus latis.*

Bright steel-blue, very shining; wings with four black crossbands, the last three of which are broad. Long. corp. 0.14—0.15; long. al. 0.14—0.15.

SYN. E. Thomæ LOEW, Berl. Ent. Zeitschr. XI, p. 306, Tab. II, f. 14.

Very like the preceding species, although very probably a distinct one, notwithstanding the great resemblance in all the plastic characters. The differences are the following: the whole coloring of the body is of a lighter and more brilliant steel-blue, which often verges on violet in the middle of the abdomen. The thoracic dorsum is much less pollinose. The second crossband of the wings is broader, approaches more the posterior margin of the wings, and is not sinuate on its inner side in the vicinity of the fifth longitudinal vein. The third crossband is much broader than in *E. annonæ*, especially its anterior portion; the fourth band crosses the fourth longitudinal vein a little, or else the gray shadow beyond the end of this vein is somewhat darker.

Hab. St. Thomas (Westermann).

10. E. abdominalis LOEW. ♂ ♀.—(Tab. IX, f. 15.) *Chalybeo-nigra, abdominis basi sordide luteâ, alarum fasciis nigris quatuor integris, ultimis duabus ad costam anguste cohærentibus.*

Bluish-black, with a dirty-yellow basis of the abdomen; wings with four complete black crossbands, the last two of which are connected by a narrow stripe at the costa. Long. corp. 0.12—0.14; long. al. 0.12—0.14.

SYN. *Euresta abdominalis* Loew, Berl. Ent. Zeitschr. XI, p. 307, Tab. II, f. 15.

Head brick-red or brownish-brickred; the small stripes running down from the vertex along the orbits of the eyes and the surroundings of the ocelli, are steel-blue, shining; almost the whole occiput is black, with a grayish-white bloom. Front rather narrow; rather dense and conspicuously coarse hairs upon the lateral borders, which are covered with white pollen; the hairs upon the remainder of the surface are very scarce. Antennae brownish-brickred, or brick-red; in the latter case the rounded-oval last joint is more or less infuscated towards its end. The face is moderately excavated, usually for the most part with a shining steel-blue reflection; its white bloom is very thin along the edge of the mouth. Clypeus only moderately projecting, with a more or less distinct steel-blue reflection on the sides. Thorax of a shining, blackish-steelblue color, which usually verges somewhat on green upon its dorsum. Scutellum and metathorax still darker greenish-black, not pollinose. Abdomen more greenish-black than bluish-, or metallic-black, at the basis always dirty clay-yellow. The coloring of the first segment of the flattened ovipositor is the same as that of the abdomen, or a more purely black one. Fore coxæ, at the tip at least, brownish-brickred, with white pollen; femora black, more or less metallescens, with a brownish-brickred tip; tibiæ blackish-brown, only the extreme tip reddish-brown; tarsi reddish-brown at the root, otherwise blackish-brown. Halteres whitish or yellowish. Wings with four not abbreviated black crossbands. The first is broader than in the two preceding species, but is likewise placed upon and immediately beyond the humeral crossvein, and extends as far as the basis of the anal cell. The second band begins at the anterior margin with the black tip of the costal cell and the black stigma; it is rather broad and gradually expands in approaching the posterior margin so that, at this place, its breadth exceeds considerably that of the other bands; the small crossvein lies exactly upon its outer margin. The third band likewise, which runs over the posterior crossvein, has a considerable breadth and a very perpendicular position. The fourth band runs along the

apex of the wing; it is also rather broad and reaches beyond the fourth longitudinal vein; its connection with the third band near the costa is rather narrow, so that the hyaline space, inclosed between them, almost reaches the costa anteriorly. The last section of the fourth longitudinal vein is gently arcuate and slightly converges in its latter half towards the third longitudinal vein (the figure does not give this quite correctly).

Hab. Cuba (Gundlach).

Observation.—The Museum at Vienna contains a couple of specimens taken in Cuba by Pöppig, which differ, however, by their distinctly smaller size, as well as by a somewhat different picture on the wings; all the four black bands are dissolved into oval black spots, covering the veins, the portions of the bands lying inside of the cells are crossed in the middle by gray stripes. A closer examination, however, proves conclusively that these specimens are incompletely colored ones of *E. abdominalis*. The small size is probably due to the greater contraction in drying of these unripe specimens.

11. *E. alternans* Loew. ♂.—(Tab. IX, f. 16.) Obscure chalybea, alarum fasciis nigris quatuor integris, omnibus separatis, tertiâ reliquis multo angustiore.

Dark steel-blue, wings with four complete black crossbands, entirely separate from each other; the third much narrower than the others. Long. corp. 0.13; long. al. 0.13.

Syn. *Euzesta alternans* Loew, Berl. Ent. Zeitschr. XI, p. 308, Tab. II, f. 16.

Head brick-red or brownish-brickred; the little stripes running down from the vertex along the orbits of the eyes, as well as the surroundings of the ocelli, of a shining steel-blue; the whole occiput blackish, with a whitish pollen. Front rather narrow, with coarse hairs which are more dense on the somewhat whitish, pollinose, lateral borders and more sparse on the remaining surface. Antennæ brick-red or yellowish-red, the third joint oval. Face very much excavated; with the exception of its lower, considerably projecting, portion, it has a steel-blue reflection, but is so thickly covered with a white pollen, that the bluish ground-color is but little apparent. Clypeus rather strongly projecting, brownish-brickred, sometimes with a steel-blue reflection on the sides. Thorax and scutellum of a rather dark, steel-blue color, which turns somewhat to greenish-blue on the thoracic

dorsum; the latter is but little pollinose. Abdomen darker steel-blue, shining, especially on the sides. Feet black; the knees and the first joint of the intermediate tarsi brick-red, the first joint of the hind tarsi brown or reddish-brown towards the basis. Halteres yellowish-white. Wings with four black unconnected bands. The first of them lies, as in the preceding species, on and immediately beyond the humeral crossvein and reaches the basis of the anal cell. The second band begins at the black stigma and runs, expanding somewhat, as far as the posterior margin, in the vicinity of which it gradually becomes fainter; the third band is narrow, perpendicular, and covers the posterior crossvein; the fourth runs along the apex of the wing, is even broader than the second and completely isolated from the third; beyond the fourth longitudinal vein, it becomes very faint. The last section of the fourth vein is rather strongly curved and its latter portion converges towards the third vein.

Hab. Brazil? Cuba? (Vienna Museum).

Observation.—The description is drawn from a male specimen in the Vienna Museum, labelled: Mann, Toscana 1846. As I have seen the same species, in other collections, marked as Brazilian, I take the designation of the Vienna Museum to be erroneous. I am confirmed in this supposition by the fact that next to the above-mentioned specimen is placed another, a female, pinned on the same kind of pin and labelled in the same manner, which, however, is a specimen of *E. stigmatias*, received hitherto from Cuba and Brazil only. Thus it appears evident that both specimens were sent by the same collector, probably from the same country; and as *E. stigmatias* is a common species in Cuba, the conclusion is not too far fetched that both specimens came from that island. This is the reason why I did not like to omit *E. alternans* in this volume.

12. *E. stigmatias* Loew. ♂ ♀.—(Tab. IX, f. 17.) Nigro-viridis. maculâ atrâ inter antennis sitâ insignis, alarum fasciis nigris quatuor, ultimis duabus ad costam conjunctis.

Blackish-green, conspicuous by a deep black spot between the antennæ, wings with four black bands, the last two of which are connected near the costa. Long. corp. 0.13—0.15; long. al. 0.14—0.15.

Syn. *Euxesta stigmatias* LOEW, Berl. Ent. Zeitschr. XI, p. 310, Tab. II, f. 18.

Head dark metallic-green or almost steel-blue. Front of a dusky-red; the little stripes running down from the vertex along

the orbits of the eyes, as well as the well-defined ocellar triangle, shining steel-blue. The lateral border of the front shining and generally with a rather distinct steel-blue reflection; immediately above each antenna, a trace of a small swelling is discernible. The hairs on the front are not conspicuous, moderately dense on the sides, very scarce on the remaining surface. The first two joints of the antennæ brownish-black, the rounded-oval third joint reddish-yellow from the basis as far as the arista, more brownish beyond it. Face very much excavated, shining steel-blue, with a whitish pollen on its upper part only; above this, just between the antennæ, is a conspicuous, velvet-black spot. Clypeus very much projecting, shining, steel-blue, pollinose on the margins only. The rather broad orbital circles of the eyes brick-red below, at the lower corner of the eyes. Thorax dark metallic-green, somewhat verging on steel-blue; the dorsum with a very thin gray pollen. Scutellum blackish-green. Abdomen of the same color as the thorax, but darker, often with a stronger steel-blue reflection; the last abdominal segments of the male sometimes more bronze-colored. The first segment of the flattened ovipositor metallic-black. Feet black; the tips of the knees and the basis of all the tarsi brownish-brickred. Halteres white-yellowish. Wings with four black crossbands. The first lies, as in several other species, on and immediately beyond the humeral crossvein and extends as far as the basis of the anal cell. The second band, which is rather broad, begins at the costa with the blackish end of the costal cell and the black stigma; it is generally very much fainter beyond the fourth longitudinal vein and disappears entirely between the fifth vein and the posterior margin of the wing; the small crossvein lies almost exactly upon the outer margin of this band. The third band, which is perpendicular, runs over the posterior crossvein and reaches the posterior margin of the wing almost completely; it is broader anteriorly than posteriorly, and is connected with the fourth band on the inside of the marginal cell, so that the hyaline interval between these bands extends exactly as far as the second longitudinal vein. The fourth band, lying along the apex of the wing, is also rather broad and extends as far as the fourth longitudinal vein. The last section of the fourth vein is distinctly curved and in its second half converges towards the third longitudinal vein.

Hab. Cuba (Gundluch); Brazil (coll. Wintham).

13. *E. eluta* LOEW. ♂ ♀.—(Tab. IX, f. 18.) Nigro-viridis, subchalybescens, maculâ atrâ inter antennis sitâ insignis, alarum fasciis nigris quatuor, secundâ latissimâ sed maximâ ex parte valde elutâ, tertiâ et quartâ in cellulâ costali per maculam hyalinam separatis.

Blackish-green, verging on steel-blue, conspicuous by a deep black spot lying between the antennæ; wings with four black bands, the second of which is the broadest, but, for the most part, very pale; the third and fourth are separated by a hyaline spot, lying in the costal cell. Long. corp. 0.14-0.15; long al. 0.14-0.15.

SYN. *Euzesta eluta* LOEW, Berl. Ent. Zeitschr. XI, p. 312, Tab. II, f. 19.

Front red or brownish-red; the little stripes, descending from the vertex along the orbits of the eyes and the well-defined ocellar triangle, are shining steel-blue; the hairs on the front are not striking, moderately dense on the but slightly pollinose lateral borders; otherwise very scarce. Occiput blackish-steelblue, with a grayish-white bloom. Antennæ ferruginous-brown or reddish-brown, more brick-red at the basis of the third joint; sometimes the second joint has the same coloring. Face rather excavated, generally steel-blue, or at least reddish along the anterior edge of the mouth only; in some rare cases it has a light steel-blue reflection on its upper part, the remainder brick-red; exactly between the antennæ is a conspicuous velvety-black spot; clypeus but little projecting beyond the edge of the mouth, reddish-brown, with a steel-blue reflection; the orbits of the eyes brick-red or brownish-red near the lower corner of the eye. Thorax dark metallic-green; in less mature specimens greenish steel-blue. Thoracic dorsum only slightly pollinose. Scutellum more blackish-green or blackish-blue. The color of the abdomen is not unlike that of the thorax, but is darker and verges on bluish; its middle sometimes almost violet; the last segments of the male abdomen sometimes bronze-colored. Front coxæ, except the root, brick-red, with white pollen. Feet black, the tips of the knees and the root of all the tarsi brick-red; the tip of the middle tibiæ likewise is generally brick-red; sometimes the extreme tip of the front tibiæ shows a brick-red coloring. Halteres yellowish-white. Wings with four black crossbands. The first lies upon and immediately beyond the humeral crossvein and extends to the extreme basis of the anal cell; it is rather narrow and often pallid. The second crossband is of considerable breadth, begins near the costa with the infuscated tip of the costal cell and the black stigma; but beyond the third,

or the fourth longitudinal vein it is so very faint that it extends to the posterior margin of the wing in the shape of a gray shadow; the small crossvein lies, when the band is not too pale, almost exactly upon its external margin. The third band passes over the posterior crossvein, is narrow and generally rather pale, except in the vicinity of the anterior margin; towards the posterior end of the crossvein it almost disappears; from the fourth band it is separated by a rather large, whitish-hyaline spot in the marginal cell; behind the second longitudinal vein fully colored specimens have, on the outer side of this third band a rather distinct gray shadow, between which and the fourth band only a narrow, whitish hyaline interval remains, from which, however, the above-mentioned hyaline spot near the costa is completely isolated. The fourth band, which lies along the apex of the wing, extends as far as the fourth longitudinal vein, or else it crosses it in the shape of a gray shadow. The last section of the fourth longitudinal vein is rather strongly curved and convergent towards the third vein.

Hab. Cuba (Gundlach).

Gen. VIII. CHÆTOPSIS LOEW.

Charact.—*Front* of medium breadth, somewhat narrower towards the vertex, with a row of bristly hairs on the lateral border; the remaining surface not hairy.

Antennæ rather short; third joint very little excised on the upper side, with a sharp anterior corner and a thin, bare arista.

Face but moderately excavated; *clypeus* but little projecting over the anterior border of the mouth.

Thorax with bristles on its posterior part only; *clypeus* convex, with four bristles.

Wings: posterior angle of the anal cell drawn out in a point; last section of the fourth longitudinal vein, towards its end, but very little convergent with the third vein; posterior crossvein perpendicular.

The species known to me are conspicuous for the striking length of the bristles, inserted on the posterior part of the thorax and on the scutellum. Their coloring is metallic; the wings are adorned with well-defined black crossbands. They cannot well be confounded with the species of the preceding genus, on account of their greater slenderness, and more especially, on account of the different shape of the third antennal joint and of the front, which is hairy on its lateral borders only. From the two next following

genera, which likewise have the third antennal joint with a sharp anterior angle, the species of the present genus are sufficiently distinguished by their less slender shape and the different structure of the anal cell, not to mention other characters.

1. *C. ænea* WIED. ♂ ♀.—(Tab. IX, f. 19.) Viridis, antennis fusconigris, basi tamen articuli tertii luteâ, alis trifasciatis.

Metallic-green; antennæ brownish-black, the basis of the third joint yellow; wings with three bands. Long. corp. 0.16—0.18; long. al. 0.17—0.18.

SYN. *Ortalis ænea* WIED. Anss. Zweifl. II, p. 462, 8.

Ortalis trifasciata SAY, Journ. Acad. Phil. VI, p. 184, 3.

Urophora fulvifrons MACQ. Dipt. Exot. Suppl. V, p. 125, Tab. VI, f. 9.

Chortopsis ænea LORW, Berl. Ent. Zeitschr. XI, p. 315, Tab. II, f. 21.

Trypeta (Acinra) ænea v. d. Wulp, Tijdschr. voor Ent. 1867, p. 137, Tab. V, f. 12—14.

Front red, on each side with a broad band, which is covered with white pollen. The ocelli rather far distant from the edge of the vertex; the region of the ocelli, as well as the little stripes descending from the vertex along the orbits of the eyes are blackish-green, only very little shining. Frontal lunule with white pollen. Antennæ rather short; the first two joints brown, the second sometimes in part brownish-yellow; the third joint rather broad, very little excised on the upper side, always with a sharp anterior angle, brownish-black, reddish-yellow at the basis. Face only little excavated, steel-bluish, but rather opaque on account of a whitish pollen; the edge of the mouth usually brick-red. The clypeus has but a small transverse diameter and is but little projecting over the anterior edge of the mouth. Thorax and scutellum shining metallic-green, upon the dorsum with a trace of a white bloom. Abdomen of the same color, or somewhat more bronze-green, the last joints of the male abdomen generally blackish-green. With less mature individuals the coloring of thorax and abdomen is more bluish-green, and at the basis of the latter an unmetallic, dirty-yellow coloring may be seen. The coloring of the feet is variable; in some specimens they are altogether pale-yellow, only a little darker at the tip of the tarsi; as this occurs in those specimens which have the basis of the abdomen yellow, one might almost be led to the conclusion that they form a distinct species; however, the absolute similarity of all the other characters renders this conclusion very improba-

ble; darker specimens have the color of the feet more brownish-yellow, the root of the front coxæ and the tip of the tarsi dark-brown; the femora of such specimens often show conspicuous black, metallic-green longitudinal stripes; the darkest specimens have the whole basal half of the femora, and even more, of this black coloring, while the tibiæ also are partly infuscated. Halteres yellowish-white. Wings with three brownish-black bands; the veins are black upon these bands, but ochre-yellow elsewhere, which gives the whole basal part of the wing an ochre-yellowish tinge. The first band begins at the costa with a short black stigma, is perpendicular and rather dark, as far as the fourth longitudinal vein and even beyond; the remainder of the band, as far as the posterior margin of the wing, is usually very faint; the small crossvein is a little beyond the margin of this band; the second band runs over the posterior crossvein and is perpendicular and rather broad; its posterior end is very pale; with the third band it is generally connected only by a dark border along the costa; sometimes, however, this border becomes broader and extends in some specimens as far as the second longitudinal vein. The third band, running along the apex, is likewise rather broad, extends as far as the fourth longitudinal vein, and even beyond it, in the shape of a gray shadow. The last section of the fourth longitudinal vein, beyond its middle, converges towards the third; near its tip, however, this convergence becomes again much less.

Hab. United States, rather common (Osten-Sacken); Louisiana (Sehann) Cuba (Gundlach).

Observation 1.—The comparison of the types in Wiedemann's collection do not allow any doubt about the determination of this species; they belong to the variety of a paler, but not of the palest, coloring. Say's good description of *Ortalis trifasciata* refers to the variety with dark feet. That Macquart's *Urophora fulvifrons* belongs here seems certain; that he placed the species in the genus *Urophora* is no objection, because he did the same with several *Ortaliidæ*; the figure of the wing, which he gives, is incorrect, as the comparison of the description shows; the latter proves conclusively that the second crossvein on the middle of the wing is an arbitrary addition; it seems that Macquart drew the small crossvein correctly on the extreme limit of the first crossband; later, however, in finishing his figure, he noticed that in

consequence of the very exaggerated breadth of the interval between the first and second bands, the position of the small crossvein with regard to the posterior one had become altogether distorted, and in order to correct this, he may have drawn the small crossvein a second time, at a correct distance from the large one. Mr. Van der Wulp has erroneously taken *Chætopsis ænea* for a *Trypeta* and, supposing it a new species, accidentally described it under the same specific name.

Observation 2.—The *Urophora ænea* Macq. (Suites, etc., Dipt. II, p. 458, 13), may be a synonym of the present species, although I do not consider this as certain. The figure of the wing, as given in Dipt. Exot. II, 3, Tab. XXX, f. 7, shows at the basis of the wing an extensive and very conspicuous black spot, of which there is no vestige in *C. ænea*. It seems certain that *Urophora ænea* Macq. is a species belonging to the present group of *Ortalidæ*.

2. *C. debilis* Loew. ♀.—(Tab. IX, f. 20.) Viridi-chalybea, antennis totis pedibusque flavis, alis trifasciatis.

Greenish-blue; the entire antennæ and the feet yellow; wings with three bands. Long. corp. 0.12; long. al. 0.11.

SYN. *Chætopsis debilis* Loew. Berl. Ent. Zeitschr. XI, p. 318, Tab. II, f. 22.

Very like the preceding species, but smaller; the white bloom forming a border on both sides of the front is comparatively a little broader; it has a single row of four bristles upon it, whereas in the preceding species these hairs are much more numerous. The antennæ are altogether yellow and their third joint upon its upper side is somewhat more excised. The stigma is comparatively smaller; the three bands have the same position, but are less pale towards the posterior margin; the last two are entirely separated from each other, which is very seldom the case with *Chætopsis ænea*; the last section of the fourth vein is much more straight and shows only a vestige of a slight convergency towards the third longitudinal vein. The coloring of the described specimen is not green, but greenish steel-blue; of a dirty-yellowish at the basis of the abdomen; but as it is a rather immature specimen, these differences cannot have much weight. The first segment of the flattened ovipositor is comparatively long.

Hab. Cuba (Gundlach)

Gen. IX. **HYPOECTA** Loew.

Charact.—*Front* of an equal, rather considerable breadth, somewhat projecting when viewed in profile; delicately hairy on the sides only.

Antennæ short; third joint very much excised on the upper side, with a very sharp anterior corner and with a thin, bare arista.

Face not excavated, somewhat retreating on the under side; *clypeus* rudimentary, not projecting over the edge of the mouth, of a very small transverse diameter.

Thorax with bristles on its hind part only; *scutellum* convex, with four bristles.

Wings: posterior angle of the anal cell pointed, open; the last section of the fourth longitudinal vein converges somewhat towards the third; the posterior crossvein perpendicular.

The species of this genus are considerably more slender than the species of *Chælopsis* and their shape is somewhat more like that of *Eumetopia*. The third antennal joint, the shape of which reminds one of *Cerorys*, the not excavated face, the rudimentary clypeus and the open anal cell, are easy to recognize. The ovipositor is conspicuously broad, and so closely joined to the abdomen that it may be easily mistaken for its last segment. The typical species is *H. longula* Loew, Berl. Ent. Zeitschr. XI, p. 319, Tab. II, f. 23, from Santos (in Brazil).

No North American species are as yet known.

Gen. X. **STENOMYIA** Loew.

Charact.—*Front* of equal breadth, somewhat projecting in profile, hairy on the sides; upon the remaining surface with two longer hairs only.

Antennæ rather short, third joint hardly excised upon the upper side, but with a sharp anterior angle; arista thin and bare.

Face not excavated, somewhat retreating, with a slight depression under each antenna; gently convex between these depressions; *clypeus* of moderate transverse diameter, somewhat projecting over the border of the mouth.

Thorax with bristles on its posterior part only.

Wings comparatively long; posterior angle of the anal cell sharp, but not pointed, last section of the fourth longitudinal vein about double the length of the preceding section, gently converging towards the third longitudinal vein; posterior crossvein rather perpendicular.

The striking slenderness of the narrow body and the metallic coloring, are points of resemblance between the species of this genus and those of *Eumetopia*; the picture of the wings is like-

wise a similar one. The former are at once distinguished, however, by the front, which is not conically projecting. They are characterized also by the shape of the wings and the venation, which it will be easier to understand from the figure than from a description.

1. *S. tenuis* LOEW. ♂.—(Tab. IX, f. 21.) Chalybeo-viridis, pedibus nigris, basi tarsorum rufâ; alis cinereis, stigmatè et plagâ permagnâ apicali nigris.

Greenish-steelblue, the feet black, the root of the tarsi red; the grayish wings have a black stigma and a large black spot at the apex. Long. corp. 0.14; long. al. 0.13.

Syn. *Stenomysia tenuis* LOEW, Berl. Ent. Zeitschr. XI, p. 321, Tab. II, f. 24.

Front brown, almost black above, rather hairy along the orbits of the eyes, upon the remaining surface only with two more elongated hairs; the little stripes running down from the vertex along the orbits of the eyes and the ocellar triangle are dark bluish-green, shining. Antennæ black; the second joint at its upper corner to a certain extent dirty-whitish; third joint rather broad, upon the upper side hardly excised, but with a sharp anterior corner. Face somewhat retreating, with a distinct depression under each antenna, longitudinally convex along its middle, dark steel-blue, shining, but on its upper half with a thin, whitish bloom. Clypeus of a very moderate transverse diameter, but distinctly projecting over the upper border of the mouth, deep steel-blue and shining. Palpi black. Eyes large and rather round, their horizontal diameter, however, is a little larger than the vertical one. Cheeks narrow. Thorax dusky blue-green, rather shining, scutellum greenish-black, but little shining, with an entirely even upper side. The narrow and long abdomen has the same coloring as the thorax; however, towards its extremity it gradually becomes more black and opaque. Feet black; femora and tibiæ with metallic, dark bluish-green reflections; the extreme tips of the tibiæ and the root of the tarsi are dark brick-red, the remainder of the feet brownish-black. Halteres whitish. Wings long and narrow, grayish, the root and a rather large spot immediately behind the stigma lighter; the rather small, narrow stigma is of a black color, which extends below it as far as the second longitudinal vein; the last third of the wings, beginning at the costa as far as the fourth longitudinal vein, is tinged with

blackish; this color, at its inner border, between the third and fourth longitudinal veins, is very pale, and extends sometimes as a gray shadow even beyond the fourth vein; the first, second, third, and fifth longitudinal veins are conspicuous for their stoutness and black color; the basis of the second vein and the portion of it lying in the clear spot beyond the stigma, are of a paler color and less stout. The small crossvein is immediately below or but little beyond the end of the stigma; but always beyond the middle of the discal cell; the posterior crossvein is perpendicular; the last section of the fourth longitudinal vein is conspicuous for its great length and converges gently towards the third; anal cell with a sharp angle, which is not, however, drawn out in a point.

Hab. Georgia.

Gen. XI. **EUMETOPIA** Macq.

Charact.—*Front* very much projecting anteriorly, so that the head, seen in profile, appears conical; upon its sides and its anterior part it is sparsely beset with short, not erect, hairs.

Antennæ of middle size; third joint oval, with a bare arista.

Face unusually retreating, almost horizontal, below each antenna distinctly excavated and with a small ridge between these impressions; *clypeus* small, but distinctly projecting over the anterior edge of the mouth.

Wings narrow and rather long; stigma very narrow, posterior angle of the anal cell acute; the last section of the fourth vein somewhat converging towards the third near the tip.

The species of this genus are always bare, very slender and have a metallic coloring; moreover, they are easily distinguished by the extraordinary projection of their foreheads and the conical profile of their heads; the picture of their wings only consists in a more or less extended black spot on the apex.

1. E. rufipes Macq. ♂.—(Tab. IX, f. 22.) Viridis, pedibus luteis; alarum apice nigro.

Green, feet dark-yellow; wings with a blackish apex. Long. corp. 0.2; long. al. 0.13.

Syn. *Eumetopia rufipes* Macq. Dipt. Exot. Suppl. II, p. 88, Tab. VI, f. 2.

Eumetopia rufipes Loew, Berl. Ent. Zeitschr. XI, p. 322, Tab. II, f. 25.

Front reddish-brown, often very dark, the projecting portion on both sides of a lighter coloring; moreover, both sides of the front have a white, pollinose margin; the sides and the anterior

portion bear some scattered, short, neither numerous nor erect hairs; the little stripes running down from the vertex along the orbits of the eyes and ocellar triangle are of a shining metallic-green; the latter is somewhat distant from the vertex. Antennæ rather deep black; face and clypeus moderately shining, bluish-black; the lower orbit, however, reddish-brown, with a narrow white border. Palpi and proboscis dark-yellow. The thorax, the moderately convex scutellum, and the abdomen shining metallic-green; the latter, however, becomes more opaque and darker towards its end. The fore coxæ altogether, the second joint of the posterior ones and the feet of a rather dark, saturate yellow coloring, but by no means red; the front tarsi altogether and the tip of the posterior ones brownish-black. Wings narrow, somewhat grayish-hyaline; the veins are tinged with yellow at the basis and in the proximity of the anterior margin, as far as the black spot on the apex; this gives to those parts of the wings a yellowish coloring; the other veins are blackish; a large brownish-black spot on the apex of the wing occupies almost one-quarter of the length of the wing and extends beyond the fourth longitudinal vein. The small and narrow stigma is yellowish. The small crossvein generally lies only a little beyond the end of the stigma and very little beyond the middle of the discal cell; the last section of the fourth longitudinal vein is perceptibly longer than the interval between both crossveins, and gently converges near its end towards the third vein; the posterior crossvein is always perpendicular; the posterior angle of the anal cell acute.

Hab. United States, not rare (Osten-Sacken).

2. *E. varipes* LOEW. ♀.—(Tab. IX, f. 23.) Viridis, femoribus nigris, genibus tibisque luteis, alarum apice nigro.

Green, femora black, knees and tibiæ yellow; wings with a blackish apex. Long. corp. 0.25; long. al. 0.12.

Syn. *Eumetopia varipes* LOEW, Berl. Ent. Zeitschr. IX, p. 181.

Eumetopia varipes LOEW, Berl. Ent. Zeitschr. XI, p. 323, Tab. II, f. 26.

Very like *E. rufipes*, but easily distinguished on account of the different coloring of the feet. Front almost black. The palpi of the only specimen in my possession seem to be yellowish-brown. Scutellum more flattened than that of *E. rufipes*. Coxæ and femora black with a bluish-green metallic reflection;

the tip of the femora and the tibiæ clay-yellow, the latter sometimes brownish-yellow; tarsi brown, the posterior ones paler at the basis. The first segment of the flattened ovipositor black. The wings of the same outline as those of *E. rufipes*, but the veins at the basis and in the vicinity of the anterior margin less yellow; the small crossvein is far beyond the middle of the discal cell, and hence it is less distant from the posterior crossvein; the fifth longitudinal vein is interrupted at a somewhat greater distance from the posterior margin of the wing and the last section of the fourth vein converges a little more towards the third; the blackish spot at the apex of the wing is perceptibly larger, so that it occupies more than one-fourth of the length of the wing. All the rest as in *E. rufipes*.

Hab. Cuba (Gundlach).

Second Section: RICHARDINA.

Gen. I. CONICEPS nov. gen.

Charact.—*Head* in shape like a long, somewhat flattened cone; *Front* rather broad, eyes rather distant from the posterior edge of the head; their horizontal diameter somewhat longer than the vertical one.

The first two antennal joints short, the third elongated and of equal breadth, *arista* bare.

The *metathoracic bristle* indicated only by a hardly perceptible little hair; *prothoracic bristle* not extant.


Scutellum with two bristles.



Abdomen slender and elongated.

Femora not incrassated, unarmed; the under side of the hind ones with some rather stiff bristles.





Wings: posterior angle of the anal cell abbreviated; crossveins not approximated; the smaller one on the middle of the discal cell; the third and fourth longitudinal veins parallel.

The present genus is very like *Eumetopia* on account of its narrow, elongated shape and its strongly projecting front. I place it here in order to bring it as near as possible to *Eumetopia*, although I am far from considering it as a typical genus of the group *Richardina*. It is distinguished from *Eumetopia* not only by the abbreviated angle of the anal cell, but also by the still more projecting forehead, by the somewhat turgid, cushion-shaped occiput, and by the comparatively shorter, but broader wings.





**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

1.43 2.8
1.32 2.5
1.24 2.2
1.20
1.18

10

1. *C. niger* n. sp. ♂ ♀.—*Modice nitens, niger, halteribus concoloribus, coxis pedibusque luteis, alis cinereis, adversus costam et apicem nigris.*

Moderately shining, black, with the halteres of the same color; coxæ and feet dark-yellow; wings gray, tinged with black along the anterior margin and towards the apex. Long. corp. ♂ 0.15; ♀ cum terebrâ 0.21; long. al. ♂ 0.11; ♀ 0.13.

Black, moderately shining, beset with short, unobtrusive, black hairs. The bristles on the sides of the vertex of medium length. Antennæ deep black; third joint comparatively long, of equal breadth, rounded at the end; the bare arista of medium length, perceptibly stouter towards the basis; the parts of the mouth comparatively small and rather hidden. Abdomen long and narrow, of almost equal breadth. The first segment of the ovipositor, at the basis, has the same breadth as the posterior margin of the last abdominal segment; it is but little narrowed towards its end; it is clothed with a black pubescence which, although by no means long, is nevertheless rather conspicuous; in some specimens its sides are turned upwards, so that it appears narrower towards its end than it really is; the second and third joints of the ovipositor are generally retracted within the first, which might produce the impression that the species is a new form of *Micropezidæ*; when they are projecting, both prove to be comparatively rather broad and the third ends in a short, but sharp point. Coxæ and feet are of a dark yellow color; the tarsi are infuscated towards the tip; the hind femora with some stiff bristles of moderate length on their under side, which, however, do not resemble spines. Halteres blackish. Wings gray with black veins; the apex of the wings blackened and the costa with a black border, beginning at the tip of the costal cell.

Hab. Texas (Belgrave).

Gen. II. **RICHARDIA** ROB. DESV.

Charact.—*Front* of most species rather broad; ocelli not far from the edge of the vertex; the anterior one more distant from the posterior ones than these from each other; in the males of several species the head is very much expanded transversely, as in the species of *Achius*.

Arista pubescent, or short-feathery.

Mesothoracic and *prothoracic* bristles present, although weak.

Scutellum with four bristles; *metathorax* steep.

Abdomen narrow, still more narrowed towards the basis.

Front femora only moderately incrassated; the intermediate ones not

at all; the hind femora very much incrassated, beset with spines on the under side.

Wings: the crossveins approximated to each other; the third longitudinal vein towards its tip is more or less curved backwards; the third and fourth veins, for this reason, appear convergent; posterior angle of the anal cell obtuse.

The characters distinguishing this genus, which is peculiar to America, are as follows: the rather equally narrow abdomen; the unarmed front and middle femora; the very much incrassated hind femora, the under side of which is beset with spines; finally, the crossveins being approximated to each other.

The rather coarse hairs upon the feet of most species of *Richardia* look somewhat like spines at the further end of the under side of the front and middle femora; although I have not observed any real spines upon the under side of the four anterior femora in any of the species which I have examined.

The mention of the presence of the prothoracic and mesothoracic bristle has been introduced among the characters of this and of the following genera, wherever I was able to do so. But, as in several cases I had only a single, perhaps not particularly well-preserved, specimen for comparison, or one in which this character could not very well be ascertained, the statement about the absence of one of these bristles is not to be taken too strictly until further confirmation.

The typical species is the well-known *Richardia podagrica* Fabr., from South America.

Gen. III. CYRTOMETOPA nov. gen.

Charact.—Front broad, very much projecting in profile.

Arista pubescent.

Femora strong, although not exactly incrassated; all are beset with spines.

Wings: posterior angle of the anal cell obtuse; crossveins not approximated to each other; the end of the fourth longitudinal vein converges very much towards the tip of the third vein.

The typical species is the *Odontomera ferruginea* Macquart (Dipt. Exot. II, 3, p. 215), in which, with tolerable certainty, I recognize an American species.

The *Odontomera maculipennis* Macquart (Dipt. Exot. Suppl. I), from Columbia, probably belongs to the genus *Cælometopia*.

I have drawn the characters of this genus, as far as it was

possible, from Macquart's statements. The characters which prevent me from uniting this genus with the following are: the front, very much projecting in profile, the much shorter and stronger femora, the wings, which are not attenuated towards their basis, and the strong convergency of the third and fourth longitudinal veins. If the auxiliary vein is really as far distant from the first longitudinal as Macquart's figure shows it, this would furnish one distinctive character more.

Gen. IV. **STENOMACRA** nov. gen.

Charact.—*General shape* almost like *Sepsis*.

Front rather broad, somewhat narrower anteriorly.

Ocelli closely approximated to each other, almost in the middle of the front.

Antennal arista with a very distinct pubescence.

No mesothoracic and, to all appearances, no prothoracic bristle.

Scutellum with two bristles; *metathorax* sloping.

Abdomen narrow, almost pedunculate.

Feet slender, femora not incrassated, the intermediate ones attenuated towards the end; the hind femora a little longer than the middle ones; all are beset with spines towards the tip.

Wings rather large, very much attenuated towards the basis; posterior angle rounded off; the auxiliary vein very much approximated to the first longitudinal, coalescing with it at the tip; the second longitudinal reaches the margin of the wing far from the apex; the small crossvein is far before the middle of the discal cell; the last section of the fourth longitudinal vein almost parallel to the third vein; posterior angle of the anal cell obtuse.

I. S. Guerini Big. ♂ ♀.—(Tab. IX, f. 25., Rufescens, pleuris, scutello, metanoto abdominisque basi nigris; alæ hyalinæ, strigulâ subbasali et maculâ magnâ apicali nigris.

Reddish, pleuræ, scutellum, metathorax and the basis of the abdomen black; wings hyaline with a little black streak at the basis and a large black spot at the apex. Long. corp. 0.20; long. al. 0.20—0.22.

Syn. *Sepsis Guerini* Bigot, De la Sagra, Hist. física, etc., p. 822, Tab. XX, f. 9.

Ferruginous-red, rather shining, the upper part of the occiput, as well as the region of the vertex and the little stripes running down from it upon the front, sometimes shining black, almost metallic. Front rather broad, somewhat narrower anteriorly; the bristles of the vertex long; the bristle in front of them, inserted upon the little stripe, is likewise long, removed to almost the middle of the front. The ocelli, near which the ordinary two

bristles are placed, are likewise removed to about the middle of the front and are close to each other. Antennæ descending to the edge of the mouth; the first two joints yellow; the third more or less infuscated; the arista with a very distinct, somewhat rare, pubescence. Face of the *Dacus*-like shape, peculiar among the *Richardina*; proboscis and palpi sometimes of a dirty reddish-yellow, sometimes more brownish-ferruginous. The thoracic dorsum somewhat ferruginous; only very dark specimens have it black; the hairs upon it are placed in four distinct longitudinal rows, the intermediate ones being very closely approximated. Scutellum convex, with two bristles, black ferruginous on the sides in very pale-colored specimens only. Pleuræ, with the exception of the humeral region, as well as the whole metathorax, black. The basis of the abdomen is black to a greater or less extent; in rare specimens only does this color reach the posterior margin of the rather considerably elongated first abdominal segment; in some specimens, however, this color extends to the very end of the abdomen, or, at least, turns here into blackish-brown. The ovipositor, which is longer than the last three abdominal segments taken together, is usually black or blackish-brown; its upper side is excavated (at least in dry specimens), and its under side convex, and hence, it is less flattened than in the other genera of the *Richardina*. Coxæ pale-yellow. Front feet pale-yellowish; the tibiæ towards the basis and the tarsi, beginning from the second joint, infuscated; femora not incrassated, beset with a few, but rather strong, spines on the under side towards its end. The anterior half of the middle femora dark-brown and somewhat incrassated; the posterior half thin and dark-yellow; the greater part of the under side sparsely spinose; middle tibiæ dark-brown, in most specimens, gradually becoming yellow towards the tip; tarsi yellowish, brownish towards the tip. Hind femora not incrassated, whitish, the last third brownish-yellow, brownish towards the tip; both shades separated by an oblique brownish-black ring; hind tibiæ and tarsi as in the intermediate pair of feet. Wings very much attenuated towards the basis, hyaline; their anal angle not projecting at all; from the tip of the costal cell a narrow black streak extends over the incrassated point, where the third longitudinal vein originates and over the crossveins, closing the little cells at the basis of the wing; the apex of the wing is occupied by a large black spot,

which runs from the anterior to the posterior margin, but is very much dilated beyond the fourth longitudinal vein. The second longitudinal vein is gently curved forward and ends some distance from the tip; the small crossvein is before the middle of the discal cell; the last section of the fourth longitudinal vein is almost parallel to the third vein; the posterior angle of the anal cell is rounded.

Hab. Cuba (Gundlach).

Observation.—Through the kindness of Dr. Gundlach, who sent me the specimens, I have been informed of the identity of this species with the one described by Bigot. I have not succeeded yet in comparing De la Sagra's work, which contains the description, and I draw the attention of those, to whom this work is accessible, to the fact, that among the Cuban species described by me, one or the other may have been previously described by Mr. Bigot in that volume.

Gen. V. **SYNTACES** nov. gen.

Charact.—*Front* moderately broad, broader above (according to Macquart's statement, his figure, on the contrary, shows a front narrower above).

Antennal arista pubescent.

Feet slender; all the femora thin and all armed.

Wings: posterior angle of the anal cell rectangular; crossveins not approximated; the last section of the fourth longitudinal vein only moderately convergent with the third.

The typical species is *Setellia apicalis* from Brazil, described by Macquart (Dipt. Exot. II, 3, p. 249). As I have not seen this species, I have borrowed the generic characters from that author's description and figure, which gives these characters a somewhat uncertain basis. The close relationship to the next following genus is, in my opinion, evident; still, it does not seem advisable to unite them, as, in the present genus, the front femora are weaker and armed with less conspicuous spines; as the hind feet are much less elongated in comparison to the front feet; as the posterior angle of the anal cell is not obtuse, but rectangular, and as the second longitudinal vein has no stump of a vein upon it; nevertheless it is not impossible that the examination of a specimen would lead to a different conclusion from that which seems warranted by Macquart's description.

Gen. VI. **EUOLENA** nov. gen.

Charact.—*Front* very broad, very little narrowed anteriorly; the excavation of its upper part very shallow; the ocelli near the vertex and closely approximated to each other.

Antennal arista with a very short pubescence.

No mesothoracic bristle, and, as it seems to me, no prothoracic one.

Scutellum with four bristles; the lateral ones weak and small.

Feet: front femora rather strong, with conspicuously long and strong spines; the four posterior feet remarkably long and slender, their femora with small spines near the tip only, otherwise these femora are thin and very long, especially the intermediate ones.

Wings: posterior angle of the anal cell obtuse; the small crossvein a little beyond the last third of the discal cell; opposite this crossvein, the second longitudinal vein emits a little stump of a vein into the submarginal cell; the last section of the fourth longitudinal vein is nearly parallel to the third.

The typical species is *Michogaster egregius*, from Columbia, described by Gerstæcker (Stett. Ent. Z. XXI, p. 179). I possess the male only. The ovipositor of the female is called sugar-loaf shaped by the author; which would indicate that it is less compressed than in the other *Richardina*; it may be somewhat of the same shape as in *Stenomacra Guérini*.

Gen. VII. **IDIOTYPA** nov. gen.

Charact.—*Front* very broad, not narrowed anteriorly; ocelli rather approximated to the edge of the vertex, and placed close to each other.

Antennal arista with a short pubescence.

No mesothoracic bristle; a weak prothoracic one.

Scutellum with two bristles; *metathorax* sloping.

Abdomen slender and elongated, almost pedunculate at the basis.

All the *femora* strong and armed with spines.

Wings: posterior angle of the anal cell quite obtuse; the small crossvein beyond the last third of the discal cell; opposite this crossvein the second vein has a stump of a vein, inside of the submarginal cell, and a second one on the opposite side, in the marginal cell, nearer to the apex of the wing; the last section of the fourth longitudinal vein almost parallel to the third.

1. *I. appendiculata* n. sp. ♂ ♀.—(Tab. IX, f. 26.) Ex ochraceo ferruginea, thorace flavo-vario, alarum dimidio anteriore ex ochraceo ferrugineo, posteriore subhyalino, dilute lutescente.

Yellowish-ferruginous, with the thorax marked with yellow; the anterior half of the wings ochre-brownish, the posterior half almost hyaline, yellowish. Long corp. 0.44; ♀ cum terebrâ 0.52; long. al. 0.4—0.41.

Of this species I possess a very well preserved, and, as it seems, particularly fully-colored female, and two much paler males, probably having faded through long exposure. This difference in coloring notwithstanding, I have not the least doubt that both sexes belong to the same species. The condition of the specimens induces me, however, to begin with the description of the female and to add afterwards those characters by which the male specimens differ from it.

Female.—Head rather dark-yellow, of the ordinary *Dacus*-like shape; the front of considerable, and altogether equal, breadth; occipital bristles rather strong; the lateral bristles in front of them are wanting; likewise the bristles generally inserted near the ocelli; the ocelli are approximated to the edge of the vertex and close to each other; a black, biarcuate band runs from the orbit of the eye on one side to that on the other, across the ocelli; immediately above the antennæ there is another black band, not reaching the orbits, the upper limit of which forms a less arcuate, the lower limit a more arcuate curve. In consequence of the very approximated position of the antennæ, the frontal lunule is more isolated from the face, than is the case in any other of the *Ortalidæ* I am acquainted with. Antennæ brownish ochraceous-yellow; the third joint comparatively long; the arista with a short, but very distinct, pubescence. The lower corners of the central portion of the face rather blackish. The short, but rather broad palpi ochraceous-yellow, brownish-black at the basis. The occiput shows, not far from the edge of the vertex, a narrow, black cross-band, not quite reaching the orbit of the eye. The thorax shows a very variegated picture; the very broad middle stripe, running from end to end, is of a brownish-ferruginous color, which changes into black towards its posterior third; this stripe is divided in two by a blackish, rather indistinct longitudinal line; it is separated from the lateral stripes by a longitudinal line of ochraceous-yellow pollen; the broad lateral stripes are crossed by the transverse suture, which is covered with pale ochraceous-yellowish pollen; the anterior portion of the lateral stripes is black and leaves exposed only the pale yellow humeral stripe; the posterior portion of the lateral stripe is black on the side turned towards

the middle stripe, otherwise brownish-ferruginous. Scutellum short, with two bristles, pale-yellow. Pleuræ black; the humeral region, including the prothoracic stigma and a broad band, running from the root of the wing to the interval between the fore and middle coxæ, pale-yellow; the suture, lying in this band and running down from the root of the wing, is margined with brownish-black. Metathorax black, separated from the pleuræ by a broad yellow stripe. The first abdominal segment rather long, very slender, considerably incrassated, however, towards its end, so that here it equals in breadth the following segment; its first third is black, the second pale-yellow, the remainder, as well as the remaining portion of the abdomen, yellowish-ferruginous, almost ochre-brownish, and beset with a short pubescence of the same coloring. Ovipositor of the color of the abdomen; quite flat; the first segment not quite so long as the last three abdominal segments taken together; rather narrow towards its end. Coxæ brownish-black; the second joint of the front coxæ, the tip of the first joint and the second joint of the middle ones, yellow. All the femora beset with spines, not incrassated, but strong, black, yellow to a small extent at the basis only, yellowish-red to a considerable extent towards the end. Front tibiæ reddish-yellow; the four posterior ones of a purer yellow with reddish-yellow tips. All the tarsi yellowish-red; the front tarsi from the second joint and beyond dark-brown; the other tarsi infuscated at the tip only. The hairs on the feet are very short, and of the same color as the ground upon which they are inserted. Wings comparatively long and narrow, with ferruginous veins; the anterior half has a yellowish rusty-brownish tinge, which is more ferruginous-yellow towards the basis, and more brownish towards the apex; the posterior limit of this coloring is almost rectilinear and reaches the fourth longitudinal vein at its root and at its tip only. The whole posterior half of the wing has a decidedly yellowish tinge, but is rather transparent. The second longitudinal vein is rather straight, gently bent forward towards its end only; it reaches the margin not far from the apex of the wing; two conspicuous stumps of veins project from it not far from each other; both are perpendicular, but placed at the opposite sides of the principal vein; one is just opposite the small crossvein, the other somewhat nearer to the apex of the wing; the small crossvein itself is a little beyond the last third

of the discal cell; the last section of the fourth longitudinal vein is almost parallel to the third vein; the posterior angle of the anal cell is quite obtuse.

Males.—The two specimens which I have before me differ from the females by the absence of the upper black crossband on the front, of the black crossband of the occiput and of the spots on the face which have a black coloring; all which in the female is described as black or blackish-brown, is of a dingy rusty-brown in the male. As, at the same time, the contrast between the yellow and the ferruginous regions is less striking, this gives these specimens a less variegated appearance than that of the above-described female. The first abdominal segment is just as narrow as in the female; but this is less apparent here, as the posterior part of the abdomen is less broad.

Hab. Cuba (Gundlach).

Gen. VIII. **STENERETMA** nov. gen.

Charact.—*Front* very broad, not attenuated anteriorly; *occiput* very convex; *cheeks* broad; *ocelli* small and rather approximate to each other.

Arista thin and bare.

A strong mesothoracic bristle; no prothoracic one.

Scutellum with two bristles; metathorax sloping.

Abdomen slender and elongate, attenuate towards the basis.

Femora of medium strength, all unarmed.

Wings but little developed, short and exceedingly narrow, attenuate in the shape of a wedge towards the basis, so that their surface beyond the fifth longitudinal vein is nothing but a narrow, veinless strip; the auxiliary vein so closely approximated to the first longitudinal vein, that they can be distinctly told apart at their end only; the two ordinary crossveins approximate to each other; the small one lies but little beyond the middle of the wing; second basal cell very small and narrow; the anal cell and the sixth longitudinal vein are wanting, with the exception of a rudiment of the latter, which does not reach beyond the axillary incision.

As the group of the *Ulidina* contains the genera with a more developed anal cell, the group of the *Richardina* on the contrary those with a less developed one, there can be no doubt that the present genus, in the incompletely developed wings of which the anal cell is altogether wanting, belongs to the *Richardina*; and that this is its true location is proved by its relationship to

Idiotypa, especially evident in the structure of the abdomen. Among the differences of these two genera I will only mention that the structure of the head of *Idiotypa* is not unlike that of *Dacus*, while the head of *Steneretma* resembles that of *Tritoxa*. As *Steneretma* and *Tritoxa* also agree in the presence of a mesothoracic bristle and in the absence of a prothoracic one, the former genus, if its first longitudinal vein showed a distinct pubescence, would have to be placed next to *Tritoxa*.

1. *S. laticauda* n. sp. ♀.—Lutea, segmentis abdominalibus singulis postice anguste et æqualiter fusco-marginatis, tarsis præter basim nigrofuscis, alis luteo cinereis, albido-bitasciatis.

Dark-yellow, the single abdominal segments on their posterior margin with a narrow infuscated border; the tarsi, with the exception of the basis, blackish-brown; wings yellowish-gray with two whitish cross-bands. Long. corp. 0.14; cum terebrâ 0.19; long. al. 0.11—0.12.

Of a dark-yellow color, shining. The broad, rather convex front bears, besides the long bristles on the vertex and in the region of the ocelli, a moderate quantity of rather long black hairs; the comparatively strong convexity of the occiput almost obliterates the usual edge between it and the vertex. The antennæ are of the same color as the rest of the body, and of more than half the length of the face; their third joint elongate, rounded at the tip; the thin and bare arista is very long. Clypeus, palpi, and proboscis likewise partake of the general coloring of the body. Thorax but little elevated and rather narrow in comparison to its length; its dorsum on the sides and on its posterior border with a few rather long black bristles; upon the remainder of its surface only with a short, black pubescence. Scutellum small, bare, with the exception of the two bristles upon its end. Plenræ glabrous; besides the mesothoracic bristle they bear only a single bristle not far below the root of the wing. The abdomen is narrow and elongate, attenuate towards the basis, not so much, however, as in the females of *Idiotypa appendiculata*; its segments have, on the posterior margin, a narrow border of equal breadth and of a brown or reddish-brown color; upon the last segment this margin becomes indistinct, or it is altogether wanting. The blackish pubescence of the abdomen is everywhere very short and not conspicuous. The ovipositor is of the same color as the remainder of the body and is strikingly

broad; its first joint is about as long as the last three abdominal segments taken together; from its basis to the middle it is exactly as broad as the abdomen itself; beyond the middle it is but little attenuate, so that the truncature at the end has a considerable breadth; the second and third joints of the ovipositor are also rather broad; the latter does not end in a sharp point, but in a narrow truncature. Feet bare, their structure ordinary; femora unarmed; the tarsi blackish-brown from about the tip of the first joint. The yellowish-gray wings have two perpendicular whitish crossbands; the first passes between the two ordinary crossveins from the anterior to the posterior margin of the wing; the second lies between the first and the apex of the wing, but much nearer the latter, is obliterated in the marginal cell and does not entirely reach the posterior margin; besides these two whitish crossbands there is, at the end of the second basal cell and in the adjoining region of the first basal cell a small, whitish spot; the coloring of the wing, on this side of the first crossband, towards the root of the wing, changes gradually into clay-yellow, while beyond the second crossband the color is almost blackish-gray; the posterior crossvein shows the trace of a delicate blackish-gray veining, while there is no such trace on the small crossvein.

Hab. Texas (Belfrage).

Gen. IX. **CCELOMETOPIA** MACQ.

Charact.—*Front* of moderate breadth, slightly narrowed anteriorly, somewhat excavated; ocelli far removed from the edge of the vertex, placed close to each other on a more or less projecting bump.

Antennal arista with a very short pubescence.

No mesothoracic and one prothoracic bristle.

Scutellum with four bristles; metathorax somewhat sloping.

Femora not incrassate, nevertheless strong, the four posterior ones considerably longer than the front pair; all are provided with spines, the fore femora, however, with a few small ones towards the tip only.

Wings: posterior angle of the anal cell quite obtuse; the crossveins not approximate to each other; the last section of the fourth longitudinal vein converges towards the third.

With *Ccelometopia* a series of genera begins which have a comparatively short, oval abdomen, not very attenuate at the basis. The type of the genus is *C. trimaculata* Fab. = *C. ferruginea* Macq. from South America, which Wiedemann placed in the genus *Trypeta*.

1. *C. bimaculata* n. sp. ♂.—(Tab. IX, f. 27.) Rufa, abdomine chalybeo vel violaceo, pedibus flavis; tibia tamen tarsorumque apice fuscis; alæ hyalinæ, nigro-bimaculatæ.

Ferruginous-reddish, the abdomen steel-blue or violet; feet yellow; tibiæ and tip of the tarsi brown; wings hyaline with two black spots. Leng. corp. 0.22—0.26; long. al. 0.21—0.22.

Head and thorax ferruginous-red, rather shining; only the hind coxæ sometimes pitch-brown. Front of very moderate breadth; narrower anteriorly, somewhat excavated; the bristles on the vertex, the very much advanced lateral bristles and the two bristles near the ocelli black and rather strong. The ocelli are placed close to each other on a flattened elevation, almost in the middle of the front; the frontal humle is rather isolated from the face, in consequence of the very approximate position of the antennæ. The third antennal joint is sometimes more brownish-red towards the tip; arista with a short pubescence. The short hairs on the thoracic dorsum are whitish, and hence easily perceptible; the ordinary bristles are black or brown, sometimes only brownish; a blackish line in the middle is only occasionally perceptible. Scutellum convex, with four brownish or brown bristles. Abdomen metallic steel-blue, shining, with more or less extensive and vivid violet reflections; sometimes ferruginous-brownish at the extreme basis; its almost whitish pubescence appears much darker, when looked at against the light. Femora yellowish, usually brownish at the tip; the foremost ones strong, with a few weak and small spines on the under side, near the tip only; the four posterior femora much longer, also strong, with spines on the under side. Tibiæ brown. Tarsi of a dirty-yellowish brown from about the tip of the second joint. Wings pure hyaline, with a rather sparse and coarse microscopic pubescence and with black veins; the black stigma is confluent with a moderately large, sharply limited spot, reaching as far as the third longitudinal vein; a larger, almost triangular black spot occupies the apex of the wing; it begins before the second longitudinal vein and ends midway between the third and fourth veins; moreover, in the environs of the humeral crossvein, there is a grayish-black spot, which is easily overlooked. The third longitudinal vein is very straight; the small crossvein lies in the middle of the comparatively short discal cell. The anterior basal cell is somewhat expanded at the expense of the discal cell, so

that the latter is much narrower before the small crossvein than beyond it; posterior crossvein straight, somewhat oblique; the last section of the fourth longitudinal vein strikingly long, distinctly converging towards the third longitudinal vein; posterior angle of the anal cell very obtuse.

Hab. Cuba (Gundlach).

Gen. X. **HEMIXANTHIA** nov. gen.

Charact.—*Front* of medium breadth, somewhat narrower anteriorly, not excavated; the posterior ocelli not very far from the edge of the vertex; the anterior one removed to about the middle of the front.

Antennal arista with a distinct pubescence.

A small prothoracic, and, as it seems, no mesothoracic bristle.

Scutellum with four bristles; *metathorax* perpendicular.

Femora not incrassate, but rather strong; the posterior ones longer than the foremost ones; all are beset with spines; the spines of the foremost ones are but very few.

Wings: posterior angle of the anal cell obtuse; crossveins conspicuously approximate; the last section of the fourth longitudinal vein is parallel to the third.

The difference from *Cœlometopia* consists principally in the peculiar position of the ocelli, the remarkably approximate crossveins and the parallelism of the third and fourth longitudinal veins.

I do not know of any described species of this genus and for this reason give the following:—

1. *H. spinipes* n. sp. ♀.—(Tab. IX, f. 28.) Lutea, metanoto epimerisque metathoracis nigris, abdomine chalybeo, violaceo-splendente; alæ subhyalinæ, apice fascisque tribus fusco-nigris; harum secundâ postice, tertiâ antice, abbreviatâ.

Clay-yellow, metanotum and epimera of the metathorax black, abdomen steel-blue, with a violet reflection; wings rather hyaline, the apex and three crossbands brownish-black; the second of these abbreviated posteriorly, the third anteriorly. Long. corp. 0.24; long. al. 0.23.

Clay-yellow, thoracic dorsum more yellowish-red. Front of medium breadth, but little narrower anteriorly, not excavated, with but a small depression on the vertex; the two posterior ocelli are placed upon a very small black spot, at a moderate distance from the vertex and close to each other; the anterior ocellus is quite unusually distant from them, and placed about

the middle of the front; the bristles on the vertex, the rather distant lateral bristles and the two ocellar bristles comparatively long and strong, black. Antennæ reaching down to the border of the mouth; the comparatively long third joint sometimes somewhat infuscated at the tip. Arista pubescent. The pubescence of the thoracic dorsum is pale-yellowish, the ordinary bristles black. Scutellum of a pure yellow, with four black bristles; its surface rather even. The middle portion of the mesonotum, the lower portion of its sides and the epimera of the metathorax brownish-black. The pubescence of the pleuræ yellowish. Abdomen elongate-oval, clay-yellow at the extreme basis, the remainder shining steel-blue with violet reflections, more greenish-blue at the posterior end. The first segment of the ovipositor large, shining black, concave above, somewhat convex below. Feet clay-yellow, the basis of the middle tibiæ and the hind tibiæ brown; the tip of the tarsi but little infuscated; femora not incrassate, although rather strong, the four posterior ones longer than the two foremost ones; the latter with a few small spines near the tip only, the former beset with spines on the whole second half of the under side. Wings almost hyaline, with a yellowish-gray tinge, which is more yellow towards the anterior border; costal cell yellowish-brown; a narrow brownish-black band runs from the humeral crossvein to the axillary incision; a second one, somewhat broader, runs from the anterior margin over the basis of the submarginal cell and over the end of the small basal cells nearly, but not quite, to the posterior margin of the wing; a third band, inclosing the two remarkably approximate crossveins, extends from the posterior margin to the middle of the submarginal cell; the apex of the wing bears a large elongate brownish-black spot, beginning before the second longitudinal vein and occupying the border of the wing as far as beyond the fourth vein. The last section of the fourth longitudinal vein is parallel to the third vein; the posterior angle of the anal cell is obtuse; the microscopic pubescence of the surface of the wing is remarkably coarse and sparse.

Hab. Brazil.

Gen. XI. **MELANOLOMA** nov. gen.

Charact.—*Front* rather broad, somewhat narrower anteriorly, not excavated; the posterior ocelli not far removed from the edge of the vertex; the anterior one at a considerable distance from them.

Antennal arista bare.

A strong mesothoracic bristle and a very weak prothoracic one.

Scutellum with four bristles; *metathorax* rather perpendicular.

Femora not incrassate, only the hindmost ones with spines near the tip.

Wings: posterior angle of the anal cell quite obtuse; the crossveins not approximate; the last section of the fourth longitudinal vein parallel to the third.

The species of this genus are distinguished by their robust thorax and short oval abdomen; the surface of the latter is not smooth, but entirely covered by shallow scars, almost chagreened. The picture of the wings of the species known to me consists of a black border of the anterior margin of the wing and of the apex, and of a narrow black streak over the small crossvein.

The typical species is a Brazilian one, described by Wiedemann as *Trypeta cyanogaster*. As, in Wiedemann's description, the plastic characters are not sufficiently taken notice of, I will give the description of a species closely related to his.

1. *M. affinis* n. sp. ♂.—(Tab. IX, f. 29.) Rufa, tibiis concoloribus, posticis tamen basim versus infuscatis, abdomine ex violaceo chalybeo; alæ hyalinæ, costâ cum apice et venâ transversâ mediâ anguste nigro-imbatis.

Red, the tibiæ of the same color, the hindmost ones infuscated towards the basis; abdomen violet steel-blue; wings hyaline, anterior margin and apex, as well as the small crossvein, with a narrow black border. Long. corp. 0.24; long. al. 0.24.

Ferruginous-red, shining; abdomen of a dark steel-blue color, somewhat verging on violet. Front rather broad, somewhat narrower anteriorly, sometimes tinged with yellow on the sides; the short and thin hairs upon it are inserted in small, very shallow, and hence hardly perceptible pits. The two superior ocelli are quite near the vertex; the anterior one is quite a distance from them, but still above the middle of the front; bristles of the vertex, the lateral ones and the two bristles near the ocelli, are present. Antennæ reaching a little beyond the border of the mouth; the third joint long, sometimes more reddish-brown.

Arista thin and apparently bare. Thorax strongly built; the fallow-yellowish pubescence of its dorsum very short; the ordinary bristles black. Scutellum convex, with four bristles. The perpendicular mesonotum, the pleuræ and the pectus of the same color as the upper side of the thorax. The mesothoracic bristle strong, black, and hence very conspicuous; the prothoracic bristle thin and fallow-yellowish, and hence easily overlooked. The metallic-blue abdomen is of a rounded-oval shape and is covered with shallow scars, which diminish its lustre; its short pubescence is whitish on the first segment only, otherwise rather blackish. Feet of a yellowish-ferruginous color, only the distinctly arcuate hind tibiæ are gradually infuscated towards the basis; the tarsi, beyond the second joint, are more or less ferruginous-brownish. Femora not incrassate, only the hindmost ones with spines near the tip. Wings hyaline; the costal cell, the stigma, and a narrow border, running from it to the fourth longitudinal vein, along the margin of the wing, black; the small crossvein likewise with a narrow black cloud; a blackish spot lies between the extreme basis of the submarginal cell and the end of the costal cell. The second longitudinal vein reaches the anterior margin rather far from the apex of the wing; the third longitudinal vein is very straight; the small crossvein is a little beyond the middle of the discal cell, which is considerably narrower before this crossvein than after it; posterior crossvein straight, a little oblique; the last section of the fourth longitudinal vein rather long, parallel to the third vein; posterior angle of the anal cell quite obtuse. The microscopic pubescence of the surface of the wing is comparatively sparse and coarse.

Hab. Brazil.

Observation.—*M. cyanogaster* Wied. is not quite as large as the above-described species; its wings are comparatively smaller and the black border along the costa is somewhat broader at the apex of the wing; the lateral bristle of the front is somewhat more removed from the bristles on the vertex; the shallow pits on the front are not perceptible; the pubescence of the thoracic dorsum is considerably longer; the pleuræ and the tibiæ are blackish-brown.

Gen. XII. **EPIPLATEA** LOEW.

Charact.—*Front* broad, narrower anteriorly; not projecting in profile; rather densely hairy upon the whole surface.

Antennæ of medium size; third joint oval, with a thin, bare arista.

Face vertical, with a depression under each antenna; longitudinally convex between these depressions; *clypeus* of a moderate transverse diameter, projecting considerably beyond the anterior edge of the mouth, which is drawn upwards; *proboscis* stout.

Thorax with bristles on its hind part only; *scutellum* convex, with four bristles.

Femora of moderate length, strong, but not incrassate; all unarmed.

Wings comparatively short; submarginal and first posterior cells broad; third longitudinal vein bent backwards towards its end; the last section of the fourth longitudinal vein does not converge towards the third; posterior crossvein perpendicular; the posterior angle of the anal cell rather acute.

The species of this genus are rather stout, not metallic, except sometimes on the abdomen. The structure of the head recalls that of some *Sciomyzidæ*, and is very like that of the two well-known species, described by Wiedemann as *Ortalis trifasciata* and *atomaria*; in their general appearance, the species of *Epiplatea* are also not unlike the two latter species, but are easily distinguished by the first longitudinal vein being bare, by the posterior angle of the anal cell not being rounded as in those species and by the absence of the erect bristle before the end of the upper side of the tibiæ, a bristle which is always present in the latter species.

1. E. erosa LOEW. ♀.—(Tab. IX, f. 24.) Fusco-testaceo vel ex ferrugine fusca, pedibus concoloribus; abdomine nigro, alis hyalinis, fasciis duabus et puncto centrali nigris.

Brownish-yellow or ferruginous-brown, with the feet of the same color and a black abdomen; wings hyaline, with two brown crossbands and in the middle with a brown dot. Long. corp. 0.17; long. al. 0.16.

SYN. *Epiplatea erosa* LOEW, Berl. Ent. Zeitschr. XI, p. 325, Tab. II, f. 25.

The coloring of the lighter shaded specimens is yellow-brownish, in darker specimens it becomes ferruginous-brown. Head of the same color. Front broad, considerably narrowed anteriorly, upon its whole surface uniformly and rather densely clothed with an erect, black pubescence; along the lateral margin with a narrow border of white pollen; the stripes running down from

the vertex along the sides of the front and the ocellar triangle are of the same color as the front and hence indistinct. Antennæ not reaching quite to the edge of the mouth; the first two joints of the color of the head, or a little lighter; the oval third joint dark-brown, often quite black; the arista thin and bare. Face excavated under each antenna, longitudinally convex between these depressions; descending vertically in profile; the anterior edge of the mouth is strongly drawn upwards, so that the clypeus projects considerably above it. Proboscis stout; palpi brown, generally paler towards the tip. The thoracic dorsum generally has, on the posterior side, an almost silvery-white transverse crossband, and before the transverse suture, on each side, a large spot of a similar pollen; these pollinose spots are very distinct, when seen by reflected light, but can easily be overlooked in any other light. Upon the pleuræ likewise there are two spots of white pollen; one of them lies over the fore coxæ, the other immediately under the longitudinal suture of the pleuræ, where the color is generally darker-brown. The front part of the coxæ is likewise covered with a white pollen, which, however, sometimes is entirely invisible. Abdomen black, somewhat glossy, generally brown at the basis, with a rather coarse pubescence, which is longer and black on the posterior margins of the segments. The flattened ovipositor is somewhat attenuate, its first two segments black, the third orange-yellow. Feet of the same color as the body; tibiæ and tarsi darker brown, in fully colored individuals brownish-black. Halteres yellowish. Wings of very moderate length, rather broad, hyaline, with brown veins; the basis of the wings as far as the humeral crossvein and the anal cell are brownish; a narrow brownish-black band begins at the costa, where it is confluent with the small black stigma and a black spot, lying at the end of the costal cell; it runs over the bases of the submarginal, discal, and third posterior cells, as far as the sixth longitudinal vein, which its end alone crosses a little; before the apex of the wing there is a broader crossband, which is sinuate on both sides, weaker, however, on the inside than on the outside; posteriorly it bifurcates in two short, obtuse branches, the inner one of which reaches the margin of the wing and covers the perpendicular posterior crossvein; the outside one is shorter and ends in the second posterior cell, some distance from the margin of

the wing; between these two crossbands is the black spot, formed by a cloud over the small crossvein; the stigma is small; the small crossvein is beyond the middle of the discal cell; the submarginal and first posterior cells are broad; the end of the third longitudinal vein is gently curved posteriorly and ends exactly in the apex of the wing; the last section of the fourth longitudinal vein does not converge towards the third; the anal cell is comparatively rather small; the crossvein, closing it, is a little arcuate, but forms nevertheless a rather acute posterior angle.

Hab. Cuba (Gundlach).

formed
 ll; the
 ic sub-
 e third
 etly in
 tudinal
 is com-
 a little
 angle.

APPENDIX,

CONTAINING THE DESCRIPTIONS OF THE SPECIES PUBLISHED BY PREVIOUS WRITERS, AND NOT IDENTIFIED BY THE AUTHOR.

1. *Say, Journ. Acad. Nat. Sciences Phil., Vol. VI, Part II.*

Page 83. ***Ortalis ligata.***

Wings quadrifasciate with fuscous.

Inhabits Mexico.

Body blackish; *head* ferruginous, tinged with glaucous behind and on the vertex; *thorax* blackish-plumbeous; *wings* white, subopaque, with four fuscous bands; the first a little oblique, across the neck of the wing; second from the tips of the mediastinal and post costal nervures, and proceeding a little obliquely, so as to be bounded posteriorly by the middle cross-nervure; third, perpendicular to the costal margin and covering the posterior cross-nervure; fourth, terminal, slightly connected on the costal edge with the third; *poisers* white; *tergum* coppery-black; *feet* black; *knees* and *tarsi* ferruginous. Length three-twentieths of an inch.

[Belongs very probably to the genus *Rivellia*, but it will be difficult to decide to which species, on account of the great similitude between the species of that genus.—*Loew.*]

2. *Rob. Desvoidy, Myodaires.*

Page 715. ***Meckelia philadelphica.***

Minor *M. eleganti*; *pedes fulvi*, *tibiis nigricantibus*; *alæ flavescentes*, *unicâ maculâ subfusca*.

Plus petite que la *Meckelia elegans*; *frontaux*, *antennes*, *face*, *rouges*; *optiques* d'un gris rougeâtre; *corselet* d'un brun-gris;

(197)

abdomen un peu moins gris et d'un noir plus luisant; cuisses fauves; tibias mélangés de noir et de fauve; tarses noirs; ailes flavescences, n'offrant que l'apparence d'une seule macule.

Originaire de Philadelphie.

(Translation.)—Smaller than *Meckelia elegans*; frontal bristles, antennæ, face, red; optical bristles of a reddish-gray; thorax brownish-gray; abdomen a little less gray and of a more shining black; femora fulvous; tibiae mixed with black and fulvous; tarsi black; wings flavescence, with the appearance of a single spot.

From Philadelphia.

[It seems hardly doubtful that this species belongs to the *Ortalina*; it is probably either an *Anacampta* or a *Ceroxys*, as Rob. Devoidy's genus *Meckelia* has the third antennal joint excised on the upper side and ending in a very sharp angle.—*Loew*.]

3. Walker, *Insecta Saundersiana*.

Page 373. ***Ortalis basalis***, Mas. et Fœm.

Nigro-cyanea, caput fulvum; antennæ luteæ; abdomen basi ferrugineum, *fœm.* apice luteum attenuatum; pedes fulvi; alæ hyalinæ, basi fulvæ, vitta antica interrupta fusca.

Ceroxys? Blackish-blue: head tawny; face with a whitish covering; epistoma prominent; mouth pitchy; feelers luteous; third joint much deeper than the second and more than twice its length; sixth black, bare, very slender, more than twice the length of the third; abdomen longer than the chest, ferruginous towards the base; abdomen of the female pale luteous towards the tip, which is much attenuated; legs tawny; wings colorless, slightly tawny at the base, adorned along the fore border with a dark-brown interrupted stripe, which is widened at the tip; veins black; fifth vein converging towards the tip of the fourth; sixth not reaching the hind border; crossveins straight, almost upright; poisers pitchy. Length of the body $1\frac{1}{2}$ —2 lines; of the wings 2—3 lines. United States.

[It is utterly improbable that this species should be a *Ceroxys*, as Mr. Walker supposes; his description rather suggests that it belongs to the *Ulidina*.—*Loew*.]

4. *Macquart, Dipt. Exot. II, III, Tab. XXIX. fig. 3.*Page 208. ***Herina mexicana.***

Viridi-cyanea. Alis limbo externo nervisque transversis fuscis.

Long. 4 lin.—Face testacée. Front noir; vertex et derrière de la tête testacés. Antennes brunes; style fauve. Thorax d'un vert brillant, à reflets bleus. Abdomen manque. Pieds noirs. Ailes jaunâtres jusqu'à l'extrémité; cellules basilaires brunes; nervures transversales bordées de brun; première oblique.

Du Mexique.

(Translation.)—Length 4 lines. Face testaceous; front black; vertex and occiput testaceous. Antennæ brown; arista fulvous. Thorax of a brilliant green, with blue reflexions. Abdomen—(wanting). Feet black. Wings yellowish, anterior margin brown from the stigmatal cell, inclusively, as far as the apex; basal cells brown; crossveins bordered with brown; the first of them oblique.

Mexico.

[Macquart very improperly placed this species in the genus *Herina*; it is a perfectly normal species of his own genus *Stenopterina*.—Loew.]

5. *Walker, List of Dipt. Ins. IV.*Page 992. ***Ortalis massyla*, n. sp., Fem.**

Viridis, capite ferrugineo, abdominis segmento quinto purpureo apice fulvo, palpis ferrugineis, antennis pedibusque nigris, tarsis fulvis, alis albis fusco trifasciatis.

Body metallic-green, slender, clothed with short black hairs: head and chest beset with black bristles: head ferruginous above and along the borders of the eyes; epistoma ferruginous, prominent, eyes red; fore part slightly convex; its facets a little larger than those elsewhere: sucker black, clothed with tawny hairs; palpi ferruginous; beset with black bristles: feelers black, much shorter than the face; third joint conical, ferruginous at the base, much longer than the second; bristle bare, very slender, more than thrice the length of the third joint; abdomen long-obconical, much longer than the chest, tapering, flat, and with a vein on each side towards the tip, which is tawny; fifth segment dark-purple: legs black, clothed with short black hairs; knees ferruginous; feet and tips of shanks dull tawny: wings white, with

three dark-brown bands; the first extends nearly to the hind border, and joins the side of the middle crossvein; the second reaches the hind border and incloses the lower crossvein; it is darkest on the fore border, and there unites with the third, which widens along the fore border and occupies the whole of the tip of the wing; wing-ribs, veins, and posers tawny; veins pitchy in the brown parts of the wings; lower crossvein nearly straight. Length of the body $1\frac{3}{4}$ —2 lines; of the wings 3—4 lines.

North America.

[This seems to be an *Euxesta*.—Loew.]

6. Walker, *List of Dipt. Ins. IV.*

Page 995. **Ortalis? diopsides**, BARNSTON'S MSS. Fem.

Nigra, obscura, capite antico fulvo, palpis antennis pedibusque piceo-ferrugineis, alis subcinereis ad costam fusco bimaculatis.

Body dull-black, clothed with very short black hairs: head beset with a few black bristles, tawny in front and beneath, where it is covered with white bloom; sides of the face without bristles; epistoma slightly prominent; eyes dark-red; facets of the fore part a little larger than those elsewhere: sucker and palpi ferruginous, partly pitchy; sucker clothed with tawny hairs; palpi beset with black bristles; feelers ferruginous, shorter than the face; third joint pitchy above, nearly round, longer than the second joint; bristle black, bare, slender, much more than twice the length of the third joint; abdomen spindle-shaped, much longer than the chest; last segment flat: legs pitchy, mostly ferruginous beneath, clothed with very short black hairs; claws black: wings slightly gray, with a narrow pitchy band at half the length of the fore border, on which, near the tip, there is a small brown spot; wing-ribs tawny; veins black, tawny at the base; longitudinal veins straight; lower crossvein straight, slightly oblique, nearly twice its length distant from the middle crossvein; posers pale tawny. Length of the body 2 lines; of the wings $2\frac{1}{2}$ lines.

St. Martin's Falls, Albany River, Hudson's Bay.

[This species seems likewise to belong to the *Ulidina*, a group which is so abundantly represented in America.—Loew.]

7. Walker, *List of Dipt. Ins. IV.*Page 995. **Ortalis? costalis**, n. s., Fem.

Nigra, abdomine nigro-aeneo, pedibus nigris, alis limpidis ad costam fusco bimaculatis, stigmatè nigro.

Head wanting: chest dull black, beset with a very few black bristles: abdomen sessile, brassy-black, shining, slightly spindle-shaped, much longer but hardly broader than the chest: legs black, clothed with very short black hairs: wings colorless, with a small brown spot just above the tip, and another at the base of the fore border, where the vein is thickened; a black band along the middle of the fore border; wing-ribs and veins black; third longitudinal vein straight, with the exception of a very slight angle at its junction with the lower crossvein, which has two very slight curves, the upper inward, the lower outward. Length of the body $1\frac{3}{4}$ line; of the wings $3\frac{1}{2}$ lines.

St. Martin's Falls, Albany River, Hudson's Bay.

[In this description, after the words "third longitudinal vein straight," something seems to be wanting, as this vein does not at all meet the posterior crossvein. The species very likely also belongs to the *Ulidina*.—Loew.]

8. Macquart, *Dipt. Exot. Suppl. IV, Tab. XXVI, fig. 17.*Page 289. **Urophora antillarum**.

Viridi-nigra. Fronte testacea, alis fasciis duabus, apiceque fuscis.

Long. $1\frac{1}{2}$ lin. ♂.—Palpes noirs. Face d'un vert noirâtre luisant, à léger duvet blanc sur les côtés. Front testacé; une tache verte sur le vertex. Antennes noirs. Thorax et abdomen d'un vert luisant noirâtre. Pieds noirs; premier article des tarsi testacé. Ailes claires, à base jaunâtre; une première bande passant sur la première nervure transversale, et n'atteignant pas le bord intérieur; la deuxième entière, passant sur la deuxième transversale; extrémité à tache brune, liée à la deuxième bande par le bord extérieur également brun.

Des Antilles.

[Almost undoubtedly an *Ulidina*.—Loew.]

9. *Bigot, Ramon de la Sagra, Hist. fis. d. l. Isla da Cuba.****Ulidia fulvifrons.***

Nigro-picco-nitens, hypostomate nigro; fronte, oculis, antennisque fulvis, occipite brunnea; thorace nigro-nitente; abdomine nigro-picco; pedibus fulvis; anticis, cruribus antice brunnescentibus; tibiis tarsisque brunneis; intermediis posticisque, femoribus basi, brunneis; tibiis postice brunneis; alis hyalinis; costa brunnea, punctoque apicali nigro.—Long. 4 mill.

[This species may belong to the *Ulidina*, but it is not probable that it is a true *Ulidia*. The *Ulidia metallica* Bigot, described in the same place, is not an *Ortalida* at all, but belongs to the *Agromyzidae*, perhaps to the genus *Agromyza*.—Loew.]

10. *Walker, Trans. of the Ent. Soc., Tom. V. 1861.*Page 326. ***Ortalis bipars.***

Nigricante viridis, capite supra antennisque rufis, harum articulo tertio longo lineari, pedibus nigris, alis albis nigro-trifasciatis et apice maculatis, vittis secundâ tertiâque postice obsolete, primâ incompletâ, halteribus pallidis.

Blackish-green: head above and antennæ red; third joint of the antennæ long, linear; wings white, with three slight black bands and an apical spot, first band very incomplete; second and third obsolete hindward; discal transverse vein straight, upright, parted by one-fourth of its length from the border and by much more than its length from the brachial transverse vein, halteres pale.

Length of the body $2\frac{1}{2}$ lines; of the wings 4 lines.

United States.

11. *Walker, Trans. of the Ent. Soc., Tom. V. 1861.*Page 324. ***Bricinnia.***

Corpus longiusculum, sat angustum. Peristoma magnum. Antennarum articulus tertius longus, gracilis, linearis; arista simplex, gracilis. Thorax longus, lateribus compressis. Abdomen longum, subfusiforme, apice attenuatum. Pedes validi. Alæ sat angustæ, venis rectis.

Fœm. Oviductus vaginæ productæ, gracilis.

Body rather long and narrow. Epistoma rather prominent; mouth large; third joint of the antennæ long, slender, linear, extending to the epistoma; arista slender, simple, nearly twice the length of the third joint. Thorax long, compressed on each side. Abdomen long, subfusiform, attenuated towards the tip. Legs stout, moderately long. Wing rather narrow; veins straight.

Female. Abdomen attenuated at the tip. Vagina of the oviduct slender, produced.


***Bricinnia flexivitta* Fom.**

Nigra, capite apud oculos albo, vittâ anticâ albâ, antennis ferrugineis basi fulvis, thorace vittis tribus albidis, pectore purpureo-cyaneo, abdomine eupreo, femoribus posticis basi flavis, tarsis fulvis, alis sub-cinereis, costâ apiceque luridis, vittâ discali angulatâ nigra, venâ discali transversâ vix arcuata.

Female. Black: head white about the eyes and with a whitish facial stripe, which is dilated towards the epistoma; antennæ ferruginous, tawny towards the base; thorax with three whitish stripes; pectus blue, varied with purple; abdomen cupreous; vagina of the oviduct attenuated; hind femora yellow towards the base; tarsi tawny; wings grayish, lurid along the costa and at the tips, and with a blackish stripe which extends from the base to and along the discal transverse vein; the latter is upright and hardly curved, and is parted by four times its length from the border, and by a little less than its length from the præbrachial transverse vein, which is oblique.

Length of the body 5 lines; of the wings 10 lines.

Mexico.



INDEX OF THE ORTALIDÆ.

- Achias, 23, 66
 Acidia, 16, 18
 Aciphoræ, 8, 16
 Aciura ænea v. d. *Wulp*, 170
 Acrosticta, 65, 151
 Acrosticta foveolata *Lw.*, 151
 scrobiculata Lw., 151
 Actora, 22
 Adapsilia, 5, 10, 12, 32, 35
 Adrana, 27
 Agastroides, 27
 Amethysta, 25
 Amethysta, 25
 Amphionephes, 44, 83
 Amphionephes pertusus *Lw.*, 84
 Anacampta, 57, 129
 Anacampta latiuscula *Lw.*, 130
 urticæ Lin., 58
 Apospasmica, 52, 131
 Ardelio, 46
 Ardelio longipennis *Lw.*, 46
 brevicornis Lw., 46
 Automola, 12, 52, 118

 Baccha capitata *Lw.*, 68
 Blainvillia, 16
 Boisduvalia, 16, 18
 Brea, 27
 Bromophila, 35

 Callopietria, 62, 140
 Callopietria annulipes *Macq.*, 62, 141
 Calobatiidæ, 30
 Campigaster, 26
 Camptoneura, 18, 24, 50, 108
 Camptoneura picta *Fbr.*, 27, 109
 Celetor, 41
 Cephalia, 7, 10, 11, 22, 23, 32, 49
 Cephalia femoralis *Wied.*, 23, 47
 myrmecoides Lw., 48, 100
 rufipes Meig., 11, 49
 Cephalina, 48, 102
 Ceroxys, 17, 25, 57, 125

 Ceroxys canus *Lw.*, 128
 cœrulea Macq., 25, 150
 crassipennis Fbr., 57
 obscuricornis Lw., 126
 ochricornis Lw., 126
 quadrifasciata Macq., 44
 similis Lw., 127
 Cestrotus, 15
 Chætopsis, 65, 169
 Chætopsis ænea *Wied.*, 170
 debilis Lw., 172
 Chloria, 63
 Chlorophora, 16, 19
 Chromatomyia, 27, 39
 Chromatomyia laeta *Wik.*, 40
 Chrysomyza, 6, 10, 12, 63
 Chrysomyza demandata *Meig.*, 65
 Cleitamia, 21
 Clidonia, 16, 18, 21
 Clitodoca, 42
 Cœlometopia, 23, 26, 68, 188
 Cœlometopia bimaculata *Lw.*, 189
 ferruginea Wik., 13, 68
 trinniculata Fbr., 66,
 188
 Cœlopa, 22
 Coniceps, 64, 70, 177
 Coniceps niger *Lw.*, 178
 Conopsidea, 10, 23
 Cordylura podagrica *Fbr.*, 14
 Cormocaris, 5, 54
 Cormocaris bucephala *Meig.*, 5, 54
 Cruphiocera, 26
 Cryphiocera, 26
 Cytometopa, 179

 Dacina, 34
 Dacus, 12
 Dacus aculeatus *Fbr.*, 158
 æneus Wied., 46
 bicolor Wied., 12
 brevicornis Fbr., 46
 costalis Fbr., 158

- Dacus flavicornis* *Wied.*, 12, 69
flavus *Fbr.*, 68
fraterculus *Wied.*, 12
macularis *Fbr.*, 47
obtusus *Fbr.*, 148
oleæ *Fbr.*, 28
parallelus *Wied.*, 12
podagricus *Fbr.*, 14, 18
serpentinus *Wied.*, 13
succinctus *Wied.*, 12, 44
Dasymetopa, 65, 145
Dasymetopa luteolenta *Lw.*, 146
Delphinia, 16, 18, 24, 50
Delphinia thoracica *Rob. Desv.*, 109
Diacrita, 27, 51, 111
Diacrita æmula *Lw.*, 114
costalis *Gerst.*, 111
Dichromyia, 13, 16, 24, 35
Dichromyia braziliensis *Rob. Desv.*,
 13, 16
caffra *Macq.*, 35
sanguiniceps *Wied.*, 24,
 35
Dictya connexa *Fbr.*, 25
decora *Fbr.*, 36
externa *Fbr.*, 14
ocellata *Fbr.*, 8
pieta *Fbr.*, 109
Diptera aciphorea, 8, 11
Dorycera, 5, 10, 17, 21, 22, 53
Dorycera graminum *Fbr.*, 53
Drosophilidæ, 24
Duomyia, 27, 41

Earomyia, 8, 10
Elaphomyia, 27
Elaphomyia Wallacei *Saund.*, 27
cervicornis *Saund.*, 27
Elaphromyia, 27
Elassogaster, 27
Empyelocera, 6, 10, 63, 65
Engistoneura, 43
Eniconeura, 24, 42
Eniconeura violacea *Macq.*, 42
Ephydrinidæ, 15
Epicausta, 46
Epicerella, 23
Epidesma, 18, 25, 44
Epidesma fascipennis *Macq.*, 45
Epiplatea, 64, 194
Epiplatea erosa *Lw.*, 194
Euchalcota, 40
Euchalcota decora *Macq.*, 40
Enmetopia, 22, 65, 175
Eumetopia rufipes *Macq.*, 175
varipes *Lw.*, 176
Euolena, 67, 183
Euolena egregia *Gerst.*, 67

Euphara, 65, 150
Euphara coerulea *Lw.*, 150
Euprosopia, 26
Eurina, 21
Euripalpus, 24
Euxesta, 65, 153
Euxesta abdominalis *Lw.*, 163
alternans *Lw.*, 165
annonæ *Fbr.*, 162
binotata *Lw.*, 160
costalis *Fbr.*, 158
eluta *Lw.*, 168
nitiliventris *Lw.*, 157
notata *Wied.*, 156
pusio *Lw.*, 155
quaternaria *Lw.*, 159
spoliata *Lw.*, 154
stigmatias *Lw.*, 166
Thomæ *Lw.*, 163

Gorgopis, 27, 65
Gymnopoda, 22

Helomyzidæ, 27
Hemixantha, 69, 190
Hemixantha spinipes *Lw.*, 190
Heniconeura, 24, 42
Heniconeura fenestralis *Macq.*, 24
Henicoptera, 21
Heramyia, 16, 17
Herina, 16, 18, 24
Herina calcarata *Macq.*, 47
chalybea *Dobesch.*, 47
metallica *v. d. Wulp.*, 93
mexicana *Macq.*, 47
quadrifasciata *Macq.*, 90
ruficeps *v. d. Wulp.*, 123
rustitarsis *Macq.*, 89

Hesiquillia, 16, 17
Hesiquillia lugubris *Rob. Desv.*, 17
Heterogaster, 24, 34
Heteromyzidæ, 24
Himeroëssa, 44, 85
Himeroëssa pretiosa *Lw.*, 85
Holodasia, 58
Holodasia fraudulosa *Lw.*, 58
Homalocephala, 23
Hypoehra *Lw.*, 57
Hypoehra albipennis *Lw.*, 57
Hypoëcta, 65, 173
Hypoëcta longula *Lw.*, 173
Hypotyphia, 33, 36

Idana, 52, 115
Idana marginata *Soy.*, 115
Idiotypa, 67, 183
Idiotypa appendiculata *Lw.*, 183

- Lamprogaster, 25, 27, 39
 Lauxaniidæ, 21, 23
 Leptopoditæ, 21, 22
 Lipara, 22
 Lonchæa, 6, 8, 9, 10, 19, 22
 Lonchæa laticornis, 22
 Loxodesma, 55
 Loxodesma lacustris Meig., 56
 Loxoneura, 13, 24, 27, 36
 Loxoneura decora Fbr., 36

 Maria, 27
 Meckelia, 16, 17
 Melanoloma, 69, 192
 Melanoloma affinis Lw., 192
 cyanogaster Wied., 69,
 193
 Melieria, 16, 17
 Meracantha, 23, 70
 Michogaster, 11, 18, 22, 23
 Micropezidæ, 16
 Mischogaster, 11, 18, 22, 47, 98
 Mischogaster diffusus Gerst., 47
 egregius Gerst., 67, 183
 purnix Gerst., 47
 Musca annonæ Fbr., 162
 costalis Fbr., 158
 octopunctata Coqueb., 8, 58
 picta Fbr., 50, 109
 stigma Fbr., 148
 Myennis, 16, 17, 58, 142
 Myennis fasciata Fbr., 59
 scutellaris Wied., 143
 Myodina, 16, 18, 152
 Myopa nigripennis Gray, 77
 Myoris, 16, 17
 Myrmecomyia, 11, 16, 18, 48, 99
 Myrmecomyia myrmecoides Lw., 48,
 100

 Notacanthina, 13, 26
 Notogramma, 12, 65, 148
 Notogramma camiciformis Lw., 149
 stigma Fbr., 148

 Odontomera, 23, 26
 Odontomera ferruginea Macq., 66, 179
 maculipennis Macq.,
 68, 179
 Oedopa, 65, 146
 Oedopa capito Lw., 146
 Omalocephala, 23
 Ortalidæ, 28, 71
 Ortalina, 52, 118
 Ortalis, 2, 32, 54
 Ortalis ænea Wied., 170
 annonæ Wied., 162
 atomaria Wied., 11, 52, 118

 Ortalis cana Lw., 123
 chalybea Wied., 51
 colon Harris, Cat., 153
 connexa Meig., 3, 10, 25
 costalis Wied., 158
 dentipes Macq., 25
 fasciata Fbr., 17, 25
 fasciata Wied., 52, 132
 frondescentiæ Lin., 25
 fulminans Meig., 3
 lamed Meig., 5
 luctuosa Meig., 7
 lugens Fbr., 43
 marginata Say, 52, 116
 moerens Fbr., 43
 notata Wied., 156
 obscura Wied., 59
 ornata Meig., 25
 Ortoeda Wlk., 44, 89
 paludum Meig., 7, 18
 parallela Wied., 43
 pœcilloptera Meig., 3, 10
 quadrifasciata Wlk., 89
 ruficeps Fbr., 55
 syngenesiæ Meig., 3, 4, 6, 32,
 44
 trifasciata Dolesch., 42
 trifasciata Say, 170
 trifasciata Wied., 11, 52, 118
 vau Say, 8, 59, 61, 138
 vibrans Lin., 3, 4, 16, 25, 32,
 153
 violacea Macq., 47
 Orygma, 21
 Oscinidæ, 22
 Oscinis, 16, 17
 Otites, 5, 17, 21
 Oxycephala, 12, 22, 24, 73
 Oxycephala fuscipennis Macq., 77
 fenestrata Macq., 83
 maculipennis Macq.,
 74, 76

 Pachycephala, 38
 Palloptera, 8, 9, 10, 19, 22
 Pallopteridæ, 9, 11, 30
 Palpomyia, 16, 17
 Palpomyia Lalandi Rob. Desv., 17
 Peronomatia 53
 Peronomatia inornata Lw., 53
 Petalophora, 42
 Physogenia, 15, 27
 Phytalmia, 27
 Phytalmia cervicornis Gerst., 27
 megalotis Gerst., 27
 Phytomyidæ myodinæ, 16
 thelidomydæ, 16
 Piara, 50

- Plagiocephala, 26
 Platycephala, 21
 Platystoma, 6, 10, 24, 32, 36
 Platystoma annulipes *Macq.*, 5, 59,
 62, 141
 asphaltina *Wied.*, 17, 37
 australis *Wlk.*, 37
 cincta *Guér.*, 38
 costalis *Wied.*, 00
 decora *Wied.*, 13, 24
 fasipennis *Macq.*, 36
 latipennis *Macq.*, 37
 lunulata *Macq.*, 37
 microcera *Macq.*, 16
 nigronotata *Lw.*, 38
 ocellata *Macq.*, 13, 36
 pectoralis *Lw.*, 37, 38
 seminationis *Fbr.*, 17
 stictica *Fbr.*, 36
 umbrarum *Fbr.*, 17, 37
 Platystomina, 36, 83
 Polistodes, 11, 16
 Polistoides, 18
 Prosopomyia, 15
 Psairoptera, 7, 10, 32, 62
 Psilomyia, 20, 22
 Psilomyia, 21
 Pterocalla, 13, 60, 132
 Pterocalla ocellata *Wied.*, 13, 60
 strigula *Wied.*, 60, 133
 Pterocallina, 58, 132
 Pterogenia, 27
 Pteropæctria, 56
 Pteropæctria palustris *Macq.*, 56
 Pteropæcila, 54
 Pteropæcila lamed *Schrk.*, 54
 Ptilonota, 54
 Ptilonota centralis *Fall.*, 54
 Pyrgota, 12, 21, 28, 36, 72
 Pyrgota millepunctata *Lw.*, 73
 pterophorina *Gerst.*, 81
 undata *Wied.*, 77
 vespertilio *Gerst.*, 79
 Pyrgotina, 33, 72

 Rhopalomera, 14, 15, 21
 Rhopalomera pleuropunctata *Wied.*,
 14
 Richardia, 14, 16, 18, 25, 63, 66, 178
 Richardia flavitarsis *Macq.*, 66
 podagrica *Fbr.*, 179
 Richardina, 66, 171
 Rioxa, 27
 Rivellia, 16, 18, 25, 44, 87
 Rivellia atra *Lw.*, 45
 basilaris *Wied.*, 44
 Boscii *Rob. Desv.*, 93
 conjuncta *Lw.*, 88

 Rivellia flavimana *Lw.*, 92
 micans *Lw.*, 94
 pallida *Lw.*, 95
 quadrifasciata *Macq.*, 90
 syngenesiæ *Lin.*, 44
 variabilis *Lw.*, 91
 viridulaus *Rob. Desv.*, 44, 88

 Sapromyza, 19
 Sapromyzidæ, 15, 30
 Scatomyzidæ, 20
 Scatophaga bispinosa *Fbr.*, 13
 fasciata *Fbr.*, 8, 17, 25,
 38, 58
 trimaculata *Fbr.*, 68
 Scholastes, 38
 Scholastes cinctus *Guér.*, 39
 nepticula *Lw.*, 39
 Scionomyza bucephala *Meig.*, 5, 10, 17
 Scionomyzidæ, 15, 30
 Scotinosoma, 45
 Seioptera, 152
 Senopterina, 12, 22, 25, 40
 Senopterina decora *Macq.*, 40
 Sceptera, 63, 151
 Sceptera colon *Lw.*, 152
 vibrans *Lin.*, 64, 153
 Sepsidæ, 7, 20, 22, 30
 Sepsis Guérini *Bigot*, 66, 180
 Setellia, 16, 18, 22, 23, 66
 Setellia apicalis *Macq.*, 67, 182
 atra *Rob. Desv.*, 67
 Sophira, 27
 Sphecomyia valida *Harris*, 74
 undata *Harris*, 77
 Sphenoprosopa, 34, 36
 Steneretma, 64, 68, 186
 Steneretma laticauda *Lw.*, 187
 Stenomacra, 66, 180
 Stenomacra Guérini *Bigot*, 180
 Stenomyia, 65, 173
 Stenomyia tenuis *Lw.*, 174
 Stenopterina, 12, 22, 46, 96
 Stenopterina ænea *Wied.*, 97
 basalis *Wlk.*, 47
 brevipes *Fbr.*, 97
 bicolor *Wlk.*, 47
 cærulescens *Lw.*, 97
 decora *Macq.*, 40, 47
 femorata *Macq.*, 46
 gigas *Macq.*, 47
 immaculata *Macq.*, 46
 nigripes *Macq.*, 47
 scutellaris *Macq.*, 47
 submetallica *Lw.*, 47
 trivittata *Macq.*, 47
 Stictiocephala, 61, 134
 Stictiocephala corticalis *Fitch*, 61, 136

- Stictiocephala cribrum *Lw.*, 62, 135
 cribellum Lw., 62, 134
 vau Say, 61, 138
 Straussia, 17
 Stranzia, 16, 17
 Strumeta, 27
 Stylophora, 16, 18
 Syntaces, 67, 182
 Syntaces apicalis *Macq.*, 67
 Systata, 55
 Systata rivularis *Fbr.*, 55


 Tachinidæ, 27
 Tephritidæ, 21, 23
 Tephritis anomæ *Fbr.*, 162
 caerulea Macq., 41
 conica Fbr., 109
 corticalis Fitch, 61
 dorsalis Macq., 25
 fasciata Meig., 17
 melliginis Fitch, 44, 89
 strigipennis Macq., 41
 violacea Gray, 36
 Tephronota, 56, 122
 Tephronota gyrans *Lw.*, 57
 humilis Lw., 123
 Terastomyia, 27
 Teromyia, 22
 Tetanocera bispinosa *Fbr.*, 13, 26
 Tetanops, 5, 10, 21, 53, 119
 Tetanops luridipennis *Lw.*, 119
 integra Lw., 121
 myopina Fall., 53
 sanguiniiceps Wied., 13, 16,
 35
 Themara, 27
 Thryophila, 56
 Thryophila frondescentiæ *Lin.*, 56
 Thyreophoridæ, 20
 Timia, 6, 10, 63, 65
 Timia apicalis *Meig.*, 6
 erythrocephala Wied., 6
 Toxoneura, 8, 9, 19, 22
 Toxotrypana, 27, 34, 36
 Toxura, 23, 35
 Traphera, 51
 Trigonometopus, 21
 Trigonostoma, 27
 Tritoxa, 50, 102
 Tritoxa cuneata *Lw.*, 107
 Tritoxa flexa *Wied.*, 102
 incurva Lw., 104
 Trypeta ænea *c. d. Walp.*, 170
 albovaria Wlk., 27, 43
 arcuata Wlk., 27, 102
 basilaris Wied., 13, 44
 cyanogaster Wied., 13, 69,
 192
 excepta Wlk., 27
 fasciata Fbr., 8, 10
 flexa Wied., 13, 50, 102
 longipennis Wied., 17
 obscura Wied., 13, 24, 60
 ocellata Wied., 13, 59
 pieta Fbr., 18
 pieta Wied., 13, 24, 109
 quadrifasciata Harris, 88
 scutellaris Wied., 13, 143
 trimaculata Wied., 26, 168
 ulula Lw., 60
 Trypetidæ, 30

 Ulidia, 6, 22, 23, 63, 65
 Ulidia ænea *Fbr.*, 12
 bipunctata Macq., 65
 demandata Meig., 6
 fulvifrons Bigot, 65
 metallica Bigot, 65
 stigma Fbr., 12, 65
 stigma Wied., 149
 Ulidina, 64, 145
 Urophora, 18, 22
 Urophora ænea *Macq.*, 172
 fulvifrons Macq., 22, 170
 quadrivittata Macq., 22,
 162

 Valonia, 27, 38
 Valonia complicata *Wlk.*, 38
 Vidalia, 16, 17

 Xangelina, 27
 Xiria, 27

 Zona, 27, 36
 Zygothrica, 23
 Zygothrica dispar *Macq.*, 24
 Zygotricha, 23
 Zygaenula, 65
 Zygaenula paradoxa *Dolensch.*, 27, 65



REVIEW
OF THE
NORTH AMERICAN TRYPETINA.

INTRODUCTION.

IN 1860, at the time of the publication of my paper on the *Trypetidæ*, contained in the first volume of these Monographs, only twenty-three North American species of this family were known. Since then, this number has reached sixty-one. Among these additions there is a number of species of previous authors, concerning which I did not possess sufficient information at the time of my earlier essay. Moreover, a number of species published by Wiedemann became accessible to me in type specimens, through the kindness of the Berlin and Vienna Museums. Since that time, also, several other authors have published new species belonging to the same group. And, finally, the systematic distribution of the group *Trypetina* has obtained, for the European species, a more solid foundation.

It would seem to be time, therefore, to undertake an entirely new work on the *Trypetina* of North America; but as the plan of the present series does not well admit of it, I have adopted the form of a supplement to my previous paper. One of the principal aims of the present essay will be, the adaptation to the American fauna, as far as it is possible, of the systematic distribution introduced among the *Trypetina* of the old continent. While I was engaged on Monographs, etc., Part I, the number of the North American species with which I was acquainted, was, as yet, too insignificant for an attempt at a subdivision in smaller groups; besides, similar attempts, undertaken for the European species

by other authors (an account of them may be found in Monographs, etc., Part I, p. 49-51), seemed to me so ill conceived, that I did not feel inclined to adopt them as a basis for further development. I perceived, on the contrary, that any attempt to subdivide exotic Trypetidae must be preceded by a rational systematic distribution of the more abundant material of the European species. In 1862, in my monograph of the European *Trypetidae*, I divided the *Trypetina* into twenty subgenera: *Platyparea*, *Euphranta*, *Aciura*, *Hemileca*, *Anomara*, *Acidia*, *Spilographa*, *Zonosema*, *Rhagoletis*, *Rhacochlæna*, *Trypeta*, *Eusina*, *Myopites*, *Urophora*, *Sphenella*, *Corphotricha*, *Oryphora*, *Oryna*, *Tephritis*, and *Urellia*. The definitions of these groups will be found in the above-quoted work. To these must be added: *Hyphenidium* (established by me since, in the *Berliner Entom. Zeitschr.*, VI, p. 87), *Orellia* (separated by Schiner, in his *Fauna Austriaca*, from *Oedaspis*) and *Chetostoma* (established by Rondani, in his *Prodromus*, Vol. I). Such is the present state of the classification of the European *Trypetina*, upon which the distribution of the known North American species is to be based. Considerable as the number of the latter is, it is certain at the same time that this number does not reach one-fifth, perhaps not one-tenth, of all the existing North American *Trypetina*. Any attempt at a distribution, therefore, would probably be modified by further discoveries. In this dilemma, the course I adopted was, to append to the description of each species the necessary remarks on its systematic position, and to give a general survey of all the results thus obtained, at the end of the volume.

Detailed descriptions of those species only are given here, which are not described in Monographs, etc., Part I, or the descriptions of which were insufficient. The descriptions contained in that volume are indicated by references; the diagnoses, however, even of those older species are reproduced here, with the modifications rendered necessary by the addition of the new species.

An important defect of the present publication is, that a considerable number of the new species are not represented on the plates. The reason is, that the plates were prepared more than four years ago, at a time when the number of the known North American species was not sufficient to fill the required number

of figures. This was done by the addition of a number of South American species, described for the sake of comparison, but the figures of which I would have preferred now to replace by those species from North America, which I received after the plates were printed.

The critical examination of the species described by other authors, appended to the first volume, p. 57-61, required several corrections and additions. I have, therefore, reproduced it, thus amended, at the end of the present volume, as Appendix I. Appendix II contains descriptions, by other authors, of species not known by me and not contained in Part I.

The materials for the present publication, as far as the North American species are concerned, are principally, almost exclusively, derived from the communications of Baron Osten-Sacken. If I had had a similar support from more than one side, my work might, of course, have been more complete and more perfect. As it is, I have been compelled to draw the descriptions of several species from single, often badly preserved, specimens, and I am afraid that these descriptions, as well as the opinions expressed by me on the systematic position of some species, may sometimes betray the incompleteness of my materials. I trust that an equitable critic will bear these circumstances in mind in framing his appreciations.

H. LOEW.

GRUBER, August, 1873.

LIST OF THE DESCRIBED SPECIES OF TRYPETA.¹

	PAGE		PAGE
1. <i>eximia</i> Wied. ♂. ♀ . . .	216	24. <i>tabellaria</i> Fitch, ♀ . . .	263
2. <i>amabilis</i> n. s. ♂ . . .	219	25. <i>pomonella</i> Walsh, ♀ . . .	265
3. <i>suspensa</i> Lw. ♂ . . .	222	26. <i>insecta</i> Lw. ♀ . . .	268
4. <i>fraterculus</i> Wied. ♂ . . .	222	<i>phoenicra</i> n. s. ♂. ♀ . . .	269
5. <i>ludens</i> n. s. ♂ . . .	223	27. <i>pœcilogastra</i> n. s. ♂ . . .	270
6. <i>tricincta</i> n. s. ♂ . . .	225	28. <i>testudinea</i> n. s. ♀ . . .	272
<i>parallela</i> Wied. ♂. ♀ . . .	229	29. <i>sparsa</i> Wied. ♂. ♀ . . .	274
<i>hamata</i> n. s. ♂. ♀ . . .	229	30. <i>rotundipennis</i> Lw. ♂ . . .	276
<i>integra</i> n. s. ♂. ♀ . . .	230	31. <i>culta</i> Wied. ♂. ♀ . . .	276
<i>consobrina</i> n. s. ♂. ♀ . . .	230	32. <i>solidaginis</i> Fitch, ♂. ♀ . . .	279
<i>pseudoparallela</i> n. s. ♂. ♀ . . .	230	33. <i>comma</i> Wied. ♀ . . .	280
7. <i>vulnerata</i> n. s. ♂. ♀ . . .	232	34. <i>latifrons</i> Lw. ♀ . . .	283
8. <i>fratria</i> Lw. ♀ . . .	234	35. <i>melanura</i> n. s. ♀ . . .	283
9. <i>suavis</i> Lw. ♂ . . .	235	36. <i>alba</i> Lw. ♂. ♀ . . .	285
10. <i>canadensis</i> n. s. ♀ . . .	235	37. <i>albidipennis</i> Lw. ♂. ♀ . . .	286
11. <i>longipennis</i> Wied. ♂. ♀ . . .	238	38. <i>Vernoniæ</i> Lw. ♂. ♀ . . .	286
12. <i>electa</i> Say, ♀ . . .	243	39. <i>seriata</i> Lw. ♂ . . .	287
13. <i>flavonotata</i> Macq. ♂ . . .	244	40. <i>circinata</i> n. s. ♂. ♀ . . .	288
14. <i>tetanops</i> n. s. ♂ . . .	245	41. <i>Lichtensteini</i> Wied. ♂ . . .	289
15. <i>sarcinata</i> Lw. ♀ . . .	247	42. <i>humilis</i> Lw. ♂. ♀ . . .	291
16. <i>discolor</i> Lw. ♂ . . .	250	<i>peregrina</i> n. s. ♂. ♀ . . .	292
17. <i>obliqua</i> Say, ♂. ♀ . . .	251	43. <i>angustipennis</i> Lw. ♂. ♀ . . .	293
<i>biseriata</i> n. s. ♀ . . .	252	44. <i>finalis</i> Lw. ♂. ♀ . . .	296
18. <i>palposa</i> Lw. ♂ . . .	253	45. <i>clathrata</i> Lw. ♀ . . .	297
19. <i>florescentiæ</i> Lin. ♂. ♀ . . .	254	46. <i>geminata</i> Lw. ♀ . . .	298
20. <i>polita</i> Lw. ♀ . . .	256	47. <i>fucata</i> Fbr. ♂ . . .	300
21. <i>atra</i> Lw. ♂. ♀ . . .	256	48. <i>albiceps</i> n. s. ♂. ♀ . . .	302
<i>nigerrima</i> n. s. ♀ . . .	258	49. <i>euryptera</i> n. s. ♀ . . .	304
22. <i>glbba</i> n. s. ♀ . . .	260	50. <i>platyptera</i> n. s. ♀ . . .	306
23. <i>cingulata</i> Lw. ♂. ♀ . . .	263	51. <i>æqualis</i> Lw. ♂. ♀ . . .	308

¹ The species from South America, described for the sake of comparison with North American species, are printed in smaller type and not numbered in this list.

LIST OF THE DESCRIBED SPECIES OF TRYPETA. 215

	PAGE		PAGE
52. <i>festiva</i> Lw. ♂. ♀ . . .	309	56. <i>mexicana</i> Wied. ♂ . . .	317
<i>spectabilis</i> n. s. ♂. ♀ . . .	309	57. <i>pura</i> n. s. ♀ . . .	320
53. <i>bella</i> Lw. ♂. ♀ . . .	311	58. <i>abstersa</i> Lw. ♂. ♀ . . .	322
54. <i>timida</i> Lw. ♂ . . .	311	59. <i>polyclona</i> n. s. ♀ . . .	324
<i>obscuriventris</i> n. s. ♀ . . .	313	60. <i>solaris</i> Lw. ♀ . . .	325
55. <i>melanogastra</i> Lw. ♂. ♀ . . .	315	61. <i>actinobola</i> n. s. ♂ . . .	326
<i>tenuis</i> n. s. ♀ . . .	316		

PAGE

263
265
268
269
270
272
274
276
276
279
280
283
283
285
286
286
287
288
289
291
292
293
296
297
298
300
302
304
306
308

ison
um-

DESCRIPTION OF THE SPECIES.

1. *T. eximia* WIED. ♂ ♀.—Lutea, abdomine nigro-fasciato; scutellum magnum, planum, setis sex validis instructum; alarum plectura fusca inde a basi maculis irregularibus variegata ad ultimum usque trientem pertinet, ubi vittam costalem et fasciam a margine antico ad posticum oblique ductam emittit; præterea in margine antico duæ macule triangone et hyaline, in postico duæ subovatæ et subhyaline conspiciuntur, ad quas in speciminibus plerisque macula rotunda hyalina in cellule discoidalis basi sita accedit.

Clay-yellow, abdomen banded with black; scutellum large, flat, with six strong bristles; the brownish-black coloring of the wings reaches from the irregularly spotted basis to the last third of the wing, where it emits two bands, one of which forms a border along the costa, the other runs obliquely from the anterior to the posterior margin; moreover, the anterior margin shows two triangular hyaline spots, the posterior margin two almost oval and less hyaline spots; most specimens have, besides, a round hyaline spot on the basis of the discal cell. Long. corp. 0.26—0.28, ♀ cum terebrâ 0.29—0.30; long. al. 0.25—0.26.

SYN. *Trypeta eximia* WIED. Zweifl. Ins. II, p. 477, 2.

Tephritis fasciventris MACQ. Dipt. Exot. Suppl. IV, p. 291. Tab. XXVII, f. 3.

Clay-yellow; head of a somewhat purer yellow, rather disciform. Front narrow, still more narrowed anteriorly, with a small, but well-defined frontal lunule. Frontal and vertical bristles black, rather long and strong; the upper half of the posterior orbit of the eyes with a row of black and blackish-brown bristles. Antennæ clay-yellowish, third joint elongated, rounded at the tip; arista very slender, with a hardly perceptible pubescence. Face perpendicular; the edge of the mouth not upturned; palpi yellowish, broad, reaching as far as the anterior edge of the mouth; their pubescence, as well as that of the mentum and of the occiput, is yellow. Thorax rather strongly built, comparatively broad between the roots of the wings; the humeral callus and a longitudinal stripe between it and the root of the wing, are yellowish-white or sulphur-yellow; a longitudinal stripe of a similar color, which is generally but little visible in dried speci-

mens, runs from the posterior corner of the thoracic dorsum to the transverse suture; in some specimens the posterior border of the thoracic dorsum also shows a trace of a lighter coloring; the dense, but very short, pubescence of the thoracic dorsum is yellowish; the macrochaetae upon it are black; there are seven of them on each side, viz.: three on each side, in a row beginning at the humerus and ending before the root of the wings; three others a little farther from the lateral margin in a row beginning at the transverse suture and ending in the vicinity of the posterior corner; finally, a single bristle between the last one of this second row and the lateral corner of the scutellum; there are only two pairs of macrochaetae on the longitudinally middle portion of the thoracic dorsum, not far from the posterior margin; the bristles of the posterior pair are at a moderate distance from each other, the distance between those of the anterior pair is perhaps three times greater. All the bristles and bristle-like hairs upon the pleurae and the pectus are black; the short pubescence upon the upper half of the pleurae is blackish, on the lower half it is pale-yellow. Scutellum comparatively large, flat, with a short, yellowish pubescence on the upper side, and with six strong macrochaetae along the edge; in life, the scutellum is probably altogether whitish-yellow or sulphur-yellow, while in dry specimens, this coloring is perceptible along the borders only. The abdomen has brownish-black bands, which do not reach the posterior margin of the segments; these bands occur upon the second, third, and fourth segments; they are often less developed upon the anterior segments than upon the posterior ones, and here sometimes interrupted; upon the rather large last abdominal segment of the male the brownish-black crossband is especially broad and more or less emarginate on its posterior side; my only female specimen has on the first abdominal segment an incompletely developed brownish-black band, situate before the posterior margin. The pile upon the abdomen is black; pale-yellowish on the upper side of the first segment and sometimes also on the basis of the second; however, all the pile upon the abdomen assumes, in a reflected light, and especially in specimens of a lighter coloring, a brownish-yellow, almost a ferruginous-yellow tinge (with the exception of the stronger, bristle-like hairs). The hypopygium is brownish-black; the brown ovipositor is conical, not flattened at all, perceptibly longer than the last

two segments taken together, but shorter than the last three. Its pile is brownish-yellow or brown, the color of the rather long bristle-like hairs on the end of the first segment is dark-brown or black. Feet clay-yellow; front femora on the upper side with short, on the under side with more elongate black bristles; front tibiae not bristly; middle femora at the end of the posterior side with a few bristles and, also, on the under side, with two longitudinal rows of short black bristles, which are more developed in the male than in the female; middle tibiae with a single row of bristles; hind femora, at the end of the upper side with elongated bristles, with shorter ones on the under side; hind tibiae with bristle-like cilia. Tegulae more than usually developed. Wings rather large and broad; the first longitudinal vein altogether beset with bristles, the third far beyond the small crossvein, the fifth upon the first and upon the beginning of its second section, bristly; the second longitudinal vein ends in the costa at an acute angle, and diverges very strongly from the third, the latter is not bent anteriorly at its end; crossveins rather approximate, the small one perpendicular and of a comparatively considerable length; the posterior one very steep and somewhat curved towards its posterior end; posterior angle of the anal cell drawn out in a rather long lobe. The brownish-black, sometimes almost black picture of the wings, is recognizable in Macquart's above-quoted figure, although not correctly rendered; the round pale spot in the discal cell should be much nearer to its basis; the pale indentation at the posterior margin, near the basis of the wing, should be much narrower; the stigma should be placed entirely in the dark portion of the coloring; the hyaline double spot near the anterior margin is seldom merely emarginate posteriorly; in most cases it is divided in two approximate triangular spots; other differences in the picture likewise occur; the most common is, that in the discal cell, a little beyond the small crossvein, there is a short, pale streak, crossing the cell, and which in some cases becomes a hyaline transverse spot. A male from Brazil in my collection has, instead of the round pale spot in the discal cell, only a somewhat paler place without any distinct outline; the agreement in the other characters being perfect, I take it for a rather unusual variety of *T. eximia*.

Hab. Brazil, especially Bahia and St. Paulo; Surinam; Mexico.

Observation 1.—Mr. Macquart, in the above quoted place, supposes that his *Tephritis fasciventris* may be only a variety of the *Tephritis major*, Dipt. Exot. Suppl. II, p. 93, Tab. VI, f. 6. However, this *Tephritis major* is identical with *Tephritis socialis* Wied., a species which is very distinct from *fasciventris* Macq. (syn. *eximia* Wied.).

Observation 2.—I have gone into more detail about the plastic characters of this species than was strictly necessary for its specific identification. I did so on account of the great resemblance in the plastic characters of *T. eximia* with *T. amabilis*, with *T. socialis* Wied., and with several other South American species. These species form a very well-defined group, for which I choose the name of *Hexachæta*, and which deserves to be considered as a separate genus. The generic character may be derived from what has been said, in the above description of *Trypeta eximia*, concerning the shape of the head and of its parts, the shape of the thorax and of the scutellum, the number and position of their macrochætæ, the bristles on the feet, as well as concerning the bristles on the wing-veins. The body and the picture of the wings of all the species of *Hexachæta* are strikingly uniform. I know of no other but American species of this group.

2. *T. amabilis* n. sp. ♂.—Lutea, thoracis dorsum sulphureo-vittatum, postice nigricans; pleuræ fusco-nigræ, sulfureo-vittatæ; scutellum magnum, planum, setis sex validis instructum, nigrum, late sulfureo-marginatum; abdomen fasciis tribus interruptis nigris ornatum; femora intermedia magnâ ex parte, postica fere tota nigra; alarum pictura fusco-nigra, præter maculam ingentem, quæ in mediâ alâ locum habet et totam ejus latitudinem explet, fasciam angustam subperpendicularem, quâ vena transversalis posterior includitur, et vittam costalem inde ab hac fasciâ usque ad summam alâ apicem pertinentem ostendit.

Clay-yellowish, thoracic dorsum with sulphur-yellow longitudinal stripes, blackish along the posterior margin; pleuræ brownish-black with sulphur-yellow longitudinal stripes; scutellum large, flat, with six macrochætæ, black, with a broad yellow border; abdomen with three interrupted black crossbands; intermediate femora partly, hind femora almost entirely brownish-black; the brownish-black picture of the wings shows, besides an unusually large spot upon the middle of the wing, occupying its whole breadth, a narrow, almost perpendicular crossband, covering the posterior crossvein, and from which a border extends along the costa as far as the apex of the wings. Long. corp. 0.26; long. al. 0.26.

Of the size of *T. eximia* Wied., and so closely allied to it in all the plastic characters, that their detailed description would be superfluous. Head and all its parts of the same coloring and the same structure as in that species, only the frontal bristles are somewhat weaker. The thoracic dorsum shows a delicate middle line, gradually fading anteriorly and expanding posteriorly into a large spot, which does not entirely reach the posterior thoracic margin, and is surrounded laterally and posteriorly by a blackish coloring; beginning at the shoulder, a sulphur-yellow stripe runs, gradually expanding, to the root of the wing; it emits, near the humeral callus, an upper branch, running towards the transverse suture; between both branches, the color changes into brownish. Pleuræ brownish-black, with a sulphur-yellow longitudinal stripe across the middle; moreover, the sulphur-yellow stripe between the humerus and the root of the wings, is prolonged under the latter as far as the posterior end of the thorax. Scutellum entirely of the same structure as in *T. eximia*, sulphur yellow, at the basis of the upper side with a large, semicircular brownish-black spot, the border of the upper side only remaining sulphur-yellow. Metathorax brownish-black, spotted with brown on the sides, and with a yellow spot on the middle of its upper side. The dense and very short pubescence of the thorax and the scutellum is more whitish-yellow than is usually the case in *T. eximia*; otherwise the hairs and bristles of both species are alike in their coloring; the number and position of the macrochaete is the same in both. Abdomen with three very broad black cross-bands, which lie on the second, third, and fourth segments, and leave uncovered only the middle line and the posterior margin of these segments. The pile on the abdomen is black; on the upper side of the first segment and along the posterior border of the second, pale-yellowish. Hypopygium brownish-black. Coxæ and feet yellow; the intermediate femora towards the basis, to a great, but variable extent, brownish-black; hind femora black, somewhat yellow towards the end, especially on the under side. The bristles on the femora and tibiæ are almost as in *T. eximia*. The shape of the wings, the venation, and the position of the bristles are exactly as in that species; the pattern of the picture is likewise a somewhat similar one; however, it differs considerably in the details; the bulk of the dark coloring extends a little beyond the small crossvein and is gently

rounded off, the curve formed by it striking the anterior margin nearly at a right, the posterior margin at an acute angle; the latter margin, however, is not quite reached, as a narrow hyaline space remains between it and the dark coloring; this curve would have been a perfect one, were it not for a small projection before the posterior crossvein and for a small excision immediately beyond it; near the anterior margin, the dark-brown coloring, immediately before its end, is interrupted by a triangular hyaline indentation, the tip of which reaches the third longitudinal vein immediately before the small crossvein; the distal side is concave, the proximal side is straight and perpendicular to the costa. The brown coloring has no distinct limit towards the base of the wing; it gradually dissolves into a system of irregular spots; the costal cell is hyaline, with the exception of a brown infuscation along the costa between the humeral crossvein and the auxiliary vein; likewise hyaline are the extreme basis of the marginal cell and the entire second basal cell with the exception of a very narrow brownish-black border along the veins inclosing it; the first basal cell at its root, as far as the humeral crossvein, is also rather hyaline; beyond this, for an almost equal distance, it is yellowish; the anal cell is of a dirty yellow, blackish-brown towards its end, which color also extends over the basis of the third posterior cell; alula, posterior angle of the wing, and the portion of the third posterior cell lying alongside of it, are hyaline; moreover, in the third posterior cell, quite near its basis, at the place where it is contiguous to the second basal cell, there is an elliptical hyaline drop; in the first basal cell, below the beginning of the third longitudinal vein, there is a longitudinal spot of a dirty ferruginous color; a somewhat larger spot of the same coloring is in the marginal cell, below the place, where the auxiliary vein diverges from the first longitudinal. The hyaline apical portion of the wing shows a narrow crossband, covering the posterior crossvein, almost perpendicular, very gently curved, of a brownish-black color; its anterior end turns towards the costa in the shape of a bow and follows it afterwards as a narrow border, as far as the tip of the fourth longitudinal vein.

Hab. Mexico (collection of Mr. v. Roeder).

3. *T. suspensa* Lw. ♂. (Tab. X, f. 5.)—Tota lutea, alarum rivulis fuscans, cellulâ basali secundâ et cellulæ discoidalis basi non hyalinis, apice venæ longitudinalis quartæ recurvo.

Altogether clay-yellow, rivulets of the wings infuscated; second basal cell and root of discal cell not hyaline, the tip of the fourth longitudinal vein curved forward. Long. corp. 0.21; long. al. 0.22—0.23.

SYN. *Trypeta suspensa* LOEW, Monogr., etc., I, 69. Tab. II, f. 5.

The present species begins a group of very closely allied species, very much resembling one another. I have nothing to add to my above-quoted description of *T. suspensa*; I will only notice that the absence of pale yellow stripes on the thorax and of a pale yellow coloring of the scutellum cannot be considered as absolutely distinctive of this species, as these marks often disappear in other species in the process of drying. The readiest distinctive mark between *T. suspensa* and the very similar, but larger *T. fraterculus* is, that in the former, the second basal cell and the root of the discal cell have a yellowish color, while in the latter they are hyaline. I regret to have to notice here, that the engraver, in figuring *T. suspensa*, has committed an error in drawing the curvature of the tip of the fourth vein; this curvature is exactly similar to that in *T. fraterculus*, that is, running forward; and although this curved tip in *T. suspensa* is a little shorter, the difference is not at all such as the figure would lead one to suppose. The second basal cell and the basis of the discal cell should be somewhat paler in the figure, as they are not brown, but only yellow.

Hab. Cuba (Poey).

4. *T. fraterculus* WIED. ♂. (Tab. X, f. 6.)—Lutea, thoracis vittis et scutello dilutius tinctis, ultimo abdominis segmento duobus præcedentibus simul sumtis paulo brevior, alarum rivulis lutescentibus, cellulâ basali secundâ et cellulæ discoidalis basi hyalinis, apice venæ longitudinalis quartæ recurvo.

Clay-yellow, longitudinal stripes of thorax and scutellum paler yellow; last abdominal segment a little shorter than the two previous ones taken together; wings with rather clay-yellow rivulets; first basal cell and root of the discal cell hyaline; the end of the fourth longitudinal vein curved forward. Long. corp. 0.26; long. al. 0.27.

SYN. *Dacus fraterculus* WIEDEMANN, ANS. ZW. II, p. 524.

Trypeta unicolor LOEW, Monogr., etc., I, p. 70. Tab. II, f. 6.

To my former description of this species, I have to add two observations. First, it contains a misprint, as the third line should read "bristle very thin," and not "bristle very short." Secondly, the examination of well-preserved specimens renders it doubtless, that the dark spots on the thoracic dorsum, mentioned in the description, were produced by the immersion of the specimens in spirits, and that the better preserved specimens do not show them.

When I described *T. unicolor*, I took it for distinct from *Dacus fraterculus* Wied., as Wiedemann describes the bristles and hairs on head and thorax as black, and says that the large triangular hyaline spot at the end of the posterior margin is connected with the S-shaped hyaline band. The comparison of Wiedemann's original specimen, however, showed that my *T. unicolor* is nothing else but *Dacus fraterculus* Wied. By the terms *hairs* and *bristles* Wiedemann understood only the stronger and weaker bristles; the remaining short pile on the head and the thorax of his specimen is entirely similar to the yellowish pubescence of *T. unicolor*. The connection between the posterior hyaline spot with the S-shaped hyaline band, which he mentions, is only an apparent one, as the rivulet separating both is not interrupted at the tip of the triangular hyaline spot, but only very much faded.

Hab. Brazil, Peru, New Granada, Cuba.

Observation.—The *Tephritis obliqua* Macq. Dipt. Exot. II, 3, p. 225, Tab. XXX, f. 11, undoubtedly belongs in the relationship of the two preceding species; it differs, however, in the picture of the wings too much to be identified with any of them.

5. *T. ludens* n. sp. ♂. (Tab. XI, f. 19.)—Lutea, thoracis vittis et scutello lætius flavis, ultimo abdominis segmento duobus præcedentibus simul sumtis multo longiore, alarum rivulis lutescentibus, cellulâ basali secundâ et cellulâ discoidalis basi hyalinis, apice venæ longitudinalis quartæ recurvo.

Clay-yellow, longitudinal stripes of thorax and scutellum of a purer yellow; the last abdominal segment much longer than the two preceding ones taken together; wings with rather clay-yellow rivulets, the second basal cell and the root of the anal cell hyaline; the end of the fourth longitudinal vein curved forward. Long. corp. 0.30; long. al. 0.31—0.32.

Pale clay-yellow. Front of a somewhat more bright yellow,

of a very moderate breadth; the usual frontal bristles black, only the upper ones rather long and strong. The yellow antenna almost as long as the face; arista long and slender, with a very short and delicate pubescence. Oral opening rather large; oral edge rather sharp. Proboscis and palpi yellow, the latter rather broad; the suctorial flaps somewhat prolonged. The upper side of the thorax of a light, bright clay-yellow; a sulphur-yellow middle stripe, gradually vanishing anteriorly, expanding posteriorly in a cuneiform shape, and nowhere well defined; scutellum sulphur-yellow; on each side, above the root of the wings, a well-marked pale-yellow longitudinal stripe, which runs from the transverse suture to the posterior margin of the thorax; quite on the lateral margin an indistinct, but broader pale yellow stripe; the humeral corner and a well-defined stripe on the upper part of the pleuræ, reaching to the root of the wings, likewise of a bright pale yellow. The very short pile on the thorax is yellowish; the usual bristles are black or blackish-brown. Scutellum with four black bristles. Metathorax clay-yellow. Abdomen with short yellowish pile and with black bristles on its posterior end; the last segment very much prolonged, much longer than the two preceding ones taken together (this character serves easily to distinguish this species from *T. fraterculus*, which is very much like it). Feet yellow; under side of the front femora with several blackish-brown bristles. Wings not very broad in comparison to their considerable length; the rivulets upon them are pale brownish-yellow with narrow, but little conspicuous, and not always perceptible brown borders; near the posterior margin and on the apex of the wing they are altogether brownish; the hyaline spaces between the rivulets are as follows: 1. An oblique band, interrupted upon the third longitudinal vein, the anterior part of which forms, immediately beyond the stigma, a spot extending from the costa to the third longitudinal vein, while the posterior part of the band occupies the portion of the basal cell which lies under the stigma, the basis of the discal cell and the second basal cell; 2. A broad S-shaped band which begins at the posterior margin, between the tips of the fifth and sixth longitudinal veins, passes between the two crossveins, reaches the second longitudinal vein, turns backwards and reaches the margin in the vicinity of the end of the fourth longitudinal vein; 3. A large triangular spot near the posterior margin, which fills a considerable part of

the second posterior cell, reaches with its tip considerably beyond the fourth longitudinal vein, and almost coalesces here with the S-shaped hyaline band. The external costal cell also is hyaline, with the exception of its basis, but has a more yellowish tinge than the other hyaline spaces. Stigma rather long, almost imperceptibly darker than its surroundings. Crossveins straight and steep; the third longitudinal vein distinctly bristly; the end of the fourth longitudinal vein turned forward; the posterior end of the anal cell drawn out in a very narrow, long lobe.

Hab. Mexico (coll. Winthem).

Observation.—The comparison of the description of *Trypeta fraterculus* and *T. ludens* shows the great resemblance of the two species and an entirely satisfactory distinctive character in the different length of the last abdominal segment. The females of these species, which unfortunately I have not seen, will probably be easy to distinguish, if attention is paid to the size, which is larger in *T. ludens*, to the somewhat broader cheeks, the longer last abdominal segment of this species, and to the course of the third and fourth longitudinal veins, which suddenly diverge here, while their divergency in *T. fraterculus* is much more gradual. In using the coloring for distinguishing the two species, a certain caution is necessary here, as well as in the other species of this group.

6. *T. tricineta* n. sp. ♂.—Intea, sentelli basi tribusque abdominis fasciis nigris, alarum rivalis nigro-fuscis, apice vene longitudinalis quartæ recurvo.

Clay-yellow; basis of the scutellum and three crossbands of the abdomen black; the end of the fourth longitudinal vein somewhat curved forward. Long. corp 0.26; long. al. 0.26—0.27.

Clay-yellow, more yellowish-red on the thoracic dorsum. Head of the same color and shape as in the three preceding species. In the middle of the thoracic dorsum there is a longitudinal sulphur-yellow stripe, proceeding from the posterior margin; it is rather broad posteriorly, gradually becomes narrower anteriorly, and finally disappears near the anterior margin; moreover each posterior corner emits a conspicuous sulphur-yellow stripe to the transverse suture; the humeral callosity and a broad longitudinal stripe reaching from it to the root of the wing and then passing under the latter to the posterior part of the thorax,

are, likewise, sulphur-yellow. The very short pile on the thoracic dorsum is pale yellowish, towards the posterior corners only it assumes a blackish tinge or at least a blackish appearance. The black macrochaetae of the thoracic dorsum are similar, in number and position, to those of the three preceding species. Scutellum sulphur-yellow, with four macrochaetae on the margin. Metathorax brownish-black, with a clay-yellow longitudinal stripe in the middle of its superior margin. Abdomen on the 2d, 3d, and 4th segments with a transverse band near the anterior margin; that of the second segment is entire and occupies only one-half of its length; those of the third and fourth segments are narrowly interrupted in the middle and cover a little more than the anterior half of the segment; the fourth segment is hardly longer than the preceding two, taken together. Hypopygium clay-yellow. The pile on the abdomen is blackish, and yellowish only on the upper side of the first and on the pale-colored portions of the upper side of the second segment; in a reflected light, the pile on the whole abdomen assumes a paler hue; the rather weak bristles at the end of the last segment are black. Feet clay-yellowish; the pile and bristles are similar to those in the three preceding species. Wings hyaline, with a rather dark-brown picture; it is not quite as brownish-black as that of *T. serpentina* Wied, figured on Tab. XI, f. 25, but it is more like it than any other species to me known. In order to form an idea of the picture of the wings of *T. tricineta*, let us represent to ourselves that the whole outer costal cell in that figure is rather hyaline, that the regions figured in gray are yellow and those represented as black are dark brown; that the S-shaped rivulet, beginning at the basis of the third posterior cell, running towards the anterior margin, and ending at the apex of the wing, is, upon its latter half, at least one-half broader than represented; that the band beginning at the posterior margin and covering the posterior crossvein is also broader than represented in the figure, and this in such a manner, that its side, looking towards the root of the wing, is a little less concave; finally, add to this picture a little streak of a saturate brown, beginning at the posterior margin and reaching somewhat beyond the fourth longitudinal vein (at the very place where Tab. XI, fig. 22, shows a similar streak, reaching only as far as the fourth longitudinal vein).

Hab. Hayti (caught on shipboard, by Mr. P. R. Uhler, sixty miles northwest of St. Nicholas, Hayti).

Observation 1.—The *Trypeta* described by Wiedemann as *Dacus serpentinus*, differs from *T. tricineta* not only in the picture of the wings, but also in the coloring. Wiedemann's original specimen, compared by me, comes from Brazil; but I have received a number of specimens of the same species from Peru. The *Urophora vittithorax* Macq. Dipt. Exot. Suppl. IV, p. 286, Tab. XXVI, f. 11, is identical with *T. serpentina* Wied. The habitat "de l'Inde," given by Macquart, is certainly erroneous, if it means the East Indies; but the species may occur in the West Indies, just as *T. fraterculus* occurs in Peru, Brazil, and Cuba.

Observation 2.—*T. suspensa* Lw., *fraterculus* Wied., *ludens* n. sp., and *tricincta* n. sp., and a considerable number of other American species, among which *T. serpentina* Wied. and *obliqua* Macq., have already been mentioned above, form a well-defined group, which well deserves to be considered as a separate genus. The character which distinguishes it from all other *Trypetina*, is the course of the fourth longitudinal vein, which, towards its end, is curved forwards in a rather striking manner, and reaches the margin at a very acute angle, being prolonged beyond as the costal vein. With reference to this character I propose to call it *Acrotaxa*. The species of this group have, moreover, the following characters in common: In the structure of the head and of all its parts they resemble the species of *Hexachata*; the thorax has a similar structure, but it is a little smaller in bulk, as compared to the rest of the body, and a little narrower between the roots of the wings; the macrochaetae of the thoracic dorsum agree with those of *Hexachata* both in their number and position. The scutellum is smaller than in the latter genus and not quite as flat, and bears not six, but four macrochaetae. Front femora on the upper side with shorter, on the under side with longer bristles and the front tibiae without bristles, as in the species of *Hexachata*. Middle femora without bristles; only the basis of the under side is sometimes provided with one or several bristle-like hairs; the two rows of bristles which, in *Hexachata*, are found on the under side of the middle femora, are replaced here by two rows of hairs. Middle tibiae without bristles. Hind femora towards the end of their upper side, more or less densely

bristly, on the under side with somewhat longer pile and moreover from the basis to a little beyond the middle, with a rather sparse row of long, almost bristle-like hairs; hind tibiae ciliated with rather weak bristlets. Tegulae almost as much developed as in *Hexachata*. Wings large, and, comparatively to their length, less broad than in *Hexachata*; the venation, with the exception of the difference in the course of the fourth vein, already adverted to, is very like that of *Hexachata*, only all the cells, and especially the stigma, are longer in comparison to their breadth; the posterior angle of the anal cell is drawn out in an equally long and pointed lobe; the whole of the first longitudinal vein and the third some distance beyond the small crossvein, are bristly. The very characteristic picture of the wings in *Acrotoxa* is sufficiently rendered by the figures 5 and 6 of Tab. X, and 19-27 of Tab. XI. The portions of this picture which could not well be called bands (*fasciæ*), or stripes (*vitæ*), I have called *rivulets* (following in this Meigen's example, who called them *rivuli* in latin, and *Bäche* in german). The same term may be applied to the species of *Acidia*. The species of *Acrotoxa* are often very much alike, and very difficult to distinguish in the male sex; the females are frequently easier to distinguish on account of the very different length of the ovipositor in different species.

Observation 3.—In view of the difficulty of this group and of the probable occurrence of species belonging to it in some portions of the North American continent and of the West Indies (besides *Trypeta fraterculus* Wied., already referred to), I deem it useful to enter into a more detailed examination of them. Most of the numerous *Acrotoxæ* occurring in the European museums come from Brazil, and pass rather indiscriminately for the *Dacus parallelus* Wied. I will give a description of this species, based upon the original specimens in the Wiedemann-Winthem and the Seckenberg collections, and of some of the species more closely allied to it, confining myself to those species only which are known in both sexes. Special mentions of coloring and picture will be omitted, as the former is clay-yellow in all the species, and the latter very probably is pretty much like that of *T. ludens*, as given above, at least in living specimens; in drying it becomes somewhat indistinct, and affords no trustworthy marks for discrimination.

a. *T. parallela* WIED. ♂ ♀. (Tab. XI, f. 20.)

Long. corp. 0.37, long. terebræ 0.20—0.21; long. al. 0.40.

Arista with a short pubescence, which is longer, however, than in the following species. The pile on the body in general is somewhat longer than in those species, which is especially perceptible on the abdomen of both sexes and on the ovipositor. Ovipositor slender, not quite as long as the thorax and the rounded abdomen of the female taken together. Wings comparatively broad and very blunt and rounded at the tip; their venation differs from the allied species in the distinct undulation of the second vein and the peculiar bend, which the last section of the third vein shows in the vicinity of the small crossvein; two characters of which there is an indication in *T. consobrina* only. Picture of the wings brownish-yellow, in some places brown, more intense than in the following species; the uninterrupted and even course of the first hyaline space from the basis of the second basal cell to the costa is especially characteristic. The picture of the wings varies sometimes in the fact that both the S-shaped and the V-shaped rivulet each emit, exactly upon the third longitudinal vein, a little pointed projection, almost forming a narrow bridge between them; sometimes the portion of the V-shaped rivulet, cut off by the fourth vein, is filled by a brownish-yellow coloring; I have observed this variety much more often in female than in male specimens.

Hab. Brazil.

b. *T. hamata* n. sp. ♂ ♀. (Tab. XI, f. 22.)

Long. corp. 0.39, long. terebræ 0.26; long. al. 0.41—0.42.

Abdomen short. The ovipositor slender, proportionally somewhat longer than in *T. parallela*. Wings comparatively narrower and less rounded towards the end; second longitudinal vein without any trace of an undulated course and the third longitudinal vein beyond the small crossvein without the curvature so characteristic in *T. parallela*. Picture of the wings paler and more yellow than in the latter species; the branch of the V-shaped rivulet which is more distant from the tip of the wing is prolonged in front beyond the third vein, without diminution of its breadth, so that it coalesces with the S-shaped rivulet between the third and the second vein; the branch of the V-shaped rivulet which is nearer the apex of the wing is either altogether wanting, or its pale yellowish tip only is visible near the posterior margin, as it is represented on Tab. XI, f. 22. The hyaline band running from the basis of the second basal cell towards the costa forms (as it also does in *T. consobrina* and *pseudoparallela*), a row of three contiguous spots. Besides the different picture of the wings, *T. hamata* differs from *T. consobrina* and still more from *T. pseudoparallela* in the shape of the wings, which are comparatively narrower and a little less obtusely rounded at the tip. Moreover, the ovipositor of the female is a little shorter and more slender towards the tip than in *T. consobrina*; but it is very much longer than that of *T. pseudoparallela*.

Hab. Brazil.

c. *T. integra* n. sp. ♂ ♀. (Tab. XI, f. 23.)

Long. corp. 0.41, long. terebræ 0.36—0.37; long. al. 0.42.

The abdomen of this species is longer and narrower than in the other species. The picture of the wings is paler and yellower than that of *Trypeta parallela*, but otherwise resembles it more than any other, as in both, the first hyaline band is not divided in three contiguous spots. However, in the present species this band becomes narrower towards the costa and stops before reaching it, neither of which is the case in *T. parallela*. Moreover, its wings are much narrower and less obtusely rounded at the tip; likewise they show no trace of the wavy course of the second longitudinal vein and of the curvature of the third, which is so well marked in *T. parallela*. The ovipositor is remarkably long in comparison to the size of the body, longer than in all the other species described here. The design of the picture might give rise to the supposition that *T. integra* and *T. obliqua* Macq. are identical. The much smaller size of *T. obliqua* Macq. and the much shorter ovipositor, however, render this impossible. From *T. consobrina* and *pseudoparallela* this species is sufficiently distinguished by the different shape of the first hyaline band of the wings.

Hab. Brazil.

d. *T. consobrina* n. sp. ♂ ♀. (Tab. XI, f.)

Long. corp. 0.31—0.32, long. terebræ 0.26- long. al. 0.38.

Abdomen short. The venation shows more analogy to that of *T. parallela* than to any other species mentioned here, as the third longitudinal vein is somewhat curved beyond the small crossvein; the second longitudinal likewise shows a vestige of a weak undulation (which is not rendered in the figure). The outline of the wings likewise resembles that of *T. parallela* especially in the obtuse rounding of the apex; but the wings are narrower in comparison to their length. The picture of the wings is considerably paler than in *T. parallela*, and resembles in outline that of *T. pseudoparallela*, so that the males of both species may easily be taken for each other, unless attention is paid to the difference in the course of the third vein. The females of both are very easily distinguished, as the ovipositor of *T. consobrina* is considerably longer than that of *T. pseudoparallela*.

Hab. Brazil.

e. *T. pseudoparallela*. ♂ ♀. (Tab. XI, f. 24.)

Long. corp. 0.35, long. terebræ 0.13—0.14; long. al. 0.38—0.39.

The wings resemble those of *T. parallela* in outline very much, differ, however, in the fact that the second and third longitudinal veins do not show the peculiar course which they have in *T. parallela*. The picture of the wings is but little paler than in *T. parallela*, but differs from it considerably in the breaking up of the first hyaline band into three con-

tiguous spots. In speaking of *T. consobrina*, I have adverted to the difference between the males of the two species, which otherwise are closely alike. The female of this species cannot easily be mistaken for that of *T. consobrina* or any other of the species described here.

Hab. Brazil.

The great importance which the comparative length of the ovipositor has for determination of the closely resembling species of the present group, induces me to give here the following figures representing the average of several measurements. The relation of the length of the ovipositor to that of the rest of the body is in *pseudoparallela* 1:2.6; in *parallela* 1:1.8—1.9; in *hamata* 1:1.5; in *consobrina* 1:1.2; in *integra* 1:1.1. Their relation to the length of the wing is in *pseudoparallela* 1:2.8; in *parallela* 1:2.1; in *hamata* 1:1.6; in *consobrina* 1:1.4; in *integra* 1:1.2.

Trypeta Ocesia Walker (List, etc., IV, p. 1016), from Jamaica, is an *Aerotoxa*, closely allied to the species described by me. Whether *Trypeta Acidusa* Walker (ibid., p. 1014) from Jamaica likewise belongs here is uncertain, as the author does not state whether the end of the third longitudinal vein is directed forwards or backwards; moreover there is no statement whatever concerning the shape of the scutellum and the number of its bristles. If this species is an *Aerotoxa*, it cannot possibly be identified with any of those described above, on account of the differences in the coloring. The same applies, in a greater measure still, to *Trypeta serpentina* Wiedemann, already alluded to above.

In order to bring together whatever I know concerning the *Trypetæ* belonging to the group *Aerotoxa*, I give on Tab. XI, f. 26, a copy of the figure of the wing of *Trypeta grandis* Macq. (Dipt. Exot. Suppl., I, p. 212. Tab. XVIII, f. 14), from New Granada, and on Tab. XI, f. 27, that of the wing of *Urophora bivittata* Macq. (Dipt. Exot., II, 3, p. 222. Tab. XXX, f. 7), of unknown *habitat*. Both wings show an outline somewhat different from the other *Aerotoxæ*, more oblique transverse veins, a more narrow first posterior cell, a weaker forward turn of the third vein, etc. I am inclined to believe that these differences do not, for the most part, exist in reality, but are only due to the usual inaccuracy in Macquart's figures; and for this reason I believe that both *T. grandis* Macq. and *Urophora bivittata* Macq. are *Aerotoxæ*. Should my supposition prove correct, then it becomes very probable that America is the *habitat* of the latter species.

7. *T. vulnerata* n. sp. ♂ ♀.—Fusco-nigra, infra fusca; caput exalbidum, fronte et facie ochraceo-vittatis, antennis, palpisque lutescentibus; scutellum subtumidum, setis quatuor praeditum; pedes lutei, femoribus tamen posterioribus fusco-nigris; alae latiusculae, cellula stigmatica brevissima, quadrata, cellula marginali lata et cellula posteriore prima adversus apicem angustata instructae, fasciis nigris inter se coherentibus similiter atque *Aciura lychnidis* F., pictae, colore tamen nigro adversus alarum basim latius diffuse.

Blackish-brown, under side brown; head whitish, front and face with an ochre-yellow longitudinal stripe, antennae and palpi more clay-yellowish; scutellum rather tumid, with four bristles; feet clay-yellowish, the posterior femora, however, brownish-black; wings rather broad, with a short, square stigmatical cell, a broad marginal cell and a first posterior cell, which is attenuated at the posterior end; the black, connected crossbands almost resemble those of *Aciura lychnidis* Fab., but the black coloring is more extended towards the basis. Long. corp., ♂, 0.18, ♀, cum terebra 0.24; long. al. 0.18.

Coloring of a rather shining brownish-black; the humeral region and the under side of thorax and abdomen brown. Head whitish, front and face with a conspicuous ochre-yellow or almost orange-yellow middle stripe. Antennae clay-yellowish, descending below the middle of the perpendicular, very little concave, face; the first two joints with short black pile; the third with an almost sharp anterior corner; arista brownish-black with an extremely short pubescence; oral opening of a medium size; the broad palpi do not extend beyond its anterior edge, which is slightly drawn upwards. Cheeks of a very moderate breadth; at the lower corner of the eye, there is an ochre-brownish spot and a black bristle. The usual frontal bristles black and of a considerable length; between the two black bristles inserted upon the little stripes, coming down from the vertical margin, there is, on each side, a short, white bristle; four similar bristles are inserted upon the posterior vertical margin; the erect pile of the occiput and the cilia of the upper posterior orbit of the eye are white. Thoracic dorsum and pleurae with a very scattered, almost stubble-shaped white pile and black bristles. Scutellum very convex, perceptibly swollen, with four long bristles. The intermediate abdominal segments have a more or less distinct pale coloring on the posterior margin; all segments, with the exception of the posterior one, have some scattered whitish pile towards the posterior margin and blackish pile on the lateral margins; the

last segment, towards its end, has several black bristles. Ovipositor flat, rather broadly truncate, hardly as long as the last three abdominal segments taken together, blackish-brown or black, with black pile. Front feet, as well as the entire fore-coxæ, clay-yellow; on the posterior feet the first joint of the coxæ and the femora are brownish-black, or dark brown, the second joint of the coxæ, the tip of the femora, and the entire tibiæ and tarsi are clay-yellow; the under side of the front femora bears a row of black bristles, while the under side of the posterior femora is without them. Halteres infuscated. Wings large, rather broad, with convex anterior and posterior margins; veins, with the exception of the first longitudinal, without bristles; the first longitudinal vein turns, not very far beyond the end of the auxiliary vein, in a sharp, rectangular fracture, perpendicularly towards the margin of the wing, which causes the stigmatal cell to assume a strikingly short and square shape; the second longitudinal vein is rather distant from the anterior margin of the wing and has a rather straight course, so that the marginal cell, although rather broad, is attenuated towards its end; the third longitudinal vein is turned backwards towards its end, so that the first posterior cell is somewhat attenuated at the end; the small crossvein is placed about the middle of the discal cell, which becomes considerably broader towards its end; the last section of the fourth longitudinal vein has a wavy course; the posterior crossvein is very steep and only very gently curved; the posterior angle of the anal cell is drawn out in a point in the usual way. The picture of the wings has somewhat the appearance of rivulets, and consists of conspicuous and rather well-defined brownish-black crossbands, which come in contact almost in the same way as in the European *Acidia lychnidis* Fab. (compare Loew, *Bohrfliegen*, Tab. III, f. 4); the picture of the present species differs, however, in the more considerable extent of the black coloring on the basis of the wings; the black bands leave two hyaline indentations on the anterior and three on the posterior margin; these hyaline spots have, in a certain light, a whitish reflection. The first of these spots on the anterior margin is a rectangular triangle, the hypotenuse of which begins on the costa a little before the end of the first longitudinal vein and runs as far as the anterior end of the small crossvein; the second hyaline spot, separated from the first by an almost perpendicular

dark band, runs from the costa over the middle of the penultimate section of the fourth vein, as far as the middle breadth of the discal cell. The first hyaline spot of the posterior margin begins at the end of the last longitudinal vein and reaches as far as the fourth vein; the second and third spots begin, as usual in the species with this kind of picture, at the posterior end of the second posterior cell; both are very pointed at their end, and while the second spot reaches only to the fourth vein, the third goes as far as the third vein. Besides these hyaline spots, there is, at the basis, a small hyaline mark, connected with the whitish tegule; in the anal angle of the wing, near the margin, there is a diluted dot. The last of the dark bands is separated from the costa, as far as the third vein, by a narrow, hyaline border; the small crossvein has a similar, very narrow, hyaline border. A peculiar mark of this species is, that the spot at which the second and third longitudinal veins diverge, forms a knot-shaped, blood-red swelling, like a drop of coagulated blood; the first longitudinal vein, near its basis, likewise shows a more or less distinct blood-red coloring.

Hab. Massachusetts (Mr. Sanborn).

Observation.—*Trypeta vulnerata* cannot be well located in any of the genera hitherto formed out of the old genus *Trypeta*. The great resemblance of the picture of its wings to that of *Acidia lychnidis* Fab. (= *discoidea* Meig.), naturally suggests its location in the same genus. A closer examination, however, proves that, although its relationship to the species of that genus is rather close, it differs very much in the structure of the head, the very much more swollen scutellum, the structure of the ovipositor, some details in the venation, and the almost stubble-shaped pile. Thus we are compelled to establish a separate genus, *Stenopa*, for it, which finds its place next to *Acidia*.

S. T. fratria Lw. ♀. (Tab. X, f. 4.)—Lutea, corpore brevi et latiusculo, scutello setas quatuor gerente; alarivnlis luteo-fuscans, maculam ovatam hyalivam in apicali cellulæ discoidalis parte sitam includentibus, apice venæ longitudinalis quartæ non recurvo.

Clay-yellow, stature short and somewhat broad, with four bristles on the scutellum; wings with yellowish-brown rivulets, which inclose an oval, hyaline spot before the end of the discal cell; the end of the fourth longitudinal vein is not curved forwards. Long. corp. 0.22; long. al. 0.22.

SYN. *Trypeta fratria* LOEW, Monographs, etc., I, p. 67. Tab. II, f. 4.

?*Trypeta liogaster* THOMSON, Eug. Resa, p. 578, No. 251.

Hab. United States (Osten-Sacken).

Observation.—I have nothing to add to the description of this species as given in the first part of these Monographs. Its close relationship to the European *T. heraclei* Lin. is a sufficient proof that this species is a true, typical *Acidia*. I believe that *T. liogaster* Thoms. is this same species, although he describes the ovipositor as darker than I find it in my specimen.

9. *T. suavis* Lw. ♂. (Tab. X, f. 10.)—Dilute lutea, corpore brevi, latiusculo, scutello setis quatuor instructo; alarivulis latissimis fuscis, in formam literæ S confluentibus, pictæ, apice venæ longitudinalis quartæ non recurvo.

Pale clay-yellowish, stature short and rather broad, scutellum with four bristles; wings with very broad brown rivulets, which coalesce in the shape of the letter S; the tip of the fourth vein is not curved forwards. Long. corp. 0.20; long. al. 0.21.

SYN. *Trypeta suavis* LOEW, Monographs, etc., I, p. 75. Tab. II, f. 10.

Hab. Middle States (Osten-Sacken).

I possess only one very badly preserved specimen, which I described in the Monographs, etc., Part I. The species is easily distinguished on account of the peculiar picture of its wings. Of all the genera hitherto established in the family *Trypetidæ*, the present species undoubtedly belongs to *Acidia*; and, as far as the imperfect preservation of my specimen allows an opinion, it agrees with the *Acidiæ* in all the important characters, except one: while all the European *Acidiæ* have the third longitudinal vein more or less bristly, I perceive no bristles, whatever, in *T. suavis*, and have no reason to suppose that they have been rubbed off. Such an agreement of characters decides me to place *T. suavis* in the genus *Acidia*; at the same time, however, the bristles of the third longitudinal vein cannot any longer be considered as characteristic of the genus *Acidia*.

10. *T. canadensis* n. sp. ♀.—Dilute lutescens, segmentis abdominalibus tertio et quarto fusco-fasciatis, corpore brevi, latiusculo, terebrâ medioeri, latâ et late truncatâ; alarum rivuli angusti, fusci, apex venæ longitudinalis quartæ non recurvus.

Pale clay-yellowish, with a brown crossband on the third and fourth abdominal segments, stature short and somewhat broad; ovipositor of

medium length, broad and broadly truncate; wings with narrow pale brownish rivulets and with a fourth longitudinal vein which is not curved forwards at the tip. Long. corp. 0.18, cum terebrâ 0.23; long. al. 0.20.

Pale clay-yellowish. The head resembles that of *T. fratria* in shape, only the front is somewhat broader and the vertical diameter of the eyes is a little smaller; the anterior edge of the mouth is more projecting. On the border of the front the described specimen bears, on each side, three long, but rather weak black bristles. Antennæ of a more saturate yellow, not reaching the edge of the mouth; their third joint is rounded at the tip; arista blackish, yellow towards the basis, with a very short pubescence. Rostrum and palpi pale yellow, the latter not reaching beyond the anterior edge of the oral opening. Thoracic dorsum with a very thin, whitish bloom, only the double middle stripe and the narrow lateral stripes not pollinose, rather shining and somewhat darker than their surroundings. The posterior end of the thoracic dorsum and the scutellum likewise without pollen, shining, very pale yellow; a not very broad yellowish stripe runs from the humeral corner to the root of the wings. The scutellum is convex and not very large; in my specimen it has three bristles on one side and only two on the other, so that I cannot say whether the normal number of the bristles of the scutellum is six or four. The bristles of the thorax and of the scutellum, as well as the short pile of the thoracic dorsum, are black. Metathorax distinctly infuscated on its superior margin and its middle line. Abdomen shining, with short black pile; the third and fourth segments have, each at its basis, a chestnut crossband, interrupted upon its middle, while upon the second segment only a lateral beginning of such a stripe is indicated by a chestnut-brown spot. The very broad ovipositor is flat, almost as long as the last three abdominal segments taken together, very broadly truncate and infuscated at the end. The front femora are sparsely beset with bristles upon the upper and under side; the middle femora are entirely without bristles; upon the hind femora, likewise, there are only a few bristle-like hairs before the end of the upper side; the upper side of the hind tibiæ is merely beset with exceedingly short bristle-like hairs. Wings of the usual shape, hyaline, with a pale-brown picture; it consists: 1. In an oblique half crossband running from the humeral crossvein

to the basis of the second basal cell; 2. Of a crossband parallel to the first, abbreviated behind, which begins at the stigma, near the anterior margin, and runs across the basis of the submarginal cell, as well as across the crossveins, which close the second and third basal cells, and thus reaches the sixth longitudinal vein; 3. Of a rivulet which begins above the posterior crossvein, near the third longitudinal vein, runs from it across the posterior crossvein as far as the posterior margin, is continued along this margin inside of the third posterior cell, but, before reaching the sixth longitudinal vein, is suddenly turned upwards, running parallel to the band which begins at the stigma, crossing the small crossvein, and thus reaching the anterior margin, where, gradually expanding, it forms a border ending a little beyond the tip of the fourth crossvein. The two crossbands, as well as the rivulet, are of moderate breadth only; the latter has, in the described specimen, the following faded spots, which, in more fully colored specimens, are probably less apparent or altogether absent: 1. A rounded spot in the marginal cell, above the origin of the rivulet; 2. Upon the longitudinal axis of the submarginal cell an indentation in the inner margin of the section bordering the apex of the wing; 3. Upon the longitudinal axis of the first posterior cell an interruption of the rivulet at its origin and an indentation in the inner margin of the portion bordering the apex of the wing; 4. Upon the longitudinal axis of the discal cell a narrow interruption of the section, running again towards the anterior margin; 5. The spot upon the posterior margin connects the first, descending, portion, with the second, which rises again upwards. The first and third longitudinal veins are bristly; the third and fourth are parallel towards their end, both very gently curved backwards; the section of the fourth vein preceding the discal cell is gently, but rather distinctly arcuated backwards, so that the shape of the discal cell somewhat reminds of that of the species of *Rivellia*; the crossveins are comparatively rather long, moderately approximated, their distance being about equal to the length of the posterior crossvein; the latter is rather steep, however, perceptibly approximated to the apex with its anterior end, more than with the posterior; the posterior corner of the anal cell is very much drawn out in a point.

Hab. Canada (Mr. Provancher). [Norway, Maine; S. J. Smith—seems to be a common species in those regions. O. S.]

Observation.—*Trypeta canadensis* resembles the species of *Acidia* in its general habitus and, at first sight, seems to differ only in the somewhat modified picture of the wings, which seems to hold the middle between the rivulet and the crossband. A closer examination shows, that in the structure of the head and of its parts, as also in the bristles upon the feet, this species is closely allied to *Acidia*, but that it also shows characters not belonging to that genus; such is the structure of the ovipositor, which is longer, quite flattened, and broadly truncate at the end; also the very peculiar course of the section of the fourth longitudinal vein preceding the small crossvein. If the scutellum is provided with six bristles in normal specimens, we would have another important distinctive character from *Acidia*. Thus the admission of *T. canadensis* in the genus *Acidia* would render the limitation of this genus too indefinite, and it becomes necessary to establish a new genus for it, which would be characterized by a modified type of the picture, a peculiar course of the fourth vein, and a different structure of the ovipositor. I will call this genus *Epochre*.

11. *T. longipennis* WIED. ♂ ♀. (Tab. X, f. 2 ♂, 3 ♀.)—Lutea capite tumido, corpore elongato et angusto; alæ longæ et angustæ, maris adhuc longiores et angustiores quam feminæ, rivulis luteo-fuscis pictæ.

Clay-yellow; head tumid; body long and narrow; wings long and narrow, those of the male still longer and more narrow than in the female, pictured with yellowish-brown rivulets. Long. corp. 0.17—0.26; long. al. 0.22—0.30.

SYN. *Trypeta longipennis* WIEDEMANN, Auss. Zweifl., II, 483, 12 (♂ ♀).

Strauzia armata R. DESVOIDY, Myod. 719, 2 (♂).

Strauzia inermis R. DESVOIDY, Myod. 718, 1 (♀).

Tephritis trimaculata MACQUART, Dipt. Exot., II, 3, p. 226, 8. Tab. XXXI, f. 3.

Trypeta cornigera WALKER, List Brit. Mus., IV, p. 1010.

Trypeta cornifera WALKER, List Brit. Mus., IV, p. 1011.

Trypeta longipennis LOEW, Monographs, etc., I, p. 65.

It cannot be doubted that *Trypeta longipennis* Wied., either is a very variable species, or that North America possesses a number of closely allied species, resembling it very much, and which, as long as they are represented only by single, often imperfectly preserved specimens, it is as difficult to distinguish

and to describe as, for instance, the majority of the European *Urophora*. It is only by observations upon the insect in life, that the question will probably have to be solved, whether we have here different species or only varieties. In writing the first part of these Monographs I surmised that I had specimens of a single, but very variable species before me. In the mean time my materials have increased considerably, and specimens have been added to it, which differ so materially from the typical *U. longipennis*, that my former conviction has been shaken, without, however, having been superseded by the opposite one. I prefer therefore to continue to treat these different forms as varieties of the same species, but, at the same time, to define these varieties with more precision than has been done in the first part of the Monographs. In order to avoid useless repetitions, I will notice in advance that in all the varieties the anterior end of the middle stripe is colored black, and that in all of them, immediately above the root of the wing, there is a small, deep-black dot, which is not visible when the wings are folded.

1. *Varietas perfecta*, ♂ ♀.—Of the four lateral bristles of the front, the two upper ones, in the male, are very much incrassated and truncated at the end. Thorax without black lateral stripes. Scutellum unicolorous; metathorax without black picture. Picture of the wings not very deep in its coloring, complete in both sexes; the male as Tab. X, f. 2.

Of this variety I have compared rather numerous specimens. Among those of my collection there is a male and two females, caught at the same time.

2. *Varietas typica* ♂ ♀.—Of the four lateral bristles on the front the two upper ones are very much incrassated in the male and truncate at the end. Thorax without black lateral stripes; scutellum upon each lateral corner with a well-defined black spot. Metathorax without any black coloring. The picture of the wings is of a rather dark shade, especially towards the tip; complete in the female, incomplete in the male, almost like Tab. X, f. 2, except that the rivulet covering the posterior crossvein does not reach the margin of the wing, but gradually becomes more attenuated and pointed and never reaches beyond the posterior end of the posterior crossvein; the branch of this rivulet which runs along the last section of the fourth vein is likewise very narrow and always disappears at a considerable distance from

the margin of the wing; the hyaline interval between it and the branch bordering the anterior margin of the wing is, in the male, comparatively longer and conspicuously narrower than in var. *perfecta*; the female shows the same difference, but very feebly.

Wiedemann's description is based upon specimens of this variety, which is a very common one. The other synonyms, quoted above, likewise belong here, with the only exception of *Trypeta cornigera* Walker. I possess of this variety four perfectly well-preserved specimens (a male and three females), caught at the same time by Mr. Auxer in Lancaster City, Penn.; the three females have, at the posterior end of the two posterior abdominal segments, longer, stronger, and somewhat more abundant pile than the females of other varieties.

3. *Varietas longitudinalis* ♂ ♀.—Of the four lateral bristles of the front the two uppermost, in the male, are very much incrassated and truncated at the end. Thorax without any black lateral stripes; scutellum on each lateral corner with a black spot; metathorax without black picture. The wings of the male comparatively narrower than in all the other varieties; their picture coalesces into a single broad longitudinal stripe, which, from the root of the wing as far as nearly the end of the posterior basal cells, has a dirty clay-yellowish coloring; beyond this point, it changes into dark-brownish. The interval between the second and fourth longitudinal veins is completely filled by this stripe, with the only exception of a small hyaline spot at the end of the fourth longitudinal vein; moreover, the stripe encroaches a little beyond the second and fourth veins in the shape of little wavy expansions. The picture of the female hardly differs from that of var. *typica*; only the spot in the costal cell, between the stigma and the humeral crossvein, which is usually wanting in var. *perfecta* and present in var. *typica*, is much darker than in the latter species; this is also the case in the male.

These statements are taken from a very fine pair of specimens from Sharon Springs, N. Y., collected by Baron Osten-Sacken. He sent me at the same time a male from Connecticut (collected by Mr. Bassett), which agrees with the former in the picture and in the shape of the wings, except that the uniformly brown part of the picture of the specimen from Sharon is clouded with yellowish-brown and dark-brown; moreover, in the latter specimen, the spot placed between the humeral crossvein and the stigma is

very much faded. The description which Mr. Walker gives of his *Trypeta cornigera* refers, if I understand it right, to this variety. [The male specimen from Sharon was caught on the same spot with the female; I possess, moreover, a couple from Connecticut, stuck on one pin, as if caught *in copula*. Thus there can hardly be a doubt as to the sexes belonging together, the very different picture notwithstanding. O. S.]

4. *Varietas vittigera*, ♂ ♀.—Of the four lateral bristles of the front, the two upper ones are very much incrassated and truncate at the end. The thoracic dorsum shows, besides the anterior end of the middle stripe, two well-marked black lateral stripes of a moderate breadth, abbreviated in front, rather broadly interrupted at the transverse suture and pointed posteriorly. Scutellum, upon each lateral corner, with a black spot; metathorax on each side with a deep black longitudinal spot. Wings of the male somewhat less elongated than in the male of the var. *perfecta*. The picture of the wings in both sexes is complete, hardly different from that of var. *perfecta*. Of this variety I possess only a male and a female from Nebraska (Dr. Heyden).

5. *Varietas intermedia* ♂.—Of the four lateral bristles of the front, the two superior ones, although strong, are not incrassated and not truncate at the tip, but end, as usual, in a point. Thorax without black lateral stripes; scutellum upon each lateral corner with a black spot; metathorax on each side with a deep black longitudinal spot. Wings of the male less elongated, and perceptibly less pointed than in the first two varieties; the picture of the wings rather intense in coloring, the design resembling that of the female of the first variety; however, the hyaline band passing between the two crossveins is rather conspicuously expanded at its posterior end. The last joint of all the feet is rather conspicuously infuscated on the sides and at its end. Of this variety I possess only a single male, without indication of the precise locality.

6. *Varietas confluens*, ♂.—Of the four lateral bristles of the front the two upper ones are rather strong, but not incrassated and not truncate at the tip, but end, as usual, in a point. The thoracic dorsum, besides the anterior end of the middle stripe, shows two well-defined black lateral stripes of a moderate breadth, which are abbreviated anteriorly, rather broadly interrupted at the transverse suture, and pointed posteriorly. Scutellum upon

each lateral corner with a black spot. Metathorax on each side with a deep-black longitudinal spot. Wings of the male comparatively less elongated, and less attenuated towards the tip, consequently comparatively broader than in the first and second variety. The picture of the wings is complete, its coloring uniform, not very saturate, seldom here and there with a trace of darker margins; the oblique hyaline crossband passing between the crossveins is comparatively narrow, reaches, however, the anterior margin completely. The brownish-yellow rivulet rising across the posterior crossvein is of a considerable breadth in all its parts, so that the branch of it which borders the margin of the wing and that which runs along the last section of the fourth longitudinal vein, coalesce in their middle.

I possess a single male only (Connecticut; Mr. Norton); it is one of the smallest specimens of this species in my collection.

7. *Varietas arcuata* ♂.—Of the four lateral bristles of the front, the two upper ones are not stronger than usual among the species of the same size; as usual, also, they end in a point. The thoracic dorsum shows, besides the anterior end of the middle stripe, two strongly marked black lateral stripes of moderate breadth, which are abbreviated anteriorly, rather broadly interrupted at the transverse suture, and end in a point posteriorly; scutellum with a black spot upon each lateral corner; metathorax on each side with a deep black longitudinal spot. The wings of the male are less attenuated towards the apex than in the males of the first and second varieties, but comparatively less broad than in the sixth variety. The picture of the wings has a rather uniform yellowish-brown coloring. It differs from that of all the other varieties in the fact that the oblique hyaline band, running between the two crossveins, does not reach the anterior margin, but suddenly ends between the second and the third longitudinal veins, so that the border of the anterior margin is not at all interrupted beyond the triangular hyaline spot near the stigma; at the same time, this hyaline band is connected with the hyaline streak in the latter portion of the first posterior cell, the rivulet crossing over the posterior crossvein being interrupted here. These modifications give the picture a very different appearance.

Of this variety I likewise possess but one specimen (Illinois; Mr. Brendel); it is but little larger than the male specimen of the sixth variety.

Observation.—*Trypeta longipennis* has no immediate relatives among the European *Trypetidæ*. From *Spilographa abrotani* Meig., and *macrochaeta* Lw., which resemble it somewhat in the peculiar shape of the frontal bristles, it differs too much, in the stature of the body, the shape of the head, as well as in the outline, the venation, and the picture of the wings, to be placed in the same genus *Spilographa*. It must be considered, therefore, as the type of a separate genus. Mr. R. Desvoidy has given it the name of *Strauzia*, which may be preserved, after being modified into the more correct form of *Straussia*. The principal characters of the genus *Straussia* are the following:—

Body long and narrow; head remarkably swollen, especially the occiput; eyes rounded and rather small for a *Trypeta*, so that in the profile the front advances much before the eyes and the cheeks are very broad. Lateral border of the front raised in the shape of a cushion, so that the whole front assumes the appearance of a basin. Antennæ short, reaching, perhaps, as far as the middle of the face; the last joint rounded at the tip. Face retreating inferiorly; oral opening small, without any sharp anterior edge; the rather broad palpi not reaching beyond this edge. Scutellum convex, with four bristles. Abdomen elongated and considerably narrower than the thorax. Ovipositor of the female not flattened. Wings comparatively long and only moderately broad, in the male narrower than in the female, especially towards their end; the picture consists of rivulets; first and third longitudinal veins distinctly bristly; the third and fourth veins towards their end somewhat divergent and rather strongly bent backwards; small crossvein placed about the beginning of the last third of the discal cell; the posterior angle of the anal cell is drawn out in a sharp point.

12. T. electa SAY. ♀. (Tab. X, f. 7.)—Lutea, vittis thoracis et scutello ex-albidis, angulis lateralibus hujus nigris; tibiæ posticæ setis nigris, proportione longis ciliatæ; alæ hyalinæ, fasciis duabus integris adversus marginem posticum convergentibus, strigulâ interjectâ a costâ ad venam longitudinalem tertiam ductâ, et costæ ipsius limbo inle a fasciâ secundâ usque ad apicem cellulæ posterioris secundæ pertinente, fuscis.

Clay-yellow, longitudinal stripes of the thorax and scutellum whitish; the latter with blackish lateral corners; posterior tibiæ ciliated with comparatively long black bristles; wings hyaline, with two complete

crossbands, converging towards the posterior margin, an incomplete band beginning at the anterior margin and running as far as the third longitudinal vein, and a border of the costa, beginning at the second crossband and ending at the tip of the second posterior cell; the whole of this picture being brown. Long. corp. 0.29; long. al. 0.29.

SYN. *Trypeta electa* SAY, Journ. Acad. Phil., VI, p. 185, 1. .

Trypeta electa LOEW, Monographs, etc., I, p. 71, 6. Tab. II, f. 7.

I have nothing to add to the description, given in the first part of these Monographs, but I must observe that, deceived by Macquart's insufficient description of his *T. flavonotata*, I have taken it to be merely a paler variety of *T. electa* Say, while a specimen received since then has convinced me that it is a very closely allied but distinct species.

Hab. Florida (Osten-Sacken).

Observation.—*Trypeta electa* belongs in the genus *Spilographa*.

13. *T. flavonotata* MACQ. ♂.—Lutea, vittis thoracis et scutello unicolore pallidioribus, tibiæ posticæ setulis brevissimis pallidis subciliatis; alæ hyalinæ, fasciis duabus postice paulo abbreviatis adversus marginem posticum convergentibus, strigulâ interjectâ a costâ prope ad venam longitudinalem tertiam ductâ et costæ ipsius limbo inde a fasciâ secundâ usque ad cellulæ posterioris secundæ apicem pertinente, fuscis.

Clay-yellow, longitudinal stripes of the thoracic dorsum and the unicolorous scutellum paler; hind tibiæ somewhat ciliated with very short, pale bristles; wings hyaline, with two crossbands, which are somewhat convergent posteriorly and interrupted a little before the posterior margin, a little crossband between them, extending from the anterior margin almost to the third longitudinal vein, and a border of the costa, running from the second crossband to the tip of the second posterior cell; the whole picture being brown. Long. corp. 0.18; long. al. 0.21—0.22.

SYN. *Tephritis flavonotata* Macq. Dipt. Exot. Suppl. V, p. 125. Tab. VII, f. 9.

This species is very like *Trypeta electa* Say, differs, however, from it as follows. It is smaller; the head is comparatively smaller and has much narrower checks. The third antennal joint ends at a much sharper angle. In what way the picture of the thorax differs from that of *T. electa* cannot be well ascertained in my specimen, in which it has become somewhat indistinct, probably in the process of drying; the whitish stripe, running from the humerus to the root of the wings, is very per-

ceptible; there is also a trace of the whitish stripe above the root of the wings; but this stripe shows no trace of the dark border on the inside, which it has in *T. electa*; nor do I see a whitish median line. Scutellum comparatively smaller and somewhat more convex, without black spot on the lateral corners. The upper border of the metathorax is marked, at each end, with a very small spot of a deep black color. The punctiform black lateral dots, which exist on the last abdominal segment of the female of *T. electa*, are not perceptible in the male of the present species. All the bristles of the body are less strong and of a paler color, especially upon the femora, and instead of the comparatively long black bristles with which the upper side of the hind tibiae of *T. electa* is fringed, there are in the present species only very short pale yellow bristlets. The third longitudinal vein of the wings has, at its basis, several little bristles, but upon the remainder of its course, is entirely bare (while the bristles extend much farther in *T. electa*). The picture of the wings is very like that of *T. electa*, with the following differences: the two crossbands in the middle of the wings do not altogether reach the posterior margin and are also less approximated, that is, they do not form the figure V; the basal portion of the submarginal cell lying before the first of these bands is hyaline; the picture in the vicinity of the root of the wing is much less extended and much paler, so that its darker portions do not, as in *T. electa*, form a kind of crossband, running almost parallel to the following band.

Hab. Yukon River, Alaska (R. Kennicott).

Observation.—*T. flavonotata* is very closely allied to those European species, which I have placed in the genus *Zonosema* (in my Monograph of the European *Trypetidae*), and should be placed in it, as long as it is separated from *Spilograpta*. Should, however, *Zonosema* be united with *Spilograpta*, which seems the best course to follow, owing to the intermediate forms, which occur among the exotic species, then, as a matter of course, *T. flavonotata* will have to be placed in the genus *Spilograpta*.

14. *T. tetanops* n. sp. ♂. (Tab. XI, f. 15.)—Mellea, capite subinflato, oculis parvis; alae hyalinae, fasciis duabus adversus marginem posticum convergentibus, strigula interjecta inde a costa ad tertiam usque venam pertinente, maculis denique duabus parvis, altera in vena longitudinalis tertiae, altera in quartae apice sita, fuscis, his maculis limbo marginis tenuissimo fusco conjunctis.

Honey-yellow, with a rather tumid head and small eyes; wings hyaline, with a brown picture, which consists of two crossbands, converging towards the posterior margin, of a little band, between both, reaching from the costa to the third longitudinal vein and of two little spots upon the third and fourth longitudinal veins, which spots are connected by a narrow infuscation along the margin of the wing. Long. corp. 0.19—0.20; long. al. 0.17—0.18.

Honey-yellow, the head of a purer yellow, somewhat tumid. Front broad, with some scattered, short, very delicate blackish pile; its lateral bristles weak. Frontal lunule very small. Eyes small, elongated, with a rather projecting anterior corner. Face descending straight; edge of the mouth blunt, somewhat swollen; the conspicuously deepened antennal furrows become narrowed below and disappear in the lateral edges of the mouth; the part of the face between them forms an acute, level triangle; the cheeks are remarkably broad, beset with a few short black hairs; oral opening very small; clypeus unusually little developed; palpi short, but considerably broad, sparsely beset with short, black hairs. Proboscis rather short and stout; the stout suctorial flaps, although somewhat long, are not prolonged, nor folded backwards. The upper side of the thorax, with the exception of the posterior and lateral margins, which are shining, is covered with a thin ochre-yellow pollen, and hence opaque; the short pile upon it and the bristles are black; the number and position of the latter is the usual one; of the two pairs of bristles in front of the scutellum, the anterior one is inserted upon very small dots of a somewhat darker color; in the proximity of the suture there are two similar dots; moreover, the trace of a slender dark middle line is perceptible. Scutellum shining honey-yellow, rather convex, sparsely beset with little black hairs and bearing four strong black bristles. Pleuræ of the same color with the scutellum, beset with black pile. Abdomen, likewise, shining honey-yellow, in the middle with a trace of an ochre-yellow dust, beset with black pile, but without longer bristles. The yellow feet have rather strong femora; the two front femora are beset with bristles upon the under and upper side. Wings hyaline, with a picture which is very like that of the two preceding species. The principal feature consists in two narrow brown transverse bands; the first, somewhat faded at its beginning, starts from the end of the stigma and runs perpendicularly over the small crossvein as far as the proximity of the posterior margin, while

the second begins at the tip of the second longitudinal vein and runs in an oblique direction over the posterior crossvein to the posterior margin; between these two bands there is a short, brown one, extending from the anterior margin to the third longitudinal vein; it follows the same direction as the second band; the tips of the third and fourth longitudinal veins bear each a small brown spot and these spots are connected by a narrow brown shade along the margin of the wing; a small brown spot covers the end of the anal cell, which is drawn out in an acute point; the inner costal cell, the beginning of the first basal cell, as far as the origin of the third vein, the basis of the submarginal cell as far as the first brown crossband, the stigma and the anal cell are tinged with yellow; a yellow coloring likewise surrounds that crossvein, which divides the second basal cell from the discal cell; the basis of the exterior costal cell is tinged with yellowish-brown. The third longitudinal vein is, in the vicinity of its origin, densely beset with bristles; more sparsely beyond that point; the third and fourth longitudinal veins somewhat diverge towards their end; the small crossvein is a little before the middle of the discal cell; the posterior crossvein is straight and steep.

Hab. Mexico (Deppe; Mus. Berol.).

Observation.—The principal difference between this species and the typical *Spilographæ* consists in the structure of the head, which has been described above; moreover, the wings are comparatively shorter and the third vein has, as far as its tip, an entirely rectilinear course, while, in all the species of *Spilographa* (comp. Tab. X, f. 7), it is gently curved backwards. Should a new genus be founded for this single species, the name *Edicarena*, alluding to the structure of its head, might be adopted for it. It would seem preferable, however, until a number of allied species becomes known, to let *T. tetanops* remain in the genus *Spilographa*, with which it is undoubtedly related on account of the great resemblance of the picture of its wings with that of *T. electa* and still more of *T. flavonotata*.

15. *T. sarcinata* Lw. ♀. (Tab. XI, f. 16.)—Sordide lutea, dorso thoracis circerscente, punctisque aliquot majusculis atris picto, scutello tumido, binnammato atro, alarum angulo axillari fascisque quatuor valde obliquis ex luteo fuscis, venis transversis obliquis et valde approximatis, cellulâ discoidali adversus basium valde angustatâ.

Dingy clay-yellow, with several deep black dots upon the gray thoracic dorsum and with a tumid bituberculate black scutellum; wings with a yellowish-brown posterior angle and four very oblique yellowish-brown crossbands, with oblique and very approximate crossveins and with a discal cell which is gradually attenuated towards its basis. Long. corp. 0.28; long. al. 0.26—0.27.

SYN. ? *Tephritis quadrifasciata* MACQUART, Dipt. Exot. II, 3, p. 226. Tab. XXX, f. 8.

Trypeta sarcinata LOEW, Berl. Entom. Zeitschr., VI, p. 218, and Dipt. Amer. Cent., I, 88.

Dark clay-yellow, almost brownish-yellow. The broad head is of a lighter color; front very broad, on the anterior part of the lateral margin with two bristles, and before them, near the orbit, with a small black dot. Antennæ yellowish, by far not reaching the edge of the mouth. Face somewhat excavated, but very little protruding towards the edge of the mouth, broad and with broad orbits along the eyes. Cheeks rather broad, with a small black spot near the lower corner of the eye. Oral opening transversely oval; proboscis and palpi yellowish, short, entirely withdrawn in the oral opening; the usual frontal bristles black; the pile on the cheeks, below the black dot which occurs upon them, blackish; the remaining pile on the head is whitish. The upper side of the thorax seems to have an almost black ground color, assumes, however, in consequence of the rather thick pollen which covers it, a gray, entirely opaque, appearance; upon the middle of the thorax, lengthways, there are three pairs of large, black, opaque dots, the largest, anterior pair being on the transverse suture, the posterior pair immediately in front of the scutellum; upon the lateral margin of the thoracic dorsum, the humeral callus, the callus in front of the root of the wings, and a rather large spot above the root of the wings are not clothed with pollen and rather shining black. The ordinary bristles are black; the bristles in pairs, along the thoracic dorsum, are inserted upon the black dots, described above, except upon the anterior pair (where they may have been rubbed off in the described specimen). Scutellum shining black, remarkably swollen, but with a strong coarctation along the longitudinal middle line, and thus appearing bituberculate; each of the tubercles bears a strong bristle, below which a second one, much weaker, seems to have existed. Metathorax and pleuræ clay-yellow; the immaculate, glabrous abdomen is of the same color. Ovipositor flat, pointed, somewhat longer

than the last four abdominal segments taken together, of the same color with the abdomen, or somewhat more reddish-yellow, black at the extreme tip only, with scattered, blackish pile. Feet dark clay-yellow. Wings rather large; their picture consists, besides the yellowish-brown posterior corner, of four oblique yellowish-brown crossbands, with dark-brown borders; the brown coloring which fills the posterior corner is separated from the first band on the posterior half of the wing only, and that by an oblique hyaline half band, lying in the third posterior cell, but which does not reach the root of this cell; a small, square hyaline spot near the humeral crossvein indicates the separation of the yellowish-brown coloring of the base of the wing from the first crossband; the first and second crossbands are completely coalescent before the third longitudinal vein; beyond this vein, they are separated by a hyaline, very oblique band, which begins below the basis of the comparatively long stigma and ends at the tip of the fifth vein; the second and third brown bands are separated by a narrow hyaline band, which crosses the whole breadth of the wing, but is almost interrupted upon the second longitudinal vein; the third and fourth brown bands, the latter of which runs along the apex of the wing, are entirely coalescent upon their anterior portion; their posterior portion is separated by a narrow, hyaline, half band, which does not reach the third longitudinal vein; upon the last section of the anterior margin the brown coloring is somewhat spotted and shows here and there a very small pale drop. The venation shows the following peculiarities; stigma rather long, third and fourth longitudinal veins curved backwards towards their end; the very approximate crossveins are very oblique and have their posterior ends nearer to the apex of the wing than the anterior ends; the discal cell is very much contracted towards the basis, and very much dilated towards the end; the posterior angle of the anal cell is drawn out in a sharp point; the third vein has scattered bristles upon nearly its whole extent.

Hab. South Carolina (Zimmerman; Mus. Berol.).

Observation 1.—In the synonymy, I have doubtfully quoted *Tephritis quadrifasciata* Macq. from Georgia. It is true that *Trypeta varcinata* is not recognizable in Macquart's description; and if Macquart's figures had the least claim to faithfulness, the synonymy of these two species would be out of question. But

with the knowledge we have of the character of Macquart's publications, we cannot but suspect that his species is after all nothing but the one we have described above. The position and direction of the crossveins, as well as the general pattern of the picture of the wings, distinctly show a certain analogy to *T. sarcinata*. The synonymy cannot be assumed as certain, as Macquart, in his description, does not mention either the black dots on the thoracic dorsum, or the black coloring and the very striking shape of the scutellum of *T. sarcinata*; moreover his figure of the wing shows important discrepancies in outline, venation, and picture. By all means, should even the identity of these species be confirmed, Macquart's name would be lost for it, as it has been preoccupied by Meigen.

Observation 2.—The great approximation of the crossveins and their oblique position indicate the relationship of the present species with *Edaspis*. It differs, however, in the peculiar shape of the scutellum, the greater length of the wings, and the shape of the discal cell, which is more attenuated towards the basis. The pattern of the picture of the wings differs from that of the European and American species, as far as they are known. For this reason, I do not think that it would be well placed in the genus *Edaspis*, and I propose for it the formation of a new genus, *Peronyma*. The position and direction of the crossveins, as well as the picture of the wings (the second crossband of which, as in *Edaspis*, incloses both crossveins), remind of *Trypeta obliqua* Say and the species related to it; however, the structure of these latter species has too little in common with *T. sarcinata* to allow their juxtaposition in the same genus.

16. *T. discolor* Lw. ♂. (Tab. X, f. 1.)—Lutea, abdomine nigro. alarum fasciis quatuor obliquis fuscis, primâ et secundâ antice, tertiâ et quartâ postice connexis, venâ longitudinali tertiâ setosâ, venisque transversis valde approximatis.

Clay-yellow, with a black abdomen; wings with four oblique infuscated bands, the first and second of which are connected anteriorly, the third and fourth posteriorly; the third longitudinal vein is beset with bristles; crossveins very much approximated. Long. corp. 0.13; long. al. 0.15.

Syn. *Trypeta discolor* Loew, Monogr., I, p. 64. Tab. II, f. 1.

Hab. Cuba.

This pretty species is so closely allied to *T. obliqua* Say, that

generically they cannot be separated; the systematic position of these two species and of some South American ones, related to them, will be discussed below (see the last observation to the next following species).

17. *T. obliqua* SAY. ♂ ♀. (Tab. XI, f. 14.)—Flava, thoracis dorso postice atro-bipunctato, abdomine maris utrinque punctis atris in seriea dispositis quatuor, fœminæ quinque notato, alæ hyalinæ, fasciis quatuor obliquis flavis et fusco-marginatis variegatæ.

Yellow, with two deep-black punctiform dots on the posterior end of the thoracic dorsum, and on each side of the abdomen with rows of four similar dots in the male, and of five in the female; wings hyaline with four oblique, yellow crossbands, bordered with brown. Long. corp. 0.12—0.14; long. al. 0.13—0.14.

Syn. *Trypeta obliqua* SAY, Journ. Acad. Phil., VI, p. 186, 3.

Trypeta obliqua LOEW, Monogr., I, p. 99.

Say's description, with the additions given by Baron Osten-Sacken in these Monographs, Vol. I, p. 100, is sufficient for the identification of this pretty species. I would only add that in all the specimens examined by me, the males had four, the females five black dots on each side of the abdomen, and that all the specimens showed three deep black dots on the posterior part of the pleuræ; one immediately above the middle coxæ, the second above the hind ones, the third crescent-shaped, surrounding the basis of the stem of the halteres. Ovipositor about as long as the last two abdominal segments taken together, of the same coloring as the abdomen, very little infuscated at the end.

Hab. Indiana (Say); Pennsylvania (Osten-Sacken; on *Vernonia* in August); Texas (Belfrage).

Observation 1.—I am in doubt whether *Trypeta obliqua* also occurs in Brazil. The specimens generally labelled with this name in the collections, seem to belong to a different, although closely resembling species. They are usually somewhat larger than the North American specimens of *T. obliqua* Say; the pile on the whole body as well as the bristles on the third vein are somewhat longer; moreover, I notice on the sides of the abdomen of the male only two, of the female only three black dots; not fully colored specimens do not show any trace of the three black spots on the posterior part of the pleuræ, as they occur in *T. obliqua*; better colored specimens have a trace of the two posterior spots

only. In all other respects the agreement with *T. obliqua* is so great, that I do not dare to decide whether this Brazilian *Trypeta* is a distinct species or merely a variety of *T. obliqua*. It is not to be confounded with another Brazilian species, which is considerably larger, and of which I possess only the female. I let its description follow:—

T. biseriata n. sp. ♀.—*Trypetæ obliquæ* Say, quam magnitudine superat, simillima, sed capite proportione majore, pilis totius corporis longioribus, pleuris immaculatis, alis minus pure hyalinis et cellulâ basali secundâ non hyalinâ, sed luteâ distincta.

Very like *T. obliqua* Say, but larger, with a comparatively larger head, longer pile on the whole body and unspotted pleuræ; wings of a less pure hyaline; second basal cell not colorless, but yellow. Long. corp. 0.17—0.18; long. al. 0.22—0.23.

Coloring and picture of the body similar to the female of *T. obliqua* Say, especially the two black dots upon the posterior portion of the thoracic dorsum and the five black dots upon each side of the abdomen; the black dots which *T. obliqua* has on the posterior portion of the pleuræ are entirely wanting here. The pile on the whole body is much longer, black upon the abdomen and especially striking upon the posterior edge of its first segment. The head is proportionally larger. The wings are comparatively somewhat broader and their surface, especially towards the posterior margin, is a little more dusky; the first and third longitudinal veins are beset with much longer bristles; the venation agrees, in the main, with that of *T. obliqua*; the picture of the wings also is very much alike, only the dark portions of it are less brownish-black and more diluted; the last two yellow bands are much less extensive; the second basal cell, which in *T. obliqua* is always hyaline, is altogether tinged with clay-yellow here. The ovipositor is about as long as the last two abdominal segments taken together, and is broadly truncate at the end.

Hab. Brazil.

Observation 2.—*Trypeta discolor* and *obliqua* Say, as well as the *T. biseriata* described in the preceding observation, are three very closely resembling species, agreeing in all the principal characters. They have no immediate relatives in Europe, with which they could be placed in the same genus; however, they are somewhat allied to *Ædaspis*, as they have the direction of the crossveins and the course of the second crossband, covering the crossveins, in common with that group; in almost all the other important characters they show striking differences. I propose, therefore, the formation of a new genus for them, which I call *Pla-*

giotoma. The characters of this genus are as follows: In the structure of the head and of its parts and of the scutellum it resembles *Acidia* very much; the scutellum, provided with four bristles, is convex, without appearing swollen; the shape of the abdomen likewise reminds one of the species of *Acidia*; the ovipositor also has a similar structure, but is longer than in *Acidia*, rather broadly truncate at the end. Wings rather large, with a distinctly convex anterior margin; the first and third veins are distinctly bristly; the crossveins are very much approximated; their posterior end is nearer to the apex of the wing than the anterior one; the last section of the fourth vein forms a bow, the convex side of which is turned towards the anterior margin, so that it distinctly diverges at the end from the end of the third vein, which is much more straight; the posterior corner of the anal cell is drawn out in an acute point. The picture of the wings consists of four very oblique crossbands, the second of which runs over both crossveins; the last crossband forms a border along the apex of the wing.

18. *T. palposa* Lw. ♂. (Tab. X, f. 9.)—Lutea, abdomine puncturum nigrorum seriebus quatuor picto; alæ hyalinæ, fasciis tribus sordide luteis, primâ et secundâ perpendicularibus et parallelis, tertiâ marginali et inde a præcedente usque ad cellulæ posterioris secundæ apicem pertinente.

Clay-yellow, with four longitudinal rows of black dots on the abdomen; wings hyaline with three crossbands of a dingy clay-yellow, the first two of which are perpendicular and parallel; the third forms a border along the margin of the wing, reaching from the second band to the end of the second posterior cell. Long. corp. 0.26—0.27; long. al. 0.26.

Syn. *Trypeta palposa* Loew, Monogr. I, p. 74, 8. Tab. II, f. 9.

The quoted description, drawn from an indifferently preserved male, is sufficient for the identification of the species. I will only notice here that in the first line of that description, *Cederh.*, must be read, instead of *Cederli*, and that on page 75, line 4, the expression "the edge of the tip" means the third band, which forms a border along the last portion of the anterior margin and the apex of the wing.

Hab. Northern Wisconsin River (Kennicott).

Observation.—The present species is a type of the genus *Trypeta*, in the narrower sense, as defined in my Monograph of the European *Trypetina*. It belongs in the group of those

species which are related to *Trypeta arctii* Deg. and are abundantly represented in the European fauna. The most salient features of *Trypeta* sensu stricto are also the shape of the head, as well as the size and position of the rather broad palpi, which reach beyond the somewhat projecting anterior edge of the mouth. As these characters are easier to perceive than to describe in a few words, the present species deserves to be studied as a type of *Trypeta* in the narrower sense.

19. *T. florescentiæ* LIN. ♂ ♀.—Ex flavo-virescens, thoracis disco nigricante, postice breviter bifido, maculis alarum hyalinarum quatuor nigris, intermediis fere contiguis, aut in fasciam perpendicularem confluentibus.

Yellowish-green; the blackish color of the thoracic dorsum which does not reach the lateral margin is slightly bifid posteriorly; the hyaline wings show four black spots, the two intermediate ones of which are almost contiguous, or confluent in a perpendicular crossband. Long. corp. ♂ 0.17, ♀ cum terebrâ 0.20—0.21; long. al. 0.18.

SYN. *Musca florescentiæ* LINNÆ, Syst. Nat. X, p. 601, 99.!

Musca ruficauda FABRICIUS, Ent. Syst. IV, p. 353, 169.

Tephritis punctata FALLEN, Act. Holm. 1814, p. 167, 12.

Trypeta florescentiæ MEIGEN, Syst. Besch. V, p. 321. Tab. XLVIII, f. 3.

Trypeta florescentiæ LOEW, Germar's Zeitschr. V, p. 338. Tab. I, f. 15.

Trypeta florescentiæ LOEW, Europ. Bohrlf. 59, 11. Tab. IX, f. 2.

Pale yellowish-green. Front, third antennal joint, and palpi usually of a much more vivid yellow. Eyes very much rounded. Face short, excavated; the anterior edge of the mouth distinctly projecting. Antennæ rather short; the longer bristle upon the second antennal joint but little conspicuous. Palpi comparatively long, reaching beyond the anterior edge of the oral opening. Thoracic dorsum blackish, with the exception, however, of the lateral border and of a cuneiform beginning of a middle stripe, starting from the posterior end, and which renders the black coloring bifid posteriorly. Scutellum immaculate, except on the under side of the lateral angles, and provided with four bristles. Metathorax black. Pleuræ more or less infuscated, sometimes rather blackish-brown, with a yellowish-green longitudinal stripe upon their upper side and another across the middle. Abdomen with four rows of conspicuous black spots; its pile, in both sexes, is usually whitish; however, along the posterior margin of the single segments, some black hairs are usually inserted; the last

segment of the abdomen of the male is often clothed with altogether black pile. Ovipositor red or brownish-red; at its basis two, sometimes confluent, black spots are visible; the extreme tip also is usually black; in length, the ovipositor hardly exceeds the last two abdominal segments; it is not very much attenuated towards the end and is beset with black or blackish pile. Feet altogether pale clay-yellow. Wings hyaline, with a black or rather blackish picture; the outlines of this picture are surrounded, in immature specimens, with a purer hyaline, in riper ones, with a more whitish-hyaline hue; beyond this pellucid border, the former kind of specimens show an indistinct, the latter ones a more pronounced gray shade; the picture of the wings consists of four spots, very variable as to their size and the intensity of their coloring; the first spot covers the stigma and usually reaches only as far as the second longitudinal vein; the second begins near the anterior margin immediately above the posterior crossvein, thus leaving the tip of the marginal cell uncovered; it becomes narrower and more faint posteriorly, thus reaching more or less completely the anterior end of the posterior crossvein; the third spot usually appears as a broad border along the posterior crossvein and is more or less coalescent with the second, forming a perpendicular crossband; the fourth spot lies upon the apex of the wing and is more or less triangular, as its inner limit runs perpendicularly from the tip of the second vein to the fourth vein, which limits it posteriorly; around the small crossvein and in the environs of the root of the third vein there is a more or less apparent, sometimes very distinct infuscation.

Hab. Canada (Mr. Provancher); common also in all Europe, where the larva inhabits the flower-heads of different species of *Cirsium*.

Observation 1.—Europe possesses, besides the variety of this species, discovered by Mr. Provancher in Canada, another form, distinguished by considerably larger and darker spots on the wings. Specimens of both varieties might easily be taken for different species; nevertheless, passages from one form to the other occur in the picture of the wings, and I am not able to discover between both the slightest plastic difference. In Germar's Zeitschrift, Part V, Tab. I, f. 15, I have figured a wing of the first variety. An extreme instance of the second variety is figured in my Monograph: *die Europäischen Bohrfliegen*, Tab.

IX, f. 2. Meigen's figure (Syst. Besch. V, Tab. XLVIII, f. 3) likewise represents the latter variety. It is probable that it will also be found in America.

Observation 2.—The present species, as well as the preceding, belongs to the genus *Trypeta* in the narrower sense.

20. *T. polita* LOEW. ♀. (Tab. X, f. 12.)—Atra, nitida, scutello tumido concolore, capite præter faciem exalbida pedibusque lutescentibus, alæ albido-hyalinæ, maculâ basali atrâ, fasciisque tribus latisimis fusco-nigris, venis transversis valde approximatis.

Deep black, shining; the tumid scutellum is concolorous; the head, with the exception of the whitish face, and the feet clay-yellowish; the whitish-hyaline wings have a deep black spot upon the basis and three very broad deep black crossbands; the crossveins are very much approximated. Long. corp. ♀ 0.17—0.18, cum terebrâ 0.22; long. al. 0.17—0.18.

Syn. *Trypeta polita* LOEW, Monogr. Vol. I, p. 77. Tab. II, f. 12.

Hab. Mississippi (Schaum); Washington, D. C.; New York; Connecticut (O. S.).¹

Observation.—I have nothing to add to the above-quoted description. The systematic position of this species will be discussed in the second remark to the following species.

21. *T. atra* LW. ♂ ♀. (Tab. XI, f. 17.)—Atra, nitida, scutello tumido, concolore, capite præter faciem albidam, femorum apice, tibiis tarsisque luteis; alæ albido-hyalinæ, maculâ basali atrâ, fasciisque tribus latis fusco-atris, venis transversis valde approximatis.

Deep black, shining; the tumid scutellum concolorous; the head, with the exception of the whitish face, the tip of the femora, the tibiae, and the tarsi clay-yellow; the whitish-hyaline wings have a deep black spot upon the basis and three broad, deep brownish-black crossbands; crossveins very approximate. Long. corp. ♂ 0.12—0.13, ♀ 0.13—0.14, cum terebrâ 0.17—0.18; long. al. 0.13—0.15.

Syn. *Trypeta atra* LOEW, Berl. Entom. Zeitschr. VI, p. 219.

Trypeta atra LOEW, Dipt. Amer. Sept. Cent. II, No. 89.

Deep black, shining. Front rather broad, of a vivid reddish-yellow; the ocellar triangle, as well as the little stripes descending from the vertex and bearing the uppermost bristles of the vertex, black, with a whitish-gray pollen; anteriorly, on the lateral

¹ This species produces the galls on *Solidago*, described by me in the Trans. Amer. Entomol. Soc. Vol. II, p. 301. O. S.

margin of the front there are on each side two black bristles. Antennæ yellow; the blackish arista distinctly incrassated at the basis. Face whitish; the anterior oral margin not at all projecting. Cheeks whitish, under the eyes with a more or less brownish-red spot. Oral opening rather round. Proboscis short. Palpi short, but broad, pale yellowish, with some short, whitish pile. The upper and middle part of the occiput for the most part black. The ordinary frontal bristles and some of the bristles on the cheeks are black; otherwise the pile upon the head consists of very scattered, bristle-like, or stubble-shaped whitish hairs, which easily drop off. The upper side of the thorax is shining black, very convex; besides the usual black bristles, it shows white, bristle-like hairs, which border the denuded stripes. Metathorax with white pollen; its lower part shining black; pleuræ shining black, with some rare, stiff, bristle-like white hairs. Abdomen short, shining black, at the root of the single segments only somewhat glossy, in consequence of a very thin grayish pollen. The scattered, very rough pile on the abdomen is whitish; only the posterior margin of the segments and partly also the middle line of the abdomen, have black hairs. Ovipositor stout, conical, not flattened, shining black, beset with black pile, somewhat longer than the last three abdominal segments taken together. Coxæ and femora shining black, only the front femora on the under side with a few black bristles; the tip of the femora, the tibiæ, and the tarsi brownish-yellow or more reddish-yellow. Wings whitish-hyaline, short and rather broad, with very much approximated and very perpendicular crossveins. The extreme root of the wings is whitish; next follows a rather large and almost deep black spot, reaching as far as the axillary excision, and not much beyond the basis of the small basal cells; the first two crossbands, which follow next, are connected near the anterior margin and strongly diverge towards the posterior one; the first of them is even a little broader than the second and altogether black, while the inner part of the second is partly brown; the third band is separated from the second, near the anterior margin, only by a very narrow hyaline spot; it borders the apex of the wing far beyond the tip of the fourth longitudinal vein, but actually touches the margin of the wing only beyond the tip of the third vein; its inner portion is brown anteriorly.

Hab. Mexico (coll. Winth.); New York (Osten-Sacken).

Observation 1.—The appended figure of the wing is taken from a Mexican specimen. The specimens which I received from New York differ from the former in being a little larger and in the circumstance that the face is somewhat more uneven; perhaps only in consequence of a stronger desiccation. Moreover, the last section of the fourth vein is a little less curved, and the posterior end of the first crossband is prolonged further along the margin towards the posterior corner of the wing. In all other respects the agreement is such that I cannot believe *T. atra* to be a different species. From *T. polita* the present species is easily distinguished by the much greater divergency of the second and third crossbands on the wings, by the absence of the pale gray border of the crossbands, which is always perceptible in *T. polita*, and by the black coloring of the femora; moreover, the anterior part of the lateral border of the front bears only two bristles in *T. atra*, while there are three in *T. polita*. The Brazilian species *T. nigerrima* Loew is very much like *T. atra*, nevertheless they are easily distinguished. In order to facilitate the comparison, I let the description of this species follow.

***T. nigerrima* Loew.** ♀. (Tab. XI, f. 18.)—*Atra*, nitida, scutello tumido concolore, thoracis maculis laterallibus utrinque binis velutinis, abdomine fasciis albido-pollinoso ornato, capite flavo, pedibus ex-ferrugineo luteis, femoribus tamen posterioribus anticorumque liturâ ex-fusco nigris; alæ albido-hyalinæ, maculâ basali atrâ, fascisque tribus fusco-atris, primâ latissimâ, reliquis minus latis, venis transversis valde approximatis.

Deep black, shining; the tumid scutellum concolorous; thoracic dorsum with two velvet black spots on each side; abdomen with crossbands of white pollen; head yellow; feet brownish-yellow, the posterior femora and a stripe on the front femora brownish-black; wings whitish-hyaline with a deep black spot on the basis and with three black crossbands, the first of which is very broad, the two others less so; crossveins very much approximated. Long. corp. 0.12—0.13; long. al. 0.12—0.13.

SYN. *Trypeta nigerrima* Loew, Berl. Ent. Zeitschr. VI, p. 219.

Trypeta nigerrima Loew, Dipt. Amer. Sept. Cent. II, p. 89.

Shining black. Head whitish-yellow; the rather narrow and steep front much darker yellow; the frontal bristles black. Antennæ dark yellow, rather large, especially the elongated third joint, which has a rather sharp anterior corner. Arista apparently bare, rather slender, not incrassated towards its root, of a pale color. Face but very little excavated, and very

little retreating; the anterior edge of the mouth distinctly projecting in the profile. Eyes elongated. Cheeks somewhat broad, with an infuscated spot near the inferior corner of the eye, and with white pile. Oral opening small, rounded. The rather broad palpi yellowish, beset with whitish pile. The short and not geniculate proboscis dark brown. Thorax shining black, with a metallic lustre in the middle; upon its lateral border, on each side, there are two large, opaque, velvet black spots, separated by the origin of the transverse suture, which is tinged with yellow. The usual bristles are black; the number of pairs which were inserted on the thoracic dorsum cannot well be ascertained. Moreover, the surface of the thoracic dorsum shows remains of stiff, yellowish hairs, which seem to have bordered the broad, bare stripes and to have also been inserted on the posterior part of the broad middle line. Scutellum turgid, shining black, with four bristles. The upper part of the metathorax is black, as in most of the allied species; the lower portion is covered with white pollen, which does not quite reach its lower margin. Femora with whitish pollen and white hairs; the humeral corner, as well as a little stripe behind it, near the upper margin, are velvet black. Abdomen shining black; a thin whitish pollen covers the whole anterior part of the first segment, forms, upon the first, second, and third segments, a band along their posterior margin which is perceptibly expanded and sharply emarginate in the middle; the posterior margin of the fourth segment has a similar, although narrower, band. The scattered pile on the abdomen is black, gray at its basis, in part yellowish-white upon the last segment. The flat, shining black ovipositor is about as long as the three last abdominal segments taken together, and is beset with delicate, black pile. Feet reddish-yellow, the middle and hind femora, with the exception of the extreme root and of the tip, brownish-black; the front femora have a brownish-black stripe upon their upper side. Wings broad, the apex but little rounded, hyaline, somewhat whitish; at their basis there is a large black spot, reaching into the basal cells; besides, there are three black cross-bands, entirely coalescent at the anterior margin of the wing and diverging posteriorly; the first of them, which is by far the broadest and is rather perpendicular, runs from the stigma, over the basis of the discal and of the third posterior cells, towards the posterior margin of the wing; the second band is the narrowest, and runs from the stigma over both crossveins, and hence, obliquely, towards the posterior margin; the third band starts from the stigma and follows the anterior margin and the apex, as far as the tip of the fourth vein, but, nevertheless, remains separated from the costal vein by a narrow, irregular, hyaline interval, which extends almost to its very end; near the submarginal cell, this interval is a little expanded and includes a punctiform dot, placed near the third vein; the first and second longitudinal veins are a little more distant from the anterior margin than in most of the related species; both crossveins are very approximate; the third longitudinal vein is beset with short bristles.

Hab. Brazil (coll. Winthem).

Observation 2.—*T. polita* and *atra*, as well as *T. nigerrima* are closely related in their organization. Among the European *Trypetes*, the species of the genus *Oedaspis* stand next to them, especially when this genus is confined to *Oedaspis multifasciata* Loew and its next congeners, at the exclusion of *Oed. Wiedemanni* Meig. and *resoviana* Costa. The American species differ from the above-mentioned European ones (*multifasciata* Lw., *dichotoma* Lw., and *fissa* Loew) in several characters, which they have in common; the most striking of these are: 1. The rather long, stubble-shaped pile; 2. The longer and more pointed ovipositor; 3. The different picture of the wings. The latter difference will be sufficiently apparent, when the figures which I give of the wings of *polita*, *atra*, and *nigerrima* are compared with the figures of the wing of *T. multifasciata*, produced in the *Europ. Bohrfliegen*, Tab. VI, f. 2. The pictures of *T. fissa* and *dichotoma* agree, in their general features, with that of *multifasciata*. These differences of the three North American species are not of sufficient importance to require the establishment of a new genus for them, and I have not the slightest hesitation in placing them in the genus *Oedaspis*, in the narrower sense, defined above.

22. *T. gibba* n. sp. ♀.—*Atra*, nitida, scutello tumido, concolore, facie albicante, pedibus subbadiis; alæ albido-hyalinae, maculâ basali atrâ fasciisque tribus latis fusco-atris, venis transversis valde approximatis, cellulâ marginali per venulam transversalem adventitiam dissectâ.

Deep black, shining; the turgid scutellum of the same color; face whitish; feet chestnut-brownish; wings whitish-hyaline, with a deep black spot at the basis, and with three brownish-black crossbands, very much approximated crossveins, and a supernumerary crossvein dividing the marginal cell. Long. corp. 0.13, cum terebrâ 0.17; long. al. 0.14—0.15.

Very like the three preceding species and closely allied to them, nevertheless, distinguished in some peculiar plastic characters. Deep black, shining. Front conspicuously broad, of an opaque, dirty, brownish, more reddish-brown on the sides; the four bristles on the posterior part of the vertex, the bristles near the ocelli, the four bristles crowded together and inserted on the small stripes running from the vertex towards the front, finally two bristles on each side, near the lateral frontal border, are all black; the latter two are inserted, one very high up, the other very low

down, so that the distance between them is remarkably large. Otherwise the head is beset with almost bristle-like white stubble-shaped pile. The very large and sharply defined frontal lunule, the face, including the cheeks, and the lower half of the occiput are whitish; the upper part of the latter blackish, although covered with whitish pollen. The perpendicular diameter of the eyes has about double the length of the horizontal one; nevertheless, the cheeks are remarkably broad; a brownish stripe runs from the lower corner of the eye perpendicularly towards the edge of the mouth; the hairs, inserted upon its lower end, are brownish-black or black. The first two antennal joints are clay-yellowish; the third joint is dark brown, rather large, short-oval in outline; arista bare, not incrassated at the basis, black. Oral opening larger than in the preceding species; its transverse diameter comparatively larger; proboscis and palpi short, brown. The very convex thorax and the turgid scutellum are deep black, shining, with a very weak metallic, violet reflection; the remarkably broad lateral stripes and the anterior end of the broad middle stripe are bare. The lateral stripes are bordered with coarse, yellowish, stubble-shaped pile, and the posterior two-thirds of the middle stripe, besides being covered with white pollen, are densely beset with similar hairs. The ordinary bristles of the thoracic dorsum are black, and more numerous than usual, as there are four pairs of them along the longitudinal middle line, the anterior pair being inserted immediately in front of the transverse suture. The shining black metathorax has, under the swelling lying immediately under the scutellum, a crossband of thick white pollen. The pleuræ show upon the greater part of their upper half, a thin, whitish-gray pollen, and are everywhere beset with stubble-like white hairs. The abdomen seems to be covered everywhere with a thin gray dust, which is somewhat more dense and more whitish-gray upon the posterior border of the single segments; its rather long stubble-like pile is white. The comparatively long and pointed ovipositor is deep black, shining, and beset with short, fine, black pile. Feet chestnut-brownish. Wings short, rather broad in proportion to their length; the altogether black venation is very similar to that of the immediately preceding species, except that the comparatively broad marginal cell is divided in two halves by a perpendicular crossvein, which touches the costa at a point perceptibly nearer

from the tip of the first than from that of the second vein. I take this crossvein to be a constant character of the species, as it exists on both wings of my specimen, and as several closely allied *Trypetidæ*, for instance *Gonygl. Wiedemanni* and *Coprom. vesuviana*, have it likewise, although incompletely developed. The picture of the wings is not unlike that of *T. atra*, in its design as well as in its coloring; the black spot upon the basis of the wings does not cover their extreme root, and extends, on the anterior margin, only very little beyond the humeral crossvein; it hardly reaches beyond the first longitudinal vein, and dissolves in several radiating points, which occupy the longitudinal middle of the marginal and of the three basal cells and almost come in contact (except the hindmost), with similar rays, meeting them from the opposite side and emitted by the first crossband; the first black crossband has almost the same position as in the three preceding species, but it is much narrower, especially towards its end, which reaches the posterior margin; its interior does not show any brownish tinge. The second band runs over both crossveins, exactly as it does in those three species, and is connected with the first on the anterior margin in the same manner as this is the case in *T. atra*; the stigma, lying within this connecting portion, is very short; the veins surrounding it have, on the inner side, a very narrow hyaline border; the interior of the second band is for the most part brownish. The last black band begins in the marginal cell somewhat beyond the supernumerary crossvein in this cell, and reaches some distance beyond the end of the fourth vein; as far as this vein, it is separated from the margin of the wing by a narrow hyaline border, which somewhat projects on the inside on the second and third veins; beyond the fourth vein the band comes in immediate contact with the margin of the wing; the inside of this band is brownish upon the anterior two-thirds of its course.

Hab. Texas (Belfrage).

Observation.—The differences between the present species and the three preceding ones are evident: they consist in an aberrant arrangement of the bristles of the front and of the thoracic dorsum, in the size and shape of the third antennal joint, and in the presence of the crossvein, dividing the marginal cell; nevertheless the agreement between those species in most of the other plastic characters, in the shape of the body and in the picture of the

wings, is convincing enough to remove all doubt as to its location in the genus *Edaspis*.

23. *T. cingulata* Lw. ♂ ♀. (Tab. X, f. 11.)—Nigra, capite pedibusque luteis, thoracis margine laterali scutelloque præter margines laterales et anticum dilute flavis, margine postico segmentorum abdominalium singulorum albido; alæ hyalinæ, maculâ parvâ apicis fascisque quatuor fusco-nigris, harum duabus primis postice abbreviatis et liberis, duabus ultimis integris et antice conjunctis.

Black, head and feet clay-yellow; lateral border of the thorax and the scutellum, the latter with the exception of the anterior and lateral border, light yellow; abdominal segments whitish on the posterior border; wings hyaline; a small spot upon the apex and four crossbands brownish-black; the first two bands abbreviated posteriorly and not connected; the two posterior bands are entire and connected on the anterior margin. Long. corp. 0.14—0.22; long. al. 0.15—0.20.

SYN. *Trypeta cingulata* LOEW, Monogr. I, 70 Tab. II, f. 11.

Hab. Middle States; Long Branch, N. J., in July (Osten-Sacken).

Observation.—The description given by me in the first part of these Monographs will easily help to identify this species. I have nothing to add to it, but must call attention to the great variation in the size of different specimens. The smallest ones which I possess, are without exception males. *T. cingulata* is closely allied to the European species of *Rhagoletis*, especially to *R. flavicincta* Loew; its systematic location is, therefore, not doubtful.

24. *T. tabellaria* FITCH. ♀.—Atra, capite, trochanteribus, tibiis tarsisque dilute luteis, thoracis margine laterali scutelloque præter margines laterales albis; segmentorum abdominalium singulorum margine postico exalbido; alæ pure hyalinæ, fasciis quatuor latis nigris, duabus primis postice, duabus ultimis antice coherentibus.

Deep black; head, second joint of the coxæ, tibiæ, and tarsi yellow; lateral border of the thorax and scutellum, with the exception of the anterior and the lateral borders, white; the posterior borders of the abdominal segments whitish; wings of a pure hyaline, with four broad, black crossbands, of which the first two are connected at the posterior, the last two at the anterior margin. Long. corp. 0.14—0.15; long. al. 0.14—0.15.

SYN. *Tephritis tabellaria* FITCH, First Report, p. 66.

Shining black; head yellowish; occiput black, with a pale yellow border; front broad, more bright yellow; only the spot upon which the ocelli are placed and the small, very narrow stripes, which run down from the vertex upon the front, are of a blackish color; the usual frontal bristles are black. Antennæ of a vivid ochre-yellow; their last joint is elongated-oval, obtuse at the end; arista blackish, with a hardly perceptible pubescence. Oral opening rather large, somewhat longer than broad; its anterior edge drawn up, but not projecting in the profile. Proboscis and palpi short, brown, the latter more clay-yellow towards the tip. The thoracic dorsum shows two longitudinal stripes, rather distant from each other, somewhat abbreviated posteriorly and covered with a thin, white pollen; upon the anterior part of the thoracic dorsum a similar pollen covers not only the interval between the stripes, but also extends beyond them. The whole of this pollen, however, is but little conspicuous and seems to be easily rubbed off. The humeral angle and a stripe running from it towards the root of the wings, are white. The flat scutellum, with the exception of its lateral border, has the same color. Metathorax without any pollen, altogether shining deep black. The usual bristles of the thorax and the four bristles of the scutellum are deep black. The other hairs on the thoracic dorsum are very short and delicate. Abdomen shining black; its first two segments are more opaque, being clothed with a brownish-black pollen. The first three segments, upon their posterior margin, have a crossband of a whitish pollen. The very short and soft hairs upon the abdomen are black; the paler crossbands upon the posterior border of the first three segments show some whitish hairs; the bristles upon the sides of the intermediate segments and upon the rather large last segment are black. Ovipositor shorter than the last abdominal segment, broad at the basis, much narrower at the end, shining black and with a black pubescence. Second coxal joint pale clay-yellowish. Femora black, only the extreme tip yellowish-brown; tibiæ and tarsi pale clay-yellowish; the former somewhat more brownish at the basis; the bristles upon the upper side of the hind tibiæ are remarkably short. Wings pure hyaline, almost whitish hyaline, with four entire black crossbands, the first of which of a medium breadth, the three others very broad. The first band is somewhat oblique and begins on the humeral cross-

vei
cou
pos
the
the
pos
bet
cro
tud
inn
ban
nea
wit
veit
the
sec
inte
the

Z
C
pre
may
how
to t

25.

fe
lo
pi
pe
ni
di
Bro
an
tl
w
tl
w
tl
d
0.

Szn

vein; the second is perpendicular and begins on the stigma; both converge posteriorly and coalesce quite a distance from the posterior margin, so that the cuneiform hyaline space between them does not reach beyond the anterior angle of the basis of the third posterior cell. The third black band runs over the posterior crossvein and is parallel to the second band, so that between both there is a somewhat irregularly limited hyaline crossband, which is perceptibly dilated between the third longitudinal vein and the anterior margin; it reaches the latter immediately behind the stigma; the posterior end of the third band shows some inclination to coalesce with the second band near the posterior margin. The fourth band completely coalesces with the third between the costa and the second longitudinal vein, and follows the margin of the wing some distance beyond the end of the fourth longitudinal vein; between the tips of the second and fourth veins, however, there is a rather broad hyaline interval between it and the margin; beyond this point, it touches the margin completely.

Hab. New York (Dr. A. Fitch); Canada (Mr. Provancher).

Observation.—In the first volume of the Monographs I expressed the supposition that the *Tephritis tabellaria* of Fitch may not be a *Trypeta* at all, but an *Ortalida*; this supposition, however, proved to be erroneous; it is a *Trypetida*, belonging to the genus *Rhagoletis*.

25. *T. pomonella* WALSH. ♀.—Fusco-nigra, capite, trochanteribus, femorum apice, tibiis, tarsisque luteis, thoracis margine laterali, scutelloque præter margines laterales et anticum albis, abdominis colore in piceum vergente, segmentorum marginibus posticis confertim albido-pollinoso, terebrâ latissimâ, sed brevi; aë hyalinæ. fasciis quatuor nigris, primâ subbasali, reliquis tribus integris, antice conjunctis, postice divergentibus.

Brownish-black; head, second joint of the coxæ, tip of the femora, tibiæ, and tarsi clay-yellowish; lateral margin of the thorax and scutellum, the latter with the exception of its basis and of its lateral margins, white; abdomen more pitch-brown, with crossbands of white pollen on the posterior margins of the segments, ovipositor very broad, but short; wings hyaline, with four black crossbands, the first of which lies near the basis, the last three are connected near the anterior margin and divergent toward the posterior one. Long. corp. 0.17, cum terebrâ 0.19; long. al. 0.17.

Syn. *Trypeta pomonella* WALSH, First Rep. Illin. etc., p. 29-33, f. 2.

I possess but a single specimen of this species. Its coloring is not fully developed, although otherwise its preservation is perfect. It is black, with a distinct brownish tinge; its abdomen is more pitch-brown and rather shining. Head pale yellowish, with a narrow dark yellow front and more ochre-yellow antennæ; the third joint of the latter is narrow and rather long, rounded at the end; the slender arista is dark brown, with a short, although distinctly discernible pubescence. The usual frontal bristles are black; behind the ocelli, however, near the lateral margin, two shorter, whitish bristles are placed. Oral opening large, broader than long. Palpi and proboscis pale yellowish, with a pale pubescence; the former do not project beyond the anterior edge of the mouth, the flaps of the latter somewhat prolonged. The thoracic dorsum shows four rather narrow longitudinal stripes, formed by a whitish pollen; these stripes, arranged in pairs, are confluent anteriorly; the outside stripes are moderately abbreviated before the posterior margin of the thorax; the inside ones reach only as far as the anterior pair of bristles, inserted upon the longitudinal middle of the thorax; each of the bristles of this pair is placed between the end of the corresponding inside stripe and the outside one; the inside stripes are separated by a broad dark interval, which shows the shining brownish-black color of the remainder of the thorax. When the thorax is viewed from the front side, the light falling in from behind, the pollinose stripes appear somewhat more broad; the interval between the inside stripes appears somewhat narrower and a little more opaque; at the same time, this point of view discloses upon the outside stripes and upon the margin of the inside ones, alongside of them, some short, snow-white pile, while the remaining pile of the thoracic dorsum is black. The humeral callosity and a stripe running from it to the root of the wing, is white. The rather flat scutellum is white, blackish on the sides and at the basis. The bristles of the thorax and the four bristles of the scutellum are black. The first four segments of the abdomen have each, on the posterior margin, a rather uniformly broad crossband, formed by whitish pollen; the last segment, which has no such band, is paler brown along the posterior margin. The comparatively scattered and not very short pile on the abdomen is black; it is white only on the pale crossband on the posterior part of the first segment. The bristles

on the sides of the middle and of the last segments are black. Ovipositor very short, about once and a half the length of the last abdominal segment, very conspicuously broad, not much attenuated towards the end, very broadly truncate and somewhat convex; its coloring is a shining brownish-black or black; the pubescence is black. In agreement with the unusual breadth of the ovipositor, the last abdominal segment is also very broad, which causes the whole abdomen to have a peculiar shape. The second coxal joint yellowish; posterior femora black with a clay-yellow tip; front femora clay-yellow, with a large, broad, brownish-black stripe upon the hind side; tibiae and tarsi clay-yellowish, the tip of the latter dark brown. Hind tibiae on the upper side beset with rather long bristles.

Hab. Illinois (Walsh); the larva, originally feeding upon the fruit of a *Crataegus*, is now frequently found upon the fruits of the apple-tree, which it damages.

Observation.—The next relatives of *T. pomonella* are found in a series of South American species, only a single one of which, as far as I know, has been previously described; it is to be found in Macquart's *Diptères Exotiques*, Suppl. IV, p. 288, Tab. XXVI, f. 15, under the name of *Urophora scutellaris*. It is not an *Urophora* however, and moreover, the name of *scutellaris* cannot be maintained, as Wiedemann has previously used it for another species. The species may, therefore, be called *Trypeta Macquartii*. Macquart's figure shows, that this Brazilian species differs in the picture of its wings from the species of *Rhagoletis* previously described, and that, in this respect, it is more like the species of *Acidia*. The structure of its body shows a corresponding approach to the species of this latter genus, while, on the other hand, coloring and picture of the body are most strikingly like those of *Rhagoletis*. As this species is also very like the North American *Rhagoletis* in the structure of its body, the question arises whether it is better to place it in the genus *Acidia* or in *Rhagoletis*. I prefer the latter course, because we thus facilitate the generic determination of the allied species. *Trypeta pomonella*, as has already been mentioned above, is among the number of such species, the picture of its wings being very like that of *T. Macquartii*. It is true that it differs not inconsiderably from *T. Macquartii* in the greater length of the third antennal joint, the considerable size and breadth of the oral opening, and

the strikingly large transverse diameter of the short ovipositor; but, like *Trypeta Macquartii*, it agrees with the true species of *Rhagoletis* in the coloring and in the picture of the body, so that I prefer, for the present, to leave it in that genus. It may be objected that, in this case, I lay a greater stress upon peculiarities of the coloring and mere differences of *habitus* than upon plastic characters. In answer to this objection I may state that I fully appreciate the value of plastic differences in matters of generic grouping of species, but that the knowledge of the exotic *Trypetæ*, as well as the existing descriptions of them, are not sufficient for their generic distribution upon plastic characters only. Most descriptions mention but very little about these characters, the more so as in most cases they have to be drawn from a few indifferently preserved specimens, which do not allow a sufficiently clear view of such characters. And thus it happens that peculiarities of coloring and other habitual characters become in many cases very useful for the generic distribution of exotic *Trypetæ*, especially in cases where the only available plastic characters are of a very delicate nature and hence more difficult to perceive. It is true that the exotic species thus treated are merely *grouped*, and not systematized; but this grouping in itself is a progress towards the determination of the species, and is one of the usual steps towards a systematic distribution.

26. *T. insecta* Lw. ♀. (Tab. X, f. 8.)—Thorace nigro, capite, abdomine pedibusque luteis, alarum nigrarum incisuris marginalibus, guttulisque inter venarum longitudinalium tertiam et quartam tribus vel quatuor pellucidis, venâ longitudinali tertiâ nudâ, setis sentelli duabus.

Thorax black; head, abdomen, and feet clay-yellow; wings black, with hyaline indentations along the margin and with three or four hyaline drops between the third and fourth veins; the third vein not bristly; sentellum with two bristles. Long. corp. 0.14; long. al. 0.14.

Syn. *Trypeta insecta* Loew, Monogr. I, p. 72. Tab. II, f. 8.

Hab. Cuba (Poey). [Hayti; P. R. Uhler.—O. S.]

Observation 1.—*T. insecta* belongs to the typical species of the genus *Acitura*, the sentellum of which bears only two bristles. The picture of the wings of this genus is characteristic.

Observation 2.—Another *Trypeta* of the same genus occurs in Brazil, which may be easily mistaken for *Trypeta insecta*. I prefer, therefore, to describe it here:—

T. phœnicura n. sp. ♂ ♀. (Tab. XI, f. 12.)—Nigra, capite pedibusque ochraceis, alarum nigrarum incisuris marginalibus guttulisque later venas longitudinales tertiam et quartam tribus pellucidis, venâ longitudinali tertiâ undâ, setis scutelli duabus.

♂. Abdomen ex ferrugineo rufum, segmento ultimo nigro.

♀. Abdomen nigrum, basi ferruginâ, terebrâ latâ late aurantiacâ.

Black, head and feet ochreous-yellow; wings black, with hyaline indentations along the margin and with three hyaline drops between the third and fourth longitudinal veins; the third longitudinal vein is not bristly; the scutellum has two bristles.

♂. Abdomen ferruginous, its last segment black.

♀. Abdomen black, ferruginous at the basis; the broad ovipositor is of a vivid orange-yellow. Long. corp. ♂ 0.14, ♀ 0.15—0.16; long. al. 0.14.

Black; head of an impure ochre-yellow; the occiput alone mostly blackish; front narrow, especially anteriorly; frontal bristles black. Eyes very large, cheeks very narrow. Face short, concave; nevertheless, the anterior oral edge not projecting in the profile. The antennæ reach down to the oral edge; their third joint is rounded at the tip; the blackish arista is long and slender, apparently bare. Oral opening of medium size, rounded; proboscis not geniculate. The thorax and the two-bristly scutellum are black, their short pile yellowish-white, their bristles rather black; the somewhat rounded abdomen of the male is of a dirty ferruginous color (in living specimens its color may be purer); its last segment is black. The extent of the black color is greater in the female abdomen, the first segment, the basis of the second, and the anterior corners of the third alone, being ferruginous. The short pile of the abdomen is paler, almost yellowish in the male, somewhat brown in the female; on the posterior border of the last segment of the abdomen of the female there are some black hairs. The flattened, comparatively broad ovipositor, attenuated towards its end, has a shining surface; its color is a very bright orange-yellow, the tip alone shows a narrow black border; its short pubescence is pale. Coxæ and feet ochreous-yellow; the extreme tip of the posterior femora is somewhat blackish. Wings comparatively long and narrow, towards the end somewhat less broad and less obtuse than those of *T. insecta*, black, with a hyaline picture; near the costa, anterior to the stigma, there are three small hyaline spots, the first anterior to the humeral crossvein, the two others in the costal cell; immediately beyond the stigma, which is altogether black, there are two conspicuous triangular hyaline spots, which, with their pointed end, do not quite reach the third longitudinal vein; on the posterior margin of the wing there are six hyaline indentations, the last of which alone ends in a point; the first two are connected with the almost hyaline posterior angle of the wing, reach as far as the fifth longitudinal vein, and are separated by a much broader black band than the other indentations; the two following indentations cross beyond the fifth vein, the first below the

small crossvein, the second immediately before the end of the discal cell; the fifth indentation follows the outer side of the great crossvein (which runs obliquely backwards); the sixth, separated from the preceding by a black band of moderate breadth, is almost triangular; the three small hyaline dots between the third and fourth veins lie, the first under the stigma, the second between the two crossveins, near the fourth vein, the third above the last of the hyaline excisions along the posterior margin.

Hab. Brazil.

The coloring of the abdomen of *T. insecta* and *phoenicera* seems to be somewhat variable, and hence not to be relied on as a specific character; the more marked are the differences in the outline and picture of the wings.

27. *T. pœcilogastra* n. sp. ♂.—Lutea, scutello setis sex instructo, abdomine nigro-variegato, alis latis fuscis, inaequaliter limbo-guttatis, venisque longitudinalibus primâ, tertîâ et quintâ confertim nigro-setosis.


Clay-yellow, scutellum with six bristles, abdomen variegated with black; wings broad, blackish-brown, with unevenly distributed hyaline drops; the first, third, and fifth longitudinal veins densely beset with black bristles. Long. corp. 0.21; long. al. 0.24.



Clay-yellow; the color of head and antennæ more ochre-yellow; the last joint of the latter elongated, rounded at the tip; the long brown arista beset with a very short pubescence. The face is rather retreating nearly as far as the vicinity of the anterior edge of the mouth; the latter is somewhat turned upwards and abruptly projecting when seen in profile. The vertical diameter of the eyes has double the length of the horizontal one; hence, the cheeks are very narrow. Proboscis tumid; palpi rather broad and short, although they project a little beyond the anterior edge of the mouth. The usual frontal bristles are black. The two pairs of bristles on the middle line of the thoracic dorsum are weak and of a blackish-brown color, like the other thoracic bristles; the anterior pair is at an unusual distance behind the transverse suture. Scutellum rather flat, with six brown bristles. Metathorax with two brown longitudinal stripes. Abdomen with a complicated black picture, the only visible portions of the ground color being an uninterrupted middle line of almost trapezoidal spots, and on both sides of it, two rows of other spots; the spots of the outer row lie on the anterior angles of the single segments; those of the inner row on the anterior

borders of the segments. Feet pale clay-yellowish. Wings broad, blackish-brown, with large and small hyaline dots, unequally distributed; the costal cell is pale brown between the extreme basis and a trifle beyond the humeral crossvein; next follows upon the costa a square brown spot, and then a square hyaline space, somewhat encroaching upon the stigmatical cell, so as to include the end of the auxiliary vein, which runs perpendicularly towards the margin of the wing; the stigmatical cell is otherwise tinged with blackish-brown and has, close to the anterior margin, two hyaline drops; immediately beyond the tip of the first longitudinal vein, near the anterior margin, there is a hyaline drop, reaching as far as the second longitudinal vein, the largest in the whole picture of the wing; in the vicinity of the apex of the wing the drops are larger than in the middle and more close together; so that a row of dots, reaching from the tip of the second vein to the posterior angle of the second posterior cell, and moreover four dots along the margin of the wing, may be discerned; among the latter, the first lies in the submarginal cell and is connected with a little drop behind the third vein; the second lies at the extreme tip of the wing; the last two in the second posterior cell; a second group of larger drops lies in the third posterior cell, immediately below the stigma; it consists of four drops, between which the black ground color is more or less faint, and of two other drops on the anterior side of the fifth vein; between this group of drops and the fifth longitudinal vein, there is, near the margin of the wing, a single larger drop; the posterior angle of the wing is brownish-gray, with several rather large limpid drops; the middle of the wing shows only small and isolated drops. The first, third, and fifth longitudinal veins are very closely beset with rather strong bristles; the second is strongly curved; the third and fourth diverge towards their end; the small crossvein is but little beyond the middle of the very broad discal cell, and the posterior crossvein has a very steep position; the anal cell is drawn out in a narrow and very long lobe.





Hab. Cuba (Gundlach).

Observation.—The six bristles upon the scutellum, as well as the dense bristles upon the first, third, and fifth longitudinal veins, distinguish *T. paucilogastra* from all the following species, provided with a reticulate picture of the wings. It is very





**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

1.5 28
1.6 32
1.8 22
1.9 20
1.8

1.5
1.6
1.8
1.9

closely allied to the species of *Hexachata*, in which, however, as far as I know them, the fifth vein has bristles upon the basis only, while in the present species the bristles almost reach to the tip. For this reason, as well as on account of the different character of the picture of the wings, I do not deem it convenient to place it in the genus *Hexachata*. Whether Mr. Saunders's genus *Dasyneura* would better answer for it, I am unable to say, as I have not been able to procure the publication which contains it. For the present therefore I set this species up as the type of a new genus, which I call *Blepharoneura*.

28. *T. testudinea* n. sp. (Tab. XI, f. 13)—Ex luteo fusca, capite, thoracis dorso, pedibusque luteis, terebrâ duobus ultimis abdominis segmentis semel sumtis paulo longiore; alæ valde dilatatæ, e nigro fuscæ, strigis duabus hyalinis inde a margine cellulæ posterioris secundæ usque ad venam longitudinalem tertiam ascendentibus, primo limbi costalis dimidio grosse nigro maculato, disco alarum guttulis minutis pellucidis confertim asperso.

Yellowish-brown, head, thoracic dorsum, and feet clay-yellow; the ovipositor only a little longer than the last two abdominal segments taken together; wings very broad, blackish-brown; two hyaline indentations reach from the posterior side of the second posterior cell to the third longitudinal vein; the anterior half of the region along the costa shows a number of large, black spots: the central portion of the wing is occupied by many small, hyaline drops. Long. corp. cum terebrâ 0.21; leng. al. 0.19.

A species very much resembling the *T. latipennis* Wied., but differing in the smaller size and the less minute dots on the central portion of the wing. The coloring of the body is yellowish-brown, but may be somewhat darker in fully colored specimens. The ground color of a great part of the upper side of the thorax is blackish, but very much concealed under a thick clay-yellow pollen. Front opaque, of a moderate breadth, still narrower anteriorly; the usual frontal bristles are brown. Eyes large, elongated; cheeks very narrow, with much pile; face short, descending rather perpendicularly, but distinctly excavated under the antennæ; the anterior edge of the mouth not projecting. Antennæ ochre-yellow, of a medium length, but, owing to the shortness of the face, reaching to the anterior edge of the mouth; the third joint has a rather rounded anterior corner; the moderately long arista thin and bare. The middle of the thoracic

dorsum shows traces of a pair of bristles. Scutellum but little convex, provided with four bristles. Metathorax blackish with a grayish-yellow pollen. The color of the pleuræ, in the described specimen, does not differ much from that of the remainder of the body; it seems, nevertheless, that, in more fully colored specimens, a considerable portion of the pleuræ may be blackish; they are thickly clothed with a clay-yellow pollen; the pile and the bristles upon them, like those on thorax and scutellum, are yellowish-brown. The abdomen shows a trace of four dark longitudinal stripes, formed by very much faded blackish spots; the pile upon it is somewhat shorter and rather blackish upon the anterior half of the single segments; upon their posterior half, it is somewhat longer and almost whitish; yet the long bristles on the posterior border of the last segment are blackish-brown. The flat ovipositor, which in the allied *T. latipennis* Wied. equals the last four abdominal segments in length, is but a little longer here than the last two segments taken together; it is of the same color with the abdomen, somewhat blackened at the root and tip, and beset everywhere with short blackish pile. Feet brownish-ochre yellow. Wings very broad, very like those of *T. latipennis* in outline, venation, and picture; proportionally, however, they are not quite as broad and not quite as convex on the anterior margin; upon the apical third of the wing there are three crossbands, connected anteriorly and separated by narrow, hyaline intervals, beginning at the posterior margin; the first band is contiguous, on its outer side, to the posterior crossvein, and expands across it near its posterior end; the second runs across the middle of the second posterior cell, the third borders the apex of the wing. The remaining portion of the surface of the wing, beyond the second longitudinal vein, has a somewhat darker brownish tinge, and is covered with a multitude of small hyaline drops, which partly coalesce into longitudinal rows, and in some places, as at both ends of the small crossvein and here and there on the longitudinal veins, leave unbroken brown spots. Upon the posterior margin, there is a broad brown border, bearing a few larger, but not very well-defined drops, which are also less hyaline than those of the centre of the wing; on the posterior angle of the wing the border is somewhat faint. The brownish-black stigma coalesces with a spot of the same color immediately behind it, which spot crosses but little the second longitudinal

vein; two large spots of the same color lie in the exterior costal cell and fill out a large portion of it; a double spot of the same color is in the marginal cell immediately beyond the stigma; finally, there are two large spots of the same kind on the second longitudinal vein, the one upon its root, the other below the double spot in the marginal cell. The basis of the exterior costal cell is irregularly reticulate with very small drops. A small hyaline spot is situated between the double spot of the marginal cell and the end of this cell, filled out by the common origin of the three crossbands which occupy the apex of the wing. The third longitudinal vein is distinctly bristly, gently curved forward before its end and as gently backwards; posterior crossvein long, but not as long as in *T. latipennis* Wied.

Hab. Cuba (Otto); in the Berlin Museum.

Observation.—The present species forms, with *T. latipennis* Wied. and a group of related species from South America, an easily recognizable genus, very well characterized by the breadth of its large wings, their outline, which reminds of *Phasia*, and their peculiar picture. These species also have the structure of the head and the bristly third vein in common. I adopt for this genus, apparently exclusively American, the name of *Acrotama*, in allusion to the most striking peculiarity of the picture of the wings.

29. *T. sparsa* WIED. ♂ ♀. (Tab. X, f. 13.)—Fusca, alæ latissimæ, subrotundatæ, nigræ, albido-guttulatæ, apice albido-marginato ornatæ.

Brown; wings very broad, almost round, black, with whitish drops, and the apex margined with white. Long. corp. ♂ 0.15—0.27; ♀ cum terebrâ 0.19—0.30; long. al. 0.16—0.26.

SYN. *Trypeta sparsa* WIEDEMANN, ANS. Zweifl. II, p. 492.

Trypeta caliptera SAY, Journ. Acad. Phil. VI, p. 187, 3.

Platystoma latipennis MACQUART, Dipt. Exot. II, 3, p. 200. Tab. XXVI, f. 8.

Acinia noveboracensis FITCH, Firs. Report, 67.

Trypeta sparsa LOEW, Monographs, etc., I, p. 78. Tab. II, f. 13.

Hab. Northern Wisconsin River (Kennicott); Texas (Bel-
frage).

Observation 1.—*Trypeta sparsa* Wied. is either a very variable species, both in its size and in the shape of its wings, or else several species are mixed up here, which, owing to the insuffi-

ciency of my materials, I am unable to distinguish. The description given in the first volume of these Monographs refers to the specimen from Northern Wisconsin River. Another specimen from the same locality, much smaller and paler and with less broad wings, has been mentioned in a note, appended to the same description. The mention concerning the size of the specimen, however, has been omitted there. The drops on the wings of that specimen are larger and more rounded than in ordinary specimens and show less tendency to form longitudinal rows; the costal cell also contains such drops, while in the larger specimens it shows at the utmost some pale drops along the auxiliary vein. Nevertheless, even now, I would not consider this specimen but as a variety of *T. sparsa*.

Observation 2.—Wiedemann's collection contains at present, under the name of *T. sparsa*, a pair of specimens, the communication of which I owe to the kindness of the Vienna Museum. In the list of species sent to me, they were marked as coming from Brazil. As Wiedemann prepared his description from a single female of unknown origin, it seems hardly probable that the female specimen now existing in his collection is the typical one. It is more likely, on the contrary, that the couple of specimens from Brazil now to be found in the collection was later added to it by Wiedemann. Both sexes most closely resemble my Wisconsin specimens, except that the wings are still broader, which is caused by the greater breadth of the costal and stigmatal cells; their anterior margin is distinctly more convex. These specimens seem therefore to belong to a South American species, very closely allied to the North American one. However, my conviction that such is the case has been somewhat shaken by a number of specimens from Texas, collected by Mr. Belfrage. The larger ones have the wings a little broader than the larger specimens from Wisconsin, and the pellucid drops are less regularly distributed; the costal and stigmatal cells are not broader; a small and incompletely colored specimen has much narrower wings than the larger specimens; yet they are broader than the wings of the above-mentioned smaller specimen from Wisconsin. Whether the specimens from Wisconsin and Texas belong to the same species, will have to be proved by further observation.

Observation 3.—The present species, together with *T. rotundipennis*, as well as the species represented by the above-

mentioned specimens from Brazil, now called *T. sparsa* in Wiedemann's collection, form a separate genus, the characters of which may be easily gathered from the descriptions of *T. sparsa* and *rotundipennis* in the first volume. I call it *Eutreta*, in allusion to the characteristic picture of the wings.

30. *T. rotundipennis* Lw. ♂. (Tab. X, f. 14.)—Fusca, alis latissimis, rotundatis, nigris, albido-guttatis, in marginibus antico et apicali maculas minutas albidas gerentibus.

Brown, wings very broad, rounded, black, dotted with white; the anterior and apical margins are beset with small whitish spots. Long. corp. 0.28; long. al. 0.26.

SYN. *Trypeta rotundipennis* LOEW, Monographs, etc., I, p. 79. Tab. II, f. 14.

Hab. Middle States (Osten-Saeken).

Observation.—Since the above-quoted description was drawn, I have not received any addition to the single, imperfect specimen in my collection, and have, therefore, nothing more to add about it. The systematic position of this species has been discussed above, in the third observation to *T. sparsa*.

31. *T. culta* WIED. ♂ ♀. (Tab. XI, f. 3.)—Ex rufo-lutea; caput nigro-maculatum; alæ luteæ, in margine antico toto, in apice et in marginis postici dimidio apicali eleganter radiatæ, in disco maculis aliquot magnis fusciscentibus, maculâ minutâ atrâ, guttisque aliquot limpidis, fusco-circumscriptis, notatæ, in angulo postico confertius limpido-guttatæ, venâ longitudinali tertiâ nudâ.

Reddish-yellow; head with black spots; wings clay-yellow, the anterior margin, the apex, and the apical portion of the posterior margin are handsomely adorned with ray-like streaks; upon the middle there are some brownish spots, a small black dot, and a moderate number of hyaline drops, margined with black; on the posterior angle numerous hyaline drops; the third longitudinal vein not bristly. Long. corp. ♂ 0.21, ♀ cum terebrâ 0.31; long. al. 0.29—0.32.

SYN. *Trypeta culta* WIEDEMANN, Auss. Zweifl. II, p. 486, 16.

Acinia fimbriata MACQUART, Dipt. Exot. II, 3, p. 228, 5. Tab. XXXI, f. 5.

Trypeta culta LOEW, Monogr. etc., I, p. 94. Tab. II, f. 29.

Reddish-yellow, opaque; the head somewhat paler yellow. The front of moderate breadth, dark yellow; the two bristles before the ocelli, directed forwards, and three strong bristles on the lateral margin of the front, are black; the other frontal bristles yellowish. The frontal lunule and the anterior part of

the lateral frontal border are shining; upon the first, almost without exception, a very small, deep black longitudinal dot is perceptible; near the antennæ, at the orbit of the eye, there is a deep black dot and a black spot in the middle of the posterior orbit. The face is deeply excavated, shining and sometimes with a distinct steel-blue reflection; upon its middle, below the antennæ, there is a rounded black spot, on each side an elongated, larger one, descending from the lower angle of the eye to the oral margin; the oral opening is very large, somewhat drawn upwards anteriorly. Palpi yellowish, broad, reaching to the anterior edge of the oral opening, with black pile at the tip, and with yellowish hairs elsewhere. Proboscis brown, sometimes yellowish-brown, rather stout, not geniculate. The thorax unicolorous, yellowish-red or reddish clay-yellow, opaque; the usual bristles, of which there are two pairs on the middle of the dorsum, are black, the short pile is pale yellowish. Scutellum somewhat paler yellow and rather shining, with erect yellowish bristle-like pile upon the middle and with four black bristles; the two apical ones are inserted upon black dots, while round the basis of the two anterior ones only a darker shade of the ground color is perceptible. The abdomen has the same coloring as the thorax and no spots, or only a trace of two longitudinal, contiguous rows of somewhat darker spots; all the pile and bristles upon it are yellowish and only a certain number of the bristles upon the posterior border of the last segment are usually blackish. The flat ovipositor is almost as long as the four posterior abdominal segments taken together, red, blackish towards the tip. Feet, as well as the bristles on the under side of the front femora, yellow; often, however, some of the bristles are black; the front femora have, a short distance before their end, on the outer side, a small black dot; the posterior femora, on the under side, have two black dots, the one before the middle, the other before the tip. The wings are rather long; their yellowish-red, almost gamboge-yellow color ends in rays along the anterior margin, the apex and the posterior portion of the hind margin; these rays are separated by hyaline intervals; between the humeral crossvein and the end of the auxiliary vein there are three narrow rays, running perpendicularly from the auxiliary vein to the costa, the first of which is less dark than the others; moreover, the extreme root and the extreme tip of that cell are marked by a blackish-brown

crossline; the short stigmatal cell, which is somewhat yellowish, is divided in two halves by a narrow dark brown line and is marked at both ends by a ray; in the marginal cell, besides a ray at the end of the first longitudinal vein, which is incompletely formed and margined with brown on its outer side only, there are three yellow rays, margined with brown and running towards the anterior margin; the first two are attenuated towards the margin and much narrower, the third is much broader; the five following rays are again so narrow, that only the first among them preserves a trace of the yellow coloring of its inner side; they gradually grow longer and end: the first at the tip of the second longitudinal vein, the next two between this and the third vein, the fourth exactly upon the tip of the third, the last a little before the tip of the fourth vein; the hyaline intervals between the last of these rays show upon their middle a faded cloud. The rays upon the latter portion of the posterior margin gradually grow shorter, are rather broad and altogether brown, but not as dark as the narrow rays of the anterior margin or the dark borders of the broader rays which follow upon the latter; they are five in number, or six if the last of them, which is very short, is counted for one; the second and third are less completely separated from each other than the rest, and the fifth, which includes the tip of the fifth vein, is the broadest of all. Upon the middle of the wing the following hyaline drops are visible: 1. Between the second and third longitudinal veins a very small one (sometimes a second one beyond it) below the end of the auxiliary vein and a second, somewhat larger one below the second ray, which runs, in the marginal cell, towards the anterior margin; 2. Between the third and fourth veins, nearer to the latter, there are three drops in a row; the middle one is nearly opposite the middle of the discal cell, the first one beyond the anterior end of this cell, and the last one at an equal distance before its posterior end; 3. In the second posterior cell only a single drop almost in its inner corner; 4. In the discal cell four or five, two of which upon its longitudinal axis (one near the anterior, the other near the posterior end) and three inconstant ones on the posterior margin of the cell (the first sometimes wanting, the second being the largest); sometimes a very small drop in the posterior corner of the discal cell is added to them. All these drops are encircled with dark brown or almost black, in such a manner, that this

dark ring becomes paler round those drops which are more distant from the anterior margin. The convex spot in the first posterior cell is rather large; it contains a comparatively small rounded-ovate deep black dot. Moreover, in the submarginal cell, in the first and second posterior cells, and in the discal cell, differently colored spots (one in each) may be noticed, which, at an oblique view, assume a dark coloring. In the third posterior cell, in the posterior angle of the wing, and on the alula, there is a number of hyaline drops, among which only those placed immediately behind the fifth vein show a trace of a brown border. The double costal spine is strong and comparatively long, the small crossvein is placed upon the last third of the discal cell; the posterior crossvein is steep, but distinctly sinuate; the third longitudinal vein is not bristly.

Hab. Savannah (Wiedemann); Carolina (Macquart); Texas (Belfrage).

Observation.—*T. culta* is closely allied to the European *T. pupillata* Fall. and *strigilata* Lw., and this relationship is sufficient to justify its location in the genus *Carpotricha*, formed by me for the reception of these species, as well as of *T. guttularis* Meig. However, in consequence of this addition, the definition of the genus, as given by me in the Monograph of the European Trypetæ, will have to be somewhat modified. In *T. culta* the scutellum is less convex, and, although smooth, it is not polished; the tip of the abdomen is not shining. The nature of the pile and the pattern of the picture of the wings, the structure of the head, and the arrangement of the frontal bristles furnish sufficient data for the modification alluded to.

32. *T. solidaginis* FREN. ♂ ♀. (Tab. X, f. 16.)—Sordide ferruginea, capite pedibusque luteis; frons latissima; scutelli valde convexi setæ duæ; alæ fusco-reticulatæ, apice incisuræque tribus. unâ marginis antici duabusque postici, hyalinis et parce fusco-maculatis.

Of a dingy ferruginous-red; head and feet clay-yellowish; front exceedingly broad; scutellum very convex, with two bristles; wings reticulate with brown; the tip and three indentations, one on the anterior and two on the posterior margin, hyaline, sparsely dotted with brown. Long. corp. ♂ 0.24—0.25, ♀ cum terebrâ 0.26—0.28; long. al. 0.25—0.26.

Syn. *Tephritis asteris* HARRIS, Ins. Injur. to Veg., 3d Edit., p. 620.

Acinia solidaginis FITCH, First Report, 66.

Trypeta solidaginis LÖEW, Monographs, etc., I, p. 82. Tab. II, f. 16.

Hab. New York (Fitch); Washington (Osten-Sacken); New England (Harris). [Canada.—O. S.]

Observation 1.—To the description of this species in the *Monographs*, Vol. I, I may add, by way of correction, that the costal spine of the wings is not altogether wanting, but that it is very short and weak, and hence, in some specimens, hardly visible. The words "the first longitudinal vein alone being hairy," in the observation to the above description, only meant that the bristles upon that vein were more like hairs, and not that this vein alone is provided with bristles; the third vein also, bears weak, hair-like bristles.

Observation 2.—Baron Osten-Sacken, having seen the original specimen of *Tephritis asteris* Harris in Mr. Harris's collection in the museum of natural history in Boston, has settled its identity with *Acinia solidaginis* Fitch. Harris's name, although based upon an error in the name of the plant upon which this fly undergoes its transformations, would have to be retained, but for the circumstance that Mr. Haliday had previously used it for another European *Trypeta*.

Observation 3.—Among the genera established for the European *Trypetina*, *Oxyphora* is the only one in which *T. solidaginis* might, perhaps, be placed. Among the European species *Oxyphora Schæfferi* Fruf. is nearest to it in its general appearance; the outline of the wings reminds somewhat of *O. Westermanni*. The much heavier body, the strikingly broad front, and the much broader checks, as well as the peculiar shape of the wings, which are broadly rounded at the tip, the heavy, conical, not at all flattened ovipositor of the female, isolate this species sufficiently to justify the formation of a new genus, for which I propose the name of *Eurosta*.

33. T. comma WIEN. ♀. (Tab. XI, f. 2.)—Sordide rufa aut fusca, capite magno, thoracis dorso, tibiis, tarsisque lutescentibus; alæ obtusæ, ex fusco nigræ, guttulis minutis modice dilutioribus adpersæ, maculâ costali trigonâ comma fuscum includente, limbo apicis angustissimo, guttulisque aliquot confertioribus prope venâ longitudinalem sextæ apicem, hyalinis; venâ longitudinalem tertiâ setosâ; scutellum setis duabus instructum; terebra conica, non depressa.

Dingy red or brown, head large, thoracic dorsum, tibiae, and tarsi clay-yellowish; wings obtuse, brownish-black, covered with small, moderately limpid drops; a triangular indentation on the costa contains a brown comma; a narrow border along the apex and a dense cluster of drops near the tip of the sixth vein, are hyaline; the third longitudinal vein is bristly; scutellum with two bristles; ovipositor conical, not flattened. Long. corp. ♀ cum terebrâ 0.32—0.34; long. al. 0.30—0.31.

SYN. *Trypeta comma* WIEDEMANN, AUSS. Zweifl. II, p. 478, 4.

Acinia comma MACQUART, Dipt. Exot. II, 3, p. 229, 6.

Trypeta comma LOEW, Monographs, etc., I, p. 93. Tab. II, f. 28.

This conspicuous species was described by Wiedemann from a very pale-colored specimen, which I have had occasion to examine. The coloring varies from a dingy brick-red almost to dark brown; the abdomen especially is often dark. The large head is yellow; the front is more than half as broad as the head, usually of a darker yellow; the usual bristles upon it are brown or brownish, weak, and rather short. Antennæ clay-yellow, very short, not even reaching to the middle of the face. Face perpendicular, very little excavated; oral opening of a very moderate size, and the anterior edge of the mouth not projecting; ocular orbits very broad. Eyes elongated, but the cheeks of a considerable breadth, although by far not equalling those of the preceding species; the pile upon them is brownish or brown, sometimes paler; proboscis short, not geniculate; the clay-yellowish palpi broad, reaching to the anterior edge of the oral opening. The upper side of the thorax covered with a thick clay-yellowish pollen and with short, dense clay-yellowish pile; the latter sometimes has a more ferruginous tinge; the usual bristles of the thoracic dorsum are brown and weak; upon its middle there are only two pairs, the anterior one very much behind the transverse suture; it is weaker and shorter than the posterior one. Scutellum dark brown, very convex, with only two bristles. Metathorax and pleuræ are sometimes brick-red, sometimes brown or blackish-brown; the darker the pleuræ are, the darker the bristles upon them. Abdomen unicolorous, brick-red, brown, or brownish-black, with rather delicate blackish or black pile. Ovipositor not compressed, conical, about as long as the last two abdominal segments taken together, with delicate black pile; in paler specimens the ovipositor is red, the extreme tip only black; in very dark specimens it is black with a reddish crossband upon the middle. Very dark specimens have blackish-brown femora; their

tip and the tibiae are yellowish-brown, the tarsi dirty yellowish; in paler specimens tibiae and femora are not much darker than the tarsi; front femora with black bristles; tarsi, especially their first joint, somewhat longer than usual, especially in *T. solidaginis*. Knob of the halteres blackish or black. The wings broad and very obtuse at the end, blackish-brown or black, including the extreme root; upon their whole surface are a very variable number of very small dots of but moderate transparency; upon the anterior margin, immediately beyond the stigma, there is a triangular hyaline spot, the tip of which does not quite reach the third longitudinal vein and which includes a blackish-brown crossline, extending from the costa to the second longitudinal vein; the end of the sixth vein is surrounded by a cluster of small, more or less coalescent drops, which extends especially on the anterior side of this vein; the extreme tip of the wing has a very narrow hyaline border, which begins a little before the tip of the third longitudinal vein and ends beyond the tip of the fourth vein; at the tip of these veins the border is very often interrupted; on the posterior margin of the wing there are often two, sometimes three or four, in such a case larger, hyaline drops. The third longitudinal vein is beset with scattered but distinct bristles; at its end, it is strongly bent backwards so that its divergency from the second vein is unusually large; the latter ends rather far from the apex of the wing; the crossveins are but little approximated, the small one is oblique, the posterior one arcuated.

Hab. Kentucky (Wiedemann); Maryland (Osten-Sacken).

Observation 1.—This species is subject to remarkable variations in the coloring of the body, as well as in the shape of the wings; the tip of the latter is sometimes more, sometimes less distinctly obtuse; all these differences certainly do not constitute specific distinctions. The figure which I have given in the first volume of these Monographs was prepared from a specimen in the Berlin Museum, and as it is based upon a rather hasty pencil sketch, made many years previously, it lays no claim upon an absolute fidelity. This figure shows some discrepancies however, which raise a suspicion that this Berlin specimen is not *Trypeta comma* at all, but a closely allied species.

Observation 2.—*Trypeta comma* differs from *T. solidaginis* in its larger eyes, a less excavated face, and a smaller and much

narrower oral opening; the shape of the body, the striking breadth of the forehead, the distribution of the bristles upon it and upon the thoracic dorsum and scutellum, the shape of the ovipositor, the outline of the wings, and the pattern of the picture are remarkably analogous in both species, so as to preclude a generic separation.

34. *T. latifrons* Lw. ♀. (Tab. X, f. 22.)—Obscura, capite, tibiis tarsisque lutescentibus, fronte latissimâ, scutello convexo, setis duabus instructo, alæ latiusculæ, colore fusco-nigro pictæ, in disco parvius et subæqualiter reticulatæ, in dimidiâ apicalis margine radiatæ.

Coloring dark; head, tibiæ, and tarsi clay-yellowish, front unusually large; the convex scutellum with two bristles only; wings rather broad, with a brownish-black picture, upon their middle somewhat sparsely and not very evenly marked with hyaline drops, their apical border radiate. Long. corp. ♀ enim terebrâ 0.30; long. al. 0.27.

SEX. *Trypeta latifrons* Loew, Monographs, etc., I, p. 89, 22. Tab. II, f. 22.

Hab. Carolina (Zimmerman); Connecticut (Norton).

Observation.—A female from Connecticut, communicated to me by Baron Osten-Sacken, is not much better preserved than the female from South Carolina, from which my description in the Monogr. Vol. I was drawn, and for this reason I am not able to give a better one here. Of the two pairs of bristles upon the thoracic dorsum the anterior one has dropped off; it seems to have been inserted rather far behind the transverse suture. The structure of thorax and abdomen, the broad front, the bisetose scutellum, and the conical, not at all flattened, ovipositor, indicate a relationship between this species and the two preceding ones, from which, however, it differs in the shape of the wings and the pattern of the picture. In the latter two points it reminds one of *Trypeta platyptera* Lw., which differs again in the more narrow front, a four-bristled scutellum, and a flattened ovipositor. Such being the case, we will be better justified in connecting this species with *T. solidaginis* and *comma*, than with *T. platyptera* and its congeners.

35. *T. melanura* n. sp. ♀. (Tab. XI, f. 6.)—Lutea, metanoto, abdominis maculis in series quatuor dispositis et terebrâ brevi, atris; caput lætius luteum, fronte latissimâ, facie modice recedente, antennis longis et acutis; femora anteriora macula minutâ nigrâ notatæ; alarum pictura fusca, guttis majusculis hyalinis reticulatâ, quarum in cellulâ posteriore secundâ tres, in tertiâ quatuor conglobatæ.

Clay-yellow; the metanotum, four rows of abdominal spots, and the short ovipositor, deep black; head of a brighter clay-yellow; front very broad, face moderately receding; antennæ long and acute; the anterior femora with a little black spot; picture of the wings brown, reticulate with rather large hyaline drops, among which three form a cluster in the second posterior cell and four in the third. Long. corp. ♀ cum terebrâ 0.13—0.14; long. al. 0.14.

Head almost ochreous-yellow, the rather level face, somewhat retreating on the under side, the moderately broad checks, and the lower portion of the occiput pale yellowish. Front more than half as broad as the whole head. Frontal lunule very flat. Third antennal joint unusually long, with a remarkably sharp anterior corner; the thin, bare arista is incrassated at its basis for a short distance only. Oral opening rather large, rounded, but somewhat broader than long; its anterior edge is neither drawn upwards, nor projecting in the profile. Proboscis and palpi yellowish, withdrawn in the oral opening. The pile on the head is ochreous-yellow; the ordinary frontal bristles are brownish or brown. The ground color of the thoracic dorsum is blackish, with the exception of the pale yellow humeral callus, but very much concealed under ochre-yellow pollen, and reddish ochre-yellow, coarse, and almost stubble-shaped pile. When the thorax is viewed from behind, several opaque black, punctiform dots become apparent, especially two on the transverse suture and two larger ones between the first and the posterior border. The bristles of the thoracic dorsum are partly pale yellow, partly brown; viewed against the light, they appear dark. The scutellum, which, in the described specimen, is much damaged, seems very convex; it is smooth and for the most part yellow; among its four bristles, the two apical ones are inserted on small black dots. The abdomen is reddish-yellow or almost honey-yellow and somewhat shining; upon the second segment there are four black dots in a row, the lateral ones of which are small; upon each succeeding segment the lateral spots become larger, and upon the fifth segment the lateral spots completely coalesce with the middle ones, only a median reddish line being left on the segment. The flat, shining black ovipositor is hardly longer than the last abdominal segment. Feet rather dark ochre-yellow; the front and intermediate femora have, upon their hind side, beyond the middle, a little black spot. The reticulate picture of the wings is brown, black-

ish-brown within the stigma; the hyaline drops, appearing in 2 different light whitish, and which perforate the brown coloring, are generally large, but not numerous; the stigma contains but a single yellowish drop; its extreme basis also has a narrow hyaline border; the triangular cluster of larger drops which occurs on the anterior margin, immediately beyond the stigma, consists here of five drops, absolutely separated from each other; the end of the marginal cell contains but a single small drop; a larger drop occurs below the end of the second longitudinal vein and a similar one under it, in the first posterior cell; between these two drops and the apex of the wing there are four smaller drops, forming a somewhat arcuated crossband; especially characteristic for the species are three conspicuous drops in the second and four similar ones in the third posterior cell, between which the brown coloring is so pale or faded, that they appear almost coalescent; (this is not well expressed in the figure, which is kept altogether in too dark a shade); upon the middle of the discal cell there is a large drop, occupying its whole breadth. The third longitudinal vein is distinctly bristly about as far as the small crossvein; this crossvein corresponds to the last third of the discal cell; the posterior crossvein is straight and very perpendicular.

Hab. Distr. Columbia (Osten-Sacken).

Observation.—In several respects this species resembles the European species of *Carphotricha*; but, on account of the striking breadth of the forehead, the unusual length of the antennæ, and the comparatively very even face, somewhat retreating below, it cannot well be placed in that genus, especially when *T. culta* Wied. is admitted in it, on account of its rather close relationship to *Carphotricha pupillata* Fall. As I know of no other species with which the present one could be generically united, I prefer to establish a separate genus for it, which I call *Acidogona*.

36. *T. alba* Lw. ♂ ♀. (Tab. XI, f. 11.)—Albida, alis concoloribus immaculatis, capite, pleuris, scutello segmentorumque abdominalium singulorum margine postico pallide sulphureis, antennis, terebrâ, pedibusque luteis.

Whitish, with whitish, altogether immaculate wings; head, pleuræ, scutellum, and the posterior margin of the single abdominal segments, sulphur-yellow; antennæ, ovipositor, and feet clay-yellow. Long. corp. ♂ 0.13, ♀ cum terebrâ 0.17; long. al. 0.15—0.16.

Syn. *Trypeta alba* Loew, Berl. Entom. Zeitschr. V, p. 345, 72, lb., Dipt. Amer. Cent. I, p. 39, 72.

Trypeta alba Loew, Monographs, etc., I, p. 100, 18.

Hab. Pennsylvania (Osten-Sacken).

Observation 1.—I have only the following remarks to make concerning this species, described in the above-quoted places and easily recognizable. The antennæ are often not clay-yellow, but more or less bright ochre-yellow, which is especially the case in the best preserved and fully colored specimens; in such specimens the face is pale sulphur-yellow, while, on the contrary, the front, probably in consequence of desiccation, shows, in other specimens, a more dark yellow, often impure, hue.

Observation 2.—This and the next following species show a striking agreement in all plastic characters, especially in the structure of the head, and the characteristic outline of the wings, so that they may be considered as the types of a new genus, distinguished from the related ones by the above-mentioned characters, and which may be called *Aspilota*.

37. T. albidipennis Lw. ♂ ♀. (Tab. XI, f. 10.)—Nigro-cinerea, thoracis dorso albicans, capite, thoracis vittâ laterali scutelloque sulphureis, alarum albidarum stigmatibus fusco, terebrâ femineâ atrâ.

Blackish-gray, thoracic dorsum whitish; head, a stripe on the lateral margin of the thorax, and the scutellum sulphur-yellow; wings whitish with a brown stigma; the ovipositor of the female black. Long. corp. ♂ 0.17, ♀ cum terebrâ 0.20; long. al. 0.18—0.19.

Syn. *Trypeta albidipennis* Loew, Berl. Entom. Zeitschr. V, p. 345, 73, and Dipt. Amer. Cent. I, p. 39, 73.

Trypeta albidipennis Loew, Monographs, etc., I, p. 100, 19

Hab. Pennsylvania (Osten-Sacken).

Observation.—The antennæ are usually more ochre-yellow than ferruginous-yellow. The generic location of this species has been mentioned in the note to the preceding one.

38. T. Vernoniæ Lw. ♂ ♀. (Tab. XI, f. 8.)—Dilute lutea, capite, thoracis vittâ marginali in pleuras dilatâ, scutelloque purius flavis, thoracis dorso subhelvo, metanoto nigro; alarum dimidium basale impictum, apicale colore subfusco grosse reticulatum, guttis magnis confluentibus, ita ut fasciæ tres valde inæquales fuscæ conspiciantur; primâ incompletâ et obsolete, secundâ integrâ, tertiâ postice abbreviatâ.

Pale clay-yellowish; head, a lateral thoracic stripe, dilated upon the pleuræ, and the scutellum of a purer yellow, thoracic dorsum more isabelle-yellow, metanotum black; the basal half of the wings is inma-

culate, the apical half shows a very coarse brownish reticulation, the large hyaline drops of which coalesce in such a manner, that three brown, irregular crossbands are formed; the first is only incompletely developed and rather faded, the second complete, the third abbreviated posteriorly. Long. corp. ♂ 0.18, ♀ cum terebrâ 0.22; long. al. 0.17—0.18.

SYN. *Trypeta Vernoniæ* LOEW, Beil. Entom. Zeitschr. V, p. 346, 74, and Dipt. Amer. Cent. I, p. 40, 74.

Trypeta Vernoniæ LOEW, Monographs, etc., I, p. 101, 20.

HAB. Pennsylvania (Osten-Sacken); on the iron-weed (*Vernonia*).

Observation.—*T. Vernoniæ* agrees in all the plastic characters, especially in the structure of the head and the shape of the wings, with the two preceding species in a very striking manner, and the presence of a picture on the wings alone is not a sufficient ground for a generic separation.

39. *T. seriata* Lw. ♂. (Tab. X, f. 18.)—Lutea, alis concoloribus, totis aequalibus et obtusis, per maculas minutas fuscas seriatim dispositas reticulatis, adversus marginem præter trientem basalem nigricantibus, venâ longitudinali tertîa setosâ.

Clay-yellow; wings of the same color, of a very equal breadth, obtuse at the end, reticulate with small brown spots arranged in rows; blackish along the margin, except on the proximal third of its extent; third longitudinal vein bristly. Long. corp. 0.24; long. al. 0.26—0.27.

SYN. *Trypeta seriata* LOEW, Monographs, etc., I, p. 84. Tab. II, f. 18.

HAB. Illinois.

Observation.—Should *T. seriata* be placed in one of the genera established for the European *Trypetina*, it would of course be the genus *Oxyphora*, the most characteristic marks of which are the reticulate wings and the bristles on the third vein. And, indeed, this species reminds one very much of *Oxyphora Westermanni* Meig. in the very peculiar shape of the wings, and even in the coloring of the body and the pattern of the picture of the wings. But when we bear in mind that this European species occupies in the genus a very isolated, in fact an artificial position,¹ it will appear more natural to withdraw *O. Westermanni* from the genus and to form a new genus of it, together with the above described as well as the next following American species. This genus may be called *Icteriva*.

¹ The European *Oxyphora Schafferi* Egger shares this exceptional position, though for other reasons.

40. *T. circinata* n. sp. ♂ ♀.—Lutea, alis concoloribus, totis æqualibus et obtusis, per circulos fuscios inter se cohærentes reticulatis, adversus marginem præter dimidium basale nigrescentibus, venâ longitudinali tertiâ setosâ.

Clay-yellow, wings of the same color, of very equal breadth, obtuse at the end, reticulate with small, brown, contiguous circles; infuscated along the margin, except upon its first half; the third longitudinal vein bristly. Long. corp. ♂ 0.24, ♀ cum terebrâ 0.27; long. al. 0.26—0.27.

The resemblance of this species to *T. seriata* is so striking that one would almost be tempted to take it for a mere variety. However, the picture of the wings, perfectly identical in both sexes, shows such differences from that of *T. seriata*, as occur in closely allied species, but not in a variety of the same species. While the reticulation of *T. seriata* consists of small, angular brown spots, arranged in double rows between each pair of longitudinal veins on the middle of the wing, in the present species the spots are replaced by small brown ringlets, mostly closed, but some of them open, and connected with each other. The infuscated portion of the anterior margin in *T. seriata* begins before the end of the auxiliary vein and fills the stigmal cell entirely, with the exception of a but little perceptible clay-yellow drop at the tip, and a similar, obsolete drop at the basis; between the ends of the first and second longitudinal veins there are, besides the somewhat hyaline spot immediately beyond the former, only two brownish-yellow drops near the anterior margin. In *T. circinata* the extreme tip of the auxiliary vein and the spot on the costa corresponding to it are black, but there is no trace of dark coloring in the costal cell before the tip of the auxiliary vein; the stigma is rather saturate yellow, and has upon its middle a considerable rectangular black spot; the pale spot which follows immediately upon the tip of the first longitudinal vein is more extensive, but less limpid, and the two drops which lie between it and the second longitudinal vein are much larger and more limpid, so that they entirely interrupt the black border along the anterior margin. A similar interruption is caused by a drop immediately beyond the tip of the second longitudinal vein, which is entirely wanting in *Trypeta seriata*. By these complete breaks in the black anterior border *Trypeta circinata* is very easily distinguished from *Trypeta seriata*, which has only one break of this kind immediately beyond the apex of the first longitudinal vein.

Hab. New York (Mr. Akhurst).

41. T. Lichtensteinii WIED. ♂. (Tab. XI, f. 9.)—Tota lutea; alae dilute cinereo-hyalinae, guttis majusculis albicantibus, maculisque tribus fusco-nigris variegatae, primâ harum reliquis minore et a stigmate oblique decurrente, secundâ quadrangulâ et venam transversam posteriorem includente, tertiâ denique primis duabus majore et apicem alae cingente.

Altogether clay-yellow; wings grayish-hyaline, with rather large whitish drops and three brownish-black spots, the first among which is smaller than the others and descends from the stigma in an oblique direction, the second is square and includes the posterior crossvein, and the third is larger than the two preceding ones and forms a border along the apex. Long. corp. 0.22; long. al. 0.23.

Syn. *Trypeta Lichtensteinii* WIEDEMANN, Auss. Zweifl. II, p. 497, 31.

Trypeta Lichtensteinii LOEW, Monographs, etc., I, p. 92. Tab. II, f. 25.

Clay-yellow, the pile on head, thorax, and feet yellowish; the bristles yellow or yellowish-brown, according to the light in which they are seen; the pile on the abdomen yellowish at the basis only, black elsewhere. Front of a more vivid yellow, rather broad, with long bristles, the eyes rather large, oval; cheeks of a medium breadth. The face rather retreating, somewhat excavated under the antennæ; the anterior edge of the mouth not projecting in the profile. Antennæ yellow, of medium length; the third joint with a rounded anterior corner; the rather long arista is much incrassated at its extreme basis, otherwise very thin and bare. Oral opening rather large, rounded; palpi and proboscis not projecting beyond it; the latter not geniculated. The middle of the upper side of the thorax seems to have borne only two pairs of bristles. The very moderately convex scutellum bears four bristles. Scutellum and abdomen are more shining than the thoracic dorsum, which is opaque in consequence of a yellowish pollen; abdomen without any picture. Wings rather long and of nearly equal breadth; the third longitudinal vein distinctly bristly for a considerable portion of its length; crossveins straight and steep; small crossvein a little beyond the middle of the discal cell. The picture of the wings is a very peculiar one; its principal feature consists of three very conspicuous brownish-black spots; the smallest among them has the shape of an oblique, somewhat irregular half-crossband; with its anterior end it covers the tip of the stigma, with its posterior end it covers the small crossvein and suddenly stops near the fourth vein; the second spot, which covers the posterior crossvein, has a square shape, is

higher than broad and reaches from the fourth vein to the posterior margin; the third spot forms a broad margin of the tip of the wing, which begins not far beyond the first longitudinal vein and, gradually increasing in breadth, reaches beyond the beginning of the second posterior cell. The outlines of these three spots are irregular and sinuate. The remaining surface of the wings is grayish-hyaline; held against the light this grayish surface shows some round, whitish spots of a rather considerable size, occurring especially within the sinuosities along the margins of the dark spots, however, without following their outline exactly. In some places the grayish tinge of the wings becomes infuscated, thus forming several other, probably very variable, spots; the typical specimen shows the following ones: a narrow little spot in the middle of the anterior margin of the costal cell; a hook-shaped spot, which begins at the anterior end of the third brown spot and runs to the second vein; a small, thimble-shaped spot, situated on the fourth vein, a little beyond the posterior crossvein and directed forwards; a little spot upon the posterior margin, in the middle between the second and third of the large brown spots; a punctiform dot upon the middle of the discal cell; a larger spot, behind the preceding one, within the third posterior cell; finally, behind the latter, upon the posterior margin, another small, faded, little spot. It is probable that, sometimes, the greatest part of the grayish surface becomes brownish, and then it may happen that, in some specimens, beyond the root of the wing, but little pale colored portions remain, except the large drops with a whitish reflection. The fact that the described specimen does not seem to be a fully matured one, serves to confirm this supposition.

Hab. Mexico (Wiedemann).

Observation 1.—Description and figure are prepared after the same specimen in the Berlin Museum, which Wiedemann had before him in drawing his description. In the figure, the engraver has represented the large whitish drops somewhat more vividly than they appear in nature. The relationship of *T. Lichtensteinii* to the two preceding species, is close enough to enable us to place it in the genus *Ictericia*.

Observation 2.—Among the species described in the sequel, *Trypeta æqualis* (Tab. X, f. 20) stands next to the species of *Ictericia* in the shape of the wings. But, besides the fact that

its wings are neither as equally broad, nor as obtuse, as those of the species united in the genus *Ictericia*, that species differs also in the absence of bristles upon the third vein.

42. *T. humilis* Lw. ♂ ♀. (Tab. X, f. 17.)—Luteo-cinerea, capite pedibusque saturate flavis, femoribus tamen nigris adversus apicam in mare late, in fœminâ latissime flavis; peristomium valde productum, proboscis geniculata, alæ rare reticulatæ, stigmatæ atro, non guttato.

Yellowish-gray; head and feet saturate yellow; the femora black, a considerable portion at their tip in the male, a still more considerable one in the female, yellow; edge of the mouth very much produced, proboscis geniculated, wings sparsely reticulate, the black stigma without pale drops. Long. corp. ♂ 0.09—0.1, ♀ cum terebrâ 0.11—0.12; long. al. 0.11—0.12.

Syn. *Acinia picciola* Bigot, R. de la Sagra, Hist. Fis. Vol. VII. Tab. XX, f. 10.

Trypeta humilis Loew, Monogr. etc. I, p. 81. Tab. II, f. 17.

Hab. Cuba (Poey, Gundlach). [Key West; communicated by Mr. Burgess. O. S.]

Observation 1.—The saturate yellow coloring of the apex of the femora in the male has a rather considerable, but at the same time variable, extent; in the female, the yellow sometimes occupies so much space, that the blackish color remains visible at the basis of the femora only. Females with the femora as pale as that, mentioned by me in the first part of these Monographs, seem to be rare, as among the numerous specimens of my collection that single one only is to be found.

Observation 2.—To recognize the present species in the *Acinia picciola* Bigot is not possible. Nevertheless the synonymy is not doubtful, as, through the kindness of Mr. Gundlach, I have been put in possession of numerous typical specimens. It is to be regretted that Mr. Bigot has given the species a name which cannot possibly be admitted, unless names like *littella*, *petitella*, *kleinella* for any small species were likewise tolerated.

Observation 3.—The strongly produced oral edge and the strikingly geniculated proboscis, with its very much prolonged flaps, reaching backwards as far as the mentum, define this species as an *Ensina*. As soon as exotic species are taken in consideration, this genus cannot be maintained within exactly the same limits which I defined for it in my Monograph of the European species. A part of the species, which I placed there under

the head of *Oryna*, as for instance *Oryna elongatula* Lw., and its congeners, will have to be admitted in the genus *Eusina*.

Observation 4.—A Brazilian species, not rare in collections, likewise belonging to *Eusina*, is so very like *humilis*, that I give here its description, in order to avoid a possible confusion.

T. peregrina n. sp. ♂ ♀. (Tab. X, f. 30.)—Luteo-cinerea, abdomine nigro-maculato, genis angustissimis, peristomio eximie producto, proboscidis geniculatæ labellis longissimis, alis elongatis et subæqualiter fusco-reticulatis; pedes lutei, basali femorum posteriorum dimidio piceo; terebra feminae atra, tribus ultimis abdominalis segmentis simul sumtis longiore.

Yellowish-gray, abdomen spotted with black; the cheeks very narrow, the oral edge very much produced, the flaps of the geniculated proboscis very much prolonged; wings comparatively long and rather uniformly reticulated with brown; feet of a saturate yellow, basal half of the hind femora black; ovipositor of the female black, larger than the last three abdominal segments taken together. Long. corp. ♂ 0.12—0.13; ♀ cum terebrâ 0.14—0.16; long. al. 0.13—0.14.

Resembles *T. sororcula* Wied. from Teneriffe and the European *T. elongatula* Lw. very much, both in the structure of the body and in general appearance. In the female sex, it differs from the latter easily by its ovipositor, which is once and a half as long; the male is easily distinguished by several features of the picture of the wings, which in other respects is very much the same: namely, the drop which lies at the tip of the submarginal cell is not present in *T. elongatula*: in the dark coloring at the extreme end of the discal cell there is only a single hyaline drop, while in *T. elongatula* there are several of them, usually three. From *T. humilis* it differs sufficiently in the scutellum, which is tinged with yellow at the tip, in the coloring of the feet and in the picture of the wings. Yellowish-gray; the head, of the same structure as in the species just compared with it, rather saturate yellow, as well as antennæ, palpi, and proboscis; the occiput alone in part gray. Front long and not very broad; along the orbit with a narrow, rather whitish border. Antennæ rather broad, not quite descending to the anterior edge of the mouth, which is somewhat drawn upwards and remarkably projecting in the profile. Eyes rounded; cheeks very narrow. Oral opening very much drawn out; the very elongated flaps of the geniculated proboscis reach backwards to the mentum. The usual bristles of the front, the thorax, and the scutellum are black; the latter is yellow at its tip only. The abdomen is of the same color as the thorax, and bears, like the latter, some short, pale yellowish pile, while the longer hairs on the posterior border of the last segments are black. The flattened and only moderately pointed ovipositor is shining black and a little longer than the last three abdominal segments taken together; its short pubescence is almost without exception black.

Feet dark yellow, only the hind femora are brownish-black beyond their middle, and the other femora somewhat infuscated near the root and with a brown stripe on the under side. The wings are elongated, hyaline, with a grayish-brown, very loose, but not disconnected, reticulation; the root of the wings is not spotted up to the end of the small basal cells; beyond this, up to the stigma, there are only three inconspicuous grayish spots. The grayish-brown stigma contains a rather conspicuous hyaline drop (represented too small on the figure); a spot adjoining it, comparatively small and not much perforated, reaches beyond the second vein with two points only, and contains a little drop immediately before the second vein. The larger and less perforated spot before the end of the second longitudinal vein always contains a considerable hyaline drop near the anterior margin; between the second and third longitudinal veins, the same spot contains two or three small drops and is variously connected with the remaining reticulation. Between these two less perforated spots, there are, in the marginal cell two, in the submarginal three, large hyaline drops, which generally assume the shape of quadrangular spots, and are only separated by grayish-brown lines, running from one longitudinal vein to the other. Upon the remainder of the surface of the wing, the reticulation is formed by rather considerable rounded drops, and is more regular; only in the proximity of the posterior crossvein there are no drops.

Hab. Brazil.

43. *T. angustipennis* Lw. ♂ ♀.—*Cinerea*, capite pedibusque flavis, femoribus inaguâ et parte nigris vel fuscis; proboscis non geniculatâ; alæ subangustatæ, nigro-reticulatæ, in basi et limbo marginis postici subinmaculatæ, stigmatæ non guttato, maculis duabus ordinariis obscurioribus mediocribus, separatis, secundâ guttulam unicam, rarius duas includente; terebra fœminæ atra, duobus ultimis abdominis segmentis simul sumtis subæqualis.

Gray; head and feet yellow; femora for the most part black or brown; proboscis not geniculated; wings reticulate with black, almost without spots at the basis and in the vicinity of the posterior margin; the two ordinary dark spots only of middle size and separated from each other; in the second, one, rarely two, hyaline drops; ovipositor black, almost as long as the last two abdominal segments taken together. Long. corp. ♂ 0.13, ♀ 0.14—0.15; long. al. 0.14.

Syn. *Tephritis Leontodontis* ZETTERSTEDT, *Ins. Lapp.* 745, 6. *Var. a.* (ex p.).

Trypeta angustipennis LOEW, *Germ. Zeitschr.* V, p. 382. Tab. II, f. 4.

Tephritis angustipennis ZETTERSTEDT, *Dipt. Scand.* VI, p. 2229, 35.

Tephritis angustipennis LOEW, *Trypetidæ*, p. 113, No. 24.

Tephritis segregata FRAUENFELD, *Verh. Zool. Bot. Ges.* XIV, p. 147.

Gray; thorax without picture; the pile upon it is whitish; the bristles black. Abdomen blackish-gray, without spots; the pile

whitish, only the bristles upon the posterior margin of the last segment are black. Ovipositor black, hardly as long as the last two segments taken together; with distinct whitish pile upon its anterior half. Feet yellow; the femora for the most part black or brown. The wings are comparatively a little longer and narrower than in most of the related species. The rather dark reticulation is loosely meshy and somewhat disconnected; it disappears almost entirely in the region of the posterior margin, with the exception of a few little spots, which distinguishes this species from the otherwise related ones; the black stigma does not include a hyaline dot; the two ordinary dark spots are of moderate size; the first is connected with the stigma and reaches from it directly backwards; the second usually contains, near the anterior margin, only a single hyaline drop, which lies immediately beyond the tip of the second longitudinal vein; this spot reaches as far as the fourth longitudinal vein; the two rays which, in the related species, run from this vein over the second posterior cell to the posterior margin, are incomplete or wanting; the posterior crossvein also has only a comparatively narrow dark border, which sometimes exists on its posterior half only; upon the posterior part of the crossvein, this border emits a short branch, characteristic for this species, and reaching into the discal cell; this branch sometimes coalesces with a second similar branch upon the posterior side of the fourth vein, so as to include a hyaline drop; otherwise the picture of the discal cell is limited to a small crossband, lying beyond its middle, or there is sometimes before it, near the anterior margin of the cell, another dark spot, which in some specimens becomes a second small crossband; upon the posterior side of the fifth vein generally two small, dark spots of variable size are observable, of which the one nearer the root of the wing is often wanting.

Hab. Yukon River (Kennicott).

Observation 1.—I cannot distinguish this species from the *T. angustipennis* occurring in Scandinavia; the typical pair after which I have described it in *Germer's Zeitschrift* has, it is true, the femora much less dark, but as the specimens seem to be immature, I do not consider this a specific difference. The figure given in *Germer's Zeitschrift* has not well succeeded in the engraving and gives only an approximate idea of the picture of the wings.

Observation 2.—Should we distribute the present and all the next following species among the subgenera which I have established for the European *Trypetidae*, they would have to be referred to the genera *Oxyphora*, *Oxyga*, *Tephritis*, and *Urellia*. The genus *Urellia* is easily distinguished from the others by the picture of the wings: it consists in a conspicuous star-shaped black design near the apex, while the rest of the wing is altogether immaculate, or is marked with only a few isolated spots, at the utmost with a very pale reticulate picture. A part of the species described in the sequel, can undoubtedly be referred to *Urellia*. Among the remaining species, those would have to be located in the genus *Oxyphora*, which have the third longitudinal vein of the wings beset with bristles. This character is of a very easy application when a number of well-preserved specimens is at hand, but it becomes of much less value when applied only to single and indifferently preserved specimens. For this reason I am not quite sure whether in all the species in which I have not been able to discern the presence of bristles on the third vein, they are really wanting; and hence, with the materials I now possess, I am not able to refer with certainty to *Oxyphora* the North American species which may belong to it. Among the North American species with a distinctly bristly third longitudinal vein, *T. geminata* alone comes near the European species of *Oxyphora*, while *T. timida* is more related not to the former, but to the European *T. guttata* Fall., and to the American *T. tennis*, *melanogastra*, and *mexicana*, in which I am unable to discern the bristles upon the third vein. Thus, the maintenance of the genus *Oxyphora* for those species only which have bristles upon the third vein, would separate from each other species most closely allied. In order, therefore, to make this genus applicable to the North American species, we should exclude from it all the species the picture of the wings of which ends in distinctly developed rays, in which case only *T. geminata* would remain in it. Theoretically there is no objection to such an arrangement; practically, however, there remains the difficulty of ascertaining positively the presence of bristles upon the third vein in all the specimens which I have at hand, and this difficulty compels me to drop entirely the genus *Oxyphora* for the present. Should we follow the suggestion already made above, of removing from the genus *Oxyga* those species which have remarkably prolonged

flaps of the proboscis, and placing them in the genus *Eusina*, then the difference between *Oxyga* and *Tephritis* is rendered so very subtle, as to become unavailable for my essay of a classification of North American *Trypetina*, based as it is upon very insufficient materials. The question arises, therefore, whether it would not be better, temporarily, to bring together all the species to be described below (with the exception of the *Urellia*) under the head of the genus *Tephritis*, or else to distribute those species in genera on some other principle. The latter course seems to me preferable, in rendering the determination of the species easier. I would propose to call *Tephritis* those species, the picture of the wings of which does not form at the apex distinctly developed rays, and those which have such rays would form a new genus *Euaesta*. Most species will then gain a position in conformity to their true relationship, as well as to their habitual affinities; and although it cannot be denied that the location of some species will thus be rendered somewhat artificial, this disadvantage cannot well be avoided as long as the knowledge of the American fauna is not more complete than it actually is.

That *Trypeta angustipennis* belongs to the genus *Tephritis* results from the foregoing explanation.

41. T. finalis Loew. ♂ ♀. (Tab. XI, f. 4.)—Cinerea, capite pedibusque luteis, proboscide non geniculatâ, alis nigro-reticulatis, fasciâ obliquâ inde a stigmate trans venas transversales ad posticum alæ marginem ductâ, maculisque duabus alterâ subapicali, costæ contiguâ et alterâ apicali non reticulatis, stigmatis nigri basi dilutissime sublavescente, venâ longitudinali tertiâ nudâ.

Cinereous; head and feet clay-yellow; proboscis not geniculated; reticulation of the wings black; a crossband running from the stigma over the crossveins, a spot near the anterior margin before the apex, and another one on the apex, are not reticulate; the basis of the black stigma is of a very faint yellow; the third longitudinal vein is not bristly. Long. corp. ♂, 0.16; cum terebrâ 0.24; long. al. 0.20—0.21.

SYN. *Trypeta finalis* Loew, Dipt. Am. Cent. II, 78.

Cinereous, thorax and abdomen without any picture. Head, antennæ, and palpi rather dark yellow, the larger part of the occiput dark brown. The front is of a very moderate breadth; its usual bristles are black. The antennæ do not reach to the anterior edge of the mouth; their second joint does not bear a longer bristlet; the anterior corner of the third joint is rounded;

the arista is but little incrassated at the basis, its pubescence is but very little perceptible. The upper side of the thorax bears some short, yellowish-white pile and black bristles, two pairs of which seem to have been inserted upon its middle. Scutellum, at the basis, of the same color with the thorax, towards the tip more or less yellowish; it bears four black bristles. The comparatively somewhat narrow abdomen is likewise of the same color with the thorax, its last segment a little elongated; its short pubescence is yellowish-white; the long bristles at the end of the last segment are usually black. The flat ovipositor of the female is somewhat longer than the last two abdominal segments taken together, red, blackened at the root and at the extreme tip only; its short and fine pile is of a very pale color. The wings are comparatively long and narrow, coarsely reticulate with brownish-black upon their whole surface; the root of the wing, up to a little beyond the end of the small basal cells, shows but some scattered spots; upon the rest of the surface the single drops are large and hence rather close together, although but little coalescent; no drops at all, or almost none, are to be found on a crossband running obliquely from the stigma over both crossveins to the posterior margin of the wing, on a spot beginning at the anterior margin near the apex of the wing, and on a smaller spot upon the apex itself; the basis of the black stigma forms a large, limpid drop, somewhat tinged with yellowish; the usual triangular cluster of drops between the stigma and the unperforated crossband before the apex consists of six drops, three quadrangular ones between the costa and the second longitudinal vein, a larger quadrangular spot and a smaller rounded one between the second and third longitudinal veins, finally a large round one beyond the third vein. The latter vein has no bristles; the small crossvein corresponds to the last third of the discal cell.

Hab. California (A. Agassiz); Texas (Belfrage).

Observation.—This species is a normal *Tephritis*.

45. *T. clathrata* Lw. ♀. (Tab. X, f. 15.)—Cana, capite pedibusque flavis, femoribus liturâ nigricante signatis, abdomine bifariam nigromaculato; alæ colore nigro rare maculato-reticulate, stigmatè atroguttan hyalinam includente, venâ longitudinali tertiâ nudâ; peristomium modice productum et proboscis breviter geniculata; terebrâ aterrima, duobus ultimis abdominis segmentis simul sumtis æqualis.

Whitish-gray, head and feet yellow, femora with a black streak, abdomen with two rows of black dots, wings with a sparse reticulation, almost reduced to spots; the stigma includes a hyaline drop; third longitudinal vein not bristly; oral edge moderately produced, proboscis short, geniculate; the deep black ovipositor is as long as the last two abdominal segments taken together. Long. corp. 0.12; long. al. 0.13.

Syn. *Trypeta clathrata* LOEW, Monographs, etc., I, p. 80. Tab. II, f. 15.

Hab. Middle States (Osten-Sacken).

Observation.—In accordance with what has been said in the second observation to *T. angustipennis*, *T. clathrata* belongs to the genus *Tephritis*. Should the distribution adopted by me in my Monograph of the European *Trypetidae* be strictly applied to this species, it would, on account of the distinctly geniculate proboscis with but moderately prolonged flaps, be referred to the genus *Oryna*; and it agrees very well with a number of European species, placed in that genus.

46. *T. geminata* Lw. ♀. (Tab. XI, f. 1.)—Ex luteo-cinerea, capite, thoracis margine laterali, scutello, abdominis dimidio basali, femorum apice, tibiis tarsisque flavis, pleuris, metanoto, abdominis maculis et apice, terebrâ femoribusque ex nigro fuscis; alæ præter basin fusce, limbo-guttatâ, guttulis disci minutis et raris, guttis marginis postici majoribus, anguli axillaris confertioribus, maculis demique duabus costalibus trigonis limpidis, venâ longitudinali tertiâ setosâ.

Yellowish-gray; head, lateral margin of the thorax, scutellum, anterior half of the abdomen, tip of the femora, tibiae, and tarsi, yellow; pleurae, metanotum, spots and posterior part of the abdomen, ovipositor, and femora blackish-brown; wings, with the exception of the basis, brown, with pale drops, which are small and scattered in the middle, larger upon the posterior margin, more dense upon the posterior angle; upon the anterior margin there are two triangular hyaline spots; the third longitudinal vein is bristly. Long. corp. 0.17; long. al. 0.20.

Syn. *Trypeta geminata* LOEW, Dipt. Am. Sept. Cent. II, 75.

Head pale yellow, only a large spot upon the occiput blackish-brown; front rather broad; the ordinary bristles pale brownish or almost yellowish. Antennae dark yellow; the short pile upon the second antennal joint pale yellowish; a single more elongate hair is black; the anterior corner of the third joint is rather sharp. Face rather concave and the anterior corner of the mouth rather conspicuously projecting. Cheeks narrow. Oral opening large, rounded; palpi and proboscis short, not reaching beyond

the anterior edge of the oral opening; proboscis not geniculated. The ground color of the upper side of the thorax is black, but, in consequence of its pulverulence and of its short, yellowish pile, it appears gray; upon its anterior margin, in the vicinity of the yellowish humeral callus, there are some blackish hairs; the ordinary bristles, of which I perceive only two pairs upon the middle of the dorsum, are brown. The lateral margin of the thoracic dorsum is yellow; scutellum yellow, with four bristles. Metanotum and pleuræ blackish-brown; the latter rather shining; the bristles upon them for the most part black. The ground color of the abdomen is yellow; it has four rows of brownish-black spots, which begin to expand upon the third segment; upon the fourth and the following segments they coalesce in such a manner that the segments appear altogether blackish-brown. The pile upon the abdomen is generally whitish-yellow, but upon the black spots it is black; the bristles upon the posterior margin of the posterior segments are generally black. The rather broad ovipositor is of a shining blackish-brown, flattened, although somewhat swollen at the basis; its short and very delicate pile is not easily discernible; it seems to be brownish. Femora brownish-black, the anterior ones with long black bristles; the extreme root and the tip dark yellow. Tibiæ and tarsi rather dark yellow; wings of the ordinary shape, blackish-brown, sparsely guttate; the root of the wings, almost as far as the tip of the small basal cells, is rather hyaline and almost altogether immaculate; the alula also, bears no spots and is without dark coloring; the brown coloring begins on the anterior margin about the middle of the costal cell, and includes before its end a rather large hyaline drop, close by the margin; a smaller hyaline drop is placed upon the tip of the brownish-black stigma; immediately beyond the stigma, on the anterior margin, there are two triangular, hyaline spots, separated only by a brown stripe; their end crosses the second longitudinal vein; the whole middle portion of the wing is perforated by a few isolated, very small hyaline drops; upon the second half of the posterior margin there are four large hyaline drops, two before and two after the end of the fifth longitudinal vein; a fifth, much smaller drop, is placed much nearer the tip of the fourth vein; the last portion of the sixth longitudinal vein is surrounded by a cluster of somewhat larger spots, which, in consequence of the more faded brown,

surrounding them, appear more coalescent; in the posterior angle of the wing the pale drops are more numerous and somewhat larger than upon the middle of the wing, and moreover, well separated from each other; the apex of the wing shows between the third and fourth veins a very narrow, hardly apparent hyaline border.

Hab. Pennsylvania (collection v. Winthem).

Observation.—In accordance with the explanations given in the second observation to *T. angustipennis* I leave *Trypeta geminata*, in spite of its distinctly bristly third vein, in the genus *Tephritis*, but I do this with the explicit understanding that this position is an unnatural one. In the above-quoted place I have already explained why one would feel tempted to place this species in the genus *Oxyphora* on account of the pattern of its picture, as well as of the bristles upon the third vein; but I must again add that this location would not be natural. Its rather stubble-shaped pile, the distribution of the bristles upon the front, and the structure of the antennæ indicate a rather close relationship to those European species which I have united in the genus *Carphotricha*; nevertheless, in some other characters it differs from those species in a measure which prevents its reception in that genus. A number of South American species stand in the same relation to the European *Carphotrichæ*, although they differ among themselves in many very striking plastic characters. A more complete study of these species will result in the breaking up of the genus *Carphotricha*, based upon too insufficient material, and then only, in all probability, *T. geminata* will find its true position.

47. *T. fucata* FABR. ♂.—Lutea, capite pedibusque flavis; setæ scutelli quatuor; alæ guttis hyalinis majusculis subraris reticulatæ, retis parte posticâ unicolore ex cinereo-fuscâ, anticâ luteo et fusco variâ, ita ut guttule luteæ guttis hyalinis interjectæ sint, margine antico strigulis quinque et maculâ subapicali fuscis notato, venâ longitudinali tertiâ setosâ; proboscis non geniculata.

Clay-yellow, head and feet of a purer yellow; scutellum with four bristles; the reticulation of the wings, formed of rather large and moderately numerous hyaline drops, is uniformly grayish-brown upon the posterior part of the wings, yellow and brown upon the anterior portion, in such a manner that yellowish drops are mixed among the hyaline ones: upon the anterior margin, there are five small brown transverse streaks and

before its end there is a brown spot; the third longitudinal vein is beset with bristles; proboscis not geniculated. Long. corp. 0.17; long. al. 0.20.

SYN. *Musca fucata* FABRICIUS, Ent. Syst. IV, p. 359, 194.

Tephritis fucata FABRICIUS, Syst. Antl. p. 321, 24.

Trypeta fucata WIEDEMANN, Auss. Zweifl. II, p. 505, 44.

Clay-yellowish, almost ochre-yellow. Head rather pale yellow. Front and sides of the face with short, unusually dense yellowish pile. Front of a medium breadth; the bristles brownish-yellow, brown towards the tip. Antennæ pale yellow, of medium length, reaching almost to the anterior edge of the mouth, which is very much drawn upwards; the short pile on the second joint is yellowish; the third joint has an indistinctly rounded anterior corner; antennal arista apparently bare, but little incrassated at the basis. Face rather narrow, somewhat excavated, distinctly carinate between the antennal foveæ; in the profile, its lower part is produced in the shape of a short snout. Eyes large, oval; cheeks narrow, with yellow pile and bristles. Oral opening large, longer than broad; the rather broad palpi yellowish and with yellowish pile, reaching to the anterior edge of the oral opening; proboscis short, not geniculated. The whole thorax is so thickly covered with yellow pollen and short, yellow pile, that its ground color, which seems to be grayish-brown, is hardly visible; the ordinary bristles, two pairs of which are inserted upon the middle of the upper side, are brownish-yellow; their tip is dark brown. The ground color of the scutellum is pale yellow, which color is, however, but little apparent, on account of a short yellow pile, similar to that on the thorax; the scutellum has four bristles. Abdomen of the same coloring as the thorax; the short hairs and bristles are all yellow. Feet yellow, with yellow pile, the anterior femora have yellowish bristles. The reticulation of the wings consists of hyaline, almost whitish, rather large, and not very numerous drops; it does not reach the extreme root of the wings; upon the posterior margin and at the extreme apex of the wing the coloring is uniformly grayish-brown; elsewhere, it is clay-yellow, with a brown picture, which partly frames in the hyaline drops, partly includes little clay-yellowish drops, so that the coarser reticulation formed by the hyaline drops, in its turn appears reticulate. Upon the anterior margin itself there are five, in part almost punctiform, brownish-black transverse streaks; upon the

end of the marginal cell a brownish-black spot; the streaks are upon the humeral crossvein, in the middle between the latter and the basis of the stigma, upon the latter, on the end of the stigma, and between that and the tip of the second longitudinal vein. The small crossvein lies a little beyond the last third of the discal cell. The third longitudinal vein is distinctly bristly.

Hab. The Antilles? (Fabricius); South America (Wiedemann); Buenos Ayres (collect. Wiedemann).

Observation 1.—Fabricius, the first describer of the species, names Dr. Pflug as the discoverer, and the South American islands as the *habitat*, which probably means the Antilles. Later, the species was described by Wiedemann, who names South America as the *habitat*. It is impossible to tell from the descriptions of both authors, whether they really meant the same species, although the descriptions contain nothing positively contrary to this assumption. As the species is easy to identify, and as Wiedemann's identification was based upon the comparison of Fabricius's specimens, it can be safely assumed that he has described the same species. My description is based upon a male, marked Buenos Ayres and communicated to me as a type from Wiedemann's collection.

Observation 2.—This species may also remain in the genus *Tephritis*, for the sake of facilitating identification, although its third vein is distinctly bristly. This character, as well as the not geniculated proboscis, recalls those species which, in my Monograph of the European Trypetidæ, I placed in the genus *Oxyphora*; in fact I know of no other American species which stands closer than *T. fucata* to the typical species of that genus, as, for instance, to *T. corniculata* Zett., *biflexa* Lw., etc. I also call attention to a peculiarity of most species of this group, that the dark spots of the picture in the female are more extensive than in the male; this may likewise be the case with *T. fucata*.

48. *T. albiceps* n. sp. ♂ ♀. (Tab. XI, f. 5.)—Ex luteo cinerea, capite albicante, fronte, antennis, scutello pedibusque luteis, abdomine bifariam nigro maculato; alæ latiusculæ, præter imam basim totæ colore fusco-nigro guttato-reticulatæ, guttis valde inæqualibus, in apice et prope venam transversam posteriorem quam in reliquâ alâ minus confertis, stigmatæ nigro uniguttato, venâ longitudinali tertiâ non setosâ; terebrâ femineæ aterrimâ, duobus ultimis abdominis segmentis simul sumtis æqualis.

Yellowish-gray; head whitish; front, antennæ, scutellum, and feet yellow; the abdomen with two longitudinal rows of black spots; wings somewhat broad, and with the exception of the extreme basis, entirely covered with a guttate brownish-black reticulation; the drops are of a very unequal size and less numerous upon the apex and in the vicinity of the posterior crossvein; the black stigma has a hyaline drop; third longitudinal vein not bristly; ovipositor of the female deep black; as long as the last two abdominal segments taken together. Long. corp. ♂ 0.13, ♀ 0.16; long. al. 0.15—0.16.

Yellowish-gray; thorax and abdomen with whitish-yellow pile; the latter with two longitudinal rows of black or blackish dots. In well-preserved specimens the head is white, and it probably has the same color in living ones; in some of the dried specimens it has assumed a yellowish hue; the front, with the exception of its lateral margins, is yellowish; the usual bristles upon it are almost without exception black; the bristles upon the vertical margin are pale yellowish. Antennæ pale yellowish; the third joint has an almost sharp anterior angle. Oral opening large, somewhat longer than broad; the anterior edge of the mouth rather drawn upwards, somewhat projecting in the profile. Palpi pale yellowish. Proboscis yellowish, short geniculate, with but moderately prolonged, comparatively stout flaps. The upper half of the occiput is gray, with the exception of the margin along the orbit. The ground color of the humeral callosities is yellow, while upon the rest of the thorax it is blackish. The bristles of the thoracic dorsum are all black, those of the pleuræ are partly black, partly pale yellowish. Scutellum pale yellow; lateral angles and sometimes also the basis darker; with four black bristles. The bristles upon the posterior margin of the last abdominal segments have the same pale yellowish tinge as the pile upon the abdomen; only exceptionally a dark bristle is sometimes found among them. The ground color of the abdomen is not quite constant; as a rule, it is blackish; I possess specimens, however, in which, upon the posterior margin of the second and third segments, it is yellowish-red. The ovipositor is shining black, rather strongly contracted towards its end, as long as the last two abdominal segments taken together; their short pile is very delicate and hence somewhat difficult to discern; it seems to have the same coloring as the pile on the abdomen. Feet saturate yellow. The wings have an almost regularly elliptical shape and are somewhat broader in the female than in

the male (the figure is made from a male specimen). The guttate reticulation, which leaves open the extreme basis only, has a brownish-black coloring, which assumes a paler hue wherever the drops are nearer together; upon the stigma, however, and upon the end of the marginal cell, it becomes nearly black; the stigma contains a rather conspicuous hyaline drop; the drops upon the remaining surface are in general large, upon the middle of the wing, however, numerous, much smaller drops are interspersed, which perforate the dark coloring between the larger drops; this also takes place between the six large drops which form the usual pyramid of drops, situated beyond the stigma; upon the portion of the wing beyond this pyramid there are generally but very few little drops, and those are usually in the proximity of the pyramid; some larger drops, rather distant from each other, are also to be found there, and among these a row of very rounded drops along the margin of the wing, sometimes a little remote from it; they are either of very unequal size (as in the figure), or of the same size; the proximity of the posterior crossvein shows a more considerable space, which is but little perforated. The third longitudinal vein is not bristly.

Hab. Canada (Couper); English River (Kennicott); Maine (Packard).

Observation.—In the distribution adopted by me for the American species, the present one would belong to the genus *Tephritis*. Should my distribution of the European *Trypetæ* be applied to it, the shape of its oral opening and of the proboscis would refer it to *Oxya*.

49. *T. euryptera* n. sp. ♀.—Ex luteo-cinerea, abdomine bifariam nigro-maculato, capite et apice scutelli flavicantibus, pedibus luteis; alæ valde dilatate, rotundato-ovate, præter imam basin totæ colore fusco-nigro guttato-reticulatæ, guttis valde inæqualibus, in apice et prope stigmâ venamque transversam posteriorem minus confertis, stigmatæ uniguttato, venâ longitudinali tertiâ non setosâ; terebrâ femine æterinâ, duobus ultimis abdominis segmentis simul sumtis æqualis.

Yellowish-gray; abdomen with two longitudinal rows of black spots; head and tip of the scutellum pale yellow; feet saturate yellow; wings very broad, rounded oval, with the exception of the extreme basis covered with a guttate, brownish-black reticulation, the drops of which are of a very unequal size and less numerous in the vicinity of the stigma, of the posterior crossvein, and on the apex of the wing; stigma with a hyaline drop; the third longitudinal vein not bristly; the ovipositor of the

female deep black, as long as the last two abdominal segments taken together. Long. corp. cum terebrâ 0.16; long. al. 0.16.

Closely allied to *T. albiceps* and very like it, but easily distinguished by its very broad wings. Yellowish-gray; thorax and abdomen with yellowish-red pile; the abdomen with two longitudinal rows of black spots. Head yellowish; front and antennæ more yellow; the usual bristles on the front black, the bristles on the vertical margin bright reddish-yellow. The third antennal joint with an almost sharp anterior corner. The oral opening longer than broad, the upper oral edge somewhat drawn upwards, distinctly projecting in the profile. The rather broad palpi and the proboscis are yellowish; the latter short geniculate, with but moderately prolonged, rather stout flaps; the occiput, in the vicinity of the point of attachment, grayish. The ground color of the humeral callus is yellowish, that of the thorax blackish; the bristles of the dorsum are black, the two pairs upon its middle are inserted upon very small black dots, easily overlooked. Scutellum yellow at the tip, with four black bristles. Ovipositor of the female shining black, about as long as the last two abdominal segments taken together (in the only specimen in my possession the shape of the ovipositor is not distinctly discernible, but it does not seem to differ from that of *T. albiceps*); its short pubescence is delicate, and hence somewhat difficult to perceive; its coloring seems to be altogether reddish. Feet saturate yellow. The wings are very broad and have a rounded elliptical shape. The guttate reticulation shows the most striking likeness to that of *T. albiceps*, so that the description of the latter may be applied to this; the only addition to be made would be, that the region immediately below the stigma is somewhat darker and a little less guttate. Thus the figure of the wing of *T. albiceps* gives quite a correct idea of the wing of the present species, except of its broader shape; moreover, the three posterior drops of the usual pyramid are smaller, and separated by larger intervals, and the intervals of all the six drops are perforated by much more numerous small drops. The third longitudinal vein is likewise not beset with bristles in this species.

Hab. West Point, N. Y. (Osten-Sacken).

Observation.—The systematic position of this species is exactly the same as that of *T. albiceps*.

50. *T. platyptera* n. sp. ♀.—Cinerea, abdomine quadrifarium nigro-maculato, capite pedibusque luteis, femoribus tamen posterioribus nigro-maculatis, scutello nigro- et flavo-variegato; alæ valde dilatate, rotundato-ovate, totæ colore nigro guttato-reticulate, venâ longitudinali tertiâ non setosâ.

Gray, abdomen with four rows of black spots, head and feet yellow, the hind femora spotted with black; scutellum variegated with yellow and black; wings very broad, rounded-ovate, covered upon their whole surface with a reticulate black picture; third vein not bristly. Long. corp. cum terebrâ 0.21; long. al. 0.16.

Of this species I possess a single badly preserved specimen, and I would not have attempted to describe it, but for the circumstance that it is distinguished by a number of very peculiar characters, which render its recognition easy, even should the description be imperfect. Head yellowish; occiput immediately above the point of attachment somewhat blackish; on each side, near the basis of the antennæ, there is, on the border of the eye, a small, almost punctiform, blackish-brown transverse streak. The breadth of the front, which is distinctly narrowed anteriorly, is comparatively considerable, as it equals half the breadth of the head; the usual frontal bristles are black, those upon the vertical margin are yellowish-white. The third antennal joint is gently excised upon its upper side, and has a rather sharp anterior angle. Checks rather broad, with a black bristle, in front of which, along the lateral edge of the mouth, there is some black pile. Oral opening very wide; its anterior edge is but little drawn up, although rather projecting in the profile. Palpi very broad, reaching beyond the anterior edge of the mouth, beset with black and whitish-yellow hairs. Proboscis short geniculate, with moderately prolonged, stout flaps. The thorax of the specimen is greasy, and it is impossible to make any positive statement about its coloring and the pile upon it; the coloring upon the dorsum seems to have been more blackish; on the sides more brown; the pile seems to have been stubble-shaped, yellowish-white; all the bristles, upon the thoracic dorsum as well as upon the pleuræ, are black. The very convex, blackish scutellum has, upon the lateral margins and upon the tip, a broad yellowish border; the four blackish bristles of the scutellum are placed inside of this border upon blackish dots; the pair of those dots which is near the tip, although smaller, is connected with the black coloring of the

scutellum. Abdomen gray, with four rows of black spots. The spots of both intermediate rows are comparatively large rectangular triangles, one cathetus of which lies along the posterior margin of the segment, the other is parallel to the longitudinal axis of the abdomen; thus between both rows of spots, only a narrow gray intermediate line remains visible; the spots of the outer rows lie upon the lateral margins and also occupy the whole length of the segments, forming broad, uninterrupted lateral stripes. The whitish pile upon the abdomen is rather stubble-shaped; the comparatively long and strong bristles upon the posterior margin of the last segment are black. Venter somewhat dirty brick-red, gradually becoming blackish towards the lateral margins. Ovipositor flattened, broadly truncate at the end, shining black on the surface; the under side bright yellowish-red, with a black tip. Feet of an impure yellowish, the posterior femora on the under side with two well-defined blackish spots, and near the tip with a faded blackish spot. Wings very broad, of the same rounded elliptical shape as in *T. euryptera*. The black, guttate reticulation covers the whole wing to the extreme basis; along the whole posterior margin as far as the apex, there is a row of hyaline drops of middle size, separated by considerable intervals; beyond the apex, along the anterior margin, these drops become larger, their intervals growing smaller; in the marginal and costal cells they coalesce with a little drop placed behind them, so that, in these cells, the reticulation emits something like little rays, running towards the anterior margin; the stigma, upon the extreme basis, has a whitish crossline and includes a hyaline drop at the end; upon the whole inner side of the surface of the wing the black color is rather sparsely perforated by drops of middle and of the very smallest size; the latter are more numerous upon the posterior than upon the anterior half of the wing. The cells of the wings are all of an unusual breadth, and the crossveins accordingly of an unusual length; the distance between them is but little shorter than the middle crossvein; the second and third longitudinal veins are considerably divergent towards the end; upon the third I do not perceive any bristles.

Hab. Connecticut (H. F. Bassett).

Observation.—I leave this species provisionally in the genus *Tephritis*; the description shows sufficiently that it is a stranger there, whose affinities point towards the genus *Eurosta*. To

found a special genus for this single form would be premature, as there are several concurrent South American species, without the knowledge of which it is difficult to choose the characters upon which to establish the genus. To place the species in the genus *Eurosta* is likewise unadvisable, as the absence of bristles upon the third vein, and the not conical but flattened ovipositor are in conflict with the chief characters of *Eurosta*.

51. *T. æqualis* Loew. ♂ ♀. (Tab. X, f. 20.)—Dilute lutea, terebrâ concolore, tribus ultimis abdominis segmentis simul sumtis longiore, pilis, setisque totius corporis exalbidas; alæ colore ex-fusco nigricante, adversus costam et apicem in nigrum mutato, æqualiter guttato-reticulata, guttis confertis plerisque majusculis, picturâ marginis antice radiatâ, marginis apicalis subradiatâ; vena longitudinalis tertia non pilosa.

Pale yellowish; ovipositor of the female likewise yellow, longer than the last three abdominal segments taken together; pile and bristles of the whole body whitish; wings with a brownish-black guttate reticulation, black near the anterior margin and the apex; the drops are crowded and the majority of them are of a considerable size; the pattern of the picture consists of rays along the anterior border, which are less well-marked along the apex; the third vein is not beset with bristles. Long. corp. ♂ 0.22, ♀ cum terebrâ 0.25—0.26; long. al. 0.24—0.25.

SYN. *Trypeta æqualis* Loew, Monogr. etc., I, p. 86. Tab. II, f. 20.

Hab. Illinois (Kennicott). [Maryland, P. R. Uhler; Ohio, H. F. Bassett.—O. S.]

Observation.—The present species shows such a peculiar structure of the head and of the parts of the mouth, that I would not have hesitated to establish a separate genus for it, if I had had better preserved specimens for examination. The general appearance reminds of the species which I have united in the genus *Icterica*, but it differs in a smaller oral opening, a different shape of the wings, and a third longitudinal vein which is not beset with bristles. Not being able to assign a better position for it at present, I had the choice of leaving it in the genus *Tephritis* or of removing it to the genus *Euaresta*, proposed in the second observation to *Tryp. angustipennis*. The choice is not a very easy one, because, although the picture of the wings is distinctly radiate along the anterior margin as far as the apex, the apex itself and the space immediately behind it are more guttate than radiate. By all means, the question is more about

an artificial than about a final location of the species, as the latter will have to depend upon the results of a future investigation. The circumstance that the pyramid of drops beyond the stigma, usually well developed in the species collected in the subgenus *Tephritis*, is not distinctly marked here, decides me to place the species in *Euaresta*, although its affinities to the types of this subgenus may be very slight.

52. *T. festiva* Loew. ♂ ♀. (Tab. X, f. 21.)—Lutea, unicolor, alæ inæqualiter guttato-reticulatæ, in margine antico et apice radiatæ, picturâ in basi et disco sordide lutescente, prope marginem anticum et in apicali alarum tridente fusco-nigrâ; terebra femine quatuor ultimis abdominalis segmentis simul sumtis subæqualis, non depressa, adversus apicem valde angusta, superne nigra vel fusco-nigra, infra adversus basin rufa.

Clay-yellow, unicolorous, the reticulation of the wings unequally guttate, radiate along the anterior margin and on the apex, more dingy clay-yellow upon the basis and in the middle; brownish-black along the anterior margin of the wing and upon the apex; the ovipositor of the female is almost as long as the last four abdominal segments taken together, not flattened, very narrow at the tip, black or brown on the upper side, the under side red towards the basis. Long. corp. ♂, 0.17—0.18; ♀ cum terebrâ 0.20—0.23; long. al. 0.22.

Syn. *Trypeta festiva* Loew, Monographs, etc., I, p. 86. Tab. II, f. 21.

Hab. Pennsylvania (Osten-Sacken); Connecticut (Norton). [New Jersey, Mr. Jung; Illinois, Dr. Brendel; Ohio, H. F. Bassett.—O. S.]

Observation 1.—*Trypeta festiva* may be considered as a typical form of the genus *Euaresta*. As the third longitudinal vein of the wings is beset with spines, this species would have to be placed in the genus *Oxyphora*, in the classification adopted by me for the European species.

Observation 2.—Brazil possesses a conspicuous species closely allied to the present one, but more approaching the next following ones in the pattern of the picture of the wings. I let its description follow:—

***T. spectabilis* n. sp.** ♂ ♀. (Tab. X, f. 27.)—Tota luteola, terebra tamen obscure ferruginea, non depressa et quatuor ultimis abdominalis segmentis subæqualis; scutellum quadrisetosum; alarum pictura nigra, in apice pulchre, sed breviter radiata, adversus angulum posticum rarior, in disco rarissime guttata, guttâ cellulæ posterioris primæ unicâ; vena longitudinalis tertia setosa.

Altogether yellowish, except the ovipositor, which is dark ferruginous, not flattened, and nearly as long as the last four abdominal segments taken together. Scutellum with four bristles; the black picture of the wings shows, on the apex, handsome, although short, rays; it is sparsely guttate towards the posterior angle, very sparsely in the middle of the wing; the first posterior cell contains but a single drop; the third longitudinal vein is bristly. Long. corp. 0.26—0.27; long. al. 0.26.

A rather conspicuous species, of the same coloring as the European *T. valida* Lw. With the exception of the ovipositor, it is altogether yellowish, only the basis of the abdomen is sometimes brownish. Front of a middle breadth and somewhat convex; its brownish-yellow or reddish-yellow bristles are comparatively strong; the frontal lunule rather large. Antennæ short, by far not reaching the edge of the mouth; the second joint bears a conspicuous bristle; the anterior edge of the mouth considerably drawn up, but not very projecting in the profile. Eyes not very high; cheeks broad. Oral opening rounded, rather large; proboscis not geniculate; palpi rather broad, reaching abundantly as far as the anterior edge of the mouth. The short pile on the thorax is partly pale ferruginous, partly pale yellowish-red; the usual bristles are pale yellow or brownish-yellow. The somewhat convex scutellum has four bristles. Metathorax and pleuræ yellow, like the rest of the body. Abdomen likewise uniformly yellow, but there are specimens the abdomen of which is infuscated at the basis; the pile on the abdomen is like that on the thorax, only its coloring is more yellowish. The stout, conical ovipositor is not flattened at all, about as long as the last four abdominal segments taken together; in paler specimens it is reddish-brown with a black tip; in darker specimens it is rather brownish-black; it is beset, as far as the tip, with comparatively long pile, which assumes a more yellowish hue near the basis, a more brownish one near the tip; in darker specimens it is sometimes blackish-brown. Feet altogether yellow. Wings hyaline with a very much expanded and very little perforated black reticulation, which is radiated at the apex of the wing. The root of the wings is not spotted nearly as far as the end of the small basal cells; the costal cell contains a gray crossline near the humeral crossvein, a brownish-black crossband upon its middle, and a crossline of the same color at its extreme end; the obliterate end of the auxiliary vein, running perpendicularly towards the margin of the wing, is rather hyaline; the stigma is altogether black and does not include any hyaline drop; immediately beyond the stigma near the anterior margin, there are two cuneiform hyaline spots, the first of which is a little broader than the second and crosses the second vein a little further; between these spots and the end of the second vein the brownish-black coloring is entirely unbroken; five short brownish-black rays of almost equal length run towards the apex; the first ends between the second and third longitudinal veins, the next two coincide with the

er: of the third and fourth veins; the last two cross the second posterior cell; the last of all is connected with the remaining brownish-black coloring by a narrow brownish-black bridge and sometimes interrupted at the basis; upon the anterior side of the fourth vein there are only two hyaline drops, the one below the stigma, the other between both cross-veins; in the third posterior cell there are six hyaline drops, the one of which is at its extreme basis and the others upon its latter half; some of the latter drops are sometimes coalescent; in the posterior angle there are, moreover, four or five hyaline drops. The small crossvein is almost perpendicular and is nearly opposite the last third of the discal cell; the posterior crossvein likewise is rather perpendicular; the third longitudinal vein is distinctly bristly.

Hab. Brazil (collection v. Winthem).

53. *T. bella* Lw. ♂ ♀. (Tab. X, f. 23.)—Luteo-cinerea, capite, pedibus, abdomineque flavis, hoc apicem versus nigricante; setæ scutelli quatuor; alarum pictura nigra, in margine antico et apice pulchre radiata, prope marginem posticum paulo confertius, in disco rarissime guttata, guttâ cellulae posterioris primæ plane nullâ; vena longitudinalis tertia setosa.

Yellowish-gray; head, feet, and abdomen yellow; the latter blackish towards the end; the black picture of the wings handsomely radiate on the anterior margin and the apex; in the vicinity of the posterior margin with numerous drops, upon the middle of the wing with very few, in the first posterior cell with none; third longitudinal vein bristly. Long. corp. ♂, 0.12—0.13, ♀ cum terebrâ 0.13—0.15; long. al. 0.11—0.12.

Syn. *Trypeta bella* Loew, Monographs, etc., I, p. 88. Tab. II, f. 23.

Hab. New York (Fitch); Washington (Osten-Sacken); Wisconsin, etc. [Rather common everywhere in the U. S.—O. S.]

Observation.—Closely related to *T. festiva*, and, as to its systematic location, the remarks appended to that species are also applicable here.

54. *T. timida* Lw. ♂. (Tab. X, f. 25.)—Lutea, metanoto pleuris que ex-nigro fascis, capite pedibusque flavis; setæ scutelli quatuor; alarum pictura nigra, in apice pulchre radiata, prope marginem posticum rare et in disco rarissime guttata; guttâ cellulae posterioris primæ unicâ; vena longitudinalis tertia setulis paucis brevissimis instructa.

Clay-yellow, metathorax and pleræ blackish-brown; head and feet yellow; four bristles upon the scutellum; the black picture of the wings is prettily radiated at the tip, in the vicinity of the posterior margin sparsely, and upon the middle of the wing very sparsely guttate, in the first posterior cell with a single drop; the third longitudinal vein is

beset with extremely short and scarce bristles. Long. corp. 0.17; long. al. 0.16.

Syn. *Trypeta timida* Loew, Dipt. Am. Cent. II, No. 76.

Clay-yellow; the coloring of the head is of a purer yellow, but the middle of the occiput is grayish. Front comparatively narrow; its pale brownish bristles are strong and long. Antennæ yellow, not reaching to the oral edge; anterior corner of the third joint rounded; arista comparatively thin, its pubescence so short, that, to the naked eye, the arista appears bare. Face excavated; the anterior edge much drawn upwards, but little projecting in the profile. Eyes elongated-rounded; cheeks very narrow. Oral opening of a middle size, rather round; the yellowish proboscis not geniculate, short; palpi short, yellowish. The upper side of the thorax is clothed with pale yellowish hairs; upon its middle there is a weak trace of a very broad grayish stripe, which, however, in less denuded specimens, may be hardly visible. The bristles upon the upper side of the thorax are pale brownish; upon its middle there are three pairs. The yellow scutellum bears four bristles. The ground color of the metathorax is blackish-brown, but assumes a grayish aspect from a thin covering of pollen. The pleuræ have a similar coloring, but towards the upper margin, it becomes more yellow, and below the root of the wings there also is a spot of dingy yellow. The clay-yellow abdomen shows, in the described specimen, upon the last two segments brownish spots, which, however, seem to be the result of some lesion. Feet yellow. Wings rather broad with a brownish black, very sparsely reticulated picture, which is radiated on the apex; the root of the wings is very sparsely spotted before the end of the two small basal cells; the costal cell, near its basis, has a blackish transverse line, a brownish-black one beyond its middle, and another brownish-black one upon its extreme end; the obliterate end of the auxiliary vein, which runs perpendicularly towards the anterior margin, is rather hyaline; stigma brownish-black with a yellow crossline in the vicinity of its end; immediately beyond the stigma there are two cuneiform hyaline indentations, which extend from the margin to the second longitudinal vein; the latter is somewhat remote from the margin; between the second of these indentations and almost the end of the second vein, the brown color is not perforated; along the apex, the

brown color emits five brown rays of almost equal length. The first of these rays ends a little before the middle of the distance between the tips of the second and third veins; the two next ones, which are a little expanded at the tip, lie on the ends of the third and fourth veins, the last two in the second posterior cell; the last of these rays, in the vicinity of its origin, is not quite well separated from the remaining brownish-black picture. The hyaline drops are rather large, but few in number; there are two between the third and fourth longitudinal veins, the first before the small crossvein, the second less far beyond it; the discal cell also contains but two drops, placed under the small crossvein and nearer to the posterior side of the cell; the third posterior cell has a drop at its extreme basis and five considerable ones in the posterior angle of the wing, which, however, are less conspicuous, because the dark coloring in that region is more faded. The distance of the first and second longitudinal veins from the margin is a little larger than usual; the second and third veins are strongly diverging towards the end; a weaker divergency exists between the third and fourth veins; the two crossveins are perpendicular and straight; the small crossvein is almost twice as far from the proximal end of the discal cell as from the distal end. In my first description of this species I said that the third vein was not beset with bristles; a more attentive examination of the specimen, however, revealed to me, on one of the wings, a few very short bristles, which are either rubbed off on the other wing, or else in a situation which does not allow their close scrutiny; the first posterior cell does not contain a conspicuous concavity, like that in *T. bella*; and the corresponding spot is not darker than its surroundings.

Hab. Mexico (collect. v. Winthem).

Observation 1.—The systematic position of *T. timida* is exactly the same as that of *T. festiva* and *bella*.

Observation 2.—The next relative of *T. timida* is a Brazilian species, which can be very easily mistaken for it; and in order to prevent this confusion, I let its description follow here:—

T. obscuriventris n. sp. ♀. (Tab. X, f. 26.)—Ex luteo cinerea, capite pedibusque lutescentibus, abdomine ex piceo nigro et nitido, terebrâ concolore, tribus ultimis abdominis segmentis simul sumptis æquali; setæ scutelli quatuor; alarum pictura nigra, in apice pulchre

radiata, prope marginem posticum raro et in disco rarissime guttata, guttâ cellulæ posterioris primæ unicâ; vena longitudinalis tertia setosa.

Yellowish-gray, head and feet yellow, abdomen shining brownish-black, ovipositor concolorous, as long as the last three abdominal segments taken together; scutellum with four bristles; the black picture of the wings handsomely radiated on the apex, sparsely guttate in the vicinity of the posterior margin, very sparsely in the middle of the wing; a single drop in the first posterior cell; the third longitudinal vein beset with bristles. Long. corp. cum terebrâ 0.20; long. al. 0.16.

Head, including palpi, proboscis, and antennæ, yellow; only the occiput for the most part grayish. Front comparatively narrow; its brown bristles are long and strong. Antennæ not reaching to the edge of the mouth; third joint rounded at the end; arista comparatively thin, appearing bare to the naked eye, as the pubescence is very short; face excavated; the oral opening hardly of middle size, round; proboscis short, not geniculate. Palpi of middle size; the ground color of the thorax is altogether black, including even the humeral callosities, but this color is so much concealed under ochre-yellow pile and pulverulence, that it assumes a yellowish-gray hue; upon the pleuræ and especially on the metanotum the dark ground color is more apparent. The scutellum, bearing four bristles, is yellow to a considerable extent at the tip; the abdomen is of a shining brownish-black and shows weak traces of a yellowish-brown pollen; the pile is short and scattered, of mixed yellow and black hairs; the latter prevail or seem to do so, as many of the yellow hairs assume a blackish hue when they do not reflect the light. The flat, not very pointed ovipositor is pitch-black, shining, about as long as the last three segments of the abdomen taken together, beset as far as the tip with a brown pubescence, appearing black in some directions. Feet yellow. The comparatively rather broad wings have a brownish-black, very sparsely guttate picture, which is handsomely radiate at the tip; the root of the wings, as far almost as the end of the small basal cells, is hardly spotted at all; the costal cell, quite near the humeral crossvein, has a grayish crossline, a brownish-black one upon the middle and one of the same color, but narrower, at the end; the obliterate end of the auxiliary vein, running perpendicularly towards the anterior margin, is rather hyaline; stigma altogether brownish or only with a trace of a very small yellowish drop in the vicinity of its apex, near the anterior margin; immediately beyond the stigma there are two hyaline indentations on the anterior margin, the first of which alone reaches the rather distant second longitudinal vein; before the end of the second longitudinal vein near the anterior margin, there always is a considerable hyaline drop, which *T. timida* does not possess; five rays of almost equal length occupy the apex; the first of them reaches the margin nearer to the end of the second than of the third vein; the two following are somewhat expanded at the tip and end upon the tips of the third and fourth veins; the last two rays

cross the second posterior cell, and the last of them is a little broader than the preceding one and generally connected in the vicinity of its root with the remaining brownish-black picture by a brownish-black bridge, which cuts off the end of the hyaline indented interval in the shape of a drop. The hyaline drops are of a considerable size, but not very numerous; two are placed between the third and fourth veins, the one before, the other less far behind the small crossvein; in the same way there are only two drops in the discal cell, placed upon its posterior side, below the small crossvein; the third posterior cell contains a drop near its extreme basis and five considerable drops upon its distal half; finally four drops are situated in the posterior corner of the wing, which, however, are less conspicuous on account of the less dark coloring surrounding them. The first and second longitudinal veins are somewhat more distant from the anterior margin than usual; the second and third are strongly divergent towards the end; a lesser divergency exists between the third and fourth; both crossveins are perpendicular and straight; the small one is twice as far from the basis as from the end of the discal cell; the third vein is distinctly bristly; there is no distinct concavity in the first posterior cell, and the spot where it occurs in some species is not darker than the surroundings.

Hab. Brazil (coll. v. Winthem).

55. *T. melanogastra* Lw. ♂ ♀. (Tab. X, f. 24).—Luteo-cinerea, abdomine nigro, capite pedibusque flavis; setæ scutelli duæ; alarum pictura nigra, in apice radiata, prope marginem posticum paulo confertius, in disco rarissime guttata, guttâ cellulæ posterioris primæ unicâ; vena longitudinalis tertia non setosa.

Yellowish-gray, abdomen black, head and feet yellow; scutellum with two bristles; the black picture of the wings with rays at the tip, more densely guttate in the vicinity of the posterior margin, very sparsely in the middle, and with a single drop in the first posterior cell; the third longitudinal vein is not bristly. Long. corp. ♂ 0.09, ♀ cum terebrâ 0.12; long. al. 0.12.

Syn. *Trypeta melanogastra* Loew, Monographs, etc., I, p. 90. Tab. II, f. 24.

Hab. Cuba (Poey).

Observation 1—Two misprints must be corrected in the description in the first volume of these Monographs: the figure of the wing is quoted fig. 23, instead of 24, and on page 91, line 19, "fifth" must be read, instead of "first." Moreover, it must be added that the figure was drawn from a female specimen. The relation of *T. melanogastra* to *T. mexicana* Wied. will be explained under the head of the latter.

Observation 2.—*T. melanogastra* belongs, together with the

preceding species, to the genus *Euaesta*; it differs from them in the presence of only two bristles upon the scutellum and in the absence of bristles upon the third vein. In the system adopted by me some time ago for the European Trypetidae, this species, on account of the somewhat prolonged flaps of its proboscis and of the bareness of the third vein, would have to be placed in the genus *Oryna*. I do not believe that its generic separation from the preceding species is to be recommended. A close relative of this species is a Brazilian one, which differs, however, in its wings being comparatively much narrower and its body more slender. I let its description follow:—

T. tenuis n. sp. ♀. (Tab. X, f. 29.)—Angusta, luteo-cinerea, capite pedibusque gracilibus flavis; setæ scutelli duæ; alæ pro portione angustæ, picturâ nigrâ in angulo postico elutâ, in apice radiatâ, prope marginem posticum confertius, in disco rarissime guttatâ, guttâ celluæ posterioris primæ unicâ.

Slender, yellowish-gray; the head and the slender feet are yellow; wings comparatively narrow, with a black picture, which is faded on the posterior angle, radiate on the apex, more densely guttate near the posterior margin, very sparsely in the middle of the wing, where the first posterior cell contains but a single drop; third longitudinal vein not bristly. Long. corp. cum terebrâ 0.13; long. al. 0.13.

Body remarkably narrow and slender. Ground color blackish, but so much covered with yellowish pile and pulverulence that thorax and abdomen have a yellowish-gray appearance. Head, including antennæ, palpi, and proboscis, yellow; occiput, on its upper half, with a large blackish-gray spot. The front a little more than of medium breadth; its usual bristles blackish. Face somewhat excavated and narrower than the front. Antennæ somewhat broad, not quite reaching the edge of the mouth, which is somewhat drawn upwards, but does not project distinctly in the profile. Eyes comparatively large and rounded; cheeks very narrow. The palpi reach to the anterior edge of the mouth. The suctorial flaps seem to be somewhat injured in the described specimen, so that I am not quite sure whether the proboscis is geniculate or not; I believe that, in uninjured specimens, it would look short-geniculate; the dark color which the flaps have in the described specimen is certainly an unnatural one. The ground color of the thorax is altogether blackish, even upon the humeral corners; its upper side has a yellowish-gray appearance, in consequence of its pulverulence and pile; on the metathorax and the pleura the coloring is more blackish-gray. The scutellum is of the same coloring with the upper side of the thorax, the extreme apex only somewhat tinged with yellow; it bears only two bristles, which, like those of the

thorax, are blackish. Abdomen narrow, but little more gray than the upper side of the thorax, without any rows of dark spots, but on each side of the second segment with a but little apparent yellow spot; the short pile as well as the longer hairs upon the posterior margin of the last segment are yellowish. The flat, shining black ovipositor is as long as the whole abdomen, and beset with dark pile. The feet are slender and yellow, as well as the coxæ. Wings rather hyaline, with a brownish-black very continuous reticulation; the root of the wings is not distinctly spotted as far as the end of the small basal cells, but somewhat dusky; upon the middle of the costal cell there is a blackish-brown crossline; the stigma does not contain any hyaline drop, but its inner basal end is very slightly tinged with yellow; immediately beyond the stigma there are, near the anterior margin, two drop-like hyaline spots; each of them has a small hyaline drop under it, below the second longitudinal vein; the second one is smaller; before the end of the second vein there is no hyaline drop; near the tip of the wing the apex shows the usual five rays, which have a considerable breadth, and the last of which is connected by a bridge with the remaining brownish-black coloring, which thus isolates the inner end of the hyaline interval in the shape of a drop; the anterior side of the fourth vein shows two conspicuous spots, one immediately before, the other not far beyond, the small crossvein; in the discal cell there are three hyaline drops along the fifth vein, the middle one being the largest and lying almost under the small crossvein; above the last of these drops there is sometimes one little drop more; the extreme basis of the discal cell also shows an indistinct, sometimes double, little drop; the third posterior cell contains but a few large drops, which are partly coalescent in couples; the posterior corner of the wing is likewise guttate, but the drops are much less apparent here, owing to the pale ground color. The small crossvein is hardly half as distant from the end of the discal cell as from the basis; the third longitudinal vein is not bristly.

Hab. Brazil (collect. v. Winthem).

56. *T. mexicana* WIED. ♂. (Tab. X, f. 28.)—Luteo-cinerea, abdomine nigro, adversus basim interdum sordide luteo, capite pedibusque flavis; setæ scutellidnæ; alarum pictura nigra in apice radiata, radiis tamen in marginem posticum excurrentibus minus explicatis et minus liberis, prope marginem posticum confertius, in disco rarissime guttata, guttâ cellulæ posterioris primæ unicâ; vena longitudinalis tertia non setosa.

Yellowish-gray, abdomen black, sometimes of a dingy clay-yellow towards the basis; head and feet yellow; scutellum with two bristles; the black picture of the wings is radiate on the apex, but the rays in the vicinity of the posterior margin are less developed and less free; the drops near the posterior margin are more numerous, those in the middle of the wing very sparse; the first posterior cell contains but a single drop;

third longitudinal vein not beset with bristles. Long. corp. 0.09—0.10; long. al. 0.12.

SYN. *Trypeta mexicana* WIEDEMANN, AUSS. Zweifl. II, p. 551.

Yellowish-gray. Front of a more vivid yellow, upon the lateral margin with a rather indistinct whitish pollen; the usual bristles upon it are black; those on the vertical margin pale yellowish. Eyes rounded ovate; cheeks very narrow. Face distinctly excavated, the anterior edge of the mouth is strongly drawn upwards and rather projecting in the profile. The bristles of the thoracic dorsum seem to be black, in reflected light they appear brown; in the middle of the dorsum there are but two pairs, the first of which is very much advanced. The short pile upon the thorax and the bristles upon the pleuræ are pale yellowish. Scutellum of a dingy-yellow at the tip, and with two bristles. Abdomen black (a male from Texas shows a dingy yellowish coloring at the basis), appearing almost grayish-black under a very thin pulverulence, which does not prevent it from retaining some lustre; its pile is almost without exception pale yellowish. Feet and coxæ rather saturate yellow, the pile and bristles upon them yellowish. Wings hyaline with a brownish-black picture, which is almost completely radiate towards the end; however, the rays ending in the posterior margin are less developed and less separated from each other than is the case in a normal pattern of this kind; the hyaline intervals between the rays distinctly show that they owe their origin to confluent drops. The root of the wings is but little spotted as far as the beginning of the stigma and the end of the small basal cells; the adjoining portion of the picture is almost without drops, so as almost to assume the appearance of an oblique crossband, running towards the posterior margin; the stigma at its basis contains a small hyaline drop; immediately beyond it, in the marginal cell, there are two square hyaline spots, separated by a brownish-black line; under the first of them the submarginal cell contains a considerable hyaline drop; the anterior side of the fourth vein shows two large drops, the one a little before, the other a little beyond the small crossvein; the discal cell, on the fifth vein, contains three drops, the first of which is the smallest and the second the largest; the third posterior cell contains, besides the small hyaline spot at the basis, four drops of considerable size, three of which are placed at the posterior side of the fifth longitudinal vein; in the poste-

rior corner likewise there are several drops. The third longitudinal vein is without bristles and the small crossvein corresponds to the second third of the discal cell.

Hab. Mexico (Berlin Museum); Texas (Belfrage).

Observation.—The above description, as well as the figure, are prepared after the specimen in the Berlin Museum, which is the original type of Wiedemann's description. Two males, sent by Mr. Belfrage from Texas, agree in all respects, with the only exception that, in one of them, the basis of the abdomen is dingy yellowish. I am in doubt whether *T. mexicana* is not the male of the Cuban species, which I described as *T. melanogastra*, and of which I possess a very imperfect soiled and faded specimen, not sufficient to enable me to form an opinion. A part of the apparent differences may be due to this condition of the specimen. The description of *T. melanogastra* in the first volume of the Monographs says that there is sometimes a clear drop immediately before the end of the second vein; I must complete this statement by saying that this drop exists in the two females of my collection, but not in the male; whether this difference in the picture of the wings is a constant, or at least an ordinary, sexual distinction, I am not prepared to say. The development of the rays ending in the posterior margin in the female of *T. melanogastra* is not even always as complete as Tab. X, f. 24 (drawn after a female specimen) represents it; and the male of my collection approaches very much in this respect the typical male of *T. mexicana*. The differences which fig. 24 and 28 show in the development of the drops in the vicinity of the posterior margin, are of not much importance for specific distinction, as the reticulation in that vicinity is very variable in many species. All these circumstances seem to militate very strongly in favor of specific identity. The only notable difference which I can perceive in the typical male of *T. mexicana* (in the Berlin Museum) as well as in the two males from Texas in my collection, when compared to my single male specimen and my two females of *T. melanogastra*, consists in the position of the hyaline drop in the submarginal cell, which in *T. mexicana* is placed under the first of the two hyaline indentations situated in front of it, while in *T. melanogastra* it is under the brown line which separates the two indentations. This difference is not important and not equally distinct in all specimens, and it is probable that the

comparison of a larger number of them will still more prove its insignificance. There will be no reason then to maintain *T. melanogastra* as a separate species.

57. *T. pura* n. sp. ♀.—Cinerea, thorace fusco-vittato, pilisque albidis instructo, abdomine nigro-piloso, capite pedibusque ex fusco-luteis; setae scutelli quatuor; alae albidæ, præter basin et angulum posticum colore nigro guttato-reticulatæ, in apice radiatæ, guttis in dimidio posteriore confertis, in anteriore rarissimis, tribus tamen majoribus ultra stigma in triangulum dispositis; vena longitudinalis tertia non pilosa.

Gray, thorax with brown longitudinal stripes and white pile, abdomen with black pile, head and feet brownish-yellow; scutellum with four bristles; wings whitish, except the basis and the posterior angle, with a black reticulation, which is radiate on the apex; it is numerously guttate upon the posterior region, sparsely on the anterior; immediately beyond the stigma there are three large drops, disposed in a triangle; the third longitudinal vein is not bristly. Long. corp. cum terebra 0.22; long. al. 0.13.

Gray; abdomen more blackish-gray. Head clay-yellowish; the front more brownish-brick color (which may be due to a discoloration of the described specimen); it is remarkably broad, almost half as broad as the whole head; the usual bristles upon it are black, the bristles on the vertical margin whitish. Antennæ almost brownish-brick color; the short pile on the second segment is whitish; that on the third is blackish; the third joint is gently excised on the upper side; arista blackish-brown. Oral opening of medium size; its anterior edge somewhat drawn upwards and a little projecting in the profile. Proboscis not geniculate; palpi not quite reaching to the anterior edge of the mouth, with black pile. Eyes rounded, their perpendicular diameter but little longer than the horizontal one. Cheeks of a moderate breadth. Thoracic dorsum with indistinctly limited, although well-marked, rather dark brown longitudinal stripes; its short pile is whitish, the bristles black. Scutellum grayish-brown, with a broad grayish border on the sides, and with four black bristles. The ground color of the abdomen is black, the posterior margin of each segment brick-red, especially the last segment, where this border is the broadest; its pile is yellowish-white in the vicinity of the basis only, elsewhere without exception black. Ovipositor flattened, rather broadly truncate at the end, shorter than the last two abdominal segments

taken together, red or brownish-red, with a black border at the tip; its pile is black, whitish on the basal corners only; on the under side there is some whitish pile, conspicuous for its greater length. The whole venter has a brick-brownish coloring; the pile upon it seems to be of the same color as that on the upper side. Feet brownish clay-yellow, almost brownish, brick color. Wings in the reflected light altogether milky-white, in a transmitted light whitish-hyaline. The picture of the wings is black; it does not cover the whole posterior corner of the wing and the longitudinal half of the third posterior cell, contiguous to it, with the only exception of a very striking black border along the sixth longitudinal vein; above this the reticulation begins a little beyond the little basal cells, and, at the anterior margin, with the black stigma, which contains no drops. The space thus left free contains but a few isolated little black spots. The pattern of the picture recalls the European *T. pulchra* Lw. (compare my *Trypetidæ*, Tab. XXIV, f. 2); in the shape and position of the two portions of it which are almost without any drops, it is still more like *T. conjuncta* Lw. (comp. *Trypetidæ*, Tab. XXIV, f. 1), only the drops upon the posterior half of the wing are much more numerous than in those two species; the first almost dropless space begins at the stigma and runs obliquely to the small crossvein; the second is limited posteriorly by the fourth vein and becomes completely confluent with the first space upon and immediately behind the third vein; upon the anterior margin both spaces are separated, immediately beyond the stigma, by a large, somewhat triangular drop, and by a rounded drop which follows it; a third drop, of considerable size, in the submarginal cell, forms a triangle with the other two; the submarginal cell contains, moreover, under the second drop near the anterior margin, another little drop; the second almost non-reticulated space contains two drops on the anterior margin, the first of which is a little distance before, the second immediately beyond the end of the second longitudinal vein, and sends four tolerably well-developed rays towards the margin; the first two of these coincide with the ends of the second and third longitudinal veins, the last two are in the second posterior cell; the first basal cell shows only a row of drops along its posterior side; the first posterior cell contains, besides a few very small drops in the vicinity of its posterior side, a large drop, placed a little before

the end of the discal cell; the numerous drops of the discal cell are of very unequal size, show an inclination to be arranged in two rows and leave more black space on the anterior than on the posterior side; in the third posterior cell the somewhat lacerated reticulation is confined in a very marked manner, to the somewhat larger longitudinal half of the cell, contiguous to the discal cell; both crossveins are perpendicular and less distant from each other than the length of the small crossvein; the third vein is not bristly.

Hab. Massachusetts (Sanborn).

Observation.—I place this species in the genus *Euaesta*, on account of the reticulation, which is radiate on the apex. While *T. festiva*, *spectabilis*, *bella*, *obscuriventris*, *mexicana*, *melanogastra*, and *tenuis*, all closely related, form the solid nucleus of the genus, the connection of *T. pura* with it is a purely artificial one, based upon a resemblance in the picture of the wings; it has more real relationship to those *Urelliæ*, the scutellum of which has four bristles. But in order to place *T. pura* in that genus, it will be necessary to modify its definition, which will have to be done in further developing the system of the *Trypetina*. According to the system adopted in my Monograph of the European *Trypetidæ*, this species would have to be placed in the genus *Tephritis*.

58. *T. abstersa* Lw. ♂ ♀. (Tab. XI, f. 7.)—Cinerea, capite, pedibus et scutello setis quatuor instructo, flavis; alarum dimidium basale colore cinereo obsolete reticulatum, apicale maculâ nigrâ, pulchre radiatâ, ornatum.

Gray, head, feet, and the four-bristly scutellum yellow; the proximal half of the wings with a faded gray reticulation, the distal half with a black, handsomely radiated spot. Long. corp. ♂ 0.12—0.13, ♀ cum terebrâ 0.13—0.14; long. al. 0.12—0.13.

SYN. *Trypeta abstersa* Loew, Dipt. Amer. Cent. II, No. 77.

The ground color of thorax and abdomen is rather variable; generally it is altogether blackish; the humeri, often also the upper side of the pleuræ, the scutellum, the basis of the abdomen, and the posterior margins of its segments usually are, to a greater or lesser extent, clay-yellowish; sometimes the yellowish color is so extended, that, except upon the thoracic dorsum and the metathorax, hardly any blackish is left; nevertheless the ground color of the thorax and of the abdomen is so covered up by a pale

pulverulence and pale yellowish pile, that thorax and abdomen assume a uniform grayish-yellow hue. Head yellow, except the middle of the occiput, which shows a large blackish-brown spot. Front rather broad, attenuated anteriorly; the usual bristles very pale yellowish. Antennæ dark yellow, not quite reaching the somewhat projecting edge of the mouth; the anterior corner of the third joint rounded. The broad oral opening rather round. Proboscis and palpi short, not reaching beyond the anterior edge of the oral opening; proboscis not geniculate. The upper side of the thorax is beset with brown or brownish bristles. Scutellum yellow, with four bristles. Ovipositor reddish-yellow, flat, rather broad, somewhat shorter than the last two abdominal segments taken together, beset with whitish pile. Feet yellow, front femora with yellowish bristles. Wings hyaline; their proximal half is somewhat less limpid than the distal one, rather uniformly pictured with a loose, gray reticulation, which is faintest near the anterior margin; the distal half of the wing is occupied by the radiated black spot, characteristic of the genus *Urellia*, which extends from the anterior margin to the fourth vein; this spot emits two narrow oblique rays, running towards the anterior margin; the first begins at the anterior end of the small crossvein and runs to the end of the colorless stigma; the second, shorter one, reaches the margin in the middle between the tip of the stigma and beginning of the black spot itself; three rays run towards the apex, of which the ends of the two posterior ones coincide with the ends of the third and fourth veins, where they are somewhat expanded; the shortest, anterior ray, sometimes separated from the body of the black spot by two drops only, reaches the anterior margin between the ends of the second and third veins; the first two of the rays running towards the posterior margin cross the middle of the second posterior cell; the narrower third ray follows the posterior crossvein and is sometimes connected with the second by a gray bridge, which divides the hyaline indentation between them into two large drops; in the first posterior cell, above and a little before the posterior crossvein there is a large hyaline drop, which, upon its proximal side, is bordered with black or blackish. The third vein is not bristly.

Hab. North America (coll. Winthelm); Cuba (Gundlach).

Observation 1.—I have described *T. abstersa* in the Dipt. Am.

Cent. II, after a North American female in the Winthem collection. I have received since several specimens of a Cuban *Trypeta* from Mr. Gundlach, which I suppose to be the same species. They are somewhat smaller, have a more extended blackish coloring, and the incomplete gray reticulation of the proximal half of the wing is considerably darker towards the posterior margin. Unfortunately, I have not the original specimen of the Winthem collection at hand for comparison, and, therefore, cannot finally decide about the specific identity. In the figure of the wing, the gray reticulation of its proximal half is represented by the engraver as too distinctly guttate, in fact more so than is the case in either the Cuban or in the typical specimen.

Observation 2.—*T. abstersa* belongs in the genus *Urellia*, and in the group of species having four bristles upon the scutellum. The more developed picture on the basal half of the wing requires, however, that it should be placed on the limit of this genus and in the close relationship of *T. pura* and similar species.

59. *T. polyclona* n. sp. ♀.—Albido-cinerea, capite squæ flavis; setæ scutelli quatuor; alæ hyalinæ, præter dimidiū apicalis maculam magnam nigram, radios novem emittente, duos in costam, duos in apicem et quinque in marginem posticum excurrentes.

Whitish-gray, head and feet yellow. Scutellum with four bristles; wings hyaline, upon their distal half with a large black spot, which emits nine rays, namely, two to the anterior margin, two to the apex, and five to the posterior margin. Long. corp. cum terebrâ 0.15; long. al. 0.14.

Of this handsome species I possess only a single, rather worn, specimen. Head yellow, of the same structure as in *T. abstersa*, only the front comparatively narrower. Thorax, scutellum, and the whole abdomen whitish-gray. The bristles on the scutellum are broken off, nevertheless it is apparent that they were four in number. Ovipositor black, somewhat longer than the last two abdominal segments taken together. Feet yellow. Wings whitish-hyaline, upon their distal half with a large spot, emitting nine rays towards the margin of the wing; the spot is a little removed from the small crossvein, near which, in the first basal cell, there is an irregular blackish spot; the first ray runs from the anterior end of the small crossvein in an oblique direction through the otherwise colorless stigma, to the costal vein, which, at the place where it is thus reached, has a conspicuously black

color; the second ray also runs obliquely to the anterior margin, which it reaches before the middle of the distance between the ends of the second and third longitudinal veins; the third and fourth rays run towards the apex and end upon the ends of the third and fourth veins; among the five rays running towards the posterior margin, the first two cross, as usual, the second posterior cell, and the third follows the posterior crossvein; the fourth originates but little beyond the posterior crossvein, exactly at the place where the spot incloses a large drop, placed on the anterior side of the fourth vein; it runs almost parallel to the preceding ray as far as the posterior margin; the last ray finally originates at the posterior end of the small crossvein, and runs in a very oblique direction, diverging from that of the preceding ray, towards the margin, in the vicinity of which the intensity of its coloring is diminished; the drops, through the coalescence of which the hyaline intervals between the last three rays are formed, are indicated by the irregular outlines of the last two rays; besides the drop already mentioned, which is situated on the anterior side of the fourth vein, the black spot contains a second drop immediately beyond the end of the second vein.

Hab. Cuba (Gundlach).

Observation 1.—*T. polycloa* is a typical *Urellia*, and belongs, as well as *T. abstersa*, to the division with four bristles on the scutellum.

Observation 2.—One would almost be tempted to recognize in this species the *T. mevarna* Walker, List, etc., IV, p. 1023, from Florida, which is an *Urellia*. But a positive identification is prevented by the circumstance that Walker mentions the feet as having black pile, which is not at all the case in my species.

60. *T. solaris* Lw. ♀. (Tab. X, f. 19.)—Albido-cinerea, capite pedibusque flavis, setæ scutelli duæ; alæ albo-hyalinæ, prope venam transversalem mediam subinfuscatæ, in dimidio apicali maculâ magnâ nigrâ ornatæ, guttas duas includente et radios septem integros, octavamque abbreviatum emittente.

Whitish-gray, head and feet yellow; scutellum with two bristles; wings whitish-hyaline, brownish in the vicinity of the small crossvein, upon the distal half with a large black spot, which contains two drops and emits eight rays, the last of which alone is shortened. Long. corp. cum terebrâ 0.17; long. al. 0.16—0.17.

Syn. *Trypeta solaris* Loew, Monogr., etc., I, p. 84. Tab. II, f. 19.

Hab. Georgia (Osten-Sacken).

Observation.—To the above-quoted description I must add, in order to facilitate the distinction from the following species, that the rays running towards the posterior margin are strongly marked, and that the last of them ends abruptly at the fifth vein (the distance is too large in the figure); that there is no trace of a blackish spot near the fifth vein, but that, in the discal cell, immediately beyond its middle, there is an exceedingly minute gray mark. This species is a typical *Urellia*, of the group with two bristles on the scutellum.

61. *T. actinobola* n. sp. ♂.—Albido-cinerea, capite pedibusque flavis, setæ scutelli duæ, alæ totæ albo-hyalinæ, præter punctum nigrum venæ quintæ oppositum in dimidio apicali maculâ magnâ nigrâ ornata, guttas duas includeute et radios septem integros, octavumque abbreviatum emittente.

Whitish-gray, head and feet yellow, scutellum with two bristles; wings altogether whitish hyaline, with the exception of a punctiform dot on the fifth longitudinal vein and of a large black spot upon the distal half of the wing; the latter contains two drops and emits eight rays, the last of which alone is abbreviated. Long. corp. 0.13—0.14; long. al. 0.15.

This species is so very like the preceding that the mention of the differences in the picture of the wings will be sufficient for its recognition. There is no trace here of the brownish coloring which, in *T. solaris*, surrounds the small crossvein, and likewise none of the minute mark in the first basal cell, near the small crossvein; the little dot beyond the middle of the discal cell which occurs in *T. solaris* is likewise wanting here; but instead of these, there is, on the posterior side of the fifth vein, nearly under the end of the first vein, a very well-marked punctiform blackish dot; the large black spot on the apex is very like that in *T. solaris*, with the following differences: the first ray is not extinguished within the stigma, but crosses it without being discolored and reaches the margin; the rays running towards the apex and the posterior margin are very much narrower; the same applies to the last ray, which, moreover, is interrupted already in the discal cell, before reaching the fifth vein.

Hab. Texas (Belfrage).

ANALYTICAL TABLE OF THE SMALLER GENERA,

ADOPTED FOR THE NORTH AMERICAN TRYPETÆ.

	A picture on the wings is extant, but it is never reticulate.	2
1	{ The picture is entirely or partly reticulate, sometimes altogether wanting. ¹	15
2	{ Scutellum with six bristles. ² I. HEXACHETA	
	{ Scutellum not with six bristles.	3
3	{ The third vein conspicuously curved forwards at the tip. II. ACROTOXA.	
	{ The third vein not curved forwards at the tip.	4
4	{ The picture of the wings is on the rivulet-pattern. VI. STRAUSSIA.	5
	{ The picture of the wings is not on the rivulet-pattern.	8
5	{ Body elongate, abdomen narrower than thorax.	6
	{ Body short, abdomen as broad as thorax.	6
6	{ Horizontal diameter of the eyes remarkably short. III. STENOPE.	
	{ Horizontal diameter of the eyes not shorter than usual.	7
7	{ The antepenultimate section of the fourth vein straight. IV. ACIDIA.	
	{ The antepenultimate section of the fourth vein curved. V. EPOCHRA.	
8	{ Coloring of the body generally light, never black.	9
	{ Coloring of the body black.	13
9	{ Upon the middle of the wing there are two crossbands converging towards the posterior margin.	10
	{ No crossbands converging posteriorly upon the middle of the wing.	11
10	{ The third longitudinal vein is gently curved backwards towards the end; head not tumid. VII. SPILOGRAPHA.	
	{ The third longitudinal vein is straight, up to its tip: head perceptibly swollen. VIII. OEDICARENA.	
11	{ Wings with four very oblique crossbands and with very oblique approximate crossveins.	12
	{ Wings with crossbands which are rather perpendicular or dissolved in spots and with very steep crossveins. X.I. TRYPETA.	

¹ Among these species is *T. Lichtensteini*, the picture of which cannot well be called reticulate, but rather spotted.

² Compare also *Epochra*.

- | | | | |
|----|---|--|-----------------------|
| 12 | { | Scutellum tumid, bituberculate. | IX. PERONYMA. |
| | | Scutellum of the ordinary structure, not swollen, although convex. | X. PLAGIOTOMA. |
| 13 | { | Crossveins conspicuously approximate, scutellum unusually swollen. | XII. OEDASPIS. |
| | | Crossveins not approximate, scutellum not swollen. | 14 |
| 14 | { | Scutellum yellow, with four bristles, wings with black crossbands. | XIII. RHAGOLETIS. |
| | | Scutellum black, with two bristles, wings black, with hyaline indentations along the margin. | XIV. ACIURA. |
| | | Fifth vein strongly bristly; scutellum with six bristles. | |
| 15 | { | | XV. BLEPHARONEURA. |
| | | Fifth vein not bristly; scutellum with six or two bristles. | 16 |
| 16 | { | Wings banded on the apex. | XVI. ACROTENIA. |
| | | Wings not banded on the apex. | 17 |
| 17 | { | Face spotted. | 18 |
| | | Face not spotted. | 19 |
| 18 | { | Wings very much dilated; pattern of the picture not radiating. | XVII. EUTRETA. |
| | | Wings not dilated; the pattern of the picture radiating along the margin (Tab. XI, f. 3). | XVIII. CARPHOTRICHIA. |
| 19 | { | Front remarkably broad. | 20 |
| | | Front narrow, or of medium breadth. | 21 |
| 20 | { | Third antennal joint short, rounded at the tip, ovipositor conical. | XIX. EUCROSTA. |
| | | Third antennal joint remarkably long, with a very sharp anterior angle, ovipositor flattened. | XX. ACIDOGOXA. |
| 21 | { | Wings without picture, or on the apical half only, with a reticulation dissolved in crossbands. | XXI. ASPILOTA. |
| | | Wings neither without picture, nor, on the apical half, with a reticulation dissolved in bands. | 22 |
| 22 | { | Wings of an evenly broad shape, and with an unusually blunt apex. (Tab. X, f. 18, and Tab. XI, f. 9.) | XXII. ICTERICA. |
| | | Wings of the usual shape or dilated. | 23 |
| 23 | { | Flaps of the proboscis very much prolonged. | XXIII. ENSINA. |
| | | Flaps of the proboscis short, or but little prolonged. | 24 |
| 24 | { | Pattern of the picture not radiating. | XXIV. TEPHRITIS. |
| | | Pattern of the picture radiating. | 25 |
| 25 | { | The whole or nearly the whole surface of the wings with a unicolorous reticulation. | XXV. EUARESTA. |
| | | A star-shaped black picture on the apex, the remaining surface immaculate, or with very few spots, at the utmost with a very faded reticulation. | XXVI. URELLIA. |

DISTRIBUTION OF THE NORTH AMERICAN TRYPETÆ
AMONG THE ADOPTED SMALLER GENERA.

- | | |
|--------------------------------|----------------------------------|
| Gen. I. HEXACHÆTA. | Gen. XI. TRYBETA. |
| 1. <i>eximia</i> Wied. | 18. <i>palposa</i> Lw. |
| 2. <i>amabilis</i> nov. sp. | 19. <i>florescentiæ</i> Lin. |
| Gen. II. ACROTOXA. | Gen. XII. OEDASPIS. |
| 3. <i>suspensa</i> Lw. | 20. <i>polita</i> Lw. |
| 4. <i>fraterculus</i> Wied. | 21. <i>atra</i> Lw. |
| 5. <i>ludens</i> nov. sp. | 22. <i>gibba</i> nov. sp. |
| 6. <i>tricincta</i> nov. sp. | Gen. XIII. RHAGOLETIS. |
| Gen. III. STENOPA. | 23. <i>cingulata</i> Lw. |
| 7. <i>vulnerata</i> nov. sp. | 24. <i>tabellaria</i> Fitch. |
| Gen. IV. ACIDIA. | 25. <i>pomonella</i> Walsh. |
| 8. <i>fratria</i> Lw. | Gen. XIV. ACIDRA. |
| 9. <i>suavis</i> Lw. | 26. <i>insecta</i> Lw. |
| Gen. V. EPOCHRA. | Gen. XV. BLEPHARONEURA. |
| 10. <i>canadensis</i> nov. sp. | 27. <i>pœcilogastra</i> nov. sp. |
| Gen. VI. STRAUSSIA. | Gen. XVI. ACROTÆNIA. |
| 11. <i>longipennis</i> Wied. | 28. <i>testudinea</i> nov. sp. |
| Gen. VII. SPILOGRAPHA. | Gen. XVII. ECTRETA. |
| 12. <i>electa</i> Say. | 29. <i>sparsa</i> Wied. |
| 13. <i>flavonotata</i> Macq. | 30. <i>rotundipennis</i> Lw. |
| Gen. VIII. OEDICARENA. | Gen. XVIII. CARPHOTRICHA. |
| 14. <i>tetanops</i> nov. sp. | 31. <i>culta</i> Wied. |
| Gen. IX. PERONYMA. | Gen. XIX. EUROSTA. |
| 15. <i>sarcinata</i> Lw. | 32. <i>solidaginis</i> Fitch. |
| Gen. X. PLAGIOTOMA. | 33. <i>comma</i> Wied. |
| 16. <i>discolor</i> Lw. | 34. <i>latifrons</i> Lw. |
| 17. <i>obliqua</i> Say. | |

Gen. XX. ACIDIGONA.

35. *melanura* nov. sp.

Gen. XXI. ASPILOTA.

36. *alba* *Lw.*37. *albidipennis* *Lw.*38. *Vernoniæ* *Lw.*

Gen. XXII. ICTERICA.

39. *seriata* *Lw.*40. *circinata* nov. sp.41. *Lichtensteinii* *Wied.*

Gen. XXIII. ENSINA.

42. *humilis* *Lw.*

Gen. XXIV. TEPHRITIS.

43. *angustipennis* *Lw.*44. *finalis* *Lw.*45. *clathrata* *Lw.*46. *geminata* *Lw.*47. *fucata* *Fbr.*48. *albiceps* nov. sp.49. *euryptera* nov. sp.50. *platyptera* nov. sp. ?

Gen. XXV. EUARESTA.

51. *æqualis* *Lw.*52. *festiva* *Lw.*53. *bella* *Lw.*54. *timida* *Lw.*55. *melanogastra* *Lw.*56. *mexicana* *Wied.*57. *pura* nov. sp.

Gen. XXVI. URELLIA.

58. *abstersa* *Lw.*59. *polyclona* nov. sp.60. *solaris* *Lw.*61. *actinobola* nov. sp.

COMPARISON BETWEEN THE EUROPEAN AND THE
AMERICAN FAUNA OF TRYPETINA.

INCOMPLETE as our knowledge of the North American Trypetina is, our scanty materials are, nevertheless, sufficient to enable us to form an approximate idea of their relation to the European fauna. Even a superficial comparison of a North American with a European collection of *Trypetæ* will show, that certain subgenera, characteristic for Europe by the number of species which represent them, are absolutely or almost wanting in America, while, on the contrary, North America possesses other, very peculiar forms, which do not occur in Europe.

We will notice, in the first place, that the subgenus *Urophora*, which, in Europe, embraces fully one-eighth of all the species, is not represented at all in North America.¹ Next to this, we become aware of the fact that the subgenus *Trypeta*, containing another eighth of all the European species, is represented in North America by *Trypeta palposa* only, besides *Trypeta florecentiæ* Lin., which is very probably introduced from Europe.

As forms peculiar to North America and entirely foreign to the circles of relationship of the European *Trypetina*, the species of the subgenera *Hexachæta*, *Acrotoxa*, *Blepharoneura*, *Acrotænia*, *Eutreta*, and *Acidogona* deserve especial attention.

Besides these two very striking differences between the two faunas, a close comparison reveals other discrepancies; as, for instance, that less characteristic European subgenera are entirely wanting in North America, while subgenera occurring in North

¹ In South America likewise, no species of *Urophora* have as yet been found; all the South American species published by European authors as *Urophora* do not belong to this genus at all; most of them are not even *Trypetidæ*, but *Ortalidæ*, with black crossbands on the wings.

America, although wanting in Europe, are found to be closely related to European forms.

Subgenera with a small number of species, occurring in Europe and wanting in North America, are: 1. *Platyparea* (two species; the larva of the typical *Pl. parviflora* lives in the stems of *Asparagus officinalis*); 2. *Euphranta* (one species, on *Asclepias* and *Vincetoxicum*); 3. *Hemilea* (one species); 4. *Hypenidium* (one species); 5. *Chaetostoma* (one species, distinguished by the bristly sides of the face); 6. *Anomæa* (one species, in the fruits of *Cratægas*); 7. *Zonosema* (two species, related to *Rhagoletis*; in the fruits of *Rosa* and *Berberis*); 8. *Rhacochlæna* (one species); 9. *Myopites* (several, but as yet not well separated species; the larvæ live in the flowers of *Inula* and of the related genera); 10. *Sphenella* (one species; larvæ in the flowers of *Senecio*). If we accept the sufficiently well-founded division of the genus *Oedaspis*, in *Oedaspis* and *Orellia*, we have, moreover: 11. *Orellia* (three species, one on *Bryonia*, another on *Zizyphus*), to add to those small European subgenera, which have no representatives in North America.

The subgenera peculiar to North America, but allied to some European forms, are: the subgenera *Straussia* and *Oedicarona*, which resemble *Spilographa*; *Epochra* and *Stenopa*, which stand very close to *Acidia*; *Aspilota*, *Plagiatora*, and *Peronyma*, which all approach *Trypeta*; *Ictericæ*, related to *Oxyphora*; and finally *Eurosta*, closely allied to some species of *Oxygma*.

Such are the differences between the two faunæ; I will now show the resemblances, as far as observed, between them.

The most striking coincidence and the most remarkable for the great number of analogous species, between the two faunas, occurs within the circle of relationship of the European species belonging to the subgenera: *Carphotricha*, *Oxyphora*, *Oxygma*, *Tephritis*, *Ensina*, *Urellia*. Another point of coincidence of the same kind, although less well represented as to the number of species, occurs within the closely related subgenera *Spilographa*, *Acidia*, and *Rhagoletis*. A third one may be noticed within the genus *Oedaspis*. Moreover, the North American species of the subgenera *Trypeta* and *Actura*, a single one in each, are very much like European species of the same subgenera in their general appearance. Two species, common to both continents, have, until now, been ascertained: *Trypeta florescentiæ* (living on

Sonchus) and *Tephritis angustipennis* (occurring in Europe on *Achillea ptarmica*). The specific identity of the American *Acidia fratria* and the European *Acidia heraclei* is not impossible, although as yet not certain.

It must be borne in mind, however, that all the comparative statements, given above, are founded upon a very imperfect knowledge of the North American fauna, and may be considerably modified with an increase of this knowledge.

If the European *Trypetina* be compared, not with those of the whole North American continent, but with the fauna occurring in America within the European latitudes, then some of the more striking differences between the two faunas at once disappear, as those subgenera which are absolutely foreign to Europe (*Hexachata*, *Acrotaxa*, *Blepharoneura*, and *Acrotænia*) do not reach so far north. The occurrence of all four of these subgenera in Brazil proves that they are South American forms, which extend to the southern portions of the North American continent.

It was to be expected that the knowledge of the North American species should exercise an influence upon the subdivision of the old genus *Trypeta* in subgenera, a subdivision hitherto based almost exclusively upon European species. Those North American subgenera, which have no relationship whatever to European forms, of course merely increase the number of subgenera, without influencing in any manner the already existing subdivision. But it is different with those subgenera which contain forms common to both continents, and here the modifying influence of the American fauna becomes apparent. Thus we can already recognize: 1. That the definition of the subgenus *Carphotracha*, founded upon European species, has to be modified, in order to include all the species belonging to it; 2. That the genus *Oxyphora*, in its present acceptation, contains, besides a number of closely allied species, several far too aberrant forms; moreover, that it can no more be separated from the neighboring subgenera merely by the presence of bristles upon the third vein, a character which hitherto has been found sufficient for the distinction of the European species; 3. That the subgenus *Ensina* must be taken in a broader sense than has been done in my Monograph of the European *Trypetæ*, especially through the addition of some species which, in the same Monograph, were placed in *Oxyina*; 4. That the remaining portion of *Oxyina*

must be united generically with the subgenus *Tephritis*, or else that it should be separated from it in some other manner than has been hitherto done; and that, in order to facilitate the identification of species, a new genus, closely allied to the two above named ones, should be founded, for which I have already proposed the name of *Euaresia*.

I reserve for another place to carry out in detail the improvements of the system of *Trypetina* of which I have here given the outline, and I intend, at the same time, to take in consideration the known species from all the other continents.

APPENDIX I.

CRITICAL ENUMERATION OF ALL THE NORTH AMERICAN TRYPETINA DESCRIBED BY OTHER AUTHORS.

1. *acidusa* Walk. very probably belongs to the subgenus *Acrotoxa*; what Mr. Walker says of the curvature of the end of the third longitudinal vein is evidently to be referred to the fourth vein, and the same remark applies to *Trypeta Ethalea* Walker, from Para, which follows upon the former in the text.
2. *acutangula* Thom., unknown to me; probably belongs to the subgenus *Tephritis*.
3. *ænea* v. d. Wulp (Tijdschr. voor Ent. 2 Ser. II, p. 157), described as *T. (Acinura) ænea*; does not belong to the *Trypetide* at all, but to the *Ortalide*, and is synonymous with *Chetopsis ænea* Wied.
4. *albiscutellata* Harr. has never been described, and, hence, is to be stricken out.
5. *antillarum* Macq., described by Macquart as *Urophora* does neither belong to this genus nor to the *Trypetide* in general, but to the *Ortalide*; figure and description agree so little that the identification will be difficult.
6. *arcuata* Walk. is synonymous with *Tritoxa flexa* Wied. (*Ortalidæ*).
7. *armata* R. Desv., published as a *Strauzia*, this is the male of *T. (Straussia) longipennis* Wied.
8. *asteris* Harris is identical with *T. (Eurosta) solidaginis* Fitch, as Baron Osten-Sacken has shown; the choice of the name depended on an erroneous assumption as to the plant on which the larva lives.
9. *aurifera* Thom., a species unknown to me, belonging to the subgenus *Ensina*.
10. *avala* Walk.; the very insufficient data given by the author do not even enable me to decide whether this is a *Trypetida* or an *Ortalida*; even the location of the species in the genus *Urophora* does not help through this dilemma, because *Myennis fasciata* Fab. is placed in the same genus, thus proving that Mr. Walker was not cognizant at all of the characters of this genus.
11. *Beauvoisii* R. Desv., described as *Prionella Beauvoisii*; unknown to me, so far that I am unable to decide whether it is a *Trypetida* or

- an *Ortalida*; the former, however, seems more probable. Its occurrence in America is uncertain, and is only supposed by R. Desvoidy, because the described specimen belonged to *Palisot de Beauvois* collection.
12. **caliptera** Say is synonymous with *T. (Entreta) sparsa* Wied.
 13. **cinotipes** Harris is to be stricken out, as undescribed.
 14. **omma** Wied.; an *Eurosta*, has been described in this volume.
 15. **cornigera** Walk., an unimportant variety of the male of *T. longipennis* Wied.
 16. **cornifera** Walk., same remark as the preceding.
 17. **cribrata** v. d. Wulp (l. c. p 158), syn. with *T. (Eurosta) latifrons* Lw.
 18. **culta** Wied., a *Carphotricha*, described above.
 19. **dinia** Walk. In the Monographs, Vol. I, I expressed the supposition that it may belong to the relationship of *T. (Acinra) tibialis*; but in doing it, I paid too little attention to the coloring of the body. I think it more probable now that this is a species closely allied to *T. (Hexacheta) eximia* Wied., perhaps even only a badly described variety of this very species.
 20. **electa** Say, a *Spilographa*; has been described in Monographs, Vol. I.
 21. **eximia** Wied., a *Hexacheta*, described above; known long ago as a Brazilian species; its occurrence in Mexico has been discovered recently.
 22. **fasciventris** Macq., synonymous with *T. (Hexacheta) eximia* Wied.
 23. **femoralis** Thoms., an unknown *Urellia* from the group with two bristles on the scutellum.
 24. **imbriata** Macq. is the same as *T. (Carphotricha) culta* Wied.
 25. **flavonotata** Macq., a species closely allied to *T. (Spilographa) electa* Say, but not a mere variety of this species, as I formerly supposed. It is described in this volume.
 26. **flexa** Wied. is a *Tritoza* (*Ortalida*).
 27. **fraterculus** Wied., described by Wiedemann as *Dacus fraterculus*, after a specimen from Brazil; occurs likewise in Peru, New Granada, and Cuba. Belongs in the genus *Acrotora*, and is the same as *Trypeta unicolor* Lw., Monographs, Vol. I. Wiedemann's description did not enable me to recognize this identity, which I have, however, found out since, by comparing the original specimens. As a matter of course, Wiedemann's name has to be maintained.
 28. **fucata** Fabr. may be referred to *Tephritis*, and has been described above.
 29. **fulvifrons** Macq. is *Chatopsis aenea* Wied. (*Ortalidæ*).
 30. **genalis** Thoms., from California; unknown to me; probably a *Tephritis*.
 31. **inermis** R. Desv., published as a *Strauzia*, is *T. (Straussia) longipennis* female.

32. *interrupta* Macq., described as *Urophora*, is not a *Trypetida* at all, but a *Rivellia* of difficult identification.
33. *latipennis* Macq., published as a *Platystoma*, is most certainly a *Trypetida*, in which I cannot recognize anything else but *T. (Eutreta) sparsa*.
34. *Lichtensteini* Wied., described above after the types of Wiedemann's work, and provisionally placed in the genus *Ictericia*, from the typical species of which, however, it is somewhat different.
35. *hogaster* Thoms. is the same as *T. (Acidia) fratria* Lw.
36. *longipennis* Wied. is the typical species of the genus *Straussia*, and, as it seems, undergoes considerable variations. In Monographs, Vol. I, I have given a description of this species, and in the present volume have enumerated the varieties which I have had occasion to see, some of which, however, may be distinct species.
37. *marginipunctata* Macq., almost certainly a *Trypetida*, but which it would be premature to identify with *T. (Carpotricha) culta* Wied. Macquart's data are so very insufficient that the identification will be very difficult.
38. *melliginis* Fitch is a *Rivellia*, under which head it has been discussed.
39. *mevarna* Walk., an *Urellia*, unknown to me.
40. *mexicana* Wied., an *Euresta*; the above description has been prepared from Wiedemann's typical specimen.
41. *narytia* Walk.; the remarks appended above to *arala* Walk. may be repeated here.
42. *novæboracensis* Fitch is synonymous with *T. (Eutreta) sparsa* Wied.
43. *nigriventris* Macq., erroneously described as *Urophora*; it is a *Trypetida*, which I do not know, and concerning the systematic position of which I am in doubt.
44. *obliqua* Macq. is an *Acrotora*; I do not know it.
45. *obliqua* Say is the type of the subgenus *Plagiotoma*; is described in Monographs, Vol. I.
46. *ocresia* Walk. belongs to the subgenus *Acrotora*; I am unable to identify it among the species known to me.
47. *picciola* Bigot (R. de la Sagra, Hist. fis. Cub. Tab. XX, f. 10). This species, described as *Acinia*, is the same as *T. (Eusina) humilis* Lw.
48. *picta* Fabr., type of the genus *Camponeura* (*Ortalida*).
49. *pomonella* Walsh, subgenus *Rhagoletis*, is described in this volume.
50. *quadrifasciata* Macq. I suppose that this species will be found to be identical with *T. (Peromyza) sarcinata* Lw.
51. *quadrivittata* Macq. is an *Ortalida*.
52. *scutellaris* Wied. is an *Ortalida*.
53. *scutellata* Wied. is a *Trypeta* the position of which cannot be made out of Wiedemann's description; Wiedemann's typical specimen unfortunately is no more in existence.
54. *septenaria* Harris is to be stricken out, as undescribed.

55. *solidaginis* Fitch, an *Eurosta*, sufficiently described in Monographs, Vol. I.
 56. *sparsa* Wied.; described in Monographs, Vol. I; typical species of the genus *Eutreta*.
 57. *tellaria* Fitch; described as a *Tephritis*. In Monographs, Vol. I, I have expressed the erroneous supposition that this species belongs to the *Ortalidæ*. It is a Trypetida of the subgenus *Rhagoletis*, and has been described in the present volume.
 58. *tribulis* Harris is not described, and hence must be stricken out.
 59. *trimaiculata* Macq. is *T. (Straussia) longipennis* Wied.
 60. *trifasciata* Harris; not described.
 61. *villosa* R. Desv.; described as *Prionella*; the remark appended above to *Prionella Beauvoisii* may be repeated here.

The result of the above remarks may be summed up as follows:—

1. *Five* of the above-quoted species named by Mr. Harris must be stricken out, as their descriptions have never been published:—

albiscutellata Harris.
cinctipes Harris.
septenaria Harris.

tribulis Harris.
trifasciata Harris.

2. *Ten* species must be transferred to the *Ortalidæ*:—

ænea v. d. Wulp.
antillarum Macq.
arcuata Walk.
flexa Wied.
fulvifrons Macq.

interrupta Macq.
melliginis Fitch.
picta Fabr.
quadrivittata Macq.
scutellaris Wied.

3. *Fifteen* species are merely synonyms of other Trypetidæ; the two marked with an interrogation are not as certain as the others:—

armata R. Desv. = *longipennis* Wied.
asteris Harr. = *solidaginis* Fitch.
calliptera Say = *sparsa* Wied.
cornigera Walk. = *longipennis* Wied.
cornifera Walk. = *longipennis* Wied.
cribrata v. d. Wulp. = *latifrons* Lw.
fasciventris Macq. = *eximia* Wied.

fimbriata Macq. = *culta* Wied.
 ? *logaster* Thoms. = *fratria* Lw.
inermis R. Desv. = *longipennis* Wied.
latipennis Macq. = *sparsa* Wied.
novæboracensis Fitch = *sparsa* Wied.
plioclola Bigot = *humilis* Lw.
trimaculata Macq. = *longipennis* Wied.
 ? *quadrifasciata* Macq. = *sarcinata* Lw.

4. Fifteen species are recognized by me and described in detail in Monographs, Vol. I, and in the present work:—

<i>comma</i> Wied.	<i>longipennis</i> Wied.
<i>culta</i> Wied.	<i>mexicana</i> Wied.
<i>electa</i> Say.	<i>obliqua</i> Say.
<i>eximia</i> Wied.	<i>pomonella</i> Walsh.
<i>flavomaculata</i> Macq.	<i>solidaginis</i> Fitch.
<i>fraterculus</i> Wied.	<i>sparsa</i> Wied.
<i>fucata</i> Fabr.	<i>tabellaria</i> Fitch.
<i>Liichtensteinii</i> Wied.	

5. Sixteen species consequently remain, which I have never seen or have not been able to identify; most of them are undoubtedly *Trypetidæ*; the doubtful ones I have marked with an interrogation:—

<i>acidusa</i> Walk.	<i>marginepunctata</i> Macq.
<i>acutangula</i> Thoms.	<i>mevarna</i> Walk.
<i>aurifera</i> Thoms.	? <i>narytia</i> Walk.
? <i>avala</i> Walk.	<i>nigriventris</i> Macq.
? <i>Beauvoisii</i> R. Desv.	<i>obliqua</i> Macq.
<i>Dinia</i> Walk.	<i>ocresia</i> Walk.
<i>femoralls</i> Thoms.	<i>scutellata</i> Wied.
<i>genalis</i> Thoms.	? <i>villosa</i> R. Desv.

APPENDIX II.

THE descriptions of North American species of *Trypeta*, published by previous authors, but not identified in the foregoing Monograph, are reprinted *verbatim* in the Monographs, etc., Vol. I, p. 94. The following five Californian species of Mr. Thomson, were published since the issue of that volume (*T. liogaster* Thoms. is left among them, as its synonymy with *T. fratria* Lw. is not quite certain).

Thomson, Eugenes Resa, etc., Zoologi, VI, p. 578.

Genus **TRYPETA.**

A. Alæ cubiti ramo submarginali setuloso, cellula anali postice angulo infero breviter sed acute producta, abscissa costali 2a spinula fere nulla. Frons serie laterali 5-setosa. Thorax setarum dorsalium pari pone medio sito. Scutellum 4-setosum. Proboscis brevis.

251. *Trypeta liogaster.*—Ferruginea nitida, abdomine glabro; alis albis, fusco-flexuoso-variegatis; postscutello macula magna didyma nigra. ♀. Long. 5 mill.

Patria. California.

T. Onopordi colore et alarum pictura simillima, abdomine glabro mox distincta. Caput rotundum, fere globosum, ferrugineum, occipite haud excavato; fronte subopaca, subtilissime puberula, serie laterali 5-setosa, setis 2 posterioribus magis ab oculis remotis; epistomate haud brevi, foveis antennalibus minus determinatis, divergentibus, genis angustis, inferne paullo latioribus; peristomio magno, rotundo, utrinque medio seta una validiore nigra instructo, proboscide brevi, capitulo crasso; oculis nudis, fere ovalibus, inferne sat longe descendentibus, orbita frontali parallela, faciali minus divergente. Antennæ subdeflexæ, basi vix distantes, articulo 3o ovali, apice haud mucronato,

epistomatis apicem haud attingente, seta nuda. Thorax ferragineus, nitidus, glaber, setarum dorsalium pari pone medium sito; scutellum subtriangulare, 4-setosum, postscutello macula magna nigra nitida utrinque ornato. Alæ longæ, margine infero vix sinuato, albo-brunneoque flexuoso-variegatæ, macula nempe oblongo-quadrata cellulam totam mediastinam fere occupante alteraque costali triangulari ad cubiti ramum submarginalem usque descendente, pone postcostæ exitum sita, cum maculis duabus disci sinubusque profundis marginis inferioris ante apicem albis, basi inferne late albida; nervis costali abscissa 2a spinula fere nulla, 3a 2a haud duplo brevior, 5a sextæ fere æquali; mediastino apice sub angulo recto costam versus abscendente ibidemque obsoleto; postcostali toto dense setuloso, medium alæ vix attingente; cubiti furca sat longe ante apicem cellule humeralis sita, ramo submarginali parce vix ultra nervum transversum ordinarium setuloso, postice lenissimo curvato et brachiali plane parallelo; humerali mox pone nervum transversum discoidalem desinente; cellula discoidali postice recta truncata nervum transversum ordinarium perpendicularem, longe pone postcostæ exitum situm, in sua tertia posteriore parte excipiente; anali inferne haud longe, sed acute producta, quam humerali brevior. Abdomen ovali-rotundum, supra leviter convexum, glabrum, nitidum, segmento 5o margine postico utrinque setis 4 ornato, 6o parvo fere triangulari, apice truncato, brunneo-nigro, terebram includente. Pedes haud validi, coxis anticis medium mesosterni vix attingentibus; femoribus anticis subtus setosis; tibiis intermediis apice calcari nigro armatis; mesosternum, ut in omnibus, seta in angulo posteriore instructum; epimeris etiam sub alis seta nigra præditis.

B. Alæ ramo cubiti submarginali nudo.

- aa.* Alæ cellula discoidali postice quam nervi transversus ordinarius longitudine vix latiore.
b. Proboscide haud hamato-reflexa.
cc. Alæ cellula discoidali angulo infero recto.
dd. Alæ minus angustæ, cellulis brachiali et humerali haud brevibus, nervo transverso discoidali margine infero alæ approximato, abscissa costali 5a 6a haud duplo longiora.
ee. Alæ albidæ vel hyalinæ, fusco-maculatæ, vel reticulatæ.
f. Scutellum bisetosum. Alæ angulo inferiore cellule analis recto. Thorax setarum dorsalium pari ante medium sito. Femora plerumque tenuia, antica setis 3-4 subtus ornata.
gg. Cellula postcostali nigra vel nigro-fusca.

258. *Trypeta femoralis*.—Nigro-fusca, glauco-pruinosa, capite cum antennis pedibusque flavis, femoribus intermediis scotus setulis 4-5 munitis; alis albidis, macula posteriore fusca apicem versus nullum ramum sed inferne ramum integrum nervum transverso-discoidalem transeuntem emittente, cum cellula postcostali per plagam obliquam connexa. ♂. Long. 4 mill.

Patria. California.

Præcedentibus¹ similis et affinis, femoribus intermediis subtus setis 4-5 validioribus munitis, alis macula posteriore nigro-fusca guttulas 3 majores includente, ramum apicalem nullum sed inferne ramulos 2 abbreviatos ante ramum nervum transverso-discoidalem transeuntem emittente, ramis 2 posterioribus basi tantum indicatis, nervo humerali longitudinaliter ultra medium infuscato distincta.

ff. Scutellum 4-setosum. Alæ angulo inferiore postico cellulæ analis acute subproducto.

261. *Trypeta acutangula*. Nigricans, cano-pruinosa, capite, scutelli apice pedibusque flavis; alis parce fusco-reticulatis, macula majore nigricante, fusco-radiata, cellula postcostali flavescenti. ♂. Long. 4 mill.

Patria. California.

Alis pictura fere *T. cometa*, sed disco et antice parce fusco-reticulatis, cellula postcostali fere tota flavescenti, serie frontali 5-setosa mox distincta. Caput thoracis latitudine, flavo-testaceum, occipite superne fusco, fere truncato; fronte fere transversa, serie utrinque 5-setosa; epistomate brevi, foveis antennalibus fere parallelis, sat discretis; peristomio subrotundo, proboscide haud geniculata; oculis sat magnis, inferne sat longe descendentibus, orbita frontali antrorsum fere convergente. Antennæ breves, subdeflexæ, articulo 3o ovali-rotundo, epistomatis apicem fere attingente, nigro-fusco, seta nudinsecula. Thorax cano-pruinosus, setarum dorsalium pari pone suturam transversam sito; scutello apice late testaceo, 4-setoso, setis apicalibus minoribus approximatis. Alæ sat late, obsenre hyalinae, parcius, disco medio evidentius, fusco-reticulatae, macula posteriore nigricante, subrotunda, guttas 2 costales includente, quarum posteriore paullo

¹ The two preceding species are: *T. glauca* from Sidney, which the author calls "*T. solari* Loew similis et affinis," and *T. meteorica* from Buenos Ayres, described as "præcedenti simillima." O. S.

ante rami marginalis exitum sita, apicem versus ramum bifurcatum, inferne ramos 3 angustos integros fuscus emittente; fuscedine nervi transversus ordinarius sat lata, postice guttis 2 majusculis a macula posteriore magna sejuncta, per strigam obliquam cum cellula postcostali flavescenti connexa et in cellulam discoidalem lobum triangularem emittente; cellula marginali postice guttis 2 magnis albidis, linea transversa fusca separatis ornata; nervis costali abscissa 2a spinula distincta armata, 5a 6a plus quam sesqui longiore; postcostali medium alae attingente; cubiti ramo submarginali postice cum brachio plane parallelo, hoc pone nervum transverso-discoidalem lenissime curvato; cellula discoidali nervum transversum ordinarium, sat longe pone postcostae exitum, nonnihil pone medium alae situm, in sua 5a posteriore parte exeipiente; anali angulo inferno postico acute subproducto. Abdomen unicolor, nigricans, cano-puberulum et pilis depressis parvis rigidis pallidis vestitum, segmento 4o praecedente plus quam duplo longiore. Pedes toti flavi, femoribus haud validis, anticis subtus et tibus 3-4 flavidis longioribus et basi nonnullis brevioribus ornatis.

bb. Proboscide hamato-reflexa. Epistomate brevi, inferne prominente; peristomio antice exciso-assurgente. Palpis prominulis.

hh. Alae fascia recta nervum transversum ordinarium transeunte haud ornata sed fusco reticulata.

i. Scutellum 4-setosum.

264. *Trypeta aurifera*.—Nigricans, capite cum antennis pedibusque testaceis, femoribus ultra medium nigris; alis subhyalinis, obsolete fusco-reticulatis, macula costali quadrata pone spiulam sita, determinate nigricante. ♂ ♀. Long. 3—4 mill.

Patria. California.

T. elongatulae simillima, femoribus ultra medium nigro-fuscis, alis adhuc obsolete fusco-reticulatis, cellula postcostali nigro-fusca, guttam albidam haud includente mox distincta. Caput haud transversum, thoracis latitudine, testaceum, occipite fusco, inferne tumido; fronte subdeclivi, latitudine sua dimidio longiore, utrinque albida 4-setosa, scuto ocelligero nigro-fusco; epistomate brevi, verticali, genis inferne haud latis, superne angustis; peristomio oblongo, antice angulato-exciso, proboscide elongata, geniculata, capitulo longissimo, tenui; oculis magnis obliquis. Antennae basi contiguae, testaceae, breves, epistomatis apicem

attingentes, articulo 3o breviter ovali, seta nuda. Thorax nigricans, cano-pruinosis et pube brevi rigida pallida vestitus, setarum dorsalium pari mox pone suturam sito; scutellum 4-setosum, setis apicalibus parvis. Alæ subhyalinæ, obsolete fusco-reticulatæ, macula quadrata pone spinulam nigricante, determinata: nervis costali abscissa 2a spinula munita, 5a 6a paullo longiore; postcostali medium alæ haud attingente; brachiali et ramo submarginali cubiti parallelis; cellula discoidali nervum transversum ordinarium, longe pone postcostæ exitum paullo pone medium alæ situm, in sua 4a posteriore parte excipiente; anali angulo inferno acuto. Abdomen subdepressum, pruinosis et pube rigida pallida vestitum, segmentis 4o et 5o apice setulis nonnullis marginatis, 6o depresso, nitido, glabro, 2 præcedentibus simul sumptis longitudine æquali. Pedes haud validi, femoribus anticis subtus setis 2-3 ornatis, omnibus nigris, apice cum tibiis tarsisque testaceis.

265. *Trypeta genalis*.—Nigricans, capite pedibusque flavis, femoribus ultra medium nigris; alis hyalinis, fusco-reticulatis, striga obliqua pone spinulam apiceque magis fuscis; abdomine bifariam fusco-maculato, terebra brevi depressa. ♂ ♀. Long. 3-4 mill.

Patria. California.

T. tessellatæ Loew, simillima genis superne angustioribus; alis obscure hyalinis, guttis minoribus, disco interiore basali magis fusco-reticulato distincta; a præcedente capituli labiis brevioribus, alis evidentius fusco-reticulatis discedens. Caput ut in præcedente, fronte paullo latiore, epistomate parum prominule, proboscide capitulo minus elongato. Thorax et scutellum ut in præcedente constructa. Alæ subhyalinæ, fusco-reticulatæ, striga nigro-fusca pone spinulam guttam albam costalem includente, oblique nervum transversum ordinarium transeunte, apice fusco, guttis pluribus majoribus albidis, quarum 5 arcum ante apicem formantibus ornato; nervis omnino ut in præcedente directis, sed postcostali medium alæ attingente, transverso ordinario paullo pone postcostæ exitum sito. Abdomen bifariam fusco-maculatum, pilis brevibus rigidis albidis in margine apicali segmentorum evidentioribus vestitum. Pedes ut in præcedente, sed femoribus anticis subtus setis 4-5 ornatis.

INDEX OF THE TRYPETINA.

- Acidia*, 234, 235, 327, 329, 332
Acidia fratria *Lw.*, 234
 suavis *Lw.*, 235
Acidogona, 285, 327, 330, 331
Acidogona melanura, 283
Acinia *comma* *Macq.*, 281
 fimbriata *Macq.*, 276
 novæboracensis *Fitch*, 274
 picciola *Bigot*, 291
 solidaginis *Fitch*, 280
Acinra, 267, 327, 329
Acinra insecta *Lw.*, 268
Acrotenia, 274, 327, 329, 331
Acrotenia testudinea *Lw.*, 272
Acrotoxa, 227, 327, 329, 331
Acrotoxa fraterculus *Wied.*, 222
 ludens *Lw.*, 223
 suspensa *Lw.*, 222
 tricincta *Lw.*, 225
Anomæa, 332
Aspilota, 286, 327, 330, 332
Aspilota alba *Lw.*, 285
 albidipennis *Lw.*, 286
 Vernoniæ *Lw.*, 286
Blepharoneura, 272, 327, 329, 331
Blepharoneura pœcilogastra *Lw.*,
 270
Carphotricha, 279, 327, 329, 333
Carphotricha culta *Wied.*, 276
Chætostoma, 332
Dacus fraterculus *Wied.*, 222
 parallelus *Wied.*, 227
 serpentinus *Wied.*, 227
Dasyneura, 272
Ensina, 291, 327, 330, 333
Ensina humilis *Lw.*, 291
Epochra, 234, 327, 329, 332
Epochra canadensis *Lw.*, 235
Euaresta, 296, 327, 330
Euaresta æqualis *Lw.*, 308
 bella *Lw.*, 311
 festiva *Lw.*, 309
 melanogastra *Lw.*, 315
 mexicana *Wied.*, 317
 timida *Lw.*, 311
 pura *Lw.*, 320
Euphranta, 332
Eurosta, 280, 327, 329, 332
Eurosta comma *Wied.*, 280
 latifrons *Lw.*, 283
 solidaginis *Fitch*, 279
Entreta, 276, 327, 329, 331
Entreta rotundipennis *Lw.*, 276
 sparsa *Wied.*, 274
Hemilea, 332
Hexachæta, 219, 327, 329, 331
Hexachæta anabilis *Lw.*, 219
 eximia *Wied.*, 216
Hypenidium, 332
Ictericæ, 287, 290, 327, 329, 332
Ictericæ circinata *Lw.*, 288
 Lichtensteini *Wied.*, 28
 seriata *Lw.*, 287
Musca fucata *Fbr.*, 301
 rudicauda *Fbr.*, 254
Myopites, 332
Oedaspis, 250, 260, 263, 328, 329
Oedaspis atra *Lw.*, 256
 polita *Lw.*, 256
 gibba *Lw.*, 260
Oedicærena, 247, 327, 329, 332
Oedicærena tetanops *Lw.*, 245
Orellia, 332
Oxyina, 295, 332, 333
Oxyphora, 280, 287, 295, 309, 332,
 333
Peronyma, 250, 328, 329, 332
 (345)

- Peronyma sarcinata* *Lo.*, 247
Plagiotoma, 252, 328, 329, 332
Plagiotoma discolor *Lo.*, 250
 obliqua *Say*, 251
Platyparea, 332
Platystoma latipennis *Macq.*, 274, 337
Prionella Beauvoisii *Rob. Desv.*, 335
 villosa *Rob. Desv.*, 338


Rhacochlæna, 332
Rhagoletis, 267, 327, 329
Rhagoletis cingulata *Lo.*, 263
 pomonella *Walsh*, 265
 tabellaria *Fitch*, 263

Sphenella, 332
Spilographa, 327, 329, 332
Spilographa electa *Say*, 243
 flavonotata *Macq.*, 244
Stenopa, 234, 327, 329, 332
Stenopa vulnerata *Lo.*, 232
Straussia, 243, 327, 329, 332
Straussia longipennis *Wied.*, 238
Strauzia armata *Rob. Desv.*, 238, 335
 inermis *Rob. Desv.*, 238

Tephritis, 295, 327, 330
Tephritis albiceps *Lo.*, 302
 angustipennis *Lo.*, 293, 333
 asteris *Harris*, 280
 clathrata *Lo.*, 297
 euryptera *Lo.*, 304
 fasciventris *Macq.*, 216, 219
 finalis *Lo.*, 296
 fucata *Fbr.*, 301
 geminata *Lo.*, 298
 Leontodontis *Zett.*, 293
 major *Macq.*, 219
 obliqua *Macq.*, 223
 platyptera *Lo.*, 000
 punctata *Full.*, 254
 quadrifasciata *Macq.*, 248, 249
 segregata *Fruh.*, 293
 socialis *Wied.*, 219
 trimaenulata *Macq.*, 238
Trypeta, 253, 256, 327, 329, 331
Trypeta
 abstersa *Lo.*, 322
 acidula *Wlk.*, 231, 335, 342
 actinobola *Lo.*, 326
 acutangula *Thoms.*, 335
 ænea *v. d. Wulp.*, 335
 æqualis *Lo.*, 308
 alba *Lo.*, 285
 albiceps *Lo.*, 302

Trypeta albidipennis *Lo.*, 286
 albiscutellata *Harris*, 335
 amabilis *Lo.*, 219
 angustipennis *Lo.*, 293, 333
 antillarum *Macq.*, 335
 arcuata *Wlk.*, 335
 armata *Rob. Desv.*, 335
 asteris *Harris*, 335
 atra *Lo.*, 256
 aurifera *Thoms.*, 335, 343
 avala *Wlk.*, 335
 bella *Lo.*, 311
 Beauvoisii *Rob. Desv.*, 335
 biseriata *Lo.*, 252
 caliptera *Say*, 274
 canadensis *Lo.*, 235
 cinctipes *Harris*, 336
 cingulata *Lo.*, 263
 circinata *Lo.*, 288
 clathrata *Lo.*, 297
 comma *Wied.*, 280
 consobrina *Lo.*, 230
 cornifera *Wlk.*, 238, 336
 cornigera *Wlk.*, 238, 336
 cribrata *v. d. Wulp.*, 336
 culta *Wied.*, 276
 dimia *Wlk.*, 336
 discolor *Lo.*, 250
 electa *Say*, 243
 ethalea *Wlk.*, 335
 euryptera *Lo.*, 304
 eximia *Wied.*, 216, 336
 fasciventris *Macq.*, 336
 femoralis *Thoms.*, 336, 342
 festiva *Lo.*, 309
 fimbriata *Macq.*, 336
 finalis *Lo.*, 296
 flavonotata *Macq.*, 244, 336
 flexa *Wied.*, 336
 florescentiæ *Lin.*, 254, 331
 fraterculus *Wied.*, 222, 336
 fratria *Lo.*, 234
 fucata *Fbr.*, 300
 fulvifrons *Macq.*, 336
 geminata *Lo.*, 298
 genalis *Thoms.*, 336, 347
 gibba *Lo.*, 260
 grandis *Macq.*, 231
 hamata *Lo.*, 229
 humilis *Lo.*, 291
 inermis *Rob. Desv.*, 336
 insecta *Lo.*, 268
 integra *Lo.*, 230
 interrupta *Macq.*, 337
 latifrons *Lo.*, 283
 latipennis *Macq.*, 337
 latipennis *Wied.*, 274
 Lichtensteini *Wied.*, 259

- Trypeta liogaster* *Thoms.*, 235, 337,
 340
longipennis *Wied.*, 238
ludens *Lw.*, 223
Macquartii *Lw.*, 267
marginepunctata *Macq.*,
 337
melanogastra *Lw.*, 315
melanura *Lw.*, 283
melliginis *Fitch*, 337
mevarna *Wlk.*, 325, 337
mexicana *Wied.*, 317
narytia *Wlk.*, 339
nigerrima *Lw.*, 258
nigriventris *Macq.*, 337
novæboracensis *Fitch*, 337
obliqua *Say*, 251
obliqua *Macq.*, 337
obscuriventris *Lw.*, 313
ocresia *Wlk.*, 231, 336
palposa *Lw.*, 253, 331
parallela *Wied.*, 229
peregrina *Lw.*, 292
phœnicura *Lw.*, 269
picciola *Bigot*, 337
picta *Fbr.*, 337
platyptera *Lw.*, 306
pœcilogastra *Lw.*, 270
polita *Lw.*, 256
polyclona *Lw.*, 324
pomonella *Walsh*, 265
pseudoparallela *Lw.*, 230
pura *Lw.*, 320
quadrifasciata *Macq.*, 337
quadrivittata *Macq.*, 337
rotundipennis *Lw.*, 276
sarcinata *Lw.*, 247
- Trypeta scutellaris* *Wied.*, 337
scutellata *Wied.*, 337
seriata *Lw.*, 287
serpentina *Wied.*, 226
solidaginis *Fitch*, 279
solaris *Lw.*, 325
sparsa *Wied.*, 274
spectabilis *Lw.*, 309
suavis *Lw.*, 235
suspensa *Lw.*, 222
tabellaria *Fitch*, 263
tennis *Lw.*, 316
testudinea *Lw.*, 272
tetanops *Lw.*, 245
timida *Lw.*, 311
tribulis *Harris*, 338
tricincta *Lw.*, 225
trimaculata *Macq.*, 338
unicolor *Lw.*, 222
Vernoniæ *Lw.*, 286
villosa *Rob. Desr.*, 338
vulnerata *Lw.*, 232
- Urella*, 295, 327, 330
Urella abstersa *Lw.*, 322
actinobola *Lw.*, 326
polyclona *Lw.*, 324
solaris *Lw.*, 325
- Urophora*, 331
Urophora antillarum *Macq.*, 335
bivittata *Macq.*, 231
interrupta, 337
nigriventris *Macq.*, 337
scutellaris *Macq.*, 267
vittithorax *Macq.*, 227
- Zonosema*, 245, 332



EXPLANATION OF THE PLATES.

ORTALIDÆ.

PLATE VIII.

1. *Amphicnephes pertusus* nov. sp.
2. *Himeroessa pretiosa* nov. sp.
3. *Rivellia conjuncta* nov. sp.
4. *Rivellia viridulans* Rob. Desv.
5. *Rivellia quadrifasciata* Macg.
6. *Rivellia variabilis* nov. sp.
7. *Rivellia flavimana* nov. sp.
8. *Rivellia pallida* nov. sp.
9. *Myrmecomymia myrmecoides* Lw.
10. *Tritoxa flexa* Wied.
11. *Tritoxa cuneata* nov. sp.
12. *Tritoxa incurva* nov. sp.
13. *Camptoneura picta* Fbr.
14. *Diacrita costalis* Gerst.
15. *Diacrita æmula* nov. sp.
16. *Idana marginata* Say.
17. *Tetanops luridipennis* nov. sp.
18. *Tetanops integer* nov. sp.
19. *Anacampta latiuscula* nov. sp.
20. *Ceroxys obscuricornis* nov. sp.
21. *Ceroxys ochricornis* nov. sp.
22. *Ceroxys canus* Lw.
23. *Ceroxys similis* nov. sp.
24. *Tephronota humilis* nov. sp.
25. *Stictocephala cribrum* nov. sp.

26. *Stictocephala cribellum* nov. sp.
27. *Callopietria annulipes* Macq.
28. *Stictocephala corticalis* Fitch.
29. *Stictocephala vau* Say.
30. *Pterocalla strigula* nov. sp.

PLATE IX.

1. *Oedopa capito* Lw.
2. *Oedopa capito* Lw.
3. *Oedopa capito* Lw.
4. *Euphara cærulea* Macq.
5. *Notogramma stigma* Fbr
6. *Scoptera colon* Lw.
7. *Euxesta spoliata* Lw.
8. *Euxesta pusio* Lw.
9. *Euxesta notata* Wied.
10. *Euxesta costalis* Fbr.
11. *Euxesta quaternaria* Lw.
12. *Euxesta binotata* Lw.
13. *Euxesta annouæ* Fbr.
14. *Euxesta Thomæ* Lw.
15. *Euxesta abdominalis* Lw.
16. *Euxesta alternans* Lw.
17. *Euxesta stigmatias* Lw.
18. *Euxesta eluta* Lw.
19. *Chætopsis ænea* Wied.
20. *Chætopsis debilis* Lw.
21. *Stenomyia tenuis* Lw.
22. *Eumetopia rufipes* Macq.
23. *Eumetopia varipes* Lw.
24. *Epiplatea erosa* Lw.
25. *Stenomacra Guerinii* Bigot.
26. *Idiotypa appendiculata* nov. sp.
27. *Cœlometopia bimaculata* nov. sp.
28. *Hemixantha spinipes* nov. sp.
29. *Melanostoma afinis* nov. sp.

TRYPETIDÆ.

PLATE X.


TRYPETA.


- | | |
|---|---|
| 1. discolor <i>Lw.</i> | 16. solidaginis <i>Fitch.</i> |
| 2. longipennis <i>Wied.</i> , ♂. | 17. humilis <i>Lw.</i> |
| 3. longipennis <i>Wied.</i> , ♀. | 18. seriata <i>Lw.</i> |
| 4. fratria <i>Lw.</i> | 19. solaris <i>Lw.</i> |
| 5. suspensa <i>Lw.</i> | 20. æqualis <i>Lw.</i> |
| 6. fraterculus <i>Wied.</i> | 21. festiva <i>Lw.</i> |
| 7. electa <i>Say.</i> | 22. latifrons <i>Lw.</i> |
| 8. insecta <i>Lw.</i> | 23. bella <i>Lw.</i> |
| 9. palposa <i>Lw.</i> | 24. melanogastra <i>Lw.</i> , ♀. |
| 10. suavis <i>Lw.</i> | 25. timida <i>Lw.</i> |
| 11. cingulata <i>Lw.</i> | 26. obscuriventris <i>nov. sp.</i> |
| 12. polita <i>Lw.</i> | 27. spectabilis <i>nov. sp.</i> |
| 13. sparsa <i>Wied.</i> | 28. mexicana <i>Wied.</i> |
| 14. rotundipennis <i>Lt</i> | 29. tenuis <i>nov. sp.</i> |
| 15. clathrata <i>Lw.</i> | 30. peregrina <i>nov. sp.</i> |


PLATE XI.


TRYPETA.


- | | |
|---|--|
| 1. geminata <i>Lw.</i> | 15. tetanops <i>nov. sp.</i> |
| 2. comma <i>Wied.</i> | 16. sarcinata <i>Lw.</i> |
| 3. culta <i>Wied.</i> | 17. atra <i>Lw.</i> |
| 4. finalis <i>Lw.</i> | 18. nigerrima <i>Lw.</i> |
| 5. albiceps <i>nov. sp.</i> , ♂. | 19. Iudens <i>nov. sp.</i> |
| 6. melanura <i>nov. sp.</i> | 20. parallela <i>Wied.</i> |
| 7. abstersa <i>Lw.</i> | 21. consobrina <i>nov. sp.</i> |
| 8. Vernoniæ <i>Lw.</i> | 22. hamata <i>nov. sp.</i> |
| 9. Lichtensteinii <i>Wied.</i> | 23. integra <i>nov. sp.</i> |
| 10. albidipennis <i>Lw.</i> | 24. pseudoparallela <i>nov. sp.</i> |
| 11. alba <i>Lw.</i> | 25. serpentina <i>Wied.</i> |
| 12. phæricura <i>nov. sp.</i> | 26. grandis <i>Mocq.</i> |
| 13. testudinea <i>nov. sp.</i> | 27. bivittata <i>Mocq.</i> |
| 14. obliqua <i>Say.</i> | |




















CORRECTIONS TO VOLUME III.

Page 283, as a synonym of *T. latifrons* insert:—

Trypeta cribrata v. D. WULF, Tijdschr. v. Entom. 2 Ser. Vol. II, p. 158.
Tab. V, f. 15.

Observation (by the Editor) to page 153.—This volume was already printed when I received from Mr. E. Burgess specimens taken near Beverly, Mass., and showing the characters of *Scoptera vibrans* Lin., as distinguished from *S. colon* Loew. Immediately afterwards I found in the Museum of Comparative Zoölogy a precisely similar specimen, apparently taken near Cambridge, Mass.—O. S.



ADDITIONS AND CORRECTIONS
TO THE PREVIOUSLY PUBLISHED VOLUME.


Corrections to Volumes I and II, furnished by Mr. Loew.

VOLUME I.

- Page 10, line 3 from bottom, instead of *Cylindrotoma Meig.* read *Macq.*
" 17, " 15 from top, instead of *wing*, read *margin of the wing.*
" 17, " 14 from bottom, *Metoponia (=Inopus)*, strike out *Inopus.*
" 18, " 20 " instead of *Aissa*, read *Antissa.*
" 19, " 15 from top, instead of *fourth cell of posterior margin*, read
fourth posterior cell.
Page 21, lines 17 and 12 from bottom, instead of *Obsebius*, read *Opsebius.*
(The same name must be corrected in the Index.)
Page 38, line 17 from bottom, instead of *legs proportionately short*, read
legs very long and slender, with the tarsi proportionately short.
Page 39, line 12 from bottom, instead of *generally*, read *mostly.*
" 40, " 12 " instead of *with no* read *without.*
" 42, " 6 " instead of *tarsi* read *tibie.*
" 47, *Asteidae*; add at the end: (*Sigaloëssa* alone has a posterior cross-
vein).
Page 55, line 4 from top, instead of *is*, read *it is.*
" 56, " 4 from bottom, instead of *and*, read *or.*
" 57, " 6 from top, instead of *and*, read *or.*
" 70, " 10 " instead of *short*, read *thin.*
" 75, " 4 " instead of *edge*, read *border.*
" 90, " 12 from bottom, instead of 23, read 24.
" 91, " 20 from top, instead of *first* read *fifth.*
" 178, " 10 from bottom, before the word "*longitudinal*", add *fourth.*
-

VOLUME II.

- Page 299, lines 7 from top and 13 from bottom, instead of *Nordhausen*,
read *Nordshausen.*



CORRECTIONS TO VOLUME IV.

(By C. R. OSTEN-SACKEN)

Page 2, line 6 from bottom, instead of *general*, read *common*.

" 16, " 5 from top, instead of p. 11, read p. 3-7.

" 23, " 8 " instead of *auxiliary*, read *subcostal* (this error occurs twice on the same line).

Page 129, line 3 from bottom, instead of *all the*, read *most*.

" 132, lines 2, 4, 14, 15 from bottom, instead of *Paratropesa*, read *Paratropesa*.

The same error occurs on page xl, line 4 from bottom.

" 49, " 18 "

" 333, " 2 from top.

" 343, " 3 from bottom, column first.

" 345, " 8 "

Page 134, line 4 from bottom, strike out *lin*.

" 159, " 15 " instead of *is*, read *are*.

" 179, " 19 from top, instead of 1822, read 1829.

" 219, " 4 from bottom, before *yellowish*, insert *femora*.

" 249, lines 15 and 16 from bottom: the quotation from Doleschall given here refers to his paper in pamphlet form; the full quotation may be found on page 16, line 5 from top, where p. 387 should be read, instead of p. 11.

Page 275, line 11 from top, instead of *paupera*, read *pauper*.

The same error occurs on page x, line 4 from top, column first.

" 277, " 5 "

" 278, " 3 "



" 344, " 4 from bottom, column sec'd.

Page 293, line 13 from top, instead of *ruficornis* Wied. and *erythrocephala* Macq., read *ruficornis* Macq. and *erythrocephala* Wied.





Page 295, line 3 from bottom, instead of p. 15, read p. 391.

" 331, " 18 from top, instead of 17, read 14.





**IMAGE EVALUATION
TEST TARGET (MT-3)**




**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

15 28 25
32 22
20
9

10



ADDITIONS TO VOLUME IV.

(By C. R. OSTEN-SACKEN.)

Page 4. **Ptychoptera.** The larvæ of this genus examined by Brauer, differ from all the known larvæ of Tipulidæ in having the head *not* imbedded up to the mouth in the first thoracic segment, but entirely free. This observation justifies the isolated position which I have given to this group in the family. Compare Verh. Zool. Bot. Ges. 1869, p. 844.

Page 23. The analytical table, given here, would be improved by being modified thus:—

I. A single submarginal cell.

Antennæ 14-jointed.

Antennæ 16-jointed.

Empodia indistinct or none.

Sect. II. **Limnobina anomala.**

Sect. I. **Limnobina.**


II. Two submarginal cells. Empodia distinct, etc. etc.

Page 49. The same modification may be made on this page.

Page 57. **Dicranomyia.** My remarks concerning the differences between this genus and *Limnobia* apply to those North American and European species which I had occasion to compare. I have accumulated as many distinctive characters as a careful comparison of the material before me could disclose; but I should not wonder at all if forms occurred the location of which remained doubtful, all the enumerated distinctive characters notwithstanding.

Page 81. Mr. Loew draws my attention to the fact, that the antennæ of *Rhipidia* cannot be properly called *pedicelled*, because the short stems, connecting the joints, are processes of the anterior part of the joint and not of the posterior one.

Page 102. **Styringomyia.** During my passage through Stockholm in 1872, I made the interesting discovery that this genus, besides its occurrence in amber and copal, is found living in Africa. I saw several specimens among the unnamed diptera from Caffraria (from Wahlberg's voyage) in the Stockholm Museum. The species was apparently different from that included in copal, which I possess.



Page 115. *Toxorhina muliebris* O. S. ♂. I found three males and one female near Tarrytown, N. Y., in July, 1871. They all have the discal cell open, which, therefore, seems to be the rule in this species. The stripes of the thorax are dark brown; the position of the great crossvein is variable, sometimes at the very basis of the discal cell, sometimes before it.

Page 138, at the bottom. *Sigmatomera*. I described this new genus, from Mexico, without adding the description of the typical species, which, as I anticipated, would be soon published in a new fascicle of Mr. Bellardi's *Saggio*, etc. This publication having been, in the mean time, indefinitely postponed, it becomes necessary to supply the above mentioned omission.

Sigmatomera flavipennis n. sp.—Yellow, antennæ long, black, except the first joint, which is yellow; front feet and middle femora yellow (the remaining feet as well as the middle tibiæ and tarsi, are wanting). Wings tinged with yellowish; central crossveins and fifth vein slightly bordered with brown. Long. corp. 0.56—0.6; long. al. 0.64.
Hab. Mexico (Sumichrast).


Page 173. *Psiloconopa*. I had occasion to examine specimens of *P. Meigenii* Zett., since the publication of Vol. IV, and have become aware that my opinion about its location was erroneous. This genus is related to *Trimicra*, and its venation is exactly like the latter genus, the subcostal crossvein being quite remote from the tip of the auxiliary vein. The two other European species, mentioned on pages 173 and 174 as *Psiloconopa*, do not belong to this genus at all, and are much better placed in the genus *Goniomyia*. The above correction will necessitate changes in all the passages, where the genus *Psiloconopa* is mentioned. Such passages are the following:—

Page 21, line 4 from bottom, instead of *Psiloconopa*?, read *Goniomyia*?.
" 36, " 7 from bottom, strike out the whole passage beginning with *is represented*.

Page 36, line 3 from bottom, add *Psiloconopa*.

" 47, modify the analytical table thus:—

	The distance, etc.	30
29	{ The distance, etc.	Gen. XXII. GNOPHOMYIA.
	Seventh longitudinal vein straight; Tab. II, f. 1.	31
30	{ Seventh longitudinal vein conspicuously bisinuated; Tab. I, fig.	
	20.	Gen. XXI. SYMPLECTA.
	{ Three terminal joints of the antennæ abruptly smaller.	
		Gen. XVIII. TRIMICRA.
31	{ Three terminal joints, etc., not abruptly smaller.	
		Gen. XX. PSILOCONOPA.



Page 49, line 14 from bottom, transfer Gen. XXII. *Psilocoenopa*, as Gen. XX, after *Chionea*.


Page 135, line 2 from bottom, strike out the passage beginning with "I believe now" and ending with "typical Eriopterina."

Page 137, line 11 from top, instead of *Psilocoenopa*, read *Goniomyia*.

" 173, line 13 from bottom, strike out the whole paragraph beginning with the words: "A genus closely allied, etc.," as well as its continuation on the next page, down to the "Description of the species."

Page 176. Gen. XXII. *Psilocoenopa* should be placed between *Chionea* and *Symplecta* as Gen. XX, with the following notice: Established by Zetterstedt in 1840 (*Fauna Lapponica*, p. 847), and later in *Dipt. Scand.* X, p. 4007, upon a single species, found in Sweden. This genus, as far as I have been able to study it upon a dry specimen, is related to *Trimicra*, and its venation is exactly the same, the subcostal crossvein being quite remote from the tip of the auxiliary vein, etc. However, it does not have the last three antennal joints abruptly smaller, and its general appearance is altogether different.

Page 177, line 10 from bottom, strike out the passage beginning with the words: "The majority" down to the bottom of the page, and read as follows instead: Some European species differ from the American ones in the following characters: in their coloring the black prevails over the yellow; only a few traces of the latter color are left; the auxiliary vein seems to extend much farther beyond the origin of the preforea than is the case in the American species; the structure of the male forceps seems also to show some differences, which, however, I have not been able to ascertain, not having had fresh specimens for comparison. Such species are the *Erioptera lateralis* Macq., *Hist. Nat. Dipt.* II, p. 653 (Syn. *Limnobia flavolimbata* Hal., in Walker's *Ins. Brit. Dipt.* III, p. 304); the *Goniomyia scutellata* Egger and *G. cincta* Egger, in Schiner's *Fauna Austriaca*, Diptera. One of the latter may be synonymous with the former, and Dr. Schiner was perfectly right in referring them to the genus *Goniomyia*. All these species are not unlike the American species of *Gnophomyia* in their general appearance; they differ, nevertheless, in the absence of the marginal crossvein, in the shortness of the first submarginal cell, in the diverging direction of the branches of the fork which form it, and in the presence of yellow in the coloring. It is not impossible, however, that forms of transition may be discovered between these two genera, as well as between them and *Empeda*.



Page 219. *Linnophila inornata* O. S. ♂.—This species was quite common near Tarrytown, N. Y., in June, 1871. Two females which I have before me have the stigma somewhat tinged with brown; the brown at the tip of the femora is more abruptly marked. In the above-quoted description, p. 219, line 4 from bottom, the word *femora* must be added before the word *yellowish*. On the following page, line 5 from top, instead of *about*, read *somewhat less than*. The fore tarsi of the females are shorter than those of the male. The length of the second posterior cell is variable.

Page 260. **Polymera.** This South American genus, never seen by me before the publication of my volume, was doubtfully mentioned among the *Malophilina*. Mr. Loew had opportunities of examining good specimens recently, and published the result in a paper entitled *Über die systematische Stellung d. Gatt. Polymera* Wied. (*Zeitschr. f. d. gesammten Naturwiss. Neue Folge*, 1871, Bd. III, Tab. V, f. 1, 2). It appears now that the antennæ of *Polymera* are not 28-jointed, as was stated by former authors, but 16-jointed, and that there cannot exist the slightest doubt about its location among the *Linnophilina*. It has peculiarities, however, which distinguish it from the ordinary *Linnophilina* of Europe and North America: a remarkably elongated third antennal joint, a structure of the following joints, in the male, which makes them appear double (hence the error of former authors), an open discal cell, and both branches of the fourth longitudinal vein forked (contrary to the rule stated on page 201, No. 2); the wingveins have a rather conspicuous pubescence. Mr. Loew ends his article with a statement of the principal characters of *Polymera*, as recognized by him, which I reproduce here, with a slight modification:—

Polymera.—The number of antennal joints is normal, 16; the first joint of the flagellum is remarkably elongated, cylindrical, beset with long, erect hairs; each of the following joints, in the male, shows two consecutive knots, or swellings, every one of which is provided with a distinct verticil of hairs; in the female, these joints are simply cylindrical, and beset with hairs like the first joint of the flagellum. Wingveins beset with a long pubescence; subcostal crossvein only a short distance from the tip of the auxiliary vein; marginal crossvein distinct, inserted on, or a little beyond the middle of the very long submarginal cell; basal cells comparatively rather short; discal cell open, coalescent with the third posterior cell; five posterior cells; the second with a petiole of a very great length; feet long and slender; tibiæ with very small but distinct spurs; ungues and empodia very small.

