

**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1998

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming are checked below.

- Coloured covers / Couverture de couleur
- Covers damaged / Couverture endommagée
- Covers restored and/or laminated / Couverture restaurée et/ou pelliculée
- Cover title missing / Le titre de couverture manque
- Coloured maps / Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black) / Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations / Planches et/ou illustrations en couleur
- Bound with other material / Relié avec d'autres documents
- Only edition available / Seule édition disponible
- Tight binding may cause shadows or distortion along interior margin / La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure.
- Blank leaves added during restorations may appear within the text. Whenever possible, these have been omitted from filming / Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- Additional comments / Commentaires supplémentaires:

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured pages / Pages de couleur
- Pages damaged / Pages endommagées
- Pages restored and/or laminated / Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed / Pages décolorées, tachetées ou piquées
- Pages detached / Pages détachées
- Showthrough / Transparence
- Quality of print varies / Qualité inégale de l'impression
- Includes supplementary material / Comprend du matériel supplémentaire
- Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image / Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.
- Opposing pages with varying colouration or discolourations are filmed twice to ensure the best possible image / Les pages s'opposant ayant des colorations variables ou des décolorations sont filmées deux fois afin d'obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below /
Ce document est filmé au taux de réduction indiqué ci-dessous.

	10x		14x		18x		22x		26x		30x	
									/			
	12x		16x		20x		24x		28x		32x	

The copy filmed here has been reproduced thanks to the generosity of:



University of British Columbia Library
Vancouver

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol \rightarrow (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

University of British Columbia Library
Vancouver

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.


Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole \rightarrow signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.


MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)



APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14619 USA
(716) 481-0300 - Phone
(716) 288-5989 - Fax



CLASSN. QE326. C2 M5

SERIES _____ No. 7

THE LIBRARY

The University of British Columbia

GEOLOGICAL SURVEY OF CANADA

ROBERT BELL, M.D., D.Sc. (CANTAB.), LL.D., F.R.S.

MINERAL RESOURCES OF CANADA

MANGANESE

*Reprint of Article in Annual Report of Section of Mines for 1902,
Part S, Vol. XV.*




OTTAWA

PRINTED BY S. E. DAWSON, PRINTER TO THE KING'S MOST
EXCELLENT MAJESTY

1904

No 858



GEOLOGICAL SURVEY OF CANADA, OTTAWA.

To Dr. ROBERT BELL, L.L.D., F.R.S., &c.,
Acting Director.


SIR:—The following pamphlet dealing with the manganese industry and deposits of Canada is reprinted from the Annual Report of the Mines Section for 1902, constituting Part S, Vol. XV, N.S., of the Annual Report of the Geological Survey Department.

Pursuant to a policy suggested some years ago and now carried out with your permission, this report is one of a series of similar bulletins intended to give in condensed and popular form, information regarding the mineral resources and possibilities of the country, together with any data regarding similar occurrences in other countries where such would seem to be of use to prospectors and operators in Canada.

I am, sir,
Your obedient servant,

ELFRIC DREW INGALL,
Mining Engineer in Charge.

MINES SECTION,
January 9, 1904.



MANGANESE

MANGANESE

Returns of the production of manganese for 1902 were incomplete Production and the figures of exports have been given as the closest approximation to the output. The exports were 172 tons valued at \$1,062.

The production since 1886 is shown in Table I below :

TABLE I
MANGANESE
ANNUAL PRODUCTION

Calendar Year	Tons	Value	Value per ton
1886	1,786	\$41,499	\$23.20
1887	1,245	43,678	35.07
1888	1,801	47,944	26.62
1889	1,435	32,737	22.80
1890	1,328	27,559	20.74
1891	975	6,694	6.86
1892	115	10,250	89.13
1893	213	14,578	68.44
1894	74	4,180	56.49
1895	125	8,461	67.71
1896	1230	5,975	32.19
1897	151	1,166	7.72
1898	50	1,600	32.00
1899	1,581	20,001	12.65
1900	50	1,800	36.00
1901	440	1,820	4.14
1902	172	1,062	6.17

* Exports

MANGANESE
Exports.TABLE 2
MANGANESE
EXPORTS OF MANGANESE ORE

CALENDAR YEAR	NOVA SCOTIA		NEW BRUNSWICK		TOTAL	
	Tons.	Value	Tons.	Value	Tons.	Value
1875			1,031	829,192	1,031	829,192
1876	6	8	779	16,961	782	16,973
1877		19	194	5,311	203	5,311
1878	21	793	291	7,316	312	8,039
1879	106	3,099	785	12,210	891	15,909
1880	106	1,889	59	5,971	626	10,860
1881	134	5,496	1,732	29,016	1,886	27,436
1882	79	3,090	2,100	31,797	2,179	31,797
1883	200	18,023	1,601	22,532	1,791	19,551
1884	123	11,529	771	11,227	894	25,747
1885	313	8,635	1,913	16,798	1,396	25,343
1886	131	1,054	69	9,025	663	20,089
1887	77	5,054	1,667	29,395	1,681	31,649
1888	441	39,854	1,377	27,484	6,4818	38,338
1889	578	14,210	857	29,562	4,415	31,842
1890	87	5,759	1,694	16,973	1,481	21,842
1891	59	3,024	1,377	26,326	1,436	29,350
1892	175	2,684	1,799	34,218	1,906	36,831
1893	22	563	53	6,131	255	6,694
1894	84	6,189	59	2,025	143	8,265
1895	123	12,069	10	112	133	12,521
1896	11	720	5	2,490	56	3,120
1897	108	6,418		3	108	6,351
1898	131	3,975			131	3,975
1899	151	4,166			151	4,166
1900	11	325			11	325
1901	57	2,528		87	79	2,410
1902					31	1,720
1903					116	4,829
1904					172	1,682

no 200 tons from Ontario should have correctly be classed under the heading of mineral payments.

TABLE 3
MANGANESE
Imports. ORE OF MANGANESE

Fiscal Year	Pounds	Value	Fiscal Year	Pounds	Value
1874	3,989	8,258	1894	101,863	84,522
1875	39,513	1,794	1895	64,151	2,751
1876	41,967	1,753	1896	168,599	1,07
1877	29,655	2,933	1897	79,663	2,741
1878	65,914	3,022	1898	139,456	5,047
1879	52,211	3,182	1899	111,356	5,569
1880	97,192	3,199	1900	126,725	4,155
1881	22,987	3,743	1901	272,151	8,476
1882	76,097	3,339	1902 (Data not)	176,331	5,360
1883	94,116	3,696			

The manganese deposits of Canada were described in a previous annual report of the Mines Section. The present more extended article, bringing our information up to date, and the table of analyses have been compiled by Mr. Theo. Denis from all the available published information.

Although Canada has not so far taken a very prominent place among the manganese producing countries of the world, the reason for this is not due to the lack of deposits of the ores of this metal. By far the greater proportion of the world's production, some 90 per cent of it, is used for the manufacture of ferro-manganese and spiegeleisen. These two alloys of iron and manganese differ from each other only in the proportion of manganese which they contain, up to 30 per cent of manganese the admixture is called spiegeleisen, whereas, when containing greater proportions it is called ferro-manganese, the standard of the latter containing 80 per cent of manganese. These alloys are manufactured to contain all degrees of proportions of the two metals, some spiegeleisen holding as little as two per cent of manganese, whereas high grade ferro-manganese contains as much as 90 per cent. They are used exclusively by steel manufacturers for the production of certain steels of great toughness used for stamp mill dies and shoes, crushing rolls, car wheels, etc.

Use of
manganese

The extension of manganese production depends greatly on the development of steel manufacture, and as Canada is now making great strides in that direction, its deposits will probably assume before long a much greater importance than heretofore.

Besides the manufacture of steel, manganese ore has several other very important uses, the main one of which is its use as an oxidizing agent in the manufacture of certain chemicals, such as bromine, chlorine, manganates and permanganates; it is also one of the elements of the Leclanche cells; it is also used as a decolorizer of glass, as a coloring material in dyeing and in the manufacture of pottery and of paints, etc.

When the manganese ore is used as an oxidizing agent in the manufacture of chemicals, certain requirements of purity and composition are necessary which are not needed in the ore consumed for spiegeleisen and ferro-manganese, and it has therefore a value three to four times greater than that used for the latter purpose. Pyrolusite is the ore of manganese which has the greatest oxidizing power, and as the ore of some of the Canadian deposits contains a large proportion of this mineral, it is specially well adapted to that use. In Canada the ores represented comprise pyrolusite, manganite, psilomelane and wad or bog manganese ore. The principal deposits of the crystalline ores of manganese of the eastern provinces are referred by Dr. Gilpin to a

MANGANESE. horizon low down in the Carboniferous marine limestones, in most cases underlying the lowest beds of gypsum, yet that these ores have a wider distribution is shown by the fact that wad or bog manganese is found in superficial deposits connected with every geological formation known in Nova Scotia and New Brunswick; moreover, occurrences of pyrolusite have been noticed in the quartzites of lower Cambrian age and in granites, also in quartzites and slates of presumably Silurian age, and in Triassic trap rocks.

Origin of
manganese
deposits

Dr. Penrose* ascribes the origin of manganese deposits to secondary action and contends that the source of the manganese ores found in the Paleozoic and later sedimentary rocks, is to be traced to the underlying archæan rocks and various igneous rocks of all ages. The fact that the largest manganese deposits in the United States and Canada are in the neighbourhood of such rocks is in itself suggestive; but when it is found that large areas of bog manganese ore occupy basins in the decayed surface of the pre-Paleozoic rocks, and that the river pebbles in areas of these rocks are frequently encrusted with a black coating of oxide of manganese, other facts are encountered which at once suggest a possible pre-Paleozoic source for manganese deposits. Moreover, when it is observed that volcanic breccias are sometimes cemented by manganese, that segregated masses of oxide of manganese are sometimes found in lava, and that the manganese nodules dredged up from the sea bottom are in intimate association with volcanic debris, the possible source of manganese in igneous rocks claims attention. When these two classes of rocks, pre-Paleozoic and igneous, especially the former are examined in their more minute details, and it is found that of the minerals composing them, those containing manganese are among the most common, the probability of their being the source of manganese in the younger rocks becomes established. The different steps of the formation of manganese deposits being as follows:

1. The derivation of the manganese from the decay of the Archæan and other pre-Paleozoic rocks and from the products of igneous action.
2. The solution and transportation of the manganese in the form of soluble organic and inorganic salts of the metal.
3. The precipitation of the manganese as oxide or carbonate.
4. The conversion of the carbonate into oxide.
5. The subsequent decay of the rocks which were deposited with the ore and an accompanying change in the nature of the ore and sometimes in its physical condition. That is to say, that the stages

*Penrose, Geol. Surv. of Arkansas, 1890.

in the history of the manganese deposits involve first, a decay of the rocks in which manganese is originally present as a constituent; secondly, a series of chemical reactions leading to a redeposition, and thirdly, a decay of the rocks of those newly formed deposits. As Dr. Penrose remarks very appropriately, the various stages in the formation of manganese deposits are similar in many respects to those known usually to have gone on in the formation of certain iron ores, but differ in minor details. Both metals have their origin in the same rocks; they go into solution in the same manner; but in the mode of redeposition, though they sometimes resemble each other, they often differ considerably in the chemical changes which go on in the subsequent alteration of the oxides. Hence manganese is often associated with iron ore deposits and sometimes is comparatively free from such accompaniments.

MANGANESE.

In the following description of the known deposits of manganese ore in Canada, the localities will be taken, as far as possible, in their geographical order from east to west, and not in the order of their relative economic importance.

NOVA SCOTIA.

Nova Scotia.

This province possesses some of the most important deposits of manganese known in Canada, and as the iron and steel industries in that region are fast developing, these may, in the near future, become important sources of ore.

In Cape Breton, manganese ore is found in the western part of the county of that name. The most important belt of manganese bearing ore crops out in the district of the head waters of the Salmon river and Loch Lomond. The rocks are of lower Carboniferous age and are met with in a valley between the felsites of the Mira and East Bay hills.

From personal observations and from notes furnished by Mr. Hugh Fletcher, of the Geological Survey of Canada, Mr. Edwin Gilpin describes the general conditions of the occurrence of the ore as follows*:—"The felsites of the Mira hills form a series of bays along which are exposed Carboniferous limestones, conglomerates, shales and grits as they were accumulated, subject to the varying conditions of the winds and currents of the period under consideration. At some points the limestones rest on the felsites; at other localities, grits and shales intervene; elsewhere, the basal conglomerates are covered directly by the mill tone grit. The manganese ores were discovered

* Gilpin, Manganese in Nova Scotia. Trans. R.S.C., Vol. II.

MANGANESE. two years ago in one of these recesses where the felsites were succeeded by shales and grits, and finally by limestones, the latter apparently extending from point to point of the ancient bay. The ores at the western mine are found in irregular bedded layers in a soft arenaceous reddish-coloured shale, which is in some places calcareous and coated with films of manganese oxide. The layers vary in thickness up to eighteen inches, and are frequently connected by cross stringers of ore. The shales when weathered present the ore in small nodules, and the disintegration of the former by water probably indicates the source of the beds of gravel manganese ore found lying on them. The ore at the eastern mine occurs as a bed immediately underlying a layer of black manganiferous limestone, with red and greenish shales and coarse grit. The thickness of the ore and limestone varies from two to eight inches, the average thickness of the two being about eight inches. The ore also occurs in this vicinity as lenticular pockets and irregular nests in conglomerate, &c., and sometimes forms the cementing material. The latter mode of occurrence is similar to that shown by the red hematites (sometimes highly manganiferous) found at various points in the Lower Carboniferous conglomerates of the island near their junction with older strata."

This deposit was first opened in 1880 by the Hon. E. T. Mosely, of Sydney, and has been worked at two places about three-quarters of a mile apart, near the head of Loch Lomond, eight miles south of the village of Big Pond on East bay. At the most easterly of these, the workings are on a vein about seven inches thick, dipping at an angle of 25° "in red fine sandstone overlying reddish and greenish grit with grains of quartz about the size of wheat, and red marly limestone, red and greenish shale, conglomerate and other rocks, blotched with calc spar. It is in lenticular layers and also intimately mixed with the limestone, being probably of the same nature and origin as the hematite, and forming at times a cover for the pebbles of the conglomerate." *

At the western working the ore is found in the bedding planes of a bright red argillaceous shale overlaid by calcareous argillaceous shale and limestone, and underlaid by conglomerate. The rocks have here a dip of 32°. Although these steep dips show the result of a general disturbance of the region, yet the rocks have not been as minutely shattered as might be expected. The ore is found both crystalline and massive, a great proportion being pyrolusite; it is very free from iron, remarkably pure, and is well adapted to chemical manufacture. Besides the ore mined from the two workings above mentioned, a large quantity was obtained as drift nodules in the rock beds. These

* Fletcher, Hazle. Geol. Survey Rep., 82-83-84.

nodules have been washed out of the original decayed rock, and on the outside are earthy, but on breaking, the interior shows the bright black surface of fresh ore. MANGANESE.
Nova Scotia.

On Boulardarie island, in the vicinity of Big harbour, a deposit of wad or bog manganese occurs. The deposit is stated to be several feet thick and extensive, but there is a great lack of uniformity in the composition of the ore in different parts of the bed.

There are other occurrences of manganese ores in Cape Breton county. In a limestone quarry at Sabm road some pockets of manganese were encountered and mined in 1897. The rocks are of Lower Carboniferous age.

Some samples of wad received at the laboratory of the Geological Survey are said to have been obtained from a deposit situated at the head of Lewis bay.

Hants County.—On the south shore of Minas Basin there is a development of Lower Carboniferous limestone which from the Shubenacadie river extends westward for a distance of about forty miles, as far as the estuary of the Avon. This belt contains a limestone band some 300 feet thick in which are found the most important deposits of manganese of the region, the largest and best known of which are the Tenny Cape mines. It underlies the gypsiferous horizon, and Mr. Fletcher says of it that "next to the gypsum, the most interesting member of this formation is the red basal limestone, along which the manganese ores are found. It is of considerable thickness, concretionary, brecciated and associated in places with red conglomerate and grit." About fifteen miles south of Tenny Cape, near Windsor and at Douglas, the manganiferous limestone reappears. The occurrence of manganese, however, is not confined to the limestone, but it has also been noticed in the Devonian sandstone which is found below the Carboniferous marine limestone, and in places it occurs in large enough quantities to be worked. In this class of deposits which are not in the immediate vicinity of the limestone, the ore occurs in veins, joints or blotches, from one-quarter of an inch to five inches, in Devonian quartzites and shales. The important manganese veins of the district, however, are in the limestone above mentioned, lying at the base of the Carboniferous formation. Work on a comparatively large scale has been performed at Tenny Cape, Walton and Cheverie.

Tenny Cape Mines.—These are the most important workings of the region, and have been worked since 1861. The quantity of ore produced is not very great but has been described as the purest and

MANGANESE. most beautifully crystallized pyrolusite found in America. It has of course been chiefly used in the manufacture of chemicals, glass decolorizing, etc. Of the deposit Mr. H. Fletcher writes as follows: "The rock, a twisted, reddish, shaly or brecciated dolomite is sometimes separated by two to four inches of hard red clay from the Devonian sandstone or quartzite which forms the foot-wall or floor of the mine. The ore occurs in veins, strings, nodules and masses. One of the latter is said to have yielded ... thousand tons associated with calcite, selenite, barite, and limonite but in some places almost entirely free from foreign matter. It occupies the lines of jointing and bedding, breaks apart the fragments of the breccia and replaces the shale and limestone. The latter dips S. 20 E. at a variable angle, beneath a mass of gypsum: it has been worked for about 200 yards on the strike and the whole distance tested is probably less than 500 yards." The workings, as may be inferred from the wavy nature of the deposit, are scattered and irregular; they consist of open cuts, tunnels and shafts, the deepest of which a few years ago was 170 feet

The Parker Mine.—This is situated to the north of the Tenny Cape mine, about three-quarters of a mile from it. The deposit is in a much disturbed limestone, forming apparently an outlier of the Carboniferous basal beds among rocks of Devonian age. In 1881 some thirty tons of excellent pyrolusite were mined, but since then the work has been mostly of a prospecting and development nature.

The Churchill or Walton Mine.—This is on the west bank of Walton river immediately above the bridge, on the shore road some twelve miles north east of Cheverie. The deposit worked here is also in an outlier of red and gray limestone filling a hollow in red Devonian quartzite and shale. The ore is mainly pyrolusite with some manganite and is associated with calcite crystals and barite. Some large masses of very high grade ore have been mined from this deposit.

In this vicinity, the main development of the manganiferous limestone is encountered a short distance south of the outlier on which is the Walton mine, and these two bear to each other the same relation as the Parker mine deposit does to the Tenny Cape mines. The limestone crosses the Walton river south of Walton, and on both sides of the river extending some distances east and west, it has been subjected to a great deal of prospecting and preliminary work. One of the more important properties is the Stephens mine.

The Stephens Mine.—Of this deposit Mr. Willimott of the Survey, as the results of observations made in 1883, writes as follows: "This mine is situated near the village of Walton, in Hants county, and

consists of an excavation of about thirty feet in depth, in a reddish shaly limestone, striking E. and W. with a southerly dip. Pockets and irregular veins of manganite and pyrolusite can be traced along the strike for about 400 yards. MANGANESE.
NOVA SCOTIA.

Nothing beyond the preliminary prospecting work, scarcely sufficient to develop this promising mine, had been done at the time of my visit. Mr. Stephens informed me that about ten tons of fair grade ore had been taken out during the progress of their investigations. The limestone belt is perhaps continuous to Hibernia where a quantity of ore was found in reddish calcareous grit interstratified with concretionary limestone."

Sturgis Mine.—At this place which is about two miles west of Walton, manganese ore occurs in somewhat large quantities, as stringers, veins and films, or impregnations and stains in large masses of both the limestone and of the underlying dirty sandstone, in both of which shafts have been sunk and tunnels driven.

Tomlinson Mine.—This is situated west of the above mine and consists of openings made in the reddish and grey quartzite which underlies the limestone. These openings show masses of pyrolusite and hematite, sometimes mixed, sometimes separated.

Lantz Mine.—Several shallow pits have been opened in limestone, and from these workings some fine specimens of pyrolusite were obtained.

The Cheverie Mine.—This is situated near the village of Cheverie and the deposit here worked underlies the gypsum of the Cheverie quarries. The ore, a mixture of pyrolusite and manganite, occurs in a reddish and grey concretionary limestone, and is associated with white calcite in a network of small veins, from an eighth of an inch to three or four inches in thickness. Frequently the calcite associated with the manganese is in long crystals standing at right angles to the wall of the veins and forming a comb structure on both sides of the manganese.

According to observations made by Mr. H. Fletcher, at Cheverie, the ore is found near the top of the manganiferous band of limestone, at Walton it is found at the base of it near the contact with the underlying quartzite, and at Tenny Cape the best development of ore is met with at some thirty-seven feet from the bottom.

Minasville or Moose Brook Mines.—This mine is situated some four miles northeast of Tenny Cape. The occurrence of manganese ore here is not in the limestone, but in the underlying Devonian quartzites,

MANGANESE. where it is found in joints, veins and blotches, varying from a quarter of an inch to five inches in thickness. A certain quantity of ore has been shipped from these workings.

Nova Scotia.

The same geological conditions are observed at Bear brook and at a deposit east of Noel river, where some preliminary work yielded small quantities of pyrolusite.

In Hants county other occurrences of crystalline manganese ores, more or less important, have been noticed at Douglas, Rawdon, Goshen and other places. Bog manganese occurs near Goshen, south of Cheverie, at the head of Bass creek.

Colchester County, East Onslow. At this place is the most important deposit of manganese now known in Colchester county. The ore occurs in the joints and bedding planes of old Devonian quartzite. In some places the ore which consists mainly of pyrolusite, is a foot thick; some manganite and psilomelane are also encountered. Operations were begun on this deposit in 1886 or 1887 and have since been carried on intermittently. The principal workings consist of a shaft fifty five feet deep, and a large irregular cut.

Valley, Manganese Mines.—At Manganese Mines near Valley, a quantity of black oxide of manganese is found in irregular veins cutting a reddish, slaty rock, which underlies the Carboniferous limestone.

Furholm's Mill brook.—From the appearance of the occurrence of manganese at this place the deposit is a contact deposit between grey, rusty, concretionary, massive Carboniferous limestone and Devonian rocks. The ore occurs mostly in pockets in the limestone which also contains disseminated hematite, giving the rock a mottled weathering appearance.

Other occurrences of manganese ores have been noted on both shores of Minas basin in Colchester county. At Black Rock mine, near Clifton, at the mouth of the Stubenacudie river some work was done on a deposit in limestone. The ore which was of a ferruginous and magnesian nature was found in small quantities.

On the north shore of Minas basin at Lower Economy several barrels of fine crystalline pyrolusite were obtained in 1891. The occurrence is similar to those of East Onslow.

Besides the above occurrence, manganese is also found in large quantities associated with the important deposits of iron ores of Londonderry iron mines. The iron ore is found as veins of brown hematite

accompanied by ochre, ankerite and sideroplesite. In places secondary MANAGNESE changes have enriched the iron ore with manganese peroxide to the Nova Scotia extent of fourteen per cent of its total constituents.

Other localities in Nova Scotia where occurrences of manganese ores have been observed, are as follows:—

In Cumberland county, Minudie, some small quantities of soft fine-grained pyrolusite were obtained from the Lower Carboniferous limestone. At Amherst (Cumberland) some manganese ore occurs in the same formation.

At Springhill (Cumberland) and Parrshoro', wad is met with in superficial deposits.

At New Ross, in Lunenburg county, a few shipments of ore are said to have been made from the college grant. The ore appears to be a mixture of psilomelane and manganite occurring in veins sometimes three feet in thickness. Wad is reported to occur at La Have and Chester, Lunenburg.

In Picton county deposits of manganese ore are met with in connection with the iron ore deposits at Bridgeville and at Springhill, where boulders and concretions of psilomelane, manganite and wad are found.

In Antigonish county, near the head of the Ohio settlement, large pieces of pyrolusite were found in the drift on a hill, and in the same county occurrences of wad have been noticed near Afton; in Pomquet river: in Sutherland's brook and on a hill west of Piedmond station.

In King's county, near Wolfville, pyrolusite is found in small masses and stringers, in slates of Devonian age.

In Halifax county, at Musquodobit and Ship Harbour, pyrolusite occurs as veinlets in granite, and at Jeddore, wad is found in the superficial deposits.

NEW BRUNSWICK.

New
Brunswick.

The geological characters of the manganese deposits in New Brunswick resemble those of Nova Scotia. They are found in rocks of pre-Carboniferous age as well as in Lower Carboniferous measures, besides the superficial deposits of wad. The most important deposits, from an economic standpoint, are, however, those found in the Lower Carboniferous limestone.

Gloucester County.—In the vicinity of Tête à Gauche Falls, some eight miles from Bathurst, a deposit of pyrolusite occurs, in the red

MANGANESE,
New Brunswick.

slates of the district which are probably of Cambrian age. The ore is found in numerous small veins, some of which are said to be as wide as eight inches; and detached masses of it are often found in the superficial deposits in the neighbouring fields. This occurrence was the first to attract attention to the manganese ores of the province; it was worked a number of years ago, and a certain quantity of ore is said to have been shipped from this place. As a result of personal examination, Dr. Bailey is of opinion that the district is worthy of closer examination than it has yet received. Unfortunately, the conditions are not very favourable to easy prospecting, as the district is flat and deeply covered with clayey soil.*

King's County.—In this county are the deposits of Markhamville, which are the most important ones of the province. These deposits were examined by Dr. Penrose in 1890, and as a result of his visit he describes them as follows:

† "The Markhamville mine is situated at the village of Markhamville near the head of Hammond river in Kings county about forty miles north east of St. John, about fifteen miles north of the shore of the Bay of Fundy, and about eight miles south of Sussex on the Inter-colonial railway. The existence of manganese was noted at the head waters of the Hammond river many years ago by Mr. Geo. F. Matthew, of the Geological Survey of Canada, but the property was first opened about 1864 under the management of Major A. Markham. Major Markham was the first to attempt to develop in a systematic manner the manganese deposits of this province, and it is due to his energy and perseverance that the ores have been introduced into the market.

"The ore occurs either as crystalline pyrolusite and manganite, or in a compact, massive, nodular or bedded form, sometimes containing psilomelane.

"The ore-bearing limestone is generally of a gray colour, but at times is pink or buff, and is associated with shaly strata. It contains veins of crystalline calcite in which masses of pyrolusite are frequently found, but the principal ore deposits are lenticular bodies interstratified with the limestone. These ores occur as irregular pockets or as flat layers more or less continuous for considerable distances, and becoming thin and thick at intervals. In some places such deposits widen out into pockets from which several hundred tons of ore have been taken and in one opening 3,000 tons are said to have been mined.

* Annual Report of the Geol. Surv. of Can., Vol. X, N. S., 1897, Part M.
† The Manganese deposits of U. S. and Canada, by Dr. Penrose. Geol. Survey of America, vol. I, 1890.

Though in places the pockets do not always adhere strictly to the bedding of the rock, yet in a general way they follow it. Sometimes veins and pockets cut directly across the bedding, but these are generally smaller than the others and are probably due to a secondary chemical action by which they have been derived from the bedded ores.

MANGANESE.
New
Brunswick.

“The surface of the limestone has often been decomposed, and a red residual clay, frequently mixed with surface gravel, has collected in considerable quantities. The ore that was originally in the part of the limestone which has decayed, is now found buried in the clay; and therefore deposits of ore bearing clay or gravel, overlying the partly decomposed surface of the limestone, are of frequent occurrence. Such deposits are rarely more than from eight to twenty feet in thickness, but the ore in them is cheaply worked and they have supplied a large part of the output of the Markhamville mine. Frequently the decomposition of the limestone has spread downward more rapidly along the outcrop of a body of ore than elsewhere, causing somewhat abrupt hollows filled with residual clay and manganese ore and containing in the bottom the outcrop of the ore in situ in the rock.

“Not only has decomposition taken place on the surface but it has also gone on to a considerable extent underground frequently causing subterranean cavities and passages. When these have intersected bodies of manganese the floors are covered with loose fragments of ore, brought there in the same way as that in the residual clay on the surface. Kidney-shaped masses of glossy, black limonite are frequently found with the cave deposits, and these also have doubtless come from the limestone.

“Though a large amount of manganese has been taken from the surface clay beds and the caves, yet the deposits of ore in the limestone have also been extensively worked, and in many places the rock is honeycombed with a network of shafts and drifts, following the erratic courses of the ore bodies in all their intricacies.

“The thickness of the limestone varies considerably: in one of the pits a depth of twelve feet was found, and a diamond drill boring in another part of the property showed a thickness of fifty-five feet. Probably a greater thickness will be found elsewhere. The bed is much disturbed and is folded into small anticlines and synclines, but at Markhamville it has a general dip to the northwest and a strike of northeast and southwest. In many places it contains fossils, and sometimes the carbonate of lime of these has been partly replaced by manganese, which has subsequently been oxidized and now exists as a black, more or less calcareous mass.

MANGANESE.
New
Brunswick.

"The Hammond river rises near Markhamville and flows south west parallel to the coast of the Bay of Fundy, until it finally turns south and empties into the bay about eight miles south east of St. John. In the region of Markhamville, and for some miles down the river, the Lower Carboniferous limestone occupies the centre of the valley; but it is only locally that manganese occurs in it, and only at the Markhamville mine that it has yet been found in large quantities.

"The limestone area is bordered on the south by a range of hills which forms the southern barrier of the Hammond river valley. According to information kindly furnished by Mr. G. E. Matthew, of the Geological Survey of Canada, these hills are composed largely of the underlying pre-Cambrian rocks, and the Carboniferous rocks dip away from them. To the north of the river the limestone is cut off in many places by an abrupt escarpment of Carboniferous conglomerate, which according to the same authority, probably belongs above the manganese-bearing limestone.

"The ore from this mine is mostly used for chemical purposes. It is prepared for market by crushing, washing and sizing with screens. Certain quantities of the lower grades, however, are shipped without previous preparation, under the name of "furnace ore" and are used in the manufacture of spiegelisen and ferro manganese."

The importance of the Markhamville mines may be realized by the fact that between 1868 and 1894, the total exports of manganese ore from New Brunswick amounted to over 23,000 tons representing a value of nearly \$410,000, almost the whole of which, was derived from the Markhamville deposits. (Dr. Bailey, Mineral Resources of New Brunswick.)

King's County, The Glebe mine.—The deposit of the Glebe Mine is situated some three miles N.N.E. of the Markhamville vein, and seven miles from Sussex station on the Intercolonial Railway.

Dr. Penrose in his bulletin on the manganese ores of America describes this occurrence as follows:—"The ore is found in a limestone resembling that at Markhamville, though it is much less disturbed than at that place and dips gently to the west. The manganese ore occurs in the limestone in nodules and thin layers, frequently associated with calcite and following the general direction of the stratification. Several shafts and tunnels have been made, the deepest shaft being 85 feet."

King's County, Jordan Mountain.—This deposit is situated on the south eastern side of Jordan mountain, about seven miles from Sussex

station on the Intercolonial Railway. According to Dr. Bailey, the geological relations here are similar to those of Markhamville; the ore is found in strata of Lower Carboniferous age near their contact with older metamorphic rocks. But instead of occurring as in the last named locality, in limestone, it is found in connection with shales and shaly conglomerate, the brecciated character of which is in contrast with the rocks at Markhamville. Work was begun on this deposit in 1882 and some good ore is said to have been extracted from an open cut. The ore is a mixture of pyrolusite and manganite occurring in lenticular interbedded masses. There are also small veins and stringers of manganese oxide penetrating the surrounding rocks.

MANGANESE.
New
Brunswick.

King's County, Hillsdale.—Some fine surface indications of manganese ore are said to have been observed at Hillsdale about five miles southeast from Elgin corner. No particulars of this deposit are available.

St. John County, Quaco Head mine.—This is situated on the north shore of the Bay of Fundy on a promontory which forms the southern boundary of Quaco Harbour. The mine is about one mile south of the village of St. Martins. The following description of the deposit is taken from the bulletin by Dr. Penrose who examined it in 1890 when it was being worked by the Brunswick Manganese Company. "The manganese is sometimes crystalline representing pyrolusite and possibly also manganite, while at other times it is hard and massive, possibly representing psilomelane, and still again porous honeycombed form. These ores are found in Lower Carboniferous shales and limestones, associated with a large conglomerate bed.

"The rocks are greatly disturbed and have been much shattered and broken by igneous intrusions. They now stand at steep angles sometimes almost vertically, exposing in different parts of the headland areas of limestone, shale, and coarse conglomerate. Masses of igneous material protrude into these beds at different points and on either side of the headland are beds of Triassic sandstone and fine conglomerate lying unconformably on the upturned edges of the older rocks.

"The manganese occurs as nodules and irregular discontinuous veins, in both the shale and the limestone, though the larger quantities are in the former. The nodules vary from a fraction of an inch to several inches in diameter, and the thickness of the veins is equally variable. The disturbed character of the rocks renders it somewhat difficult to determine the thickness of the main ore bearing bed but it is probably not over thirty feet though smaller quantities of manganese are found in the rocks on either side. The ore is scattered through this thickness in very variable quantities.

MANGANESE
New
Brunswick

"The igneous rock is a hard light gray, close grained material of a texture somewhat like trap. The limestone is like that of Markhamville, though it is much reddened at the contact with the igneous rock. The conglomerate bed is composed of coarse pebbles of metamorphic rocks. It dips steeply to the south and forms a bold bluff on which the light house of Quaco Head is situated. The sandstones and conglomerates at each end of the section are of a brick red colour and vary from coarse sandstone to a fine conglomerate, with pebbles from a quarter of an inch to one inch in diameter, both sand and pebbles being composed of white quartz stained by a ferruginous cement. Sometimes these beds contain small irregular seams or nodules of manganese ore, which, however, are in very limited quantity, and have doubtless been derived during the deposition of the beds, from the erosion of the Lower Carboniferous rocks.

"The ore-bearing rocks can be traced back on the promontory at intervals for almost a mile, to a place where an opening has been made on the farm of Mr. Mohaskey. On the north side of the head, small scattered nodules of manganese ore are found in the gravel drift that lines that part of Quaco Harbour, and extends inland over the Lower Carboniferous rocks. They have doubtless been derived from the latter rocks during deposition of the gravel, in the same way that the red sandstone just mentioned obtained its manganese contents at an earlier date."

Subsequent to Dr. Penrose's visit the Brunswick Manganese Company erected a well equipped mill, consisting mainly of a crusher, rolls, screens, two pigs, etc. The mine is exceptionally well situated for shipping by water. Operations have, however, been discontinued for several years.

Albert County, Shepody mines—Shepody mountain is one of the highest eminences in southern New Brunswick. The lower part of this mountain is composed of chloritic hydro-mica schists, and the upper part consist of Lower Carboniferous strata; at the contact between these two sets of rocks are found the manganese occurrences.

Dr. Ellis in his report for 1885* writes as follows:—"Shepody mountain, 1,050 feet high, is a rugged peak which forms a prominent landmark for many miles in all directions, and was one of the signal stations for the Admiralty survey of the Bay of Fundy. It is composed largely of red conglomerates, which are well exposed on the east flank in Robertson's brook and its branches. A deposit of reddish

* Geol. Surv. Can., Report of Progress, 1885.

impure limestone has been opened up at this place for a marble quarry, but the rock was found to be too much shattered to be of great value. The limestone contains a small quantity of manganese. The rocks of the mountain rest upon a small outlier of the talc-chloritic schists which show on the road to the north, leading to Curryville, and are flanked on the east by the gray sandstones of the millstone grit. On the north-west side a large deposit of manganese was worked for some years, a tunnel being driven into the mountain along the contact with the underlying schist for nearly 1,000 feet. The ore, which consisted of pyrolusite and psilomelane occurred at the base of the conglomerate in irregular pockets.

The Shepody mountain mines were first opened in 1860, and it is said that 500 tons of ore was extracted. This was a compact black oxide, less crystallized than the ores of Madburyville, but of high grade. It was found both in veins and in interbedded masses.

Albert County, Elgin, Garland Mountain.—On the north-east side of the mountain some exploratory work revealed occurrences of pyrolusite and psilomelane in a very broken and decomposed granite of post-Auriferian age.

In the same county, on the east side of Salisbury bay, a small deposit of manganese occurs near a contact of Lower Carboniferous and Triassic sandstones. This deposit was worked many years ago but shortly abandoned.

Albert County, Dawson Settlement.—At this place occurs a very important deposit of wad or bog manganese. Dr. Barley, who visited the deposit in 1899, gives the following description, which is the latest and fullest available:

"This very remarkable deposit is located about five miles and a half from the town of Hillsborough, on the slope of a hill inclining north-easterly at a low angle towards a small brook, flowing thence to the Petiteodac river, and whose opposite slope is occupied by the settlement above named. The upper part of the first ridge is wooded, but between the edge of the latter and the brook the ground is cleared and upon removal of a thin coating of vegetable matter, usually not more than two inches in depth, is found to be everywhere covered with a very fine black powdery deposit consisting essentially of manganese oxide.

"The property, as leased, embraces an area of about 150 acres, and upon about eighteen or twenty acres, or as far as searched for, there has been found the deposit varying in depth from a few inches to

MANGANESE.
New
Brunswick.

thirty feet. In a survey recently made by a Crown land surveyor, seventy three borings were made, in squares of 100 feet, over a space of seventeen acres, showing an average depth of six feet seven and three-quarter inches, equal to 1,900 pounds to the cubic yard. There is accordingly already in sight and available for use:—

	Tons.
In situ on hill-side, 17 acres	173,176
In drying house and sheds	400
Total	173,576

According to the statements of the manager of the property Mr. R. P. Hoyt, to whom I am indebted for assistance and valuable information, the iron rods used in the above borings, in many of the deepest places, failed to go down over twenty-five or thirty feet, and then struck what was apparently hard manganese ore, so that the above results indicate the minimum quantity. These ores are, in comparison with those of Markhamville, low grade ores and would be of little or no value for the uses to which the latter are chiefly put, nor in their natural condition would they have commercial value of any kind. It is, however, proposed to subject them to a briquetting process whereby the pulverulent and absorbent mass shall be rendered solid, non-absorbent, and capable of easy handling, in which condition it may be advantageously used in the manufacture of spiegeleisen and ferro-manganese. For this purpose an extensive plant embracing drying furnaces, compressors, briquetting machines, etc., has been erected close by the manganese deposits, and also near to the track of a branch railway one mile and a half in length, built by the company, and connecting with the Harvey and Salisbury Railway at a point eleven miles from Salisbury, whence, over the Intercolonial Railway the product may be readily shipped to all Canadian and United States points. The shipping point by sea is five miles and a half by rail from the mine to Hillsborough, with direct landing at wharf for vessels of 800 to 1,000 tons capacity. This company, "The Mineral Products Company of New York," sent the briquettes to Bridgeville, Nova Scotia, where the smelting plant of "The Picture Charcoal Iron Company" had been secured for the manufacture of ferro-manganese. For some unknown reason, after a period of apparently successful operations this company have discontinued work.

Besides this deposit of wad, other deposits have been noticed at different places, among which are Queensbury, York county; north branch of S. W. Miramichi; in gravelly bank near government house

at Fredericton; near Harvey, Albert county; Bull Moose Hill, Kings county; Moore's Mills, Charlotte county and other places.

MANGANESE.
New
Brunswick.

As to the probable origin and mode of formation of the deposits of wad or bog manganese, Dr. Chalmers* assigns it to the action of springs.

In the case of the Dawson Settlement deposit, the bed of wad lies in a valley at the northern base of a hill, and springs are trickling down the hillside; doubtless the process of formation of bog manganese is still going on. Dr. Bailey on the same subject writes as follows:

"An interesting question in connection with these deposits is that of their probable origin. Upon this point the locality throws very little light, there being absolutely no exposures of rocks anywhere in the vicinity or any visible source from which the manganese may have come. The nearest rocks are indeed those of the millstone grit, though these are doubtless underlain as at Hillsborough and about the Albert mines by Lower Carboniferous rocks, including limestones. None of these however, are markedly manganimiferous. It is also a little singular that the deposit should have such a decided slope instead of being, as usual with bog ores, nearly horizontal. Finally, the abruptness with which the deposits end along the line of the brook referred to above, towards which it inclines while no such material is to be found on the opposite slope, is also remarkable, and seems to suggest that the ores are the result of deposition from springs originating on the one slope but wanting on the other, while the brook has carried off the excess of the solvent water. In support of this view it may be observed that the hillside on which the ore beds rest, is remarkable for the number of springs which issue from its surface, in the waters of which both iron and manganese may be readily detected.

QUEBEC

Quebec

In this province the only occurrences of crystalline manganese ore are those of the Magdalen islands in the Gulf of St. Lawrence. From an examination of these islands Mr. Jas. Richardson in the report of the Geological Survey for 1879-80 writes as follows: "Immediately under Demoiselle hill, on Anherst island, numerous blocks charged with peroxide of manganese or pyrolusite, occur among the debris of the fallen cliffs. They are in pieces varying from one pound to ten or fifteen pounds in weight. There can be little doubt that they are derived from a deposit more or less regular in the hill side, but which

MANGANESE is now completely concealed by the fallen debris. At a place bearing
Quebec nearly due west from Cap aux Meules, at the distance of about a mile, and close to the English Mission church, similar pieces to those above described are very frequently picked up." These deposits have lately attracted some attention, and in 1903 were purchased by a syndicate which intend working them.

Wad or bog manganese has been observed at a great number of points in this province but the quality is poor as a rule, and of small commercial value. Of these deposits one of the most considerable is in the township of Stanstead, lot 9, range X: This deposit is stated to cover an area of about twenty acres with a maximum thickness of about twelve inches. Some of this ore after undergoing a washing to free it from the sand, gave 37 per cent of peroxide.

Another deposit of several hundred square yards in extent with a maximum thickness of six inches was observed in the township of Bolton, lot 20, range XII.

Mr. A. P. Low, mentions the occurrence of wad on the St. Louis road some four miles and a half from Quebec. The deposit here is about sixty yards by five with a maximum observed thickness of twelve inches.


The following other occurrences were compiled from the reports of the Geological Survey. The majority are of limited extent.

On the road from Lambton to St. Francis, Beauce county, near the eastern boundary of the township of Tring; on the west side of the Chaudiere river, opposite the mouth of the Emain river; in the seignory of the Mesy; in the seignory of Ste. Anne de la Poetiere in rear of the church; in Cleveland township, county of Richmond, on lot 16, range XIII; in St. Sylvester lot 9, range St. Charles; in Gaspé seignory, half a mile west of St. Apollinaire church.

Ungava
Territory

UNGAVA TERRITORY.

In his report on the east coast of Hudson bay, Dr. Bell mentions the occurrence of very important deposits of spathic iron ore in the Nastapoca chain of islands. These ores are in places very rich in carbonate of manganese: an average specimen from Flint island yielded 25.44 per cent metallic iron and over 24 per cent of carbonate of manganese. These deposits are very accessible and may some day be worked profitably. The high contents of manganese in these ores would render them valuable in the manufacture of spiegeleisen.



ANALYSES OF CANADIAN

Locality	W.	Mo.	Al.	Fe.	Ca.	Mg.	Si.	Sp. of Fe.	Ca.	Al.	Other constituents
NOVA SCOTIA.											
<i>Cape Breton County.</i>											
Grand Mira, Head of Lewis Bay	41.99										
Mira Hills, two miles east of Loch Lomond	41.00										
"	8.96										
Morrison Mine, near Loch Lomond	6.84										
Bonaventure Island	41.43										
"	25.17										
"	41.64										
<i>Colchester County.</i>											
Londonderry	67.49	10.67	1.00								1.14
<i>Charlottetown County.</i>											
Amlurst	97.01										
<i>Halifax County.</i>											
	36.00										
<i>Halifax County.</i>											
Tenny Cape	85.51										31
"	1.33										
"	21.71										
"	8.64										
Cheyne	90.15	61.97									Trace
"	86.81										
"	47.79										
Douglas	84.02	51.41									
<i>Prince County.</i>											
Springville	41.49	9.46									
"	92.31	61.89									1.78
NEW BRUNSWICK.											
<i>Abbotsford County.</i>											
Hillsborough	47.09										
Dawson Settlement, 12 miles Edge of Hastings	71.07	2.71									6
<i>Essex County.</i>											
Gowland Mountain	50.21										
<i>Kent County.</i>											
Markhamville	8.76										
"	91.25										
"	94.62										
"	71.71										
Jordan Mountain	95	4.17									1.40
"	92										
"	77.97										0.01
<i>St. John.</i>											
Quaco Hill	71.04										
"	11.00										0.01
QUEBEC.											
Ambroist Island, Magdalen Islands	45.61										
One of the Magdalen Islands		61.62	71.70								

* Contains also small proportions Ni, Co and Cu. † From peroxide and ferric. ‡ Oxides of iron and aluminum. § MANGANESE - Page 12, 24.

ADIAN ORES OF MANGANESE.

Locality	Latitude	Longitude	Altitude	Weight of sample	Weight of residue	Weight of loss	Analyst	Reference	
1	45° 30' N.	70° 00' W.	1000	100.00	12.20	0.00	Kenrick & Adams	Geol. Surv. of Can., 1886, pt. T.	
2	45° 30' N.	70° 00' W.	1000	100.00	2.91	0.00	"	" 1879-80, " H.	
3	45° 30' N.	70° 00' W.	1000	100.00	10.00	0.00	Gilpin	" 1889-81-82, " H.	
4	45° 30' N.	70° 00' W.	1000	100.00	57.76	0.00	Kenrick & Adams	Trans. Roy. Soc., Can., Vol. II	
5	45° 30' N.	70° 00' W.	1000	100.00	1.98	0.00	Kenrick & Adams	Geol. Surv. of Canada.	
6	45° 30' N.	70° 00' W.	1000	100.00	9.61	0.00	Kenrick & Adams	"	
7	45° 30' N.	70° 00' W.	1000	100.00	8.04	0.00	Kenrick & Adams	Mineral Res., U.S., 1885.	
8	45° 30' N.	70° 00' W.	1000	100.00	2.65	0.00	Kenrick & Adams	Erleb. and Dublin Phil. Mag., 1866.	
9	45° 30' N.	70° 00' W.	1000	100.00	1.11	0.00	Kenrick & Adams	Nova Scotia Inst. of Sc., 1865.	
10	45° 30' N.	70° 00' W.	1000	100.00	1.728	0.00	Kenrick & Adams	Trans. Roy. Soc., Can., Vol. II.	
11	45° 30' N.	70° 00' W.	1000	100.00	12.030	0.00	Gilpin	Trans. Roy. Soc., Can., Vol. II.	
12	45° 30' N.	70° 00' W.	1000	100.00	1.460	0.00	Gilpin	Nova Scotia Inst. of Sc., 1865.	
13	45° 30' N.	70° 00' W.	1000	100.00	2.731	0.00	Gilpin	Trans. Roy. Soc., Can., Vol. II.	
14	45° 30' N.	70° 00' W.	1000	100.00	71.85	0.00	Kenrick & Adams	Trans. Roy. Soc., Can., Vol. II.	
15	45° 30' N.	70° 00' W.	1000	100.00	33.78	0.00	Kenrick & Adams	Trans. Roy. Soc., Can., Vol. II.	
16	45° 30' N.	70° 00' W.	1000	100.00	55	0.00	Kenrick & Adams	Trans. Roy. Soc., Can., Vol. II.	
17	45° 30' N.	70° 00' W.	1000	100.00	2.50	0.00	Kenrick & Adams	Trans. Roy. Soc., Can., Vol. II.	
18	45° 30' N.	70° 00' W.	1000	100.00	2.86	0.00	Kenrick & Adams	Trans. Roy. Soc., Can., Vol. II.	
19	45° 30' N.	70° 00' W.	1000	100.00	9.70	0.00	Kenrick & Adams	Trans. Roy. Soc., Can., Vol. II.	
20	45° 30' N.	70° 00' W.	1000	100.00	0.23	0.00	Kenrick & Adams	Trans. Roy. Soc., Can., Vol. II.	
21	45° 30' N.	70° 00' W.	1000	100.00	8.37	0.00	Kenrick & Adams	Trans. Roy. Soc., Can., Vol. II.	
22	45° 30' N.	70° 00' W.	1000	100.00	10	0.00	Kenrick & Adams	Trans. Roy. Soc., Can., Vol. II.	
23	45° 30' N.	70° 00' W.	1000	100.00	31.63	80	1.10	Kenrick & Adams	Trans. Roy. Soc., Can., Vol. II.

§ Baryte and silica. Also trace of Cu. and Co.

MAS

Quel

Ung
Terr

ONTARIO.

MANGANESE.

There are very few occurrences of ores of manganese in this province, and none of these has been worked. One of the first discoveries is that of Bachewanung bay on Lake Superior, which is described as follows in the "Geology of Canada 1863."

"At Bachewanung bay, near the southwest end of the Upper Canada Mining Company's location, and not far from the shore is a large vein of manganese ore, running north and south, and from fifty to sixty feet wide. It is described as presenting the aspect of a succession of small knobs, in which, mixed with a reddish trappan rock, are numerous strings of the ore, associated with quartz and calc spar, and occasionally with octahedral crystals of fluor. The ore, which is massive with small geodes of crystals, is described by Prof. Hudley as manganite or hydrous sesquioxide of manganese, which for manufacturing purposes is inferior to the peroxide. A specimen was found by assay to be equal to sixty per cent of peroxide of manganese."

In the Rainy lake district manganese has been discovered associated with the iron ore deposits to the north of Gunflint lake: an occurrence is described as follows by Mr. Connee in the second report of the Ontario Bureau of Mines: "This range (iron ore) is near Sand lake, four miles from the Port Arthur, Duluth, and Western railway. The deposit has been found to be a large one. A pit has been sunk about 15 feet, and as far as the pit has shown up the vein, it seems to be very much decomposed. The ore assayed 65½ per cent. of iron and carried also a good percentage of manganese. The manganese appears to be dispersed among the iron, but it also occurs in pockets: they have taken out small quantities of manganese almost pure."

Wad or bog manganese occurs at several places in Ontario but has not attracted any attention, so that very little is known about these occurrences. An extensive deposit is said to occur in Hastings county, Madoc tp., lot 4, range V.

NORTH-WEST TERRITORIES.

North-west
Territories.

In Assiniboia a deposit of manganese ore is reported to occur on the north bank of the north fork of Willow creek. Tp. 5, R. 1, west of 4th Mer.

Mr. Pearce gives the following description of the deposit: "The manganese is found in pockets in a honeycombed formation four or five feet thick, composed of clay and sand with no sulphur or lime. Taking the deposit as a whole, manganese is estimated to run 5 per cent."

1903

MANGANESE. BRITISH COLUMBIA.

British
Columbia.

The only record available of discovery of ores of manganese in British Columbia, is that of important occurrences of bog manganese in Nicola valley, but only a comparatively small part of the province has been thoroughly prospected, and it is very probable that rich deposits exist which have not yet been found.

References.

REFERENCES.

- BAILEY, L. W. The Mineral Resources of New Brunswick, Report Geol. Survey, Vol. X, 1897, Part M.
- BELL, DR. R. — Report on an Exploration of the East Coast of Hudson Bay. Report of Progress, Geol. Survey of Canada, 1877-78.
- BRUMMEL, H. P. H. Report Geological Survey of Canada, 1890, Part S, Vol. V.
- HAWSON, SIR J. W. — Acadian Geology.
- ELLS, DR. R. W. The Mineral Resources of the Province of Quebec. Report Geol. Survey of Canada, Vol. IV, Part K.
- FLETCHER, H. — Report on the Geology of Picton and Colchester counties. Geol. Survey of Canada, Vol. V, Part P.
- FLETCHER, H. — Summaries of operations of Geol. Survey for years 1893 and 1901-02.
- GILPIN, ETC. — Mines and Minerals of Nova Scotia, 1880.
- GILPIN, ETC. — Notes on Manganese ores of Nova Scotia, Trans. Roy. Soc. Canada, Vol. II.
- GILPIN, ETC. — The Minerals of Nova Scotia, 1900.
- HOW, PROF. — Trans. Nova Scotia Inst. of Sc. Vol. I.
- HOW, PROF. — How's Mineralogy of Nova Scotia.
- JENNISON, W. F. Manganese deposits of Nova Scotia. Trans. Can. Mining Inst., 1898.
- MINERAL STATISTICS AND MINES. — Geological Survey of Canada. Annual Report, Mines Section, 1890-1891-1897-1898.
- PENROSE, DR. — Manganese Deposits of America, Geol. Survey of Arkansas, Vol. I, 1890.
- WILLMOTT, C. W. — Notes on Mines and Minerals. Geol. Survey of Canada, 1882-83-84, Part L.

TABLE OF CONTENTS.

	Page
Letter of transmittal	3
Statistics, table of jurisdiction of manganese	5
" " " exports of manganese	6
" " " imports of manganese	6
Manganese deposits of Canada	7-26
Uses of manganese	7
Origin of deposits	8
Deposits of Nova Scotia	9
Deposits of New Brunswick	15
Occurrences in Province of Quebec	23
Occurrences in Ungava	24
Table of analyses of Canadian manganese ores	Oppp. 24
Occurrences in Ontario	25
" North West Territories	25
" British Columbia	26
References, List of	26

This book is for use in
Library Building ONLY
and as a
"Reserved" Loan

