

193

*Flora Americæ Septentrionalis;*

OR, A

SYSTEMATIC ARRANGEMENT

QK

AND

581.970

110

DESCRIPTION

P986

P986

OF

1

THE PLANTS

OF

NORTH AMERICA.

CONTAINING, BESIDES WHAT HAVE BEEN DESCRIBED BY  
PRECEDING AUTHORS, MANY NEW AND RARE  
SPECIES, COLLECTED DURING TWELVE  
YEARS TRAVELS AND RESIDENCE  
IN THAT COUNTRY,

BY

FREDERICK PURSH.

---

IN TWO VOLUMES.

WITH TWENTY-FOUR ENGRAVINGS.

---

VOL. I.

---



LONDON:

PRINTED FOR WHITE, COCHRANE, AND CO.,  
FLEET STREET.

---

1814.

Bibliothèque,  
Le Séminaire de Québec  
3, rue de l'Université,  
Québec 4, QU

AYLMER

VICE-PRI

SIR,

I  
small tributary  
be permitted  
with your  
meet, with  
have, in a  
for their ex-  
and assistance  
my own fee  
inadequate  
portant an i

PRINTED BY RICHARD AND ARTHUR TAYLOR,  
SHOE-LANE, LONDON.



Should  
any degree i

TO

AYLMER BOURKE LAMBERT, Esq.

F.R.S. F.A.S.

VICE-PRESIDENT OF THE LINNEAN SOCIETY.

SIR,

*I AM conscious of offering but a small tribute of gratitude, when I beg to be permitted to inscribe the following pages with your name. Should they happily meet with approbation, the Public will have, in a great measure, to thank you for their existence; as, without the aid and assistance you so liberally afforded me, my own feeble exertions would have been inadequate to the performance of so important an undertaking.*

*Should the perusal of this work in any degree add to the pleasure you feel in*

*the cultivation of your favourite science, and in encouraging the efforts of others for its advancement, the utmost of the wishes are attained of,*

*Sir,*

*Your obedient and much devoted  
humble Servant,*

**FREDERICK PURSH.**

AMONG the natural history Continent, not than the vegeta forests produce timber trees ; tal flowering sh grounds ; and exceedingly hand possessing val those of other c able of being ad greater part of which circumst priety, the first

A country so favourite pursui life, a strong de natural soil and and to make s throw in my wa in the year 179 I had received i

e science,  
of others  
st of the

ed

PURSH.

## PREFACE.

AMONG the numerous useful and interesting objects of natural history discovered on the vast extent of the New Continent, none claim our attention in a higher degree than the vegetable productions of North America. Her forests produce an endless variety of useful and stately timber trees; her woods and hedges the most ornamental flowering shrubs, so much admired in our pleasure grounds; and her fields and meadows a number of exceedingly handsome and singular flowers (many of them possessing valuable medicinal virtues), different from those of other countries. All these are more or less capable of being adapted to an European climate, and the greater part of easy cultivation and quick growth; which circumstances have given them, with much propriety, the first rank in ornamental gardening.

A country so highly abundant in all the objects of my favourite pursuits, excited in me, at an early period of life, a strong desire to visit it, and to observe in their natural soil and climate the plants which I then knew; and to make such discoveries as circumstances might throw in my way. This plan I carried into execution in the year 1799; when I left Dresden, the place where I had received my education, and embarked for Balti-

more in Maryland, with a determination not to return to Europe until I should have examined that country to the utmost extent of my means and abilities. In 1811, after an absence of nearly twelve years, I returned to Europe, with an ample stock of materials towards a *FLORA OF NORTH AMERICA*, an attempt at which I now venture to lay before the public, with the flattering hope, that a generous allowance will be made for its unavoidable imperfections, when the extent of the undertaking is considered; and that it will be accepted as it really is intended, as only the groundwork of some future more perfect work upon the subject.

As it may be interesting to some readers to have a sketch of the particulars of my stay in America, and to be informed of the means by which I became possessed of so general a collection of its plants, I shall give a few outlines thereof, previous to the necessary remarks respecting the execution of the work itself.

My first object, after my arrival in America, was to form an acquaintance with all those interested in the study of Botany. Among these I had the pleasure to account one of the earliest, and ever after the most valuable, the Rev. Dr. Mühlenberg of Lancaster, in Pennsylvania, a gentleman whose industry and zeal for the science can only be surpassed by the accuracy and acuteness of his observations; I only regret, that his extensive and interesting materials towards a Flora of the United States, in which he has been engaged for a number of years, should not before this have been communicated to the public.

I next visited the old established gardens of Mr. Marshall, author of a small "Treatise on the Forest-trees

of North A  
advanced i  
ducted me  
ing trees  
then new t  
and applic  
vigour of i  
establishm  
happened  
kept up by  
cline, only  
mento of  
spectable b

Near P  
Messrs. Jo  
an old esta  
the late D  
Bartrams.  
of the Del  
and probab  
John Bart  
tion, and  
Mr. Willia  
vels throug  
intelligent,  
and from  
about the  
ing the ha  
trees. It  
call to mi  
man's com  
bourhood.

not to return  
that country to  
es. In 1811,  
I returned to  
ials towards a  
ot at which I  
the flattering  
ude for its un-  
of the under-  
accepted as it  
of some future  
ers to have a  
merica, and  
became poss-  
lants, I shall  
he necessary  
rk itself.  
merica, was  
rested in the  
e pleasure to  
the most va-  
ster, in Pen-  
zeal for the  
y and acute-  
t his exten-  
lora of the  
for a num-  
en commu-

f Mr. Mar-  
Forest-trees

of North America." This gentleman, though then far advanced in age and deprived of his eye-sight, conducted me personally through his collection of interesting trees and shrubs, pointing out many which were then new to me, which strongly proved his attachment and application to the science in former years, when his vigour of mind and eye-sight were in full power. This establishment, since the death of Mr. Marshall, (which happened a few years ago,) has been, in some respects, kept up by the family, but is now very much on the decline, only a few old established trees being left as a memento of what formerly deserved the name of a respectable botanic garden.

Near Philadelphia I found the botanic garden of Messrs. John and William Bartram. This is likewise an old establishment, founded under the patronage of the late Dr. Fothergill, by the father of the now living Bartrams. This place, delightfully situated on the banks of the Delaware, is kept up by the present proprietors, and probably will increase under the care of the son of John Bartram, a young gentleman of classical education, and highly attached to the study of botany. In Mr. William Bartram, the well known author of "Travels through North and South Carolina," I found a very intelligent, agreeable, and communicative gentleman; and from him I received considerable information about the plants of that country, particularly respecting the habitats of a number of rare and interesting trees. It is with the liveliest emotions of pleasure I call to mind the happy hours I spent in this worthy man's company, during the period I lived in his neighbourhood.

Not far from the latter place are also the extensive gardens of William Hamilton, Esq., called the Woodlands, which I found not only rich in plants from all parts of the world, but particularly so in rare and new American species. Philadelphia being a central situation, and extremely well calculated for the cultivation of plants from all the other parts of North America, I found this collection particularly valuable for furnishing me with a general knowledge of the plants of that country, preparatory to more extensive travels into the interior, for the discovery of new and unknown subjects. Mr. John Lyon, (of whom I shall have an opportunity to speak hereafter,) who had the management of these gardens, was then about to give them up: having the offer of being appointed his successor, I embraced it, and accordingly in 1802 I entered upon the situation. During my stay in this place, which was until 1805, I received and collected plants from all parts of North America; and when Michaux's "Flora Boreali-Americanæ" appeared, which was during that time, I was not only in possession of most of his plants, but had then a considerable number not described by him.

Within this period I had also formed a connection with Dr. Benjamin S. Barton, Professor of Botany in the University of Pennsylvania, &c. whose industrious researches in all the different branches of natural history are so well known to the literary world. He likewise, for some time previous, had been collecting materials for an American Flora. As I was now very anxious to explore the more remote parts of the country, particularly the interesting ranges of the Allegany Mountains, I was enabled, by the kind assistance of this gentleman, to

take a more  
which durin  
fined within  
sary attentio  
mitting me

According  
the mountai  
States, begin  
rolinas, (in v  
Virginia and  
and returnin  
countries al  
lowing seas  
Northern St  
sylvania and  
which tract l  
ing country e  
as before by

Both thes  
appropriate  
in mountaine  
more than th  
other compa  
ing up my lo  
impenetrable  
men. The  
course of the  
Dr. Barton,  
covery of ma  
history in ge  
acquired of t  
plants of tha

extensive  
e Wood-  
from all  
and new  
al situa-  
vation of  
merica, I  
urnishing  
at coun-  
he inter-  
subjects.  
ortunity  
of these  
ring the  
raced it,  
tuation.  
1805, I  
f North  
-Ameri-  
was not  
l then a

nection  
tany in  
ious re-  
history  
kewise,  
ials for  
to ex-  
icularily  
, I was  
an, to

take a more extensive range for my botanical excursions, which during my stay at the Woodlands had been confined within a comparatively small compass, the necessary attention to the duties of that establishment not permitting me to devote more time to them.

Accordingly, in the beginning of 1805, I set out for the mountains and western territories of the Southern States, beginning at Maryland and extending to the Carolinas, (in which tract the interesting high mountains of Virginia and Carolina took my particular attention,) and returning late in the autumn through the lower countries along the sea-coast to Philadelphia. The following season, 1806, I went in like manner over the Northern States, beginning with the mountains of Pennsylvania and extending to those of New Hampshire, (in which tract I traversed the extensive and highly interesting country of the Lesser and Great Lakes,) and returning as before by the sea-coast.

Both these tours I principally made on foot, the most appropriate way for attentive observation, particularly in mountainous countries; travelling over an extent of more than three thousand miles each season, with no other companions than my dog and gun, frequently taking up my lodging in the midst of wild mountains and impenetrable forests, far remote from the habitations of men. The collections and observations made in the course of these journeys, all of which I communicated to Dr. Barton, were considerable, in respect to the discovery of many new and interesting subjects of natural history in general. But the knowledge which I thereby acquired of the geography, soil, and situation of the plants of that country, (points of the greatest interest

\* \* \*  
PREFACE.

to the practical botanist) was of the greatest importance to me.

Soon after my return from the last-mentioned journey, I had the pleasure to form an acquaintance with Meriwether Lewis, Esq., then Governor of Upper Louisiana, who had lately returned from an expedition across the Continent of America to the Pacific Ocean, by the way of the Missouri and the great Columbia rivers, executed under the direction of the Government of the United States. A small but highly interesting collection of dried plants was put into my hands by this gentleman, in order to describe and figure those I thought new, for the purpose of inserting them in the account of his Travels, which he was then engaged in preparing for the press. This valuable work, by the unfortunate and untimely end of its author, has been interrupted in its publication; and although General Daniel Clark, the companion of Mr. Lewis, (to whom I transmitted all the drawings prepared for the work,) undertook the editorship after his death, it has not, to my knowledge, yet appeared before the public, notwithstanding the great forwardness the journals and materials were in when I had the opportunity of perusing them.

The collection of plants just spoken of was made during the rapid return of the expedition from the Pacific Ocean towards the United States. A much more extensive one, made on their slow ascent towards the Rocky mountains and the chains of the Northern Andes, had unfortunately been lost, by being deposited among other things at the foot of those mountains. The loss of this first collection is the more to be regretted,

when I consider to me, the dimensions, compared with me to be entirely different, at least six times the size of the idea of the continent of America, and the various regions in which it is productive.

The dimensions we find in their words "had an average some being larger and others smaller than the plants in

Here we have the future botany of the interesting specimens which the collector calls "the best of the species." resembling the genus *Thlaspi*; the leaves their bases rounded in axillary clusters, a depression in the middle, an orange color, and a genus *Microseris*.

when I consider that the small collection communicated to me, consisting of about one hundred and fifty specimens, contained not above a dozen plants well known to me to be natives of North America, the rest being either entirely new or but little known, and among them at least six distinct and new genera. This may give an idea of the discerning eye of their collector, who had but little practical knowledge of the Flora of North America, as also of the richness of those extensive regions in new and interesting plants, and other natural productions.

The descriptions of those plants, as far as the specimens were perfect, I have inserted in the present work in their respective places, distinguishing them by the words "*v. s. in Herb. Lewis.*" Several of them I have had an opportunity of examining in their living state, some being cultivated from seeds procured by Mr. Lewis, and others since my arrival in England from seeds and plants introduced by Mr. Nuttall.

Here I cannot refrain from drawing the attention of future botanists travelling those regions, to two highly interesting plants, of which I have only seen imperfect specimens. The first is what Mr. Lewis in his journals calls "the Osage Apple," or "Arrowwood of the Missouris." This is a tree, or rather shrub, with leaves resembling those of a pear-tree, but broader in proportion; they are alternate, and have a recurved thorn near their base; the flowers are of separate sexes, and appear in axillary, peduncled, globular catkins, which produce a depressed globular fruit, in size and colour resembling an orange, in interior structure approaching near to the genus *Morinda*. This shrub, on account of its fruit

and handsome foliage, must be highly ornamental : the wood, being excessively hard, is preferred by the natives to any other for making their arrows ; and as it does not spontaneously grow in their neighbourhood, they travel annually to a considerable distance South-west, to procure it. About the village of the Osage Indians a few trees have been planted, from which one has been introduced into one of the gardens at St. Louis on the Mississippi. Perfect seeds from the last-mentioned tree were given by Mr. Lewis to Mr. M'Mahon, nursery and seedsman, at Philadelphia, who raised several fine plants from them, and in whose possession they were when I left America. The other plant alluded to is called by Mr. Lewis "A bulbous Rush, which serves the Indians for bread." Of this I have only seen the root, which is of a yellowish-brown colour, about the thickness of a finger, and jointed ; on each joint, which are about three inches asunder, several tubers of about an inch and a half long are produced, of an oblong ovate shape, tapering into a point towards the end ; these tubers are filled with an exquisite white and fine farina, resembling starch ; these roots are used in the same manner as those of *Psoralea esculenta*, vid. vol. ii. p. 475. I take this to be a larger species of tuberous Cyperus.

While I was thus engaged in describing and figuring those new acquisitions to the American Flora, another opportunity offered to augment my resources. Mr. Aloysius Enslen, who had been sent to America by Prince Lichtenstein of Austria, as a collector of new and interesting subjects of natural history, returned to Philadelphia from his extensive travels through the Western Territories and Southern States. This gentle-

man, with  
timate fri  
ly valua  
I had u  
debted fo  
up a desi  
plants of  
exclusivel  
guished b

At the  
seeing th  
Mr. John  
and skill  
been intr  
channel v  
from his  
*Lyon.*"

By the  
unnecessa  
sional ex  
United S  
herbarium  
of North  
nearly de  
chaux's e  
tremely se  
pendium  
Flora Ge  
acceptabl  
cultivator  
ford and  
me every

man, with whom I had previously been on terms of intimate friendship, was now in possession of an extremely valuable collection of living and dried plants, to which I had unrestrained access. To his liberality I am indebted for many new and scarce specimens, which filled up a desideratum in my collection, particularly in the plants of Lower Louisiana and Georgia. Those species exclusively received from his collection I have distinguished by "*v. s. in Herb. Enslen.*"

At the same time I had frequent opportunities of seeing the herbarium and collection of living plants of Mr. John Lyon, a gentleman through whose industry and skill more new and rare American plants have lately been introduced into Europe than through any other channel whatever. Those plants particularly adopted from his collection I have marked with "*v. s. in Herb. Lyon.*"

By these and several other connections, which it is unnecessary to mention here, added to my own occasional excursions through the different parts of the United States, I was put in possession of an extensive herbarium, containing plants from all the different parts of North America; which when summed up would nearly double the number of those described in Michaux's excellent Flora. As that work was then extremely scarce in America, I determined to publish a Compendium of it, executed on the plan of Hoffmann's Flora Germanica, which work I had no doubt would be acceptable to the botanist, and particularly so to the cultivator. I communicated my design to Messrs. Bradford and Inskeep, booksellers at Philadelphia, who gave me every encouragement towards the prosecution thereof.

While I was engaged in arranging my materials for this publication, I was called upon to take the management of the Botanic Garden at New York, which had been originally established by the arduous zeal and exertions of Dr. David Hosack, Professor of Botany, &c., as his private property, but has lately been bought by the Government of the State of New York for the public service. As this employment opened a further prospect to me of increasing my knowledge of the plants of that country, I willingly dropped the idea of my intended publication for that time, and in 1807 took charge of that establishment.

Here I again endeavoured to pay the utmost attention to the collection of American plants, as the establishment was principally intended for that purpose. In this I was supported by my numerous botanical connections and friends, among whom I must particularly mention John Le Conte, Esq., of Georgia, whose unremitting exertions added considerably to the collection, particularly of plants from the Southern States.

The additions to my former stock of materials for a Flora were now considerable, and in conjunction with Dr. D. Hosack I had engaged to publish a periodical work, with coloured plates, all taken from living plants, and if possible from native specimens, on a plan similar to that of Curtis's Botanical Magazine; for which a great number of drawings were actually prepared. But at this period I was attacked by a serious and obstinate intermitting fever, which made a change of air and climate absolutely necessary to me; I therefore, in 1810, took a voyage to the West Indies, visiting the islands of Barbadoes, Martinique, Dominique, Guadaloupe, and

St. Bartholomew in autumn of 1811. The vessel in which I had never been with all possible safety too far advanced gained considerable knowledge of plants, interesting to New York, I Peck of Cambridge highly interested tour to the Hampshire. I was in those mountains to me.

On my return very unfurnished, the pul war with Great all my material should not on most celebrated with that enclosed works of science there.

These expected rival in London ing a circle of a science of Botany.

ials for  
anage-  
ch had  
nd ex-  
ry, &c.  
ight by  
public  
rospect  
of that  
d pub-  
f that  
ention  
abish-  
In this  
ections  
ention  
ng ex-  
xularly  
  
for a  
with  
odical  
lants,  
imilar  
ich a  
  
But  
tinate  
d cli  
1810,  
ds of  
and

St. Bartholomew's, from which I returned in the autumn of 1811 in a perfect state of recovered health. The vessel in which I returned landed at the town of Wiscasset in the province of Main. This being a country I had never visited before, I examined its vegetation with all possible attention; and although the season was too far advanced for making any new discoveries, I gained considerable information respecting the geography of plants, a point I always considered highly interesting to the science. During my journey towards New York, I had an opportunity of visiting Professor Peck of Cambridge College near Boston, and seeing his highly interesting collection of plants, collected on a tour to the alpine regions of the White Hills of New Hampshire. As the season was too far advanced when I was in that country to suffer me to think of ascending those mountains, this collection was highly gratifying to me.

On my return to New York, I found things in a situation very unfavourable to the publication of scientific works, the public mind being then in agitation about a war with Great Britain. I therefore determined to take all my materials to England, where I conceived I should not only have the advantage of consulting the most celebrated collections and libraries, but also meet with that encouragement and support so necessary to works of science, and so generally bestowed upon them there.

These expectations I found amply realized on my arrival in London. I had very soon the pleasure of forming a circle of acquaintance among those attached to the science of Botany, by whom I was gradually introduced

to the most eminent botanical characters of our age Sir Joseph Banks, with his accustomed liberality, supported my undertaking by giving me access to his extensive library and herbarium. A. B. Lambert, Esq. favoured me not only with the same assistance, but very handsomely offered me the entire use of his library and collection during the compilation of the work. To this gentleman I am in the highest degree indebted for his strenuous support under all the difficulties I had to encounter during its progress; which by his generous exertions he endeavoured to remove whenever they occurred.

Among the many new sources of information now opened to me, I must particularly mention the following original collections, by the assistance of which I have been enabled to correct many errors of former publications, particularly respecting the synonyms.

I. Clayton's Herbarium.—The whole of this interesting collection, from which the "*FLORA VIRGINICA*" of Gronovius was compiled, is preserved in the Banksian Museum. This has been particularly useful to me in tracing the plants which Linnaeus had adopted from that work.

II. Professor P. S. Pallas's Herbarium.—This extensive collection, now in the possession of A. B. Lambert, Esq. was highly useful to me, in comparing the plants of North America with those of the North of Asia, to some of which they have a great affinity, and others are common to both continents.

III. Plukenet's Herbarium.—Preserved in the British Museum, as also part of

IV. Catesby's Herbarium.

V. Walter from which compiled, i Sloane Squ Mr. Walter use of it w I am als who very ob Plants coll collection c been collect fore: those in the Supp. Besides t ber of intere collected by them I foun Menzies, I Vancouver, several of t Lewisian co zies to adop my plan, w

The num series and p contributed highly indep particularly sive collectio

The accou Sherardian i

2 VOL. I.

V. Walter's Herbarium. This interesting collection, from which Walter's "FLORA CAROLINIANA" was compiled, is in possession of the Messrs. Frasers of Sloane Square, whose father was the intimate friend of Mr. Walter. They very kindly permitted me to make use of it whenever I had occasion to refer to it.

I am also highly indebted to William Roscoe, Esq., who very obligingly communicated to me Mr. Bradbury's Plants collected in Upper Louisiana. This valuable collection contains many rare and new species, having been collected in a tract of country never explored before: those which were entirely new I have described in the Supplement to the present work.

Besides these general collections, there were a number of interesting new plants in the Banksian Herbarium collected by different persons in North America. Among them I found a number of those collected by Archibald Menzies, Esq., during the famous expedition under Vancouver, on the north-west coast of America. As several of them had been described by me from the Lewisian collection, I requested permission of Mr. Menzies to adopt such as were immediately connected with my plan, which he very obligingly complied with.

The number of American plants in the different nurseries and private collections in the vicinity of London, contributed considerably to my resources, and I am highly indebted to those who gave me their assistance, particularly to Messrs. Lee and Kennedy, whose extensive collections exceed any other in Europe.

The accounts I had frequently heard of the celebrated Sherardian Herbarium at Oxford, made me particularly

anxious to examine it. Dr. G. Williams, professor of botany in that University, received me with that kindness so congenial to the true friend of science, and supported my researches to the utmost of his power. Besides the entire Herbarium of Morison, and other valuable collections, I found a general collection of North American plants, among which I was particularly gratified to find all those sent by Mr. M. Catesby to Dr. Sherard. This collection, though made a century ago, is in very fine preservation, and contains specimens of most of the new species of Walter, Michaux, &c. and even some entirely new ones, with the remarks and synonyms of Dr. Sherard, and frequently accompanied with the original letters of M. Catesby. In the same collection are also a number of rare specimens collected by a Mr. Tilden at Hudson's Bay, and those of John Bartram and other correspondents of Dr. Sherard. In fine, I consider this the most complete collection of North American plants now extant. The notes and remarks made here are given in the Appendix and Addenda, as it was not until the body of the work was printed that I had the opportunity of seeing this valuable collection.

Having thus given a general view of the means and resources I had in composing the present work, I shall only add a few remarks respecting its execution.

On a slight inspection of the work, it will be readily perceived that my chief object has been, throughout, to comprise as much matter as possible in the smallest space which attention to perspicuity would allow.

In respect to nomenclature I have followed Willdenow's "SPECIES PLANTARUM" and Michaux's "FLORA

BOREALI-A  
carefully to  
faulty, unless  
more approp  
*Fraxinus ac*  
*americana*,  
Michaux fo  
have never a  
and *cineraria*  
*thartica*, as  
tiers" has do  
is full of sue  
long establis  
the least hin

The specif  
delled anew  
ever any alte  
quote the aut  
in a separate  
the specific  
same with  
avoid this su  
as possible,  
to the end, th  
the specific c  
so that *Stipa*  
scribed by W  
new or aug  
description w  
*giganteum*, l  
racter has bee

"BOREALI-AMERICANA" as near as possible, avoiding carefully to change old established names, however faulty, unless in cases where good authority furnished a more appropriate one. As for example, I have adopted *Fraxinus acuminata* of Lamarck, instead of *Fraxinus americana*, Linn., and *Cynoglossum amplexicaule* of Michaux for *Cynoglossum virginicum*, Linn. But I have never approved of such changes as *Juglans alba* and *cinerea* of Linnaeus to *Juglans squamata* and *cathartica*, as the younger Michaux in his "Arbres Forestiers" has done. This otherwise very useful publication is full of such new names, imposed on well known and long established species, and sometimes even without the least hint respecting their names in other authors.

The specific characters in most instances have been modelled anew or altered as it was found necessary. Wherever any alteration had been made it was necessary to quote the author from which the species had been taken in a separate line, as page 16 *Monarda didyma*; after the specific description I had to repeat that it was the same with *M. didyma*, Willd. sp. pl. 1, p. 125. To avoid this superfluous repetition, and save as much room as possible, I found it expedient to adopt, from page 72 to the end, the following mark (—), to be placed after the specific character whenever it had been amended; so that *Stipa avenacea*, page 72, is the same species described by Willd. sp. pl. 1. p. 442, but the character is new or augmented. Wherever the author follows the description without this mark, as page 73, *Saccharum giganteum*, Pers. encr. l. p. 103, in that case the character has been adopted without alteration.

In some of the more intricate genera, as *Monarda*, *Vaccinium*, *Andromeda*, *Salix*, &c. the specific characters have been extended to a rather unusual length, which may be objected to by some, but I am persuaded will be approved of by most. The beauty of a short and perspicuous character is without doubt preferable to a long one, which the more it is extended the more ambiguous it will become; but the nature of certain genera will absolutely not admit this conciseness, and the works of some of the most celebrated authors have lately proved that it is inconvenient to be limited by a certain number of words in forming the specific character. Among many others I only need to mention Willdenow's last volume of the Species Plantarum, particularly the genus *Aspidium*, and Mr. Robert Brown's excellent "Treatise on the Proteaceæ," in the tenth volume of the Transactions of the Linnean Society. In both these works examples will be found, where the length of the specific character equals if not exceeds in number of words the longest in the present work.

It would have been needless to encumber the work with a superfluous number of synonyms, which can be found in the authors referred to; but those of a late date, or any way illustrative toward the true determination of the species, have been carefully collected, as the means I was possessed of were particularly favourable to the ascertainment of those points, which may be of the highest utility to future publications on the same subject.

In respect to the figures referred to, care has been taken to quote the most correct whenever there was a choice. The plates which accompany the work have

been carefull  
new plants,  
as some of  
the plates  
work throug  
be inserted  
themselves r  
easy to supp

Particular  
soil, situatio  
found in, wl  
cal botanist,  
plants.

After the  
flowering, e  
whether I h  
or only in a  
name of the  
tioned. Th  
myself in eit  
marked (+)

To each s  
ing the colou  
plant, or som  
species.

Very full  
of the most  
work would  
references to  
this deficienc

In the alte

been carefully selected, either from highly interesting new plants, or those which will not admit of cultivation, as some of the *Rhexias*, *Lupinus*, *Gerardia*, &c. As the plates were executed during the progress of the work through the press, references to them could not be inserted in the proper places; but as the plates themselves refer to the pages of the work, it will be easy to supply this deficiency.

Particular attention has been paid in describing the soil, situation, and range of country each species is found in, which will be found very useful to the practical botanist, and especially so to the cultivator of those plants.

After the usual sign of duration, and the time of flowering, each species has been faithfully marked whether I have seen it myself in a living state (*v. v.*), or only in a dried specimen (*v. s.*); in the latter case the name of the herbarium I had it from is generally mentioned. Those I have adopted without seeing them myself in either state, and which are but few, I have marked (+).

To each species a short note has been added, respecting the colour and size of the flower, the habit of the plant, or some other interesting subject illustrative of the species.

Very full descriptions have been given of only some of the most interesting new species, as the limits of the work would not admit of it in general. The careful references to authors and good figures will amply supply this deficiency.

In the alteration made in the systematic arrangement,

I have been guided by the hints given by the learned Dr. J. E. Smith, in his excellent "Introduction to Botany." The classes Dodecandria and Polyadelphia are entirely omitted, and their genera inserted in their proper places in the other classes. *Asarum* next to *Aristolochia*, in Gynandria. *Bejaria*, *Portulacca*, *Talinum*, *Calligonum*, and *Hypericum*, in Polyandria. *Hudsonia* next to *Helianthemum*, with which it is closely related in the same class, *Decumaria*, *Lythrum*, *Cuphea*, and *Agrimonia*, in Icosandria. *Halesia* near *Styrax*, in Monadelphia. *Euphorbia* in Diclinia, where it forms, with its other genera, a natural as well as artificial order. The nineteenth class, Diclinia, contains the natural families of Tricoccæ or Euphorbia, Amentaceæ, and Coniferæ. The other genera of the classes Monoecia, Dioecia, and Polygamia, having been referred to their proper classes. The genus *Diotis* more properly belongs to Tetrandria, but had been omitted in that class. *Najas* and *Valisneria* are genera of an order of this class, to which probably might be added all the Aroideæ. How far these alterations may meet the approbation of those best qualified to judge, I have yet to learn, but I am strongly persuaded they will facilitate the study of this lovely science to the young beginner.

In the last class, Cryptogamia, I have gone no further than the family of the *Filices*, according to the formation of the plan of this work, which only included plants which either are, or can be cultivated. The other orders, *Musci*, *Algæ*, and *Fungi*, will form a distinct work from the present, and may appear at some future

period in a  
work would  
illustrate th  
out which i

At the e  
malous Fl  
it as corre  
service, pa  
purpose I st  
the young b  
lium, happi  
he would in  
referring to  
will very ea  
amining a f  
consider th  
and look fo  
sistance of

For the a  
given a dou  
tion has b  
Index, in  
to the num  
The synony

Notwith  
the correct  
that it will  
unavoidable  
tions of thi  
rities are to  
self with t

the learned  
tion to Bo-  
delphia are  
in their pro-  
ct to *Aris-*  
, *Talinum*,  
*Hudson-*  
*is closely*  
*hrum, Cu-*  
*ilesia near*  
*inia, where*  
well as ar-  
a, contains  
ia, Amen-  
the classes  
been re-  
*notis more*  
omitted in  
era of an  
be added  
may meet  
ge, I have  
they will  
the young  
  
no further  
the forma-  
included  
The other  
a distinct  
one future

period in a more perfect shape than the limits of this work would admit; it being absolutely necessary to illustrate those minute objects with correct figures, without which mere words are useless in describing them.

At the end of the work I have given a Table of Anomalous Flowers; great pains have been taken to make it as correct as possible. This will be found of infinite service, particularly to the young botanist. For this purpose I shall give an example of the use of it. Should the young beginner, unacquainted with the genus *Galium*, happen to find *Galium trifidum*, vol. i, p. 103, he would in vain look for its genus in Triandria; but by referring to this table among the anomalous genera, he will very easily find its proper genus. Again, by examining a flower of *Asclepias*, he might very probably consider the ten pollen masses for as many anthers, and look for its genus in Decandria, which by the assistance of this table he will trace to be Pentandria.

For the accommodation of the English reader, I have given a double Latin and English Index. Great attention has been paid to the correctness of the General Index, in which the first number in parentheses refers to the number of the genus, the other to the page. The synonyms are distinguished by being in Italics.

Notwithstanding the unremitting attention paid to the correctness of this work, the author is well aware that it will not be found free from those errors which are unavoidable in (nay almost inseparable from) publications of this nature, where so great a number of authorities are to be referred to. But he still consoles himself with the idea of having given to the world a work

which, with all its imperfections, will be found useful, and contribute to the general advancement and progress of the science. This has been the only aim, and will be the highest reward of the labours of

the author, who, in his desire to have his work published, has not been able to find any publisher willing to take it up.

**THE AUTHOR,**

wishes to thank all those who have assisted him in the preparation of this work.

**LONDON,**

December 1813.

## REFERI

### Tab. / Page.

1.	17.	Mo
2.	62.	Mil
3.	175.	Clay
4.	219.	Ber
	1.	1
	2.	tl
	3.	p
	4.	3.
	5.	4, 5
5.	219.	Ber
	1.	1
	2.	A
6.	224.	Con
7.	156.	Sola
8.	228.	Lili
	1.	1
	2.	A
9.	243.	Hel
10.	258.	Rhe
	by	
	pr	
	or	
	2.	A
11.	260.	Clar
12.	283.	Gaul
13.	283.	Gaul
	ma	
	21.	Cera
	siz	
	178.	The
	2.	
	3.	
14.	302.	Myl
	an	
	vie	
	sti	

## REFERENCES TO THE PLATES.

### Tab. / Page.

1. 17. *Monarda Kalmiana*.
2. 62. *Milium amphicarpon*.
3. 175. *Claytonia lanceolata*.
4. 219. *Berberis Aquifolium*.
  1. A single pinna of a full grown leaf, from the under side.
  2. An expanded flower, with its component parts viewed from behind, natural size.
  3. A petal with its stamen, magnified.
  - 4, 5. A stamen and pistillum, more magnified.
5. 219. *Berberis nervosa*.
  1. The calyx, natural size.
  2. A petal with its stamen, magnified.
6. 224. *Conostylis americana*.
7. 156. *Solanum heterandrum*.
8. 228. *Lilium pudicum*.
  1. The pistil.
  2. A stamen, both natural size.
9. 243. *Helonias tenax*.
10. 258. *Rhexia ciliosa*, and *Rhexia lutea*. Both species, by a mistake of the engraver, have been represented with five petals, which ought to be only four.
  2. A variety of *Rhexia lutea*.
11. 260. *Clarkia pulchella*.
12. 283. *Gaultheria Shallon*; with a stamen magnified.
13. 283. *Gaultheria serpyllifolia*; a flowering branch, magnified.
14. 21. *Ceratiola ericoides*; a female branch, natural size.
178. The fruit of *Hamiltonia oleifera*, natural size.
  2. A longitudinal cut of the same.
  3. The kernel.
14. 302. *Mylocarium ligustrinum*. The dissections are: an opened corolla with the stamens; a back view of the calyx and the germ with its stigma; all magnified.

## REFERENCES TO THE PLATES.

## Tab. Page.

15. 333. *Tigarea tridentata*.  
 1. A flower divested of its petals.  
 2. A fruit.  
 3. The same longitudinally cut to show the seed.
16. 248. *Rubus spectabilis*.
17. 390. *Caltha flabellifolia*.
18. 423. *Gerardia fruticosa*.
19. 423. *Gerardia quercifolia*.
20. 427. *Mimulus Lewisii*.
21. 468. *Lupinus villosus*.
22. 475. *Psoralea esculenta*.  
 1. The calyx.  
 2. The vexillum.  
 3. The carina with the alæ.  
 4. A fruit.
23. 479. *Lupinaster macrocephalus*.
24. 499. *Prenanthes Serpentaria*.

IN

*Abbot. insect.*—

terous Ins

vations of

1797. 41

*Act. holm.*—K

Stockholm

*Act. petrop.*—A

politanæ P

1786. 41

*Act. soc. amer.*—

cal Society

Knowledge

*Act. soc. hist. n*

turelle de

*Act. soc. nat. sc*

bel. Kio

*Act. soc. linn. l*

London.

*Act. ups.*—Acta

Stockholm

*Ait. kew.*—Hor

cultivated i

Aiton. V

*Ait. kew. ed.* 2

Townsend

*Allion. fl. ped.*—

Augustæ T

*Alp. exot.*—Pros

tii, 1629.

*Amcn. acad.*—C

10. Holm

8vo.

*Amm. rath.*—St

to show the

## INDEX AUCTORUM.

---

- Abbot. insect.*—The Natural History of the rarer Lepidopterous Insects of Georgia, collected from the Observations of J. Abbot, by J. E. Smith. 2 vols. London, 1797. 4to.
- Act. holm.*—Kongl. Vetenskaps Academiens Handlingar. Stockholm, 1739 et seq. 8vo.
- Act. petrop.*—Acta Academiae Scientiarum Imperialis Petropolitanae pro Annis 1777—1782. Petropoli, 1778—1786. 4to.
- Act. soc. amer.*—Transactions of the American Philosophical Society, held at Philadelphia, for promoting useful Knowledge. Philad. 1771 et seq. 4to.
- Act. soc. hist. nat. par.*—Actes de la Société d'Histoire Naturelle de Paris. Tom. 1, 1 partie. Paris, 1792. fol.
- Act. soc. nat. scrut. haf.*—Skrivter of Naturhistorie Selskab. Kiobenhaven, 1790 et seq. 8vo.
- Act. soc. linn. lond.*—Transactions of the Linnean Society of London. London, 1791 et seq. 4to.
- Act. ups.*—Acta Societatis Regiae Scientiarum Upsaliensis. Stockholmiae, 1744 et seq. 4to.
- Ait. kew.*—Hortus Kewensis; or, A Catalogue of the Plants cultivated in the Royal Botanic Garden at Kew, by W. Aiton. Vol. 1—3. London, 1789. 8vo.
- Ait. kew. ed. 2.*—The second edition enlarged, by W. Townsend Aiton. Vol. 1—5. London, 1810.
- Allion. fl. ped.*—Car. Allioni Flora Pedemontana. Tomi 3. Augustae Taurinorum, 1785. fol.
- Alp. exot.*—Prosop. Alpini de Plantis exoticis Libri 2. Venetiis, 1629. 4to.
- Amoen. acad.*—Carolii Linnæi Amoenitates academicæ. Vol. 10. Holmiae, 1749—1769. Erlangæ, 1785. 1790. 8vo.
- Amm. rath.*—Stirpium rariorum in Imperio Rutheno sponte

- provenientium Icones et Descriptiones collectæ a Io. Ammano. Petropoli, 1739. 4to.
- Annales du mus. d'hist. nat.* *Ann. mus. hist. nat.*—Annales du Museum d'Histoire Naturelle, par les Professeurs de cet Etablissement. Paris, 1802 et seq. 4to.
- Annals of bot.*—Annals of Botany, by C. Konig and J. Sims. 2 vols. London, 1805, 1806. 8vo.
- Ard. spec.* *Arduin. spec.*—Petr. Arduini Animadversionum botanicarum Specimen 1. Patavii 1759. Specimen 2. Venetiis, 1764. 4to.
- Aubl. guj.* *Aubl. fl. guj.*—Histoire des Plantes de la Gujane Francoise, par M. Fusée Aublet. Tom. 4. Paris, 1775. 4to.
- Barrel. ic.* *Barr. rar.*—Plantæ per Galliam, Hispaniam et Italiam observatæ, Iconibus æneis exhibitæ a Jac. Barrellero. Parisiis, 1714. fol.
- Bartr. it.* *Bartr. trav.*—Travels through North and South Carolina, Georgia, East and West Florida, &c., by W. Bartram. London, 1794. 8vo.
- Blackw.*—Herbarium Blackwellianum emendatum et auctum, cum Praefatione D. Chr. Trew. Cant. 1—6. Novimber. 1750—1772. fol.
- Bocc. sic.*—Icones et Descriptiones rariorū Plantarū Siciliæ, Melitæ, Galliæ, et Italiae; Auctore Paulo Boccone. Oxonii, 1674. 4to.
- Bolton filic.*—Jam. Bolton, Filices Britanniæ, an History of the British proper Ferns, partes 1—2. Leeds and Huddersfield, 1785—1790. 4to.
- Bot. mag.*—Curtis's Botanical Magazine, continued by J. Sims. London, 1801 et seq. 8vo.
- Bot. rep.*—The Botanist's Repository for new and rare Plants, by H. Andrews. London, 1797 et seq. 4to.
- Breyn. cent.*—Jac. Breynii Exoticarum aliarumque minus cognitarum Plantarum Centuria prima. Gedani, 1678. fol.
- Brown. jam.*—The Civil and Natural History of Jamaica, by Patr. Browne. London, 1756. fol.
- Bullet. de la soc. philomat.*—Bulletin des Sciences, par la Société Philomatiique. Paris, 1797 et seq. 4to.
- Cam. epit.*—De Plantis Epitome Petr. Andr. Matthioli, aucta a Joach. Camerario. Francof. ad Mœn. 1586. 4to.
- Catesb. car.*—The Natural History of Carolina, Florida, and the Bahama Islands, by Mark Catesby. 2 vols. London, 1747, 1748. fol.
- Cav. diss.*—Ant. Jos. Cavanilles, Monadelphiae Classis Dissertationes 4to.
- Cav. ic.*—Eiusdem aut sponte iuntur. Ve
- Clayt. fl. virg.*—
- Col. ephr.*—Tab nostro Cœle 1613. Par
- Com. græt.*—Co Gættingensi
- Comm. hort.*—F Plantarum L lino. Amst Commelin
- Comm. præl.*—C Batav. 1703
- Corn. canad.*—Ja rumque non Crantz fl. austr.—Austriacarun
- Curt. lond.*—Cur nensis. 2 v Delaroche eryng. roche. Par
- Deff. atl.*—Flora Tomi 2. P
- Diar. hist. nat.*—Paris, 1792.
- Dill. elth.*—Joh. Londini, 17
- Dill. musc.*—Eju 4to.
- Dod. mem.* *Dod stoire des Pla*
- Donn. catal.* *Don James Donn.*
- Duhamel. arb.*—I et Arbustes Tomes 2. P
- Du Roi harbk.*—J Baumzucht. ed.
1795. 8vo.

llectæ a Io.  
st.—Annales  
Professeurſ  
4to.  
onig and J.  
vo.  
adversionum  
Specimen 2.  
de la Gujane  
4. Paris,  
Iſpaniam et  
ibitæ a Jac.  
h and South  
&c., by W.  
itum et auc-  
. 1—6. No-  
lantarum Si-  
Paulo Boc-  
n History of  
eds and Hud-  
tinued by J.  
d rare Plants,  
4to.  
mque minus  
ledani, 1678.  
of Jamaica,  
ences, par la  
4to.  
Latthioli, auc-  
. 1586. 4to.  
Florida, and  
t vols. Lon-  
Classis Dis-

- Sertationes decem. Parisiis et Matriti, 1785—1790.  
4to.  
*Cav. ic.*—Eiusdem Icones et Descriptiones Plantarum quæ aut sponte in Hispania crescunt, aut in Hortis hospitantur. Vol. 1—6. Matriti, 1791—1801. fol.  
*Clayt. fl. virg.*—vid. *Gron. fl. virg.*  
*Col. ecphr.*—Tab. Columnæ minus cognitarum rariorumque nostro Cœlo orientium Stirpium Ecphrasis. Romæ, 1613. Pars 2da. *ibid.* 1616. 4to.  
*Com. gret.*—Commentarii Societatis Regiæ Scientiarum Göttingensis. Göttingæ, 1751 et seq. 4to.  
*Comm. hort.*—Horti Medici Amstelodamensis rariorum Plantarum Descriptio et Icones, Auctore Io. Commelinno. Amstelodamii, 1697. Pars altera, Auctore Casp. Commelinno, 1701. fol.  
*Comm. præl.*—Casp. Commelinii Præludia Botanica. Lugd. Batav. 1703. 4to.  
*Corn. canad.*—Jac. Cornuti Canadensium Plantarum aliarumque nondum editarum Historia. Parisiis, 1635. 4to.  
*Crantz fl. austr.*—Henr. Joan. Nepom. Crantz, Stirpium Austriacarum. Pars 1, 2. Vien. 1769. 4to.  
*Curt. lond.* *Curt. fl. lond.*—William Curtis's Flora Londinensis. 2 vols. London, 1777. 1798. fol.  
*Delaroche eryng.*—Eryngiorum Historia, Auctore F. Delaroche. Parisiis, 1808. fol.  
*Desf. atl.*—Flora Atlantica, Auctore Renato Desfontaines. Tomi 2. Parisiis. 4to.  
*Diar. hist. nat.*—Journal d'Histoire Naturelle. Tom. 1, 2. Paris, 1792. 8vo.  
*Dill. elth.*—Joh. Jac. Dillenii Hortus Elthamensis. Vol. 2. Londini, 1732. fol.  
*Dill. musc.*—Eiusdem Historia Muscorum. Oxon. 1741. 4to.  
*Dod. mem.* *Dodart. mem.*—Mémoires pour servir à l' Histoire des Plantes, par M. Dodart. Paris, 1676. fol.  
*Donn. catal.* *Donn. hort. cant.*—Hortus Cantabrigiensis, by James Donn. Cambridge, 1812. 8vo.  
*Duhamel. arb.*—Du Hamel du Monceau, Traité des Arbres et Arbustes qui se cultivent en France en pleine Terre. Tomes 2. Paris, 1755. 4to.  
— *ed. nov.*—Eiusdem Operis Editio nova. *ib.* fol.  
*Du Roi harbk.*—Joh. Phil. Du Roi, Die Harbkesche wilde Baumzucht. 1. 2. Theil. Braunschw. 1771, 1772. 8vo.  
— *ed. 2.*—Eiusdem Operis Editio nova. *ib.* 1795. 8vo.

## INDEX AUCTORUM.

- Ehret. pict.*—Plantæ et Papilioes rariores, depictæ et æri incisæ a Ge. Dion. Ehret. Londini, 1748, 1759. fol.
- Ehrh. beitr.*—Frid. Ehrhart Beiträge zur Naturkunde. 1—7 Band. Hann. 1787—1792. 8vo.
- Engl. bot.*—English Botany, by J. E. Smith, the Figures by J. Sowerby. London, 1790 et seq. 8vo.
- Exot. bot.*—Exotic Botany, by J. E. Smith, the Figures by J. Sowerby. London, 1804 et seq. 4to.
- Feuil. peruv.*—Journal des Observations faites sur les Côtes Orientales de l'Amérique Méridionale, par le R. P. Louis Feuillée. Tomes 3. Paris, 1714. 1725. 4to.
- Fl. dan.*—Icones Plantarum sponte nascentium in Regnis Daniae et Norwegiae, editæ a G. C. Oeder, O. F. Müller, et Mart. Vahl. Hafniæ, 1761 et seq. fol.
- Fl. græc.*—Flora Græca, sive Plantarum rariorum Historia, quas in Provinciis aut Insulis Græciæ legit, investigavit, et depingi curavit Joh. Sibthorp; Characteres omnium, Descriptiones et Synonyma elaboravit J. E. Smith. Londini, 1806 et seq. fol.
- Fl. lapp.*—Car. Linnæi Flora Lapponica. Amstelod. 1738. 8vo.
- Fl. per.* *Fl. peruv.*—Flora Peruviana et Chilensis: sive Descriptiones et Icones Plantarum Peruvianarum et Chilensem, Auctoribus Hipp. Ruiz et Jos. Pavon. Matrixi, 1798—1802. fol.
- Gært. de fr. et sem.*—Jos. Gærtner, de Fructibus et Seminibus Plantarum. Vol. 1 et 2. Stutgard, 1788—1791. 4to.
- Gært. carp.*—G. F. Gærtner, Carpologia: seu Continuatio Operis de Fructibus et Seminibus Plantarum. Vol. 3. Tubingæ, 1805. 4to.
- Gen. pl.*—Car. a Linné, Genera Plantarum, Ed. 8vo. curante J. C. D. Schreber. Vol. 2. Francof. 1789. 1791. 8vo.
- Gmel. sib.* *Gmel. fl. sib.*—J. G. Gmelin, Flora Sibirica. Tom. 1—4. Petrop. 1747—1769. 4to.
- Gmel. syst. nat.* *Gmel. syst. veg.*—Car. a Linné, Systema Naturæ, Editio Gmelini. Lips. 1788—1793. 8vo.
- Gron. virg.* *Gron. fl. virg.*—J. F. Gronovius, Flora Virginica, exhibens Plantas quas J. Clayton in Virginia colligit. Lugd. Batav. 1762. 4to.
- Häll. helv.* *Hall. fl. helv.*—Alb. v. Haller, Historia Stirpium indigenarum Helvetiae inchoata. Tomi 3. Bernæ, 1768. fol.

- Herm. lugdb.*—Lugduno-  
*Herm. parad.*  
*Hoffm. phytop* 1 Band.  
*Hoffm. sal.*—Ej siæ, 1787.  
*Houttuyn. linn.* Linné, V Nurnb. 171  
*Hort. angl.*—A Flowers wl near Lond  
*Hort. cliffort.*—lodami, 17  
*Hort. schœnbr.*—brunnensis Vol. 1—3.  
*Host gram.*—N num Aust 1 803. fol  
*Huds. angl.*—G. 1778. 8vo  
*Humb. et Bonpl.* Bonpland P  
*Jacq. austr.* Ja aca. Vol. t  
*Jacq. collect.* J njicam, Che Vol. 4 et Si 4to.  
*Jacq. hort.* Jac Vindobonen  
*Jacq. hort. scheen*  
*Jacq. ic.* Jacq. i rum. Vol.  
*Jacq. obs. bot.*—Partes 4. V  
*Jacq. oxal.*—Eju trata. Vien  
*Journ. hist. nat.*  
*Juss. gen. pl.*—A Ordines Natu 8vo.  
*Knorr delic.* Kn

## INDEX AUCTORUM.

xxx

- epictæ et æri  
s, 1759. fol.  
rkunde. 1—7  
ne Figures by  
o.  
Figures by  
sur les Côtes  
par le R. P.  
1725. 4to.  
n in Regnis  
, O. F. Mul-  
fol.  
im Historia,  
investigavit,  
res omnium,  
Smith. Lon-  
telod. 1738.  
sis : sive De-  
rum et Chi-  
'avon. Ma-  
s et Semini-  
788—1791.  
Continuatio-  
m. Vol. 3.  
8vo. curante  
1789. 1791.  
ra Sibirica.  
né, Systema  
93. 8vo.  
Flora Virgin-  
in Virginia  
ia Stirpium  
i. Bernæ,

- Herm. lugdb.*—Pauli Hermanni Catalogus Horti Academicus  
Lugduno-Batavi. Lugd. Batav. 1687. 8vo.  
*Herm. parad.*—Eiusdem Paradisus Batavus, ib. 1698. 4to.  
*Hoffm. phytogr.*—G. F. Hoffmann Phytographische Blätter.  
1 Band. Götting. 8vo.  
*Hoffm. sal.*—Eiusdem Historia Salicum. Tom. 1, 2. Lipsiæ, 1787. fol.  
*Houttuyn. linn. syst.*—Mart. Houttuyn des Ritters von  
Linné, Volständiges Pflanzensystem. 1—14 Theils.  
Nurnb. 1771—1788. 8vo.  
*Hort. angl.*—A Catalogue of Trees, Shrubs, Plants, and  
Flowers which are propagated for Sale in the Gardens  
near London. London, 1730. fol.  
*Hort. cliffort.*—Car. Linnæi Hortus Cliffortianus. Amste-  
lodami, 1737. fol.  
*Hort. schœnbr.*—Plantarum riariorum Horti Cæsarei Schœn-  
brunnensis Descriptiones et Icones, opera N.J. Jacquin.  
Vol. 1—3. Viennæ, 1797—1798. fol.  
*Host gram.*—N. V. Host, Icones et Descriptiones Grami-  
num Austriacorum. Vol. 1—3. Vindobonæ, 1801—  
1803. fol.  
*Huds. angl.*—G. Hudson, Flora Anglicæ. Tom. 1, 2. Lond.  
1778. 8vo.  
*Humb. et Bonpl. equinoct.*—Alex. de Humboldt et Amati  
Bonpland Plantæ Æquinoctiales. Parisiis, 1805. fol.  
*Jacq. austr.* *Jacq. fl. austr.*—R. J. Jacquin, Flora Austria-  
ca. Vol. 5. Viennæ, 1773—1778. fol.  
*Jacq. collect.* *Jacq. coll.*—Eiusdem Collectanea, ad Botan-  
icam, Chemiam et Historiam Naturalem spectantia.  
Vol. 4 et Supplementum. Vindobonæ, 1786—1796.  
4to.  
*Jacq. hort.* *Jacq. hort. vind.*—Eiusdem Hortus Botanicus  
Vindobonensis. Tomi 3. Viennæ, 1770—1776. fol.  
*Jacq. hort. schœnbr.* *Jacq. schœnbr.*—Vid. *Hort. schœnbr.*  
*Jacq. ic.* *Jacq. ic. rar.*—Eiusdem Icones Plantarum rario-  
rum. Vol. 3. Vindobonæ, 1781—1793. fol.  
*Jacq. obs. bot.*—Eiusdem Observationum Botanicarum  
Partes 4. Vindobonæ, 1764—1771. fol.  
*Jacq. oxal.*—Eiusdem Oxalis, Monographia Ieonibus illus-  
trata. Viennæ, 1794. 4to.  
*Journ. hist. nat.*—Vid. *Diar. hist. nat.*  
*Juss. gen. pl.*—A. L. Jussieu, Genera Plantarum secundum  
Ordines Naturales disposita, Editio Usteri. Turici, 1791.  
8vo.  
*Knorr delic.* *Knorr thes.*—G. Wolffg. Knorr, Thesaurus

## INDEX AUCTORUM:

- Rei Herbariae Universalis.* Vol. 2. Nurnberg. 1772.  
fol.
- Lam. encycl.*—Encyclopédie Méthodique; Botanique, par  
M. le Chevalier de Lamarck. Paris, 1783 et seq. 4to.
- Lam. illustr.*—Eiusdem Illustration des Genres. Paris,  
1791 et seq. 4to.
- Lambert monogr.*—A Description of the Genus *Pinus*, by  
Aylmer Bourke Lambert. London, 1803. fol.
- Lapeyr.*—Figures de la Flore des Pyrénées, par P. Picot La-  
peyrouse. Tom. 1. Paris, 1795. fol.
- Leers fl. herb.*—J. D. Leers, Flora Herbornensis: Herbornæ,  
1775. 8vo.
- L'Her. Corn.*—C. L. L'Heritier *Cornus*. Parisiis, 1788.  
fol.
- L'Herit. sert. angl.*—Eiusdem *Sertum Anglicum*. Parisiis,  
1788. fol.
- L'Herit. stirp.*—Eiusdem *Stirpes novæ aut minus cognitæ*.  
Parisiis, 1784, 1785. fol.
- Linn. fl. lapp.*—Vid. *Fl. lapp.*
- Linn. fil. dec.*—Car. a Linné Filius Decas 1–2 Plantarum  
rariorum Horti Upsaliensis. Stockholmæ, 1762, 1763.  
fol.
- Linn. hort. cliff.*—Vid. *Hort. Cliffort.*
- Linn. mant.*—C. a Linné Mantissa Plantarum. Holmiæ,  
1767—1771. 8vo.
- Linn. sp. pl.*—Eiusdem *Species Plantarum*. Holmiæ, 1753.  
8vo.
- Linn. suppl.*—C. a Linné Filius *Supplementum Plantarum*.  
Brunswigæ, 1781. 8vo.
- Linn. syst. veg.*—C. a Linné *Systema Vegetabilium*, curante  
Murray. Gottingæ, 1784. 8vo.
- Linn. trans.* *Linn. soc. trans.*—Vid. *Act. soc. linn. lond.*
- Lob. ic.*—Petro Pena et Matthia de Lobel *Plantarum seu  
Stirpium Icones*. Tom. 1, 2. Antwerp. 1581—1591.  
4to. dbl.
- Loes. pruss.*—J. Loeselii *Flora Prussica*. Regiomonti, 1703.  
4to.
- Magn. hort.*—Petr. Magnol *Hortus Regius Monspeliensis*.  
Monspelii, 1697. 8vo.
- Marsh. arb.*—Humph. Marshall *Arbustum Americanum*.  
Philadelphiae, 1758. 8vo.
- Mart. dec.* *Mart. cent.*—J. Martyn, *Historia Plantarum  
rariorum*. Londini, 1728. fol.
- Mem. de l'instit. sc. phys.*—Mémoires de l'Institut National  
des Sciences et Arts. Paris, an 6 et seq. 4to.

*Meerb. ic.*—A  
Nic. Mee  
Pl  
Nic. Mee  
*Merian surin.*—  
morphosil  
Comitum,  
*Mich. arb.*—H  
Septentrio  
ris, 1810.  
*Mich. fl. amer.*  
Tom. 2.  
*Mich. querc.*—  
Paris, 186  
*Mich. gen.* M  
tore Petros  
*Mill. ic.* *Mill*  
Gardener's  
don, 1760  
*J. F. Mill. ic.*—  
by J. F. M  
*Mœnch meth.*  
Botanici e  
burg, 179  
*Mœnch weiss.*  
Bäume un  
Cassel. F  
*Moris. hist.*—F  
lis. Pars  
*Murray com. g*  
*Ort. dec.*—C. C  
Horti Re  
Matriti, 1  
*Pall. astrag.*—  
scriptæ et  
*Pall. fl. ross.*—  
trop. 1784  
*Pall. illustr.*—  
vel nondu  
*Pall. itin.*—Ej  
Provinzen  
1771—17  
*Parad. lond.*—  
cultivated  
scriptions  
Hooker.

- Meerb. ic.*—Afbeeldingen van Zeldzaame Gewassen, door Nic. Meerburgh: Leyden, 1775. fol.
- Plantarum selectarum Icones pictæ, editæ a Nic. Meerburgh: Lügd. Batav. 1798. fol.
- Merian surin.*—M. Sibylla Merian de Generatione et Metamorphosibus Insectorum Surinamensium. Hagæ Comitum, 1726. fol.
- Mich. arb.*—Histoire des Arbres Forestiers de l'Amérique Septentrionale, par F. André Michaux. Tom. 3. Paris, 1810—1812. 8vo.
- Mich. fl. amer.*—Andr. Michaux Flora Eoreali-Americanæ. Tom. 2. Paris, 1803. 8vo.
- Mich. querc.*—Ejusdem Histoire des Chênes de l'Amérique. Paris, 1801. fol.
- Mich. gen.* Michel. gen.—Nova Plantarum Genera, Auctore Petro A. Michelio. Florentiæ, 1729. fol.
- Mill. ic.* Mill. icon.—Figures of Plants, described in the Gardener's Dictionary; by Phil. Miller. 2 vols. London, 1760. fol.
- J. F. Mill. ic.*—60 Coloured Plates of Plants and Animals, by J. F. Miller. 1776—1794. fol.
- Mœnch meth.*—C. Mœnch, Methodus Plantarum Horti Botanici et Agri Marburgensis a Staminum Situ. Marburg. 1794. 8vo.
- Mœnch weissenst.*—Ejusdem Verzeichnis Ausländischet Bäume und Stauden des Lustschlosses Weissenstein by Cassel. Francf. 1785. 8vo.
- Moris. hist.*—Rob. Morison, Plantarum Historiæ Universalis. Pars 2, 3. Oxon. 1680—1699. fol.
- Murray com. gætt.*—Vid. *Com. gætt.*
- Ort. dec.*—C. G. Ortegæ novarum aut rariorū Plantarum Horti Reg. Botan. Matrit. Descriptionum Decades. Matriti, 1797 et seq. 4to.
- Pall. astrag.*—Petr. Sim. Pallas Species Astragalorum de scriptæ et Iconibus coloratis illustratae. Lipsiæ, 1800. fol.
- Pall. fl. ross.*—Ejusdem Flora Rossica. Tom. 1, 2. Petrop. 1784—1788. fol.
- Pall. illust.*—Ejusdem Illustrationes Plantarum imperfecte vel nondum cognitarum. Lipsiæ, 1803. fol.
- Pall. itin.*—Ejusdem Reise durch verschiedene Russische Provinzen des Russischen Reichs. P. 1—3. Petersb. 1771—1776. 4to.
- Parad. lond.*—The Paradisus Londinensis, containing Plants cultivated in the Vicinity of the Metropolis; the Descriptions by R. A. Salisbury, the Figures by William Hooker. London, 1806 et seq. 4to.

- Pers. enh.* — *Pers. syn.* — Synopsis Plantarum, seu Enchiridium Botanicum, curante C. H. Persoon. Pars 1, 2. Parisis, 1805. 12mo.
- Petiv. gaz.* — Jac. Petiver, Gazophylacium Naturæ et Artis. fol.
- Philad. philos. trans.* — Vid. *Act. soc. amer.*
- Pl. rar. hung.* — Franc Comitis Waldstein et Pauli Kitaibel Descriptiones et Icones Plantarum rariorum Hungariae. Tomi 1, 2. Viennæ, 1802—1805. fol.
- Pluk. alm.* — Leon. Plukanett, Almagestum Botanicum. Londini, 1696. 4to.
- Pluk. amalh.* — Ejusdem Almaltheum Botanicum. Londini, 1705. 4to.
- Pluk. mant.* — Ejusdem Mantissa Almagesti Botanici. Londini, 1700. 4to.
- Pluk. phyt.* — Ejusdem Phytographia. Londini, 1691, 1692. 4to.
- Plum. amer.* — Carl. Plumier, Descriptions des Plantes de l'Amérique. Paris, 1693. fol.
- Plum. filic.* — Ejusdem des Fougères de l'Amérique. Paris, 1705. fol.
- Plum. ic.* — Ejusdem Plantarum Americanarum Fasciculi decem, continentes Plantas quas olim C. Plumierius detectis: edidit J. Burmannus. Amstel. 1755—1760. fol.
- Pollich fl. palat.* — J. A. Pollich, Historia Plantarum in Palatinatu Electorali sponte crescentium. Tom. 1—3. Mannh. 1776—1778. 8vo.
- Redouté lil.* — Les Liliacées, par P. J. Redouté. Paris, 1802 et seq. fol.
- Retz scand.* — Andr. Joh. Retz, Floræ Scandinaviae Prodromus. Lipsiæ, 1795. 8vo.
- Retz obs.* — Ejusdem Observationes Botanicæ. Lipsiæ, 1779—1791. fol.
- Rheed. mal.* — Hortus Indicus Malabaricus, adornatus per Henr. van Rheede van Drakenstein. Tomi 12. Amstelodami, 1678—1703. fol.
- Rivin. monop.* — A. Q. Rivini Ordo Plantarum quæ sunt Flore irregulare monopetalio. Lipsiæ, 1691. fol.
- Rob. ic.* — 319 Plates of Plants, engraved by Nic. Robert, A. Bosse, and L. de Chastillon. fol.
- Rostk. junc.* — F. G. Rostkovii Dissertatio Botanica de Junco. Halæ, 1801. 8vo.
- Roth. catalect.* — A. W. Roth, Catalecta Botanica, Fascicul. 1, 2. Lipsiæ, 1797—1800. 8vo.
- Roth. germ.* — Ejusdem Tentamen Floræ Germanicæ. Tom. 3. Lipsiæ, 1798. 1800. 8vo.

*Roth. neue bei*  
Bremen.  
*Rottb. gram.* —  
rum pro  
1786. fo  
*Sabb. hort.* — fortii, &c.  
tius Sabbatius  
*Salisb. stirp. ra*  
Descriptio  
*Scheuchz. gra*  
Graminum  
Historia.  
*Schk. car.* — C. tenb. 1801  
*Schkahr handb*  
Wittenber  
*Schkahr filic.* —  
*Schmidt arb.* — Baumzuel  
*Schreb. gram.* — Leipzig, 1  
*Scop. del. insul*  
subricæ.  
*Sloan. hist. jan*  
Madeira,  
maica, wi  
Islands, b  
1725. fo  
*Smith brit.* — S  
Vol. 1—3.  
*Smith exot. bot.*  
*Smith ic. ined.*  
tore J. E. 1  
*Smith ic. pict.* — Londini, 1  
*Smith spicil.* — 1791 et seq.  
*Sp. pl.* — Vid. 1  
*Swartz fl. ind.* — lis. Tom.  
*Swartz ic.* — Ej  
in India O  
Erlangæ,  
*Swartz observ.* langæ, 179  
*Swartz prodr.* —

- Roth* *neue Beitr.*—Eiusdem *Neue Beiträge zur Botanik.*  
Bremen. 8vo.
- Rottb. gram.*—C. Friis Rottböl *Descriptiones et Icones rario-*  
*rum pro maxima parte novarum Plantarum.* Hafn.  
1786. fol.
- Sabb. hort.*—*Hortus Romanus secundum Systema Tourne-*  
*fortii, &c.* Species suppeditabat et describēbat Liber-  
tius Sabbati. *Tomi 4.* Romæ, 1772—1777. fol.
- Salisb. stirp. rar.*—R. A. Salisbury, *Icones Stirpium rariorum*  
*et Descriptionibus illustratae.* Londini, 1791. fol. max.
- Scheuchz. gram.*—Joh. Scheuchzer, *Agrostographia, sive*  
*Graminum, Juncorum, Cyperorum, & isque affinum*  
*Historia.* Tiguri, 1719. 4to.
- Schk. car.*—C. Schkuhr, *Abbildungen der Riedgräser.* Wit-  
tenb. 1806. 8vo.
- Schkahr handb.*—Eiusdem *Botanisches Handbuch.* 3 Theile.  
Wittenberg, 1791—1803. 8vo.
- Schkahr filic.*—Eiusdem *Abbildungen der Farrnkräuter.* 4to.
- Schmidt arb.*—Franz Schmidt, *Oesterreichs Allgemeine*  
*Baumzucht.* Wien, 1792 et seq. fol.
- Schreb. gram.*—J. C. D. Schreber's *Beschreibung der Gräser.*  
Leipzig, 1769 et seq. fol.
- Scop. del. insub.*—J. A. Scopoli *Deliciæ Floræ et Faunæ In-*  
*subricæ.* Pars 1—3. Ticini, 1786—1788. fol.
- Sloan. hist. jam.* *Sloan. jam.*—*A Voyage to the Islands of*  
*Madeira, Barbadoes, Nevis, St. Christopher's and Ja-*  
*mica, with the Natural History of the last of those*  
*Islands, by Hans Sloane.* 2 vols. London, 1707—  
1725. fol.
- Smith brit.* *Smith fl. brit.*—J. E. Smith, *Flora Britannica.*  
Vol. 1—3. Londini, 1800—1804. 8vo.
- Smith exot. bot.*—Vid. *Exot. bot.*
- Smith ic. ined.*—*Plantarum Icones hactenus ineditæ, Auc-*  
*tore J. E. Smith.* Londini, 1789 et seq. fol.
- Smith ic. pict.*—Eiusdem *Icones pictæ Plantarum rariorum.*  
Londini, 1790 et seq. fol.
- Smith spicil.*—Eiusdem *Spicilegium Botanicum.* Londini,  
1791 et seq. fol.
- Sp. pl.*—Vid. *Linn. sp. pl.*
- Swartz fl. ind. occid.*—Ol. Swartz, *Flora Indiæ Occiden-*  
*alis.* Tom. 3. Erlangæ, 1797. 1806. 8vo.
- Swartz ic.*—Eiusdem *Icones Plantarum incognitarum, quas*  
*in India Occidentali detexit atque delineavit.* Fasc. 1, 2.  
Erlangæ, 1788. fol.
- Swartz observ.*—Eiusdem *Observationes Botanicæ.* Er-  
langæ, 1791. 8vo.
- Swartz prodr.*—Eiusdem *Prodromus Descriptionum Ve-*

- gefabilium quæ sub Itinere in Indiam Occidentalem  
digessit. Holm. 1788. 8vo.  
*Thuil. fl. par.*—Thuillier, La Flore des Environs de Paris.  
Paris, l'an 7. 8vo.  
*Trew ehr.*—Plantæ Selectæ, quarum Imagines pinxit G.  
D. Ehret, collegit et illustravit C. J. Trew. Norim-  
bergæ, 1750—1773. fol.  
*Trew pl. rar.*—Plantæ rariores, quas depingendas ærique  
incidentas curavit C. J. Trew. Norimbergæ, 1763. fol.  
*Usteri annal.*—Paul Usteri, Annalen der Botanik. 1—24  
Stuck. Zurich, 1792 et seq. 8vo.  
*Vahl. enum. pl.*—Mart. Vahlii Enumeratio Plantarum vel  
ab aliis vel ab ipso observatarum. Vol. 1, 2. Havniæ,  
1804, 1805. 8vo.  
*Vahl. symb.*—Eiusdem Symbolæ Botanicæ. Partes 3. Hav-  
niæ, 1790—1794. fol.  
*Vaill. par.*—Seb. Vaillant, Botanicum Parisiense. Leide et  
Amsterdam, 1727. fol.  
*Vandell. fasc.*—Domin. Vandelli Fasciculus Plantarum,  
cum novis Generibus et Speciebus. Olisip. 1771. 4to.  
*Vent. cels.* *Vent. hort. cels.* *Vent. jard. cels.*—Description  
des Plantes nouvelles et peu connues, cultivées dans le  
Jardin de J. M. Cels, avec Figures, par E. P. Ventenat.  
Paris, 1800. fol.  
*Vent. choix de plant.*—Eiusdem Choix de Plantes, dont la plus  
part sont cultivées dans le Jardin de Cels. Paris, 1803.  
*Vent. malmais.*—Eiusdem Jardin de la Malmaison. Paris,  
1803 et seq. fol.  
*Waldst. hung.*—Vid. *Pl. rar. hung.*  
*Walt. fl. car.*—Flora Caroliniana, Auctore Thom. Walter.  
Londini, 1788. 8vo.  
*Walth. hort.*—Designatio Plantarum quas Hortus Aug.  
Trid. Waltheri complectitur: accedunt novæ Plantarum  
Icones 24. Lipsiæ, 1735. 8vo.  
*Wangh. amer.*—F. A. J. von Wangenheim Anpflanzung  
North Amerikanischer Holzarten. Gotting. 1787. fol.  
*Wendl. hort. herrenh.*—J. C. Wendland, Hortus Herrenhu-  
sianus. Fasc. 1—4. Hannov. 1798—1801. fol.  
*Willd. amaranth.*—C. Willdenow, Historia Amaranthorum.  
Turici, 1790. fol.  
*Willd. enum.*—Eiusdem Enumeratio Plantarum Berolinens-  
ium. 8vo.  
*Willd. hort. ber.*—Eiusdem Hortus Berolinensis. Fasc. 1—3.  
Berol. 1803, 1804.  
*Willd. sp. pl.*—Eiusdem Species Plantarum. Berolini, 1797  
et seq. 8vo.

I.

1. ZOSTERA  
1-sperma.
2. CAULINI  
filiformis.  
*noicum.*
3. SALICORI  
*Stam.* 1—2.
4. HIPPURIS  
*Sem.* 1.
5. PODOSTE  
furcatum.  
2-loc. 2-v.
6. CALLITRI  
2-locularis.
7. BLITUM.
8. CORISPER

9. ZANNICH  
*Styl.* 3—5
10. CHARA. C

# I. MONANDRIA.

---

## I. MONOGYNIA.

1. ZOSTERA. *Cal. et Cor. nulla. Anth. sessilis. Caps. 1-sperma. Monoicum.*
2. CAULINIA. *Cal. et Cor. nulla. Anth. sessilis. Styl. filiformis. Stigm. 2-fidum. Caps. 1-sperma. Monoicum.*
3. SALICORNIA. *Cal. ventricosus, integer. Cor. o. Stam. 1-2. Sem. 1. in fundo calycis.*
4. HIPPURIS. *Cal. margo bilobus, superus. Cor. o. Sem. 1.*

## II. DIGYNIA.

5. PODOSTEMUM. *Cal. squamæ 2-3. Cor. o. Fil. furcatum. Anth. 2. Styl. o. Stigm. reflexa. Caps. 2-loc. 2-valv. polysperma.*
6. CALLITRICHÈ. *Cal. 2-partitus. Cor. o. Caps. 2-loctularis, 4-sperma.*
7. BLITUM. *Cal. 3-fidus, baccatus. Cor. o. Sem. 1.*
8. CORISPERMUM. *Cal. 2-partitus. Cor. o. Sem. 1.*

## III. POLYGYNIA.

9. ZANNICHELLIA. *Cal. 1-phyllus; turbinatus. Cor. o. Styl. 3-5. Stigm. peltata. Sem. 3-5. nuda.*
10. CHARA. *Cal. et Cor. o. Stigm. 5. Bac. polysperma.*

## I. MONOGYNIA.

1. ZOSTERA. *Gen. pl.* 1390. Grass-wrack.*marina.*

1. *Z. foliis integerrimis subtrinerviis, caule teretiusculo.*  
*Vahl. enum. pl. 1. p. 14.*

*Icon. Engl. lot. 467. Fl. dan. t. 15.*

In salt-marsh ditches, in the bay of Delaware, scarce.

 $\gamma$ . Aug. v. v. Leaves long; flowers very small.

The leaves are narrower and more obtuse than in the European plant; in every other respect it appears to be the same.

*fragilis.**flexilis.*2. CAULINIA. *Willd. in act. ac. ber.* 1798. p. 87.

1. *C. foliis ternis oppositivis linearis-subulatis recurvatis aculeato-dentatis rigidis.* *Willd. sp. pl. 4. p. 182.*

*Icon. Willd. l. c. t. 1. f. 2.—Michel. gen. t. 8. f. 3.*In slow-flowing waters; Susquehanna, Kenhaway, &c. rivers.  $\odot$ . Aug. v. v. Immersed, floating.

2. *C. foliis senis linearibus apice denticulatis patentibus.*  
*Willd. sp. pl. 4. p. 183.*

*Icon. Willd. l. c. t. 1. f. 1.*

In small ponds and old ditches: New York to Carolina.

 $\odot$ . Aug. v. v. Immersed, floating; flowers very small.*herbacea.*3. SALICORNIA. *Gen. pl.* 14. Samphire or Glasswort.*annua.*  
*simplex.*  
*gracilis.**virginica.*

1. *S. herbacea patula, articulis apice compressis exargi-nato-bifidis, spicis axillaribus oppositis pedunculatis, squamis obtusis.* *Willd. sp. pl. 1. p. 23.*

 $\alpha$  *Engl. bot. 415.* $\beta$  ramis simplicissimis, caule prostrato. $\gamma$  caule stricto ramoso, ramis laxis.Var.  $\alpha$  and  $\beta$  are common along the sea-coast: var.  $\gamma$  I observed on the Salt-lake of Onondago, New York.  
 $\odot$ .  $\delta$ . Aug. v. v. Fleshy, smooth; salt taste.

2. *S. herbacea erecta, ramis simplicissimis indivisis: spicis longis articulatis.* *Willd. sp. pl. 1. p. 25.*

Frequent on the coast of Virginia and Carolina.  $\odot$ . July, Aug. v. v.; v. s. in *Herbario Gronoviano in Museo Banksiano asservato*. Smaller than No. 1.  
I consider this no more than a sub-species, and probably the same with *S. herbacea*  $\gamma$ .

3. *S. fruticul amer. 1.*  
In sedgy sa July, At taller tha

## 4. HI

1. *H. foliis pl p. 1.*  
In ditches rare, 2 new spec

## 5. PODO

1. *Podostemus*  
Icon. *Mic*  
Closely att  
the rivers  
v. v.  
I take the li  
will be m  
this head,

## 6. CALLIT

1. *C. foliis sup nearibus C. verna.*  
Icon. *Fl. d*  
In springs  
July. v.  
floating.  
2. *C. foliis om C. autumnal*  
In similar  
June. v.  
somewhat  
3. *C. foliis omr matis.*

## MONANDRIA MONOGYNIA. Salicornia.

3

3. S. fruticulosa assurgens inordinate ramosa. *Mich. fl. amer. 1. p. 2.* *ambigua.*

In sedgy salt-meadows; New Jersey to Carolina. ♂. July, Aug. v. v. Resembles *S. fruticosa* very much; taller than either No. 1. or 2; pale green.

4. HIPPURIS. *Gen. pl. 15.* Mare's-tail.

1. H. foliis plerumque senis linearibus. *Mich. fl. amer. 1. p. 1.* *vulgaris.*

In ditches and small ponds: Canada to Pennsylvania, rare. U. v. s. s. fl. Upright, simple; may be a new species.

## II. DIGYNIA.

5. PODOSTEMUM. *Mich. fl. amer. 2. p. 164.*

1. Podostemum. *Mich. l. c. p. 165.*

*ceratophyl-*  
*lum.*

Icon. *Mich. l. c. t. 44.* Closely attached to loose stones in shallow water of the rivers Ohio, Holston, Kenhaway, &c. ♀. July. v. v.

I take the liberty to place this plant in Monandria, as it will be more easily found by a young Botanist under this head, than under *Monococcia Diandria*.

6. CALLITRICHE. *Gen. pl. 17.* Water-starwort.

1. C. foliis superioribus spathulato-obovatis, inferioribus li- *heterophylla.*  
nearibus obtusis emarginatisque.

C. verna. *Willd. sp. pl. 1. p. 28?*

Icon. *Fl. dan. 129?*

In springs and rivulets, very common. ⊖ May—  
July. v. v. Flowers white, axillary; upper leaves floating.

2. C. foliis omnibus linearibus apice bifidis.

*linearis.*

C. autumnalis. *Willd. sp. pl. 1. p. 29?*

In similar situations, not so common. ⊖ May,  
June. v. v. More slender than No. 1. Upper leaves somewhat spathulate-linear.

3. C. foliis omnibus linearibus truncatis brevibus approxi-  
matis.

*brevifolia.*

4

**MONANDRIA DIGYNIA.** Callitrichæ.

In sandy barren grounds, near mill-ponds, occasionally overflowed: New York to Virginia. ☽. Aug. v. v. Small, creeping along the ground, with a moss-like appearance; flowering seldom.

*capitatum.*

7. BLITUM. *Gen. pl. 18.* Strawberry-spinage.

1. B. foliis acute triangulo-hastatis, capitulis alternis in racemum superne aphyllum approximatis. *Mich. fl. amer. 1. p. 2.* *Willd. sp. pl. 1. p. 30.*  
In woods and about houses, in Canada and the western parts of New York and Virginia. ☽. June, July. v. v. Flowers herbaceous; berries red.

*hyssopifoli-*  
*uni.*

8. CORISPERMUM. *Gen. pl. 16.* Tick-seed.

1. C. spicis terminalibus, foliis linearibus enerviis muticis. *Willd. sp. pl. 1. p. 28.*  
Icon. *Pall. fl. ross. 2. t. 98.*  
In open plains, called Prairies, of Canada and on the banks of the Missouri River; found by M. Lewis, Esq. ☽. Aug. Sept. v. s. in *Herbario Lewisiano*. Small shrub, of a gray aspect: flowers small, in axillary clusters.

---

**III. POLYGYNIA.**

9. ZANNICHELLIA. *Gen. pl. 1391.*

*palustris.*

1. Z. anthera quadriloculari, stigmatibus integerrimis, seminibus dorso denticulatis. *Willd. sp. pl. 4. p. 181.*  
Icon. *Michel. gen. t. 34. f. 1.*  
In ditches and small rivulets, in Virginia near the Sweet springs. ☽. July, Aug. v. v. Resembles *Potamogeton marinum*; flowers very small.

*vulgaris.*

10. CHARA. *Gen. pl. 1397.*

1. C. caulis ramulisque basi nudis, ramulis teretibus, articulis foliosis, foliolis oblongo-subulatis, bracteis bacca brevioribus. *Willd. sp. pl. 4. p. 183.*  
Icon. *Engl. bot. 336.* *Feather-beds.*  
Floating in old ponds and ditches: Canada to Carolina, common. ☽. July, Aug. v. v. Incrusted; smells disagreeably.

2. C. caulis  
basi et  
bacca bre  
Icon. *Wu*  
In clear wa  
&c. ☽  
3. C. caulis  
lis comp  
pl. 4. p.  
Icon. *Eng*  
In muddy p  
sylvania,  
der, trans

- occasionally  
Aug. v.v.  
a moss-like
- spinage.
- ternis in ra-  
Mich. fl.
- the western  
June, July.
- ck-seed.
- viis muticis.
- and on the  
M. Lewis,  
*Lewisiano*.  
nall, in axil-
- 
- 1.
- errimis, se-  
. 4. p. 181.
- ar the Sweet  
ables *Pota*-
- teretibus, ar-  
bracteis bac-
- Feather-beds*.  
to Carolina,  
ited; smells
2. C. caulis nudis superne papillosis, ramulis teretibus  
basi et in articulis foliosis, foliis linearibus, bracteis  
bacca brevioribus. *Willd. sp. pl. 4. p. 184.*  
Icon. *Willd. in act. ac. ber. t. 1. f. 2.*  
In clear waters and small ponds: Pennsylvania, Virginia,  
&c. ☺. July. v.v. Green, shining, fragile.
3. C. caulis diaphanis nudis, ramulis inarticulatis aphyll-  
lis compressis, baccis nudis lateribus. *Willd. sp.*  
*pl. 4. p. 187.*  
Icon. *Engl. bot. 1070.*  
In muddy pools and ditches in the western part of Pen-  
sylvania, &c. ☺. Aug. v.v. Green, flexible, ten-  
der, transparent.

*foliolosa.**flexilis.*

## II. DIANDRIA.

### I. MONOGYNIA.

\* *Fl. completi, inferi, 1-pet. regulares.*

12. OLEA. *Cor. 4-fida : laciniis subovatis. Drupa 1-sperma.*
13. CHIONANTHUS. *Cor. 4-fida : laciniis longissimis. Drupa 1-sperma : nuce striata.*
14. ORNUS. *Cor. 4-partita : petalis longis ligulatis. Fil. longa. Nux alata.*
11. LIGUSTRUM. *)Cor. 4-fida. Bac. 2-loc. : loculis 2-spermis.*

\*\* *Fl. completi, inferi, 1-pet. irregulares, angiospermi.*

18. VERONICA. *Cor. limbo 4-partito : lacinia inferiore angustiore. Caps. 2-loc.*
  17. MICRANTHEMUM. *Cor. subcampanulata, 4-partita : lacinia superiore minore. Caps. 2-valv. 1-loc. polysperma.*
  19. GRATIOLA. *Cor. irregularis, resupinata. Stam. 2. sterilia. Stigm. 2-lob. Caps. 2-loc.*
  16. CATALPA. *Cor. irregularis, 5-fida. Cal. 2-phyllo. Caps. 2-locularis.*
  20. ELYTRARIA. *Cor. 5-fida, subæqualis. Stam. 2. sterilia. Caps. 2-loc.*
  21. JUSTICIA. *Cor. irregularis. Caps. 2-loc. 2-valvis. Anth. geminæ.*
  22. PINGUICULA. *Cor. ringens, calcarata. Caps. 1-loc. Cal. 2-lab. 5-fidus.*
  23. UTRICULARIA. *Cor. ringens, calcarata. Caps. 1-loc. Cal. 2-phyllo, æqualis.*
- \*\*\* *Fl. completi, inferi, 1-pet. gymnospermi.*
24. LYCOPUS. *Cor. tubulosa, subæqualis, 4-fida : lacinia superiore latiore emarginata. Sem. 4. retusa.*
  25. MONARDA. *Cor. ringens : labium superius lineare, filamenta involvens.*

26. SALVIA.  
affixa.

27. COLLIN.  
multifl  
orienti

28. CIRCÆ.  
spermii

15. FRAXIN.  
Polyga

29. CERATI.

30. BORYA.

31. LEMNA.  
spermia

11. LIG.

1. L. foliis 1  
Willd. s  
Icon. En  
In hedges  
woods :  
Flowers

12.

1. O. foliis la  
tatis, br  
Willd. s  
Icon. Ca  
On the sea  
rida, &c  
white ;

13. CHIO

1. C. panicul  
acutis.

## DIANDRIA MONOGYNIA.

7

26. SALVIA. *Cor. ringens. Filam. transverse pedicello affixa.*  
 27. COLLINSONIA. *Cor. inæqualis: labio inferiore multifido, capillari. Stam. 2—4. Sem. 4. 3. abortientia.*  
 } *\*\*\*\* Fl. completi, superi.*  
 28. CIRCÆA. *Cor. 2-petala. Caps. 2-loc. : loculis 1-spermis.*  
 } *\*\*\*\*\* Fl. incompleti.*

15. FRAXINUS. *Cor. 0. vel 4-pet. Samara 2-loc. Sem. 1. Polygamum.*  
 29. CERATIOLA. *Cal. gemma. Cor. 0. Bac. 2-sperma.*  
 30. BORYA. *Cal. 4-phyllus. Cor. 0. Bac. 1-sperma.*  
 31. LEMNA. *Cal. 1-phyllus. Cor. 0. Caps. 1-loc. 2-sperma.*
- 

## I. MONOGYNIA.

11. LIGUSTRUM. *Gen. pl. 23. Privet. Prim.*  
 1. *L. foliis lanceolatis acutiusculis, panicula coarctata. Willd. sp. pl. 1. p. 41.* *vulgare.*  
*Icon. Engl. bot. 764.*  
*In hedges on road sides, frequent; scarce in remote woods: Canada to Virginia. H. May—July. v. v. Flowers in white panicles; berries black.*
12. OLEA. *Gen. pl. 25. Olive-tree.*  
 1. *O. foliis lanceolato-ellipticis integerrimis, racemis angustatis, bracteis omnibus persistentibus connatis parvis. Willd. sp. pl. 1. p. 45.* *americana.*  
*Icon. Catesb. car. 1. t. 61. Devil's-wood.*  
*On the sea-coast, in sandy boggy places: Carolina, Florida, &c. H. April, May. v. v. Flowers small, white; berries globular, purple, eatable.*
13. CHIONANTHUS. *Gen. pl. 26. Fringe-tree.*  
 1. *C. panicula terminali trifida, pedunculis trifloris, foliis acutis. Willd. sp. pl. 1. p. 46.* *virginica.*

- Icon. *Catesb. car.* 1. t. 68.  
*montanus.*  $\alpha$  foliis ovali lanceolatis coriaceis glabris, paniculis densis, drupis ovalibus.  
*maritimus.*  $\beta$  foliis obovato-lanceolatis membranaceis pubescentibus, paniculis laxissimis, drupis ellipticis.  
*C. v. angustifolia* *Ait. kew.* 1. p. 23.  
 Var.  $\alpha$  in the mountains of Pennsylvania and Carolina.  
 $\beta$  on the sea-coast in boggy woods from New Jersey to Carolina.  $\frac{1}{2}$ . May, June. v. v. Flowers white, very handsome; berries purplish-blue.  
 It is probable that those, as yet considered subspecies, are really distinct species, which further observations will decide.

14. ORNUS. *Dalech.* Flowering-ash.

- americana.* 1. O. foliolis late ovatis serratis: impari fere obcordato. *Pers. enc.* 2. p. 205.  
 In moist shady woods: Maryland and Virginia, rare.  $\frac{1}{2}$ . May. v. v. Flowers resembling those of *Chionanthus*; seeds small, winged.

15. FRAXINUS. *Gen. pl.* 1597. Ash-tree.

- sambucifolia.* 1. F. foliolis sessilibus ovato-lanceolatis serratis rugoso-nitidis, basi rotundatis inaequalibus, axillis venarum subtus villosis, floribus nudis. *Willd. sp. pl.* 4. p. 1099.  
 Icon. *Mich. f. arb.* t.... *Black-ash, Water-ash.*  
 In stony soil, on river sides from Canada to Virginia.  $\frac{1}{2}$ . April. v. v.  
*quadrangulata.* 2. F. foliolis subsessilibus lanceolato-ellipticis serratis, subtus pubescentibus, samaris utrinque obtusis. *Willd. sp. pl.* 4. p. 1102.  
 Icon. *Mich. f. arb.* t.... *Blue-ash.*  
 In rich bottoms of the western states: Ohio, Kentucky and Tennessee.  $\frac{1}{2}$ . May. v. v. s. fl.; v. s. c. fr.  
*epiptera.* 3. F. foliolis lanceolato-ellipticis subserratis, samaris cuneatis apice obtusis emarginatis inferne teretibus. *Vahl. enum. pl.* 1. p. 50.  
 F. discolor. *Mich. f. arb.* 33.  
 F. canadensis. *Gärt. de fr. et sem.* 1. p. 222: t. 49.  
 Icon. *Mich. f. arb.* t....  
 On water-sides and rich bottom lands, from Canada to Carolina, frequent.  $\frac{1}{2}$ . April. v. v.

4. F. foliolis mis sub cycl. 2. |  
 F. america F. concolo  
 Icon. *Mi* Common i April.  
 5. F. foliolis ramulis p. 1103.  
 F. serratifol F. excelsior  
 Icon. *Mi* In rocky si  $\frac{1}{2}$ . May  
 6. F. foliolis lanceolat Icon. *Ca* In boggy May.  
 7. F. foliolis ramulisq sp. pl. 4 encycl. 2 F. nigra.  
 F. tomento Icon. *Mi*  
 $\beta$  foliolis ova  $\gamma$  foliolis ova  
 ratis.  $\delta$  foliolis ellip ferne sub  
 Pers. enc Common in  $\frac{1}{2}$ . April  
 8. F. foliolis p axillaris ve bus calyc F. carolinia  
 Icon. *Mic* In shady w chiefly in

4. *F. foliolis petiolatis oblongis nitidis acuminatis integerimis subtus glaucis, floribus calyculatis.* *Lam. encycl. 2. p. 542.*  
*F. americana.* *Willd. sp. pl. 4. p. 1102.*  
*F. concolor.* *Mich. f. arb. 33.*  
*Icon. Mich. l. c. . . . .* *White-ash. Green-ash.*  
*Common in most woods from Canada to Carolina.*  $\frac{1}{2}$ .  
*April. v. v.*
5. *F. foliolis petiolatis lanceolatis serrulatis nitidis glabris, caroliniana.*  
*ramulis glabris, floribus calyculatis.* *Willd. sp. pl. 4. p. 1103.*  
*F. serratifolia.* *Mich. f. arb. p. 33.*  
*F. excelsior.* *Walt. fl. car. p. 254.*  
*Icon. Mich. f. arb. t. . . . .* *Shining-ash.*  
*In rocky situations, from Pennsylvania to Carolina, scarce.*  $\frac{1}{2}$ .  
*May. v. v. s. fl.*
6. *F. foliolis subsessilibus extrorsum serratis samarisque platycarpa.*  
*lanceolato-ellipticis.* *Willd. sp. pl. 4. p. 1103.*  
*Icon. Catesb. car. t. 80.* *Mich. f. arb. t. . . . .*  
*In boggy woods of Virginia and Carolina.*  $\frac{1}{2}$ .  
*April, May. v. s.*
7. *F. foliolis petiolatis elliptico-ovatis serratis subtus petiolis ramulisque tomentosis, floribus calyculatis.* *Willd. sp. pl. 4. p. 1103.* *Walt. fl. car. p. 234.* *Lam. encycl. 2. p. 543.*  
*F. nigra.* *Du Roi, ed. 2. t. 1. p. 398.*  
*F. tomentosa.* *Mich. f. arb. p. 33.*  
*Icon. Mich. l. c. t. . . . .* *Red-ash. Black-ash.*  
 $\beta$  *foliolis ovato-lanceolatis attenuatis subserratis.*  
 $\gamma$  *foliolis ovatis latioribus obsolete ad medium usque serratis.*  
 $\delta$  *foliolis elliptico oblongis acuminatis argute serratis inferne subpubescentibus, petiolo communis glaberrimo.*  
*Pers. encycl. 2. p. 605.*  
*Common in the woods of Canada and the United States.*  $\frac{1}{2}$ .  
*April, May. v. v.*
8. *F. foliolis petiolatis ovatis opacis serratis subtus glaucis juglandifolia.*  
*axillæ venarum pubescentibus, ramulis glabris, floribus calyculatis.* *Willd. sp. pl. 4. p. 1104.*  
*F. caroliniana.* *Wangh. amer. 81.*  
*Icon. Mich. f. arb. t. . . . .* *Western Black-ash.*  
*In shady wet woods, from Canada to North Carolina; chiefly in the western districts.*  $\frac{1}{2}$ .  
*May. v. v.*

## DIANDRIA MONOGYNIA.

16. CATALPA. *Juss. gen. 138. Catalpa-tree.**syringæfolia.*

1. *C. foliis cordatis planis.* *Sims in bot. mag. 1094.*  
*Bignonia Catalpa.* *Willd. sp. pl. 3. p. 289.*  
*C. bignonoides.* *Walt. fl. car. p. 64.*

*Icon. Schmidt, arb. 1. t. 41.*

On the banks of Ohio, Mississippi and Delaware, in  
gravelly soil.  $\frac{1}{2}$ . July, Aug. *v. v.* Flowers white,  
speckled with purple and yellow.

17. MICRANTHEMUM. *Mich. fl. amer. 1. p. 10.**orbiculatum.*

1. *Micranthemum.* *Mich. l. c.*  
*Anonymos umbrosa.* *Walt. fl. car. p. 63.*  
*Globifera umbrosa.* *Gmel. syst. nat. t. 2. p. 1. p. 32.*  
*Icon. Mich. l. c. t. 2.*
- In shady wet woods of Carolina and Georgia.  $\frac{1}{4}$ . Aug.  
*v. v.* Stem creeping; leaves orbicular; flowers axillary, very small.

18 VERONICA. *Gen. pl. 32. Speedwell.**virginica.*

1. *V. spicis terminalibus, foliis quaternis quiniske.* *Willd. sp. pl. 1. p. 34.*  
*Icon. Pluk. alm. t. 70. f. 2.*  
On calcareous hills, in sunny exposures.  $\frac{1}{4}$ . July—  
Sept. *v. v.* Spikes long, white, or blush-colour.  
On the mountains of Virginia I observed a very tall-growing variety, with purple flowers, extremely beautiful.

*officinalis.*

2. *V. spicis lateralibus pedunculatis, foliis oppositis obovato-subrotundis pilosis annuis, caule procumbente hirto.* *Willd. sp. pl. 1. p. 59.*  
*Icon. Engl. bot. 765. Fl. dan. 248.*  
In barren, dry woods, very common from Canada to  
Carolina.  $\frac{1}{4}$ . May—July. *v. v.*

*reniformis.*

3. *V. spicis lateralibus pedunculatis, foliis oppositis reniformi-cordatis inciso-crenatis, caule repente.*  
Collected by Messrs. Lewis and Clark in boggy soil, on  
the banks of the Missouri.  $\frac{1}{4}$ . June. *v. s. in Herb. Lewis.*  
*Caulis repens, filiformis, ad geniculos radicans. Folia opposita, longe petiolata, reniformi-cordata, inciso-crenata, glabra. Pedunculi axillares, alterni, longitudine foliorum, teretes, glabri, superne unibracteati. Bractea oblonga, crenata. Spica oblonga.*

DIA

*brevis.*  $\frac{1}{4}$   
4-fidus : 1  
bus linear  
simus.  $\frac{1}{2}$   
lyce triplo  
corollæ.

4. *V. racemo t*  
natis. *W*  
*Icon. Fl. a*  
On road sic  
—Aug.  
stripes.

5. *V. racemis 1*  
*Willd. sp*  
*Icon. Engl*  
In rivulets  
 $\frac{1}{4}$ . June,  
flowers la

6. *V. racemis*  
erecto.  $\frac{1}{2}$   
*Icon. Engl*  
In ditches at  
rolina.  $\frac{1}{2}$

7. *V. racemis 1*  
linearibus.  
*Icon. Engl*  
Var. foliis in  
In wet mead  
The Americ  
European

8. *V. floribus*  
hrito. *W*  
*Icon. Engl*  
Common in  
Sept. *v.*

9. *V. floribus*  
integerrim  
*Vahl. enu*  
*V. romana.*  
*V. maryland*  
*V. caroliniae*  
*V. carnosula*  
*Icon. Mur*

- brevis. *Flores magni, conferti, pallide cærulei. Cal. 4-fidus : laciniis 2. superioribus oblongis; 2. inferioribus linearibus, multo minoribus. Cor. Tubus brevissimus. Limbus planus : laciniis oblongis, acutis, calyce triplo longioribus, infima linearis. Fil. longitudine corollæ.*
4. V. racemo terminali subspicato, foliis ovatis glabris cre- *serpyllifolia*.  
natis. *Willd. sp. pl. 1. p. 64.*  
Icon. *Fl. dan. 492.*  
On road sides and in dry woods, common. ♀. May—  
Aug. v. v. Creeping; flowers pale with blue  
stripes.
5. V. racemis lateralibus, foliis ovatis planis, caule repente. *Beccabunga.*  
*Willd. sp. pl. 1. p. 64.*  
Icon. *Engl. bot. 655. Fl. dan. 511.*  
In rivulets and springs; Pennsylvania to Virginia, rare.  
♀. June, July. v. v. Leaves shining green, blunt;  
flowers large, blue.
6. V. racemis lateralibus, foliis lanceolatis serratis, caule  
erecto. *Willd. sp. pl. 1. p. 65.*  
Icon. *Engl. bot. 781. Fl. dan. 903.*  
In ditches and rivulets, more common: Canada to Ca-  
rolina. ♀. June—Aug. v. v. Leaves acute. *Anagallis.*
7. V. racemis lateralibus alternis : pedicellis pendulis, foliis  
linearibus. *Willd. sp. pl. 1. p. 65.*  
Icon. *Engl. bot. 782. Fl. dan. 209.*  
Var. foliis integerrimis et serratis.  
In wet meadows on gravelly soil. ♀. July. v. v.  
The American plant has longer leaves than any of the  
European specimens I have seen.
8. V. floribus solitariis foliisque cordatis subsessilibus, caule  
hirtio. *Willd. sp. pl. 1. p. 73.*  
Icon. *Engl. bot. 734. Fl. dan. 515.*  
Common in fields and cultivated grounds. ♂. May—  
Sept. v. v. Flowers blue, small. \**arvensis.*
9. V. floribus sessilibus, foliis linear-lanceolatis dentatis  
integerrimisque, caule erecto. *Willd. sp. pl. 1. p. 76.*  
*Vahl. enum. pl. 1. p. 85.*  
V. romana. *Linn. mant. 317.*  
V. marylandica. *Murr. in com. gæt. 1782. p. 11, t. 3.*  
V. caroliniana. *Walt. fl. car. p. 61.*  
V. carnosula. *Lam. illustr. 1. p. 47.*  
Icon. *Murr. l. c. Fl. dan. 407.* *peregrina.*



In cultivated ground, common. ♂. May—Aug. v. v.; v. s. in *Herbario Walteriano*, necnon *Gronoviano*. Very variable in habit. Flowers small, white.

19. GRATIOLA. *Gen. pl. 37.*

*aurea.*

1. G. foliis lato-linearibus sessilibus integerrimis 3 nerviis supra punctatis, pedunculis oppositis vix longitudine foliorum; filamenta castrata nullis.  
*G. officinalis.* *Mich. fl. amer.* 1. p. 6.  
*G. carolinensis.* *Pers. enc.* 1. p. 14.  
In sandy wet places, in the pine-barrens of New England, New Jersey and Carolina. ♀. Aug. v. v.; v. s. in *Herbario Banksiano*. *Golden-pert.*  
*Caulis* bisulcatus, quadrangulus, basi ramosus. *Folia* basi dilatata, sessilia, acuta, glabra. *Pedunculi* longitudine foliorum, villosi. *Cal.* laciinis linearibus, apice glandulosis. *Cor. aurea.* *Tubus* hirsutus, medio ventricosus, obliquus. *Lab. superius* subrotundum, emarginatum: *inferius* æqualiter trifidum, laciinis oblongis: intermedia emarginata. *Faux* hirsuta. *Fil.* 2. longitudine tubi corollæ. *Stylus* staminibus longior. *Stigma* infundibuliforme, obliquum. *Capsula* ovata, vix longitudine calycis.

*virginica.*

2. G. foliis obovato-lanceolatis inferne attenuatis remote dentatis nervosis glabris, pedunculis alternis brevissimis, capsulis acuminatis calyce longioribus.

*G. foliis lanceolatis obtusis subdentatis.* *Gron. virg.* 3.  
*G. virginica.* *Willd. sp. pl.* 1. p. 104.  
*G. acuminata.* *Vahl. enum. pl.* 1. p. 92. *exclus. syn.*  
In old ditches and pools from Pennsylvania to Carolina. ♀. June, July. v. v.; v. s. in *Herb. Gronov.* Stem rooting at the joints; flowers yellowish white.

*quadridentata.*

3. G. foliis lineari lanceolatis extrosum remote paucidentatis, pedunculis longitudine foliorum, capsula calyce subulato triplo breviore. *Mich. fl. amer.* 1. p. 6.

*G. ramosa.* *Walt. fl. car.* p. 61.  
*G. virginica.* *Lam. illustr.* 1. p. 54. t. 16. f. 2. *Vahl. enum. pl.* 1. p. 89.  
In sandy wet fields, edges of ponds, &c. Carolina. ♀. June—Aug. v. v.; v. s. in *Herb. Walt.*

*acuminata.*

4. G. foliis lanceolatis serratis pubescentibus, pedunculis oppositis folio longioribus, calycis foliolis linearibus longitudine capsulae subglobosæ.

*G. acuminata*  
In ditches  
July, Au  
pale yello  
5. *G. foliis cor*  
cauleque  
*G. peruvian*  
In pine-bar  
August.  
*G. anagalloi*  
*G. repens* S

20. ELY

1. *E. squamis f*  
lanceolati  
p. 106. I  
*Justicia acar*  
Anonymos c  
*Tubiflora ca*  
*Icon. Mic*  
In wet sand  
Aug. v.

2

1. *J. spicis axi*  
gatis alte  
p. 146. su  
*Dianthera ar*  
*Dianthera et*  
*Justicia line*  
*Icon. Pluk*  
In bays and  
rence, Os  
v. v.; v.

2. *J. spicis axil*  
distantibu  
longis. I  
*Dianthera or*  
In swamps  
v. s. in *H*

3. *J. pedunculi*  
cellis gen  
acutis, pe  
sissimo.

- G. acuminata. *Walt. fl. car. p. 61.* nec *Vahl. enum.*  
In ditches and pools: Pennsylvania to Carolina. ♀.  
July, Aug. v. v.; v. s. in *Herb. Walt.* Flowers  
pale yellow, large. Caps. larger than any other.
5. G. foliis cordato-ovatis, floribus axillaribus subsessilibus  
cauleque erecto pilosis. *Vahl. enum. pl. 1. p. 94.* *pilosa.*
- G. peruviana. *Walt. fl. car. p. 62.*  
In pine-barrens: Carolina and Georgia. ♀. June—  
August. v. v.; v. s. in *Herb. Walt. et Banks.*  
G. anagalloidea *Mich. &c. vid. Lindernia.*  
G. repens *Swartz. et Monnieria Mich. vid. Herpestis.*

20. ELYTRARIA. *Mich. fl. amer. 1. p. 8.*

1. E. squamis floriferis ovatis margine villosis, scapi foliisque  
lanceolatis integerrimis glabris. *Vahl. enum. pl. 1.* *virgata.*  
*p. 106. Mich. fl. amer. 1. p. 9.*  
*Justicia acaulis* γ. *Willd. sp. pl. 1. p. 84.*  
*Anonymos carolinensis.* *Walt. fl. car. 60.*  
*Tubiflora carolinensis.* *Gmel. syst. nat. 2. p. 1, 27.*  
Icon. *Mich. l. c. t. 1.*  
In wet sandy woods of Lower Carolina. ♀. July,  
Aug. v. v. Flowers white, imbricated.

21. JUSTICIA. *Gen. pl. 35.*

1. J. spicis axillaribus, floribus confertis, pedunculis elongatis alternis, foliis lanceolatis. *Vahl. enum. pl. 1.* *pedunculosa.*  
*p. 146. sub J. americana. Mich. fl. amer. 1. p. 7.*  
*Dianthera americana.* *Linn. syst. veg. 64.*  
*Dianthera ensiformis.* *Walt. fl. amer. 63.*  
*Justicia linearifolia.* *Lam. illustr. 1. p. 41.*  
Icon. *Pluk. alm. t. 423. f. 5.* *Water-willow.*  
In bays and slow-flowing waters of the rivers St. Lawrence, Oswego, Ohio, Kenhaway, &c. July, Aug.  
v. v.; v. s. in *Herb. Walt. et Gronov.*
2. J. spicis axillaribus terminalibusque paucifloris, floribus distantibus, bracteis linearibus, foliis lanceolato-oblongis. *Vahl. enum. pl. 1. p. 143.* *humilis.*  
*Dianthera ovata.* *Walt. fl. car. p. 63.*  
In swamps of Carolina and Florida. ♀. July, Aug.  
v. s. in *Herb. Walt.*
3. J. pedunculis axillaribus racemoso-vorticillatis: pedicellis geminis, bracteis obovatis trifloris, foliis ovatis acutis, petiolis longissimis, caule sexangulari ramosissimo. *brachiata.*

On the islands of Roanoak river in North Carolina. 4.  
July, Aug. v. v.

*Caulis* plures, geniculati, hexagoni: angulis oppositis sulcatis, brachiato-ramosissimi, glabri. *Folia* opposita, longe petiolata, ovato-oblonga, sensim acuta, integerrima, glabriuscula; nervis, venis, petiolisque tenui pilosa. *Ramuli floriferi* axillares, breves, foliosi. *Pedunculi* axillares gemini vel terni. *Bracteæ* duæ, obovatæ, altera minor. *Flores* 2-3. sessiles, rubelli. *Cal.* simplex, 5-fidus: laciniis subulatis. *Cor.* extus pilosa. *Tubus* tortuosus. *Limbus* resupinatus, 2-labiatus: labium superius obliquum, oblongum, acutum, plerumque emarginatum: *inferius* truncatum, 3-4-dentatum, fauce striatum. *Antheræ* 2. in singulo filamento. *Stigm.* bifidum. *Caps.* oblonga, compressa, mucronata. *Sem.* compressa, hispida, atrofusca.

22. PINGUICULA. Gen. pl. 40.

*elatior.*

1. P. nectario subulato obtuso corolla breviore, superne tubo ventricoso, scapo inferne villosso. *Vahl. enum. pl. 1. p. 19.* *Mich. fl. amer. 1. p. 11.*

*P. cærulea.* *Walt. fl. car. p. 63.*

In open swamps on the sands of Carolina and Georgia. 4. May. v. v. Flowers of a beautiful amethystine colour.

*lutea.*

2. P. nectario subulato recurvo corolla campanulata breviore, fauce barbata, labiis dentatis, scapo subvilloso. *Vahl. enum. pl. 1. p. 93.* *Walt. fl. car. p. 63.* *Mich. fl. amer. 1. p. 11.*

*P. campanulata.* *Lam. in diar. hist. nat. 1. n. 9. p. 336. t. 18. f. 1.*

In pine-barrens of Lower Carolina. 4. April, May. v. s. Flowers yellow; leaves obovate.

*pumila.*

3. P. nectario tubo breviore, corolla oblongiuscula tubulosa, scapo brevi glabello. *Mich. fl. amer. 1. p. 11.*

In open swamps of Georgia. 4. May. +. Flowers small, purple.

*acutifolia.*

4. P. glaberrima, foliis erectis ovalibus acutissimis. *Vahl. enum. pl. 1. p. 194.* *Mich. fl. amer. 1. p. 11.*

In shady woods, near rivers about Lake Mistassins. 4. July. +.

23.

1. U. scapi foliaceis. *V.* p. 12.

*U. inflata.* Floating in Carolina. Flowers yellow.

2. U. nectario guttidine p. 198. *Icon. Fl. a* In stagnant and Pens yellow, v

3. U. nectario *Vahl. enum.* *Icon. Fl. a* In swamps Jersey. ]

4. U. nectario *Vahl. enum.* In morasses v. s. in H

5. U. nectario scapo subt p. 200. *U. pumila.* On the marg in Herb. II

6. U. nectario c floro. *U. nectario s* *U. setacea.* ] p. 201.

In sandy wet to Carolina in Herb. G

7. U. nectario datis, scap Floating on th

Carolina. 2.

gulis oppositis  
*Folia* oppositissim acuta, in-  
 petiolisque te-  
 breves, foliosi.  
*Bracteæ* duæ,  
 ssiles, rubelli.  
 i. *Cor.* extus  
 pinatus, 2-la-  
 longum, acu-  
 s truncatum,  
 e 2. in singu-  
 blonga, com-  
 hispida, atro-

0.

ore, superne  
*Vahl. enum.*

and Georgia.  
 amethystine

lata breviore,  
 illoso. *Vahl.*  
 . *Mich. fl.*

n. 9. p. 336.

April, May.

ascule tubu-  
 . 1. p. 11.

. Flowers

imis. *Vahl.*

11.

Mistassins.

23. UTRICULARIA. *Gen. pl. 41.*

1. U. scapi foliis vesiculosus cylindricis divisis apice ramen- *ceratophylla.*  
*taceis. Vahl. enum. pl. 1. p. 197. Mich. fl. amer. 1.*  
*p. 12.*  
*U. inflata. Walt. fl. car. p. 64.*  
 Floating in ponds and lakes of Virginia and Lower Ca-  
 rolina. June, July. v. v.; v. s. in *Herb. Walt.*  
 Flowers yellow.
2. U. nectario conico, labio superiore lateribus reflexo lon-  
 gitudine palati, scapo stricto. *Vahl. enum. pl. 1.*  
*p. 198. Willd. sp. pl. 1. p. 112.*  
*Icon. Fl. dan. 138.*  
 In stagnant waters in the western parts of New York  
 and Pennsylvania. 2. July. v. v. Flowers large,  
 yellow, with red veins.
3. U. nectario gibbo carinato, corollarum fave aperta. *vulgaris.*  
*Vahl. enum. pl. 1. p. 199. Willd. sp. pl. 1. p. 112.*  
*Icon. Fl. dan. 128.*  
 In swamps and ditches, in the pine-barrens of New  
 Jersey. 2. Aug. v. v. Flowers small, deep yellow.
4. U. nectario obtuso, scapo subunifloro foliisque setaceis. *minor.*  
*Vahl. enum. pl. 1. p. 199. Walt. fl. car. p. 64.*  
 In morasses of the pine-barrens of Carolina. July.  
*v. s. in Herb. Walt.* Stems purple; flowers orange.
5. U. nectario subulato recto labio superiore subæquante,  
 scapo subbifloro, foliis setaceis. *biflora.*  
*Vahl. enum. pl. 1. p. 200. Lam. illustr. 1. p. 50.*  
*U. pumila. Walt. fl. car. p. 64.*  
 On the margin of ponds in Lower Carolina. July. v. s.  
*in Herb. Walt.* Flowers small, yellow.
6. U. nectario obtuso labio inferiore breviore; scapo subbi-  
 flor. *subulata.*  
*U. nectario subulato. Gronov. virg. 5. exclus. syn. Clayt.*  
*U. setacea. Mich. fl. amer. 1. p. 12. Vahl enum. pl. 1.*  
*p. 201.*  
 In sandy wet places near ponds and rivers, from Canada  
 to Carolina, common. ①. July, Aug. v. v.; v. s.  
*in Herb. Gronov.* Flowers small, bright yellow.
7. U. nectario carinato brevissimo, laciiniis corollæ rotun-  
 datis, scapo 2-3-floro. *purpurea.*  
*Walt. fl. car. p. 64.*  
 Floating on the lakes and ponds in Carolina; I observed

## DIANDRIA MONOGYNIA. Utricularia.

it likewise in Pennsylvania, on the Broad-mountains.  
♂. Aug. v. v. Flowers bright purple, small.

- cornuta*. 8. U. nectario subulato porrecto, corollæ labium inferius amplissimum, scapo subsessiliter bifloro. *Mich. fl. amer. 1. p. 12.*

Near mountain lakes from Canada to Virginia. July.  
v. v. Flowers large, bright yellow.

- gibba*. 9. U. nectario gibboso, laciniis corollæ rotundatis, scapo subunifloro. *Willd. sp. pl. 1. p. 113.*  
In boggy soil of New Jersey and Carolina. July. v. v.;  
v. s. in *Herb. Gronov.* Very small; flowers yellow.

24. LYCOPUS. *Gen. pl. 44.* Water-horehound.

- europaeus*. 1. L. foliis inferioribus incisis, superioribus lanceolatis serratis, calycibus subspinescenti-acuminatis. *Mich. fl. amer. 1. p. 14.* *Willd. sp. pl. 1. p. 120.*

Near rivers and ponds from Canada to Virginia. ♀.  
July, Aug. v. v. Flowers white, small, in whorls.

- virginicus*. 2. L. foliis lato-lanceolatis serratis basi integra angustatis, calycibus brevissimis muticis. *Mich. fl. amer. 1. p. 14.*

- quercifolius*. β foliis sinuato-pinnatifidis.  
In wet places from New England to Carolina; var. β  
in high mountain woods of Virginia. ♀. June,  
Aug. v. v. Flowers white; leaves smooth.

- pumilus*. 3. L. foliis lanceolatis subserratis glabris, stolonibus procumbentibus. *Vahl. enum. pl. 1. p. 211.*

- L. uniflorus. *Mich. fl. amer. 1. p. 14.*  
In Canada. ♀. Flowers solitary; stems low.

- obtusifolius*. 4. L. foliis lanceolatis remote obtuse serratis. *Vahl. enum. pl. 1. p. 212.*  
At Hudson's Bay. *Michaux.* ♀. +.

25. MONARDA. *Gen. pl. 48.*

- didyma*. 1. M. glabriuscula, capitulis magnis proliferis, bracteis exterioribus coloratis magnis oblongis acuminatis subintegerrimis: calycibus corollisque longis glabriusculis, foliis lato-ovatis subcordatis acuminatis serratis rugosis: serraturis mucronatis, caule acutangulo glabro.

- M. didyma. *Willd. sp. pl. 1. p. 125.*  
M. purpurea. *Lam. encycl. 4. p. 256.*

ilaria.

road-mountains,  
e, small.

labium inferius  
floro. Mich. fl.

Virginia. July.

otundatis, scapo

a. July. v. v.;  
flowers yellow.

-horehound.

lanceolatis ser-  
atis. Mich. fl.  
20.

Virginia. 2.  
mall, in whorls.

egra angustatis,  
i. fl. amer. 1.

arolina; var. β  
ia. 2. June,  
mooth.

onibus procum-

is low.

. Vahl. enum.

3.

is, bracteis ex-  
minatis subin-  
glabriusculis,  
serratis rugo-  
gulo glabro.



*Monarda balmifera.*

M. coccinea.  
Icon. Mill. in  
In rivulets o  
Michaux.  
ers long.

2. M. sparse hi  
exteriorib  
corollisque  
oblongis a  
tis, caule a  
In boggy wo  
Oswego, N  
anum in M  
Flowers vi  
what fragr

3. M. glabriusc  
teis exterio  
liis ovatis  
subtus colo  
M. rugosa.  
In Canada.  
Flowers sm

4. M. glabra,  
bracteis ext  
briusculis,  
centibus te  
ratis pilosiu  
M. clinopodi  
Gronov.  
M. glabra.  
In dry gravell  
on the mo  
yellow with  
The specimen  
*Clinopodium*

5. M. glaberrim  
terminalibu  
tis, calycib  
tenuissimis  
serratis glat  
In the mount  
*v. s. in Her*

6. M. glabriusc  
terioribus r

VOL. I.

- M. coccinea. *Mich. fl. amer.* 1. p. 16.  
*Icon. Mill. ic. t. 183. f. 1. Bot. mag. 546.* Oswego-tea.  
 In rivulets on the high mountains of North Carolina.  
*Michaux.* 2. June—Sept. v. v. in Hortis. Flowers long, scarlet.
2. M. sparse hirsuta, capitulis magnis simplicibus, bracteis exterioribus coloratis lanceolatis attenuatis, calycibus corollisque pubescentibus, corollis longissimis, foliis oblongis attenuatis serratis undique pilis sparsis hirsutis, caule acutangulo petiolisque ciliato-piloso.  
 In boggy woods in black rich soil, near Onondago and Oswego, New York. 2. v. v.; v. s. spec. *Kalmianum in Museo Banksiano, sub nomine M. didymæ.* Flowers very long, beautiful crimson; leaves somewhat fragrant; stems double the size of *M. didyma.*
3. M. glabriuscula, capitulis mediocribus simplicibus, bracteis exterioribus oblongis, calycibus glabriusculis, foliis ovatis subcordatis acutis rugosis glabris, nervis subtus coloratis pilosis, caule acutangulo hirto.  
*M. rugosa. Ait. kew. ed. 2. 1. p. 51.*  
 In Canada. 2. July—Sept. v. s. in *Museo Banksiano.* Flowers small, white.
4. M. glabra, capitulis parvis simplicibus terminalibus, bracteis exterioribus lato-ovatis acutis integerrimis glabriusculis, calycibus brevibus ciliatis, corollis pubescentibus tenuibus, foliis ovato-oblongis acuminatis serratis pilosiusculis, caule obtusangulo glabro.  
*M. clinopodia. Willd. sp. pl. 1. p. 125. excl. syn. Gronov.*  
*M. glabra. Lam. encycl. 4. p. 256?*  
 In dry gravelly soil from Canada to Carolina, particularly on the mountains. 2. July—Sept. v. v. Flowers yellow with purple specks.  
 The specimen of Grönovius in the Banksian Museum is *Clinopodium incanum.*
5. M. glaberrima, capitulis parvis nudiusculis lateralibus terminalibusque, bracteis exterioribus linearibus ciliatis, calycibus pubescentibus ciliatis, corollis brevibus tenuissimis glabris, foliis linear-lanceolatis acuminatis serratis glabris, caule obtusangulo glabro.  
 In the mountains of South Carolina and Virginia. v. v.; v. s. in *Herb. Lyon.* Flowers yellowish white.
6. M. glabriuscula, capitulis magnis foliosis, bracteis exterioribus magnis coloratis lato-ovatis acuminatis ser-
- Kalmiana.*  
*rugosa.*  
*clinopodia.*  
*gracilis.*  
*purpurea.*

- ratis, calycibus coloratis fauce barbatis, corollis longis  
glabriusculis, foliis ovato-oblongis acutis grosse serra-  
tis: nervis undique pilosis, caule acutangulo glabro.  
*M. clinopodia.* *Vahl. enum. pl. 1. p. 218. descriptio.*  
*M. fistulosa* var. crimson. *Bot. mag. 145.*  
Icon. *Bot. mag. 145.*  
In the mountains of Virginia.  $\frac{1}{4}$ . July—Sept. v. v.;  
v. s. in *Herb. Lambert.* Flowers crimson, large.
- oblongata.* 7. *M. undique hirsuta*, capitulo simplicibus, bracteis ex-  
terioribus ovatis acutis, calycibus brevibus fauce bar-  
batis: dentibus divaricatis, caule obtusangulo superne  
hirsuto.  
*M. oblongata.* *Ait. kew. ed. 2. 1. p. 51.*  
*M. longifolia.* *Lam. encycl. 4. p. 255.*  
Frequent in the mountains of Pennsylvania and Virginia.  
 $\frac{1}{4}$ . July, August, v. v. Flowers pale purple.
- mollis.* 8. *M. cano-pubescentis*, capitulo simplicibus, bracteis ex-  
terioribus cordato-ovatis acutis, calycibus brevibus  
fauce dense barbatis, corollis hirsutis: galea apice  
longissime barbata, foliis oblongis attenuatis basi rotun-  
datis serratis utrinque cano-pubescentibus, caule acu-  
tangulo hirsuto.  
*M. mollis.* *Lin. amoen. acad. 3. p. 399.*  
*M. fistulosa*  $\beta$ . *Willd. sp. pl. 1. p. 124.*  
*M. allophylla.* *Mich. fl. amer. 1. p. 16.*  
In the mountains of Virginia and Carolina.  $\frac{1}{4}$ . July—  
Sept. v. v. Flowers pale purple.
- fistulosa.* 9. *M. villis sparsis hirsuta*, capitulo simplicibus proliferis-  
que foliosis, bracteis exterioribus oblongis acutis gla-  
briusculis, calycibus longis barbatis, corollis hirsutis  
mediocribus, foliis ovatis acuminatis serratis, petiolis  
longis ciliatis, caule obtusangulo glabro.  
*M. fistulosa.* *Willd. sp. pl. 1. p. 124.*  
Icon. *Mill. icon. t. 122. f. 2.*  
In hedges and woods, from Canada to Carolina, common.  
 $\frac{1}{4}$ . July—Sept. v. v. Flowers purplish.
- punctata.* 10. *M. glabriuscula*, floribus mediocribus verticillatis, brac-  
teis lanceolatis nervosis coloratis verticillo longioribus,  
foliis lanceolato-oblongis remote serratis glabris, caule  
obtusangulo villoso-canescente.  
*M. punctata.* *Willd. sp. pl. 1. p. 126.*  
*M. lutea.* *Mich. fl. amer. 1. p. 16.*  
Icon. *Plumb. alm. t. 24 f. 1. Bot. rep. t. 546.*

- In sandy fie-  
—Augus-
11. *M. hirsuta*  
glabris vi-  
to-oblong  
caule act  
*M. ciliata.*  
On slate hil-  
v. v. Flo-
12. *M. undique*  
vis vertic-  
tis, calyc-  
serratis lo-  
*M. ciliata.*  
In high mo-  
August.

1. *S. foliis line-*  
cauleque  
*S. azurea.*  
*S. acuminat-*  
*S. angustifol-*  
*S. mexicana.*  
Icon. *Vent-*  
 $\beta$  *Lam. encycl.*  
In open san-  
July—Se-  
times whi-
2. *S. foliis cord-*  
duplo lon-  
*p. 141.*  
Icon. *Muri-*  
On the bank  
v. r. Flo-
3. *S. foliis lance-*  
libus oppo-  
caule brac-  
In open plai-  
*v. s. in He-*  
*Caulis erectu-*  
*Folia oppa-*  
cemis folio-  
gitudine p-

la.  
corollis longis  
grossae serratae.  
gulio glabro.  
*descriptio.*

—Sept. v. v.;  
on, large.

i, bracteis ex-  
ibus fauce bar-  
ingulo superne-

and Virginia.  
purple.

bracteis ex-  
ibus brevibus  
galea apice  
itis basi rotun-  
as, caule acu-

July—

bus proliferis-  
gis acutis gla-  
borris hirsutis  
ratis, petiolis

lina, common.  
h.

ciliatis, brac-  
o longioribus,  
glabris, caule

546.

In sandy fields from New Jersey to Carolina. 4. June—August. v. v. Flowers yellow, punctated.

11. *M. hirsuta*, floribus parvis verticillatis, bracteis ovatis glabris venosis ciliatis calycem aequantibus, foliis ova-to-oblongis attenuatis subsessilibus serratis subpilosis, caule acutangulo hirsuto.

*M. ciliata*. Willd. sp. pl. 1. p. 126.

On slate hills in Virginia and Carolina. 4. June—Aug. v. v. Flowers small, blue.

12. *M. undique pilis longis albidis hirsutissima*, floribus parvis verticillatis, bracteis brevissimis oblongis acuminate, calycibus longe aristatis, foliis ovatis acuminatis serratis longe petiolatis, caule acutangulo hirsuto.

*M. ciliata*. Mich. fl. amer. 1. p. 16.

In high mountains of North Carolina and Virginia. 4. August. v. v. Flowers small, very pale purple.

#### 26. SALVIA. Gen. pl. 50. Sage.

1. *S. foliis linear-lanceolatis inferioribus extrorsum serratis*, cauleque glabris, calyce pubescente brevissime trifido.

*S. azurea*. Lam. encycl. 6. p. 625.

*S. acuminata*. Vent. cels. p. 50.

*S. angustifolia*. Mich. fl. amer. 1. p. 15.

*S. mexicana*. Walt. fl. car. p. 65.

Icon. Vent. cels. p. 50.

β *Lam. encycl. 6. p. 625.*

In open sandy situations of Carolina and Georgia. 4. July—Sept. v. v. Flowers beautiful blue, sometimes white.

2. *S. foliis cordatis acutis tomentosis serratis*, corollis calyce duplo longioribus angustioribus. Willd. sp. pl. 1. p. 141.

Icon. Murr. in com. gæt. 1778. p. 86. t. 1.

On the banks of rivers in Florida. 4. July—Sept. v. v. Flowers scarlet.

3. *S. foliis lanceolatis serratis*, floribus in racemis terminibus oppositis, corollis calyce trifido subaequantibus, caule brachiato-ramosissimo.

In open plains of the Missouri river. *M. Lewis*. ①. v. s. in Herb. Lewis. Flowers small, sky-blue.

*Caulis erectus ramosissimus*: ramuli tetragoni 4-sulcati.

*Folia opposita*, brevi-petiolata. *Flores oppositi*, in racemis foliosis. *Bracteæ* vel *folia floralia lineares*, longitude pedicellarum. *Cal.* tubuloso-campanulatus,

*ciliata*.

*hirsuta*.

*azurea*.

*elata*.

*coccinea*.

*trichostem-  
moides*.

striatus, hispidus. *Lab.* superius subrotundum, integrum: inferius 2-fidum: lacinias oblongis, acutis. *Corollæ labium* superius subrotundum, concavum: inferius 3-lobum, lacinias lateralibus brevissimis: intermedia major, dilatata, undulatim crenata. *Stylus* corollæ duplo longior. *Sem.* 3-gona, glabra, venosa. *Habitus Trichostemmæ dichotomæ.*

*urticifolia.*

4. *S. villos-viscosa*, foliis ovatis oblongis dentatis, per petiolum decurrentibus. *Vahl.* enum. pl. 1. p. 261. *Willd.* sp. pl. 1. p. 131.

Icon. *Moris. hist.* 3. s. 11. t. 13. f. 31. In the mountains of Virginia and Carolina.  $\frac{1}{4}$ . July, Aug. v. v. Flowers blue, viscous.

*lyrata.*

5. *S. foliis radicalibus lyratis dentatis, corollarum galea brevissima, caule subaphyllum retrorsum piloso.* *Vahl.* enum. pl. 1. p. 257. *Willd.* sp. pl. 1. p. 128.

Icon. *Moris. hist.* 3. s. 11. t. 13. f. 27.

*obovata.*

- $\beta$  foliis obovatis repando-dentatis.

Icon. *Dill. eth.* 219. t. 175. f. 216.

In meadows from Canada to Carolina.  $\frac{1}{4}$ . May. v. v. Flowers blue.

A number of varieties are comprehended under this species, which probably may prove to be distinct species. In the Banksian Museum, among the plants of Gronovius, is one of those under the name of *S. verbena-naca*.

## 27. COLLINSONIA. Gen. pl. 51. Horse-weed.

*canadensis.*

1. *C. foliis lato-cordato-ovatis, glabris, calycis dentibus brevi-subulatis, panicula terminalis composita.*

*C. canadensis.* *Willd.* sp. pl. 1. p. 152.

Icon. *Linn. hort. cliff.* 14. t. 5.

*cordata.*

- $\beta$  foliis lato-cordatis, superioribus integerrimis,

*ovata.*

- $\gamma$  foliis ovatis basi attenuatis omnibus serratis.

In shady woods, rocky rich soil from Canada to Carolina.  $\frac{1}{4}$ . July—Sept. v. v.; var.  $\beta$  v. s. in *Herb. Lambert.* Flowers yellow, large.

*scabra.*

2. *C. foliis minoribus ovatis sub-cordatis pilosiusculis, calycis dentibus brevi-subulatis, panicula terminalis simpliuscula, caule piloso scabro.*

*C. scabriuscula.* *Ait. kew. ed. 2.* 1. p. 66.

*C. præcox.* *Walt. fl. car.* p. 65.

In Carolina and Florida.  $\frac{1}{4}$ . July—Sept. v. s. in *Museo Banks.* Flowers small, reddish-yellow.

DIA

In the coll  
lied to t  
stamina.

3. *C. foliis  
petiolis l  
cula tern*

In South  
*Lambert*

4. *C. foliis s  
calycis c  
composit*

*C. tuberosa*

*C. serotina*

In shady  
small, ye

5. *C. foliis ov  
tus pube  
subæqua  
floribus t*

Icon. *Bo*

In the mo

v. v. Fl

28. CIRCAE

1. *C. caule er  
culis.*

Icon. *Eng*

In shady w

July v

2. *C. caule ad  
lyce men*

Icon. *Eng*

In high mo

Canada t

very smo

29. CER

1. *Ceratiola.*

In gravelly

plenty on

ver. h

Leaves ve

Heath ve

*tundum, interlongis, acutis, concavum; inissimis: interta.* *Stylus co-*

*glabra, venosa.*  
*entatis, per pe-*  
*pl. 1. p. 261.*

*ina. 4. July,*  
*rollarum galea*  
*piloso. Vahl.*  
*p. 128.*

*4. May. v. v.*

*under this spe-*  
*distinct species.*  
*plants of Gro-*  
*ne of S. verbe-*

*Horse-weed.*  
*alycis dentibus*  
*imposita.*

*nis,*  
*ada to Carolina.*  
*Herb. Lambert.*

*susculis, calycis*  
*minalis simpli-*

*6.*  
*v. s. in Museo*  
*w.*

In the collection of Mr. Lyons is a specimen, nearly allied to the present one, with larger flowers and four stamens.

3. *C. foliis oblongo-ovalibus utrinque subacutis glabris, petiolis longissimis, calycis dentibus brevissimis, panicula terminalis simplex nudiuscula, caule glabro.*

In South Carolina. *Fraser.* 4. Aug. v. s. in *Herb. Lambert.* Flowers very small, yellow.

4. *C. foliis subrhomboido-ovalibus utrinque acutis glabris, calycis dentibus setaceis tubo longioribus, panicula composita foliosa, caule ramoso subvilloso.*

*C. tuberosa. Mich. fl. amer. 1. p. 17.*

*C. serotina. Walt. fl. car. p. 65.*

In shady wet woods of Carolina. 4. Sept. Flowers small, yellow?

5. *C. foliis ovatis cordatis rugosis glabriusculis, nervis subtus pubescentibus, calycis dentibus linearibus tubum subaequantibus, panicula composita foliosa pubescente, floribus tetrandris, caule ramoso pubescente.*

Icon. *Bot. mag. t. 1213.*

In the mountains of Georgia. *Lyon.* 4. July, Aug. v. v. Flowers large, pale yellow.

## 28. CIRCÆA. Gen. pl. 31. Enchanter's nightshade.

1. *C. caule erecto, foliis ovatis denticulatis opacis glabriusculis. Vahl. enum. pl. 1. p. 301.*

Icon. *Engl. bot. 1056?*

In shady woods from Canada to Carolina. 4. June, July. v. v. Flowers small, reddish white.

2. *C. caule adscendente, foliis cordatis dentatis nitidis, calyce membranaceo. Vahl. enum. pl. 1. p. 301.*

Icon. *Engl. bot. 1057.*

In high mountain woods, on rotten trees and rocks, from Canada to Carolina. 4. July, Aug. v. v. Smaller, very smooth; flowers reddish-white.

## 29. CERATIOLA. Mich. fl. amer. 2. p. 221.

1. Ceratiola. *Willd. sp. pl. 4. p 712. Mich. l. c. p. 222.*

In gravelly dry soil of Georgia and Florida; in great plenty on the Islands in the mouth of St. Mary's River. 4. — v. v. s. fl.; v. s. c. fl. in *Herb. Fraser.*

Leaves verticillate; flowers very small; resembles a Heath very much.

*ovalis.*

*tuberosa.*

*anisata.*

*lutetiana :*  
*canadensis.*

*alpina.*

*ericoides.*

## 30. BORYA, Willd. sp. pl. 4. p. 711.

*porulosa.*

1. B. foliis oblongo-lanceolatis obtusis sessilibus coriaceis, margine revolutis subitus punctatis. Willd. l. c.

*Adelia porulosa.* Mich. fl. amer. 2. p. 224.On the coast of Georgia and Florida.  $\frac{1}{2}$ . v. v.*ligustrina.*

2. B. foliis ovato-lanceolatis acutis subsessilibus submembranaceis. Willd. l. c.

*Adelia ligustrina.* Mich. l. c.In woods on river-sides in the Illinois country.  $\frac{1}{2}$ . July, Aug. v. s. in Herb. Lyon.*acuminata.*

3. B. foliis ovato-lanceolatis utrinque attenuatis petiolatis membranaceis serrulatis. Willd. l. c.

*Adelia acuminata.* Mich. l. c. p. 225.

Icon. Mich. l. c. t. 48.

On banks of rivers in Carolina and Georgia.  $\frac{1}{2}$ . v. s. in Herb. Lyon.*minor.*

## 31. LEMNA. Gen. pl. 1400. Duck-weed.

*gibba.*

1. L. foliis ellipticis utrinque planis basi cohærentibus, radicibus solitariis. Willd. sp. pl. 4. p. 195.

Icon. Engl. bot. t. 1095.

In ditches and stagnant waters, common. ○. June, July. v. v. Floating; flowers exceeding small.

*polyrhiza.*

2. L. foliis ellipticis subtus bullato-convexis basi cohærentibus, radicibus solitariis. Willd. l. c.

Icon. Engl. bot. t. 1233.

In similar situations, rare. ○. June, July. v. v. c. fl. Observed near Liverpool in the western part of New York State.

3. L. foliis ellipticis planis basi cohærentibus, radicibus fasciculatis. Willd. sp. pl. 4. p. 195.

Icon. Vail, par. t. 20. f. 2.

Common in ditches, &amp;c. ○. July, Aug. v. v. s. fl.

III

32. VALERI

calyce c

33. FEDIA.

coronata

34. TRIPTEI

nutis ;

35. IXIA.

C

36. IRIS.

nis cont

37. DILATR

dum.

38. SISYRIN

connata

39. BOERHA

Sem. 1.

40. COMME

Nectar.

41. SYENA.

locularis

42. HETERA

losa : li

tubulos

43. SERPICI

bo 6-pa

44. XYRIS.

3-petalis

### III. TRIANDRIA.

---

#### I. MONOGYNIA.

\* *Flores superi, completi.*

32. VALERIANA. *Cor. 1-petala, s<sup>æ</sup>pe calcarata. Sem. calyce corticatum, coronatum.*
33. FEDIA. *Cal. 3-4-dentatus. Cor. 5-fida. Caps. coronata, 3-locularis : loculamento 1. fertili.*

\*\* *Flores superi, incompleti.*

34. TRIPTERELLA. *Cor. 6-fida : laciniis alternis minutis ; tubo basi ventricoso.*
35. IXIA. *Cor. 6-partita, patens,  $\approx$ equalis tubo recto.*
36. IRIS. *Cor. 6-partita : petalis alternis reflexis, alternis conniventibus. Stigm. 3. petaliformia.*
37. DILATRIS. *Cor. 6-petaloidea, hirsuta. Stigm. 3-fidum.*
38. SISYRINCHIUM. *Spatha. Petala 6. plana. Fil. connata.*

\*\*\* *Flores inferi.*

39. BOERHAVIA. *Cal. tubulosus. Cor. 1-petala. Sem. 1.*
40. COMMELINA. *Cal. 3-phyllos. Cor. 3-petala. Nectar. staminiforme. Caps. 3-locularis.*
41. SYENA. *Cal. 3-phyllos. Cor. 3-petala. Caps. 1-locularis, 3-valvis,*
42. HETERANTHERA. *Spatha 1-flora. Cor. tubulosa : limbo 6-partito : laciniis sub $\approx$ equalibus. Stigm. tubulosum.*
43. SERPICULA. *Spatha 1-flora. Cor. tubulosa : limbo 6-partito : laciniis  $\approx$ equalibus. Stigm. 3. ligulata.*
44. XYRIS. *Cal. cartilagineo glumaceus, 3-valvis. Cor. 3-petala. Caps. 1-locularis, 3-valvis.*

## TRIANDRIA.

\*\*\*\* Flores glumacei.

45. SPARGANIUM. *Amentum* subrotundum. *Cal.* 3-phylus. *Cor.* O. *Stigm.* 2-fidum. *Drupa* 1-sperma. *Monoicum.*
46. TYPHA. *Amentum* cylindricum. *Cal.* obsoletus. *Sem.* 1. pedicellatum basi papposum. *Monoicum.*
47. CAREX. *Amentum* imbricatum. *Cal.* squama. *Stigm.* 2-3. *Semen* tunicatum, 3-quetrum. *Monoicum.*
48. SCLERIA. *Cal.* gluma 2-6-valvis. *Stigm.* 1-3. *Nux* colorata subglobosa. *Monoicum.*
49. ZEA. *Cal.* gluma 2-flora, 2-valvis. *Spadix* simplex, cylindricus: gluma 2-valvis, 1-flora. *Sem.* subrotundum. *Monoicum.*
50. KYLLINGIA. *Flores* distincti, in spicam subimbricati. *Cal.* 2-valvis, 1-florus. *Cor.* 2-valvis, calyx major. *Stylus* bifidus.
51. MAPANIA. *Flores* capitati. *Cal.* 6-valvis. *Cor.* O. *Sem.* 1.
52. DICHROMENA. *Cal.* squamæ in spicam undique imbricatae, inferiores vacuae. *Cor.* O. *Styl.* basi persistens. *Setæ* O.
53. SCHÖENUS. *Cal.* squamæ in spicam fasciculatae, inferiores vacuae. *Cor.* O. *Styl.* deciduus.
54. RYNCHOSPORA. *Cal.* squamæ in spicam fasciculatae, inferiores vacuae. *Cor.* O. *Styl.* basi persistens. *Setæ* basi seminis.
55. FIMBRISTYLIS. *Cal.* squamæ in spicam undique imbricatae. *Cor.* O. *Stylus* basi bulbosus, ciliatus, deciduus.
56. CYPERUS. *Cal.* squamæ in spicam distiche imbricatae. *Cor.* O. *Styl.* deciduus.
57. DULICHIUM. *Cal.* squamæ in spicam distiche imbricatae: spiculis racemosis, ex axillis foliorum. *Cor.* O. *Stylus* longissimus, 2-fidus.
58. SCIRPUS. *Cal.* squamæ in spicam undique imbricatae. *Cor.* O. *Styl.* filiformis, imberbis, deciduus. *Sem.* 1. nudum s. setulis cinctum.
59. TRICHOPHORUM. *Cal.* squamæ in spicam undique imbricatae. *Cor.* O. *Styl.* deciduus. *Sem.* setis (6-9) longissimis calyculatum.
60. ERIOPHORUM. *Cal.* squamæ in spicam undique imbricatae. *Cor.* O. *Sem.* lana longissima calyculatum.
61. VAGINARIA. *Cal.* squamæ in spicam undique im-

- bricatae.  
tim trip  
62. FUIREN  
dique in  
forme,  
63. MARISC  
cati.  
ves, infi  
64. MIEGIA  
mutica.  
*Sem.* nu  
65. LIMNET  
imbricata  
2-valv.  
66. CENCHI  
4 florun  
mutica.  
67. ORYZO  
vis : ex  
68. ZIZANI  
aristata.  
69. MUHLE  
*Cor.* 2-  
tata.  
70. TRICHO  
acutis,  
sessilia.  
71. LEERSI  
compre  
72. MILIUM  
majorib  
*Stigm.*  
73. AGROST  
vioribus  
74. ANTHO  
acumin  
75. PHALAI  
libus.  
76. PHLEUI  
sessilis.  
77. ALOPEC  
tata.

- bricatae. *Stigm. 3.* *Seminis* involucellum alternatim tripaleaceum et trisetum.
62. FUIRENA. *Cal.* squamæ mucronatae in spicam unidique imbricatae. *Sem.* involucellum latum, paleæ forme, saepius aristatum.
63. MARISCUS. Flores distincti, in spicam subimbricatae. *Cal.* 2-valvis, subtriflorus. *Corollulae* 1-valvis, inferior superiorum amplectens. *Styl.* 3-fidus.
64. MIEGIA. *Cal.* multiflorus, 2-valvis. *Cor.* 2-valvis, mutica. Appendices bini, lato-lanceolati. *Stigm. 3.* *Sem.* nudum, maximum.
65. LIMNETIS. *Spica lateriflora*: flosculis subbifariam imbricatis. *Cal.* 2-valvis : valv. altera minore. *Cor.* 2-valv. mutica, compressa, carinata. *Styl.* longus.
66. CENCHRUS. *Involuer.* laciniatum, echinatum, 3-4 florum. *Cal.* 2-valvis, 2-florus. *Cor.* 2-valvis, mutica. *Styl.* 2-fidus.
67. ORYZOPSIS. *Cal.* 2-valvis, 1-florus. *Cor.* 2-valvis : exteriore aristata. *Nectar* 2-phylum.

## II. DIGYNIA.

\* *Flores uniflori, vagi.*

68. ZIZANIA. *Cal.* 0. *Cor.* 2-valvis : valvæ exteriore aristata. *Stam.* 6. *Monoicum.*
69. MUHLENBERGIA. *Cal.* minutissimus, 1-valvis. *Cor.* 2-valvis, basi pilosa : valvæ exteriore apice aristata.
70. TRICHODIUM. *Cal.* 2-valvis : valvis subæqualibus acutis. *Cor.* 1-valvis, brevior, mutica. *Stigm.* subsessilia.
71. LEERSIA. *Cal.* 0. *Cor.* 2-valvis, clausa : valvæ compressis, muticis.
72. MILIUM. *Cal.* 2-valvis : valvis ventricosis, corolla majoribus, subæqualibus. *Cor.* 2-valv. brevissima. *Stigm.* villosa.
73. AGROSTIS. *Cal.* 2-valvis : valvis acutis corolla brevioribus. *Cor.* 2-valvis.
74. ANTHOXANTHUM. *Cal.* 2-valvis. *Cor.* 2-valvis, acuminata, aristata. *Stam.* 2.
75. PHALARIS. *Cal.* 2-valvis : valvis carinatis, æquilibus. *Cor.* 2-valvis, inclusa, basi pilosa.
76. PHLEUM. *Cal.* 2-valvis, truncatus, mucronatus, sessilis. *Cor.* inclusa.
77. ALOPECURUS. *Cal.* 2-valvis. *Cor.* 1-valvis, aristata. *Spica.*

78. PANICUM. *Cal.* 3-valvis : valvula tertia dorsali minima. *Cor.* 2-valvis.
79. DIGITARIA. *Cal.* 2-3-valvis, concavus. *Cor.* 2-valvis, oblongo-ovata, teres, mutica. *Nectar.* fissum. *Spicæ digitatæ* : flores gemini.
80. CYNODON. *Cal.* 2-valvis, patens, lanceolatus. *Cor.* major, 2-valvis : valv. exteriore maxima ovoidea. *Nectar.* truncatum. *Spicæ digitatæ* : floribus solitariis.
81. PASPALUM. *Cal.* 2-valvis : valvis subrotundis, figura corollæ. *Flores* unilaterales.
82. ARISTIDA. *Cal.* 2-valvis. *Cor.* 1-valvis : aristis tribus terminalibus.
83. STIPA. *Cal.* 2-valvis. *Cor.* 2-valvis : valvis involutis truncatis. *Arista* terminalis, basi contorta, longissima.
84. SACCHARUM. *Cal.* 2-valvis : valvis basi villis densis involucrata. *Cor.* 2-valvis : valva interiore longe aristata. *Monandra.*
85. ANDROPOGON. *Flores* gemini : altero masculo. *Cal.* 2-valvis. *Cor.* 3-valvis : valv. secunda minore aristata, tertia intima minuta. *Recept.* plerumque villosum. *Involucrum* villis fasciculatis.

\*\* *Flores biflori, vagi.*

86. ATHEROPOGON. *Cal.* 1-valvis, biflorus. *Cor.* 2-valvis : valvula exteriore 3-aristata. *Polygamum.*
87. TRISETUM. *Cal.* 2-3-florus, acuminatus, carinatus. *Cor.* aristis 2. terminalibus, 1. dorsali recta nec contorta, *flosc.* glabris.
88. ANTHISTIRIA. *Cal.* 4-valvis, 2-4-flora. *Cor.* 2-valvis. *Arista* tortilis e fundo corollæ. *Polygamum.*
89. AIRA. *Cal.* 2-valvis. *Cor.* 2-valvis. *Flosculi* absque rudimento tertii.
90. MELICA. *Cal.* 2-valvis, coloratus, obtusus. *Cor.* 2-valvis, ventricosa. *Rudimentum* tertii inter flores.
91. HOLCUS. *Cal.* 2-valvis. *Cor.* minor : valv. altera aristata *Nectar.* 2-partitum. *Polygamum.*
92. SORGHUM. *Flores* gemini : altero pedicellato, masculo. *Cal.* 2-valvis. *Cor.* 3-valvis : valv. secunda aristata, tertia nectaria villosa connectens. *Polygamum.*

93. POA. *Cal.* 2-valvis : valvula exteriore 3-aristata, marginata. *Cor.* 2-valvis.
94. BRIZA. *Cal.* 2-valvis, ventricosa.
95. UNIOLA. *Cal.* 2-valvis, ovata ; tertiorem.
96. FESTUCA. *Cal.* 2-valvis, tenuiscula.
97. DACTYLIS. *Cal.* 2-valvis, longiore.
98. BROMUS. *Cal.* 2-valvis, cha : arvensis.
99. KOELEB. *Cal.* 2-valvis.
100. AVENA.
101. ARUNDINACEA.
- \*\*\*\*
102. ELEUSIS. *Cal.* multo.
103. CHLORIS. *Cal.* 2-valvis, 2-ovata. *Polygonum.*
104. ROTTBOELIA. *Cal.* ovata, minor, glabra. *Polygonum.*
105. TRIPSA. *Cal.* 2-valvis, mascula gluma perforata.
106. ELYMUS. *Cal.* 2-valvis.
107. HORDEUM. *Cal.* 2-valvis, luciflorum, libus strobiliferus.
108. SECALIS. *Cal.* 2-valvis, solitaria.
109. TRITICUM. *Cal.* 2-valvis, *Floribus* solitariis.

## \*\*\* Flores multiflori, vagi.

93. POA. *Cal.* 2-valvis. *Spicula* ovata mutica : valvulis margin'e scariosis.
94. BRIZA. *Cal.* 2-valvis. *Spiculae* distichæ : valvulis ventricosis, cordatis, obtusis : interiore minuta.
95. UNIOLA. *Cal.* multivalvis. *Spiculae* ancipes ovatae : valvulis muticis, interiore minore, intra exteriorem recumbente.
96. FESTUCA. *Cal.* 2-valvis. *Spicula* oblonga, teretiuscula : valvulis mucronatis.
97. DACTYLIS. *Cal.* 2-valvis, carinatus : valvula altera longiore. *Cor.* 2-valvis. *Spic.* aggregatæ.
98. BROMUS. *Cal.* 2-valvis. *Spicula* oblonga, disticha : arista infra apicem sæpius bifidum.
99. KOELERIA. *Cal.* 2-valvis, compresso-carinatus. *Cor.* 2-valvis, brevi-aristata : glumis nervosis.
100. AVENA. *Cal.* 2-valvis : arista dorsali, contorta.
101. ARUNDO. *Cal.* 2-valvis, glaber. *Cor.* lana cincta.

## \*\*\*\* Flores spicati : receptaculo subulato.

102. ELEUSINE. *Spicæ* digitatae. *Flor.* secundi, mutici. *Cal.* multiflorus. *Cor.* 2-valvis, mutica.
103. CHLORIS. *Spica* digitata, unilateralis. *Cal.* 2-valvis, 2—6-florus : valv. carinatae muticæ aut aristatae. *Polygamum.*
104. ROTTBOELLIA. *Rachis* articulata, teretiuscula. *Cal.* ovato-lanceolatus, planus, uni- vel bivalvis. *Cor.* minor, 2-valvis. *Flosc.* alterni in rachi flexuoso. *Polygamum.*
105. TRIPSACUM. *Mas.* *Cal.* gluma 2-flora, exteriore masculo. *Cor.* gluma membranacea. *Fem.* *Cal.* gluma 1-flora, involucro tecto monophyllo, sinibus perforato. *Cor.* gluma 2-valvis.
106. ELYMUS. *Cal.* involucriformis 4-phyllo, lateralis, 2-valvis, aggregato ternus multiflorus.
107. HORDEUM. *Cal.* 1—2-florus, hexaphyllus, involucriformis. *Flosc.* terni, intermedio sessili, lateribus stipitatis. *Cor.* 2-valvis.
108. SECALE. *Cal.* oppositus, 2-valvis, aut multiflorus, solitarius.
109. TRITICUM. *Cal.* 2-valvis, solitarius, multiflorus. *Floribus* obtusiusculis.

## III. TRIGYNYIA.

110. HOLOSTEUM. *Cor. 5-petala. Cal. 5-phyllo. Caps. apice dehisces.*  
 111. POLYCARPON. *Cor. 5-petala. Cal. 5-phyllo. Caps. 3-valvis.*  
 112. LECHEA. *Cor. 3-petala. Cal. 5-phyllo. Caps. 3-cocca.*  
 113. EROCAULON. *Cor. 3-petala. Cal. compositus. Sem. 1. coronatum corolla.*  
 114. MOLLUGO. *Cor. O. Cal. 5-phyllo. Caps. 3-lo-cularis.*  
 115. PROSERPINACA. *Cor. O. Cal. 3-partitus. Nux 3-locularis, infera.*

## IV. POLYGYNIA.

116. EMPETRUM. *Cal. 3—5-part. Cor. 3-petala. Styl. 9. Bac. 9-sperma.*

## I. MONOGYNIA.

32. VALERIANA. *Gen. pl. 60.*

*pauciflora.*

1. *V. foliis radicalibus pinnatis, caulinis ternatis: foliolis ovalibus acutis serratis, paniculis laxis paucifloris.* *Mich. fl. amer. 1. p. 18.*

In shady woods on the Allegany mountains: Tennessee.  $\frac{1}{2}$ . June, July. v. v. Flowers white.

33. FEDIA. *Gært. de fr. et sem. 2. p. 36.*

*radiata.*

1. *F. foliis spatulato-oblongis subintegris, fructu pubescente subtetragono apice nudo.* *Mich. fl. amer. 1. p. 18.*

*Valeriana locusta radiata.* *Linn. sp. pl. 48.*

In fields from Pennsylvania to Carolina; very plentifully in Virginia.  $\odot$ . June, July. v. v.

34. TRIPTERELLA. *Mich. fl. amer. 1: p. 19.*

*capitata.*

1. *Tripterella.* *Mich. l. c.*

*Anonymos capitatus.* *Walt. fl. car. 68.*

Vogelia capital  
Icon. *Mich.*

In wet places,  
 $\odot$ . May—J  
in a small h

1. *I. foliis linea-*

*viore.* *W.*  
Icon. *Bartr.*  
Found by Mr.  
gia.  $\frac{1}{2}$ .

36.

1. *I. barba cristi-*  
floro, foliis  
larum æqua-

*I. cristata.*  
*p. 225.*

*I. virginiana p*

Icon. *Bot. n*

On the banks  
tains of Vir-  
ers blue, ha-

The roots of  
able; they  
which in a

by far more

Virginia, n

frequently t

2. *I. imberbis,*  
liis ensiformi-

oribus, cap-

*I. virginica.*

*I. hexagona.*

Icon. *Bot. i*

*elth. t. 155*

On river sides

$\frac{1}{2}$ . June,

with the be

3. *I. imberbis,*  
ensiformi-

tibus, caps-

*I. versicolor.*

*Vogelia capitata.* *Gmel. syst. nat. 2. p. 107.*

Icon. *Mich. l. c. t. 3.* In wet places, borders of ponds, &c. in Lower Carolina.

⊖. May—July. v. v. Stem filiform; flowers white, in a small head.

35. IXIA. *Gen. pl. 76.*

1. *I. foliis linearis subulatis scapo unifloro multoties breviore.* *Willd. sp. pl. 1. p. 200.*

Icon. *Bartr. it. 152. t. 3.*

Found by Mr. W. Bartram in South Carolina and Georgia. ♀. Flowers bright blue.

*cælestina.*

36. IRIS. *Gen. pl. 97. Flag.*

1. *I. barba cristata, caule brevissimo compresso subuni-*  
*floro, foliis gladiatis longitudine flororum, laciniis corol-*  
*larum æqualibus.*

*I. cristata.* *Ait. kew. ed. 2. 1. p. 117. Willd. sp. pl. 1.*  
*p. 225.*

*I. virginiana pumila.* *Pluk. alm. 198. t. 196. f. 6.*

Icon. *Bot. mag. 412.* *Ladies' Calamus.*  
On the banks of rivers in rocky situations in the mountains of Virginia and Carolina. ♀. May. v. v. Flowers blue, handsomely crested with yellow.

The roots of this species when chewed are very remarkable; they at first occasion a pleasant sweet taste, which in a few minutes turns to a burning sensation, by far more pungent than Capsicum. The hunters of Virginia, notwithstanding those properties, use it very frequently to alleviate thirst.

*cristata.*

2. *I. imberbis, caule ancipiti multifloro foliis altiore, foliis ensiformibus, stigmatibus petalis interioribus brevioribus, capsulis oblongis: angulis sulcatis.*

*I. virginica.* *Willd. sp. pl. 1. p. 232.*

*I. hexagona.* *Walt. fl. car. 66.*

Icon. *Bot. mag. 703. Jacq. ic. rar. 2. t. 223. Dill.*  
*elth. t. 155. f. 188.*

On river sides and near ponds from New York to Florida.

♀. June, July. v. v. Flowers beautiful purple, with the bottom of the outside petals yellow.

*virginica.*

3. *I. imberbis, caule tereti flexuoso folia æquante, foliis ensiformibus, stigmatibus petalis interioribus æquantiibus, capsulis ovatis: angulis obtusis.*

*I. versicolor.* *Willd. sp. pl. 1. p. 233.*

*versicolor.*

- Icon. *Bot. mag.* 21. *Dill. elth.* t. 155. f. 187.  
Very common in wet places, and on the banks of ditches  
from Canada to Carolina.  $\frac{1}{4}$ . June, July. v. v.  
Flowers purple, stigmata white.
- cuprea.* 4. *I. imberbis*, caule tereti flexuoso folia aequante, foliis  
lato-ensiformibus, stigmatibus linearibus brevibus,  
petalis omnibus reflexis emarginatis obovatis, interiori-  
bus brevioribus, capsulis magnis hexagonis.  
On the banks of the Mississippi near New Orleans; dis-  
covered by Mr. Enslen, Collector to the Prince Lich-  
tenstein of Austria.  $\frac{1}{4}$ . July. v. v. Flowers of a  
beautiful copper colour, veined with purple.
- tridentata.* 5. *I. imberbis*, caule tereti foliis longiore, foliis ensiformi-  
bus, rudimentis petalorum interiorum tridentatis, den-  
ticulo medio acuminato, capsulis trigonis.  
*I. tripetala.* *Walt. fl. car.* 66.  
On the sea-shore of Lower Carolina. *Walter.*  $\frac{1}{4}$ .  
Flowers blue, with purple veins.
- verna.* 6. *I. imberbis*, acaulis, uniflora, foliis longissimis gramineis  
glaucis, tubo longissimo, petalis subæqualibus, capsulis  
subrotundis ad terram sessilibus.  
*I. verna.* *Willd. sp. pl.* 1. p. 235. excl. syn. *Pluk.*  
On high mountains of Virginia and Kentucky; plenti-  
fully on the Cacapon mountains near Winchester.  
 $\frac{1}{4}$ . April, May. v. v. Flowers pale blue, striped  
with purple.
- sibirica.* 7. *I. imberbis*, caule fistuloso tereti foliis altiore subtrifloro,  
foliis linearibus; capsulis abbreviato-trigonis utrinque  
obtusis.  
*I. sibirica.* *Willd. sp. pl.* 1. p. 237.  
Icon. *Bot. mag.* 50. *Jacq. austr.* 1. t. 3.  
On the banks of the Missouri. *M. Lewis.*  $\frac{1}{4}$ . July.  
v. v.; v. s. in *Herb. Lewis.* Flowers blue.
- prismatica.* 8. *I. imberbis*, caule solido tereti folia aequante, foliis an-  
gustissimis longis, capsulis elongato-prismaticis utrin-  
que acutis.  
In deep swamps of New Jersey, near Egg harbour.  $\frac{1}{4}$ .  
July. v. v. Flowers pale purple.
37. DILATRIS. *Gen. pl.* 82.
- tinctoria.* 1. *D. petalis lanceolatis extus villosis, panicula corymbosa*  
tomentosa, foliis longis nudis linearibus.  
*D. caroliniana.* *Lam. illustr.* 1. p. 127.

TRIA  
Anonymos tir  
Heritiera Gm  
Icon. Mich.  
In sandy swa  
Florida. 1  
corymb ve

## 38. SIS

1. *S. foliis scap  
rata : valva  
Pers. encl.*  
In wet meade-  
nia.  $\frac{1}{4}$ .  
smaller tha
2. *S. scapo simp  
rantibus.*  
*S. gramineu*  
Icon. *Bot. i*  
On dry hills  
 $\frac{1}{4}$ . July,

## 39.

1. *B. caule tet*  
*corymboso*  
Icon. *Pluk.*  
On the sea-  
v. v.; v. s.

## 40.

1. *C. foliis line*  
*involucris*  
*p. 165.* 1  
In sandy wo  
in this as v
2. *C. foliis lanc*  
*duplicatis,*  
*libus.* *Va*  
*C. longifolia*  
In shady roc  
July. v. 1
3. *C. foliis lang*  
*subpubesc*  
*pl. 2. p. 1*

187.  
anks of ditches  
July. v. v.  
quante, foliis  
us brevibus,  
atis, interiori-  
nis.  
Orleans; dis-  
Prince Lich-  
Flowers of a  
ple.  
is ensiformi-  
dentatis, den-  
s.

*Anonymos tinctoria.* *Walt. fl. car.* 68.  
*Heritiera Gmelini.* *Mich. fl. amer.* 1. p. 21.  
Icon. *Mich. fl. amer.* 1. t. 4.  
In sandy swamps of New Jersey, and from Carolina to  
Florida. ♀. June, July. v. v. Root red; stem and  
corymb very tomentose; petals yellow inside.

38. SISYRINCHIUM. *Gen. pl.* 1101.

1. S. foliis scapisque simplicibus subsetaceis, spatha colo- *mucronatum.*  
rata: valva altera in mucrone longum desinente.  
*Pers. enc.* 2. p. 50. *Mich. fl. amer.* 2. p. 33.  
In wet meadows of short grass in Pennsylvania and Virgi-  
nia. ♀. June, July. v. v. Flowers beautiful blue,  
smaller than the following species.
2. S. scapo simplici alato, spathis inaequalibus flores supe-  
rantibus. *Pers. enc.* 2. p. 50 *Cav. diss.* 6. p. 345.  
*S. gramineum.* *Lam. encycl.* 1. p. 408.  
Icon. *Bot. mag.* 464. *Cav. l. c. t.* 190. f. 2.  
On dry hills and grass plots from Canada to Carolina.  
♀. July, Aug. v. v. Flowers blue.

39. BOERHAVIA. *Gen. pl.* 13.

1. B. caule tetragono glabro, internodiis viscosis, floribus  
corymboso-paniculatis. *Willd. sp. pl.* 1. p. 19.  
Icon. *Pluk. alm. t.* 226. f. 7.  
On the sea-coast of Carolina and Florida. ♀. July.  
v. v.; v. s. in *Herb. Walter.*

40. COMMELINA. *Gen. pl.* 86.

1. C. foliis lineari-lanceolatis glabris, vaginis subciliatis,  
involutris cordatis conduplicatis. *Vahl. enum. pl.* 2.  
p. 165. *Mich. fl. amer.* 1. p. 24.  
In sandy woods of Carolina. ♀. July. v. v. Flowers  
in this as well as all the following species sky-blue.
2. C. foliis lanceolatis petiolatis cauleque erecto-pilosus con-  
duplicatis, involutris lateralibus terminalibusque sessi-  
libus. *Vahl. enum. pl.* 2. p. 166.  
*C. longifolia.* *Mich. fl. amer.* 1. p. 23.  
In shady rocky situations in Virginia and Carolina. ♀.  
July. v. v. Stems long, branched.
3. C. foliis lanceolatis subpetiolatis supra glabris, vaginis  
subpubescentibus, caule erecto simplici. *Vahl. enum.*  
pl. 2. p. 167. *Willd. sp. pl.* 1. p. 251.

## TRIANDRIA MONOGYNIA. Commelinaceae.

*communis.**erecta.**fluvialis.**acuta.**limosa.**graminea.*

In shady woods from Pennsylvania to Carolina. 24. July, Aug. v. v. Flowers large.

4. C. foliis ovato-lanceolatis subsessilibus acutis cauleque repente glabris, involucris cordatis conduplicatis. Vahl. enum. pl. 2. p. 170. Willd. sp. pl. 1. p. 249.

Icon. Dill. eth. t. 78. f. 89.

C. caroliniana. Walt. fl. car. 68.

In sandy fields in Carolina. Walter. 24. July. v. s. in *Herb. Walter.*

5. C. foliis ovato-lanceolatis scabris, involucris cucullatis turbinatis, caule erecto. Vahl. enum. pl. 2. p. 172. Willd. sp. pl. 1. p. 251.

C. communis. Walt. fl. car. 68.

Icon. Dill. eth. t. 77. f. 88?

In sandy shady places in Virginia and Carolina. 24. July. v. v. Pet. 2. large.

## 41. SYENA. Gen. pl. 88.

1. Syena. Willd. sp. pl. 1. p. 254.

Mayaca fluvialis. Aubl. guj. 1. p. 42.

Icon. Aubl. guj. lyc. t. 15. Lam. illustr. f. t. 36

In small rivulets of Virginia and Carolina. 24. July. v. v. Flowers white, solitary, small.

## 42. HETERANTHERA. Beauvois in act. soc. amer. 4.

1. H. foliis subrotundo-reniformibus acutis, spathis subtriangularibus glabris. Vahl. enum. pl. 2. p. 42. Willd. in act. soc. am. h. nat. 3. p. 438.

Leptanthus reniformis. Mich. fl. amer. 1. p. 25.

Icon. Beauvois in act. soc. amer. 4. p. 173. H. reniformis.

In small rivulets and ditches, on a clay soil, in Pennsylvania, Virginia, &c. 24. June, July. v. v. Flowers white; leaves floating.

2. H. foliis ovato-oblongis, spathis lanceolatis unifloris. Vahl. enum. pl. 2. p. 44.

Pontederia limosa. Swartz. fl. ind. occid. 1. p. 611.

Leptanthus ovalis. Mich. fl. amer. 1. p. 25.

Icon. Mich. l. c. t. 5. f. 1. Sloan. hist. 1. t. 149. f. 1.

On the banks of rivers and small rivulets in the southwestern parts of the United States. 24. July. v. s. Flowers blue; leaves erect.

3. H. foliis omnibus linearibus. Vahl. enum. pl. 2. p. 45.

Leptanthus gramineus. Mich. fl. amer. 1. p. 25.

Commelina dubia. Jacq. obs. bot. 3. p. 9.

## TRIAND

Anonymos aqu.

Icon. Mich.

Floating, in th

July. v. v.

## 43. S

1. S. floribus tria reflexis bifidis

Elodea canade

In stagnant w

24. July.

white; leav

Michaux de

which is onl

at flowering

acute.

## 44

1. X. foliis longis

tulo globoso

X. indica. W

Icon. Pluk. a

In overflowed

Virginia. 1

2. X. foliis linear

acuto: squa

X. caroliniana.

X. Jupicai. M

In low grassy

Florida. 2

flowers yell

3. X. foliis subula

so. Mich. J

In low meadow

July. v. s.

45. SPARGA

1. S. foliis basi

communi ra

p. 961. W

S. erectum. S

Icon. Curt. le

VOL. I.

*Anoëmös aquatica graminifolia. Clayt. num. 814.  
Icon. Mich. fl. amer. 1. t. 5. f. 2. Jacq. l. c. t. 59.  
Floating, in the waters of Pennsylvania, Virginia, &c. 24.  
July. v. v.; v. s. in Herb. Gronov. Flowers yellow.*

## 43. SERPICULA. Gen. pl. 1413.

1. *S. floribus triandris hermaphroditis, stigmatibus ligulatis occidentalis.  
reflexis bifidis, foliis ternatis linearibus acutis.  
Elodea canadensis. Mich. fl. amer. 1. p. 20.  
In stagnant waters, frequent; from Canada to Virginia.  
24. July. v. v. Flowers very small and delicate,  
white; leaves when magnified very finely serrulate.  
Michaux describes the leaves to be oblong and obtuse,  
which is only the case in the early part of the season:  
at flowering time they certainly are long, linear, and  
acute.*

## 44. XYRIS. Gen. pl. 89.

1. *X. foliis longissime-gramineis scapisque tortuosis, capitulo globoso: squamis subrotundis obtusis.  
X. indica. Willd. sp. pl. 1. p. 254?  
Icon. Pluk. alm. t. 416. f. 4?  
In overflowed meadows and small ponds: Pennsylvania to  
Virginia. 24. June, July. v. v. Flowers yellow.*
2. *X. foliis linear-gramineis, scapo ancipiti, capitulo ovato  
acuto: squamis obtusis.  
X. caroliniana. Lam. illustr. 1. p. 132. Walt. fl. car. 69.  
X. Jupicai. Mich. fl. amer. 1. p. 23.  
In low grassy fields, in sandy soil, from New Jersey to  
Florida. 24. June—Aug. v. v. Heads small;  
flowers yellow; it is extremely variable.*
3. *X. foliis subulato-gladiatis, scapo tereti, capitulo globo-  
so. Mich. fl. amer. 1. p. 23.  
In low meadows of Lower Carolina and Georgia. 24.  
July. v. s. The smallest species; flowers yellow.*

## 45. SPARGANIUM. Gen. pl. 1402. Bur-reed.

1. *S. foliis basi triquetris lateribus concavis, pedunculo  
communi ramoso, stigmate lineari. Smith. brit. 3.  
p. 961. Willd. sp. pl. 4. p. 199.  
S. erectum. Sp. pl. 1378.  
Icon. Curt. lond. 5. t. 66. Engl. bot. 744.*

## TRIANDRIA MONOGYNIA. Sparganium.

- In ditches and ponds from Canada to Virginia, rare.  $\frac{1}{4}$ . July, August. v. v. Flowers in round heads, white.
- simplex.* 2. S. foliis basi triquetris lateralibus planis, pedunculo communi simplici, stigmate linearis. *Smith. brit.* 3. p. 962. *Willd. sp. pl.* 4. p. 199.  
*S. erectum*  $\beta$ . *Sp. pl.* 1378.  
 Icon. *Curt. lond.* 5. t. 67. *Engl. bot.* 745.  
 In similar situations with the former; more frequent.  $\frac{1}{4}$ . July, Aug. v. v.
- natans.* 3. S. foliis decumbentibus planis, pedunculo communi simplici, stigmate ovato brevissimo, capitulo masculino subsolitario. *Smith. brit.* 3. p. 962. *Willd. sp. pl.* 4. p. 200.  
 Icon. *Engl. bot.* 273. *Fl. dan.* 260.
- angustifoli- um.*  $\beta$ . S. foliis planis angustissimis, caulem longissime superantibus, debiliter fluitantibus. *Mich. fl. amer.* 2. p. 189.  
 In rivulets of Canada, and very frequent on the outlets of the small lakes of New York.  $\frac{1}{4}$ . Aug. v. v.  
 Floating; heads small.
46. TYPHA. *Gen. pl.* 1401. Cat's-tail. Reed-mace.
- latifolia.* 1. T. foliis linearibus planis, spica mascula femineaque approximatis, utraque cylindrica. *Willd. sp. pl.* 4. p. 197.  
 Icon. *Bot. mag.* 1455. *Fl. dan.* 645. Coopers-reed.  
 In marshes and pools very common.  $\frac{1}{4}$ . July, Aug. v. v. Flowers in a long cylindrical spike.
- angustifolia.* 2. T. foliis linearibus canaliculatis, spica mascula femineaque remotis, utraque cylindrica. *Willd. sp. pl.* 4. p. 198.  
 Icon. *Bot. mag.* 1456.  
 In similar situations, only more rare. Pennsylvania and Virginia.  $\frac{1}{4}$ . July, Aug. v. v.
47. CAREX. *Gen. pl.* 1407. Sedge-grass.
- A. *Stigmatibus binis.*  
 $\dagger$  *Spicis dioicis.*
- scirpoidea.* 1. C. spica solitaria imbricato-cylindrica, capsulis dense pubescentibus, foliis planis. *Mich. fl. amer.* 2. p. 171.  
 Near Hudson's Bay. *Mich.*  $\frac{1}{4}$ . +.
- sterilis.* 2. C. spicis subsenis, fructibus ovatis compresso-triquetris acuminatis, apice recurvis bicuspidiatis, margine ciliato-serratis. *Willd. sp. pl.* 4. p. 208.

Icon. *Schk.*  
 In wet mead.  
 $\frac{1}{4}$ . May.

3. C. spicis in f  
 tis compre  
 tis. *Will*  
*C. typhina.*  
 Icon. *Schk.*  
 In shady wo  
 v. v.

- \*\*
4. C. spiculis al  
 ferioribus  
 sulis ovatis  
 curvo. *W*  
 Icon. *Schk.*  
 On the sea-coa  
 $\frac{1}{4}$ . June, .
5. C. spiculis ot  
 sulis oblon  
 oblongis m  
 pl. 4. p. 21  
 Icon. *Schk.*  
 In dry fields
6. C. spiculis s  
 bidentatis  
 oblongo-la  
 Icon. *Schk.*  
 On dry hills
7. C. spiculis su  
 tibus ovatis  
 vosis, culn  
 p. 233.  
 Icon. *Schk.*  
 In bogs and s  
 $\frac{1}{4}$ . June.
8. C. spiculis ov  
 ovatis plan  
 gine ciliatc  
 Icon. *Schk.*

Icon. *Schk. car. t. Mmm. f. 146.*  
In wet meadows; frequent from Canada to Virginia.  
*4. May. v. v.*

*†† Spicis androgynis.**\* Spica unica apice mascula.*

3. C. spicis in formam ellipticam aggregatis, fructibus ovatis compressis bifidis marginatis superne ciliato-serratis. *Willd. sp. pl. 4. p. 220.*

*cephalophora.*

*C. typhina. Mich. fl. amer. 2. p. 169.*

Icon. *Schk. car. t. Hhh. f. 133.*

In shady woods: Pennsylvania, Virginia, &c. *4. June. v. v.*

*\*\* Spicis pluribus apice masculis.*

4. C. spiculis alternis confertis, superioribus masculis, inferioribus feminineis, intermediis apice masculis, capsulis ovatis marginatis bifidis ciliato-serratis, culmo in curvo. *Willd. sp. pl. 4. p. 223.*

*arenaria.*

Icon. *Schk. car. t. B. Dd. f. 6. Host. gram. 1. t. 49.*  
On the sea-coast in loose sands; New York to Virginia.  
*4. June, July. v. v.*

5. C. spiculis oblongis alternis remotiusculis sessilibus, capsulis oblongis acuminatis rostratis bicuspitatis, squamis oblongis mucronatis. *Schk. car. 2. p. 8. Willd. sp. pl. 4. p. 258.*

*bromoides.*

Icon. *Schk. car. t. Xxx. f. 176.*  
In dry fields and woods. *4. May. v. v.*

6. C. spiculis subquaternis remotiusculis, fructibus ovatis bidentatis margine glabris reflexo-patentibus, squamis oblongo-lanceolatis. *Schk. car. 2. p. 11.*

*retroflexa.*

Icon. *Schk. car. t. Kkk. f. 140.*  
On dry hills: Pennsylvania, &c. *4. v. v.*

7. C. spiculis subquinis oblongis aggregatis, fructibus patentibus ovatis acuminatis bicuspitatis convexo-planis nervosis, culmo triquetro asperrimo. *Willd. sp. pl. 4. p. 233.*

*stipata.*

Icon. *Schk. car. t. Hhh. f. 132.*  
In bogs and swamp meadows: New England to Virginia.  
*4. June. v. v.*

8. C. spiculis ovatis approximatis, fructibus horizontalibus ovatis plano-convexis bidentatis, basi marginatis, margine ciliato-serratis. *Willd. sp. pl. 4. p. 234.*

*muricata.*

Icon. *Schk. car. t. E. Dd. f. 22.*

- In wet woods and meadows: New York, &c. ¼. May, June. v. v.
- Muhlenbergii.*
9. C. spiculis subquinis ovatis alternis approximatis, fructibus subrotundo-ovatis marginatis compressis bidentatis ciliato-serratis, squamis mucronatis. *Willd. sp. pl. 4. p. 231.*  
*C. vulpinoidea. Mich. fl. amer. 2. p. 169.*  
*Icon. Schk. car. t. Ygg. f. 178.*
- On dry hills and in barren woods. ¼. May, June. v. v.
- multiflora.*
10. C. spicis anguste paniculatis oblongis obtusis, fructibus ovatis acuminatis bicuspitatis, squamis ovatis mucronatis, bracteis foliaceis filiformibus. *Willd. sp. pl. 4. p. 243.*  
*Icon. Schk. car. t. Lll. f. 144.*  
 In wet meadows: Pennsylvania, Virginia, &c. ¼. v. v.
- sparganoides.*
11. C. spiculis multifloris suboctonis ovatis subapproximatis, fructibus ovatis compressis marginatis bifidis margine ciliato-serratis horizontalibus. *Willd. sp. pl. 4. p. 237.*  
*Icon. Schk. car. t. Lll. f. 142.*  
 On the edges of ponds: New England, Pennsylvania, &c. ¼. May—July. v. v.
- divulsa.*
12. C. spica basi subramosa, spiculis ovatis approximatis, inferioribus remotis, fructibus erectiusculis ovatis planis convexis bidentatis marginatis, margine glabris. *Willd. sp. pl. 4. p. 235.*  
*Icon. Schk. car. t. Dd. Ww. f. 89.*  
 In shady boggy places: New York, Virginia, &c. ¼. June, July. v. v.
- rosea.*
13. C. spiculis subquaternis remotis, fructibus ovatis acuminatis bidentatis margine ciliato-serratis horizontalibus, squamis ovatis obtusis, bractea foliacea ad basin spiculae inferioris. *Willd. sp. pl. 4. p. 237.*  
*C. echinata β rosea. Wahlb. act. holm. 1803. p. 147.*  
*Icon. Schk. car. t. Zxx. f. 179.*  
 In dry woods: Canada, Pennsylvania, Virginia, &c. ¼. v. v.
- paniculata.*
14. C. spicis paniculatis, fructibus ovatis superne marginatis bidentatis margine ciliato-serratis, culmo triquetro. *Willd. sp. pl. 4. p. 244.*  
*Icon. Schk. car. t. D. f. 20. et var. t. Ttt. f. 163.*  
 In bog-meadows: Virginia, &c. ¼. July. v. v.
- \*\*\* *Spicis pluribus apice semineis.*
- leporina.*
15. C. spiculis tribus subrotundo-ellipticis alternis congestis,

TRIA

- fructibus ell. *Willd. sp.*  
*C. Lachenali. Icon. Schk.*  
 In the west. ¼. June.
16. C. spiculis sul ovatis bide erectis, squ p. 237.  
*C. triceps. A*  
*Icon. Schk.*  
 In wet meado
17. C. spiculis du tis, fructibl bractea foli Willd. sp.  
*C. Richardi.*  
*Icon. Schk.*  
 In swamp me
18. C. spiculis su proximatis tis bidentat C. leporina.  
*Icon. Schk.*  
 In wet places
19. C. spiculis sul matis, fruc datis, bract p. 230.  
*C. viridula.*  
*Icon. Schk.*  
 In wet and ¼. May—
20. C. spiculis su fructibus ov erectis, sqt pl. 4. p. 24 C. canescens.  
*Icon. Schk.*  
 In wet woods
21. C. spiculis alt suffultis, siusculis.

- fructibus ellipticis compressis acuminatis ore integris.  
*Willd. sp. pl. 4. p. 229.*
- C. Lachenalii. *Schk. car. 1. p. 51.*  
Icon. *Schk. car. t. Y.f. 29. Fl. dan. 294.*  
In the western parts of New York, Pennsylvania, &c.  
 $\frac{4}{4}$ . June. v. v.
16. C. spiculis subquaternis approximatis ellipticis, fructibus ovatis bidentatis compressis margine ciliato-serratis erectis, squamis ellipticis obtusis. *Willd. sp. pl. 4. p. 237.*  
*C. triceps. Mich. fl. amer. 2. p. 170?*  
Icon. *Schk. car. t. Zxx. f. 180.*  
In wet meadows, common.  $\frac{4}{4}$ . May—July. v. v.
17. C. spiculis duodenis alternis ellipticis obtusis approximatis, fructibus ovato-lanceolatis marginatis bicuspidiatis, bractea foliacea longissima ad basin spicæ ultimæ. *Willd. sp. pl. 4. p. 230.*  
*C. Richardi. Mich. fl. amer. 2. p. 170.*  
Icon. *Schk. car. t. Yyy. f. 177.*  
In swamp meadows, common.  $\frac{4}{4}$ . June, July. v. v.
18. C. spiculis subsenis subrotundo-ellipticis alternis subapproximatis inferne masculis, fructibus ovatis marginatis bidentatis ciliato-serratis. *Willd. sp. pl. 4. p. 229.*  
*C. leporina. Wahlb. act. holm. 1803. p. 145.*  
Icon. *Schk. car. t. B.f. 8.*  
In wet places: New York, &c.  $\frac{4}{4}$ . June. v. v.
19. C. spiculis subquinis alternis ellipticis obtusis subapproximatis, fructibus ovato-lanceolatis marginatis bicuspidiatis, bracteis oblongis mucronatis. *Willd. sp. pl. 4. p. 230.*  
*C. viridula. Mich. fl. amer. 2. p. 170?*  
Icon. *Schk. car. t. Xxx. f. 175.*  
In wet and sometimes even very dry places, common.  
 $\frac{4}{4}$ . May—July. v. v.
20. C. spiculis subsenis alternis approximatis cylindraceis, fructibus ovatis plano-convexis acutiusculis ore integris erectis, squama ovata acuta longioribus. *Willd. sp. pl. 4. p. 241.*  
*C. canescens. Wahlb. act. holm. 1803. p. 148.*  
Icon. *Schk. car. t. C. f. 13.*  
In wet woods: New York, &c.  $\frac{4}{4}$ . June. v. v.
21. C. spiculis alternis remotis, bracteis foliaceis longissimis suffultis, fructibus ovatis acuminatis bifidis compres-siusculis. *Willd. sp. pl. 4. p. 239.*

*scirpoides.**lagopodioides.**ovalis.**scoparia.**curta.**remota.*

## TRIANDRIA MONOGYNIA. Carex.

- Icon. Schk. car. t. E. t. 23.  
In shady woods: Pennsylvania mountains, &c. ♀. June. v. v.
- festucacea.* 22. C. spiculis suboctonis subapproximatis alternis cylindraceis, fructiferis clavatis, fructibus subrotundo-ovatis rostratis bidentatis margine ciliato-serratis, squama lanceolata mucronata majoribus. Willd. sp. pl. 4. p. 242.  
Icon. Schk. car. t. Ww. f. 173.  
On dry hills, frequent. ♀. May, June. v. v.
- ††† *Spicis sequi distinctis.*
- \* *Spica mascula solitaria, reliquis femineis ex parte androgynis.*
- saxatilis.* 23. C. spicis femineis geminis inferiore pedunculata oblongis, fructibus ellipticis obtusis squamam oblongam obtusam æquantibus, bracteis oblongis amplexicaulibus apice subfoliaceis, culmo glabro. Willd. sp. pl. 4. p. 272.  
Icon. Schk. car. t. I. et Tt. f. 40.  
In hemlock woods of Vermont and New Hampshire. ♀. June. v. v.
- cæspitosa.* 24. C. spicis femineis cylindraceis obtusis subternis distantibus infima brevissime exerte pedunculatis, fructibus ovatis obtusis ore pertusis, squama oblonga obtusa majoribus, foliis patulis. Willd. sp. pl. 4. p. 287.  
C. polyandra. Schk. car. 1. p. 59. t. Dd. f. 90.  
Icon. Schk. car. t. Aa. et Bb. f. 85.  
In bog-meadows, very common. ♀. May, June. v. v.
- crinita.* 25. C. spicis masculis geminis, femineis quaternis distantibus pedunculatis pendulis cylindraceis, fructibus subrotundo-ellipticis ventricosis brevissime rostellatis ore integris squama oblonga arista brevioribus. Willd. sp. pl. 4. p. 300. Lam. encycl. 3. p. 379.  
Icon. Schk. car. t. Eee. f. 125. et t. Ttt. f. 164.
- paleacea.* β. C. spicis cylindraceis, masculis pluribus, pedunculis longis reclinatis, bracteis foliaceis distantibus, squamis cuspide longa serrata terminatis, capsulis suborbicularis, ore emarginato, culmo laxe. Wahlb. act. holm. 1803. p. 164.  
In wet woods, common. ♀. June, July. v. v.
- acuta.* 26. C. spicis masculis binis ternisve, femineis subquaternis subpedunculatis subnutantibus cylindraceis remotis, fructibus oblongis brevissime rostellatis, ore integro

TRIAN

pertuso, squ  
Willd. sp. pl  
Icon. Schk. c  
t. 95.  
In swamps, very

† Spic

27. C. spica simpli integris squa undulatis cre  
Icon. C. Fras  
On the moun  
April. v. v.
28. C. spica simpli questris scabri infima apice  
Icon. Schk. c  
In barren wet
29. C. spica simpli so-triquetris tuis mucro  
C. leptalea. 1  
C. microstachy  
Icon. Schk. c  
In low wet pla
30. C. spicis subq bus obovatis inuncronatis.  
Icon. Schk. c  
On dry hills :  
&c. ♀. M
31. C. spicis subq bus obovat capsulam æ  
C. ovata. Ru  
Icon. Rudge  
In Canada and
- †† *Spicis ter*
32. C. spica andro  
femineis su

pertuso, squamam oblongam acutam subæquantibus.  
Willd. sp. pl. 4. p. 304.

Icon. Schk. car. t. Ee. et Ff. f. 92. Host. gram. 1.  
t. 95.

In swamps, very common. 24. June, July. v. v.

B. *Stigmatibus tribus.*

† *Spicis androgynis, apice masculis.*

27. C. spica simplici ovata, fructibus ovato-subglobosis ore integris squama oblonga longioribus, foliis lanceolatis undulatis crenulatis, scapo basi vaginato.  
Icon. C. Fraseri. Bot. mag. 1391. Bot. rep. 639.  
On the mountains of North Carolina. Fraser. 24.  
April. v. v.
28. C. spica simplici, fructibus alternis oblongis tereti-tri-*Willdenowii.*  
queritis scabris acuminatis, squamis ovatis acuminatis,  
infima apice foliacea. Willd. sp. pl. 4. p. 211.  
Icon. Schk. car. t. Mmm. f. 145.  
In barren wet soil, common. 24. May, June. v. v.
29. C. spica simplici, fructibus oblongo-lanceolatis compres-*polytrichoi-*  
so-triquetris obtusis emarginatis, squamis oblongis ob-*des.*  
tusis mucronatis. Willd. sp. pl. 4. p. 213.  
C. leptalea. Wahlb. act. holm. 1803. p. 139.  
C. microstachya. Mich. fl. amer. 2. p. 169.  
Icon. Schk. car. t. Iii. f. 138.  
In low wet places: Pennsylvania, &c. 24. June. v. v.
30. C. spicis subquaternis pedunculatis ramosissimis; fructi-*pedunculata.*  
bus obovatis triquetris obtusis, squamis oblongis obtusis  
inucronatis. Willd. sp. pl. 4. p. 222.  
Icon. Schk. car. t. Ggg. f. 131.  
On dry hills: New England, New York, Pennsylvania,  
&c. 24. May—July. v. v.
31. C. spicis subquinis pedunculatis ovatis pendulis, fructi-*ovata.*  
bus obovatis utrinque acutis, squamis ovatis acutis  
capsulam æquantibus.  
C. ovata. Rudge in linn. trans. 7. p. 96.  
Icon. Rudge l. c. t. 9. f. 1.  
In Canada and Newfoundland. 24. +.
- †† *Spicis terminalibus masculis, reliquis androgynis.*
32. C. spica androgyna linearis-pedunculata inferne mascula,  
femineis subapproximatis binis subpedunculatis linea-*virescens.*

- ribus, fructibus globoso-triquetris obtusis pubescentibus. *Willd. sp. pl. 4. p. 251.*  
 Icon. *Schk. car. t. Mmm. f. 147.*  
 On barren dry soil, common. ♀. May. v. v.
- hirsuta.* 33. *C. spica androgyna oblonga obovata inferne mascula, femineis remotiusculis subternis subsessilibus oblongis, fructibus ovatis obtusissimis obtuse triquetris, foliis vaginisque hirsutis.* *Willd. sp. pl. 4. p. 252.*  
 Icon. *Schk. car. t. Ww. f. 172.*  
 On dry hills, frequent. ♀. May, June. v. v.
- Buxbaumii.* 34. *C. spica androgyna pedunculata obovata inferne mascula, femineis subternis remotis subpedunculatis, fructibus ellipticis triquetris obtusis obsolete bidentatis squamam oblongam mucronatam subæquantibus.* *Willd. sp. pl. 4. p. 252.*  
*C. polygama. Schk.*  
 Icon. *Schk. car. t. X. Gg. f. 76.*  
 In swamps and wet woods: New York, Pennsylvania, &c. ♀. July, Aug. v. v.
- trichocarpa.* 35. *C. spicis androgynis tribus, femineis binis pedunculatis erectis cylindraceis remotis: fructibus ovatis acuminate bicuspidatis pilosis squama ovato-lanceolata articulata longioribus.* *Schk. car. 2. p. 47.*  
*C. lanuginosa. Mich. fl. amer. 2. p. 175.*  
 Icon. *Schk. car. t. Nnn. f. 148.*  
 In woods and wet fields: New England, Pennsylvania, &c. ♀. June. v. v.
- ††† *Spicis sexu distinctis.*
- \* *Spica mascula solitaria, femineis sessilibus seu inclusopedunculatis.*
- varia.* 36. *C. spicis femineis subternis subapproximatis sessilibus subglobosis, fructibus subgloboso-triquetris rostratis bidentatis pubescentibus squama oblonga brevioribus, culmo erecto.* *Willd. sp. pl. 4. p. 259.*  
 Icon. *Schk. car. t. Uuu. f. 167.*  
 On dry hills, frequent. ♀. May, June. v. v.
- subulata.* 37. *C. spicis femineis subquaternis remotissimis sessilibus, mascula sessilis, fructibus subulatis divaricatis reflexis, culmo tenui, foliis planis.* *Mich. fl. amer. 2. p. 173.*  
 In Canada. *Michaux. +.*
- marginata.* 38. *C. spicis femineis subgeminis approximatis subglobosis subsessilibus, fructibus globosis tomentosis bidentatis*

- squama ob  
annotinis cu  
Icon. *Schk.*  
On dry hills a
39. *C. spica mas  
sessilibus a  
obliquis pu  
æquantibus*  
Icon. *Schk.*  
In wet meado
40. *C. spicis femi  
approximat  
ovatis vent  
dentatis s  
Willd. sp. 1*  
Icon. *Schk.*  
β. Icon. *Schk.*  
In swamps ar  
v. v.
41. *C. spica femi  
tea setace  
amer. 2. p. p.*  
In bogs and s  
July. v. v.
42. *C. spicis fen  
approximat  
ovatis ven  
ore bicuspi  
longioribus*  
Icon. *Schk.*  
In swamps ar
43. *C. spicis fen  
inclusa pec  
rostro curv  
gioribus.*  
C. Oederi.  
Icon. *Schk.*  
1. t. 65.  
In wet fields
44. *C. spicis fer  
dunculata,  
rostellatis  
gioribus.*

- pubescens.  
v. v.  
ne mascula,  
bus oblongis,  
uetris, foliis  
252.  
v. v.  
ferne mascula-  
culatis, fruc-  
te bidentatis  
oæquantibus.  
Pennsylvania,  
pedunculatis  
vatis acuminata-  
lanceolata aristata.  
Pennsylvania,  
seu inclusa-  
tis sessilibus  
tris rostratis  
brevioribus,  
v. v.  
is sessilibus,  
catis reflexis,  
er. 2. p. 173.  
s subglobosis  
sis bidentatis
- squama oblongo-ovata majoribus, foliis radicalibus annotinis culmo longioribus. *Willd. sp. pl. 4. p. 261.*  
Icon. *Schk. car. t. Lll. f. 143.*  
On dry hills and rocks, common. 4. April, May. v. v.
39. C. spica mascula lanceolata, femineis geminis ovatis sessilibus approximatis, fructibus ovatis rostratis ore obliquis pubescentibus squamam ovatam acutam subæquantibus. *Willd. sp. pl. 4. p. 263.*  
Icon. *Schk. car. t. Bbb. f. 182.*  
In wet meadows : Pennsylvania, &c. 4. June. v. v.
40. C. spicis femineis tribus inclusa-pedunculatis ovatis subapproximatis, bracteis longissimis foliaceis, fructibus ovatis ventricosis nervosis longissime rostratis ore bidentatis squama lanceolata mucronata longioribus. *Willd. sp. pl. 4. p. 266.*  
Icon. *Schk. car. t. Ggg. f. 130.*  
β. Icon. *Schk. car. t. Hhh. f. 134.*  
In swamps and wet woods, common. 4. June, July.  
v. v.
41. C. spica feminea plerumque solitaria sessili-ovata, bractea setacea, fructibus globosis glabris. *Mich. fl. amer. 2. p. 174.*  
In bogs and shady woods : Canada, Vermont, &c. 4. July. v. v.
42. C. spicis femineis tribus incluse pedunculatis oblongis approximatis, bracteis longissimis foliaceis, fructibus ovatis ventricosis nervosis longissime conico-rostratis ore bicuspidatis, squama oyata mucronata multoties longioribus. *Willd. sp. pl. 4. p. 266.*  
Icon. *Schk. car. t. Ddd. f. 123. et t. Iii. f. 194.*  
In swamps and woods, frequent. 4. June—Aug. v. v.
43. C. spicis femineis subternis subapproximatis ellipticis incluse pedunculatis, fructibus ovatis reflexis rostratis, rostro curvato bidentato, squama ovato-lanceolata longioribus. *Willd. sp. pl. 4. p. 268.*  
C. Oederi. *Schk. car. 1. p. 67.*  
Icon. *Schk. car. t. H. f. 36. et t. F. f. 26. Host. gram. 1. t. 65.*  
In wet fields and meadows, frequent. 4. June. v. v.
44. C. spicis femineis binis tri- s. quadrifloris inferiore pendulata, fructibus subrotundo-triquetris obovatis rostellatis ore integris squama oblonga mucronata longioribus. *Willd. sp. pl. 4. p. 279.*
- vestita.  
tentaculata.  
rostrata.  
miliaris.  
lupulina.  
flava.  
oligocarpa.

- C. paupercula.* *Mich. fl. amer.* 2. p. 172?  
Icon. *Schk. car. t. Vvv. f. 170.*  
In wet places: New York, Pennsylvania, &c. 4. May.  
v. v.
- folliculata.* 45. *C. spica feminea subsolitaria subsexflora subexerta pendunculata, fructibus ovatis ventricosis nervosis rostratis ore bipartitis squama ovata longioribus.* *Willd. sp. pl. 4. p. 281.*  
Icon. *Schk. car. t. N.f. 52.*
- major. 46. *C. spicis femineis quaternis sessilibus infima exerte pendunculata, fructibus ovato-triquetris pubescentibus rostratis bidentatis, squama oblonga mucronata longioribus, culmo foliis pubescentibus.* *Willd. sp. pl. 4. p. 281.*  
Icon. *Schk. car. t. Eee. f. 126.*  
In shady woods: Pennsylvania, Virginia, &c. 4. May.  
v. v.
- pubescens.* 47. *C. spicis pedunculatis femineis quaternis distantibus, fructibus ellipticis triquetris pedicellatis glabris squama ovata cuspidata brevioribus, bracteis vaginatis apice subfoliaceis, radicalibus lanceolatis nervosis.* *Willd. sp. pl. 4. p. 257.*  
*C. latifolia,* *Wahlb. act. holm.* 1803. p. 156.  
Icon. *Schk. car. t. U. f. 70. et t. Kkkk. f. 195.*  
In dry woods, very common. 4. May, June. v. v.
- anceps.* 48. *C. spicis femineis tribus remotis, inferioribus pedunculatis, fructibus ovatis nervosis ore membranaceis squama oblonga mucronata longioribus.* *Willd. sp. pl. 4. p. 278.*  
*C. striatula,* *Mich. fl. amer.* 2. p. 173.  
Icon. *Schk. car. t. Fff. f. 128.*  
In wet fields on the sides of ditches, from New Jersey to Carolina. 4. June. v. v.
- granularis.* 49. *C. spicis femineis tribus remotis, binis inferioribus pedunculatis, fructibus globoso-ovatis nervosis ventricosis brevissime rostellatis ore obsolete emarginato squama ovato-lanceolata longioribus.* *Willd. sp. pl. 4. p. 279.*

*C. lenticularis*  
Icon. *Schk. c*  
In barren mead  
4. May—

50. *C. spicis femi  
fima longe  
tus squam*  
*p. 280.*  
Icon. *Schk. c*  
In dry woods,

51. *C. spicis femi  
fima longe  
curvis ore i  
bus.* *Schk.*  
Icon. *Schk. c*  
In woods and  
v. v.

52. *E. spicis femi  
remote pe  
obtusis squa  
sp. pl. 4. p.*  
*C. paupercula*  
Icon. *Schk.*  
On dry grassy

53. *C. spica mas  
mucronatis;  
dunculata,  
bifidis squa  
sp. pl. 4. p.*  
*C. erinacea.*  
Icon. *Schk.*  
In swamps an  
v. v.

54. *C. spicis fen  
sessili, reli  
natis bicusp  
æquantibus*  
Icon. *Schk.*  
In wet mead

55. *C. spicis femi  
dunculus ce  
rostratis bil  
ribus.* *W*  
*C. debilis.* 1

24. May.  
bexerta per-  
ervosis ros-  
us. Willd.  
p. 97. t. 9.  
v. v.  
exerte pe-  
bescensibus  
eronata lon-  
d. sp. pl. 4.  
24. May.  
culatis, va-  
distantibus,  
bris squama  
inatis apice  
sis. Willd.  
95.  
ne. v. v.  
us peduncu-  
naceis squa-  
ld. sp. pl. 4.  
New Jersey  
rioribus pe-  
sis ventrico-  
ginato squa-  
l. sp. pl. 4.
- C. lenticularis. Mich. fl. amer. 2. p. 172.  
Icon. Schk. car. t. Vvv. f. 169.  
In barren meadows and woods, from Canada to Carolina.  
24. May—July. v. v.
50. C. spicis femineis binis remotis, suprema subsessili, infima longe pedunculata, fructibus oblongo-conicis obtusis squamam aristam æquantibus. Willd. sp. pl. 4. p. 280.  
Icon. Schk. car. t. Vvv. f. 168.  
In dry woods, common. 24. June. v. v.
51. C. spicis femineis binis remotis, suprema subsessili, infima longe pedunculata, fructibus obovatis apice recurvis ore integris squamam obtuso-ovatam longioribus. Schk. car. 2. p. 68.  
Icon. Schk. car. t. Ggg. f. 100. et t. Oooo. f. 207.  
In woods and dry meadows, frequent. 24. May, June.  
v. v.
52. E. spicis femineis tribus 6—8-floris distantibus, infima remote pedunculata, fructibus oblongis ventricosis obtusis squama ovata mucronata majoribus. Willd. sp. pl. 4. p. 281.  
C. paupercula. Mich. fl. amer. 2. p. 172.  
Icon. Schk. car. t. Kkk. f. 141.  
On dry grassy hills, common. 24. April, May. v. v.
53. C. spica mascula solitaria, squamis ovato-oblongis submucronatis; femineis cylindraceis binis, infima pedunculata, fructibus ovatis multinerviis rostratis ore bifidis squama oblonga aristata longioribus. Willd. sp. pl. 4. p. 282.  
C. erinacea. Muhl. in litt....  
Icon. Schk. car. t. Fff. f. 127.  
In swamps and boggy woods, frequent. 24. June, July.  
v. v.
54. C. spicis femineis tribus oblongis distantibus, suprema sessili, reliquis pedunculatis, fructibus ovatis acuminatis bicuspidatis squamam ovatam acuminatam subæquantibus. Willd. sp. pl. 4. p. 271.  
Icon. Schk. car. t. T. et Yy. f. 63.  
In wet meadows, frequent. 24. June, July. v. v.
55. C. spicis femineis subquaternis remotis filiformibus pedunculis cernuis, fructibus distantibus alternis oblongis rostratis bifidis squama ovata mucronata duplo longioribus. Willd. sp. pl. 4. p. 297.  
C. debilis. Mich. fl. amer. 2. p. 172.
- conoidea.  
tetanica.  
laxiflora.  
hystericina.  
distans.  
flexuosa.

## TRIANDRIA MONOGYNIA. Carex.

- C. tenuis. *Rudge in linn. trans. t. 7. p. 97.*  
Icon. *Schk. car. t. Ddd. et Aaaa. f. 124.*  
In sandy wet meadows: Canada to Carolina. ♀. June.  
v. v.
- digitalis.* 56. C. spicis feminineis subternis remotis filiformibus pedunculatis cernuis, fructibus ellipticis obtusis squama oblongo-lanceolata longioribus. *Willd. sp. pl. 4. p. 298.*  
In Pennsylvania. *Muhl.* ♀. +.
- \*\*\* *Spica mascula solitaria, feminineis pedunculatis, vaginæ subnullæ.*
- umbellata.* 57. C. spicis feminineis tribus pedunculatis octofloris ovatis verticillatis, fructibus ovatis pubescentibus rostratis ore integris squamam ovato-lanceolatam æquantibus. *Willd. sp. pl. 4. p. 290.*  
Icon. *Schk. car. t. Wwww. f. 171.*  
On dry sunny hills: New York to Virginia. ♀. May.  
v. v.
- miliacea.* 58. C. spicis feminineis tribus filiformibus, summa subsessili, reliquis pedunculatis, fructibus ovatis triquetris breve rostratis ore integris squama oblonga emarginata aristata longioribus. *Willd. sp. pl. 4. p. 290.*  
Icon. *Schk. car. t. Ooo. f. 151.*  
In wet meadows: Canada to Carolina. ♀. July. v. v.
- Pseudo-Cyperus.* 59. C. spicis feminineis quaternis geminatis pedunculatis pendulis cylindricis, fructibus ovato-lanceolatis bicuspidatis reflexis squamam setaceam æquantibus. *Willd. sp. pl. 4. p. 295.*  
Icon. *Schk. car. t. Mm. f. 102.*  
In ditches and swamps, frequent. ♀. June, July. v. v.
- \*\*\* *Spicis masculis pluribus.*
- recurva.* 60. C. spicis masculis geminis, feminineis cylindricis subternatis pedunculatis pendulis, fructibus ellipticis obtusis saepe scabriusculis squamam ovato-cuspidatam æquantibus. *Willd. sp. pl. 4. p. 298.*  
C. flaccia. *Schk. car. 1. p. 117.*  
C. glauca. *Scop.*  
Icon. *Schk. car. t. O. P. f. 57. et t. Zz. f. 113.*  
In bogs and wet woods, common. ♀. June. v. v.
- pellita.* 61. C. spicis masculis geminis, feminineis geminis cylindraceis erectis remotis, superiore sessili, fructibus ovatis bifidis pilosis squama oblonga aristata brevioribus. *Willd. sp. pl. 4. p. 302.*

C. striata. *I*  
Icon. *Schk.*  
In wet mead  
v. v.

62. C. spicis mas  
draceis pe  
rostratis bi  
ibus. *W*  
Icon. *Schk.*  
β. *Rudge in l*  
In deep swa  
Carolina.

63. C. spicis mas  
latis cylind  
bicuspidat  
triquetro.  
Icon. *Schk.*  
In swamps a  
v. v.

64. C. spicis ma  
dunculatis  
tratis bifur  
joribus.  
Icon. *Schk.*  
In bogs, ve

48

1. S. culmo fo  
ralibus pa  
4. p. 314  
Carex lithos  
In sandy b  
July, Au
2. S. culmo s  
glomerat  
mis glab  
rugoso-va  
In wet fie  
July. v.
3. S. culmo si  
spica glo  
glumis se

C. striata. *Mich. fl. amer.* 2. p. 174?  
 Icon. *Schk. car. t. Nnn. f.* 149 et 150.  
 In wet meadows and woods, frequent. 4. June, July.  
 v. v.

62. C. spicis masculis quaternis, femineis binis erectis cylindraceis pedunculatis, fructibus oblongis multinerviis rostratis bifurcatis squama oblonga mucronata longioribus. *Willd. sp. pl.* 4. p. 306.

Icon. *Schk. car. t. Ooo. f.* 152.

$\beta$ . *Rudge in linn. trans. t. 7. p. 99. t. 10. f. 2.*  
 In deep swamps and on the edges of ponds: Canada to Carolina. 4. June, July. v. v.

63. C. spicis masculis tribus, femineis subgeminis pedunculatis cylindraceis, fructibus oblongis inflatis rostratis bicuspidatis squama lanceolata majoribus, culmo acute triquetro. *Willd. sp. pl.* 4. p. 307.

Icon. *Schk. car. t. Ss. f.* 106.

In swamps and wet meadows, common. 4. May—July.  
 v. v.

64. C. spicis masculis tribus, femineis binis cylindraceis pedunculatis erectiusculis, fructibus ovato-globosis rostratis bifurcatis, rostris hispidis squama lanceolata majoribus. *Willd. sp. pl.* 4. p. 309.

Icon. *Schk. car. t. Uuu. f.* 166.

In bogs, very common. 4. June. v. v.

*lacustris.*

*gigantea.*

*vesicaria.*

*bullata.*

*reticularis.*

*verticillata.*

*interrupta.*

#### 48. SCLERIA. *Gen. pl.* 1408.

1. S. culmo foliisque glabris, paniculis terminalibus et lateralis paucifloris, nucibus reticulatis. *Willd. sp. pl.* 4. p. 314. *Mich. fl. amer.* 2. p. 167.

*Carex lithosperma.* *Walt. fl. car.* 227.

In sandy barren fields: New Jersey to Carolina. 4. July, Aug. v. v.

2. S. culmo simplicissimo triquetro foliisque glabris, spica glomerata nuda, glomerulis alternis distantibus, glumis glabris, nucibus globosis mucronatis transversim rugoso-verrucosis. *Willd. sp. pl.* 4. p. 317.

In wet fields: Virginia, Carolina. *Muhl.* 4. June, July. v. v.

3. S. culmo simplicissimo triquetro foliisque pubescentibus, spica glomerata nuda, glomerulis alternis distantibus, glumis setosis, nucibus globosis mucronatis transver-

4. June.  
 ibus pedun-  
 squama ob-  
 ol. 4. p. 298.

*itis, vaginæ*  
 floris ovatis  
 rostratis ore  
 æquantibus.

4. May.

*ia subsessili,*  
*juetris breve*  
*rginata aris-*

July. v. v.  
 iculatis pen-  
 itis bicuspis-  
 us. *Willd.*

July. v. v.

*tricis subter-*  
*pticis obtusis*  
*itam æquan-*

113.  
 ie. v. v.  
 cylindraceis  
 s ovatis bi-  
 bus. *Willd.*

sim rugoso-verrucosus. Willd. sp. pl. 4. p. 317.  
Mich. fl. amer. 2. p. 168.

In wet meadows of Lower Carolina and Florida. ♀. +.

*hirtella.*

4. S. culmo simplicissimo triquetro foliisque pubescentibus,  
spica glomerata nuda, glomerulis alternis distantibus,  
glumis setosis, nucibus elliptico-globosis mucronatis  
lævibus nitidis. Willd. sp. pl. 4. p. 318. Mich. fl.  
amer. 2. p. 168.

In dry woods on slate rocks: Virginia, Carolina. ♀.  
June. v. v. The smallest species.

*ciliata.*

5. S. culmo erecto glabriuscule, foliis ciliatis fasciculo ter-  
minali ciliato, nucibus globosis scabriusculis. Willd.  
sp. pl. 4. p. 318. Mich. fl. amer. 2. p. 167.

In dry woods: Virginia, Carolina, &c. ♀. July. v. v.

*pauciflora.*

6. S. culmo erecto simplici triquetro scabro, foliis anguste  
linearibus canaliculatis margine scabris, fasciculis pau-  
cifloris binis terminalibus unico remotissimo axillari,  
nucibus globosis acutiusculis transversim rugosis.  
Willd. sp. pl. 4. p. 318.

S. oligantha. Mich. fl. amer. 2. p. 167.

In low grassy places in the woods of Carolina. ♀. +.

*triglomerata.*

7. S. culmo erecto simplici triquetro scabro, foliis margine  
scabris, fasciculis paucifloris terminalibus, glumis  
ovatis mucronatis scabris, nucibus globosis acutis ru-  
gosus. Mich. fl. amer. 2. p. 168.

In dry swamps and old fields: Pennsylvania to Carolina.  
♀. June, July. v. v.

#### 49. ZEA. Gen. pl. 1403. Indian Corn.

*Mays.*

1. Z. foliis integerrimis. Willd. sp. pl. 4. p. 200.

Cultivated throughout all America, and sometimes found  
as it were in a native state. It would take up too  
much room to mention all the numberless varieties;  
but one of those brought by M. Lewis, Esq. from the  
Mandan nation, on the Missouri, deserves particular  
attention, as it ripens sooner and produces as excel-  
lent ears as any sort I know. It would be calculated  
to cultivate in climates not quite so warm as all the  
other sorts require.

#### 50. KYLLINGIA. Gen. pl. 91.

*pumila.*

1. K. capitulo globoso sessili solitario, involucro brevi, cul-

TRIAN

mo setace  
p. 380. A

In the west  
♀. June.

2. K. capitulis s  
phylo, gl  
flosculis m  
Scirpus cepha  
In inundated  
v. s.

51. MA

1. Mapania. A  
In remote pa  
v. s. A be  
made by N  
every respe

52. DICHI

1. D. glabra, in  
pl. 2. p. 24  
In wet places  
v. s.

2. D. pilosa, in  
pl. 2. p. 24  
Schœnus stell  
Icon. Sloan  
In inundated  
v. v.

53. SCHŒN

1. S. pedunculi  
culmo trig  
p. 219.  
Icon. Rottb  
In Carolina.

2. S. pedunculi  
cis subglo  
dis. Vahl  
In Carolina.

4. p. 317.  
orida. 4. +.  
abescentibus,  
s distantibus,  
mucronatis  
Mich. fl.  
orolina. 4.  
fasciculo ter-  
alis. Willd.  
67.  
July. v. v.  
foliis anguste  
fasciculis pau-  
cimo axillari,  
im rugosissi-
- mo setaceo, flosculis diandris. *Vahl. enum. pl. 2.*  
p. 380. *Mich. fl. amer. 1. p. 28.*  
In the western parts of South Carolina and Georgia.  
4. June. v. v.
2. K. capitulis subternis sessilibus glomeratis, involucro 3-  
phylo, glumis oblongis: valvulis cuneato-lanceolatis,  
flosculis monandris. *Mich. fl. amer. 1. p. 29.*  
*Scirpus cephalotes. Walt. fl. car. 71.*  
In inundated places: Carolina, Florida, &c. 4. July.  
v. s.

*maculata.*

51. MAPANIA. *Aubl. fl. gui. 1. p. 47.*
1. Mapania. *Aubl. fl. gui. 1. p. 47. t. 17.*  
In remote parts of the Cherokee mountains. 4. July.  
v. s. A beautiful grass, which I found in a collection  
made by Mr. Kinn of Philadelphia, corresponded in  
every respect with Aublet's plant.
52. DICHROMENA. *Mich. fl. amer. 1. p. 37.*
1. D. glabra, involucris basi utrinque coloratis. *Vahl. enum. leucocephala.*  
*pl. 2. p. 240. Mich. fl. amer. 1. p. 37.*  
In wet places in Lower Carolina and Georgia. 4. July.  
v. s.
2. D. pilosa, involucris supra basin coloratis. *Vahl. enum.* *ciliata.*  
*pl. 2. p. 240.*  
*Schoenus stellatus. Swartz. fl. ind. occid. 1. p. 102.*  
Icon. *Sloan. hist. jam. 1. t. 78. f. 1.*  
In inundated places of Georgia and Florida. 4. July.  
v. v.

*sylvatica.*

53. SCHCENUS. *Linn. et Vahl. enum. pl. 2. p. 207.*
1. S. pedunculis axillaribus terminalibusque subtrifloris,  
culmo trigono foliisque setaceis. *Vahl. enum. pl. 2.*  
*p. 219.*  
Icon. *Rottb. gram. t. 21.*  
In Carolina. *Vahl. 4. +.*
2. S. pedunculis axillaribus terminalibusque tristachyis, spi-  
cis subglobosis pedicellatis, foliis filiformibus hispi-  
dis. *Vahl. enum. pl. 2. p. 219.*  
In Carolina. *Lam. 4. +.*

*setaceus.**hispidulus.*

- TRIANE
54. RHYNCHOSPORA. *Vahl. enum. pl. 2. p. 229.*
- sparsa.* 1. R. corymbis diffusis axillaribus compositis, terminalibus decompositis, pedunculis capillaribus, seminibus rostro longioribus. *Vahl. enum. pl. 2. p. 230.*  
*Schœnus sparsus. Mich. fl. amer. 1. p. 35.*  
*Schœnus miliaceus. Lam. illustr. 1. p. 137.*  
*In shady woods: Virginia—Georgia. June. v. v.*
- micrantha.* 2. R. corymbis diffusis axillaribus simplicibus, terminalibus compositis, pedunculis capillaribus, culmo foliisque filiformibus glabris. *Vahl. enum. pl. 2. p. 231.*  
*Schœnus rariflorus. Mich. fl. amer. 1. p. 35.*  
*In Georgia. Mich. 4. +.*
- laxa.* 3. R. corymbis terminalibus compositis, spicis solitariis, rostello seminum longissimo, culmo acutangulo scabro. *Vahl. enum. pl. 2. p. 231.*  
*Schœnus longirostris. Mich. fl. amer. 1. p. 35.*  
*Schœnus corniculatus. Lam. illustr. 1. p. 132.*  
*In wet woods and near ponds in Virginia and Carolina. 4. July. v. v.*
- inexpansa.* 4. R. corymbis coarctatis axillaribus simplicibus compositisque, terminalibus decompositis, culmo filiformi. *Vahl. enum. pl. 2. p. 232.*  
*Schœnus inexpansus. Mich. fl. amer. 1. p. 35.*  
*In Carolina. Mich. 4. +.*
- fascicularis.* 5. R. spicis corymboso-fasciculatis, foliis linearibus latitudine culmi. *Vahl. enum. pl. 2. p. 234.*  
*Schœnus fascicularis. Mich. fl. amer. 1. p. 37.*  
*In Carolina. 4. v. s.*
- glomerata.* 6. R. spicis corymboso-fasciculatis remotissimis geminatis, culmo obtusangulo, foliis linearibus. *Vahl. enum. pl. 2. p. 234.*  
*Schœnus glomeratus. Willd. sp. pl. 1. p. 266.*  
*In wet meadows along ditches: Pennsylvania—Carolina. Walter. 4. June, July. v. v.*
- distans.* 7. R. spicis corymboso-fasciculatis remotissimis solitariis, culmo obtusangulo, foliis filiformibus. *Vahl. enum. pl. 2. p. 235.*  
*Schœnus distans. Mich. fl. amer. 1. p. 36.*  
*Schœnus axillaris. Lam. illustr. 1. p. 137.*  
*In ponds which are dry in summer: Virginia, Carolina. 4. July, Aug. v. v.*

8. R. spicis conbatis, bracteatis, Vahl. enum. Schœnus capillaris. In Carolina.
9. R. spicis conbatis foliisque. Schœnus ciliatus. In Carolina.
10. R. spicis corporeo. Vahl. enum. Schœnus albus. Icon. Engl.  $\beta$  Schœnus fuscus. Schœnus setiger. In bog-meadow.
55. FIMBI
1. F. spicis ovata, diphylloidea. Vahl. enum. Scirpus pubescens. In low ground. v. v.
2. F. spicis ovata, umbellata. p. 92. Scirpus castaneus. Scirpus carolinianus. Schœnus umbrosus. In Carolina and Florida.
3. F. spicis cylindrica, phyllo rigidula. enum. pl. 2. Scirpus palustris. In North America.
4. F. spicis oblonga, rigida umbellata. pl. 2. p. 26. Scirpus spadiceus. Icon. Sloaniana. On the banks. v. s.

2. p. 229.  
terminalibus  
inibus rostro

June. v. v.  
terminalibus  
no foliisque  
231.

cis solitariis,  
tangulo sca-  
35.

132.  
ind Carolina.

bus composi-  
mo filiformi.

35.

aribus latitu-

37.

nis geminatis,  
l. enum. pl. 2.

66.  
ia—Carolina.

nis solitariis,  
Vahl. enum.

nia, Carolina.

8. R. spicis corymboso-fasciculatis approximatis subgemini-  
natis, bracteis glabris, culmo filiformi acutangulo.  
*Vahl. enum. pl. 2. p. 235.*  
*Schoenus capitellatus. Mich. fl. amer. 1. p. 36.*  
In Carolina. *Mich. 4. +.*
9. R. spicis corymboso-fasciculatis terminalibus, bracteis  
foliisque obtusiusculis ciliatis. *Vahl. enum. pl. 2.*  
*p. 235.*  
*Schoenus ciliaris. Mich. fl. amer. 1. p. 36.*  
In Carolina and Florida. *4. v. s.*
10. R. spicis corymboso-fasciculatis, culmo foliisque setaceis.  
*Vahl. enum. pl. 2. p. 236.*  
*Schoenus albus. Willd. sp. pl. 1. p. 267.*  
Icon. *Engl. bot. 985. Fl. dan. t. 320.*  
 $\beta$  *Schoenus fuscus. Willd. sp. pl. 1. p. 262.*  
*Schoenus setaceus. Thouil. fl. par. 19.*  
In bog-meadows, common. *4. July, Aug. v. v.*
55. FIMBRISTYLIS. *Vahl. enum. pl. 2. p. 285.*
1. F. spicis ovatis pubescentibus, involucro involucellisque  
diphyllis umbella composita umbellulisque longioribus.  
*Vahl. enum. pl. 2. p. 289.*  
*Scirpus puberulus. Mich. fl. amer. 1. p. 31.*  
In low grounds: Virginia to Florida. *4. July, Aug.*  
*v. v.*
2. F. spicis ovato-oblongis obtusis, involucro diphyllo rigido  
umbellam subcompositam æquante. *Vahl. enum. pl. 2.*  
*p. 92.*  
*Scirpus castaneus. Mich. fl. amer. 1. p. 31.*  
*Scirpus carolinianus. Lam. illustr. 1. p. 142.*  
*Schoenus umbellatus. Walt. fl. car. 70.*  
In Carolina and Florida. *4. July. v. v.*
3. F. spicis cylindricis obtusissimis, involucro submono-  
phyllo rigido umbellam simplicem æquante. *Vahl.*  
*enum. pl. 2. p. 293.*  
*Scirpus palustris altissimus, &c. Gron. fl. virg. 11.*  
In North America. *Bosc. 4. +.*
4. F. spicis oblongis teretibus acutis, involucro diphyllo  
rigido umbellam decompositam æquante. *Vahl. enum.*  
*pl. 2. p. 294.*  
*Scirpus spadiceus. Willd. sp. pl. 1. p. 305.*  
Icon. *Sloan. hist. 1. t. 76. f. 2.*  
On the banks of rivers in Carolina and Florida. *4. July.*  
*v. s.*

## 56. CYPERUS. Gen. pl. 93.

## \* Culmo tereti.

- articulatus.* 1. C. spiculis compressis elongatis nitidiusculis, involucro subnullo, culmo articulato. *Vahl. enum. pl. 2. p. 301.*  
Icon. *Sloan. hist. jam. 1. t. 81. f. 1.*  
In small slow streams: Carolina and Georgia. 24. July.  
*v. v.*

## \*\* Culmo triquetro.

## || Spiculis aggregatis.

- Kyllingæoides.* 2. C. capitulo globoso, spiculis oblongis convexis suboctofloris, involucro tetraphyllo foliisque carinatis laxis. *Vahl. enum. pl. 2. p. 312.*  
Icon. *Rottb. gram. t. 4. f. 5.* C. dubius, p. 20.  
In wet pine-woods: New Jersey, Carolina, &c. 24. June. *v. v.*

- uncinatus.* 3. C. capitulo subgloboso glomerato, spiculis linearibus suboctofloris: valvulis lanceolatis recurvato-acuminatis, involucro subdiphylio longissimo.  
On overflowed gravelly banks of rivers and lakes: Canada, in New York, on Lake Champlain. 25. June.  
*v. v.*

*Planta* pusilla. *Folia* linearia glabra, longitudine culmi. Culmus setaceus, acute triquetrus, biuncialis. *Involucrum* foliola 2. interdum 3. unum culmo longius, structura foliorum. *Capitulum* solitarium, conglomeratum, ratione plantæ magnum. *Spiculæ* numerosæ, divaricatæ, linearis-oblongæ, 6—8-floræ: valvulae lanceolato-ovatae, longe acuminatae, apice divaricato quasi uncinatim reflexæ, nervosæ, pallide castaneæ, margine albidae. C. *pygmæo* affinis.

- pœfornis.* 4. C. spiculis oblongis complanatis fasciculato-corymbosis: fasciculis sessilibus pedunculatisque, involucro triphylio longissimo.

In sandy fields: South Carolina, &c. 24. July. *v. v.*

*Planta* spithamea. *Folia* angusto-linearia, glabra, vix longitudine culmi. Culmus trigonus, glaber. *Involucrum* triphyllum; foliolis 2. longissimis. *Fasciculis* florum ovatis, sessilibus, uno alterove brevi pedunculato. *Ochreæ* truncatæ, coloratae. *Spiculæ* ovato-oblongæ, circiter octo, breves, quatuor—sex-floræ: valvulae ovatae, obtusæ, carinatae, flavescentes. An C. *difformis*?

|| Umbellæ

5. C. spiculis crum dij p. 318.  
Scirpus aut Cyperus jur Cyperus co Icon. Rott On the edge lina. 24 valves pu lindric.

6. C. spiculis c vulvis mu enum. pl. Icon. Rott f. 1. In low mea v. v.

7. C. spiculis aphyllis, p. 106.  
In swamps c

8. C. spiculis valvulis o giore. V p. 283.

- C. compressi C. glomeratu C. Eragrosti C. declinatus C. monandru Icon. Jacq. On the bord

9. C. spiculis o aphylllo lor p. 327.  
In Louisiana

10. C. spiculis umbella su setaceis.  
In Carolina.

|| | Umbellæ radiis elongatis : spiculis terminalibus, umbellatis.

5. C. spiculis linearibus digitato-subternis, umbella involucrum diphylum subæquante. *Vahl. enum. pl. 2. p. 318.*

*Scirpus autumnalis. Rottb. gram. p. 58.*

*Cyperus juncoides. Lam. illustr. 1. p. 147.*

*Cyperus complanatus. Willd. sp. pl. 1. p. 270.*

Icon. *Rottb. gram. t. 17. f. 3.*

On the edge of ponds and ditches in Virginia and Carolina. 24. July, Aug. v. v. Spikes very slender; valves purple with a green carina; stem nearly cylindric.

6. C. spiculis digitatis subquaternis lanceolato-oblongis valvulis mucronatis latoribus membranaceis. *Vahl. enum. pl. 2. p. 324. Willd. sp. pl. 1. p. 282.*

Icon. *Rottb. gram. t. 9. f. 3. Sloan. hist. jam. 1. t. 17. f. 1.*

In low meadows: Pennsylvania to Carolina. 24. July. v. v.

7. C. spiculis oblongo-ovatis obtusis, umbellulis confertis aphyllis, foliis angustis. *Rich. in act. soc. linn. par. p. 106.*

In swamps of Carolina. 24. July. v. v.

8. C. spiculis lanceolatis subgloboso-capitatis compactis, valvulis ovatis uninervibus, involucro umbella longiore. *Vahl. enum. pl. 2. p. 326. Willd. sp. pl. 1. p. 283.*

*C. compressus. Jacq. hort. vind. 3. t. 12.*

*C. glomeratus. Walt. fl. car. 70.*

*C. Eragrostis. Lam. illustr. 1. p. 146.*

*C. declinatus. Mœnch.*

*C. monandrus. Roth. catalect. 1. p. 3.*

Icon. *Jacq. l. c.*

On the borders of ponds, frequent. 24. July, Aug. v. v.

9. C. spiculis ovato-lanceolatis capitatis, involucro subhexaphyllo longissimo margine scabro. *Vahl. enum. pl. 2. p. 327.*

In Louisiana. *Jussieu. 24. +.*

10. C. spiculis linearibus globoso-capitatis patentissimis, umbella subuniradiata, involucris triphyllis culmisque setaceis. *Vahl. enum. pl. 2. p. 328.*

In Carolina. *Vahl. 24. +.*

- ||| Umbella radiis elongatis: spiculis alternis.
- filicinus.* 11. C. spiculis linear-lanceolatis remotiusculis laxe imbricatis, involucro triphylo umbella longiore foliisque laxis. *Vahl. enum. pl. 2. p. 332.*  
In swamps of Virginia and North Carolina. 4. July.  
v. v.
- flavescens.* 12. C. spiculis linear-lanceolatis confertis, involucro triphylo umbella trifida longiore. *Vahl. enum. pl. 2. p. 335. Willd. sp. pl. 1. p. 279.*  
Icon. *Fl. graeca. t. 47.*
- castaneus.* β C. pusillus, spiculis brevioribus castaneis.  
Common in low grounds. 4. June, July. v. v.
- Hydra,* 13. C. spicis corymbosis, spiculis linear-lanceolatis distantibus, foliis recurvatis, tuberibus oblongo-subrotundis. *Vahl. enum. pl. 2. p. 344. Mich. fl. amer. 1. p. 27.*  
In cultivated fields and barren woods: Virginia to Florida. 4. July. v. v.
- tuberosus.* 14. C. spiculis linear-lanceolatis convexiusculis, involucro triphylo umbella quinqueradiata longiore. *Vahl. enum. pl. 2. p. 340.*  
Icon. *Rottb. gram. t. 7. f. 1.* Nut-grass  
On the banks of Delaware and other rivers: Pennsylvania to Carolina. 4. June, July. v. v. The roots are very sweet, and eaten by children.
- tenuiflorus.* 15. C. spicis corymbosis, spiculis linearibus convexiusculis, involucro umbella longiore, culmo folioso. *Vahl. enum. pl. 2. p. 347.*  
Icon. *Rottb. gram. t. 14. f. 1.*  
In wet fields in Carolina and Georgia. 4. July. v. v.
- virens.* 16. C. spicis corymbosis, spiculis ovalibus, involucro longissimo. *Vahl. enum. pl. 2. p. 353. Mich. fl. amer. 1. p. 28.*  
In wet meadows and woods: Pennsylvania to Carolina. 4. July, Aug. v. v.
- odoratus.* 17. C. spicis corymbosis, spiculis subulatis remotis distichis, valvulis subdistantibus, umbellis patentissimis involucellum subaequantibus. *Vahl. enum. pl. 2. p. 356. Willd. sp. pl. 1. p. 284.*  
Icon. *Sloan. hist. jam. 1. t. 74. f. 1. et t. 8. f. 1.*  
On the banks of rivers: Pennsylvania to Florida, rare. 4. Aug. v. v.
- strigosus.* 18. C. spicis oblongis laxis, spiculis subulatis patentibus re-

motiusculis,  
ternis,  
p. 281.

In swamps

19. C. spiculis fructiferis,  
*enum. p.*

C. elegans,

In boggy v.

Aug. t

20. C. spicis d  
culis dis  
p. 362.

Icon. *Ro*

In sandy w

21. C. spicis c  
lulis inv  
acutang

C. virginic

In Virginii

22. C. spicis c  
lis nume  
valvulis  
lam sub

In ditches

Aug.

*Folia radi*

*rinaque*

*triquet*

*teriora*

*Umbelle*

*oblonga*

(circiter

*ciosi?*

I have na

through

put in

plants

57.

1. D. culmo  
*Pers. e*  
Pleuranth  
*Cyperus s*

motiusculis, involucellis subnullis, umbellulæ radiis alternis. *Vahl. enum. pl. 2. p. 358.* *Willd. sp. pl. 1. p. 281.*

In swamps and ditches, common.  $\frac{4}{4}$ . July. v. v.

19. C. spiculis linear-lanceolatis, valvulis oblongis obtusis, fructiferis patentibus, involucro longissimo. *Vahl. enum. pl. 2. p. 360.* *Mich. fl. amer. 1. p. 27.* *flavicomus.*  
C. elegans. *Walt. fl. car. 70.*

In boggy woods: Pennsylvania to Carolina.  $\frac{4}{4}$ . July, Aug. v. v:

20. C. spicis distichis, spiculis filiformibus patentibus, flosculis distantibus, umbella stricta. *Vahl. enum. pl. 2. p. 362.* *Willd. sp. pl. 1. p. 289.* *distans.*  
Icon. *Rottb. gram. t. 10.* C. elatus.

In sandy wet woods: Carolina and Georgia.  $\frac{4}{4}$ . v. s.

21. C. spicis corymbosis, spiculis subulatis distichis, umbellulis involucello brevioribus, ochreis biaristatis, culmo acutangulo. *Vahl. enum. pl. 2. p. 364.*

C. virginicus. *Jacq. in Herb. Lambert.*  
In Virginia. *Vahl.  $\frac{4}{4}$ . v. s. in. Herb. Lambert.*

22. C. spicis corymbosis oblongis basi ramosis nudis, spiculis numerosis divaricatis confertis linearibus subsexfloris, valvulis oblongis striatis, involucro octophyllo umbellam subæquante.

In ditches and near ponds: Virginia to Florida, rare.  
Aug. v. v.

*Folia* radicalia, linearia, 3-nervia, glabra, margine carinaque scabra, culmo breviora. *Culmus* laevigatus, triquetrus. *Involucrum* suboctophyllum: foliola interiora breviora, exteriora radiis umbellæ longiora. *Umbella* composita: radii octo vel decem. *Spicæ* oblongæ, basi ramosæ, magnæ. *Spiculæ* numerosæ (circiter 100), horizontales, castaneæ. An var. *C. speciosi*?

I have named this species in memory of Mr. Enslen, through whose generous communication I have been put in possession of a number of grasses and other plants growing in the southern States.

### 57. DULICHIUM. Pers. enc. 1. p. 65.

1. D. culmo triquetro folioso, spiculis patulis subsexfloris. *spathaceum.*  
*Pers. enc. l. c.*

*Pleuranthus floribundus.* Rich.

*Cyperus spathaceus.* Willd. sp. pl. 1. p. 289.

## TRIANDRIA MONOGYNIA. Dulichium.

- Schœnus angustifolius. *Vahl. enum. pl. 2. p. 225.*  
*Scirpus spathaceus. Mich. fl. amer. 1. p. 32.*  
*Icon. Pluk. alm. t. 301. f. 1.*  
 In swamps and bogs from Canada to Florida. ♀. July,  
 Aug. v. v.
- canadense.* 2. D. racemis simplicibus pedunculatis, spiculis paucis  
 erectis subdecemfloris. *Pers. l. c.*  
 In Canada and New York, in similar situations, rare.  
 ♀. Aug. v. v. Lower growth and shorter leaves  
 than No. 1.
58. SCIRPUS. *Gen. pl. 94.*
- \* *Culmo monostachyo.*
- acicularis.* 1. S. spica ovata, squamis duabus infimis majoribus, cul-  
 mis tetragonis setaceis, vaginis muticis. *Vahl. enum.*  
*pl. 2. p. 245. Willd. sp. pl. 1. p. 295.*  
*Icon. Pluk. alm. t. 40. f. 7. Host. gram. 3. t. 60.*  
 In shallow ponds and pools, under water, common. ♀.  
 July, Aug. v. v.
- pusillus.* 2. S. spica oblonga acuta (pauciflora) : squamis linearibus  
 acutis, culmis angulatis capillarisbus. *Vahl. enum. pl. 2.*  
*p. 246.*  
*S. capillaris. Mich. fl. amer. 1. p. 30.*  
 In springs and ditches, frequent. ♀. June, July. v. v.  
 The smallest of the genus.
- palustris.* 3. S. spica ovali, squamis lanceolatis acutis, culmis tereti-  
 bus, vaginis muticis lanceolatis acutis, radice repente.  
*Vahl. enum. pl. 2. p. 247. Willd. sp. pl. 1. p. 291.*  
*Icon. Fl. dan. t. 273. Host. gram. 3. t. 55.*  
 In overflowed fields and ditches, frequent. ♀. July.  
 v. v.
- tuberculatus.* 4. S. spica ovata acuta, squamis subrotundis, seminibus tu-  
 berculo ovato coronatis, culmis teretibus. *Vahl. enum.*  
*pl. 2. p. 248. Mich. fl. amer. 1. p. 30.*  
 In Lower Carolina. *Michaux.* ♀. +.
- filiformis.* 5. S. spica cylindrica oblonga obtusa, squamis subrotundis,  
 seminibus vertice nudis, culmis filiformibus teretibus.  
*Vahl. enum. pl. 2. p. 248. Lam. illustr. 1. p. 138.*  
 In wet fields, near ditches: New Jersey to Carolina.  
 ♀. July, Aug. v. v.
- ovatus.* 6. S. spica ovata, squamis oblongis, floribus subdiandris,  
 culmis subcompressis, vaginis muticis. *Vahl. enum.*  
*pl. 2. p. 249. Willd. sp. pl. 1. p. 294.*

S. a  
Icon  
t.  
Nea7. S. s  
va  
p.  
Icon  
In si8. S. s  
m  
pl  
Icon  
On  
v. i9. S. s  
m  
a  
In C10. S. s  
lar  
Icon  
S  
In v  
211. S. s  
su  
In v  
Cul  
c  
d12. S. s  
c  
p  
Icon  
In s

18. S. :

- S. compressus. Mœnch. meth. 394.  
Icon. Moris. hist. 3. s. 8. t. 10. f. 34. Host. gram. 3.  
t. 56.  
Near ponds and ditches : Pennsylvania. 4. July. v. v.
7. S. spica subglobosa, culmis sulcato-angulatis setaceis,  
vaginis muticis, spica subglobosa. Vahl. enum. pl. 2.  
p. 250. Willd. sp. pl. 1. p. 294.  
Icon. Rottb. gram. t. 15. f. 3.  
In small ponds, very common. 4. Aug. v. v.
8. S. spica ovato-oblonga, squamis ovato-subrotundis, cul- geniculatus.  
mis teretibus approximate interstinctis. Vahl. enum.  
pl. 2. p. 250. Willd. sp. pl. 1. p. 291.  
Icon. Sloan. hist. 1. t. 83. f. 3.  
On the sea-shore of Virginia and Carolina. 4. July.  
v. v.
9. S. spica cylindrica, squamis oblongo-subrotundis, cul- quadrangula-  
mis tetragonis. Vahl. enum. pl. 2. p. 252. Mich. fl. tus.  
amer. 1. p. 30.  
In Carolina. Michaux. 4. +.
- \*\* Culmo polystachyo : spicis lateralibus.
10. S. spicis ovatis acutis subsolitariis, culmis foliisque capil- minimus.  
laribus curvatis. Vahl. enum. pl. 2. p. 253.  
Icon. Pluk. alm. t. 300. f. 5. Rottb. gram. t. 15. f. 4.  
S. setaceus.  
In wet sandy fields and near ponds : Virginia to Carolina.  
4. July. v. v.
11. S. spicis brevi-ovatis conglomerato-sessilibus, squamis debilis.  
subrotundis membranaceis, culmis striatis.  
In wet meadows : Pennsylvania. 4. July, Aug. v. v.  
*Culmi* teretes, striati, pedales et ultra. *Spiculæ* 4—6.  
crassæ. *Squamis* latis, acutiusculis. *Setulæ* semine  
duplo longiores. Affinis S. setaceo.
12. S. aphyllus, spicis oblongis, squamis integerrimis mu- mucronatus.  
cronato-acuminatis, culmo triquetro. Vahl. enum.  
pl. 2. p. 256. Willd. sp. pl. 1. p. 303.  
Icon. Host. gram. 3. t. 66.  
In swamps and salt-marshes : Canada to Carolina. 4.  
Aug. v. v.
- \*\*\* Culmo polystachyo : spicis terminalibus.
13. S. aphyllus, spicis oblongis, squamis glabris mucronu- lacustris.

## TRIANDRIA MONOGYNIA. Scirpus.

- latis, stylis trifidis, umbella decomposita, culmo tereti.*  
*Vahl. enum. pl. 2. p. 267. Willd. sp. pl. 1. p. 296.*  
*Icon. Fl. dan. t. 1142. Host. gram. 3. t. 61.*  
*In stagnant waters: Canada to Carolina. 4. July. v. v.*
- validus.* 14. *S. aphyllus, spicis ovato-oblongis, squamis dorso villosis,*  
*stylis bifidis, umbella decomposita, involucro brevis-*  
*simo apice tereti.* *Vahl. enum. pl. 2. p. 268.*  
*In lakes and ponds: Canada to Carolina. 4. Aug. v. v.*  
*From four to ten feet high.*
- robustus.* 15. *S. spicis oblongis, corymbo composito, involucris poly-*  
*phyllis longissimis, involucellis ovatis longissime aristatis,*  
*squamis acutis aristatis, apice laceris, culmo tri-*  
*quetro folioso.*  
*S. maritimus β macrostachyos.* *Mich. fl. amer. 1. p. 32.*  
*In salt-marshes and on the banks of rivers, common.*  
*4. Aug. v. v. Certainly specifically distinct from*  
*S. maritimus, with which I carefully compared it.*
- americanus.* 16. *S. spiculis sessilibus conglomeratis oblongo-ovatis mu-*  
*crone brevioribus, culmo acute triquetro nudo.* *Pers.*  
*ench. 1. p. 68.*  
*S. triqueter.* *Mich. fl. amer. 1. p. 30.*  
*In salt marshes, frequent. 4. July, Aug. v. v.*
- sylvaticus.* 17. *S. spicis oblongis congestis, corymbo supradecomposito*  
*folioso, squamis oblongis obtusiusculis viridibus, cul-*  
*mo folioso trigono.*  
*Icon. Host. gram. 3. t. 68. Fl. dan. t. 307.*  
*In Canada. Michaux. 4. +.*
- exaltatus.* 18. *S. spicis brevi-ovatis congestis, corymbo composito,*  
*squamis rhomboideis carinatis mucronatis coloratis,*  
*culmo folioso obtuse trigono.*  
*S. brunneus.* *Muhl. in litt.*
- viviparus.* β *S. culmo altissimo subscandente, corymbis viviparis basi*  
*ramulorum floriferis.*  
*In shady woods: New York to Carolina. 4. July,*  
*Aug. v. v. The viviparous variety grows in deep shady*  
*swamps, and frequently acquires the height of ten*  
*feet and more.*
- nitens.* 19. *S. spicis ovatis pedicellatis, corymbis subcompositis ax-*  
*illaribus terminalibusque, culmo tereti.* *Vahl. enum.*  
*pl. 2. p. 272.*  
*In Virginia and Carolina. 4. July. v. v.*
- lineatus.* 20. *S. spicis oblongo-ovatis pedicellatis, corymbis axillaribus*  
*terminalibusque supradecompositis, pedunculis elon-*
- gatis,  
Mich.  
Trichoph  
In sandy  
v. v.  
chopho  
21. S. spicis  
termin  
Schoenus  
In shady  
22. S. spicis  
Vahl. e  
S. pusilli  
In Virgin  
23. S. spicis  
compos  
Willd.  
S. foliosu  
S. mucroi  
In sandy  
v. v.;
59. TR
1. *T. culmo*  
*culis co*  
*Scirpus E*  
*Eriophoru*  
*Icon. Pi*  
*In ditches*  
*4. Jul*
2. *T. culmo*  
*spica si*  
*Eriophoru*  
*In bogs o*  
*4. Aug*
- 60
1. *E. culmis*  
*scariosis*  
*Icon. H*  
*On high r*  
*June, J*

- gatis, culmo trigono. *Vahl. enum. pl. 2. p. 273.*  
*Mich. fl. amer. 1. p. 32.*
- Trichophorum lineatum. *Pers. enc. 1. p. 69.*  
 In sandy wet woods : New Jersey to Carolina. 4. July.  
 v. v. The seeds are naked, consequently it is no Trichophorum.
21. S. spicis capitulisque subglobosis glomeratis, corymbo *polyphyllus*.  
 terminali, culmo folioso. *Vahl. enum. pl. 2. p. 274.*  
*Scheenus cymosus. Willd. sp. pl. 1. p. 265?*  
 In shady woods : Virginia and Carolina. 4. July. v. v.
22. S. spicis ovatis, umbella biradiata, culmo capillari. *capillaris.*  
*Vahl. enum. pl. 2. p. 276. Linn. mant. 321.*  
 S. pusillus autumnalis, &c. *Gron. fl. virg. 11.*  
 In Virginia. 4. July. v. s. in Herb. Gronov.
23. S. spicis oblongis acutis, involucro diphyllo umbella composita breviore, culmo ancipi, foliis linearibus. *autumnalis.*  
*Willd. sp. pl. 1. p. 301. excl. syn. Rottboellii.*  
 S. foliosus pusillus autumnalis, &c. *Gron. fl. virg. 10.*  
 S. mucronulatus. *Mich. fl. amer. 1. p. 32.*  
 In sandy wet fields : New Jersey to Carolina. 4. Aug.  
 v. v.; v. s. in Herb. Gronov.

59. TRICHOPHORUM. *Pers. enc. 1. p. 69.*

1. T. culmo triquetro folioso, panicula decomposita, spiculis confertis ovatis. *Pers. l. c.* *cyperinum.*  
 Scirpus Eriophorum. *Mich. fl. amer. 1. p. 33.*  
 Eriophorum cyperinum. *Willd. sp. pl. 1. p. 313.*  
 Icon. *Pluk. mant. t. 419. f. 3.*  
 In ditches and near ponds, from Canada to Carolina.  
 4. July, Aug. v. v. Very tall.
2. T. culmo triquetro nudo, foliis canaliculato-setaceis, spica simplici. *alpinum.*  
 Eriophorum hudsonianum. *Mich. fl. amer. 1. p. 34.*  
 In bogs on the mountains of Pennsylvania, Vermont, &c.  
 4. Aug. v. v.

60. ERIOPHORUM. *Gen. pl. 95.*

1. E. culmis superne trigonis, spica oblonga unica, squamis scariosis. *Host. gram. 1. p. 33.*  
 Icon. *Host. l. c. t. 39.* *cespitosum.*  
 On high mountain meadows : Canada to Virginia. 4.  
 June, July. v. v.

- polystachium.* 2. E. culmis foliosis teretibus, foliis planis, spicis pedunculatis, seminibus ovatis. *Roth. neue Beytr.* 1. p. 94. *Willd. sp. pl.* 1. p. 312.  
Icon. *Host. gram.* 1. t. 37. *Engl. bot.* 563.  
In bog-meadows: common. ♀. May. v. v.
- angustifoli-  
um.* 3. E. culmis foliosis teretibus, foliis canaliculato-triquetris, spicis pedunculatis, seminibus ovatis. *Roth. neue Beytr.* 1. p. 94. *Willd. sp. pl.* 1. p. 313.  
Icon. *Engl. bot.* 564.  
In mountain meadows, frequent. ♀. May. v. v.
- virginicum.* 4. E. culmo tereti folioso, foliis planis, spicis congestis subsessilibus involucro brevioribus. *Vahl. enum. pl.* 2. p. 390. *Willd. sp. pl.* 1. p. 313.  
Icon. *Pluk. alm.* t. 299. f. 4.  
Common in bog-meadows, from Canada to Georgia. ♀. April, May. v. v.

61. VAGINARIA. *Pers. enc.* 1. p. 70.

- Richardi.* 1. V. culmo aphylo vaginato, spica ovata subsolitaria, squamis lanceolato-spathulatis. *Pers. l. c.*  
*Fuirena scirpoidea.* *Mich. fl. amer.* 1. p. 38.  
Icon. *Mich. l. c.* t. 7.  
In overflowed fields: Georgia to Florida. ♀. v. s.

62. FUIRENA. *Gen. pl.* 90.

- squarrosa.* 1. T. umbellis simplicibus, spiculis ovatis, culmo glabro, foliis ciliatis, vaginis pilosis. *Vahl. enum. pl.* 2. p. 385. *Mich. fl. amer.* 1. p. 37.  
In bog-meadows: New Jersey, Carolina, &c. ♀. July. v. v. A very handsome little grass.

63. MARISCUS. *Vahl. enum. pl.* 2. p. 372.

- ovularis.* 1. M. spicis ovali-subrotundis, involucris polyphyllis. *Vahl. enum. pl.* 2. p. 374.  
*Scheuenus umbellatus.* *Jacq. collect.* 1. p. 113.  
*Kyllingia ovularis.* *Mich. fl. amer.* 1. p. 29.  
Icon. *Pluk. alm.* t. 91. f. 4. *Jacq. ic. rar.* 1. t. 10?  
In sandy wet places: Georgia and Carolina. ♀. v. s.
- retrofractus.* 2. M. spicis retrorsum subimbricatis, spiculis subulatis retrofractis, involucro triphyllio. *Vahl. enum. pl.* 2. p. 375.

TRI.

*Scirpus retr.*  
Icon. *Plu.*  
In rocky sh  
ginia, Ca  
leaves ve

3. M. spicis cy  
lyphyllis.  
*Scirpus cyp*  
*Kyllingia u*  
Icon. *Rot.*  
In shady w  
The Amer  
in every

64.

1. *Miegia.* 1  
*Arundinaria*  
*Arundo gig*  
In extensiv  
Cane bra  
Walter &  
♀. Marc

65. LIMI

1. L. foliis dis  
paucis re  
peris. I  
*Trachynotia*  
*Dactylis pa*  
In dry salt-  
2. L. subglau  
secundis,  
*Pers. enc*  
*Trachynotia*  
On rocky se

3. L. foliis lat  
linearibu  
*Trachynotia*  
*Dactylis cy*  
Common in  
Aug. v.

Scirpus retrofractus. Willd. sp. pl. 1. p. 304.

Icon. Pluk. alm. t. 415. f. 4.

In rocky shady situations, on the banks of rivers: Virginia, Carolina, &c. ♀. July. v. v. Stems and leaves very rough.

3. M. spicis cylindricis retrorsum imbricatis, involucris polyphyllis. Vahl. enum. pl. 2. p. 376. umbellatus.

Scirpus cyperoides. Linn. mant. 181.

Kyllingia umbellata. Linn. suppl. 105.

Icon. Rottb. gram. t. 4. f. 2.

In shady woods: Virginia and Carolina. ♀. Aug. v. ♀.

The American specimen is without partial involucres, in every other respect it agrees with the description.

#### 64. MIEGIA. Pers. enc. 1. p. 101.

1. Miegia. Pers. enc. 1. p. 102. macrosperma.

Arundinaria macrosperma. Mich. fl. amer. 1. p. 74.

Arundo gigantea. Walt. fl. car. 81.

In extensive tracts on the Mississippi and Floridas, called

Cane brakes, almost impenetrable. Arundo tecta of Walter seems to be only a variety of this species.

♀. March, April. v. v.

#### 65. LIMNETIS. Richard in Pers. enc. 1. p. 72.

1. L. foliis distichis breviusculis setaceo-convolutis, spicis paucis remotis patulis, calyce acuminato: carinis asperis. Pers. enc. 1. p. 72. juncea.

Trachynotia juncea. Mich. fl. amer. 1. p. 64.

Dactylis patens. Donn. hort. cant. 25.

In dry salt-meadows, common. ♀. July. v. v.

2. L. subglaucia, foliis longissimis, spicis plurimis alternis cynosuroides, secundis, calyce aristato acuminato: carinis aculeatis.

Pers. enc. 1. p. 72.

Trachynotia cynosuroides. Mich. fl. amer. 1. p. 64.

On rocky sea-shores, frequent. ♀. July. v. v.

3. L. foliis latis planis, spicis pluribus undique versis longis polystachya. linearibus, carinis aculeatis. Pers. enc. 1. p. 72.

Trachynotia polystachya. Mich. fl. amer. 1. p. 64.

Dactylis cynosuroides. Willd. sp. pl. 1. p. 407.

Common in salt-marshes, from Canada to Florida. ♀.

Aug. v. v.

66. CENCHRUS. *Gen. pl. 1574.**tribuloides.*

1. *C. spica* spiculis alternis, glumis feminineis globosis mucato-spinosis hirsutis. *Willd. sp. pl. 1. p. 317.*

*C. carolinianus. Walt. fl. car. 79.*Icon. *Sloan. hist. 1. t. 65. f. 1.*

Near the sea-shore, from New Jersey to Florida. ○

June, July. *v. v.; v.s. in Herb. Walter.*

7. July.

69. MI

1. *M. culmo* arctata, c  
Dilepycum i

On dry grass

2. *M. culmo* f  
nicula lax

ench. 1. p

- Dilepycum

In dry shad

*v. v.*

70. TRIC

1. *T. culmis* asperiusc p. 42.

Cornucopiae

Agrostis sca

Agrostis lax

Icon. *Mic*

Common in

2. *T. culmis* glumis li amer. 1.

Cornucopia

Agrostis an

Agrostis co

Icon. *Fra*

On moist h

Carolina

species a

Monogr.

3. *T. culmis* valvis me

Cornucopia

Agrostis dis

## II. DIGYNIA. -

68. ZIZANIA. *Gen. pl. 1433.* Indian Rice.*aquatica.*

1. *Z. panicula pyramidata*, inferne divaricata mascula, superne spicata feminine, fulcris florum clavatis aristis longis, semine elongato.

*Z. aquatica. Lambert. in linn. trans. 7. p. 264.**Z. palustris. Linn. mant. 295. Gron. virg. 148.**Z. clavulosa. Mich. fl. amer. 1. p. 75.*Icon. *Lambert. l. c. t. 13.* Tuscarora or Wild Rice.

In all the waters from Canada to Florida, common. 1.

July, Aug. *v. v.**miliacea.*

2. *Z. panicula effusa*, glumis brevi-aristatis: masculis et feminineis mixtis, semine ovato laevi. *Mich. fl. amer. 1. p. 74.*

*Z. aquatica. Willd. sp. pl. 4. p. 394?*Icon. *Sloan. hist. 1. t. 67?*On the tide water meadows and ditches: Pennsylvania and Carolina. 2. Aug. *v. v.*

*fluitans.*

3. *Z. pusilla*, culmis gracilibus ramosis, foliis linearibus planis, spicis solitariis axillaribus setaceis subquadri-floris, glumis muticis. *Mich. fl. amer.* 1. p. 75.  
On the banks of Lake Champlain and St. Laurence.  
*v. v.*

69. MUHLENBERGIA. *Gen. pl.* 103.

1. *M. culmo debili ramoso foliisque glabris*, panicula co-arctata, calyce univalvi. *Pers. enc.* 1. p. 73.  
*Dilepycum minutiflorum.* *Mich. fl. amer.* 1. p. 40.  
On dry grassy hills, frequent. *v. v.*  
*erecta.*
2. *M. culmo firmo simplici foliisque pubescentibus*, panicula laxa, calyce bivalvi, arista longissima. *Pers. enc.* 1. p. 73.  
*Dilepycum aristosum.* *Mich. fl. amer.* 1. p. 40.  
In dry shady woods, from Canada to Carolina. *v. v.*

70. TRICHODIUM. *Mich. fl. amer.* 1. p. 41.

1. *T. culmis erectis, foliis angustis breviusculis, vaginis asperiusculis, glumis lanceolatis.* *Mich. fl. amer.* 1. p. 42.  
*Cornucopiæ hyemalis.* *Walt. fl. car.* 73.  
*Agrostis scabra.* *Willd. sp. pl.* 1. p. 370.  
*Agrostis laxa.* *Schreb. gram.*  
*Icon.* *Mich. fl. amer.* 1. t. 8.  
Common in all grassy situations. *v. v.*
2. *T. culmis decumbentibus, foliis patulis, vaginae lœvi, glumis linear-lanceolatis sensim acutis.* *Mich. fl. amer.* 1. p. 42.  
*Cornucopiæ perennans.* *Walt. fl. car.* 74.  
*Agrostis anomala.* *Willd. sp. pl.* 1. p. 370.  
*Agrostis cornucopiæ.* *Fraser. monogr. cum icon.*  
*Icon.* *Fraser. monogr.*  
On moist hills and short grass meadows, from Canada to Carolina. *v. v.* This and the first species are highly valuable grasses. *Vide* Fraser's Monogr. and Walter's Flora.
3. *T. culmis erectis, foliis angustis scabris, vaginis glabris valvis membranaceis ovato-lanceolatis acuminatis.* *elatum.*  
*Cornucopiæ altissima.* *Walt. fl. car.* 74.  
*Agrostis dispit.* *Mich. fl. amer.* 1. p. 52.

## TRIANDRIA DIGYNIA. Trichodium.

In sandy deep swamps : New Jersey, Carolina, &c. 1.  
July, Aug. v. v. Near three feet high ; panicle  
close, coloured.

## 71. LEERSIA. Gen. pl. 105.

*oryzoides.*



*lenticularis.*

1. L. panicula effusa, spiculis triandris patulis, glumis oblongo-ovalibus carina ciliatis. Willd. sp. pl. 1. p. 325.
  2. L. virginica. Willd. l. c. Icon. Host. gram. 1. t. 35.
- In wet woods and ditches, frequent, 1. July. v. v.

Fly-catch-grass.  
In wet gravelly woods in Illinois and Virginia. 1. July,  
v. v. This singular and elegant grass I found on the  
islands of Roanoak river in North Carolina, and ob-  
served it catching flies in the same manner as *Dionaea*  
*muscipula* : the valves of the corolla are nearly of the  
same structure as the leaves of that plant. I communi-  
cated specimens with this particular circumstance to  
Dr. B. S. Barton of Philadelphia, who has made men-  
tion of it in a paper on the irritability of plants.

## 72. MILIUM. Gen. pl. 110. Millet-grass.

*amphicarpon.* 1. M. paniculae ramis strictis simpliciusculis, floribus mas-  
culis alternis pedunculatis ; floribus femineis in scapis  
unifloris radicalibus vaginatis, demum subterraneis.

In light sandy fields of New Jersey near Egg-harbour.  
July, Aug. v. v.

*Culmi* plures teretes, vaginati, sesquipedales et ultra.  
*Folia* lato-linearia, striata, geniculata breviora, undique  
pilis longis albidis rigidis tuberculo insidentibus tecta.  
*Vaginæ* teretes, striati : suprema aphylla, more folio-  
rum pilosæ. *Ligula* barbata.

*Panicula* in summitate culmi, pauciflora, stricta : ramis  
simpliciusculis ; pedicellis florum clavatis. *Glumis*  
oblongis, acutis, nervosis, æqualibus, glabriusculis.  
*Floribus* omnibus masculis.

*Flores* feminei in scapis radicalibus unifloris basi vagina-  
tis. *Flores* ante anthesin erecti, dein reflexi, ad ma-  
turitatem semenis magni subrotundi subterranei.

This singular grass deserves particular examination in  
the living plants, as I had no opportunity to describe



*Milium amphicarpon.*

it on the  
of its ca

73. AG

1. A. petalo  
cula pat  
Icon. *H*  
In rye-field  
by seed.
2. A. panicu  
tatis, *pl.*  
*p. 364.*  
On the t  
July.
3. A. panicu  
calyce i  
losis.  
On high  
*v. v.*  
*Culmus se*  
*villosa,*  
*brevis.*  
*floris.*  
*tis.* *C*  
*duplo l*
4. A. panicu  
arista e  
*pl. 1. f*  
In good  
July.
5. A. calyci  
culmis  
Icon. *F*  
In wet  
*2. Ju*
6. A.. panic  
lycibu  
tuso.  
A. hispic  
A. capill

it on the spot where I first found it; not being aware of its curiosity, until I examined the dried specimens.

73. AGROSTIS. Gen. pl. 111. Bent-grass.

\* *Aristatae.*

1. A. petalo exteriore arista recta stricta longissima, panicula patula. *Willd. sp. pl. 1. p. 361.*

Icon. *Host. gram. 3. t. 47. Engl. bot. 951.*

In rye-fields in Pennsylvania and Virginia; brought there by seeds of grain from Europe. ☉ June, July. v.v.

2. A. paniculae simplicissimae ramis adpressis, corollis aristatis, arista flore longioribus. *Willd. sp. pl. 1. p. 364.*

On the banks of rivers: Virginia and Carolina. 4. July. v.v.

3. A. paniculae ramis unifloris filiformibus, corollis aristatis calyce inaequali duplo longioribus hirsutis vaginis villosis.

On high mountains in Virginia and Carolina. 24. July. v.v.

*Culmus* sesquipedalis, glaber. *Folia* longissime-linearia, villosa, nervosa. *Vaginæ* striatæ, villosæ. *Ligula* brevis. *Panicula* simplicissima, ramis elongatis unifloris. *Cal.* valvis inaequalibus, villosis, brevi-aristatis. *Cor.* oblonga, villosa, calycis valvula interiore duplo longior, brevi-aristata.

4. A. panicula elongata stricta, corollis calyce minoribus: arista e basi petali tortili flore longiore. *Willd. sp. pl. 1. p. 366.*

In good meadow grounds, very frequent. 4. June, July. v.v.

5. A. calycibus elongatis, petalorum arista dorsali recurva, culmis prostratis subramosis. *Willd. sp. pl. 1. p. 367.*

Icon. *Fl. dan. 161.*

In wet meadows, frequent. Brought from Europe. 4. July. v.v.

*Spica venti.*

*tenuiflora.*

*pauciflora.*

*stricta.*

*canina.*

*vulgaris.*

\*\* *Muticæ.*

6. A. panicula patente, ramulis divaricatis capillaribus, calycibus æqualibus, petalo interiore duplo breviore retuso. *Smith. fl. brit. 1. p 79.*

A. hispida. *Willd. sp. pl. 1. p. 370.*

A. capillaris. *Retz. scand. no. 95.*

## TRIANDRIA DIGYNIA. Agrostis.

- Icon. *Fl. dan.* 161.  
Common in all grassy fields. 4. July. v. v.
- aspera.* 7. A. paniculis coarctatis lateralibus terminalibusque, vaginis barbatis, foliis recurvatis asperis, glumæ valvis interioribus triplo majoribus. *Mich. fl. amer.* 1. p. 52.  
A. barbata. *Pers. enc.* 1. p. 75.  
A. littoralis. *Lam. illust.* 1. p. 161.  
In the western districts of the United States, in rocky situations. *Michaux.* On the sea-shore of Florida. *Lam.* 4. July. v. s. in *Herb. Enslin.*
- alba.* 8. A. panicula laxa, calycibus muticis æqualibus, culmo repente. *Willd. sp. pl.* 1. p. 371.  
On the banks of rivulets and springs, frequent. 4. July. v. v.
- pungens.* 9. A. panicula contracta spicæformi, foliis involutis rigidis pungentibus superioribus oblique oppositis, culmo ramoso. *Schreb. gram.* 2. p. 46.  
A. virginica. *Willd. sp. pl.* 1. p. 373.  
Icon. *Schreb. L. c. t.* 27. f. 3.  
On dry gravelly hills in Virginia. 4. July, Aug. v. v.
- Cinna.* 10. A. panicula magna debili, foliis lato-linearibus glabris, valvula altera infra apicem subaristata, caule glabro.  
*Cinna arundinacea.* *Willd. sp. pl.* 1. p. 31.  
On rivers and islands, under the tide waters, common; from Canada to Carolina. 4. Aug. v. v.
- mexicana.* 11. A. panicula oblonga congesta, calycibus corollisque acuminatis subæqualibus muticis. *Willd. sp. pl.* 1. p. 374.  
In wet meadows: Virginia, &c. 4. June. v. v.
- indica.* 12. A. panicula contracta mutica, racemis lateralibus erectis alternis, glumæ valvis exterioribus longissimis, foliis longissimis, sensim setaceo-convolutis. *Mich. fl. amer.* 1. p. 52. *Willd. sp. pl.* 1. p. 375.  
In Carolina, Florida, &c. *Michaux.* ⓠ. +. v. s. in *Herb. Lambert.*
- junccea.* 13. A. panicula oblongo-pyramidata, floribus muticis: valvula extima duplo minore, foliis stricte erectis convoluto-setaceis rigidulis. *Mich. fl. amer.* 1. p. 52.  
In barren sandy places: New Jersey to Florida. 4. July, Aug. v. v.
- lateriflora.* 14. A. paniculis lateralibus terminalibusque coarctatis, floribus muticis, valvulis acutissimis: interioribus ma-

goribus  
amer.  
In rich s  
Florid  
July.  
of the  
15. A. racei  
exteri  
bus, §  
amer.  
On the

74.  
1. A. spica  
longi  
Icon. E  
Commo  
seeds

1. P. panic  
natis.  
Arundo  
Icon.  
On rive  
Aug.

2. P. panic  
Mich  
In wood

76. P  
1. P. spic  
Will  
Icon.  
In field  
v. v.

1. A. cult  
sp. 1  
Icon.  
In field  
4. 1

VOL. I.

isque, va-  
mæ valvis  
. 1. p. 52.  
  
in rocky  
f Florida.  
  
culmo re-  
4. July.  
  
is rigidis  
ulmo ra-  
  
g. v. v.  
glabris,  
labro.  
  
mon ;  
  
que acu-  
pl. 1.  
  
. v.  
  
s erectis  
is, foliis  
tich. fl.  
  
v. s.  
  
is : val-  
is con-  
52.  
a. 4.  
  
is, flo-  
us ma-

goribus basi barbatis, foliis planis brevibus. *Mich. fl. amer.* 1. p. 53.

In rich soil on the edges of woods, from New York to Florida, especially in the western countries. ♀. June, July. v. v. Apparently a grass worth the attention of the cultivator.

15. A. racemo oblongo, e spicis densifloris interrupto, glumæ exterioris valvis in mucronem aristiformem desinentibus, gluma interiore mutica longioribus. *Mich. fl. amer.* 1. p. 53.

On the overflowed banks of the river Mississippi. +.

#### 74. ANTHOXANTHUM. *Gen. pl. 58.*

1. A. spica oblonga ovata, flosculis subpedunculatis arista longioribus. *Willd. sp. pl. 1. p. 156.*

Icon. *Host. gram. 1. t. 5. Fl. dan. 666. Engl. bot. 647.*

Common in meadows; probably brought with grass seeds from Europe. ♀. May—July. v. v.

*racemosa.*

*odoratum.*

#### 75. PHALARIS. *Gen. pl. 106.*

1. P. panicula oblonga ventricosa coarctata, glumis acumi- *arundinacea.*  
natis. *Pers. enc. 1. p. 78. Willd. sp. pl. 1. p. 326.*

*Arundo colorata. Ait. kew. ed. 2. 1. p. 174.*

Icon. *Host. gram. 2. t. 33. Engl. bot. 402.*

On river banks, frequent: Canada to Virginia. ♀. July, Aug. v. v.

2. P. panicula multiflora stricta, floribus ovatis villosis. *Mich. fl. amer. 1. p. 43.*

In woods of Carolina. *Michaux.* +.

*villosa.*

#### 76. PHLEUM. *Gen. pl. 128. Timothy-grass.*

1. P. spica cylindrica longissima ciliata, culmo erecto. *Willd. sp. pl. 1. p. 354.*

Icon. *Host. gram. 3. t. 9.*

In fields and meadows, common. ♀. ♂. June—Sept. v. v.

*pratense.*

#### 77. ALOPECURUS. *Gen. pl. 102.*

1. A. culmo erecto, glumis villosis, corollis muticis. *Willd. sp. pl. 1. p. 357.*

Icon. *Host. gram. 2. t. 31. Engl. bot. 759.*

In fields and gardens, frequent, introduced from Europe.

♀. May—Aug. v. v.

*pratensis.*

- subaristatus.* 2. A. culmo erecto, glumis ovalibus obtusis, arista vix exerta. *Mich. fl. amer.* 1. p. 43.  
 A. carolinianus, culmo radicante, panicula subspicata, glumis laevibus, corollis aristatis. *Walt. fl. car.* 74?  
 In swamps of Canada. *Michaux.* 4. v. s. in *Herb. Walter.*

78. PANICUM. *Gen. pl.* 107.\* *Spicata.*

- glaucum.* 1. P. spica tereti subflavida, involucellis bifloris fasciculato-setosis, seminibus transversim rugosis. *Lam. encycl.* 4. p. 736. *Willd. sp. pl.* 1. p. 335.  
 Icon. *Host. gram.* 2. t. 16.  
 Common in fields and on road sides; probably from Europe. ☽. June—Aug. v. v.
- viride.* 2. P. spica tereti subcomposita indivisa, involucellis setosis mitibus, seminibus nervosis. *Lam. encycl.* 4. p. 737. *Willd. sp. pl.* 1. p. 335.  
 Icon. *Host. gram.* 2. t. 14. *Engl. bot.* 875.  
 In sandy fields: Pennsylvania and Virginia. ☽. July. v. v.
- verticillatum.* 3. P. spica verticillata, racemulis quaternis, involucellis unifloris bisetis, culmis diffusis. *Willd. sp. pl.* 1. p. 334.  
*P. alopecuroideum.* *Walt. fl. car.* 72.  
 Icon. *Host. gram.* 2. t. 13.  
 In sandy woods: New Jersey to Carolina. ☽. June, July. v. v.; v. s. in *Herb. Walter.*
- hirtellum.* 4. P. spica composita: spiculis adpressis alternis, calycibus geminis: valvulis omnibus aristatis: extima longissima. *Willd. sp. pl.* 1. p. 340.  
 Icon. *Host. gram.* 3. t. 52.  
 In shady places: Carolina and Florida. *Michaux.* +.
- Crus galli.* 5. P. spicis alternis conjugatisque crassis squarrosis, glumis hispidis aristatis, rachi angulato vaginis glabris. *Lam. encycl.* 4. p. 744. *Willd. sp. pl.* 1. p. 337.  
 Icon. *Host. gram.* 2. t. 19. *Engl. bot.* 876. *Fox-tail.*  
 $\alpha$ . P. glumis omnibus aristatis: aristis praelongis.  
 $\beta$ . P. glumis omnibus muticis mucronatis.  
 $\gamma$ . P. glumis mucronatis coloratis.  
 In cultivated grounds, frequent. ☽. July. v. v.
- aristatum.* 6. P. spicis alternis erectis solitariis simplicibus, glumis ovatis, muricato-hispidis aristatis: arista altera lon-

gissi  
glab  
P. mur  
P. hirte  
Near t  
Sept  
7. P. spic  
dens  
muti  
phro  
In sedg

8. P. race  
pilis  
Icon.  
In rice

9. P. pani  
pulci  
In Car

10. P. pani  
tis n  
P. virg  
P. colo  
On rive  
Aug

11. P. pan  
tibus  
aribi  
P. nitic  
p. 49

$\beta$ . P. omn  
Comm  
P. l  
spec

12. P. pani  
foliis  
barb  
P. dich  
In dry

13. P. pau

- gissima, rachi trigona, vaginis hispidissimis, foliis  
glabris.  
*P. muricatum.* *Mich. fl. amer.* 1. p. 47.  
*P. hirtellum.* *Walt. fl. car.* 72.  
 Near the salt-water: Canada and New York. ☽ July—  
 Sept. v. v.; v. s. in *Herb. Lambert.* nec non *Walter.*  
 7. *P. spicis* solitarii alternis simplicissimis, rachi glumisque  
dense pubescentibus, glumis solitariis alternis ovatis  
muticis bifloris: flore altero masculo, altero herma-  
phrodito. *Mich. fl. amer.* 1. p. 47.  
 In sedgy meadows: Florida. *Michaux.* +.

molle.

- \*\* *Paniculata.*
8. *P. racemis* linearibus virgatis, glumis clavatis coloratis, *fusco-rubens.*  
 pilis sub panicula divisuris. *Lam. encycl.* 4. p. 737.  
 Icon. *Sloan. hist.* 1. t. 72 f. 2?  
 In rice fields in Georgia. 4. Aug. v. s.
9. *P. panicula* oblonga, glumis majusculis glabris viridibus      *striatum.*  
 pulchre striatis. *Lam. encycl.* 4. p. 748.  
 In Carolina. *Fraser.* +.
10. *P. panicula* virgata ramosissima, glumis ovatis acumina-  
 tis muticis bifloris, foliis longissimis planis.      *virgatum.*  
*P. virgatum.* *Mich. fl. amer.* 1. p. 352.  
*P. coloratum.* *Walt. fl. car.* 73.  
 On river sides: New England to Carolina. 4. July,  
 Aug. v. v. Very tall.
11. *P. panicula* capillacea ramosa, glumis striatis pubescenti-  
 bus, seminibus nitidis, foliis remotis lanceolato-line-  
 aribus collo barbatis, caule glabro.      *nitidum.*  
*P. nitidum.* *Lam. encycl.* 4. p. 748. *Mich. fl. amer.* 1.  
 p. 49.
- β. *P. omnibus partibus multo majus.*      *majus.*  
 Common in woods and fields. ☽ June—Aug. v. v.  
*P. barbulatum* *Mich.* I consider a variety of this  
 species.
12. *P. paniculata* simplicissimis paucifloris, glumis obovatis, *dichotomum.*  
 foliis linear-lanceolatis divaricatis glabris basi et collo  
 barbatis, caule dichotomo.  
*P. dichotomum.* *Willd. sp. pl.* 1. p. 346.  
 In dry woods, frequent. 4. July, Aug. v. v.
13. *P. panicula* capillari ramosissima laxa, floribus minutis      *capillare.*

## TRIANDRIA DIGYNIA. Panicum.

omnibus pedicellatis solitariis oblongo-ovatis acumina-tis muticis, foliis vaginisque hirsutissimis. *Mich. fl. amer.* 1. p. 47. *Willd. sp. pl.* 1. p. 349.

In sandy fields: New Jersey to Florida. ♂. June—  
Aug. v. v. This grass is remarkably variable in size.  
I have seen specimens from three inches to five feet  
high.

*diffusum.*

14. *P. panicula simpliciuscula capillari patente, flosculis remotiusculis acutis, culmo simplici, foliis longis linearibus collo vaginisque villosis.* *Swartz. prodr.* 23.

*P. cæspititum.* *Lam. encycl.* 4. p. 743.

In dry sandy woods: Virginia, Carolina, &c. ♂. July.  
v. v.

*scoparium.*

15. *P. panicula erecta composita setaceo-ramosissima, glumis obovatis pubescentibus, foliis lanceolatis villosis.* *Mich. fl. amer.* 1. p. 49.

In dry swamps of Carolina. *Michaux.* ♂. +.

*nodiflorum.*

16. *P. paniculis minimis lateralibus terminalibusque, glumis ovatis pubescentibus, foliis angustis breviusculis collo barbatis.* *Lam. encycl.* 4. p. 744.

*P. ramulosum.* *Mich. fl. amer.* 1. p. 30.

In dry fields: Pennsylvania to Carolina. ♂. July. v. v.

*proliferum.*

17. *P. glaberrimum, paniculis oblongis erectis lateralibus terminalibusque, glumis oblongis acutis striatis, caule ramoso-dichotomo.* *Lam. encycl.* 4. p. 747.

*P. dichotomiflorum.* *Mich. fl. amer.* 1. p. 48.

In rich soil, in woods, on edges of ditches: Pennsylvania to Carolina. ♀. July, Aug. v. v.

*pubescens.*

18. *P. erectum ramosissimum pubescens, paniculis parvis paucifloris laxis sessilibus, glumis globoso-ovatis sub-pedicellatis pubescentibus.* *Lam. encycl.* 4. p. 748. *Mich. fl. amer.* 4. p. 49.

In shady rich woods: Virginia to Carolina. ♂. July. v. v.

*laxiflorum.*

19. *P. panicula patente laxa pilosa, glumis rariss obtusis pubescentibus.* *Lam. encycl.* 4. p. 748.

In North America. *Lamarck.* ♂. +.

*latifolium.*

20. *P. panicula patenti mediocri, glumis ovatis acutiusculis subpubescentibus, foliis ovato-lanceolatis amplexicaulis glabris, vaginis pilosis.*

*P. latifolium.* *Willd. sp. pl.* 1. p. 350.

*clandestinum.*

β. *P. paniculis inclusis paucifloris.* *Willd. sp. pl.* 1. p. 351.

In shady woods: New England to Georgia. ♀. July.

v. v.

21. *P. erecta mulcivis pinguis*  
In shad.
22. *P. debilis libus americanus*  
P. nudata In Car.
23. *P. glaberrimus*  
In high.
24. *P. glaber gemmatus*  
mul-tis to,  
In di-  
♀. five
25. *P. paniculatis bus subsp.*  
On th.
1. *D. sp. folia amara*  
Panicum  
Paspalum  
Synthemis  
Iconocarpus  
Comandra
2. *D. sp.*

21. *P. erectum*, paniculæ ramis simplicibus interrupte racemulosis, flore accessorio neutro : hermaphrodito valvis primariis semibreviore obtusissimo, foliis longis, vaginis compressis pilosis. *Mich. fl. amer.* 1. p. 48.  
In shady wet woods : Carolina. *Michaux.* +.
22. *P. debile*, glaberrimum, panicula gracili longa : ramulis raris adpressis, glumæ membranaceæ valvis subæquilibus lanceolatis, foliis angustis longis. *Mich. fl. amer.* 1. p. 50.  
*P. nudum*. *Walt. fl. car.* 73 ?  
In Carolina and Georgia. ¼. July, Aug. v. v.
23. *P. glabrum*, erectum ; panicula pyramidata, ramis paucis dissitis, plerisque geminis simpliciusculis divaricatis reflexisque, floribus muticis binis : altero pedicellato neutro, foliis angusto-longis. *Mich. fl. amer.* 1. p. 50.  
In high mountains of Carolina. *Michaux.* +.
24. *P. glabrum*, elatum ; paniculis pyramidatis lateralibus geminis elongato-pedunculatis terminalibusque : ramulis alternis divaricatis, glumis alternis oblongis acutis pedicellatis coloratis, foliis longis : collo subbarbato, caule compresso.  
In ditches and near ponds : New Jersey to Virginia.  
¾. July. v. v. A very handsome grass, sometimes five feet high; panicle a dark purple mixed with green.
25. *P. panicula solitaria* folio terminali breviora : ramis simplicibus flexuosis, glumis alternis pedunculatis obovatis turgidis : valvulis multistriatis acutis, foliis linearibus strictis sensim acutissimis, supra striatis scabris subtus pilosis, vaginis longissime pilosis.  
On the banks of Delaware : Pennsylvania. ⊙. July. v. v.

79. DIGITARIA. *Hall.* Crop-grass.

1. *D. spicis pluribus*, floribus imbricatis geminis, vaginis foliisque papilloso-pilosis, culmo repente. *Mich. fl. amer.* 1. p. 45.  
*Panicum sanguinale*. *Willd. sp. pl.* 1. p. 342.  
*Paspalum sanguinale*. *Lam. encycl.* 5. p. 33.  
*Syntherisma præcox*. *Walt. fl. car.* 76.  
*Icon. Host. grum.* 2. t. 17. *Curt. lond.* 4. t. 7.  
Common in cultivated ground. ⊙. June—Aug. v. v.
2. *D. spicis pluribus* (2—5) : erectis filiformibus, floribus

*Sanguinale.**pilosa.*

TRIANDRIA DIGYNIA. *Panicum.*

- remotis hirsutis geminis : altero vix pedicellato, foliis brevibus, vaginis pilosis, culmo erecto glabro superne nudo.* *Mich. fl. amer. 1. p. 45.*
- Syntherisma villosa.* *Walt. fl. car. 77.*  
*Panicum filiforme.* *Willd. sp. pl. 1. p. 343.*  
*Paspalum debile.* *Lam. encycl. 5. p. 34.*  
 In sandy dry fields and on hills; New Jersey to Georgia.  
 ☉ May—Aug. v. v.
- paspaloides.* 3. *D. spicis geminis basi subvillosis, rachi latiuscula glabra, floribus solitariis bifarium imbricatis glabris, foliis patentibus, vaginis glabris collo villosis, culmo repente.* *Mich. fl. amer. 1. p. 46.*  
*Paspalum supinum.* *Bosc. in lam. encycl. 5. p. 29.*  
 On hills of red clay and iron ore: Virginia. ☉ July.  
 v. v.
- serotina.* 4. *D. spicis pluribus setaceis, floribus binis vel ternis minutis pedicellatis, foliis vaginisque villosis, culmo decumbente.* *Mich. fl. amer. 1. p. 46.*  
*Syntherisma serotina.* *Walt. fl. car. 76.*  
 In low wet places: Carolina. ☉ June—Aug. v. v.
80. CYNODON. *Richard in Pers. enc. 1. p. 85.*
- Dactylon.* 1. *C. spicis digitatis patentibus, basi interiori villosis, floribus solitariis, sarmentis repentibus.* *Pers. l. c.*  
*Panicum Dactylon.* *Willd. sp. pl. 1. p. 342.*  
*Digitaria Dactylon.* *Allion. fl. ped. 11. 238.*  
*Paspalum Dactylon.* *Lam. encycl. 5. p. 32.*  
*Paspalum præcox.* *Walt. fl. car. 75.*  
 On road sides and in cultivated ground, frequent. ♀.  
 v. v.; v. s. in *Herb. Walter.*
81. PASPALUM. *Gen. pl. 81.*
- setaceum.* 1. *P. spica unica gracili, glumis geminis suborbiculato-obovatis glabris, foliis villosis, culmo setaceo erecto.* *Mich. fl. amer. 1. p. 43.*  
*P. dissectum.* *Walt. fl. car. 75.*  
 In dry sandy fields in Lower Carolina. ☉ June, July.  
 v. s.
- debile.* 2. *P. spica plerumque unica tenui, glumis uniseriatis, contingue alternis solitariis pubescentibus, brevi-obovatis, foliis hirsutis, culmo debili setaceo.* *Mich. fl. amer. 1. p. 43.*  
 On the sea-shores of Carolina and Georgia. *Michaux.* +.
3. *P. spicis orbiculare linearifolia americana.* In clay soil.  
 4. *P. spicis to-ovalis vaginis glabri p. 44.* P. lentiformis  
P. membranacea  
In dry soil.  
 5. *P. spicis glabri glabri americanum.* P. virga-aurea  
In sandy soil.  
 6. *P. glaberrima subtriangularis valvula longipetiolata.* P. paniculata  
In Georgia.  
 7. *P. spicigera glumosa, stratiformis.* P. stolonifera  
P. racemosa  
Milium iconocarpum  
Iconocarpus  
In cedar woods.  
 I found  
plants  
ty, I  
collected  
very  
same  
as I  
collected

3. *P. spicis alternis subbinis, glumis subtriseriatis geminis orbiculato-ovatis obtusis glabris, foliis lanceolato-linearibus serrulato-ciliatis, culmo decumbente.* *Mich. fl. amer. 1. p. 44.*  
In clay soil where iron ore abounds : Virginia and Carolina. 2. July. v. v.
4. *P. spicis pluribus alternis, glumis biseriatis suborbiculato-ovatis glabris laevibus, foliis glabris, ligula ciliata, vaginis compressis, caule suberecto.* *Mich. fl. amer. 1. p. 44.*  
*P. lentiferum.* *Lam. encycl. 5. p. 30.*  
*P. membranaceum.* *Walt. fl. car. 75.*  
In dry meadows and grassy hills : Pennsylvania to Carolina. 2. June—Aug. v. v.
5. *P. spicis paucis erectis, glumis biseriatis suborbiculatis glabris laevibus, foliis inferioribus pilosis, superioribus glabris, vaginis longissimis, caule erecto.* *Mich. fl. amer. 1. p. 44.*  
*P. virgatum.* *Walt. fl. car. 75?*  
In sandy fields : Carolina to Georgia. 2. June. v. s.
6. *P. glaberimum; spicis pluribus alternis erectis, glumis subtriseriatis brevibus ovatis rufescensibus glabris, valvula plana juxta margines transverse plicata, foliis longis, caule erecto.* *Mich. fl. amer. 1. p. 45.*  
*P. paniculatum.* *Walt. fl. car. 75?*  
In Georgia and Florida. *Michaux.* +.
7. *P. spiculis elongato-racemosis subverticillatis pretentibus, stoloniferum.*  
glumis alternis ovatis serrulato-ciliatis transverse rugosis, foliis brevibus basi subcordatis, caule basi prostrato.  
*P. stoloniferum.* *Bosc. in linn. trans. 2. p. 83.*  
*P. racemosum.* *Jacq. ic. rar. t. 302.*  
*Milium latifolium.* *Cav. ic. 3. p. 37. t. 273.*  
*Icon.* *Bosc. l. c.* *Jacq. l. c.*  
In cedar swamps of New Jersey. 2. July, Aug. v. v.  
I found this plant among a parcel of grasses and other plants collected in autumn 1811, in Monmouth county, New Jersey. As I had no time to examine the collection then, I was much surprised on finding this very interesting grass among them. That it is the same with *P. stoloniferum* of Peru, I have no doubt, as I carefully compared it with the specimens in the collection of A. B. Lambert, Esq.

82. ARISTIDA. *Gen. pl. 125.**dichotoma.*

1. A. cespitosa, culmis dichotomis, floribus subspicatis, subuniaristatis: arista contorta gluma breviore: lateralibus brevissimis. *Mich. fl. amer. 1. p. 41.*

On dry gravelly soil: New England to Carolina, common. ♂. May—July. *v. v.* Resembles *Nardus stricta*.

*stricta.*

2. A. culmis foliisque stricte erectis, racemo longo spicato coarctato: aristis gluma longioribus, foliis convoluto-filiformibus, glabris. *Mich. fl. amer. 1. p. 41.*

In shady rocky situations, on river sides: Virginia, Carolina, &c. ♀. June, July. *v. v.*

*oligantha.*

3. A. culmo stricto subramoso, foliis convoluto-filiformibus, floribus distantibus solitariis, aristis longis divaricatis. *Mich. fl. amer. 1. p. 41.*

A. adscensionis. *Walt. fl. car. 74.*

On the banks of rivers, in rocky places: Illinois, Virginia, &c. ♀. June, July. *v. v.*

*juncea.*

1. S. foliis convoluto-filiformibus, panicula laxa, calycibus subaristatis semine longioribus, aristis longis vix pubescentibus.

S. juncea. *Willd. sp. pl. 1. p. 440.*

Icon. *Desf. atl. 1. t. 28.*

On the banks of the Missouri. *M. Lewis.* ♀. Aug. *v. s. in Herb. Lewis.* By a careful comparison of this plant and one from the north-west coast of America, collected by Mr. Menzies, in possession of A. B. Lambert, Esq. and a specimen from the south of Europe in the same collection, I could find no other distinction from *S. juncea* than the seeds being more pubescent, and the arista not so much as figured in Desfontaines' Flora.

*canadensis.*

2. S. foliis setaceis, panicula parva, calycibus glabris obtuse-ovatis semine pubente æquantibus, aristis crassis brevibus. *Lam. encycl. 7. p. 452.*

A. juncea. *Mich. fl. amer. 1. p. 54.*

In rocky places: Canada, Hudson's Bay, &c. +.

*avenacea.*

3. S. foliis striatis glabris, panicula patente subunilateralis: ramis verticillatis ramulosis, calycibus acutis membranaceis semine glabro æquantibus, aristis nudis, tortuosos.—*Willd. sp. pl. 1. p. 442.*

In barren  
*v. v.*

4. S. foliis comis simi-  
natis sen-  
barbato.

*S. barbata.*

*S. avenacea.*

*S. virginica.*  
Icon. *Cav-*

In shady *v.*  
*v. v.*

5. S. foliis ti-  
tenuissimis  
viore, a-

*S. capillaris.*  
*S. diffusa.*

In sandy f-

Aug.  
panicles  
deviate  
short gl-

separate

6. S. foliis t-

panicul-

cibus c-

*Lam. e*

In Carolin-

7. S. foliis duncul-

uousis.

In Caroli-

8. *S. vagini-*  
ribus v

*Erianthus*  
*Anthoxa-*

*Andropo-*

In low p-

da, ra-

high,

are ex-

In barren dry woods: Virginia, Carolina, &c. 4. June.  
v. v.

4. S. foliis convoluto-filiformibus, panicula pauciflora: ramis simplicibus 1-2-floris, calycibus oblongis acuminate semine vix longioribus, corollis stipitatis: stipite barbato.—*Vahl. symb. 2. p. 24.*

*S. barbata. Mich. fl. amer. 1. p. 53.*

*S. avenacea. Walt. fl. car. 77.*

*S. virginica. Pers. engr. 1. p. 99.*

Icon. *Cav. ic. 5. t. 466. f. 2?* *Winter-green.*

In shady woods: Virginia to Georgia. 4. July, Aug.  
v. v.

5. S. foliis tereti-filiformibus longissimis, panicula effusa tenuissime capillacea, calyce corolla parva triplo breviore, aristis nudis strictis. *Mich. fl. amer. 1. p. 54.*

*S. capillaris. Lam. encycl. 7. p. 450.*

*S. diffusa. Walt. fl. car. 78.*

In sandy fields: New Jersey to Carolina. 4. June—Aug. v. v. This elegant grass, whose silky purple panicles exceed in beauty any other I know, seems to deviate from the rest of this genus, particularly in the short glumes of the calyx, and almost deserves to be a separate genus.

6. S. foliis striatis glabris glaucescentibus, spicis alternis paniculatis expansis, floribus sessilibus remotis, calycibus corolla longioribus, arista brevissima nuda.—*Lam. encycl. 7. p. 453.*

In Carolina. *Bosc. +.*

7. S. foliis arundinaceis, panicula elongata aristata, pedunculis articulatis strictissimis, aristis nudis subflexuosis. *Lam. encycl. 7. p. 453.*

In Carolina. *Fraser. +.*

*bicolor.*

*sericea.*

*expansa.*

*stricta.*

#### 84. SACCHARUM. Gen. pl. 104.

1. S. vaginis lanatis, gluma villis breviore, valvis exteriores villosis. *Pers. engr. 1. p. 103.*

*giganteum.*

*Erianthus saccharoides. Mich. fl. amer. 1. p. 55.*

*Anthoxanthum giganteum. Walt. fl. car. 65.*

*Andropogon culmo paniculato. Gron. fl. virg. 133.*

In low places generally among rocks: Virginia to Florida, rare. 4. July, Aug. v. v. From five to nine feet high, with very large panicles, which when in flower are extremely handsome.

- brevibarbis.* 2. *S. vaginis collo subvillosis, panicula stricta, gluma villosa longiore, valvis acutissimis nudis.* Pers. enc. 1. p. 103.

*Erianthus brevibarbis.* Mich. fl. amer. 1. p. 55.

On dry hills, in mountains: Virginia, Tennessee, Carolina, &c.  $\frac{1}{4}$ . July. v. v. Not above two feet high.

### 85. ANDROPOGON. Gen. pl. 1566.

#### \* *Panicula terminali.*

- nutans.* 1. *A. panicula ramosa coarctata hirtante, floribus geminatis aristatis, calycibus hirsutis, hermaphrodito sessili, masculo pedicellato caduco.* Willd. sp. pl. 4. p. 906. In barren dry situations: New England to Florida.  $\frac{1}{4}$ . June, July. v. v.

- avenaceus.* 2. *A. foliis asperis vaginisque glabris, panicula nuda subpyramidaliter oblonga multiflora, floribus rufescensibus triandris, glumæ valva altera villosa, arista valde contorta.* Mich. fl. amer. 1. p. 58.

In the prairies of Illinois. Michaux. +.

- ambiguus.* 3. *A. foliis amplexicaulibus subcordato-lanceolatis, panicula ramis promissa setaceis simplicibus erectis, glumis solitaria alternis unifloris, flore imberbi longissime aristato triandro, flosculi accessorii incremento aristiformi.* Mich. fl. amer. 1. p. 58.

In sandy woods: Carolina. Michaux. +.

- alopecuroides.* 4. *A. panicula simplici oblonga elongata, spicis verticillatis simplicibus, floribus aristatis geminatis, altero sessili, altero pedicellato, rachi lanata, lana floribus longiore.* Willd. sp. pl. 4. p. 911.

Icon. Sloan. hist. 1. t. 70. f. 1.

In North America. Linn. sp. pl. 1045. +.

This species is very doubtful as a North American plant; the synonym of *Gronovius* evidently belongs to *Saccharum giganteum*. Neither Michaux nor any other botanist (as far as I could learn) collected it in North America.

- Scoparium.* 5. *A. spicis simplicibus pedunculatis geminatis stricte paniculatis, floribus geminis, hermaphrodito sessili lanceolato aristato, neutro pedicellato aristato, rachi ciliata.* — Mich. fl. amer. 1. p. 57.

*A. purpurascens.* Willd. sp. pl. 4. p. 913.

*A. divaricatum.* Willd. sp. pl. 4. p. 905?

TRIAN

Common in grass.  $\frac{1}{4}$ .

\*\*

6. *A. vaginis in latim poly tuplo glu Mich. fl. e*

Cinna glome In low mead

7. *A. vaginis g sitis, flor terioribus Cinna latera In dry wood mountain*

8. *A. ramis re tristachyis lucri glu terioribus gam cont In the mou July. v*

9. *A. panicula rachi lat Willd. s*

In Virginia

10. *A. spicis co giato-cor hermaph rachi lor In Virginia*

11. *A spicis d maphroc rachi pil In rocky s*

July. +

### 86. AATHERO

1. *Atheropo*

In North

leaves n

Common in old fields and dry meadows ; called *Broom-grass*. ♀. June—Aug. v. v. A very obnoxious weed.

\*\* *Spicis conjugatis s. fasciculatis.*

6. A. *vaginis marginie villosis, paniculis lateralibus fasciculatis polystachyis, floribus monandris, arista 4-5-tuplo glumæ longitudine, valvis anterioribus glabris.* *Mich. fl. amer. 1. p. 56.*  
*Cinna glomerata. Walt. fl. car. 59.*  
In low meadows and woods : Carolina and Florida. ♀. +.
  7. A. *vaginis glabris, culmis erectis, spicis lateralibus distitis, floribus monandris, arista longa recta, valvis interioribus glabris.* *Mich. fl. amer. 1. p. 57.*  
*Cinna lateralis. Walt. fl. car. 59.*  
In dry woods and old fields, frequently found on high mountains : Pennsylvania to Florida. ♀. July. v. v.
  8. A. *ramis remote alternis solitariis simplicibus plerisque tristachyis, spicis distanter alternis bifidis, villis involucri gluma brevioribus, floribus triandris, valvis interioribus subvillosis, intima ex incisura aristam longam contortam emitente.* *Mich. fl. amer. 1. p. 57.*  
In the mountains of Virginia and North Carolina. ♀.  
July. v. v.
  9. A. *paniculae spicis conjugatis, pedunculis simplicibus, rachi lanata, flosculis muticis : mascula tabescente.* *Willd. sp. pl. 4. p. 916.*  
In Virginia. *Gronov.* ♀. +.
  10. A. *spicis conjugatis paniculatis, panicula bracteata fastigiatocorymbosa, floribus ternis muticis, intermedio hermaphrodito sessili, lateralibus neutris pedicellatis, rachi longissime ciliata.* *Willd. sp. pl. 4. p. 916.*  
In Virginia. *Gronov.* ♀. +.
  11. A. *spicis digitatis subquaternis, floribus geminatis, hermaphrodito sessili aristato, masculo mutico pedicellato, rachi pilosa.* *Willd. sp. pl. 4. p. 919.*  
In rocky situations : Canada to Virginia, rare. ♀. June, July. v. v.
86. AATHEROPOGON. *Muhl. in Willd. sp. pl. 4. p. 937.*
1. *Atheropogon.* *Willd. l. c.*  
In North America. *Muhl.* +. About a foot high ; leaves narrow ; flowers alternate in a spike.

*macrourus.*

*dissitiflorus.*

*ternarius.*

*virginicus.*

*bicornis.*

*furcatus.*

*apludoilles.*

## 87. TRisetum. Pers. enc. 1. p. 97.

*pratense.*

1. T. panicula subcontracta, spicis nitidis, vaginis longis striatis glabris, foliis superne villosis. Pers. l. c.  
*Avéna flavescens*. Willd. sp. pl. 1. p. 449.  
 Icon. Schreb. gram. t. 9. Host. gram. 3. t. 38.  
 In meadows: Pennsylvania and Virginia, rare. 4. July.  
 v. v.

## 89. AIRA. Gen. pl. 112.

*pumila.*

1. A. panicula parva fastigiata (pauciflora): pedicellis brevibus, floribus muticis obtusis calyce duplo longioribus: valvulis margine membranaceis, foliis planis glabris, culmo erecto vix foliis altiore.

On barren clay soil, near brick-yards: Pennsylvania. 4. June. v. v. Scarce an inch high; in close tufts.

*aquatica.*

2. A. panicula patente semiverticillata, floribus muticis obtusis laevibus calyce longioribus, foliis planis, culmo repente.—Willd. sp. pl. 1. p. 376.

Icon. Host. gram. 2. t. 41. Engl. bot. 1557.  
 On water sides and sometimes floating in water, frequent.  
 4. May—July. v. v.

*melicoides.*

3. A. panicula parva subracematum coarctata: gluma communi partialibusque majusculis linear-lanceolatis muticis: floribus basi villosis, rudimento flosculi tertii, foliis planis glabris, culmo erecto. Mich. fl. amer. 1. p. 62.

In shady woods: Canada to Pennsylvania. 4. July. v. v.

*obtusata.*

4. A. panicula in racemum oblongum confertiflorum coarctata, floribus muticis apice compressis, glumæ valva altera angusto-lineari acuta: altera lato-obovali rotundato-obtusissima, foliis planis, culmo gracili erecto.

*Mich. fl. amer. 1. p. 62.*

In dry fields: New England to Florida. 4. June, July. v. v.

*brevifolia.*

5. A. panicula semiverticillata patente: ramis simpliciusculis alternifloris, floribus muticis calycem æquantiibus, glumis omnibus lanceolatis acutis, foliis planis, culmo stricto glabro nudiusculo.

In the plains of Missouri. *M. Lewis.* 4. June, July. v. s. in *Herb. Lewis.* This grass is the most common in those plains.

*capillacea.*

6. A. panicula capillacea effusa, flosculis muticis calyce

- T
- longioribus  
asperis, foliis  
A. capillacea  
On slate hill  
4. July.  
purple.
7. A. foliis seta  
spicatis, 1  
p. 380.  
Icon. Fl. c  
On dry sand
8. A. panicula  
recta brev  
Icon. Host.  
In meadow
- β. A. foliis sub  
in Canada.  
pitosa.
9. A. panicula  
aristis gr  
Willd. sp.  
Icon. Hos  
dan. 157  
In shady w  
July. v.
10. A. panicula  
altera bi  
concava,  
car. 78.  
In Carolina
1. M. panicu  
bus obtus  
fl. amer.  
M. mutica  
Icon. Mo  
In shady p
2. M. panicu  
bus, ca  
M. altissim  
In sandy s

- longioribus : altero pedicellato, glumis acutis carina  
asperis, foliis angustis glabris brevibus, culmo erecto.  
A. capillacea. *Lam. illustr.* 1. p. 177.  
On slate hills and in sandy fields : Virginia to Carolina.  
4. July. v. v. About eight inches high, panicle  
purple.
7. A. foliis setaceis, vaginis angulatis, floribus paniculato-  
spicatis, flosculis basi aristatis. *Willd. sp. pl.* 1.  
p. 380.  
Icon. *Fl. dan.* 383.  
On dry sandy hills : Virginia, &c. ①. June. v. v.
8. A. panicula patente, petalis aristatis basi villosis : arista  
recta brevi. *Willd. sp. pl.* 1. p. 378.  
Icon. *Host. gram.* 2. t. 42. *Engl. bot.* 1153.  
In meadows, frequent. 4. June, July. v. v.
- β. A. foliis subsetaceis. *Mich. fl. amer.* 1. p. 61.  
In Canada. +. Appears to be only a variety of A. ces-  
pitosa.
9. A. panicula patente trichotoma : pedunculis flexuosis,  
aristis geniculatis, foliis setaceis, culmo subnudo.—  
*Willd. sp. pl.* 1: p. 378.  
Icon. *Host. gram.* 2. t. 43. *Engl. bot.* 1519. *Fl.*  
*dan.* 157.  
In shady woods and meadows : Canada to Carolina. 4.  
July. v. v.
10. A. panicula sparsa purpurea, floribus aristatis, glumis  
altera bifida nervo aristata : altera integra plumosa  
concava, foliis subulatis, culmo cresto.—*Walt. fl.*  
*car.* 78.  
In Carolina. *Walter.* +.

90. MELICA. *Gen. pl.* 113.

1. M. panicula laxa pauciflora : ramulis simplicibus, flori-  
bus obtusis imberibus, caule erecto glabro.—*Mich.*  
*fl. amer.* 1. p. 62.  
M. mutica. *Walt. fl. car.* 78.  
Icon. *Moris. hist.* 3. s. 8. t. 7. f. 51.  
In shady places : Virginia to Florida. 4. July. v. s.
2. M. panicula diffusa ramosissima, floribus acutis imberbi-  
bus, caule erecto pubescente.  
M. altissima. *Walt. fl. car.* 78.  
In sandy swamps : Virginia and Carolina. 4. June. v. s.

91. HOLCUS. *Gen. pl.* 1565.*striatus.*

1. *H. panicula conferta oblonga*: pedunculis brevissimis confertis ramosis, glumis bifloris striatis muticis acuminatis, foliis planis longis.—*Willd. sp. pl.* 4. p. 935.  
In swamps of Virginia. *Clayt.* +. This species appears to belong to *Aira*.

*fragrans.*

2. *H. panicula patula*, glumis trifloris muticis, flosculis confertis, hermaphrodito intermedio diandro glabro, masculis triandris, valvula exteriore levissime ciliata, dorso laevibus. *Willd. sp. pl.* 4. p. 936.

*H. odoratus.* *Mich. fl. amer.* 1. p. 56.

In low meadows on the borders of lakes: Canada to Virginia. ♀. July. v. v. A very fine grass.

*lanatus.*

3. *H. glumis bifloris*, hermaphrodito mutico, masculi aristata flore multo breviore recurva. *Willd. sp. pl.* 4. p. 933.

Icon. *Host. gram.* 1. t. 2. *Schreb. gram.* t. 20. f. 1.  
In low meadows, frequent. ♀. June, July. v. v.

92. SORGHUM. *Pers. enc.* 1. p. 101. Broom-grass.

- saccharatum.* 1. *S. panicula subverticillata patentissima*, seminibus ellipticis glumis villosis persistentibus tectis. *Pers. enc.* 1. p. 101.

Icon. *Mich. in act. helv.* 8. t. 4. f. 1.

Cultivated for the purpose of making brooms, and frequently found in an apparently wild state. ♂. July. v. v.

93. POA. *Gen. pl.* 114. Meadow-grass.*trivialis.*

1. *P. panicula diffusa*, spiculis trifloris, glumis lanceolatis quinque nervibus basi villo connexis, ligula elongata. *Smith. fl. brit.* 1. p. 103. *Willd. sp. pl.* 1. p. 387.

Icon. *Host. gram.* 2. t. 62. *Engl. bot.* t. 1072.

Common in meadows and fields. ♀. June—Aug. v. v.

*pratensis.*

2. *P. panicula diffusa*, spiculis quadrifloris, glumis lanceolatis quinque nervibus villo connexis, ligula abbreviata obtusa. *Smith. fl. brit.* 1. p. 104. *Willd. sp. pl.* 1. p. 388.

Icon. *Host. gram.* 2. t. 61. *Engl. bot.* t. 1073.

In grassy fields, not so frequent. ♀. May, June. v. v.

3. *P. panicula lanceolata gram.*  
In all mead Aug. v  
4. *P. panicula trifloris, brevissim*  
Icon. *Ho*  
In grassy s  
5. *P. panicula tiusculis presso.*  
Icon. *Ho*  
In fields a  
6. *P. panicula ovatis su*  
*Smith fl*  
Icon. *En*  
On the All  
7. *P. panicula sp. pl.* 1  
In Pensylv  
8. *P. panicula presso, nesis.* S  
Icon. *Ho*  
In dry fiel  
v. v.  
9. *P. panicula floribus subangu*  
*Willd.*  
*P. striata.*  
*P. lineata.*  
In meadow  
10. *P. panicula queflori collo p*  
p 394.  
Icon. M  
In fields a  
♂. Jun

3. *P. panicula diffusa, spiculis ovatis subquadriifloris, glumis lanceolatis trinervibus, ligula brevi truncata.* Schreb. *gram.*  
In all meadows and grass plots, common. ♀. June—Aug. v. v.
4. *P. panicula foliisque attenuatis, spiculis lanceolatis subtrifloris, glumis acutis obsolete quinquenervibus, ligula brevissima crenata.* Smith *fl. brit.* 1. p. 106.  
Icon. *Host. gram.* 2. t. 71. *Engl. bot.* 1265.  
In grassy shady places, rare. ♀. June, July. v. v.
5. *P. panicula divaricata, spiculis ovatis, flosculis remotiusculis quinquenervibus liberis, culmi obliquo compresso.* Smith *fl. brit.* 1. p. 105. Willd. *sp. pl.* 1. p. 390.  
Icon. *Host. gram.* 2. t. 64. Leers *fl. herb.* t. 6. f. 1.  
In fields and grass plots, frequent. ♂. April—Sept. v. v.
6. *P. panicula diffusa, spiculis quadrifloris cordatis, glumis ovatis subfalcatis liberis, ligulis inferioribus brevissimis.* Smith *fl. brit.* 1. p. 100. Willd. *sp. pl.* 1. p. 386.  
Icon. *Engl. bot.* 1003. Scheuchz. *gram.* t. 3.  
On the Allegany mountains, frequent. ♀. July. v. v.
7. *P. panicula diffusa, spiculis ovato-oblongis nitidis.* Willd. *sp. pl.* 1. p. 390.  
In Pennsylvania and Virginia. ♀. v. s. in *Herb. Walter.*
8. *P. panicula secunda coarctata, culmo adscendente compresso, flosculis angulosis basi villo complicato connessis.* Smith *fl. brit.* 1. p. 99. Willd. *sp. pl.* 1. p. 397.  
Icon. *Host. gram.* 2. t. 70. *Engl. bot.* 365. Blue-grass.  
In dry fields and meadows, common. ♀. June—Aug. v. v.
9. *P. panicula æquali diffusa, spiculis ovatis quinquefloris, floribus liberis septemnervibus obtusis, culmo sulcato subangulato, radice subrepente.* Willd. *enum.* 105.  
Willd. *sp. pl.* 1. p. 389.  
*P. striata.* Mich. *fl. amer.* 1. p. 69.  
*P. lineata.* Pers. *ench.* 1. p. 89.  
In meadows and fields, common. ♀. June—Aug. v. v.
10. *P. panicula laxa patentissima capillari, spiculis tri-quinquefloris, floribus ovatis acutis pubescentibus, foliis collo pilosis, culmo ramosissimo.*—Willd. *sp. pl.* 1. p. 394.  
Icon. *Moris. hist.* 3. s. 8 t. 6. f. 33.  
In fields and sandy woods: Canada to Florida, common.  
♂. June—Aug. v. v. Very variable.

*viridis.**nemoralis.**annua.**alpina.**flava.**compressa.**nervata.**capillaris.*

- tenella.* 11. *P. panicula subverticillata capillari, pedunculis flexuosis, spiculis linearibus subsexfloris pedicellatis, floribus oblongis acutis minutis glabris, foliis brevibus planis apice subulatis glabris, vaginis collo barbatis culmo decumbente.*  
*P. tenella. Willd. sp. pl. 1. p. 395?*  
*In sandy fields: New Jersey to Carolina. ♂. July. v. v.; v. s. in Herb. Lambert. sub nomine *P. pilosæ*.*
- erecta.* 12. *P. panicula laxa patula, spiculis ovatis quadri-quinquefloris pedicellatis, floribus oblongis subpubescentibus, foliis glabris, vaginis collo nudo, culmo erecto glabro.*  
*Mich. fl. amer. 1. p. 68.*  
*In Canada, Hudson's Bay, and Lake Mistassins. Michaux. +.*
- hirsuta.* 13. *P. panicula ramosissima capillari, spiculis sparsis longe pedicellatis subquinquefloris, floribus oblongis acutis glabris, foliis culmo longioribus glabris, vaginis hirsutissimis, culmo erecto crasso.*  
*Mich. fl. amer. 1. p. 68.*  
*In sandy fields: New Jersey to Carolina, frequent. ♂. July. v. v.*
- aquatica.* 14. *P. panicula erecta semiverticillata: ramulis flexuosis, spiculis ovatis quinquefloris crassis, floribus obtusis glabris septemnervibus, culmo erecto glabro tereti.*  
*P. aquatica. Willd. sp. pl. 1. p. 385?*  
*In ditches, frequently in salt-marshes: Canada to Virginia. ♀. June—Aug. v. v. It approaches *P. aquatica* very nearly, but seems to differ in *habitus* and other characters considerably.*
- parviflora.* 15. *P. panicula diffusa capillari semiverticillata ramosissima spiculis parvis subquadrifloris, floribus oblongis subacutis exquisite striatis, foliis distichis culmum superantibus glabris, ligulis lanceolatis acuminatis, culmo exerto vaginato.*  
*P. striata. Lam. encycl. 5. p. 84?*  
*In close copses and shady places: New York to Virginia. ♀. July. v. v. About a foot high; very slender.*
- subverticilla-* 16. *P. panicula laxa basi subverticillata, pedicellis elongatis, spiculis rarissimis subquinquefloris, calycibus crassiusculis subangulatis obtusis.*  
*Pers. encycl. 1. p. 94.*  
*P. laxa. Lam. illustr. 1. p. 183.*  
*In Virginia. Lamarck. +.*
- Eragrostis.* 17. *P. panicula patente pyramidata, ramulis divaricatis, spiculis linearibus decemfloris, floribus ovatis acutis tri-*

nervib  
p. 39  
Icon. I  
In low sa  
June, I  
and lon  
18. *P. panic  
culis c  
rum v  
losa ap  
nicula  
P. sesleri  
In mour  
rolina  
v. v.  
meado  
twice  
withou  
ing, 1  
the le  
being  
19. *P. panic  
cimflo  
ribus  
natis  
pilosis  
1. p.  
In sandy  
ern te  
20. *P. panic  
nearit  
dorsal  
pilosis  
Poa am  
In dry b  
July.  
purpl  
21. *P. panic  
multi  
vibus  
fl. an  
P. hyprr  
Icon. I  
In mos****

- nervibus glabris, foliis brevibus.—*Willd.* sp. pl. 1.  
p. 392.  
Icon. *Host. gram.* 2. t. 69?  
In low sandy grass fields: Virginia, Carolina, &c. ¼.  
June, July. v. v. The spikes are of a greenish colour  
and long.
18. *P. panicula erecta*: ramis inferioribus inferne nudis, spi-  
culis oblongis alternis subsessilibus sexfloris, flosculo-  
rum valva exteriore dorso inferne marginibusque vil-  
losa apice quinquefida, vaginarum collo axillisque pa-  
niculae pilosis, foliis culmoque erecto glabris. *Mich. fl. amer.* 1. p. 68.  
In mountain meadows, frequent: New England to Ca-  
rolina. Commonly called *Red-top*. ¼. July, Aug.  
v. v. A most excellent grass. I have seen mountain  
meadows in Pennsylvania, where they mow this grass  
twice a year, producing most excellent crops each time,  
without manure, or any other trouble than the mow-  
ing, lasting for the space of sixteen years, without  
the least decline in the crops, the soil at the same time  
being a very indifferent one.
19. *P. panicula laxa patula erecta*, spiculis linearibus duode-  
cimfloris: rachi deflorata valvulis flosculorum interio-  
ribus persistentibus pectinata, floribus ovatis acumi-  
natis trinervibus, vaginarum collo axillisque paniculae  
pilosis, foliis culmoque erecto glabris.—*Mich. fl. amer.*  
1. p. 69.  
In sandy fields: New Jersey to Carolina, and the west-  
ern territories, frequent. ⊖. July. v. v.
20. *P. panicula divaricata ramosissima*, spiculis pendulis li-  
nearibus decemfloris, floribus ovatis acutis margine  
dorsoque scabris, vaginarum collo axillisque paniculae  
pilosis, foliis culmoque erecto brevi glabris.  
*Poa amabilis*. *Walt. fl. car.* 80.  
In dry barren sand-fields: New York to Carolina. ⊖.  
July. v. v. A beautiful grass: the large panicle is  
purple mixed with green stripes.
21. *P. paniculus fasciculatis*, spiculis subsessilibus longissimis  
multifloris, floribus oblongis acutis laxis, foliis bre-  
vibus pubescentibus, culmo ramoso repente.—*Mich.*  
*fl. amer.* 1. p. 69.  
*P. hypnoides*. *Lam. encycl.* 5. p. 87.  
Icon. *Mich. l. c. t. 11.*  
In mossy swamps and shady places: Pennsylvania, Vir-

ginia, and on the banks of the Mississippi. ☎ July. v. v. This is the most delicate grass in North America.

94. BRIZA. *Gen. pl. 115.**Eragrostis.*

1. B. spiculis ovali-lanceolatis vigintifloris, floribus subacutis, collo vaginarum piloso, culmo geniculato decumbente. *Mich. fl. amer. 1. p. 72.* *Willd. sp. pl. 1. p. 405.*

*B. caroliniana. Walt. fl. car. 79.**Icon. Schreb. gram. 2. t. 39.*

In wet fields and swamps: Virginia to Carolina. ☎ June. v. v.

*canadensis.*

2. B. panicula laxa, spiculis erectis 4—10-floris, gluma communi parvula, florum valva exteriore acuta ovata, foliis longis, culmo erecto. *Mich. fl. amer. 1. p. 71.*  
In Canada. *Michaux. +.*

*virens.*

3. B. spiculis ovatis, calyce flosculis (7) æquali, pedunculis longis. *Walt. fl. car. 79.*  
In Carolina. *Walter. +.*

*latifolia.*95. UNIOLA. *Gen. pl. 116.**paniculata.*

1. U. panicula laxa, spiculis omnibus longe pedicellatis, calyce trivalvi, floribus monandris subfalcatis carina pilosis, foliis latis planis. *Mich. fl. amer. 1. p. 70.*  
On the Allegany mountains, in shady woods, among rocks. ¼. June. v. v. A very handsome grass.

*gracilis.*

2. U. panicula longa, spiculis subsessilibus, calyce multivalvi, floribus triandris carina glabris, foliis convolutis, culmo altissimo.—*Willd. sp. pl. 1. p. 406.*

*U. maritima. Mich. fl. amer. 1. p. 71.**Icon. Catesb. car. 1. t. 32. Pluk. alm. t. 32. f. 6.*

On sandy sea-shores: Virginia to Florida. ¼. June, July. v. v.

3. U. panicula longissima subspicata, ramis brevibus adpressis, spiculis subsessilibus, floribus monandris di-varicatis acuminatis glabris, calyce trivalvi, foliis planisculis, vaginis culmoque glabris compressis.—*Mich. fl. amer. 1. p. 71.*

*U. virgata. Bartram. in Herb. Banks.**Holcus laxus. Willd. sp. pl. 4. p. 934.*

In shady rocky situations: Virginia to Georgia. ¼. July. v. v.

1. F. sp. florib. foliis  
On the June
2. F. pan. florib. eulin. 113. F. bro. F. octo. Comm
3. F. pan. florib. setae F. qua. In dry App
4. F. pan. tere brit. Icon. In dry
5. F. pan. ovat. nerv. p. 4 Icon. In we intr. v. v
6. F. pan. oblong. 117 In dr Illin
7. F. pan. spic. acu. per. Icon.

## 96. FESTUCA. Gen. pl. 119.

1. *F. spiculis alternis sessilibus erectis subquinquefloris, floribus subulatis glabriusculis, aristis longis scabris, foliis linearibus culmoque glabris.* *spicata.*  
On the waters of the Missouri and Columbia rivers.  
June. v. s. in *Herb. Lewis.*
2. *F. panicula simplicissima secunda, spiculis subnoven-floris aristatis, foliis linear-setaceis, ligula biaurita, culmo superne tetragono, basi ramosa.* *Willd. enum.*  
*113. Willd. sp. pl. 1. p. 419.*  
*F. bromoides. Mich. fl. amer. 1. p. 66.*  
*F. octoflora. Walt. fl. car. 81.*  
Common in sandy fields. ☽. May—July. v. v.
3. *F. panicula coarctata simplici, spiculis subquadrifloris, floribus subulatis aristatis, culmo foliisque tenuissime setaceis.* — *Mich. fl. amer. 1. p. 66.* *Myurus.*  
*F. quadriflora. Walt. fl. cur. 81.*  
In dry fields: Virginia and Carolina. ☽. June. v. v.  
Appears to be different from *F. Myurus* of Linnæus.
4. *F. panicula secunda diffusa, flosculis aristatis, culmo tereti, foliis caulinis planis, radice fibrosa.* *Smith fl. brit. 1. p. 115. Willd. sp. pl. 1. p. 421.*  
Icon. *Host. gram. 2. t. 83. Engl. bot. 470.*  
In dry pastures, frequent. ♀. July. v. v.
5. *F. panicula subdiffusa nutante ramosissima laxa, spiculis ovato-lanceolatis acutis, flosculis cylindricis obsolete nervosis.* *Smith fl. amer. 1. p. 124. Willd. sp. pl. 1. p. 425.* *duriuscula.*  
Icon. *Host. gram. 2. t. 79. Engl. bot. 1593.*  
In wet meadows: Pennsylvania, New England, &c. introduced with grass seeds from Europe. ♀. July. v. v.
6. *F. panicula elongata, ramis approximatis strictis, spiculis oblongis 8—10-floris adpressis aristatis.* *Willd. enum.*  
*117. Mich. fl. amer. 1. p. 66.*  
In dry meadows: Virginia and Carolina mountains, Illinois, &c. ♀. July. v. v.
7. *F. panicula stricte contracta, ramis simplicibus sparsis, spiculis tereti-lanceolatis subquinquefloris, floribus acuminatis diandris, foliis lati planis, caule erecto asperimo.* *Mich. fl. amer. 1. p. 67.* *diandra.*  
Icon. *Mich. fl. amer. 1. t. 10.*

In shady woods, in the western parts: Kentucky, Tennessee, &c. ♀. July. v. v.

- grandiflora.* 8. F. panicula simplici erecta, spiculis perpaucis subseptemfloris, flosculis acutis distantibus. *Lam. illustr.* 1. p. 191.  
In Carolina. *Fraser.* ♀. +.

- fluitans.* 9. F. panicula longa laxa, ramis simplicibus, spiculis adpressis tereti-linearibus multifloris, flosculis muticis striatis, culmo decumbente foliisque glaberrimis.—  
*Willd. sp. pl.* 1. p. 426. *Icon. Host. gram.* 2. t. 77. *Engl. bot.* 1520.  
Icon. *Host. gram.* 2. t. 77. *Engl. bot.* 1520.  
In ditches and slow-flowing waters, frequent. ♀. June—Aug. v. v.

- distichiphyl-la.* 10. F. panicula racemoso-coarctata, spiculis approximatis adpressis subsexfloris muticis, foliis convoluto-subulatis rigidis alternis distichis, culmo eretto glabro.—  
*Mich. fl. amer.* 1. p. 167.

*F. triticea.* *Lam. illustr.* 1. p. 190.

*Uniola spicata.* *Willd. sp. pl.* 1. p. 406.

Icon. *Pluk. alm.* t. 33. f. 4?

In salt-meadows, common along the coast from Canada to Florida. ♀. July, Aug. v. v.

- poooides.* 11. F. panicula subconferta, ramis inferioribus simplicibus, spiculis alternis oblongis subsessilibus muticis, foliis planis glaberrimis.—*Mich. fl. amer.* 1. p. 67.  
On the banks of the river St. Laurence. ♀. +.

12. F. panicula nutante patula, spiculis oblongo-lanceolatis compressis octofloris muticis, foliorum vaginis apice barbatis, radice fibrosa. *Willd. enum.* 115.

Icon. *Willd. hort. ber.* t. 3.

In rich bottom lands: Pennsylvania to Carolina. ♀. July, Aug. v. v.

- nutans.* 13. F. paniculæ ramis secundis nutantibus scabris, spiculis ovatis compressis subsexfloris, spiculis acutis muticis, foliis linear-lanceolatis. *Willd. enum.* 116.  
In shady woods: New England to Carolina. ♀. July. v. v.

#### 97. DACTYLIS. Gen. pl. 117. Orchard-grass.

- glomerata.* 1. D. panicula secunda glomerata. *Willd. sp. pl.* 1. p. 408.  
Icon. *Host. gram.* 2. t. 94. *Schreb. gram.* t. 8 f. 2.  
In meadows and woods: Pennsylvania, &c. Probably originally brought from Europe. ♀. June. v. v.

98. BROMUS. *Gen. pl. 120.*

1. *B. panicula nutante, spiculis ovatis compressis, glumis nudis distinctis, aristis subulatis brevioribus flexuoso-rectis.* Willd. *sp. pl. 1. p. 428.*  
Icon. *Host. gram. 1. t. 12.*

In rye- and wheat-fields, frequent; introduced from Europe. ♂. July. v. v.

2. *B. panicula erecta coarctata, pedunculis ramosis, spiculis ovatis, flosculis imbricatis depressis nervosis pubescentibus.* Smith *fl. brit. 1. p. 126.* Willd. *sp. pl. 1. p. 429.*

Icon. *Host. gram. 1. t. 19.* *Engl. bot. 1078.*  
In dry fields and pastures; probably from Europe. ♂.  
June. v. v.

3. *B. panicula nutante, spiculis lanceolatis teretibus, flosculis aristatis pilosis, aristis erectis, foliis utrinque glabris, vaginis pilosis.* Willd. *enum. 120.* Willd. *sp. pl. 1. p. 431.*

In old fields and on road-sides: Canada to New England.  
♀. May—Aug. v. v.

4. *B. panicula nutante, spiculis oblongis compressis 6—8-floris, flosculis aristatis margine ciliato-villosis, foliis utrinque vaginisque subpilosis, culmo ad nodos barbato.* Willd. *sp. pl. 1. p. 433.*

*B. canadensis.* Mich. *fl. amer. 1. p. 65?*  
In barren and sandy soil: Canada to Carolina. ♀. July.  
v. v.

*secalinus.**mollis.**purgans.**ciliatus.*99. KOELERIA. *Pers. enc. 1. p. 97.*

1. *K. spica laxiuscula, spiculis divergentibus 3—4-floris subaristatis rugulosis carina subciliatis.* Pers. *t. c.*

*cristata.*

*Aira cristata.* Sp. *pl. 2. p. 24.*

*Poa cristata.* Willd. *sp. pl. 1. p. 402.*

Icon. Moris. *hist. 3. s. 8. t. 4. f. 7.*

On the plains of Columbia river. M. Lewis. ♀. July.  
v. s. in *Herb. Lewis.*

100. AVENA. *Gen. pl. 122. Oats.*

1. *A. panicula attenuata, calycibus bifloris, seminibus vil-pensylvanica.* losis, aristis calyce duplo longioribus. Willd. *sp. pl. 1. p. 445.*

In fields and open woods: New England to Carolina,  
frequent. ♂. June. v. v.

*palustris.*

2. *A. panicula conferta, calycibus bifloris, floribus glabris : altero mutico, altero tenuissime aristato, foliis planis linearibus.* *Mich. fl. amer.* 1. p. 72.

*A. caroliniana.* *Walt. fl. car.* 81.

In low grassy fields and woods : Virginia to Florida.  $\frac{1}{2}$ . May - July. v. v.

*striata.*

3. *A. panicula coarctata rariflora. calycibus quinquefloris, floribus calyce longioribus subimberibus, valva exteriore striata, dorso longe aristata : semine nudo.* *Mich. fl. amer.* 1. p. 73.

On the mountains near Hudson's Bay. *Michaux.* +.

*mollis.*

4. *A. racemo confertiuscula adpresso-spicato, calycibus bifloris, floribus imberibus aristatis : aristis longitudine florum strictis, foliis culmoque molliter pubescentibus.* — *Mich. fl. amer.* 1. p. 72.

In Canada. *Michaux.* +.

*spicata.*

5. *A. panicula subspicata pauciflora, calycibus spiculis sex-floris pubescentibus longioribus, valva exteriore ex apice bicorni aristata : arista basi spirali, foliis subsecundis collo villosis.* — *Willd. sp. pl.* 1. p. 453.

*A. glumosa.* *Mich. fl. amer.* 1. p. 72.

On dry hills, on the banks of rivers : New England to Carolina.  $\frac{1}{2}$ . June—Aug. v. v.

101. ARUNDO. *Gen. pl.* 124. Reed.*Phragmites.*

1. *A. calycibus quinquefloris, panicula laxa.* *Willd. sp. pl.* 1. p. 454.

*Icon.* *Engl. bot.* 401. *Leers. herb.* t. 7. f. 1.

On the banks of rivers and in large salt-marshes : Canada to Virginia, common.  $\frac{1}{2}$ . July. v. v.

*airoides.*

2. *A. panicula subcoarctata incurva, calycibus bifloris glabris inaequalibus, corollis membranaceis longitudine calycis, pilis corollam aequantibus, foliis planis scabris.* — *Lam. encycl.* 6. p. 270.

In North America. *Michaux.*  $\frac{1}{2}$ . +.

*agrostoides.*

3. *A. panicula laxa diffusa, calycibus unifloris acuminatis glabris, corollis calyce brevioribus membranaceis, valva exteriore apice incisa, dorso brevissima aristata, interiore minima, pilis corollam aequantibus, foliis planis scabriusculis, culmis vaginisque glabris.*

In bogs of New Jersey and Pennsylvania.  $\frac{1}{2}$ . July. v. v.

*canadensis.*

4. *A. panicula oblonga, calycibus unifloris lanceolatis carina scabris pubescentibus, corollis dorso aristatis :*

arista  
foliis c  
p. 73.

On mou  
v. v.

5. *A. panic  
coroll  
exced*

In Nort

6. *A. pan  
bus,  
bus.*

Icon.

On the  
v. v.

102.

7. *E. glab  
gitati  
floris*

In san

①. J

the

the C

2. *E. vag  
latis  
flor*

In cul

①. J

103.

1. *C. spi  
mai  
glar*

Aegile

In wo

July

bea

2. *C. sp  
glar  
mo*

Cyno

Agro

Icon.

aris :  
anis  
  
4.  
  
oris,  
ex-  
udo.  
  
·  
· bi-  
dine  
bus.  
  
sex-  
: ex-  
bse-  
  
d to  
  
. pl.  
  
Ca-  
  
gla-  
dine  
bris.  
  
natis  
alva  
nt-  
lanis  
  
v. v.  
ca-  
itis :

arista longitudine corollæ, pilis corollam æquantibus, foliis culmisque glabris angustis.—*Mich. fl. amer. 1. p. 73.*

On mountain meadows: Canada to Carolina. 4. July.  
v. v. Resembles *A. Calamagrostis* very much.

5. *A. panicula patente, calycibus unifloris acuminatis, pilis corollam æquantibus, arista dorsali geniculata calycem excedente.* *Willd. enum. p. 127.*

In North America. *Willdenow.* +.

6. *A. panicula spicata, calycibus unifloris corolla longioribus, floribus erectis muticis, foliis involutis pungentibus.* *Smith fl. brit. 1. p. 148.*

*Icon. Fl. dan. 917.*

On the sea-coast: Canada to New England. 4. July.  
v. v.

### 102. ELEUSINE. *Gärt. de fr. et sem. 1. p. 7.*

1. *E. glabra; vaginis compressis, collo pilosis: spicis digitatis strictis longo-linearibus rectis, spiculis subsexfloris, floribus lanceolatis.* *Mich. fl. amer. 1. p. 64.*

In sandy cultivated grounds: New Jersey to Florida.

◎. June—Aug. v. v. In Virginia it is known by the name of *Wire-grass*, as a weed very noxious to the cultivator.

2. *E. vaginis pilosis, spicis numerosis filiformibus paniculatis, spiculis minutissimis, calyce mucronato quadriphoro, floribus muticis.* *Mich. fl. amer. 1. p. 65.*

In cultivated fields: Virginia, Carolina, Illinois, &c.  
◎. June, July. v. v.

### 103. CHLORIS. *Swartz. fl. ind. 1. p. 189.*

1. *C. spica unica, spiculis subsexfloris, floribus aristatis monostachya. margine barbatis, glumis calycis exteriore scrobiculata-glandulosa dorso aristata.* *Mich. fl. amer. 1. p. 59.*

*Aegilops aromaticum.* *Wall. fl. car. 249.*

In woods and cultivated fields: Virginia to Florida. ◎ July. v. v. The spike of this grass is most singularly beautiful.

2. *C. spicis 4—6. strictis erectis, flosculis imbricatis subglabris muticis, glumis calycis exteriore aristata, culmo compresso glaberrimo.* *Swartz. fl. ind. 1. p. 194.*

*Cynosurus paspaloides.* *Vahl. symb. 1. p. 21.*

*Agrostis complanata.* *Ait. kew. 1. p. 96.*

*Icon. Vahl. l. c. t. 27.*

*confinis.*

*arenaria.*

*indica.*

*mucronata.*

*petræa.*

## TRIANDRIA DIGYNIA. Chloris.

- On the gravelly sea-shores of Georgia and Florida. ♂. June. v. v.
- mucronata*. 3. C. culmo repente, spicis plerumque 4, rachi triangulares mucronatae, spicis subquadriifloris, gluma exteriore aristata.—*Mich. fl. amer.* 1, p. 59.  
*Aegilops saccharinum*. *Walt. fl. car.* 249.  
*Icon. Pluk. alm. t. 300, f. 8.* Wire-grass.  
 In cultivated grounds: Virginia to Georgia. ♂. June.  
 v. v.
- curtipendula*. 4. C. culmo adscendente racemoso, spiculis plurimis brevibus distichis alternis pendulis sexfloris, calycibus quadrifloris.—*Mich. fl. amer.* 1, p. 59.  
 In dry situations in the western parts of America, Illinois, &c. *Michaux.* 4. +.

104. ROTTBOELLIA. *Gen. pl.* 1572.

- dimidiata*. 1. R. spica dimidiata compressa linearis, latere exteriore aggregato-flosculosa, interiori laevi nudo, vaginis compressis, culmo decumbente. *Willd. sp. pl.* 1, p. 466.  
*Ischaemum secundum*. *Walt. fl. car.* 249.  
 On the sea-shores of Carolina and Florida. v. s. in *Herb. Walter.*

105. TRIPSACUM. *Gen. pl.* 1134.

- dactyloides*. 1. T. spicas subtenuis aggregatis superne masculis inferne femineis. *Willd. sp. pl.* 4, p. 201.  
*Ischaemum glabrum*. *Walt. fl. car.* 249.  
*Icon. Pluk. alm. t. 190, f. 2.*  
 On the banks of rivers and along the sea-shore: Virginia, Carolina, Illinois. ♀. Aug. v. v.
- monostachyon*. 2. T. spica solitaria superne mascula inferne feminea. *Willd. sp. pl.* 4, p. 202.  
*Icon. Willd. hort. ber.* 1, t. 1.  
 In salt meadows and ditches: New York to Carolina.  
 ♀. Aug. v. v. Not sufficiently distinct from the former.
- cylindricum*. 3. T. spica solitaria hermaphrodita cylindrica, spiculis contiguis in articulos secedentibus. *Willd. sp. pl.* 4, p. 202. *Mich. fl. amer.* 1, p. 60.  
 In sandy barren fields in Florida. *Michaux.* +.

106. El

1. E. spica per enum. 13  
In shady w.
  2. E. spica numeribus t *Willd. s.*  
In rocky si
  3. E. spica numeratis, fo  
In the mor  
v. v.
  4. E. spica er calycibus  
131.  
In dry rocl  
v. v.
  5. E. spica eratis, caly  
*Willd.* e  
In the mor
  6. E. spica ei natis, ca  
fere sup  
enum. 1  
In shady  
June.
107. 1
1. H. floscul duobns  
Icon. *Hos*  
In fields i  
introdu
  2. H. aristis  
pl. 1, p  
On the isl  
July. I  
I conside  
the spe  
Esq., I  
the Ho

106. ELYMUS. *Gen. pl.* 128. Wild Rye.

1. *E. spica pendula patula, spiculis sexfloris aristatis. Willd. philadelphicus. enum. 131. Willd. sp. pl. 1. p. 468.*  
In shady woods : Canada and New York. 4. July. v. v.
2. *E. spica nutante patula, spiculis sexfloris aristatis, inferioribus ternatis, superioribus binatis. Willd. enum. 131. Willd. sp. pl. 1. p. 468.*  
In rocky situations : Canada, &c. 4. July. +.
3. *E. spica nutante patula, spiculis sexfloris aristatis geminatis, foliis subtus glaucis. Willd. enum. 131. glaucifolius.*  
In the mountains of Virginia, Pennsylvania, &c. 4. July.  
v. v.
4. *E. spica erecta, spiculis trifloris aristatis villosis ternatis, calycibus aristatis spiculis superantibus. Willd. enum. 131. villosus.*  
In dry rocky soil : Pennsylvania, Virginia, &c. 4. June.  
v. v.
5. *E. spica erecta, spiculis trifloris aristatis glabris geminatis, calycibus lanceolatis nervosis spiculis æquantibus. Willd. enum. 131. Willd. sp. pl. 1. p. 469. virginicus.*  
In the mountains very common. 4. May—July. v. v.
6. *E. spica erecta, spiculis bifloris aristatis hispidis geminatis, calycibus linearibus nervosis aristatis spiculis fere superantibus, foliis vaginisque glabris. Willd. enum. 131. Willd. sp. pl. 1. p. 470. striatus.*  
In shady situations : Pennsylvania, Virginia, &c. 4. June. v. v.

107. HORDEUM. *Gen. pl.* 129. Barley.

1. *H. flosculis omnibus hermaphroditis aristatis, ordinibus duobus erectioribus. Willd. sp. pl. 1. p. 472. vulgare.*  
Icon. *Host. gram. 3. t. 34.*  
In fields and hedge-rows, apparently in a wild state ; introduced from Europe. 4. July. v. v.
2. *H. aristis involucrisque setaceis longissimis. Willd. sp. pl. 1. p. 476. excl. synon. Ait. kew. ed. 2. p. 180. jubatum.*  
On the islands of the Missouri river. *M. Lewis.* ♂.  
July. v. s. in *Herb. Lewis. et Lambert.* This species I considered at first a new one; but on examination of the specimens in the Herbarium of A. B. Lambert, Esq., I found it to be the same as the *H. jubatum* of the *Hortus Kewensis.*

108. SECALE. *Gen. pl. 127.* Rye.*cereale.*

1. *S. glumarum ciliis scabris.* *Willd. sp. pl. 1. p. 471.*  
Icon. *Host. gram. 2. t. 48.*  
Frequently found apparently in a wild state. ♂. June.  
*v. v.*

109. TRITICUM. *Gen. pl. 130.* Wheat.*œstivum.*

1. *T. calycibus quadrifloris ventricosis glabris imbricatis aristatis.* *Willd. sp. pl. 1. p. 476.*  
Icon. *Host. gram. 3. t. 26.*  
In corn-fields; introduced from Europe. ○. June. *v. v.*

## III. TRIGYNI.

110. HOLOSTEUM. *Gen. pl. 136.**succulentum.*

1. *H. foliis ellipticis carnosis, petalis subtrifidis calyce minoribus.* — *Willd. sp. pl. 1. p. 489.*  
*Polycarpon uniflorum.* *Walt. fl. car. 83.*  
A doubtful plant, which I never have been able to find, either in New York or Carolina.

111. POLYCARPON. *Gen. pl. 138.**stipulifidum.*

1. *P. erectum ramosissimum, ramis setaceis, foliis (radicibus) spatulatis, floribus terminalibus fasciculatis, stipulis setaceo-multifidis.* *Pers. encycl. 1. p. 111.*  
*Stipulicida setacea.* *Mich. fl. amer. 1. p. 26.*  
Icon. *Mich. l. c. t. 6.*  
In barren gravelly soil of Lower Carolina. ○. May, June. *v. s. in Herb. Lyon.* The style is so very short, that it scarcely can be called monogynous.

112. LECHEA. *Gen. pl. 142.**major.*

1. *L. undique hirsuta : foliis oblongo-lanceolatis mucronatis, panicula foliosa pyramidata : ramis apice floriferis, floribus fasciculato-racemosis secundis brevissime pedicellatis, caule erecto* — *Willd. sp. pl. 1. p. 495.*  
*L. minor Linnaei.* *Smith in encyclop. ed. nov.*  
On sandy barren soil: Canada to Florida. ♀. July, Aug. *v. v.* I have frequently observed from three to nine stamens in this species, which I suppose is the case with the following one likewise.

2. *L. glabriuscula foliosa, rami brevi-pedicatae*  
*p. 495.*

Icon. *Lam. i.*  
On dry gravelly soil  
Aug. *v. v.*  
than No. 1.

3. *L. undique a ciliatis, paniculis nudi caule erecto*  
In sandy fields

4. *L. undique a panicula fasciculatis la simis, flo*  
*L. thymifolia*  
In barren dry soil  
July. *v. v.*  
species of resembling

5. *L. sparse pilosus nudiusculus ricatis, c*  
On dry gravelly soil  
July. *v. v.*  
linear leaves  
the rest.

1. *E. scapo distico magnum libus acutis*  
*amer. 1. p. 1.*  
*E. serotinum*  
Icon. *Plu.*  
In deep soil  
*v. v.* F.

2. *E. scapo distico bulato-obcordato convexum scariosum*  
*E. decangula*

471.  
♂. June.  
it.  
mbricatio  
ne. v.v.  
rice mino  
e to find,  
s (radica  
sciculatis,  
11.  
  
○. May,  
ery short,  
mucrona  
floriferis,  
ssime pe  
495.  
  
♀. July,  
a three to  
se is the
2. L. glabriuscula ; foliis linear-lanceolatis acutis, panicula foliosa, ramis elongatis undique floriferis, floribus brevi-pedicellatis, caule assurgente.—*Willd. sp. pl. 1.* p. 495.  
Icon. *Lam. illustr. t. 52 f. 1.*  
On dry gravelly hills : Canada to Pennsylvania. ♀. July, Aug. v. v. Lower in growth and larger in fruit than No. 1.
3. L. undique adpresso-pubescent; foliis linearibus acutis ciliatis, panicula gracilis ramosissima pyramidata, racemulis nudiusculis, floribus parvis alternis pedicellatis, caule erecto.—*Mich. fl. amer. 1. p. 77.*  
In sandy fields : New Jersey to Carolina. ♀. July. v. v.
4. L. undique adpresso-cano-villosa ; foliis linearibus acutis, panicula foliosa elongata : ramis brevibus, floribus fasciculatis lateralibus terminalibusque, pedicellis brevissimis, floribus parvis cano-tomentosis, caule erecto.  
L. thymifolia. *Mich. fl. amer 1. p. 77?*  
In barren dry woods, on slate hills : Virginia, &c. ♀. July. v. v. The lower branches, which in most species of this genus trail on the ground, have a great resemblance to *Thymus Serpyllum*.
5. L. sparse pilosa ; foliis angustissimis, panicula divaricata nudiuscula, ramulis alternis, pedicellis elongatis divaricatis, caule erecto.—*Mich. fl. amer. 1. p. 77.*  
On dry gravelly hills : Virginia to Georgia. ♀. May—July. v. v. The lower branches in this species have linear leaves, by which it is easily distinguished from the rest.

113. ERIOCAULON. *Gen. pl. 132.*

1. E. scapo decemstriato, foliis ensiformibus glabris, capitulo magno depresso-globoso, squamis involuci ovalibus acutis, paleis receptaculi mucronatis.—*Mich. fl. amer. 1. p. 165. Willd. sp. pl. 1. p. 485.*  
E. serotinum. *Walt. fl. car. 83. Lam. encycl. 3. p. 276.*  
Icon. *Pluk. amalthe. t. 409. f. 5.*  
In deep swamps : New Jersey to Carolina. ♀. Aug. v. v. From three to four feet high.
2. E. scapo subcompresso decemstriato, foliis brevibus subulato-ensiformibus glabris : capitulo hemisphaerice convexo, involuci squamis ovalibus rotundato-obtusis scariosis argenteo-lucidis. *Mich. fl. amer. 1. p. 165.*  
E. decangulare. *Walt. fl. car. 83.* *gnaphalodes/*

## TRIANDRIA TRIGYNIA. Eriocaulon.

*E. compressum. Lam. encycl. 3. p. 276.*

In sandy mossy swamps: Virginia to Carolina. Ȑ. July, v. v. About a foot high.

*pellucidum.*

3. *E. scapo tenuissime subseptemstriato, foliis subulato-linearibus canaliculatis glabris pellucidis quinque nervis transverse striatis, capitulo parvo subgloboso, squamis involuci ovalibus obtusis.—Mich. fl. amer. 1. p. 166.*

In pools of water, floating: Canada to New Jersey. Ȑ. Aug. v. v. About a foot high; very slender.

*villosum.*

4. *E. scapis aggregatis compressis subquadrisulcis villosa, foliis brevibus subulato-linearibus pilosis, capitulo sphæroideo parvo, flosculis subfuliginosis.—Mich. fl. amer. 1. p. 166.*

*E. enceps. Walt. fl. car. 83.*  
On the edges of ponds, in sandy soil: Virginia to Carolina. July. v. v. v. s. in *Herb. Walter.*

*flavidulum.*

5. *E. scapis aggregatis subseptemstriatis subpubescentibus, foliis brevibus subulato-ensiformibus nervosis, capitulo convexo, squamis involuci suborbiculatis, flosculis vix papilliosis.—Mich. fl. amer. 1. p. 166.*

On the banks of rivers, below high-water mark: Pennsylvania to Carolina. Ȑ. July. v. v. From one to two inches high.

## 114. MOLLUGO. Gen. pl. 139.

*verticillata.*

1. *M. foliis verticillatis cuneiformibus acutis, caule subdviso decumbente, pedunculis unifloris. Willd. sp. pl. 1. p. 492.*

*Icon. Pluk. mant. t. 332. f. 5. Ehret. pict. t. 6. f. 3.*  
In cultivated grounds: Canada to Georgia. Ȑ. May-Aug. v. v.

## 115. PROSERPINACA. Gen. pl. 134.

*palustris.*

1. *P. foliis linear-lanceolatis serratis, infimis pinnatifidis. Lam. illustr. 1. p. 214. Willd. sp. pl. 1. p. 488.*

In ditches and pools: Canada to Carolina. Ȑ. July, v. v.

*pectinata.*

2. *P. foliis omnibus pectinato-pinnatifidis. Lam. illustr. 1. p. 214.*

*P. palustris β. Mich. fl. amer. 1. p. 76.*

In overflowed places and ditches: New Jersey to Carolina. Ȑ. July, Aug. v. v. This is certainly a distinct species, as I never found them growing promiscuously, or near one another.

## 116. EMI

1. *E. procumbens. Mich. Icon. Engl. i. On rocks near May, June species.*

a. 4. July

## IV. POLYGYNIA.

## 116. EMPETRUM. Gen. pl. Crake-berry.

1. E. procumbens; foliis oblongis obtusis margine revolutis. Mich. fl. amer. 2. p. 255. Willd. sp. pl. 4. p. 713.  
Icon. Engl. bot. 526. Fl. dan. 975.

On rocks near the sea-coast in Canada. Michaux. t.  
May, June. v. s. May be distinct from the European  
species.

nigrum.



\*\*\* Flo

128. MITCHE  
4-dentati  
coalitione  
129. OLDENL  
*Caps.* di

\*\*\*\*

130. RUBIA.  
131. GAIJUM  
132. SPERMA  
133. DIODIA.  
locularis

\*\*\*\*

134. HOUSTO  
minifera  
sperma.

135. AMMAM  
*Caps.* 4  
136. PTELEA  
*Stigm.*  
137. CORNU  
2-biloc  
138. LUDWI  
tetrago

139. ALCHE  
inclusu  
140. BOEHM  
tar O.  
141. URTIC  
tar. cy  
tum.  
142. PARIE  
multip  
Polyg  
143. VISCU  
tatum  
144. CEPHA  
Recep

## IV. TETRANDRIA.

### I. MONOGYNIA.

\* *Flores monopetalii, superi, monospermii. Aggregate.*

117. DIPSACUS. *Cal. communis foliaceus. Recept. co-  
nicum, paleaceum. Sem. columnaria.*  
118. ALLIONIA. *Cal. communis 5-fidus, 3-florus : pro-  
prius obsoletus, superus. Corollulae irregulares.  
Recept. nudum.*

\*\* *Flores monopetalii, monocarpi, inferi.*

119. LYCIUM. *Cor. tubulosa, fauce clausa filamentorum  
barba. Cal. 4-fidus. Bac. 2-locularis, polysperma.*  
120. CALICARPA. *Cor. tubulosa. Cal 4-fidus. Bac.  
4-sperma.*  
121. CENTUNCULUS. *Cor. rotata. Cal. 4-fidus. Stam  
brevia. Caps. 1-locularis, circumscissa.*  
122. PLANTAGO. *Cor. limbo reflexo. Cal. 4 fid s.  
Stam. longissima. Caps. 2-locularis, circumscissa.*  
123. POLYPREMUM. *Cor. rotata, fauce barbata. Cal.  
4-partitus. Stam. inclusa. Caps. 2-locularis, obcur-  
data.*  
124. CENTAURELLA. *Cor urceolata. Cal. 4-partitus,  
adpressus. Stigm. crassum, glandulosum, subbifidum.  
Caps. 1-locularis, 2-valvis, calyce corollaque persi-  
tentibus involucrata.*  
125. EXACUM. *Cor. subcampylata. Cal. 4-phyllo.  
Caps. 2-locularis, compressa.*  
126. SWERTIA. *Cor. rotata. Nectariferi pori ad basin  
laciniarum corollæ. Caps. oblonga 1-locularis, 2-  
valvis.*  
127. FRASERA. *Cor. 4-fida, patens : petala medio glan-  
dula barbata. Cal. 4-fidus. Caps. compressa, sub-  
marginata, 1-locularis. Sem. pauca, imbricata, mar-  
ginata.*

## \*\*\* Flores monopetalī, monocarpi, superi.

128. MITCHELLA. *Cor. binæ, infundibuliformes Cal. 4-dentatus. Stigm. 4-fidum. Baccæ ex germinum coalitione didyma, 4-sperma.*  
 129. OLDENLANDIA. *Cor. tubulosa. Cal. 4-partitus. Caps. didyma, polyspermia.*

## \*\*\*\* Flores monopetalī, dicocci, superi.

130. RUBIA. *Cor. campanulata. Baccæ 2., monospermæ.*  
 131. GALLIUM. *Cor. plana. Sem. 2., subrotunda.*  
 132. SPERMACOCE. *Cor. tubulosa. Sem. 2., bidentata.*  
 133. DIODIA. *Cor. infundibuliformis. Caps. 4-gona, 2-locularis, 2 sperma.*

## \*\*\*\*\* Flores monopetalī, dicocci, inferi.

134. HOUSTONIA. *Cor. infundibuliformis. Caps. seminifera, subglobosa, emarginata, 2-locularis, 2-sperma.*

## \*\*\*\*\* Flores tetrapetalī.

135. AMMANNIA. *Cal. inferus, tubulosus, 8-dentatus. Caps. 4-locularis.*  
 136. PTELEA. *Cal. inferus, 4 partitus. Pet. coriacea. Stigm. 2. Samara monosperma.*  
 137. CORNUS. *Cal. superus, 4-dentatus. Drupa nuce 2-biloculari.*  
 138. LUDWIGIA. *Cal. 4-partitus. Caps. 4-locularis, tetragona.*

## \*\*\*\*\* Flores incompleti.

139. ALCHEMILLA. *Cal. inferus, 4-fidus. Sem. 1. calyce inclusum.*  
 140. BOEHMERIA. *Cal. inferus, 4-partitus, ♀. O. Nectar O. Sem. 1. Monoica.*  
 141. URTICA. *Cal. inferus, 4-phyllos, ♂. 2. valvis. Nectar. cyathiforme. Stigm. villosum. Sem. 1. ova-tum. Monoica.*  
 142. PARIETARIA. *Cal. inferus, 4-fidus. Involucrum multipartitum. Sem. 1. calyce sicco elongato tectum. Polygania.*  
 143. VISCIUM. *Cal. superus, 4-partitus. Stigm. capitatum. Baccæ monosperma. Dioicum.*  
 144. CEPHALANTHUS. *Cal. superus, infundibuliformis. Recept. globosum, nudum. Sem. 1. lanuginosum.*

## TETRANDRIA.

145. ELÆAGRUS. *Cal.* basi persistens, superne coarctatus et quasi superus, campanulatus, 4-fidus. *Drupa* 1-sperma.  
 146. HIPPOPHAE. *Cal.* deciduus, inferus, tubulosus, 2-fidus. *Bacca* globosa. *Divica.*

## II. DIGYNIA.

147. PLANERA. *Cal.* campanulatus, 4-fidus. *Cor.* 0. *Nux* 1-sperma, squamulosa. *Polygamia.*  
 148. HAMAMELIS. *Cal.* 4-fidus, extus 2-squamosus. *Pet.* 4. longissima. *Nux* 2-locularis, 2-cornis.  
 149. CUSCUTA. *Cal.* 4-fidus. *Cor.* ovata, 4-fida. *Caps.* 2-locularis circumscissa.  
 150. SANGUISORBA. *Cal.* 2-phylloides, inferus. *Cor.* plana, 4-fida, supera. *Caps.* 4-gona, intra calycem et corollam.

## III. TRIGYNIA.

151. PACHYSANDRA. *Cal.* 4-phylloides. *Cor.* 0. *Caps.* 3-locularis. *Monoica.*

## IV. TETRAGYNIA.

152. ILEX. *Cal.* 4-dentatus. *Cor.* rotata. *Styl.* 0. *Bac.* 4-sperma.  
 153. SAGINA. *Cal.* 4-phylloides. *Cor.* 4-petala. *Caps.* 4-locularis polysperma.  
 154. POTAMOGETON. *Cal.* 4-phylloides. *Cor.* 0. *Sem.* 4. sessilia.  
 155. RUPPIA. *Cal.* et *Cor.* 0. *Sem.* 4. pedicellata.

## I. MONOGYNIA.

## 117. DIPSACUS. Gen. pl. 148.

*sylvestris.*

1. D. foliis connatis sinuatis, paleis rectis, involucro capitulo brevioribus patentibus.—*Willd.* sp. pl. 1. p. 544?  
*Icon.* *Jacq. fl. aust.* 5. t. 403.  
 On the edges of fields and on water sides: Pennsylvania to Virginia. ♂. July. v. v. Probably introduced from Europe, but sometimes found in very remote parts from cultivation.

1. A. caule erectum dunculis involucro amer. 1. f. On the bank 4. July, very muc  
 2. A. caule digitatum pedunculis ovatis In Lower C Walter.  
 3. A. caule nudis, floribus volucro & usculis r. On the pla v. s. in 1

1. L. inerme fl. amer. L. salsum. In salt marsh Flowers 120  
 1. C. foliis ovoidibus.—Icon. C. In dry grass Carolinian small a whorl 1

1. C. foliis c perior simplic In wet t

VOL. I.

## 118. ALLIONIA. Gen. pl. 151.

1. A. caule erecto, foliis lato-cordatis acutis, glabris, pedunculis solitariis, calycibus fructiferis sub-hirsutis, involucro ampliato multo minoribus.—*Mich. fl. amer. 1. p. 100.*  
On the banks of the river Tennessee and its branches.
2. A. caule diffuso, foliis lanceolatis scabriusculis ciliatis, pedunculis solitariis, involucro subpentaphyllo : foliolis ovatis reticulato-venosis.—*Walt. fl. car. 84.*  
In Lower Carolina. *Walter.* ♂. July. v. s. in *Herb. Walter.*
3. A. caule diffuso dichotomo, foliis ovatis scabriusculis nudis, floribus terminalibus fasciculato-racemosis, involucro semiquinquefido : laciniis subrotundis acutiusculis reticulato-venosis.  
On the plains of the Missouri. *M. Lewis.* ♂. Aug. v. s. in *Herb. Lewis.*

## 119. LYCIUM. Gen. pl. 343.

1. L. inerme ; foliis anguste spathulato-oblongis. *Mich. fl. amer. 1. p. 95. Walt. fl. car. 84.* *carolinianum.*  
*L. salsum. Bartr. trav. 59.*  
In salt marshes of Carolina and Georgia. ♀. July. v. v.  
Flowers purple, the size of *L. barbarum.*

## 120. CALICARPA. Gen. pl. 175.

1. C. foliis ovalibus serratis subtus tomentosis, cymis sessilisibus.—*Willd. sp. pl. 1. p. 619.*  
*Icon. Catesb. car. 2. t. 47. Pluk. alm. t. 136. f. 3.*  
In dry gravelly soil, below the mountains in Virginia and Carolina. ♀. June, July. v. v. Flowers red, in small axillary bunches ; berries red, apparently in a whorl round the branches, hang on very long.

## 121. CENTUNCULUS. Gen. pl. 189.

1. C. foliis deorsum angustatis : inferioribus ovalibus, superioribus lanceolatis, laciniis calycis subulatis, caule simplici.—*Mich. fl. amer. 1. p. 193.*  
In wet barren fields: Lower Carolina, &c. ♂. May.

## TETRANDRIA MONOGYNA. Centunculus.

*v. s. in Herb. Nuttall.* A very small inconspicuous plant with axillary reddish flowers. The specimens of Mr. Nuttall's were collected on the Missouri, they are about four inches high: and according to his observations, very obligingly communicated to me, this species is generally pentandrous.

122. PLANTAGO. *Gen. pl.* 186.*data.*

1. *P. foliis ovatis cordatis latissimis subdentatis glabris, spica longissima, floribus subimbricatis, inferioribus sparsis, bracteis ovatis obtusis.* — *Lam. encycl.* 5. p. 369.

*P. kentuckensis. Mich. fl. amer.* 1. p. 94.*P. canadensis. Hort. par.*

On river sides: Canada, Kentucky, Tennessee, and other western parts.  $\frac{1}{4}$ . June—Aug. v. v.

*major.*

2. *P. foliis ovatis subdentatis glabriusculis, scapo tereti, spica oblonga imbricata.* *Lam. encycl.* 5. p. 368. *Willd. sp. pl.* 1. p. 641.

*Icon. Engl. bot.* 1558. *Fl. dan.* 461.

In fields and on road sides, common.  $\frac{1}{4}$ . May—Sept. v. v.

*cucullata.*

3. *P. foliis ovatis concavo-cucullatis novemnerviis subtus pubescentibus, spica cylindrica imbricata, scapo tereti alto.* *Lam. encycl.* 5. p. 370.

*P. maxima. Jacq. ic.* 1. t. 26. *Willd. sp. pl.* 1. p. 642.*Icon. Jacq. l. c.*

In wet rocky situations: Canada and Province of Main.  $\frac{1}{4}$ . July, Aug. v. v.

*lanceolata.*

4. *P. foliis lanceolatis villosis, spica brevi cylindrica, bracteis ovatis acuminatis, scapo angulato adpresso piloso.* *P. lanceolata. Willd. sp. pl.* 1. p. 643?

In fields and pastures, common.  $\frac{1}{4}$ . June—Aug. v. v.  
May be a new species. *P. lanceolata* of Europe, according to the specimens examined in the Herbarium of A. B. Lambert, Esq. is *bracteis oblongis acutis*.

*virginica.*

5. *P. undique canescenti-pubescentis; foliis lanceolato-ovatis subdenticulatis, floribus remotis, scapo angulato.* *Willd. sp. pl.* 1. p. 643.

*Icon. Moris. hist.* 3. s. 8. t. 15. f. 8.

In dry barren fields, common: New England to Florida.  $\odot$ . May—Aug. v. v. There are a number of varieties in respect to size and pubescence.

*caroliniana.*

6. *P. undique glabra; foliis lanceolatis integerrimis longis, floribus remotis, caule tereti.* — *Walt. fl. car.* 85.

T  
P. angusti  
In sandy g  
July.

7. *P. foliis o  
longa  
cycl. 5.*

P. sparsifl  
In shady w  
v. v.

8. *P. foliis li  
po tere  
bracteai*

On the  
 $\frac{1}{4}$ . Aug  
Esq. ar  
respect

9. *P. undiqu  
tegerrim  
longior  
ribus l*

In dry sit  
 $\odot$ . Ju  
plant i

10. *P. foliis  
tereti p  
Willd.*

Icon. *I*  
On the b  
York.

11. *P. foliis  
bracte  
amer.*

In the na

12. *P. polyp  
Icon. L*

In dry p

v. v.

124. C

1. *C. caule  
gatis,*

- P. angustifolia glabra, &c. *Clayt. fl. virg.* 753.  
In sandy grassy woods: Virginia to Carolina. ♂. June,  
July. v.v.; v. s. in *Herb. Walter*.
7. P. foliis ovato-lanceolatis integerrimis villosiusculis, spica  
longa gracili interrupta, floribus glabris.—*Lam. en-*  
*cycl. 5. p. 375.*
- P. sparsiflora. *Mich. fl. amer. 1. p. 94.*  
In shady woods: Virginia to Carolina. ♀. July, Aug.  
v. v.
8. P. foliis linear-lanceolatis integerrimis glabriusculis, sca-  
po tereti foliis breviore, spica pauciflora interrupta,  
bracteis ovatis acutis glabris.  
On the sea-coast of New England and New Jersey.  
♀. Aug. v.v. In the Herbarium of A. B. Lambert,  
Esq. are specimens from Labrador, agreeing in every  
respect with this species.
9. P. undique argenteo-villosa; foliis linear-lanceolatis in-  
tegerrimis supra nudiusculis, scapis teretibus foliis vix  
longioribus, spica cylindrica imbricata, bracteis linea-  
ribus longissime ciliato-villosis.  
In dry situations on the banks of the Missouri. *Nuttall.*  
○. July. v. s. in *Herb. Nuttall*. This beautiful little  
plant is very variable in size.
10. P. foliis carnosis subulato-linearibus basi pilosis, scapo  
tereti pubescente, spica cylindrica, bracteis obtusis.—  
*Willd. sp. pl. 1. p. 647.*  
Icon. *Engl. bot. 175. Fl. dan.* 243.  
On the banks of rivers near the sea: Canada to New  
York. ♀. Aug. v. v.
11. P. foliis subsetaceo-linearibus, spica oblongo-cylindrica,  
bracteis subulato-aristatis flore longioribus. *Mich. fl.*  
*amer. 1. p. 95.*  
In the natural meadows of Illinois. *Michaux.* +.

123. POLYPREMUM. *Gen. pl. 178.*

1. P. polypremum. *Willd. sp. pl. 1. p. 623.* *procumbens.*  
Icon. *Lam. illustr. 1. t. 71.*  
In dry pastures: Virginia to Georgia. ○. May—Aug.  
v. v. Flowers white, very small.

124. CENTAURELLA. *Mich. fl. amer. 1. p. 97.*

1. C. caule simplici, pedunculis sparsis subcorymboso-elon-  
gatis, lacinii corollæ oblongis acutis calyce duplo

## TETRANDRIA MONOGYNIA. Centaurella.

longioribus, stylo germine longiore. *Mich. fl. amer.* 1. p. 98.

*uniflora.*

Icon. *Mich. l. c. t. 12. f. 2.*

*aestivalis.*

$\beta$ . C. pusilla, caule unifloro.

In mossy swamps: Virginia to Georgia.  $\odot$ . May—July. v. v. Var.  $\beta$  v. s. in *Herb. Lyon.* About five inches high; the flowers white, and larger than the rest. The variety  $\beta$  may be a distinct species?

*autumnalis.*

2. C. caule simplici, pedunculis oppositis simplicibus, laci- niis corollæ spathulatis abrupte acutis calyce triplo longioribus, stylo longitudine germinis.

In mossy swamps: Carolina to Georgia.  $\odot$ . June—Aug. v. s. in *Herb. Enslen.* From two to four inches high; flowers smaller than No. 1; the calyx very short.

3. C. caule subramoso, pedunculis oppositis, inferioribus ramosis, laciis corollæ ovalibus acutis calyce vix lon- gioribus, stylo brevissimo.

C. paniculata. *Mich. fl. amer.* 1. p. 98.

Bartonia tenella. *Muhl. in litt.*

Sagina virginica. *Willd. sp. pl.* 1. p. 719.

Icon. *Mich. l. c. t. 12. f. 1.*

In mossy swamps, in inundated grassy places: New England to Carolina, common.  $\odot$ . July—Sept. v. v.; v. s. in *Herb. Gronov. sub nomine Sagina, caule erecto subnudo, floribus oppositis. Saginæ affinis planta mi- nima, floribus albis. Clayt. MSS.* 649. About a foot high, sometimes quite simple and very small; flowers small, greenish-white.

125. EXACUM. *Gen. pl.* 185.

*pulchellum.*

1. E. corollis quadrifidis, calycibus quadripartitis: laciis subulatis, panicula corymbosa, pedunculis filiformibus. Near the sea-coast of New Jersey; rare.  $\odot$ . Aug. v. v. Flowers rose-coloured; resembling *E. filiforme* very much.

126. SWERTIA. *Gen. pl.* 449.

*corniculata.*

1. S. corollis campanulatis corniculatis: cornibus deflexis, foliis ovatis, ramis brevibus.

S. corniculata. *Willd. sp. pl.* 1. p. 1330?

In swamps of Canada and round the lakes of the western part of the United States.  $\delta$ . July. v. v. Flowers greenish yellow. The Siberian plant differs from the

America  
bus, foli  
America

2. S. corollis  
bus fast  
pedicelli  
caule ra

On the M  
Lewis.  
large;

3. S. corollis  
cissimo

On the al  
shire.  
above a  
leaves,  
of the  
In the  
dor, in  
shire p

1. Frasera.  
Frasera c

In swan  
the la  
v. v.  
whorl  
green  
whole  
gener  
withc  
cies c

1. Mitche  
Icon. (

In shad  
Canad  
ing |

- American in being *cornibus divaricatis adscendentibus, foliis lanceolatis et ramis elongatis*; by which the American plant appears to be a distinct species.
2. *S. corollis campanulato-rotatis longitudine calycis, floribus fastigiato-aggregatis axillaribus terminalibusque, pedicellis geminis, foliis spathulato-obovatis nervosis, caule ramoso.* On the Missouri flats, near the Rocky mountain. *M. Lewis.* 4. July. *v. s. in Herb. Lewis.* Leaves large; flowers sky-blue.
3. *S. corollis rotatis calyce duplo longioribus, caule simplissimo unifloro, foliis oblongis.* On the alpine regions of the White-hills of New Hampshire. ① June. *v. v.* The whole plant scarcely above an inch high, with one or two pair of small leaves, and a considerable-sized blue flower. Divisions of the corolla oblong, acuminate; of the calyx obtuse. In the Banksian Museum are specimens from Labrador, in every respect agreeing with the New Hampshire plant.

127. FRASERA. *Mich. fl. amer. 1. p. 96.*

1. *Frasera. Mich. l. c. p. 97.*  
*Frasera carolinensis. Walt. fl. car. 88.*  
 In swamps of Lower Carolina and on the borders of the lakes in Pennsylvania and New York. ♂. July. *v. v.* Stem from three to six feet high; leaves in whorls or opposite; flowers in aggregate clusters, greenish-yellow, sometimes mixed with purple. The whole of the plant has a very stately appearance; its generic character approaches so near to *Swertia*, that, without seeing the fruit, it might be considered a species of that genus.

*Walteri.*128. MITCHELLA. *Gen. pl. 174.*

1. *Mitchella. Willd. sp. pl. 1. p. 617.*  
 Icon. *Catesb. car. 1. t. 20. Pluk. amalh. t. 444. f. 2.*  
 In shady situations on rocks and round the roots of trees: Canada to Georgia. ♀. June. *v. v.* A small creeping plant; flowers white, hairy inside; berries red.

*repens.*

their feath  
of a beauti

4. G. caulinis  
foliis caulinis  
natis marginatis  
divaricatis.—*Mich. fl. amer.* 1. p. 83.  
In swamps a  
July.

5. G. caulinis  
foliis quinis,  
nervoque  
pedicellis  
*sp. pl.* 1.  
Aparine flor  
*G. Claytoni.*  
In swamps  
July. v.

6. G. caulinis  
lineari-lanceolata  
carinaque  
liorum  
*Willd. sp.*

Icon. *Fl. de*  
In hedges a  
June—A

7. G. caulinis  
foliis brevibus  
carinaque  
losis, petiolaris  
In mountai  
v. v. F  
small.  
bles G.

8. G. caulinis  
dis: rauca  
acuminate  
ramis flexu  
tomis:  
In meadow  
England  
white.

129. OLDENLANDIA. *Gen. pl. 205.**glomerata.*

1. O. procumbens; foliis lanceolato-ovatis, floribus axillaris terminalibusque conglobatis pedicellatis, germinibus hispidis. *Mich. fl. amer.* 1. p. 83.  
O. uniflora. *Willd. sp. pl.* 1. p. 675.  
*Hedyotis auricularia.* *Walt. fl. car.* 85.  
*Hedyotis uniflora.* *Lam. illustr.* 1. p. 271.  
On sandy wet fields: New Jersey to Carolina. ○. May—July. v. v. A small creeping plant; flowers greenish-white.

130. RUBIA. *Gen. pl. 164.* Madder.*Brownii.*

1. R. hispida; foliis quaternis ovalibus, pedunculis solitariis unifloris, caule decumbente. *Mich. fl. amer.* 1. p. 81.  
R. subhirsuta scandens vel reclinata, foliis cruciatis, floribus singularibus ad alas. *Brown. jam.* p. 141.  
R. peregrina. *Walt. fl. car.* 86.  
*Valantia hypocarpa.* *Linn.*  
In shady woods: Carolina and Florida. ¼. +. Flowers yellow; berries purple, smooth.

131. GALIUM. *Gen. pl. 162.*\* *Fructu glabro.**latifolium.*

1. G. caulinis erectis laevibus, foliis quaternis ovatis acutis trinervibus: margine nervisque hispidis, pedunculis oppositis lateralibus terminalibusque dichotomis divaricatis laxe multifloris.—*Mich. fl. amer.* 1. p. 79.  
In dry slaty mountains of Virginia and Carolina. ¼. June, July. v. v. Leaves broad; flowers purple. It generally has but one seed.

*uniflorum.*

2. G. caulinis flaccidis laevibus, foliis quaternis linearibus acutis glabris, pedunculis axillaris solitariis unifloris brevissimis.—*Mich. fl. amer.* 1. p. 79.  
In Carolina. *Michaux.* +. Flowers white.

*tinctum.*

3. G. caulinis diffusis laeviusculis, foliis caulinis senis, ramulis quaternis linearibus acutiusculis, margine nervo tenuissime aculeatis, pedunculis terminalibus elongatis plerumque trifloris.—*Willd. sp. pl.* 1. p. 586.  
In wet woods and along hedges: Canada to Virginia. ¼. June—Aug. v. v. Flowers white. This is the plant with which the Indians of North America dye

their feathers, porcupine quills, and other ornaments, of a beautiful red.

4. G. caulinis diffusis ramosissimis retrorsum aculeatis, foliis caulinis senis, rameis quaternis lanceolatis acuminateis margine nervo que aculeatis, ramulis floriferis divaricatis multifloris, pedicellis brevibus.—*Mich. fl. amer. 1. p. 78.*

In swamps and among low bushes: Canada to Virginia.  
4. July. v. v. Flowers white, very small.

5. G. caulinis procumbentibus retro-scabris, foliis caulinis quinis, rameis quaternis linearibus obtusis margine nervo que scabris, fasciculis terminalibus paucifloris, pedicellis brevibus, corollis plerumque trifidis.—*Willd. sp. pl. 1. p. 585.*

*Aparine* floribus albis, &c. *Gron. fl. virg. 18.*

*G. Claytoni. Mich. fl. amer. 1. p. 78.*

In swamps and wet fields: Canada to Carolina.  
July. v. v. Flowers white, very small.

\*\* *Fructu hirsuto.*

6. G. caulinis flaccidis retrorsum scabris, foliis subocononis linear-lanceolatis mucronatis supra hispidis, margine carinaque aculeatis, ramulis floriferis longitudine foliorum subtrifloris, fructibus uncinato-hispidis.—*Willd. sp. pl. 1. p. 597.*

*Icon. Fl. dan. 495. Engl. bot. 816.*

In hedges and small woods: Pennsylvania, &c. ♂.  
June—Aug. \* v. v. Flowers small, white.

7. G. caulinis ramosissimis divaricatis retrorsum aculeatis, foliis brevibus lanceolatis mucronatis glabris, margine carinaque aculeatis, ramulis floriferis divaricato-ramulosis, pedicellis subbifloris, fructibus hispidis.

In mountain swamps: Canada to New York.  
4. July. v. v. Flowers white, very numerous and exceeding small. It produces generally only one seed. Resembles *G. uliginosum* very much.

8. G. caulinis flaccidis elongatis brachiato-ramosis hispidis: ramis brevibus, foliis senis oblongo-lanceolatis acuminatis glabris, margine carinaque setaceo-ciliatis, ramis floriferis verticillis longioribus divaricato-dichotomis: pedicellis bifloris, fructibus uncinato-hispidis.

In meadows and on the edges of woods, frequent: New England to Virginia.  
4. June—Aug. v. v. Flowers white.

*asprellum.*

*trifidum.*

*Aparine.*

*brachiatum.*

*triflorum.*

9. *G. caulis procumbentibus glabriusculis, foliis senis quinisque obovato-lanceolatis mucronatis glabris margine vix ciliatis, ramulis floriferis elongatis apice trifloris, floribus pedicellatis, fructibus parvis hispidis.*  
*Mich. fl. amer. 1. p. 80.*

Icon. *Wendl. hort. t. 66.*In shady rocky places: Pennsylvania to Carolina.  $\frac{1}{4}$ . July. v. v. Flowers white, very small.*hispidum.*

10. *G. caulis procumbentibus ramosissimis hispidis, foliis quaternis ovali-lanceolatis acuminatis margine revolutis rugosis utrinque hispidis, floribus subsolitarie terminalibus, fructibus hispidis.* — *Mich. fl. amer. 1. p. 79.*  
*G. lappaceum. Fl. per. 1. p. 59?*

In Lower Carolina. *Michaux.* +. Flowers white.*bermudianum.*

11. *G. caulis ramosissimis vix pubescens, foliis quaternis ovatis obtusis glabris, margine nervisque pubescens, pellucido-punctatis, ramulis floriferis elongatis, fructibus uncinato-hispidis.* — *Willd. sp. pl. 1. p. 596.*  
*G. purpureum. Walt. fl. car. 87.*

*G. puncticulatum. Mich. fl. amer. 1. p. 80.*In wet places: Virginia to Carolina.  $\frac{1}{4}$ . July. v. v. Flowers purple.*pilosum.*

12. *G. caulis subsimplicibus elongato-adscendentibus remote-geniculatis hispidis, foliis quaternis ovalibus brevissime mucronatis utrinque hirsutissimis enervibus, ramulis floriferis elongatis subsimplicibus apice subtrifloris, fructibus pilosis.* *Willd. sp. pl. 1. p. 599.*  
In dry woods: Pennsylvania to Carolina.  $\frac{1}{4}$ . June, July. v. v. Flowers brownish purple.

*circæzans.*

13. *G. caulis erectis laevibus, foliis quaternis ovalibus obtusis glabris, margine nervisque ciliolatis, pedunculis divaricatis paucifloris, floribus remotis subsessilibus, fructibus cernuis uncinato-hispidis.* *Mich. fl. amer. 1. p. 80.*  
*G. boreale. Walt. fl. car. 87.*

In dry barren woods and mountains: New England to Carolina.  $\frac{1}{4}$ . June—Aug. v. v. Flowers yellowish-white; stem not above six inches high.*boreale.*

14. *G. caulis erectis laevigatis, foliis quaternis linearibus acutis trinervibus glabris, margine nervoque scabris, panicula terminali divaricata, fructibus tenuissime muricato-setosis.* — *Willd. sp. pl. 1. p. 595.*  
Icon. *Fl. dan. 1024? Engl. bot. 105?*

On the banks of the Susquehanna near Wilkesbarre, among rocks: Pennsylvania and about the lakes of Ca-

nada and  
very num  
I doubt  
found so  
tion of  
Russia, t  
tion betw

1. *S. caule er  
tus glabi  
tis.* — *W*  
Icon. *Sc*  
*f. 1.*

In dry grav  
—Aug.

2. *S. caule  
utrinque  
bris.* — *J*  
On the ba  
*○. Jul*  
No. 1.

3. *S. caule  
glabrius  
pulis lo  
bus sol  
amer. 1*  
In dry ba  
lina.  
small.

4. *S. caule  
latis a  
capitul*  
In Caro  
a foot  
flower

1. *D. glab  
lance  
longis*  
*D. teres*  
Icon. *J*

nada and New York. ♀. July. v. v. Flowers white, very numerous, collected in a regular panicle. Though I doubt of its being the *G. horeale* of Linnaeus, I found so many intermediate specimens in the collection of A. B. Lambert, Esq., collected by Pallas in Russia, that I could draw no line of essential distinction between them.

132. SPERMACOCE. *Gen. pl. 155.*

1. S. caule erecto glabro, foliis lanceolatis supra scabris subtus glabriuseulis, floribus verticillatis, seminibus hirtis.—*Willd. sp. pl. 1. p. 568.*

*tenuior.*

Icon. *Schkuhr handb. 1. t. 22. Lam. illustr. 1. t. 62. f. 1.*

In dry gravelly situations: Virginia to Carolina. ☽. June—Aug. v. v. Flowers white, very small.

2. S. caule procumbente glabro, foliis ovato-lanceolatis utrinque glabris, floribus verticillatis, seminibus glabris.—*Mich. fl. amer. 1. p. 82.*

*glabra.*

On the banks of rivers in most of the western territories. ☽. July, Aug. v. v. Flowers white, longer than No. 1.

3. S. caule diffuso tereti hirsuto, foliis linear-lanceolatis glabriusculis, margine carinaque serrulato-scabris, stipulis longissime multisetis, floribus axillaribus sessilibus solitariis alternis, seminibus hirsutis.—*Mich. fl. amer. 1. p. 82.*

*diodina.*

In dry barren soil, on iron ore hills: Virginia to Carolina. ☽. July, Aug. v. v. Flowers white, very small.

4. S. caule alterne ramoso hispidissimo, foliis ovato-lanceolatis acuminatis utrinque hirsutis, stipulis multisetis, capitulis terminalibus involucratis, staminibus exertis.

*involucrata.*

In Carolina. *Fraser. v. s. in Herb. Lambert.* About a foot high; leaves broad and somewhat oblique; flowers white, with a very long tube.

133. DIODIA. *Gen. pl. 158.*

1. D. glabra; caulibus procumbentibus teretiusculis, foliis lanceolatis, corollis intus glabriusculis, fructibus oblongis glabris.—*Willd. sp. pl. 1. p. 580.*

*virginica.*

D. teres. *Walt. fl. car. 87.*

Icon. *Jacq. ic. rar. 1. t. 29. bona.*

In sandy wet places, on road sides : Virginia to Carolina.  
*v. v.* Stem smooth, purple; flowers white.

*hirsuta.*

2. *D. undique hirsutissima*; caulis procumbentibus teragonis, foliis linearis-ob lanceolatis, corollis intus dense-barbatis, fructibus ovatis hirsutis.

In sandy fields : Georgia near Savannah. *v. v.* Stems very branching; flowers white, larger than the former species.

*cærulea.*

1. *H. caule erecto setaceo dichotomo, foliis radicalibus spatulatis, caulinis oblanceolatis oppositis, pedunculis unifloris elongatis.* — *Willd. sp. pl. 1. p. 583.*

*Poiretia erecta. Gmel. syst. 263. Anonymos. Walt. fl. car. 86.*

*Icon. Bot. mag. 370. Pluk. alm. t. 97. f. 9.*

*a. H. caule pedunculisque erectis, floribus magnis cœruleis.*

*β. H. caule divaricato, floribus plerumque albis.*

From New England to Florida: *α* in rocky situations;  
*β* along the sandy sea-coast. *v. v.* Flowers a beautiful blue, sometimes white.

*serpyllifolia.*

2. *H. caule procumbente cæspitoso, foliis spathulatis obtusis, pedunculis terminalibus unifloris longis.* — *Mich. fl. amer. 1. p. 85.*

In high mountains, near rivulets and springs: Virginia to Carolina. *v. v.* Flowers blue.

*tenella.*

3. *H. caule repente filiformi, foliis orbiculatis acutis nervosis, pedunculis terminalibus unifloris longissimis.*

On high mountains: North Carolina. *Lyon. 24. June. v. s. in Herb. Lyon.* Flowers purple; smaller than the second species, which it resembles very much.

*rotundifolia.*

4. *H. caule repente, foliis suborbiculatis abrupte petiolatis crassi sculis, pedunculis axillaribus solitariis unifloris.* — *Mich. fl. amer. 1. p. 85.*

*Poiretia procumbens. Gmel. syst. 263. Anonymos. Walt. fl. car. 86.*

Along the sea-coast: Virginia to Florida. *v. v.* Flowers white; leaves evergreen.

*angustifolia.*

5. *H. caule erecto ramosissimo glabro, foliis linearibus, floribus terminalibus fasciculatis subsessilibus, saepè ternis.* — *Mich. fl. amer. 1. p. 85.*

TETRAN

*H. longifolia.*

Along the sea-  
da *v. v.* Ma

6. *H. caule erecto  
foliis sessilibus  
fasciculis te  
p. 584.*

*H. varians. M  
Hedyotis umb  
Knoxia purpu  
In dry woods  
*v. v.* Flow*

135.

1. *A. caule erec  
semi-ample  
— Willd. s*

In swamps an  
the western  
ers purple.

2. *A. caule pro  
gustatis, fl  
fl. amer. 1*

*A. ramosior.*  
In gravelly w  
lina. ◎.

136.

1. *P. foliis tern  
Icon. Schmi*

*a. P. foliis qui*

*b. P. foliis pub*

In shady mo  
rocks: N  
small tre  
Flowers s  
The varie  
a new spe

137.

1. *C. herbacea  
cris ovata  
pl. 1. p.*

- olina,  
flow-  
is te-  
intus  
v. v.  
in the  
is spa-  
nculus  
ult. fl.  
co-  
ions :  
v. v.  
s ob-  
Mich.  
rginia  
ervous-  
June.  
than  
sh.  
foliatis  
floris.  
Walt.  
rch—  
ribus,  
sæpe
- H. longifolia. Willd. sp. pl. 1. p. 583 ?  
Along the sea-coast, in sandy fields: Carolina to Florida.  $\frac{1}{2}$ . May—July. v. v. Flowers purple.
6. H. caule erecto superne ramoso ad geniculis pubescente, foliis sessilibus ovatis lanceolatis vel linear-lanceolatis, fasciculis terminalibus corymbosis. Willd. sp. pl. 1. p. 584.
- H. varians. Mich. fl. amer. 1. p. 86.  
Hedyotis umbellata. Walt. fl. car. 85.  
Knoxia purpurea. Lam. illustr. 1. p. 259.  
In dry woods: Pennsylvania to Carolina.  $\frac{1}{2}$ . June, July.  
v. v. Flowers purple.

## 135. AMMANNIA. Gen. pl. 206.

1. A. caule erecto, foliis lanceolato-linearibus basi dilatata semi-amplexicaulibus, floribus inferioribus verticillatis.—Willd. sp. pl. 1. p. 678.
- In swamps and inundated fields: Virginia, Carolina, and the western territories.  $\odot$ . July, Aug. v. v. Flowers purple.
2. A. caule procumbente, foliis lanceolatis in petiolum angustatis, floribus solitariis oppositis sessilibus.—Mich. fl. amer. 1. p. 99.
- A. ramosior. Walt. fl. car. 88.  
In gravelly wet places, along rivers: Virginia to Carolina.  $\odot$ . Aug. Sept. v. v. Flowers red.

*purpurea.**ramosior.**humilis.**trifoliata.**pentaphylla.*  
*pubescens.*

In shady moist hedges and on the edges of woods among rocks: New York to Carolina.  $\frac{1}{2}$ . June. v. v. A small tree or shrub from ten to fifteen feet high. Flowers small; greenish white, in corymbous clusters. The variety  $\beta$  grows in Pennsylvania, and appears to be a new species.

## 136. PTELEA. Gen. pl. 1505.

1. P. foliis ternatis. Willd. sp. pl. 1. p. 670.  
Icon. Schmidt arb. 2. t. 76. Mill. ic. 2. t. 211.
- a. P. foliis quinatis. Du Roi.
- $\beta$ . P. foliis pubescentibus.
- \* Involucratæ capitatæ.
1. C. herbacea; foliis summis verticillatis venosis, involucris ovatis acuminatis, drupis globosis. Willd. sp. pl. 1. p. 661.

*canadensis.*

- Icon. *L'Her. corn. t. 1.* *Bot. mag.* 880.  
In shady woods of Canada, and on high mountains in boggy ground, from New England to Carolina.  $\frac{1}{2}$ .  $\frac{1}{2}$ . May. *v. v.* About six inches high; flowers purplish-white; involucra white; berries red, ripe in July.
- florida.* 2. *C. arborea*; foliis ovatis acuminatis, involucris magnis quasi obcordatis, drupis ovatis.—*Willd. sp. pl.* 1. p. 661.  
Icon. *Catesb. car. t. 27.* *Schmidt arb. t. 62.* *Bot. mag.* 526. *Dog-wood.*  
Frequent in woods from Canada to Carolina.  $\frac{1}{2}$ . May. *v. v.* A beautiful small tree; flowers greenish-yellow; involucrum white, tinged with red; berries scarlet, ripe in August. The wood is exceedingly hard, and of a very fine texture.
- \*\* *Nudæ cymosæ.*
- circinata.* 3. *C. ramis verrucosis*, foliis lato-ovalibus acuminatis subtus albo-tomentosis, cymis patentissimis.—*L'Herit. corn.* p. 7.  
*C. tomentulosa.* *Mich. fl. amer.* 1. p. 91.  
*C. rugosa.* *Lam. encycl.* 2. p. 115.  
*C. virginiana.* *Hort. par.*  
Icon. *L'Herit. l. c. t. 3.* *Schmidt arb. 2. t. 69.*  
On the banks of rivers: Canada to Virginia.  $\frac{1}{2}$ . June, July. *v. v.* A shrub from six to ten feet high; leaves broad, waved on their edges; flowers white, as in all the following species; berries blue.
- sericea.* 4. *C. ramis patulis*: ramulis lanuginosis, foliis ovatis acuminatis subtus ferrugineo-pubescentibus, cymis depressis lanuginosis.—*Willd. sp. pl.* 1. p. 663.  
*C. lanuginosa.* *Mich. fl. amer.* 1. p. 92.  
*C. alba.* *Walt. fl. car.* 88.  
*C. cærulea.* *Lam. encycl.* 2. p. 116.  
*C. amomum.* *Du Roi harbk.* 1. p. 165.  
*C. rubiginosa.* *Ehrh. beitr.* 4. p. 15.  
*C. ferruginea.* *Hort. par.*  
Icon. *L'Herit. corn. t. 2.* *Schmidt arb. 2. t. 64.*  
In swampy woods and on river banks: Canada to Carolina.  $\frac{1}{2}$ . June, July. *v. v.* From five to eight feet high; berries bright blue.
- asperifolia.* 5. *C. ramis erectis*: ramulis cymisque pubescentibus, foliis brevi-petiolatis ovalibus acuminatis supra asperis subtus subtomentosis.—*Mich. fl. amer.* 1. p. 93.  
In shady woods of Lower Carolina. *Michaux.*  $\frac{1}{2}$ .

- June, July.  
former species
6. *C. ramis strictis* bris subconcretae.—*L'Herit. corn.*  
*C. fastigiata.* *M*  
*C. sanguinea.* *C*  
*C. canadensis.* *C*  
Icon. *L'Herit.*  
On the banks of rivers.  $\frac{1}{2}$ . June. *v. v.* blue: anthers yellow
7. *C. ramis strictis* brescentibus, *C*  
Icon. *Schmidt*  
Near the lake. July. *v. v.* yellow; berries yellow
8. *C. ramis recurvata* pubescentibus.—*corn.* p. 6.  
*C. stolonifera.* *C*  
*C. tartarica.* *I*  
Icon. *Schmid*  
On the banks of rivers.  $\frac{1}{2}$ . May. branches red
9. *C. ramis erectis* canis, cymis *C*  
*C. racemosa.* *C*  
*C. citrifolia.* *C*  
*C. foliis ellipticis* *C*  
Icon. *L'Herit.*  
In swamps and on the banks of rivers: Canada to Carolina.  $\frac{1}{2}$ . four to six feet high.
10. *C. ramis verticillata* canis, cymis *C*  
*Willd. sp.*  
Icon. *L'Herit.*  
In shady woods.  $\frac{1}{2}$ . May. twenty feet high.

- June, July. +. Probably only a variety of the former species.
6. *C. ramis strictis fastigiatis, foliis ovatis acuminatis glabris subconcoloribus, cymis convexis subpaniculatis.* *L'Herit. corn. p. 8.* *stricta.*
7. *C. fastigiata. Mich. fl. amer. 1. p. 92.*  
*C. sanguinea. Walt. fl. car. 88.*  
*C. canadensis. Hort. par.*  
 Icon. *L'Herit. corn. t. 4. Schmidt arb. 2. t. 67.*  
 On the banks of rivers: Canada to Carolina, frequent.  
 h. June. v. v. From six to ten feet high; berries blue; anthers blue.
7. *C. ramis strictis, foliis ovatis concoloribus utrinque pubescentibus, cymis patentibus. L'Herit. corn. p. 5.* *sanguinea.*  
 Icon. *Schmidt arb. 2. t. 66. Fl. dan. 481.*  
 Near the lakes of Canada and New York. h. June, July. v. v. From eight to twelve feet high; anthers yellow; berries dark brown.
8. *C. ramis recurvatis, ramulis glabris, foliis ovatis acutis pubescentibus subtus canis, cymis depressis. L'Herit. corn. p. 6. Willd. sp. pl. 1. p. 662.*  
*C. stolonifera. Mich. fl. amer. 1. p. 92.*  
*C. tartarica. Mill. dict. n. 7.*  
 Icon. *Schmidt arb. 2. t. 65.*  
 On the banks of rivers and lakes: Canada to Virginia.  
 h. May—July. v. v. From six to ten feet high; branches red; berries blueish-white.
9. *C. ramis erectis, foliis ovatis acuminatis glabris subtus canis, cymis thyrsoides. L'Herit. corn. p. 9.* *paniculata.*  
*C. racemosa. Lam. encycl. 2. p. 116.*  
*C. citrifolia. Hort. par.*  
 Icon. *L'Herit. corn. t. 5. Schmidt arb. 2. t. 68.*  
 In swamps and near rivulets among other bushes: Canada to Carolina, rare. h. July, Aug. v. v. From four to six feet high; berries white, depressed globous.
10. *C. ramis verrucosis, foliis alternis ovatis acutis subtus canis, cymis depressis patentibus. L'Herit. corn. p. 10. Willd. sp. pl. 1. p. 664.* *alternifolia.*  
 Icon. *L'Herit. corn. t. 6. Schmidt arb. 2. t. 70.*  
 In shady woods, on river banks: Canada to Carolina.  
 h. May—July. v. v. A small tree from fifteen to twenty feet high; berries purple.

138. LUDWIGIA. *Gen. pl. 204.*\* *Corollatæ.*

- macrocarpa.* 1. *L. erecta*, ramosa, glabriuscula; foliis alternis lanceolatis subtus canis, pedunculis unifloris axillaribus, capsulis globoso-tetragonis, laciinis calycinis magnis coloratis coronatis. *Mich. fl. amer.* 1. p. 89.  
*L. alternifolia.* *Willd. sp. pl.* 1. p. 672.  
*Icon. Lam. encycl.* 3. p. 77. *Trew ehr.* 2. t. 2.  
In wet pastures and swamps: New England to Florida.  
 $\frac{1}{2}$ . July. v. v. Stem purple, two feet high; flowers yellow.
- hirsuta.* 2. *L. erecta*, ramosa, hirsuta; foliis alternis oblongis sessilibus utrinque hirsutis, pedunculis unifloris axillaribus, capsulis globoso-tetragonis coronatis basi bibracteatis. *Lam. encycl.* 3. p. 587.  
*L. pilosa.* *Walt. fl. car.* 89.  
In ditches and ponds on sandy soil: New Jersey to Carolina.  $\frac{1}{2}$ . July, Aug. v. v. The same size and habit as No. 1.
- linearis.* 3. *L. erecta*, virgata, glabra; foliis alternis linearibus acutis, floribus axillaribus solitariis sessilibus, capsulis oblongis turbinatis, laciinis calycinis semilanceolatis. — *Walt. fl. car.* 89.  
*L. angustifolia.* *Mich. fl. amer.* 1. p. 88.  
Near ditches and ponds in sandy soil: Virginia to Carolina.  $\frac{1}{2}$ . July, Aug. v. v. From two to five feet high, very branched; flowers small, yellowish-brown.
- virgata.* 4. *L. erecta*, virgata, glabra; foliis alternis linearibus obtusis, floribus terminalibus subspicatis pedicellatis, capsulis globoso-tetragonis ecoronatis. *Mich. fl. amer.* 1. p. 89.  
*L. ramosissima.* *Walt. fl. car.* 89.  
In dry sandy woods of Lower Carolina. *Michaux.*  $\frac{1}{2}$ . May, June. v. s. Flowers small.
- decurrens.* 5. *L. erecta*, ramosissima, glabra; foliis alternis linearilanceolatis decurrentibus, floribus axillaribus subsessilibus solitariis alternis, capsulis clavatis coronatis, laciinis calycinis ovali-lanceolatis. — *Walt. fl. car.* 89.  
*L. jussiaeoides.* *Mich. fl. amer.* 1. p. 89.  
In shady woods near ponds and ditches: Virginia and Lower Carolina.  $\odot$ . July, Aug. v. v. About a foot high, very branched; flowers large, yellow.

6. *L. erecta*, g  
basi rotun  
calyce bre  
ciniis caly  
p. 90.  
*L. suffrutico*  
In swamps c  
Sept. v.  
have short
7. *L. repens*, p  
glabris, l  
capsulis c  
lanceolatis  
*L. arcuata.*  
In swamps i  
jina.  $\frac{1}{2}$ .  
large, yell
8. *L. repens*; l  
latis, flor  
subovatis  
*L. apetala.*  
*L. repens.* i  
*Isnardia pal*  
*Icon. Lam.*  
In ditches  
 $\frac{1}{2}$ .  $\odot$ . M  
flowers e  
of petals.
9. *L. procumb*  
obovalib  
sessilibus  
natis, la  
*fl. car.* 8  
*L. microcat*  
In swamps  
Flowers
10. *L. erecta*, i  
oblongis,  
gestis, ca  
*L. rudis.* N  
In swamps

6. *L. erecta*, glabra; foliis alternis lato-linearibus acutis basi rotundatis, capitulis spicatis terminalibus, petalis calyce brevioribus, capsulis subglobosis coronatis, laciniis calycinis dilatatis brevibus. *Mich. fl. amer. 1.* p. 90.

*L. suffruticosa*. *Walt. fl. car. 90.*  
In swamps of North and South Carolina. ♀. July—  
Sept. v. v. The root is creeping; infertile branches  
have short obovate leaves; flowers small, yellow.

7. *L. repens*, pubescens; foliis oppositis linear-lanceolatis *pedunculosa*.  
glabris, pedunculis axillaribus unifloris longissimis,  
capsulis clavato-oblongis coronatis, laciniis calycinis  
lanceolatis. *Mich. fl. amer. 1. p. 88.*

*L. arcuata*. *Walt. fl. car. 89.*  
In swamps near the sea-coast: Virginia to South Caro-  
lina. ♀. May—July. v. s. A small plant; flowers  
large, yellow.

\*\* *Apetalæ*.

8. *L. repens*; lucida; foliis oppositis ovato-lanceolatis petio-  
latis, floribus axillaribus solitariis sessilibus, capsulis  
subovatis brevi-coronatis. *Mich. fl. amer. 1. p. 87.*

*L. apetala*. *Walt. fl. car. 89.*

*L. repens*. *Swartz. fl. ind. occid. 1. p. 273.*  
*Isnardia palustris*. *Willd. sp. pl. 1. p. 680.*  
Icon. *Lam. encycl. 3. t. 77.*

In ditches and ponds, common: Canada to Georgia.  
♀. ○. May—Sept. v. v. Floating in the water;  
flowers exceeding small, frequently with rudiments  
of petals.

9. *L. procumbens*, glaberrima; foliis alternis spatulato-  
obovalibus superne glanduloso-marginatis; floribus  
sessilibus axillaribus solitariis, capsulis minimis coro-  
natis, laciniis calycinis rotundis acuminatis. *Walt.*  
*fl. car. 88.*

*L. microcarpa*. *Mich. fl. amer. 1. p. 88.*  
In swamps of Lower Carolina. *Michaux. ♀. July. v. s.*  
Flowers small; leaves acuminate, small.

10. *L. erecta*, ramosa, pubescens; foliis alternis lanceolato-  
oblongis, floribus sessilibus alternis superioribus con-  
gestis, capsulis subrotundis. *Mich. fl. amer. 1. p. 90.*

*L. rudis*. *Walt. fl. car. 89.*  
In swamps of Lower Carolina. *Michaux. July. +.*

*capitata*.

*nitida*.

*glandulosa*.

*mollis*.

139. ALCHEMILLA. *Gen. pl.* 222. Ladies-mantle.

*alpina.*

1. *A. foliis digitatis apice dentatis sericeis.* *Willd. sp. pl.* 1. p. 698.

*Icon. Fl. dan.* 49. *Engl. bot.* 244.

On the peaks of high mountains in Vermont and New Hampshire. ♀. June, July. v. s. Whether the American species is the true *A. alpina* or not, I am not able to determine, as I am at present in want of specimens to compare them; but the plate in the *Flora Danica* represents the American plant fully, as far as recollection can decide.

*Aphanes.*

2. *A. foliis tripartitis, laciiniis trifidis pubescens, floribus axillaribus glomeratis monandris.* *Willd. sp. pl.* 1. p. 699.

*Aphanes arvensis. Linn. sp. pl.* 179.

*Icon. Engl. bot.* 1011. *Fl. dan.* 973. *Lam. illustr.* 1. t. 87.

In fields of Virginia. *Clayton.* ♂. May—Aug. v. s. in *Herb. Gronov.*

140. BOEHMERIA. *Gen. pl.* 1421.

*cylindrica.*

1. *B. foliis oppositis ovato-oblongis acuminatis dentatis glabris, floribus dioicis, spicis masculis glomeratis interrufatis, femineis cylindricis, caule herbaceo.* *Willd. sp. pl.* 4. p. 340.

*Urtica cylindrica. Linn. sp. pl.* 1396. *Mich. fl. amer.* 2. p. 179.

*Icon. Sloan. hist. jam.* 1. t. 82. f. 2.

In shady wet places: Canada to Florida. ♀. June—Aug. v. v. About two feet high.

*lateriflora.*

2. *B. foliis alternis ovato-lanceolatis acuminatis serratis scabris, floribus glomeratis lateralibus, caule herbaceo.* *Willd. sp. pl.* 4. p. 342.

In rocky shady woods: Pennsylvania to Virginia. ♀. July. v. v. Leaves on long petioles, trinerved.

141. URTICA. *Gen. pl.* 1422. Nettle.

*pumila.*

1. *U. foliis oppositis ovatis acuminatis trinerviis serratis, petiolis inferioribus longitudine folii, floribus monoicis triandris capitato-corymbosis petiolo brevioribus.* *Willd. sp. pl.* 4. p. 348.

In shady wood. July. v. size.

2. *U. foliis oppositis serratis, sp. p. 352.*

*Icon. Fl. dan.* In cultivated May—July

3. *U. foliis oppositis, floribus minatis petioli.* *Icon. Fl. dan.* On road sides Carolina.

4. *U. foliis oppositis ciliatis ratis geminis pl. 4. p. 35*  
*U. gracilis.*

On water sides vania. ♀ *tica gracilis* agrees in e

5. *U. foliis opp. gosis, gloflexis, cat.* On the island v. s. in He white and

6. *U. foliis alte nervibus spicis sol caule nud.* In shady w. June, Ju

7. *U. foliis alte petiolis ledivaricato muloso.—* In shady w. lina. ♀ to the fo

In shady woods among rocks: Canada to Carolina. ♂.  
July. *v. v.* Smooth and shining; very variable in size.

2. *U. foliis oppositis ellipticis subquinquenervibus argute serratis, spicis glomeratis geminatis.* *Willd. sp. pl. 4. p. 352.*

Icon. *Fl. dan.* 739. *Engl. bot.* 1236.

In cultivated grounds: Canada to Virginia; rare. ♂.  
May—July. *v. v.* Probably introduced from Europe.

3. *U. foliis oppositis cordatis ovato-lanceolatis grosse serratis, floribus dioicis, spicis paniculatis glomeratis geminatis petiolo longioribus.* *Willd. sp. pl. 4. p. 352.*

Icon. *Fl. dan.* 746.

On road sides and in waste places, frequent: Canada to Carolina. ♀. June—Aug. *v. v.*

4. *U. foliis oppositis cordatis ovato-lanceolatis serratis, petiolis ciliatis, floribus dioicis, spicis subramosis glomeratis geminis petiolum subæquantibus.* *Willd. sp. pl. 4. p. 353.*

*U. gracilis.* *Ait. kew.* 3. p. 341?

On water sides in rocky situations: Canada to Pennsylvania. ♀. July, Aug. *v. v.* The specimen of *Urtica gracilis* in the Herbarium of A. B. Lambert, Esq. agrees in every respect with the present species.

5. *U. foliis oppositis subsessilibus ovatis serratis subtus stri-gosis, glomerulis axillaribus sessilibus subglobosis reflexis, caule stimuloso.*

*chamædroi-des.*

On the islands of Georgia: St. Simon's, &c. ♂. May.  
*v. s. in Herb. Lyon.* The leaves are small: the stings white and very conspicuous.

6. *U. foliis alternis cordato-ovatis acuminatis serratis tri-nervibus petiolo duplo longioribus, glomerulis spicatis, spicis solitariis folio brevioribus, superne foliosis, caule nudo.* —*Willd. sp. pl. 4. p. 363.*

In shady woods, near rocks; Canada to Carolina. ♀.  
June, July. *v. v.* Resembles *U. dioica*.

7. *U. foliis alternis ovatis acuminatis serratis, glabriusculis, petiolis longis ciliatis, paniculis axillaribus solitariis divaricato-ramosissimis petiolo longioribus, caule sti-muloso.* —*Willd. sp. pl. 4. p. 365?*

In shady woods, in rocky situations: Canada to Carolina. ♀. July, Aug. *v. v.* This species is very near to the following, but is sufficiently distinct in the

*urens.*

*dioica.*

*procera.*

*capitata.*

*divaricata.*

## TETRANDRIA MONOGYNIA. Urtica.

leaves being not cordate and smooth, the panicles solitary and mixed with female flowers, and the general appearance of the plant.

*canadensis.*

8. *U. foliis alternis cordato-ovatis acuminatis serratis utrinque hispidis, paniculis axillaribus plerumque geminatis, divaricato-ramosissimis, inferioribus masculis petiolo longioribus, superioribus elongatis feminineis, caule hispidissimo stimuloso.* — *Willd. sp. pl. 4. p. 365.*

Icon. *Pluk. alm. t. 237. f. 2.*

Near rivulets, in rocky or sandy situations : Canada to Carolina, especially in the mountains.  $\frac{1}{2}$ . July, Aug. v. v. From five to six feet high ; leaves large.

142. PARIETARIA. *Gen. pl. 1576.* Pellitory.

*pensylvanica.*

1. *P. foliis oblongo-lanceolatis venosis opaco-punctatis, involucro triphylo floribus longiore.* — *Willd. sp. pl. 4. p. 955.*

On moist rocks : Pennsylvania, &c.  $\circ$ . June. v. v.; v. s. in *Herb. Nuttall.*

143. VISCUM. *Gen. pl. 1504.* Mistletoe.

*flavescens.*

1. *V. foliis lanceolatis obtusis, spicis axillaribus aggregatis.* — *Willd. sp. pl. 4. p. 740?*

*V. album. Walt. fl. car. 241.*

Parasitic on oaks and other trees ; rare.  $\frac{1}{2}$ . May. v. v. It is doubtful whether this is truly *V. flavescens* ; but the specimens observed by me in the West Indies were exactly the same as those of North America ; berries white.

144. CEPHALANTHUS. *Gen. pl. 147.* Button-wood.

*occidentalis.*

1. *C. foliis oppositis ternisque.* — *Willd. sp. pl. 1. p. 543.*

Icon. *Schmidt arb. t. 45. Pluk. alm. t. 77. f. 4.*

In deep swamps and morasses, common : Canada to Florida.  $\frac{1}{2}$ . July, Aug. v. v. Flowers white, in a round ball.

145. ELÆAGRUS. *Gen. pl. 213.* Oleaster.

*argentea.*

1. *E. inermis, foliis oblongis utrinque acutis argenteo-lepidotis, floribus subsolitariis nutantibus.*

In the extensive plains on the banks of the Missouri. *M. Lewis and T. Nuttall.*  $\frac{1}{2}$ . July. v. s. in *Herb.*

TET

Lewis.

A. B. La under the near to the and but on the up on both s gate : it cies, diff I do not

## 146. HIPPO

1. *H. foliis ov argenteo- gineis.* —

Near lakes v. v.

2. *H. foliis ov lepidotis.*

On the bank *Herb. L argentea* one for t bears rea other gen from Mr Elæagr

## 147. P

1. *Planera. W P. Gmelini*

*Anonymos* On the ban

A tree re the leave serrated end of the a greeni

*Lewis.* *Missouri Silver-tree.* In the collection of A. B. Lambert, Esq., are specimens from Dr. Pallas, under the name of *E. latifolius*, which approach very near to the present species; but the leaves are longer, and but slightly covered with that silvery substance on the upper surface, which in the present one is equal on both sides: besides, the flowers are generally aggregate: it is probable that both belong to the same species, different from *E. latifolius* of Linnæus, which I do not know.

146. HIPPOPHAE. *Gen. pl. 1509.* Sea Buckthorn.

1. *H. foliis ovatis acutiusculis, supra glabriusculis, subtus argenteo-pilosis lepidotisque: squamis sparsis ferrugineis.* — *Willd. sp. pl. 4. p. 744.*

Near lakes and rivers: Canada to New York.  $\frac{1}{2}$ . May.  
*v. v.*

2. *H. foliis ovatis obtusis utrinque glabris dense argenteo-lepidotis.*

On the banks of the Missouri. *M. Lewis.*  $\frac{1}{2}$ . *v. s. in Herb. Lewis. c. fr.* This shrub resembles *Elæagrus argentea* so much, that they might easily be mistaken one for the other when without fruit; but this species bears really a berry, different from the drupa of the other genus. The flowers are according to information from Mr. Nuttall four-cleft, and resemble those of *Elæagrus* very much.

*canadensis.**argentea.*

## II. DIGYNIA.

147. PLANERA. *Mich. fl. amer. 2. p. 247.*

1. *Planera.* *Willd. sp. pl. 4. p. 967.*

*P. Gmelini.* *Mich. fl. amer. 2. p. 248.*

*Anonymous aquatica.* *Walt. fl. car. 230.*

On the banks of rivers in both Carolinas.  $\frac{1}{2}$ . May. *v. v.*

A tree resembling an Elm in its foliage very much; the leaves are small, equal on their base and equally serrated; the flowers appear before the leaves at the end of the branches in close bunches, are small and of a greenish-brown colour.

*aquatica.*

148. HAMAMELIS. *Gen. pl. 226.* Witch-hazel.*virginica.*

1. *H. foliis obovatis acute dentatis sinu parvo cordatis.* —  
*Willd. sp. pl. 1. p. 701.*

*Icon. Catesb. car. 3. t. 2. Duham. arb. 1. t. 114.*

In stony and dry situations, but frequently near water : Canada to Florida. ♀. Octob.—Feb. v. v. This singular shrub, which grows about ten or twelve feet high, begins to show its yellow flowers in autumn, when all others have dropt their fruit, continues to flower all winter, and forms its fruit the spring following.

- macrophylla.* 2. *H. foliis suborbiculatis cordatis grosse obtuse-dentatis subtus scabro-punctatis.*

On river sides in the western part of Georgia. *J. Lyon.* ♀. v. s. in *Herb. Lyon.* The large leaves punctated on their lower side, with rough tubercles and other marks, give sufficient reasons to consider it a distinct species : besides this, it is very probable that under *H. virginica* are two distinct species, as the northern plant seems to differ very much from the southern in its outside appearance.

149. CUSCUTA. *Gen. pl. 227.* Dodder.*americana.*

1. *C. floribus pedunculatis umbellatis quinquefidis.* *Willd. sp. pl. 1. p. 702.*

*Icon. Sloan. hist. 1. t. 128. f. 4.*

Parasitic in hedges and on other plants, in moist situations. ♂. July. v. v. Twining round other plants ; leafless ; flowers in clusters ; the whole plant has a reddish-yellow appearance.

150. SANGUISORBA. *Gen. pl. 190.**canadensis.*

1. *S. spicis cylindricis longissimis, staminibus corolla multoties longioribus.* *Willd. enum. pl. & sp. pl. 1. p. 654.*

*Icon. Corn. canad. t. 174.*

In wet meadows : Canada to New York. ♀. July, Aug. v. v. Flowers in very long white spikes ; leaves resembling Burnet.

*media.*

2. *S. spicis cylindricis, staminibus corolla longioribus.* *Willd. enum. pl. & sp. pl. 1. p. 654.*

In wet meadows, principally on the mountains : Canada to Carolina. ♀. July, Aug. v. v. The spikes shorter, and tinged with red.

## 151. PACH

1. *P. caule pro crenato-d puberula.*

*Icon. Mich*

In shady ro  
♀. June.

## IV

## 152

1. *I. foliis ova basin ram p. 708.*

*I. Aquifoliu*

*Icon. Mees*

In sandy w  
in the s

An ever  
high and

2. *I. foliis ov peduncu gregatis*

*I. Aquifoli*

In shady

v. v. ; v.  
lower gr  
lowish-r

3. *I. foliis media su ramosis.*

*Icon. Cat*

In shady s  
v. v. A  
high.

4. *I. foliis la flexis, co*

## III. TRIGYNIA.

151. PACHYSANDRA. *Mich. fl. amer. 2. p. 177.*

1. *P. caule procumbente, foliis brevibus ovalibus superne procumbens. crenato-dentatis, calycibus minutim ciliatis, capsula puberula. Mich. fl. amer. 2. p. 178.*

Icon. *Mich. l. c. t. 45.*

In shady rocky situations on the Allegany mountains.  
1/2 June. v. v. Evergreen : flowers greenish-white.

## IV. TETRAGYNIA.

152. ILEX. *Gen. pl. 232. Holly.*

1. *I. foliis ovatis acutis spinosis glabris planis, floribus ad basin ramulorum annotinorum sparsis. Willd. sp. pl. 1. p. 708.*

*opaca.*I. *Aquifolium baccis rubris. Walt. fl. car. 241.*Icon. *Meerb. ic. 2. t. 5.*

In sandy woods, Canada to Carolina ; particularly plenty  
in the state of New Jersey. 1/2 May, June. v. v.  
An evergreen beautiful tree, sometimes eighty feet  
high and four in diameter ; berries scarlet.

2. *I. foliis ovatis sinuato-dentatis spinosis nitidis planis, pedunculis supra axillaribus in ramulis junioribus aggregatis laxe-ramosis. Lam. encycl. 3. p. 147.*

*laxiflora.*I. *Aquifolium baccis flavis. Walt. fl. car. 241.*

In shady sandy woods : Carolina. 1/2 May, June.  
v. v. ; v. s. in *Herb. Fraser.* An evergreen shrub of  
lower growth than the former species ; berries yellowish-red.

3. *I. foliis ovato-lanceolatis argute serratis planis, costa media subtus glabra, pedunculis lateralibus corymboso-ramosis. Ait. kew. 1. p. 170.*

*Cassine.*Icon. *Catesb. car. 1. t. 31.*

In shady swamps : Lower Carolina to Florida. 1/2 June.  
v. v. An evergreen shrub from eight to twelve feet  
high.

4. *I. foliis lanceolato-ellipticis subintegerrimis margine reflexis, costa media subtus villosa, pedunculis lateralibus*

*Dahoon.*

TETRANDRIA TETRAGYNIA. *Ilex.*

terminalibusque corymboso-paniculatis. — *Walt. fl. car. 241. Mich. fl. amer. 2. p. 228.*

In open swamps: Carolina to Florida.  $\frac{1}{2}$ . May, June. v. v. A beautiful evergreen shrub and sometimes tree; the leaves are very coriaceous, and resemble those of *Laurus Borbonia*.

*angustifolia.* 5. I. foliis linear-lanceolatis apice serratis margine reflexis, costa media subitus glabra, cymis lateralibus pedunculatis paucifloris. — *Willd. enum. pl.*

I. *myrtifolia. Walt. fl. car. 241.*

I. *rosmarinifolia. Lam. illustr. 1. p. 356.*

*ligustrifolia.*  $\beta$ . I. foliis latioribus omnibus integerrimis. An species distincta?

In deep swamps: Virginia to Georgia.  $\frac{1}{2}$ . June. v. v. An evergreen shrub from six to ten feet high.

*vomitoria.* 6. I. foliis oblongis utrinque obtusis crenato-serratis glabris umbellis lateralibus subsessilibus. *Ait. kew. 1. p. 170. Willd. sp. pl. 1. p. 709.*

I. *Cassine. Walt. fl. car. 241.*

*Ilex ligustrina. Jacq. collect. 4. p. 105.*

*Cassine Peragua. Mill. ic. t. 83. f. 2.*

Icon. *Jacq. ic. rar. 2. t. 310. Wendl. hort. t. 31.*

Along the sea-coast: Carolina to Florida.  $\frac{1}{2}$ . June, July. v. v. The leaves of this handsome shrub are used by the Indians to make their *Black drink*, so much in use among them, not only as a medicine, but also as a drink of etiquette on their councils, when matters of consequence are to be transacted.

*prinoides.* 7. I. foliis *deciduis* lanceolatis in petiolum attenuatis leviter serratis, costa media subitus villosa, pedunculis unifloris aggregatis, fertilibus solitariis. *Ait. kew. 1. p. 169. Willd. sp. pl. 1. p. 709.*

I. *decidua. Walt. fl. car. 241.*

On rocky shady banks of rivers: Virginia to Georgia.  $\frac{1}{2}$ . June, July. v. v. A small shrub; berries red.

*canadensis.* 8. I? foliis *deciduis* ovatis integerrimis seu apice rariter serratis glabris, pedunculis subsolitariis longissimis unifloris, fructu obtuse subtetragono. *Mich. fl. amer. 2. p. 229.*

Icon. *Mich. l. c. t. 49.*

On high mountains, near the edges of ponds: Canada to Carolina.  $\frac{1}{2}$ . April, May. v. v. Not above three feet high; berries large, beautiful crimson, very ornamental.

9. I? foliis *opposita*, postice *glabrum*, brevissimis *Pedunculis* terni, basi *Cal.* 4-fidata, 4-fid corollæ *rum*, ovali lobum, *ovata*, *alpisticum*. This very served a Esq., com mused in Mygind dered it only the not ser plant. would be cold mo same w sent, ti genus it itself.

1. S. *caulibus* *Smith.* Icon. *En* In barren *and Cl*

9. I? foliis *oppositis* oblongis obtusis serratis margine reflexis glabris sempervirentibus, pedunculis axillaribus brevissimis subsolitariis unifloris basi bibracteatis. *myrsinites.*

On the Rocky-mountain and near the Pacific Ocean.

*M. Lewis.*  $\frac{1}{2}$ . July, Aug. v. s. in *Herb. Lewis.*

*Frutex* sempervirens, glaberrimus, 4-pedalis, ramosissimus: *rami* teretes, fusi, dense foliosi. *Folia* parva opposita, oblonga seu obovata, obtusa, superne serrata, postice integerrima, margine revoluta, vix petiolata. *Pedunculi* axillares, brevissimi, uniflori, solitarii seu terni, basi opposite bibracteati. *Flores* minutti, albi. *Cal.* 4-fidus: laciniis subrotundis, concavis. *Cor.* rotata, 4-fida: laciniis oblongis. *Filam.* 4. ad incisuras corollæ inserta? longitudine calycis. *Germen* superum, ovale. *Styl.* brevissimus, clavatus. *Stig.* 4-lobum, crassum. *Drupa* maturitate monosperma, ovata, atro purpurea, magnitudine pisi. *Sem.* 1. ellipticum.

This very curious shrub, of which I have likewise observed a specimen in the museum of A. B. Lambert, Esq., collected by A. Menzies, deserves to be examined in the living plant. It approaches so near to *Myginda Rhacoma* of Jacquin, that at first I considered it to be the same plant. It likewise has great affinity to *Paltozia ovalis* *Flor. per.* 1. p. 44. t. 84. only the leaves are not opposite, but sparse and crenate not serrated, which is the case always in the present plant. As I do not know the *Myginda Rhacoma*, it would be a curious fact if this plant, from the high cold mountains of the north, should turn out to be the same with the tropical plant. I refer it to *Ilex* at present, till further observations will decide to which genus it belongs, or whether it may not form one by itself.

### 153. SAGINA. Gen. pl. 236.

1. S. caulinibus procumbentibus glabris, petalis brevissimis. *procumbens.*

*Smith fl. brit.* 1. p. 199. *Willd. sp. pl.* 1. p. 718.

*Icon. Engl. bot.* 880. *Pluk. alm.* t. 74. f. 2.

In barren sandy fields: Virginia and Carolina. *Walter* and *Clayton.* ☺. May—Aug. v. s. in *Herb. Walter.*

154. POTAMOGETON. *Gen. pl. 234.**natans?*

1. *P. foliis longe petiolatis natantibus sublanceolato-ovalibus, primariis nonnullis subcordatis.* *Mich. fl. amer. 1. p. 101.*

In ponds and slow-flowing waters, common.  $\mathcal{U}$ . June, July. *v. v.* Floating in the water; flowers, as all the following, in spikes axillary or sometimes terminal, small and of a greenish colour. It is doubtful whether this is not a new species, but I had no opportunity to compare it in a living or dried state with the European plant.

*fluitans.*

2. *P. foliis inferioribus longissimis linearibus, superioribus lanceolatis nervosis coriaceis, omnibus petiolatis.* — *Willd. sp. pl. 1. p. 713.*

*P. foliis lanceolato-oblongis, petiolis longis.* *Gron. fl. virg. 139.*

In ponds on the barrens of New Jersey and Carolina.  $\mathcal{U}$ . July. *v. v.*

*heterophyl-lum.*

3. *P. foliis superioribus petiolatis ellipticis utrinque attenuatis, inferioribus confertis sessilibus linearibus.* *Willd. sp. pl. 1. p. 713.*

*P. hybridum.* *Mich. fl. amer. 1. p. 101.*

In slow flowing waters of Virginia and Carolina.  $\mathcal{U}$ . June. *v. v.*

*setaceum.*

4. *P. foliis superioribus oppositis lanceolatis quinquenervibus brevi-petiolatis, inferioribus alternis filiformibus, spicis densis alternis folio brevioribus.* — *Willd. sp. pl. 1. p. 716?*

In small lakes and rivers, on the pine barrens of Jersey.  $\mathcal{U}$ . July, Aug. *v. v.* The upper leaves are scarcely half an inch long; the whole plant very slender.

*perfoliatum.*

5. *P. foliis cordato-ovatis amplexicaulibus omnibus immersis, spicis terminalibus, floribus alternis.* — *Mich. fl. amer. 1. p. 101.* *Willd. sp. pl. 1. p. 713.*

*Icon. Fl. dan. 196.* *Engl. bot. 168.*

In ponds and rivers, frequent: Canada to Pennsylvania.  $\mathcal{U}$ . July, Aug. *v. v.* Spike of flowers only floating, brown.

*lucens.*

6. *P. foliis lanceolatis subsessilibus, costa media crassa, spicis longis cylindricis.* *Mich. fl. amer. 1. p. 101.* *Willd. sp. pl. 1. p. 714.*

*Icon. Fl. dan. 195.* *Engl. bot. 376.*

In ponds and lina.  $\mathcal{U}$ . J the other s

7. *P. foliis alterri serrat, sp*

*Icon. Fl. dan*

In rivers and

July. *v. v.*

8. *P. foliosum;*

*spicis capit*

*P. gramineus*

In rivers nea

9. *P. foliis prael*

*ginantibus*

*fructibus*

*P. marinum.*

In ponds and

nia.  $\mathcal{U}$ .

water; fl

1. *Ruppia.* *W*

leon. *Engl*

On the mou

In ponds and rivers on sandy soil : New Jersey to Carolina. 2. July, Aug. v.v. Leaves larger than any of the other sorts ; spike greenish brown.

7. *P. foliis alternis oppositivis lanceolato-ellipticis undulatis serratis, spicis paucifloris.* — *Willd. sp. pl. 1. p. 714.*  
Icon. *Fl. dan. 927. Curt. lond. 5. t. 15.*  
In rivers and ponds : Canada to Virginia. 2. June, July. v.v. Spikes greenish red, small.
8. *P. foliosum ; foliis sessilibus angusto-linearibus planis, pauciflorum.*  
spicis capitatim subquadrifloris.  
*P. gramineum.* *Mich. fl. amer. 1. p. 102.*  
In rivers near the sea-coast of Lower Carolina. 2.
9. *P. foliis praelongo-setaceis approximatis distichis basi vaginantibus, spicis terminalibus interruptis verticillatis, fructibus ovatis turgidis.* — *Smith fl. brit. 1. p. 197.*  
*P. marinum.* *Mich. fl. amer. 1. p. 102.*  
In ponds and slow-flowing waters : Canada to Pennsylvania. 2. July. v.v. Only the spikes appear above water ; flowers seldom.

*crispum.**pectinatum.*155. RUPPIA. *Gen. pl. 235.*

1. *Ruppia.* *Willd. sp. pl. 1. p. 717.*  
Icon. *Engl. bot. 136. Mich. gen. 35. Fl. dan. 364.*  
On the mouth of most rivers in North America. 2. v.v.

*maritima.*

## V. PENTANDRIA.

### I. MONOGYNIA.

\* *Flores monopetali, inferi, tetraspermi. Asperifoliae.*

156. ECHIUM. *Cor. fauce nuda, irregularis, campanula-*

*lata.*

157. HELIOTROPIUM. *Cor. fauce nuda, hypocrateri-*

*formis, lobis dente interjectis.*

158. PULMONARIA. *Cor. fauce nuda, infundibuliformis.*

*Cal. prismaticus.*

159. LITHOSPERMUM. *Cor. fauce nuda, infundibuli-*

*formis. Cal. 5-partitus.*

160. BATSCHIA. *Cor. fauce nuda, hypocrateriformis,*

*intus basi annulo barbato cinctus : laciniis rotundatis.*

*Cal. 5-fidus.*

161. ONOSMODIUM. *Cor. fauce nuda, subcampanulata,*

*limbo ventricoso : laciniis conniventibus acutis.*

*Anth. sessiles inclusae. Cal. 5-partitus.*

162. LYCOPSIS. *Cor. fauce fornicata, infundibuliformis,*

*tubo curvato.*

163. CYNOGLOSSUM. *Cor. fauce fornicata, infundibu-*

*liformis. Sem. depressa, latere affixa.*

164. MYOSOTIS. *Cor. fauce fornicata, hypocrateri-*

*formis, lobis emarginatis.*

\* *Flores monopetali, inferi, angiospermi.*

165. HYDROPHYLLUM. *Caps. 1-locularis, 2-valvis.*

*Cor. campanulata, interne striis 5. melliferis longitu-*

*dinalibus. Stigm. 2-fidum.*

166. ANAGALLIS. *Caps. 1-locularis, circumscissa. Cor.*

*rotata. Stigm. capitatum.*

167. LYSIMACHIA. *Caps. 1-locularis, 10-valvis. Cor.*

*rotata. Stigm. obtusum.*

168. DODECATHEON. *Caps. 1-locularis, oblonga. Cor.*

rotata, re-  
tusum.

169. PRIMULA  
dricus, o

umbellula

170. ANDROS  
riformis, f

171. SABBATI  
limbo 5-  
spiralis

172. HOTTON  
formis.  
bosum.

173. MENYAN  
rotata : l

gine plan

174. VILLARS  
tata : lac

2-lobum

175. SPIGELL  
dibulifor

176. OPHIOR  
infundib

177. PHACEL  
sub-cam

178. ELLISIA  
spermii,

liformis,

Cal. pe

179. DATURA  
dibulifo

180. HYOSC  
Cor. inf

Stigm.

181. NICOTI  
infundi

Stigm.

182. VERBAS  
æqualis

183. CALYST  
Cor. c

Cal. ba

184. CONVO  
panula

aut bit

- rotata, reflexa. *Stam.* tubo insidentia. *Stigm.* obtusum.
169. PRIMULA. *Caps.* 1-locularis. *Cor.* tubus cylindricus, ore patulo. *Stigm.* globosum. *Involucr.* umbellulæ.
170. ANDROSACE. *Caps.* 1-locularis. *Cor.* hypocrateriformis, fauce coarctata. *Stigm.* globosum.
171. SABBATIA. *Caps.* 1-locularis. *Cor.* tubo urceolato, limbo 5—12-partito. *Stigm.* 2-partitum: laciniis spiralibus. *Anth.* demum revoluta.
172. HOTTONIA. *Caps.* 1-locularis. *Cor.* hypocrateriformis. *Stam.* tubo corollæ imposita. *Stigm.* globosum.
173. MENYANTHES. *Caps.* 1-locularis, 2-valvis. *Cor.* rotata: laciniis disco longitudinaliter barbatis, margine planis. *Stigm.* 2-fidum.
174. VILLARSIA. *Caps.* 1-locularis, evalvis. *Cor.* rotata: laciniis basi barbatis, margine inflexis. *Stigm.* 2-lobum. *Glandulæ* 5. staminibus alternæ.
175. SPIGELIA. *Caps.* 2-locularis, didyma. *Cor.* infundibuliformis. *Stigm.* simplex.
176. OPHIORRHIZA. *Caps.* 2-locularis, 2-partita. *Cor.* infundibuliformis. *Stigm.* 2-fidum.
177. PHACELIA. *Caps.* 2-locularis, 4-sperma. *Cor.* sub-campanulata. *Stam.* exerta. *Styl.* brevis.
178. ELLISIA. *Caps.* 2-locularis, 2-valvis: loculi 2-spermi, semine uno supra alterum. *Cor.* infundibuliformis, angusta. *Stam.* inclusa. *Stigm.* 2-fidum. *Cal.* persistens.
179. DATURA. *Caps.* 2-locularis, 4-valvis. *Cor.* infundibuliformis. *Cal.* deciduus.
180. HYOSCYAMUS. *Caps.* 2-locularis, operculata. *Cor.* infundibuliformis, irregularis. *Stam.* inclinata. *Stigm.* capitatum.
181. NICOTIANA. *Caps.* 2-locularis, 2-valvis. *Cor.* infundibuliformis, limbo plicato. *Stam.* inclinata. *Stigm.* emarginatum.
182. VERBASCUM. *Caps.* 2-locularis. *Cor.* rotata, inaequalis. *Fil.* barbata. *Stigm.* obtusum.
183. CALYSTEGIA. *Caps.* semibiloculare, sub-4-sperma. *Cor.* campanulata, 5-plicata. *Stigm.* 2. globosa. *Cal.* basi bibracteatus.
184. CONVOLVULUS. *Caps.* 2—3-locularis. *Cor.* campanulata, 5-plicata. *Stigm.* 2. filiformia. *Cal.* nudus aut bibracteatus.

## PENTANDRIA.

185. IPOMCEA. *Caps.* 2-3-locularis. *Cor.* infundibuliformis, 5-plicata. *Stigm.* capitatum, 2-3-lobum. *Cal.* nudus.
186. CANTUA. *Caps.* 3-locularis, 3-valvis. *Cor.* infundibuliformis. *Stigm.* 3-fidum. *Sem.* alata.
187. DIAPENSIA. *Caps.* 3-locularis, 3-valvis, polysperma. *Cor.* hypocrateriformis: tubo brevi. *Cal.* 5-partitus, basi bracteatus. *Styl.* brevis. *Stigm.* 3-lobum.
188. PHLOX. *Caps.* 3-locularis, 1-sperma. *Cor.* hypocrateriformis. *Fil.* inaequalia. *Stigm.* 3-fidum. *Cil.* prismaticus.
189. POLEMONIUM. *Caps.* 3-locularis. *Cor.* 5-partita, fundo clausa valvis staminiferis. *Stigm.* 3-fidum.
190. AZALEA. *Caps.* 5-locularis. *Cor.* infundibuliformis aut campanulata. *Stam.* receptaculo inserta. *Stigm.* obtusum.
191. BUMELIA. *Drupa* 1-sperma. *Cor.* hypocrateriformis limbo dentibus interjecto. *Nect.* 5-phylum.
192. SOLANUM. *Bacca* 2-locularis. *Cor.* rotata, plicata. *Antheræ* biperforatae, subcoalitæ. *Cal.* persistens.
193. PHYSALIS. *Bacca* 2-locularis intra calycem inflatum. *Cor.* campanulato-rotata. *Antheræ* approximatæ.
194. NICANDRA. *Bacca* 3-5-locularis, exsucca. *Cor.* campanulata. *Stam.* incurva. *Cal.* 5-angulatus: laciniis sagittatis,

\* *Flores monopetali, superi.*

195. SAMOLUS. *Caps.* 1-locularis, apice 5-valvis. *Cor.* hypocrateriformis. *Stigm.* capitatum.
196. PINKNEYA. *Caps.* 2-locularis: valvis medio septiferis. *Cor.* tubulosa. *Cal.* laciniis 1-2. bracteæformibus. *Fil.* basi tubi inserta. *Sem.* alata.
197. CAMPANULA. *Caps.* 2-5-locularis, poris lateribus dehiscens. *Cor.* campanulata, fundo clauso valvis staminiferis. *Stigm.* 3-5-fidum.
198. CHIOCOPCA. *Bacca* 2-sperma, didyma. *Cor.* infundibuliformis. *Stigm.* simplex.
199. CAPRIFOLIUM. *Bacca* 3-locularis, polysperma, distincta. *Cor.* tubulosa, longa, 5-fida. *Cal.* 5-dentatus.
200. XYLOSTEUM. *Baccæ* duæ basi connatæ, 2-locu-

lares, polyqualis.

201. SYMPHOE

nata, locu

brevis, 5-

202. DIERVILI

ma. *Cor.*

oblongus,

203. TRIOSTE

calyce vi

oblongum

204. RIBES.

bifidus.

205. RHAMN

corollifer

206. CEANO

losus, c

207. CELAST

calypt

208. EVONY

Sem. cal

209. VITIS.

emarcid

210. CISSUS.

reflexo-

gens.

211. CYRILL

Petala

binatus

212. ITEA.

introfle

Cal. 5

lobum.

213. IMPATI

regular

hærent

214. VIOLA

laris, c

215. CLAYT

valvis.

dibuli-  
lobum.  
infun-  
lysper-  
Cal. 5.  
gmt. 3.  
hypo-  
1. Cil.  
partita,  
um.  
formis  
Stigm.  
erifor-  
im.  
plica-  
l. per-  
infla-  
prox-  
Cor.  
latus :  
  
Cor.  
septi-  
eæfor-  
latera-  
clauso  
or. in-  
erma,  
i-den-  
-locu-

lares, polyspermæ. *Cor.* infundibuliformis, subæqualis. *Cal.* 5-dentatus.

201. SYMPHORIA. *Bacca* 4-locularis, 4-sperma, coronata, loculis interdum 2. abortivis. *Cor.* tubulosa, brevis, 5-fida, subæqualis. *Cal.* 4-dentatus.
202. DIERVILLA. *Caps.* 4-locularis, oblonga, polysperma. *Cor.* infundibuliformis, 5-fida, patens. *Cal.* oblongus, 5-fidus.
203. TRIOSTEUM. *Bacca* 3-locularis, 3-sperma. *Cor.* calyce vix longior, tubulosa. *Cal.* 5-fidus. *Stigm.* oblongum.

\* *Flores pentapetalii, superi.*

204. RIBES. *Bacca* polysperma. *Cal.* corollifer. *Styl.* bifidus.

\* *Flores pentapetalii, inferi.*

205. RHAMNUS. *Bacca* 3-4-sperma. *Cal.* tubulosus, corollifer. *Pet.* oris 5, convergentia. *Dioicus.*
206. CEANOTHUS. *Bacca* 3-cocca, sicca. *Cal.* tubulosus, corollifer. *Petala* fornicata.
207. CELASTRUS. *Caps.* 3-cocca, 3-locularis. *Sem.* calyprata. *Cor.* 5-petala, patens.
208. EVONYMUS. *Caps.* 5-gona, 5-valvis, colorata. *Sem.* calyprata. *Cor.* 5-petala, plana.
209. VITIS. *Bacca* 5-sperma. *Petala* apice cohærentia, emarcida. *Styl.* 0. *Dioica.*
210. CISSUS. *Bacca* 2-locularis, 1—4-sperma. *Petala* reflexo-patula, decidua. *Nectar.* discus germen cingens.
211. CYRILLA. *Bacca* 2-locularis, 2-sperma, exsiccabilis. *Petala* 5. stellatum patentia. *Cal.* minimus subturbinatus. *Stylus* 2-fidus.
212. ITEA. *Caps.* 2-locularis, 2-valvis : valvis margine introflexo seminiferis. *Petala* 5. reflexo-patentia. *Cal.* 5-fidus, campanulatus. *Stigm.* capitatum, 2-lobum.
213. IMPATIENS. *Caps.* 1-locularis, 5-valvis. *Cor.* irregularis, calcarata. *Cal.* 2-phillus. *Anther.* cohærentes.
214. VIOLA. *Caps.* 1-locularis, 3-valvis. *Cor.* irregularis, calcarata. *Cal.* 5-phillus. *Anther.* cohærentes.
215. CLAYTONIA. *Caps.* 1-locularis, 3-valvis. *Cal.* 2-valvis. *Stigm.* 3-fidum.

*Cal. inferus, 1-phyllo, coloratus, 5-lobus.  
Caps. 1-locularis, 5-valvis, 5-sperma, calyce cincta.*

\* *Flores incompleti.*

216. GLAUX. *Cal. inferus, 1-phyllo, coloratus, 5-lobus.  
Caps. 1-locularis, 5-valvis, 5-sperma, calyce cincta.*
217. ANYCHIA. *Cal. inferus, connivens, 5-partitus:  
laciiniis oblongis, apice subsaccatis. Stigm. 2. Ultra-  
culus 1-spermus.*
218. THESIUM. *Cal. superus, staminifer. Sem. 1. co-  
ronatum.*
219. NYSSA. *Cal. superus, 5-partitus. Drupa. Mas.  
Stam. 10. Polygama.*
220. HAMILTONIA. *Cal. superus, 5-fidus. Nect. dis-  
cus 5-dentatus. Drupa. Dioica.*

229. DICHONI

*locularis:  
Stigm. P  
subspatu*

230. EVOLVUI

*Cor. rotat  
laciiniis ca*

231. HEUCHE

*calyci ins*

232. ZIZYPHI

*nous, o  
sus, coro*

\* *Flores pe*

233. ERYNGI

*Invol. p  
perus.*

234. HYDRO

*phyllum  
pressa.*

235. PANAX.

*sperma.*

236. SANICU

*tus acul*

237. DAUCU

*tivi. I*

*tracta,*

238. AMMI.

*Fructu*

239. SELINU

*nus, ir*

*data, a*

240. FERUL

*phyllu*

*utrinq*

241. HERAC

*compr  
flexo-*

242. LIGUS

*que.*

*5-den*

## II. DIGYNIA.

\* *Flores monopetalii, inferi.*

221. ECHITES. *Folliculi 2. longi, recti. Sem. papposa.  
Cor. infundibuliformis, fauce nuda. Anther. conni-  
ventes in conuris.*
222. GONOLOBUS. *Folliculi 2. utplurimum costati seu  
angulati. Sem. papposa. Cor. rotata, profunde 5-  
partita. Appendix brevissima, inclusa. Styl. dis-  
coideo 5-gonus.*
223. APOCYNUM. *Folliculi 2. longi, lineares. Sem.  
papposa. Cor. campanulata. Filamenta 5. cum sta-  
minibus alterna. Styl. subnullus. Stigm. latum.*
224. ASCLEPIAS. *Folliculi 2. Sem. papposa. Cor. ro-  
tata, plerumque reflexa. Nectar 5. ovata, concava,  
corniculum exserentia. Anther. cornæ, longitudinaliter  
dehiscentes.*
225. AMSONIA. *Folliculi 2. Sem. nuda, teretia, ob-  
lique truncata. Cor. infundibuliformis fauce clausa.*
226. GELSEMINUM. *Caps. 2-locularis, 2-valvis, com-  
presso-plana. Sem. plana, valvularum marginibus  
annexa. Cor. infundibuliformis, limbo patente, sub-  
æquali. Cal. 5-dentatus.*
227. GENTIANA. *Caps. 1-locularis, 2-valvis. Recep-  
taculis 2. longitudinalibus. Cor. 1-petala, indeter-  
minata.*
228. HYDROLEA. *Caps. 2-locularis, 2-valvis. Cor.  
rotata. Filam. basi cordata. Stigm. capitato-peltata.  
Cal. 5-phyllo.*

- lobus.  
cincta,  
titus :  
Utri.  
1. co.  
Mas.  
t. dis.  
ppos.  
conni.  
ti seu  
nde 5.  
l. dis.  
Sem.  
m sta.  
im.  
or. ro-  
ncava,  
gitudi-  
a, ob-  
ausa.  
com-  
inibus  
, sub-  
Recep-  
deter-  
Cor.  
eltata.
229. DICHONDRA. *Caps.* subcompressa, didyma, 2-locularis: loculis 1-spermis. *Cor.* brevi-campanulata. *Stigm.* peltato-capitata. *Cal.* 5-partitus: laciniis subspathulatis.
230. EVOLVULUS. *Caps.* 1-locularis, 4-valvis, 1-sperma. *Cor.* rotato-campanulata. *Styli* 2, profunde 2-fidi: laciniis capillaribus, divergentibus. *Stigm.* simplicia.

\* *Flores pentapetali, inferi.*

231. HEUCHERA. *Caps.* 2-locularis, 2-rostris. *Pet.* 5. calyci inserta.
232. ZIZYPHUS. *Drupa* nuce 2-loculari. *Discus* carnosus, orbicularis, pistillum ambiens. *Cal.* tubulosus, corollifer. *Dioicus.*

\* *Flores pentapetali, superi dispermi. Umbellatae.*

233. ERYNGIUM. *Flores* capitati. *Recep.* paleaceum. *Invol.* polyphyllum. *Cal.* proprius 5-phillus, superus.
234. HYDROCOTYLE. *Umbella* simplex. *Invol.* 4-phylum. *Pet.* integra. *Sem.* semioberculato-compressa.
235. PANAX. *Umbella* simplex. *Bacca* cordata, 2-sperma. *Polygama.*
236. SANICULA. *Umbellæ* confertæ, subcapitatæ. *Fructus* aculeatus. *Flores* disci abortientes.
237. DAUCUS. *Corollæ* subradiatæ. *Flosculi* disci abortivi. *Fructus* pilis hispidus. *Umbella* fructus contracta, infundibuliformis. *Involucra* pinnatifida.
238. AMMI. *Corollæ* radiatæ: omnes hermaphroditæ. *Fructus* lœvis. *Umbella* laxa. *Involucra* pinnatifida.
239. SELINUM. *Fructus* ovali-oblongus, compresso-planus, in medio striatus. *Invol.* reflexum. *Pet.* cordata, æqualia. *Cal.* integer.
240. FERULA. *Invol.* universale caducum: partiale polyphyllum. *Fructus* ovalis, compresso-planus, striis utrinque 3.
241. HERACLEUM. *Fructus* ellipticus, emarginatus, compressus, striatus, marginatus. *Cor.* difformis, inflexo-emarginata. *Invol.* caducum.
242. LIGISTICUM. *Fructus* oblongus. 5-sulcatus utrinque. *Corollæ* æquales. *Pet.* involuta, integra. *Cal.* 5-dentatus.

243. ANGELICA. *Fructus* subrotundus, solidus, utrinque 3-alatus, stylis reflexis. *Corollæ* æquales : petalis incurvis.
244. SIUM. *Fructus* subovatus, compressus, striatus. *Invol.* polypyllum. *Pet.* cordata.
245. CENANTHE. *Flosculi* difformes : in disco sessiles, steriles. *Fructus* striatus, retusus, suberoso-corticatus, coronatus calyce et pistillo. *Umbellæ* globosæ.
246. PHELLANDRIUM. *Flosculi* disci minores. *Fructus* ovatus, lœvis, coronatus calyce et pistillo.
247. CICUTA. *Fructus* subovatus, sulcatus. *Invol.* 0; partiale 3—5-phylum.
248. CONIUM. *Fructus* subglobosus, 5-striatus, utrinque crenatus. *Involucella* dimidiata, subtriphylla.
249. CHÆROPHYLLUM. *Invol.* reflexum, concavum. *Pet.* inflexo-cordata. *Fructus* oblongus, lœvis, aut striatus, glaberimus.
250. PASTINACA. *Fructus* ellipticus, compresso-planus. *Pet.* involuta, integra.
251. SMYRNIUM. *Fructus* subcompressus, gibbosus, striatus. *Pet.* acuminata, carinata.
252. SESELI. *Umbellæ* globosæ. *Invol.* foliolo uno altero. *Fructus* ovatus, striatus.

## \* Flores incompleti.

253. SAL SOLA. *Sem.* 1. cochleatum, tectum. *Cal.* 5-phillus.
254. CHENOPODIUM. *Sem.* 1. lenticulare, superum. *Cal.* 5-phillus, 5-gonus.
255. ATRIPLEX. *Sem.* 1. compressum seu depresso. *Cal.* 5-phillus. *Flor.* feminei intermixti : quibus *Cal.* 2-phillus. *Polygama.*
256. CANNABIS. *Nux* 2-valvis, intra calycem clausum. *Mas.* *Cal.* 5-partitus. *Fem.* *Cal.* 5-phillus, integer, latere hians. *Dioica.*
257. HUMULUS. *Sem.* 1. intra calycem foliatum (in strobilum). *Mas.* *Cal.* 5-phillus. *Fem.* *Cal.* 1-phillus, oblique patens, integer. *Dioica.*
258. ULMUS. *Samara* compressa, membranaceo-alata. *Cal.* campauulatus, 5-fidus, marcescens.
259. CELTIS. *Drupa* 1-sperma. *Cal.* 5—6-partitus. *Stam.* 5—6. *Styl.* divaricati, crassiusculi. *Polygama.*

260. VIBUR fida.
261. SAMBU Bacca
262. RHUS. sperm
263. STAPH Caps. catrice
264. TURN Pet. 5 1-loct
265. CROTC Stigm Mono
266. KOCH simpli cinctu
267. AMAR Stam. scissa.
268. ACNII silia.
269. PARN corda 4-val
270. ARAI coroi
271. ZANT Caps
272. LINU coali
273. DROS locul

## III. TRIGYNIA.

260. VIBURNUM. *Cal. superus, 5-partitus. Cor. 5-fida. Drupa 1-sperma.*
261. SAMBUCUS. *Cal. superus, 5-partitus. Cor. 5-fida. Bacca 3-sperma.*
262. RHUS. *Cal. inferus, 5-partitus. Pet. 5. Bacca 1-sperma.*
263. STAPHYLEA. *Cal. inferus, 5-partitus. Pet. 5. Caps. inflatæ, connatæ. Nuces 2. globosæ cum cicatrice.*
264. TURNERA. *Cal. inferus, 5-fidus, infundibuliformis. Pet. 5. calyce inserta. Stigmata multifida. Caps. 1-locularis, 3-valvis.*
265. CROTONOPSIS. *Cal. inferus, 5-partitus. Pet. 5. Stigmata bifida. Caps. 1-sperma, non dehiscens. Monoica.*
266. KOCHIA. *Cal. inferus, 5-fidus. Cor. 0. Stigm. simplicia. Sem. 1. orbiculatum, calyce persistente cinctum.*
267. AMARANTHUS. *Cal. 3—5-phyllo. Cor. 0. Stam. 3—5. Caps. 1-locularis, 1-sperma, circumscissa. Monoicus.*
268. ACNIDA. *Cal. 3—5-partitus. Cor. 0. Stigm. sessilia. Caps. 1-sperma. Dioica.*

## IV. TETRAGYNIA.

269. PARNASSIA. *Cal. 5-partitus. Pet. 5. Nectar. 5. cordata, ciliata, apicibus globosis. Caps. 2-locularis, 4-valvis.*

## V. PENTAGYNIA.

270. ARALIA. *Cal. superus, 5-dentatus. Pet. 5. Bacca coronata, 5-sperma. Involucr. umbellulæ.*
271. ZANTHOXYLUM. *Cal. inferus, 5-partitus. Cor. 0. Caps. 3—5. 1-spermæ. Dioicum.*
272. LINUM. *Cal. inferus, 5-phyllo. Pet. 5. Fil. basi coalita. Caps. 10-locularis, 5-valvis. Sem. solitaria.*
273. DROSERA. *Cal. inferus, 5-fidus. Pet. 5. Caps. 1-locularis, apice 3—5-valvis. Sem. plurima.*

274. SIBBALDIA. *Cal.* inferus, 10-fidus. *Pet.* 5. *calyci inserta.* *Styli* e latere germinis. *Sem.* 5.  
 275. STATICE. *Cal.* inferus, 1-phyllo, tubulosus, pli-catus, 10-dentatus. *Cor.* 1-petala, hypocrateriformis, 5-fida. *Caps.* 1-locularis, basi 5-valvis. *Sem.* 1. teres.

## VI. POLYGYNIA.

276. ZANTHORHIZA. *Cal.* 0. *Pet.* 5. *Nectar.* 5. pedicellata. *Caps.* 1-sperma.  
 277. SCHISANDRA. *Cal.* triplici serie 9-phyllo. *Cor.* 0. *Anther.* subsessiles, apice cohaerentes. *Baccæ numerosæ, 1-spermæ, receptaculo filiformi elongato insertæ.* *Monoica.*

## I. MONOGYNIA.

156. ECHIUM. *Gen. pl.* 267.*vulgare.*

1. *E.* caule tuberculato-hispido, foliis caulinis lanceolatis hispidis, floribus spicatis lateralibus. *Willd. sp. pl.* 1. p. 787.  
 Icon. *Fl. dan.* 445. *Lam. illustr.* 94. f. 1.  
 On dry slate-hills: New York to Virginia, rare. Introduced from Europe? ♂. July—Sept. v. v.

157. HELIOTROPIUM. *Gen. pl.* 239.*indicum.*

1. *H.* foliis cordato-ovatis acutis scabriusculis, spicis solitariis, fructibus bifidis. *Willd. sp. pl.* 1. p. 740.  
 Icon. *Pluk. phyt.* t. 295. f. 4.  
 On road sides: Virginia to Carolina, rare. ○. June—Aug. v. v.

158. PULMONARIA. *Gen. pl.* 244.*virginica.*

1. *P.* glabra, erecta; foliis lanceolato-ovatis obtusiusculis, floribus fasciculato-terminalibus, corollis calyce multe longioribus.—*Willd. sp. pl.* 1. p. 769.  
 Icon. *Mill. ic.* 212. *Bot. mag.* 160.

PENT.

On gravelly

4. Ma-

2. *P.* subhirsu-

floribus

titis.—N

On Hudson

or white

3. *P.* glaberr

rupe

ad

floris,

amer.

P. maritim

On the sea

July.

159.

1. *L.* seminib

foliis o

Willd. s

Icon. En

In waste f

probably

v. v. I

2. *L.* seminib

tis, caly

foliis o

p. 131.

In shady v

v. v. I

3. *L.* seminib

tatis, fl

presso-p

amer. 1

In shady v

July, A

4. *L.* seminib

bracteis

Willd.

Icon. Col

In dry v

May—

On gravelly shores of rivers: Pennsylvania to Carolina.  
 ♀. March—May. v. v. Flowers beautiful blue.

2. *P. subhirsuta*, erecta; foliis ovato-oblongis acuminatis, floribus paniculatis, calycibus abbreviatis quinquepartitis.—*Willd. sp. pl. 1. p. 769.*

On Hudson's Bay. *Aiton.* May, June. +. Flowers blue or white.

3. *P. glaberrima*, procumbens; foliis ovali-spathulatis abrupte acuminatis carnosus, pedunculis lateralibus unifloris, corollis calyce vix duplo longioribus.—*Mich. fl. amer. 1. p. 131.*

*P. maritima?* *Willd. sp. pl. 1. p. 770.*

On the sea shore of New England and Canada. ♀. July. v. s. Flowers blue.

### 159. LITHOSPERMUM. Gen. pl. 241.

1. *L. seminibus rugosis*, corollis vix calycem superantibus, foliis obtusis aveniis. *Smith fl. brit. 1. p. 213.*  
*Willd. sp. pl. 1. p. 751.*

Icon. *Engl. bot. t. 123. Fl. dan. 456.*

In waste fields, on dry ground: frequent in Pennsylvania, probably introduced from Europe. ♂. May—July.  
 v. v. Flowers pale white; seeds rough, brownish.

2. *L. seminibus turgide ovatis lucidis undique cavo-punctatis*, calycibus fructiferis patulis corollam superantibus, foliis ovato-oblongis nervosis.—*Mich. fl. amer. 1. p. 131.*

In shady woods: Virginia to Kentucky. ♀. June, July.  
 v. v. Flowers pale yellow.

3. *L. seminibus turgide ovatis nitidis undique cavo-punctatis*, floribus passim lateralibus, foliis linearibus adpresso-pubescentibus, caule procumbente.—*Mich. fl. amer. 1. p. 130.*

In shady woods: on the river Ohio. *Michaux.* ♂.  
 July, Aug. v. s. Flowers white.

4. *L. seminibus muricatis*, spicis terminalibus secundis, bracteis lanceolatis, foliis linear-lanceolatis acutis.  
*Willd. sp. pl. 1. p. 752.*

Icon. *Col. ecphr. 1. t. 192.*  
 In dry woods: Virginia, Ohio and Mississippi. ♂.  
 May—Aug. v. v. Flowers yellow, very small.

*paniculata.*

*parviflora.*

*arvense.*

*latifolium.*

*angustifoli-  
um.*

*apulum.*

## PENTANDRIA MONOGYNIA.

160. BATSCHIA. *Mich. fl. amer.* 1. p. 129.*Gmelini.*

1. *B. hirsuta*; foliis floralibus ovatis, calycibus longis sub-lanceolatis.—*Mich. fl. amer.* 1. p. 130.  
*Anonymos carolinensis*. *Walt. fl. car.* p. 91.  
In dry sunny woods of Lower Carolina.  $\frac{1}{4}$ . May—July.  
*v. s. in Herb. Lyon.* Flowers pale yellow.

*canescens.*

2. *B. candicanti-villosa*; foliis omnibus oblongis, calycibus brevissimis, laciiniis corollæ integris.—*Mich. fl. amer.* 1. p. 130.

*Anchusa floribus sparsis, caule glabro.* *Gron. fl. virg.* p. 24.

*Icon. Mich. fl. amer.* 1. t. 14.

On dry sunny hills, in sandy soil: Virginia, Tennessee, &c.  $\frac{1}{4}$ . June, July. *v. v.* Flowers of a deep golden yellow; the root is covered with a red substance, which is the true *Puccoon* of the Indians, and paints a beautiful red.

*longiflora.*

3. *B. sericeo-villosa*; foliis linearibus, calycibus longis linearibus, laciiniis corollæ crenato-incisis, tubo elongato, On the banks of the Missouri. *Nuttall.*  $\frac{1}{4}$ . July. *v. s. in Herb. Nuttall.* Flowers yellow.

161. ONOSMODIUM. *Mich. fl. amer.* 1. p. 132.*hispidum.*

1. *O. hispidum*; foliis obovali-lanceolatis acutis papilloso-punctatis, laciiniis corollæ subulatis.—*Mich. fl. amer.* 1. p. 133.

*Lithospermum virginianum.* *Willd. sp. pl.* 1. p. 752.  
On dry lime-stone hills: New York to Carolina.  $\frac{1}{4}$ . July, Aug. *v. v.* Flowers yellowish-white, similar to those of *Sympyrum*.

*molle.*

2. *O. candicanti-vilosum*; foliis oblongo-ovalibus subtriplinerviis, laciiniis corollæ subovalibus.—*Mich. fl. amer.* 1. p. 133.

*Lithospermum carolinianum.* *Lam. illustr.* 1. p. 397.  
*Icon. Mich. l. c. t. 15.*  
In the western countries from Pennsylvania to Tennessee.  $\frac{1}{4}$ . June—Aug. *v. v.* Flowers white.

162. LYCOPSIS. *Gen. pl.* 250.*arvensis.*

1. *L. hispida*; foliis lanceolatis repando-dentatis, racemis geminis, floribus sessilibus.—*Willd. sp. pl.* 1. p. 780.

Icon. *Fl.* c  
In fields :  
○. Jun  
2. *L. pusilla*,  
periorib  
tariis, fl  
In dry wo  
○. Ma

163.

1. *C. mollis*  
racemis

*Icon. Cu*

In dry wo  
June, J

2. *C. nudius*  
tus scal

*C. virgin*

*Icon. En*

In simila

Flower

3. *C. hirsut*  
plexia

culato

*C. foliis*

*C. virg*

In shady

May—

*Willd*  
allude

which the pi

1. *M. pere*  
glabri

foliis

*M. scor*

*Icon. J*

In rivul

2. *M. ann*  
natis

- Icon. *Fl. dan.* 435. *Engl. bot.* 938. *Lam. illustr.* t. 92.  
In fields: Pennsylvania to Virginia. Probably introduced.  
♂. June, July. v. v. Flowers blue.
2. *L. pusilla*, hispidissima; foliis inferioribus spathulatis, superioribus lineariforme-oblongis integerrimis, racemis solitariis, floribus pedunculatis.—*Willd. sp. pl.* 1. p. 781.  
In dry woods and on road sides: Pennsylvania to Virginia.  
♂. May—July. v. v. Flowers white, very small.

*virginica.*163. CYNOGLOSSUM. *Gen. pl.* 243.

1. *C. mollissime pubescens*; foliis lato-lanceolatis sessilibus, racemis paniculatis.—*Willd. sp. pl.* 1. p. 760.  
Icon. *Curt. lond.* 4. t. 16. *Bot. mag.*  
In dry woods, in the lime-stone countries, frequent. ♂.  
June, July. v. v. Flowers brownish-red.
2. *C. nudiusculum*; foliis spathulato-lanceolatis lucidis subtus scabris, racemis sparsis. *Smith fl. brit.* 1. p. 216.  
*C. virginicum*. *Willd. sp. pl.* 1. p. 760.  
Icon. *Engl. bot.* 1642.  
In similar situations in Virginia. ♂. June, July. v. v.  
Flowers blueish-brown.
3. *C. hirsutissimum*; foliis ovali-oblongis, superioribus amplexicaulis, corymbo terminali aphylo longe pedunculato. *Mich. fl. amer.* 1. p. 132.  
*C. foliis amplexicaulis*. *Gron. fl. virg.* p. 24.  
*C. virginicum*. *Linn. sp. pl.* 134.  
In shady moist woods: Pennsylvania to Virginia. ♀.  
May—July. v. v. Flowers beautiful blue and white.  
Willdenow's *C. virginicum* is certainly not the plant alluded to by Linnaeus in his *Species Plantarum*, by which he refers to Clayton's plant, which evidently is the present species.

*officinale.**sylvaticum.*164. MYOSOTIS. *Gen. pl.* 240.

1. *M. perennis*; seminibus laevibus, calycibus subovalibus glabris tubo corollæ fere longitudine, caule subramoso, foliis lanceolatis. *Roth. germ.* i. 87. ii. 221.  
*M. scorpioides*. *Willd. sp. pl.* 1. p. 746.  
Icon. *Fl. dan.* 583.  
In rivulets and ditches, frequent. ♀. May—Sept. v. v.
2. *M. annua*; seminibus laevibus, calycibus ovalibus acuminatis hirsutissimis tubo corollæ longioribus, caule ra-

*palustris.**arvensis.*

PENTANDRIA MONOGYNIA. *Myosotis.*

mosissimo, racemis conjugatis, foliis ovato-lanceolatis.  
*Willd. sp. pl. 1. p. 747.*

In fields and shady woods: New York, Pennsylvania, &c. ☽. June, July. v. v. Flowers pale blue, very small.

*virginiana.*

3. M. pilosum; seminibus aculeato-glochidibus, foliis ovato-lanceolatis acuminatis, racemis divaricatis.—*Willd. sp. pl. 1. p. 748.*

In dry open woods: Virginia, Carolina, &c. ☽. June, July. v. v. Flowers small, pale blue; leaves large.

*Lappula.*

4. M. hispidum; seminibus aculeis glochidibus, foliis linear-i-oblengis, caule stricte ramosissimo.—*Willd. sp. pl. 1. p. 749.*

On way sides and among rubbish: Canada to Virginia, ☽. May—Aug. v. v. Flowers blue.

160

1. A. foliis o  
Smith fl  
Icon. Eng  
In cultivat  
duced fi  
red.

161

1. L. glaberr  
longo-li  
laciniis  
In Lower  
as the  
2. L. glaber  
puncta  
corolla  
L. vulga  
L. stricta  
L. bulbit  
Icon. Bo  
In wet  
Virgin  
quentl  
axillæ

3. L. glabr  
sessilis  
lateral  
L. thyrs  
In salt-  
Thou  
tainly

4. L. pubes  
ovalit  
unifl  
sp. p  
L. punct  
Icon. I  
In dry  
v. v.  
verti

5. L. glal

165. HYDROPHYLLUM. *Gen. pl. 267.*

*appendicula-  
tum.*

1. H. hirsutissimum; foliis radicalibus subpinnatifidis, caulinis sublobato-angulosis, calycis sinubus appendiculatis, fasciculis florum subpaniculatis.—*Mich. fl. amer. 1. p. 134.*

On shady rocks, near springs: Virginia, near Harper's Ferry, Tennessee, &c. ♀. May. v. v. Flowers pale blue; the calyx extremely hispid, and almost the length of the corolla.

*virginicum.*

2. H. glabriuscultum; foliis pinnatifidis pinnatisque, laciniis ovali-lanceolatis inciso-serratis, fasciculis florum conglomeratis.—*Willd. sp. pl. 1. p. 814.*

Icon. *Lam. illustr. t. 97. f. 1.*

In shady rocky situations: Canada to Carolina. ♀. May, June. v. v. Flowers beautiful blue.

*canadense.*

3. H. hirsutum; foliis lobato-angulosis, fasciculis florum confertis.—*Willd. sp. pl. 1. p. 815.*

Icon. *Lam. illustr. t. 97. f. 2.*

On wet rocks and shady mountains: Canada to Pennsylvania. ♀. June, July. v. v. Flowers white, shaded with purple.

*lineare.*

4. H. pilosum; foliis linearibus, racemis elongatis.

On the banks of the Missouri. *M. Lewis* ♀. April. v. s. in *Herb. Lewis.*

166. ANAGALLIS. *Gen. pl. 270.*

1. *A. foliis ovatis subtus punctatis, caule procumbente.* *arvensis.*  
*Smith fl. brit. 1. p. 230. Willd. sp. pl. 1. p. 821.*

*Icon. Engl. bot. 529. Fl. dan. 88.*

In cultivated grounds: Pennsylvania to Virginia; introduced from Europe? ♂. June, July. v.v. Flowers red.

167. LYSIMACHIA. *Gen. pl. 269.*

1. *L. glaberrima, ramulosa; foliis oppositis verticillatisque angustifolia.*  
*longo-linearibus punctatis, racemo terminali brevi,*  
*laciniis corollæ oblongis.—Mich. fl. amer. 1. p. 128.*

In Lower Carolina. *Michaux.* ♀. +. Flowers yellow, as the other species, very small.

2. *L. glaberrima, elata; foliis ovali-lanceolatis oppositis punctatis, racemo terminali longissimo laxo, laciniis corollæ oblongo-ovalibus.—Mich. fl. amer. 1. p. 128.* *racemosa.*

*L. vulgaris. Walt. fl. car. p. 92.*

*L. stricta. Ait. kew. 1. p. 199.*

*L. bulbifera. Bot. mag. 104.*

*Icon. Bot. mag. 104. Pluk. alm. t. 428. f. 4.*

In wet meadows, on the side of ditches: Canada to Virginia. ♀. July, Aug. v.v. This species is frequently found viviparous, having oblong bulbs in the axillæ of the leaves; but this is not a general case.

3. *L. glabra; caule simplicissimo punctato, foliis oppositis sessilibus late-lanceolatis acutis punctatis, pedunculis lateralibus elongatis, floribus capitatis congestis.*

*L. thyrsiflora? Mich. fl. amer. 1. p. 127.*

In salt-marshes, near New York. ♀. June, July. v.v. Though nearly approaching to *L. thyrsiflora*, it is certainly a distinct species.

4. *L. pubescens; foliis subsessilibus quaternis quiniske ovalibus acuminatis punctatis, pedunculis quaternis unifloris, corollæ laciniis ovalibus integris.—Willd.*

*sp. pl. 1. p. 818.*

*L. punctata. Walt. fl. car. p. 92.*

*Icon. Pluk. alm. t. 428. f. 4. Lam. illustr. t. 101. f. 2.*

In dry woods: Canada to Carolina. ♀. June, July. v.v. It varies sometimes with smooth leaves and verticillated flowers.

5. *L. glaberrima, ramosissima; foliis oppositis sessilibus li-*

*capitata.*

*quadrifolia.*

*longifolia.*

## PENTANDRIA MONOGYNIA. Lysimachia.

nearibus longissimis : superioribus quaternis, pedunculis quaternis unifloris, corollæ laciniis ovatis acuminatis serrulatis.

*L. quadriflora. Bot. mag. 660.*

In wet woods, near ponds : Pennsylvania to Virginia.

♀. June—Aug. v. v. A handsome plant, with showy flowers.

*heterophylla.* 6. *L. glabra, gracilis ; foliis oppositis linearibus sessilibus, basi ciliatis, imis suborbiculatis brevi-petiolatis, floribus cernuis. Mich. fl. amer. 1. p. 127.*

Icon. *Pluk. alm. t. 333. f. 1.*

In wet meadows : Virginia to Georgia ; rare. ♀. July. v. v. Flowers resembling those of *L. ciliata*.

*hybrida.*

7. *L. glabra ; foliis oppositis petiolatis lanceolatis utrinque acutis, petiolis ciliatis, floribus cernuis, corollis calyce brevioribus : laciniis crenulatis.—Mich. fl. amer. 1. p. 126.*

*L. ciliata. Walt. fl. car. p. 92.*

*L. angustifolia. Lam. illustr. 1. p. 440.*

On banks of rivers, where the tide overflows : Pennsylvania to Carolina ; rare. ♀. June—Aug. v. v.

*ciliata.*

8. *L. subpubescens ; foliis oppositis longe petiolatis, subcordato-ovalibus margine pubescentibus, petiolis ciliatis, pedicellis subgeminis, floribus cernuis, corollæ laciniis rotundatis, acuminatis crenatis.—Mich. fl. amer. 1. p. 126.*

Icon. *Walth. hort. t. 12.*

In shady woods, in rich soil. ♀. June—Aug. v. v.

## 168. DODECATHEON. Gen. pl. 261.

*Meadia.*

1. *D. foliis oblongo-ovalibus repando-dentatis, umbellis multifloris laxis, bracteis ovalibus.—Willd. sp. pl. 1. p. 808.*

Icon. *Bot. mag. 12. Lam. illustr. t. 99.*

On river sides, in rocky shady situations : Pennsylvania to Virginia, and on the Missouri. ♀. May. v. v. Flowers extremely handsome, purple.

*integrifolium.*

2. *D. foliis subspathulatis integerrimis, umbellis paucifloris strictis, bracteis linearibus. Mich. fl. amer. 1. p. 123.*

Icon. *Pluk. alm. t. 79. f. 6.*

In the Allegany mountains, in shady woods near rivers. ♀. June. v. v. Flowers pale blue, smaller than the former species.

169

1. *P. pusilla, g. scapo elong. flexo, lacini oblonga ex On lake Mist*

170.

1. *A. tenuissim spathulatis calyce bre On the bank Herb. Nuc cording-to A. B. Lan*

171. SABB

1. *S. debilis; ra ticus, calyc laciniis co t. 32.*

*Chironia gra Chironia car In wet pastu Pensylvan rose-colou*

2. *S. erecta ; ceolatis ac laciniis co Icon. Bartr In salt ma Aug. v. rose colou which is is extrem for that p*

3. *S. erecta ; gato-cory lanceolat Chironia an In shady w ♂. July*

169. PRIMULA. *Gen. pl. 258.*

1. *P. pusilla*, glabra ; foliis ovali-spathulatis subdentatis, *mistassinica*.  
scapo elongato, umbella pauciflora, corollæ limbo reflexo, laciniis cuneato-oblongis obtuse bifidis, capsula oblonga exerta. *Mich. fl. amer. 1. p. 124.*  
On lake Mistassins, and Hudson's Bay. *Michaux. 4. +.*

170. ANDROSACE. *Gen. pl. 257.*

1. *A. tenuissime pubescens* ; foliis involucrivo oblongo-spathulatis integerrimis, perianthiis angulatis, capsulis calyce brevioribus. *occidentalis.*

On the banks of the Missouri. *Nuttall. ♂. v. s. in Herb. Nuttall.* Approaches near to *A. elongata*, according to specimens of Pallas's in the Herbarium of A. B. Lambert, Esq.

171. SABBATIA. *Adanson. Parad. lond. t. 32.*

1. *S. debilis* ; ramis laxis elongatis 1-floris, foliis linear-ellipticis, calycis laciniis linearibus corollam subæquantibus, laciniis corollæ obovatis, caule angulato.—*Parad. lond. t. 32.*

*Chironia gracilis. Mich. fl. amer. 1. p. 146.*

*Chironia campanulata. Linn. sp. pl. 272.*

In wet pastures, on the banks of rivers : Canada. *Kalm.* Pennsylvania to Carolina. ♂. July, Aug. v. v. Flowers rose-coloured, segments of the corolla obovate.

2. *S. erecta* ; ramis dichotomis elongatis 1-floris, foliis lanceolatis acutis, calyce subulato corollæ semibreviore, laciniis corollæ obovatis, caule tereti.

Icon. *Bartram ic. ined. t. 13. in Musæo Banksiano.*

In salt marshes : New York, New Jersey, &c. ♂.  
Aug. v. v. The flowers are large and of a beautiful rose colour, with an elegant yellow star in the centre, which is surrounded by a deep red border ; this plant is extremely shy of cultivation, though very desirable for that purpose. It varies with white flowers.

3. *S. erecta* ; foliis ovatis amplexicaulibus, pedunculis elongato-corymbosis, calyce corollæ semibreviore, laciniis lanceolatis, caule marginato-quadrangulo.

*Chironia angularis. Willd. sp. pl. 1. p. 1067.*

In shady woods, on grassy hills : Canada to Carolina. ♂. July, Aug. v. v. It varies with longer and

*gracilis.*

*stellaris.*

*angularis.*

PENTANDRIA MONOGYNIA. *Sabbatia.**calycosa.*

- shorter, broader and narrower leaves. Flowers a purplish red. It is used by the inhabitants in the same way as *Chironia Centaurium* is used in Europe.
4. *S. erecta, foliosa; foliis oblongo-obovalibus, floribus solitariis subseptempartitis, calyce foliaceo corollam superante, laciis oblanceolatis.* *Mich. fl. amer.* 1. p. 147.

*Chironia dichotoma?* *Walt. fl. car.* 95.*Icon. Bartram. ic. ined. t. 10. in Musæo Banksiano.*

In wet meadows: New York to Carolina. ♂. Aug. v. v. Flowers dark rose-coloured, segments of the corolla obovate.

*chloroides.*

5. *S. debilis; foliis lanceolatis erectis, ramis paucis unifloris, floribus 7—13-partitis amplioribus, laciis calycis linearibus corollæ brevioribus.*  
*Chironia chloroides.* *Mich. fl. amer.* 1. p. 147.  
*Chironia dodecandra.* *Walt. fl. car.* 95.  
*Chlora dodecandra.* *Linn. syst. veg.* 299.  
In bogs: New York to Carolina. ♂. Aug. Sept. v. v. Flowers rose-coloured, large.

*paniculata.*

6. *S. erecta; foliis lanceolato-linearibus, panicula multiflora brachiata subfastigiata, calyce subulato corolla triplo-breviore, caule quadrangulo.*  
*Chironia paniculata.* *Mich. fl. amer.* 1. p. 146.

*Swertia difformis.* *Willd. sp. pl.* 1. p. 1330.*latifolia.*

- a. *S. foliis brevi-ovatis, paniculis paucifloris.*  
*Chironia lanceolata.* *Walt. fl. car.* 95.

*angustifolia.*

- β. *S. foliis linearibus, panicula corymbosa multiflora.*  
In bogs and cedar swamps: New Jersey and Carolina. ♀. July, Aug. v. v. Flowers white. This certainly is the long lost *Swertia difformis*, as the specimens in the Herbarium of Clayton, now in possession of Sir Joseph Banks, sufficiently prove.

172. HOTTONIA. *Gen. pl.* 265.*palustris.*

1. *H. floribus verticillatis subsessilibus, caule geniculato, geniculis inflatis.* *Willd. sp. pl.* 1. p. 812.  
In stagnant waters of New Jersey and Virginia. ♀. July. v. v. Whether this is only an accidental variety or a distinct species, I cannot now decide; but certainly the habit of the plant, as well as its smaller flowers, indicate that it may, by further examination, turn out a new species.

173. MI

1. *M. foliis ternata.*  
*Icon. Engl. b. f. 1.*  
In bogs and stada, ♀. J handsome.

174. VI

1. *V. foliis reniantibus, choix de pl.*  
*V. aquatica.*  
*Menyanthes*  
*Anonymos a.*  
In ponds and ♀. June,

175

1. *S. caule tetricum.*  
*Icon. Bot.*  
On the edge Carolina. nial, with under the shops, an children,

176.

1. *O. caule ac dunculatum.*  
*pl. 1. p.*  
*Cynoconium*  
*Anonymos*  
β. *C. petiolatum.*  
*Icon. Swart*  
In dry grav to Carol I strongly are two

173. MENYANTHES. *Gen. pl.* 299.

1. *M. foliis ternatis.* *Willd. sp. pl.* 1. p. 811.  
Icon. *Engl. bot.* 495. *Fl. dan.* 541. *Lam. illustr.* t. 100.  
*f. 1.*  
In bogs and stagnant waters: New Jersey, Virginia, Canada,  $\mathcal{U}$ . June, July. *v. v.* Flowers pale red, very handsome.

*trifoliata.*174. VILLARSIA. *Gmel. syst. veg.* 447.

1. *V. foliis reniformibus subpeltatis, subtus lacunosis nantibus, petiolis floriferis, corollis glabris.* *Vent. choix de plant.* p. 9.  
*V. aquatica.* *Gmel. syst. veg.* 447.  
*Menyanthes trachysperma.* *Mich. fl. amer.* 1. p. 126.  
*Anonymos aquatica.* *Walt. fl. amer.* 109.  
In ponds and lakes, floating: New Jersey to Carolina.  
 $\mathcal{U}$ . June, July. *v. v.* Flowers white.

*lacunosa.*175. SPIGELIA. *Gen. pl.* 272.

1. *S. caule tetragonum, foliis omnibus oppositis.* *Willd.* *marilandica.*  
Icon. *Bot. mag.* 80.  
On the edges of woods, in rich moist soil: Maryland to Carolina.  $\mathcal{U}$ . July, Aug. *v. v.* A beautiful perennial, with large scarlet flowers. The roots are sold, under the name of *Worm-grass* or *Pink-root*, in the shops, and are really a most excellent medicine for children, when troubled with worms.

176. OPHIORRHIZA. *Gen. pl.* 273.

1. *O. caule acutangulo, foliis sessilibus, cyma longius pendunculata.* *Mich. fl. amer.* 1. p. 148. *Willd. sp. pl.* 1. p. 826.  
*Cynoctonum sessilifolium.* *Gmel. syst. veg.* 443.  
*Anonymos sessilifolia.* *Walt. fl. car.* 108.  
2. *C. petiolatum.* *Gmel.* & *Anonymos petiolata.* *Walt.*  
Icon. *Swartz obs.* t. 3. f. 2. *Lam. illustr.* t. 107. f. 1.  
In dry gravelly fields, on wood sides: Lower Virginia to Carolina.  $\mathcal{U}$ . July. *v. v.* Flowers pale blue, small.  
I strongly suspect that Swartz's plant and Michaux's are two distinct species.

*Mitreola.*

177. PHACELIA. *Juss. Gen. pl. 144.*

*bipinnatifida.* 1. *P. erecta*; foliis pinnatifidis, laciniis inciso-lobatis, racemis plerumque bifidis oblongis multifloris, laciniis corollæ integerrimis.—*Mich. fl. amer. 1. p. 134.*

Icon. *I. c. t. 16.*

In the western countries, principally on the Allegany mountains: Pennsylvania to Virginia. May, June. v. v. Flowers blue.

*heterophylla.* 2. *P. erecta*, hispida; foliis petiolatis pinnatifidis, laciniis lanceolatis integerrimis nervosis: impari elongata: floralibus simplicibus lanceolatis petiolatis, racemis bifidis densifloris, laciniis corollæ integerrimis.

On dry hills on the banks of the Kooskoosky. *M. Lewis. J. June, July. v. s. in Herb. Lewis.* Flowers pale blue.

*Caulis* erectus, ramosus: rami elongati. *Folia* petiolata: inferiora pinnatifida: laciniis lateralibus lanceolatis, 1-2-jugis: lacinia terminali longo-lanceolata: superiora et ramea simplicia, lanceolata. *Petoli* marginati. *Racemi* terminales pedunculati dichotomi, spirales, secundi. *Pedicelli* brevissimi. *Calyx* laciniis linearibus. *Corolla* subcampanulata, calyx duplo longior: laciniis oblongis, obtusis, integerrimis. *Filamenta* corollæ duplo longiora. *Stylus* staminibus longior, bifidus.

*fimbriata.* 3. *P. assurgens*, pilosa; foliis sessilibus pinnatifidis: laciniis lanceolatis acutis integris supra hispido-pilosis, subtus glabris: racemis solitariis, pedicellis elongatis, laciniis corollæ ciliato-fimbriatis.—*Mich. fl. amer. 1. p. 134. excl. syn. Pluk.*

On high mountains in Carolina. *Michaux.* Near Harper's Ferry, on the Potowmac; collected by myself. ☺. May, June. v. v. Flowers beautiful blue. Michaux mentions white flowers, which I consider a mistake, though there may be a white variety.

*parviflora.* 4. *P. diffusa*, pubescens; foliis subsessilibus pinnatifidis: laciniis oblongis obtusiusculis integris, racemis solitariis, pedicellis brevibus, laciniis corollæ rotundatis integerrimis.

*Polemonium* foliis inferioribus hastatis, superioribus lanceolatis. *Gronov. fl. virg. p. 29.*

*Polemonium dubium.* *Willd. sp. pl. 1. p. 887.*  
Icon. *Pluk. alni. t. 245. f. 5.*

PENTA

On rocks, ne  
May. v. v.  
size of the

17

1. *Ellisia. Willd.*  
Icon. *Lam. 1.*  
On the bank  
June. v.

17

1. *D. pericarpii*  
*Willd. sp.*  
Icon. *Fl. da*  
Among rub  
July. v.  
2. *D. pericarpii*  
dentatis.  
On road side  
July. v.

180.

1. *H. foliis a*  
*Willd. sp.*  
Icon. *Eng*  
On way sid  
York, ne  
from Eu  
low, wit

181

1. *N. foliis la*  
ribus par  
Icon. *Blac*  
In cultivate  
introduc  
2. *N. foliis o*  
ramulor  
bus, lac  
4-valvib  
Cultivated  
among  
v. v.; v  
white, v

On rocks, near Harper's Ferry, on the Potowmac. ⊖.  
May. v. v. Flowers pale blue, not one fourth the size of the former species.

## 178. ELLISIA. Gen. pl. 268.

1. Ellisia. Willd. sp. pl. 1. p. 815. *Nyctelea.*  
Icon. Lam. illustr. t. 97. Moris. hist. 3, s. 11. t. 28. f. 3.  
On the banks of the Potowmac : Virginia. ⊖ May, June. v. v. Flowers white, very small.

## 179. DATURA. Gen. pl. 332.

1. D. pericarpiis spinosis erectis ovatis, foliis ovatis glabris. *Stramonium.*  
Willd. sp. pl. 1. p. 1008.  
Icon. Fl. dan. 436. Engl. bot. 1288. Lam. illustr. 123.  
Among rubbish and on road sides : common. ⊖ June, July. v. v. Flowers white.
2. D. pericarpiis spinosis erectis ovatis, foliis cordatis glabris dentatis. *Tatula.*  
Willd. sp. pl. 1. p. 1008.  
On road sides ; not so frequent as the former. ⊖ June, July. v. v. Flowers blue.

## 180. HYOSCYAMUS. Gen. pl. 333.

1. H. foliis amplexicaulis sinuatis, floribus sessilibus. *niger.*  
Willd. sp. pl. 1. p. 1010.  
Icon. Engl. bot. 52.  
On way sides and among rubbish : Pennsylvania, New York, near Oswego town, &c. Probably introduced from Europe. ⊖ June, July. v. v. Flowers yellow, with purple veins.

## 181. NICOTIANA. Gen. pl. 334.

1. N. foliis lanceolato-ovatis sessilibus decurrentibus, floribus paniculatis acutis. *Tabacum.*  
Willd. sp. pl. 1. p. 1014.  
Icon. Blackw. t. 146. Lam. illustr. 113.  
In cultivated grounds, frequently found in a wild state ; introduced. ⊖ July. v. v. Flowers red.
2. N. foliis oblongo-ovatis petiolatis, floribus in summitate quadrivalvis. *ramulorum sparsis solitariis, corollis infundibuliformibus, laciis oblongis acutiusculis, capsulis subglobosis 4-valvibus.*  
Cultivated and spontaneous on the Missouri ; principally among the Mandan and Ricara nations. ⊖ July. v. v.; v. s. in Herb. Lewis. nec non Nuttall. Flowers white, with a tinge of blue. The tobacco prepared

## PENTANDRIA MONOGYNIA. Nicotiana.

from it is excellent. The most delicate tobacco is prepared by the Indians from the dried flowers.

## 182. VERBASCUM. Gen. pl. 331.

*Thapsus.*

1. V. foliis decurrentibus utrinque tomentosis, caule simplici. Willd. sp. pl. 1. p. 1001.

Icon. Engl. bot. 549. Fl. dan. 631.

On road sides and in neglected fields; common: probably introduced. ♂. July, Aug. v. v. A singular circumstance frequently struck my attention, respecting this plant, as it appears in great abundance in fields newly cleared and burnt in the most remote part of the country, where sometimes not a plant of it is found within the compass of more than a hundred miles. How the seed is brought there I cannot imagine.

*Lychnitis.*

2. V. foliis ovatis acutis crenatis subtus tomentoso-incanis, spicis laxis lateralibus et terminalibus.—Willd. sp. pl. 1. p. 1003.

Icon. Fl. dan. 586.

On the banks of the Delaware near Philadelphia. ♂. June, July. v. v. Flowers yellow, as all the rest here mentioned, smaller than the former species.

*Blattaria.*

3. V. foliis amplexicaulis obovato-oblongis serratis glabris, pedicellis solitariis unifloris.—Willd. sp. pl. 1. p. 1005.

Icon. Engl. bot. 393.

In fields and on road sides; frequent. ♂. June, July. v. v. Flowers yellow. There frequently occurs a variety with white flowers, which are larger and sweet scented.

*Claytoni.*

4. V. foliis amplexicaulis sublanceolato-oblongis: caulinis inciso-crenatis: crenis crenulatis, pedicellis solitariis unifloris.—Mich. fl. amer. 1. p. 148.

In woods and fields: Virginia to Carolina. ♂. June, July. v. v. Flowers yellow, and larger than the third species.

## 183. CALYSTEGIA. Brown fl. nov. holl. p. 483.

*Sepium.*

1. C. volubilis; foliis sagittatis acutissimis: postice obtusis truncatis integris, bracteis acutis calyce longioribus dimidio corollæ brevioribus, pedunculo angulato petiolum superante. Brown l. c.

## PENTAD

Convolvulus  
Convolvulus  
Brown.  
Convolvulus  
Convolvulus  
perenni. G  
Icon. Engl. l  
In hedges and  
Canada to  
pale red,  
son, as figu

2. C. erecta, to  
natis cord  
elongatis,  
ceolatis, c  
Convolvulus  
On dry rocl  
June, July  
3. C. erecta, pi  
pedunculi  
acutis, cat  
Convolvulus  
On dry hills  
July. v.

## 184. C

1. C. humifusi  
aribus ob  
dunculis  
campanul  
p. 137.

α. C. filiformi  
peduncul  
1. p. 453

C. aquaticu

β. C. volubilis  
subsessili  
Lam. illi

C. humistrata

Icon. Pluk

In sandy w

v. v. F

2. C. repens,

- Convolvulus Sepium. *Willd. sp. pl. 1. p. 844.*  
 Convolvulus repens. *Linn. sp. pl. p. 158.* inonente Cl.  
 Brown.  
 Convolvulus repens. *Mich. fl. amer. 1. p. 137.*  
 Convolvulus lactescens foliis sagittatis, radice longa alba  
 perenni. *Gron. fl. virg. p. 28.*  
 Icon. *Engl. bot. 313.* *Fl. dan. 458.*  
 In hedges and bushes, sometimes trailing on the ground:  
 Canada to Carolina. 2. June, July. v. v. Flowers  
 pale red, white with a red border or a beautiful crim-  
 son, as figured in the Botanical Magazine, No. 732.  
 2. C. erecta, tomentosa; foliis oblongo-lanceolatis acumi-  
 natis cordatis: postice obtusis, pedunculis unifloris  
 elongatis, bracteis ovatis acutis, laciniis calycis lan-  
 ceolatis, caule inferne florifero. *tomentosa.*  
 Convolvulus stans. *Mich. fl. amer. 1. p. 136.*  
 On dry rocky hills: Canada to Virginia; rare. 2.  
 June, July. v. v. Flowers white, large.  
 3. C. erecta, pubescens; foliis subcordato-ovalibus obtusis,  
 pedunculis unifloris foliis brevioribus, bracteis ovatis  
 acutis, caule superne florifero. *spithamea.*  
 Convolvulus spithameus. *Willd. sp. pl. 1. p. 873.*  
 On dry hills: Pennsylvania to Carolina; rare. 2. June,  
 July. v. v. Flowers white.

184. CONVOLVULUS. *Gen. pl. 287.*

1. C. humifusus, pubescens; foliis lanceolato-oblongis line- *trichosanthes.*  
 aribus oblongis obtusis mucronatis subsessilibus, pe-  
 dunculis foliis longioribus 1—5-floris, corolla brevi  
 campanulata, stylo bipartito.—*Mich. fl. amer. 1.*  
 p. 137.  
 a. C. filiformis, suberectus; foliis linearibus mucronatis,  
 pedunculis patentissimis folio longioribus. *Lam. illustr.*  
 1. p. 453. *patens.*  
 C. aquaticus. *Walt. fl. car. 94.*  
 β. C. volubilis; foliis oblongo-ellipticis obtusis mucronatis  
 subsessilibus, pedunculis foliis longioribus subbifloris.  
*Lam. illustr. 1. p. 459.*  
 C. humistratus. *Walt. fl. car. p. 94.*  
 Icon. *Pluk. alm. t. 160. f. 4.*  
 In sandy woods: Virginia to Georgia. ①. June, July.  
 v. v. Flowers white, small.  
 2. C. repens, tuberosa; foliis cordatis hastatis anguloso-lo- *Batatas.*

## PENTANDRIA MONOGYNIA. Convolvulus.

- batis quinquenervibus glabriusculis, pedunculis longis, floribus fasciculatis, calycis glabri laciiniis lanceolatis acuminatis, corolla campanulata.—*Mich. fl. amer.* 1. p. 138. *Willd. sp. pl.* 1. p. 853.  
Cultivated throughout America. ♀. July. v. v. Flowers white, red, and sometimes purple. Vid. *Catesb. car.* 2. t. 60. There are a number of varieties, as is the case with most cultivated plants.
- obtusilobus.* 3. *C. prostratus*, glaber; foliis crassis sinuato-lobatis, lobis rotundato-obtusis: impari majore emarginato, pedunculis unifloris, laciiniis calycis oblongo-ovalibus, corolla brevi-campanulata.—*Mich. fl. amer.* 1. p. 139.  
On the sandy beach of Georgia and Florida. *Michaux.* ♀. +.
- sagittifolius.* 4. *C. volubilis*, glaberrimus; foliis oblongo-sagittatis: sing. profundissimo; auriculis subacuminatis, pedunculis unifloris, laciiniis calycis rotundato-ovalibus, corolla infundibuliformi-campanulata.—*Mich. fl. amer.* 1. p. 138.  
*C. speciosus.* *Walt. fl. car.* p. 93.  
Icon. *Catesb. car.* 1. t. 35.  
In wet situations, among bushes: Virginia to Carolina. ♀. June, July. v. v. Flowers of a beautiful rose-colour, large.
- panduratus.* 5. *C. volubilis*, pubescens; foliis lato-cordatis integris lobatisve panduriformibus, pedunculis longis, floribus fasciculatis, calycibus glabris muticis, corollis subulato-campanulatis. *Mich. fl. amer.* 1. p. 138. *Willd. sp. pl.* 1. p. 850.  
Icon. *Dill. elth. t.* 85. f. 99.  
In sandy fields and among bushes: Canada to Carolina. ♀. June—Aug. v. v. Flowers white, with a red bottom. There is a variety with double flowers, cultivated in the gardens of America, which is a singular circumstance in this genus. The root in some instances grows to an enormous size; I have seen one the thickness of a man's thigh, and more than two feet long. Taken in the place of Jalap, it has the same effect, only in a less degree.
- arvensis.* 6. *C. volubilis*, glaber; foliis sagittatis utrinque acutis, pedunculis subunifloris, bracteis minutis a flore remotis. *Smith fl. brit.* 1. p. 232.  
Icon. *Engl. bot.* 312. *Fl. dan.* 459.  
In meadows: Virginia. *Gron. fl. virg.* 27. ♀. June, July. Flowers rose-coloured. I never observed this

PENTAN

species in  
on the aut

18

1. *I. foliis pin-*  
rollis tubi-  
Icon. *Bot.* 1.  
In cultivated  
lina. ◎  
a most de
2. *I. pubescen-*  
dunculis  
bulosis :  
*Willd. sp.*  
Icon. *Bot.*  
On banks c  
◎. June
3. *I. caule,* P  
bris 7-pa  
floris, la  
Icon. *Jacq*  
Convolvuli  
*fl. amer.*  
On calcare  
♀. h.
4. *I. glabrius-*  
tegrisve,  
lanceola  
Convolvuli  
Icon. *Cate*  
In hedges  
pale pu
5. *I. glabra;*  
angulat  
pilosis,  
*Mich. f*  
Icon. *Dil*  
In Caroli  
v. v. 1
6. *I. glaberr*  
dunculi  
visis : 1

is longis,  
incolatis  
amer. 1.  
  
Flowers  
b. car. 2.  
is the  
, lobis ro-  
peduncu-  
, corolla  
9.  
Michaux.  
  
tis : sing  
culis uni-  
corolla in-  
1. p. 138.  
  
Carolina,  
rose-co-  
  
gris loba-  
ibus fas-  
subulato-  
Willd. sp.  
  
Carolina,  
th a red  
ers, cul-  
singular  
ome in-  
seen one  
han two  
the same  
  
e acutis,  
re remo-  
  
4. June,  
rved this

species in any part of North America, and take it only  
on the authority of Gronovius.

185. IPOMŒA. *Gen. pl. 288.*

1. I. foliis pinnatifidis linearibus, floribus subsolitariis, corollis tubulosis. *Willd. sp. pl. 1. p. 879.* *Quamoclit.*  
Icon. *Bot. mag. 244.*  
In cultivated grounds, near gardens: Virginia and Carolina. ☽. July, Aug. v. v. Flowers scarlet or white; a most delicate and handsome plant.
2. I. pubescens; foliis cordatis acuminatis subangulatis, pedunculis sub-5-floris, calycibus aristatis, corollis tubulosis: limbo subintegro.—*Mich. fl. amer. 1. p. 140.* *Willd. sp. pl. 1. p. 880.*  
Icon. *Bot. rep. t. 99. Bot. mag. 221.*  
On banks of rivers: Virginia to Georgia, and Louisiana. ☽. June, July. v. v. Flowers yellowish-scarlet.
3. I. caule, petiolis pedunculisque pilosissimis; foliis glabris 7-partito-lobatis: lacinij sinuatis, pedunculis unifloris, lacinij calycis ovalibus, corollis campanulatis.  
Icon. *Jacq. hort. t. 159. Willd. phytog. 1. t. 2. f. 3.*  
*Convolvulus dissectus.* *Willd. sp. pl. 1. p. 864. Mich. fl. amer. 1. p. 139.*  
On calcareous hills: Georgia and Florida. *Michaux.* ♀. ½ July. v. v. in *Hortis.*  
*dissecta.*
4. I. glabriuscula; foliis profunde sublanceolato-3-lobis integris, pedunculis subbifloris, lacinij calycis oval-lanceolatis pilosis, corollis brevibus, capsulis pilosis.  
*Convolvulus carolinus.* *Mich. fl. amer. 1. p. 139.*  
Icon. *Catesb. car. t. 87?*  
In hedges: Carolina. ☽. June—Aug. v. v. Flowers pale purple.  
*carolina.*
5. I. glabra; foliis cordatis acuminatis scrobiculatis basi angulatis, pedunculis brevibus subunifloris, calycibus pilosis, corollis tubulosis brevibus, capsulis pilosis.—*Mich. fl. amer. 1. p. 140. Willd. sp. pl. 1. p. 881.*  
Icon. *Dill. eth. t. 87. f. 102.*  
In Carolina and Florida. *Michaux.* ☽. June, July. v. v. Flowers white, with a purple rim.  
*lacunosa.*
6. I. glaberrima; foliis cordatis integris seu angulatis, pedunculis 1—3-floris, calycibus aristatis, corollis indivisis: tubo longissimo.—*Willd. sp. pl. 1. p. 882.*  
*Bona nox.*

- Icon. *Bot. mag. t. 752.* *Jacq. hort. 1. t. 36.*  
On river sides: Carolina and Florida. ☽. July, Aug.  
*v. v.* Flowers white, very large.
- tamnifolia.* 7. I. *hirsuta*; foliis cordatis acuminatis, floribus aggregato-capitatis involucratis: bracteis linearibus, corollis brevi-tubulosis: limbo patente, capsulis glabris.—  
*Willd. sp. pl. 1. p. 885.*  
Icon. *Dill. elth. t. 318. f. 414.*  
In hedges: Carolina. ☽. July, Aug. *v. v. in Hortis.*  
Flowers blue, very small.
- purpurea.* 8. I. *pubescens*; foliis cordatis integris, pedunculis 2—5-floris: pedicellis cernuis incrassatis, laciniis calycis lanceolatis, capsulis glabris.  
*Convolvulus purpureus.* *Willd. sp. pl. 1. p. 852.*  
Icon. *Bot. mag. 113.*  
About gardens, frequent. ☽. June—Aug. *v. v.* Flowers very variable in colour.
- Nil.* 9. I. *hirsuta*; foliis ventricoso-trilobis, pedunculis brevibus 1—2-floris, bracteis subulatis, calycibus villosissimis longissime acuminatis.  
*Convolvulus Nil.* *Willd. sp. pl. 1. p. 851.* *Mich. fl. amer. 1. p. 139.*  
Icon. *Bot. mag. t. 188.* *Dill. elth. t. 80. f. 91 et 92.*  
Near gardens and in hedges, on river sides: Virginia to Carolina. ☽. June—Aug. *v. v.* Flowers beautiful pale blue, only open early in the morning, from which it has been called *Morning-glory.*
10. I. *glabra*; foliis eximie cordatis margine subciliatis, pedunculis longis unifloris, calycis ebracteati laciniis lato-ovalibus obtusis, corollis tubulato-campanulatis.  
*Convolvulus ciliosus.* *Mich. fl. amer. 1. p. 137.*  
Near Knoxville in Tennessee. *Michaux.* +.
- ciliosa.* 11. I. *pubescens*; foliis cordatis integris lobatisque plicato-rugosis subtus subtomentosis, pedunculis 1—3-floris, foliolis calycis ovalibus minutis, corollis campanulatis, seminibus prolixe lanuginosis, radice crassissimo.  
*Convolvulus Jalapa.* *Willd. sp. pl. 1. p. 860.*  
*Ipomoea macrorhiza.* *Mich. fl. amer. 1. p. 141.*  
Icon. *Desfont. in ann. mus. hist. nat. 2. t. 40 et 41.*  
Near the sea shore of Georgia and Florida. ♀. June, July. *v. v.* Flowers white, very large.  
*Obs.* — This very interesting plant was first discovered in Georgia by Michaux. After this I frequently have received seeds and roots from Georgia myself; but

little di  
lapa, t  
The fir  
nales du  
and the  
raised c  
ico, in  
every  
Linnae  
with o  
a light  
family

1. C. glabri  
laxe n  
subrotu  
sp. pl.  
Polemoni  
Ipomoea  
Ipomops  
Gilia cor  
Icon. Ba  
f. 312.  
On river  
*v. v.*
2. C. pubes  
minali  
silibus  
tis, st  
On the b  
*v. s.* is  
than t  
not sp  
the tu
1. D. foliis  
*Salisb*  
D. lappo  
Icon. F  
On the  
shire.  
Flow

little did I suspect it to be the true *Convolvulus Jalapa*, till two circumstances convinced me thereof. The first was seeing a paper on this subject in the *Annales du Muséum d'Histoire Naturelle*, by Desfontaines; and the other, on seeing the living plant recently raised out of a collection of seeds brought from Mexico, in possession of A. B. Lambert, Esq., which in every respect proved to be *Convolvulus Jalapa* of Linnæus, as well as *Ipomoea macrorhiza* of Michaux, with only the small difference of colour, which was a light purple: but this is of no consequence in this family of plants.

186. CANTUA. *Juss. gen. pl. 152.*

1. *C. glabriuscula*; foliis pinnatifidis, floribus terminalibus *coronopifolia*.  
laxe racemosis pedunculatis, lacinij corollæ amplis  
subrotundis brevi-acuminatis, stylo exerto.—*Willd.*  
*sp. pl. 1. p. 879.*

*Polemonium rubrum*. *Linn. sp. pl. ed. 3. p. 231.*

*Ipomoea rubra*. *Linn. syst. veg. 171.*

*Ipomopsis elegans*. *Mich. fl. amer. 1. p. 142.*

*Gilia coronopifolia*. *Pers. encycl. 1. p. 187.*

*Icon. Bot. rep. 415. Exot. bot. 13. Dill. eth. t. 241.  
f. 312.*

On river sides, in sandy soil: Carolina. ♂. June, July.  
*v. v.* Flowers beautiful red.

2. *C. pubescens*; foliis tenuissime pinnatifidis, floribus terminalibus lateralibusque aggregato-subcapitatis subsessilibus, laciniis corollæ angustatis oblongis sensim acutis, stylo inclusio brevissimo.

On the banks of the Mississippi. *M. Lewis*. ♂. June.  
*v. s. in Herb. Lewis.* Flowers scarlet, more slender than the former species; the segment of the corolla not spreading, but erect, and small in proportion to the tube.

*aggregata.*

187. DIAPENSIA. *Gen. pl. 255.*

1. *D. folijs spatulatis glabris, antheris obliquis muticis.* *obtusifolia.*  
*Salisb. parad. lond. 104.*
  2. *D. lapponica*. *Willd. sp. pl. 1. p. 795.*  
*Icon. Fl. dan. 47. Bot. mag. 1108.*
- On the highest mountains, among moss: New Hampshire. 4. May—July. *v. s. in Herb. Prof. Peck.*  
Flowers white.

*cuneifolia.*

2. *D. foliis lanceolato-cuneatis inferne pubescentibus, antheris horizontalibus basi rostratis.* *Salisb. parad. lond. 104.*  
*D. americana. Mus. Banks.*  
*Pyxidanthera barbulata. Mich. fl. amer. I. p. 152.*  
*Icon. Mich. l. c. t. 17. Salisb. parad. lond. 104.*  
 On high mountains in North Carolina. *Michaux.* On the White mountains of New Hampshire. *Prof. Peck.* Plentifully on the pine-barrens of New Jersey near the sea coast, &c.  $\frac{1}{2}$ . June—Aug. *v. v.* Flowers small, white.

*panicalata.*

1. *P. erecta, glabra; foliis lanceolatis sensim angustatis planis margine asperis, corymbis paniculatis, corollæ laciniis rotundatis, calycibus aristatis.* — *Willd. sp. pl. I. p. 839.*  
 Icon. *Mill. ic. t. 205. f. 2. Dill. elth. t. 166. f. 203.*  
 In rich moist meadows: Virginia and Lower Carolina.  
 $\frac{1}{2}$ . July—Sept. *v. v.* Flowers purple; stem from three to four feet high.

*undulata.*

2. *P. erecta, glabra; foliis oblongo-lanceolatis subundulatis margine scabris, corymbis paniculatis, corollæ laciniis obovatis subretusis, calycibus aristatis.* — *Willd. sp. pl. I. p. 840.*  
*P. paniculata. Mich. fl. amer. I. p. 144.*  
 In high mountains and meadows of Virginia and Carolina.  $\frac{1}{2}$ . July, Aug. *v. v.* A beautiful plant, resembling the former species, but taller; flowers the same colour.

*pyramidalis.*

3. *P. erecta, glabra; caule scabro, foliis cordato-ovatis acutis, panicula fastigiata pyramidalis, corollæ laciniis cuneato-truncatis, dentibus calycinis suberectis lanceolatis acutis.* — *Smith exot. bot. 2. p. 55.*  
*P. carolina. Walt. fl. car. 96.*  
 Icon. *Smith l. c. t. 87.*  
 In mountain meadows: Pennsylvania to Carolina.  $\frac{1}{2}$ . June—Aug. *v. v.* Flowers beautiful purple; plant not so tall as the two former species.

*latifolia.*

4. *P. erecta, glabra; caule lœvi, foliis cordato-ovatis, floribus fastigiato-corymbosis, corollæ laciniis suborbiculatis, dentibus calycinis lanceolatis vix acuminatis.* — *Mich. fl. amer. I. p. 143.*  
 An eadem cum priori?

In shady w  
Carolina5. *P. erecta;*

latis glat

tiflora,

acutis re

Icon. *Jac*

In wet me

Aug.

spotted

6. *P. erecta;*

lanceola

læ lacu

acutis e

In meadow

Flowers

the fore

7. *P. erecta*

lævibus

corollæ

ceolatis

*P. triflora*Icon. *Bot*

In sandy

Flower

olens.

8. *P. cæspito*

lævibus

ciniis

*Willd.*Icon. *Di*

In wet n

tains

deep p

9. *P. erecta,*

bus,

sinuclat

emargi

tibus.

On the p

*v. s. in*

purple

*alba.*

the wh

In shady wet places, mossy woods, among Sphagnum : Carolina. *Michaux.* 4. +. Flowers purple, large.

5. *P. erecta* ; caule scabro maculato, foliis oblongo-lanceolatis glabris margine asperis, panicula oblonga confertiflora, corollæ laciniis rotundatis, dentibus calycinis acutis recurvatis.—*Willd. sp. pl. 1. p. 840.*

Icon. *Jacq. hort. t. 127.*

In wet meadows : New England to Carolina. 4. July, Aug. v. v. Flowers purplish-violet or purple ; stem spotted with brown dots.

6. *P. erecta* ; caule glaberrimo immaculato, foliis ovato-lanceolatis undique lœvibus, racemo paniculato, corollæ laciniis rotundatis, dentibus calycinis lanceolatis acutis erectiusculis.—*Willd. sp. pl. 1. p. 840.*

In meadows : Canada to Virginia. 4. July, Aug., v. v. Flowers white, sweet-scented ; plant not so tall as the foregoing species.

7. *P. erecta* ; caule pubescente, foliis ovato-lanceolatis lœvibus, corymbis subfastigiatis, ramiulis subtrifloris, corollæ laciniis rotundatis, dentibus calycis glabri lanceolatis erectis.—*Willd. sp. pl. 1. p. 841.*

*P. triflora?* *Mich. fl. amer. 1. p. 143.*

Icon. *Bot. mag. 1344. Mart. dec. 1. t. 10.*

In sandy wet meadows : Carolina. 4. July—Sept. v. v. Flowers beautiful pale purple, the size of *P. suaveolens*.

8. *P. cæspitosa*, assurgens, glabra ; foliis linear-lanceolatis lœvibus, corymbo terminali subfastigiato, corollæ laciniis rotundatis, dentibus calycis mucronatis.—*Willd. sp. pl. 1. p. 841.*

Icon. *Dill. elth. t. 166. f. 202?*

In wet meadows of Lower Carolina, and on the mountains of Virginia. 4. June—Aug. v. v. Flowers deep purple.

9. *P. erecta*, glabra, frutescens, ramosissima ; foliis linearibus, superioribus alternis basi dilatatis, racemis paniculato-corymbosis, corollæ laciniis cuneato-oblongis emarginatis, dentibus calycis subulatis tubum æquantibus.

On the plains of Columbia. *M. Lewis.* 5. May, June. v. s. in *Herb. Lewis.* Flowers white, with a red or purple centre, similar to those of *Vinca rosea*, var. *alba*. They appear in such abundance, that they cover the whole shrub.

*maculata.*

*suaveolens.*

*carolina.*

*glaberrima.*

*speciosa.*

*aristata.*

10. *P. debilis*, erecta, viscidulo-pubescent; foliis linear-lanceolatis, panicula laxa fastigiata, pedicellis subgeminatis, corollæ laciniis obovalibus: tubo curvato pubescente, dentibus calycinis longissimis subulatis.—*Mich. fl. amer.* 1. p. 144.

*virens.*

- P. pilosa*. *Bot. mag.* 1307?  $\alpha$ . corolla roseo-purpurascente.  $\beta$ . corolla albido-rosea.

*canescens.*

In sandy fields: Pennsylvania to Carolina.  $\mathcal{U}$ . May—July. v. v. Flowers red, or sometimes white.

*pilosa.*

11. *P. humilis*, decumbens, pubescens; foliis linear-lanceolatis brevibus asperiusculis: summitatibus subsessiliter fasciculifloris, corollæ laciniis rotundato-obtusis: tubo stricto glabro, dentibus calycis subulato-acuminatis. *Mich. fl. amer.* 1. p. 145. *Walt. fl. car.* 96. *Willd. sp. pl.* 1. p. 840?

*amœna.*

- Icon. *Pluk. alm.* t. 98. f. 1.  $\beta$ . *Bot. mag.* 1308.

In sandy fields: Carolina and on the mountains of Georgia.  $\mathcal{U}$ . May—July. v. v. Flowers purple, sometimes white, smaller than the foregoing species. Var.  $\beta$ . is most likely only a luxuriant garden production of this species.

*divaricata.*

12. *P. humilis*, decumbens, pubescens; foliis ovali-lanceolatis: superioribus alternis, ramulis divaricatis laxis paucifloris, corollæ laciniis subobcordatis, dentibus calycis subulato-linearibus.—*Willd. sp. pl.* 1. p. 841.

Icon. *Bot. mag.* 163. *Mill. ic. t.* 203. f. 1.

On the mountains: Pennsylvania to Virginia.  $\mathcal{U}$ . May—July. v. v. Flowers purplish-blue; plant of low growth.

*ovata.*

13. *P. erecta*, glabra; foliis radicalibus ovatis acutis subcarnosis: caulinis lanceolatis, corymbis subfastigiatis, corollæ laciniis undulatis retusis, dentibus calycis linearibus acutis.—*Bot. mag.* 528, *Willd. sp. pl.* 1. p. 842.

Icon. *Bot. mag. l. c.*

In North America,  $\mathcal{U}$ . May—July. v. v. in Hortis. Flowers beautiful reddish-purple.

*stolonifera.*

14. *P. repanti-stolonifera*, pubescent; foliis radicalibus spatulato-bovatis, caulinis ovali-lanceolatis, corymbo divaricatum paucifloro, corollæ laciniis obovatis; dentibus calycis linearibus reflexis.—*Bot. mag.* 563,

*P. reptans*. *Mich. fl. amer.* 1. p. 144.

Icon. *Bot.*  
In the big  
June—  
centre,

15. *P. cæspit*  
pungen  
trifidis,  
calycis  
 $\text{sp. pl.}$

Icon. *Bot.*  
On dry sa  
 $\mathcal{U}$ . Ap  
a hand

16. *P. cæspit*  
punger  
umbell  
tibus c  
—Wil

Icon. *Bo*  
On rocks  
 $\mathcal{U}$ . Ma  
with a

18  
1. *P. foliis*  
bus.

Icon. *M*  
In the m  
to Car  
blue;

1. *A. subn*  
adulti  
dentil  
viore.

$\alpha$ . *A. floril*  
 $\beta$ . *A. floril*

$\alpha$ . On the  
*Mich*

$\beta$ . On the  
*h. A*

foliis linearibus  
ellis subge-  
curvato pu-  
subulatis.

May—July.

lineari-lan-  
bus subes-  
ndato-obtu-  
s subulato-  
fl car. 96.

mountains of  
ers purple,  
ing species.  
leu produc-

vali-lanceo-  
catis laxis  
, dentibus  
1. p. 841.

4. May  
ant of low

atits subcar-  
bstigiatits,  
s calycis li-  
l. sp. pl. 1.

in Hortis.

calibus spa-  
orymbo di-  
tis ; denti-  
563,

Icon. Bot. mag. 563.  
In the high mountains of Virginia and Carolina. 2.  
June—Sept. v. v. Flowers blue, with a purple  
centre, very handsome.

15. P. cæspitosa, candicanti-pubescentis ; foliis linearibus  
pungentibus ciliatis, corymbis paucifloris : pedicellis  
trifidis, corollæ laciniis cuneatis emarginatis, dentibus  
calycis subulatis tubo corollæ vix brevioribus.—Willd.  
sp. pl. 1. p. 842.

Icon. Bot. mag. 411.  
On dry sandy hills and rocks: New Jersey to Carolina.  
2. April—June. v. v. Flowers pink-coloured, with  
a handsome purple star in the centre.

16. P. cæspitosa, pubescens ; foliis fasciculatis subulatis  
pungentibus ciliatis, pedicellis paucis terminalibus sub-  
umbellatis, corollæ laciniis cuneatis emarginatis, den-  
tibus calycis subulatis tubo corollæ triplo brevioribus.  
—Willd. sp. pl. 1. p. 842.

Icon. Bot. mag. 415.  
On rocks and barren sandy hills: Virginia to Carolina.  
2. May—July. v. v. Flowers pale rose-coloured,  
with a purple star in the centre.

subulata.

setacea.

reptans.

#### 189. POLEMONIUM. Gen. pl. 289.

1. P. foliis pinnatis septenis, floribus terminalibus nutanti-  
bus. Willd. sp. pl. 1. p. 886.

Icon. Mill. ic. 2. t. 209.  
In the mountains, on the side of rich hills: Pennsylvania  
to Carolina. 2. April, May. v. v. Flowers sky-  
blue; there is a variety with white flowers.

#### 190. AZALEA. Gen. pl. 277.

1. A. subnudiflora ; foliis oblongis utrinque pubescentibus : *calendulacea*.  
adultis hirsutis, floribus amplis non viscosis, calycis  
dentibus oblongis, corolla tubo hirsuto laciniis bre-  
viore.—Mich. fl. amer. 1. p. 151.

a. A. floribus flammeo-calendulaceis.

β. A. floribus croceis. Bot. mag. . . .

α. On the banks of the Savannah river, near Two-sisters.  
Michaux. On the Flint river, Georgia, &c.

β. On the highest mountains of Virginia and Carolina.  
2. April, May. v. v. Flowers of var. α. a beautiful

flammea.

crocea.

canescens.

flame colour, sometimes deep yellow mixed with scarlet, as in *Mirabilis Jalapa*; var.  $\beta$ . bright yellow, resembling *A. pontica*. It is, without exception, the handsomest shrub in North America. Vid. *Bartram's Travels*, p. 321.

periclymenoides.

2. *A. subnudiflora*; foliis obovato-oblongis supra pubescentibus subtus tomentosis, nervo non setigero, floribus non viscosis, tubo lacinii vix breviore, calycis dentibus brevissimis rotundato-obtusis, staminibus vix exsertis.—*Mich. fl. amer.* 1. p. 150.

On the banks of rivers in Lower Carolina. *Michaux*. On the mountains of Capoone springs near Winchester, Virginia.  $\frac{1}{2}$ . May, June. *v. v.* Flowers rose-coloured.

coccinea.  
rutilans.

3. *A. subnudiflora*; foliis lanceolato-oblongis utrinque glabriusculis, concordibus, nervo supra lanuginoso, subtus setigero, margine ciliatis, floribus amplis non viscosis, tubo lacinii longiore, calycis dentibus brevibus ovali-subrotundis, staminibus longissime exsertis.—*Mich. fl. amer.* 1. p. 151.

*A. nudiflora*. *Willd. sp. pl.* 1. p. 831.

$\alpha$ . *A. floribus coccineis*, foliis lanceolatis. *Bot. mag.* 180.  
 $\beta$ . *A. floribus saturate rubris*, calycibus minutis. *Ait. kew. 1.* p. 319.

$\gamma$ . *A. floribus pallide rubicundis*, tubo basi rubro, calycibus foliaceis. *Ait. l. c.*

$\delta$ . *A. floribus albidis*, calycibus mediocribus. *Ait. l. c.*

$\epsilon$ . *A. floribus rubicundis*: lacinia infima alba, calycibus foliaceis. *Ait. l. c.*

$\zeta$ . *A. floribus carneis usque ad basin quinquepartitis*. *Ait. l. c.*

$\eta$ . *A. floribus roseis brevibus 10—20-andris*.

On the side of hills: Canada to Georgia. Var.  $\alpha$ . near Savannah, Georgia. Var.  $\eta$ . near Philadelphia, Pennsylvania.  $\frac{1}{2}$ . April—June. *v. v.* This beautiful shrub has a great many varieties, besides the above-mentioned, among which is one with double flowers; but the var.  $\eta$ . is the most remarkable. This variety proves clearly, that *Azalea* and *Rhododendron* are one genus, *A. procumbens* excluded, as *Rhododendron* is frequently found with less than ten, and all the species of *Azalea* with more than five stamens.

arborescens.

4. *A. foliosiflora*; foliis obovatis obtusiusculis utrinque glabris subtus glaucis: nervo vix setigero, margine cilia-

tis, flor  
giore, c  
mentis

*A. arbore*  
On rivule

May  
knowle  
I have  
den of  
father  
to twe  
liege a  
finest  
so muc  
the sc  
brown

5. *A. nudif*  
pubese  
non vi  
vissim  
giore,  
*A. nudif*

*Icon. T*  
On bar

May,  
the re  
deep-

6. *A. foli*  
acutis  
marg  
longi  
jame

*A. 83*

In swa  
July,  
Ther  
speci

7. *A. foli*  
ceola  
nitid  
tis, t  
lycib

In dee  
to V

xed with  
it yellow,  
tion, the  
Bartram's  
pubescen-  
, floribus  
cis denti-  
as vix ex-

*Michaux.*  
Winches-  
vers rose-

unque gla-  
ioso, sub-  
oplis non  
bus brevi-  
s exertis.

ig. 180.  
tit. kew. 1.

calycibus

*L. c.*  
calycibus

nepartitis.

r. α. near  
bia, Pen-  
beautiful  
e above-  
; flowers;  
is variety  
m are one  
dendron is  
I all the  
ia.

unque gla-  
rine cilia-

*A. arborea. Bart. catal.*

On rivulets, near the Blue mountains : Pensylvania.  $\frac{1}{2}$ . May - July. *v. v.* — This beautiful species has, to my knowledge, not yet been introduced into the gardens. I have only seen it in its native place and in the garden of Mr. John Bartram, near Philadelphia, whose father introduced it many years ago. It rises from ten to twenty feet high, and forms, with its elegant foliage and large abundant rose-coloured flowers, the finest ornamental shrub I know. The flowers are not so much pubescent as the rest of the other species ; the scales of the flower-buds are large, yellowish-brown, surrounded with a fringed white border.

5. *A. nudiflora* : foliis oblongis utrinque tenuissime cano-pubescentibus : nervo non setigero, floribus parvis non viscosis : tubo laciniis vix longiore, calycibus brevissimis : lacinia unica linearis reliquis quadruplo longiore, filamentis exertis, ramulis piloso-hispidis.

*A. nudiflora bicolor. Ait. kew. 1. p. 319.*

*Icon. Trew ehret. t. 48.*

On barren, sandy hills : Carolina and Georgia.  $\frac{1}{2}$ . May, June. *v. v.* Flowers slender, and smaller than the rest, of a pale rose colour or nearly white, with a deep-red-coloured tube.

6. *A. foliosiflora* ; ramis hispidis, foliis oblongo-ovatis acutis utrinque glabris concoloribus : nervo setigero, margine ciliatis, floribus viscosis : tubo laciniis duplo longiore, calycis dentibus brevissimis rotundatis, filamentis corollae vix longioribus. — *Willd. sp. pl. 1. p. 831.*

In swamps and shady woods : Canada to Georgia.  $\frac{1}{2}$ . July, Aug. *v. v.* Flowers white, sweet-scented. There are a number of varieties and even intermediate species between this and the *A. periclymenoides*.

7. *A. foliosiflora* ; ramis glabriusculis, foliis parvis oblongo-ovatis submucronatis coriaceis utrinque glabris supra nitidis : nervo subtus setigero, margine revoluto-ciliatis, floribus viscosis : tubo laciniis paulo longiore, calycibus brevissimis, filamentis exertis.

In deep mossy swamps on the mountains : New York to Virginia.  $\frac{1}{2}$ . June, July. *v. v.* Flowers white

*bicolor.*

*viscosa.*

*nitida.*

## PENTANDRIA MONOGYNIA. Azalea.

*glaуca.*

with a red tinge; leaves very dark green and shining; they are smaller than in any other species.

8. A. foliosiflora; ramulis hispidis, foliis oblanceolatis acutis utrinque glabris subitus glaucis: nervo setigero, margine ciliatis, floribus viscosissimis: tubo lacinij duplo longiore, calyce brevissimo, filamentis lacinij corollæ subæquantibus.—*Lam. illustr.* p. 493.

*A. viscosa floribunda.* *Ait. kew.* 1. p. 319.

In swamps of a clayey soil: New England to Virginia.  
J. June, July. v. v. Flowers white, very fragrant; the shrub lower than the rest, and flowering in great abundance.

*hispida.*

9. A. foliosiflora; ramis strictis hispidissimis; foliis longe lanceolatis: supra hispidis, subitus glabris; utrinque glaucis: nervo setigero, margine ciliatis, floribus viscosissimis: tubo lacinij amplis vix longiore, calycis dentibus oblongis rotundatis, filamentis exsertis.

*A. viscosa glauca.* *Ait. kew.* 1. p. 319?

On the borders of lakes on the highest part of the Blue mountains: New York and Pennsylvania. J. July, Aug. v. v. Flowers white with a red border, and a tinge of red on the tube, which makes them appear to be of a rose-colour before open. They have frequently ten stamina, as in var.  $\eta.$  of *A. periclymenoides*. This shrub generally grows to the height of from ten to fifteen feet, very upright, and has a blueish appearance, by which it may be distinguished from all others at a great distance.

*procumbens.*

10. A. foliosiflora; ramis diffuso-procumbentibus, foliis oppositis ellipticis glabris margine revolutis, corollis campanulatis glabris, filamentis inclusis æqualibus.—*Willd. sp. pl.* 1. p. 832.

*Icon. Fl. dan.* 9. *Engl. bot.* 865.

In the alpine regions of the White mountains, New Hampshire. *Prof. Peck.* On the Grandfather mountain, Carolina. *Lyon.* J. July. v. s. in *Herb. Peck.* nec non *Lyon.* v. v. in *Hortis.* Flowers small, rose-coloured. This plant has so much affinity to *Ledum buxifolium*, Ait. that I have scarcely been able to persuade myself that they are distinct plants. Comparing specimens of different varieties of the latter, with those of *A. procumbens* from different countries, in the herbarium of A. B. Lambert, Esq. I could find no other distinction between both, than that of the

*Ledum*  
flower  
has p  
It inc  
or el  
dendr

1. B. spin  
que  
corol  
*Siderox*  
*Siderox*  
Icon. i  
In shad  
v. v.  
flow

2. B. spin  
diva  
Per  
*Sidero*  
Icon.  
On th  
July

3. B. su  
foli  
sis.  
*Sidero*  
*Sidero*  
In we

4. B. sul  
tus  
*Wi*

*Sidero*  
*Sidero*  
*Sidero*  
In he  
Ge  
ha

5. B. ir  
ut  
ax  
On t  
in

shining;  
olatis acu-  
setigero,  
laciiniis  
tis laciiniis  
493.

Virginia.  
fragrant;  
in great

foliis longe  
utrinque  
ribus vis-  
e, calycis  
is.

the Blue  
July,  
er, and a  
m appear-  
have fre-  
clymeno-  
height of  
nd has a  
inguished

foliis op-  
corollis  
alibus.—

ns, New  
er moun-  
erb. Peck.  
all, rose.  
to *Ledum*  
le to per-  
omparing  
er, with  
ntries, in  
ould find  
at of the

*Ledum* being an upright little shrub, with decandrous flowers which are white, whereas the present species has procumbent stems and pentandrous red flowers. It most certainly ought to be taken from this genus, or else all the rest but this one united with *Rhododendron*.

## 191. BUMELIA. Gen. pl. 1736.

1. B. spinosa, erecta; foliis deciduis lato-lanceolatis utrinque glabris, pedunculis aggregatis unifloris, laciiniis corollæ trifidis.—*Mich. fl. amer.* 1. p. 122.

*Sideroxylum lycioides*. *Willd. sp. pl.* 1. p. 1090.

*Sideroxylum læve*. *Walt. fl. car.* 100.

Icon. *Duham. arb.* 2. t. 68.

In shady woods: Carolina and Georgia.  $\frac{1}{2}$ . July, Aug.  
*v. v. s. fl.* A small tree, with minute greenish flowers.

2. B. spinosa, dumosa, diffuse reclinata; ramis sterilibus divaricato-ramosis, foliis parvis obovalibus glaberrimis.  
*Pers. syn.* 1. p. 237.

*Sideroxylum reclinatum*. *Mich. fl. amer.* 1. p. 122.

Icon. *Vent. choix de plant.* t. 22.

On the banks of rivers in Georgia. *Michaux.*  $\frac{1}{2}$ .  
July, Aug. *v. v. s. fl.* A small straggling shrub.

3. B. subspinosa; ramulis patentissimis pubescensibus, foliis ovali-lanceolatis, supra glabris, subtus lanuginosiss.  
*Pers. syn.* 1. p. 237.

*Sideroxylum lanuginosum*. *Mich. fl. amer.* 1. p. 122.

*Sideroxylum tenax*. *Walt. fl. car.* 100.

In wet copses of Georgia. *Michaux.*  $\frac{1}{2}$ . +.

4. B. subspinosa; foliis cuneato-lanceolatis plerumque obtusis, subtus sericeo-nitentibus. *Pers. syn.* 1. p. 237.  
*Willd. sp. pl.* 1. p. 1085.

*Sideroxylum tenax*. *Linn. syst. veg.*

*Sideroxylum sericeum*. *Walt. fl. car.* 100.

*Sideroxylum chrysophylloides*. *Mich. fl. amer.* 1. p. 123.

In hedges and copses near the sea-coast of Carolina and Georgia.  $\frac{1}{2}$ . June. *v. v.* A small tree; leaves very handsome.

5. B. inermis; foliis sempervirentibus oblongo-lanceolatis utrinque acutis glabris subaculeato-serratis, racemis axillaribus sessilibus multifloris.

On the banks of the Missouri. *Nuttall*  $\frac{1}{2}$ . July. *v. s.*  
*in Herb. Nuttall.* Berries black, large.

*lycioides*.

*reclinata*.

*lanuginosa*.

*chrysophyl-  
loides*.

*serrata*

192. SOLANUM. *Gen. pl. 337.*

- Dulcamara.* 1. *S. caule inermi fruticoso scandente, foliis cordatis glabris : superioribus auriculatis, corymbis oppositifoliis.* Willd. *sp. pl. 1. p. 1028.*  
 Icon. *Fl. dan. 607. Schmidt arb. 146. Engl. bot. 565.*  
 On banks of rivers, in hedges : Pennsylvania. Bartram.  
 ½ June, July. v. v. Flowers violet-blue ; berries red.
- nigrum.* 2. *S. caule inermi herbaceo, ramis angulatis dentatis, foliis ovatis reparatis glabris, racemis distichis nutantibus.* Willd. *sp. pl. 1. p. 1035.*  
 Icon. *Dill. elth. t. 275. f. 256.*  
 On cultivated grounds, common. ○. June—Aug. v. v.  
 Flowers white ; berries black.
- mammosum.* 3. *S. caule aculeato herbaceo, foliis cordatis angulato-lobatis utrinque villosis aculeatis.* Willd. *sp. pl. 1. p. 1040.*  
 Icon. *Merian. surin. t. 27. Pluk. alm. t. 226. f. 1.*  
 On the sea-coast of Virginia and Carolina. ○. June—  
 Aug. v. v. Flowers pale blue ; fruit yellow, shaped  
 like a pear.
- virginianum.* 4. *S. caule erecto aculeato, foliis pinnatifidis undique aculeatis, lacinias sinuatis obtusis, margine ciliatis, calycibus aculeatis.* Willd. *sp. pl. 1. p. 1041.*  
 Icon. *Dill. elth. t. 267. f. 346. Pluk. alm. t. 62. f. 3.*  
 On road sides and sandy fields : Virginia to Carolina.  
 ○. July. v. v. Flowers blue ; berries small, greenish-white.
- carolinense.* 5. *S. caule aculeato annuo, foliis hastato-angulatis aculeatis utrinque tectis, racemis laxis.* Willd. *sp. pl. 1. p. 1043.*  
 Icon. *Dill. elth. 269. f. 348. Jacq. ic. 2. t. 331.*  
 On road sides and old fields : Virginia to Georgia. ¼.  
 June—Aug. v. v. Flowers large, pale blue ; fruit yellow.
- heterandrum.* 6. *S. caule aculeato annuo, foliis bipinnatifidis utrinque hirsuto-tomentosis aculeatisque : lacinias obtusiusculis, racemis lateralibus, antheris patulis corniformibus : infima maxima longitudine corollæ, bacca subquadri-loculare calyce echinato inclusa.*  
 On the banks of the Missouri. Nuttall. ○. July. v. s.  
 in *Herb. Nuttall.* ; v. v. in *Hortis.* Flowers large,



*Solanum heterandrum*

Engraved by F. Brush

Engraved by J. Hooker

yellow. I  
nearly allie  
hist. nat. 3  
a new gen  
amined spe

1. P. foliis gemi  
tomentosis  
cibus frue  
p. 1021.  
Icon. Jacq. I  
On road sid  
rolina. 24  
ries viscou
2. P. foliis qua  
liter denta  
ramis ang  
a. P. summitat  
P. angulata.  
β. P. viscido-p  
In sandy fie  
Flowers yo  
thers.
3. P. foliis gen  
sensim ai  
chotomo,  
In Lower C  
Walt.
4. P. foliis ova  
herbaceo  
longioribu  
P. tomentosa  
On road side  
24. June,
5. P. foliis oval  
ramosissim  
vioribus.  
In dry place  
○. July.  
species, y
6. P. foliis villa

yellow. This singular species of *Solanum* is very nearly allied to *S. cornutum*, Juss. in *Annales de Mus. hist. nat.* 3. p. 120. t. 9. and they probably will form a new genus, to which some other not properly examined species may belong.

193. PHYSALIS. *Gen. pl. 336.*

1. *P. foliis geminis subcordato-ovalibus repandis obtusis sub-tomentosis, caule herbaceo superne paniculato, calycibus fructiferis pubescentibus.* — *Willd. sp. pl. 1. p. 1021.*

*Icon. Jacq. hort. 2. t. 136. Dill. elth. t. 10. f. 10.*

On road sides and among rubbish: Pennsylvania to Carolina.  $\frac{1}{2}$ . July—Aug. *v. v.* Flowers yellow; berries viscous.

*viscosa.*

2. *P. foliis quasi cordato-suborbiculatis acuminatis inaequaliter dentatis, caule herbaceo divaricato-ramosissimo: ramis angulatis.* — *Mich. fl. amer. 1. p. 149.*

- a. P. summitatibus petiolisque vix pubescentibus.*

*P. angulata. Walt. fl. car. 99.*

*obscura.*

- b. P. viscido-pubescentis.*

In sandy fields of Lower Carolina.  $\frac{1}{2}$ . July, Aug. *v. v.* Flowers yellow, with brown spots and blueish anthers.

*glabra.*

3. *P. foliis geminis ovali-lanceolatis subintegris, in petiolum sensim angustatis pubescentibus, caule herbaceo dichotomo, calyce villoso.* — *Mich. fl. amer. 1. p. 149.*

In Lower Carolina. *Michaux. 1/2. July. v. s. in Herb. Walt.*

*pubescens.*

4. *P. foliis ovatis subrepandis obtusis nudiusculis, caule pensylvanica.* herbaceo ramoso, pedunculis solitariis petiolis paulo longioribus. — *Willd. sp. pl. 1. p. 1021.*

*P. tomentosa. Walt. fl. car. 99?*

On road sides, among rubbish: Pennsylvania to Carolina.  $\frac{1}{2}$ . June, July. *v. v.* Flowers entirely yellow.

*lanceolata.*

5. *P. foliis ovatis repando-dentatis glabris, caule herbaceo ramosissimo, pedunculis solitariis petiolo multo brevioribus.* — *Lam. encycl. 2. p. 101.*

In dry places on river sides; New England to Virginia.

$\odot$ . July. *v. v.* Flowers larger than the foregoing species, yellow, with brown stripes.

*philadelphi-ca.*

6. *P. foliis villoso-viscosis cordatis, caule herbaceo ramo-*

*pubescens.*

PENTANDRIA MONOGYNIA. *Physalis.*

sissimo, floribus solitariis pendulis, calycibus fructiferis subrotundo-globosis angulatis. Willd. sp. pl. 1. p. 1023.

Icon. *Fevil. peruv.* 3. t. 1.

On the sandy shores of Carolina. ☺. June, July. v. v. Flowers yellow, with brown spots; berries round, yellow.

194. NICANDRA. *Juss. gen. pl.* 140.

*physaloides.*

1. Nicandra. *Pers. syn.* 1. p. 219.

*Atropa physaloides.* Willd. sp. pl. 1. p. 1017.

*Calydermos erosus.* *Fl. peruv.* 2. p. 44.

Icon. *Fevil. peruv.* t. 16.

In cultivated ground, near gardens: Pennsylvania to Virginia; introduced. ☺. July, Aug. v. v. Flowers blue.

195. SAMOLUS. *Gén. pl.* 294.

*Valerandi.*

1. S. foliis obovatis, racemis elongatis, pedicellis medio bracteatis.—Willd. sp. pl. 1. p. 927.

Icon. *Fl. dan.* 198. *Engl. bot.* 703.

In marshes and near springs: Canada to Carolina; frequent. ☺. July, Aug. v. v. Flowers white, small.

196. PINCKNEYA. *Mich. fl. amer.* 1. p. 103.

*pubescens.*

1. P. foliis ovalibus utrinque acutis subtus subtomentosis. *Mich. fl. amer.* 1. p. 105.

Icon. *Mich. l. c. t.* 13.

On the banks of the river St. Mary, Georgia. ☺. July, Aug. v. v. Flowers blush-red, marked with purple lines, in large clusters. Each flower has one of the divisions of the calyx grown out to a large white leaf, tinged with red, which makes this shrub highly ornamental. It is only to be regretted, that it is very shy of cultivation. The small difference of its fruit from *Cinchona* only makes it excusable to keep it separate from that genus; but I have no hesitation in saying, that if *Pinckneya* is not united with *Cinchona*, there will be some of the latter genus found belonging to *Pinckneya*.

1. C. glab  
caulin  
florib  
Icon. F  
On rock  
nada.  
consid
2. C. glab  
liosis,  
viter  
On high  
Sept.
3. C. glab  
serrat  
bus n  
On slate  
Aug.  
ful pe
4. C. glab  
inferi  
rollis  
sp. p  
In shad  
July.  
in th  
two
5. C. glab  
gum  
mult  
p. 10  
C. nitid  
Icon.   
On th  
4..  
semi
6. C. gra  
que  
lanc  
tate  
In low  
Ver  
July

197. CAMPANULA. *Gen. pl. 889.*

1. *C. glabra*; foliis radicalibus reniformi-cordatis crenatis; caulinis linearibus integris, panicula laxa pauciflora, floribus nutantibus.—*Willd. sp. pl. 1. p. 892.*

*Icon. Fl. dan. 855. Engl. bot. 866.*

On rocks on the banks of the Susquehanna and in Canada. ♀. July, Aug. v. v. Flowers azure blue, of considerable size.

2. *C. glabra*; caule erecto-flexuoso, ramulis confertis foliosis, foliis lanceolatis subremote serratis, floribus breviter pedicellatis nutantibus.—*Mich. fl. amer. 1. p. 109.*  
On high mountains: Virginia and Carolina. ♀. Aug. Sept. v. v. Flowers small, pale blue.

*flexuosa.*

3. *C. glabra*, nitida, erecta; foliis membranaceis remote serratis, panicula ramulis subaphyllis divaricatis, floribus nutantibus.—*Mich. fl. amer. 1. p. 109.*

On slate mountains in Virginia and Carolina. ♀. July, Aug. v. v. Flowers small, azure blue; a very graceful perennial.

*divaricata.*

4. *C. glabra*; foliis cordatis lanceolatisque serratis, petiolis inferioribus ciliatis, floribus axillaribus sessilibus, corollis 5-partitis planis, stylis corolla longioribus. *Willd. sp. pl. 1. p. 898.*

In shady wet places: Pennsylvania to Virginia. ♂. June, July. v. v. Flowers small, pale blue, two or three in the axillæ of the leaf, nearly sessile; plant from two to three feet high.

*americana.*

5. *C. glabriuscula*, erecta; foliis ovato-lanceolatis in longum acumen desinentibus subserratis, spica fasciculata-multiflora, corollis subrotatis.—*Mich. fl. amer. 1. p. 108.*

*C. nitida. Ait. kew. 1. p. 346.*

*Icon. Dodart. mem. 4. t. 18.*

On the high mountains of Pennsylvania and Virginia. ♀. July, Aug. v. v. Flowers in size and colour resemble the fourth species.

*acuminata.*

6. *C. gracilis*; caule simplici angulato: angulis foliorum que margine et nervo retrosum aculeatis, foliis linear-lanceolatis supra glabris, pedunculis paucis in summitate caulis flexuosis axillaribus unifloris filiformibus.

In low inundated meadows, generally in company with *Veronica scutellata*: Pennsylvania to Virginia. ♂. July. v. v. Flowers small, pale blue.

*aparinoides.*

*perfoliata.*

7. C. caule simplici erecto, foliis cordatis crenatis amplexicaulis, floribus axillaribus sessilibus glomeratis. *Willd. sp. pl. 1. p. 915.*  
 C. amplexicaulis. *Mich. fl. amer. 1. p. 108.*  
*Icon. Moris. hist. 2. s. 5. t. 2. f. 23. Barr. rar. t. 1133.*  
 In cultivated grounds: common from North to South.  
 ♂. May—Aug. v. v. Flowers small, pale blue.

198. CHIOCOCCA. *Gen. pl. 315.**racemosa.*

1. C. foliis ovato-oblongis acutis planis, racemis axillaribus pedunculatis simplicibus. — *Willd. sp. pl. 1. p. 975.*  
*Icon. Dill. eth. t. 228. f. 295. Bot. rep. t. 284.*  
 On the sea-shore of Georgia and Florida. *Michaux.*  
 ♂. July. v. v. Flowers yellowish-white. It appears to differ in some respects from the West Indian plant, but can be considered nothing more than a variety.

199. CAPRIFOLIUM. *Juss. gen. pl. 235.**semperfiriens.*

1. C. spicæ verticillis subnudis distantibus, corollis subæqualibus: tubo superne ventricoso, foliis obovatis ovatisque subtus glaucis: summis connato-perfoliatis.— *Willd. sp. pl. 1. p. 983. sub Lonicera.*  
*Icon. Bot. mag. 781. Schmidt arb. 104.*  
 In stony dry woods: New York to Carolina. ♂. May—Aug. v. v. Flowers a beautiful scarlet outside and yellow inside; there are several varieties of it, particularly one with an almost upright stem.

*ciliatum.*

2. C. spicæ verticillis approximato-capitatis subsessilibus, corollis subæqualibus: tubo hirsuto medio ventricoso, foliis subamplexicaulibus sessilibus petiolatisque ovatis subtus glaucis, margine conspicue ciliatis: summis connato-perfoliatis.  
 On the banks of the Kooskoosky. *M. Lewis.* ♂. June. v. s. in *Herb. Lewis.* Flowers of a deep yellow.

*Fraseri.*

3. C. spicæ verticillis capitatis, corollis subringentibus: laciiniis oblongis obtusis, foliis ovatis subtus glaucis cartilagineo-marginatis: summis connato-perfoliatis. *Lonicera flava. Bot. mag. 1318.*  
 On the Paris mountains, South Carolina. *Fraser.* On the Catskill mountains, New York, &c. ♂. June, July. v. v. Flowers bright yellow.

4. C. spicæ tubo crona mis co Lonic Icon. H On the situati Sept.  
 5. C. spicæ usculi foliis bracte cycl. 1 Lonicera Lonicera Caprifoli Icon. M In rocky ♂. Ju in any with I never

- 200  
 1. X. baccinum ciliatiss calcarat acutis, Xylosteu β. baccis al. On mou Virgin ♂. Ju of red from t X. tan utrinq brevi  
 2. X. ba ovalib pedunc. On rock +.

PENTANDRIA MONOGYNIA. Caprifolium.

161

gratum.

4. C. spicæ verticillis approximatis, corollis ringentibus : tubo elongato, foliis perennantibus obovatis submucronatis subtus reticulato-venosis pallidioribus : summis connato-perfoliatis.—*Willd. sp. pl. 1. p. 984. sub Lonicera:*

Icon: *Hort. angl. 15. n. 10. t. 8.*

On the mountains, rambling among rocks in shady moist situations : New York to Carolina ; rare.  $\frac{1}{2}$ . June—Sept. v. v. Flowers inclining to scarlet.

5. C. spicæ verticillis capitatis subsessilibus, corollis breviusculis ringentibus basi gibbis, filamentis barbatis, foliis deciduis subtus glaucis : omnibus connatis : bractea perfoliata floribus multo ampliore.—*Lam. en-cycl. 1. p. 728.*

*Lonicera dioica. Willd. sp. pl. 1. p. 983.*

*Lonicera media. Murr. com. gætt. 1776. p. 28. t. 3.*

*Caprifolium bracteosum. Mich. fl. amer. 1. p. 105.*

Icon: *Murr. com. gætt. t. 3. Schmidt arb. 109.*

In rocky shady situations : New England to Carolina.

$\frac{1}{2}$ . June, July. v. v. Flowers yellow, smaller than in any of the foregoing species. There is a variety with purplish flowers, according to Michaux, which I never observed myself.

200. XYLOSTEUM. *Juss. gen. pl. 235.*

1. X. baccis distinctis, foliis ovatis subcordatisque margine ciliatis : junioribus subtus villosis, corollis conspicue calcaratis : tubo superne ventricoso : laciniis brevibus acutis, stylo exerto.

*Xylosteum tartaricum. Mich. fl. amer. 1. p. 106?*

- β. baccis albis.

On mountains, among rocks, in rich soil : Canada to Virginia. β. On the Rocky-mountain. *M. Lewis.*  $\frac{1}{2}$ . June, July. v. v. Flowers white, with a tinge of red or yellow. *X. tartaricum* differs materially from this species, and the diagnosis might be thus : *X. tartaricum*, baccis distinctis, foliis ovatis acutis utrinque glabris, corollis basi vix gibbosus : tubo brevi : laciniis oblongis obtusis, stylo inclusus.

2. X. baccis distinctis, ramis villosis, foliis oblongo-ovalibus obtusis utrinque subtomentoso-villoso-villosum, pedunculis brevibus. *Mich. fl. amer. 1. p. 106.*

On rocks in the mountains of Canada. *Michaux.*  $\frac{1}{2}$ . +. Berries blue.—Whether this is really a distinct

ciliatum.

album.

villosum.

species from *X. vulgare*, I cannot ascertain, having never seen the plant.

## 201. SYMPHORIA. Juss. Pers. syn. 1. p. 214.

*glomerata.*

1. S. floribus axillaribus subcapitato-globoseis. Pers. syn. l. c.  
*S. vulgaris*. Mich. fl. amer. 1. p. 106.  
*Lonicera symphoricarpos*. Willd. sp. pl. 1. p. 989.  
Icon. Schmidt arb. 115. Dill. eth. t. 278. f. 360.  
In sandy dry fields: Virginia and Carolina. ♀. July—Sept. v. v. Flowers small, red and yellow; berries purple.

*racemosa.*

2. S. racemo terminali, corolla intus barbata. Mich. fl. amer. 1. p. 107.  
On mountains, near lake Mistassins. Michaux. On the banks of the Missouri. M. Lewis. ♀. Aug. v. v. in Herb. Lewis.

## 202. DIERVILLA. Juss. gen. pl. 235.

*lutea.*

1. D. pedunculis axillaribus et terminalibus dichotomis trifloris, foliis ovatis serratis acuminatis.  
*D. Tournetortii*. Mich. fl. amer. 1. p. 107.  
*D. humilis*. Pers. syn. 1. p. 214.  
*Lonicera Diervilla*. Willd. sp. pl. 1. p. 989.  
Icon. Schmidt arb. 116. Duham. arb. t. 87.  
On rocks and the highest mountains: Canada to Carolina. ♀. June, July. v. v. Flowers yellow; fruit a brown dry capsule. There are a number of varieties, in respect to size, flowers and leaves.

## 203. TRIOSTEUM. Gen. pl. 300.

*perfoliatum.*

1. T. foliis ovalibus acuminatis connatis, floribus sessilibus verticillatis.—Willd. sp. pl. 1. p. 990.  
Icon. Dill. eth. t. 293. f. 378.  
In rich rocky grounds, principally lime-stone soil: New England to Carolina; rare. ♀. June—Aug. v. v. Flowers and berries dark purple.

*angustifoli-  
um.*

2. T. foliis ovali-lanceolatis subconnatis, pedunculis oppositis unifloris, caule hispido.—Willd. sp. pl. 1. p. 991.  
*T. minus*. Mich. fl. amer. 1. p. 107.  
Icon. Pluk. atm. t. 104. f. 2.  
On the borders of woods, in sandy soil: Virginia and

Carries  
1. R. in  
net  
Mi  
On la  
ski  
v. i  
2. R. in  
pu  
pla  
the  
p.  
In C  
&c  
lov  
3. R. in  
ret  
rig  
p.  
On l  
sy  
rie  
4. R. ir  
br  
er  
te  
R. gl  
Icon  
In re  
sy  
ti  
5. R. ir  
te  
ti  
bi  
st  
On t  
M  
6. R. ir  
co

Carolina.  $\frac{1}{2}$ . June, July. v. v. Flowers and berries yellow.

## 204. RIBES. Gen. pl. 390.

- R. inerme; foliis abbreviatis acute lobatis glabriusculis: *albinervium*. nervis albidis, racemis recurvatis, baccis glabris.—*Mich. fl. amer.* 1. p. 110.

On lake Mistassins: Canada. *Michaux*. On the Catskill mountains: New York, &c.  $\frac{1}{2}$ . April, May. v. v. Flowers small, greenish-yellow; berries red.

- R. inerme; foliis modice lobatis supra glabris subtus pubescentibus, racemis laxis pubescentibus, floribus planiusculis, laciniis calycinis substrifidis, petalis spatulatis obtusis, baccis hirsutis.—*Mich. fl. amer.* 1. p. 110.

In Canada. *Michaux*. On the Pennsylvania mountains, &c.  $\frac{1}{2}$ . April, May. v. v. Flowers greenish-yellow; petals purple; berries red.

- R. inerme; ramis rectis, foliis acute lobatis dentatisque reticulato-rugosis subtus pubescentibus, racemis laxis rigescenti-erectis, baccis hispidulis.—*Mich. fl. amer.* 1. p. 110.

On lake Mistassins: Canada. *Michaux*. In the Pennsylvania mountains, &c.  $\frac{1}{2}$ . May, June. v. v. Berries red, erect as well as the flowers.

- R. inerme; ramis reclinato-prostratis, foliis lobatis glabriusculis: junioribus pubescentibus, racemis suberectis, calycibus planiusculis, petalis deltoideis, bracteis minutis, baccis hispidis.—*L'Herit. stirp.* 1. p. 3.

*R. glandulosum*. *Ait. kew.* 1. p. 41.

Icon. *L'Herit. l. c. t. 2. Schmidt arb.* 95.

In rocky moist places: Newfoundland, Canada, Pennsylvania.  $\frac{1}{2}$ . April, May. v. v. Flowers yellow tinged with red; berries red.

- R. inerme; omnibus partibus pilis resinoso-glandulosis tectum; foliis 3—5-lobatis subrotundis, racemis erectis, calycibus planiusculis, petalis obtuse-rhomboideis, bracteis linearibus pedicello longioribus, baccis hirsutis.

On the mountains of North America. *Fraser*.  $\frac{1}{2}$ . April, May. v. v. in *Hort.* Flowers green.

- R. inerme; omnibus partibus pilis viscidis tectum; foliis cordatis obtuse-trilobis serratis, racemis erectis brevi-

*trifidum*.

*rígens*.

*prostratum*.

*resinosum*.

*viscosissimum*.

bus, calycibus tubulatis, petalis oblongis, bracteis linearis-spathulatis pedicello duplo brevioribus, germinibus hirsutis.

On the Rocky-mountain in the interior of North America. *M. Lewis.* ♀. June. *v. s. in Herb. Lewis.* Flowers large, yellow; the whole plant covered with viscous hair. This species approaches near to *R. glandulosum*. *Fl. peruv.* 3. p. 13. t. 233. f. b. It differs principally in the leaves being equally lobated, not having the middle lobe projecting; its long slender pedicels, and its petals.

- sanguineum.* 7. *R. inerme*; foliis cordatis trilobis serratis venoso-lineatis supra glabris, subtus tenui tomento albicanibus, racemis laxis pubescentibus foliis duplo longioribus, calycibus tubulatis, petalis oblongis longitudine calycis, bracteis obovato-spathulatis longitudine pedicellorum, germinibus hirsutis.

On the Columbia river. *M. Lewis.* ♀. March. *v. s. in Herb. Lewis.* Flowers beautiful, of a blood red or purple; branches purple. It approaches near to *R. albinervium*. *Fl. peruv.* 3. p. 12. t. 232. f. b.

- aureum.* 8. *R. inerme*; glaberrimum; foliis trilobis: lobis divaricatis inciso-pauci-dentatis petiolo basi ciliato brevioribus, racemis laxis dense-multifloris, calycibus tubulatis pedicellis longioribus: tubo gracili: laciinis oblongis obtusis, petalis linearibus laciinis calycis duplo brevioribus, bracteis linearibus longitudine pedicellorum, baccis glabris.

On the banks of the rivers Missouri and Columbia. *M. Lewis.* ♀. April. *v. s. in Herb. Lewis.*; *v. v. in Hort.* Flowers in close racemes, beautiful golden-yellow; berries red or brown, of an exquisitely fine taste, and considerably larger size than any of the garden currants. The shrub before flowering has the appearance of a species of *Crataegus*.

- recurvatum.* 9. *R. inerme*; ramis recurvatis, foliis latiusculis acute-latis pubescentibus glanduloso-punctatis, racemis reflexis, calycibus tubulatis glabris.

Near Hudson's Bay, Canada. *Michaux.* ♀. +. Berries black.

- floridum.* 10. *R. inerme*; foliis utrinque punctatis, racemis pendulis, calycibus cylindraceis, bracteis pedicellis longioribus. *Willd. sp. pl. 1. p. 1156.*

*R. pensylvanicum.* *Lam. encycl.* 3. p. 47.

Icon.  
In he  
Ma  
11. *R. spi*  
bus  
lim  
p. 1  
On th  
12. *R. spi*  
mit  
cis  
Amor  
Vir  
13. *R. spi*  
put  
cap  
lati  
On re  
Can  
or  
14. *R. sp*  
der  
bra  
hir  
hor  
Icon.  
On th  
Ma  
wh  
15. *R. ac*  
rib  
pec  
sp.  
Icon.  
On re  
v.  
16. *R. sp*  
ac  
ba  
R. o  
In sv  
Ap  
be

bracteis  
 , germi-  
 nis  
 b Ame-  
 Lewis.  
 red with  
 R. glan-  
 It differs  
 ted, not  
 slender  
 o-lineatis  
 cantibus,  
 gioribus,  
 ine caly-  
 edicello-  
 h. v. s.  
 blood red  
 near to  
 b.  
 divarica-  
 vioribus,  
 ulatis pe-  
 oblongis  
 slo brevi-  
 tellorum,  
 Columbia.  
 ; v. v. in  
 l golden-  
 itely fine  
 ry of the  
 g has the  
 acute-lo-  
 cemis re-  
 +. Ber-  
 pendulis,  
 ngioribus.

- Icon. *Dill. elth. t. 244. f. 315.* *Schmidt arb. 92.*  
 In hedges and woods : Canada to Virginia.  $\text{h}.$  April,  
 May. *v. v.* Flowers pale yellow ; berries black.
11. *R. spina subaxillari* ; foliis suborbiculatis subpubescenti-  
 bus : lobis subrotundo-obtusis, pedunculis unifloris,  
 limbo calycis tubuloso, baccis glabris. *Mich. fl. amer. 1.*  
*p. 110.*  
 On the high mountains of Carolina. *Michaux.*  $\text{h}.$  +.
12. *R. spinula subaxillari*, ramis subhispidis foliis parvis se-  
 mitrifidis : lobis subdentatis, pedunculis unifloris, bac-  
 cis glabris.—*Mich. fl. amer. 1. p. 111.*  
 Among rocks, in the Allegany mountains : Canada to  
 Virginia.  $\text{h}.$  May, June. *v. v.* Berries red.
13. *R. spinula subaxillari*, foliis petiolis gracilibus utrinque  
 pubescentibus : lobis acutis dentato-incisis, pedunculis capillaris subbifloris, calycibus tubulato-campanu-  
 latis, baccis glabris.—*Mich. fl. amer. 1. p. 111.*  
 On rocks and in mountain meadows : New York to  
 Carolina.  $\text{h}.$  April—June. *v. v.* Berries purple  
 or blue, of excellent taste.
14. *R. spina subaxillari*, foliis glabris 3—5-lobatis inciso-  
 dentatis, pedunculis subtrifloris : pedicellis elongatis,  
 bracteis brevissimis, petalis spatulatis undulatis, stylo  
 hirsuto semibifido exerto, baccis glabris.—*Willd.*  
*hort. 1. p. 61.*  
 Icon. *Willd. l. c. t. 61.*  
 On the Blue mountains : Pennsylvania to Virginia.  $\text{h}.$   
 May, June. *v. v.* Flowers yellowish-green ; petals  
 white ; berries pale red, small.
15. *R. aculeis majoribus et subsolitariis ad gemmas, mino-  
 ribus undique sparsis, foliis glabris* : lobis dentatis,  
 pedunculis brevibus subbifloris, baccis glabris.—*Willd.*  
*sp. pl. 1. p. 1159.*  
 Icon. *Dill. elth. t. 139. f. 166.*  
 On rocks : Canada, New York, &c.  $\text{h}.$  April, May.  
*v. v.*
16. *R. spina subaxillari multiplici, caule undique hispido-  
 aculeato*, foliis ultra medium lobatis, petiolis villosis,  
 baccis racemosis hispidis.—*Pers. syn. 1. p. 252.*  
*R. oxyacanthoides.* *Mich. fl. amer. 1. p. 111.*  
 In swamps, on the mountains : Canada to Virginia.  $\text{h}.$   
 April—June. *v. v.* Flowers small, greenish-yellow ;  
 berries amber-coloured or brown.

- Cynosbati.* 17. *R. passim* subgeminis 2-aculeatum, foliis brevi-lobatis inciso-dentatis molliter pubescentibus, racemis nutanti-paucifloris, calycibus erecto-campanulatis, baccis aculeatis.—*Mich. fl. amer. 1. p. 111. Willd. sp. pl. 1. p. 1159.*

Icon. *Jacq. hort. 2. t. 123. Schmidt arb. 98.*

On the sides of hills and rocks in the Allegany mountains, and in Canada.  $\text{h}.$  April—June. *v. v.* Flowers green; berries dark brown and full of thorns.

### 205. RHAMNUS. Gen. pl. 358.

- franguloides.* 1. *R. inermis*; foliis ovalibus acuminatis serrulatis subtus ad nervos pubescentibus, floribus dioicis, pedunculis unifloris aggregatis, calycibus acutis, fructibus turbatis.—*Mich. fl. amer. 1. p. 153.*  
*R. alnifolius.* *Willd. sp. pl. 1. p. 1100.*  
 On dry hills, near rivers: Canada to Virginia.  $\text{h}.$  June, July. *v. v.; v. s. in Herb. Lambert.* Flowers green; berries black; leaves small.

- alnifolius.* 2. *R. inermis*; foliis ovalibus denticulatis brevi-acuminatis basi sinu parvo cordatis subtus ad nervos pubescentibus, pedunculis bis bifidis, baccis depresso-globosis.  
 On the banks of the river Kooskoosky. *M. Lewis.*  $\text{h}.$  May. *v. s. in Herb. Lewis.* Berries purple, very highly esteemed by the Indians of that country.

- carolinianus.* 3. *R. inermis*; foliis ovali-oblongis integriusculis glabris, umbellis pedunculatis, floribus hermaphroditis 4-andris 1-gynis, stigmate bilobo, fructibus globosis.—*Mich. fl. amer. 1. p. 153.*  
 In woods and swamps: Virginia and Carolina.  $\text{h}.$  May—July. *v. v.* Berries black.

- lanceolatus.* 4. *R. inermis*, arborescens; foliis lanceolatis serrulatis utrinque acutis subtus pubescentibus.  
 On the side of hills: Tennessee. *Lyon.*  $\text{h}.$  *v. s. in Herb. Lyon.* Berries black.

- minutiflorus.* 5. *R. inermis*; foliis suboppositis ovalibus serrulatis, floribus minutissimis dioicis spicatis in rachi alterne sessilibus, stylo 3-fido, bacca 3-sperma. *Mich. fl. amer. 1. p. 154.*  
 On the sea-coast of Carolina and Florida. *Michaux.*  $\text{h. +.}$

- 2  
 1. *C. foliis nervis longe pl. 1. Icon. L In dry June called tually*  
 2. *C. foliis tripli bus laxis Icon. In the Leaf*  
 3. *C. foli axill celli Near sou Leu lon*  
 4. *C. fol deis On ro v. i the do*  
 5. *C. su sul M In sa v. lik w*

206. CEANOTHUS. *Gen. pl. 361.*

1. *C. foliis ovatis acuminatis serratis triplinervibus subtus nervis venisque pubescentibus, paniculis axillaribus longe pedunculatis, pedicellis corymbosis.* *Willd. sp. pl. 1. p. 1114.*  
Icon. *Duham. arb. 51.* *Schmidt arb. 132.*  
In dry woods; common: Canada to Florida.  $\frac{1}{2}$ . May, June. v. v. Flowers small, white. It is commonly called *New Jersey-tea* or *Red-root*; having been actually used in the American war as a substitute for tea.
2. *C. foliis oblongo-ovalibus acutis mucronato-serrulatis triplinervibus subtus pubescentibus, paniculis axillari- bus longissime pedunculatis, pedicellis corymbosis laxis.*  
Icon. *Pluk. alm. t. 28. f. 6.*  
In the woods of Tennessee. *Lyon.*  $\frac{1}{2}$ . June, July. v. v. Leaves not one fourth the size of the foregoing species.
3. *C. foliis obovatis serratis subtus pubescentibus, paniculis axillaribus thyrsoideis brevissime pedunculatis, pedicellis aggregatis.*  
Near the Rocky mountains on the banks of the Missouri. *M. Lewis.*  $\frac{1}{2}$ . May, June. v. s. in *Herb. Lewis.* Branches blood-red or purple; panicles not longer than the leaves.
4. *C. foliis ovalibus subserrulatis glabris, paniculis thyrsoideis terminalibus axillaribusque.*  
On rocks near rivers: Virginia and Carolina.  $\frac{1}{4}$ . May. v. v. Leaves the size of the first species, but smooth; the stems scarcely ever remain over winter, but die down to the root.
5. *C. subdecumbens; foliis minutis subfasciculatis oblongis microphyllus. subintegerrimis glabriusculis, corymbulis terminalibus.*  
*Mich. fl. amer. 1. p. 154.*  
In sandy woods: Carolina to Florida.  $\frac{1}{2}$ . May, June. v. v. A very delicate little shrub, the root of which, like all the foregoing species, is large and red, from which they derive the name of *Red-root*.

207. CELASTRUS. *Gen. pl. 372.*

1. *C. inermis, scandens; foliis oblongis acuminatis serratis, racemis terminalibus.* *Willd. sp. pl. 1. p. 1125.*  
*scandens.*

Icon. *Duham. arb.* 95. *Schmidt arb.* 140.

In hedges and woods, among rocks: Canada to Virginia.  
h. May, June. v. v. Flowers pale yellow; berries  
orange-scarlet.

*bullatus.*

2. C. inermis; foliis ovatis acutis integerrimis, panicula  
terminali. *Willd. sp. pl.* 1. p. 1123.

Icon. *Pluk. alm. t.* 28. f. 5.

In Virginia. h. +. Notwithstanding all my diligent  
researches at the place of its supposed nativity, as well  
as in the Herbariums of Plukenet and Banister, now at  
the British Museum, I have not been able to find any  
information respecting this species: consequently I  
strongly suspect its not being an American plant.

### 208. EVONYMUS. *Gen. pl.* 1. p. 373.

*americanus.*

1. E. ramis 4-angulatis, foliis subsessilibus elliptico-lan-  
ceolatis acutis serratis, pedunculis subtrifloris, floribus  
omnibus 5-fidis, fructibus verrucoso-muricatis.—  
*Willd. sp. pl.* 1. p. 1132.

Icon. *Pluk. alm. t.* 115. f. 5. *Schmidt arb.* 75.

In hedges and shady woods, among rocks, and on the  
edges of swamps: New England to Carolina. h.  
June, July. v. v. Flowers yellow, tinged with red;  
fruit scarlet, resembling those of *Arbutus Unedo* at  
first sight. They are a great ornament to this almost  
evergreen shrub. The common name in America is  
*Burning-bush*.

*angustifolius.*

2. E. ramis 4-angulatis, foliis subsessilibus elongato-lineari-  
ellipticis subfalcatis subintegerrimis, pedunculis ple-  
rumque unifloris, floribus omnibus 5-fidis, fructibus  
verrucoso-muricatis.

In shady woods: Georgia. Lyon. h. June, July.  
v. v. Flowers and fruit the same as the first species.  
Though nearly related to it, it is propagated by seeds,  
and keeps its own distinction, according to informa-  
tion from Mr. Lyon.

*atropurpu-  
reus.*

3. E. foliis petiolatis oblongo-lanceolatis acuminatis serratis,  
pedunculis divaricatis multifloris, floribus 4-fidis, fruc-  
tibus laevibus.—*Willd. sp. pl.* 1. p. 1132.

Icon. *Jacq. hort. 2. t.* 120. *Schmidt arb.* 73.

On the banks of rivulets: New York to Carolina. h.  
June, July, v. v. Flowers dark purple; fruit red.

1. V. foliis  
tomen  
Mich.

V. taurin

Icon. Ja

In shady

v. v. I

comm

white

2. V. foliis  
rufo-t  
parvis

V. vulpi

V. Labr

Icon. Ja

β. V. foliis  
boidei

In field

June.

and f

wine.

3. V. foliis  
bris,  
amer

V. incis

V. vulp

Icon. .

On the  
h. J

small

com

4. V. foliis  
petio  
amer

V. odo

On the  
sylva

of a

seda

nort

very

5. V. utr

## 209. VITIS. Gen. pl. 396.

- o Virginia,  
n; berries  
panicula
- V. foliis lato-cordatis sublobato-angulatis subtus incano-tomentosis, racemis fertilibus parvis, baccis majoribus. — *Mich. fl. amer. 2. p. 230. Willd. sp. pl. 1. p. 1181.*

*Labrusca.*

- V. taurina. *Walt. fl. car. 242.*  
Icon. *Jacq. schaenbr. 426.*

In shady woods: Canada to Florida.  $\frac{1}{2}$ . June, July.  
*v. v.* Berries black, large, of a disagreeable fox-smell, commonly called *Fox-grape*. There is a variety with white berries called *Bland's-grape*.

- V. foliis lato-cordatis 3—5 lobis: junioribus subtus rufo-tomentosis, racemis fertilibus oblongis, baccis parvis. — *Mich. fl. amer. 2. p. 230.*

*festivalis.*

- V. vulpina. Willd. sp. pl. 1. p. 1181.*  
V. *Labrusca. Walt. fl. car. 242.*  
Icon. *Jacq. schaenbr. 425.*

- β.* V. foliis sinuato-palmatis grosse dentatis: sinubus rhomboideis. An species distincta?

*sinuata.*

In fields and woods: Virginia to Carolina.  $\frac{1}{2}$ . May, June. *v. v.* Berries dark blue, very agreeable to eat, and frequently converted into very good home-made wine. It is known by the name of *Summer-grape*.

- V. foliis cordatis acuminatis inciso-dentatis utrinque glabris, racemis laxe multifloris, baccis parvis. — *Mich. fl. amer. 2. p. 231.*

*cordifolia.*

- V. incisa. Jacq. schaenbr. 427.*  
*V. vulpina. Walt. fl. car. 243.*

- Icon. *Jacq. l. c.*

On the edges of rivers and in woods: Canada to Florida.  $\frac{1}{2}$ . June, July. *v. v.* Berries green or amber-coloured, small, and ripen extremely late, of a very tart taste: commonly called *Winter-grape* or *Chicken-grape*.

- V. foliis inæqualiter inciso-dentatis breviuscule 3-fidis: petiolo nervis margineque pubescentibus. — *Mich. fl. amer. 2. p. 231.*

*riparia.*

- V. *odoratissima. Donn catal. 66.*

On the gravelly shores and islands of the rivers: Pennsylvania to Carolina,  $\frac{1}{2}$ , May—July. *v. v.* Flowers of an exquisitely fine smell, somewhat resembling *Reseda odorata*. Female plants are very seldom found north of the Potowmac river, though the male extend very far beyond it.

- V. utrinque lucidis reniformi-cordatis subæqualiter den-

*rotundifolia.*

## PENTANDRIA MONOGYNIA. Vitis.

tatis, racemorum floribus plures capitulatis, baccis magnis. *Mich. fl. amer.* 2. p. 231.

On river sides and islands: Virginia to Florida.  $\text{h}.$  June, July. *v. v.* Berries very large, dark blue, agreeable, commonly called *Bull-* or *Bullet-grapes.*

*palmata.*

6. *V. foliis cordatis palmatis glabris: laciniis lanceolatis incisis, umbellis racemosis.* *Willd. sp. pl.* 1. p. 1180.  
In Virginia. *Vahl.*  $\text{h. +.}$

210. CISSUS. *Gen. pl.* 655.

*Ampelopsis.*

1. *C. foliis truncato-cordatis acuminatis dentatis angulatisque: nervis subtus pubescentibus, racemis duplicato-bifidis.* — *Pers. syn.* 1. p. 142.

*Ampelopsis cordata.* *Mich. fl. amer.* 1. p. 159.

On river sides and among hedges, in the Allegany mountains: Pennsylvania to Carolina.  $\text{h.}$  June, July. *v. v.* Berries pale red.

*hederacea.*

2. *C. radicans, scandens; foliis quinato-digitatis utrinque glabris: foliolis petiolatis oblongis acuminatis mucronato-dentatis, pedunculis dichotomis: racemis corymbosis.* — *Pers. syn.* 1. p. 143.

*Ampelopsis quinquefolia.* *Mich. fl. amer.* 1. p. 160.

*Vitis hederacea.* *Willd. sp. pl.* 1. p. 1182.

*Hedera quinquefolia.* *Linn. hort. cliff.* 74.

*Icon. Corn. canad. t.* 100.

*hirsuta.*

- β. *C. foliis utrinque pubescentibus: foliolis ovatis acuminatis grosse-dentatis.*

*Ampelopsis hirsuta.* *Donn catal.* 166.

On the Allegany mountains: Pennsylvania to Virginia.  $\text{h.}$  *v. v.* The variety β. appears to be a distinct species; but, as I never have seen its flowers, a sufficient character is wanting.

*stans.*

3. *C. foliis bipinnatis: foliolis inciso-serratis, racemis pedunculatis duplicato-bifidis.* — *Pers. syn.* 1. p. 143.

*Ampelopsis bipinnata.* *Mich. fl. amer.* 1. p. 160.

*Vitis arborea.* *Willd. sp. pl.* 1. p. 1183.

*Icon. Pluk. mant. t.* 412. *f. 2.*

In shady woods, on river sides: Virginia and Carolina.  $\text{h.}$  June, July. *v. v.* Stem upright.

211. CYRILLA. *Linn. Mich. fl. amer.* 1. p. 157.

*caroliniana.*

1. *C. foliis cuneato-lanceolatis acutis membranaceis nervosis, spicis gracilibus, petalis pedicello longioribus.* *Mich. fl. amer.* 1. p. 158.

C. racemifl  
It ea Cyrrilla  
Icon. Jacq  
In sandy  
A very

## 212. ITEA

1. *I. foliis ob*  
*sp. pl.* 1.  
Icon. *Du*  
In swamp  
*v. v.* 1

21  
1. *I. pedunc*  
*dentati*  
*pl. 1. 1*  
Icon. *Fl.*  
Near spr  
gland  
plain

2. *I. pedun*  
*argute*  
*car. 2*  
In wet  
July.  
on th

1. *V. folii*  
*latis*  
*Wil*  
Icon.  
On dry  
na.  
blue  
petal

2. *V. fol*  
*nua*  
In Vir  
pale

latis, baccis  
lorida. h.  
dark blue,  
let-grapes.  
ncoelatis in-  
1. p. 1180.

is angulatis-  
is duplicato-

59.

eganey moun-  
, July. v. v.

itis utrinque  
natis mucro-  
remis corym-  
p. 160.

vatis acumi-

to Virginia,  
distinct spe-  
s, a sufficient

racemis pe-  
1. p. 143.  
. 160.

ind Carolina.

1. p. 157.  
naceis nervo-  
longioribus.

C. racemiflora. Linn. *mant.* 50.  
Itea Cyrilla. Willd. *sp. pl.* 1. p. 1146.  
Icon. *Jacq. ic.* 1. t. 47. *L'Herit. stirp.* 1. t. 66.  
In sandy swamps of Carolina. h. June, July. v. v.  
A very elegant shrub; flowers white.

212. ITEA. *Gen. pl. 381.* *Mich. gen. pl.* 1. p. 156.  
1. I. foliis oblongis serratis, spicis pubescentibus.—Willd. *sp. pl.* 1. p. 1146.  
Icon. *Duham. arb.* 1. t. 126.  
In swamps: Pennsylvania to Carolina. h. June, July.  
v. v. From four to six feet high: flowers white.

213. IMPATIENS. *Gen. pl. 1365.*  
1. I. pedunculis multifloris solitariis, foliis ovatis obtuse *Nolitangere.*  
dentatis, geniculis caulinis tumentibus.—Willd. *sp. pl.* 1. p. 1176.  
Icon. *Fl. dan.* 588. *Engl. bot.* 937.  
Near springs and rivulets, in shady places: New England to Carolina. O. June, July. v. v. Flowers plain yellow.  
2. I. pedunculis plerumque bifloris solitariis, foliis ovatis  
argute dentatis.—Willd. *sp. pl.* 1. p. 1175. *Walt. fl. car.* 219.  
In wet shady places: Canada to Carolina. O. June,  
July. v. v. Flowers dark yellow, with red spots  
on the inside.

214. VIOLA. *Gen. pl. 1364.*  
\* *Acaules.*  
1. V. foliis subpedato-multipartitis: laciiniis linear-lanceo-  
latis subintegris, calycis laciiniis linearibus acutis.—  
Willd. *sp. pl.* 1. p. 1160.  
Icon. *Bot. mag.* 89. *Bot. rep.* 153.  
On dry sandy hills and fields: New England to Caroli-  
na. 4. May, June. v. v. Flowers large, beautiful  
blue. There is a very handsome variety with pale blue  
petals, ornamented with a dark purple velvet at the  
bottom, similar to *Viola tricolor*.  
2. V. foliis palmato-5—7-lobatis postice in petiolum atte-  
nuatis: lobis integris.  
In Virginia. *Leconte in litt.* 4. May. v. s. Flowers  
pale blue.

*palmata.*

3. V. pubescens ; foliis cordatis hastato-lobatis palmatisive ; laciniis dentatis, calycis laciniis lanceolatis glabris, petalis 2. oppositis basi barbatis.—*Willd. sp. pl. 1. p. 1159.*

Icon. *Bot. mag. 535. Plak. alm. t. 447. f. 1.*

On dry hills and pastures, generally in sandy soil. ♀. April—June. v. v. Flowers blue ; early in spring the leaves are generally kidney-shaped, and after the first flowers are gone they become palmated ; sometimes it occurs with smooth leaves.

*sagittata.*

4. V. pubescens ; foliis oblongis acutis cordato-sagittatis subserratis basi incisis, pedunculis foliis longioribus, calycis laciniis linearibus glabris, petalis 3. inferioribus basi barbatis.—*Willd. sp. pl. 1. p. 1160.*

On dry hills : New England to Virginia. ♀. April—June. v. v. Flowers blue ; lower petal white towards the bottom, with purple veins ; the rest longer, narrower, and white towards the base.

*dentata.*

5. V. glabra ; foliis oblongis acutis basi truncatis serratis inferne subhastato-grandidentatis, pedunculis foliis brevioribus, calycis laciniis linearibus, petalis 3. inferioribus basi barbatis.

In wet meadows and woods : Pennsylvania. ♀. May, June. v. v. Flowers of nearly the same colour and shape as the preceding species.

*lanceolata.*

6. V. glabra ; foliis lanceolatis subserratis, pedunculis vix foliis longioribus, petalis imberbibus : 2. superioribus subrotundis.—*Willd. sp. pl. 1. p. 1161.*

In overflowed meadows : Canada to Pennsylvania. ♀. June, July. v. v. Flowers white ; the upper petal painted with purple veins. There are several varieties of this species, some with ovate leaves attenuated in a petiole, some exceeding small, &c.

*blanda.*

7. V. glabra ; foliis cordatis acutiusculis planiusculis remote serratis, pedunculis longitudine foliorum, petalis imberibus : 2. lateralibus brevioribus, intimo reliquo longiore lanceolato —*Willd. hort. berol. 1. t. 24.*

In wet places, bog-meadows : New York to Carolina. ♀. April—June. v. v. Flowers yellowish white ; lower petal marked with blue stripes and veins.

*obliqua.*

8. V. glabra ; foliis cordatis acutis crenato-serratis planiusculis, floribus erectis, pedunculis longitudine foliorum, petalis oblique flexis : lateralibus angustioribus et lon-

PEN

gioribus  
p. 1161.

In shady we  
—June.

low vei

9. V. pubescen  
marginati  
obovatis  
p. 1162.

On dry hil  
v. v. Fl

10. V. glabra ;  
pedunculi  
flexis : 1

In grassy w  
Flowers

11. V. foliis co  
centibus  
infimo t

In overflo  
—June.

lower p

12. V. foliis t  
glabrius  
obovatis  
ventibu

Near Phil  
Flower  
yellow

13. V. glabri  
clauso  
stoloni

giorbu

V. rotund

On the h

woods  
♀. Ju

from  
under

wood  
small

groun

inhab

used

- palmatisve;  
latis glabris,  
*l. sp. pl. 1.*
1.  
ly soil. 2.  
ly in spring  
nd after the  
ated; some.
- lato-sagittatis  
longioribus,  
. inferioribus
2. April—  
al white to  
rest longer,
- icatis serratis  
anclus foliis  
alis 3. inferi.
4. 2. May,  
e colour and
- dunculus vix  
superioribus
- Pennsylvania. 2.  
upper petal  
eral varieties  
attenuated in
- culis remote  
, petalis im-  
imo reliquis  
*l. t. 24.*  
to Carolina,  
wishes white;  
veins.
- ratis planius.  
ine foliorum,  
ribus et lon-
- gioribus infra medium barbatis.—*Willd. sp. pl. 1.*  
*p. 1161.*
- In shady wet places: Pennsylvania to Virginia. 2. April—June. *v. v.* Flowers white with purple and yellow veins.
9. *V. pubescens*; foliis ovatis subcordatis crenatis: petiolis *primulifolia*.  
marginatis, calycibus ciliatis longis, petalis omnibus obovatis: 2. lateralibus barbatis.—*Willd. sp. pl. 1.*  
*p. 1162.*
- On dry hills: Canada to Virginia. 2. April—June.  
*v. v.* Flowers blue.
10. *V. glabra*; foliis cordatis glabris serratis basi cucullatis, pedunculis longitudine petiolorum, petalis oblique flexis: lateralibus barbatis.—*Willd. sp. pl. 1. p. 1162.*
- In grassy wet places: common. 2. May, June. *v. v.*  
Flowers blue, white at their base.
11. *V. foliis cordatis crenato-serratis obtusis subtus pubescentibus, pedunculis foliis brevioribus, petalis oblongis: infimo basi barbato.*—*Willd. hort. berol. 1. t. 72.*
- In overflowed meadows: Pennsylvania, &c. 2. April—June. *v. v.* Flowers blue, white at the bottom; lower petal veined.
12. *V. foliis trianguli-cordatis acutis crenatis subcucullatis papilionacea.*  
glabriusculis, pedunculis longitudine foliorum, petalis obovatis: 3. inferioribus infra medium barbatis conniventibus, 2. superioribus reflexis.
- Near Philadelphia, in wet places. 2. May, June. *v. v.*  
Flowers blue, elegantly striated and bearded with yellow down.
13. *V. glabriuscula*; foliis suborbiculatis obtusiusculis sinu clauso cordatis crenato-serratis: serraturis glandulosis, stolonibus floriferis, petalis linearibus calyce vix longioribus.
- V. rotundifolia. Mich. fl. amer. 2. p. 150?*
- On the high mountains of Pennsylvania, in shady beechwoods, among rotten wood and rich vegetable mould.  
2. June—Sept. *v. v.* This singular species differs from all the rest, in producing its flowers as it were under ground, as they always are covered with rotten wood or leaves; they are of a chocolate brown, very small; the seed-vessel buries itself still deeper in the ground, and is large in proportion to the plant. The inhabitants know it by the name of *Heal-all*, being used by them in curing all kinds of wounds or sores.

## \*\* Caulescentes.

- canadensis.* 14. *V. glabriuscula*; foliis subcordatis acuminatis serratis, pedunculis longitudine foliorum, stipulis brevibus integris.—*Willd. sp. pl. 1. p. 1166.*  
In shady woods, in rich moist situations: Canada to Carolina, on the mountains. ♀. June—Aug. v. v. Flowers outside purplish-blue, inside white and elegantly veined, sweet scented.
- striata.* 15. *V. glabra*; foliis cordatis acuminatis serratis planiusculis, pedunculis longissimis, stipulis lanceolatis serrato-ciliatis.—*Willd. sp. pl. 1. p. 1166.*  
In shady woods: Pennsylvania to Virginia. ♀. May—July. v. v. Flowers white, with purple veins.
- debilis.* 16. *V. glabra*; foliis reniformi-cordatis brevissime acuminatis crenatis basi cucullatis, pedunculis foliis duplo longioribus, stipulis serrato-ciliatis.—*Mich. fl. amer. 2. p. 150.*  
*V. canina* *Walt. fl. car. 219.*  
In low grounds: Pennsylvania to Carolina. ♀. May—July. v. v. Flowers not one half the size of the preceding species, pale blue.
- rostrata.* 17. *V. glabra*; foliis cordatis acutis serratis, pedunculis foliis duplo longioribus, stipulis lanceolatis serrato-ciliatis, nectariis rostratis corolla longioribus.  
On shady rocks: near Eastown, Pennsylvania. ♀. May, June. v. v. Flowers blue.
- pubescens.* 18. *V. viloso-pubescent*; caule erecto superne folioso, foliis lato-cordatis, stipulis oblongis apice serratis.—*Willd. sp. pl. 1. p. 1166.*  
*V. pensylvanica*. *Mich. fl. amer. 2. p. 149.*  
In shady woods, among rocks, particularly limestone rocks: New York to Virginia. ♀. May, June. v. v. Flowers yellow.
- hastata.* 19. *V. glabra*; caule simplici superne folioso, foliis hastatis brevissime petiolatis, stipulis minutis denticulatis. *Mich. fl. amer. 2. p. 149.*  
On high mountains: Pennsylvania to Carolina. ♀. May, June. v. v. Flowers yellow.
- Nuttallii.* 20. *V. pubescens*; caule simplici erecto, foliis ovato-oblongis acutis obsolete-dentatis nervosis, in petiolum longum attenuatis, stipulis lanceolatis integris, pedunculis longitudine foliorum.  
} On the banks of the Missouri. ♀. June. v. s. in *Herb Nuttall.* Flowers yellow.

*latis serratis,  
brevibus in-*

*Canada to Ca.  
-Aug. v. v.  
hite and ele-*

*planiusculis,  
is serrato-cili-*

*♀. May—  
e veins.*

*ime acuminata  
foliis duplo  
h. fl. amer. 2.*

*♀. May—  
size of the*

*Iunculus foliis  
errato-ciliatis,*

*a. ♀. May,*

*folioso, foliis  
ratis.—Willd.*

*ly limestone  
, June. v. v.*

*foliis hastatis  
denticulatis,*

*na. ♀. May,*

*ovato-oblongo.  
petiolum lon-  
is, pedunculis*

*v. s. in Herb*

21. V. stricte-t  
ceolato-l  
geminis  
t. 28.

V. stricta.  
On lime-st  
About t

22. V. subpub  
ribus r  
lanceola  
cibus a  
In fields  
v. v.  
Nearly  
collect  
barium

2

1. C. foliis  
liolis a  
berosa  
Icon. B  
In wet  
Engla  
ers w

2. C. foliis  
tusis,  
Para  
C. caro  
Icon. I  
On hi  
Mar  
spec

3. C. foli  
solit  
mis  
Ms  
On th  
in I  
the  
A.  
by  
agr

4. C. fo  
stir



*Claytonia lanceolata*

PENTANDRIA MONOGYNIA. Viola.

175

1. *V. stricte-erecta*; foliis lato-lanceolatis stipulisque lanceolato-linearibus integerrimis, pedunculis axillaribus geminis brevissimis.—*Forster in Linn. trans. 6. p. 309. t. 28.*

*concolor.*

*V. stricta. Muhl. MSS.*

On lime-stone rocks: Pennsylvania. 24. June, July. v. v.  
About two feet high; flowers small, green.

2. *V. subpubescens*; caule angulato simplici, foliis inferioribus rotundato-spathulatis subdentatis, superioribus lanceolatis integris, stipulis pectinato-pinnatifidis, calycibus acutis corolla duplo brevioribus.

*bicolor.*

In fields of Pennsylvania and Virginia. 24. May—July.  
v. v. Flowers small white, with a few purple veins.  
Nearly approaches to *V. tricolor*. Specimens in the collection of A. B. Lambert, Esq. from Pallas's Herbarium, are the same with this species.

215. CLAYTONIA. Gen. pl. 402.

1. *C. foliis linear-lanceolatis, racemis solitariis, calycis foliolis acutiusculis, petalis obovatis retusis, radice tuberosa.*—*Willd. sp. pl. 1. p. 1185.*

*virginica.*

Icon. *Bot. mag. 941.*

In wet woods, round the stumps of rotten trees: New England to Carolina. 24. March—May. v. v. Flowers white, with red veins.

2. *C. foliis spathulatis, racemo solitario, calycis foliolis obtusis, petalis subrotundis retusis, radice tuberosa.*—*Parad. lond. 71.*

*spathulaefolia.*

*C. caroliniana. Mich. fl. amer. 1. p. 160.*

Icon. *Parad. lond. 71. Pluk. alm. t. 102. f. 3.*

On high mountains of Virginia and Carolina. 24. March, April. v. v. Flowers smaller than the first species, rose-coloured with purple veins.

3. *C. foliis lanceolatis: caulinis ovatis sessilibus, racemo solitario elongato, calycis foliolis brevibus obtusissimis, petalis cuneatis bifidis, radice tuberosa.*—*Pall. MSS.*

*lanceolata.*

On the Rocky-mountains. *M. Lewis.* 24. June. v. s. in *Herb. Lewis*. Flowers white, nearly the size of the first species, without veins. In the collection of A. B. Lambert, Esq. I found a specimen collected by Pallas in the eastern parts of Siberia, perfectly agreeing with the present species.

4. *C. foliis radicalibus spathulato-ovatis, caulinis ovatis distinctis, racemis subgeminatis, floribus secundis, brac-*

*alsinoides.*

## PENTANDRIA MONOGYNIA. Claytonia.

teis ovatis linearibusque, petalis emarginatis, radicibus fibrosis.—*Sims in bot. mag.*

*C. sibirica. Hortul.*

Icon. *Bot. mag.* 1309.

On the Columbia river. *M. Lewis.* ☽. May, June. v. v. in *Hortis.* Flowers small, white.

*perfoliata.*

5. *C. foliis radicalibus spathulato-rhomboideis : caulin suborbiculato perfoliato, racemis subsolitariis ebracteatis, petalis emarginatis, radicibus fibrosis.—Donn catal. 50. Willd. sp. pl. 1. p. 1186.*

*C. cubensis. Humb. & Bonpl. equinoct. t. 26. Annales du mus. d'hist. nat. 7. p. 82. t. 6.*

Icon. *Bot. mag.* 1336.

On the Rocky-mountains. *M. Lewis.* ☽. April, May. v. v. in *Hort.* Flowers smaller than any of the foregoing species, white.

216. GLAUX. *Gen. pl.* 408.

*maritima.*

1. *Glaux. Willd. sp. pl. 1. p. 1216.*

Icon. *Fl. dan.* 548.

In overflowed marshes, near the sea : Canada. ♀. July. v. v. Leaves oblong, smooth. Flowers axillary, sessile, flesh-coloured.

217. ANYCHIA. *Mich. fl. amer.* 1. p. 112.

*dichotoma.*

1. *A. caule patulo dichotomo ramosissimo, foliis lanceolatis glabris erectis, stipulis floralibus calyce glabro subæqualibus. Mich. fl. amer. 1. p. 113.*

*Queria canadensis. Willd. sp. pl. 1. p. 494.*

Icon. *Ort. dec.* t. 15. f. 2.

On dry limestone hills: New York to Kentucky. ♀. June—Aug. v. v. Flowers exceeding small; very variable in the number of filaments, generally from two to five.

*herniarioides.*

2. *A. humifusa, conferta, undique pubescens ; foliis oblongo-ovalibus ciliatis mucronatis, laciinis subulatis : acumine setaceo patuloque. Mich. fl. amer. 1. p. 113.*

On the dry sands of North Carolina. *Mihaux.* ♀. +.

*argyrocoma.*

3. *A. cæspitosa, procumbens ; caulinibus pubescentibus, foliis linearibus acutissimis subpilosis, capitulis fasciculato-terminalibus argenteis, calycis pilosi apicibus barbatis et longe acuminatis. Mich. fl. amer. 1. p. 113.*

On rocks in Upper Carolina and Virginia. ♀. June, July. v. v.

1. T. erectu  
rymos  
T. corym  
Icon. *Pl*  
On dry h  
June, .

1. N. foliis  
costa n  
subtrifl  
*Willd.*  
N. multi  
N. mont  
Icon. *M*  
In all the  
May.  
the size  
of *Sor*

2. N. foliis  
bris, p  
vatis :  
N. aquat  
N. integri  
Icon. *M*  
In deep  
May.  
cies, t

3. N. foliis  
basi c  
neis  
p. 111  
N. capit  
Icon. *M*  
On the  
river  
orang  
from

4. N. foli  
serrat  
ris, d

VOL. I.

218. THESIUM. *Gen. pl. 410.*

1. *T. erectum* : foliis ovali-lanceolatis, fasciculis florum co-  
rumboso-terminalibus.—*Willd. sp. pl. 1. p. 1214.*  
*T. corymbulosum*. *Mich. fl. amer. 1. p. 112.*  
Icon. *Pluk. mant. t. 342. f. 1.*  
On dry hills and fields; New York to Carolina. 2.  
June, July. v. v. Flowers white.

219. NYSSA. *Gen. pl. 1599.*

1. *N. foliis oblongis integerrimis utrinque acutis, petiolo,  
costa media margineque villosis, pedunculis feminineis  
subtrifloris, nucibus brevi-obovatis obtuse striatis.*—  
*Willd. sp. pl. 4. p. 258.*  
*N. multiflora*. *Walt. fl. car. 253.*  
*N. montana*. *Hortul.*  
Icon. *Mich. arb. 21. Wangh amer. t. 16. f. 39.*  
In all the woods from New England to Carolina. 2.  
May. v. v. Flowers small, greenish; berries black,  
the size of a pea. This tree is known by the name  
of Sour-gum.
2. *N. foliis ovato-oblongis integerrimis utrinque acutis gla-  
bris, pedunculis feminineis bifloris, drupis brevi-obo-  
vatis: nuce obtuse-striata.*—*Willd. sp. pl. 4. p. 1113.*  
*N. aquatica*. *Sp. pl. 1511.*  
*N. integrifolia*. *Ait. kew. 3. p. 446.*  
Icon. *Mich. arb. 22. Catesb. car. 1. t. 41.*  
In deep swamps: Virginia and Carolina. 2. April,  
May. v. v. Berries the size of the preceding spe-  
cies, blueish-black.
3. *N. foliis brevissime petiolatis oblongis sub-integerrimis  
basi cuneatis subtus candicantibus, pedunculis femi-  
neis unifloris, drupis oblongis.*—*Willd. sp. pl. 4.  
p. 1113.*  
*N. capitata*. *Walt. fl. car. 253.*  
Icon. *Mich. arb. 20.*  
On the banks of rivers: Carolina, principally on the  
river Ogeechee. 2. May, June. v. v. Fruit large,  
orange-coloured, full of an acid similar to a lime,  
from which it is known by the name *Ogeechee-lime*.
4. *N. foliis longe petiolatis oblongis acuminatis remote  
serratis subtus tomentosis, pedunculis feminineis uniflo-  
ris, drupis oblongis.*—*Willd. sp. pl. 4. p. 1113.*

*villosa.**biflora.**candicans.**tomentosa.*

*N. grandidentata. Mich. arb. t. 19.*

On the banks of the river St. Mary, and in Florida.  
½ May. v. s. Fruit the size of the preceding, dark blue.

*denticulata.*

5. *N. foliis longe petiolatis oblongis acuminatis remote serratis undique glabris, pedunculis femineis unifloris, drupis oblongis. Willd. sp. pl. 4. p. 1114.*

*N. uniflora. Walt. fl. car. 253.*

*N. angulans. Mich. fl. amer. 2. p. 259.*

Icon. Wangh amer. t. 27. f. 57. Catesb. car. 1. t. 60.  
In swamps: Virginia and Carolina. ½ v. v. s. fl.

## 220. HAMILTONIA. Willd. sp. pl. 4. p. 1114.

Pyrularia. Mich. fl. amer. 2. p. 231.

*oleifera.*

1. *Hamiltonia. Willd. l. c.*

Pyrularia pubera. Mich. l. c.

In shady woods, in the mountains of Pennsylvania,  
Virginia near the Sweet springs, and in Carolina. ½  
May, June. v. v. A shrub from four to six feet high,  
very pubescent; leaves oblong, entire, acuminate;  
flowers small, greenish-yellow, in terminal racemes;  
fruit known in the country by the name of Oil-nut,

## II. DIGYNIA.

### 221. ECHITES. Gen. pl. 421.

*difformis.*

1. *E. volubilis; foliis ovali-lanceolatis acuminatis basi acutis: infimis linearibus, floribus fasciculato-corymbosis.*  
—Walt. fl. car. 98.

*E. puberula. Mich. fl. amer. 1. p. 120.*

In woods of Lower Virginia and Carolina. ½ July.  
v. v. Flowers small, dirty yellow; seed pods very slender and long.

### 222. GONOLOBIUM. Mich. fl. amer. 1. p. 119.

*macrophyllum.*

1. *G. sarmentis petiolisque hirsutis, foliis amplis abrupte acuminatis puberulis, corollæ latiniis ovali-lanceolatis, folliculis costato-angulosis.—Mich. fl. amer. 1. p. 119.*

Vincetox  
In shady  
v. v.

2. *G. sarmentis minatis longo-  
Mich.*

Vincetox  
In hedges  
June,

3. *G. sativis sensim corolla levibus*

Cynanch  
In hedge  
v. v. s

Cynanch  
atithem  
in flo  
never  
I shall

1. *A. caule  
bris,  
limbo*  
Icon. 1  
In field  
rolina  
stripe

2. *A. caul  
dicar  
lis lin*  
Icon. 1  
In sand  
v. v.  
gene

3. *A. cau  
mis  
A. sibi  
On th  
gini*

1. Florida, ling, dark  
remote ser- unifloris,  
1. t. 60.  
v. s. fl.  
. 1114.  
  
ensylvania, rolinia. h.  
feet high, cùminate;  
racemes;  
Oil-nut,
1. Florida, ling, dark
  2. G. sarmentis petiolisque hirsutissimis, foliis sensim acuminatis utrinque pubescentibus, corollæ laciniis oblongo-ovalibus obtusis, folliculis oblongis muricatis.—*Mich. fl. amer. 1. p. 119.*

*Vincetoxicum acanthocarpos. Walt. fl. car. 104.*  
In hedges, near rivulets: Pennsylvania to Carolina. 4. June, July. v. v. Flowers dark purple.

3. G. sarmentis glabriusculis, foliis subconoideo-cordatis sensim acutis nervis subtus puberulis, floribus glabris, corollæ laciniis ovali-oblongis obtusiusculis, folliculis laevibus. *Mich. fl. amer. 1. p. 119.*

*Cynanchum carolinense. Jac. ic. 2. t. 342?*  
In hedges on the banks of the Mississippi. 4. June.  
v. v. s. fr. Flowers dark purple.

*Cynanchum obliquum. Jacq. ic. 2. t. 241.* probably is another species of this genus. I have seen the plant in flower, but without fruit, in a cultivated state, but never observed it growing spontaneously; consequently I shall omit it, as a doubtful North American plant.

### 223. APOCYNUM. Gen. pl. 426.

1. A. caule rectiusculo patente, foliis ovatis utrinque glabris, cymis lateralibus et terminalibus glabris, corollis limbo patentibus.—*Willd. sp. pl. 1. p. 1259.*

Icon. *Dod. mem. t. 50.*  
In fields and along the edges of woods: Canada to Carolina. 4. July. v. v. Flowers pale red with darker stripes.

2. A. caule rectiusculo, foliis oblongo-ovalibus subtus canalicante-pubescentibus, paniculis pubescentibus, corollis limbo erectis.—*Willd. sp. pl. 1. p. 1259.*

Icon. *Pluk. alm. t. 13. f. 1.*  
In sandy fields and woods; common. 4. June, July.  
v. v. Flowers yellowish-green, small. The plant is generally called *Indian Hemp*.

3. A. caule erectiusculo, foliis oblongis cordatis glabris, cymis folio brevioribus. *Willd. sp. pl. 1. p. 1260.*

*A. sibiricum. Jacq. hort. 3. t. 66.*  
On the gravelly shores of rivers: New York to Virginia; scarce. 4. July, Aug. v. v. Flowers very

*hirsutum.*

*læve.*

*hypericifo-  
lium.*

## PENTANDRIA DIGYNIA. Apocynum.

small; plant lower than the preceding species, and very smooth.

224. ASCLEPIAS. *Gen. pl. 429.*\* *Foliis oppositis.**syriaca.*

1. A. caulis simplicissimus, foliis lanceolato-oblongis sensim acutis subtus tomentosis, umbellis subnutantibus tomentosis.—*Willd. sp. pl. 1. p. 1265.*

In fields and on the banks of rivers, very common: New England to Virginia.  $\frac{1}{4}$ . July, Aug. v. v. From three to five feet high; flowers in large close clusters, pale purple, sweet-scented. The nectaries act as fly-traps.

*phytolacoides.*

2. A. caule erecto simplici, foliis lato-ovato-oblongis acutis glabris subtus pallidioribus, umbellis compositis laterilibus solitariis longe pedunculatis nutantibus.—*Gron. virg. 38.*

On the mountains of Virginia and Carolina.  $\frac{1}{4}$ . July, Aug. v. v. Stem about three feet or more high, spotted with purple; leaves large, middle rib purple; flowers sweet-scented; petals reflexed, very dark purple. This plant was introduced into England in 1812, by Mr. Lyon, under the above name.

*debilis.*

3. A. glabriuscula; caule debili errecto simplici, foliis petiolatis ovali-lanceolatis utrinque acutis membranaceis, umbellis terminalibus laxifloris, pedicellis capillaribus.—*Mich. fl. amer. 1. p. 116.*

In shady rocky situations, near rivulets: New York to Carolina; rare.  $\frac{1}{4}$ . July, Aug. v. v. Flowers white; leaves large. The fibres of the stem produce a kind of flax, as all the rest of this genus do, but far superior, being of a beautiful silky gloss and extreme strength.

*parviflora.*

4. A. foliis lanceolatis acuminatis glabris oppositis basi attenuatis, umbellis lateralibus solitariis, caule suffruticoso errecto. *Willd. sp. pl. 1. p. 1267.*

A. perennis. *Walt. fl. car. 107.*

In sandy fields: Carolina and Florida.  $\frac{1}{4}$ . July, Aug. v. v. Flowers small, white.

*nivea.*

5. A. foliis ovato-lanceolatis glabriusculis, umbellis erectis lateralibus solitariis, caule simplici. *Willd. sp. pl. 1. p. 1266.*

Icon. *Dil.*  
On the ba-  
nia to  
white.

6. A. caule  
lati s ut  
ribus o  
*Mich.*

Icon. *Jac.*  
*Willd.* l.  
In swampy  
Aug.  
in ever-  
the lea-  
upper

7. A. caule  
oblongo-  
minal  
*Willd.*

Icon. *D.*  
In wet  
New  
Flowe

8. A. caule  
erecti  
Icon. *L.*  
In shade  
Aug.

9. A. caule  
obtuse-  
umbra-  
subgr.  
In dry  
v. v.

10. A. caule  
nudica-  
*Wil-*

A. hyb.  
Icon.  
On dry  
Yor  
some

Icon. Dill. elth. t. 29. f. 32. Bot. mag. 1187.

On the banks of rivers in sandy or gravelly soil : Virginia to Carolina. 4. July, Aug. v. v. Flowers white.

6. A. caule erecto superne ramoso tomentoso, foliis lanceolatis utrinque subtomentoso-lanuginosis, umbellis pluribus origine geminis, corniculis appendicis exsertis. Mich. fl. amer. 1. p. 115. Willd. sp. pl. 1. p. 1267.

Icon. Jacq. hort. t. 107.

- $\beta$ . Willd. t. c.

In swamps and on river banks ; common. 4. June — Aug. v. v. Flowers red. The variety  $\beta$ . is more hairy, in every other respect no ways different ; sometimes the leaves are quite smooth, and only pubescent on the upper part of the stem.

7. A. caule simplici bifarium puberulo, foliis subsessilibus oblongo-ovalibus subtus pubescentibus, umbellis terminalibus nectariisque erectis, appendicibus exsertis. — Willd. sp. pl. 1. p. 1265.

Icon. Dill. elth. t. 27. f. 30.

In wet meadows, on the edges of woods and ditches : New England to Virginia. 4. July, Aug. v. v. Flowers beautiful purple.

8. A. caule simplici, foliis ovatis subtus villosis, umbellis erectis, nectariis resupinatis. Willd. sp. pl. 1. p. 1265.

Icon. Dill. elth. t. 28. f. 31.

In shady swamps : Virginia and Carolina. 4. July, Aug. v. v. Flowers purple.

9. A. caule simplici erecto hirsuto, foliis lanceolato-oblongis obtusiusculis subsessilibus utrinque tomentoso-hirsutis, umbellis lateralibus solitariis subsessilibus nutantibus subgloboso-densifloris, appendicibus nullis.

In dry fields : Pennsylvania to Virginia. 4. June, July. v. v. Flowers green.

10. A. caule simplici erecto, foliis ovatis petiolatis rugosis nudis, umbellis subsessilibus pedicellis tomentosis. Willd. sp. pl. 1. p. 1265. Walt. fl. car. 104.

A. hybrida. Mich. fl. amer. 1. p. 115.

Icon. Pluk. alm. t. 77. f. 1. Bot. mag. 1182.

On dry sunny hills, on the borders of woods : New York to Carolina. 4. July, Aug. v. v. A very handsome species ; flowers in dense umbels ; petals and nectaries white ; fructification red.

incarnata.

pulchra.

amaena.

purpurascens

viridiflora.

variegata.

- obtusifolia.* 11. *A. foliis amplexicaulibus oblongis rotundato-obtusis undatis, umbella terminali longius pedunculata multiflora glabra, corniculis exsertis.* *Mich. fl. amer. 1. p. 115.*  
*A. purpurascens.* *Walt. fl. car. 105.*  
In sandy fields: New Jersey to Carolina.  $\frac{1}{4}$ . June-Aug. v. v. Flowers large, purple.
- amplexicaulis.* 12. *A. tota glauca; caulis decumbentibus simplicibus, foliis cordatis amplexicaulibus venosis, corniculis delitescentibus compresso-ligulatis.* *Mich. fl. amer. 1. p. 115.*  
*A. humistrata.* *Walt. fl. car. 105.*  
In barren sandy fields: Carolina and Georgia.  $\frac{1}{4}$ . June, July. v. v. Leaves red-veined, thick, flowers handsome, red.
- acuminata.* 13. *A. caule erecto glabro simplici, foliis ovatis subcordatis acuminatis brevi-petiolatis, superioribus sessilibus; umbellis lateralibus solitariis erectis.*  
*A. cordata.* *Walt. fl. car. 105.*  
In deep cedar-swamps, and on the banks of rivers: New Jersey.  $\frac{1}{4}$ . July. v. v. Flowers red and white; stem about two feet high; roots very large, resembling those of *A. tuberosa*.
- laurifolia.* 14. *A. caule erecto glabro, foliis subsessilibus ovali-lanceolatis sensim angustatis acutissimis glabris; supra ad oras asperiusculis.* *Mich. fl. amer. 1. p. 117.*  
On the river Althamaha, Georgia. *Michaux.*  $\frac{1}{4}$ . July. v. s. Flowers purple.
- paupercula.* 15. *A. caule erectiusculo glabro oligophyllo, sursum nudo, foliis linearibus longissimis glabriusculis, umbellis terminalibus paucifloris.* *Mich. fl. amer. 1. p. 118.*  
*A. lanceolata.* *Walt. fl. car. 105.*  
In sandy wet meadows: Virginia and Lower Carolina.  $\frac{1}{4}$ . June, July. v. v. Flowers red, with yellow appendices.
- pedicellata.* 16. *A. caule simplici erecto piloso, foliis elliptico-linearibus pubescentibus supra punctato-asperis, umbella lateralis solitaria pauciflora, petalis erectis calyce multoties longieribus, nectariis lunulatis patentibus corniculis nullis.* *Walt. fl. car. 106.*  
On dry sandy fields of South Carolina. *Walter.* Near Fort Barrington. *Lyon.*  $\frac{1}{4}$ . June. v. s. in Herb. Lyon, nec non *Walter.* The whole plant not above

- itus una multi-  
amer. 1.  
1. June—  
nplibus,  
iculis de-  
amer. 1.  
vers hand-  
ubcordatis  
essilibus ;  
vers: New  
d white ;  
esembling  
lanceola-  
ora ad oras  
4. July.  
sum nudo,  
abellis ter-  
118.  
Carolina,  
yellow ap-  
-linearibus  
lla lateralis  
multoties  
corniculis  
ter. Near  
in Herb.  
not above
- three inches high ; flowers large in proportion, greenish yellow ; fructification on a pedicel.
17. *A. caulis erectis, foliis petiolatis oblongis obtusis laevibus, umbellis paucis terminalibus, petalis erectis calyce multo longioribus, nectariis integris patulis brevibus.* —Walt. fl. car. 107.

South Carolina. Walter. 4. v. s. in Herb. Lyon.

Flowers large ; petals green ; nectaries purple ; fructification white. This and the preceding species deviate from the general character of *Asclepias*, and probably constitute a distinct genus ; but having had no opportunity to examine the living plant, I refer them to this genus until another opportunity may be had to investigate them.

18. *A. caulis erectis simplicibus glabris, foliis ovatis acuminatis petiolatis : in medio cauli majoribus quaternis, umbellis duabus terminalibus laxifloris, pedicellis filiformibus.* —Jacq. obs. 2. t. 23.

On lime-stone rocks : New York to Virginia. 4. May —July. v. v. From nine inches to a foot high ; flowers small, sweet-scented ; petals white ; nectaries rose-coloured.

*vividis.*

*quadrifolia.*

- \*\* *Folii sparsis.*
19. *A. caule erecto simplicissimo lineatum puberulo, foliis angustissime linearibus strictis glabris plerumque verticillatis, corniculis exsertis.* Mich. fl. amer. 1. p. 116. Wild. sp. pl. 1. p. 1272.

Icon. Pluk. mant. t. 336, f. 4.

On dry hills and fields : New Jersey to Carolina. 4. June, July. v. v. Petals yellowish-green ; nectaries white.

20. *A. caule decumbente foliisque sparsis prælongo-linearibus puberulis, appendicibus corollæ corpusculo genitilifero brevioribus et corniculo vacuis.* Mich. fl. amer. 1. p. 116.

*A. incarnata.* Walt. fl. car. 106.

*A. floridana.* Lam. encycl. 1. p. 284.

In shady wet woods : Carolina and Georgia. 4. July, Aug. v. s. Flowers pale purple ; fructification yellow.

*verticillata.*

*longifolia.*

21. *A. caule erectiusculo summitate divaricato-ramoso hirsutissimo, foliis sparsis oblongo-lanceolatis hirsutis,*

*tuberosa.*

- umbellis subcorymboso-terminalibus.—Willd. sp. pl. 1. p. 1273.*  
Icon. *Dill. elth. t. 30. f. 34.*
- decumbens.* *β. A. caule decumbente, foliis sublinearibus hirsutissimis umbellis lateralibus. Willd. sp. pl. 1. p. 1268.*  
In stony or sandy fields and woods; common. *β.* In barren fields of Carolina. *♀. June, July. v. v.* Flowers very showy; petals and nectaries bright orange; fructification yellow.
224. *b. CEROPEGIA. Gen. pl. 431.*
- palustris.* 1. *C. foliis linearibus integerrimis, floribus umbellatis, pedunculis longitudine foliorum, calycis laciniis linearibus.*  
In salt marshes, winding round Scirpi and Junci: Carolina. *Lyon. ♀. v. s. in Herb. Lyon. nec non Walter.* The whole habit of the plant is that of *C. tenuiflora. Willd. sp. pl. 1. p. 1276.* the flowers only considerably smaller.
225. *AMSONIA. Walt. fl. car. 98.*
- tatifolia.* 1. *A. caule glabriusculo, foliis ovali-lanceolatis superioribus promisse acuminatis subtus ad nervos pubescensibus.—Mich. fl. amer. 1. p. 121.*  
Tabernæmontana Amsonia. *Willd. sp. pl. 1. p. 1246.*  
Icon. *Pluk. phyt. t. 115. f. 3.*  
In shady wet woods: Carolina. *♀. May. v. v.* Flowers in compound corymbose racemes, blue.
- salicifolia.* 2. *A. caule laevigato, foliis linear-lanceolatis utrinque acutis glaberrimis.*  
In Carolina and Georgia. *Lyon. ♀. May. v. v.* Flowers the same as the first species, more abundant.
- angustifolia.* 3. *A. caule pubescente, foliis angusto-linearibus crebris erectis pubescensibus. Mich. fl. amer. 1. p. 121.*  
Amsonia ciliata. *Walt. fl. car. 98.*  
In sandy barrens of Carolina and Georgia. *♀. May, June. v. s. in Herb. Walter. &c.* Flowers of the same disposition and colour as both the foregoing species.
226. *GELSEMINUM. Juss. gen. pl. p. 168.*
- nitidum.* 1. *G. scandens, glaberrimum; foliis lanceolatis, fasciculis axillaribus paucifloris. Mich. fl. amer. 1. p. 120.*

Bignonia  
AnonymoIcon. *Cate*

On the se

June, J  
grant;22  
1. *G. caule latis ac ciliatis, p. 1352.**G. fimbri*In dry pa  
♂. Au

elegant

2. *G. caule floribus peduncul rotund Willd.*Icon. *Fl.*

On high lets:

Flowe

3. *G. caule floribus que, obtusi sp. pl.**G. fimbri**G. Cates*Icon. *C**rep. 4*

In low r

Aug.

4. *G. caule tis as nalib cinii Will**G. Sap**p. 17**G. villa*

*Bignonia sempervirens*. Willd. sp. pl. 3. p. 291.  
*Anonymos sempervirens*. Walt. fl. car. 98.  
 Icon. Catesb. car. 1. t. 53. Pluk. alm. t. 112. f. 5.  
 On the sea-coast near rivers : Virginia to Florida. ♂.  
 June, July. v. v. Flowers large, deep yellow, fragrant ; known by the name *Carolina Jessamine*.

## 227. GENTIANA. Gen. pl. 450.

1. *G. caule tereti, ramis elongatis unifloris, foliis lanceolatis acutis, corollis 4-fidis, laciniis obovatis inciso-ciliatis, plicis interioribus simplicibus*.—Willd. sp. pl. 1. p. 1352. *crinita*.  
*G. fimbriata*. Bot. rep. 509.  
 In dry pastures and woods : New York to Carolina. ♂. Aug. Sept. v. v. Flowers a beautiful sky-blue, elegantly fringed on their edges.
2. *G. caule tereti, foliis linear-lanceolatis obtusiusculis, floribus terminalibus fasciculatis, lateralibus solitariis pedunculatis, corollis 5-fidis campanulatis : laciniis rotundatis, plicis interioribus acute-unidentatis*.—Willd. sp. pl. 1. p. 1336. *Pneumonanthe*.  
 Icon. Fl. dan. 269. Bot. mag. 1101.  
 On high mountains, near the borders of lakes and rivulets : Canada to Pennsylvania. ♀. Aug. Sept. v. v. Flowers large, dark-blue.
3. *G. caule tereti glabro, foliis oblongo-lanceolatis trinerviis, floribus sessilibus fasciculatis terminalibus axillaribusque, corollis 5-fidis campanulatis ventricosis : laciniis obtusis, plicis interioribus laciniis dentatis*.—Willd. sp. pl. 1. p. 1338. *Saponaria*.  
*G. fimbriata*. Vahl 3. p. 47.  
*G. Catesbaei*. Walt. fl. car. 109.  
 Icon. Catesb. car. 1. t. 70. Bot. mag. 1039. Bot. rep. 418.  
 In low meadows and woods : Canada to Carolina. ♀. Aug. Sept. v. v. Flowers very crowded, fine blue.
4. *G. caule subangulato scabriusculo, foliis ovato-lanceolatis asperiusculis, floribus sessilibus fasciculatis terminalibus, corollis 5-fidis campanulatis ventricosis : laciniis acutis, plicis interioribus simplicibus acutis*.—Willd. sp. pl. 1. p. 1338. *ochroleuca*.  
*G. Saponaria*. Walt. fl. car. 109. Mich. fl. amer. 1. p. 176.  
*G. villosa*. Willd. sp. pl. 1. p. 1338.

Icon. Pluk. alm. t. 186. f. 1.

In dry sandy fields and on gravelly hills: Pennsylvania to Florida; more scarce. ♀. Aug. Sept. v. v. Not so tall as the preceding species; flowers outside of a yellowish-green, inside, blue and striped with purple. That the present plant is the same with *G. villosa*, there remains no doubt, having seen the Gronovian specimen in the Banksian Herbarium, which is the same with *G. Saponaria* of Walter, and *G. ochroleuca* of Frölich.

*linearis.*

5. *G. caule asperiusculo, foliis linear-lanceolatis undulosis uti calyces ciliolatis, floribus 5-fidis campanulatis terminalibus sessilibus congestis, laciniis brevibus obtusis, plicis interioribus laciniis denticulatis.* — Willd. sp. pl. 1. p. 1339.

*G. puberula.* Mich. fl. amer. 1. p. 176.

In valleys, on the mountains of Pennsylvania and Virginia. ♀. Aug. v. v. Flowers large, blue.

*amareloides.*

6. *G. caule quadrangulo ramoso, foliis semiamplexicaulibus ovalibus acutis trinervibus, floribus 5-fidis tubulosos campanulatis fauce glabris terminalibus et lateralibus ternis quinisve subpedicellatis, laciniis lanceolatis acuminatis, plicis simplicibus, calycibus brevissimis angustis.* — Mich. fl. amer. 1. p. 175.

*G. quinqueflora.* Willd. sp. pl. 1. p. 1339.

On hills, in shady dry places: New York to Kentucky. ♂. July, Aug. v. v. Flowers small, pale blue, not yellow as they are represented by Michaux.

*acuta.*

7. *G. caule 4-gono, foliis sessilibus subamplexicaulibus erectis oblongo-ovatis acutissimis, floribus fasciculatis terminalibus et lateralibus, corollis 5-fidis, laciniis linear-lanceolatis, plicis ciliatis.* Mich. fl. amer. 1. p. 177.

On the high mountains of Carolina and in Canada: Michaux. +. Flowers very small, greenish-yellow.

*angustifolia.*

8. *G. caule tereti simplici unifloro, foliis angusto-linearibus patulis, corollis 5-fidis, laciniis ovalibus, plicis interioribus laciniis laceris.* — Mich. fl. amer. 1. p. 177.

*G. purpurea.* Walt. fl. car. 109.

In meadows and near rivers, in sandy soil: New Jersey to Carolina. ♀. Aug. Sept. v. v. Flowers large, a beautiful azure blue.

1. *H. spinosa,*  
axillaris  
*Walt. fl.*  
*H. carolinia*  
In stagnant  
Flowers

1. *D. pubescens*  
concolor  
*amer. 1*  
Anonymo  
Demidofia  
In sandy  
South  
creepin

1. *E. erectus*  
pedunculata  
On the b.  
yellow

1. *H. viscidum*  
liis m  
tatis o  
choto launc  
sertis.

*H. amei*  
*H. Cort*  
Icon. P  
In shad  
frequ  
simpl  
with

2. *H. pulchellum*  
tus g  
rotui confi

228. HYDROLEA. *Gen. pl. 445.*

1. *H. spinosa*, pilosa; foliis longo-lanceolatis, pedunculis quadrivalvis. axillaribus brevissimis 1—5-floris, capsulis glabris.—  
*Walt. fl. car. 110.*  
*H. caroliniana*, *Mich. fl. amer. 1. p. 177.*  
In stagnant waters of Lower Carolina.  $\frac{1}{2}$ . July. v. s.  
Flowers blue.

229. DICHONDRA. *Gen. pl. 451. Mich. fl. amer. 1.*  
*p. 135.*

1. *D. pubescens*; foliis absque emarginatura reniformibus *carolinensis*. concoloribus, calycibus villoso-ciliatis.—*Mich. fl. amer. 1. p. 136.*  
*Anonymos repens*. *Walt. fl. car. 110.*  
*Demidofia repens*. *Gmel. syst. nat. 458.*  
In sandy fields, on road sides, about Charlestown, South Carolina.  $\frac{1}{2}$ . June—Sept. v. v. A small creeping plant, with small herbaceous coloured flowers.

230. EVOLVULUS. *Gen. pl. 524.*

1. *E. erectus*; foliis oblongis utrinque sericeo-tomentosis, *argenteus*. pedunculis unifloribus brevibus.  
On the banks of the Missouri. *Nuttall. v. s.* Flowers yellow.

231. HEUCHERA. *Gen. pl. 447.*

1. *H. viscidio-pubescent*; scapo foliisque asperiusculis, foliis modice rotundato-lobatis dentatis: dentibus dilatatis obtusis mucronatis, pedunculis paniculæ tres dichotomis divaricatis, calycibus brevibus obtusis, petalis lanceolatis longitudine calycis, staminibus longe exsertis.  
*H. americana*. *Willd. sp. pl. 1. p. 1328.*  
*H. Cortusa*. *Mich. fl. amer. 1. p. 171.*  
*Icon. Pluk. alm. t. 58. f. 3.*  
In shady rocky situations: New England to Carolina; frequent.  $\frac{1}{2}$ . May—July. v. v. Flowers in a long simple panicle, small; petals red. It varies sometimes with nearly smooth leaves.  
2. *H. pulverulento-pubescent*; scapo inferne foliisque subtus glabris, foliis subacuto-lobatis dentatis: dentibus rotundatis mucronatis, pedunculis paniculæ brevibus confertifloris, calycibus campanulatis majoribus, pe-

*viscida.**pubescens.*

## PENTANDRIA DIGYNIA. Heuchera.

talis spathulatis calyce longioribus, staminibus vix exertis.

On the Blue mountains of Pennsylvania and Virginia. 2d. May, June. v. v. Flowers larger than the rest of the genus ; petals pale red with yellow.

*hispida.*

3. *H. hispida*, scabra ; scapo petiolis foliisque subtus glabris, foliis acute-lobatis, supra hispido-pilosis dentatis : dentibus brevissimis subretusis mucronatis, pedunculis paniculæ paucifloris, calycibus mediocribus subacutis, petalis spathulatis longitudine calycis, staminibus exertis.

On high mountains of Virginia and Carolina. 2d. May, June. v. v. Petals purple ?

*villosa.*

4. *H. villosissima* ; foliis acute lobatis dentatis : dentibus acutis mucronatis, fasciculis paniculæ capillaribus laxifloris, calycibus parvis, staminibus exertis.—*Mich. fl. amer. 1. p. 172.*

On high mountains of North Carolina and Virginia ; collected on the Peaks of Otter. 2d. June. v. v. Flowers small ; petals white.

*caulescens.*

5. *H. basi suffruticosa* ; scapo basi petiolisque pilosis, foliis supra glabris subtus ad nervos pilosis acute-lobatis ciliatis dentatis ; dentibus acutis mucronatis, calycibus brevibus villosis, petalis linearibus calyce duplo longioribus, staminibus exertis.

On high mountains of Carolina. *Lyon.* 2d. May, June. v. v. Petals white. In the Herbarium of A. B. Lambert, Esq. are specimens of a *Heuchera*, collected by Pallas in Kamtschatka, which appear to be the same with the present species.

232. ZIZYPHUS. *Lam. encycl. 3. p. 303.*

*vulbilis.*

1. *Z. inermis* ; foliis ovatis costato-venosis integerrimi subundulatis, umbellis pedunculatis axillaribus, caule volubili. *Willd. sp. pl. 1. p. 1102.*

*Rhamnus vulbilis.* *Linn. suppl. 152.* *Walt. fl. car. 101.* *Icon. Jacq. ic. 2. t. 336.* *Schmidt arb. 153.*

In deep swamps, near the sea-coast : Virginia to Carolina. 2d. June. v. v. Flowers small, greenish-yellow ; berries oblong, violet-coloured. It ascends the highest trees of *Cupressus disticha*, in the Dismal swamp, near Suffolk in Virginia, and is known there by the name of *Supple-Jack*.

## 233. ERYNGIUM

1. *E. praealtum*  
ratis : ra-

capitulo  
petuluis pa-

eryng. p.

*E. aquatitic*  
*amer. 1.*  
*Icon. Dela-*

*On the bar-*  
*Pensylv-*  
*hollow ;*  
*found a*

flowers,

2. *E. simplex*  
capitulis  
leisque

—*Lam.*

*E. ovalifol*

*E. integrif*

*Icon. Del-*

*In wet me-*  
*July.*  
*round,*

3. *E. pumila*  
calibus  
drincis s-

lati.—

*Icon. Br-*

*In dry fie-*  
*Flower*  
*teemed*

4. *E. elatun*  
floralib  
globos-

tomo.—

*Icon. Ja-*

*In Swa-*  
*Somet*  
*flower*

233. ERYNGIUM. Gen. pl. 456. Delaroche monog.  
eryng.

1. E. praealtum ; foliis longissime lanceolato-linearibus serratis : rameis linearis-multipartitis, involucri foliolis capitulo ovoideo longioribus paleisque 3—5-fidis, capitulis paniculatis.—Lam. encycl. 4. p. 759. Delaroche eryng. p. 48.  
E. aquaticum β. Willd. sp. pl. 1. p. 1357. Mich. fl. amer. 1. p. 163.  
Icon. Delaroche eryng. t. 19. Pluk. alm. t. 396. f. 3.  
On the banks of rivers, in overflowed tide-meadows : Pennsylvania to Virginia. 4. July, Aug. v. v. Stems hollow ; flower-heads green or pale blue. I once found a beautiful proliferous variety with pale purple flowers, one on the top of the others.
2. E. simplex ; foliis ovato-cordatis brevissime petiolatis, capitulis omnibus pedunculatis, involucri foliolis paleisque 3-fidis, caule virgato superne tantum ramoso.—Lam. encycl. 4. p. 757. Delaroche eryng. p. 49.  
E. ovalifolium. Mich. fl. amer. 1. p. 163.  
E. integrifolium. Walt. fl. car. 112.  
Icon. Delaroche eryng. t. 20.  
In wet meadows and woods : Carolina and Georgia. 4. July. v. v. About a foot high ; flower-heads small, round, amethystine-blue.
3. E. pumilum, graveolens ; foliis serrato-spinosis : radicibus lanceolatis, floralibus palmatis, capitulis cylindricis subsessilibus, paleis longissimis linearis-lanceolatis.—Willd. sp. pl. 1. p. 1356.  
Icon. Brown. jam. 1. t. 156. f. 3. 4.  
In dry fields : Georgia to Florida. 4. July, Aug. v. v. Flowers in small heads, herbaceous. Very much esteemed among the natives as a medicine.
4. E. elatum ; foliis late-linearibus remote ciliato-spinosis, floralibus lanceolatis dentatis, foliolis involucri capitulo globoso brevioribus paleisque integris, caule subdichotomo.—Willd. sp. pl. 1. p. 1357.  
Icon. Jacq. ic. 2. t. 347. Pluk. alm. t. 396. f. 3.  
In swamps : Virginia to Carolina. 4. Aug. v. v. Sometimes more than six feet high ; leaves very long ; flower-heads pale green.

234. HYDROCOITLE. *Gen. pl. 457.**americana.**flexicaulis.**gracilis.**vulgaris.**umbellata.**repanda.**lineata.**composita.*

1. *H. glabra, tuberosa; foliis subpeltato-orbiculatis dupli-cato-crenatis, glomerulis paucifloris sessilibus.* — *Willd. sp. pl. 1. p. 1361.*

*a. H. caulibus firmioribus brevioribusque quasi angulatum flexuosis.* *Mich. fl. amer. 1. p. 162.*

*β. H. gracilior, filiformis, non flexuosa; foliis minoribus.* *Mich. l. c.*

In bogs and sphagnous swamps: *α.* Canada to Virginia. *β.* In the mountains of Carolina, Virginia, and in Canada. *4.* June—Aug. *v. v.* Flowers exceeding small, and of a greenish colour, as is the case with all the following species.

2. *H. foliis orbiculatis peltatis leviter crenatis, scapo interrumpo-spicato paucifloto.* — *Willd. sp. pl. 1. p. 1360.*

*Icon. Fl. dan. 90. Engl. bot. 751.*

In swamps and overflowed places: Canada to Georgia. *4.* May—July. *v. v.*

3. *H. foliis peltatis crenatis basi emarginatis, umbellis pendulatis multifloris, floribus pedicellatis.* *Willd. sp. pl. 1. p. 1361.*

In wet fields and mossy swamps: Virginia and Carolina. *4.* May—July. *v. v.*

4. *H. foliis rotundato-cordatis subangulato-repandis, petiolis nervisque pilosis, capitulo pedunculato trifloro villosa.* *Pers. syn. 1. p. 302.*

*H. ficaroides.* *Mich. fl. amer. 1. p. 161.*

*H. reniformis et cordata.* *Walt. fl. car. 113.*

On overflowed banks of rivers in Lower Carolina and Georgia. *4.* June, July. *v. v.* Seeds larger than the preceding species.

5. *H. minuta, glaberrima; foliis crassiusculis linear-i-cuneatis transversim lineatis, umbella pedunculata.* — *Mich. fl. amer. 1. p. 161.*

In overflowed places of Lower Carolina. *4.* April, May. +.

6. *H. radice globoso-bulbosa, caule simplici diphyllo bifido, folio laterali 3-partito, partitionibus subpinnatis, pinnulis divisis: terminali sub-biternato, umbellis terminalibus 3—5-floris.*

Sison bulbosum. *Mich. fl. amer. 1. p. 169.*

On rocks. in wet shady places, near Knoxville, Kentucky. *Michaux. 4.* March, April. *v. s. in Herb.*

Enslen.

dried spe

belongs.

The flow

2

1. P. radice fr

bus acu

p. 1124.

Icon. Cate

mag. 13

In the m

shady w

the fan

by the

tion in

for it.

2. P. radice

natives

ratis.—

Icon. Bo

In low sh

sylvani

easily e

1. S. foliis

bus se

sis.—

Icon. Ja

In dry w

July.

bur.

2. S. foliis

pl. 1.

In Virg

1. D. sem

sp. p

The W

dry 1

*Enslen.* As far as I have been able to learn from a dried specimen, there remains no doubt that this plant belongs to this genus rather than any other I know. The flowers have white petals and dark purple anthers.

235. PANAX. *Gen. pl. 1604.*

1. *P. radice fusiformi, foliis ternis quinatis, foliolis ovali-quinquefoli-  
bus acuminatis-petiolatis serratis.* — *Willd. sp. pl. 4. p. 1124.*

*Icon. Catesb. car. 3. t. 16. Trew ehret. t. 6. f. 1. Bot.  
mag. 1333.*

In the mountains from Canada to Tennessee, in rich shady woods.  $\frac{1}{2}$ . May. v. v. This plant produces the famous root called *Ginseng*, so much esteemed by the Chinese; it formerly was an article of exportation in America, but at present there is little demand for it.

2. *P. radice subrotundo-bulbosa; foliis ternis ternatis qui-  
natise, foliolis oblongo-lanceolatis subsessilibus ser-  
ratis.* — *Willd. sp. pl. 4. p. 1124.*

*Icon. Bot. mag. 1334.*

In low shady woods, near rivulets: New York and Pennsylvania.  $\frac{1}{2}$ . May. v. v. A small delicate plant, easily overlooked even when in search of it.

*trifolium.*

236. SANICULA. *Gen. pl. 458.*

1. *S. foliis digitatis, foliolis oblongis incisis, floribus fertili-  
bus sessilibus subternis, sterilibus pedicellatis numero-  
sis.* — *Willd. sp. pl. 1. p. 1367.*

*Icon. Jacq. ic. 2. p. 348.*

In dry woods, frequent: Canada to Carolina.  $\frac{1}{2}$ . June, July. v. v. Flowers greenish-white; fruit a little bur.

2. *S. foliis radicalibus compositis, foliolis ovatis.* *Willd. sp. pl. 1. p. 1366.*

In Virginia.  $\frac{1}{2}$ . A doubtful species.

237. DAUCUS. *Gen. pl. 466.*

1. *D. seminibus hispidis, petiolis subtus nervosis.* *Willd. sp. pl. 1. p. 1389.*

The *Wild Carrot* is one of the most common weeds in dry fields.

*Carota.*

*pusillus.*

2. D. retrosum-hispidus ; foliolis laciinis minimis sublan-  
ceolatis, umbellis parvis, umbellulis paucifloris, semi-  
nibus 8-fariam cristato-muricatis.—*Mich. fl. amer. I.*  
*p. 164.*  
In dry fields of Carolina. *Michaux.* +.

238. AMMI. *Gen. pl. 467.**capillaceum.*

1. A. caule ramoso, foliis omnibus composite capillaceo-  
multipartitis, involucris umbella brevioribus multipartitis,  
seminibus glabratris.  
*A. majus. Walt. fl. car. 113.*  
In sandy fields of Carolina and Georgia. ☀. June, July.  
*v. s.* The plant small, with very finely divided leaves.

239. SELINUM. *Gen. pl. 470.**canadense.*

1. S. glaberrimum, lucidum ; foliis bipinnatis, foliolis mul-  
tipartitis, laciinis lanceolatis, fructibus ovalibus. *Mich.*  
*fl. amer. I. p. 165.*  
*Apium bipinnatum. Walt. fl. car. 115.*  
On the mouths of large rivers from Canada to Carolina.  
¶. July. v. v. Flowers white.

240. FERULA. *Gen. pl. 475.**villosa.*

1. F. foliis supradecomposito-ternatis, foliolis ovatis serratis  
rigidis venosis, caule pedunculis umbellisque villosis.  
*Walt. fl. car. 115.*  
*Cicuta venenata. Philad. philos. trans.*  
On dry hills : New York to Carolina. ¶. July, Aug.  
*v. v.* Flowers white. In the Philosophical Transactions  
of Philadelphia, a description of this plant has  
been given, and a full account of its poisonous qua-  
lities.

241. HERACLEUM. *Gen. pl. 477.**lanatum.*

1. H. foliorum petiolo nervisque subtus lanato-vilosissimis,  
foliis trifoliatis, foliolis omnibus petiolatis amplis sub-  
rotundo-cordatis subpalmato-lobatis, seminibus orbicu-  
latis. *Mich. fl. amer. I. p. 166.*  
In shady woods, on rich moist soil : Canada to Pensyl-  
vania ; rare. ¶. June, July. v. v. Flowers white.

- 24
1. L. foliis *ralibus strictis. Mich.*  
On the ri
  2. L. foliis *dariis 2 tis, in subula*  
On the ri  
Staunt  
feet hi
  3. L. ? cau  
foliis p  
fructib  
*fl. ame*  
In Upper  
I suspe  
bably

1. A. petiol  
foliolis  
sessili,  
*p. 167*  
In Cana  
July.
2. A. foliis  
subses  
lato.—  
In Cana  
Aug.  
grant;
3. A. foliol  
*pl. 1.*  
In shad  
*v. v.*

242. LIGUSTICUM. *Gen. pl. 478.*

1. *L. foliis caulinis binternatis, summis trifoliatis, foliis lateribus trapeziformibus, terminali rhombeo, umbellis strictis, involucro involucellisque linearibus polyphyllis.* *Mich. fl. amer. 1. p. 166.* *scoticum.*  
On the river St. Laurence. *Michaux.*  $\frac{1}{4}$ . +. July.
2. *L. foliis petiolo duplato-tripartito, partitionibus secundariis 2-3-foliolatis, foliolis ovalibus æqualiter dentatis, involucellis setaceis, fructibus oblongo-ovalibus subulato-10-costatis.* *Mich. fl. amer. 1. p. 166.* *actæifolium.*  
On the river St. Laurence. *Michaux.* In Virginia near Staunton, &c.  $\frac{1}{4}$ . July. v. v. More than three feet high.
3. *L. ? caule lævi glabroque, nodis confertim barbulatis, foliis plerisque binternatis margine nervisque puberulis, fructibus ovalibus utrinque margine subdipteris.* *Mich. fl. amer. 1. p. 167.* *barbinode.*  
In Upper Carolina. *Michaux.*  $\frac{1}{4}$ . v. s. in *Herb. Lyon.*  
I suspect this species to belong to another genus, probably *Sium.*

243. ANGELICA. *Gen. pl. 479.*

1. *A. petiolo tripartito, partitionibus 5-foliolatis, foliolis inciso-dentatis : terminalium impari rhombeo sessili, lateralibus decursivis.* *Mich. fl. amer. 1. p. 167.* *triquinata.*  
In Canada and on the mountains of Virginia.  $\frac{1}{4}$ . June, July. v. v. Smooth.
2. *A. foliis compositis, foliolis oblongis sublobatis serratis subsessilibus : extimo pari coadunato : terminali petiolato.* — *Willd. sp. pl. 1. p. 1430.* *atropurpurea.*  
In Canada and on the mountains of Virginia.  $\frac{1}{4}$ . July, Aug. v. v. From three to six feet high ; root very fragrant; petals purple.
3. *A. foliolis æqualibus ovatis inciso-serratis.* *Willd. sp. pl. 1. p. 1430.* *lucida.*  
In shady woods : Canada to Pennsylvania.  $\frac{1}{4}$ . June, July. v. v. Flowers white.

244. SIUM. *Gen. pl. 480.**rigidius.*

1. *S. foliis pinnatis, foliolis lanceolatis integerrimis vel superne dentatis.* *Willd. sp. pl. 1. p. 1433.*

In wet meadows : Pennsylvania to Virginia. 24. July, Aug. v. v. Flowers small. This and the following are considered very poisonous plants, especially in respect to horned cattle ; and every farmer who knows their bad qualities is busily employed to destroy them.

*lineare.*

2. *S. foliis pinnatis, foliolis longis sublanceolato-linearibus remotiuscule serratis, involucro oligophyilo, involucellis linearibus polyphyllis, umbella breviter radiata.* *Mich. fl. amer. 1. p. 167.*

*S. suave.* *Walt. fl. car. 115.*

In wet meadows and along ditches : Canada to Pennsylvania. 24. July. v. v.

*longifolium.*

3. *S. foliis pinnatis, foliolis inferioribus longissimis falcato-linearibus rarer dentatis, caule oligophyllo superne nudo, umbellis subgeminis nudiusculis.*

In ditches and bogs : New Jersey. 24. Aug. v. v. Very slender.

244. b. SISON. *Gen. pl. 481.**pusillum.*

1. *S. foliis binternatim multipartitis, umbellulis 3—5-floris, seminibus scaberulis.* *Mich. fl. amer. 1. p. 168.*

*Daucus divaricatus.* *Walt. fl. car. 114.*

In dry sandy fields : Carolina. ① April, May. v. s.

*trifoliatum.*

2. *S. foliis omnibus trifoliatis, foliolis dentatis, inferioribus ovalibus 2—3-lobis, superioribus ovali-lanceolatis, umbella terminali solitaria pedunculata, seminibus subrotundis.* *Mich. fl. amer. 1. p. 168.*

In Upper Carolina. *Michaux.* +.

*marginatum.*

3. *S. foliis pinnatis, superioribus quinatis, foliolis omnibus sessilibus lanceolatis integerrimis albido-marginatis, involucro involucellisque nullis.* *Mich. fl. amer. 1. p. 168.*

In wet meadows : Virginia to Carolina. 24. July, Aug. v. v.

245. CENANTHE. *Gen. pl. 484.**carolinensis.*

1. *O. foliis omnibus simplicibus filiformibus.* *Walt. fl. car. 113.*

South Carolina. *Walter.* +.

## 246. P

1. *P. foliorum*

p. 1444.

On the wat

July. v

country

1. *C. foliorum*

apice bila

Icon. *Pluk*

In wet mea

24. Jul

2. *C. foliis fa**Willd. s*

In overflow

Jersey.

## 249. C

1. *C. seminib*

procumb

Scandix pr

On shady

① May

served it

2. *C. foliis te*

culis gen

Sison canad

Common i

v. v.

3. *C. petiolo*

oblongo-

ci-radiat

vigatis.—

246. PHELLANDRIUM. *Gen. pl. 485.*

1. *P. foliorum ramificationibus divaricatis. Willd. sp. pl. 1. aquaticum. p. 1444.*

On the waters of the Rocky-mountain. *M. Lewis.* 24 July. *v. s. in Herb. Lewis.* The Indians of that country use it as a medicine in the diseases of horses.

247. CICUTA. *Gen. pl. 486.*

1. *C. foliorum serraturis mucronatis, petiolis membranaceis apice bilobis. Willd. sp. pl. 1. p. 1446.*

*Icon. Pluk. alm. t. 76. f. 1.*

In wet meadows and ditches : New England to Carolina. 24 July, Aug. *v. v.*

2. *C. foliis fœniculaceis : ramis verticillatim bulbiferis.— Willd. sp. pl. 1. p. 1445. exclus. syn. Gronov.*

In overflowed meadows and pastures : Canada to New Jersey. 24 July. *v. v.* A very singular small plant.

248. CONIUM. *Gen. pl. 469.*

1. *C. seminibus striatis. Willd. sp. pl. 1. p. 1395.*

*Icon. Jacq. austr. 156.*

In cultivated grounds and near habitations ; probably introduced : New York and Virginia. 3. June—Aug. *v. v.* Very poisonous.

249. CHÆROPHYLLUM. *Gen. pl. 490.*

1. *C. seminibus nitidis lœvibus, foliis decompositis, caule procumbente, umbellis paucifloris. Pers. syn. 1. p. 320.*

*Scandix procumbens. Willd. sp. pl. 1. p. 1452.*

On shady wet hills, on the foot of mountains : Virginia. 25 May. *v. v.* A small and scarce plant ; I observed it near Hager's Town, Maryland.

2. *C. foliis ternatis, foliolis ovatis acutis subincisis, pedunculis geminis, fructibus oblongis.—Pers. syn. 1. p. 320.*

*Sison canadense. Willd. sp. pl. 1. p. 1436.*

Common in the woods : Canada to Virginia. 24 July. *v. v.*

3. *C. petiolo tripartito, partialibus 3—5-foliolatis, foliolis oblongo-ovalibus subpinnatifido-lobatis, umbella pauci-radiata divaricata, fructibus elongatis teretibus lœvigatis.—Pers. syn. 1. p. 320.*

*Claytoni.*

## PENTANDRIA DIGYNIA. Chærophyllo.

*Scandix dulcis. Muhl. MSS.*

*Myrrhis Claytoni. Mich. fl. amer. 1. p. 170.*

In the Allegany and other mountains; New England to Carolina.  $\frac{1}{4}$ . June, July. v. v. The root has a very agreeable scent and sweet taste.

## 250. PASTINACA. Gen. pl. 494.

*sativa.*

1. *P. foliis simpliciter pinnatis. Willd. sp. pl. 1. p. 1466.*  
Common Parsnep is found frequently in a wild state apparently as a native, but certainly originated from European seeds.

## 251. SMYRNIUM. Gen. pl. 495.

*cordatum.*

1. *S. foliis radicalibus suborbiculato-cordatis crenatis, caulinis petiolatis trifoliatis, supremis 3-partitis, umbella radiis brevibus.—Walt. fl. car. 114.*

*Thapsia trifoliata. Willd. sp. pl. 1. p. 1465.*

In woods, frequently on the mountains: New England to Carolina.  $\frac{1}{4}$ . June, July. v. v. Flowers yellow, as are all the following.

*aureum.*

2. *S. foliis biternatis, partialium intermedio saepe subquincato; foliolis ovali-lanceolatis serrulatis, umbella brevi-radiata.—Willd. sp. pl. 1. p. 1468.*

On the mountains and hills, near banks of rivers: Pennsylvania to Carolina.  $\frac{1}{4}$ . June, July. v. v.

*atropurpure-um.*

3. *S. foliis omnibus ternatis, foliolis ovatis acutis serratis. Lam. encycl. 3. p. 667.*

*S. barbinode. Muhl. MSS.*

On dry slate hills: Virginia and Carolina.  $\frac{1}{4}$ . May—July. v. v. Flowers dark purple.

*integerri-mum.*

4. *S. glaberrimum; foliis glaucis, inferioribus 3-ternatis, superioribus 2-ternatis, foliolis ovalibus integris, umbella radiis paucis setaceo-elongatis.—Willd. sp. pl. 1. p. 1468.*

On dry hills: New England to Carolina; scarce.  $\frac{1}{4}$ . June. v. v.

*nudicaule.*

5. *S. foliis radicalibus 3-ternatis, foliolis inaequaliter paucidentatis, scapo radicali, umbella radiis elongatis, involucro involucellisque subnullis.*

On the Columbia river. *M. Lewis.*  $\frac{1}{4}$ . April, May. v. s. in *Herb. Lewis.* The natives eat the tops of this plant and boil it in their soups, the same as we use celery.

1. S. foliis 3-t  
misphaeri  
longitudi  
On the wa  
May. v

The fusif  
vegetable

2

1. S. herbacea  
lineatis,  
pl. 1. p.  
Icon. Pall

Near the  
plentiful  
Flowers

2. S. herbacea  
spinesce  
calycibu  
p. 174.  
In Caroli  
of the j

3. S. herbacea  
muticis  
lycibus  
pl. 1. p.  
Icon. Jac  
On the m  
v. s.

4. S. herba  
foliis si  
illaribu  
Sueda pro  
On the vo  
June.

254.

1. C. foliis  
situs ap  
Icon. En  
In Virgir

252. SESELI. *Gen. pl. 492.*

1. *S. foliis 3-ternatis, foliolis longo-linearibus, umbellis hemisphaericis, involucellis polyphyllis: foliolis linearibus longitudine umbellularum.* *triternatum.*

On the waters of Columbia river. *M. Lewis.* 4. April, May. *v. s. in Herb. Lewis.* Flowers deep yellow. The fusiform root of this species is one of the grateful vegetables of the Indians: they use it baked or roasted.

253. SALSOLA. *Gen. pl. 437.*

1. *S. herbacea diffusa: foliis triquetro-subulatis mucronatis lineatis, calycibus marginatis axillaribus.* — *Willd. sp. pl. 1. p. 1310.*

*Rali.*

Icon. *Pall. illustr. t. 28 & 29.*

Near the sea-coast, in gravelly dry soil: frequent; plentifully in the city of New York. ⊖ July. *v. v.* Flowers very small; seeds in a membranaceous calyx.

2. *S. herbacea, decumbens, glabra; foliis dilatato-subulatis spinescentibus, gemmis fructificantibus valde turgidis, calycibus fructiferis explanato-alatis.* *Mich. fl. amer. 1. p. 174. Walt. fl. car. 111.*

*caroliniana.*

In Carolina. *Michaux.* +. Probably only a variety of the preceding species.

3. *S. herbacea, erectiuscula, ramosissima; foliis linearibus muticis carnosis, glomerulis confertis subspicatis, calycibus fructiferis depresso-subrotundis.* — *Willd. sp. pl. 1. p. 1312. Mich. fl. amer. 1. p. 174.*

*salsa.*

Icon. *Jacq. hort. 3. t. 83.* On the mouth of the river St. Laurence. *Michaux.* ⊖ *v. s.*

4. *S. herbacea, humifusa, ramosissima; ramulis distichis, foliis succulentis linearibus acutis glabris, floribus axillaribus sessilibus, staminibus exsertis.*

*depressa.*

*Sueda prostrata. Pall. illustr. 55. t. 47.*

On the volcanic plains of the Missouri. *Nuttall.* ⊖ June. +.

254. CHENOPODIUM. *Gen. pl. 435.*

1. *C. foliis triangulari-sagittatis integerrimis, spicis compo-* *Bonus Hen-*  
*sitis aphylis axillaribus.* *Willd. sp. pl. 1. p. 1299.* *ricus.*

Icon. *Engl. bot. 717.*

In Virginia. *Gronov.* 4. June—Aug. *v. v.* I never

PENTANDRIA DIGYNIA. *Chenopodium.*

- murale.* observed this species myself in America, but it is very probable it may be found there, as Gronovius asserts.
2. *C. foliis ovatis inæqualiter dentatis acutis nitidis, racemis ramosis cymosis aphyllis.* — *Willd. sp. pl. 1. p. 1301.* Frequent on road sides, among rubbish and in gardens. ⊖ July—Sept. v. v.
  3. *C. foliis rhomboideo-ovatis erosis postice integris: superioribus oblongis integerrimis, seminibus laevibus.* *Smith fl. brit. 1. p. 273.* *Willd. sp. pl. 1. p. 1302.*  
Icon. *Engl. bot. 722.*  
β. *Willd. sp. pl. 1. p. 1303.*
  4. *C. foliis cordatis angulato-dentatis acuminatis, racemis ramosissimis subcymosis divaricatis aphyllis.* *Smith fl. brit. 1. p. 275.* *Willd. sp. pl. 1. p. 1303.*  
Icon. *Curt. lond. 4. t. 23.*  
In cultivated grounds, near roads and fences: common.  
⊖ July, Aug. v. v.
  5. *C. foliis oblongis sinuatis, racemis nudis multifidis.* *Willd. sp. pl. 1. p. 1304.*  
On the banks of the Susquehanna, Pennsylvania, and about Onondago, New York. ⊖ July, Aug. v. v.  
Smell strong, but not disagreeable.
  6. *C. foliis lanceolatis dentatis, racemis foliatis simplicibus.* *Willd. sp. pl. 1. p. 1304.*  
Icon. *Moris. hist. 2. s. 5. t. 35. f. 8.*  
In sandy fields: Georgia. ⊖ July. v. v. Very sweet scented.
  7. *C. foliis ovato-oblongis dentatis, spicis longis interruptis aphyllis.* — *Willd. sp. pl. 1. p. 1304.*  
Icon. *Dill. elth. t. 66. f. 76.*  
On road sides: Pennsylvania and Virginia; plentifully in the streets of Philadelphia. ⊖ June—Aug. v. v.  
Smells very strong, but not disagreeable to some persons.
  8. *C. foliis subulatis carnosis semiteretibus, floribus glomeratis axillaribus.* *Willd. sp. pl. 1. p. 1307.*  
Icon. *Moris. hist. 2. s. 5. t. 33. f. 3.*  
On the sea-coast of Pennsylvania and Virginia. ⊖ July, Aug. v. v.

1. *A. caule fr rhombei*  
On the sea  
Hortis.  
Gronovi  
America
2. *A. caule e concolor*  
gerrimis  
Frequent i
3. *A. caule e dentatis nerviis*  
Icon. Mo  
Near the  
Virgini
4. *A. caule funde dentati*  
p. 963.  
Frequent

1. *C. foliis Common  
indigen  
duced*

1. *Humulu*  
In deep  
in the  
genou

1. *U. rami acum  
sissim*  
Icon. A

255. ATRIPLEX. *Gen. pl.* 1577.

1. *A. caule fruticoso, foliis alternis oppositis oblongo-subrhombeis integerrimis.* *Willd. sp. pl.* 4. p. 957.

On the sea-coast of Virginia. *Gronovius.* ½. v. v. in *Hortis.* I take this species up only on the authority of Gronovius, having never seen it myself as a native of America.

2. *A. caule erecto herbaceo, foliis triangularibus dentatis concoloribus, calycibus fructus ovatis reticulatis integerrimis.* *Willd. sp. pl.* 4. p. 961.

Frequent in fields and about gardens. ☽. July. v. v.

3. *A. caule erecto herbaceo, foliis triangularibus profunde dentatis subtus albidis, calycibus fructus rhombeis trinerviis denticulatis.* *Willd. sp. pl.* 4. p. 963.

*Icon. Moris. hist.* 2. s. 5. t. 32. f. 17.

Near the sea-coast, on banks of rivers: New York to Virginia. ☽. June—Aug. v. v. Leaves gray.

4. *A. caule herbaceo erecto, foliis triangulari-hastatis profunde dentatis concoloribus, calycibus fructus sinuato-dentatis, dente intermedio elongato.* *Willd. sp. pl.* 1. p. 963.

Frequent about gardens and in fields. ☽. July. v. v.

*Halimus.*

*hortensis.*

*laciniata.*

*hastata.*

*sativa.*

256. CANNABIS. *Gen. pl.* 1522.

1. *C. foliis digitatis.* *Willd. sp. pl.* 4. p. 768.

Common hemp is frequently found in fields, apparently indigenous, though it is one of those plants introduced from Europe. ☽.

257. HUMULUS. *Gen. pl.* 1523.

1. *Humulus.* *Willd. sp. pl.* 4. p. 769.

In deep shady valleys on the mountains and on river sides in the lower countries, frequent. It is really indigenous. ¼.

*Lupulus.*

258. ULMUS. *Gen. pl.* 443.

1. *U. ramis lăvibus basi inaequalibus: serraturis uncinato-acuminatis, floribus pedicellatis, fructibus villo densissimo fimbriatis.* — *Willd. sp. pl.* 1. p. 1325.

*Icon. Mich. arb.* t....

*americana.*

*pendula.*

- β. *U. foliis oblongis glabriusculis, ramis pendulis.* *Ait. kew. 1. p. 319.*

Common in all low lands and woods: New England to Carolina.  $\frac{1}{2}$ . April. *v. v.* Known by the name of *White Elm.*

*nemoralis.*

2. *U. foliis oblongis glabriusculis æqualiter serratis basi subæqualibus, floribus sessilibus.* *Ait. kew. 1. p. 319.*  
*U. campestris. Walt. fl. car. 111?*

On the banks of rivers: New England to Virginia, scarce.  $\frac{1}{2}$ . April, May. *v. v.* *River Elm.*

*fulva.*

3. *U. ramis scabris albidis, foliis ovato-oblongis acuminatis basi subæqualibus, inæqualiter serratis utrinque pubescentibus scaberrimis, gemmis lana densa fulva tomentosis, floribus sessilibus.* —*Mich. fl. amer. 1. p. 172.*

Icon. *Mich. arb. t. . . .*

On mountains from Canada to Pennsylvania.  $\frac{1}{2}$ . May. *v. v.* *Red Elm.* *Slippery Elm.* The viscous inside bark is used by the natives as a healing salve.

*alata.*

4. *U. ramis utrinque alato-suberosis, foliis oblongo-ovalibus sensim acutis basi subæqualibus, fructu pubescente cilioso.* —*Mich. fl. amer. 1. p. 173.*

*U. pumila. Walt. fl. car. 111.*

Icon. *Mich. arb. t. . . .*

In sandy low woods: Virginia and Carolina.  $\frac{1}{2}$ . April. *v. v.* *Whahoo.*

### 259. CELTIS. *Gen. pl. 1591.*

*occidentalis.*

1. *C. foliis ovatis acuminatis æqualiter serratis basi inæqualibus supra scabris, subtus hirtis, fructu solitario.* —*Willd. sp. pl. 4. p. 994.*

Icon. *Duham. arb. 1. t. 53.* *Mich. arb. t. . . .*

In woods and near rivers: Canada to Carolina.  $\frac{1}{2}$ . May. *v. v.* *Nettle-tree.* Berries dark purple.

*crassifolia.*

2. *C. foliis ovatis acuminatis inæqualiter serratis basi inæqualiter cordatis subcordatis utrinque scabris, pedunculis subbifloris.* —*Willd. sp. pl. 4. p. 995.*

*C. cordifolia. Duham. 2. t. 9.*

In woods and near rivers: Virginia, Kentucky, and Tennessee.  $\frac{1}{2}$ . May. *v. v.* Known by the name of *Hagberry* or *Hoop-ash.*

*pumila.*

3. *C. foliis ovatis acuminatis æqualiter serratis basi inæqua-*

libus ut  
bescentil  
On the ba  
May.  
black.

26

1. *V. glabru  
nato-sei  
bus rotu  
Icon. Plu  
Common  
lina. E  
ers, as*

2. *V. glabru  
læyibus  
latis.—  
On the b  
h. Ma  
not so*

3. *V. glabru  
ratis, p  
Willd.  
Frequent  
Engla  
clined  
black.*

4. *V. glabe  
integre  
culatis  
Icon. M  
In swan  
gia.  
South  
branc  
their*

5. *V. glabi  
rimis  
rotan*

5. *Ait.*  
gland to name of  
basi sub-  
319.  
Virginia,  
uminatis  
que pu-  
fulva to-  
zmer. 1.  
  
h. May.  
us inside  
-ovalibus  
abescente  
  
h. April.  
  
ii inaequa-  
politario.—  
  
h. May.  
basi inae-  
is, pedun-  
ucky, and  
the name  
isi inaequa-

libus utrinque glabriusculis : junioribus tantum pubescentibus, pedunculis subtrifloris, fructu solitario. On the banks of rivers : Maryland and Virginia. h. May. v. v. A small straggling bush ; berries ovate, black.

## III. TRIGYNIA.

## 260. VIBURNUM. Gen. pl. 503.

1. V. glabrum ; ramis patentissimis, foliis subrotundis crenato-serratis, petiolis laevibus, cymis sessilibus, fructibus rotundatis.—*Willd. sp. pl. 1. p. 1847.*  
Icon. *Pluk. alm. t. 46. f. 2. Duham. 2. t. 38.*  
Common in hedges and fields : New England to Carolina. h. May, June. v. v. Berries dark blue ; flowers, as all the following species, white.
2. V. glabrum ; foliis ovatis subacutis subserratis, petiolis laevibus, fructibus ovato-oblongis cymis subpedunculatis.—*Lam. encycl. 8. p. 653.*  
On the banks of rivers : Pennsylvania, New Jersey, &c.  
h. May, June. v. v. Resembles the former, but is not so straggling in its growth ; berries black.
3. V. glabrum ; foliis late-ovatis acuminatis uncinato-serratis, petiolis marginatis undulatis, cymis sessilibus.—*Willd. sp. pl. 1. p. 1491.*  
Frequent among hedges and on borders of woods : New England to Carolina. h. July. v. v. Is more inclined to grow to a tree than any of the rest ; berries black.
4. V. glaberrimum ; foliis ovalibus margine revoluto subintegerrimis, petiolis laevibus, cymis ebracteatis pedunculatis.—*Willd. sp. pl. 1. p. 1487.*  
Icon. *Mill. ic. 274.*  
In swamps, particularly on sandy soil : Canada to Georgia. h. May, June. v. v. Berries black ; in the Southern States it becomes an evergreen ; the young branches are ferruginous and sometimes the leaves on their under surface.
5. V. glabrum ; foliis obovatis crenato-dentatis seu integerimis obtusis, umbellis sessilibus, fructibus ovato-subrotundis.—*Lam. encycl. 8. p. 658. Walt. fl. car. 116.*

*prunifolium.**Lentago.**nudum.**obovatum.*

- V. *cassinoides*. *Mich. fl. amer.* 1. p. 179.  
*punicifolium*.  $\beta$ . V. foliis obovatis integris seu apice laxe crenatis obtusis.  
*Hort. par.* 115.  
 In shady woods : Carolina and Georgia.  $\frac{1}{2}$ . May, June.  
 v. s.
- cassinoides*. 6. V. glabrum ; foliis ovato-lanceolatis utrinque acutis cre-  
 natis margine subrevolutis, petiolis carinatis eglandula-  
 tis.—*Willd. sp. pl.* 1. p. 1491.  
 In swamps : New York to Carolina,  $\frac{1}{2}$ . June, July.  
 v. v. Berries blueish-black.
- lævigatum*. 7. V. foliis lanceolatis lævibus remote-serratis basi inte-  
 gerrimis, ramulis ancipitibus.—*Willd. sp. pl.* 1.  
 p. 1492.  
 Icon. *Mill. ic. t.* 83. f. 1.  
 Near the sea-coast : Virginia and Carolina.  $\frac{1}{2}$ . June,  
 July. v. v. Berries black.
- nitidum*. 8. V. glaberrimum ; foliis linear-lanceolatis supra nitidis  
 obsolete serratis integris ; ramis tetragonis.—*Willd.*  
*sp. pl.* 1. p. 1492.  
 In sandy barren woods : Carolina and Georgia.  $\frac{1}{2}$ . v. v.  
*s. fl.* A low shrub with small leaves.
- dentatum*. 9. V. glabriuscum ; foliis suborbiculatis ovatisque acutis  
 dentato-serratis plicato-sulcatis utrinque glabris, cynis  
 pedunculatis, fructibus subglobosis.—*Willd. sp. pl.* 1.  
 p. 1488.  
 Icon. *Jacq. hort. t.* 36.  
 In mountain woods ; frequent : New York to Carolina.  
 $\frac{1}{2}$ . June, July. v. v. Berries dark blue ; known by  
 the name of *Arrow-wood*.
- pubescens*. 10. V. pubescens ; foliis ovalibus acuminatis dentato-serratis  
 plicato-sulcatis subtus villos-tomentosis, cymis pe-  
 dunculatis, fructibus oblongis.  
 V. *dentatum pubescens*. *Ait. kew.* 1. p. 168.  
 V. *dentatum tomentosum*. *Mich. fl. amer.* 1. p. 179.  
 In the lower parts of Virginia and Carolina.  $\frac{1}{2}$ . June,  
 v. v. The whole of the shrub smaller than the pre-  
 ceding one.
- lantanoides*. 11. V. petiolis nervisque pulverulento-tomentosis, foliis am-  
 plis suborbiculato-cordatis abrupte acuminatis inæqua-  
 liter serratis, serraturis muticis, cymis arcte sessilibus,  
 fructibus ovatis.—*Mich. fl. amer.* 1. p. 179.  
 V. *Lantana grandifolium*. *Ait. kew.* 1. p. 168.  
 In shady woods on high mountains : Canada to Virginia ;  
 principally in the forests called *Beech-woods*.  $\frac{1}{2}$ .

PEN

June, J  
known12. V. *ramuli*  
*dato-ova*  
*pubesce*  
*pl.* 1. p.  
 Icon. *Ven*  
 In rocky  
 rolinia.13. V. *foliis*  
*subtus*  
*radiatis*  
*p.* 180.  
 V. *alnitol*  
 In hedges  
 nassee  
 Berries14. V. *foliis*  
*ricatis*  
*glandu*  
 V. *Opul*  
 V. *Opult*  
 V. *trilob*  
 In swam-  
 tains c  
 Berrie  
 Cranb  
 are a v15. V. *foliis*  
*bis bi*  
*natis,*  
 V. *Opul*  
 On the l  
 v. v.  
 ceding  
 when1. S. *rami*  
*oblon*  
*medi*  
*cymi*  
 Icon. S

- June, July. v. v. Berries red, but when ripe black ; known by the name of *Hobble-bush*.
12. V. ramulis petiolisque pilosis eglandulosis, foliis subcordato-ovatis trilobis acuminatis argute-serratis subitus pubescentibus, cymis longe-pedunculatis.—*Willd. sp. pl. 1. p. 1489.*
- Icon. *Vent. hort. cels. t. 72.*  
In rocky mountainous situations : New England to Carolina. ♀. May, June. v. v. Berries black.
13. V. foliis suborbiculato-cordatis plicato-sulcatis dentatis subitus pubescentibus, petiolis subglandulosis, cymis radiatis, fructibus oblongo-ovatis. *Mich. fl. amer. 1. p. 180.*
- V. alnifolium. *Marsh. arb. 162.*  
In hedges : Kentucky near Danville. *Michaux.* Tennessee and Upper Carolina. ♀. June, July. v. s.  
Berries red ; resembles the following species.
14. V. foliis trilobatis postice acutis 3-nervibus : lobis divaricatis promisse acuminatis rariter dentatis, petiolis glandulosis, cymis radiatis.
- V. *Opulus americanum*. *Ait. kew. 1. p. 168.*
- V. *Opulus Pimina*. *Mich. fl. amer. 1. p. 180.*
- V. *trilobum*. *Marsh. arb. 162.*  
In swamps and shady woods : Canada ; on the mountains of New York and New Jersey. ♀. July. v. v.  
Berries red, of an agreeable acid ; resembling that of Cranberries, *Vaccinium macrocarpon*, for which they are a very good substitute.
15. V. foliis trilobatis, postice obtusiusculis 3-nervibus : lobis brevissimis denticulato-serratis, serraturis acuminatis, petiolis glandulosis, cymis radiatis.
- V. *Opulus edule*. *Mich. fl. amer. 1. p. 180.*  
On the banks of rivers : Canada to New York. ♀. July.  
v. v. A smaller and more upright shrub than the preceding species ; berries the same colour and size, but, when completely ripe, more agreeable to eat.

*molle.**Oxycoccus.**edule.**canadensis.*261. SAMBUCUS. *Gen. pl. 505.*

1. S. ramulis petiolisque glaberrimis, foliolis sub-4-jugis oblongo-ovalibus nitidis glabris acuminatis : nervo medio subpubescente, basi interdum appendiculatis, cymis laxis 5-fidis.—*Willd. sp. pl. 1. p. 1494.*
- Icon. *Schmidt arb. 142.*

## PENTANDRIA TRIGYNYIA. Sambucus.

- subescens.* In swamps and near hedges : Canada to Carolina.  $\frac{1}{2}$ . June, July. *v. v.* Berries dark purple or black.
2. *S. cortice verrucoso, foliolis 2-jugis ovali-lanceolatis subtus pubescentibus, cymis confertis racemosis.* — *Mich. fl. amer. 1. p. 181.* On the highest mountains from Canada to Carolina.  $\frac{1}{2}$ . July. *v. v.* Berries red ; resembles *S. racemosa* very much.
- typhinum.* 262. RHUS. *Gen. pl. 502.*
1. *R. subarborescens ; ramis petiolisque villosissimis, foliis pinnatis multijugis, foliolis lanceolato-oblongis argute serratis subtus subtomentosis.* — *Willd. sp. pl. 1. p. 1478.* *Icon. Duham. 2. t. 47.* In rocky dry situations : Canada to Virginia.  $\frac{1}{2}$ . July, Aug. *v. v.* The purple velvety berries appear in large close clusters.
2. *R. undique glabrum ; foliis pinnatis multijugis, foliolis lanceolato-oblongis serratis subtus albanticibus, fructibus holosericeis.* — *Willd. sp. pl. 1. p. 1478.*
- $\beta$ . *R. elegans. Ait. kew. 1. p. 162. Catesb. car. appen. t. 4.* *Icon. Dill. elth. t. 243. f. 314. Catesb. car. 3. t. 4.* Common in old fields and along fences : New England to Carolina.  $\frac{1}{2}$ . July, Aug. *v. v.* Berries red ; flowers greenish-red ; in var.  $\beta$ . scarlet.
3. *R. glabriusculum ; foliis pinnatis multijugis, foliolis lanceolato-oblongis serratis subtus subtomentosis, racemis erectis herbaceis.* — *Lam. encycl. 7. p. 504.* *R. canadense. Mill. dict. no. 5.* On the edges of woods, in dry sunny situations : Pennsylvania and Virginia.  $\frac{1}{2}$ . July, Aug. *v. v.* Flowers yellowish-green ; probably not much more than a variety of the foregoing species.
- pumilum.* 4. *R. humile ; ramis petiolisque pubescentibus, foliis pinnatis multijugis, foliolis ovalibus vix acuminatis aut muticis inciso-dentatis subtus tomentosis, fructibus holosericeis.* — *Mich. fl. amer. 1. p. 182.* In Upper Carolina.  $\frac{1}{2}$ . July. *v. s. in Herb. Lyon.* ; *v. v. in Hortis.* Not above a foot high : it is the most poisonous of the genus, according to information from Mr. J. Lyon, who, by collecting the seed of this species, got poisoned all over his body, and was lame for a considerable time.

5. *R. arborescens foliolis ovali-lanceolatis subtus pubescentibus, cymis confertis racemosis.* — *Mich. fl. amer. 1. p. 181. Icon. Dill. elth. t. 243. f. 314.* In low damp situations. Berries red.

6. *R. ramosus foliolis lanceolatis subtus pubescentibus.* — *Willd. sp. pl. 1. p. 1478. Icon. Hor. 1. t. 1. f. 1.* In dry fields. Jersey. The leaves of Indians.

7. *R. radicalis integris foliis subtus pubescentibus.* — *Mich. fl. amer. 1. p. 182. Icon. Dill. elth. t. 243. f. 314.* Common to Georgia known as no only loc.

$\alpha$ . *R. radicans dentatis.* — *R. radicans dentatis.*

$\beta$ . *R. erecta floratio.*

$\gamma$ . *R. foliolis boideis.* Common to Georgia known as no only loc.

8. *R. fruticosa foliis subtentib.* — *R. canadensis. Mill. dict. no. 5. Icon. Theophrast. In rocky places, yellow pear-shaped fruit. foliata is not Berries.*

*Vernix.*

5. *R. arborescens*, glaberrimum ; foliis pinnatis multijugis, foliolis ovalibus abrupte acuminatis integris, panicula laxa, floribus dioicis, fructibus glabris.—*Willd. sp. pl. 1. p. 1479.*

Icon. *Dill. elth. t. 292. f. 377. Pluk. alm. t. 145. f. 1.*  
In low copses : Canada to Carolina. ♂. July. v. v.  
Berries white.

6. *R. ratnís punctatis*, foliis pinnatis multijugis (5—6), foliolis ovali-lanceolatis integerrimis supra lucidis, panicula foliosa, racemis sessilibus, floribus dioicis.—*Willd. sp. pl. 1. p. 1480.*

Icon. *Hort. schœnbr. 341. Pluk. alm. t. 56. f. 1.*  
In dry fields and woods, particularly sandy soil : New Jersey to Carolina. ♂. July. v. v. Berries red.  
The leaves of this species are used as tobacco by the Indians of the Missouri and Mississippi.

7. *R. radicans* ; foliis ternatis, foliolis ventricoso-ovalibus integris aut sinuate-crenatis, racemis rameis et axillaribus sessilibus dioicis.—*Willd. sp. pl. 1. p. 1481.*  
*Mich. fl. amer. 1. p. 183.*

8. *R. radicante-scandens* ; foliolis amplis integris seu rariter dentatis. *Mich. l. c.*

*R. radicans. Willd. sp. pl. 1. p. 1481.*

- β. *R. erectum*, humile ; foliolis varie sinuato-lobatis, sub quercifolium. floratione tomentosis. *Mich. l. c.*

- γ. *R. foliolis oblongo-ovalibus longe acuminatis subrhomboidibus*, fructu multo minore. *Mich. l. c.*

Common in all woods, fields, and along fences : Canada to Georgia. ♂. June, July. v. v. Berries white ; known by the name *Poison-oak* or *Poison-vine*. There is no doubt but *R. radicans* and *Toxicodendron* are only local varieties.

8. *R. fruticosum*, amentaceum, nudiflorum ; foliis ternatis, foliolis rhombo-ovalibus dentatis subtus pubescens, floribus dioicis.—*Willd. sp. pl. 1. p. 1482.*

*R. canadense. Marsh. arb. 129.*

- Icon. *Turpin in annal. du mus. hist. nat. 5. p. 445. t. 30.*  
In rocky situations, about springs : Pennsylvania, Carolina, and Kentucky. ♂. May, June. v. v. Flowers yellow, in catkins, which give it quite a different appearance from the habitus of the genus. *Myrica trifoliata* Hort. is the same with *R. suaveolens* Ait. and is nothing more than the male plant of this species. Berries brown.

*Toxicoden-dron.**vulgare.*

263. STAPHYLEA. *Gen. pl. 507.**trifolia.*

1. S. foliis trifoliatis : racemis pendulis, petalis inferne ciliatis, fructu ovato.—*Willd. sp. pl. 1. p. 1498.*

Icon. *Schmidt arb. 81.*

On the side of dry hills, in rocky situations : New York to Upper Carolina.  $\frac{1}{2}$ . May, June. *v.v.* Flowers white.

264. TURNERA. *Gen. pl. 514.**cistoides.*

1. T. hirsuta ; foliis linear-lanceolatis obtuse dentato-serratis, pedunculis axillaribus folio brevioribus.—*Willd. sp. pl. 1. p. 1505.*

Icon. *Plum. ic. t. 150. f. 1.* *Sloan. jam. 1. t. 127. f. 7.*  
In dry sandy woods, near the town of Savannah, Georgia. *Enslen.*  $\odot$ . July. *v.s. in Herb. Enslen.* Flowers yellow, resembling those of *Cistus carolinianus*.

265. CROTONOPSIS. *Mich. fl. amer. 2. p. 185.**argentea.*

1. C. caule erecto dichotome ramosissima, foliis supra stellato-pilosis, subtus argenteo-lepidotis.

$\alpha$ . C. foliis linear-lanceolatis. *Willd. sp. pl. 4. p. 380.* *Mich. fl. amer. 2. p. 186.*

$\beta$ . C. foliis ellipticis utrinque obtusis. *Willd. l.c.*

In sandy soil near the sea-coast, Illinois country and on the Missouri.  $\odot$ . July. *v.v.* Flowers very minute.

266. KOCHIA. *Roth in Schrad. journ. 1800. p. 307.**dentata.*

1. K. foliis lanceolatis sinuato-dentatis, caule erecto ramosissimo.—*Willd. hort. berol. 1. t. 28.*

*Salsola radiata. Desf. in annal. mus. nat. p. 28. t. 34.*

*Salsola platyphylla. Mich. fl. amer. 1. p. 174.*

On the banks of the Mississippi, in the Illinois country.

$\odot$ . June. *v.v.* This plant has the aspect of a *Chenopodium* in every part.

267. AMARANTHUS. *Gen. pl. 1431.**albus.*

1. A. glomerulis axillaribus triandris, foliis obovatis retusis, caule tetragono simplici. *Willd. sp. pl. 4. p. 382.*

Icon. *Willd. amaranth. t. 1. f. 2.*

A common weed in the gardens.  $\odot$ . May—Sept. *v.v.*

2. A. glomerulatis, obovatis  
*sp. pl. 4.*

Icon. *Wil...*  
In cultivate

3. A. glomerulatis reticulatis, *t. 1. f. 1.*

In similar

4. A. glomerulatis, *foliis ov.*  
*p. 387.*

Frequently  
June—

5. A. glomerulatis, *tridulatis.*

Frequentl  
July, A

6. A. racemulatis, *foliis ov.*  
*f. 17.*

Common  
 $\odot$ . Ju

7. A. racemulatis, *bus p.*  
*amara*

Among  
to Vir

8. A. racemulatis, *patent amara*

In culti  
*v.v.*

9. A. racemulatis, *ra amara*

In culti  
Virgi

10. A. racemulatis, *oblonga*

Comm  
Leav

2. *A. glomerulis axillaribus triandris, floribus trifidis, foliis obovatis emarginatis, caule teretiusculo ramoso.* Willd. *græcizans.*  
*sp. pl. 4. p. 382.* *Icon. Willd. amaranth. t. 4. f. 7.*  
 In cultivated grounds : Virginia. ⊙. Aug. v. v.
3. *A. glomerulis triandris subspicatis rotundatis, foliis ellipticis retusis, caule erecto.* Willd. *amaranth. p. 20.*  
*t. 1. f. 1.* *lividus.*  
 In similar situations : Virginia. ⊙. June—Aug. v. v.
4. *A. glomerulis triandris subspicatis, floribus triphyllis, foliis ovatis retusis, caule diffuso.* Willd. *sp. pl. 4. p. 387.* *Blitum.*  
 Frequently a very troublesome weed in gardens. ⊙.  
 June—Sept. v. v.
5. *A. glomerulis axillaribus geminis triandris, floribus masculis triphyllis, foliis ellipticis emarginatis margine undulatis.* Willd. *amaranth. 18. t. 8. f. 16.* *viridis.*  
 Frequently found on road sides and among rubbish. ⊙.  
 July, Aug. v. v.
6. *A. racemis pentandris decompositis congestis erectis, foliis ovato-lanceolatis.* Willd. *amaranth. p. 26. t. 9.*  
*f. 17.* *hybridus.*  
 Common in cultivated grounds ; New York to Carolina.  
 ⊙. June—Sept. v. v.
7. *A. racemis pentandris supradecompositis, ramis patentibus pubescentibus, foliis ovato-lanceolatis.* Willd. *amaranth. 32. t. 2. f. 2.* *paniculatus.*  
 Among rubbish and in cultivated grounds : Pennsylvania to Virginia. ⊙. Aug. v. v.
8. *A. racemis pentandris supradecompositis erectis, ramis patentibus glabris, foliis oblongis acutis.* Willd. *amaranth. 31. t. 2. f. 3.* *sanguineus.*  
 In cultivated fields : Virginia and Carolina. ⊙. July.  
 v. v.
9. *A. racemis pentandris supradecompositis confertis erectis, ramis pubescentibus, foliis ovatis undulatis.* Willd. *amaranth. 33. t. 11. f. 21.* *retroflexus.*  
 In cultivated grounds and on road sides : Pennsylvania to Virginia. ⊙. June—Aug. v. v.
10. *A. racemis pentandris compositis confertis erectis, foliis oblongo-lanceolatis mucronatis.* Willd. *amaranth. 80.* *hypochondriacus.*  
 Common in fields of Virginia. ⊙. July, Aug. v. v.  
 Leaves red.

*spinosa*. 14. A. racemis pentandris terminalibus compositis, axillis spinosis. *Willd. amaranth.* 38. t. 4. f. 8.

On road sides and among rubbish; frequent. Ⓛ. June—Sept. v. v.

268. ACNIDA. *Gen. pl.* 1521. *Mich. fl. amer.* 2. p. 233.

*cannabina*. 1. A. foliis lanceolatis, capsulis laeviusculis acutangulis. *Willd. sp. pl.* 4. p. 767.

On the banks of rivers and in marshes near the sea-coast: New England to Carolina. Ⓛ. July, Aug. v. v.

*ruscocarpa*. 2. A. foliis ovali-lanceolatis, capsulis obtusangulis rugosis. *Willd. sp. pl.* 4. p. 767.

Icon. *Mich. fl. amer.* 2. t. 50.

In close swamps, on the banks of rivers among bushes: Virginia. Ⓛ. July, Aug. v. s. in *Herb. Lyon.* Very tall and robust.

#### IV. TETRAGYNIA.

269. PARNASSIA. *Gen. pl.* 523.

*palustris*. 1. P. foliis radicalibus cordatis, nectariis multisetis.—*Willd. sp. pl.* 1. p. 1516.

Icon. *Fl. dan.* 584. *Engl. bot.* 82.

In bog meadows: Pennsylvania and Virginia. ¼. July, Aug. v. v. Flowers white, with netted veins of green or pale purple.

*caroliniana*. 2. P. foliis radicalibus suborbiculatis, nectariis trisetis. *Mich. fl. amer.* 1. p. 184.

Icon. *Bot. mag.* 1459.

In swamps and mosses: New York to Virginia. ¼. July, Aug. v. v. Flower the same as No. 1.

*asarifolia*. 3. P. foliis radicalibus reniformibus, petalis unguiculatis, nectariis trifidis. *Vent. malmais.* 39. t. 39.

On high mountains in Virginia and Carolina. ¼. July, Aug. v. v. Larger in leaves and flowers than the preceding species.

1. A. subaca  
foliolis  
umbell  
Icon. *Plu*  
In shady  
June.  
such us

2. A. divari  
3—5-f  
illaribu  
perne a  
Icon. *Co*  
In rocky  
June—  
nard, a

3. A. humili  
pidis, 1  
serratis  
duncul  
Icon. *Ba*  
In stony  
mounta  
v. v.

4. A. caule  
pinnati  
umbell

5. A. caule  
Icon. *Sci*  
In fertile  
likewis  
Know  
used in  
medici  
was fo  
not see

## V. PENTAGYNYIA.

## 270. ARAlia. Gen. pl. 525.

1. A. subacaulis, unifoliata ; folio triternato vel triquinato : foliolis oblongo-ovalibus, scapo nudo folio breviore, umbellulis paucis.—Willd. sp. pl. 1. p. 1521.  
Icon. Pluk. alm. t. 238. f. 5.  
In shady rocky woods : Canada to Carolina. 4. May, June. v. v. Commonly called *Sarsaparilla*, and as such used in the shops.
2. A. divaricato-ramosa ; petiolis 3-partitis, partitionibus 3—5-foliolatis, foliolis plerisque cordatis, ramulis axillaribus foliosis, umbellulis pluribus subpaniculatis superne aphyllis.—Willd. sp. pl. 1. p. 1521.  
Icon. Cold. canad. t. 75. Moris. hist. 1. s. 1. t. 2. f. 9.  
In rocky and shady situations : Canada to Virginia. 4. June—Sept. v. v. Known by the name of *Spike-nard*, and highly esteemed as a medicine.
3. A. humilis, suffruticosa ; caulis petiolisque setis hispidis, foliis duplicato-pinnatis, foliolis ovatis inciso-serratis inermibus glabris, umbellulis longissime penduculatis.—Mich. fl. amer. 1. p. 185.  
Icon. Bot. mag. 1085. Vent. hort. cels. 41.  
In stony woods : Canada, New England, &c. On high mountains : Pennsylvania, Virginia. ½. June, July. v. v. Called *Wild Elder*.
4. A. caule arborescente foliisque spinosis, foliis duplicato-pinnatis, foliolis leviter serratis, panicula ramosissima, umbellis numerosis.—Willd. sp. pl. 1. p. 1520.
5. A. caule spinoso glauco, foliis inermibus.  
Icon. Schmidt arb. 102 & 103. Comm. hort. 1. t. 47.  
In fertile low woods of Lower Virginia and Carolina, likewise in the Illinois country. ½. Aug. Sept. v. v. Known by the name of *Angelica-tree*. The berries used in an infusion of wine or spirits are a remarkable medicine for relieving rheumatic pains. The var. β. was found near Charlestown, South Carolina ; I have not seen its flowers.

## 271. ZANTHOXYLUM. Gen. pl. 1512.

1. Z. aculeatum ; foliis pinnatis, foliolis ovali-lanceolatis subintegerrimis sessilibus basi æqualibus, petiolo com-

muni inermi, umbellis axillaribus.—*Willd. sp. pl. 4. p. 754.*

*Z. Clava Herculis*  $\beta$ . *Linn. sp. pl. 1455.*

*Z. ramiflorum*. *Mich. fl. amer. 2. p. 235.*

Icon. *Duham. arb. 1. t. 97.*

In shady woods, near rivers: Canada to Virginia and Kentucky.  $\frac{1}{2}$ . April, May. *v. v.* A tincture of the bark and capsules is recommended in rheumatism and tooth-ach, from which its name *Tooth-ach-tree*.

*tricarpon.*

2. *Z. aculeatum*; foliis pinnatis, foliolis petiolatis oblongo-ovalibus acuminatis serrulatis basi obliquis, petiolo communi aculeato, capsulis ternis sessilibus.—*Mich. fl. amer. 2. p. 235.*

In woods of Carolina and Florida. *Michaux.*  $\frac{1}{2}$ . July. *v. v. in Hortis.*

### 272. LINUM. Gen. pl. 528.

*usitatissi-  
mum.*

1. *L. foliolis calycinis ovatis acutis trinervibus, petalis cre-  
natis, foliis lanceolatis alternis, caule subsolitario.*—  
*Willd. sp. pl. 1. p. 1533.*

Icon. *Engl. bot. 1357.*

Common flax is frequently found in old fields, apparently indigenous.  $\odot$ . June, July. *v. v.* Flowers blue.

*Lewisii.*

2. *L. foliolis calycinis ovatis acuminatis, petalis cuneatis apice rotundatis, foliis sparsis lanceolato-linearibus mucronatis, caulis altis numerosis.*

In the valleys of the Rocky-mountains and on the banks of the Missouri. *M. Lewis.*  $\frac{1}{2}$ . July. *v. v.* Flowers large, blue; a very good perennial, and it probably might become an useful plant if cultivated.

*virginicum.*

3. *L. foliolis calycinis acutis panicula terminali, floribus remote alternis, foliis linear-lanceolatis sparsis: radi-  
calibus ovatis.*—*Willd. sp. pl. 1. p. 1538.*

On dry sunny hills and fields: New York to Virginia.  $\odot$ . July, Aug. *v. y.* Flowers small, yellow.

*rigidum.*

4. *L. foliolis calycinis ovatis acuminatis trinervibus ciliatis, petalis oblongis angustissimis, foliis rigide-erectis line-  
aribus brevibus.*

On the Missouri. *Nuttall.* *v. s.* Flowers sulphur-yellow.

### 273. DROsera. Gen. pl. 531.

*rotundifolia.*

1. *D. scapis radicatis simplicibus, foliis suborbiculatis basi attenuatis, petiolis elongatis pilosis.*—*Willd. sp. pl. 1. p. 1543.* *Mich. fl. amer. 1. p. 186.*

Icon. *Fl.*  
In swamp  
species,  
July, A  
and curi

2. *D. scapis  
petiolis  
Mich. fl.*  
Icon. *Eng*  
In similar

3. *D. pusilla  
cuneatis*  
In sandy s  
in Herb  
known;

4. *D. scapis  
simis.*

In the pi  
 $\frac{1}{2}$ . Jun

*Radix:* E  
lanuginos  
subulata  
angustis  
bris visc  
pedales  
bulbifer  
simplex  
purpure

This eleg  
place at  
spersed  
ral bota  
with th

1. *S. erecta,*  
*p. 1567*  
Icon. *Am*

On the *l*  
*v. s. in*

2. *S. procu*  
*p. 1567*  
Icon. *Fl.*

On the M  
nada ar

sp. pl. 4.

rginia and  
in cture of  
eumatism  
ach-tree.  
oblongo-  
is, petiolo-  
as.—Mich.

July.

petalis cre-  
solitario.—

ields, appa-  
owers blue.  
lis cuneatis  
o-linearibus

n the banks  
v. v. Flow-  
it probably  
1.

nali, floribus  
parsis : radi-

to Virginia.  
allow.

vibus ciliatis,  
erectis line-

vers sulphur-

ribiculatis basi-  
illd. sp. pl. 1.

Icon. *Fl. dan.* 1028.

In swamps filled with sphagnum, as all the following species, generally on sandy or gravelly ground. ♀. July, Aug. v. v. Flowers white; a very delicate and curious plant, as are all the rest of this genus.

2. *D. scapis radicatis simplicibus, foliis spathulato-ovatis, petiolis elongatis nudis.* — *Willd. sp. pl. 1. p. 1544.*  
*Mich. fl. amer. 1. p. 186.*

Icon. *Engl. bot.* 868. *Moris. hist.* 1. s. 15. t. 4. f. 2.

In similar situations: Canada to Carolina. ♀. July. v. v.

3. *D. pusilla ; scapis radicatis simplicibus, foliis brevibus cuneatis vix petiolatis, petalis ovalibus.*

In sandy swamps of Georgia. *Enslen.* ♂. June. v. s. in *Herb. Enslen.* The smallest of all the species known; flowers rose-coloured.

4. *D. scapis radicatis subramosis, foliis filiformibus longis-*  
*simis.*

In the pine-barrens of New Jersey near Tuckerton. ♀. June. v. v. Flowers large, purple.

*Radix:* *Hybernaclum squamosum, extus ferrugineo-*  
*lanuginosum. Folia primordia glabra, dilatato-*  
*subulata; secundaria radicalia spiraliter erumpentia,*  
*angustissime linearia subtus glabra, supra pilis ru-*  
*bris viscidis tecta, 9—12 pollices longa. Scapi erecti,*  
*pedales et ultra, teretes, glabri, interdum vivipari,*  
*bulbiferi. Racemus spicatus, unilateralis, recurvatus,*  
*simplex seu ramosus. Flores pedicellati, erecti, magni,*  
*purpurei. Styli sex, interdum novem.*

This elegant species I discovered about 1805, in the place above mentioned: this extensive swamp, interspersed with barren sand hills, might be called a natural botanic garden, uniting a number of southern plants with those of the north, as mentioned in the preface.

274. SIBBALDIA. *Gen. pl.* 536.

1. *S. erecta, foliis linearibus multifidis.* *Willd. sp. pl. 1.*  
*p. 1567.*

Icon. *Amm. ruth.* t. 15.

On the banks of the Missouri. *Nuttall.* ♀. July. v. s. in *Herb. Nuttall.* Flowers small, flesh-coloured.

2. *S. procumbens, foliis tridentatis.* *Willd. sp. pl. 1.* *procumbens.*  
*p. 1567.*

Icon. *Fl. dan.* 32. *Pluk. alm.* t. 212. f. 3.

On the Missouri. *Nuttall.* On high mountains of Canada and Vermont. ♀. April. v. s. Flowers yellow.

*longifolia.**brevifolia.**filiformis.**erecta.*

275. STATICE. *Gen. pl. 527.*

- Armeria.* 1. *S. scapo simplici capitato, foliis linearibus planis obtusis.* Willd. sp. pl. 1. p. 1522.  
Icon. Engl. bot. 226. Thrift. Sea Gilliflower. On rocks near the sea-shore: Pennsylvania to Virginia, July, Aug. v. v. Flowers rose-coloured.
- caroliniana.* 2. *S. scapo tereti, panicula divaricato-ramosissima, calycibus acutis, foliis lanceolato-oblongis obtusiusculis glabris margine planis.* —Walt. & car. 118.  
In salt-marshes along the sea-coast: New Jersey to Carolina. ♀. Aug. Sept. v. v. Flowers blue; resembles *St. gmelini* in some respects.

## VI. POLYGYNIA.

276. ZANTHORHIZA. *Gen. pl. 1581.*

- apiifolia.* 1. *Zanthorhiza.* Willd. sp. pl. 1. p. 1568.  
*Zanthorhiza simplicissima.* Marsh. arb. 168.  
Icon. L'Herit. stirp. 1. t. 38.  
On shady banks of rivers: Virginia to Georgia. ♂. May. v. v. A small shrub with compound leaves, and small dark-purple flowers in panicles.

277. SCHISANDRA. *Mich. fl. amer. 2. p. 218.*

- coccinea.* 1. *S. glabra, foliis lanceolato-ovalibus utrinque acutis rarer subdentatis.* Mich. fl. amer. 2. p. 219.  
Icon. Mich. l. c. t. 47. Bot. mag. 1413.  
In shady woods of Carolina and Georgia. ♂. June. v. v. A handsome climbing plant; flowers scarlet.

VI

278. BURN  
cus,  
minu  
279. TILL  
Cor.  
com  
280. TRAD  
Filas  
281. DIPH  
Pet.  
apice  
2-3.  
282. CAUL  
calyc  
Dru  
283. BERE  
teati  
Stig  
284. PRIN  
tat  
285. GLEI  
men  
phy

286. PAN  
tub  
Sta  
287. AMA  
inæ

nis obtusis.

Gilliflower.  
Virginia.

na, calyci-  
nusculis gla-

rsey to Ca-  
blue; re-

81.

Georgia. h.  
ound leaves,

p. 218.

e acutis rari-  
9.

1. h. June.  
vers scarlet.

## VI. HEXANDRIA.

### I. MONOGYNIA.

#### \* *Flores calyculati.*

278. BURMANNIA. *Cal.* inferus, 1-phillus, prismatis-  
cus, coloratus. *Pet.* 3. *Caps.* 3-locularis. *Sem.*  
minuta.
279. TILLANDSIA. *Cal.* inferus, 3-fidus, persists.  
*Cor.* 3-fida, tubulosa. *Caps.* 1—3-locularis. *Sem.*  
comosa.
280. TRADESCANTIA. *Cal.* inferus, 3-phillus. *Pet.* 3.  
*Filam.* villis articulatis. *Caps.* 3-locularis.
281. DIPHYLLEJA. *Cal.* inferus, 3-phillus, deciduus.  
*Pet.* 6. *calyce* opposita. *Anth.* membrana a basi ad  
apicem solubili dehiscentes. *Bac.* 1-locularis. *Sem.*  
2—3. subrotunda.
282. CAULOPHYLLUM. *Cal.* inferus, 6-phillus. *Pet.* 6.  
*calyce* opposita. *Anth.* loculis margine dehiscentes.  
*Drupa* 1-sperma.
283. BERBERIS. *Cal.* inferus, 6-phillus, extus 3-brac-  
teatus. *Pet.* 6. ad unguis glandulis 2. *Styl.* 0.  
*Stigm.* umbilicatum. *Bac.* 1-locularis, 2—4-sperma.
284. PRINOS. *Cal.* inferus, 6-fidus. *Cor.* 1-petala, ro-  
tata, 3—7-fida. *Bac.* 6-sperma. *Dioicus.*
285. GLEDITSCHIA. ♂. *Cal.* 4-fidus. *Pet.* 4. *Legu-*  
*men.* ♂. *Cal.* 3-phillus. *Pet.* 3. ♀. *Cal.* 5-  
phillus. *Pet.* 5. *Legumen.* *Polygama.*

#### \*\* *Flores spathacei.*

286. PANCRATIUM. *Cor.* supera, infundibuliformis :  
tubo longo. *Nectar.* campanulatum, 12-fidum.  
*Stam.* nectario imposta.
287. AMARYLLIS. *Cor.* supera, campanulata. *Stam.*  
inæqualia. *Pist.* declinatum.

288. ALLIUM. *Cor. infera, 6-partita, patens. Spatha multiflora. Umbella congesta. Capsula.*
289. BRODIAEA. *Cor. infera, campanulata, 6-partita. Filam. fauci inserta. Germ. pedicellatum. Caps. 3-locularis : loculis polyspermis.*
290. PONTEDERIA. *Cor. infera, 6-fida, bilabiata. Stam. 3. summo, 3. medio tubo corollæ inserta. Caps. 3-locularis.*
291. HYPOXIS. *Cor. supera, 6-partita, persistens. Spatha 2-valvis. Caps. elongata. Sem. subrotunda, nuda.*
- \*\*\* *Flores nudi.*
292. CONOSTYLIS. *Cor. semi-supera, 6-fida, persistens, lanata pilis ramulosis. Anth. erectæ. Stylus conicus, tripartibilis. Stigm. simplex. Caps. apice dehisces, 3-locularis, polysperma.*
293. ALETRIS. *Cor. semi-supera, tubulosa, 6-fida, rugosa, persistens. Stam. summo tubo inserta. Stylus triqueter, tripartibilis. Caps. apice dehisces, 3-locularis, polysperma.*
294. AGAVE. *Cor. supera, 6-fida : limbo erecto. Filam. corolla longiora, erecta.*
295. PHALANGIUM. *Cor. infera, 6-partita. Stam. filamenti filiformibus, nudis. Sem. angulata.*
296. NARTHECIUM. *Cor. infera, 6-petala, patens. Filam. filiformia, hirsuta. Caps. prismatica, 3-locularis, polysperma. Sem. utrinque arillo subulato instructa.*
297. ORNITHOGALUM. *Cor. infera, 6-partita. Filam. 3. alterna basi dilatata. Caps. subrotunda, 3-locularis. Sem. subrotunda, nuda.*
298. YUCCA. *Cor. infera, campanulata, laciniis non nectariferis. Filam. clavata. Styl. O. Caps. oblonga, obtuse 3-gona. Sem. plana.*
299. LILIUM. *Cor. infera, 6-petala, campanulata : linea longitudinali nectarifera. Stam. stylo breviora. Stigm. indivisum. Caps. 3-gona. Sem. plana.*
300. FRITILLARIA. *Cor. infera, 6-petala, campanulata, laciniis rectis supra unguis cavitate nectarifera. Stam. longitudine corollæ. Sem. plana.*
301. ERYTHRONIUM. *Cor. infera, 6-petala, campanulata, laciniis a medio reflexis, 3. interioribus basi intus 2. callosis. Caps. subglobosa. Sem. oblonga.*
302. UVULARIA. *Cor. infera, 6-petala, campanulata, laciniis rectis, unguibus fovea nectarifera excavatis.*

Filan  
3-val  
globe  
303. STRE  
lata.  
gata,  
304. CONV  
Stan  
bosa,  
cem  
305. SMIL  
dive  
3-loc  
bella  
306. POLY  
Fila  
laris  
307. ASPA  
talis  
poly

308. ORO  
Spa  
lus  
309. ACO  
6-pa  
locu  
310. JUN  
3-lo  
311. PEPI  
poly

312. NEC  
peti  
spei

313. SAB  
cras  
314. CHA

*Spatha*  
5-partita.  
*Caps.* 3.  
a. *Stam.*  
*Caps.* 3.  
ns. *Spa-*  
rotunda,

persistens,  
conicus,  
dehis-

ida, rugo-  
1. *Stylus*  
ens, 3-lo-

o. *Filam.*

*Stam.* fi-  
ta.  
, patens.  
ca, 3-locu-  
subulato

ta. *Filam.*  
, 3-locula-

s non nec.  
s. oblonga,

lata : linea  
breviora.  
plana.  
mpanulata,  
fera. *Stam.*

la, campa-  
ioribus basi  
i. oblonga.  
mpanulata,  
a excavatis.

*Filam.* brevissima. *Stigm.* reflexa. *Caps.* trigona,  
3-valvis : valvis medio septiferis. *Sem.* plura, sub-  
globosa, ad hilum arillata.

303. STREPTOPUS. *Cor.* infera, 6-petala, subcampanu-  
lata. *Stigm.* brevissima. *Bacca* subglobosa, lævi-  
gata, carthacea, 3-locularis. *Sem.* pauca, hilo nudo.
304. CONVALLARIA. *Cor.* infera, 6-fida, campanulata.  
*Stam.* corolla breviora, ad basin inserta. *Bacca* glo-  
bosa, 3-locularis : loculis 1-2-spermis. *Scapus* ra-  
cemosus.
305. SMILACINA. *Cor.* infera, 6-partita, patens. *Filam.*  
divergentia, laciniarum basi infixa. *Bacca* globosa,  
3-locularis. *Flores* terminales paniculati s. subum-  
bellati.
306. POLYGONATUM. *Cor.* infera, 6-fida, cylindrica.  
*Filam.* tubo superne inserta. *Bacca* globosa, 3-locu-  
laris : loculis 2-spermis. *Flores* axillares.
307. ASPARAGUS. *Cor.* infera, 6-partita, erecta : pe-  
talis 3. interioribus apice reflexis. *Bacca* 3-locularis,  
polysperma. *Folia* fasciculata.

\*\*\*\* *Flores incompleti.*

308. ORONTIUM. *Spadix* cylindricus, flosculis tectus.  
*Spatha* O. *Cal.* 6-partitus. *Stigm.* 2-fidum. *Follicu-*  
*lus* 1-spermus.
309. ACORUS. *Spadix* cylindricus, flosculis tectus. *Cal.*  
6-partitus. *Stigm.* punctum prominens. *Caps.* 3-  
locularis, 3-sperma.
310. JUNCUS. *Cal.* inferus, 6-partitus, æqualis. *Stigm.*  
3-lobum. *Caps.* 1-locularis, polysperma.
311. PEPLIS. *Cal.* inferus, 12-fidus. *Caps.* 2-locularis,  
polysperma.

## II. DIGYNIA.

312. NECTRIS. *Cal.* 6-partitus : laciniis 3. interioribus  
petaloideis minoribus obtusis. *Utriculi* bini 1-2-  
spermii. *Sem.* globosa.

## III. TRIGYNIA.

313. SABAL. *Spathæ* partiales. *Filam.* libera, basi in-  
crassata. *Bacca* 1-sperma. *Sem.* osseum.
314. CHAMÆROPS. *Spatha* compressa. *Spadix* ramo-

- sus. *Cal.* 3-partitus. *Pet.* 3. *Filam.* submonadelpha. *Drupæ* 3. monospermæ. *Polygama.*
315. NOLINEA. *Cor.* 6-partita, patens: laciniis subæqualibus. *Styl.* brevissimus. *Stigm.* recurva. *Caps.* 3-gona, membranacea, 3-locularis, dissepimentis bipartilibus dehiscens. *Sem.* solitaria, hinc convexo-incurva.
316. CALOCHORTUS. *Cor.* 6-partita, patens: laciniis 3. interioribus majoribus supra lanatis, basi macula glabra subrotunda notatis. *Filam.* brevissima, basi petalorum inserta. *Anth.* erectæ, sagittatæ. *Stigm.* reflexa. *Caps.* 3-locularis.
317. MELANTHIUM. *Cor.* 6-partita, patens: laciniis æqualibus unguiculatis: unguiculis staminiferis, laciniis basi biglandulosis. *Caps.* subovata, apice subtrifida, 3-locularis. *Sem.* plura, alato-membranacea. *Polygamum.*
318. ZIGADENUS. *Cor.* 6-partita, patens: laciniis subæqualibus, supra basin angustatam biglandulosis. *Stam.* ad contactum ovarii inserta. *Styli* 3. contigui. *Caps.* membranacea, 3-locularis. *Sem.* plurima, aptera.
319. VÉRATRUM. *Cor.* 6-partita, patens: laciniis sessilibus, eglandulosis. *Stam.* receptaculo inserta. *Caps.* 3. polyspermæ. *Polygamum.*
320. HELONIAS. *Cor.* 6-partita, patens: laciniis sessilibus eglandulosis. *Caps.* 3-locularis, 3-cornis seu 3-cocca. *Sem.* pauca (1-2).
321. MEDEOLA. *Cor.* 6-partita, revoluta. *Bacca* 3-sperma.
322. TRILLIUM. *Cor.* 6-petala, patens: laciniis 3. exterioribus angustioribus calycinis. *Bacca* 3-locularis.
323. TOFIELDIA. *Cor.* 6 petala, æqualis, calyculo minimo 3-partito cincta. *Caps.* 3. polyspermæ, basi junctæ.
324. TRIGLOCHIN. *Cal.* 6-phylloides: laciniis 3. interioribus petaloideis. *Cor.* 0. *Styl.* 0. *Caps.* basi dehiscens.
325. SCHEUCHZERIA. *Cal.* 6-partitus. *Cor.* 0. *Styl.* 0. *Stigm.* lateralia. *Caps.* 3. inflatæ, distinctæ, 1-spermæ.
326. RUMEX. *Cal.* 3-phylloides. *Pet.* 3. conniventia. *Sem.* 1. triquetrum.

327. SMILA  
laris328. DIOSC  
locula329. SAUR  
Bacc330. WEN  
succ

mor

331. ALIS  
moi1. B. flor  
Wil

In swa

nov.

I ne

1. T. cæ  
seta

sp.

Icon.

Parasi

v. i

2. T. fil  
bu

Icon.

f. 1

327. SMILAX. *Cal.* 6-phyllo. *Cor.* 0. *Bacca* 3-locularis. *Sem.* 2. *Dioica*.  
 328. DIOSCOREA. *Cal.* 6-partitus. *Cor.* 0. *Caps.* 2-locularis, compressa. *Sem.* 2. membranacea.

## IV. TETRAGYNIA.

329. SAURURUS. *Cal.* amentum squamis 1-floris. *Cor.* 0. *Baccæ* 4. monospermae.

## V. POLYGYNIA.

330. WENDLANDIA. *Cal.* 6-phyllo. *Cor.* 6-petala, succulenta. *Styl.* reclinati. *Caps.* 6. uniloculares, monospermae.  
 331. ALISMA. *Cal.* 3-phyllo. *Pet.* 3. *Caps.* plures, monospermae.
- 

## I. MONOGYNIA.

278. BURMANNIA. *Gen. pl.* 542.

1. B. flore gemino; alis semicuneatis apice truncatis.—  
*Willd. sp. pl. 2. p. 16.*

*biflora.*

In swamps of Virginia. ♀. Sept. v. s. in *Herb. Gronov.* Flower pale purple. This delicate little plant I never was lucky enough to find myself.

279. TILLANDSIA. *Gen. pl.* 541.

1. *T. cæspitosa*, pruinosa; foliis subulatis recurvatis, scapo setaceo foliis longiore summitate subbifloro.—*Willd. sp. pl. 2. p. 14.*

*recurvata.*

Icon. *Sloan. hist.* 1. t. 121. f. 1.  
 Parasitic, on old trees: Florida and Georgia. ♀. June.  
 v. v. Flowers purple.

2. *T. filiformis*, intorta, pruinosa; floribus solitariis sessili- bus.—*Willd. sp. pl. 2. p. 15.*

*usneoides.*

Icon. *Sloan. jam.* 1. t. 122. f. 2 & 3. *Pet. gaz.* t. 62.  
 f. 12.

## HEXANDRIA MONOGYNIA. Tillandsia.

Parasitic, on old trees, in shady woods: Virginia to Florida. ♀. July. v. v. Flowers yellowish-green. This singular plant, known by the name of *Long-moss*, does not extend further north than the neighbourhood of the Dismal Swamp in Virginia. The fibres of it, when divested of the outside coating, make excellent mattresses, and are a very good substitute for horse-hair for that purpose.

## 280. TRADESCANTIA? Gen. pl. 543.

*virginica.*

1. T. erecta; foliis lanceolatis elongatis glabris, floribus sessilibus umbellato-congestis pubescentibus.—Willd. sp. pl. 2. p. 16.

Icon. Bot. mag. 105. Moris. hist. 3. s. 5. t. 2. f. 4.  
In shady woods: Pennsylvania to Carolina. ♀. May, June. v. v. Flowers blue or purple.

*rosea.*

2. T. erecta; foliis linear-gramineis prælongis, pedunculis elongatis umbellatis paucifloris, calycibus glabris.—Mich. fl. amer. 1. p. 193.

Icon. Vent. hort. cels. t. 24.

In wet sandy fields: Carolina and Georgia. ♀. June, July. v. v. Flowers rose-coloured, small, but very handsome.

## 281. DIPHYLLEJA. Mich. fl. amer. 1. p. 203.

*cymosa.*

1. D. glaberrima; foliis subpalmatis angulato-lobatis serratis: lobis acuminatis, cyma multiflora. Mich. l. c.

Icon. Mich. l. c. t. 19 ♂ 20.

Near rivulets on the high mountains of Virginia and Carolina. ♀. May, June. v. v. Flowers white; berries dark blue: the whole of the plant has the appearance of *Podophyllum peltatum*.

## 282. CAULOPHYLLUM. Mich. fl. amer. 1. p. 204.

*thalictroides.*

1. C. glaberrimum; foliis supradecompositis, foliolis ovalibus, inferioribus petiolatis lobatisque, extremo trilobato.—Mich. l. c. t. 21.

Leontice thalictroides. Willd. sp. pl. 2. p. 149.

In shady woods, on the side of fertile hills: New England to Virginia. ♀. May, June. v. v. Flowers yellowish-green; berries large, blueish-black; called by the Indians *Co-hosh*, and esteemed as a medicinal plant among them.

inia to Flo-  
wishes-green.  
*Long-moss,*  
ighbourhood  
ibres of it,  
e excellent  
r horse-hair

3.

ris, floribus  
us.—*Willd.*

2. f. 4.

4. May,

pedunculis  
glabris.—

4. June,  
ll, but very

p. 203.

*lobatis serra-*  
*Mich. l. c.*

Virginia and  
vers white;  
lant has the

1. p. 204.

foliolis ova-  
extremo tri-

19.

: New En-  
v. Flowers  
black; called  
a medicinal



*Berberis aquifolium.*

Tut.





*Berberis nervosa.*

Drawn & Engraved by

1. *B. erecta*  
foliis si  
vibus s  
*B. vulgar*  
amer. 1  
On fertile  
h. Ap  
lous ra
2. *B. sarme*  
jugis o  
tatis.  
*B. pinna*  
On the g  
in rich  
v. s. in  
ters ;  
compa  
*Caulis* f  
procun  
data, i  
lia, im  
cata,  
utrinco  
*Petiol*  
gemm  
ducæ.  
ceæ.  
nimus  
triplo  
nervo  
nacei  
incise  
6. co  
oppo  
ovatu  
3-spe
3. *B. sarm*  
ovato  
lis in  
In the  
The sp  
prec  
and

## 283. BERBERIS. Gen. pl. 595.

1. *B. erecta*; ramis confertim punctatis, aculeis triplicibus, foliis simplicibus obovatis remote serratis, racemis brevibus subcorymbosis, drupis vix carnosis.

*B. vulgaris* var. *Willd. sp. pl. 2. p. 227. Mich. fl. amer. 1. p. 205.*

On fertile hills, among rocks, &c. Canada to Virginia.

↳ April, May. v. v. Flowers yellow, in pendulous racemes; berries red, of an agreeable acid.

2. *B. sarmentosa*, inermis; foliis pinnatis: foliolis sub-3-jugis oblongis repando-dentatis venosis, petalis bidentatis.

*B. pinnata. Herb. Banks.*

On the great rapids of Columbia River, among rocks, in rich vegetable soil. *M. Lewis.* ↳ April, May. v. s. in *Herb. Lewis.* Flowers yellow, in large clusters; berries dark purple, eatable; called by Lewis's company *Mountain-holly.*

*Caulis* fruticosus, laxe ramosus: ramis sarmentosis, procumbentibus. *Folia* sempervirentia alterna, petiolata, impari-pinnata. *Foliola* 3-juga, opposita, sessilia, impari-petiolata, oblongo-ovata, basi oblique truncata, margine cartilaginea repando-dentata, coriacea, utrinque glabra, laevigata, nitida: dentibus aculeatis. *Petioles* teretes, glabri. *Racemi* congesti, bracteati, e gemma precedentis anni. *Flores* aurei. *Bracteæ* caducæ, solitariae, subcordatae, acuminatae, membranaceæ. *Calyx* triplex, deciduus, patens: exterior minimus, 3-phyllos: foliolis ovatis, acutis; medius tripli longior: foliolis suborbiculatis, membranaceis, nervosis; interior longior: foliolis ovalibus, membranaceis, nervosis. *Petala* 6. suberecta, oblonga, apice inciso-bidentata, vix longitudine calycis. *Filamenta* 6. corollæ breviora, crassa, medio bidentata: dentibus oppositis. *Antheræ* bilobæ, crassæ. *Germen* superum, ovatum. *Stigma* sessile, 3-lobum. *Bacca* 3-locularis, 3-sperma, abortione interdum monosperma.

3. *B. sarmentosa*, inermis; foliis pinnatis: foliolis 6-jugis ovato-oblongis repando-serratis, sub-5-nervibus, petalis integris.

In the same situations. v. s. in *Herb. Lewis.*

The specific difference excluded, the description of the preceding species is applicable in every other respect, and together with another in the collection of A. B.

*canadensis.*

*Aquifolium.*

*nervosa.*

## HEXANDRIA MONOGYNIA. Berberis.

Lambert, Esq., collected in Napaul by Mr. Buchanon, forms a new division of the genus, with pinnated leaves; which probably may become a new genus, whenever the fruit is perfectly known, as the statement I have given of it was taken from a single and imperfect berry.

## 284. PRINOS. Gen. pl. 594.

- verticillatus.* 1. P. foliis deciduis ovalibus serratis acuminatis subtus pubescentibus, fasciculis florum masc. axillaribus umbelliformibus; femineis aggregatis utrinque 6-partitis.—*Willd. sp. pl. 2. p. 225.*  
*P. Gronovii. Mich. fl. amer. 2. p. 236.*  
*P. padifolius. Willd. enum. 394.*  
*Icon. Duham. arb. 1. t. 23.*  
 In wet woods, on the banks of ditches; common: Canada to Virginia.  $\frac{1}{2}$ . June, July. v. v. Flowers small, white; berries red or crimson.
- amliguus.* 2. P. foliis deciduis ovalibus utrinque acuminatis mucronato-serrulatis subtus pubescentibus, floribus 4-5-fidis, masculis ad imos ramulos congestis, femineis solitariis.—*Mich. fl. amer. 2. p. 236.*  
*Cassine caroliniana. Walt. fl. car. 242.*  
 In sandy wet woods and on the borders of swamps: New Jersey to Carolina.  $\frac{1}{2}$ . July, Aug. v. v. Flowers white; berries red, larger than No. 1.
- lævigatus.* 3. P. foliis deciduis lanceolatis adpresso-serratis acuminatis utrinque glabris supra nitidis, subtus ad nervos vix pubescentibus, floribus femineis axillaribus solitariis subsessilibus: masculis sparsis: omnibus 6-fidis.  
 On the Allegany-mountains: New York to Virginia.  $\frac{1}{2}$ . July. v. v. Berries large, dark-red.
- lanceolatus.* 4. P. foliis deciduis lanceolatis tenuissime et remote serrulatis utrinque acutis utrinque glabris, floribus femineis sparsis subgeminis pedunculatis 6-fidis, masculis aggregatis 3-andris.  
 In the lower counties of Carolina and Georgia.  $\frac{1}{2}$ . June. v. s. Berries small, scarlet.
- glaber.* 5. P. foliis sempervirentibus cuneato-lanceolatis coriaceis glabris nitidis superne pariter subdentatis, pedicellis axillaribus subsolitariis plerumque 3-floris.—*Willd. sp. pl. 2. p. 226.*  
 In sandy shady woods: Canada to Florida.  $\frac{1}{2}$ . July,

Aug.  
 black;  
 6. P. foliis  
 glabri;  
 vissim;  
 a. P. foliis  
 β. P. foliis  
 In sand  
 Ensle  
 shrub  
 Prinos l  
 specie  
 2  
 1. G. spin  
 oblor  
 sp. p  
 G. Mel  
 Icon.  
 β. G. ram  
 Near t  
 v. v  
 kno  
 2. G. spin  
 obtu  
 G. tria  
 In the  
 gini  
 stin  
 tim  
 bra  
 3. G. ra  
 leg  
 W  
 G. ca  
 G. aq  
 Icon.  
 In the  
 and  
 1. P. sp  
 se  
 pl

uchanon,  
ed leaves;  
whenever  
ent I have  
imperfect

subtus pu-  
ribus um-  
jue 6-par-

mon : Ca-  
. Flowers

mucronato-  
4-5-fidis,  
ineis solita-

of swamps:  
Aug. v. v.  
o. 1.  
; acuminatis  
nervos vix  
bus solitariis  
6-fidis.  
to Virginia.

note serrula-  
nas feminineis  
masculis ag-

orgia. h.

itis coriaceis  
is, pedicellis  
ris.—Willd.

h. July,

Aug. v. v. A low and handsome shrub; berries black; called in Jersey *Ink-berries*.

6. P. foliis sempervirentibus cuneato-lanceolatis coriaceis glabris nitidis integerrimis, corymbis axillaribus brevissimis sessilibus multifloris, floribus 6-fidis.

α. P. foliis obovato-lanceolatis acuminatis.

β. P. foliis lanceolatis acutis.

In sandy woods, near the banks of rivers: Georgia.

*Enslen.* h. June, July. v. s. A handsome tall shrub, of the appearance of *Ilex Dahoone*.

*Prinos lucidus* Ait. is *Ilex canadensis* Mich. according to specimens in the Herbarium of A. B. Lambert, Esq.

*coriaceus.*

*latifolia.*

*angustifolia.*

#### 285. GLEDITSCHIA. Gen. pl. 1526.

1. G. spinis robustis cruciatim ramosis, foliolis linearioribus oblongis, leguminibus longissimis compressis.—*Willd.* sp. pl. 4. p. 1097.

*G. Meliloba.* *Walt. fl. car.* 254.

*Icon. Duham. arb.* 1. t. 195. *Mich. arb. t.* . . . . .

β. G. ramis subinermibus.

Near the sea-coast: Virginia and Carolina. h. July.

v. v. A large tree; the pods contain a sweet pulp; known by the name *Honey-locust*.

*triacanthos.*

*inermis.*

2. G. spinis crassis brevibus subternatis, foliolis oblongis obtusis, leguminibus oblongis brevibus.

*G. triacanthos* β. *brachycarpos.* *Mich. fl. amer.* 2. p. 257.

In the Allegany mountains: on the Indian river, Virginia. h. July. v. v. This species is certainly distinct from No. 1, having short curved spines, sometimes solitary, sometimes by threes, on the old branches.

3. G. ramis subspinosus, foliolis ovato-oblongis acutis, *monosperma* leguminibus ovalibus mucronatis submonospermis.—*Willd. sp. pl. 4. p. 1097.*

*G. carolinensis.* *Lam. encycl.* 2. p. 461.

*G. aquatica.* *Marsh. arb.* 95.

*Icon. Mich. arb.* . . . . .

In the woods and on river banks: Carolina to Florida, and Illinois. h. July. v. v. *Swamp Locust-tree.*

#### 286. PANCRATIUM. Gen. pl. 551.

1. P. spatha subbiflora, foliis lanceolatis, nectariis dentibus sex staminiferis, intermediis simplicibus. *Willd. sp. pl. 2. p. 42.*

*mexicanum.*

Icon. *Dill. eth. t. 222. f. 289.*

On the sea-coast of Carolina and Georgia. ♀. July, Aug. v. v. Flowers white, as are all the following.

*rotatum.*

2. P. spatha multiflora, foliis linear-lanceolatis, nectariis hypocrateriformibus, inferne tubulosis, dentibus sex staminiferis, intermediis inciso-dentatis, staminibus nectario duplo longioribus.—*Ker in bot. mag.* 827.

On the sea-coast: Virginia to Florida. ♀. July, Aug. v. v.

*maritimum.*

3. P. spatha multiflora, foliis linear-lanceolatis, nectariis dentibus duodecim, non staminiferis. *Willd. sp. pl. 2. p. 42.*

P. carolinianum. *Linn. sp. pl. 418.*

P. verecundum. *Ait. kew. 1. p. 412.*

Icon. *Catesb. car. 2. app. 5. t. 5. Salisb. in linn. trans. 2. t. 9.*

On the coast of Carolina and Georgia. ♀. July, Aug. v. v.

### 287. AMARYLLIS. *Gen. pl. 554.*

*Atamasco.*

1. A. spatha bifida acuta, flore pedicellato, corolla campanulata subæquali erecta, basi brevi tubulosa, staminibus declinatis æqualibus. *Willd. sp. pl. 2. p. 51.*

Icon. *Bot. mag. 239. L'Herit. sert. angl. 10. Catesb. car. app. t. 12.*

In shady woods and swamps: Virginia and Carolina. ♀. June, July. v. v. Flowers rose-coloured.

### 288. ALLIUM. *Gen. pl. 557.*

*vineale.*

1. A. caule teretifolio bulbifero, staminibus tricuspidatis. *Willd. sp. pl. 2. p. 73.*

Icon. *Lob. ic. 156.*

In old fields; common. ♀. June, July. v. v.

*fragrans.*

2. A. scapo nudo subtereti, foliis linearibus planis acutis subcarinatis, corollis turbinato-campanulatis, laciniis obovatis, staminibus linearibus acuminatis tubo adnatis.—*Vent. malmais. 26. Redouté lil. 2. t. 68.*

A. inodorum. *Bot. mag. 1129.*

A. mutabile. *Mich. fl. amer. 1. p. 195?*

On the mountains of Virginia and Carolina. ♀. June. v. v. Flowers white, with red veins.

*striatum.*

3. A. scapo nudo subtriquetro, foliis linearibus convoluto-concavis obtusiusculis dorso striatis, corollis patentibus,

- laciniis oblongo-lanceolatis, staminibus alternis sub
  - latis, alternis compressis.—*Bot. mag.* 1524. *Jacq.*  
*ic.* 2. *t.* 366. *Willd. sp. pl.* 2. *p.* 77.
  - A. ornithogalooides.* *Walt. fl. car.* 121.  
*Ornithogalum bivalve.* *Sp. pl. ed.* 1. *p.* 306.  
In Virginia and Carolina. ♀. June, July. *v. s. in*  
*Herb. Walt.* Flowers large, white, with a red carina.
  - 4. *A. scapo nudo ancipi, foliis linearibus canaliculatis sub-*  
*tus subangulatis, umbella fastigiata.* *Willd. sp. pl.* 2.  
*p.* 76.  
Icon. *Gmel. sib* 1. *t.* 14. *f.* 2. *Jacq. austr.* 5. *t.* 425.  
On the banks of the Missouri. *M. Lewis* and *Nuttall.*  
♀. July. *v. s. in Herb. Lewis.* Flowers white.
  - 5. *A. scapo nudo tereti foliis breviore, foliis lanceolatis ner-*  
*vosis, umbella pauciflora.*  
In shady woods, on the high mountains of Pennsylvania.  
♀. May, June. *v. v.* *Mountain Leeks.*
  - 6. *A. scapo nudo tereti, foliis linearibus, capitulo bulbifero.* *Willd. sp. pl.* 2. *p.* 78.  
In fields and woods: Canada to Carolina. ♀. June.  
*v. v.*
  - 7. *A. scapo nudo semitereti, foliis lanceolato-oblongis pla-*  
*nis glabris, umbella globosa, seminibus solitariis.*  
*Willd. sp. pl.* 2. *p.* 81.  
In shady woods: Pennsylvania to Virginia. ♀. July.  
*v. v.*

289. BRODIÆA. *Smith in linn. trans.* 10. *p.* 2.

1. *B. umbella multiflora, floribus pedicellatis, staminibus* *grandiflora.*  
alternis margine membranaceis.—*Smith l. c.*  
*Hookera coronaria.* *Salisb. par. lond.* *t.* 98.  
On the plains of the Columbia and Missouri rivers.  
*M. Lewis.* ♀. April, May. *v. v.* This elegant  
bulbous plant, which M. Lewis, Esq. used to call  
*Missouri Hyacinth*, is certainly hexandrous; Dr. Smith  
having been misled, by dried specimens, to consider it  
to be only triandrous.

290. PONTEDERIA. *Gen. pl.* 545.

1. *P. foliis oblongo-cordatis obtusis, spica confertim-multi-*  
flora, corollæ laciniis oblongis.—*Willd. sp. pl.* 2.  
*p.* 23. *cordata.*

Icon. *Pluk. mant. t. 349. f. ult. Bot. mag. 1156.*

On the edges of rivers in the lower countries, from Canada to Florida.  $\frac{1}{2}$ . July. v. v. Flowers blue.

- angustifolia.* 2. *P. foliis elongate-triangularibus sensim acutis basi truncato-subcordatis, corollæ laciniis linear-lanceolatis.*  
In high mountain-lakes: New York to Carolina.  $\frac{1}{2}$ . July. v. v. Flowers blue, smaller than the first species.

291. HYPOXIS. *Gen. pl. 565.*

*erecta.*

1. *H. pilosa; scapo 2-3-floro foliis linear-lanceolatis breviore, corollæ laciniis lanceolato-oblongis.* — *Willd. sp. pl. 2. p. 106.*

Icon. *Pluk. adm. t. 315. f. 12. Bot. mag. 710.*

In pastures and on grassy hills: Canada to Virginia.  $\frac{1}{2}$ . June, July. v. v. Flowers yellow: the root is reckoned one of the antidotes against the bite of the rattle-snake.

*graminea.*

2. *H. pilosa; scapo 4-floro foliis longissimis gramineis breviore, corollæ laciniis linear-lanceolatis.*

*H. carolinensis. Mich. fl. amer. 1. p. 118.*

In dry sandy fields and woods: New Jersey to Carolina.  $\frac{1}{2}$ . June, July. v. v. Leaves very narrow, and double the length of the scape.

*junccea.*

3. *H. pilosa; scapo uniflоро, foliis filiformibus canaliculatis.* — *Smith spicil. t. 16.*

On pine-barrens of South Carolina.  $\frac{1}{2}$ . May—July. v. v. *H. sessilis Willd. sp. pl. 2. p. 110.* figured by Dillenius t. 220. is only an accidental variety of *H. erecta*, as I frequently have found this species at an early part of the season to flower very near to the ground.

292. CONOSTYLIS. *Brown fl. nov. holland. 1. p. 300.*

*americana.*

1. *C. corollis intus lanatis, scapis corymboso-paniculatis, foliis gramineo-ensiformibus glaucis, filamentis aquilibus.*

*Argolasia aurea. Pursh catal.*

In boggy soil, on the pine-barrens of New Jersey and Carolina.  $\frac{1}{2}$ . July. v. v. Flowers golden-yellow. *Radix fibrosa, repens. Folia radicalia, scapo breviora, irideo-equitantia, angusto-ensiformia, acuta, glabra, laevigata, glaucescentia. Scapus erectus, teres, lato-tomentosus, folio uno altero brevi instructus.*



*Conostylis americana*.

Drawn & engraved by J. C. Green

*Coryn*  
extus  
florua  
interie  
luteis.  
reis.  
viora.  
super  
3-par  
plex.

This s  
about  
*Argo*  
had t  
mus,  
any g  
genus  
super

### 293. ALE

1. A. florif  
tiferis  
crona  
A. farin  
Icon. A  
In dry g  
gland  
white  
grass  
consist

2. A. florif  
tis, co  
lato-  
Wurmbr  
In sand  
jina.  
a me  
divisi  
are n  
this p  
but  
and c

*Corymbus* dense multiflorus, lanato-tomentosus. *Flores* extus albo-lanati, intus flavi. *Pedicelli* longitudine florum. *Corolla* 6-fida: laciniis oblongis acutis, 3. interioribus vix angustioribus, superne glabris fusco-luteis, inferne densissime lanatis: pilis pluinosis aureis. *Filamenta* 6. filiformia, glabra, corollæ vix breviora. *Antheræ* ovatæ, erectæ, aurantiacæ. *Germen* superum, subrotundum, glabrum. *Stylus* subulatus, 3-partibilis, longitudine filamentorum. *Stigma* simplex.

This singular and beautiful perennial I discovered about the year 1805, and considered it as a species of *Argolasia* of Jussieu or *Lanaria* of Aiton, until I had the opportunity of seeing Mr. Brown's *Prodromus*, in which the genus *Conostylis* is the nearest of any genus I know, though it even differs from that genus materially by the germ being almost entirely superior, and not half inferior.

293. ALETRIS. *Gen. pl. 579. Mich. fl. amer. 1. p. 189.*

1. *A. floribus pedicellatis oblongo-tubulosis, corollis fructiferis læviusculis farinosis, foliis lato-lanceolatis mucronatis.* — *Mich. fl. amer. 2. p. 189.*

*alba.*

*A. farinosa. Willd. sp. pl. 2. p. 183.*

Icon. *Pluk. amalh. t. 437. f. 2. Bot. mag. 1418.*

In dry gravelly woods, on the side of hills: New England to Carolina. ♀. June, July. v. v. Flowers white, in a long spike; known by the name of *Star-grass* or *Colic-root*; the last name, from its being considered an excellent remedy in that disease.

2. *A. floribus subsessilibus breviter tubulatis subcampanulatis, corollis fructiferis rugosis scaberrimis, foliis lanceo-lato-ensiformibus acutis.* — *Mich. fl. amer. 1. p. 190.*

*auræa.*

*Wurmbea bullata. Willd. hort. berol. t. 8.*

In sandy fields and woods: New Jersey to South Carolina. ♀. July, Aug. v. v. Flowers covered with a mealy substance, as in the first species, but the divisions are tipped with golden-yellow; the leaves are narrower and longer. There is not any doubt of this plant properly belonging to *Hexandria Trigynia*; but the styles are glued together during flowering, and only divide on the ripening of the fruit.

294. AGAVE. *Gen. pl. 582.*

- virginica.* 1. A. acaulis, herbacea; foliis cartilagineo-serratis, scapo simplicissimo. *Willd. sp. pl. 2. p. 193.*  
*Icon. Jacq. ic. rar. 2. t. 378. Bot. mag. 1157.*  
 On the rocky and fertile banks of rivers: Virginia to Carolina. ♀. July. v. v. Flowers greenish-yellow, very fragrant.

295. PHALANGIUM. *Tournef. Juss. gen. pl. 59.*

- croceum.* 1. P. bulbosum; scapo foliis gramineis multo brevioribus, spica pyramidali, bracteis convoluto-amplexantibus obtusis, seminibus subglobosis laevigatis.—*Mich. fl. amer. 1. p. 196.*  
 In low grounds of Georgia. *Michaux. ♀. +.* Flowers saffron-coloured; seeds black.  
*Quamash.* 2. P. bulbosum; scapo foliis linearibus carinatis longiore, spica racemosa, bracteis linearibus, petalis 5. adscendentibus: infimo deflexo, stigmate obtuso.  
 On the upper part of the Missouri, near the Rocky mountains. *M. Lewis. ♀. June. v. s. in Herb. Lewis.* Flowers large, pale blue.

*Bulbus tunicatus, subrotundus. Scapus simplex, nudus, erectus, teres, glaber, pedalis et ultra. Folia radicalia, paucia, longe linearia, subtus carinata, glabra, semiuncinata lata. Flores spicato-racemosi, bracteati. Bracteae marcescentes, solitariae, lineares, membranaceae. Pedunculi solitarii, uniflori, teretes, bracteis breviores. Corolla marcescens, infera, hexapetalopartita. Petala subæqualia, lineari-lanceolata: quinque adscendentia: sextum deflexum. Filamenta subulata, æqualia, petalis breviora, ad basin petalorum inserta. Germen trigonum. Stylus longitudine corollæ, subclavatus. Stigma obtusum. Capsula subrotundo-triquetra, 3-locularis: loculis polyspermis. Semina oblonga, nigra.*

It deviates from the character of *Phalangium* in some respects, particularly in its ascending petals, but not sufficiently to give materials to frame a new genus.

This plant is known among the natives by the name *Quamash*, and the bulbs are carefully collected by them and baked between hot stones, when they assume the appearance of baked pears, and are of an

agn  
win  
Lev  
eate

## 296. NA

1. N. rac  
cau  
cell  
Ke  
In bo  
Jer  
app  
diff  
dist  
Foi  
N.  
of  
ing

2. O. ce  
fila  
Icon.  
In gra  
♀.

1. Y. ac  
fila  
W  
Icon.  
On t  
we  
wk  
fee

2. Y. ac  
fila  
On t  
in  
ve

agreeable sweet taste. They form a great part of their winter stores. Though an agreeable food to Governor Lewis's party, they occasioned bowel complaints if eaten in any quantity.

296. NARTHECIUM. *Smith fl. brit. 1. p. 368. Ker in bot. mag. 1505.*

1. *N. racemo interdum interrupte spicato laxo, bractea una americanum.*  
caulina pedicellum amplexante, altera inferne pedicellari paleaceo-setacea, lana filamentorum brevissima.  
*Ker l. c.*

In boggy fields and woods, on the pine-barrens of New Jersey.  $\frac{1}{4}$ . June, July. v. v. Flowers yellow. It approaches very near to *N. Ossifragum*, and is only different in the situation of its bracteas, which nice distinction was very ingeniously observed by Mr. Ker. For a considerable time I considered this plant to be *N. glutinosum* of Michaux; but the close investigation of Mr. Ker on this subject has convinced me of having been in an error. *Vid. Bot. mag. 1505.*

297. ORNITHOGALUM. *Gen. pl. 566.*

1. *O. corymbo paucifloro, pedunculis bracteis longioribus, umbellatum.*  
filamentis subulatis. *Willd. sp. pl. 2. p. 116.*  
Icon. *Jacq. austr. 4. t. 343.*  
In grass-plots and near gardens; probably not indigenous.  
 $\frac{1}{4}$ . May. v. v. Flowers white.

298. YUCCA. *Gen. pl. 580.*

1. *Y. acaulis; foliis lato-lanceolatis integerrimis margine filamentosis, stigmatibus recurvato-patentibus.—*  
*Willd. sp. pl. 2. p. 184.*  
Icon. *Trew ehret. t. 37.*  
On the shores of Virginia and Carolina, and in the western parts thereof.  $\frac{1}{4}$ . July, Aug. v. v. Flowers white, very showy; and the plant is from four to five feet high.
2. *Y. acaulis; foliis longo-linearibus rigidis margine raro angustifolia.*  
filamentosis, capsulis magnis obovato-cylindraceis.  
On the banks of the Missouri.  $\frac{1}{4}$ . July, Aug. v. s.  
*in Herb. Nuttall.* From two to three feet high; leaves very narrow; capsules large.

*recurvifolia.* 3. *Y. caulescens*; foliis linear-lanceolatis ~~viridibus~~ recurvo-deflexis margine raro filamentosis, petalis interioribus latioribus.—*Salisb. in parad. lond.* 31.

On the sandy shores of Georgia. *Leconte.* ♀. July, Aug. v. v. Flowers greenish-yellow, with a tinge of purple; stem about three feet high: this species has been confounded with the following.

*gloriosa.* 4. *Y. caulescens*, *ramosa*; foliis lato-lanceolatis plicatis integerrimis, petalis lanceolatis.—*Willd. sp. pl.* 2. p. 183.

*Icon. Bot. rep.* 473. *Bot. mag.* 1260.

On the sea-shore of Carolina. ♀. July, Aug. v. v. Flowers white; plant about ten feet high.

*aloifolia.* 5. *Y. caulescens*, *ramosa*; foliis linear-lanceolatis callosocrenulatis strictis.—*Willd. sp. pl.* 2. p. 184.

*Icon. Dill. elth.* 323. f. 416. *Comm. præl. t.* 14.

On the coast of Carolina and Florida. ♀. Aug. v. v. Flowers white.

### 299. LILIUM. Gen. pl. 558.

*Catesbeii.* 1. *L. foliis sparsis linear-lanceolatis*, *caule unifloro*, *corolla erecta*, *petalis longe unguiculatis* *margine undulatis* *apice reflexis*. *Willd. sp. pl.* 2. p. 86. *Walt. fl. car.* 259.

*L. spectabile*. *Salisb. icon. rar.* 5. t. 5.

*L. carolinianum*. *Lam. encycl.* 3. p. 554.

*Icon. Catesb. car.* 2. t. 58. *Bot. mag.* 259.

In low sandy meadows: Virginia to Carolina. ♀. June—Aug. v. v. Flower very handsome, scarlet, spotted with yellow and brown; stem about a foot high.

*pudicum.* 2. *L. ? foliis sparsis linear-lanceolatis*, *caule unifloro*, *corolla campanulata cernua*, *petalis erectis sessilibus* *spatulato-ovatis obtusis intus planis*.

On the head-waters of the Missouri. *M. Lewis.* ♀. May. v. s. in *Herb. Lewis*. Flower pale yellow; stem about ten inches high.

This doubtful species is at first sight more related to *Fritillaria*; but the style, which is the length of the petals, with an obtuse stigma, associates it more closely to *Lilium*. The fruit I am unacquainted with; the root is a flat biscuit-shaped bulb.



*Silium pudicum.*

3. L. foliis sp  
cillatis  
subalato  
turbinate  
ovali-lan

Icon. *Cate*

On the n  
July, At  
with da  
much.

4. L. foliis sp  
peduncu  
tis, peta  
oblongis

L. concolor

On the bar

24. July  
by the a  
on the M  
figured i  
of an un  
namenta

5. L. foliis ve  
ro, corol  
latis ung

Icon. *Mill*

In woods a  
Aug. v  
bottom,  
with onl  
or more.

6. L. foliis re  
ad nervis  
gatis plei  
panulatis

*Willd. s*

Icon. *Cate*

$\alpha$ . Bot. mag. 8  
 $\beta$ . Bot. mag. 8

On mounta

Aug. v

7. L. foliis pl  
obovalibi

3. *L. foliis sparsis linear-i-lanceolatis : supremis 4.5. verticillatis floremque subæquantibus, caule subunifloro subalato-pentagono, pedunculo lanato, corolla erécta turbinato-patente extus floccoso-lanata, lacinias infimis ovali-lanceolatis.* *Ker in bot. mag. 872.*

Icon. *Catesb. car. 3. t. 8.*

On the mountains of Pennsylvania and Virginia.  $\frac{4}{4}$ .

July, Aug. *v. v.* Flower scarlet, yellow at the base with dark spots. Resembles *L. bulbiferum* very much.

4. *L. foliis sparsis linearibus brevibus : supremis verticillatis pedunculis brevioribus, floribus 3—5. umbellatis erectis, petalis recurvo-patentibus subæqualibus ovato-oblongis stibunguiculatis.* *umbellatum.*

*L. concolor. Parad. lond. 47?*

On the banks of the Missouri. *M. Lewis and Nuttall.*

$\frac{4}{4}$ . July. *v. s.* By all appearance the plant found by the above gentlemen, among the Mandan nation on the Missouri, appears to be the same with the one figured in the *Paradisus Longdinensis*; the flowers are of an uniform deep scarlet colour, and are highly ornamental.

5. *L. foliis verticillatis lanceolato-linearibus, caule subbifloro, corolla erecta campanulata patente, petalis lanceolatis unguiculatis.* — *Willd. sp. pl. 2. p. 90.*

Icon. *Mill. ic. t. 165. f. 1. Bot. mag. 519.*

In woods and meadows: Canada to Virginia.  $\frac{4}{4}$ . July, Aug. *v. v.* Flowers large, scarlet, yellow near the bottom, with dark red spots; most generally it is found with only one flower, but sometimes even with three or more.

6. *L. foliis remote verticillatis lanceolatis trinerviis, subtus ad nervos subhirsutis, pedunculis terminalibus elongatis plerumque ternis, corollis cernuis turbinato-campanulatis recurvo-patentibus, lacinias lanceolatis.* — *Willd. sp. pl. 2. p. 89.*

Icon. *Catesb. car. 3. t. 11.*

$\alpha$ . *Bot. mag. 800.*

$\beta$ . *Bot. mag. 858.*

On mountain meadows: Canada to Virginia.  $\frac{4}{4}$ . July, Aug. *v. v.*

7. *L. foliis plerumque verticillatis cuneato-lanceolatis seu obovalibus è nervibus glaberrimis, ramis floriferis crassis*

*pensylvani-*  
*cum.*

*philadelphi-*  
*cum.*

*canadense.*

*flavum.*  
*coccineum.*

*carolinia-*  
*num.*

*ternato-terminalibus, floribus reflexis, corollis revolutis.* — *Mich. fl. amer. 1. p. 197.*

1. *L. Martagon.* *Walt. fl. car. 123.*

Icon. *Catesb. car. 2. t. 56.*

In fertile meadow grounds : Lower Virginia and Carolina. ♀. June, July. v. v. Resembles *L. Martagon* very much ; flowers larger and maculated.

*superbum.*

8. *L. foliis linear-lanceolatis trinervibus glabris, inferioribus verticillatis intermediis duplo longioribus superioribus sparsis, floribus racemoso-pyramidalis reflexis, corollis revolutis.* — *Willd. sp. pl. 2. p. 88.*

Icon. *Trew ehret. 2. t. 11. Bot. mag. 936.*

In low copses and swamps, on blue clay soil : Canada to Virginia. ♀. July, Aug. v. v. The most stately plant of this family in North America ; the number and elegance of its flowers are astonishing. I have, in favourable situations, seen stems of about seven feet high, with from thirty to fifty flowers, disposed in a graceful manner in form of a chandelier.

### 300. FRITILLARIA. *Gen. pl. 559.*

*lanceolata.*

1. *F. caule folioso 1—2-floro, foliis linear-lanceolatis obtusiusculis : inferioribus verticillatis, petalis lanceolatis.*

*Lilium camschatcense.* *Willd. sp. pl. 2. p. 89.*

On the head-waters of the Missouri and Columbia.

*M. Lewis.* ♀. July. v. s. Flowers dark purple with brown spots.

In the tenth volume of the Linnean Transactions, A. B. Lambert, Esq. in his account of the Pallasian Herbarium, has given an excellent figure of this plant, and suspects it very justly to be a Fritillaria : when I examined the Lewisian Herbarium I had no idea of finding it under *Lilium*, and made a drawing and description under the above name, for the Travels of Lewis and Clark.

### 301. ERYTHRONIUM. *Gen. pl. 562.*

*lanceolatum.*

1. *E. petalis lanceolatis basi dilatatis obtusiusculis, germine subgloboso, foliis lanceolatis.*

*E. Dens canis.* *Mich. fl. amer. 1. p. 198.*

In woods, near the roots of old trees, and on the banks of rivulets : Canada to Virginia, and on the Missouri.

♀. Ma  
with br

2. *E. petalis  
longo,*

On the K  
Flower  
yellow

### 302. UVUL

1. *U. foliis p  
intus sc  
p. 94.*

*U. perfoli*

Icon. *Sm*  
In shady  
Canada  
pale ye

2. *U. foliis  
tis, co  
pidatis.*

Anonymous  
Icon. *Bo*

In shady  
Carolin  
No. 1,

3. *U. foliis  
anther  
exot. b*

*U. perfo*

*U. lance*

Icon. *Co*  
On shad  
to Ca  
other  
*lata o*

*Smith*  
*Herba*

4. *U. caule  
subtu  
vibus.*

Icon. *S*  
In shad  
v. v.

1. May. v. v. Flowers yellow: there is a variety with brown spots at the bottom of the flower.
2. E. petalis laeceolatis subunguiculatis acutis, germine oblongo, foliis linear-lanceolatis.
- On the Kooskoosky. M. Lewis. 1. May, June. v. s. Flowers double the size of the preceding, and of a pale yellow colour.

*grandiflorum.*

## 302. UVULARIA. Gen. pl. 560. Mich. fl. amer. 1. p. 198.

1. U. foliis perfoliatis ellipticis obtusis, corolla campanulata intus scabrate, antheris cuspidatis.—Willd. sp. pl. 2. p. 94. Smith exot. bot. 1. p. 95.  
U. perfoliata minor. Mich. fl. amer. 1. p. 199.  
Icon. Smith l. c. t. 49.  
In shady woods, among rocks, in rich vegetable mould: Canada to Carolina. 1. May, June. v. v. Flowers pale yellow.
2. U. foliis perfoliatis elliptico-oblongis obtusis basi undulatis, corolla basi attenuata intus scabrate, antheris cuspidatis. Smith exot. bot. 1. p. 97. t. 50.  
Anonymos pudica. Walt. fl. car. 123.  
Icon. Bot. mag. 955. Smith l. c.  
In shady woods, in sandy soil: New Jersey to Lower Carolina. 1. May, June. v. v. Flowers larger than No. 1, of a deeper yellow.
3. U. foliis perfoliatis oblongis acutis, petalis intus laevibus, antheris obtusiusculis, nectariis subrotundis. Smith exot. bot. 1. p. 99. t. 51.  
U. perfoliata major. Mich. fl. amer. 1. p. 199.  
U. lanceolata. Willd. sp. pl. 2. p. 94.  
Icon. Corn. canad. t. 39. Smith l. c.  
On shady hills, in fertile soil, and among rocks: Canada to Carolina. 1. June. v. v. Flowers larger than the other species, of a beautiful yellow. The *U. lanceolata* of the Hortus Kewensis is the same with Dr. Smith's *U. grandifolia*, according to specimens in the Herbarium of A. B. Lambert, Esq.
4. U. caule glabro, foliis sessilibus lanceolato-ovalibus subtus glaucescentibus, corollæ laciniis planis intus laevibus, capsula stipitata.—Willd. sp. pl. 2. p. 95.  
Icon. Smith exot. bot. 1. p. 101. t. 52. Bot. mag. 1402.  
In shady woods: Canada to Carolina. 1. May, June. v. v. Flowers pale yellow.

*perfoliata.*

*flava.*

*grandiflora.*

*sessilifolia.*

*puberula.*

5. *U. caule puberulo, foliis subamplexicaulibus ovalibus basi rotundatis utrinque concoloribus, corollæ lacinias acutis intus lævibus, capsula sessili.* — *Mich. fl. amer. 1. p. 199.*

On the mountains of Carolina. *Michaux. 4. +.* Resembles the foregoing species; flowers larger.

303. STREPTOPUS. *Mich. fl. amer. 1. p. 200.**distortus.*

1. *S. glaber; foliis amplexicaulibus, pedicellis solitariis medio distorto-geniculatis.* *Mich. fl. amer. 1. p. 200.*  
*Uvularia amplexifolia. Willd. sp. pl. 2. p. 93.*

*Icon. Pl. rar. hung. t. 167.*

In shady woods: Canada to Pennsylvania on high mountains. *4. May, June. v. v.* Flowers greenish-yellow, the size of *Convallaria majalis.*

*roseus.*

2. *S. glaber lucidus; foliis amplexicaulibus serrulato-ciliatis, antheris brevibus bicornibus.* *Mich. fl. amer. 1. p. 201.*

*Icon. Mich. l. c. t. 18.*

On the high mountains of Carolina, Pennsylvania, and in Canada. *4. May—July. v. v.* Flowers rose-coloured.

*lanuginosus.*

3. *S. subcandicanti-lanuginosus; foliis sessilibus basi subcordatis acuminatis, pedicellis in brevissimo stipite geminatis.* *Mich. fl. amer. 1. p. 201.*

On the high mountains: Pennsylvania to Carolina. *4. June. v. v.* Flowers large, yellowish-green; berries red.

304. CONVALLARIA. *Gen. pl. 575. Desfont. in annal. mus. hist. nat. 9. p. 46.**majalis.*

1. *C. scapo nudo lævi, foliis ovatis.* *Willd. sp. pl. 2. p. 160.*

*Icon. Fl. dan. 854.*

On the highest mountains of Virginia and Carolina. *4. May. v. v.* Flowers white, fragrant.

305. SMILACINA. *Desfont. in annual. mus. hist. nat. 9. p. 51.**umbellata.*

1. *S. foliis radicalibus oblongo-ovalibus margine et carina ciliatis, scapo pubescente, umbella terminali, pedicellis bracteatis.* — *Desfont. l. c. 53.*

HEXANDRIA

Convallaria um  
S. borealis. Ke  
Icon. Desfont.  
On the Allega  
rare. 4. 1

2. S. subcaulesce  
tis, scapo p  
prolifera, pe  
Dracæna bore  
Icon. Ait. keu  
On the Allega  
New Engla  
greenish-ye

3. S. caule bif  
utrinque gl  
ribus tetra  
Convallaria bi  
Majanthemur

β. S. foliis ovatis  
γ. S. caule trifol  
Icon. Pluk. a  
In shady woo  
bogs on hig  
small, whit

4. S. caule alte  
lanceolatis,  
Convallaria t  
Icon. Gmel.  
On the high  
places: Pe  
June, July

5. S. caule alte  
seu ovali  
Desfont. l.  
Convallaria s  
Icon. Corn.  
Bot. mag.  
In Canada,  
banks as f  
About a f

6. S. caule alt  
ciliatis, p  
p. 53.  
Icon. Desfo

- ovalibus  
æ lacinii  
amer. 1.  
+. Re-  
r.  
200.  
solitariis  
p. 200,  
  
gh moun-  
enish-yel-  
  
lato-cilio-  
l. amer. 1.  
  
nia, and in  
's rose-co-  
  
basi sub-  
mo stipite  
  
olina. ¼.  
; berries  
  
t. in annal.  
  
sp. pl. 2.  
  
arolina. ¼.  
  
hist. nat. 9.  
  
ine et carina-  
ali, pedice-  
-
- Convallaria umbellata. *Mich. fl. amer.* 1. p. 202.  
*S. borealis.* *Ker in bot. mag.*  
*Icon. Desfont. l. c. t. 8.* *Bot. mag.* 1155.  
 On the Allegany mountains: Pensylvania to Carolina;  
 rare. ¼. May, June. v. v. Flowers white, fragrant.
2. *S. subcaulescens*; foliis elliptico-obovatis margine cilia-  
 tis, scapo pubescente, umbella corymbosa interdum  
 prolifera, pedicellis nudis nutantibus.—*Bot. mag.* 1403.  
*Dracæna borealis.* *Willd. sp. pl.* 2. p. 158.  
*Icon. Ait. kew.* 1. t. 5. *Bot. mag.* 1403. *Bot. repos.* 206.  
 On the Allegany mountains, in bogs, and in Canada and  
 New England. ¼. June, July. v. v. Flowers large,  
 greenish-yellow: fruit beautiful amethystine blue.
3. *S. caule bifolio*, foliis cordato-oblongis subsessilibus  
 utrinque glaberrimis, racemo simplici terminali, flo-  
 ribus tetrandris.  
*Convallaria bifolia.* *Mich. fl. amer.* 1. p. 201.  
*Majanthemum canadense.* *Desfont. l. c. p.* 54.  
*S. foliis ovatis basi vix cordatis.*  
*S. caule trifoliato*, foliis lato-cordatis.  
*Icon. Pluk. alm.* t. 434. f. 4.  
 In shady woods, round the roots of old trees and in the  
 bogs on high mountains. ¼. May. v. v. Flowers  
 small, white.
4. *S. caule alterne trifoliato pubescente*, foliis oblongo-  
 lanceolatis, racemo terminali laxo.—*Desfont. l. c. p.* 52.  
*Convallaria trifolia.* *Willd. sp. pl.* 2. p. 163.  
*Icon. Gmel. fl. sibir.* 1. t. 6.  
 On the highest Allegany mountains, in deep sphagnumous  
 places: Pensylvania, New York, and in Canada. ¼.  
 June, July. v. v. Flowers small, white.
5. *S. caule alterne folioso*, foliis amplexicaulibus lanceolatis  
 seu ovali-lanceolatis, racemo simplici terminali.  
*Desfont. l. c. p.* 52.  
*Convallaria stellata.* *Willd. sp. pl.* 2. p. 163.  
*Icon. Corn. canad.* t. 33. *Moris. hist.* 3. s. 15. t. 4. f. 7.  
*Bot. mag.* 1043.  
 In Canada, and along the tract of mountains on river  
 banks as far south as Virginia. ¼. May, June. v. v.  
 About a foot high; flowers in a close spike, white.
6. *S. caule alterne folioso arcuato*, foliis sessilibus ovatis  
 ciliatis, panicula terminali conferta.—*Desfont. l. c.*  
*p.* 53.  
*Icon. Desfont. l. c. t. 9.*

*borealis.**canadensis.**ovalis.*  
*trifolia.**trifolia.**stellata.**ciliata.*

- racemosa.* In Virginia and Canada. *Desfontaines.* 4. +. Flowers very small, white.
- angustifolium.* 7. S. caule alterne folioso, foliis sessilibus oblongo-ovalibus acuminatis nervosis pubescentibus, floribus terminalibus racemoso-paniculatis.—*Desfont. l. c. p. 51.*  
Icon. *Corn. canad. t. 37.* *Ptuk. alm. t. 311. f. 2. Bot. mag. 899.*  
In mountain woods: Canada to Carolina; common.  
4. June, July. v. v. Flowers small, yellowish.
- canaliculatum.* 306. POLYGONATUM. *Desfont. in mus. hist. nat. 9.*  
*p. 48.*
- pubescens.* 1. P. caule tereti, foliis alternis sessilibus elliptico-lanceolatis sensim acutis subtrinervibus glabris, pedunculis axillaribus elongatis subbifloris.  
Convallaria biflora. *Walt. fl. car. 122.*  
In rocky situations: Pennsylvania to Virginia. 4. May, June. v. v. Flowers yellowish-white, with green tips, as most of this genus.
- hirtum.* 2. P. caule canaliculato, foliis alternis amplexicaulibus oblongis margine pubescentibus, pedunculis axillaribus bifloris.  
Convallaria canaliculata. *Willd. hort. berol. 45.*  
In shady woods: Pennsylvania to Virginia. 4. June.  
v. v.
- multiflorum.* 3. P. caule teretiusculo sulco exarato, foliis alternis amplexicaulibus ovatis subtus pubescentibus, pedunculis axillaribus subbifloris.  
Convallaria pubescens. *Willd. hort. berol. 45. cum icon.*  
On rocks, near water: New England to Carolina. 4.  
May, June. v. v.
4. P. caule angulato hispido, foliis alternis subamplexicaulibus ovatis obtuse acuminatis, pedunculis axillaribus trifloris.  
Convallaria hirta. *Lam. encycl. 4. p. 369.*  
In North America. *Bosc.* 4. +.
5. P. caule tereti, foliis alternis amplexicaulibus oblongo-ovalibus, pedunculis axillaribus multifloris.—*Desfont. l. c. p. 50.*  
Convallaria multiflora. *Willd. sp. pl. 2. p. 162.*  
Icon. *Fl. dan. 152.* *Abbot 77. Engl. bot. 279.*  
In dry hilly woods: Canada to Carolina. 4. June, July.  
v. v.

ALEXANDR

6. P. caule angula-dunculis uni-Convallaria latifl. Icon. *Jacq. au* On the sandy a situations: 1  
v. v. The feet high.

307. A

1. A. caule herba-mollibus, sti-  
*Willd. sp. 1*  
Icon. *Fl. dan.*  
Common *Aspi* wild state,

308. C

1. O. foliis lanc. *Willd. sp.*  
Icon. *Amaen.*  
In rivulets ar  
4. April,  
cylindric s  
a variety v  
near New

30

1. A. scapo mu-  
*p. 199.*  
Icon. *Abbot*  
In overflow  
to Caroli  
Sweet-fla  
roots.

3

1. J. culmo n  
phylo sp  
fl. brit  
J. maritim

- + Flowers  
ongo-ovalibus  
us terminali-  
p. 51.  
1. f. 2. Bot.  
ia; common.  
yellowish.  
  
. hist. nat. 9.  
liptico-lanceo-  
is, pedunculis  
  
nia. 4. May,  
e, with green  
exicaulibus ob-  
ulis axillaribus  
  
L. 45.  
ia. 4. June,  
is alternis am-  
bus, pedunculis  
  
L. 45. cum icono.  
Carolina. 4.  
subamplexicau-  
culis axillaribus  
).  
  
alibus oblongo-  
floris.—Desfont.  
p. 162.  
t. 279.  
4. June, July.
6. P. caule angulato, foliis sessilibus ovatis acuminatis, pe-  
dunculis uni- aut multifloris. Desfont. l. c. p. 50. *latifolium.*  
Convallaria latifolia. Willd. sp. pl. 2. p. 162.  
Icon. Jacq. austr. 3. t. 232.  
On the sandy and gravelly banks of rivers, in shady wet  
situations: Pennsylvania and Virginia. 4. June, July.  
v. v. The tallest of all the species, sometimes four  
feet high.

## 307. ASPARAGUS. Gen. pl. 573.

1. A. caule herbaceo inermi suberecto tereti, foliis setaceis  
mollibus, stipulis subsolitariis. Smith fl. brit. 1. p. 369.  
Willd. sp. pl. 2. p. 150.  
Icon. Fl. dan. 803. Engl. bot. 339.  
Common Asparagus is frequently found in an apparently  
wild state, but has been introduced from Europe.

## 308. ORONTIUM. Gen. pl. 587.

1. O. foliis lanceolato-ovatis, scapo cylindrico spicato.— *aquaticum.*  
Willd. sp. pl. 2. p. 199.  
Icon. Amoen. acad. 3. t. 1. f. 3.  
In rivulets and low stagnant waters: Canada to Florida.  
4. April, May. v. v. Yellow flowers in a close  
cylindric spike, of a most singular smell. I observed  
a variety with almost linear leaves, in the Salt marshes  
near New York.

## 309. ACORUS. Gen. pl. 586.

1. A. scapo mucrone longissimo foliaceo. Willd. sp. pl. 2. *Calamus.*  
p. 199.  
Icon. Abbot 77. Engl. bot. 356.  
In overflowed places and old ditches; frequent: Canada  
to Carolina. Has the appearance of a flag, and called  
Sweet-flag, from the aromatic smell and taste of its  
roots.

## 310. JUNCUS. Gen. pl. 590.

\* *Culmis nudis.*

1. J. culmo nudo tereti, panicula terminali, involuero di-  
phylo spinoso, capsulis subrotundis mucronatis. Smith  
fl. brit. 1. p. 374. Willd. sp. pl. 2. p. 204.  
J. maritimus. Lam. encycl. 3. p. 253.  
  
acutus.

## HEXANDRIA MONOGYNIA. Juncus.

- conglomeratus.* Icon. *Engl. bot.* 1614.  
On the sandy sea-coast : New Jersey, &c. 24. July. v. v.
- effusus.* 2. J. culmo nudo stricto, panicula laterali conglobata, capsulis retusis, floribus triandris. *Smith fl. brit.* 1. p. 376. *Willd. sp. pl.* 2. p. 205.  
Icon. *Host. gram.* 3. t. 82. *Leers fl. herb.* t. 13. f. 1.  
In wet pastures, frequent. 24. June, July. v. v.
- filiformis.* 3. J. culmo nudo stricto, panicula laterali effusa supradecomposita, capsulis obtusis. *Smith fl. brit.* 1. p. 376. *Willd. sp. pl.* 2. p. 205.  
Icon. *Host. gram.* 3. t. 83.  
Frequent in low grounds, on road sides, &c. 24. June, July. v. v.
- bicornis.* 4. J. culmo nudo filiformi nutante, panicula laterali bracteata subsimplici, capsulis subrotundis. *Smith fl. brit.* 1. p. 377. *Willd. sp. pl.* 2. p. 207.  
Icon. *Host. gram.* 3. t. 84. *Pluk. phyt.* t. 40. f. 8.  
In boggy mountain meadows ; frequent. 24. July, Aug. v. v.
- setaceus.* 5. J. culmo erecto nudo, foliis subsetaceis canaliculato-planis, involucro setaceo diphylo erecto paniculam superante, floribus distinctis, calyce linear-lanceolato acutissimo.  
—*Mich. fl. amer.* 1. p. 191.  
J. tenuis. *Rostk. junc.* 21. t. 1. f. 3. *Moris. hist.* 3. s. 8. t. 9. f. 15.  
In sandy wet fields, near ponds : Carolina and Georgia. 24. July. v. v.
- marginatus.* 6. J. culmo nudo filiformi nutante, umbella laterali composita pauciflora, pedunculis multifloris, calyce subulato. *Rostk. junc.* 13. t. 1. f. 2.  
In low grounds : Canada to Virginia. 24. July. v. v.
- nodosus.* 7. J. foliis planis glabris, corymbo terminali simplici proliferis, capitulis sub-10-floris, calyce capsulam obtusam æquante. *Rostk. junc.* 30. t. 2. f. 3.  
On dry shady hills : Pennsylvania. 24. June. v. v.
8. J. foliis nodoso-articulatis teretibus, corymbo simplici, capitulis globosis, capsulis acuminatis calyce longioribus. *Rostk. junc.* t. 2. f. 2.  
J. scirpoidea. *Lam. encycl.* 3. p. 267.  
Icon. *Pluk. alm.* t. 417. f. 3.  
In inundated fields and ditches : Canada to Virginia. 24. July, Aug. v. v.
9. J. culmo eretum capitulis glabris linearibus tenuibus. *Icon. Pluk.* t. 417. f. 3.  
J. major, foliis subfusiformibus. In inundated fields : July. v. v.
10. J. culmo eretum culis supra rioribus longis. *J. acutiflorus.* *Icon. Host.* t. 417. f. 3.  
In inundate fields : Virginia.
11. J. culmo eretum ratis verticillatis. *Icon. Pluk.* t. 212. f. 3.  
*J. uliginosus.* *J. fluitans.* *Icon. Fl. daurica.* In rivulets : July, Aug.
12. J. culmo foecundum culicula terminali dunculatis 3-and 5-and 7-and 9-and 11-and 13-and 15-and 17-and 19-and 21-and 23-and 25-and 27-and 29-and 31-and 33-and 35-and 37-and 39-and 41-and 43-and 45-and 47-and 49-and 51-and 53-and 55-and 57-and 59-and 61-and 63-and 65-and 67-and 69-and 71-and 73-and 75-and 77-and 79-and 81-and 83-and 85-and 87-and 89-and 91-and 93-and 95-and 97-and 99-and 101-and 103-and 105-and 107-and 109-and 111-and 113-and 115-and 117-and 119-and 121-and 123-and 125-and 127-and 129-and 131-and 133-and 135-and 137-and 139-and 141-and 143-and 145-and 147-and 149-and 151-and 153-and 155-and 157-and 159-and 161-and 163-and 165-and 167-and 169-and 171-and 173-and 175-and 177-and 179-and 181-and 183-and 185-and 187-and 189-and 191-and 193-and 195-and 197-and 199-and 201-and 203-and 205-and 207-and 209-and 211-and 213-and 215-and 217-and 219-and 221-and 223-and 225-and 227-and 229-and 231-and 233-and 235-and 237-and 239-and 241-and 243-and 245-and 247-and 249-and 251-and 253-and 255-and 257-and 259-and 261-and 263-and 265-and 267-and 269-and 271-and 273-and 275-and 277-and 279-and 281-and 283-and 285-and 287-and 289-and 291-and 293-and 295-and 297-and 299-and 301-and 303-and 305-and 307-and 309-and 311-and 313-and 315-and 317-and 319-and 321-and 323-and 325-and 327-and 329-and 331-and 333-and 335-and 337-and 339-and 341-and 343-and 345-and 347-and 349-and 351-and 353-and 355-and 357-and 359-and 361-and 363-and 365-and 367-and 369-and 371-and 373-and 375-and 377-and 379-and 381-and 383-and 385-and 387-and 389-and 391-and 393-and 395-and 397-and 399-and 401-and 403-and 405-and 407-and 409-and 411-and 413-and 415-and 417-and 419-and 421-and 423-and 425-and 427-and 429-and 431-and 433-and 435-and 437-and 439-and 441-and 443-and 445-and 447-and 449-and 451-and 453-and 455-and 457-and 459-and 461-and 463-and 465-and 467-and 469-and 471-and 473-and 475-and 477-and 479-and 481-and 483-and 485-and 487-and 489-and 491-and 493-and 495-and 497-and 499-and 501-and 503-and 505-and 507-and 509-and 511-and 513-and 515-and 517-and 519-and 521-and 523-and 525-and 527-and 529-and 531-and 533-and 535-and 537-and 539-and 541-and 543-and 545-and 547-and 549-and 551-and 553-and 555-and 557-and 559-and 561-and 563-and 565-and 567-and 569-and 571-and 573-and 575-and 577-and 579-and 581-and 583-and 585-and 587-and 589-and 591-and 593-and 595-and 597-and 599-and 601-and 603-and 605-and 607-and 609-and 611-and 613-and 615-and 617-and 619-and 621-and 623-and 625-and 627-and 629-and 631-and 633-and 635-and 637-and 639-and 641-and 643-and 645-and 647-and 649-and 651-and 653-and 655-and 657-and 659-and 661-and 663-and 665-and 667-and 669-and 671-and 673-and 675-and 677-and 679-and 681-and 683-and 685-and 687-and 689-and 691-and 693-and 695-and 697-and 699-and 701-and 703-and 705-and 707-and 709-and 711-and 713-and 715-and 717-and 719-and 721-and 723-and 725-and 727-and 729-and 731-and 733-and 735-and 737-and 739-and 741-and 743-and 745-and 747-and 749-and 751-and 753-and 755-and 757-and 759-and 761-and 763-and 765-and 767-and 769-and 771-and 773-and 775-and 777-and 779-and 781-and 783-and 785-and 787-and 789-and 791-and 793-and 795-and 797-and 799-and 801-and 803-and 805-and 807-and 809-and 811-and 813-and 815-and 817-and 819-and 821-and 823-and 825-and 827-and 829-and 831-and 833-and 835-and 837-and 839-and 841-and 843-and 845-and 847-and 849-and 851-and 853-and 855-and 857-and 859-and 861-and 863-and 865-and 867-and 869-and 871-and 873-and 875-and 877-and 879-and 881-and 883-and 885-and 887-and 889-and 891-and 893-and 895-and 897-and 899-and 901-and 903-and 905-and 907-and 909-and 911-and 913-and 915-and 917-and 919-and 921-and 923-and 925-and 927-and 929-and 931-and 933-and 935-and 937-and 939-and 941-and 943-and 945-and 947-and 949-and 951-and 953-and 955-and 957-and 959-and 961-and 963-and 965-and 967-and 969-and 971-and 973-and 975-and 977-and 979-and 981-and 983-and 985-and 987-and 989-and 991-and 993-and 995-and 997-and 999-and 1001-and 1003-and 1005-and 1007-and 1009-and 1011-and 1013-and 1015-and 1017-and 1019-and 1021-and 1023-and 1025-and 1027-and 1029-and 1031-and 1033-and 1035-and 1037-and 1039-and 1041-and 1043-and 1045-and 1047-and 1049-and 1051-and 1053-and 1055-and 1057-and 1059-and 1061-and 1063-and 1065-and 1067-and 1069-and 1071-and 1073-and 1075-and 1077-and 1079-and 1081-and 1083-and 1085-and 1087-and 1089-and 1091-and 1093-and 1095-and 1097-and 1099-and 1101-and 1103-and 1105-and 1107-and 1109-and 1111-and 1113-and 1115-and 1117-and 1119-and 1121-and 1123-and 1125-and 1127-and 1129-and 1131-and 1133-and 1135-and 1137-and 1139-and 1141-and 1143-and 1145-and 1147-and 1149-and 1151-and 1153-and 1155-and 1157-and 1159-and 1161-and 1163-and 1165-and 1167-and 1169-and 1171-and 1173-and 1175-and 1177-and 1179-and 1181-and 1183-and 1185-and 1187-and 1189-and 1191-and 1193-and 1195-and 1197-and 1199-and 1201-and 1203-and 1205-and 1207-and 1209-and 1211-and 1213-and 1215-and 1217-and 1219-and 1221-and 1223-and 1225-and 1227-and 1229-and 1231-and 1233-and 1235-and 1237-and 1239-and 1241-and 1243-and 1245-and 1247-and 1249-and 1251-and 1253-and 1255-and 1257-and 1259-and 1261-and 1263-and 1265-and 1267-and 1269-and 1271-and 1273-and 1275-and 1277-and 1279-and 1281-and 1283-and 1285-and 1287-and 1289-and 1291-and 1293-and 1295-and 1297-and 1299-and 1301-and 1303-and 1305-and 1307-and 1309-and 1311-and 1313-and 1315-and 1317-and 1319-and 1321-and 1323-and 1325-and 1327-and 1329-and 1331-and 1333-and 1335-and 1337-and 1339-and 1341-and 1343-and 1345-and 1347-and 1349-and 1351-and 1353-and 1355-and 1357-and 1359-and 1361-and 1363-and 1365-and 1367-and 1369-and 1371-and 1373-and 1375-and 1377-and 1379-and 1381-and 1383-and 1385-and 1387-and 1389-and 1391-and 1393-and 1395-and 1397-and 1399-and 1401-and 1403-and 1405-and 1407-and 1409-and 1411-and 1413-and 1415-and 1417-and 1419-and 1421-and 1423-and 1425-and 1427-and 1429-and 1431-and 1433-and 1435-and 1437-and 1439-and 1441-and 1443-and 1445-and 1447-and 1449-and 1451-and 1453-and 1455-and 1457-and 1459-and 1461-and 1463-and 1465-and 1467-and 1469-and 1471-and 1473-and 1475-and 1477-and 1479-and 1481-and 1483-and 1485-and 1487-and 1489-and 1491-and 1493-and 1495-and 1497-and 1499-and 1501-and 1503-and 1505-and 1507-and 1509-and 1511-and 1513-and 1515-and 1517-and 1519-and 1521-and 1523-and 1525-and 1527-and 1529-and 1531-and 1533-and 1535-and 1537-and 1539-and 1541-and 1543-and 1545-and 1547-and 1549-and 1551-and 1553-and 1555-and 1557-and 1559-and 1561-and 1563-and 1565-and 1567-and 1569-and 1571-and 1573-and 1575-and 1577-and 1579-and 1581-and 1583-and 1585-and 1587-and 1589-and 1591-and 1593-and 1595-and 1597-and 1599-and 1601-and 1603-and 1605-and 1607-and 1609-and 1611-and 1613-and 1615-and 1617-and 1619-and 1621-and 1623-and 1625-and 1627-and 1629-and 1631-and 1633-and 1635-and 1637-and 1639-and 1641-and 1643-and 1645-and 1647-and 1649-and 1651-and 1653-and 1655-and 1657-and 1659-and 1661-and 1663-and 1665-and 1667-and 1669-and 1671-and 1673-and 1675-and 1677-and 1679-and 1681-and 1683-and 1685-and 1687-and 1689-and 1691-and 1693-and 1695-and 1697-and 1699-and 1701-and 1703-and 1705-and 1707-and 1709-and 1711-and 1713-and 1715-and 1717-and 1719-and 1721-and 1723-and 1725-and 1727-and 1729-and 1731-and 1733-and 1735-and 1737-and 1739-and 1741-and 1743-and 1745-and 1747-and 1749-and 1751-and 1753-and 1755-and 1757-and 1759-and 1761-and 1763-and 1765-and 1767-and 1769-and 1771-and 1773-and 1775-and 1777-and 1779-and 1781-and 1783-and 1785-and 1787-and 1789-and 1791-and 1793-and 1795-and 1797-and 1799-and 1801-and 1803-and 1805-and 1807-and 1809-and 1811-and 1813-and 1815-and 1817-and 1819-and 1821-and 1823-and 1825-and 1827-and 1829-and 1831-and 1833-and 1835-and 1837-and 1839-and 1841-and 1843-and 1845-and 1847-and 1849-and 1851-and 1853-and 1855-and 1857-and 1859-and 1861-and 1863-and 1865-and 1867-and 1869-and 1871-and 1873-and 1875-and 1877-and 1879-and 1881-and 1883-and 1885-and 1887-and 1889-and 1891-and 1893-and 1895-and 1897-and 1899-and 1901-and 1903-and 1905-and 1907-and 1909-and 1911-and 1913-and 1915-and 1917-and 1919-and 1921-and 1923-and 1925-and 1927-and 1929-and 1931-and 1933-and 1935-and 1937-and 1939-and 1941-and 1943-and 1945-and 1947-and 1949-and 1951-and 1953-and 1955-and 1957-and 1959-and 1961-and 1963-and 1965-and 1967-and 1969-and 1971-and 1973-and 1975-and 1977-and 1979-and 1981-and 1983-and 1985-and 1987-and 1989-and 1991-and 1993-and 1995-and 1997-and 1999-and 2001-and 2003-and 2005-and 2007-and 2009-and 2011-and 2013-and 2015-and 2017-and 2019-and 2021-and 2023-and 2025-and 2027-and 2029-and 2031-and 2033-and 2035-and 2037-and 2039-and 2041-and 2043-and 2045-and 2047-and 2049-and 2051-and 2053-and 2055-and 2057-and 2059-and 2061-and 2063-and 2065-and 2067-and 2069-and 2071-and 2073-and 2075-and 2077-and 2079-and 2081-and 2083-and 2085-and 2087-and 2089-and 2091-and 2093-and 2095-and 2097-and 2099-and 2101-and 2103-and 2105-and 2107-and 2109-and 2111-and 2113-and 2115-and 2117-and 2119-and 2121-and 2123-and 2125-and 2127-and 2129-and 2131-and 2133-and 2135-and 2137-and 2139-and 2141-and 2143-and 2145-and 2147-and 2149-and 2151-and 2153-and 2155-and 2157-and 2159-and 2161-and 2163-and 2165-and 2167-and 2169-and 2171-and 2173-and 2175-and 2177-and 2179-and 2181-and 2183-and 2185-and 2187-and 2189-and 2191-and 2193-and 2195-and 2197-and 2199-and 2201-and 2203-and 2205-and 2207-and 2209-and 2211-and 2213-and 2215-and 2217-and 2219-and 2221-and 2223-and 2225-and 2227-and 2229-and 2231-and 2233-and 2235-and 2237-and 2239-and 2241-and 2243-and 2245-and 2247-and 2249-and 2251-and 2253-and 2255-and 2257-and 2259-and 2261-and 2263-and 2265-and 2267-and 2269-and 2271-and 2273-and 2275-and 2277-and 2279-and 2281-and 2283-and 2285-and 2287-and 2289-and 2291-and 2293-and 2295-and 2297-and 2299-and 2301-and 2303-and 2305-and 2307-and 2309-and 2311-and 2313-and 2315-and 2317-and 2319-and 2321-and 2323-and 2325-and 2327-and 2329-and 2331-and 2333-and 2335-and 2337-and 2339-and 2341-and 2343-and 2345-and 2347-and 2349-and 2351-and 2353-and 2355-and 2357-and 2359-and 2361-and 2363-and 2365-and 2367-and 2369-and 2371-and 2373-and 2375-and 2377-and 2379-and 2381-and 2383-and 2385-and 2387-and 2389-and 2391-and 2393-and 2395-and 2397-and 2399-and 2401-and 2403-and 2405-and 2407-and 2409-and 2411-and 2413-and 2415-and 2417-and 2419-and 2421-and 2423-and 2425-and 2427-and 2429-and 2431-and 2433-and 2435-and 2437-and 2439-and 2441-and 2443-and 2445-and 2447-and 2449-and 2451-and 2453-and 2455-and 2457-and 2459-and 2461-and 2463-and 2465-and 2467-and 2469-and 2471-and 2473-and 2475-and 2477-and 2479-and 2481-and 2483-and 2485-and 2487-and 2489-and 2491-and 2493-and 2495-and 2497-and 2499-and 2501-and 2503-and 2505-and 2507-and 2509-and 2511-and 2513-and 2515-and 2517-and 2519-and 2521-and 2523-and 2525-and 2527-and 2529-and 2531-and 2533-and 2535-and 2537-and 2539-and 2541-and 2543-and 2545-and 2547-and 2549-and 2551-and 2553-and 2555-and 2557-and 2559-and 2561-and 2563-and 2565-and 2567-and 2569-and 2571-and 2573-and 2575-and 2577-and 2579-and 2581-and 2583-and 2585-and 2587-and 2589-and 2591-and 2593-and 2595-and 2597-and 2599-and 2601-and 2603-and 2605-and 2607-and 2609-and 2611-and 2613-and 2615-and 2617-and 2619-and 2621-and 2623-and 2625-and 2627-and 2629-and 2631-and 2633-and 2635-and 2637-and 2639-and 2641-and 2643-and 2645-and 2647-and 2649-and 2651-and 2653-and 2655-and 2657-and 2659-and 2661-and 2663-and 2665-and 2667-and 2669-and 2671-and 2673-and 2675-and 2677-and 2679-and 2681-and 2683-and 2685-and 2687-and 2689-and 2691-and 2693-and 2695-and 2697-and 2699-and 2701-and 2703-and 2705-and 2707-and 2709-and 2711-and 2713-and 2715-and 2717-and 2719-and 2721-and 2723-and 2725-and 2727-and 2729-and 2731-and 2733-and 2735-and 2737-and 2739-and 2741-and 2743-and 2745-and 2747-and 2749-and 2751-and 2753-and 2755-and 2757-and 2759-and 2761-and 2763-and 2765-and 2767-and 2769-and 2771-and 2773-and 2775-and 2777-and 2779-and 2781-and 2783-and 2785-and 2787-and 2789-and 2791-and 2793-and 2795-and 2797-and 2799-and 2801-and 2803-and 2805-and 2807-and 2809-and 2811-and 2813-and 2815-and 2817-and 2819-and 2821-and 2823-and 2825-and 2827-and 2829-and 2831-and 2833-and 2835-and 2837-and 2839-and 2841-and 2843-and 2845-and 2847-and 2849-and 2851-and 2853-and 2855-and 2857-and 2859-and 2861-and 2863-and 2865-and 2867-and 2869-and 2871-and 2873-and 2875-and 2877-and 2879-and 2881-and 2883-and 2885-and 2887-and 2889-and 2891-and 2893-and 2895-and 2897-and 2899-and 2901-and 2903-and 2905-and 2907-and 2909-and 2911-and 2913-and 2915-and 2917-and 2919-and 2921-and 2923-and 2925-and 2927-and 2929-and 2931-and 2933-and 2935-and 2937-and 2939-and 2941-and 2943-and 2945-and 2947-and 2949-and 2951-and 2953-and 2955-and 2957-and 2959-and 2961-and 2963-and 2965-and 2967-and 2969-and 2971-and 2973-and 2975-and 2977-and 2979-and 2981-and 2983-and 2985-and 2987-and 2989-and 2991-and 2993-and 2995-and 2997-and 2999-and 3001-and 3003-and 3005-and 3007-and 3009-and 3011-and 3013-and 3015-and 3017-and 3019-and 3021-and 3023-and 3025-and 3027-and 3029-and 3031-and 3033-and 3035-and 3037-and 3039-and 3041-and 3043-and 3045-and 3047-and 3049-and 3051-and 3053-and 3055-and 3057-and 3059-and 3061-and 3063-and 3065-and 3067-and 3069-and 3071-and 3073-and 3075-and 3077-and 3079-and 3081-and 3083-and 3085-and 3087-and 3089-and 3091-and 3093-and 3095-and 3097-and 3099-and 3101-and 3103-and 3105-and 3107-and 3109-and 3111-and 3113-and 3115-and 3117-and 3119-and 3121-and 3123-and 3125-and 3127-and 3129-and 3131-and 3133-and 3135-and 3137-and 3139-and 3141-and 3143-and 3145-and 3147-and 3149-and 3151-and 3153-and 3155-and 3157-and 3159-and 3161-and 3163-and 3165-and 3167-and 3169-and 3171-and 3173-and 3175-and 3177-and 3179-and 3181-and 3183-and 3185-and 3187-and 3189-and 3191-and 3193-and 3195-and 3197-and 3199-and 3201-and 3203-and 3205-and 3207-and 3209-and 3211-and 3213-and 3215-and 3217-and 3219-and 3221-and 3223-and 3225-and 3227-and 3229-and 3231-and 3233-and 3235-and 3237-and 3239-and 3241-and 3243-and 3245-and 3247-and 3249-and 3251-and 3253-and 3255-and 3257-and 3259-and 3261-and 3263-and 3265-and 3267-and 3269-and 3271-and 3273-and 3275-and 3277-and 3279-and 3281-and 3283-and 3285-and 3287-and 3289-and 3291-and 3293-and 3295-and 3297-and 3299-and 3301-and 3303-and 3305-and 3307-and 3309-and 3311-and 3313-and 3315-and 3317-and 3319-and 3321-and 3323-and 3325-and 3327-and 3329-and 3331-and 3333-and 3335-and 3337-and 3339-and 3341-and 3343-and 3345-and 3347-and 3349-and 3351-and 3353-and 3355-and 3357-and 3359-and 3361-and 3363-and 3365-and 3367-and 3369-and 3371-and 3373-and 3375-and 3377-and 3379-and 3381-and 3383-and 3385-and 3387-and 3389-and 3391-and 3393-and 3395-and 3397-and 3399-and 3401-and 3403-and 3405-and 3407-and 3409-and 3411-and 3413-and 3415-and 3417-and 3419-and 3421-and 3423-and 3425-and 3427-and 3429-and 3431-and 3433-and 3435-and 3437-and 3439-and 3441-and 3443-and 3445-and 3447-and 3449-and 3451-and 3453-and 3455-and 3457-and 3459-and 3461-and 3463-and 3465-and 3467-and 3469-and 3471-and 3473-and 3475-and 3477-and 3479-and 3481-and 3483-and 3485-and 3487-and 3489-and 3491-and 3493-and 3495-and 3497-and 3499-and 3501-and 3503-and 3505-and 3507-and 3509-and 3511-and 3513-and 3515-and 3517-and 3519-and 3521-and 3523-and 3525-and 3527-and 3529-and 3531-and 3533-and 3535-and 3537-and 3539-and 3541-and 3543-and 3545-and 3547-and 3549-and 3551-and 3553-and 3555-and 3557-and 3559-and 3561-and 3563-and 3565-and 3567-and 3569-and 3571-and 3573-and 3575-and 3577-and 3579-and 3581-and 3583-and 3585-and 3587-and 3589-and 3591-and 3593-and 3595-and 3597-and 3599-and 3601-and 3

\*\* *Culnis foliosis.*

4. July. v. v.  
anglobata, cap-  
h fl. brit. 1.  
  
t. 13. f. 1.  
v. v. v.  
ffusa suprade-  
rit. 1. p. 376.  
  
c. 4. June,  
laterali brac-  
mth fl. brit. 1.  
  
. 40. f. 8.  
4. July, Aug.  
  
liculato-planis,  
lam superante,  
ato acutissimo.  
  
is. hist. 3. s. 8.  
and Georgia.  
  
laterali com-  
, calyce subu-  
  
. July. v. v.  
li simplici pro-  
capsulam obtu-  
3.  
ine. v. v.  
ymbo simplici,  
alyce longiori-  
  
da to Virginia.
9. J. culmo erecto oligophyllo, foliis nodoso-articulatis, *polycephalus*.  
capitulis globosis multifloris subpaniculatis, calycibus  
linearibus triandris.—*Mich. fl. amer. 1. p. 192.*  
Icon. *Plak. alm. t. 92. f. 9.*
- z. J. major, foliis crassioribus compressis. *crassifolius.*  
β. J. foliis subfiliformibus. *tenuifolius.*
- In inundated places : Pennsylvania to Carolina. 4. June,  
July. v. v.
10. J. culmo erecto, follis nodoso-articulatis teretibus, pani-  
cula supradecomposita, foliolis calycinis aristatis inte-  
rioribus longioribus. *Willd. sp. pl. 2. p. 211.*  
J. acutiflorus. *Hoffm. germ. 125.*  
Icon. *Host. gram. 3. t. 86.*
- In inundated woods and on river sides : Pennsylvania to  
Virginia. 4. July. v. v.
11. J. culmo decumbente, foliis setaceis, floribus glome- *verticillatus.*  
ratis verticillatis glomerulis foliosis. *Willd. sp. pl. 2.*  
p. 212.
- J. uliginosus. *Roth fl. germ. 1. p. 155.*  
J. fluitans. *Lam. encycl. 3. p. 26.*  
Icon. *Fl. dan. 817. Host. gram. 3. t. 88.*
- In rivulets and ponds : Canada, New York, &c. 4.  
July, Aug. v. v.
12. J. culmo folioso erecto, foliis subnodoso-articulatis, pa-  
nicula terminali composita, glomerulis subtrifloris pe-  
dunculatis sessilibusque, calycibus gracilibus acumina-  
tis 3-andris. *Mich. fl. amer. 1. p. 192.*  
In stagnant waters and on river banks : Pennsylvania to  
Carolina, 4. July, Aug. v. v. Resembles *J. articulatus* very much.
13. J. radice bulbosa, culmo folioso erecto compresso, foliis  
angustis subcanaliculatis, panicula composita, glome-  
rulis 3-floris, calycibus 3-andris bracteisque aristatis.  
—*Mich. fl. amer. 1. p. 192.*  
In low grounds of Georgia and Carolina. 4. July. v. s.  
Resembles the following species.
14. J. culmo subcompresso indiviso, foliis linearibus canali-  
culatis, corymbo terminali, calycibus obtusis capsula  
subrotunda obtusa brevioribus. *Willd. sp. pl. 2.*  
p. 213.
- J. compressus. *Jacq. vind. 235.*  
Icon. *Host. gram. 3. t. 89. Fl. dan. 431.* *bulbosus.*

## HEXANDRIA MONOGYNIA. Juncus.

- On dry barren soil, on road sides; frequent. ♀. June, July. v. v.
- tenuis.* 15. *J. culmo folioso simplici teretiusculo, foliis canaliculatis corymbo terminali dichotomo bracteis breviore, capsula oblonga obtusa calyce breviore.* *Rostk. junc. 24. t. 1. f. 3.* *Willd. sp. pl. 2. p. 214?*  
In Georgia and Carolina. ♀. +.
- bufonius.* 16. *J. culmo folioso dichotomo, foliis angulatis subsetaceis, floribus oblongis solitariis sessilibus.* — *Willd. sp. pl. 2. p. 214.*  
Icon. *Fl. dan. 1098.* *Host. gram. 3. t. 90.* *Engl. bot. 802.*  
In overflowed fields, common. July, Aug. v. v.
- campestris.* 17. *J. foliis planis pilosis, spicis pedunculatis ovatis subcerebruis, intermedia sessili, calycinis foliolis mucronatis capsula obtusa longioribus.* *Willd. sp. pl. 2. p. 221.*  
Icon. *Engl. bot. 672.*  
On dry sunny hills: Pennsylvania to Carolina. ♀. April, May. v. v.
- melanocarpus.* 18. *J. culmo folioso, foliis sub lanceolatis glabris, panicula capillari laxa, floribus pedicellatis.* — *Mich. fl. amer. 1. p. 190.*  
*Luzula melanocarpa.* *Desvoux in journal de botanique 1. p. 142. t. 5. f. 2.*  
In the western part of Canada. *Michaux.* +. Seeds black.
- repens.* 19. *J. culmis repentibus foliosis, foliis planis angustis, capitulis paucis sessilibus pedunculatisque paucifloris, calycibus subulatis 3-andris.* — *Mich. fl. amer. 2. p. 191.*  
*Cephaloxis flabellata.* *Desvoux in journal de botanique 1. p. 324. t. 11. f. 2.*  
On sandy hills in Carolina and Georgia. ♀. June. v. s.

## 311. PEPLIS. Gen. pl. 605.

- americana.* 1. *P. floribus axillaribus solitariis, foliis crassis spatulato-obovatis, floribus apetalis.*  
Inundated during its flowering time, in slow-flowing places of rivers, in Pennsylvania; the flowers so diminutive that to examine them it requires a strong microscope.

312.

1. *N. foliis demissis alternis orbicularibus.*  
*N. aquatica.*  
*Cabomba aquatica.*  
In the streams.  
v. s.
2. *N. foliis omnino nato-pinnatis.*  
*Floerkea prostrata.*  
In ditches and streams.  
Virginia. ♀. both species.  
character  
Aublet's;  
his plant is trifling that

313. SAEKI

1. *Sabal.* *Pechumia*  
*Raphis acaule*  
*Chamærops*  
*Corypha macrocarpa*  
*Corypha pubescens*  
Icon. *Bot. Soc. philadelph.*  
Along the sea-shore. v. v. A.

314.

1. *C. caudicea*  
*dibus pallens*  
*amer. 1.*  
On the coast.

ent. 24. June,

iis canaliculatis,  
breviore, cap.  
Rostk. junc. 24.atis subsetaceis,  
Willd. sp. pl. 2.

t. 90. Engl.

ug. v. v.

is ovatis subcer.  
iolis mucronatis  
pl. 2. p. 221.

olina. 24. April,

glabris, panicula  
Lich. fl. amer. 1.

l de botanique 1.

aux. +. Seeds

is angustis, capi.  
que paucifloris,  
L. amer. 2. p. 191.  
l de botanique 1.

24. June. v. s.

5.

rassis spatulato-

in slow-flowing  
flowers so dimi-  
nires a strong mi-

## II. DIGYNIA.

## 312. NECTRIS. Gen. pl. 610.

1. N. foliis demersis oppositis capillaceo-multifidis : emersis alternis orbiculatis peltatis integerrimis.

N. aquatica. Willd. sp. pl. 2. p. 248.

Cabomba aquatica. Aubl. Gui. 1. p. 321. t. 124.

In the streams of Carolina and Georgia. 24. July, Aug.

v. s.

2. N. foliis omnibus alternis demersis ternatis ; emersis quinato-pinnatis.

Floerkea proserpinacoides. Muhl. mss.

In ditches and slow-flowing rivers : Pennsylvania and Virginia. 24. Aug. v. v. Flowers greenish-yellow in both species, on long axillary solitary petioles. The character of the last species deviates a little from Aublet's; but as I never had an opportunity of seeing his plant in a living state, I consider the difference so trifling that they will be better to remain united.

peltata.

pinnata.

## III. TRIGYNIA.

## 313. SABAL. Adanson. Pers. syn. 1. p. 399.

1. Sabal. Pers. l. c.

Raphis acaulis. Willd. sp. pl. 4. p. 1093.

Chamærops acaulis. Mich. fl. amer. 1. p. 207.

Corypha minor. Jacq. vind. 3. t. 8.

Corypha pumila. Walt. fl. car. 119.

Icon. Bot. mag. 1434. Guersent. obs. in bullet. de la soc. philomat. no. 87.

Along the sea-coast of Carolina and Georgia. 24. Aug. v. v. A species of palm of but little beauty or use.

Adansoni.

## 314. CHAMÆROPS. Gen. pl. 1688.

1. C. caudice repente, stipitibus aculeato-serrulatis, frondibus palmatis.—Willd. sp. pl. 4. p. 1155. Mich. fl. amer. 1. p. 206.

On the coast of Georgia and Florida. 24. v. v. s. fl.

serrulata.

*Hystris.*

2. C. caudice repente, stipitibus aculeis longissimis intermixtis, frondibus palmatis.

Near the town of Savannah, Georgia. *h. v. v. s. fl.*

This singular palm grows in company with others, and distinguishes itself by its long aculei resembling porcupine quills, and often growing to the length of more than fifteen inches.

*Palmetto.*

3. C. caudice arboreo, stipitibus inermibus, spathis duplicatis, frondibus palmatis. *Willd. sp. pl. 4. p. 1155.*

*Corypha Palmetto. Walt. fl. car. 119.*

On the sea-coast of Carolina and Florida. *h. v. v. s. fl.*

A fine and useful plant; the wood used for forming almost everlasting piles for building wharfs, and the leaves can be manufactured into light and very durable hats.

315. NOLINA. *Mich. fl. amer. 1. p. 207.**georgiana.*

1. N. foliis angustissimis longis aride-gramineis, scapo laxo-ramoso foliis subulatis sparsim vestito, floribus racemosis, pedicellis aggregatis.—*Mich. fl. amer. 1. p. 208.*

*Phalangium virgatum. Lam. encycl. 5. p. 246.*

In barren sandy woods: Georgia. *Michaux and Fraser.*

*Y. July. v. s. in Herb. Enslen.* Root a tunicated bulb; stem two feet and more high; flowers white and small.

316. CALOCHORTUS. *Pursh in linn. trans. v. 11. p. . . .**elegans.*

1. C. scapo subtrifloro folio unico breviore, petalis interioribus lanatis.

On the head-waters of the Kooskoosky. *M. Lewis. Y.*

May. *v. s. in Herb. Lewis.* Flowers very elegant, white; the three inside petals are covered with a long down, and have a purple smooth spot on their base. The roots are eaten by the natives.

*Bulbus solidus, subglobosus. Foliū unicum, radicale, longe-gramineum, glabriuscum, plicato-nervosum, scapo longiore. Scapus simplex, teres, glaber, bi-aut triflorus. Bracteæ tot quod pedicelli, lineari-lanceolatae. Pedicelli filiformes, bracteis vix longiores. Flores nutantes, magnitudine Hypoxidis erecti.*

317. MELANTHIUM. *Gen. pl. 618.**virginicum.*

1. M. elatum; panicula pyramidata, petalis ovalibus subhastatis planis bimaculatis, floribus plerunque her-

maphroditic  
cycl. 4. p. 1

*Helonias virg*

In low groun

to Carolina  
ish-white a

ering for se

2. M. panicula  
sa, petalis  
culatis, st

*car. 125.*

*M. polygam*

On the mour

*v. v. Flow*

3. M. panicula  
tundis ung

*extus hirsu*

*M. latifolium*

*M. racemosu*

On the side c

*Carolina.*

*pale white*

## 318. ZIGA

1. Z. scapo foli  
minatis.

*Icon. Mich.*

In low mead

*June, Jul*

*trum albu*

2. Z. scapo sub

On the wat

mountains

*Flowers w*

*Folia radicali*

*ato-nervos*

*simplex,*

*dalis. Ra*

*Bracteæ l*

*pedicellor*

*virginici.*

*glandulis*

*breviora.*

maphroditis.—*Willd. sp. pl. 2. p. 266. Lam. encycl. 4. p. 25.*

*Helonias virginica. Bot. mag. 985.*

In low grounds, among luxuriant herbage : New York to Carolina. ♀. June, July. v. v. Flowers greenish-white at first, and turn to a dark brown after flowering for some time.

2. *M. panicula inferne mascula, superne feminea racemosa, petalis oblongis planis brevi-unguiculatis bimaculatis, stylis germine duplo brevioribus.—Walt. fl. car. 125.*

*M. polygamum. Lam. encycl. 4. p. 25.*

On the mountains of Virginia and Carolina. ♀. July. v. v. Flowers smaller, and not so apt to change colour.

3. *M. panicula superne racemosa feminea, petalis subrotundis unguiculatis plicato-undulatis vix maculatis, extus hirsutis.—Walt. fl. car. 125.*

*M. latifolium. Lam. encycl. 4. p. 25.*

*M. racemosum. Mich. fl. amer. 1. p. 251.*

On the side of hills, in moist fertile soil : Virginia and Carolina. ♀. June, July. v. v. Flowers small, pale white, and do not change.

*monoicum.*

*hybridum.*

### 318. ZIGADENUS. *Mich. fl. amer. 1. p. 213.*

1. *Z. scapo folioso, bracteis ovatis acuminatis, petalis acuminatis, glaberrimus.*

*Icon. Mich. fl. amer. 1. p. 214. t. 22.*

In low meadows : Virginia and Lower Carolina. ♀. June, July. v. v. Flowers white, the size of *Veratrum album*; leaves long, recurved and channelled.

2. *Z. scapo subnudo, bracteis linearibus, petalis acutis.*

On the waters of Cokahlaishkit river, near the Rocky-mountains. *M. Lewis. ♀. July. v. s. in Herb. Lewis.* Flowers white.

*elegans.*

*Folia radicalia, erecta, longissime-linearia, glabra, lineato-nervosa, plana, scapo breviora. Scapus teres, simplex, uno altero folio brevi instructus, subbipidalis. Racemus multiflorus, interdum basi ramosus. Bracteæ lineares, nervosæ, membranaceæ, longitudine pedicellorum. Flores albidi, magnitudine *Melanthi virginici*. Petala ovata, acuta, subunguiculata, basi glandulis 2. cinnabarinis notata. Filamenta corolla breviora. Stigmata 3. reflexa.*

319. VERATRUM. *Gen. pl. 1564.**viride.*

1. *V. racemis paniculatis, bracteis ramorum oblongo-lanceolatis, partialibus pedunculo subpubescente longioribus, foliis lato-ovatis plicatis.* — *Willd. sp. pl. 4. p. 896.*  
*V. album.* *Mich. fl. amer. 2. p. 249.*

In swamps and on mountain bogs : Canada to Carolina.  
2. July. *v. v.* A stately plant, from three to six feet high ; flowers greenish-yellow.

*parviflorum.*

2. *V. racemis paniculatis, ramis filiformibus, floribus pedicellatis stellatis, petalis ovali-lanceolatis utrinque acutis ungue staminiferis, foliis ovalibus lanceolatis planis glabris.* — *Mich. fl. amer. 2. p. 250.*

*Melanthium bracteolare.* *Lam. encycl. 4. p. 25.*

On high mountains of Carolina. 2. July. *v. v. et v. s. in Herb. Lyon.* Flowers small, green.

*angustifolium.*

3. *V. altum, dioicum; panicula simplici, petalis linearibus, foliis longissimis linearibus carinatis.*

On high mountains of Virginia and Carolina. 2. June.  
*v. v.* Flowers greenish-yellow.

320. HELONIAS. *Gen. pl. 622. Mich. fl. amer. 1. p. 211.**latifolia.*

1. *H. scapo subaphyllo, spica ovata conferta, bracteis linear-lanceolatis, foliis lanceolatis mucronatis nervosis.* — *Mich. fl. amer. 1. p. 212.*

*H. bullata.* *Willd. sp. pl. 2. p. 273.*

*Icon. Bot. mag. 747. Bot. rep. 352. Pluk. alm. t. 174. f. 5.*  
In shady swamps on sandy soil : New Jersey to Virginia.  
2. May. *v. v.* Flowers purple ; anthers blue.

*erythrosperma.*

2. *H. scapo folioso, racemo oblongo, bracteis brevibus oblongis, foliis glabris lanceolato-linearibus, seminibus ovatis legumine carnoso rubentibus.* — *Mich. fl. amer. 1. p. 212.*

*H. læta.* *Bot. mag. 803.*

*Melanthium lætum.* *Willd. sp. pl. 2. p. 267.*

*Melanthium phalangioides.* *Lam. encycl. 4. p. 28.*

*Melanthium Muscætoxicum.* *Walt. fl. car. 125.*

*Anthericum subtrigynum.* *Jacq. ic. rar. 2. t. 419.*

In shady moist situations on river banks, and on high mountains : Pennsylvania to Carolina. 2. June, July.  
*v. v.* Root a kind of bulb ; flowers first white, then changing to green ; anthers white : seeds covered with a red substance.

*angustifolia.*

3. *H. scapo inferne folioso, racemo oblongo laxo, bracteis brevibus, foliis longissimis angustissimisque, capsulis*

Jongo-lan-  
nante longio-  
l. 4. p. 896.

o Carolina,  
three to six

loribus pe-  
is utrinque  
aneolatisve

25.

y. v. v. et  
n.

s linearibus,

t. 24. June.

tr. 1. p. 211.

bracteis li-  
atis nervosis.

n. t. 174. f. 5.  
y to Virginia.  
ers blue.

brevibus ob-  
is, seminibus  
ch. fl. amer. 1.

7.

p. 28.

125.

t. 419.

and on high  
t. Jane, July.  
t white, then  
s covered with

laxo, bracteis  
isque, capsulis

*Helonias tenax.*

HE  
oblongis,  
amer. 1.  
In shady w  
v. s.

4. *H. scapo fo  
ceis, fila  
foliis sub  
Icon. Bot.  
On the san  
May, Ju  
of bulb :  
of some*

5. *H. scapo fo  
branacei  
rollam :  
simis.  
On high la  
4. Jun  
approach  
ers are t  
distincti  
not seer  
out of it  
tight b  
tuals in.*

6. *H. scapo  
brevissim  
bus, pe  
H. lutea.  
H. pumila  
Veratrum  
Melanthi  
Melanthic  
In dry sha  
4. Ju  
of Sta  
Flower  
which  
Jacquin's  
spike c  
germer  
being c  
appear  
rudime*

oblongis, seminibus angusto-linearibus.—*Mich. fl. amer. I. p. 212.*

In shady wet woods of Carolina and Georgia. 4. July.

v. s.

4. *H. scapo folioso, racemo oblongo conferto, bracteis seta-asphodeloides-*  
*ceis, filamentis basi lato-ribus corollam aequalibus,*  
*foliis subulato-setaceis.—Willd. sp. pl. 2. p. 275.*

*Icon. Bot. mag. 748. Plak. mant. t. 342. f. 3.*

On the sandy plains of New Jersey and Carolina. 4.

May, June. v. v. Flowers white, small; root a kind  
of bulb: the leaves form large tufts resembling those  
of some *Carices*.

5. *H. scapo folioso, racemo specioso laxo, bracteis triem-  
branaceis, petalis ellipticis, filamentis filiformibus co-  
rollam superantibus, foliis subulato-setaceis longis-  
simis.*

On high lands near the Rocky-mountains. *M. Lewis.*

4. June. v. s. Flowers white; anthers yellow. It  
approaches near to the foregoing species, but its flow-  
ers are three times larger, and probably an essential  
distinction may lie in the seed-vessel, which I have  
not seen. This plant is very useful to the natives:  
out of its very tenacious leaves they weave thier water-  
tight baskets, which they use for cooking their vic-  
tuals in.

6. *H. scapo folioso, racemo spicato nutante, pedicellis  
brevissimis subebracteatis, filamentis corolla longiori-  
bus, petalis linearibus, foliis oblongo-lanceolatis.*

*H. lutea. Bot. mag. 1062.*

*H. pumila. Jacq. ic. rar. 2. t. 453.*

*Veratrum luteum. Willd. sp. pl. 4. p. 897.*

*Melanthium dioicum. Walt. fl. car. 126.*

*Melanthium densum. Lam. encycl. 4. p. 26.*

In dry shady and hilly situations: Pennsylvania to Georgia.

4. July. v. v. Root bulbous, known by the name  
of *Star-wort*, and considered a remedy in colic.  
Flower white, male and female on different plants,  
which is a singularity in this family of plants.

Jacquin's *H. pumila* is without doubt an unexpanded  
spike of a male of this species: the appearance of a  
germen inferum is occasioned by the male flower  
being divested of its petals; and no female fructification  
appearing, the small pedicel was considered to be the  
rudiments of a germen.

*tenax.*

*dioica.*

*dubia.*

7. H. ? scapo nudo, spica gracili, floribus sessilibus, foliis longissimis angustissimumisque. — *Mich. fl. amer.* 1. p. 213.

On the sandy fields of Georgia and Florida. *Michaux.* 4. +. Scape nearly three feet high; flowers small.

321. MEDEOLA. *Gen. pl. 619.**virginica.*

1. M. foliis in medio caule verticillatis, summitate ternis, ovali-lanceolatis, pedicellis aggregatis terminalibus. — *Willd. sp. pl. 2. p. 270.*

*Icon. Bot. mag.* 1316. *Pluk. alm. t. 328. f. 4.*

In fertile shady woods, near rivulets: Canada to Georgia. 2d. April, May. v. v. Flowers yellowish-green; anthers brown. This plant is known by the name *Indian Cucumber*: the roots have a strong resemblance in taste and flavour to cucumbers, and are eaten by the natives.

322. TRILLIUM. *Gen. pl. 620.**sessile.*

1. T. flore sessili erecto, petalis lanceolatis erectis calyce duplo longioribus, foliis sessilibus lato-ovalibus acutis. — *Willd. sp. pl. 2. p. 272.*

*Icon. Bot. mag.* 40. *Catesb. car.* 1. t. 50. *Pluk. alm. t. 111. f. 6.*

On the side of fertile hills, in shady and rocky grounds: Pennsylvania to Carolina. 2d. April, May. v. v. Flower dark chocolate brown; berry purple. I have seen a specimen with yellow flowers, brought from the Cherokee nation, which probably may be a distinct species.

*petiolatum.*

2. T. flore sessili erecto, petalis linear-lanceolatis erectis calyce paulo longioribus, foliis longissime petiolatis ovali-lanceolatis acutis.

On the waters of the Kooskoosky. *M. Lewis.* 2d. June. v. s. This singular species, whose flowers resemble those of the first species, has leaves very much like *Plantago major*.

*pictum.*

3. T. pedunculo erectiusculo, petalis ovali-lanceolatis acutis recurvatis calyce angusto subduplo longioribus, foliis ovatis acuminatis basi rotundatis abrupte petiolatis.

T. erythrocarpum. *Mich. fl. amer.* 1. p. 216.

T. undulatum. *Willd. hort. ber.* 1. p. 55.

In sphagnum bogs, on the high mountains of Pensylva-

nia, Cara  
Flowers  
berry scar

4. T. peduncu  
calyce lin  
acutis arc  
On the rapic  
v. s. Flc

5. T. peduncu  
foliis ov  
amer. 1. 1  
In the pine  
May. v.

6. T. peduncu  
planis re  
dilatato-r  
petiolatis.  
Icon. Bot.  
In shady ro  
ticularly  
2d. May.

may be c  
seen it i  
nine inch  
cies is, i  
roneously  
sion to q

7. T. pedunc  
acuminat  
multo la  
sessilibus

$\alpha$ . T. flore ma  
T. rhombo

T. fœtidum  
Icon. Bot.

$\beta$ . T. flore mi  
T. erectum  
On the mo  
rolina.  
or white  
almost b

8. T. pedunc  
patentib  
ovato-rh

- nia, Carolina, and Canada.  $\frac{1}{2}$ . May, June. v. v. Flowers white, with purple veins at the bottom; berry scarlet.
4. T. pedunculo erecto, petalis oblongis acutis patentibus calyce linearis paulo longioribus, foliis ovatis sensim acutis arcte sessilibus. *On the rapids of Columbia river. M. Lewis.*  $\frac{1}{2}$ . April. v. s. Flowers pale purple. *ovatum.*
5. T. pedunculo erecto, petalis calyce vix longioribus, foliis ovali-oblongis obtusis sessilibus.—*Mich. fl. amer. 1. p. 215.* In the pine-woods of Lower Carolina. *Michaux.*  $\frac{1}{2}$ . May. v. s. Flowers deep flesh-colour. *pumilum.*
6. T. pedunculo recurvato, petalis lanceolatis acuminatis planis reflexis longitudine et latitudine calycis, foliis dilatato-rhomboideis abrupte acuminatis brevissime petiolatis.—*Willd. sp. pl. 2. p. 271.* *Icon. Bot. mag. 954. Smith spicil. 4.* In shady rocky situations: Pennsylvania to Carolina; particularly on the banks of Schuylkill near Philadelphia.  $\frac{1}{2}$ . May. v. v. Petals white; berries purple. This may be called the largest of the genus, as I have often seen it near two feet high, with leaves measuring nine inches in diameter. Catesby's figure of this species is, according to Dr. Smith's observation, so erroneously represented that it would only create confusion to quote it here. *cernuum.*
7. T. pedunculo inclinato, flore nutante, petalis ovatis acuminatis planis patentibus calyce vix longioribus multo latioribus, foliis lato-rhomboideis acuminatis sessilibus.—*Willd. sp. pl. 2. p. 271.* *erectum.*
- a. T. flore majore, petalis atropurpureis. *T. rhomboideum z. Mich. fl. amer. 1. p. 215.* *atropurpureum.*
- T. flore minore, petalis albis, germine rubro. *T. foetidum. Parad. Lond. 35.* *album.*
- Icon. Bot. mag. 470.*
- b. T. erectum  $\beta$ . *Bot. mag. 1027.* On the mountains, in boggy soil: Pennsylvania to Carolina.  $\frac{1}{2}$ . May. v. v. Flowers either dark purple or white, changing to red with a red germ; berries almost black.
8. T. pedunculo erecto, petalis obovatis obtusiusculis planis patentibus calyce vix longioribus et latioribus, foliis ovato-rhombeis acuminatis arcte sessilibus. *obovatum.*

## ALEXANDRIA TRIGYNA. Trillium.

T. camtschaticum. *Pallas in herb. Lambert.*

In Canada near Montreal. ♀. v. s. Flowers dark rose-coloured, probably white when first opening. The specimens in the herbarium of A. B. Lambert, Esq. agree in every respect with those from Canada.

*pendulum.*

9. T. pedunculo inclinato, flore pendulo, petalis ovatis brevi-acuminatis planis patentibus calycem ovatum acuminatum subaequantibus, foliis subrotundo-rhomboideis acuminatis subsessilibus.—*Willd. hort. berol.* 1. t. 35.  
On the mountains of Pennsylvania. ♀. April, May. v. v. Petals of a dirty white, with netted veins.

*grandiflorum.*

10. T. pedunculo inclinato, flore subcernuo petalis spathulato-lanceolatis basi erectis, limbo patentibus calyce multo majoribus, foliis late rhomboe-ovatis.—*Parad. lond. t. 1. Bot. mag.* 835.

T. rhomboideum γ. *grandiflorum.* *Mich. fl. amer.* 1. p. 216.

On the mountains and rocky banks of rivers: Virginia and Carolina. ♀. May. v. v. Flowers large, white; berries dark purple. The roots of the whole genus are considered as possessing strong medicinal powers.

323. TOFIELDIA. *Huds. fl. angl.* 157.

Narthecium. *Juss. Mich. fl. amer.* 1. p. 209.

*pusilla.*

1. T. tota glabra, foliis brevissimis, scapo filiformi, spica pauciflora conglobata, calyculo rachi adnato, capsula globosa.—*Mich. fl. amer.* 1. p. 209.

In bogs on the mountains, near lakes, and particularly Lake Mistassins.—♀. June, July. v. v. Flowers greenish-white.

*pubescens.*

2. T. scapo, rachi pedicellisque pubescenti-asperis, spica oblonga interrupta, capsula subglobosa vix calycem superante.—*Mich. fl. amer.* 1. p. 209.

Melanthium racemosum. *Walt. fl. car.* 126.

Anthericum calyculatum. *Gronov. nec Linn.*

In swamps and sphagnous woods: Virginia and Carolina. ♀. July. v. v. Flowers white; anthers yellow.

*glutinosa.*

3. T. scapo pedicellisque glutinoso-scabro, spica fasciculis paucis alternantibus, capsula ovoidea calyce duplo longiore.—*Mich. fl. amer.* 1. p. 210.

From Quebec to Lake Mistassins. *Michaux.* ♀. +.

It has the appearance of *Anthericum Ossifragum*, according to Michaux's observation; and as I have never

seen any  
cium am  
as syno  
mag. 15  
seems to  
definitio  
subject.

324

1. T. floribus

loculari

Icon. *Fl.*

On the in

♀. Jul

flowers

2. T. floribus

laevibus

Icon. *Fl.*

In marsh

York.

3. T. floribus

trigono-

subaequ

In overfi

*Michau*

325.

1. Scheuchz

Icon. *Fl.*

In swamp

Albany

1. R. herm:

ca pra

*Willd.*

Icon. *En*

In shady

Virgin

veins;

2. R. herm:

niferis,

*pl.* 2. ;

seen any other plant approaching to it than *Narthecium americanum*, p. 227, I was induced to take them as synonymous; but after investigation,—vid. *Bot. mag.* 1505,—a specimen in the Banksian Herbarium seems to apply better to Michaux's plant, if his short definition of the species is sufficient to decide on the subject.

## 324. TRIGLOCHIN. Gen. pl. 616.

1. *T. floribus 6-andris, capsulis brevibus ovatis sulcatis 6-locularibus.*—*Willd. sp. pl. 2. p. 265.*  
Icon. *Fl. dan. 306. Engl. bot. 255.*  
On the inundated sea-coast of Canada and New England.  
*4. July. v. v.* A plant of a rush-like appearance; flowers very small.
2. *T. floribus 3-andris, capsulis linearibus basi attenuatis laevibus 3-locularibus.*—*Willd. sp. pl. 2. p. 264.*  
Icon. *Fl. dan. 490. Engl. bot. 366.*  
In marshes round the Salt-lake of Onondago, New York. *4. July. v. v.*
3. *T. floribus 3-andris 3-fidis breviter pedicellatis, capsulis trigono-subrotundis, foliis subsetaceis scapum spicamve subæquantibus.* *Mich. fl. amer. 1. p. 208.*  
In overflowed marshes, near Charlestown, Carolina. *Michaux. 4. +.*

*maritimum.**palustre.**triandrum.*

## 325. SCHEUCHZERIA. Gen. pl. 615.

1. *Scheuchzeria.* *Willd. sp. pl. 2. p. 263.*  
Icon. *Fl. dan. 76.*  
In swamps and on the borders of lakes, New York near Albany. *4. June. v. v.* Flowers greenish-yellow.

*palustris.*

2. *R. hermaphroditus; valvulis integerrimis oblongis, unicæ præcipue granifera, foliis cordato-lanceolatis.*—*Willd. sp. pl. 2. p. 250.*  
Icon. *Engl. bot. 1533.*  
In shady woods and moist meadows: Pennsylvania to Virginia. *4. June, July. v. v.* Leaves with red veins; flowers, as in all the rest, very inconspicuous.
2. *R. hermaphroditus; valvulis ovatis integris, omnibus graniferis, foliis lanceolatis undulatis acutis.*—*Willd. sp. pl. 2. p. 251.*

*sanguineus.**crispus.*

- Icon. *Curt. lond.* 2. t. 20.  
In dry fields and pastures, common. ♀. July. v. v.
- verticillatus.* 3. *R. hermaphroditus*; *valvulis integerimis, omnibus graniferis, spicis aphyllis, foliis lanceolatis, vaginis cylindricis.*—*Willd. sp. pl. 2. p. 250.*  
In rivulets and shady woods: Canada to Virginia. ♀. July. v. v.
- britannicus.* 4. *R. hermaphroditus*; *valvulis integerimis, omnibus graniferis, paniculæ spicis aphyllis, vaginis obsolete laceris, foliis lato-lanceolatis planis lævibus.*—*Willd. sp. pl. 2. p. 250.*  
Near rivulets: Virginia and Carolina. ♀. June, July. v. v.
- persicarioides.* 5. *R. hermaphroditus*; *valvulis dentatis, omnibus graniferis, foliis lanceolatis undatis integris.*—*Willd. sp. pl. 2. p. 252.*  
In shady wet woods, on the banks of ditches: Virginia and Carolina. ♂. July. v. v.
- crispatus.* 6. *R. hermaphroditus*; *valvulis obtuse cordatis utrinque cristatis tridentatis, una nuda, binis inæqualiter graniferis, spicis aphyllis, foliis inferioribus ovalibus, superioribus lanceolatis, omnibus undulatis.*—*Mich. fl. amer. 1. p. 217.*  
In Kentucky. *Michaux.* +. *Affinis R. persicarioideo.*
- obtusifolius.* 7. *R. hermaphroditus*; *valvulis dentatis, unica præcipue granifera, foliis radicalibus cordatis obtusis, caule scabriusculo.*—*Willd. sp. pl. 2. p. 254.*  
Icon. *Engl. bot.* 1999.  
In old pastures and gardens, as a common weed; probably introduced. ♀. June, July. v. v. *Common Dock.*
- aquaticus.* 8. *R. hermaphroditus*; *valvulis ovatis integris obsolete graniferis, foliis cordato-lanceolatis acutis.*—*Willd. sp. pl. 2. p. 255.*  
Icon. *Engl. bot.* 2104.  
In small ponds and ditches: Pennsylvania to Virginia; rare. ♀. July, Aug. v. v.
- digynus.* 9. *R. hermaphroditus, digynus*; *valvulis ovatis integris grano destitutis, foliis radicalibus reniformibus longius petiolatis.*—*Willd. sp. pl. 2. p. 258.*  
Icon. *Fl. dan.* 14.\* *Pluk. alm. t. 252. f. 2.*  
In Labrador. *Colmaster.* ♀. June, July. v. s. in *Herb. Dickson.*

10. *R. dioica*; *foliis* *tutis.*—*Willd.*  
Icon. *Engl. bot.*  
Common in all c.—Aug. v. v.

327. \*

*\* Caul.*

1. *S. subinermis*; *lato hastatis* *baecis globosi*  
*S. Bona nox.*  
*car. 245.*  
2. *S. aspera* ♂. *La*  
Icon. *Pluk. alm.*  
♂. *S. foliis longis a*  
On the sea-coast  
July. v. v.
2. *S. inermis*; *foliis* *7-nervibus.*  
*S. variegata.* *W*  
Icon. *Pluk. alm.*  
In woods of Ca
3. *S. aculeata*; *ca*  
*mibus ovatis*  
*pl. 4. p. 775*  
In dry woods,  
Carolina. ♀
4. *S. aculeata*; *1*  
*baccis acumi*  
*S. China?* *Wa*  
In the lower s  
on river side
5. *S. aculeata*; *fo*  
*sub-5-nervit*  
*communi pe*  
*S. glauca.* *Mic*  
In hedges and
6. *S. subinermis*; *3*  
*nervibus*  
*breviore.*  
Near Savanna  
Leaves very

- R. dioica ; foliis lanceolato-hastatis, valvulis grano destitutis.—Willd. sp. pl. 2. p. 260.

Icon. Engl. bot. 1574.

Common in all dry fields and gravelly hills. 24. June—Aug. v. v.

## 327. SMILAX. Gen. pl. 1528.

\* Caule fruticoso ; ramis angulatis.

1. S. subinermis ; foliis lanceolatis acuminatis basi auriculato hastatis margine ciliato-aculeatis 3—5-nervibus, baecis globosis.—Willd. sp. pl. 4. p. 782.

S. Bona nox. Mich. fl. amer. 2. p. 237. Walt. fl. car. 245.

S. aspera ♂. Lam. encycl. 6. p. 465.

Icon. Pluk. alm. t. 111. f. 3.

- ♂ S. foliis longis angustis lanceolatis. Walt. fl. car. 245.

On the sea-coast of Carolina and Florida. ½. June, July. v. v.

2. S. inermis ; foliis cordato-ovatis acutis ciliato-aculeatis 7-nervibus.—Willd. sp. pl. 4. p. 781.

S. variegata. Walt. fl. car. 244.

Icon. Pluk. alm. t. 111. f. 1.

In woods of Carolina and Georgia. ½. July. v. v.

3. S. aculeata ; caule tetragono superne inermi, foliis inerme quadrangulatis ovatis subcordatis acutis 5-nervibus.—Willd. sp. pl. 4. p. 775.

In dry woods, on the edges of ponds : Pensylvaniato Carolina. ½. June, July. v. v. Berries black.

4. S. aculeata ; foliis cordato-ovatis laevibus 3-nervibus, baccis acuminatis.

S. China ? Walt. fl. car. 245.

In the lower sandy countries of Virginia and Carolina, on river sides. ½. July. v. v. Berries red.

5. S. aculeata ; foliis inermibus ovato-lanceolatis cuspidatis sub-5-nervibus subtus glaucescentibus, pedunculo communi petiolo longiore.—Willd. sp. pl. 4. p. 776.

S. glauca. Mich. fl. amer. 2. p. 237. Walt. fl. car. 245. In hedges and swamps, common. ½. June, July. v. v.

6. S. subinermis ; foliis inermibus ovatis acutis cuspidatis 3-nervibus concoloribus, pedunculo communi petiolo breviore.

Near Savannah in Georgia. Enslen. ½. July. v. s. Leaves very shining on both sides ; berries black.

## Acetosella.

hastata.

lanceolata.

Bona nox.

Walteri.

ovata.

- alba.* 7. *S. subinermis*; caule obsolete angulato, foliis elongato-lanceolatis coriaceis glabris integerrimis 3-nervibus, umbellis paucifloris brevissime pedunculatis.  
In sandy grounds, on the edge of rivulets: Carolina. Walter. ½. June. v. s. in *Herb. Walter.* Berries white.  
\*\* *Caule fruticoso; ramis teretibus.*
- lanceolata.* 8. *S. inermis*; foliis inermibus lanceolatis, umbellis multifloris longe pedunculatis.—*Willd. sp. pl. 4. p. 783.*  
Icon. *Catesb. car. 2. t. 84.*  
Near the sea-coast of Virginia and Carolina. ½. May, June. v. v. Berries red.
- pubera.* 9. *S. inermis*; foliis oblongis acutis cordatis sub-5-nervibus subtus molliter pubescentibus, umbellis breviter pedunculatis, pedicellis brevissimis, baccis oblongis acutis.—*Willd. sp. pl. 4. p. 785. Mich. fl. amer. 2. p. 238.*  
*S. pumila. Walt. fl. car. 244.*  
In shady woods: Carolina and Georgia. ½. May. v. s. in *Herb. Lyon.* According to Walter, the berries are white.
- Pseudo-China.* 10. *S. inermis*; foliis inermibus, caulinis cordatis, rameis ovato-oblongis 5-nervibus, pedunculis longissimis.—*Willd. sp. pl. 4. p. 785.*  
Icon. *Sloan. jam. 1. t. 143. f. 1.*  
In sandy fields and dry woods: New Jersey to Carolina. ½. May, June. v. v. The roots very large, and are searched after by hogs.
- rotundifolia.* 11. *S. sparse aculeata*; foliis subrotundo-ovatis acuminatis levissime cordatis 5-nervibus, baccis sphaericis.—*Willd. sp. pl. 4. p. 779.*  
Common in hedge-rows: Canada to Carolina. ½. June. v. v.
- caduca.* 12. *S. aculeata*; foliis ovatis mucronatis 5-nervibus, pedunculo communi vix petiolo longiore.—*Willd. sp. pl. 4. p. 780.*  
In dry fields: Canada to Virginia. ½. June. v. v.
- laurifolia.* 13. *S. aculeata*; ramis inermibus, foliis coriaceis ellipticis vel elliptico-lanceolatis obtusis recurvato-acutis 3-nervibus, umbellis brevissime pedunculatis.—*Willd. sp. pl. 4. p. 779.*  
Icon. *Catesb. car. 1. t. 15.*  
In sandy boggy woods: New Jersey to Georgia. ½.

- July, Aug.  
thick smooth
14. *S. aculeata*; 3-nervibus, giore.  
In sandy woc v. v. Leav
15. *S. caule teret lato-cordati sp. pl. 4. p. 785.*  
Icon. *Catesb.*  
In sandy wet ¼. July.
16. *S. caule ang ovalibus 7- pedunculo Willd. sp.*  
Icon. *Pluk.*  
In dry fields, roline. 2
17. *S. caule tere tis acumini culatis.—*  
In old fields vania. 2
- 328.
1. *D. foliis ver natis utri bifidis.—*  
In hedges at July. v.
2. *D. foliis alte natis sub bus simpl D. quinata.*  
D. panicula  
Icon. *Jacq.*  
In woods at May, Ju

- July, Aug. v. v. A very handsome vine, with very thick smooth leaves.
14. *S. aculeata*; foliis ovato-panduræformibus acuminatis *panduratus*.  
3-nervibus, pedunculo communi petiolo duplo longiore.  
In sandy woods: New Jersey to Carolina. ½. July.  
v. v. Leaves smooth and shining on both sides.

\*\*\* *Caule herbaceo*.

15. *S. caule tereti scandente, foliis longe petiolatis triangulo-cordatis rotundato-obtusis, nervis glabris*.—Willd. sp. pl. 4. p. 780.  
Icon. *Catesb. car. 1. t. 52*.  
In sandy wet woods and bogs: Virginia and Carolina.  
¾. July. v. v.
16. *S. caule angulato erecto simplici, foliis longe petiolatis ovalibus 7-nervibus, umbellis longissime pedunculatis, pedunculo compresso, baccis depresso-globosis*.—Willd. sp. pl. 4. p. 782.  
Icon. *Pluk. alm. t. 225. f. 4*.  
In dry fields, on the edges of woods: New York to Carolina. ¾. June, July. v. v.
17. *S. caule tereti scandente, foliis subrotundo-ovatis cordatis acuminatis 9-nervibus, umbellis longissime pedunculatis*.—Willd. sp. pl. 4. p. 786.  
In old fields on the edges of woods: Canada to Pennsylvania. ¾. May—July. v. v.

*tamnoides*.*herbacea*.328. DIOSCOREA. *Gen. pl. 1530*.

1. *D. foliis verticillatis quaternis alternisve cordatis acuminatis utrinque glabris 9-nervibus, nervis lateralibus bifidis*.—Walt. fl. car. 246.  
In hedges and old fields: Virginia and Carolina. ¾.  
July. v. v.
2. *D. foliis alternis oppositis verticillatisque cordatis acuminatis subtus pubescentibus 9-nervibus, nervis laterali- bus simplicibus*.—Willd. sp. pl. 4. p. 796.  
D. quinata. Walt. fl. car. 240.  
D. paniculata. Mich. fl. amer. 2. p. 239.  
Icon. Jacq. sic. t. 626. Pluk. alm. t. 375. f. 5.  
In woods and hedges common: Canada to Carolina. ¾.  
May, June. v. v.

*quaternata*.*villosa*.

lliis elongato-  
3-nervibus,  
tis:  
is : Carolina.  
ter. Berries

nbellis multi-  
4. p. 783.

a. ½. May,

ab-5-nervibus  
breviter pe-  
oblongis acu-  
fl. amer. 2.

May. v. s.  
he berries are

rdatis, rameis  
longissimis.—

ey to Carolina.  
large, and are

itis acuminatis  
sphæricis.—

Carolina. ½.

rribus, pedun-  
Willd. sp. pl. 4.

June. v. v.

iaceis ellipticis  
o-acutis 3-ner-  
tis.—Willd. sp.

Georgia. ½.

## IV. TETRAGYNIA.

329. SAURURUS. *Gen. pl. 632.**cernuus.*1. *S. caule folioso polystachyo, foliis sagittato-cordatis.* — *Willd. sp. pl. 2. p. 292.**Anonymos aquatica. Walt. fl. car. 127.**Mattuschkea aquatica. Gmel. syst. nat. 589.**Icon. Pluk. alm. t. 117. f. 3 & 4.*

In swamps and shady wet woods, frequent: Canada to Carolina. ♀. Aug. Sept. v. v.

## V. POLYGYNIA.

330. WENDLANDIA. *Willd. sp. pl. 2. p. 275.**populifolia.*1. *Wendlandia. Willd. l. c.**Androphylax scandens. Wendl. obs. 38.**Menispermum carolinianum. Walt. fl. car. 248. Mich. fl. amer. 2. p. 242.**Cissampelos smilacina. Willd. sp. pl. 4. p. 863.**Icon. Wendl. hort. t. 16. Jacq. ic. rar. 3. t. 629. Catesb. car. 1. t. 51.*In hedges and woods: Carolina to Florida. ½ June, July. v. v. Flowers very small, greenish-white; berries red. I venture to take *Cissampelos smilacina* as a synonym, although Jacquin's figure deviates in character very much from the present genus; but as I have had the opportunity of seeing the *Menispermum* of the above authors frequently in flower, I am convinced that all the above synonyms are correct: the number of stamens being variable, brings the genus so near to *Menispermum*, that I should not have taken it from it, if it were not on the authority of Willdenow.331. ALISMA. *Gen. pl. 625.**trivialis.*1. *A. foliis ovalibus cordatis obtusissimis 9-nervibus, umbellis composite verticillato-paniculatis, fructibus obtuse-trigonis.*A. *Plantago. M.*  
In ditches and p  
July. v. v.2. *A. foliis ovalibus  
bus umbellis  
cellis filiformes*  
In salt-marshes,  
sylvania. ♀  
small. Both  
but are suffic  
species.3. *A. pusilla; fol  
bus.—Willd.*  
In overflowed  
plentifully o  
adelphia, to  
v. v. A sma4. *A. foliis ovatis*  
*pl. 2. p. 27*  
In stagnant v  
♀. July.

A. *Plantago*. *Mich. fl. amer.* 1. p. 218.

In ditches and ponds, common : Canada to Florida.  $\frac{1}{4}$ .

July. v. v.

2. A. foliis ovalibus subcordatis abrupte acuminatis 5-nervi-  
bus umbellis composite verticillato-paniculatis, pedi-  
cellis filiformibus.

In salt-marshes, on the coast of New Jersey and Pen-  
sylvania.  $\frac{1}{4}$ . July, Aug. v. Laves and flowers  
small. Both species approach near to *A. Plantago*,  
but are sufficiently distinct to admit them to be called  
species.

3. A. pusilla ; foliis linearis-subulatis, umbellis subsimplici-  
bus.—*Willd. sp. pl. 2. p. 279.*

In overflowed places : New York to Florida, scarce ;  
plentifully on the shores of the Delaware near Phila-  
delphia, to the edge of low water mark.  $\odot$ . Aug.  
v. v. A small plant, with flowers in proportion large.

4. A. foliis ovatis obtusis, pedunculis solitariis. *Willd. sp.*  
*pl. 2. p. 278.*

In stagnant waters, on the banks of the St. Laurence.  
 $\frac{1}{4}$ . July. v. s.

*parviflora.*

*subulata.*

*natans.*

—cordatis.—

: Canada to

p. 275.

248. *Mich.*

863,

629. *Catesb.*

h. June,  
enish-white ;  
elos smilacina  
e deviates in  
enus ; but as  
*Menispernum*  
er, I am con-  
correct : the  
ings the ge-  
ould not have  
e authority of

ervibus, um-  
fructibus ob-

2. A. foliis quinatis  
inæqualiter serratis,  
multifloris, c  
unguis longioribus,  
In Georgia, p.  
1/2 May. v.  
yellow, whit

3. A. foliis quinatis  
liter serrulatis,  
ris, corollis  
bus calyce longis  
fructibus inaequatis.  
A. lutea. *Micromeria*  
Icon. *Schmidt arb.*  
*Mich. arb.*  
In the mountains  
Kentucky.  
tree; flower

4. A. foliis quinatis  
unguis longioribus, fructibus  
In the western  
June. v. i

5. A. foliis quinatis  
calyce brevibus,  
fructibus inaequatis.  
In the forests

6. A. foliis quinatis  
longissimo  
talis patulis  
*amer.* 1. p.  
*A. parviflora.*  
On the bank  
Augustin.  
flowers w  
genus is k  
of *Buck's.*

## VII. HEPTANDRIA.

---

### I. MONOGYNIA.

332. TRIENTALIS. *Cal.* 7-phylloides. *Cor.* 7-partita,  
plana. *Bac.* 1-locularis sicca.  
333. ÆSCULUS. *Cal.* 5-dentatus. *Cor.* 5-petala, in-  
æqualis, calyci inserta. *Caps.* 3-locularis.
- 

### I. MONOGYNIA.

#### 332. TRIENTALIS. *Gen. pl.* 626.

*americana.*

1. T. foliis angusto-lanceolatis acuminatis obliquis.  
T. europaea. *Mich. fl. amer.* 1. p. 220.

In cedar swamps and other sphagnum places on high  
mountains: Canada to Virginia. 1/2 July, Aug.  
v. v. Flowers white. This delicate little plant differs  
considerably from the European sort, although Mi-  
chaux considers it to be the same.

#### 333. ÆSCULUS. *Gen. pl.* 628.

*Pavia.*

1. A. foliis quinatis glabris inæqualiter serratis, racemo-  
laxo, fasciculis subtrifloris, corollis 4-petalis, petalo-  
rum conniventium unguibus longitudine calycis, sta-  
minibus 7. corolla brevioribus, fructibus inermibus.—  
*Willd. sp. pl.* 2. p. 286.

*Icon.*, *Schmidt arb.* 39. *Trew ehret.* t. 15. *Mich. arb.* . . .

In fertile valleys, on the mountains of Virginia and Ca-  
rolina. 1/2 May, June. v. v. Flower scarlet; ge-  
nerally only a shrub or small tree.

*discolor.*

2. *A. foliis quinatis utrinque acuminatis subtus tomentosis inaequaliter serrulatis, racemo thyrsideo, fasciculis multifloris, corollis 4-petalis, petalorum conniventium unguibus longitudine calycis, staminibus 7. corolla brevioribus, fructibus inermibus.*

In Georgia, principally the western territory. *Lyon.*  
*h. May. v. v.* Not above four feet high; flowers yellow, white, and purple variegated.

3. *A. foliis quinatis subtus ad costam pubescentibus aequaliter serrulatis, racemo thyrsideo, fasciculis multifloris, corollis 4-petalis, petalorum conniventium unguibus calyce longioribus, staminibus corolla brevioribus, fructibus inermibus.* — *Willd. sp. pl. 2. p. 286.*

*A. lutea. Mich. fl. amer. 1. p. 219.*  
*Icon. Schmidt arb. 40. Act. soc. nat. berol. 8. t. 6.*

*Mich. arb. . . .*  
 In the mountains of Virginia and Carolina, and woods of Kentucky. *h. May, June. v. v.* Grows to a large tree; flowers yellow.

4. *A. foliis quinatis glaberrimis, corollis 4-petalis patulis, unguibus longitudine calycis, staminibus corolla longioribus, fructibus spinosis.* — *Willd. enum. pl. 405.*

In the western counties of Pennsylvania and Virginia. *h. June. v. v.*

5. *A. foliis quinatis, corollis 4-petalis patulis, unguibus calyce brevioribus, staminibus corolla duplo longioribus, fructibus spinosis.* — *Willd. enum. 406.*

In the forests of Kentucky. *h. June. v. s.* A tall tree.

6. *A. foliis quinatis subtus subtomentosis serrulatis, racemo longissimo laxo, fasciculis subtrifloris, corollis 4-petalis patulis, staminibus longissime exertis.* — *Mich. fl. amer. 1. p. 220.*

*A. parviflora. Walt. fl. car. 128.*

On the banks of rivers: Georgia, principally near St. Augustin. *h. May, June. v. v.* A small shrub; flowers white, very ornamental. The whole of this genus is known in their native countries by the name of *Buck's-eye-tree.*

*flava.**glabra.**pallida.**macrostachya.*

[A.]

r. 7-partita,

5-petala, in-

aris.

16.

iquis.

laces on high  
f. July, Aug.  
le plant differs  
although Mi-

S.

rratis, racemo  
petalis, petalo-  
ne calycis, sta-  
s inermibus.—t. 15. *Mich.*irginia and Ca-  
ver scarlet; ge-

## VIII. OCTANDRIA.

### I. MONOGYNIA.

\* *Germen inferum.*

334. RHEXIA. *Cal. 4-fidus. Cor. 4-petala, calyci inserata. Anth. declinatae. Caps. 4-locularis, intra ventrem calycis.*
335. EPILOBIUM. *Cal. 4-fidus, tubulosus. Cor. 4-petala. Caps. oblonga, 4-locularis. Sem. papposa.*
336. GAURA. *Cal. 4-fidus, tubulosus. Cor. 4-petala, ascendens. Nux 1-sperma, 4-angula.*
337. CLARKIA. *Cal. 4-fidus, tubulosus. Cor. 4-petala : petala cruciatim 3-loba. Filam. 4. castrata. Caps. 4-locularis.*
338. CENOTHERA. *Cal. 4-fidus, tubulosus. Cor. 4-petala : petalis integris. Filam. omnia fertilia. Anth. lineares. Caps. 4-locularis.*
339. OXYCOCCUS. *Cal. 4-fidus. Cor. 4-partita : lacinii sublinearibus, revolutis. Filam. contiventia. Anth. tubulose, bipartitae. Bacc. polysperma.*

\*\* *Germen superum.*

340. MENZIESIA. *Cal. 1-phyllo. Cor. 1-petala, ovata. Fil. receptaculo inserta. Caps. 4-locularis, dissepimenta e marginibus inflexis valvularum. Sem. numerosa, oblonga.*
341. DIOSPYROS. *Cal. 4-fidus. Cor. urceolata, 4-fida. Styl. 4-fidus. Bacc. 8-sperma. Polygama.*
342. ACER. *Cal. 5-fidus. Pet. 5. Samarae 2, basi untae, 1-spermæ, alatae. Polygamum.*
343. DIRCA. *Cal. 0. Cor. tubulosa, limbo obsoleto. Stam. tubo longiore. Bacc. 1-sperma.*
344. JEFFERSONIA. *Cal. 5-phyllo, deciduus. Pet. 8. incurvo-patentia. Caps. obovata, substipitata, 1-lo-*

cularis, infi  
ad basin ar  
345. PISTIA. C  
*Cor. 0. F  
locularis, 1*

346. CHRYSOPI  
*Caps. birost*

347. POLYGON  
*Sem. 1. an*

348. BRUNNIC  
*coriaceus.*

349. CARDIOSI  
*tar. 4-ph  
flatae.*

350. SAPINDUS  
*nosæ, con*

351. MYRIOPH  
*Stam. 4. 0*

I.

334.

1. R. caule rufe  
latis, ovali-  
longo tubi  
p. 221. I  
Icon. Pluk. I  
a. R. foliis. ang  
rate purpu  
VOL. I.

cularis, infra apicem dehiscens. Sem. plura, oblonga,  
ad basin arillata.

345. **PISTIA.** Cal. spatha tubuloso-cucullata, lingulata.  
Cor. O. Filam. laterale. Anther. 3-8. Caps. 1-  
locularis, polysperma.

## II. DIGYNIA.

346. **CHRYSOLENIUM.** Cal. 4-fidus, coloratus. Cor. O.  
Caps. birostris, 1-locularis, polysperma.

## III. TRIGYNIA.

347. **POLYGONUM.** Cal. O. Cor. 5-partita, calycina.  
Sem. 1, angulatum, tectum.

348. **BRUNNICHIA.** Cal. angulatus, 5-fidus, demum  
coriaceus. Cor. O. Caps. 1-sperma, inclusa calyce  
valde aucto, pedicello ensiformi-dilatato suffulto.

349. **CARDIOSPERMUM.** Cal. 4-phyllo. Pet. 4. Nec-  
tar. 4-phyllo, inaequale. Caps. 3. connatæ, in-  
flatae.

350. **SAPINDUS.** Cal. 4-phyllo. Pet. 4. Caps. car-  
nosæ, connatæ, ventricosæ.

## IV. TETRAGYNIA.

351. **MYRIOPHYLLUM.** Cal. 4-fidus. Pet. 4. caduca.  
Stam. 4. 6. 8. Styl. O. Sem. 4. corticata.

## I. MONOGYNIA.

### 334. RHEXIA. Gen. pl. 636.

1. R. caule rufescens hirsutissimo, foliis hirsutis subpetio-  
latis, ovali-lanceolatis, lanceolatis linearibusve, calyce  
longo tubuloso glabriusculo.—Mich. fl. amer. 1.  
p. 221. Willd. sp. pl. 2. p. 301.

- Icon. Pluk. mant. t. 428. f. 1. Lam. illustr. 283. f. 1.  
a. R. foliis angusto-lanceolatis vel oblongis, floribus satu-  
rate purpureis. Mich. l. c.

mariana:

purpurea.

- rubella.*       $\beta.$  R. foliis ovali-lanceolatis vel oblongis, floribus dilute rubellis. *Mich. l. c.*
- exalbida.*       $\gamma.$  R. foliis linearibus, caule hirsutiore, floribus pallidis. *Mich. l. c.*  
R. lanceolata. *Walt. fl. car.* 129.  
In bogs and sandy woods, near waters: New Jersey to Carolina.  $\frac{4}{4}$ . July, Aug. v. v. From one to three feet high; flowers handsome.
- ciliosa.*      2. R. caule subquadragulo glabro, foliis parvis subpetiolatis ovalibus subtus glabris, supra rarer hispidis, margine distincte ciliatis, floribus subsolitariis involucratis, antheris breviusculis.—*Mich. fl. amer.* 1. p. 221.  
R. petiolata. *Walt. fl. car.* 130?  
In bogs of Lower Carolina.  $\frac{4}{4}$ . July, Aug. v. s. in *Herb. Enslen.* Flowers large, beautiful purple.
- virginica.*      3. R. caule alato-angulato pilis rarioribus adperso, foliis sessilibus ovali-lanceolatis denticulis setaceis distinctis, que ciliato-serratis, hirsutie sparsa utrinque hispidulis. *Mich. fl. amer.* 1. p. 222.  
Icon. *Bot. mag.* 968. *Pluk. alm.* t. 202. f. 8. *Lam. illustr.* t. 283. f. 2.
- $\gamma$ -*nervia.*       $\beta.$  R. caule 4-5-pedali. *Walt. fl. car.* 130.  
On the edges of ponds, in wet woods and bogs: New York to Carolina.  $\frac{4}{4}$ . July, Aug. v. v. Flowers large, purple, and very handsome.
- glabella.*      4. R. caule cylindraceo glabro, foliis sessilibus erectis lanceolatis laevibus glabris tenuissime denticulatis, calycibus brevissime hirsutie glutinosis. *Mich. fl. amer.* 1. p. 222.
- Alifanus.*       $\beta.$  R. foliis subovatis. *Walt. fl. car.* 130.  
In sandy moist woods: Carolina and Georgia.  $\frac{4}{4}$ . July. v. s. in *Herb. Enslen.* Flowers the largest of the North American sorts, deep purple.
- stricta.*      5. R. caule stricte-erecto alato-tetragono glabro ad genicula brevi-barbato, foliis sessilibus erectis angusto-lanceolatis attenuato-acuminatis trinervibus utrinque glabris, corymbo dichotomo, calycibus glabriusculis.  
In the bogs of Lower Carolina and Georgia.  $\frac{4}{4}$ . June, July. v. s. in *Herb. Enslen.* Flowers very handsome, purple.
- lutea.*      6. R. caule quadrangulo hirsuto, foliis rarer longiusculeque hirsutis, inferioribus cuneato-oblongis obtusis, superioribus lanceolatis, antheris breviusculis. *Mich. fl. amer.* 1. p. 222. *Walt. fl. car.* 130.

s dilute ru-

us pallidis.

w Jersey to  
one to three

subpetiolata  
spidis, mar-  
involucratis,  
p. 221.

ug. v. s. in  
purple.

perso, foliis  
distinctis.  
te hispidulis.

f. 8. Lam.

bogs : New  
v. Flowers

erectis lan-  
culatis, caly-  
l. fl. amer. 1.

Georgia. 4.  
the largest of

o ad genicula  
gusto-lanceo-  
inque glabris,  
ulis.

a. 4. June,  
ery handsome,

ongiusculeque  
obtusis, supe-  
lis. Mich. fl.



Phacelia villosa.



Khamia heterophylla.

OCTAVI

In pine-woods  
v. s. in Her

7. R. caule cylindricus oblongis floribus subciliatis

In Carolina.

335.

1. E. foliis sparsis inaequilibus p. 313.

Icon. Fl. dan.  
In wet spring  
shire, New  
July.  
white; the

2. E. foliis alterna- centibus ave-

p. 314.  
Icon. Fl. dan.  
β. E. foliis angustis Pallas in E.

In Labrador.

v. s. in Her

3. E. caule tetrapetalum alternis lanceolatis illaribus altitudine p. 4-lobo.

On the north  
Lambert.  
folium.

4. E. caule lineari positis, superfluous sp. pl. 2. p.

Icon. Fl. dan.  
In low ground.  
Flowers sn

5. E. caule terebrantis caulinis gerrimis, dunculatis.

In pine-woods of Georgia and Florida. ♂. June, July.  
v. s. in Herb. Leconte. Flowers small, yellow.

7. R. caule cylindrico subpubescente, foliis alternis linearibus oblongis obtusis sessilibus utrinque pubescentibus, floribus subsolitariis.—*Lam. encycl. 6. p. 2.*  
In Carolina. Bosc. +. Flowers yellow.

*linearifolia.*

335. EPILOBIUM. Gen. pl. 639.

1. E. foliis sparsis linearis-lanceolatis venosis glabris, floribus inaequalibus, staminibus declinatis.—*Willd. sp. pl. 2. p. 313.*

*angustifoli-*  
*um.*

Icon. *Fl. dan. 289.* *Curt. fl. lond. 2. t. 24.*  
In wet springy ground in the mountains of New Hampshire, New York, and Pennsylvania, also in Canada.  
♀. July. v. v. Flowers beautiful purple, sometimes white; the plant often above five feet high.

2. E. foliis alternis lanceolato-ovatis subintegerrimis pubescens aeniis, floribus inaequalibus. *Willd. sp. pl. 2. p. 314.*

*latifolium.*

- Icon. *Fl. dan. 565.*  
β. E. foliis angusto-lanceolatis integerrimis, caule pumilo, tetrapetalum. *Pallas in Herb. Lambert.*

In Labrador. β. on the north-west coast. ♀. July.  
v. s. in Herb. Lambert. Flowers purple, large.

3. E. caule tetragono glabriusculo, foliis oppositis, supremis alternis lanceolatis acuminatis serrulatis, pedunculis axillaribus alternis elongatis, calycis laciniis angustis longitudine petalorum, stylo exerto, stigmate crasso 4-lobo.

On the north-west coast. *Pallas.* ♀. v. s. in Herb. Lambert. Flowers yellow, the size of *E. angustifolium.*

*luteum.*

4. E. caule lineis prominulis subquadrangulatis, foliis oppositis, supremis alternis lanceolatis serrulatis.—*Willd. sp. pl. 2. p. 317.*

*tetragonum.*

Icon. *Fl. dan. 1029.*  
In low grounds: Canada to Carolina. ♀. July. v. v.  
Flowers small, pale red.

5. E. caule tereti pubescente superne virgatim ramoso, foliis caulinis oppositis, rameis alternis, linearibus integerrimis, floribus paucis terminalibus longissime pendulatis.

## OCTANDRIA MONOGYNIA. Epilobium.

- E. oliganthum*. *Mich. fl. amer.* 1. p. 223. In Canada and on the high mountains of New York and Pennsylvania. ♀. July. v. v. Flowers very small, pale red or white.
- coloratum*. 6. *E. caule tereti pubescente, foliis lanceolatis serrulatis petiolatis oppositis, superioribus alternis glabris rubro-venosis*. *Willd. enum.* 411. In Pennsylvania. *Muhlenberg.* ♀. July. v. v.
- palustre*. 7. *E. caule tereti, foliis sessilibus lanceolatis subdenticulatis, stigmate indiviso*.—*Willd. sp. pl. 2. p. 317.* *Icon. Engl. bot.* 346. In low grounds: Pennsylvania to Virginia. ♀. July. v. v.
- alpinum*. 8. *E. caule simplici subtereti 1-2-floro, foliis oppositis ellipticis integerrimis, floribus sessilibus*.—*Willd. sp. pl. 2. p. 318.* *Icon. Fl. dan.* 322. In Labrador. *Colmaster.* ♀. May, June. v. s. in Herb. *Dickson.* The smallest species, not above two inches high; flowers pale purple.

336. GAURA. *Gen. pl.* 638.

- biennis*. 1. *G. foliis lanceolatis dentatis, spica conferta, fructibus subrotundo-4-gonis pubescentibus*.—*Willd. sp. pl. 2. p. 311.* *Icon. Bot. mag.* 389. *Pluk. amalh. t. 428. f. 2.* On the edges of woods in fertile stony soil: Pennsylvania to Carolina. ♂. July, Aug. v. v. Flowers rose-coloured, large.
- angustifolia*. 2. *G. foliis crebris linearibus repando-undulatis, spica fructibus dissitis oblongo-4-gonis utrinque acutis subcandicantibus*. *Mich. fl. amer.* 2. p. 226. In dry old fields and woods: Virginia to Carolina. ♂. July. v. v. Flowers scarcely half the size of the foregoing, pale red.

337. CLARCKIA. *Pursh in linn. soc. trans.* v. 11.

- pulchella*. 1. *Clarckia*. *Pursh l. c.* On the Kooskoosky and Clark's rivers. *M. Lewis.* ♂. June. v. s. Flowers beautiful rose-coloured or purple. *Caulis erectus, teres, superne subramosus, pedalis et ultra. Folia alterna, linearia, integerrima, glabra*.





Clarkia pulchella.

*Flores ex magni, amp unguiculata  
Stamina 4 duplo brevi longitudine  
4-partitum*

## 338. C

1. *O. caule vill dentatis, fl staminibus p. 306.*  
Icon. *Fl. dan*  
Common in c June, July  
I have frequ and it mig into its cat objects can stance, thi a great dis which prol perties of
2. *O. caule purp staminibus p. 807.*  
Icon. *Murra*  
In old fields ♂. July, 1 species.
3. *O. caule læv staminibus p. 306.*  
Icon. *Mill. i*  
In fields and July. v. 1
4. *O. caule gl glabris, st In woods and v. v. Flo American*
5. *O. caule diff*

*Flores ex axillis superioribus, solitarii, subsessiles, magni, amœne purpurei. Calyx oenotheræ. Petala unguiculata, 3-partita, laciñis lateralibus divaricatis. Stamina 4: antheris linearibus, involutis: altera 4. duplo breviora: antheris castratis, subrotundis. Stylus longitudine fere petalorum. Stigma pallide luteum, 4-partitum: laciñis subrotundis.*

338. OENOTHERA. *Gen. pl. 637.*

1. *O. caule villoso scabro, foliis ovato-lanceolatis planis dentatis, floribus terminalibus subspicatis sessilibus, staminibus\* corolla brevioribus.* — *Willd. sp. pl. 2. p. 306.*

Icon. *Fl. dan. 446. Alp. exot. t. 324.*

Common in old fields: New England to Carolina. ♂.  
June, July. v. v.

I have frequently observed a singularity in this plant, and it might be interesting to make further inquiry into its cause; it is, that in a dark night, when no objects can be distinguished at an inconsiderable distance, this plant when in full flower can be seen at a great distance, having a bright white appearance, which probably may arise from some phosphoric properties of the flowers.

2. *O. caule purpurascente muricato, foliis lanceolatis planis, staminibus longitudine corollæ.* *Willd. sp. pl. 2. p. 807.*

Icon. *Murray com. gætt. 7. t. 1.*

In old fields and along fences: Pennsylvania to Carolina.  
♂. July, Aug. v. v. Flowers smaller than the first species.

3. *O. caule lævi subvilloso, foliis ovato-lanceolatis planis, staminibus corolla longioribus.* *Willd. sp. pl. 2. p. 306.*

Icon. *Mill. ic. 189. f. 1. Meerb. ic. 34.*

In fields and woods; Canada to Virginia; rare. ♂.  
July. v. v.

4. *O. caule glabriusculo ramoso, foliis ovato-lanceolatis glabris, staminibus declinatis.* *Willd. sp. pl. 2. p. 306.*

In woods and fields of Carolina. *Bartram.* ♂. July.  
v. v. Flowers larger than any other of the North American species, and of an agreeable scent.

5. *O. caule diffuso molliter pubescente, foliis ovali-oblone-*

*biennis.*

*muricata.*

*parviflora.*

*grandiflora.*

*sinuata.*

## OCTANDRIA MONOGYNA. Oenothera.

- gis dentato-sinuatis, floribus axillaribus villosis, capsulis prismaticis.* — Willd. sp. pl. 2. p. 309.  
Icon. Murray com. gætt. 5. t. 9. Pluk. alm. t. 203. f. 3.  
In dry fields of Virginia and Carolina. ♂. June, July.  
v. v.
- minima.* 6. *O. caule simplici unifloro, foliis pusillis lanceolatis integrerrimis pilosis, flore sessili hirsuto, germine prismatico.*  
In barren pine-woods of Georgia. Enslen. ♂. June.  
v. s. Not above an inch high; flower smallest of all the genus.
- glaуca.* 7. *O. glaberrima; foliis lato-ovalibus repando-subdentatis лævigatis glaucis, capsula ovato-tetragona pedicellata.*  
— Mich. fl. amer. 2. p. 224.  
On the banks of the Mississippi, in Illinois, and on the Peaks of Otter, Virginia. ♀. July. v. v. A handsome species with considerable large flowers.
- fruticosa.* 8. *O. glabriuscula; foliis lanceolatis subdentatis acutis, capsulis pedicellatis oblongo-clavatis angulatis.* Willd. sp. pl. 2. p. 310.  
Icon. Bot. mag. 332. Meerb. ic. 33.  
In shady woods, on the edges of rivulets and springs: Pennsylvania to Carolina. ♀. July. v. v. Commonly called *Sun-drops*.
- pumila.* 9. *O. glabra; caulis adscendentibus, foliis lanceolatis integrerrimis obtusis, capsulis subsessilibus elliptico-obovatis angulatis.* — Willd. sp. pl. 2. p. 310.  
Icon. Bot. mag. 355. Mill. dict. t. 188.  
In dry fields of Virginia and Carolina. ♀. July. v. v.
- hybrida.* 10. *O. caule erecto villoso, foliis utrinque pubescentibus lanceolatis remote subdentatis undulatis, capsulis subspicatis breviter stipitatis ovato-tetragonis.* Mich. fl. amer. 1. p. 225.  
*O. mollissima.* Walt. fl. car. 129.  
*Onagra floribus speciosis luteis &c.* Gron. fl. virg. 59.  
In sandy fields of Virginia and Carolina. ♀. June, July. v. v.
- linearis.* 11. *O. tota pubescens, gracilis; foliis linearibus integris, capsulis longiuscule stipitatis subrotundo-tetragonis villosis.* Mich. fl. amer. 2. p. 225.  
a. *Pluk. mant. t. 426. f. 6.*  
β. *Onagra 491. Gron. fl. virg. 59.*  
In dry barren fields; Virginia and Carolina. ♀. July.
- angustifolia?*  
*brevifolia.*

## OCTANDRIA

v. v. Flowers la  
plant.

12. *O. caule debili pub  
planis integris, c  
capsula clavata a  
p. 225.*  
*O. biennis.* Walt. j  
In barren lands of  
sylvania and Car  
small, golden-ye
13. *O. subpubescens;*  
ceolato-oblongis  
mitatem axillari  
tis subæqualiter  
On rocks near L  
mountains of I  
Aug. v. v. Pl
14. *O. scap's radicalib  
repando-denticu  
staminibus coro  
On the falls of t  
v. s. Flowers l  
bles *O. avulsi*  
All the species,  
flowers.*

## 339. OXYC

1. *O. repens;* foli  
subacutis glabi  
gatis, corollæ l  
*O. palustris.* Per  
Vaccinium Oxy  
p. 228.  
Vaccinium Oxy  
Icon. Fl. dan. 80  
In bogs and mos  
da to Pensylva  
evergreen cree  
purple, smalle
2. *O. repens;* cau  
tegerrimis pla  
bus, pedicellis  
Pers. syn. 1. j

- v. v. Flowers large, in proportion to the very low plant.
12. *O. caule debili pubescente, foliis lanceolatis obtusiusculis planis integris, calycis tubo laciniis duplo breviore, capsula clavata acutangula sessili.* — *Mich. fl. amer. 1. p. 225.*
- O. biennis. Walt. fl. car. 129?*  
In barren lands of Canada and on the mountains of Pennsylvania and Carolina. 4. July, Aug. v. v. Flowers small, golden-yellow.
13. *O. subpubescens; caule pumilo subsimplici, foliis lanceolato-oblongis obtusiusculis integris, floribus ad summitatem axillaribus, capsulis sessilibus clavato-turbinate subæqualiter 8-gonis.* — *Mich. fl. amer. 1. p. 225.*  
On rocks near Lake Mistassins. *Michaux.* On high mountains of Pennsylvania and Virginia. 4. July, Aug. v. v. Plant upright, flowers small.
14. *O. scap's radicalibus unifloris, foliis oblongo-lanceolatis repando-denticulatis decurrentibus, petalis obcordatis, staminibus corolla brevioribus.*  
On the falls of the Missouri. *M. Lewis.* 4. July.  
v. s. Flowers large, purple, with dark veins; resembles *O. daulis* Cavanill.  
All the species, excepting the last one, have yellow flowers.

330. OXYCOCCUS. *Pers. syn. 1. p. 419.*

1. *O. repens; foliis ovalibus integerrimis margine revolutis subacutis glabris subtus albicantibus, pedicellis elongatis, corollæ laciniis ovalibus.*  
*O. palustris. Pers. syn. 1. p. 419.*  
*Vaccinium Oxycoccus a. ovalifolius. Mich. fl. amer. 1. p. 228.*  
Vaccinium Oxycoccus. *Willd. sp. pl. 2. p. 354.*  
Icon. *Fl. dan. 80. Lam. illustr. t. 286. f. 3.*  
In bogs and mosses on the mountains, frequent: Canada to Pennsylvania. 5. May—July. v. v. A small evergreen creeping plant; flowers red; berries light purple, smaller than the following species.
2. *O. repens; caulibus adscendentibus, foliis oblongis integrerrimis planiusculis obtusis glabris subtus albicantibus, pedicellis elongatis, corollæ laciniis lanceolatis.* — *Pers. syn. 1. p. 419.*

- Vaccinium macrocarpon.* Willd. sp. pl. 2. p. 355.  
*Vaccinium Oxyccus* β. oblongifolius. Mich. fl. amer. 1. p. 228.  
 Icon. Wangh. amer. t. 30. f. 67. Pluk. alm. t. 326. f. 6.  
 In bogs, principally on sandy soil and high mountains, frequent: Canada to Virginia. ♂. May—July. v. v. Berries larger, bright red, known by the name of *Cranberries*, as a very agreeable tart, and collected in great abundance for that purpose.
- erectus.* 3. O. erectus; foliis ovalibus acuminatis serrulatis ciliatisque, pedicellis axillaribus, corolla longa demum revoluta.
- Vaccinium erythrocarpum.* Mich. fl. amer. 1. p. 227.  
 On high mountains of Virginia and Carolina. ♂. June. v. v. Berries scarlet and quite transparent, of an exquisite taste.
- ferruginea.* 340. MENZIESIA. Smith ic. inedit. 56.
1. M. foliis obovato-lanceolatis subtus extra nervos glabris, calycibus sub-4-fidis, floribus urceolatis octandris.—Willd. sp. pl. 2. p. 355.  
*M. urceolaris.* Parad. lond. 44.  
 Icon. Smith ic. inedit. 1. t. 56.  
 On the Columbia river. M. Lewis. ♂. June. v. s. in Herb. Lambert. A considerable tall shrub; flowers of a ferruginous colour, as in the following species. Previous to my seeing Mr. Salisbury's account and the specimens in the herbarium of A. B. Lambert, Esq. I considered this and the following species to be the same, though I had seen specimens of the present one in the collection of M. Lewis, Esq.
- globularis.* 2. M. foliis lanceolatis subtus glaucis extra nervos pubescens, calycibus 4-fidis, floribus globosis octandris.—Salisb. in parad. lond. 44.  
*M. Smithii.* Mich. fl. amer. 1. p. 235.  
*Azalea pilosa.* Lam. illustr. 1. p. ....  
 Icon. Parad. lond. 44.  
 On high mountains of Virginia and Carolina; plentifully on the Cacapon mountains near Winchester, Virginia. ♂. May, June. v. v. Not above four feet high. Flowers yellowish-brown.
- empetrifor-  
mis.* 3. M. foliis linearibus serrulatis subtus concavis, pedunculis terminalibus aggregatis, floribus campanulatis decan-

dris, calyci  
p. 380.

On the north  
mountains  
M. Lewis.

4. M. foliis spars  
denticulatis  
floris, florit  
tissimis.—S

Andromeda ta  
Erica cærulea  
Phyllodoce ta  
On the Whit  
On the nor  
♂. July.

1. D. foliis ovat  
noscis, peti  
sp. pl. 4. 1  
Icon. Mill. i  
In woods an  
May. v.  
low; fruit  
of a com  
taste, wh  
before tha  
not eatabl
2. D. foliis obl  
longis, fr  
In the lowe  
♂. April  
*forestiers*  
to take it  
bescence  
the fruit

1. A. foliis pa  
ter incise  
bellis se  
bris.—H

355. dris, calycibus obtusis. *Smith in linn. trans.* 10.  
fl. amer. 1. p. 380.

On the north-west coast. *Menzies.* On the Rocky-mountains and near the mouth of Columbia river. *M. Lewis.* h. July. v. s. Flowers pale red.

4. M. foliis sparsis confertis linearibus obtusis cartilagineo-denticulatis, pedunculis terminalibus aggregatis unifloris, floribus campanulatis 10-andris, calycibus acutissimis.—*Swartz in linn. trans.* 10. p. 377. t. 30. f. 1.

*Andromeda taxifolia.* *Pall. fl. ross.* t. 72. f. 2.

*Erica cærulea.* *Willd. sp. pl.* 2. p. 393.

*Phylodoce taxifolia.* *Parad. lond.* 36.

On the White Hills of New Hampshire. *Prof. Peck.*

On the north-west coast and Labrador. *Herb. Banks.*

h. July. v. s. Flowers red.

*cærulea.*

### 341. DIOSPYROS. *Gen. pl.* 1598.

1. D. foliis ovato-oblongis acuminatis glabris reticulato-venosis, petiolis pubescentibus, gemmis glabris.—*Willd. sp. pl.* 4. p. 1107.

*Icon. Mill. ic.* 126. *Catesb. car.* 2. t. 76. *Mich. arb.* 12. In woods and old fields: New York to Louisiana. h. May. v. v. A middle-sized tree; flowers pale yellow; fruit known by the name of *Persimon*, the size of a common plum, golden-yellow, of an agreeable taste, when perfectly ripe, but extremely astringent before that time. In the Northern States the fruits are not eatable till the frost has mellowed them.

2. D. foliis oblongis acutis subtus pubescentibus, petiolis longis, fructibus oligospermis.

In the lower counties of Virginia, Carolina and Georgia.

h. April. v. v. Though Michaux in his *Arbres forestiers* considers this only as a variety, I am inclined to take it as a distinct species; not only shape and pubescence of the leaf, but difference in the structure of the fruit and seeds, indicate it sufficiently.

*virginiana.*

*pubescens.*

### 342. ACER. *Gen. pl.* 1590.

1. A. foliis palmato subquinquelobis basi cordatis inæqualiter inciso-dentatis subtus glaucis, incisuris acutis, umbellis sessilibus, pedicellis elongatis, germinibus glabris.—*Willd. sp. pl.* 4. p. 984.

*rubrum.*

Icon. *Mich. arb.* 14. *Schmidt arb.* 1. t. 6. *Catesb. car.* 1. t. 62.

In low woods: Canada to Florida.  $\frac{1}{2}$ . April, May. v. v. A middle sized tree: flowers and seeds red; commonly known under the name of *Red-maple* or *Swamp-maple*.

*dasycarpum.* 2. A. foliis palmato-5-lobis basi truncatis inaequaliter inciso-dentatis subtus glabris glaucisque, sinibus obtusis, floribus conglomeratis, pedicellis brevissimis, germinibus tomentosis.—*Willd. sp. pl.* 4. p. 985.

A. eriocarpum. *Mich. fl. amer.* 2. p. 253.

A. tomentosum. *Hort. paris.*

Icon. *Mich. arb.* 13. *Schmidt arb.* 1. t. 7.

On the banks of rivers: New England to Georgia.  $\frac{1}{2}$ . April, May. v. v. A large tree; flowers and seed greenish-yellow; known by the name of *White-* or *Soft-maple*.

*barbatum.* 3. A. foliis ovato-cordatis breviter trilobis inaequaliter serratis subtus ad nervos pubescentibus glaucisque, pedunculis pilosis masculis ramosis, femineis simplicibus, calycibus intus barbatis, capsulae alis erectis.—*Mich. fl. amer.* 2. p. 252.

A. carolinianum, *Walt. fl. car.* 251.

In deep pine and cedar swamps: New Jersey to Carolina.  $\frac{1}{2}$ . April, May. v. v. A small tree; flowers pale-green; leaves small.

*saccharinum.* 4. A. foliis palmato-5-lobis basi subcordatis acuminatis sinuato-dentatis subtus glaucis, pedunculis corymbosis nutantibus.—*Willd. sp. pl.* 4. p. 985.

Icon. *Mich. arb.* 15. *Schmidt arb.* 1. t. 8.

In low rich valleys: Canada to Pennsylvania.  $\frac{1}{2}$ . April, May. v. v. A large timber tree; flowers yellow. The sugar prepared from the sap of this tree is one of the greatest conveniences for the inhabitants of the western countries, is equal to any other sugar, and procured with little trouble.

*nigrum.* 5. A. foliis palmato-5-lobis sinu clauso cordatis, lobis divaricatis sinuato-subdentatis subtus pubescentibus, floribus corymbosis, capsulis turgide subglobosis.—*Mich. arb.* p. 238.

Icon. *Mich. arb.* 16.

On mountain lands: New York to Carolina.  $\frac{1}{2}$ . April, May. v. v. This large tree produces sugar similar

to the foregoing species where *maple*.

6. A. foliis digitatis trilobatis repens erectis, tressimis.

On the great river  $\frac{1}{2}$ . April. In any other kind of long compound *Arbor grandis*.

*Folia magnificæ* Bracteæ geniculæ, sericeæ elongatæ. Folia, glabri. rioribus vix medium batissimum. filamentis ciliis.

7. A. foliis orbiculatis acute-dentatis pilosis.

On the great river  $\frac{1}{2}$ . v. s. s. the size of

8. A. foliis inferioribus datus argutus dulcis.—*Mich.*

*A. pensylvanica* *A. canadense* *Icon. Mich.* t. 12. f. 28

In mountainous regions v. v. A. known by

9. A. foliis subcordatis, pubescens, centibus, p. 988.

*A. pensylvanica* *A. parviflora* *A. spicatum*. *Icon. Schm.*

to the foregoing species, and occupies the same situations where the other is not found; called *Black-maple*.

6. A. foliis digitato-5-lobis, sinubus rotundatis, lobis sub-trilobatis repando-dentatis subtus pubescentibus, racemis erectis, filamentis 9. hirsutis, germinibus hirsutissimis.

On the great rapids of the Columbia river. *M. Lewis.*

$\frac{1}{2}$ . April. v. s. A very tall tree; leaves larger than any other known species; flowers greenish-yellow, in long compound racemes.

*Arbor grandis. Ramuli juniores purpurascentes, glabri.*

*Folia magna longe petiolata, juniora pubescens. Bracteæ gemmarum longissimæ, lineares, apice incisæ, sericeo-villosæ. Racemi terminales, erecti, elongati. Pedicelli inferne aggregati, superne solitarii, glabri. Calyx 9-fidus: laciniis obovatis, interioribus vix latioribus. Filamenta 9. subulata, supra medium bârbata, calyce vix longiora. Germen hirsutissimum. Stylus glaber. Flores masculi decandri: filamentis calyce duplo longioribus.*

7. A. foliis orbiculatis basi subcordatis 7-lobis inæqualiter acute-dentatis utrinque glabris, nervis venisque ad axillas pilosis.

On the great rapids of Columbia river. *M. Lewis.*

$\frac{1}{2}$ . v. s. s. fl. This beautiful species has leaves of the size of *A. rubrum*.

8. A. foliis inferne rotundatis, superne acuminato-tricuspidatis argute-serratis glabris, racemis simplicibus pendulis.—*Mich. fl. amer. 2. p. 252.*

*A. pensylvanicum. Willd. sp. pl. 4. p. 989.*

*A. canadense. Marsh. arb. . . .*

*Icon. Mich. arb. 17. Schmidt arb. 10. Wangh. amer. t. 12. f. 28.*

In mountain woods: Canada to Virginia.  $\frac{1}{2}$ . May.

v. v. A small tree with an elegant striped bark; known by the name of *Striped-maple* or *Moose wood*.

9. A. foliis subquinqüelobis acutis dentatis subtus pubescentibus, racemis compositis erectis.—*Willd. sp. pl. 4. p. 988.*

*A. pensylvanicum. Wangh. amer. 82. t. 12. f. 30.*

*A. parviflorum. Ehrh. beitr. 4. p. 25.*

*A. spicatum. Lam. encycl. 2. p. 377.*

*Icon. Schmidt arb. 11. Du Roi harbk. 1. t. 2.*

*macrophyllum.*

*circinatum.*

*striatum.*

*montanum.*

## OCTANDRIA MONOGYNIA. Acer.

In Canada and on the Allegany mountains; frequent.  
½ April, May. v. v. Flowers very small, greenish-yellow.

*Negundo.*

10. A. foliis pinnatis ternatisve inaequaliter serratis, floribus dioicis. Willd. sp. pl. 4. p. 992.  
Icon. Mich. arb. 18. Schmidt arb. 12. Wangh. amer. t. 12. f. 29.

On the banks of rivers: Pennsylvania to Carolina. ½ May, June. v. v. A large tree, commonly called *Box Elder* or *Ash-leaved Maple*.

## 343. DIRCA. Gen. pl. 665.

*palustris.*

1. Dirca. Willd. sp. pl. 2. p. 424.

Icon. Duham. arb. 1. t. 212. Amoen. acad. 3. t. 1. f. 7.  
In shady and boggy woods: New York to Carolina.  
½ March, April. v. v. A low shrub; flowers yellow; known by the name of *Leather-wood*, on account of the extraordinary toughness of its branches.

## 344. JEFFERSONIA. Barton in act. philad. Mich. fl. amer. 1. p. 236.

*diphylla.*

1. Jeffersonia. Bart. l. c.

J. Bartonis. Mich. fl. amer. 1. p. 237.

Podophyllum diphyllum. Willd. sp. pl. 2. p. 1141.

On the side of hills, in rich soil, particularly limestone: Virginia and Tennessee; plentifully about Harper's Ferry and the Sweet-springs, Virginia. ¼ April, May. v. v. Flowers white, resembling those of *Sanguinaria*; the capsule is covered with a lid, which opens with elasticity and scatters the ripe seeds.

## 345. PISTIA. Gen. pl. 1112.

*spathulata.*

1. P. foliis in petiolum abrupte angustatis superne dilatatis rotundato-obtusis. Mich. fl. amer. 2. p. 162.

In Carolina. Michaux. +. Flowers white, axillary.

346. CHI

1. C. foliis oppo  
levissime c  
Icon. Pl. da  
On rocks in t  
situations :  
Flowers br  
if compare

347.

1. P. floribus s  
branaceis,  
volutis ca  
p. 449.

Icon. Cam.  
β. P. maritimu  
On the sand

Aug. v.  
white or

2. P. floribus  
scabris,  
herbaceo

Icon. Engl.  
α. P. foliis pu  
p. 237.

β. P. foliis lat  
l. c.

On pasture  
○. Apr  
small.

3. P. floribus  
abbrevia

## II. DIGYNIA.

## 346. CHRYSOSPLENIUM. Gen. pl. 763.

1. C. foliis oppositis subrotundis basi in petiolum attenuatis levissime crenatis.—*Willd. sp. pl. 2. p. 638.* *oppositifoli-*  
*um.*

Icon. *Pl. dan. 365.* *Engl. bot. 490.*  
On rocks in rivulets and springs, near cascades; in shady situations: Canada to Carolina. 2. May. v. v. Flowers brownish-yellow. Probably a distinct species, if compared in the living plants.

## III. TRIGYNIA.

## 347. POLYGONUM. Gen. pl. 677.

\* *Floribus axillaribus.*

1. P. floribus 8-andris 3-gynis, ochreis obtusis laceris membranaceis, foliis linear-lanceolatis acutis margine revolutis carnosis, caule suffruticoso.—*Willd. sp. pl. 2. p. 449.*

Icon. *Cam. epit. 691.*  
β. *P. maritimum floribus carneis speciosis. Clayt. n. 497.*  
On the sandy beach: Pennsylvania and Virginia. h. 4.  
Aug. v. v. A small prostrate evergreen plant, with white or rose-coloured flowers.

2. P. floribus 8-andris 3-gynis, foliis lanceolatis margine scabris, ochreis nervis remotis, caule procumbente, herbaceo. *Willd. sp. pl. 2. p. 449.*

Icon. *Engl. bot. 1252.* *Mart. fl. rust. t. 91.*  
α. *P. foliis pusillis lanceolato-oblongis. Mich. fl. amer. 1. p. 237.*

β. *P. foliis lato-ovalibus obtusis quasi buxifolium. Mich. l. c.*

◆ On pastures and road sides, in gardens, &c. common.  
◎. April—Oct. v. v. Flowers sessile, white, very small.

3. P. floribus 3-andris 3-gynis, foliis lanceolatis, ochreis abbreviatis margine laceris, ramis interrumpere multiflo-

*marinum.*

*roseum.*

*aviculare.*

*angustifoli-*  
*um.*

*latifolium.*

*ramosissi-*  
*mum.*

## OCTANDRIA TRIGYNIA. Polygonum.

OCTANDRIA

ris, caulibus stricte-ramosissimis striatis.—*Mich. fl. amer.* 1. p. 237.

P. erectum: *Willd. sp. pl. 2. p. 450?*

In cultivated grounds: Pennsylvania and Kentucky. ☀. June—Aug. v. v. Flowers on short pedicels, greenish-white.

4. P. pumilum; foliis longe-linearibus strictis acuminatis, ochreis tubulosis apice villosis, caule gracili erecto ramoso acutangulo, floribus alternis subsolitariis.—*Mich. fl. amer.* 1. p. 238.

P. barbatum: *Walt. fl. car. 131?*

In sandy fields and pine-woods: Canada, New Jersey, and Carolina? ☀. July, Aug. v. v. Not above six inches high, very slender; flowers small, white.

\*\* *Spica gracili.*

*Hydropiperoides.*

5. P. floribus 8-andris semi-3-gynis, foliis lanceolatis glabris, ochreis laxis glabris apice ciliatis maculatis, spicis filiformibus debilibus subcernuis, bracteis remotiusculae alternis.—*Mich. fl. amer.* 1. p. 238. sub *P. Hydropiperide.*

In inundated and exsiccated places; common: Canada to Carolina. ☀. Aug. Sept. v. v. Flowers white; taste and appearance of *P. Hydropiper*, but different in the flowers.

6. P. floribus 8-andris semi-3-gynis, foliis anguste lanceolatis subhirsutis, ochreis hirsutis promisso ciliatis, spicis linearibus, bracteis ciliatis subimbricatis.—*Pers. syn. 1. p. 446.*

*P. hydropiperoides.* *Mich. fl. amer.* 1. p. 239.

In inundated places, along ditches and ponds: Pennsylvania to Carolina. ☀. June—Sept. v. v. Flowers pale purple; leaves not acrid.

*mite.*

7. P. floribus 8-andris semi-3-gynis approximatis, foliis sessilibus lanceolatis, pedunculo elongato distachyo, spicis linearibus, caule adscendente hirsuto.—*Walt. fl. car. 132.* *Mich. fl. amer.* 1. p. 239.

In low wet places and in ditches: Lower Carolina and Virginia. ☀. July—Sept. v. v.

*hirsutum.*

8. P. floribus 5-andris 2-gynis inaequalibus, foliis lato-ovalibus, spicis longissime virgatis, floribus remotis.—*Willd. sp. pl. 2. p. 442.*

In shady woods: Canada to Florida. ¼. June, July. v. v.

*virginianum.*

9. P. caule simplici  
tiolatis, bracte-

In low ground  
*Quamash*-flat-  
white tinged v-

*Caulis* teres, gla-  
luni decurrent  
subtus glauca,  
longa. *Bracte-*  
valvula exten-  
*Calyx* subinflo-  
*Filamenta* 7—

10. P. caule simplici  
bus glabris m-  
ovatis acumin-

In Canada. ¼  
This almost  
cimens comit  
in the collec-  
Labrador.

11. P. floribus 5-  
ochreis trunc-  
429.

*P. amphibium.*

α. P. foliis ovato-e-  
β. P. foliis ovato-  
*fl. car. 131?*

α. In lakes and  
overflowed p-  
July, Aug.

12. P. floribus 8-an-  
culis hispidis  
que, caule  
p. 448.

On the edge  
grounds; co-  
cipally in the  
Flowers larg-

13. P. floribus 6-  
erectis, pedi-  
glabriusculis  
Icon. *Fl. dan.*

\*\*\* *Spica confertiflora*.

9. P. caule simplici monostachyo, foliis ellipticis planis petiolatis, bracteis unifloris 2-3-valvis. *bistortoides*.  
In low grounds on the banks of the Missouri, called Quamash-flats. M. Lewis. 4. June. v. s. Flowers white tinged with red.  
*Caulis* teres, glaber. *Folia* elliptico-oblonga, in petiolum decurrentia, integerrima, plana, utrinque glabra, subitus glauca, superiora linearia, sessilia. *Spica* oblonga. *Bracteæ* membranaceæ, unifloræ, 2-3-valves; valvula exterior subrotunda, acuminata, carinata. *Calyx* subinsundibuliformis: laciniis oblongis acutis. *Filamenta* 7—8. corolla longiora. *Styli* 3.
10. P. caule simplici monostachyo, foliis lanceolato-lineari- bus glabris margine revolutis, spica linearis, bracteis β. subacaule. *viviparum*.  
In Canada. 4. June, July. v. s. in *Herb. Lambert*. This almost stemless variety I have seen in dried specimens coming from Lower Canada. The specimen in the collection of A. B. Lambert, Esq. came from Labrador.
11. P. floribus 5-andris semi-2-gynis, spica cylindracea, ochreis truncatis glabris, foliis ovatis. *Willd. enum.* *coccineum*.  
429. P. amphibium. *Mich. fl. amer.* 1. p. 240.  
α. P. foliis ovato-ellipticis obtusis.  
β. P. foliis ovato-oblongis acuminatis. P. *Bistorta*. *Walt.* *fl. car.* 131?  
α. In lakes and ponds: Canada to Pennsylvania. β. In overflowed places on the Ohio and Mississippi. 4. July, Aug. v. v. Flowers dark red.
12. P. floribus 8-andris semdigynis, spicis oblongis, pedunculis hispidis, foliis lanceolatis, ochreis glabris nudisque, caule tumide-geniculato.—*Willd. sp. pl.* 2. p. 448.  
On the edge of ponds and ditches, in overflowed grounds; common: New York to Carolina, and principally in the western counties. ⊖. June—Aug. v. v. Flowers large, rose-coloured.
13. P. floribus 6-andris semi-2-gynis, spicis ovato-oblongis erectis, pedunculis lœvibus, foliis lanceolatis, ochreis glabriusculis apice ciliatis.—*Willd. sp. pl.* 2. p. 446.  
Icon. *Fl. dan.* 702. *Engl. bot.* 756. *Persicaria*.

## OCTANDR

272

## OCTANDRIA TRIGYNIA. Polygonum.

In overflowed and wet places; common. ☽. July, Aug. v. v. Flowers rose-coloured; leaves generally marked with a black spot in shape of a half moon; known by the name of *Ladies-thumb*.

*orientale*. 14. P. floribus 7-andris 2-gynis, foliis ovatis, caule erecto, ochreis hirtis hypocrateiformibus. *Willd. sp. pl. 2. p. 448.*

Icon. *Bot. mag. 213. Mill. ic. 201.*

Near gardens and in old fields: Virginia. ☽. July, Aug. v. v. Sometimes six feet high; flowers large, crimson.

\*\*\*\* *Floribus paniculatis.*

*articulatum*. 15. P. floribus 8-andris 3-gynis, spicis paniculatis filiformibus, floribus solitariis pedunculatis, bracteis imbricatis truncatis, foliis linearibus, ochreis vaginalibus truncatis, caule herbaceo.—*Willd. sp. pl. 2. p. 450.*

In sandy barren fields: Canada to New Jersey. ☽. July, Aug. v. v. A handsome, delicate plant, about a foot high; flowers rose-coloured, mixed with white.

*polygamum*. 16. P. floribus 8-andris 3-gynis, spicis paniculatis, floribus solitariis, calycis laciniis obovatis, foliis spathulatis, ochreis integris latere altero lanceolatis, caule fruticoso. *Vent. hort. cels. 65.*

*Polygonella parvifolia. Mich. fl. amer. 2. p. 241.*

In the barrens of Carolina. ♂. July. +. Flowers white. I strongly suspect this to be only a variety of *P. articulatum*, as the description and figure of Venetian strongly indicate; but the northern plant is certainly only an annual.

\*\*\*\*\* *Floribus subracemosis; foliis basi incisis vel cordatis.*

*sagittatum*. 17. P. floribus 8-andris semi-3-gynis capitatis, foliis sagittatis, caule retrorsum aculeato.—*Willd. sp. pl. 2. p. 453.*

Icon. *Gmel. sib. 3. t. 13. f. 2. Hort. cliff. t. 12. Pluk. mant. t. 398. f. 5.*

In low wet meadows and ditches, in open situations: New York to Florida. ☽. June—Aug. v. v. Flowers white.

*arifolium*. 18. P. floribus 6-andris semi-2-gynis distinctis, spicis paucifloris, foliis hastatis, caule retrorsum aculeato.—*Willd. sp. pl. 2. p. 453.*

Icon. *Pluk. amalh. t. 398. f. 3.*

In low shady p  
New York to  
white, tinged

19. P. floribus 8-and  
cordato-sagitta  
angulis æquali  
Icon. *Engl. bot.*  
*Common Buck-w*  
is frequently fo  
in a wild state

20. P. floribus 8-and  
cordatis, caule  
calycinis obtus  
Icon. *Fl. dan. 74*  
In cultivated gro  
rolina. ☽. J

21. P. floribus 8-and  
subacutis basi  
caule angulos  
laciniis calycis  
p. 241.

In hedges and fa  
v. v.

22. P. floribus 8-and  
truncatis nudis  
tiferis tripteris  
Icon. *Pluk. alm*  
In shady woods  
Aug. v. v.  
considerable l

## 348. BRUN

1. Brunnichia. *R*  
*Rajania ovata.* *I*  
Icon. *Gært. car*  
In wet shady pl  
Georgia; in  
A smooth vi  
in one-sided

## 349. CARI

1. C. glabrum; f  
deis.—*Willd*  
VOL. I.

- In low shady places, principally on the mountains : New York to Carolina. ⊖ July, Aug. v. v. Flowers white, tinged with red.
19. P. floribus 8-andris 3-gynis, racemis paniculatis, foliis cordato-sagittatis, caule erectiusculo inermi, seminum angulis æqualibus.—*Willd. sp. pl. 2. p. 455.*  
Icon. *Engl. bot. 1044.*  
*Common Buck-wheat*, though not originally indigenous, is frequently found in old fields and woods, as it were in a wild state.
20. P. floribus 8-andris semi-3-gynis, foliis oblongis hastato-cordatis, caule volubili angulato asperiusculo, laciiniis calycinis obtuse carinatis.—*Willd. sp. pl. 2. p. 455.*  
Icon. *Fl. dan. 744.*  
In cultivated grounds, along hedges : Pennsylvania to Carolina. ⊖ July, Aug. v. v.
21. P. floribus 8-andris semi-3-gynis, foliis cordatis, stipulis subacutis basi extrorsum serie ciliorum circumdati, caule anguloso prostrato aut scandente asperiusculo, laciiniis calycis obtuse carinatis.—*Mich. fl. amer. 1. p. 241.*  
Icon. *Pluk. alm. t. 177. f. 7. Sloan. hist. 1. t. 90. f. 1.*  
In hedges and fields : Canada to New York. ⊖ July, v. v.
22. P. floribus 8-andris 3-gynis, foliis late-cordatis, stipulis truncatis nudis, caule volubili glabro, calycibus fructiferis tripteris.—*Willd. sp. pl. 2. p. 456.*  
Icon. *Pluk. alm. t. 177. f. 7. Sloan. hist. 1. t. 90. f. 1.*  
In shady woods : New York to Kentucky. ♀ July, Aug. v. v. Stems generally red, climbing to a considerable height.

348. BRUNNICHIA. *Gært. Gen. pl. 777.*

1. Brunnichia. *Willd. sp. pl. 2. p. 731.*  
*Rajania ovata. Walt. fl. car. 247.*  
Icon. *Gært. carp. 1. t. 45. f. 2.*  
In wet shady places, climbing among bushes : islands of Georgia ; in Tennessee, &c. ♀ June, July. v. v.  
A smooth vine ; leaves cordate, acuminate ; flowers in one-sided racemes.

*cirrhosa.*349. CARDIOSPERMUM. *Gen. pl. 680.*

1. C. glabrum ; foliolis inciso-lobatis, imparibus rhomboideis.—*Willd. sp. pl. 2. p. 467.*

Icon. *Bot. mag.* 1049.

On the banks of the Mississippi, near Kaskaskias. ○.

July. v. v. A small climbing plant; flowers small, herbaceous.

### 350. SAPINDUS. *Gen. pl.* 681.

*Saponaria.*

1. *S. foliis glabris abrupte pinnatis, foliolis ovali-lanceolatis, rachi alata, fructibus sphæricis.*—*Willd. sp. pl.* 2. p. 468.

Icon. *Comm. hort. t.* 94.

On the sea-coast of Georgia and Carolina: probably introduced. h. v. v. s. fl. A small tree.

## IX. E

## IV. TETRAGYNYIA.

### 351. MYRIOPHYLLUM. *Gen. pl.* 1440.

*spicatum.*

1. *M. foliis omnibus pinnatis capillaceis, spica nuda interrupta, floribus masculis polyandris.*—*Willd. sp. pl.* 4. p. 406.

Icon. *Fl. dan.* 681. *Engl. bot.* 83.

In slow-flowing waters and ponds: Canada to New Jersey. 4. July, Aug. v. v. A sub-aquatic small plant.

*verticillatum.*

2. *M. foliis pinnatis capillaceis, superioribus pectinato-pinnatifidis, floribus omnibus axillaribus verticillatis, inferioribus femineis, superioribus masculis aut hermaphroditis 8-andris.*—*Willd. sp. pl.* 4. p. 407.

Icon. *Fl. dan.* 1046. *Engl. bot.* 218.

In stagnant waters and ditches: Canada to Pennsylvania. 4. July, Aug. v. v.

*scabratum.*

3. *M. foliis pinnatifidis, floribus omnibus verticillatis axillaribus, superioribus masculis 4-andris, inferioribus femineis, fructu 8-angulato.*—*Mich. fl. amer.* 2. p. 190.

*Potamogeton pinnatum.* *Walt. fl. car.* 90.

In stagnant waters of Carolina and Georgia. 4. July. v. s.

*heterophyl-*  
*lum.*

4. *M. foliis inferioribus capillaceo-pinnatis, superioribus floralibus ovalibus argute serratis, floribus masculis 6-andris.* *Mich. fl. amer.* 2. p. 191.

*Potamogeton verticillatum.* *Walt. fl. car.* 90.

In the waters of Carolina and Georgia. 4. July. v. s. in *Herb. Walt.*

352. LAURUS.  
bisetis, ger-  
rum 3. alte-  
*Polygama.*

353. ERIOGON-  
*Cor. O.* S-  
lucrum car-

354. PLEEA. C  
trigona, 3-  
gini valvula-

## I.

### 352.

1. *L. foliis pere-  
bris nitidis,  
bus, calycis  
libus decid-  
p. 244.*

*Laurus baccis  
tibus.* *Clay-*

*Icon. Catesb.*

On the sea-  
red pedunc-

## IX. ENNEANDRIA.

---

### I. MONOGYNIA.

352. LAURUS. *Cal. 4*—6-partitus. *Nectar. glandulis 3*,  
bisetis, germen cingentibus. *Filam. interiora quo-*  
*rūm 3*. alterna sterilia glandulifera. *Drupa 1-sperma.*  
*Polygama.*
353. ERIOGONUM. *Cal. subcampanulatus, 6-fidus.*  
*Cor. O. Sem. 1. triquetrum, calyce tectum. Invo-*  
*lucrum campanulatum.*

### II. TRIGYNIA.

354. PLEEA. *Cor. 6-partita, patens. Caps. subrotundo-*  
*trigona, 3-locularis. Sem. numerosa, oblonga, mar-*  
*gini valvularum adnata. Spica spathis 1-floris.*
- 

### I. MONOGYNIA.

#### 352. LAURUS. *Gen. pl. 698.*

1. *L. foliis perennantibus ovato-lanceolatis ramulisque gla-*  
*bris nitidis, racemis corymboso-paniculatis terminali-*  
*bus, calycis patuli laciniis oblongis obtusis subæqua-*  
*libus deciduis, baccis ovatis.—Mich. fl. amer. 1.*  
*p. 244.*
- Laurus baccis atro-purpureis, pedunculis rubris insiden-*  
*tibus. Clayt. n. 485.*
- Icon. Catesb. car. 2. t. 28.*
- On the sea-coast of Georgia and Florida. 1<sup>st</sup>. May.*  
*v. v. s. fl. Flowers white; berries black, on thick*  
*red peduncles.*

*carolinensis.* 2. L. foliis perennantibus ovali-lanceolatis opacis subtus glaucis, pedunculis simplicibus, fasciculo subcapitato paucifloro terminatis, calycis laciniis exterioribus duplo brevioribus, baccis globosis.—*Mich. fl. amer.* 1. p. 245.

L. *Borbonia. Hortul.*

L. foliis acuminatis, flore albicante, baccis cæruleis, pedunculis rubris insidentibus. *Clayt. ult.*

Icon. *Catesb. car. t. t. 63. Mich. arb. . . . .*

$\alpha$ . L. foliis subtus glabris.

$\beta$ . L. foliis subtus pubescentibus.

$\gamma$ . L. foliis obovatis obtusis.

In deep cedar and cypress swamps: Virginia to Louisiana.  $\frac{1}{2}$ . May. v. v. A small tree; flowers greenish-yellow; berries dark-blue, on red peduncles; known by the name of *Red Bay*: the wood is very valuable.

*Benzoin.* 3. L. ramis virgatis sub floratione aphyllis, foliis deciduis cuneato-ovalibus subtus albicantibus subpubescentibus, floribus glomerato-umbellatis, gemmis pedicellisque glabris.—*Willd. sp. pl. 2. p. 485.*

L. *Pseudo-Benzoin. Mich. fl. amer.* 1. p. 243.

Icon. *Comm. hort. t. 97.*

In low grounds and on the banks of rivulets: Canada to Florida.  $\frac{1}{2}$ . March, April. v. v. A large shrub; flowers yellowish-green; berries scarlet; known by the name of *Spice-wood*.

*Diospyrus.* 4. L. humilis, virgata, nudiflora; foliis deciduis oblongo-ovalibus subtus venosis subtomentosis, floribus glomerato-umbellatis, gemmis pedicellisque villosis.—*Mich. fl. amer.* 1. p. 243.

L. *mellissæfolia. Walt. fl. car. 134.*

Icon. *Bot. mag. 1470.*

In low grounds and near rivers: Virginia and Carolina.  $\frac{1}{2}$ . April. v. v. A low shrub, resembling the preceding very much; flowers greenish-yellow; berries small, scarlet or purple.

*geniculata.* 5. L. ramis divaricatis flexuosis, foliis deciduis ovalibus obtusiusculis glabris, basi subtus barbatis, sub floratione nullis, umbellulis terminalibus paucifloris, antheris 4-locularibus.—*Mich. fl. amer.* 1. p. 244. *Walt. fl. car. 133.*

L. *aestivalis. Willd. sp. pl. 2. p. 484.*

L. foliis lanceolatis enervibus annuis, *Gronov. virg.* 159.

EN  
Icon. Bot.  
In low gro  
 $\frac{1}{2}$ . Apr  
yellow;

6. L. foliis in  
Icon. Tre  
*Mich. ar*  
In woods a  
in old fir  
v. v. I  
tree, in  
large siz  
trees tw  
wards it  
stragglin

353. ER

1. E. caule ei  
solitariis  
tomento  
quaterni  
*fl. amer.*

Icon. *Mi*  
In pine-ba  
The wh  
the leav  
coat; fl  
or invol

2. E. caule s  
umbella  
laciniis  
petiolat  
On the p  
v. s. in  
well as  
loose te  
vered w  
low. I  
discover  
give a c

Icon. *Bot. mag.* 1471.

In low grounds, near rivers and on islands : Carolina.

↳ April. *v. v.*; *v. s.* in *Herb. Gronov.* Flowers yellow; berries globous, scarlet.

6. *L. foliis integris lobatisque.* *Willd. sp. pl. 2. p. 485.*

Icon. *Trew chret. t. 69 & 70.* *Catesb. car. 1. t. 55.*

*Mich. arb. t. . . .*

In woods and along the banks of large rivers, frequent in old fields : Canada to Florida. ↳ March, April. *v. v.* Flowers yellow; berries blue. This valuable tree, in respect to its medicinal properties, grows to a large size in the southern regions, where I have seen trees two feet in diameter; but from Virginia northwards it decreases in size, until it becomes a mere straggling shrub.

*Sassafras.*

353. ERIOGONUM. *Mich. fl. amer. 1. p. 246.*

1. *E. caule erecto dichotomo, florum fasciculis axillaribus solitariis sessilibus, calycis laciniis subrotundis obtusis tomentosis, foliis ad genicula caulis sessilibus ternis quaternis cuneato-obovalibus supra glabris.* — *Mich. fl. amer. 1. p. 246.*

Icon. *Mich. l. c. t. 24.*

In pine-barrens of Carolina and Georgia. 4. July. *v. v.* The whole of this plant, excepting the upper side of the leaves, is covered with a close tomentum or woolly coat; flowers from twelve to fifteen collected in a cup or involucrum, white.

*tomentosum.*

2. *E. caule simplici nudo, florum fasciculis terminalibus umbellatis pedunculatis, umbella involucrata, calycis laciniis oblongis acutis sericeo-lanatis, foliis radicalibus petiolatis lanceolato-oblongis supra villosis.*

On the prairies of the Missouri. *Nuttall.* 4. July.

*v. s.* in *Herb. Nuttall.* The whole of this plant, as well as the foregoing, is tomentose, but of a more loose texture, approaching to villous; the calyx is covered with long white silky down; flowers bright yellow. In the same country a third species has been discovered, of which I have not materials enough to give a description.

*sericeum.*

cis subtus  
subcapitato  
ioribus du-  
l. amer. 1.

eruleis, pe-

nia to Loui-  
wers green-  
peduncles;  
ood is very

iiis deciduis  
pubescenti-  
nis pedicel-

3.

: Canada to  
arge shrub;  
known by

uis oblongo-  
ribus glome-  
rosis.—*Mich.*

and Carolina.  
ding the pre-  
ow; berries

ovalibus ob-  
ub floratione  
ris, antheris  
*Walt. fl.*

*virg. 159.*

## II. TRIYGYNIA.

354. PLEEA. *Mich. fl. amer.* 1. p. 247.*tenuisolia.*

1. *P. glaberrima*; foliis angustissime ensiformibus sensim acutissimis; spicæ vaginis 1-floris.—*Mich. fl. amer.* 1. p. 248.  
Icon. l. c. t. 25.  
In open wet woods of Lower Carolina. *Michaux.* 4. v. s. Flowers of a yellowish-brown, the size of *Ornithogalum umbellatum*.

¶

X. D

355. ARBUTUS  
pellucida.356. GAULTHI  
*Cor.* ovata  
cato.357. VACCINI  
campanula  
mini inse  
sperma.358. ANDROM  
5-fido, re  
dissepimenta359. KALMIA,  
limbo sub  
cumbunt.360. EPIGÆA,  
hypocrate  
partita.361. RHODOD  
fundibuli  
5-locular362. RHODOR  
declinata363. PYROLA  
nibus lor364. CHIMAP  
sessile

## X. DECANDRIA.

### I. MONOGYNIA.

#### \* *Monopetalæ.*

355. ARBUTUS. *Cal. 5-partitus. Cor. ovata: basi ore pellucida. Bacca supera, 5-locularis.*
356. GAULTHERIA. *Cal. 5-fidus, basi bibracteatus. Cor. ovata. Caps. 5-locularis, vestita calyce bacato.*
357. VACCINIUM. *Cal. 4-5-fidus. Cor. urceolata aut campanulata, 4-5-fida: laciniis reflexis. Filam. germini inserta. Bacca infera, 4-5-locularis, polysperma.*
358. ANDROMEDA. *Cal. 5-partitus. Cor. ovata: ore 5-fido, reflexo. Caps. supera, 5-locularis, valvulis dissepimento contrariis.*
359. KALMIA. *Cal. 5-partitus. Cor. hypocrateriformis: limbo subtus 5-corni, in quorum foveis antheræ incombunt. Caps. 5-locularis.*
360. EPIGÆA. *Cal. 5-partitus, extus tribracteatus. Cor. hypocrateriformis. Caps. 5-locularis. Recept. 5-partita.*
361. RHODODENDRON. *Cal. 5-partitus. Cor. subinfundibuliformis, inæqualis. Stam. declinata. Caps. 5-locularis.*
362. RHODORA. *Cal. 5-dentatus. Cor. 3-petala! Stam. declinata. Caps. 5-locularis.*

#### \*\* *Polypetalæ; Regulares.*

363. PYROLA. *Cal. 5-partitus. Pet. 5. Stylus staminibus longior. Caps. 5-locularis, angulis dehiscens.*
364. CHIMAPHILA. *Cal. 5-partitus. Pet. 5. Stigma sessile crassum, orbiculatum, germine immerso.*

- Antheræ rostratæ, foramine subbivalvi dehiscentes.  
Caps. 5-locularis, angulis dehiscens.*
365. LEDUM. *Cal. minimus, 5-dentatus. Cor. plana,  
5-partita. Caps. 5-locularis, basi dehiscens.*
366. AMMYRSINE. *Cal. profunde 5-partitus. Pet. 5.  
Stamina exerta. Caps. 5-locularis, apice dehiscens.*
367. CLETHRÆA. *Cal. 5-partitus, persistens. Pet. 5.  
Styl. apice 3-fidus, persistens. Caps. 3-locularis,  
3-valvis.*
368. MYLOCARIUM. *Cal. 5-dentatus. Pet. 5. Stigm.  
capitatum, trigonum, sessile. Caps. 3-4-alata, 3-  
locularis.*
369. MONOTROPA. *Cal. Q. Pet. 10. horum 5. exte-  
riora basi excavata, mellifera. Caps. 5-valvis, poly-  
sperma.*
370. DIONÆA. *Cal. 5-phillus. Pet. 5. Stigm. fimbria-  
tum. Caps. 1-locularis, gibba, polysperma.*
371. JUSSIÆA. *Cal. 4-5-partitus, superus. Pet. 4-5.  
Caps. 4-5-locularis, oblonga, angulis dehiscens, calyce  
coronata. Sem. numerosa, minuta.*
372. GYMNOCLADUS. *Cal. tubulosus, 5-fidus. Pet. 5.  
aequalia, tubo inserta. Stam. non exerta. Legumen  
1-loculare, intus pulposum. Flores racemosi. Dioicus.*
373. ACACIA. *Cal. tubulosus, 5-dentatus. Pet. 5. Stam.  
5-10. exerta. Legumen 1-loculare, 2-valve. Po-  
lygama.*
374. SCHRANKIA. *Cal. tubulosus, 5-dentatus. Pet. 5.  
Stam. 8-10. exerta. Siliqua 4-valvis. Polygama.*
- \*\*\* *Polypetalæ; Irregularæ.*
375. CASSIA. *Cal. 5-phillus. Pet. 5. subæqualia. Anth.  
supremæ 3. steriles, infinæ 3. rostratæ, in filamentis  
longioribus incurvis. Legumen membranaceum,  
2-valve.*
376. PODALYRIA. *Cal. subbilabiatus, 5-fidus. Cor.  
papilionacea, alæ vexillii longitudine. Legumen lave,  
ventricosum, polyspermum.*
377. CERCIS. *Cal. 5-dentatus, inferne gibbus. Cor. pa-  
pilionacea : vexillo sub alis brevi. Legumen. Folia  
simplicia.*
378. VIRGILIA. *Cal. 5-dentatus, inferne gibbus. Cor.  
papilionacea : alæ vexillo breviores. Legumen com-  
pressum, polyspermum. Folia pinnata.*
379. HYDRA  
*Caps. 2  
dehisce-*
380. SAXIFR.  
*rostris,*
381. TIAREL  
*inserta.  
majore*
382. MITELL  
*inserta.  
libus.*
383. SAPON.  
*guicula*
384. DIANT.  
*basi squ.  
lindrica*
385. SCLERA  
*inserta.*
386. CUCUB.  
*absque*
387. SILENE  
*ronata*
388. STELLA  
*partita.  
6-denta*
389. ARENA  
*tegra.*
390. MICRO  
*minuta  
ovata,*
391. SPERGI  
*ovata,*
392. CERAS

## II. DIGYNIA.

379. HYDRANGEA. *Cal. superus, 5-dentatus. Pet. 5.*  
*Caps. 2-locularis, 2-rostris, foramine inter cornua dehiscens.*
380. SAXIFRAGA. *Cal. 5-partitus. Pet. 5. Caps. 2-rostris, 1-locularis, polysperma.*
381. TIARELLA. *Cal. 5-partitus. Pet. 5. integra, calyci inserta. Caps. 1-locularis, 2-valvis: valvula altera majore*
382. MITELLA. *Cal. 5-fidus. Pet. 5. pinnatifida, calyci inserta. Caps. 1-locularis, 2-valvis; valvulis æquilibus.*
383. SAPONARIA. *Cal. tubulosus, nudus. Pet. 5. unguiculata. Caps. oblonga, 1-locularis.*
384. DIANTHUS. *Cal. cylindricus, longus, coriaceus, basi squamis 4—8. Pet. 5. unguiculata. Caps. cylindrica, 1-locularis, apice dehiscens.*
385. SCLERANTHUS. *Cal. 1-phyllo, cui Stamina inserta. Cor. 0. Sem. 1-2. calyce inclusa.*

## III. TRIGYNIA.

386. CUCUBALUS. *Cal. inflatus. Pet. 5: unguiculata, absque corona ad faucem. Caps. 3-locularis.*
387. SILENE. *Cal. cylindricus. Pet. 5. unguiculata, coronata ad faucem. Caps. 3-locularis.*
388. STELLARIA. *Cal. 5-phyllo, patens. Pet. 5. bipartita. Caps. ovata, 1-locularis, polysperma, apice 6-dentata.*
389. ARENARIA. *Cal. 5-phyllo, patens. Pet. 5. integra. Caps. 1-locularis, polysperma.*

## IV. TETRAGYNIA.

390. MICROPETALUM. *Cal. 5-phyllo, patens. Pet. 5. minuta, integra aut nulla. Stigm. 4. sessilia. Caps. ovata, calyce longior, 4-valvis.*

## V. PENTAGYNIA.

391. SPERGULA. *Cal. 5-phyllo. Pet. 5. integra. Caps. ovata, 1-locularis, 5-valvis.*
392. CERASTIUM. *Cal. 5-phyllo. Pet. 5. bifida aut*

- emarginata. *Caps.* 1-locularis, apice dentatim de-biscens.
393. AGROSTEMMA. *Cal.* 1-phyllo, coriaceus. *Pet.* 5. unguiculata : limbo obtuso, indiviso. *Caps.* 1-locularis.
394. OXALIS. *Cal.* 5-phyllo. *Pet.* 5. unguibus connexa. *Stamina* inaequalia, 5. breviora, exteriora basi con-nata. *Caps.* angulis elastice dehicens, 5-gona. *Sem.* subarillata.
395. PENTHORUM. *Cal.* 5-fidus. *Pet.* 0. aut 5. *Caps.* 5. cuspidata, 5-locularis.
396. SEDUM. *Cal.* 5-fidus. *Pet.* 5. *Squamæ* nectari-feræ 5. ad basin germinis. *Caps.* 5.

## VI. DECAGYNIA.

397. PHYTOLACCA. *Cal.* 0. *Pet.* 5. calycina. *Bacca* supera, 10-locularis, 10-sperma.

## I. MONOGYNIA.

355. ARBUTUS. *Gen. pl.* 750.

- laurifolia.* 1. A. arborea ; foliis oblongis utrinque acuminatis acute serratis glabris, racemis axillaribus secundis sessilibus solitariis. *Willd. sp. pl.* 2. p. 617. In North America. *Linn. suppl.* 238. h. +. What part of North America this tree comes from, I am un-acquainted with ; probably from the west coast, and very likely may be found on the Columbia, as well as the foregoing species. Nothing can be found in the Linnaean Herbarium respecting the plant. *J. E. Smith.*
- Menziesii.* 2. A. arborescens ; foliis lato-ovalibus integerrimis glabris, petiolis longis, racemis axillaribus et terminalibus pa-niculatis densifloris. On the north-west coast of America. *A. Menzies.* h. v. s. in *Herb. Banks.*
- tomentosa.* 3. A. frutescens ; ramis hispidis, foliis ovalibus acutis basi subcordatis subtus albido-tomentosis, petiolis brevibus, pedunculis axillaribus longitudine foliorum subcapi-tato-racemifloris, floribus campanulato-urceolatis brac-teatis.

ntatim de.

eus. Pet. 5.  
aps. 1-locu-

us connexa.  
a basi con-  
gona. Sem.

ut 5. Caps.

mæ nectari-

cina. Bacca

ninatis acute  
idis sessilibus

+. What  
om, I am un-  
st coast, and  
ia, as well as

erbarium re-

rimis glabris,  
minalibus pa-

Menzies. h.

us acutis basi  
iolis brevibus,  
orum subcapi-  
rceolatis brac-



*Gaultheria shallon.*

Drown & Engelm.

Tab.



On the north-w.  
v. s. in Herb.

4. A. caulis pro-  
sis serratis, 1  
p. 618.

Icon. Fl. dan.  
In Canada. 1  
berries black

5. A. caulis pro-  
integerrimis  
ciculatis, bac  
Icon. Schmidt  
In the pine-ba  
and rocky s  
h. June.  
On the plai  
the leaves u  
sider them c


## 356. GA

1. G. caule proct  
perne confe  
nuissime cili  
nutantibus.-

Icon. Bot. rep  
In dry woods,  
nada to Virg  
plant, resem  
flowers wh  
by the na  
properly cu  
reason it is  
tun-tea.

2. G. caule reper  
tis, floribus  
corollis bre  
Vaccinium his  
Arbutus filifor  
Icon. Mich. J  
In sphagnum  
other everg  
April, May  
white, very

3. G. erecta frut



*Hamiltonia oleifera*. p. 178.

On the north-west coast of America. *A. Menzies.* ♂.  
*v. s. in Herb. Banks.*

4. *A. caulinibus procumbentibus, foliis obovatis acutis rugosis serratis, racemis terminalibus.* — *Willd. sp. pl. 2. p. 618.*

*Icon. Fl. dan. 73. Engl. bot. 2030.*

In Canada. ♂. May, June. v. v. Flowers white; berries black.

5. *A. caulinibus procumbentibus, foliis cuneato-obovalibus integerrimis margine convexis coriaceis, floribus fasciculatis, baccis 5-spermis.* — *Willd. sp. pl. 2. p. 618.*

*Icon. Schmidt arb. t. 138. Fl. dan. t. 33. Engl. bot. 714.*

In the pine-barrens of New Jersey, and in mountainous and rocky situations of Canada and New England. ♂. June. v. v. Flowers pale red; berries scarlet. On the plains of the Mississippi the Indians smoke the leaves under the name of *Sacacommis*, and consider them of great medicinal virtue.

*alpina.*

*Uva ursi.*

### 356. GAULTHERIA. *Gen. pl. 749.*

1. *G. caule procumbente, ramis erectis, inferne nudis superne confertim foliosis, foliis obovatis basi acutis tenuissime ciliato-dentatis, floribus paucis terminalibus nutantibus.* — *Willd. sp. pl. 2. p. 616.*

*Icon. Bot. rep. 116. Duham. arb. 1. t. 113.*

In dry woods, on mountains, and in sandy plains: Canada to Virginia. ♂. May. v. v. A little shrubby plant, resembling seedling plants of *Kalmia latifolia*; flowers white; berries red, eatable, and known by the name of *Partridge-berries*. The leaves, if properly cured, make a most excellent tea, for which reason it is likewise known by the name of *Mountain-tea*.

2. *G. caule repente hispido, foliis subrotundo-ovalibus acutis, floribus solitariis axillaribus subsessilibus octandris, corollis brevi-campanulatis.*

*Vaccinium hispidulum. Willd. sp. pl. 2. p. 355.*

*Arbutus filiformis. Lam. encycl. 1. p. 228.*

*Icon. Mich. fl. amer. 1. t. 23.*

In sphagnum swamps, particularly where cedars and other evergreens abound: Canada to Pennsylvania. ♂. April, May. v. v. A small creeping plant; berries white, very sweet, and agreeable to eat.

3. *G. erecta fruticosa; foliis ovatis subcordatis serrulatis,*

*Shallon.*

## DECANDRIA MONOGYNIA. Arbutus.

racemis secundis bracteatis, pedicellis medio bibracteolatis.

On the falls of Columbia and near the Western Ocean.  
*M. Lewis.* On the North-west coast. *A. Menzies.*

May, June. *v. s. in Herb. Lewis. nec non Banks.*

*Frutex sempervirens.* *Rami teretes, verrucosi, glabri : juniores ferrugineo-pubescentes. Folia alterna, brevissime petiolata, ovata vel lato-ovalia, interdum basi subcordata, abrupte acuminata, serrulata, coriacea, utrinque glabra, subtus nervis venisque reticulatis prominentibus. Racemi in apice ramulorum axillares et terminales, simplices, ferrugineo-pubescentes. Flores secundi, bracteati, albi, rore rubro pubescentes. Pedicelli solitarii, medio bibracteolati. Bracteae lanceolatae. Corolla urceolata, limbo subclauso, 5-dentata. Bacc subglobosa, acuta, carnosa, purpurea, extus hirsuta, intus semi-5-valvis : loculis polyspermis. Semina ovato-subtriangulatae.*

This elegant evergreen shrub is in high esteem among the natives, on account of its berries, which they call *Shallon*, and which name I have adopted here. Mr. A. Menzies, who was the first discoverer of this shrub, observes one peculiarity, which is its growing under the shade of close pine forests, where scarcely any other plant thrives: this makes it so much more desirable for our plantations, as a highly ornamental shrub, since but few will bear such a situation. I was first inclined to take this species to belong to *Arbutus*, having only seen imperfect specimens; but through the liberal communication of Mr. A. Menzies, I have been enabled to class it more properly with this genus, though it deviates from both in some respects.

## 357. VACCINIUM. Gen. pl. 658.

+ *Foliis deciduis.*

\* *Corollis campanulatis.*

*stanineum.*

1. V. foliis ovalibus acutis integerrimis subtus glaucis, pedicellis solitariis axillaribus filiformibus, corollis patulo-campanulatis, laciinis oblongis acutis, antheris exertis, dorso aristatis, baccis pyriformibus.—*Willd. sp. pl. 2. p. 349.*

Icon. *Pluk. alm. t. 339. f. 3. bona.*

In dry woods; common: New England to Florida. May, June. v. v. About two feet high; flowers white: berries green, or white when per-

## DECANDRI

fectly ripe. I name of *Deer*.

2. V. foliis ovalibus glaucis, nervis litariis axillari panulatis, laci ertis dorso ari pl. 2. p. 350?

Icon. *Bot. rep. 2*

In dry sandy v April, May. berries white whether this i never could i species, I ven Banksian *He vatum.*

3. V. foliis petiolatis glanduloso-ser subtus subput bus, pedicellis cylindrico-can clusis dorso at V. diffusum. In dry woods, orolina to Flori sometimes tw white, tinged dry.

4. V. ramulis, foli nosis irroratis cronatis integ pedicellis bre bracteatis, co antheris inclu V. hirtellum.

*Banks.*

V. frondosum. In dry sandy Jersey to Flc shrub; flowe

5. V. foliis obovat rimis glabris irroratis, race

fectly ripe. In the mountains they are known by the name of *Deer-berries*.

2. *V. foliis ovalibus seu obovatis acutis integerrimis subtus glaucis, nervis venisque pubescentibus, pedicellis solitariis axillaribus filiformibus, corollis patulo-campanulatis, lacinij ovatis subacuminatis, antheris exsertis dorso aristatis, baccis subglobosis.* — *Willd. sp. pl. 2. p. 350?*

Icon. *Bot. rep. 263. V. stamineum.*

In dry sandy woods: New Jersey to Carolina.  $\frac{1}{2}$ . April, May. v. v. Flowers white, larger than No. 1; berries white or greenish-white. I am not certain whether this is the *V. album* of Linnaeus or not. As I never could find any certainty about the authentic species, I venture to adopt this species for it. In the Banksian Herbarium it is under the name of *V. elevatum*.

3. *V. foliis petiolatis obovalibus utrinque acutis mucronatis glanduloso-serrulatis, supra nitidis, reticulato-venosis, subtus subpubescentibus, racemis bracteatis nutantibus, pedicellis axillaribus solitariis filiformibus, corollis cylindrico-campanulatis; laciñiis acutis, antheris inclusis dorso aristatis.* — *Mich. fl. amer. 1. p. 230.*

*V. diffusum. Ait. kew. 2. p. 11. Herb. Banks.*

In dry woods, on the rocky banks of rivers: North Carolina to Florida.  $\frac{1}{2}$ . May, June. v. v. A large shrub, sometimes twenty feet high, very elegant: flowers white, tinged with red; berries globular, black, almost dry.

4. *V. ramulis, foliis racemisque hispidiusculis atomis resinosis irroratis, foliis obovato-oblongis basi acutis mucronatis integerrimis concoloribus, racemis bracteatis, pedicellis brevibus axillaribus subsolitariis medio bi-bracteatis, corollis campanulatis, laciñiis rotundatis, antheris inclusis.* *Bot. mag. 1106. Bot. rep. 112.*

*V. hirtellum. Ait. kew. ed. 2. v. 2. p. 357. Herb. Banks.*

*V. frondosum. Mich. fl. amer. 1. p. 230.*

In dry sandy woods, particularly pine-forests: New Jersey to Florida.  $\frac{1}{2}$ . June, July. v. v. A low shrub; flowers large, white; berries black, globular.

5. *V. foliis obovato-oblongis obtusiusculis muticis integerimis glabris subtus glaucis copiose atomis resinosis irroratis, racemis laxis bracteatis, pedicellis longis fili-*

*album.*

*arboreum.*

*dumosum.*

*frondosum.*

## DECANDRIA MONOGYNIA. Vaccinium.

- formibus bracteolatis, bracteis linearibus, corollis ovato-campanulatis, lacinia acutis, antheris inclusis.—*Willd. sp. pl. 2. p. 352.*
- V. glaucum. Mich. fl. amer. 1. p. 231.*  
Icon. *Bot. rep. 140.*
- lanceolatum. β. V. foliis lanceolatis utrinque acutis.*  
*V. venustum. Ait. kew. 2. p. 11. Herb. Banks.*  
In open woods, sandy soil : New Jersey to Carolina.  $\frac{1}{2}$ . May, June. *v. v.; v. s. in Herb. Banks.* About three feet high ; flowers small, almost globular, white ; berries large, blue, globular, eatable ; called by the country people *Blue-tangles*.
- pallidum. 6. V. foliis ovatis acutis serrulatis glabris, racemis bracteatis, corollis cylindraceo-campanulatis. Ait. kew. 2. p. 10.*  
In North America. *Aiton.  $\frac{1}{2}$ . May, June. +.* A doubtful species. I have not been able to ascertain it, even after a strict examination of the Banksian Herbarium.
- \*\* *Corollis urceolatis; Antheris inclusis.*  
|| *Floribus racemosis seu fasciculatis.*
- resinosum. 7. V. foliis tenuibus petiolatis oblongo-ovalibus plerumque obtusis muticis integerrimis subtus atomis resinosis irroratis, racemis lateralibus secundis, pedicellis brevibus subbracteolatis, corollis ovato-conicis 5-gonis.—Willd. sp. pl. 2. p. 352.*  
*Andromeda baccata. Wangh. amer. 2. t. 30. f. 69.*
- viridescens. α. V. corollis luteo-viridescentibus.*  
*rubescens. β. V. corollis rubescens. Bot. mag. 1288.*  
*lutescens. γ. V. foliis lanceolatis, corollis rubro-lutescentibus.*  
*V. parviflorum. Bot. rep. 125.*  
In woods and mountains ; frequent : Canada to Carolina.  $\frac{1}{2}$ . May, June. *v. v.* From two to four feet high ; berries black, eatable.
- corymbosum. 8. V. ramis floriferis subaphyllis, foliis oblongo-ovalibus utrinque acutis mucronatis subintegerrimis, junioribus utrinque pubescentibus subtus subtomentosis, adultis supra glabris, subtus nervis venisque pubescentibus, racemis brevibus sessilibus squamoso-bracteatis, corollis cylindraceo-ovatis, calycibus erectis, stylo sub-exerto.—Willd. sp. pl. 2. p. 351.*  
*V. disomorphum. Mich. fl. amer. 1. p. 231.*  
*V. album. Lam. encycl. 1. p. 73.*  
In swamps and wet woods : Canada to Virginia.  $\frac{1}{2}$ . June. *v. v.; v. s. in Herb. Banks.* A shrub, rising

DECANDRIA

sometimes  
ers white,  
insipid.9. *V. ramis flor  
integerrimi  
racemis co  
acutis, caly  
pl. 2. p. 35**V. disomorph  
Icon. *Bot. re*  
In low grou  
 $\frac{1}{2}$ . May,  
shrub wit  
red tinge ;  
number of*10. *V. ramis flor  
oblongis tu  
silibus co  
fauce coar  
stylo inclu  
p. 12.*  
Icon. *Bot. re*  
In swamps c  
*v. v.* A  
crimson,11. *V. foliis obla  
gatis term  
gis nutant  
vibus, cal  
Willd. sp.**V. formosun  
β. V. foliis ang  
In swamps c  
June. *v.*  
shrub ; fl*12. *V. foliis sess  
sis pubesc  
vissimis,  
valde coa  
p. 232.*  
In shady w  
May, Ju  
berries sn

DECANDRIA MONOGYNIA. Vaccinium.

s, corollis inclusis.—  
sometimes to the height of seven or eight feet ; flowers white, sometimes tinged with red ; berries black, insipid.

9. V. ramis floriferis aphyllis, foliis oblongis utrinque acutis integerrimis glabris nervis subtus subpubescentibus, racemis confertis bracteatis, corollis cylindraceis, laciiniis acutis, calycibus reflexis, stylo incluso.—Willd. sp. pl. 2. p. 353. Ait. kew. 2. p. 12.

V. disomorphum var. Mich. fl. amer. 1. p. 231.

Icon. Bot. rep. 138.

In low grounds and swamps : New Jersey to Virginia.  
h. May, June. v. v.; v. s. in Herb. Banks. A tall shrub with red twigs : flowers large, white with a red tinge ; berries black, insipid. This species has a number of varieties in size, shape, and colour.

10. V. ramis floriferis elongatis subaphyllis, foliis lanceolato-oblongis utrinque acutis serrulatis glabris, racemis sessilibus corymbosis bracteatis, corollis cylindraceis, fauce coarctato, laciiniis brevissimis, calycibus reflexis, stylo incluso.—Willd. sp. pl. 2. p. 353. Ait. kew. 2. p. 12.

Icon. Bot. rep. 181.

In swamps of Virginia and Carolina. h. May, June. v. v. A middle-sized shrub ; flowers tinged with crimson, very handsome.

11. V. foliis oblongis acutis serrulatis glabris, racemis aggregatis terminalibus corymbosis bracteatis, pedicellis longis nutantibus, corollis cylindraceis, laciiniis erectis brevibus, calycibus acutis erectis, stylo subexerto.—Willd. sp. pl. 2. p. 351. Ait. kew. 2. p. 11.

V. formosum. Bot. rep. 97.

- β. V. foliis angusto-lanceolatis utrinque acuminatis.

In swamps of Lower Carolina and Georgia. h. May, June. v. v.; v. s. in Herb. Banks. An elegant small shrub ; flowers red and white striped ; calyx brown.

12. V. foliis sessilibus cuneato-lanceolatis subserrulatis venosis pubescentibus, fasciculis sessilibus, pedicellis brevissimis, calycibus acuminatis, corollis ovatis fauce valde coarctatis, stylo exerto.—Mich. fl. amer. 1. p. 232.

In shady woods and swamps : Virginia and Carolina. h. May, June. v. v. Flowers small, yellowish-white ; berries small, globular, black.

amoenum.

virgatum.

fuscatum.

angustifoli-  
um.

galeans.

*ligustrinum.* 13. V. ramis angulatis, foliis subsessilibus erectis lanceolatis mucronatis serrulatis venosis pubescentibus, fasciculis gemmaceis sessilibus, pedicellis vix ialis, corollis oblongiuscule-ovatis.—Willd. sp. pl. 2. p. 352. Mich. fl. amer. 1. p. 233.

In dry woods; Pennsylvania to Virginia; common on the mountains.  $\frac{1}{2}$ . May, June. v. v. An upright straight shrub; flowers purplish-red, berries black. This species varies extremely in the shape and size of the leaves.

*tenellum.* 14. V. ramis angulatis viridibus, foliis sessilibus ovato-lanceolatis mucronatis serrulatis utrinque lucidis, fasciculis confertifloris subterminalibus sessilibus, corollis ovatis.—Willd. sp. pl. 2. p. 353. Ait. kew. 2. p. 12. V. pensylvanicum. Lum. encycl. 1. p. 72. Mich. fl. amer. 1. p. 232.

On dry hills, in gravelly soil: New England to Virginia.  $\frac{1}{2}$ . May. v. v.; v. s. in Herb. Banks. A low, very branching shrub; flowers pale red; calyx green; berries large, blueish-black, extremely sweet and agreeable to eat. The mountains of Pennsylvania produce an immense variety of this species, in size, and shape of the fruit, leaves and flowers.

### ||| Floribus solitariis.

*uliginosum.* 15. V. foliis parvis obovatis obtusis integerrimis supra glabris subtus venosis pubescentibus glaucisque. floribus sub-solitariis, corollis brevi-urceolatis.—Willd. sp. pl. 2. p. 350. Mich. fl. amer. 1. p. 235.

Icon. Fl. dan. 23 t. Engl. bot. 581.

In bogs of the more northern regions of America.  $\frac{1}{2}$ . April, May. v. s. From one to two feet high; flowers rose-coloured; stamens from eight to ten; berries blueish-black, edible.

*myrtilloides.* 16. V. foliis angusto-lanceolatis membranaceis integerrimis subtus juxta nervos et margine pubescentibus, floribus sparsis subsolitariis brevissime-pedicellatis.—Mich. fl. amer. 1. p. 234.

V. angustifolium. Willd. sp. pl. 2. p. 351. Ait. kew. 2. p. 11.

In Canada about Hudson's Bay and Labrador.  $\frac{1}{2}$ . April, May. v. s. in Herb. Banks. Berries large, blueish-black, known by the name of *Bluets*.

### DECANDRI.

17. V. pusillum, ces obovalibus rot branaceis, flori rolla breviter p In the more nor son's Bay. M three inches hi cimens of A. the Banksian 1 species, but ap

++

\*

18. V. humile; cau obovatis emarg dis, subtus pt bus, corollis cyl p. 354. Mich Icon. Fl. dan. 40 On rocks near th  $\frac{1}{2}$ . May, June red, generally acid. The A European, and

19. V. repens, glabe dis revolutis ra subsessilibus pa latis brevissime Mich. fl. amer In Carolina. M

20. V. diffusum; ra lanceolatis utri mis terminali floribus nutan tulo-campanul Bot. rep. 105. In Carolina. Fr pink red, very

\*

21. V. erectum, rar dis obovato-ell cernis termina corollis cylind

17. *V. pusillum*, cespitosum, glaberrimum; foliis cuneato-  
obovalibus rotundato-obtusis conspicue serratis mem-  
branaceis, floribus rameis solitariis subsessilibus, co-  
rolla breviter pedicellata. *Mich. fl. amer.* 1. p. 234.  
*cespitosum.*

In the more northern regions, particularly about Hudson's Bay. *Michaux.*  $\frac{1}{2}$ . A little shrub not above three inches high; berries black. Among the specimens of A. Menzies from the north-west coast in the Banksian Herbarium is one very like the present species, but apparently evergreen.

++ *Foliis sempervirentibus.*

\* *Corollis campanulatis.*

18. *V. humile*; caulis repens, ramulis erectis, foliis  
obovatis emarginatis revolutis subserrulatis supra luci-  
dis, subtus punctatis, racemis terminalibus nutanti-  
bus, corollis cylindrico-campanulatis.—*Willd. sp. pl.* 2.  
p. 354. *Mich. fl. amer.* 1. p. 229.

*Icon. Fl. dan.* 40. *Engl. bot.* 593.

On rocks near the sea-coast: Canada to New England.  
 $\frac{1}{2}$ . May, June. v. v. A low shrub; flowers pale red, generally 4-cleft with four stamens; berries red, acid. The American plant is more robust than the European, and the leaves are considerably larger.

19. *V. repens*, glaberrimum; foliis petiolatis ovalibus luci-  
dis revolutis rariter denticulatis, fasciculis axillaribus  
subsessilibus paucifloris, corollis subgloboso-campanu-  
latis brevissime 5-denticulatis, antheris dorso muticis.  
*Mich. fl. amer.* 1. p. 229.

In Carolina. *Michaux.*  $\frac{1}{2}$ . +. Berries small, black.

20. *V. diffusum*; ramis adscendentibus laxis, follis oblongo-  
lanceolatis utrinque acutis serratis rigidis glabris, race-  
mis terminalibus corymbosis bracteatis paucifloris,  
floribus nutantibus, calycibus adpressis, corollis pa-  
tulo-campanulatis profunde acuteque 5-dentatis.—  
*Bot. rep.* 105.

In Carolina. *Fraser.*  $\frac{1}{2}$ . May, June. v. v. Flowers  
pink red, very ornamental.

\*\* *Corollis urceolatis.*

21. *V. erectum*, ramosissimum; ramis distichis, foliis niti-  
dis obovato-ellipticis utrinque acutis glabris serratis, ra-  
cemis terminalibus corymbosis bracteatis nutantibus,  
corollis cylindraceis.—*Bot. rep.* 480.

*Vitis Idaea.*

*myrtifolium.*

*crassifolium.*

*nitidum.*

## DECANDRIA MONOGYNIA. Vaccinium.

- In Carolina.  $\text{h}$ . May, June.  $v.v.$  Flowers pink-red. This species approaches near to No. 20, but differs in the shape of the corolla materially; it is also near to the following, but sufficiently different.
- myrsinitis.* 22. *V. erectum, ramosissimum, microphyllum; foliis sessilibus ovalibus mucronatis subserrulatis supra levigatis lucidis subtus subhirsutis scabro-punctatis, fasciculis gemmaceis terminalibus lateralibusque, corollis oblongo-ovatis.* — *Mich. fl. amer. 1. p. 233.*
- lanceolatum.*  $\beta$ . *V. foliis lanceolatis utrinque acutis.*
- obtusum.*  $\gamma$ . *V. foliis subrotundo-ovatis.*  
In dry sandy woods: Carolina and Florida.  $\text{h}$ . May, June.  $v.v.$  A beautiful little shrub; calyx scarlet; corolla purple.
- luxifolium.* 23. *V. pumilum; foliis obovatis crenato-dentatis glabris; fasciculis congestis subsessilibus axillaribus et terminalibus, corollis brevi-ovatis, filamentis glandulosis, stigmate capitato.* — *Bot. rep. 4. Bot. mag. 928.*  
*V. brachycerum.* *Mich. fl. amer. 1. p. 234.*  
In dry woods, on limestone rocks: western parts of Virginia, near Winchester and the Sweet-springs.  $\text{h}$ . June.  $v.v.$  A handsome little shrub; flowers white, delicately tinged with red.
- ovatum.* 24. *V. foliis petiolatis ovatis acutis revolutis serratis glabris coriaceis, racemis axillaribus et terminalibus bracteatis brevibus, corollis cylindraceis, calycibus acutis.*  
On the Columbia river. *M. Lewis.* On the north-west coast. *Menzies.*  $\text{h}$ . May, *v.s. in Herb. Banks.*
- obtusum.* 25. *V. repens; foliis parvis ovalibus utrinque rotundato-obtusis mucronatis integerrimis glabris coriaceis, pedunculis axillaribus solitariis unifloris.*  
On the north-west coast. *Menzies.*  $\text{h}$ . *v.s.s.fl. in Herb. Banks.*

358. ANDROMEDA. *Gen. pl. 747.*\* *Foliis perennantibus.*

- tetragona.* 1. *A. foliis imbricatis carinatis margine nudis, pedunculis axillaribus solitariis filiformibus unifloris, corollis nutantibus campanulatis.* — *Willd. sp. pl. 2. p. 607.*  
*Icon. Pall. fl. ross. 2. t. 73. f. 4. Fl. lapp. t. 1. f. 4.*  
In Canada; Labrador. *Colmaster.* On the north-west coast. *Nelson.*  $\text{h}$ . May, June.  $v.s.$  Flowers white.

- DECANDRIA
2. *A. foliis imbricatis terminalibus uncampanulatis.* — *Icon. Fl. dan. 10.*  
On the north-west coast. *Herb. Banks.* moss; flowers white.
3. *A. foliis linear-lanceolatis bicanti-glaucis aggregatis terminalibus versus apicem.* — *Mich. fl. amer. 1. p. 234.*  
 $\alpha$ . *A. foliis lanceolatis rubris.* *Ait. herbarium.*  
*Icon. Pall. fl. ross. 2. t. 73. f. 4.*  
 $\beta$ . *A. foliis oblongis, cinis patentibus.* *Ait. kew. 2. p. 607.*  
In bogs of Canada; mountain-lakes, and Pennsylvania; about one foot high. I strongly suspect which might be.
4. *A. foliis lanceolatis subrevolutis neis, racemis terminalibus brevibus solitariis, bracteatis, bracteolatis suboblango-cylindraceis.* — *Icon. Pall. fl. ross. 2. t. 73. f. 4.*  
In bogs and swamps in mountains: Canada.  $v.v.$  Flowers white.
5. *A. foliis linear-lanceolatis revolutis squamulatis, racemis terminalibus brevibus solitariis, bracteatis, bracteolatis, ovalibus.* — *A. calyculata*  $\gamma$ . *andromeda.* *Icon. Pall. fl. ross. 2. t. 73. f. 4.*  
In open swamps in Canada. May.  $v.v.$ ; very small. Approaches near to *A. colmaster*, but is distinct to admiring eyes.

*hypnoides.*

2. A. foliis imbricatis subulatis glabris, pedunculis solitariis terminalibus unifloris, corollis nutantibus globosocampanulatis.—Willd. sp. pl. 2. p. 608.

Icon. Fl. dan. 10. Pall. fl. ross. 2. t. 73. f. 2.

On the north-west coast. Nelson. ½. May. v. s. in Herb. Banks. A small creeping shrub, resembling a moss; flowers white, tinged with red.

3. A. foliis linear-lanceolatis convexis revolutis subtus albicanti-glaucis supra glauco-canescensibus, floribus aggregatis terminalibus, corollis subglobosis, antheris versus apicem aristatis.—Willd. sp. pl. 2. p. 610. Mich. fl. amer. 2. p. 254.

- $\alpha$ . A. foliis lanceolato-linearibus, laciniis calycinis oblongis rubris. Ait. kew. 2. p. 68.

Icon. Pall. fl. ross. 2. p. 53. t. 70. f. B.

- $\beta$ . A. foliis oblongis, corollis ovatis incarnatis, laciniis calycinis patentibus ovatis albis interdum apice rubicundis. Ait. kew. 2. p. 68.

In bogs of Canada and Labrador, and on the borders of mountain-lakes, in sphagnous grounds: New York and Pennsylvania. ½. June. v. v. A small shrub, about one foot high; flowers white, tinged with red. I strongly suspect the variety  $\alpha$ . to be a distinct species, which might be called *A. rosmarinifolia*.

4. A. foliis lanceolato-oblongis obtusiusculis obsolete serrulatis subrevolutis squamoso-punctatis subtus ferrugineis, racemis terminalibus foliosis secundis, pedicellis brevibus solitariis axillaribus, calycibus acutis basi bibracteatis, bracteis lato-ovatis acuminatis, corollis oblongo-cylindraceis. Willd. sp. pl. 2. p. 614.

Icon. Pall. fl. ross 2. p. 53. t. 71. f. 1.

In bogs and swamps; frequent, particularly on the mountains: Canada to Virginia. ½. April, May. v. v. Flowers white.

5. A. foliis linear-lanceolatis acutis margine subundulato-revolutis squamoso-punctatis subtus subferrugineis, racemis terminalibus foliosis secundis, pedicellis brevibus solitariis axillaribus, calycibus acuminatis basi bibracteatis, bracteis minutis acutis, corollis oblongovalibus.

*A. calyculata*  $\gamma$ . *angustifolia*. Ait. kew. 2. p. 70.

In open swamps: Carolina and Georgia. ½. April, May. v. v.; v. s. in Herb. Lambert. This species approaches nearly to the preceding, but is sufficiently distinct to admit it to be more than a variety.

*polifolia.**angustifolia.**latifolia.**calyculata.**angustifolia.*

*ferruginea.*

6. A. fruticosa ; ramis subflexuosis, foliis coriaceis distan-  
tibus longe petiolatis obovatis plerumque obtusis inte-  
gerrimis planis margine revolutis subtus squamoso-  
farinosis reticulato-venosis, pedicellis aggregatis axil-  
laribus simplicibus unifloris paucis furfuraceo-ferrugi-  
neis, corollis subglobosis, antheris muticis.

A. ferruginea. *Walt. fl. car.* 138.

A. ferruginea  $\beta$ . fruticosa. *Mich. fl. amer.* 1. p. 252.

Icon. *Vent. malm.* 80.

In the pine-woods of Georgia and Florida.  $\frac{1}{2}$ . June.  
*v. v.* A shrub from three to five feet high. Flowers  
small, white inside and ferruginous outside.

*rigida.*

7. A. arborescens ; ramis strictis, foliis coriaceis rigidis con-  
fertis brevi-petiolatis cuneato-lanceolatis acutis inte-  
gerrimis convexis margine revolutis subtus squamoso-  
tomentosis subaveniis, pedicellis aggregatis axillaribus  
simplicibus unifloris numerosis furfuraceo-ferrugineis,  
corollis subglobosis, antheris muticis.

A. ferruginea. *Willd. sp. pl.* 2. p. 609. *Ait. kew.* 2.  
p. 67.

A. ferruginea  $\alpha$ . arborescens. *Mich. fl. amer.* 1. p. 252.  
In barren sandy woods : Carolina to Florida.  $\frac{1}{2}$ . April.  
*v. v.* A shrub almost twenty feet high, or sometimes  
forming a small tree : though nearly allied to the  
former species, their habits, and particularly their flow-  
ering time, differ so materially, that I have always con-  
sidered them to be distinct species.

8. A. glaberrima ; ramis subtriquetris, foliis coriaceis ova-  
libus acuminatis integerrimis laevigatis, costa marginem  
deflexum percurrente, fasciculis axillaribus, co-  
rollis cylindraceis, laciniis calycis elongatis linearibus  
coloratis, antheris basi bicorniculatis.—*Mich. fl. amer.*  
1. p. 252.

A. coriacea. *Willd. sp. pl.* 2. p. 613. *Ait. kew.* 2. p. 70.

A. lucida. *Lam. encycl.* 1. p. 157.

A. marginata. *Duham. arb.* ....

A. mariana. *Jacq. ic.* 3. t. 465.

Icon. *Bot. mag.* 1095.

In the sandy forests of Carolina and Florida.  $\frac{1}{2}$ . June,  
July. *v. v.* A handsome small shrub ; flowers pale  
red ; the calyx long, of a dark-red colour ; leaves  
marked with very fine punctures.

*axillaris.*

9. A. foliis oblongo-ovalibus basi acutis brevi-acuminatis  
cartilagineo-serratis lucidis glabris coriaceis, racemis

## DECANDI

spicatis axill  
dique confe  
theris mutici  
p. 69.

β. A. foliis lineari-  
On the mounta  
June. *v. v.*  
sules depresso

10. A. foliis petiola  
sensim angu  
adpresso-spin  
catis axillari  
dis laxiuscul  
antheris muti

A. Catesbaei,  
p. 613.

In Lower Caroli  
the foregoing  
*Catesb. car.* :  
there can be  
belongs to A.

11. A. glaberrima ;  
angustatis ac  
serratis glabri  
cenis axillari  
floribus pedic  
ovatis, anther  
pl. 2. p. 613.

A. lucida. *Jacq*

A. populifolia.

A. reticulata. *W*

A. formosissima

A. laurina. *Mic*

Icon. *Smith exa*

In sandy swam  
*v. v.* Flowers  
this shrub a f

side, and is u  
stems of it,

12. A. glaberrima ;  
serrulatis ad  
secundis axil  
latis, pedicel  
In the mountain

spicatis axillaribus sessilibus squamoso-bracteatis un-dique confertifloris, corollis cylindraceo-ovatis, an-theris muticis.—Willd. sp. pl. 2. p. 613. Ait. kew. 2. p. 69.

$\beta$ . A. foliis lineari-lanceolatis longissimis.

On the mountains of Virginia to Georgia.  $\frac{1}{2}$ . May, June. v. v. Flowers in short spikes, white; capsules depressed, globular.

10. A. foliis petiolatis ovato-oblongis basi rotundatis superne sensim angustatis acuminatis subserratulatis, serraturis adpresso-spinulosis glabris coriaceis, racemis subspicatis axillaribus sessilibus squamoso-bracteatis secundis laxiusculis, corollis brevibus cylindraceo-ovatis, antheris muticis.

A. Catesbaei. Walt. fl. car. 137. Willd. sp. pl. 2. p. 613.

In Lower Carolina.  $\frac{1}{2}$ . May, June. v. v. Resembles the foregoing one in several respects. The figure of Catesb. car. 2. t. 43. is so very slightly done, that there can be no reference made to it, and probably belongs to A. racemosa.

11. A. glaberrima; foliis ovato-lanceolatis superne sensim angustatis acuminatis integerrimis seu inaequaliter-serratis glabris nitidis reticulato-venosis coriaceis, racemis axillaribus brevissimis corymbosis nudiusculis, floribus pedicellatis nutantibus, corollis cylindraceo-ovatis, antheris muticis postice gibbis.—Willd. sp. pl. 2. p. 613. Ait. kew. 2. p. 70.

A. lucida. Jacq. ic. 1. t. 79.

A. populifolia. Lam. encycl. 1. p. 159.

A. reticulata. Walt. fl. car. 137.

A. formosissima. Bartr. catal. ....

A. laurina. Mich. fl. amer. 1. p. 253.

Icon. Smith exot. bot. 89. bona. Jacq. ic. 79. mala.

In sandy swamps of Georgia and Florida.  $\frac{1}{2}$ . July. v. v. Flowers white, in great abundance; which gives this shrub a fine appearance. The stem is hollow inside, and is used by the natives for making their pipe-stems of it, from which the name Pipe-stem wood.

12. A. glaberrima; foliis oblongo-ovatis acutis tenuissime serrulatis adpresso-ciliatis glabris coriaceis, racemis secundis axillaribus et terminalibus congesto-paniculatis, pedicellis bibracteatis.

In the mountains of Georgia. Lyon.  $\frac{1}{2}$ . May, June.

longifolia.

spinulosa.

acuminata.

floribunda.

## DECANDRIA MONOGYNIA. Andromeda.

v. v.; v. s. in *Herb. Lyon.* Flowers clear white, in great abundance.

\*\* *Foliis deciduis.*

- mariana.* 13. A. foliis ovalibus utrinque subacutis integerrimis glabris subcoriaceis subtus pallidioribus, ramis floriferis subaphyllis, pedunculis fasciculatis, corollis ovato-cylindraceis, calycibus foliosis, antheris muticis, capsula conoidea.—*Willd. sp. pl. 2. p. 609. excl. syn. Jacquinii.*
- angustifolia.* β. A. foliis angusto-lanceolatis.  
Icon. *Pluk. mant. t. 448.*  
In woods and dry swamps, particularly in sandy soil: New England to Florida. h. May—Aug. v. v. A low shrub; flowers large, white, sometimes tinged with red. This plant has a number of very striking varieties.
- speciosa.* 14. A. foliis ovalibus obtusis mucronatis crenatis serratis reticulato-venosis, ramis floriferis denudatis aggregato-racemifloris, corollis campanulatis, antheris apice geminatum 4-aristatis.—*Mich. fl. amer. 1. p. 256.*
- nitida.* α. A. foliis oblongo-ovatis serratis utrinque viridis.
- A. *cassinæfolia.* *Vent. malm. 79.*
- pulverulenta.* β. A. foliis rotundato-ovatis remote crenatis ramisque albo pulverulentis.  
A. *pulverulenta.* *Bartr. it. 476. Bot. mag. 667.*  
A. *cassinæfolia* β. *Vent. hort. cels. 60.*  
A. *ovata.* *Solander mss. in mus. Banks.*  
In swamps of North Carolina. h. June. v. v. A very ornamental little shrub; flowers large, white. I certainly must coincide with Michaux's idea of *A. pulverulenta* being only a variety, as I very frequently have seen intermediate varieties, and even had the trace of both on one plant.
- racemosa.* 15. A. foliis ovali-lanceolatis acutis serrulatis membranaceis glabris, spicis terminalibus secundis elongatis simplicibus ramosis, bracteis linearibus acutis, corollis cylindraceis, calycibus acutis basi bibracteatis, antheris apice geminatum quadriaristatis.—*Willd. sp. pl. 2. p. 612.*  
A. *paniculata.* *Walt. fl. car. 138.*  
A. *racemis simplicibus, foliis lanceolatis alternis, &c.* *Gronov. virg. 67.*  
In bogs and swamps; frequent: Canada to Florida. h. June, July. v. v. A middle-sized shrub, which may

## DECANDRIA

be reckoned only for the fine odour th.

16. A. ramis terete nato-serratis chiis, corolla antheris lineata. Icon. Bot. mag. arb. ....

In the valleys of Florida. h. from forty to leaves are often has been called use of by thirst.

17. A. pubescens; gerrimis, radicisculis, glandulibus pubescentibus. *pl. 2. p. 612.*

A. *globulifera.*  
A. *racemosa.*  
Common in all parts. h. June, July. are a number of pubescence, s.

18. A. hispido-pubescentis longis acutis, rulis axillaris, basis hispida.

A. *racemosa.*  
A. *paniculata.*  
In the lower parts. May—July sufficiently

359

1. K. ramis, foliis oppositis alterniis, illaribus solitariis. *sp. pl. 2. p. 257.*

be reckoned one of the finest in North America, not only for the graceful appearance of its flowers, but the fine odour they have.

16. *A. ramis teretibus, foliis oblongis acuminatis mucronato-serratis glabris, paniculis terminalibus polystachyiis, corollis ovoideo-cylindraceis pubescentibus, antheris linearibus muticis.* — Willd. sp. pl. 2. p. 612.

*Icon. Bot. mag. 905. Catesb. car. 1. t. 71. Mich. arb. ....*

In the valleys of the Allegany-mountains: Pennsylvania to Florida.  $\frac{1}{2}$ . June, July. v. v. A beautiful tree, from forty to sixty feet high. Flowers white. The leaves are of a very pleasant acid taste, from which it has been called *Sorrel-tree*. They are frequently made use of by hunters in those mountains to alleviate thirst.

17. *A. pubescens; foliis obovato-lanceolatis acutis subintegerrimis, ramis floriferis terminalibus paniculatis nudiusculis, glomerulis pedunculatis, corollis subglobosis pubescentibus, antheris obtusis muticis.* — Willd. sp. pl. 2. p. 612.

*A. globulifera. Hortul.*

*A. racemosa. Lam. encycl. 1. p. 158.*

Common in all swamps and woods: Canada to Carolina.

$\frac{1}{2}$ . June, July. v. v. Flowers small, white. There are a number of varieties of this shrub, in size, pubescence, shape of leaves, &c.

18. *A. hispido-pubescentis; foliis obovato-lanceolatis seu oblongis acutis serrulatis, ramis floriferis foliosis, glomerulis axillaribus paucifloris subsessilibus, corollis globosis hispidis, antheris aristatis.*

*A. racemosa. Walt. fl. car. 138.*

*A. paniculata foliosiflora. Mich. fl. amer. 1. p. 225.*

In the lower counties of Virginia and Carolina.  $\frac{1}{2}$ . May—July. v. v. Nearly allied to the former; but sufficiently distinct to form a species.

*arborea.*

*paniculata.*

*frondosa.*

*hirsuta.*

1. *K. ramis, foliis calycibusque hirsutissimis, foliis oppositis alternisque subsessilibus ellipticis, pedunculis axillaribus solitariis unifloris foliis longioribus.* — Willd. sp. pl. 2. p. 601. *Walt. fl. car. 138. Mich. fl. amer. 1. p. 257.*

### 359. KALMIA. Gen. pl. 743.

- K. ciliata. *Bartr. it.* 18.  
Icon. *Bot. mag.* 138.  
In barren pine-woods: South Carolina and Georgia.  $\frac{1}{2}$ . June—Aug. v. v. A beautiful little shrub, but difficult to cultivate; leaves small, like thyme; flowers large, red.
- glaucæ.* 2. K. ramulis ancipitibus, foliis oppositis subsessilibus oblongis laevibus subtus glaucis margine revolutis, corymbis terminalibus bracteatis, pedunculis calycibusque glaberrimis.—*Ait. kew. ed. 2. v. 3. p. 47.* *Willd. sp. pl. 2. p. 601.*  
K. polifolia. *Wangh. act. soc. berol.* 8. p. 129. t. 5.  
Icon. *Bot. mag.* 177. *Ait. kew. 2. t. 8.* *Lam. illustr.* 363.  
 $\beta$ . K. foliis linearibus magis revolutis subtus viridibus.  
In bogs of Canada and on the borders of mountain-lakes of New York and Pennsylvania.  $\frac{1}{2}$ . April, May. v. v. An upright, small shrub; flowers pale red, very handsome. The variety  $\beta$ . I discovered in a bog near Albany, New York, and am inclined to think it a distinct species.
- cuneata.* 3. K. foliis sparsis sessilibus cuneato-oblongis subtus pubescentibus apice minutum aristatis, corymbis lateralibus paucifloris. *Mich. fl. amer.* 1. p. 257.  
On the mountains of Carolina. *Michaux.*  $\frac{1}{2}$ . +. Flowers white, with red at the bottom. Resembles the following one in some respects, but is distinct; the leaves are like those of *Azalea*. *Mich.*
- angustifolia.* 4. K. foliis ternis petiolatis oblongis obtusis subtus subferrugineis, corymbis lateralibus, bracteis linearibus, pedunculis calycibusque glanduloso-pubescentibus.—*Ait. kew. ed. 2. v. 3. p. 47.* *Willd. sp. pl. 2. p. 601.*  
Icon. *Bot. mag.* 331. *Catesb. car.* 3. t. 17. f. 1. *Trew ehret.* t. 38. f. 2.  
 $\beta$ . K. foliis laticribus subovatis, caule altiore.  
In bogs and swamps, and sometimes in dry mountain lands: Canada to Carolina.  $\beta$ . New Jersey mountains.  $\frac{1}{2}$ . May—July. v. v. A shrub from one to two feet high; flowers dark red; known by the name of *Sheep Laurel*, being considered very poisonous when fed upon by sheep.
- latifolia.* 5. K. foliis longe petiolatis sparsis ternisque ovalibus utrinque laevigatis et subconcoloribus, corymbis terminalibus viscidio-pubescentibus.—*Willd. sp. pl. 2. p. 600.*

Icon. *Bot. mag.*  
car. 2. t. 98.  
On the sides of  
June, July.  
high, very ele  
in the moun  
genus are ever

1. E. ramis petiolis  
ovatis integer  
*pl. 2. p. 615.*  
Icon. *Bot. rep.*  
*alm. t. 107. f.*  
On shady rocks  
 $\frac{1}{2}$ . May—Jul  
plant; flower

1. R. arborescens;  
umbellis term  
corollis campa  
 $\alpha$ . R. foliis ellipti  
subtus albican  
rotundis, styl  
do-roseis.  
R. maximum. *l*  
*Lam. illustr.*  
 $\beta$ . R. foliis cuneato  
tus pallidiorib  
stylo stamin  
albidis.  
 $\gamma$ . R. altissimum;  
niusculis basi  
laciniis oblon  
giore, floribu  
 $\alpha$ . In the mount  
Carolina,  $\beta$   
and Delaware  
ginia and Ca  
 $\beta$ . July, Aug  
shrub, with  
feet high; fl

Icon. Bot. mag. 175. Wangh. amer. t. 24. f. 50. Catesb.  
car. 2. t. 98.

On the sides of stony hills : Canada to Carolina.  $\text{h}$ .  
 $\text{J}$ . June, July.  $v. v.$  A shrub from three to eight feet  
high, very elegant when in flower ; called *Laurel* or  
in the mountains *Callico-bush*. The whole of this  
genus are evergreen.

## 360. EPIGAEA. Gen. pl. 748.

1. E. ramis petiolis nervo que hirsutissimis, foliis cordato-  
ovatis integerrimis, corollis cylindricis.—Willd. sp.  
pl. 2. p. 615.

Icon. Bot. rep. 102. Lam. illustr. t. 367. f. 1. Pluk.  
alm. t. 107. f. 1.

On shady rocks and in stony woods : Canada to Carolina.  
 $\text{h}$ . May—July.  $v. v.$  A small trailing and creeping  
plant ; flowers white, tinged with red, very fragrant.

*repens.*

## 361. RHODODENDRON. Gen. pl. 746.

1. R. arborescens ; foliis oblongis acutis subtus discoloribus,  
umbellis terminalibus, calycis laciniis ovalibus obtusis,  
corollis campanulatis.—Willd. sp. pl. 2. p. 606.

*maximum.*

- $\alpha$ . R. foliis elliptico-oblongis convexis basi obtusiusculis  
subtus albicantibus vel ferrugineis, corollæ laciniis sub-  
rotundis, stylo staminibus vix longiore, floribus albi-  
do-roseis.

*roseum.*

- R. maximum. Schmidt arb. t. 121. Bot. mag. 951. bona.  
Lam. illustr. 364.

*album.*

- $\beta$ . R. foliis cuneato-lanceolatis planis basi sensim acutis sub-  
tus pallidioribus, corollæ laciniis rotundato-oblongis,  
stylo staminibus vix longiore, floribus minoribus  
albidis.

- $\gamma$ . R. altissimum ; foliis majoribus oblongo-ellipticis pla-  
niusculis basi obtusiusculis utrinque viridibus, corollæ  
laciniis oblongis obtusis, stylo staminibus paulo lon-  
giore, floribus purpurascensibus.

*purpureum.*

- $\alpha$ . In the mountains, near rivulets and lakes : Canada to  
Carolina,  $\beta$ . In shady cedar-swamps of New Jersey  
and Delaware.  $\gamma$ . On the highest mountains of Vir-  
ginia and Carolina, near lakes.  $\text{h}$ .  $\alpha$ . June, July.  
 $\beta$ . July, Aug.  $\gamma$ . May, June.  $v. v.$  This elegant  
shrub, with its varieties, grows from four to twenty  
feet high ; flowers large and in great abundance. The

## DECANDRIA MONOGYNIA. Rhododendron.

variety  $\gamma$ . approaches near to *R. ponticum*, but is different in its foliaceous calyx and otherwise; it occupies the borders of lakes on high mountains, and grows to an immense size; its stem is often found eighteen inches and more in diameter, and its foliage triple the size of any other species: it might be considered as a distinct species. The only plant of this sort I have seen in England is in the garden of James Vere, Esq. at Kensington Gore. The leaves of var.  $\beta$ . are so distinct from var.  $\alpha$ . that it also has claims to be a species.

*punctatum.*

2. *R. foliis ovali-lanceolatis utrinque acutis glabris subtus ferrugineis resinoso-punctatis, umbellis terminalibus, pedicellis brevibus, dentibus calycis brevissimis, corollis infundibuliformibus, laciniis ovatis subundulatis, capsulis elongatis.* — Willd. sp. pl. 2. p. 607.

*R. minus.* Mich. fl. amer. 1. p. 258.

Icon. Bot. rep. 36. Vent. hort. cels 15.

On the mountains of Carolina, particularly on the head-water of the Savannah river.  $\frac{1}{2}$ . July, Aug. v. v. An elegant shrub, not near the size of No. 1; flowers pink-red and smaller.

*catawbiense.*

3. *R. foliis brevi-ovalibus utrinque rotundato-obtusis glabris subtus discoloribus, umbellis terminalibus, laciniis calycis elongato-oblongis, corollis campanulatis.* — Mich. fl. amer. 2. p. 258.

On the high mountains of Virginia and Carolina, particularly on the head-waters of the Catawba river.  $\frac{1}{2}$ . May, June. v. v. s. fl. The flowers are said to be scarlet.

## 362. RHODORA. Gen. pl. 745.

*canadensis.*

1. *Rhodora.* Willd. sp. pl. 2. p. 603.

Icon. Bot. mag. 474. Lam. illustr. 364. L'Herit. stirp. 1. t. 68.

In bogs of Canada, and on the high mountains of New York and Pennsylvania.  $\frac{1}{2}$ . April, May, v. v. A shrub about two feet high, of the appearance of *Azalea*. Leaves elliptic, entire, of a glaucous hue, and pubescent underneath. Flowers in terminal umbels, pale purple.

1. *P. foliis rotund serrulatis, sp. clinato.* — W Icon. Fl. dan. f. 1.

In dry stony or June, July. state, with fi

2. *P. foliis renif latus vaginante versis,* p. 251.

In Canada. M vania, in be yellowish-g

3. *P. foliis subrot spica florib sp. pl. 2. p.* Icon. Fl. dan.

In Canada and New York. foregoing sp peltated.

4. *P. foliis rotund secundis,* p. Icon. Fl. dan.

In sandy barre July. v. v. gibbous.

5. *P. foliis subrot Willd. sp.* Icon. Fl. dan.

In shady woo York. 4 large white radiated. by the nam

363. PYROLA. *Gen. pl. 572.*

1. *P. foliis rotundatis sive subrotundo-ovalibus obsolete rotundifolia.*  
serrulatis, spica floribus undique versis, pistillo de-  
clinato.—*Willd. sp. pl. 2 p. 621.*  
Icon. *Fl. dan. t10. Engl. bot. 213. Lam. illustr. t. 367.*  
*f. 1.*  
In dry stony or sandy woods : Canada to Carolina. ♀.  
June, July. v. v. Flowers white; the stigma is ro-  
tate, with five tubercles.
2. *P. foliis reniformibus, scapo squamis nonnullis convo-  
lutis vaginantibus remote vestito, spica floribus undi-  
que versis, pistillo declinato.* *Mich. fl. amer. 1.*  
*p. 251.*  
In Canada. *Michaux.* On the mountains of Pensyl-  
vania, in beech-woods. *P.* ♀. July. v. v. Flowers  
yellowish-green.
3. *P. foliis subrotundo-ovalibus serrulatis, scapo subnudo,*  
spica floribus undique versis, pistillo recto.—*Willd.*  
*sp. pl. 2. p. 621.*  
Icon. *Fl. dan. 55. Engl. bot. 158.*  
In Canada and Labrador ; rare in the western parts of  
New York. ♀. July. v. v. Flowers smaller than the  
foregoing species, white, with a tinge of red; stigma  
peltated.
4. *P. foliis rotundato-ovatis acutis serratis, spicæ floribus*  
secundis, pistillo recto.—*Willd. sp. pl. 2. p. 621.*  
Icon. *Fl. dan. 402. Engl. bot. 517.*  
In sandy barren woods : Canada to New Jersey. ♀.  
July. v. v. Flowers greenish-white; stigma peltate,  
gibbous.
5. *P. foliis suborbiculatis serratis, scapo unifloro, stylo recto.*  
*Willd. sp. pl. 2. p. 622.*  
Icon. *Fl. dan. t. 8. Engl. bot. 146.*  
In shady woods and sphagnum swamps : Canada to New  
York. ♀. July. v. v. This pretty little plant has  
large white and fragrant flowers; the stigma is five-  
radiated. All these species are evergreen, and known  
by the name of *Winter-green.*

*asarifolia.**minor.**secunda.**uniflora.*

## 364. CHIMAPHILA.

*maculata.*

1. C. foliis lanceolatis basi rotundatis remote serratis fascia longitudinali discolore notatis, scapo 2-3-floro, filamentis lanuginosus.

*Pyrola maculata. Willd. sp. pl. 2. p. 622.**Icon. Bot. mag. 897. Pluk. mant. t. 349. f. 4.*

In shady gravelly or sandy woods: Canada to Carolina.

$\frac{1}{4}$ . July. v. v. Flowers white; leaves handsomely variegated. This plant is in high esteem for its medicinal qualities among the natives, they call it *Sip-si-sewa*. I have myself been witness of a successful cure made by a decoction of this plant, in a very severe case of hysterics. It is a plant eminently deserving the attention of physicians.

*corymbosa.*

2. C. foliis cuneato-lanceolatis basi acutis serratis concoloribus, scapo corymbifero, filamentis glabris.

*P. umbellata. Willd. sp. pl. 2. p. 622.**Icon. Bot. mag. 778. Lam. illustr. t. 367. f. 2.*

In dry woods, frequent: Canada to Virginia.  $\frac{1}{4}$ . July. v. v. Flowers greenish-white, tinged with red; anthers purple. Both species are handsome evergreens, and known by the name of *Winter-green*. I have ventured to form a new genus of these two species of *Pyrola*, so very distinct in habit as well as character. The name I have given is taken from the common name *Winter-green*:  $\chi\mu\mu\alpha\ \chi\mu\mu\alpha$ , and  $\varphi\lambda\omega\varsigma\ amicus$ .

365. LEDUM. *Gen. pl. 744.**palustre.*

1. L. foliis linearibus margine revolutis subtus ferrugineo-tomentosis, staminibus denis corolla longioribus. *Lam. encycl. 3. p. 459. Willd. sp. pl. 2. p. 602.*

*Icon. Fl. dan. 1031. Duham. arb. 1. t. 67. Schmidt arb. 163.**decumbens.*

- $\beta$ . L. spithameum decumbens. *Ait. kew. ed. 2. v. 3. p. 48.*  
In swamps of Canada and round mountain-lakes of New York.  $\beta$ . Near Hudson's Bay.  $\frac{1}{2}$ . April, May. v. v.  
Flowers white.

*latifolium.*

2. L. foliis linear-oblongis margine replicatis subtus ferrugineo-tomentosis, staminibus subquinis corollam æquantibus. *Lam. encycl. 3. p. 459. Willd. sp. pl. 2. p. 602.*

*L. grœnlandicum. Retz. obs. 4. p. 26.*

*L. palustre. M*  
*Icon. Schmidt*  
*lustr. t. 363*  
*In sphagnum s*  
*A larger and*  
*one; comm*

1. A. foliis pusilli  
bis terminali  
*Ledum buxifol*  
*p. 65.*  
*Ledum thymif*  
*Icon. Bergius*  
*illustr. t. 36*  
*In pine-barren*  
*Carolina. A*  
*shrub grows*  
*sometimes a*  
*dance of its*  
*propose it a*  
*lated to *Le**  
*to *Azalea* p*  
*been derive*  
**myrtus pumi**  
**myrtle amor**

367.

1. C. foliis cuneat  
utrinque gla  
 $\Delta$  cibus bracte  
*p. 619.*

*C. alnifolia den*  
*Icon. Schmidt*  
*Catesb. car.**In swamps, fr*  
*July—Sept.*  
*high; flowe*  
*species are.*

2. C. foliis cuneat  
subtus albo  
bracteatis vi

*C. alnifolia pul*

*L. palustre*. *Mich. fl. amer.* 1. p. 259.  
*Icon. Schmidt arb.* 164. *Jacq. ic.* 3. t. 464. *Lam. illustr.* t. 363. f. 1.  
 In sphagnum swamps of Canada.  $\frac{1}{2}$ . April, May. v. v.  
 A larger and broader leaved plant than the foregoing one; commonly called *Labrador Tea*.

## 366. AMMYRSINE.

1. *A. foliis pusillis convexis ovalibus glabris lucidis, corymbis terminalibus congestis.* *buxifolia.*

*Ledum buxifolium*. *Willd. sp. pl.* 2. p. 602. *Ait. kew.* 2. p. 65.

*Ledum thymifolium*. *Lam. encycl.* 3. p. 459.  
*Icon. Bergius act. petrop.* 1777. p. 1. t. 3. f. 2. *Lam. illustr.* t. 363. f. 2.

In pine-barrens of New Jersey and on the mountains of Carolina.  $\frac{1}{2}$ . May, June. v. v. This elegant little shrub grows to the height of about six inches, and sometimes a foot; the delicacy of its leaves and abundance of its white flowers are highly ornamental. I propose it as a distinct genus, being on one side related to *Ledum*, on the other approaching very near to *Azalea procumbens*, vid. p. 154. The name has been derived from *αμυος sabulum*, and *μυρτινη*, *myrtus pumilus*; being known by the name of *Sand-myrtle* among the inhabitants of New Jersey.

367. CLETHRÁ. *Gen. pl.* 751.

1. *C. foliis cuneato-obovatis acutis superne grosse-serratis utrinque glabris concoloribus, racemis spicatis simplicibus bracteatis cano-tomentosis.* —*Willd. sp. pl.* 2. p. 619.

*C. alnifolia denudata*. *Ait. kew.* 2. p. 73.  
*Icon. Schmidt arb.* 47. *Lam. illustr.* 369. *Mill. ic.* 28. *Catesb. car.* 1. t. 66.

In swamps, frequent: New England to Virginia.  $\frac{1}{2}$ . July—Sept. v. v. A shrub from three to four feet high; flowers white, as likewise all the following species are.

2. *C. foliis cuneato-obovatis acutis superne tenue-serratis subtus albo-tomentosis, racemis spicatis simplicibus bracteatis villosso-tomentosis.* —*Lam. encycl.* 2. p. 46.

*C. alnifolia pubescens*. *Ait. kew.* 2. p. 73.

*alnifolia.*

*tomentosa.*

## DECANDRIA MONOGYNIA. Clethra.

In swamps of Virginia and Carolina. ♀. July, Aug. v. v. Though generally considered a variety of No. 1, I take it to be a genuine species, as I have seen seedling plants of both species keeping constantly their distinction.

- scabra.* 3. C. foliis lato-cuneato-obovatis acutis utrinque scabris grosse-serratis; serraturis uncinatis, racemis spicatis subpaniculatis bracteatis tenui-tomentosis.—*Pers. syn. 1. p. 483?*  
In the western parts of Georgia. Lyon. ♀. July. v. s. in *Herb. Lyon.*
- paniculata.* 4. C. foliis anguste cuneato-lanceolatis acutis acuminato-serratis utringue glabris, panicula terminali elongata racemiflora albo-tomentosa.—*Willd. sp. pl. 2. p. 620. Ait. kew. 2. p. 73.*  
In Carolina. Bartram. ♀. July, Aug. v. s. in *Herb. Banks.*
- acuminata.* 5. C. foliis ovalibus acuminatis basi obtusiusculis serratis utrinque glabris subtus subglauca, racemis spicatis subsolitariis bracteatis albo-tomentosis.—*Mich. fl. amer. 1. p. 260.*  
C. montana. *Bartr. catal.*  
On the high mountains of Carolina. ♀. July, Aug. v. v. Grows to the size of a tree; leaves large; flowers resembling those of No. 1.

## 368. MYLOCARIUM. Willd. enum.

- ligustrinum.* 1. Mylocarium. Willd. l. c.  
Cliftonia. *Herb. Banks.*  
On the dry borders of swamps: Georgia. ♀. May, June. v. v. An elegant evergreen shrub, growing to the height of eight or twelve feet; flowers white, sweet-scented, in terminal spikes; seeds resembling buck-wheat, from which it has been called *Buck-wheat Tree.*  
*Frutex sempervirens, 8-pedalis et ultra. Caulis ramosissimus, cortice cinereo. Rami sparsi, glabri, fusi. Folia sparsa in summitate rami lorum conferta, patentia, subsessilia, cuneato-lanceolata, acutiuscula, integerrima, utrinque glabra, laevigata, plana, coriacea, subtus glauca. Racemi terminales, solitarii, simplices seu interdum basi ramosi, erecti, bracteati. Bracteæ solitariae, oblongæ, acutæ, membranaceæ, deciduae,*

Aug.  
o. 1, I  
edling  
air di-

scabris  
picatis  
- Pers.

v. s.

inato-  
ongata  
o. 020.

Herb.

erratis  
picatis  
h. fl.

, Aug.  
large;

May,  
owing  
white,  
mbling  
-wheat

ramo-  
fusci.  
paten-  
la, in-  
riacea,  
splices  
*racteæ*  
ciduæ,

## DECANDRI

pedicello bre  
celli filiforme  
læ subulatæ.  
nimus, viridi  
*Petala oblonga*  
guibus erectis  
receptaculo i  
breviora, inf  
*Antheræ sub*  
4-gonum, vi  
magnitudine  
4-locularis.

369. M

1. *M. scapo spiciflora* late p. 573.  
Icon. *Fl. dan.* t. 16. f. 20.  
Parasitic, on the moist places v. v. The appearance; it
2. *M. scapo spiciflora* nosis.—*Mic*  
In similar place v. v. Of a
3. *M. scapo breviflora* cernuo syn. *Moris.*  
Icon. *Pluk. alh.* In similar situa June. v. v. It ing together
4. *M. scapo elongata* bus, flore eructa  
Icon. *Moris. h.* In shady woods. The flower twelve stami



Hylocarpium ligustrinum.

pedicello breviores, albidae, apice ferrugineae. *Pedicelli* filiformes, rubelli, medio bibracteolati. *Bracteoleae* subulatae. *Flores* albi, suaveolentes. *Calyx* minus, viridis, albo-marginatus : laciniis subrotundis. *Petala* oblongo-obovata, obtusa, subunguiculata : unguibus erectis : laminae concavae, patentes. *Filamenta* receptaculo inserta, 5. alterna longiora, corolla vix breviora, inferne dilatata, crassa, superne subulata. *Antherae* subrotundae, flavae. *Germen* prismatico-3-4-gonum, viride. *Stigma* sessile, crassum. *Capsula* magnitudine et figura *Polygoni Fagopyri*, 3-4-alata, 4-locularis. *Semina* solitaria.

369. MONOTROPA. *Gen. pl. 737.*

1. *M. scapo spicifloro, squamis floribusque extus glabris ; floribus lateralibus octandris.* — *Willd. sp. pl. 2. p. 573.*  
Icon. *Fl. dan. 232. Engl. bot. 69. Moris. hist. 3. s. 12. t. 16. f. 20. Pluk. alm. t. 209. f. 5.*  
Parasitic, on the roots of beech and other trees, in shady moist places : Canada to Pennsylvania.  $\frac{1}{2}$ . June, July.  
*v. v.* The whole plant is of a brownish-yellow appearance ; it is smaller than the European species.
2. *M. scapo spicifloro, bracteis floribusque undique lanuginosae.* — *Mich. fl. amer. 2. p. 266.*  
In similar places : Pennsylvania to Carolina.  $\frac{1}{2}$ . July.  
*v. v.* Of a light tan-colour.
3. *M. scapo breviore crasso unifloro, squamis approximatis, flore cernuo decandro.* — *Willd. sp. pl. 2. p. 573. excl. syn. Moris.*  
Icon. *Pluk. alm. t. 209. f. 2. Catesb. car. 1. t. 36. bona.*  
In similar situations : New York to Carolina.  $\frac{1}{2}$ . May, June. *v. v.* In general it appears in large clusters growing together ; the whole plant is white.
4. *M. scapo elongato rectissimo unifloro, squamis distantibus, flore erecto decandro.* — *Mich. fl. amer. 1. p. 266.*  
Icon. *Moris. hist. 3. s. 12. t. 16. f. 5. bona.*  
In shady woods : Virginia and Carolina.  $\frac{1}{2}$ . June. *v. v.*  
The flower is larger than No. 3, frequently with twelve stamens.

370. DIONÆA. *Gen. pl. 729.**muscipula.*

1. *Dionæa*. *Willd. sp. pl. 2. p. 574.*

*Icon. Bot. mag. 785. Vent. malm. 29.*

In bogs of North Carolina, near Wilmington.  $\frac{1}{4}$ . July, Aug. v. v. This singular and interesting plant, known by the name of *Venus's Fly-trap*, is fully described by the authors above quoted.

371. JUSSIEUA. *Gen. pl. 741.**grandiflora.*

1. *J. erecta*; foliis integerrimis, inferioribus spathulatis, superioribus lanceolatis, floribus decandris, pedunculis calycibusque villosis.—*Mich. fl. amer. 1. p. 267.*

*Icon. Abbot. insect. 1. t. ....*

In swamps and ponds of Carolina and Georgia, near Savannah.  $\frac{1}{4}$ . July. v. s. Roots creeping; flowers large, yellow.

*erecta.*

2. *J. erecta*, glabra; foliis lanceolatis, floribus octandris sessilibus.—*Willd. sp. pl. 2. p. 578.*

*Icon. Sloan. jam. 1. t. 11. f. 1.*

In swamps of Virginia and Carolina, about the Dismal swamp.  $\frac{1}{4}$ . Aug. v. v. Flowers yellow, smaller than the foregoing one.

*subcaulis.*

3. *J. repens*, glabra; foliis linear-lanceolatis repando-dentatis, floribus solitariis octandris pedunculatis, filamenti alternis brevissimis, petalis obovatis.

On the banks of the Missouri. *M. Lewis.*  $\frac{1}{4}$ . June. v. s. Flowers small, yellow.

372. GYMNOCLADUS. *Gen. pl. 1537.**canadensis.*

1. *G. foliis bipinnatis*, foliolis ovalibus acuminatis pubescentibus.—*Willd. sp. pl. 4. p. 816.*

*Guilandina dioica. Sp. pl. 546.*

*Icon. Mich. fl. amer. 1. p. 51. Mich. arb. 23. Lam. illustr. 823. Duham. arb. 1. t. 103.*

In the forests of Canada, western parts of New York, Tennessee, Ohio, and Kentucky.  $\frac{1}{2}$ . May, June. v. v. A large tree, known by the name of *Kentucky Coffee Tree*, and among the Canadians *Chicot*. Flowers white. Seed-pods large, dark brown.

## 373. ACAC

1. *A. inermis*; foliis multijugis spicis globosis minibus lanceolatis. *Mimosa illinoensis*. In extensive naturalized areas.  $\frac{1}{4}$ . June, July. Five stamens.

2. *A. inermis*; foliis multijugis spicis globosis guttulibus falcata. *Mimosa glandulosa*. *Icon. Vent. choix*. On the banks of rivers.  $\frac{1}{4}$ . July. v. v.

## 374. SCHRAN

1. *S. aculeata*; foliis multijugis, leguminibus. *Willd. sp. pl. 4*. *Mimosa horridula*. *Mimosa Intsia*. In old fields and pastures. June, July. v. It is known by

## 375. C

1. *C. glabra*; foliis majoribus, glandulis paucifloris rectis.—*Willd. Dill. elth. t.* In sandy fields of v. v. The flowers are yellow.

2. *C. glabra*; foliis scabris, exterioribus dunculis multifidis.

373. ACACIA. *Willd. sp. pl. 4. p. 1049.*

- 4. July,*  
*g plant,*  
*fully de-*
- thulatis,*  
*dunculus*  
*37.*
- ia, near*  
*flowers*
- octandris*
- Dismal-*  
*smaller*
- do-den-*  
*filamen-*
- . June.*
- pubes-*
- Lam.*
- York,*  
*, June.*
- Kentucky*  
*Flow-*
1. A. inermis ; foliis bipinnatis, partialibus 8-jugis, propriis multijugis, glandula inter par infimum partialium, spicis globosis solitariis pedunculatis axillaribus, leguminibus lanceolatis rectis. *Willd. sp. pl. 4. p. 1071.* *Mimosa illinoensis. Mich. fl. amer. 2. p. 254.* In extensive natural meadows of Illinois and Kentucky.  
*4. June, July. v. v.* Flowers white, with only five stamens.
  2. A. inermis ; foliis bipinnatis, partialibus 12-jugis, propriis multijugis, glandula inter omnia paria partialium, spicis globosis pedunculatis solitariis axillaribus, leguminibus falcatis. *Willd. sp. pl. 4. p. 1071.* *Mimosa glandulosa. Mich. fl. amer. 2. p. 254.* Icon. *Vent. choix 27.* On the banks of the rivers Tennessee and Mississippi.  
*4. July. v.v.* Flowers white, pentandrous.

374. SCHRANKIA. *Willd. sp. pl. 4. p. 1041.*

1. S. aculeata ; foliis bipinnatis, partialibus 6-jugis, propriis multijugis, leguminibus acuminatis, caule pentagono. *Willd. sp. pl. 4. p. 1043.* *Mimosa horridula. Mich. fl. amer. 2. p. 254.* *Mimosa Intsia. Walt. fl. car. 252.* In old fields and meadows : Virginia to Florida. *4. June, July. v. v.* Flowers in globular spikes, red.  
*It is known by the name of Sensitive Briar.*

375. CASSIA. *Gen. pl. 700.*

1. C. glabra ; foliis 3-jugis obovatis subretusis, exterioribus majoribus, glandula subulata inter inferiora, pedunculis paucifibris axillaribus, leguminibus subtetragonis rectis.—*Willd. sp. pl. 2. p. 515.* Icon. *Dill. elth. t. 63. f. 73. Mill. dict. t. 82.* In sandy fields of Georgia, near Savannah. *◎. July. v. v.* The flowers of all the species mentioned here are yellow.
2. C. glabra ; foliis sub-5-jugis ovato-lanceolatis margine scabris, exterioribus majoribus, glandula basilari, pedunculis multifloris axillaribus et subpaniculato-termi-

*Tora.**occidentalis.*

- nalibus, leguminibus linearibus falcatis. *Willd. sp. pl. 2.* p. 518.  
 Icon. *Comm. hort. 1. t. 26.* *Sloan. jam. 2. t. 175. f. 3*  
~~E~~ 4.  
 In old fields: Virginia to Carolina. 4. h. June—  
 Aug. v. v.
- ligustrina.* 3. C. tenuissime pubescens; foliis 7-jugis lanceolatis, ex-  
 timis minoribus, glandula basilari, pedunculis terminalibus subpaniculatis, leguminibus oblongis subfalcatis.—*Willd. sp. pl. 2. p. 523.*  
 Icon. *Dill. elth. t. 259. f. 338. Mart. cent. 21.*  
 In cultivated grounds: Virginia to Georgia. h. 4.  
 July, Aug. v. v.
- linearis.* 4. C. glabra; foliis 6-jugis ovali-lanceolatis angustatis acutissimis, glandula basilari, pedunculis subbifloris axillaribus et terminalibus, legumine lineari. *Mich. fl. amer. 1. p. 261.*  
 In Carolina. *Michaux. 4. May, June. +.*
- marilandica.* 5. C. glabriuscula; foliis 8-jugis lanceolato-oblongis mucronatis subæqualibus, glandula petiolari obovata, racemis axillaribus et paniculato-terminalibus, leguminibus linearibus arcuatis glabris.—*Willd. sp. pl. 2. p. 524.*  
 Icon. *Dill. elth. t. 260. f. 339. Mart. cent. 23.*  
 In low gravelly soil, on the banks of rivers: New York to Carolina. 4. June—Aug. v. v.
- Chamæcris-  
 ta.* 6. C. glabriuscula; foliis multijugis linearibus, glandula petiolari subpedicellata, floribus subgeminis, pedicellis longis, petalis duobus maculatis, leguminibus pubescentibus.—*Willd. sp. pl. 2. p. 528.*  
 Icon. *Bot. mag. 107. Comm. hort. 1. t. 37.*  
 In cultivated grounds of Virginia and Carolina. ①.  
 July, Aug. v. v. The flowers large and handsome, with two purple spots at the bottom. There is a variety, the stem of which is covered with long brown hair.
- fasciculata.* 7. C. glabriuscula; foliis ad summum 9-jugis, glandula petiolari sessili submedia, fasciculis multifloris lateribus, petalis staminibusque concoloribus, leguminibus glabris arcuatim adscendentibus.—*Mich. fl. amer. 1. p. 262.*  
 In dry fields: New York to Carolina. ①. June—Aug.  
 v. v. The flowers are smaller than the foregoing species.
8. C. patula, pub-  
 dula petiolari  
 axillaribus 2-  
~~pl. 2. p. 529.~~  
 Icon. *Pluk. aln*  
 In sandy fields :  
 Aug. v. v.  
 sitive.
9. C. foliis multi-  
~~Willd. sp. pl.~~  
 In dry fields of

## 376. POD.

1. P. glaberrima; i-  
 integrerimis,  
 leguminibus s-  
*Crotalaria perfoli*  
*Sophora perfolia*  
*Rafnia perfoliata*  
 Icon. *Dill. elth.*  
 In dry barren fie-  
 v. v. Flower
2. P. caule foliis ca-  
 sessilibus, foli-  
 minatis, floril-  
*Mich. fl. amer*  
*Sophora lanceola*  
 In Carolina and  
 yellow.
3. P. caule foliis su-  
 tibus, foliis tem-  
 obtusis, stipul-  
 cato, calycibu-  
 p. 264.  
*Sophora villosa.*  
 In low sandy g-  
 4. June, Jul-  
 lupine very m-
4. P. glabra; foliis oblongo-cunea-  
 petiolo duplo  
 leguminibus a-  
*P. australis. Wil*

8. C. patula, pubescens; foliis multijugis linearibus, glandula petiolaris pedicellata, pedunculis brevibus supra axillaribus 2-3-floris, floribus pentandris.—Willd. sp. nictitans.  
pl. 2. p. 529.  
Icon. Pluk. alm. t. 314. f. 5.  
In sandy fields: New England to Carolina. ☽. June—  
Aug. v. v. The leaves are in a slight degree sensitive.
9. C. foliis multijugis eglandulosus, caule procumbente. procumbens.  
Willd. sp. pl. 2. p. 530.  
In dry fields of Virginia. ☽. +. Flowers small.

## 376. PODALYRIA. Lam. illustr. t. 327.

1. P. glaberrima; foliis simplicibus perfoliatis suborbiculatis integerrimis, floribus axillaribus solitariis unifloris, leguminibus subglobosis.—Mich. fl. amer. 1. p. 263.  
Crotalaria perfoliata. Sp. pl. 1003.  
Sophora perfoliata. Walt. fl. car. 135.  
Rafnia perfoliata. Willd. sp. pl. 3. p. 949.  
Icon. Dill. elth. t. 102. f. 122.  
In dry barren fields: Carolina and Georgia. ♀. July.  
v. v. Flowers small, yellow.
2. P. caule foliis calycibusque pubescentibus, foliis ternatis sessilibus, foliolis lanceolatis obtusis, stipulis setaceis minutis, floribus axillaribus solitariis pedicellatis.—Mich. fl. amer. 1. p. 263.  
Sophora lanceolata. Walt. fl. car. 135.  
In Carolina and Georgia. Michaux. ♀. +. Flowers yellow.
3. P. caule foliis subtus et margine calycibusque pubescentibus, foliis ternatis subsessilibus, foliolis ovali-oblongis obtusis, stipulis linearibus, racemo terminali subspicato, calycibus subsemi-4-fidis.—Mich. fl. amer. 1. p. 264.  
Sophora villosa. Walt. fl. car. 134.  
In low sandy grounds: Virginia and North Carolina.  
♀. June, July. v. v. Flowers yellow; resembles a lupine very much.
4. P. glabra; foliis ternatis brevissime petiolatis, foliolis oblongo-cuneatis obtusis, stipulis lanceolatis acutis petiolo duplo longioribus, racemis spicatis elongatis, leguminibus acuminatis.—Mich. fl. amer. 1. p. 264.  
P. australis. Willd. sp. pl. 2. p. 503.

x 2

- Sophora australis.* *Syst. veg.* 325.  
Icon. *Bot. mag.* 509. *Vent. hort. cels* 46. *Trew pl. rar.*  
t. 14.  
On the banks of rivers: Virginia and Carolina, particularly in the western districts. ♀. June, July. v. v. Flowers blue.
- alba.* 5. *P. ramis divaricatis flexuosis glabris, foliis ternatis petiolatis, foliolis oblongis obtusis glabrioculis, stipulis subulatis petiolo brevioribus, racemo spicato elongato, leguminibus obovatis.* — *Willd. sp. pl. 2. p. 503.*  
*Sophora alba.* *Syst. veg.* 325.  
*Crotalaria alba.* *Sp. pl. 2. p. 1006.*  
Icon. *Bot. mag.* 1177. *Mart. cent.* 44.  
In the western parts of Virginia and Carolina, on river banks. ♀. June, July. v. v. Flowers white.
- mollis.* 6. *P. caule foliis calycibusque minutim pubescentibus, foliis ternatis, foliolis subrhomboideo-lanceolatis, stipulis foliaceis lanceolatis, racemo spicato terminali, calycis dentibus acutis.* — *Mich. fl. amer. 1. p. 264.*  
In the county of Mecklenburg in Upper Carolina. *Michaux.* ♀. +.
- tinctoria.* 7. *P. glaberrima, ramosissima, microphylla; foliis ternatis subsessilibus, foliolis cuneato-obovatis rotundato-obtusis, stipulis obsoletis oblongis acutis petiolo multo brevioribus, racemis spicatis terminalibus, leguminibus ovatis longe-stipitatis.* — *Willd. sp. pl. 2. p. 503.*  
*Sophora tinctoria.* *Sp. pl.* 534.  
Icon. *Bot. mag.* 1099. *Lam. illustr. t. 327. f. 1.*  
In woods, on dry hills: Canada to Carolina. ♀. July, Aug. v. v. Flowers yellow; called *Wild Indigo*.

## 377. CERCIS. Gen. pl. 696.

- canadensis.* 1. *C. foliis subrotundo-cordatis acuminatis, ad axillas nervorum villosis, stipulis minutis, leguminibus brevi-stipitatis.* — *Willd. sp. pl. 2. p. 508.*  
Icon. *Schmidt arb. 21. Mill. ic. 2.*
- pubescens.* β. *C. foliis subrotundis acutis subtus pubescentibus.*  
On the banks of rivers: Canada to Virginia. β. In Georgia. h. May. v. v. A handsome tree, generally called *Judas-tree*. Flowers crimson.

1. *V. foliis pinnatis, glabris, racem tiolatis planis.*—  
On the mountair June. v. v. s. A handsome tree much. Its ba

1. *H. foliis oblong subtus glabris,*  
*H. arborescens.* —  
Icon. *Bot. mag.* 4 ic. 251.  
On the banks of nia to Virginia. five feet high ;
2. *H. foliis lato-ovati tatis subtus gla*  
*H. vulgaris.* *Mich.*  
On the mountain This, though n distinct species, robust appearai
3. *H. foliis ovatis ac tosis: serraturi fl. amer. 1. p. 2*  
*H. radiata.* *Walt.*  
Icon. *Lam. illust.*  
On the head-wat see, &c. h. shrub.
- 4 *H. foliis oblongis ac tosis, cymis ra*  
*sp. pl. 2. p. 634*  
*H. radiata.* *Smith*



378. VIRGILIA. *Lam. illustr. t. 326.*

1. *V. foliis pinnatis, foliolis alternis ovatis brevi-acuminatis glabris, racemis elongatis pendulis, leguminibus petiolatis planis.*—*Mich. arb. t. ....*

*lutea.*

On the mountains between Georgia and Tennessee.  $\frac{1}{2}$ . June. *v. v. s. fl.; v. s. c. fl. et fr. in Herb. Lyon.* A handsome tree, resembling *Cytisus Laburnum* very much. Its bark gives a beautiful yellow dye.

## II. DIGYNIA.

379. HYDRANGEA. *Gen. pl. 760.*

1. *H. foliis oblongo-ovatis basi obtusis acuminatis dentatis subtus glabris, cymis nudis.*

*vulgaris.*

*H. arborescens. Willd. sp. pl. 2. p. 633.*

*Icon. Bot. mag. 437. Lam. illustr. t. 370. f. 1. Mill. ic. 251.*

On the banks of rivulets, in shady situations: Pennsylvania to Virginia.  $\frac{1}{2}$ . July, Aug. *v. v.* A shrub about five feet high; flowers white.

2. *H. foliis lato-ovatis basi subcordatis acuminatis grosse dentatis subtus glabris, cymis subradiatis.*

*cordata.*

*H. vulgaris. Mich. fl. amer. 1. p. 268.*

On the mountains of Carolina.  $\frac{1}{2}$ . June, July. *v. v.*

This, though nearly allied to the preceding, is a very distinct species, and easily known from that by its more robust appearance and larger leaves.

3. *H. foliis ovatis acuminatis dentatis subtus niveo-tomentosis: serraturis mucronatis, cymis radiatis.*—*Mich. fl. amer. 1. p. 268.*

*nivea.*

*H. radiata. Walt. fl. car. 251.*

*Icon. Lam. illustr. t. 307. f. 2.*

On the head-waters of the Savannah river; in Tennessee, &c.  $\frac{1}{2}$ . July, Aug. *v. v.* A very ornamental shrub.

4. *H. foliis oblongis sinuato-lobatis dentatis subtus tomentosis, cymis radiatis thyrsideo-paniculatis.*—*Willd. sp. pl. 2. p. 634.*

*quercifolia.*

*H. radiata. Smith ic. pict. 12.*

Icon. *Bartr. it. ed. germ. t. 7.*

On banks of rivers: Georgia and Florida.  $\frac{1}{2}$ . July,  
Aug. v. v. A handsome shrub; flowers in large  
clusters, white, changing to red.

380. SAXIFRAGA. *Gen. pl. 764.*

\* *Foliis indivisis radicalibus; caule subnudo.*

*Aizoon.*

1. S. foliis radicalibus aggregatis spathulatis acutiusculis gla-  
bris cartilagineo-dentatis, caule simplici folioso piloso,  
calycibus glabris.—*Willd. sp. pl. 2. p. 639.*

Icon. *Jacq. austr. 5. t. 438. Fl. lap. t. 2. f. 2.*

In Labrador and probably in Canada. *Colmaster.*  $\frac{1}{4}$ .  
June, July. v. s. in *Herb. Dickson.* Not above three  
inches high.

*stellaris.*

2. S. foliis cuneatis serratis margine pilosis, caule nudo  
simplici, petalis acutis, capsula supera.—*Willd. sp.  
pl. 2. p. 644.*

Icon. *Engl. bot. 167. Fl. dan. 23. Jacq. coll. 1. t. 13.*  
In Labrador and Canada. *Colmaster.*  $\frac{1}{4}$ . June, July.  
v. s. in *Herb. Dickson.* Flowers white.

*serpyllifolia.*

3. S. microphylla, erecta; foliis ovalibus glabris, caule  
unifloro oligophyllo, petalis obovatis.  
On the north-west coast. *Nelson.*  $\frac{1}{4}$ . v. s. in *Herb.  
Banks.* Flowers large.

*androsacea.*

4. S. pubescens; foliis spatulato-linearibus petiolatis, caule  
folioso 1-2-floro.

S. androsacea. *Jacq. austr. 389?*

On the north-west coast. *Nelson.*  $\frac{1}{4}$ . v. s. in *Herb.  
Banks.* Flowers white, small.

*bronchialis.*

5. S. stolonifera; foliis imbricatis subulatis planis mucro-  
natis spinoso-ciliatis, caule subnudo paniculato.—  
*Willd. sp. pl. 2. p. 644.*

On the north-west coast. *Nelson.*  $\frac{1}{4}$ . v. s. in *Herb.  
Banks.*

*nivalis.*

6. S. foliis subrotundo-cuneatis antice crenatis in petiolum  
decurrentibus, caule nudo simplici, floribus congestis  
racemosis.—*Willd. sp. pl. 2. p. 645.*

Icon. *Fl. lapp. t. 2. f. 5. Fl. dan. t. 12.*

In Labrador and Canada.  $\frac{1}{4}$ . May, June. v. s. in  
*Herb. Dickson.* Not above two inches high; flowers  
white.

*virginiensis.*

7. S. tota minutim pubescens; foliis ovalibus obtusis cre-  
natis in petiolum decurrentibus, caule subaphyllo pa-

## DECANDRIA DIGYNIA. Saxifraga.

- niculato, ramulis dichotomis, floribus subsessilibus.—  
*Mich. fl. amer. 1. p. 269.*
- S. pilosa. *Herb. Banks.*  
 Icon. *Pluk. alm. t. 222. f. 5.*  
 On rocks and dry hills: New England to Virginia, and  
 on the Carolina mountains.  $\frac{1}{2}$ . May, June. v. v.  
 Variable in size; flowers white.
8. S. foliis reniformibus dentatis utrinque pilosis aveniis,  
 caule nudo paniculato.—*Willd. sp. pl. 2. p. 648.*  
 Icon. *Lapeyr. t. 14. Magn. hort. 88. Moris. hist. 3.*  
 p. 12. t. 9. f. 12.  
 On the north-west coast. *Nelson.*  $\frac{1}{2}$ . v. s. in *Herb. Banks.* Flowers small.
9. S. hirsutissima; foliis elongato-spathulatis acute dentatis,  
 caulibus divaricato-dichotomis, paniculis capillaribus  
 laxis, calyce reflexo, petalis inaequalibus. *Mich. fl. amer. 1. p. 268.*  
 On high mountains of Carolina. *Michaux.* On the  
 Peaks of Otter, Virginia. P.  $\frac{1}{2}$ . June. v. v. About  
 eight inches high; flowers white, elegantly red and  
 yellow punctated. This and the *S. sarmentosa* ought  
 to form a separate genus.
10. S. pubescens; foliis oblongo-lanceolatis utrinque acutis ob- *pensylvanica.*  
 solete denticulatis, caule nudo, panicula oblonga, ramis  
 summitate fasciculifloris.—*Willd. sp. pl. 2. p. 640.*  
 Icon. *Dill. elth. t. 253. f. 3'28.*  
 In wet meadows: New York to Virginia.  $\frac{1}{2}$ . June.  
 v. v. Above a foot high; flowers small, yellowish-green.
11. S. glabriuscula; foliis oblongo-lanceolatis acutis erosio-  
 dentatis, caule nudo, panicula oblonga, ramis divari-  
 cato-ramosissimis laxifloris, pedicellis filiformibus.  
 In stony rivulets on the high mountains of Virginia and  
 Carolina.  $\frac{1}{2}$ . June, July. v. v. Resembles the  
 foregoing one in some respects very much.
- \*\* *Foliis indivisis; caule folioso.*
12. S. cæspitosa; foliis caulinis oppositis imbricatis ovatis *oppositifolia.*  
 obtusis, summis ciliatis, floribus terminalibus solitariis  
 sessilibus.—*Willd. sp. pl. 2. p. 648.*  
 Icon. *Engl. bot. 9. Fl. dan. 34. Fl. lapp. t. 2. f. 1.*  
 In Newfoundland and Labrador.  $\frac{1}{2}$ . May. v. v. v. s.  
 in *Herb. Banks.* Flowers purple, large.

- aizoides.* 13. *S. decumbens*, cæspitosa; foliis caulinis linearibus alternis dentato-ciliatis, panicula pauciflora, petalis longitudine calycis.—*Willd. sp. pl. 2. p. 650.*  
Icon. *Engl. bot. 39. Fl. dan. 72.*  
In Newfoundland and Labrador. *U. v. s. in Herb. Banks.* Flowers yellow.
- setigera.* 14. *S. pubescens*; foliis radicalibus aggregatis spathulatis acutis spinoso-ciliatis, caule folioso subbifloro, calycibus hispidis, flagellis setaceis longissimis in axillis foliorum.  
On the north-west coast. *Nelson. U. v. s. in Herb. Banks.* Flowers small, white. A very singular species.
- \*\*\* *Foliis lobatis.*
- rivularis.* 15. *S. erecta*; foliis palmatis petiolatis: summo spathulato, caule paucifloro, radice fibrosa, germine semi-infero. *Smith fl. brit. 2. p. 454.*  
Icon. *Fl. dan. 118. Fl. lapp. 2. f. 7.*  
In Labrador. *Chateaux. U. June. v. s. in Herb. Banks.* Flowers white, small.
- sibirica.* 16. *S. adscendens*; foliis reniformibus palmatis pilosis, petiolis hirsutis, caulinis sessilibus, pedunculis longissimis filiformibus bifidis nudis.—*Willd. sp. pl. 2. p. 653.*  
Labrador and Newfoundland. *U. v. s. in Herb. Banks.* Flowers white, large.
- pectinata.* 17. *S. cæspitosa*, multicaulis; foliis 3-lobatis, laciniis 3-fidis linearibus acutis glabris, nervo medio exarato, caule folioso racemifloro.  
On the north-west coast. *Menzies. U. v. s. in Herb. Banks.* Resembles *S. petræa*; but the leaves are narrow, and the divisions divaricated.
- cæspitosa.* 18. *S. foliis radicalibus aggregatis carnosis linearibus obtusis integris trifidisve subtus nervosis, caule erecto nudiusculo paucifloro, petalis calyce duplo longioribus.—Willd. sp. pl. 2. p. 456.*  
Icon. *Waldst. hung. 88. Engl. bot. 794.*
- grœnlandica.* β. *S. foliis caulinis palmato-multifidis, laciniis acutis.—Willd. l. c.*  
Icon. *Dill. eth. 353. f. 3 & 9.*  
On the north-west coast. *U. June. v. s. in Herb. Banks.* Flowers pale yellow.
- tricuspidata.* 19. *S. foliis radicalibus aggregatis cuneiformibus ciliatis*

acute tridentatis, caule adscendente racemoso, petalis lanceolatis calyce triplo longioribus. *Willd. sp. pl. 2. p. 657.*

*Icon. Fl. dan. 976.*

On Hudson's Bay. ♀. June. v. s. in Herb. Banks.

Flowers large, white, handsomely punctated.

### 381. TIARELLA. *Gen. pl. 765.*

1. *T. foliis cordatis acute-lobatis dentatis, dentibus mucronatis, scapo racemoso.* — *Willd. sp. pl. 2. p. 659.*

*Icon. Herm. parad. 129. Lam. illustr. 373.*

In shady woods of Canada, and on the high mountains of New York and Pennsylvania. ♀. April, May. v. v. Flowers white.

*cordifolia.*

2. *T. foliis ovatis cordatis acutis brevi-lobatis dentatis, caulinis alternis distantibus, racemo filiformi subspicato, calycibus tubulosis.*

On the north-west coast. *Menzies.* ♀. v. s. in Herb. Banks. More than a foot high, with five or six alternate leaves on the stem.

*Menziesii.*

3. *T. foliis ternatis, foliolis subrhomboideis serratis pilosis, racemis terminalibus, corymbulis florum alternis, calycibus campanulatis.* — *Willd. sp. pl. 2. p. 659.*

On the north-west coast. *Menzies.* ♀. v. s. in Herb. Banks.

*trifoliata.*

4. *T. foliis biternatis, foliolis ovato-cordatis obliquis inciso-lobatis dentatis, caule folioso, panicula terminali divaricato-spiciflora.* *Vent. malm. 54.*

On the mountains of South Carolina. ♀. May. v. v. This plant resembles *Spiraea Aruncus* in habit; flowers yellowish-white. The specimens I had an opportunity to examine were destitute of petals, though they are so distinctly described by Ventenat. It is however probable that some individuals may be petaliferous, while others are apetalous.

*biternata.*

### 382. MITELLA. *Gen. pl. 766.*

1. *M. foliis sublobato-acutangulis dentatis, caule erecto superne opposite diphyllo.* — *Willd. sp. pl. 2. p. 659.*

*Icon. Rob. ic. 80. Schkuhr handb. 1. t. 120. Lam. illustr. t. 373. f. 1.*

*diphylla.*

In shady places, among rocks, near springs. ♀. May.

## DECANDRIA DIGYNIA. Mitella.

v. v. Flowers white, which is the colour of all the following species.

*cordifolia.*

2. *M. foliis orbiculato-reniformibus subduplicato-crenatis lucidis, scapo setaceo aphylo.* *Mich. fl. amer.* 1. p. 270. *Lam. encycl.* 4. p. 185.

Icon. *Lam. illustr.* t. 373. f. 3.

In Canada. *Michaux.* On high mountains of New York and Pennsylvania. ♀. May, June. v. v.

*reniformis.*

3. *M. foliis reniformibus repandis ciliatis, scapo nudo.* — *Lam. encycl.* 4. p. 185.

*M. nuda.* *Willd. sp. pl.* 2. p. 660.

Icon. *Lam. illustr.* t. 373. f. 2. *Gmel. sib.* 4. t. 68. f. 2. On the mountains of New Hampshire and Vermont, and probably in Canada. ♀. June. v. v.

*prostrata.*

4. *M. radice repente, caulis prostratis, foliis alternis rotundato-cordatis subacutis obtuse sublobatis.* *Mich. fl. amer.* 1. p. 270.

In the most southern parts of Canada. *Michaux.* On the mountains of Virginia, near the Sweet-springs. ♀. May, June. v. v.

*grandiflora.*

5. *M. hirsutissima ; foliis rotundato-cordatis obtuse-lobatis dentatis, scapo folio uno altero instructo, floribus pedicellatis, calycibus campanulatis.*

On the north-west coast. *Menzies.* ♀. v. s. in *Herb. Banks.* The flowers are more than four times the size of the other species.

383. SAPONARIA. *Gen. pl. 769.*

*officinalis.*

1. *S. calycibus cylindricis, foliis ovato-lanceolatis.* *Willd. sp. pl.* 2. p. 667.

Icon. *Fl. dan.* 543. *Engl. bot.* 1060. \*

Among rubbish, near buildings and cultivated grounds; sometimes in gravel-pits and on dry hills; probably introduced from Europe. ♀. July, Aug. v. v. Flowers white, frequently semi-double.

384. DIANTHUS. *Gen. pl. 770.*

*Armeria.*

1. *D. floribus aggregatis fasciculatis, squamis calycinis lanceolatis villosis tubum æquantibus.* *Willd. sp. pl.* 2. p. 673.

Icon. *Fl. dan.* 230. *Engl. bot.* 317.

In sand  
Coop  
rose-  
2. D. flori  
dimic  
In Carr

1. S. caly  
Smit  
Icon. J  
In dry  
Jerse

1. C. glab  
acuti  
*Mic*  
Silene  
Icon. J  
In cult  
♀.

2. C. pub  
lance  
p. 6  
Icon. J  
In hill  
♀.  
are

1. S. hirs  
petal  
erect  
Icon.  
On the  
June

- In sandy fields and pine-woods: New Jersey, near Cooper's Ferry. ♂. July. v. v. Flowers small, rose-coloured.
2. D. floribus aggregatis, pedunculis longis, squamis tubo *carolinianus*. dimidio minoribus. *Walt. fl. car.* 140.  
In Carolina. *Walter.* +.

385. SCLERANTHUS. *Gen. pl. 767.*

1. S. calycibus fructus patulis acutis, caulinibus patentibus. *Smith fl. brit.* 2. p. 458. *Willd. sp. pl.* 2. p. 660. *annuus.*  
*Icon. Fl. dan.* 504. *Engl. bot.* 351.  
In dry sandy fields, on the shores of the Delaware, New Jersey opposite Philadelphia. ♂. June, July. v. v.
- 

## III. TRIGYNIA.

386. CUCUBALUS. *Gen. pl. 771.*

1. C. glaber, glaucus, decumbens; foliis oblongo-ovalibus acutis enerviis, calycibus inflato-membranaceis venosis. *Mich. fl. amer.* 1. p. 271. *Willd. sp. pl.* 2. p. 684. *Behen.*  
*Silene inflata. Smith fl. brit.* 467.  
*Icon. Fl. dan.* 914. *Engl. bot.* 164.  
In cultivated grounds, near Quebec, Canada. *Michaux.*  
♀. July. v. v. Flowers white.
2. C. pubescens, erectus; foliis quaterno-verticillatis oval-lanceolatis longissime acuminatis.—*Willd. sp. pl.* 2. p. 686. *stellatus.*  
*Icon. Bot. mag.* 1107.  
In hilly and shady woods: New England to Virginia.  
♀. July, Aug. v. v. Flowers white; the petals are without a corona.

387. SILENE. *Gen. pl. 772.*

1. S. hirsuta; foliis cuneato-oblongis, summis linearibus, petalis subrotundis integerrimis, calycibus fructiferis erectis subspicatum alternis.—*Willd. sp. pl.* 2. p. 691. *quinquevulnera.*  
*Icon. Rob. ic.* 183. *Engl. bot.* 86.  
On the sea-coast of Carolina; near Charlestown. ♂.  
June, July. v. v. Flowers small, crimson.

- pensylvanica*. 2. *S. viscido-pubescent*; foliis cuneatis, caulinis lanceolatis, caulinis in summitate paucifloris, petalis obtusissimis leviter emarginatis subcrenatis. *Mich. fl. amer.* 1. p. 272.  
*S. virginica*, *Willd. sp. pl.* 2. p. 702.  
*S. caroliniana*, *Walt. fl. car.* 142.  
 In dry sandy woods, and on rocks: New York to Virginia. ♀. May, June. v. v. Flowers purple, very handsome.
- virginica*. 3. *S. decumbens*, tota viscosa-pubescent; foliis oblongis margine asperiusculis, panicula dichotoma, petalis bifidis, genitalibus exsertis. *Mich. fl. amer.* 1. p. 272.  
*S. Catesbaei*. *Walt. fl. car.* 141. *Willd. sp. pl.* 2. p. 706.  
 Icon. *Catesb. car.* 2. t. 4. *Pluk. alm.* t. 203. f. 1.  
 In the western parts of Virginia and Carolina, and in the Illinois country. ♀. June, July. v. v. A beautiful species; flowers dark crimson.
- ovata*. 4. *S. foliis ovato-lanceolatis acuminatis glabriusculis, racemo terminali composito, calycibus ovatis, genitalibus exsertis, caule simplici*.  
*Cucubalus polypetalus*. *Walt. fl. car.* 141 ?  
 In the western part of Georgia and Carolina. ♀. July.  
 v. s. in *Herb. Banks*. Flowers white or pale red.
- antirrhina*. 5. *S. foliis lanceolatis subciliatis, pedunculis trifidis, petalis emarginatis, calycibus ovatis*. *Willd. sp. pl.* 2. p. 702.  
 Icon. *Dill. elth.* t. 313. f. 403.  
 In old fields and on the banks of rivers: Pennsylvania to Carolina. ♂. June, July. v. v. Flowers small, white or greenish.
- nocturna*. 6. *S. floribus spicatis alternis sessilibus secundis, petalis bifidis*. *Willd. sp. pl.* 2. p. 692.  
 Icon. *Dill. elth.* t. 310. f. 400.  
 On the banks of the Potowmac, near Harper's Ferry, Virginia. ♂. June. v. v. Flowers white.
- acaulis*. 7. *S. acaulis depressa; foliis linearibus carinatis, pedunculis terminalibus solitariis erectis unifloris, petalis emarginatis*.—*Willd. sp. pl.* 2. p. 709.  
 Icon. *Fl. dan.* t. 21. *Dill. elth.* t. 167. f. 206.  
 In Labrador. ♀. June, July. v. s. in *Herb. Lambert*. Flowers rose-coloured.

1. *S. puberula*  
 erecta  
 p. 272  
 In shade  
 ♀. 1  
 color  
 2. *S. foliis*  
 laterales  
*Alsine*  
 Icon. ♀  
 In culti  
 A co  
 The  
 3. *S. cæspitosa*  
 petalis  
 p. 71  
 Icon.  
 f. 1.  
 In Lab  
*Herb*  
*laria*
1. *A. caulis*  
 bus a  
 Icon. C  
 In Lab  
 2. *A. caulis*  
 nervis  
 fidis,  
 longi  
 About  
*Mar*
3. *A. caulis*  
 calycis  
 —W  
 Icon. 1  
 In culti  
 Caro
4. *A. caulis*

388. STELLARIA. *Gen. pl. 773.*

1. *S. pubescens*; foliis sessilibus ovatis ciliatis, pedicellis erectis, petalis calyce longioribus.—*Mich. fl. amer. 1.* p. 273.

In shady woods, on rich soil : Pennsylvania to Carolina.  
2. *S. foliis ovatis glabris, caulis procumbentibus linea lateralib. pilosa alterna.* *Smith fl. brit. 2.* p. 473.

*Alsine media.* *Sp. pl. 389.*  
Icon. *Engl. bot. 573.* *Fl. dan. 438* ♂ 525.

In cultivated grounds : common. ○. April—Sept. v. v.  
A common weed, known by the name of *Chickweed*.  
The number of stamens is variable, three, five, or ten.

3. *S. cæspitosa*; foliis subulatis congestis, ramis bipartitis, petalis emarginatis, calycibus striatis.—*Willd. sp. pl. 2.* p. 715.

Icon. *Fl. dan. 12.* *Swartz. nov. act. holm. 1788.* t. 1. f. 1.

In Labrador and Canada. ♀. May, June. v. s. in *Herb. Dickson.* This is rather an *Arenaria* than a *Stellaria*.

389. ARENARIA. *Gen. pl. 774.*

\* *Foliis oblongis.*

1. *A. caule dichotomo, foliis ovatis acutis carnosis, calyci- bus acuminatis.* —*Willd. sp. pl. 2.* p. 716.

Icon. *Gmel. sib. 4.* t. 64. *Loes. pruss. t. 2.*

In Labrador. ♀. June, July. v. s. in *Herb. Banks.*

2. *A. caule filiformi simplici, foliis ovatis obtusis subtriplici- nervibus, pedunculis lateralibus solitariis elongatis bi- fidis, pedicello altero medio bibracteato, corollis calyce longioribus.* —*Willd. sp. pl. 2.* p. 718.

About Hudson's Bay. In the state of New York. *Dr. Martin.* ♀. July. v. s. in *Herb. Banks.*

3. *A. caule dichotomo diffuso, foliis ovatis acutis subefficiatis, calycibus acutis substriatis, petalis calyce brevioribus.* —*Willd. sp. pl. 2.* p. 720.

Icon. *Fl. dan. 977.*

In cultivated grounds and on road-sides: New York to Carolina. ○. May—Aug. v. v.

4. *A. caule dichotomo diffuso, foliis ovalibus subacutis, pe-* *thymifolia.*

*pubera.*

*media.*

*biflora.*

*peploides.*

*lateriflora.*

## DECANDRIA TRIGYNIA. Arenaria.

dunculis alternis axillaribus solitariis elongatis, calycibus acutis, petalis longitudine calycis.

On the sea-shore of Labrador.  $\textcircled{O}$ . v. s. in *Herb. Banks.*

\*\* *Foliis subulato-linearibus.*

- macrocarpa.* 5. A. cæspitosa; foliis congestis subulato-linearibus planis margine ciliatis, pedunculis terminalibus unifloris foliosis, calycis foliolis linearibus, petalis ovatis calyce duplo longioribus, capsulis elongato-oblongis calyce triplo longioribus.  
On the north-west coast of America. *Nelson.* 4. v. s. in *Herb. Banks.* This approaches near to *A. grandiflora*, but differs in the long capsule and the linear leaves of the calyx materially.
- patula.* 6. A. pubescens; caulis filiformibus, paniculatis multifloris, foliis setaceo-subulatis patentibus, petalis subemarginatis calyce acutissimo et striato paulo longioribus.—*Mich. fl. amer.* 1. p. 273.  
On rocks, near Knoxville, Kentucky. *Michaux.* 4. +. Resembles *A. tenuifolia*.
- squarrosa.* 7. A. cæspitosa; foliis imis squarroso-imbricatis canaliculatis glabris, caulinis simplicissimis oligophyllis, floribus terminalibus paucis erectis, petalis calyce subrotundo multo majoribus. *Mich. fl. amer.* 1. p. 273.  
A. imbricata. *Herb. Banks.*  
A. caroliniana. *Walt. fl. car.* 141.  
In sandy pine-bartens: New Jersey and Carolina. 4. July, Aug. v. v. Resembles *A. saxatilis*.
- stricta.* 8. A. glabra, erecta, multicaulis; foliis subulato-linearibus erectis, panicula rariflora, petalis calyce ovali-lanceolato conspicue striato multo longioribus. *Mich. fl. amer.* 1. p. 274.  
On rocks of Canada, New England, and on the high mountains of Carolina. 4. May, June. v. v.; v. s. in *Herb. Lyon. specimen carolinianum.* Resembles *A. striata*, but is a larger and more upright plant.
- glabra.* 9. A. glaberrima, erectiuscula, filiformi-multicaulis; foliis subulato-linearibus planis patulis, pedicellis unifloris elongatis divaricatis, calycis foliolis ovalibus obtusiusculis laevibus petalis brevioribus. *Mich. fl. amer.* 1. p. 274.  
*Stellaria uniflora. Walt. fl. car.* 141.
- juniperina.* On rocks in North Carolina. *Michaux.* 4. +.
10. A. subpubescens, erecta, multicaulis, foliis subulatis

subtriquetris pungentibus, panicula terminali dichotome-multiflora, calycis foliolis ovato-lanceolatis striatis mucronatis petalis vix duplo brevioribus.—Willd. sp. pl. 2. p. 725.

Icon. Smith ic. ined. 1. t. 35.

In Labrador and Newfoundland. ♀. v. s. in Herb. Banks.

11. A. pubescens, cæspitoso-multicaulis; foliis setaceis, caulinis 1—3-floris erectis, calycis foliolis linearibus sub-hirsutis petalis duplo brevioribus.—Willd. sp. pl. 2. p. 726.

Icon. Jacq. austr. 3. t. 272.

On the north-west coast of America. Menzies. ♀. v. s. in Herb. Banks.

12. A. glabriuscula, cæspitosa; caulinis stricto-erectis, foliis subulatis pungentibus striatis, floribus dense-fasciculatis, calycibus subulatis striatis, petalis brevisimis.—Willd. sp. pl. 2. p. 727.

Icon. Jacq. austr. 2. t. 182.

In Canada. ♂. v. s. in Herb. Lambert. specimen Hort. Kew.

13. A. glabra, humifusa; foliis linearibus carnosis interno-diis multo longioribus, stipulis membranaceo-connatis vaginantibus, staminibus quinque, capsulis globosis.—Pers. syn. 1. p. 504.

A. rubra β. Mich. fl. amer. 1. p. 274.

A. marina. Smith fl. brit. 2. p. 480?

On the sea-coast, in salt-marshes: Canada to Carolina.

♂. June, July. v. v. Flowers pale-red.

*laricifolia.*

*fasciculata.*

*canadensis.*

#### IV. TETRAGYNYA.

##### 390. MICROPETALUM. Pers. syn. 1. p. 509.

Spergulastrum. Mich. fl. amer. 1. p. 275.

1. M. dense pubescens; foliis lanceolatis in petiolum an-*lanuginosum*. gustatis, pedunculis subsolitariis elongatis demum reflexis, floribus apetalis.—Pers. l. c. Mich. l. c.

In the mountains of Virginia and Carolina. ♀. June, July. v. v.

- lanceolatum.* 2. *M. glabrum*; foliis lanceolatis utrinque angustatis, floribus paniculatis, petalis ovatis brevissimis.—*Pers. l. c.* *Mich. l. c.*  
On moist rocks: Canada to Pennsylvania.  $\mathcal{U}$ . July. v. v.
- gramineum.* 3. *M. glaberrimum*; foliis linearibus erectis, panicula terminali laxa, pedicellis filiformibus, petalis lanceolatis longitudine calycis.—*Pers. l. c.* *Mich. l. c.*  
Near springs and shady rocks: New York to Virginia.  
 $\mathcal{U}$ . June, July. v. v.  
*A. fasciculata* probably belongs to this genus.

## V. PENTAGYNIA.

## 391. SPERGULA. Gen. pl. 798.

- arvensis.* 1. *S.* foliis filiformibus verticillatis, panicula dichotoma, pedunculis fructiferis reflexis, seminibus reniformibus. *Willd. sp. pl. 2. p. 818.*  
Icon. *Fl. dan.* 1033. *Engl. bot.* 1535.  
In sandy fields: Pennsylvania, New Jersey, &c.  $\odot$ . June, July. v. v.
- saginoides.* 2. *S.* foliis oppositis subulatis nudis, pedunculis solitariis longissimis glabris. *Smith fl. brit. 2. p. 504.*  
Icon. *Engl. bot.* 2105.  
In sandy fields: Carolina.  $\odot$ . June, July. v. s. in *Herb. Enslen.*

## 392. CERASTIUM. Gen. pl. 797.

- vulgatum.* 1. *C. hirsutum*, viscidum, cæspitosum; foliis ovatis, petalis oblongis calyce subæqualibus, floribus pedunculo longioribus. *Smith fl. brit. 2. p. 496.* *Willd. sp. pl. 2. p. 811.*  
Icon. *Engl. bot.* 789. *Vaill. par. t. 30. f. 3.*  
In dry fields and among rocks; frequent: New England to Pennsylvania.  $\odot$ . April—June. v. v. The flowers of the whole genus are white.
- viscosum.* 2. *C. hirsutum*, viscosum, diffusum; foliis lanceolato-oblongis obtusiusculis, petalis obovatis calyce vix longioribus, floribus pedunculo brevioribus.—*Willd. sp. pl. 2. p. 812.*

Icon.  
In past  
May  
3. *C. hirs*  
talis  
long  
p. 8  
Icon.  
On dry  
and  
4. *C. pub*  
tusis  
dati  
—  
Icon.  
On dry  
May  
5. *C. ter*  
linea  
ema  
On the  
4.  
very  
6. *C. hir*  
vari  
tüm  
ealy  
On the  
v. s.

1. *A. hir*  
nud  
Icon.  
Com  
but  
v. v

1. *L. gla*  
bifid  
Icon. I

VOL. I.

Icon. Engl. bot. 790. Vaill. par. t. 30. f. 1.  
In pastures and on old walls: Canada to Carolina. ¼.  
May—Sept. v. v.

3. C. hirsutum, viscidum; foliis ovato-oblongis acutis, petalis calyce brevioribus emarginatis, pedunculis calyce longioribus, floribus pentandris.—Willd. sp. pl. 2. p. 812.

semidecan-  
drum.

Icon. Engl. bot. 1630. Vaill. par. t. 30. f. 2.

On dry, barren, and sunny hills; frequent: Pennsylvania and Virginia. ☺. May, June. v. v.

4. C. pubescens, caespitosum; foliis linear-lanceolatis obtusis basi ciliatis internodiis brevioribus, petalis obcordatis calyce duplo longioribus, calycis foliolis obtusis.—Willd. sp. pl. 2. p. 813.

arvense.

Icon. Fl. dan. 626. Engl. bot. 93. Vaill. par. t. 30. f. 4.

On dry hills and rocks; frequent: Pennsylvania, &c. ¼.

May—Aug. v. v. Flowers large.

5. C. tenuissime pubescens, caespitosum; foliis angusto-linearibus internodiis longioribus, petalis obovatis emarginatis calyce acuto subtriplo longioribus.

tenuifolium.

On the banks of Schuylkill and Delaware, Pennsylvania.

¼. May, June. v. v. Resembles the foregoing one very much.

6. C. hirsutum; foliis linearibus internodiis longioribus divaricatis, pedunculis terminalibus elongatis bis 3-chotomis, bracteis oppositis ovatis, petalis emarginatis calyce acuto duplo longioribus, capsulis subglobosis.

elongatum.

On the plains of Columbia river. M. Lewis. ¼. April.

v. s. in Herb. Lewis.

### 393. AGROSTEMMA. Gen. pl. 795.

1. A. hirsuta; calyce corollam superante, petalis integris nudis. Willd. sp. pl. 2. p. 805.

Githago.

Icon. Fl. dan. 576. Engl. bot. 741.

Common Cockle is frequently found among the grain, but is not indigenous in America. ☺. June, July.  
v. v.

### 394. LYCHNIS. Gen. pl. 796.

1. L. glabra; floribus densis umbellato-capitatis, petalis bifidis, floribus tetragynis.—Willd. sp. pl. 2. p. 809.

alpina.

Icon. Fl. dan. 65. Hall. helv. 1. t. 7.

In Labrador and Hudson's Bay.  $\text{♀}$ .  $v. s.$  in Herb. *Dickson. nec non Banks.* About a span high; flowers red, with membranaceous bracteas.

395. OXALIS. *Gen. pl. 794.* Wood-Sorrel.

*Acetosella.*

1. *O. acaulis*; scapo unifloro foliis longiore, foliis ternatis dilatato-obcordatis, lobis rotundatis, stylis longitudine staminum interiorum, radice dentata.—*Willd. sp. pl. 2. p. 780.*

Icon. *Fl. dan. 980. Engl. bot. 762.*

In shady woods: Canada. *Michaux.* On the high mountains of Pennsylvania and New York, round the roots of old trees.  $\text{P.}$   $\text{♀. May. v. v.}$  Flowers large, white, with red veins and yellow at the bottom: the petals are cuneate, emarginate, and narrower than the European species.

*violacea.*

2. *O. acaulis*; scapo umbellifero, pedicellis subpubescens, floribus nutantibus, foliis ternatis obcordatis glabris, calycis laciniis apice callosis, stylis staminibus exterioribus brevioribus.—*Willd. sp. pl. 2. p. 786.*

Icon. *Jacq. oxal. t. 80. f. 2. Jacq. hort. 2. t. 180. Pluk. alm. t. 102. f. 4.*

In shady woods, on the side of hills, in fertile soil: New England to Carolina.  $\text{♀. April, May. v. v.}$  Flowers pink-coloured.

*Lyonii.*

3. *O. tota sericeo-pilosa*; caule ramoso decumbente, pedunculis bifloris petiolis longioribus, foliis ternatis bilobo-obcordatis, laciniis rotundatis divaricatis, petalis cuneatis, siliquis tomentosis calyce lanceolato duplo longioribus.

On Cumberland Island, Georgia. *Lyon.*  $\text{♀. June. v. s. in Herb. Lyon.}$  Flowers yellow.

*corniculata.*

4. *O. tota pubescens*; caule ramoso diffuso seu procumbente, pedunculis umbellatis petiolis brevioribus, foliis ternatis obcordatis, petalis obtuse-rotundatis apice erosis, stylis longitudine staminum interiorum.—*Willd. sp. pl. 2. p. 800.*

Icon. *Jacq. oxal. t. 5. Salisb. in act. soc. linn. lond. 2. t. 23. f. 5. O. pusilla.*

In cultivated grounds: Canada to Carolina.  $\text{O.}$   $\text{♀. May — Oct. v. v.}$  Flowers yellow; the American plant has larger flowers than the European; in every other respect they are very much alike.

5. *O. tota belliflora*

petali

rum.

*O. ambigua*

*O. cornuta*

Icon. *J*

In cult

*June,*

6. *O. hirsutissima*

belliflora

corolla

Icon. *L*

In Caro

than

confus

1. *P. caerulea*

inæqu

tis al

*Willd.*

Icon. *A*

In ditch

*♀. J*

In th

cond

Chin

ters:

elong

serra

corner

1. *S. erecta*

tiuosa

bus

In No

east

*♀. J*

stam

2. *S. glabra*

nigra

sessilis

5. *O. tota hirsuta*; caule erecto ramoso, pedunculis umbelliferis petiolis brevioribus, foliis ternatis obcordatis, petalis obovatis, stylis longitudine staminum interiorum.—*Willd. sp. pl. 2. p. 800.*

*stricta.*

*O. ambigua*. *Salisb. in act. soc. linn. lond. 2. t. 23. f. 4.*

*O. corniculata*. *Fl. dan. t. 873.*

Icon. *Jacq. oxal. t. 4.* *Dill. elth. t. 221. f. 4.*

In cultivated grounds: Pennsylvania to Virginia. ♀. June, July. v. v. Flowers yellow.

6. *O. hirsuta*; caule folioso erecto hirsuto, pedunculis umbelliferis foliis longioribus, foliis ternatis obcordatis, corollis emarginatis. *Willd. sp. pl. 2. p. 799.*

*Dillenii.*

Icon. *Dill. elth. t. 221.*

In Carolina. *Dillenius*. ♂. +. Flowers yellow, larger than the preceding one, with which it generally is confounded.

396. PENTHORUM. *Gen. pl. 790.*

1. *P. caule ramoso, angulato*; foliis lanceolatis subsessilibus inaequaliter dense-serratis, spicis terminalibus paniculatis alternis cymosisque, seminibus scrobiformibus.—*Willd. sp. pl. 2. p. 770.*

*sedoides.*

Icon. *Act. ups. 1744. t. 2.*

In ditches and small ponds: New England to Carolina. ♀. June, July. v. v. Flowers white or pale yellow. In the Herbarium of A. B. Lambert, Esq. is a second species, brought by Sir George Staunton from China, which I distinguish by the following characters: *Penthorum chinense*, caule simplici tereti, foliis elongato-linearis-lanceolatis subpetiolatis inaequaliter serratis, spicis cymosis terminalibus, seminibus ovatis corneis.

397. SEDUM. *Gen. pl. 789.*

1. *S. erectum, glabrum, 1-2-unciale*; foliis alternis teretiusculis oblongis, floribus ad summitatem paucioribus alternis subpedicellatis. *Mich. fl. amer. 1. p. 276.*

*pusillum.*

In North Carolina, on flat rocks. *Michaux*. On the east banks of the Shanadoah River, Virginia. *P.* ♀. June, July. v. v. Flowers white, with eight stamens.

2. *S. glabrum*; caulinibus assurgentibus, foliis sparsis planiuscula linearibus obtusis, cyma polystachya, floribus sessilibus octandris.—*Mich. fl. amer. 1. p. 277.*

*pulchellum.*

- On rocks near Knoxville. *Michaux.*  $\frac{4}{4}$ . July. +. Flowers purple. Resembles *S. reflexum*.
- stenopetalum.* 3. *S. glabrum*; caulis assurgentibus, foliis sparsis congestis adnato-sessilibus compresso-subulatis acutis, cyma terminali trichotoma dichotoma, spicis recurvatis, floribus sessilibus 10-andris, petalis linearibus calyce multo longioribus.
- On rocks on the banks of Clark's river and Kooskoosky.  $\frac{4}{4}$ . June, July. v. s. in *Herb. Lewis.* Flowers golden yellow; has some affinity with *S. reflexum*.
- ternatum.* 4. *S. pumilum*, repens; foliis planis rotundato-spathulatis ternis, cyma subtristachya, floribus sessilibus 8-andris, centro 10-andro.—*Mich. fl. amer.* 1. p. 277.  
*S. saxatilis* floribus albis &c. *Clayt. virg.* 891.  
*S. americanum*. *Herb. Banks.*  
 On rocks in the western parts of Pennsylvania, Virginia, and Carolina.  $\frac{4}{4}$ . May, June. v. v. An elegant little species; flowers white.
- telephiooides.* 5. *S. foliis planis ovalibus utrinque subacutis dentatis, cymbo multiplici fasciculato.* *Mich. fl. amer.* 1. p. 277.  
 On rocks on high mountains: Virginia and Carolina.  $\frac{4}{4}$ . June, July. v. v. Flowers pale purple.

## VI. DECAGYNIA.

## 398. PHYTOLACCA. Gen. pl. 800.

- decandra.* 1. *P. foliis ovatis utrinque acutis, floribus decandris decagynis.* —*Willd. sp. pl.* 1. p. 822.  
*Icon. Dill. elth. t. 339. f. 399. Pluk. alm. t. 225. f. 3.*  
 In open woods and old fields: New England to Carolina.  
 $\frac{4}{4}$ . June, July. v. v. Flowers white, on red peduncles; berries black, and give an elegant purple colour to any liquor. The young sprouts in spring give a fine vegetable for the table, resembling asparagus. The plant is known by the name of *Poke-weed*.

399. CAC  
Co  
cul  
400. BAR  
tala  
3—  
sen  
401. DEC  
—  
402. PHI  
fidu  
laris  
403. CHR  
5-fi  
5-f  
404. PRU  
cidu  
sute  
405. TIG  
Pet  
cen  
406. LYT  
bas  
lysp  
407. CUP  
bas  
laris  
408. FOT  
cren  
2-lo

## XI. ICOSANDRIA.

### I. MONOGYNIA.

399. CACTUS. *Cal.* superus, monophyllus, imbricatus. *Cor.* multiplex. *Stigm.* multifidum. *Bacca* 1-lo-  
cularis, polysperma.
400. BARTONIA. *Cal.* superus, 5-fidus. *Cor.* polype-  
tala. *Caps.* cylindrica, 1-loclaris, apice operculatim  
3—5-valvis. *Recept.* 3—5. parietalia, dupli serie  
seminifera.
401. DECUMARIA. *Cal.* superus, 8—12-fidus. *Pet.* 8  
—12. *Caps.* 7—10-loclaris, polysperma.
402. PHILADELPHUS. *Cal.* superus, turbinatus, 4—5-  
fidus. *Pet.* 4—5. *Styl.* 4-fidus. *Caps.* 4—5-lo-  
claris, polysperma.
403. CHRYSOBALANUS. *Cal.* inferus, campanulatus,  
5-fidus. *Pet.* 5. *Stylus* lateralis. *Drupæ* nux ovata,  
5-falcata, 5-valvis, 1-sperma.
404. PRUNUS. *Cal.* inferus, campanulatus, 5-fidus, de-  
ciduous. *Pet.* 5. *Stylus* terminalis. *Drupæ* nux  
suturis prominulis.
405. TIGAREA. *Cal.* inferus, campanulatus, 5-partitus.  
*Pet.* 5. *Caps.* folliculiformis, sutura lateral dehis-  
cens. *Sem.* unicum.
406. LYTHRUM. *Cal.* inferus, tubulosus, 12-dentatus,  
basi æquali. *Pet.* 6. æqualia. *Caps.* 2-loclaris, po-  
lysperma.
407. CUPHEA. *Cal.* inferus, tubulosus, 6—12-dentatus,  
basi hinc gibbus. *Pet.* 6. inæqualia. *Caps.* 1-lo-  
claris, hinc cum calyce longitudinaliter dehiscens.

### II. DI-PENTAGYNIA.

408. FOTHERGILLA. *Cal.* inferus, truncatus, obsolete  
crenatus. *Cor.* 0. *Germen* bifidum. *Styli* 2. *Caps.*  
2-loclaris. *Sem.* solitaria; ossea.

409. AGRIMONIA. *Cal.* inferus, 5-dentatus, altero obvallatus. *Pet.* 5. *Styli* 2. *Sem.* 2. in fundo calycis.
410. CRATAEGUS. *Cal.* superus, 5-fidus. *Pet.* 5. *Styli* 1—5. *Bacca* seu *Pomum* 2—5-spermum. *Sem.* ossea.
411. PYRUS. *Cal.* superus, 5-fidus. *Pet.* 5. *Styli* 5. *Pomum* 5-loculare, polyspermum. *Sem.* cartilaginea.
412. SORBUS. *Cal.* superus, 5-fidus. *Pet.* 5. *Styli* 3. *Bacca* 3-sperma. *Sem.* cartilaginea.
413. SPIRAEA. *Cal.* inferus, 5-fidus. *Pet.* 5. *Styli* 5. *Caps.* polyspermæ.

## III. POLYGYNIA.

414. ROSA. *Cal.* urceolatus, collo coartatus, 5-fidus. *Pet.* 5. *Sem.* plurima, hispida, calycis interioris lateri affixa.
415. RUBUS. *Cal.* patens, 5-fidus. *Pet.* 5. *Bacca* composita, acinis monospermis.
416. DALIBARDA. *Cal.* patens, 5-fidus. *Pet.* 5. *Pistilla* 5—8. *Styli* longi, filiformes, decidui. *Sem.* exsucca.
417. DRYAS. *Cal.* patens, 8-fidus. *Pet.* 8. *Sem.* plurima: arista lanata.
418. GEUM. *Cal.* 10-fidus. *Pet.* 5. *Sem.* aristata: arista plerumque geniculata.
419. POTENTILLA. *Cal.* 10-fidus. *Pet.* 5. *Sem.* subrotunda, receptaculo parum exsucco affixa.
420. COMARUM. *Cal.* 10-fidus. *Pet.* 5. calyce minora. *Recept.* seminum ovatum, spongiosum, persistens.
421. FRAGARIA. *Cal.* 10-fidus. *Pet.* 5. *Recept.* seminum ovatum, baccatum, deciduum.
422. CALYCANTHUS. *Cal.* 1-phyllo, urceolatus, squarrosus, foliolis coloratis. *Cor.* calycina. *Styli* plurimi. *Stigm.* glandulosa. *Sem.* plurima, caudata, intra calycem succulentum.

altero  
fundoStyli  
Sem.Styli 5.  
aginea.  
Styli 3.

Styli 5.

5-fidus.  
oris la-

ce com-

i. Pis-  
lem. ex-

n. plu-

ristata :

em. sub-

minora.  
tens.

cept. se-

ceolatus,

Styli  
caudata,

## I. MONOGYNIA.

## 399. CACTUS. Gen. pl. 838.

1. C. articulato-prolifer; articulis compressis ovatis, spinis setaceis.—Willd. sp. pl. 2. p. 943.

*Opuntia.*

Icon. Knorr thes. 1. t. F. a.

In pine-barrens and sandy fields: New Jersey to Carolina.  $\frac{1}{2}$ . June, July. v. u. Flowers large, yellow; berries red, eatable, known by the name of *Prickly Pears*.

There are two or three more species growing on the Missouri; but having had no opportunity to see them myself, I only mention it here, to draw the attention of the botanists in those regions towards them.

## 400. BARTONIA. Sims in bot. mag. 1487.

1. B. germe folioso, seminibus nudis.

*ornata.*

B. decapetala. Sims l. c.

On chalky soil and arid volcanic grounds, on the borders of the Missouri. M. Lewis. ♂. July, Aug. v. s. in Herb. Lewis.

*Radix* bietinis, fibrosa. *Caulis* ramosus, teres. *Rami* patentes, scabri. *Folia* alterna, semi-amplexicaulia, oblonga, basi dilatata, superne sensim angustiora, inciso-dentata seu subpinnatifida, utrinque scabra, glaucescentia. *Flores* magni, albi, ex axillis superioribus, solitarii, pedunculati. *Pedunculi* foliosi, folio breviores. *Calyx* superus campanulatus, 5-partitus; laciniis ovatis, longissime acuminatis, petalis paulo brevioribus. *Petala* 10. alba, lanceolato-oblonga, acuta, unguiculata. *Stamina* numerosa, filiformia, petalis breviora. *Antheræ* luteæ, oblongæ, biloculares. *Germen* inferrum, oblongum, foliis pinnatifidis calyce brevioribus ornatum. *Stylus* staminibus longior, filiformis. *Stigma* simplex. *Capsula* cylindrica, calyce coronata, apice operculatim 5-valvis. *Semina* oblonga, plana.

This beautiful plant, whose large white flowers open during the night and spread a most agreeable odour, was discovered in the year 1804, on the white bluffs near the Maha village, by the late M. Lewis, Esquire. In 1807 I made a drawing and description of it, for

## ICOSANDRIA MONOGYNIA. Bartonia.

the publication of that gentleman's Tour across the Continent of America to the Pacific Ocean. In 1812, Mr. Nuttall, on his return from a journey in those parts, brought seeds and specimens of this and another species to London; and having by those means the living plants, I agreed with Mr. Nuttall to dedicate it to the memory of Dr. B. S. Barton, of Philadelphia, our mutual friend; under which name it was published in the Botanical Magazine. Since that publication, Mr. Nuttall, whose name has occurred in several pages of this work, with all the credit due to his valuable discoveries, has found himself rather offended at not having given him all the exclusive credit of discovery, which with justice and propriety to the memory of M. Lewis, Esq. I never could do.

*nuda.*2. B. germinæ nudo, seminibus alatis. *Sims l. c.*

On the banks of the Missouri. *Nuttall.* ♂. July, Aug. v. v. This species has smaller flowers, and the leaves are not so glaucous as the foregoing: in every other respect the above description is applicable to the present one, with little deviation. The number of petals is variable, from ten to fifteen.

*barbara.*1. D. foliis ovato-oblongis utrinque acutis obsolete serratis. *Willd. enum. p. 516. Willd. sp. pl. 2. p. 850.*

In South Carolina. ♀. July, Aug. v. v. Stem climbing; flowers in corymbous panicles, white, very sweet-scented.

*sarmentosa.*2. D. foliis ovatis acutis basi rotundatis, apice serratis. *Willd. enum. 516. Willd. sp. pl. 2. p. 850.*

*Forsythia scandens. Walt. fl. car. ....*

*Icon. Bosc. in act. hist. nat. paris. 1. t. 13.*

In deep cedar-swamps of Virginia and Carolina; particularly plentiful in the Dismal-swamp, near Norfolk. ♀. July, Aug. v. v. This climbing shrub resembles the first species, but grows taller, and is in every other respect larger in all its parts. It ascends, in company of *Zizyphus volubilis*, the tallest cypress trees in those swamps.

- 4  
1. P. folii acuti quat  
Icon. On the July  
2. P. folii calyc fido,  
On th Her cedi  
3. P. folii subt styl lines  
P. inod On riv Jun  
gard far a

## 401. DECUMARIA. Gen. pl. 815.

- 403  
1. C. foli subt bus,  
Mic  
Icon. In the Jun  
sma the run  
any bra as a mig

1. P. flor cidi

402. PHILADELPHUS. *Gen. pl. 840.*

1. *P. foliis ovatis acuminatis integerrimis, calycis laciniis acutis, stylo staminibus longiore indiviso, stigmatibus quatuor oblongis.* — *Willd. sp. pl. 2. p. 948.*  
Icon. *Catesb. car. 2. t. 84.* *Bot. mag. 1478.*  
On the banks of rivers in South Carolina.  $\text{h}$ . June, July. *v. v.* An elegant shrub : flowers large, white.
2. *P. foliis ovatis acutis subintegerrimis, margine ciliatis, calycis laciniis acutis, stylo longitudine staminum 3-fido, stigmatibus tribus.*  
On the waters of Clark's river.  $\text{h}$ . July. *v. s. in Herb. Lewis.* The flowers are smaller than the preceding species.  
*Lewisii.*
3. *P. foliis ovatis acuminatis denticulatis, axillis veharum grandiflorus.*  
subtus fasciculato-pilosus, calycis laciniis acuminatis, stylo staminibus longiore indiviso, stigmatibus quatuor linearibus. — *Willd. enum. 511.*  
*P. inodorus. Hortul.*  
On river banks in South Carolina and Georgia.  $\text{h}$ . June, July. *v. v.* This is generally found in most gardens, under the name of *P. inodorus*, which is by far a more scarce plant.

403. CHRYSOBALANUS. *Gen. pl. 850.*

1. *C. foliis subcuneatim oblongis vel etiam oblanceolatis oblongifolius.*  
subtus lanuginoso-incanis, petalis subrotundo-ob-ovalibus, staminibus glabris, fructibus oblongo-ovatis. — *Mich. Fl. amer. 1. p. 283.*  
Icon. *Bartr. iter, t. . . .*  
In the barren sands of Georgia and Florida.  $\text{h}$ . May, June. *v. s. in Herb. Lyon.* Flowers white, very small, in large panicles ; they are generally dioicous : the fruit is the size of a common plum. This shrub runs with its branches under ground, without making any fibres, for a considerable distance ; and its side branches appear from one to two feet above ground, as a separate small shrub, by which more properly it might be considered as a tree under ground.

404. PRUNUS. *Gen. pl. 849.*

1. *P. floribus racemosis, racemis erectis elongatis, foliis deciduis ovali-oblongis acuminatis inaequaliter duplicato-*  
*virginiana.*

- dentatis utrinque glabris, petiolis subquadriglandulosis.*  
—Willd. sp. pl. 2. p. 985.
- P. rubra. Ait. kew. 2. p. 162.*  
Icon. Willd. arb. t. 5. f. 1. Mich. arb. t. ....  
In woods, frequent: Canada to Florida. ♀. April, May.  
*v. v.* A handsome tree: flowers white, as all the following are; berries red. The wood is very useful and handsome, and is manufactured into elegant furniture.
- serotina.* 2. *P. floribus racemosis, racemis laxis, foliis deciduis simpliciter serratis, serraturis infimis subglandulosis, costa media basin versus barbata.*—Willd. sp. pl. 2. p. 986.
- P. virginiana. Mill. dict. n. 3. Du Roi harbk. 2. p. 191.*  
Icon. Willd. arb. t. 5. f. 2. Wangh. amer. t. 14. f. 33.  
In mountain woods: Pennsylvania to Carolina. ♀. May, June. *v. v.* Resembles the first species very much.
- canadensis.* 3. *P. floribus racemosis, foliis deciduis eglandulosis latolanceolatis rugosis argute serratis utrinque subpubescentibus, in petiolum attenuatis.*—Willd. sp. pl. 2. p. 986.  
Icon. Pluk. alm. t. 158. f. 4.  
In North America. Sp. pl. ♀. +. I strongly suspect this to be nothing more than *P. hiemalis*, No. 9.
- caroliniana.* 4. *P. floribus racemosis, foliis sempervirentibus oblongolanceolatis mucronatis serratis eglandulosis lævigatis.*—Willd. sp. pl. 2. p. 987.  
In South Carolina and Georgia. ♀. May. *v. v.* A handsome evergreen shrub, resembling *P. lusitanica*.
- semperflorens.* 5. *P. floribus racemosis, racemis foliosis pendulis, calycibus serratis, foliis ovatis serratis glabris basi glandulosis.*—Willd. sp. pl. 2. p. 992.  
P. serotina. Roth catalect. 1. p. 58.  
In the western parts of Virginia. ♀. May. *v. v.* A small shrub.
- borealis.* 6. *P. floribus corymbosis, pedicellis elongatis, foliis ovalioblongis acuminatis eroso-denticulatis membranaceis glabris, fructu subovato.*—Mich. fl. amer. 1. p. 286.  
Icon. Mich. arb. t. ....  
In Canada and on the high mountains of New England and Pennsylvania. ♀. May, June. *v. v.* A very handsome small tree; the wood exquisitely hard and fine grained; the cherries hang in clusters, are small, red, and agreeable to the taste, but occasion an astringency in the mouth, from which they are called *Choke-cherries*.

7. *P. umbraculifera. pani glab sp. 1*  
*P. lancifolia. In wood gini rasu to ea*

8. *P. umbellata. ovata glab p. 9*  
Icon. In Can June

9. *P. umbellata. niis long mina*  
In Can Carc extre Blac

10. *P. umbellata. ellip glan*  
In the May the pea.

11. *P. umbellata. calyculata. rula nuce*  
P. sphaerocarpa  
On the west P. than ple,

12. *P. umbellata. acutifolia*

7. *P. umbellis subsessilibus aggregatis multifloris tandem pensylvanica.*  
paniculæformibus, foliis oblongo-lanceolatis serratis  
glabris basi biglandulosis, ramulis punctatis.—*Willd.*  
*sp. pl. 2. p. 993.*  
*P. lanceolata. Willd. arb. t. 3. f. 3.*  
In woods and near plantations : New England to Virginia.  
h. April, May. v. v. This resembles *P. Ceratas* very much: the cherries are small, and agreeable to eat.
8. *P. umbellis sessilibus solitariis paucifloris, foliis deciduis*  
ovatis acuminatis inæqualiter argute-serratis utrinque  
glabris, petiolis biglandulosis.—*Willd. sp. pl. 2.*  
*p. 993.*  
Icon. Bot. mag. 1117.  
In Canada and on the Allegany mountains. h. May,  
June. v. v.
9. *P. umbellis sessilibus aggregatis paucifloris, calycis laci-*  
niis lanceolatis, stipulis setaceo-compositis, foliis ob-  
longo-ovalibus obovalibusve abrupte promisseque acu-  
minatis, fructu subovato.—*Mich. fl. amer. 1. p. 284.*  
In Canada and on the western mountains of Virginia and  
Carolina. h. May. v. v. The fruit small, black,  
extremely astringent, but eatable in winter; called  
*Black Choke-cherry.*
10. *P. umbellis sessilibus aggregatis paucifloris, foliis ovato-*  
ellipticis acutis utrinque glabris argute serratis basi bi-  
glandulosis.—*Willd. sp. pl. 2. p. 993.*  
In the western parts of Pennsylvania and Virginia. h.  
May. v. v. A shrub about four or five feet high;  
the fruit very indifferent, black, the size of a large  
pea.
11. *P. umbellis sessilibus aggregatis paucifloris, pedunculis*  
calycibusque pubescentibus, foliis brevi-ovalibus ser-  
rulatis basi plerumque biglandulosis, drupa sphærica,  
nuce subrotundo-ovoidea.  
*P. sphærocarpa. Mich. fl. amer. 1. p. 284.*  
On the sea-coast of New England. *Michaux.* In the  
western parts of Pennsylvania on the borders of lakes.  
*P. h. May. v. v.* A small shrub; flowers smaller  
than any other American species; fruit brownish-purple,  
very astringent.
12. *P. umbellis sessilibus aggregatis paucifloris, calycibus*  
acutis, ramis virgatis teretibus, foliis angusto-lanceo-

*nigra.**hiemalis.**pygmæa.**pubescens.**pumila.*

- latis superne serratis subtus pallidis.* — *Willd. sp. pl. 2.*  
*p. 990.* *Icon. Mill. ic. t. 89. f. 2.* glabrid, i. e. smooth.  
 In low grounds and swamps : Pennsylvania and Virginia.  
 h. May. v. v. A small shrub of the appearance of  
*Amygdalus nana*; fruit red, very acid.
- depressa.* 13. *P. umbellis sessilibus aggregatis paucifloris, calycibus obtusis, ramis angulatis depresso-prostratis, foliis cu- neato-lanceolatis rariter serratis glbris subtus glaucis, fructu ovato.*  
*P. pumila. Mich. fl. amer. 1. p. 286.*  
 On the sandy shores of rivers and lakes : Canada to Virginia. h. May. v. v. This low shrub, which spreads its branches very much, and does not rise above one foot from the ground, is known by the name of *Sand-cherries*. The fruit is black, small, and agreeably tasted.
- Chicasa.* 14. *P. gemmis aggregatis bifloris, pedicellis brevissimis, ca-lycibus glbris, lacinii obtusis, foliis oblongo-ovalibus acutis seu acuminatis serrulatis, fructu subgloboso, ramis spinescentibus glaberrimis.* — *Mich. fl. amer. 1. p. 284.*  
*P. insititia. Walt. fl. car. 146.*  
 In Virginia and Carolina. h. April, May. v. v. Known by the name of *Chicasaw Plum*. The fruit is yellow and agreeably tasted. It is mentioned in Michaux's Flora, that it was introduced by the Indians; which probably may be the case, as it generally only occurs where ancient camps of Indians have been.
- maritima.* 15. *P. pedunculis subsolitariis, foliis ovato-oblongis acumini-natis duplicato-serratis.* *Willd. enum. 519.*  
*P. acuminata, fruticosa; ramulis glabris, foliis oblongo-ovalibus longiuscule acuteque acuminatis, calyce glabro, drupa longiuscule pedunculata ovata acuminata.* *Mich. fl. amer. 1. p. 284?*  
 On the sea-coast : New Jersey to Carolina. h. May. v. v. Fruit the size of a pigeon's egg, very good to eat, dark purple.
- Susquehan-na.* 16. *P. pedunculis solitariis, foliis obovato-oblongis subtus glaucis serratis basi integerrimis.* *Willd. enum. 519.*  
 On the banks of the Susquehannah, Pennsylvania, h. May. +. It appears to me that this species approaches very near to No. 13.

*sp. pl. 2.*

Virginia.  
arance of  
nt

calycibus  
foliis cu-  
glaucis,

anada to  
, which  
not rise  
the name  
all, and

mis, ca-  
ovalibus  
globoso,  
amer. 1.

Known  
s yellow  
llichaux's  
; which  
y occurs

s acumi-  
oblongo-  
e glabro,  
a. Mich.

. May.  
good to

subtus  
n. 519.  
a, h.  
proaches

17. P. pedunculata  
Near ha  
is the  
from

18. P. pedunculata  
pubescent  
sp. pl.  
The con  
in he  
been

## 405. TIG

1. T. foliis  
tatis,  
termi

In the p  
lumb  
ers ye  
Frutex

simi,  
squar  
mia,  
3-fida  
ginati  
cano-  
dunc  
lutei,  
hirsu  
guicu  
20. lo  
cular  
stami

Capsa  
This sin  
Aubl  
be o  
Aubl  
tainly  
gener

*Tigaera tridentata.**Coumthera minima.*

## ICOSANDRIA MONOGYNIA. Prunus.

333

17. P. pedunculis solitariis, foliis ellipticis glabris, fructibus pendulis, ramis subinermibus. *Willd. sp. pl. 2. p. 997.*  
Near habitations, frequent.  $\frac{1}{2}$ . April, May. v. v. This is the common *Myrabolan Cherry*, and had its origin from America, according to authors.
18. P. pedunculis solitariis, foliis elliptico-lanceolatis subtus pubescentibus, fructibus rectis, ramis spinosis. *Willd. sp. pl. 2. p. 997.*  
The common *Black Thorn*, or *Sloe*, is frequently found in hedge-rows, particularly in Pennsylvania, but has been introduced from Europe.

*cerasifera.**spinosa.*405. TIGAREA. *Aubl. guj. 2. p. 917. Pursh in linn.*  
*trans. v. 11.*

1. T. foliis in apice ramulorum confertis, obtuse cuneatis, 3-dentatis, supra villosis subtus cano-tomentosis, floribus terminalibus solitariis brevissime pedunculatis.  
In the prairies of the Rocky-mountains and on the Columbia river.  $\frac{1}{2}$ . July. v. s. in *Herb. Lewis.* Flowers yellow.  
*Frutex ramosissimus. Cortex cinerea. Ramuli brevissimi, gibbosi, apice fasciculatim foliosi. Gemmæ: squamis subrotundis, glabris, fuscis. Folia cuneiformia, quasi in petiolum angustata, apice 3-dentata seu 3-fida, laciniis lateralibus acutis, intermedia subemarginata, margine revoluta, supra strigoso-villosa, subtus cano-tomentosa, subtriplinervia, semiuincialia. Pedunculi brevissimi, terminales, solitarii, uniflori. Flores lutei, magnitudine *Crataegi oxyacanthi*. Calyx extus hirsutissimus, laciniis acutis. Petala obovata, subunguiculata, calyce paulo longiora. Filamenta circiter 20. longitudine petalorum. Antheræ oblongæ, bilobulares. Germen oblongum, hirsutum, longitudine staminum. Stylus brevis, obliquus. Stigma simplex. Capsula immatura calyce persistente cincta.*

This singular shrub approaches the nearest to *Tigarea* of Aublet: the seeming difference in character may only be owing to the imperfect state of the specimens Aublet's description seems to be made of. It certainly deserves to be separated from *Tetracera*, though generally united with that genus.

*tridentata.*

406. LYTHRUM. *Gen. pl. 825.*

*Salicaria.*  $\beta.$  *pubescens.*

1. L. pubescens; foliis oppositis ternatisque sessilibus lanceolatis basi cordatis, floribus terminalibus verticillatospicatis 12-andris, capsulis oblongis.—*Willd. sp. pl. 2. p. 865.*

Icon. *Fl. dan. 671. Engl. bot. 1061.*

In wet meadows: Canada and New England.  $\frac{1}{4}$ . July, Aug. v. v. About two feet high; flowers very showy, purple.

*verticillatum.*

2. L. pubescens; foliis oppositis lanceolatis petiolatis, floribus axillaribus corymbosis subverticillatim aggregatis decandris, fructibus globosis.—*Willd. sp. pl. 2. p. 866.*

*Decodon aquaticum. Gmel. syst. 677.*

*Anonymos aquatica. Walt. fl. car. 137.*

In swamps: Canada to Florida; not common.  $\frac{1}{4}$ . July, Aug. v. v. A showy perennial, from two to four feet high; the stems are declinate, and strike frequently roots on their summits; flowers large, fine purple.

*virgatum.*

3. L. foliis oppositis lanceolatis glabris, caule paniculato, floribus axillaribus ternis pedicellatis 12-andris.—*Willd. sp. pl. 2. p. 865.*

*L. virgatum. Walt. fl. car. 120.*

In low sandy meadows and swamps. *Bosc.*  $\frac{1}{4}$ . July, Aug. v. s. in *Herb. Enslen.* Flowers pale purple. Walter in his *Flora Caroliniensis* has it as an hexandrous plant; which probably may be the case.

*alatum.*

4. L. glaberrimum; foliis oppositis ovato-oblongis acutis basi subcordatis arcte sessilibus, ramis virgatis 4-gonatalatis, floribus axillaribus solitariis sessilibus hexandris.

In Lower Georgia. *Enslen.*  $\frac{1}{4}$ . June, July. v. s. in *Herb. Enslen.* From three to four feet high; flowers small, purple.

*lineare.*

5. L. glabrum, virgato-paniculatum; foliis suboppositis linearibus, floribus axillaribus solitariis 6-andris.—*Willd. sp. pl. 2. p. 868.*

Near the sea-coast of Virginia and Carolina.  $\frac{1}{4}$ . July, Aug. v. v. Three or four feet high; flowers small, white.

1. C. visco  
ribus  
culati  
Balsamo  
Lythrur  
Lythrur  
Icon. J  
In the n  
July.  
purple

II  
40  
1. Fotherg  
F. Gard  
 $\alpha.$  F. foliis  
Icon. B  
 $\beta.$  F. foliis  
dental  
Icon. B  
 $\gamma.$  F. foliis  
F. Gard  
In shad  
lina.  
spike  
menti  
specie

1. A. hirsut  
foliolis  
spicis  
plo lo  
basi su  
Icon. F  
Near the  
Caroli

407. CUPHEA. *Jacq. hort. 2. p. 83.*

1. *C. viscosa*; foliis oppositis petiolatis ovato-oblongis, floribus 12-andris laterali bus solitariis brevissime pedunculatis.—*Willd. sp. pl. 2. p. 870.*  
*Balsamona Pinto. Vandell. fasc. t. 3.*  
*Lythrum Cuphea. Linn. suppl. 249.*  
*Lythrum petiolatum. Willd. sp. pl. 2. p. 867.*  
*Icon. Jacq. hort. 2. t. 177.*  
 In the mountains of Virginia and Kentucky. ☽. June,  
 July. v. v.; v. s. in *Herb. Gronov.* Flowers, large,  
 purple.

## II. DI-PENTAGYNIA.

408. FOTHERGILLA. *Gen. pl. 922.*

1. *Fothergilla. Willd. sp. pl. 2. p. 1224.* *ulnifolia.*  
*F. Gardeni. Mich. fl. amer. 1. p. ....*  
 $\alpha$ . *F. foliis cuneato-obovatis superne crenato-dentatis.* *obtusa.*  
*Icon. Bot. mag. 1341. Lam. illustr. 480.*  
 $\beta$ . *F. foliis ovato-oblongis basi subcordatis superne crenato-dentatis.* *major.*  
*Icon. Bot. mag. 1342.*  
 $\gamma$ . *F. foliis ovatis acutis subintegerrimis.* *acuta.*  
*F. Gardeni. Jacq. ic. rar. 1. t. 100.*  
 In shady woods, on the side of hills: Virginia to Carolina. ♀. March, April. v. v. Flowers in close spikes or catkins, white. The different varieties here mentioned are rather to be considered as distinct species.

409. AGRIMONIA. *Gen. pl. 830.*

1. *A. hirsuta*; foliis interrupte-pinnatis, impari petiolato, foliolis obovatis inciso-argute-dentatis glabriusculis, spicis virgatis, floribus subsessilibus, petalis calyce duplo longioribus, fructibus obconicis divaricato-hispidis basi sublaevibus.—*Willd. sp. pl. 2. p. 875.*  
*Icon. Fl. dan. 580. Curt. lond. 32.*  
 Near the edge of woods on sunny hills: Pennsylvania to Carolina. ♀. June, July. v. v. Flowers yellow.

*pariflora.*

2. A. hirsuta ; foliis interrupte pinnatis, impari sessili, foliolis plurimis linear-lanceolatis inciso-serratis, spicis virgatis, floribus brevissime pedicellatis, petalis calyce sesquilonioribus, fructibus subrotundis divaricato-hispidis.—*Willd. sp. pl. 2. p. 876.*

In woods, on the side of hills : Pennsylvania to Virginia.  
J. June, July. v. v. Flowers small, yellow.

*suaveolens.*

3. A. hirsutissima ; foliis interrupte-pinnatis, impari sessili, foliolis plurimis lanceolatis argute serratis subtus hirsutis, spicis virgatis viscosis, floribus brevi-petiolatis, petalis calyce duplo longioribus, fructibus obconicis divaricato-hispidis.

On high mountain-meadows : Virginia and Carolina. J. July, Aug. v. v. Grows to the height of five feet ; is covered with very long brownish hairs : the flower spikes are likewise very closely covered with viscous hairs, and have a very agreeable balsamic scent : the flowers are small and pale yellow.

*striata.*

4. A. spicis virgatis, fructibus reflexis turbinatis sulcato-striatis apice tantum et quasi coronatim hispidis. *Mich. fl. amer. 1. p. 287.*

In Canada. *Michaux.* J. +. Flowers white.

#### 410. CRATÆGUS. *Geh. pl. 854.*

*apiifolia.*

1. C. spinosa ; foliis deltoideis inciso-lobatis, lobis acutis inciso-dentatis, corymbi pedicellis subsimplicibus, calyx tubo oblongo, lacinia subserratis.—*Mich. fl. amer. 1. p. 287.*

C. Oxyacantha. *Walt. fl. rar. 147.*

In low woods and on the banks of rivers : Virginia and Carolina ; plentifully on the banks of Roanoke-river, Southampton. J. May, June. v. v. This shrub, with its elegant foliage and scarlet berries, would be one of the finest ornaments to be introduced into the gardens, particularly for the purpose of making hedges ; which object has been so much neglected in North America.

*spathulata.*

2. C. ramulis spinescentibus fasciculatum foliosis, foliis parvis longissime deorsum angustatis subspatulatis trifidis, corymbis paucifloris, pedicellis brevibus, calycibus tomentosis.—*Mich. fl. amer. 1. p. 288.*

In dry woods, near rivers : Virginia and Carolina. J. May, June. v. v. The thorns are very large and crooked ; leaves small.

3. C. sp.  
ac  
gl  
W  
Mesp  
Icon  
In w  
M  
4. C. sp.  
and  
flo  
C. co  
C. co  
Mesp  
Mesp  
Mesp  
In he  
Vit  
sca

5. C. sp.  
tis  
line  
pl.  
C. leu  
C. tor  
C. ed  
Mesp  
Mespi  
In ro  
Pen

6. C. sp.  
peti  
bacc  
C. pru  
In cop  
App

7. C. sp.  
bris  
bacc  
C. san  
Mespi  
In Ca  
May  
Esq

3. *C. spinosa*; foliis cordato-ovatis inciso-angulatis glabris acute-serratis, petiolis calycibusque pubescentibus glandulosis, petalis orbiculatis, floribus 5-gynis.—  
*Willd. sp. pl. 2. p. 1000.*

*Mespilus æstivalis. Walt. fl. car. 148.*

*Icon. Pluk. alm. t. 46. f. 4.*

In woods and hedges: Canada to Carolina.  $\frac{1}{2}$ . April, May. v. v. Berries red, large, and agreeable to eat.

4. *C. spinosa*; foliis cordato-ovatis basi truncatis inciso-angulatis glabris, petiolis calycibusque eglandulosis, floribus pentagynis. *Walt. fl. car. 147.*

*C. corallina. H. P.*

*C. cordata. Willd. sp. pl. 2. p. 1000.*

*Mespilus acerifolia. Lam. encycl. 4. p. 442.*

*Mespilus Phœnopyrum. Linn. suppl. 254.*

*Mespilus cordata. Mill. ic. t. 179.*

In hedge rows and on the banks of rivers: Canada to Virginia.  $\frac{1}{2}$ . May, June. v. v. Berries small, scarlet.

5. *C. spinosa* inermis; foliis ovato-ellipticis inciso-serratis subuplicatis subhirtis, calycibus villosiusculis, foliolis linear-laticeolatis serratis, floribus trigynis. *Willd. sp. pl. 2. p. 1001.*

*C. leucophleos. Mænch. weissenst. p. 31. t. 2.*

*C. tomentosa. Du Roi harbk. 1. p. 183.*

*C. edulis. Loddiges catal.*

*Mespilus calpodendron. Ehrh. beitr. 2. p. 67.*

*Mespilus latifolia. Lam. encycl. 4. p. 444.*

In rocky and gravelly woods, and on banks of rivers: Pennsylvania to Carolina.  $\frac{1}{2}$ . June. v. s.

6. *C. spinosa*; foliis ellipticis inæqualiter serratis glabris, petiolis calycibusque glandulosis, laciniis calycis obtusis, baccis globosis 5-spérmiss.—*Willd. sp. pl. 2. p. 1002.*

*C. prunifolia. Lam. encycl. 4. p. 443.*

In copses and dry swamps: Canada to Carolina.  $\frac{1}{2}$ . April, May. v. v. Fruit red, small.

7. *C. spinosa*; foliis obovato-cuneiformibus angulatis glabris nitidis, petiolis stipulis calycibusque glandulosis, baccis ovalibus 5-spérmiss. *Willd. sp. pl. 2. p. 1002.*

*C. sanguinea. Pall. fl. ross. 1. p. 25. t. 11.*

*Mespilus rotundifolia. Ehrh. beitr. 3. p. 20.*

In Canada and on the Allegany mountains.  $\frac{1}{2}$ . April, May. v. v. Berries middle-sized, scarlet. M. Lewis, Esq. collected it on the Rocky-mountains.

*coccinea.*

*populifolia.*

*pyrifolia.*

*elliptica.*

*glandulosa.*

## ICOSANDRIA DI-PENTAGYNIA. Crataegus.

*flava.*

8. *C. spinosa*; foliis obovato-cuneiformibus sublobatis crenato-serratis, petiolis brevibus, stipulis cordatis calycibusque glandulosis, floribus subsolitariis, baccis turbinatis tetraspermis.—*Willd. sp. pl. 2. p. 1002.*

*C. viridis. Walt. fl. car. 147.**C. glandulosa. Mich. fl. amer. 1. p. 288.**C. Michauxii. Pers. syn. 2. p. 38.*  
*Mespilus flexispina. Moench. weissenst. p. 62. t. 4.**Mespilus caroliniana. Lam. encycl. 4. p. 442.*

In sandy, shady places: Virginia to Carolina.  $\frac{1}{2}$ . May.  
v. v. Berries large, yellow, eatable; flowers large.

*parvifolia.*

9. *C. spinosa*; foliis cuneiformi-ovatis incisis serratis subtomentosis, foliolis calycinis lanceolatis incisis longitudine fructus, floribus solitariis 5-gynis, fructibus subturbinatis punctato-verrucosis.—*Willd. sp. pl. 2. p. 1002.*

*C. tomentosa. Sp. pl. 682.**C. uniflora. Du Roi harbk. 1. p. 184.**Mespilus Xanthocarpus. Linn. suppl. 254.**Mespilus laciniata. Walt. fl. car. 147.*

In sandy, shady woods: New Jersey to Carolina.  $\frac{1}{2}$ .  
May, June. v. v. A low shrub; leaves small; fruit  
large, greenish-yellow.

*punctata.*

10. *C. spinosa inermisve*; foliis obovato-cuneiformibus subplicatis glabris incisis serratis in petiolum decurrentibus, calycibus villosis, laciniis subulatis integerrimis, baccis subglobosis umbilico depresso.—*Willd. sp. pl. 2. p. 1004.*

*C. Crus galli. Du Roi harbk. 1. p. 195.**Mespilus cuneifolia. Ehrh. beitr. 3. p. 21.**Mespilus cornifolia. Lam. encycl. 4. p. 444.**Icon. Jacq. hort. 1. t. 28.**α. C. fructibus rubris. Ait. kew. ed. 2. t. 3. p. 202.**β. C. fructibus flavis. Ait. l. c.*

In the woods and swamps of Virginia and Carolina.  $\frac{1}{2}$ .  
May, June. v. v. Is inclined to grow to a handsome tree; berries yellow, with dark spots.

*Crus galli.*

11. *C. spinosa*; foliis obovato-cuneiformibus subsessilibus nitidis coriaceis, corymbis compositis, foliolis calycinis lanceolatis subserratis, floribus digynis. *Willd. sp. pl. 2. p. 1004.*

*C. hyemalis. Walt. fl. car. 147.**C. lucida. Wangh. amer. p. 53. t. 17. f. 42.**Mespilus lucida. Ehrh. beitr. 4. p. 17.*

Icon. *Trew ic. var. 2. t. 17.* foliis lanceolatis crenato-dentatis subtus tomentosis, rachi supra glandulosa, floribus corymbosis, calycibus tomentosis.—*Willd. sp. pl. 2. p. 102.*

- $\alpha$ . C. foliis obovato-cuneiformibus. *Ait. kew. ed. 2. t. 3.* splendens.  
 $\beta$ . C. foliis oblongo-lanceolatis subcuneiformibus. *Ait. l. c. pyracanthifolia.*  
 $\gamma$ . C. foliis lanceolatis. *Ait. l. c. salicifolia.*

In woods and hedges, on the banks of rivers, common : Canada to Carolina.  $\frac{1}{2}$ . April, May. v. v. Berries small, scarlet.

#### 411. PYRUS. Gen. pl. 858.

1. P. inermis; foliis obovato-oblongis acutis crenato-dentatis subtus tomentosis, rachi supra glandulosa, floribus corymbosis, calycibus tomentosis.—*Willd. sp. pl. 2. p. 1012.*

*Crataegus pyrifolia. Lam. encycl. 1. p. 83.*

*Mespilus arbutifolia. Sp. pl. 685.*

*Mespilus arbutifolia erythrocarpa. Mich. fl. amer. 1. p. 291.*

*Mespilus pumila. Schmidt arb. 88.*

Icon. *Mill. ic. 109.*

In low copses and swamps; common : Canada to Carolina.  $\frac{1}{2}$ . May, June. v. v. A low shrub; berries red: leaves more or less tomentose.

2. P. inermis; foliis obovato-oblongis acuminatis serratis *melanocarpa.* subtus glabris, rachi supra glandulosa, floribus corymbosis, calycibus glabris. *Willd. enum. 525.*

*Crataegus arbutifolia. Lam. encycl. 1. p. 83.*

*Mespilus arbutifolia. Schmidt arb. 86.*

*Mespilus arbutifolia melanocarpa. Mich. fl. amer. 1. p. 292.*

In bogs of Canada and on the high mountains of Pennsylvania, Virginia, and Carolina.  $\frac{1}{2}$ . May, June. v. v. Berries large, black, and resembling in taste those of *Vaccinium pensylvanicum.*

3. P. inermis; foliis oblongo-ellipticis cuspidatis adultis *Botryapium.* glabris, floribus racemosis, petalis linearibus lanceolatis, germinibus pubescensibus, calycinis segmentis glabris. *Willd. sp. pl. 2. p. 1013.*

*Crataegus racemosa. Lam. encycl. 1. p. 84.*

*Mespilus canadensis. Sp. pl. 685.*

*Mespilus canadensis  $\beta$ . cordata. Mich. fl. amer. 1. p. 291.*

Icon. *Schmidt arb. 84.*

- In woods and hedges; frequent: Canada to Carolina.  
*h.* April, May. *v. v.* A small tree; leaves covered with a white tomentum, when young; berries black, very agreeably tasting.
- ovalis.*
4. *P. inermis*; foliis subrotundo-ellipticis acutis glabris, floribus racemosis, petalis obovatis, germinibus calycinisque segmentis pubescentibus. *Willd. sp. pl. 2. p. 1014.*
  - Crataegus spicata. *Lam. encycl. 1. p. 84.*
  - Mespilus Amelanchier. *Walt. fl. car. 184.*
  - Mespilus canadensis  $\alpha$ . obovalis. *Mich. fl. amer. 1. p. 291.*
- In swamps: New Jersey to Carolina. *h.* April, May. *v. v.* A small shrub; berries black and eatable.
- sanguinea.*
5. *P. inermis*; foliis ovalibus utrinque obtusis mucronatis tenuissime serratis basi subcordatis, racemis paucifloris, calycibus glabris, petalis linearibus obtusis.
  - Mespilus canadensis  $\gamma$ . rotundifolia. *Mich. fl. amer. 1. p. 291.*
- In Canada, and on the banks of the Columbia. *h.* April, May. *v. v.* A small tree with blood-red branches; berries red, eatable.
- Malus.*
6. *P. umbellis* sessilibus, foliis ovato-oblongis acuminatis serratis glabriusculis, unguibus calyce brevioribus, stylis glabris. *Willd. sp. pl. 2. p. 1017.*
  - The Common Apple-tree frequently occurs in woods apparently wild, but has been seminated by birds.
- coronaria.*
7. *P. foliis lato-ovalibus* basi rotundatis subangulatis serratis laevibus, pedunculis corymbosis.—*Willd. sp. pl. 2. p. 1018.*
  - In woods and near plantations: Pennsylvania to Carolina. *h.* May. *v. v.* The Sweet-scented Crab-tree, the fruit of which is well known as a most excellent preserve for the table, is a very fine ornamental tree, not only for the beauty but particularly for the fine violet-scent of its flowers.
- angustifolia.*
8. *P. foliis lanceolato-oblongis* basi acutis leviter crenato-dentatis nitidis, pedunculis corymbosis.—*Willd. sp. pl. 2. p. 1020.*
  - P. coronaria.* *Wangh. amer. t. 21. f. 47.*
  - In low woods of Carolina. *h.* May. *v. v.* Resembles the foregoing species; the fruit very small.

## 412. SORBUS. Gen. pl. 855.

1. *S. foliis pinnatis, foliolis acutis subæqualiter serratis petioloque communi glaberrimis.* — *Willd. enum. 520.* *americanæ.*

*S. Aucuparia β.* *Mich. fl. amer. 1. p. 290.*

In Canada and on some of the northern mountains.  $\frac{1}{2}$ . May, June. v. v. Berries purple, not scarlet as in the European species.

2. *S. foliis pinnatis, foliolis acuminatis inæqualiter inciso-serratis petioloque communi glabris, serraturis setaceo-mucronatis.* *microcarpa.*

*S. Aucuparia α.* *Mich. fl. amer. 1. p. 290.*

On the peaks of high mountains: New Jersey to Carolina.  $\frac{1}{2}$ . June. v. v. A large shrub; the young branches covered with a shining, dark brown gloss; berries small, scarlet. This species is very distinct from the Canadian *Sorbus*.

## 413. SPIRÆA. Gen. pl. 862.

\* *Fruticosæ.*

1. *S. foliis lanceolatis argute-serratis glabris, racemo terminali composito paniculato, floribus 5-gynis.* — *Willd. sp. pl. 2. p. 1055.* *salicifolia.*

- α. S. foliis lanceolatis, racemis paniculatis divaricatis, petalis albis cortice ramorum rubris.* *Ait. kew. 2. p. 198.*

*S. alba.* *Ehrh. beitr. 7. p. 137.*

Icon. *Mill. ic. t. 257. f. 2.*

- β. S. foliis ovato-oblongis, racemis paniculatis, petalis albis. cortice ramorum rufescente.* *Ait. l. c.*

In swamps and on the edge of ditches and slow-flowing rivers: Canada to Carolina; particularly on the mountains.  $\frac{1}{2}$ . June—Aug. v. v. This shrub is very variable in size, shape, and colour.

2. *S. foliis lanceolatis inæqualiter-serratis subtus tomentosis, racemo terminali composito confertifloro, floribus 5-gynis.* — *Willd. sp. pl. 2. p. 1056.* *tomentosa.*

Icon. *Schmidt arb. 57. Pluk. alm. t. 321. f. 5.*  
In Canada and on the Allegany mountains: New York to Carolina.  $\frac{1}{2}$ . July—Sept. v. v. A small shrub; flowers red.

3. *S. foliis obovatis integerrimis seu apice dentatis glabris, umbellis sessilibus.* — *Willd. sp. pl. 2. p. 1057.* *hypericifolia.*

Icon. *Schmidt arb. 56 & 57.*

- chamædrifolia.* In dry swamps of Canada and New York.  $\text{h}.$  May, v. v. Flowers white, small, and in great abundance.
4. *S. foliis obovatis apice inciso-dentatis glabris, corymbis pedunculatis.* —Willd. sp. pl. 2. p. 1058.  
Icon. Schmidt arb. 53.
- media.*  $\beta.$  *S. foliis utrinque subvillosis paudentatis.* Schmidt arb. 54.  
In Canada and on the north-west coast.  $\text{h}.$  May, June. v. v.
- betulifolia.* 5. *S. foliis lato-ovatis inciso-serratis glabris, corymbis terminalibus compositis fastigiatis foliosis.* —Pall. fl. ross. 1. p. 33. t. 16.  
In the mountains of Virginia.  $\text{h}.$  May—July. v. v. Not above a foot high; flowers tinged with red. This species is very distinct from *S. ulmifolia* Willd. sp. pl. 2. p. 1058, to which it has been referred as a synonym.
- ulmifolia.* 6. *S. foliis ovatis lobatis duplicato-dentatis crenatis glabris, corymbis terminalibus confertifloris, floribus 3-gynis, capsulis inflatis.* —Willd. sp. pl. 2. p. 1059.  
Icon. Schmidt arb. 32. Rob. ic. 134. Comm. hort. 1. t. 87.  
On the banks of rivers, particularly in the mountains: Canada to Carolina.  $\text{h}.$  June, July. v. v. Flowers white; generally known by the name of Nine-bark.
- capitata.* 7. *S. foliis ovatis sublobatis duplicato-dentatis subtus reticulatis tomentosis, corymbis terminalibus congestis subcapitatis longissime pedunculatis, calycibus tomentosis.*  
On the north-west coast. Menzies.  $\text{h}.$  June. v. s. in Herb. Banks. An imperfect specimen of this species I observed in the Lewisian Herbarium, gathered on the Columbia.
- discolor.* 8. *S. foliis ovatis lobatis dentatis subplicatis subtus niveo-tomentosis, paniculis terminalibus pedunculatis ramosissimis.*  
On the banks of the Kooskoosky.  $\text{h}.$  June, July. v. s. in Herb. Lewis. A shrub about five feet high.
- sorbifolia.* 9. *S. foliis pinnatis, foliolis uniformibus serratis, floribus paniculatis.* Willd. sp. pl. 2. p. 1060.  
Icon. Schmidt arb. 58. Pall. fl. ross. 1. t. 24.  
On the north-west coast, and probably in Canada.  $\text{h}.$  July, Aug. v. v. A handsome small shrub; flowers white.
10. *S. foliis dioides.* Icon.  $\beta.$  *S. foliis ma-* In th Jul  
11. *S. foliis bus p.* S. pal. Icon. In fe Jul  
in 1  
12. *S. foliis line lati sp.* Gille Icon. In sha cip Jul  
wh prc  
13. *S. foliis stipinal* —  
 $\beta.$  *S. foliis in sha* Jul  
Th not stir

\*\* *Herbaceæ.*

10. S. foliis 2-3-pinnatis, spicis paniculatis, floribus 3-gynis dióicis.—*Willd. sp. pl. 2. p. 1060.*  
 Icon. *Pall. fl. ross. 1. t. 26.*  
 β: S. foliis lucidis, panicula oblonga, spicis gracilibus hermaphroditis.  
 In the mountains: Pennsylvania to Georgia. Ȑ. June, July. v. v. Flowers white; a very elegant plant.
11. S. foliis pinnatis glabris, impari majore 7-lobo, laterali bus 3-lobis, corymbis proliferis. *Willd. sp. pl. 2. p. 1062.*  
*S. palmata. Linn. suppl. 262.*  
 Icon. *Jacq. hort. 1. t. 88.*  
 In fertile wet meadows: Virginia and Carolina. Ȑ. July, Aug. v. v. A beautiful perennial; flowers red, in large clusters.
12. S. foliis ternatis lanceolatis serratis subæqualibus, stipulis linearibus integris, floribus terminalibus laxè paniculatis 5-gynis, calyce tubuloso campanulato.—*Willd. sp. pl. 2. p. 1063.*  
*Gillenia trifoliata. Mænch. meth. suppl. p. 286.*  
 Icon. *Mill. ic. 256. Bot. mag. 489.*  
 In shady woods and on bogs: Canada to Florida; principally in the mountainous parts thereof. Ȑ. June, July. v. v. A very fine perennial; flowers large, white. It may, with the following species, with all propriety form a distinct genus.
13. S. foliis ternatis lanceolatis inciso-serratis subæqualibus, stipulis foliaceis ovatis inciso-dentatis, floribus terminalibus laxè paniculatis 5-gynis, calyce campanulato.—*Willd. enum. p. . . .*  
 β: S. foliis ternatis, foliolis pinnatifidis inciso-dentatis.  
 In shady woods: Kentucky and Tennessee. Ȑ. June, July. v. v. Flowers resembling the former species. The variety β. appears very different; but, as I have not seen it in flower, I did not venture to make it a distinct species.

*Aruncus.**americana.**lobata.**trifoliata.**stipulacea.**incisa.*

## III. POLYGYNIA.

414. ROSA, *Gen. pl.* 863.*blanda.*

1. *R. germinibus globosis glabris, caulibus adultis pedunculisque lævibus inermibus, foliolis (7) oblongis subæqualiter serratis glabris, petiolis glabris subaculeatis.*—*Willd. sp. pl. 2. p. 1065.*

Icon. *Andrews's roses. Miss Lawr. roses* 27.

In Canada, near Hudson's Bay, and on the river St. Lawrence. ♀. May—Aug. v. v. The younger branches are covered with straight red spines; the older ones unarmed and shining red.

*parviflora.*

2. *R. germinibus depresso-globosis pedunculisque hispidis, petiolis pubescentibus subaculeatis, caule glabro, aculeis stipularibus rectis, foliolis elliptico-lanceolatis simpliciter serratis glabris, floribus subgeminatis.*—*Willd. sp. pl. 2. p. 1068.*

*R. carolina. Du Roi herbk. 2. p. 335. Mich. fl. amer. 1. p. 295.*

*R. pensylvanica. Wangh. amer. p. 113.*

*R. humilis. Marsh. arb. 285.*

In woods on the side of hills: New York to Carolina. ♀. June, July. v. v. About two feet high. There are a number of varieties of this species.

*nitida.*

3. *R. germinibus globosis, calycibus pedunculis ramisque hispidis, petiolis pilosiusculis inermibus, foliolis (7) oblongo-lanceolatis utrinque nitidis glaberrimis.*—*Willd. enum. 544.*

In Pennsylvania and Virginia. ♀. June—Aug. v. v. Petals red, obcordate.

*lucida.*

4. *R. germinibus depresso-globosis pedunculisque subhispidis, petiolis glabris subaculeatis, caule glabro, aculeis stipularibus rectis, foliolis ovato-lanceolatis obtusiusculis grosse serratis glabris nitidis, floribus subgeminis, calycinis foliolis integris.*—*Willd. sp. pl. 2. p. 1068.*

*R. carolina. Walt. fl. car. 149.*

Icon. *Dill. elth. t. 245. f. 316.*

On the borders of swamps: New York to Carolina. ♀. July, Aug. v. v. From four to six feet high.

*gemella.*

5. *R. germinibus depresso-globosis pedunculisque glabris, floribus subgeminatis, foliis oblongis acutis opacis, pe-*

tiolis venisque subtus pubescentibus, aculeis uncinatis infra axillaribus caulinis geminatis.—Willd. enum. 544.

On dry sunny hills: New England to Carolina.  $\frac{1}{2}$ . July, Aug. v. v. A low shrub; flower red, large.

6. R. germinibus subglobosis glabriusculis, pedunculis hispidis, petiolis subaculeatis, caule glabro, aculeis sparsis rectis, foliolis (3—5) ovato-oblongis acutis serratis, supra glabriusculis, subtus tomentosis, superioribus simplicibus, floribus subternatis, stipulis linearibus, calycis laciniis tomentosis linearibus vix laciniatis.

In Tennessee. Lyon.  $\frac{1}{2}$ . July, v. s. in Herb. Lyon. Flowers pale red; leaves small, with coloured veins.

7. R. germinibus globosis, petiolis nervisque aculeatis, ramis glabris, aculeis geminis sparsisque, foliolis (3—5) acuminate glabris, calycinis foliolis subpennatim setigeris.—Mich. fl. amer. 1. p. 295.

In swamps of Virginia and Lower Carolina.  $\frac{1}{2}$ . June, July. v. v. c. fr. From five to eight feet high.

8. R. germinibus globosis pedunculisque subhispida, petiolis pilosis subaculeatis, caule glabro, aculeis stipularibus subuncinatis, foliolis (5—7) oblongo-lanceolatis acutis argute serratis subtus glaucis, floribus corymbosis.—Willd. sp. pl. 2. p. 1009.

R. corymbosa. Ehrh. beitr. 4. p. 21.

R. virginiana. Du Roi harbk. 2. p. 353.

R. pensylvanica. Mich. fl. amer. 1. p. 296.

Icon. Andrews's roses. Miss Lawr. roses 3. 24. 36. 54. 66. 68.

In swamps and on the banks of ponds: New England to Virginia.  $\frac{1}{2}$ . June, July. v. v. From five to six feet high. There are a great number of varieties of this species.

9. R. germinibus globosis pedunculisque hispidulis, calycibus inexpansis muticis, foliis ternatis subtus pubescens, petiolis glandulosis aculeatisque, caule glabro, aculeis stipularibus sparsisque subaduncis, floribus corymbosis. Ait. kew. ed. 2. t. 3. p. 260.

In North America. Mason.  $\frac{1}{2}$ . June, July. +.

10. R. germinibus ovatis promisse hispidissimis; calycis laciniis integris, aculeis geminatis recurvis, petiolis subaculeatis, foliolis (3—5) lanceolato-ovalibus subener-

Lyonii.

setigera.

carolina.

rubifolia.

levigata.

- vibus lævigatis, stipulis angustis subulato-mueronatis.  
—*Mich. fl. amer. 1. p. 295.*

In shady woods of Georgia.  $\text{h. v. v. s. fl.}$  An ever-green; climbing to a great height.

*suaveolens.* 11. R. germinibus ovatis, pedunculis petiolisque glanduloso-hispidis, petiolis subaculeatis, caule glabro, aculeis sparsis rectis tenuibus, foliolis (5—7) ovatis serratis subtus vix glanduloso-pilosus, ramulis unifloris, calycis laciniis integris.  
R. *Eglanteria americana*. *Andrews's roses*, cum icono.

*pendulina.* 12. R. inermis; germinibus oblongis, pedunculis petiolisque hispidis, caule ramisque glabris, fructibus pendulis.  
*Willd. sp. pl. 2. p. 1076.*  
Icon. *Miss. Lawr. ros. 9. Dill. elth. t. 245. f. 317.*  
In North America.  $\text{h. May, June. v. v. in Hortis.}$   
Known by the name of *Rose without Thorns*. I never have seen this plant in its indigenous state.

415. RUBUS. *Gen. pl. 864.*

\* *Fruticosi.*

*Idæus.* 1. R. foliis quinato-pinnatis ternatisque rhomboe-ovatis acuminatis subtus tomentosis, petiolis canaliculatis, caule aculeato-hispido, floribus subpaniculatis.—*Willd. sp. pl. 2. p. 1081.*  
Icon. *Fl. dan. 788.*  
In hedge-rows: Canada to Pennsylvania.  $\text{h. May, June. v. v.}$  The *Wild Raspberry* has a number of varieties.

*villosus.* 2. R. pubescens, hispidus aculeatusque; foliis 3—5-digitatis, foliolis ovato-oblongis acuminatis serratis utrinque pubescentibus, caulibus petiolisque aculeatis, calyce brevi acuminato, racemo laxo, pedicellis solitariis.—*Willd. sp. pl. 2. p. 1085.*  
In old fields and commons; frequent: New England to Carolina.  $\text{h. June, July. v. v.}$  Known by the name of *Black-berries*.

*strigosus.* 3. R. inermis, rigide hispidissimus; foliolis 3. aut pinnato-quinatis ovalibus basi obtusis acuminatis subtus lineatis et candido-tomentosis, calycibus acuminatis, floribus in apice ramulorum axillaribus solitariis, pedunculis calycibusque hispidis.—*Mich. fl. amer. 1. p. 297.*  
R. *pensylvanicum*. *Lam. encycl. 6. p. 246.*  
On the mountains: Canada to Virginia.  $\text{h. June, July. v. v.}$  An upright shrub; berries very agreeable to eat.

4. *R. glabriuscula*; foliis digitatis denis quinis ternatisque, foliolis lanceolatis utrinque nudis argute serratis, caule inermi, bracteis lanceolatis.—*Willd. sp. pl. 2. p. 1085.*  
In rocky barren woods: Canada and New England. ♂. June, July. v. v. Stem and old branches purple.
5. *R. ramis petiolis pedunculisque tomentosis sparsim recurvato-aculeatis*, foliis 3—5-digitatis, foliolis cuneato-ovatis superne inaequaliter dentatis plicatis margine integerrimo revolutis subtus tomentosis, racemis terminalibus paniculatis, pedicellis divaricatis nudiculis.  
*R. parvifolius*. *Walt. fl. cur. 149.*  
In sandy fields and woods: New Jersey and Carolina. ♂, June, July. v. v. A straggling briar of a gray aspect; the berries hard and dry.
6. *R. ramis petiolisque glaucis et aculeatis*, foliis ternis ovoidibus acuminatis sublobatim duplicato-serratis subtus cano-tomentosis, petiolis teretibus, aculeis recurvatis, racemis terminalibus.—*Willd. sp. pl. 2. p. 1082.*  
Icon. *Dill. elth. t. 247. f. 319.*  
In rocky and mountainous situations: Canada to Carolina. ♂. May, June. v. v. The stem is smooth, and covered with a blueish dew. Berries black or red; known in the gardens by the name of *Virginian Raspberry*.
7. *R. sarmentoso-procumbens*; caulis petiolis pedunculisque setis rigidis hispidissimis, foliis ternatis inciso-serratis nudis, intermedio pedicellato.—*Willd. sp. pl. 2. p. 1083.*  
In Canada. *Kalm.* ♂. +. Resembles some of the following species, particularly No. 9.
8. *R. sarmentoso-procumbens*; petiolis pedunculisque recurvato-aculeato-hispidis, stipulis subulatis, foliis ternatis quinatisve oblongo-ovalibus acutis inaequaliter serratis subpubescentibus, pedicellis solitariis elongatis, petalis obovatis calyce triplo longioribus.—*Mich. fl. amer. 1. p. 296.*  
In old fields, common: New England to Carolina. ♂. May, June. v. v. Flowers large; berries black, large, and very agreeably tasting; known by the name of *Dewberries*.
9. *R. sarmentoso-procumbens*; caule tereti petiolisque recurvato-aculeatis, foliis ternatis glabris inaequaliter

*canadensis.**cuneifolius.**occidentalis.**hispidus.**trivialis.**flagellaris.*

serratis, intermedio ovato basi cuneato, lateralibus rhombeis.—*Willd. enum.* 549.

*R. trivialis. Ait. kew. ed. 2. t. 3. p. 269.*

In fields and sandy woods: Virginia and Carolina. ♂. June, July. v. v. The leaves are smaller, smooth, and equal on the base; petals more orbicular than No. 8.

*inermis.*

10. *R. caule petiolis pedunculisque inermibus, foliis ternatis ovatis acutis inæqualiter serratis subtus tomentosis, lateralibus subincisis, stipulis setaceo-subulatis.*—*Willd. enum.* 548.

*R. hispidus, Walt. fl. car. 149?*

In Pennsylvania. ♂. +.

*spectabilis.*

11. *R. inermis glabra; foliis ternatis ovatis acutis duplicato-inæqualiter serratis subtus pubescentibus, pedunculis terminalibus unifloris solitariis, petalis ovatis.*

On the banks of the Columbia. *M. Lewis.* On the north-west coast, *Menzies.* ♂. April, May. v. s. in *Herb. Lewis. nec non Banks.* The figure was taken from a specimen in the collection of A. B. Lambert, Esq.

*Frutex elegans, 4-5-pedalis. Rami geniculati, subflexuosi, teretes, lœvigati. Gemmæ alternæ, ferrugineæ, squamæ obtusæ. Folia e gemmis subfasciculata, ternata. Foliola ovato-oblonga, acuta, inciso-duplicato-serrata, supra glabra, subtus pubescentia, lateralia basi obliqua, intermedio petiolato. Petoli pubescentes, aculeo parvo recurvato uno alterove instructi. Stipulæ lineares. Pedunculi terminales uniflori solitarii. Flores magnitudine *R. odorati*, profunde et amœne purpurascentes. Calyx laciniis oblongis, brevi-acuminatis, pubescentibus. Petala ovata, calyce plusquam duplo longiora.*

*odoratus.*

12. *R. inermis, erectus, viscidо-hispidus; foliis simplicibus acute 3—5-lobatis, corymbis terminalibus divaricatis, calycibus appendiculatis, petalis suborbiculatis.*—*Willd. sp. pl. 2. p. 1085.*

*Icon. Bot. mag. 323. Mill. ic. 223. Corn. canad. t. 159.*

In the woods of Canada and on the Allegany mountains: New York to Carolina. ♂. June, July. v. v. A fine ornamental shrub; flowers rose-coloured or crimson; berries yellow, of a very fine flavour and large size, but scarcely ever produced in the gardens.



*Rubus spectabilis.*

Drawn & Engraved by W. Hooke

- 100
13. *R. herbaceus*  
natis rho-  
tiolato, fl.  
*fl. amer.*  
In Canada,  
ginia. 2
14. *R. fruticosa*  
tundato-c  
censis flo  
ovatis, pe  
In swamps,  
York to  
with only
15. *R. herbaceus*  
ternatis  
emarginata  
Icon. Engl.  
In Labrador  
*v. s. in H*  
beautiful
16. *R. herbaceus*  
natis glab  
stylis app  
*R. acaulis.*  
In bogs of  
June, Jul  
foregoing
17. *R. herbaceus*  
incisis, P  
lycibus gl  
t. 63.  
On the nort  
*Banks.*  
belong to
18. *R. herbaceus*  
foliis sim  
talis lance  
*ined. f. 3.*  
On the nort  
*Banks.*
19. *R. herbaceus*

## \*\* Subherbacei.

13. *R. herbaceus*, pubescens; flagellis reptantibus, foliis ternatis rhomboides acutis inciso-dentatis nudis, impari petiolato, floribus subternis, pedicellis elongatis.—*Mich. fl. amer. 1. p. 298.*

In Canada, and on the mountains of New York to Virginia.  $\frac{1}{2}$ . June, July. v. v. Berries small, black.

14. *R. fruticosens*, pilis rigidis hispidis; foliis ternatis rotundato-obovalibus serratis nudis, stipulis setaceis, racemis floriferis subcorymbosis paucifloris, bracteis ovatis, pedicellis elongatis.—*Mich. fl. amer. 1. p. 298.*

In swamps, among sphagnum, on the mountains: New York to Carolina.  $\frac{1}{2}$ . May—July. v. v. Berries with only a few large grains black and sweet.

15. *R. herbaceus*, pusillus; caule inermi subunifloro, foliis ternatis glabris obtuse serratis, petalis subrotundis emarginatis.—*Willd. sp. pl. 2. p. 1088.*

Icon. *Engl. bot. 1585. Bot. mag. 132.*

In Labrador and near Hudson's Bay.  $\frac{1}{2}$ . May, June. v. s. in *Herb. Lambert.*; v. v. in *Hortis.* Flowers of a beautiful rose colour; berries amber, very delicious.

16. *R. herbaceus*, pusillus; caule inermi unifloro, foliis ternatis glabris argute serratis, petalis oblongis integris, stylis approximatis. *Smith exot. bot. 2. p. 53. t. 86.*

*R. acaulis. Mich. fl. amer. 1. p. 298.*

In bogs of Canada, and on the north-west coast.  $\frac{1}{2}$ . June, July. v. s. in *Herb. Dickson.* Resembles the foregoing one very much.

17. *R. herbaceus*, pusillus, repens; foliis pedato-quinatis incisis, pedunculis filiformibus medio bracteatis, calycibus glabriusculis reflexis. *Smith ic. ined. f. 3. p. 63. t. 63.*

On the north-west coast. *Menzies.*  $\frac{1}{2}$ . v. s. in *Herb. Banks.* Petals white? This species may probably belong to the genus *Dalibarda.*

18. *R. herbaceus*, pusillus; caule inermi unifloro erecto, foliis simplicibus cordatis trilobis rugoso-venosis, petalis lanceolatis.—*Willd. sp. pl. 2. p. 1089. Smith ic. ined. f. 3. t. 64.*

On the north-west coast. *Menzies.*  $\frac{1}{2}$ . v. s. in *Herb. Banks.* Flowers purple.

19. *R. herbaceus*, pusillus; caule inermi unifloro erecto,

*saxatilis.*

*obovalis.*

*arcticus.*

*pistillatus.*



*pedatus.*

*stellatus.*

*Chamæmor-*  
*rus.*

foliis simplicibus subreniformibus rotundato-lobatis,  
petalis oblongis.—Willd. sp. pl. 2. p. 1090.

Icon. Engl. bot. 716. Fl. dan. 1. Linn. fl. lapp. t. 5.  
f. 1.

In sphagnous bogs of Canada, and on high mountains of  
New England. ♀. May, June. v. v. Flowers  
white; berries yellow or amber-coloured.

**416. DALIBARDA.** Mich. fl. amer. 1. p. 299.

*repens.*

1. D. villosa; stolonibus reptantibus, foliis simplicibus cor-  
datis crenatis, pedunculis unifloris. Lam. encycl. 6.  
p. 250.

D. violacoides. Mich. fl. amer. 1. p. 299.

Rubus Dalibarda. Willd. sp. pl. 2. p. 1090.

Icon. Mich. l. c. t. 27. Smith ic. ined. t. 20. Lam. il-  
lustr. t. 441. f. 3.

In shady woods and bogs of Canada, and on the high  
mountains of New England and Pennsylvania. ♀. May,  
June. v. v. Flowers white.

*fragaroides.*

2. D. foliis ternatis, foliolis cuneatis serrato-incisis ciliatis,  
pedunculis multifloris, calycis tubo obconico.—Mich.  
fl. amer. 1. p. 300. t. 28.

Dryas trifoliata. Pallas in Herb. Lambert.

In shady beech-woods of Canada, and on the Allegany  
mountains. ♀. May, June. v. v. Flowers yellow.

**417. DRYAS.** (Gen. pl. 868.

*octopetala.*

1. D. foliis ovato-oblongis grosse-dentatis rugosis subtus  
niveo-tomentosis, pedunculis unifloris.—Willd. sp.  
pl. 2. p. 1118.

Icon. Engl. bot. 451. Fl. dan. 31.

On high mountains in the north-west parts of Canada.  
♀. July, Aug. v. s. Flowers large, white.

*tenella.*

2. D. microphylla; foliis ovatis acutis basi cordatis integer-  
rimis planis subtus niveo-tomentosis, pedunculis uni-  
floris.

Icon. Egede, descrip. of Greenland, p. 42. t. B. f. 3.

On the white hills of New Hampshire. Prof. Peck.  
♀. July. v. s. in Herb. Banks. Flowers not half  
the size of the former.

418. GEUM. *Gen. pl. 867.*

1. *G. hirsutum*; foliis omnibus interrupte pinnatis, impari ampliore, foliolis ovatis dentatis, stipulis incisis, calycis laciniis 5. alternis linearibus brevibus, floribus erectis, petalis suborbiculatis calyce longioribus, aristis nudis uncinatis.—*Willd. sp. pl. 2. p. 1113.*  
*G. canadense. Murr. in comment. gött. 5. p. 33. f. 4. B.*  
*G. aleppicum. Jacq. ic. rar. 1. t. 93.*  
 In wet meadows and bogs : Canada and New York. 2.  
 May, June. v. v. Flowers large, yellow.
2. *G. hirsutissimum*; foliis omnibus pinnatis, foliolis sub- *agrimonoides.*  
 æqualibus inæqualiter inciso-dentatis, stipulis ovatis subintegris, floribus erectis, calycis laciniis subæqualibus, petalis ovalibus longitudine calycis.  
*G. jacinatum. Murr. in comm. gött. 5. p. 36. t. 2?*  
 On the rocky banks of the Susquehanna, Pennsylvania,  
 and on the upper parts of the Missouri. 2. June,  
 July. v. v. Flowers white.
3. *G. pubescens*; foliis radicalibus pinnatis, caulinis ternatis, summis simplicibus trifidis, stipulis inferioribus incisis, floribus erectis, petalis longitudine calycis, aristis uncinatis nudis apice pilosis.—*Willd. enum. 556.*  
*G. canadense. Jacq. hort. 2. p. 82. t. 175.*  
*G. carolinianum. Walt. fl. car. 150?*  
 On the banks of rivers : Pennsylvania to Carolina. 2.  
 June, July. v. v. Flowers white.
4. *G. pubescens*; foliis radicalibus caulinisque infimis ternatis, superioribus lanceolatis, stipulis ovatis subintegris, floribus erectis, petalis calyce brevioribus, aristis uncinatis nudis apice pilosis tortuosus.—*Willd. sp. pl. 2. p. 1113.*  
*Icon. Murr. in comm. gött. 5. p. 32.*  
 In shady woods, common : Canada to Carolina. 2.  
 July, Aug. v. v. Flowers white, very small.
5. *G. subpaniculatum multiflorum*; foliis caulinis subsessilibus tripartitis, stipulis integris, petalis cuneato-obcordatis, aristis prorsus pubescentibus medio geniculatis.—*Mich. fl. amer. 1. p. 300.*  
 In Canada. *Michaux. 2. +.*
6. *G. pubescens*; caule simplici, foliis radicalibus interrupte-pinnatis, caulinis trifidis, floribus nutantibus, petalis longitudine calycis, aristis plumosus superne nudius-

*strictum.**album.**rivale.*

## ICOSANDRIA POLYGYNIA. Geum.

- culis minutissime uncinatis.—*Willd. sp. pl. 2. p. 1115.*  
*Mich. fl. amer. 1. p. 301.*  
*Icon. Fl. dan. 722. Engl. bot. 106.*
- In wet boggy meadows : Canada and on the New York and Pennsylvania mountains.  $\frac{1}{2}$ . June, July. v. v. Flowers yellowish purple. The flowers are smaller, the petals more rounded on the top, and the leaves with deeper incisions, than the European plant : compared in a living state, they may be found different species.
- ciliatum.* 7. *G. pubescens* ; caule simplici, foliis glabriusculis margine ciliatis, inferioribus pinnatis, caulinis pinnatifidis, superioribus palmatis : laciniis linearibus incisis, floribus corymbosis.  
 On the banks of the Kooskoosky.  $\frac{1}{2}$ . June. v. s. in *Herb. Lewis.* An elegant species ; flowers resembling those of No. 6.
- radiatum.* 8. *G. hirsutissimum* ; caule simplici, foliis radicalibus pinnatis, impari amplissimo reniformi radianti-nervoso dentato, caulinis amplexicaulibus inciso-laceris, pedunculis axillaribus solitariis, aristis glabris apice simplicibus.—*Mich. fl. amer. 1. p. 300.*
- G. camtschaticum.* *Pallas in Herb. Lambert.*  
 On high mountains of Carolina. *Michaux.* On the north-west coast. *Menzies.* Likewise a handsome plant ; flowers yellow.
- Peckii.* 9. *G. glabriusculum* ; caule unifloro, foliis radicalibus reniformibus circinatis basi subtruncatis inciso-dentatis, petiolis longissimis interdum appendiculatis, petalis longitudine calycis.  
 On the white hills of New Hampshire. *Prof. Peck.*  
 $\frac{1}{2}$ . July. v. s. in *Herb. Banks.*
- anemonoides.* 10. *G. glabrum* ; caule unifloro, foliis pinnatis, foliolis cu-neiformibus apice dentatis, aristis rectis villosis.—*Willd. sp. pl. 2. p. 1117.*
- Dryas anemonoides.* *Pall. itin. 3. app. n. 92. nota 2.*  
*t. Ee. f. 4.*  
*Dryas pentapetala.* *Sp. pl. 717.*  
*Caryophyllata Kamtschatica.* *Lam. encycl. 1. p. 395.*  
*Icon. Egede, descrip. of Greenland, p. 42. t. 2.*  
 On the north-west coast and the Kuril Islands. *Merk.*  
 $\frac{1}{2}$ . July. v. s. in *Herb. Lambert. specimen Pallasia.*  
*num.*

419. POTENTILLA. *Gen. pl. 866.*\* *Foliis ternatis.*

1. *P. assurgens*, glabriuscula; stipulis subulatis, foliis ternatis oblongo-cuneiformibus apice tridentatis adpresso-pilosiusculis, corymbo laxo paucifloro, petalis oblongo-obovatis calyce longioribus.—*Willd. sp. pl. 2. p. 1110.*

Icon. *Ait. kew. 2. t. 9. Engl. bot. ....*

On rocks in Canada and on the highest peaks of the Allegany mountains, particularly at Grandfather-mountain, Carolina.  $\frac{1}{2}$ . June, July. *v. s. in Herb. Dickson.*; *v. v. in Hortis.* Flowers white. According to Willdenow, and the figure of *P. retusa* in the Flora Danica, which is quoted as a synonym, the flowers are yellow; but this figure, if correct, presents most certainly a different plant, as the petals are round, and only the length of the calyx.

2. *P. assurgens*, hirsuta; stipulis ovatis integerrimis, foliis ternatis, foliolis sessilibus approximatis ambitu inciso-dentatis utrinque hirsutis, pedicellis paucis terminalibus elongatis unifloris, petalis cuneato-oblongis emarginatis calyce duplo longioribus.

In Labrador. *Colmaster.*  $\frac{1}{2}$ . *v. s. in Herb. Dickson.*  
A small species; flowers large in proportion.

3. *P. assurgens*, lanato-tomentosa; stipulis ovatis integerrimis, foliis ternatis, foliolis sessilibus approximatis ambitu inciso-dentatis supra hirsutis subtus niveo-tomentosis, pedunculis paucis terminalibus unifloris, petalis lato-obcordatis calyce vix longioribus.—*Willd. sp. pl. 2. p. 1109.*

Icon. *Gmel. sib. 3. t. 36. f. 1. Gunn. norv. t. 3. f. 1.*

In Labrador. *Colmaster.*  $\frac{1}{2}$ . June, July. *v. s. in Herb. Dickson.* Flowers yellow.

4. *P. assurgens*, sericeo-villosissima; stipulis latis membranaceis integerrimis, foliis ternatis, foliolis sessilibus approximatis ambitu inciso-dentatis, supra villis nitidis adpresso-villosis, subtus cano-tomentosis, pedunculis brevibus aggregatis, petalis obcordatis calyce longioribus.

*P. villosa. Pallas in Herb. Lambert.*

On the north-west coast.  $\frac{1}{2}$ . *v. s. in Herb. Lambert.*  
This species approaches in general character very near to the foregoing one, but in habit they differ strikingly; the last is a more robust plant, covered with shin-

*tridentata.**emarginata.**nivea.**villosa.*

- ing silky hairs, and the flowers are more than double the size of the former, which is a small plant covered with a white woolly tomentum. It approaches also to *P. subacaulis* Willd. sp. pl. 2. p. 1111. according to description, but Gmelin's figure is extremely different from it.
- hirsuta.* 5. *P. erecta, simplex, hirsutissima ; stipulis lanceolatis integris, foliis ternatis obovalibus laciniato-incisis, panicula pauciflora, pedicellis brevibus, petalis calyce minoribus.* — *Mich. fl. amer. 1. p. 303.*  
In Canada and the western parts of New York. 4.  
July, Aug. v. v. Flowers white, small.
- norwegica.* 6. *P. erecta, ramosa, pubescens ; stipulis ovalibus dentatis, foliis ternatis rhombeo-lanceolatis inciso-dentatis, ramis dichotomis, pedicellis brevibus axillaribus solitariis, petalis calyce brevioribus.* — *Willd. sp. pl. 2. p. 1109.*  
Icon. *Fl. dan. 171.*  
In fields of Canada and New York. ④. June, July.  
v. v. Flowers pale yellow.
- \*\* *Foliis digitatis.*
- pumila.* 7. *P. erecta, subacaulis, pubescens ; foliis quinatis cuneiformibus apice dilatatis incisis subtus lanuginosis, pedunculis petiolo brevioribus simplicibus unifloris, petalis rotundatis calyce vix longioribus.* — *Lam. encycl. 5. p. 594.*  
In dry fields and pastures : Canada to Virginia. 4.  
May—July. v. v. It has a strong resemblance to *P. verna* ; flowers pale yellow.
- canadensis.* 8. *P. procumbens, subramosa, candidanti-sericea ; stipulis ovatis incisis, foliis quinatis cuneato-obovalibus inciso-dentatis, caule adscendente hirsuto, pedunculis solitariis elongatis, calycis laciniis linearis-lanceolatis, petalis orbiculatis subintegris longitudine calycis.* — *Willd. sp. pl. 2. p. 1106.*  
In fields and pastures, common : Canada to Carolina.  
4. May—Aug. v. v. Has the appearance of a strawberry plant ; flowers bright yellow.
- simplex.* 9. *P. erecta, simplex, hirsuta ; stipulis incisis, foliis quinatis oblongo-ovalibus grosse-serratis superioribus sessilibus, pedunculis axillaribus solitariis elongatis unifloris, calycis laciniis linearis-lanceolatis, petalis rotundato-obcordatis calyce longioribus.* — *Mich. fl. amer. 1. p. 303.*

- P. sarmentosa. *Willd. enum.* 554.  
 P. caroliniana. *Lam. encycl.* 5. p. 595.  
 In fields, meadows, and dry woods: Canada to Carolina.  
 Ʌ. May—Aug. v. v. Flowers yellow.
10. P. decumbens, filiformis, hirsuta; stipulis linearis-oblongis integris, foliis radicalibus septenatis quinatisque linearis-cuneiformibus dentatis, pedunculis axillaribus solitariis unifloris, petalis retusis calycis longitudine.—  
*Willd. sp. pl. 2. p. 1103.*  
 Icon. *Jacq. ic. rar.* 91.  
 In Labrador. Ʌ. June, July. v. s. in *Herb. Banks.*  
 Flowers yellow, small.
11. P. erecta, ramosa, glabriuscula; foliis quinatis, foliolis pinnatifidis, lacinias integris acutis, floribus terminalibus subcorymbosis.—  
*Near Hudson's Bay.* Ʌ. v. s. in *Herb. Banks.*
12. P. erecta, ramosa, niveo-tomentosa; stipulis ovatis acutis, foliis quinatis cuneatis inciso-dentatis subtus tomentosis, floribus terminalibus corymbosis, petalis retusis vix calycē longioribus.—*Willd. sp. pl. 2. p. 1101.*  
 Icon. *Fl. dan.* 865. *Engl. bot.* 89.  
 In Canada, and on rocks in New Hampshire and Vermont. Ʌ. June, July. v. v. Flowers small, yellow.
- \*\*\* *Foliis pinnatis.*
13. P. erecta, ramosissima, hirsuta; stipulis ovatis integris, foliis quinato-pinnatis, foliolis linearis-oblongis planis, petiolis longis, ramulis 1-2-floris, calycis lacinias 5: alternis linearibus, cæteris lato-ovatis acutis, petalis obovatis calycē longioribus.—*Willd. sp. pl. 2. p. 1094.*  
 Icon. *Engl. bot.* 88. *Amm. ruth. t.* 17.  
 On the banks of rivers in Canada, and on the waters of the Rocky-mountains. *Lewis.* June—Aug. v. v. in *Hortis; v. s. in Herb. Lewis.* Flowers large, yellow. This shrub is from three to five feet high.
14. P. erecta, ramosissima, hirsutissima; stipulis ovatis integris, foliis quinato-pinnatis, foliolis linearis-oblongis margine revolutis, petiolis brevibus, corymbis terminalibus dichotomis dense-multifloris, calycis lacinias subæqualibus, petalis subrotundis longitudine calycis.  
 P. fruticosa alpestris. *Pallas in Herb. Lambert.*  
 Icon. *Amm. ruth. t.* 18. f. 1.

ICOSANDRIA POLYGYNIA. *Potentilla.*

In bog meadows and on the borders of lakes : Canada and on the mountains of New York and New Jersey.  $\frac{1}{2}$ . July, Aug. v. v. This is a low-growing shrub, not above eighteen inches high. Though it appears to be no more than a variety of the preceding, the difference of habit is so great, that I was persuaded to propose it as a distinct species : both cultivated near one another, will decide whether it shall be adopted as such or not.

- Anserina.* 15. *P. repens*; foliis interrupte-pinnatis numerosis inciso-serratis sericeis subtus niveo-tomentosis, pedunculis solitariis unifloris.—*Willd. sp. pl. 2. p. 1095.*

Icon. *Engl. bot. 861.*

On the banks of rivers and lakes : Pennsylvania, New York, and Canada.  $\frac{1}{2}$ . May—Sept. v. v. Flowers yellow, large. Specimens from Labrador in the Herbarium of A. B. Lambert, Esq. seem to differ in the form of its petals from the European plant ; they are obovate, and longer than the calyx.

- pensylvanica.* 16. *P. erecta*, mollissime subcandicanti-villosa ; foliis pinnatis, foliolis oblongis obtusis subsemi-pinnatifidis tomentosis, panicula stricta multiflora, calycis tomentosi lacinii semiovalibus. *Mich. fl. amer. 1 p. 304. Willd. sp. pl. 2. p. 1099.*

Icon. *Jacq. hort. t. 189.*

$\beta$ . *P. cano-pubescent* ; foliis pectinato-dentatis margine revolutis, floribus corymbosis.

*P. strigosa. Pallas in Herb. Lambert.*

Icon. *Gmel. sib. 3. t. 34. f. 1.*

In Canada and New England.  $\beta$ . on the Missouri.  $\frac{1}{2}$ . June, July. v. v.  $\beta$ . v. s. in *Herb. Lewis.* Flowers yellow. The variety  $\beta$ . appears to be a distinct species.

- supina.* 17. *P. procumbens*, dichotoma ; foliis pinnatis, foliolis oblongis profunde dentatis, pedunculis lateralibus solitariis unifloris patulis.—*Willd. sp. pl. 2. p. 1099.*

Icon. *Gmel. it. 1. t. 27. f. 2. Pluk. alm. t. 106. f. 7.*

In overflowed places, on the banks of rivers : Canada to Pennsylvania, principally in the western parts thereof.  $\odot$ . June—Aug. v. v. Flowers small, yellow.

420. COMARUM. *Gen. pl. 869.*

- palustre.* 1. *Comarum. Willd. sp. pl. 2. p. 1119.*  
Icon. *Fl. dan. 636. Engl. bot. 172.*

In swamps : Canada and on the banks of rivers and lakes  
in the western parts of New York and Pennsylvania.  
4. June, July. v. v. Flowers dark purple.

## 421. FRAGARIA. Gen. pl. 865.

1. F. calyce fructus reflexo, pubescentia petiolorum paten-tissima, peduncularum adpressa. Willd. sp. pl. 2. p. 1090.  
Icon. Engl. bot. 1524.  
In Canada and New England. 4. April, May. v. v.  
The common *Strawberry* is frequently found wild.
2. F. calyce fructus patente, pubescentia petiolorum erecta, peduncularum adpressa, foliis supra glabriusculis. Willd. sp. pl. 2. p. 1091.  
F. glabra. Duham. arb. 1. t. 5.  
In woods and fields, frequent : Canada to Carolina. 4.  
April, May. v. v.
3. F. major; foliolis amplio-ovalibus, lateralibus manifeste petiolatis, pedicellis longis recurvo-pendulis, receptaculis seminum globos favoso-scrobiculatis villosis. Mich. fl. amer. 1. p. 299.  
In shady wet woods : Canada and on the Allegany mountains. 4. April, May. v. v.

## 422. CALYCANTHUS. Gen. pl. 870.

1. C. laciniis calycis lanceolatis, foliis lato-ovalibus acutis subtus tomentosis, ramis patentibus.—Willd. sp. pl. 2. p. 1119. Willd. enum. 559.  
C. sterilis. Walt. fl. car. 151.  
Icon. Bot. mag. 503.  
On the banks of shady rivulets : Carolina. 4. May—July. v. v. The fragrant flowers are, as all the following, dark purple. This so much esteemed shrub is generally known by the name of *Carolina Allspice* or *Sweet-scented Shrub*.
2. C. laciniis calycis lanceolatis, foliis ovato-lanceolatis acuminatis subtus glaucis pubescentibus, ramis patentibus.—Willd. enum. 559.  
C. fertilis. Walt. fl. car. 151.  
Icon. Bot. rep. 539. Schmidt arb. 129.  
In the mountains of Carolina. 4. May—Aug. v. v.

## ICOSANDRIA POLYGYNIA. Calycanthus.

The scent of this species is not quite so agreeable as the foregoing one, and more faint.

*lævigatus.*

3. C. laciniis calycis lanceolatis, foliis oblongis vel ovalibus sensim acuminatis subrugosis utrinque glabris viridibus, ramis stricte erectis.—Willd. enum. 559.

C. ferax. Mich. fl. amer. 1. p. 305.

On the mountains of Pennsylvania, Virginia, and Carolina. ♀. May—July. v. v.

END OF THE FIRST VOLUME.