


6"



Photographic Sciences Corporation

**23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503**

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1982

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

☒ Coloured covers/
Couverture de couleur

☐ Covers damaged/
Couverture endommagée

☐ Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée

☐ Cover title missing/
Le titre de couverture manque

☐ Coloured maps/
Cartes géographiques en couleur

☐ Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)

☐ Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur

☒ Bound with other material/
Relié avec d'autres documents

☐ Tight binding may cause shadows or distortion along interior margin/
La reliure serrée peut causer de l'ombre ou de la distortion le long de la marge intérieure

☐ Blank leaves added during restoration may appear within the text. Whenever possible, these have been omitted from filming/
Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.

☐ Additional comments:/
Commentaires supplémentaires:

☐ Coloured pages/
Pages de couleur

☐ Pages damaged/
Pages endommagées

☐ Pages restored and/or laminated/
Pages restaurées et/ou pelliculées

☐ Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées

☐ Pages detached/
Pages détachées

☐ Showthrough/
Transparence

☐ Quality of print varies/
Qualité inégale de l'impression

☐ Includes supplementary material/
Comprend du matériel supplémentaire

☐ Only edition available/
Seule édition disponible

☐ Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image/
Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

| | | | | | |
|--------------------------|--------------------------|--------------------------|-------------------------------------|--------------------------|--------------------------|
| 10X | 14X | 18X | 22X | 26X | 30X |
| <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input checked="" type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 12X | 16X | 20X | 24X | 28X | 32X |

The copy filmed here has been reproduced thanks to the generosity of:



Library of Congress
Photoduplication Service

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol → (meaning "CONTINUED"), or the symbol ▼ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

Library of Congress
Photoduplication Service

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole → signifie "A SUIVRE", le symbole ▼ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

ails
du
odifier
une
image

rrata
to

pelure,
on à

32X

431866

R11
M14

PAPERS BY THE STAFF

OF THE

MEDICAL FACULTY,

✓
McGILL UNIVERSITY.

1899

AORTIC DISEASE WITH ANOMALOUS SIGNS DUE TO
ABERRANT CHORDÆ TENDINEÆ.

BY

W. F. HAMILTON, M.D.,

Demonstrator of Clinical Medicine, McGill University ; Assistant Physician Royal
Victoria Hospital, Montreal.

Reprinted from the Montreal Medical Journal, July, 1899.

AORTIC DISEASE WITH ANOMALOUS SIGNS DUE TO
ABERRANT CHORDE TENDINEÆ.

BY

W. F. HAMILTON, M.D.,

Demonstrator of Clinical Medicine, McGill University; Assistant Physician, Royal
Victoria Hospital, Montreal.

The patient whose disease is about to be reported, first came under my notice in the Out-patient Department of the Royal Victoria Hospital early in the winter of 1896. He was a well-developed labourer of forty years of age. He complained of pain in the chest, which first began to trouble him after an attack of acute rheumatism in 1893. During the three years since that illness he had felt thoracic pain only occasionally, but in July of 1896 it became more severe and frequent, extending widely from beneath the sternum to the right and left and through to the back. Dizziness was sometimes felt. He was admitted to the ward on December 14th, 1896.

In his past history the following points are to be noted. He had done much heavy work. In 1880 he had a chancre, or, at least, the available history concerning an infection at that time, would lead one to conclude it was a chancre. In 1892 he was treated for a right-sided popliteal aneurism by Dr. Shepherd (the femoral artery was ligated). In 1893 he suffered, as already noted, an attack of acute rheumatism, and about eighteen months before seeking advice, at the Royal Victoria Hospital, he sustained a severe strain in wheeling a heavily loaded wheelbarrow. It will be seen that sufficient cause for disease of the circulatory system will be found in the history.

His condition when first examined was that of a well-nourished man in a state of comparative comfort and well-being, with the complaints as above described. The skin showed signs of smallpox many years before. There was slight pallor of the mucous membranes; the sclerotics were icteroid. There was no œdema or dyspnoea; the finger nails showed slight clubbing; the inguinal glands were enlarged. There was a scar on the glans penis near the corona. The right thigh showed the cicatrix of a surgical wound made for the treatment of the popliteal aneurism, already mentioned. The lungs, nervous system, abdomen, and urine gave negative results on examination. The chief interest centred about the circulatory system. The arteries were slightly sclerosed. The pulse was somewhat collapsing, of good tension, regular rhythm and volume. There was visible pulsation of the vessels of the neck: the carotids springing forward in systole. Palpation revealed a diffuse precordial pulsation with a prolonged diastolic thrill palpable on the right as far as the nipple line. The thrill was also palpable in the suprasternal notch, but its maximum intensity was about the third and fourth cartilages to the left of the sternum. The apex of the heart showed some displacement to the left, being located in the fifth inter-

space one inch outside the nipple. The transverse dulness encroached upon the left edge of the sternum. The sounds at the apex were decidedly weaker than normal; at the base one could not discover any accentuation of the second sound either pulmonary or aortic. At the apex a faint diastolic murmur and a systolic murmur were audible. In addition to these one could hear a musical murmur, diastolic in rhythm, widely propagated over the chest, having its point of maximum intensity, however, at the third left interspace. This murmur was of rather high pitch and was audible to the patient. It could be distinctly heard at a distance of from 18 to 24 inches from the chest wall, and on one occasion in a quiet room the murmur was audible at a distance of fully five feet, the patient dressed and sitting in a chair. The diagnosis of aortic and mitral regurgitation was made. Various speculations, however, were made concerning the origin of the murmur just described.

The patient was discharged, and for several months he passed from under our observation. He returned in May of 1898. About sixteen months of pretty active life with freedom from distress had been enjoyed, but in April he became very short of breath and his sleep was greatly disturbed by hideous dreams and attacks of dyspnoea. The complaints made on this occasion of admission to the hospital differed with those at first noted. Precordial pain was the prominent feature of the case at first; now he complained of sleeplessness and dyspnoea,—dyspnoea even when at rest, and sometimes amounting to orthopnoea.

Along with these allied complaints one found on examining the heart that the cardiac dulness had increased. The apex was now in the sixth space, the transverse dulness $1\frac{1}{2}$ to 2 inches greater. There was epigastric pulsation. A thrill was uncertain. The musical diastolic murmur heard so widely was of a lower pitch, yet retaining its musical quality, but was no longer audible away from the chest wall. A "to-and-fro" murmur was heard at the xiphoid cartilage.

The patient wished to go home and was discharged on the first of June. He remained in bed until October 6th, when he was finally readmitted to the hospital, where he died on the 31st of the same month. The course of the case during the summer months was marked by hæmoptysis with signs of dulness (infaret) over the right lung at the base, and subsequent hæmorrhagic effusion, gradually increasing oedema of the extremities and body, the occurrence of hæmorrhoids, a diminution in the amount of urine with albuminuria, enlargement and tenderness of the liver with ascites, further increase of the cardiac dulness, and an occasional presystolic murmur at the cardiac apex. The musical diastolic murmur was constantly present, though not so plainly heard.

The treatment was directed towards the failing compensation of the heart, and consisted in digitalis, morphine, strophanthus, calomel, stry-

ness encroached
the apex were de-
discover any ac-
tic. At the apex
audible. In ad-
stolic in rhythm,
maximum inten-
sity was of rather
could be distinctly
chest wall, and on
at a distance of
r. The diagnosis
ous speculations,
just described.
he passed from
3. About sixteen
ness had been en-
and his sleep was
of dyspnoea. The
hospital differed
prominent feature
ness and dyspnoea,—
g to orthopnoea.
on examining the
apex was now in-
es greater. There
The musical dia-
s, yet retaining its
the chest wall. A
ilage.

ged on the first of
he was finally re-
of the same month.
his was marked by
the right lung at the
y increasing oedema
orrhoids, a diminution
largement and ten-
the cardiac dulness,
the apex. The musi-
ugh not so plainly

compensation of the
anthus, calomel, stry-

nine, potassium iodide, in various combinations and according to indications. The right pleura was twice aspirated. The oedema was relieved by multiple puncture of the skin under antiseptic precautions. He died very suddenly on the evening of the 31st of October.

The diagnosis of this case finally made was aortic insufficiency, mitral insufficiency with stenosis (?), myocarditis, arterio-sclerosis, secondary nephritis, right pulmonary infarct of lower lobe, and hydrothorax.

The autopsy confirmed the diagnosis with but one exception and revealed the cause of the anomalous signs. That condition which was always doubtful in this case, viz., mitral stenosis, was not found to be present and the presystolic murmur sometimes heard as compensation was more completely lost and must evidently have been the murmur of Flint associated with aortic regurgitation. But little interest attaches to the autopsy, apart from that which describes the cardiac condition.

Cor bovinum was found, the measurement being 14.25 cm. from side to side. The chambers were greatly enlarged, the right auricular walls were muscular; the tricuspid orifice admitted four fingers; the right ventricular wall was greatly hypertrophied. The mitral orifice admitted easily the tips of four fingers. The valve showed no thickening, the only abnormal condition about it, apart from the enlargement of the mitral orifice, being found in the insertion of one of the chordae tendineae arising apparently from a papillary muscle and attached fully one-half an inch from the aortic cusp upon its ventricular surface and somewhat to the left of the median line. Springing from the left side of the ventricle another fine tendon was seen to pass towards the muscle already described, and to join it just as it passes beneath the margin of this valve cusp as shown in the photograph. (Fig. 1.) Another shorter and somewhat coarser fibrous band united this papillary muscle to the ventricular surface at a point adjacent to the first. The aortic valve was incompetent, the cusps being greatly thickened and shortened.

The anomalous signs in this case give it the interest it possesses:—

(1) It is anomalous to find *two diastolic murmurs*. We had the high-pitched musical murmur and the low-pitched characteristic diastolic murmur of aortic insufficiency.

(2) It is anomalous to find a diastolic thrill so widely palpable. According to Gibson,—“They are rarely felt except in the precordial region, more especially towards the lower part of the sternum and in the neighborhood of the apex.”

(3) A murmur audible so far from the chest-wall is rarely observed.

(4) With equal truth it may be said, further, that a musical diastolic murmur is an anomalous sign.

Concerning the cause of this quality of the murmur (4), we thought upon diverse possibilities, and chief among these was a condition of rup-

tured or distorted aortic cusp occurring in an atheromatous subject at that time when he sustained a heavy strain with an overloaded wheelbarrow. It is well known that the aortic valve segments are those injured most frequently by an over-strain, and some years ago C. Theodore Williams reported a case having a musical diastolic murmur which was subsequently shown to be due to a damaged valve-cusp, the border of which "was retroverted into the ventricle and vibrated in the regurgitant stream." Another possibility was that some fibrous band was floating in the blood stream, one end attached, or, perhaps both ends fixed. How true this last possible condition was the autopsy has shown. It would appear that this arrangement of the chordæ tendineæ is a congenital anomaly. They may find their point of valvular insertion at the edges or upon the ventricular surface of the valve, but rarely is it found so high up. Such an anomaly was without signs while the blood stream flowed in the normal direction; once regurgitation took place through the aortic orifice the stream of blood set these cords in vibration and, under the increased tension induced by a dilating ventricle and before myocardial changes became pronounced, the musical murmur was produced. Then followed the period of dilatation with muscular changes inducing loss of tone and both thrill and murmur became less pronounced.

Aberrant chordæ tendineæ, though comparatively rare, have been observed by many writers, and in some instances a diagnosis of their presence has been made during life. H. Huchard, in the *Revue de Médecine*, 1893, describes five cases, three of which were diagnosed during life. The greater number of such cases, Huchard believes, are congenital anomalies, while but a few are due to pathological changes, principally those due to atrophy of the left ventricular trabeculæ. They are very rarely found in any other chamber of the heart. Murmurs produced by them are associated with those found in the upper part of the ventricle, and are almost always systolic in rhythm and possess a musical quality.

Another point among many in cardiac cases, which the observation of this case teaches, is that relief of cardiac pain and the agonising features incident to the disease when the heart shows moderate hypertrophy, are greatly relieved when dilatation occurs, and, then, too, the other features of the case change. If you will remember, pain was the chief feature at first, when no cedema and no dyspnoea were present and the apex was in the fifth interspace; that when he came again for advice dyspnoea was great and the apex was in the sixth interspace and cedema supervened. This point was discussed by Musser in 1897.

BIBLIOGRAPHY.

- Gibson—Diseases of the heart and aorta.
 H. Huchard—Tendons aberrant du cœur, *Revue de Médecine*, 1895.
 Peacock—Transactions of The Pathological Society, London, pp. 155, Vol. vi.
 Burnay Neo—Transactions of The Clinical Society, London, pp. 155, Vol. vii.
 Duckworth—British Medical Journal, Vol. ii., 1871.
 Henry F. Lewis—The Clinical Review, December, 1898, and The Philadelphia Medical Journal, July 16, 1898.

ous subject at
rloaded wheel-
s are those in-
s ago C. Theo-
murmur which
usp, the border
in the regurgi-
ous band was
haps both ends
ne autopsy has
hordæ tendineæ
t valvular inser-
valve, but rare-
out signs while
ce regurgitation
d set these cords
y a dilating ven-
ced, the musical
dilatation with
and murmur be-

rare, have been
agnosis of their
in the *Revue de*
were diagnosed
hard believes, are
hological changes,
r trabeculæ. They
heart. Murmurs
the upper part of
thun and possess a

ch the observation
the agonising fea-
moderate hyper-
and, then, too, the
mber, pain was the
a were present and
came again for ad-
xth interspace and
usser in 1897.

ine, 1895.
on, pp. 155, Vol. vi.
n. pp. 155, Vol. vii.
s, and The Philadelphia



