

**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1997

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming are checked below.

- Coloured covers / Couverture de couleur
- Covers damaged / Couverture endommagée
- Covers restored and/or laminated / Couverture restaurée et/ou pelliculée
- Cover title missing / Le titre de couverture manque
- Coloured maps / Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black) / Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations / Planches et/ou illustrations en couleur
- Bound with other material / Relié avec d'autres documents
- Only edition available / Seule édition disponible
- Tight binding may cause shadows or distortion along interior margin / La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure.
- Blank leaves added during restorations may appear within the text. Whenever possible, these have been omitted from filming / Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- Additional comments / Commentaires supplémentaires:

Various pagings.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured pages / Pages de couleur
- Pages damaged / Pages endommagées
- Pages restored and/or laminated / Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed / Pages décolorées, tachetées ou piquées
- Pages detached / Pages détachées
- Showthrough / Transparence
- Quality of print varies / Qualité inégale de l'impression
- Includes supplementary material / Comprend du matériel supplémentaire
- Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image / Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.
- Opposing pages with varying colouration or discolourations are filmed twice to ensure the best possible image / Les pages s'opposant ayant des colorations variables ou des décolorations sont filmées deux fois afin d'obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below / Ce document est filmé au taux de réduction indiqué ci-dessous.

10x	14x	18x	22x	26x	30x	32x
12x	16x	20x	24x	28x	✓	

The copy filmed here has been reproduced thanks to the generosity of:



Seminary of Quebec
Library

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol → (meaning "CONTINUED"), or the symbol ▽ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:


Séminaire de Québec
Bibliothèque

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole → signifie "A SUIVRE", le symbole ▽ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.



174

RESULTS

or

METEOROLOGICAL OBSERVATIONS

MADE AT THE

MAGNETICAL OBSERVATORY, TORONTO,

CANADA WEST,

DURING THE YEARS 1860, 1861, & 1862.



TORONTO:

PRINTED BY W. C. CHEWETT & CO., 17 & 19 KING STREET EAST.

1864.

PRINTED AT THE STEAM PRESS ESTABLISHMENT OF W. C. CHEWETT & CO.,
17 & 19 KING STREET EAST, TORONTO.

C O N T E N T S .

	PAGE
INTRODUCTION
General monthly meteorological abstracts for 1860 to 1862 inclusive, giving the daily mean values of the meteorological elements, the daily resultant directions, resultant velocities and mean velocities of the wind, the daily extremes of temperature, with the amount and approximate duration of rain and snow for each day on which rain or snow fell.	1
TABLES.	
I.—Monthly means of the temperature of the air, at each of the six observation hours, from 1860 to 1862 inclusive..	37
II.—Monthly and annual means of the temperature of the air, furnished by six daily observations from 1860 to 1862 inclusive	38
III.—Monthly means of the temperature of the air at each of the six observation hours, for the period 1860 to 1862 inclusive	38
IV.—Monthly mean abnormal variations of temperature without regard to sign, or mean differences without regard to sign between the normal temperature of the day and hour, and the observed temperature of the same day and hour, for each month of the years 1860 to 1862.....	38
V.—Monthly mean abnormal variations of temperature, without regard to sign, at each observation hour, for the period 1860 to 1862.....	39
VI.—Half-yearly and yearly mean abnormal variations of temperatures, without regard to sign, for each of the six observation hours.....	39
VII.—Probable variability of the monthly means of the temperature at each of the six observation hours, in a single year, together with their half-yearly and yearly averages, from the years 1854 to 1862 inclusive.....	39
VIII.—Mean abnormal variations of temperature with their proper signs, arranged according to the direction of the wind at the hours of observation, for each year, and for the winter and summer half years, from 1860 to 1862; the summer being considered to begin April 1st and end September 30th	40
IX.—Mean abnormal variations of temperature, with their proper signs, arranged according to the direction of the wind at the hour of observation, in each month and in the year, for the period 1860 to 1862.....	40
X.—Shewing for the period 1853 to 1862, the number of times in each month in which the temperature at the hour of observation differed from the normal to the extent of 15° and upwards, with the average value of the corresponding deviation.	40
XI.—Relative frequency of the several winds during extraordinarily high temperatures, when the temperature at the hour of observation exceeded the normal by 15° and upwards, with the average excess of temperature for each wind, deduced from observations in the ten years 1853 to 1862	41
XII.—Relative frequency of the several winds during extraordinarily low temperatures, when the temperature at the hour of observation was below the normal by 15° and upwards, with the average defect of temperature for each wind, deduced from observations in the ten years 1853 to 1862.....	42
XIII.—Mean differences without regard to sign between the temperatures of the air at 6 A. M. on consecutive days, for each month in the years 1860 to 1862, the effect of annual variation being eliminated.....	43
XIV.—Mean changes of temperature with their proper signs, from 6 A. M. to 6 A. M. for the period 1860 to 1862, arranged according to the daily resultant direction of the wind, the effect of annual variation being eliminated.....	43
XV.—Highest and lowest temperatures in each month, for the years 1860, '61, and '62.....	43
XVI.—Monthly means of the barometric pressure at each of the six observation hours, from 1860 to 1862 inclusive.....	44
XVII.—Monthly and annual means of the barometric pressures furnished by six daily observations from 1860 to 1862.....	45
XVIII.—Monthly means of the barometric pressure at each of the six observation hours, for the period 1860 to 1862.....	45
XIX.—Monthly mean abnormal variations of the barometer without regard to sign, for each month of the years 1860 to 1862. 45	45
XX.—Monthly mean abnormal variations of the barometer without regard to sign, at each of the six observation hours, for the period 1860 to 1862.....	46

CONTENTS.

TABLES.		PAGE
XXI.—Probable variability of the monthly means of the barometer at each hour in a single year, derived from observations in the nine years 1854 to 1862.....		46
XXII.—Mean abnormal variations of the barometer with their proper signs, arranged according to the direction of the wind, for each year and for the winter and summer half years, from 1860 to 1862.....		46
XXIII.—Mean abnormal variations of the barometer, with their proper signs arranged according to the direction of the wind, in each month and in the year, for the period 1860 to 1862.....		47
XXIV.—Mean differences, without regard to sign, between the barometric pressures at 6 A. M., on consecutive days, for each month in the years 1860 to 1862.....		47
XXV.—Mean changes in the barometer, with their proper signs, from 6 A. M. to 6 A. M., for the period 1860 to 1862, arranged according to the daily resultant direction of the wind.....		47
XXVI.—Highest and lowest readings of the barometer in each month, for the years 1860, '61, and '62.....		47
XXVII.—Monthly means of the pressure of dry air at each of the six observation hours, from 1860 to 1862 inclusive.....		48
XXVIII.—Monthly and annual means of the pressure of dry air, furnished by six daily observations, from 1860 to 1862.....		49
XXIX.—Monthly means of the pressure of dry air at each of the six observation hours, for the period 1860 to 1862.....		49
XXX.—Monthly mean abnormal variations of the pressure of dry air, without regard to sign, for each month of the years 1860 to 1862.....		49
XXXI.—Monthly mean abnormal variations of the pressure of dry air, without regard to sign, at each of the six observation hours, for the period 1860 to 1862.....		50
XXXII.—Mean abnormal variations of the pressure of dry air, with their proper signs, arranged according to the direction of the wind, for each year, and for the winter and summer half years, from 1860 to 1862, the summer being considered to begin April 1st, and end September 30th.....		50
XXXIII.—Mean abnormal variations of the pressure of dry air, with their proper signs, arranged according to the direction of the wind, in each month and in the year, for the period 1860 to 1862.....		50
XXXIV.—Mean differences, without regard to sign, between the pressures of dry air at 6 A. M., on consecutive days, for each month in the years 1860 to 1862.....		51
XXXV.—Mean changes in the pressure of dry air, with their proper signs, from 6 A. M. to 6 A. M., for the period 1860 to 1862, arranged according to the daily resultant direction of the wind.....		51
XXXVI.—Highest and lowest pressures of dry air in each month, for 1860, '61, and '62.....		51
XXXVII.—Monthly means of the pressure of vapour, at each of the six observation hours, from 1860 to 1862 inclusive.....		52
XXXVIII.—Monthly and annual means of the pressure of vapour, furnished by six daily observations, from 1860 to 1862.....		53
XXXIX.—Monthly means of the pressure of vapour at each of the six observation hours, for the period 1860 to 1862.....		53
XL.—Monthly mean abnormal variations of the pressure of vapour, without regard to sign, for each month of the years 1860 to 1862.....		53
XLI.—Monthly mean abnormal variations of the pressure of vapour, without regard to sign, at each of the six observation hours, for the period 1860 to 1862.....		54
XLII.—Mean abnormal variations of the pressure of vapour, with their proper signs, arranged according to the direction of the wind, for each year, and for the winter and summer half years, from 1860 to 1862.....		54
XLIII.—Mean abnormal variations of the pressure of vapour, with their proper signs, arranged according to the direction of the wind, in each month and in the year, for the period 1860 to 1862.....		54
XLIV.—Mean differences, without regard to sign, between the pressures of vapour at 6 A. M. on consecutive days, for each month in the years 1860 to 1862.....		55
XLV.—Mean changes in the pressures of vapour, with their proper signs, from 6 A. M. to 6 A. M., for the period 1860 to 1862, arranged according to the daily resultant directions of the wind.....		55
XLVI.—Highest and lowest pressures of vapour in each month, for 1860, 1861, and 1862.....		55
XLVII.—Monthly means of the relative humidity at each of the six observation hours, from 1860 to 1862 inclusive.....		56
XLVIII.—Monthly and annual means of the relative humidity, furnished by six daily observations, from 1860 to 1862.....		57
XLIX.—Monthly means of the relative humidity, at each of the six observation hours, for the period 1860 to 1862.....		57
L.—Monthly mean abnormal variations of the relative humidity, without regard to sign, for each month of the years 1860, 1861, and 1862.....		57
L1.—Monthly mean abnormal variations of the relative humidity, without regard to sign, at each of the six observation hours, for the period 1860 to 1862.....		57

CONTENTS.

TABLES.	PAGE.
ations 46	
wind, 46	
wind, 46	
..... 47	
or each 47	
2, ar- 47	
..... 47	
..... 48	
..... 49	
..... 49	
years 49	
vation 50	
lion of 50	
idered 50	
lion 50	
or each 51	
1862, 51	
..... 51	
..... 52	
..... 53	
years 53	
vation 54	
lion of 54	
tion of 54	
or each 55	
1862, 55	
..... 55	
..... 56	
..... 57	
1860, 57	
ervation 57	
LII.—Mean abnormal variations of the relative humidity, with their proper signs, arranged according to the direction of the wind, for each year, and for the winter and summer half years, from 1860 to 1862. 58	
LIII.—Mean abnormal variations of the relative humidity, with their proper signs, arranged according to the direction of the wind, in each month, and in the year, for the period 1860 to 1862. 58	
LIV.—Mean difference, without regard to sign, between the relative humidity at 6 A. M., on consecutive days, for each month in the years 1860 to 1862. 58	
LV.—Mean changes in the relative humidity, with their proper signs, from 6 A. M. to 6 A. M., for the period 1860 to 1862, arranged according to the daily resultant direction of the wind. 59	
LVI.—Comparative view of the unusual variations of certain meteorological elements, derived from the series 1842-48, 1854-59, and 1860-62. 59	
LVII.—Comparative view of the annual means of the diurnal variations at the six observation hours, for the same three series. 59	
LVIII.—Monthly means of the extent of sky clouded at each of the six observation hours, for the years 1860 to 1862. 60	
LIX.—Monthly and annual means of the extent of sky clouded, from six daily observations, for 1860 to 1862. 61	
LX.—Monthly means of the extent of sky clouded, at each observation hour, for the period 1860 to 1862. 61	
LXI.—Mean clouded sky, arranged according to the direction of the wind at the hour of observation, in each month and in the year, for the period 1860 to 1862. 61	
LXII.—Number of hours that each wind blew in each month, and in the year, for the periods 1853-57, 1858-62, and 1853-62. 62	
LXIII.—Number of hours that each wind blew in each quarter, including December of the same year, for 1853-57, 1858-62, and 1853-62. 63	
LXIV.—Ratios shewing for each month separately the comparative duration of the different winds, being the numbers in table LXII, expressed in terms of their respective means for all winds excluding calms. 64	
LXV.—Ratios shewing for each quarter the comparative duration of the different winds, being the numbers in table LXIII, expressed in terms of their respective means for all winds excluding calms. 65	
LXVI.—Ratios for each quarter, including calms, for the period 1853-62. 65	
LXVII.—Ratio shewing the annual distribution of each wind, being the numbers in table LXIV, expressed in terms of their respective yearly arithmetic means. 66	
LXVIII.—Quarterly averages of the ratios in table LXVII. 67	
LXIX.—Absolute number of the several upper currents in each month, in the two half years, and in the year; the summer half-year being considered to extend from April to September inclusive. 68	
LXX.—Ratio shewing for each month separately, as well as for the winter, summer, and year, the relative frequency of the several upper currents, including the cases of calm sky and clear sky. 68	
LXXI.—Ratios shewing for each month separately, as well as for the two half years and year, the relative frequency of the several upper currents, excluding the cases of calm sky and clear sky. 68	
LXXXI. & LXXXII.—Comparison of the half-yearly and yearly ratios in table LXX., & LXXI., with the ratios derived in a similar manner from the observations made at 8 A. M., 2 P. M., and 4 P. M., as well as those derived from the observations at 10 P. M., midnight, and 6 A. M. 69	
LXXXIV.—Ratios shewing the annual distribution of each upper wind, and comparing the frequency of each wind in the six winter months with its frequency in the six summer months. 69	
LXXXV.—Absolute number of the several upper currents at each of the six observation hours. 70	
LXXXVI.—Ratios shewing for each hour separately, the relative frequency of the different upper currents, including the cases of calm and clear sky. 70	
LXXXVII.—Ratios shewing for each hour separately, the relative frequency of the different upper winds, excluding the cases when the sky was calm and when it was clear. 70	
LXXXVIII.—Ratios shewing the diurnal distribution of each upper wind, and comparing their frequency at the hours 2 ^h , 4 ^h , & 20 ^h , with their frequency at the hours 10 ^h , midnight, and 18 ^h 70	
LXXXIX. } Winds in the upper strata from six daily observations in the years 1860 to 1862, considered with reference to the to } simultaneous surface winds. 71	
LXXXI. } On the relative frequency of the different winds during days of rain or snow, from the hourly records of ten years. 72	
LXXXII. } LXXXVI. }	
LXXXVII.—Number of days on which rain fell, with its depth in inches, for each month of the years 1860, 1861, and 1862. 82	

CONTENTS.

TABLES.	PAGE.
LXXXVIII.—Number of days in which snow fell, with its depth in inches, for each month of the years 1860, 1861, and 1862.....	83
LXXXIX.—Mean monthly and annual temperatures in each of the twenty-three years from 1841 to 1863 inclusive, with the average monthly and annual temperatures in the whole period. Also the probable variability of a single year, and of each month in a single year, together with the probable errors of the monthly and annual averages for the whole period.....	83
XC.—Mean monthly and annual barometric pressures in each of the twenty-three years from 1841 to 1863 inclusive, with the average monthly and annual pressures in the whole period. Also the probable variability of a single year, and of each month in a single year, together with the probable errors of the monthly and annual averages for the whole period.....	84

INTRODUCTION.

THE Toronto Magnetical and Meteorological Observatory is situated in the grounds of the University of Toronto, in latitude $43^{\circ} 39' 4''$ N., longitude $5^{\circ} 17' 33''$ W., 108 feet above Lake Ontario, and approximately 342 feet above the level of the sea.

The circumstances attending the establishment of the Observatory by the British Government in 1839-'40, and its transfer to the care of the Provincial Government in 1853, have been described in the introduction to the "Results of Meteorological Observations for the years 1854-'59."

The duties of the Observatory are carried on by the Director, G. T. Kingston, M.A., assisted by Messrs. Walker, Menzies, and Stewart, formerly non-commissioned officers of the Royal Artillery, and observers under the Imperial Government, together with Mr. W. F. Davidson, an additional observer who has been connected with the establishment since the commencement of 1857.

The instruments and the system of observation during the years 1860-'62 continued the same as in the previous years. The regular observation hours for reading the barometer, standard thermometer, and wet-bulb thermometer were 6 A.M., 8 A.M., 2 P.M., 4 P.M., 10 P.M., and midnight; excepting on Sundays, Christmas Day, and Good Friday, when these instruments were read at 6 A.M. and 2 P.M. only. These latter readings, though recorded in the daily register, are not included in the hourly means of the month. The hygrometric tables employed for deducing the pressure of vapour and the relative humidity were those calculated by Professor Coffin, of Lafayette College, Pennsylvania, from the more recent experiments of Regnault. The differences between the barometric pressures and the pressures of vapour have been recorded, in conformity with ordinary usage, as the "Pressures of Dry Air."

At the regular observation hours a record was always made of the general appearance of the sky, including the form, distribution, and motions of the clouds.

The maximum self-registering thermometer was read at 6 A.M., the reading being entered for the meteorological day which terminates at that hour. The minimum self-registering thermometer was read at 2 P.M., the entry being made as belonging to the actual day.

From Robinson's anemometer a continuous record was kept of the direction of the wind during each hour of every day, (Sundays and holidays included,) and its mean velocity, or number of miles passed over by the wind during the hour, the space of each hour being always designated by the point of time with which it commenced. For each of the six observation hours, the instantaneous direction and approximate velocity at the hour were also recorded, such velocity being the distance travelled by the wind during the half-hours preceding and following the hour of observation.

INTRODUCTION.

The resultant direction and resultant velocity for any day or other group of hours were calculated from the directions and velocities in the several hours composing the group, by means of the formulæ

$$\tan \bar{\theta} = \frac{\sum (V \sin \theta)}{\sum (V \cos \theta)}; \quad V = \frac{\sum (V \cos \theta)}{n \cos \bar{\theta}};$$

where θ represents the angular distance of the point from which the wind blew during any hour, measured from the north towards the right, V the corresponding velocity, $\bar{\theta}$ and V the values of θ and V corresponding to the resultant, and n the number of hours in the group under consideration.

The depth of rain or snow recorded for any day was that which fell between 6 A.M. that commenced and 6 A.M. that terminated the day in question.

The rain-gauge in ordinary use gives only the amount of rain that falls between two successive examinations, and therefore affords no means of determining the distribution of the rain among the several hours of the day. To supply this defect, an hourly rain-gauge, of which the following is a description, was constructed and brought into use in April, 1861.

The rain is received, in the first instance, by an ordinary gauge placed on the summit of a chimney, with its orifice at a height of 19 feet above the ground, and sufficiently removed from the influence of eddy winds. The rain is thence conveyed by a pipe into the room beneath, where it is discharged into a small copper cylinder, two inches in diameter, which, by means of a band connected with a clock, is made to revolve about its axis, placed vertically, once in twenty-four hours. From the lower part of the copper cylinder a tube extends slightly inclining outwards from the horizontal, whose outer extremity, bent downwards, is directed successively into twenty-four equal compartments of a large zinc vessel, bounded by two cylindrical surfaces, whose common axis coincides with that of the small copper distributing cylinder. The diameters of the bounding cylindrical surfaces of the vessel are 32 inches and 20 inches respectively, and its height 14 inches. The compartments are separated by thin vertical partitions radiating from the common axis, and are each furnished with a small discharge pipe by which the rain may be drawn off and measured at the convenience of the observer. By this arrangement a record can be made of the depth of rain that falls within each separate hour as long as the instrument is in operation. The time of its action however is limited, to those parts of the year when severe frosts are not likely to occur, namely, from April to October, inclusive, with sometimes a portion of November.

GENERAL METEOROLOGICAL REGISTER.

In the monthly abstracts on pp. 1 to 36, the numbers entered in the six columns headed "Daily Means," are the daily averages of the observations made at the six observation hours, and are uncorrected for diurnal variation. The resultant directions and velocities and the mean velocities of the wind are derived from the directions and velocities during each of the twenty-four hours. The numbers at the bases of the columns of daily means, as well as those of the mean velocities of the wind and the extremes of temperature, are the averages of the numbers under which they stand, Sundays and other holidays being excluded only in the first six columns. Under the columns containing the daily resultant directions and velocities of the wind, are placed the monthly resultant direction and resultant velocity, and under the columns for the rain and snow, are entered the sums of the numbers which these columns respectively contain. In the rain or snow columns the occurrence of a star (*) indicates that the amount was too small for measurement, or the duration less than half an hour. In the column for rain and melted snow combined, ten inches of snow are reckoned as equivalent to one inch of rain.

REMARKS ON THE TABLES.

TEMPERATURE TABLES.

The normals referred to in the temperature tables (Tables I. to XV.) are the normal temperatures proper to Toronto in its actual circumstances. They have been deduced from the table of twenty-four-hour daily means (*a*) given by General Sabine in his paper on the periodic and non-periodic variations of temperature at Toronto, by applying the diurnal variations (*b*) contained (*though with contrary signs*,) in the same paper. The normals, thus computed, are used as standards of reference, the abnormal variations being entered in the daily register side by side with the several observed temperatures.

Table I. gives the monthly means of the temperature of the air at each of the six observation hours in the three years, 1860, 1861, and 1862. The numbers in the final column of each month are the means of those in the six preceding columns, uncorrected for diurnal variation.

The final columns of the several monthly parts of Table I. are exhibited in one view in Table II., which gives the monthly and annual means of temperature furnished by six daily observations for each of the three years, together with the averages of the monthly and annual means for the three years. For the purpose of comparison, the normal monthly means derived from the same six observation hours are also introduced, together with the monthly and annual means for the period 1854 to 1859.

The numbers at the foot of each of the monthly parts of Table I. are collected in Table III. which shews, on the average of the three years, 1860 to 1862, the monthly means of temperature at each of the six observation hours.

From Table IV. it appears, from the observations of nine years, that the observed temperature is in excess or defect from the normal proper to the day and hour by a difference whose mean value is $6^{\circ}.5$. This abnormal digression has an annual period, being greater in the cold and less in the warm months, its value in February being about twice as great as in August. The quarterly averages in the two series are as follows:

	Winter.	Spring.	Summer.	Autumn.
1860-62	8.5	6.0	4.6	5.5
1854-59	9.4	6.2	5.2	5.9
1854-62	9.1	6.1	5.0	5.8

In Table V. a comparison is made for each month between the abnormal digressions of temperature, without regard to sign, at the six observation hours.

(a) Philosophical Transactions for 1853, pp. 154, 159.

(b) Philosophical Transactions for 1853, pp. 145, 146.

Having regard to the annual means alone, there appears in both series to be but little difference between the abnormal digressions at the different hours, the greatest difference being only $0^{\circ}.4$ in the earlier and $0^{\circ}.3$ in the later series; an excess as compared with other hours occurring, in both series, at 6 A.M. The observations of the years 1860-62 corroborate the statement made in the preceding volume respecting an annual period in the position of the hours of greater abnormal digression. The variations at 10^h , 12^h , 18^h , and 20^h , in the six winter months, (October to March) are found in most instances to be greater, number for number, than at 2^h and 4^h ; while in the six summer months this relation is reversed. From the comparison of the half-yearly means at each hour and for the two groups of hours given in Table VI., there appears in both series to be in winter a greater uncertainty respecting the temperature during the night and morning than during the hours of the day, whereas in summer the warmer hours are more subject to irregularity. In other words, the warm hours are most liable to disturbances of temperature in the warm months, and the cold hours in the cold months; the differences between the average extent of a disturbance (if it be so called) in the two groups of hours being about $0^{\circ}.8$ in both half-years.

It has been seen in the two preceding tables that the average abnormal digressions of the observed temperatures, considered singly, are systematically greater, for a given month, at certain hours, and that the hours of greater digression have themselves an annual periodic movement. This circumstance suggests the enquiry whether the digressions of the monthly means in *single years* from the means derived from a series of years are affected by changes of a like periodic character. With a view to this question Table VII. has been formed, giving the probable variability of the monthly means for each month, and for each of the six observation hours. The number expressing the probable variability for any month and hour was computed from the squares of the differences between the means for that month and hour in each single year, from 1854 to 1862, and the mean for the same month and hour on the average of the nine years. The three final columns contain the half-yearly and yearly averages of the variabilities for each of the six hours, and the three lowest lines give the monthly, half-yearly, and yearly averages of the variabilities for 2^h and 4^h , for 10^h to 20^h , and for the six hours collectively.

In the diurnal period the progression is regular on the average of the year, as well as in each half-year. For the whole year 2 p.m. is the most variable hour with respect to its monthly means, and 10 p.m. the most regular hour. In the summer half-year 2 p.m. is also the most variable hour, but the most regular hour is midnight, while in the winter half-year the most variable hour is 6 A.M. and the most regular hour 4 p.m.

With respect to the annual period the average variability in the six winter months is about 50 per cent. greater than in the summer months; the average variability also at 2^h and 4^h , namely, at the warmer hours, is greater in each of the six summer months than the average variability at the hours 10^h , 12^h , 18^h , and 20^h . The reverse of this takes place in January, February, and March, but not in October, November, and December, when the variability is very slightly in excess at the warmer hours. But comparing the winter and summer averages for the above named combinations of hours, it is found, as in the case of the abnormal variations, that in the warm months the temperature is most variable at the warm hours, and that in the cold months the temperature at the colder hours is most variable.

The resemblance, with respect to the relation just described, between the abnormal variations of single

observations and the probable variability of monthly means in a single year, is best seen by placing the results together as follows:

	Mean Abnormal Variation.				Probable Variability.			
	$\frac{h}{2}$	$\frac{h}{4}$	$\frac{h}{10}$	$\dots 20$	$\frac{h}{2}$	$\frac{h}{4}$	$\frac{h}{10}$	$\dots 20$
Winter	7.22	<	7.99		2.41	<	2.69	
Summer.....	5.84	>	5.01		1.99	>	1.48	
Year	6.50	>	6.50		2.20	>	2.08	

In Tables VIII. and IX. the abnormal variations of temperature, with their proper signs, are arranged according to the eight principal points of the direction of the wind. In Table VIII. the half-yearly and yearly means are given for the three years separately as well as collectively. It is seen that with a N.E. wind and a west wind the temperature was above the normal in some half-years and below it in others, without reference to the season; with east and S.E. winds the temperature was above the normal in each winter and below the normal in each summer, and with winds from S. and S.W. the temperature was above the normal in each separate half-year.

Table IX. gives the monthly and yearly means of the abnormal digressions of temperature for the several winds, on the average of the three years, together with the yearly means from 1853 to 1859. The variations which accompany the N.E. and west winds have different signs in the two series. This contrariety of sign is partly occasioned by the proximity of these points to the line, which, in the earlier series, was found to separate the relatively warm from the relatively cold winds. Another cause of this disagreement is the fact that many winds which were considered, in the later series, as belonging to the N.E. group, blew from points that were included in the N.N.E. and E.N.E. groups in the earlier series. A similar remark is applicable to the west winds.

In Table X. a classification has been made of the larger deviations of temperature, including all those cases during the ten years, 1853 to 1862, wherein the observed temperature differed to the extent of 15° and upwards from the normal proper to the day and hour. It will be seen, taking one month with another, that the excessively low temperatures are more than twice as numerous, as well as of greater average deviation, than those which are relatively high.

The general excess in the number of low temperatures is due to their large preponderance in the winter months; for in summer and autumn the high temperatures exceed in number the low temperatures. The quarterly and annual numbers, and average extent of the extraordinary deviations, of both signs, as well as when considered without regard to sign, are as follows:

	Winter.	Spring.	Summer.	Autumn.	Year.
High temperatures—Number.....	176	91	68	110	445
Average deviation... 18.2	17.6	16.7	18.9	18.0	
Low temperatures—Number.....	645	184	29	52	910
Average deviation... 21.9	19.6	17.0	17.7	21.0	
Both signs—Number	821	275	97	162	1355
Average deviation... 21.1	18.9	16.8	18.5	20.0	

INTRODUCTION.

In Table XI. the temperatures that exceeded the normal to the extent of 15° and upwards during the ten years, 1853 to 1862, are classified according to the simultaneous direction of the wind.

In the four columns marked (1) are entered, for each of the four seasons, the absolute number of times that each wind accompanied an excessively high temperature. Now as some winds, other things being the same, are more numerous than others, it is necessary that the numbers in (1) should be divided by numbers proportional to the relative frequency of the several winds, without reference to temperature, during the same period of ten years. The requisite divisors, derived from Table LXVI.,* are contained in the four columns marked (2). The quotients arising from the division of the numbers in (1) by those in (2), and which are entered in (3), are expressed severally in terms of their respective means for all winds in column (4).

The cases embraced in the ten years are not sufficiently numerous to yield very decided results; but as far as they go, they shew that with excessively high temperatures the south and S.S.E. winds are most frequent in winter, the S.S.W. and S.W. winds in spring and autumn, while in summer the west and W.S.W. winds most commonly accompany the excessively high temperatures.

Table XII. shews the results similarly obtained for the excessively low temperatures.

In Table XIII. are given, for each month and for each of the three years, the mean changes in the temperature, without regard to sign, that take place in 24 hours, between 6 A.M. and 6 A.M. on consecutive days. Taking the three years together, the average change of temperature in 24 hours is $6^{\circ}.1$, the greatest change ($9^{\circ}.6$) occurring in February, and the least ($4^{\circ}.1$) in July. The quarterly means for the nine years, 1854 to 1862, are $9^{\circ}.13$ in winter, $5^{\circ}.27$ in spring, $4^{\circ}.07$ in summer, and $6^{\circ}.50$ in autumn; the general annual mean for the nine years being $6^{\circ}.25$.

In Table XIV. the mean diurnal changes of temperature, with their proper signs, are arranged according to the resultant direction of the wind during the day in which the change took place.

It is seen that with a resultant wind from N.W. or W., the temperature is lowered in every month, and that with a resultant wind from any other point the temperature is raised on the average of the year as well as in most months separately. The exceptions may be traced to the circumstance that the resultants in certain directions are not sufficiently numerous in some months to allow the effect of accidental anomalies to be overpowered and rendered inappreciable.

In Table XV. the highest and lowest temperatures that occurred in each month are given for each year separately, as well as for the average of the three years, 1860 to 1862, accompanied by the corresponding averages for the six previous years, 1854 to 1859.

* TABLE LXVI.

Relative frequency of the several winds, including calms, in each of the four quarters, for the period 1853-'62—(See pages xvi. and 65.)

	N.	N.N.E.	N.E.	E.N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.S.W.	S.W.	W.S.W.	W.	W.N.W.	N.W.	N.N.W.	Calm.
WINTER.....	1.18	0.88	0.87	0.75	0.85	0.40	0.23	0.21	0.24	0.71	1.46	2.29	1.97	1.31	1.23	1.40	0.99
SPRING.....	1.03	0.63	0.76	1.45	1.73	0.75	0.47	0.42	0.66	1.01	0.88	0.79	1.13	1.45	1.54	1.51	0.70
SUMMER.....	1.01	0.67	0.62	0.98	1.34	0.78	0.58	0.72	1.29	1.53	0.97	0.57	0.79	1.06	1.41	1.57	1.11
AUTUMN.....	1.07	0.77	0.77	1.02	1.13	0.56	0.42	0.48	0.74	1.14	1.15	1.23	1.26	1.37	1.21	1.25	1.43

INTRODUCTION.

BAROMETRIC TABLES.

The means or approximate normals employed as standards of comparison in the barometric tables (Tables XVI. to XXVI.) are the hourly means in each month, derived from the observations of eighteen years.

Table XVI. gives the monthly means of the barometer corrected to temperature 32° Faht., at each of the six observation hours, in the years 1860, '61, and '62. The final columns of each monthly part of this table, which contain the means of the six preceding columns, are exhibited in one view by Table XVII.; while in Table XVIII. are collected the hourly means at the foot of each monthly part of Table XVI.

Table XIX. shews the monthly averages of the extent of the barometric abnormal oscillations. The numbers whose averages are given in this table are the differences, without regard to sign, between the observed height of the barometer and the assumed normal proper to the month and hour.

The annual distribution is in general accordance with that given in the comparative table for the period 1854 to 1859, the quarterly means for the two series being as follows :

	Spring.	Summer.	Autumn.	Winter.
1860-62191	.128	.175	.234
1854-59190	.122	.191	.230
1854-62190	.124	.186	.231

It thence appears that the extent of the barometric oscillations in winter is nearly twice as great as in summer, and that in the spring and autumn the oscillations are nearly equal.

In Table XX. are exhibited the mean abnormal variations of the barometric pressure in the different hours. On the average of the year, although the range in the extent of the abnormal digressions is small the progression is regular for the three years, 1860-'62, as well as for the six preceding years; the most disturbed hour being 8 A.M. and the most tranquil hour 10 P.M. A somewhat similar conclusion results from Table XXI., in which is given, for each month and hour, the probable variability of an hourly mean in a single year, computed from the squares of the differences between the hourly means of any month in each year, from 1854 to 1862, and the average for the same hour and month derived from nine years. By comparing the quarterly and yearly means of the numbers in Table XXI. with those of the abnormal variations for the same nine years, it is seen that the progression, as respects the hours of greater and less barometric disturbance, unlike that which has been noticed with regard to the disturbances of temperature, is not subject to any well marked change of character in the different seasons. The quarterly and yearly means are contained in the annexed table.

MEAN ABNORMAL VARIATIONS.					MEANS OF PROBABLE VARIABILITIES.						
N.N.W.	Calm.	QUARTERS.			Year.	QUARTERS.			Year.		
		Winter.	Spring.	Summer.		Winter.	Spring.	Summer.			
1.40	0.99	.236	.194	.127	.188	.186	.0382	.0416	.0311	.0394	.0376
1.51	0.79	.232	.189	.122	.182	.181	.0366	.0409	.0299	.0381	.0364
1.57	1.11	.226	.182	.115	.181	.176	.0318	.0391	.0294	.0389	.0355
1.25	1.43	.226	.184	.116	.184	.177	.0348	.0396	.0294	.0401	.0360
		.233	.194	.131	.189	.187	.0408	.0426	.0337	.0407	.0395
		.235	.199	.130	.192	.189	.0415	.0432	.0332	.0402	.0395

Tables XXII. and XXIII. give the mean abnormal variations of the barometer that accompany winds from the eight principal points.

From comparing the yearly means derived from the three years, 1860-'62, with those for the seven years, 1853-'59, it appears that the general character of the two results are the same, and the means nearly identical in some cases. In both series the barometer is above the normal when the wind is from N., N.E., E., S.E., and S., and below the normal when the wind is from S.W., W., and N.W.; the greatest elevation accompanying a north wind and the greatest depression occurring with a S.W. wind. The half-yearly means shew, as far as the observations of three years are competent to shew it, that the south wind accompanies a low barometer in winter and a high barometer in summer, and that the reverse is the case with the N.W. wind. In judging of the resemblance between the earlier and later series, it is to be remembered that in the more recent series, many winds are included among the eight principal points, which, for the series 1853-'59, were reckoned as belonging to the intermediate points.

Tables XXIV. and XXV. are derived from the differences between the corrected readings of the barometer at 6 A.M. on consecutive days, Sundays and other holidays being included. The results in Table XXIV. are obtained by dividing the aggregate change, without regard to sign, in each month by the number of changes, *i. e.* by the number of days in the month.

The correspondence in the two series is on the whole tolerably close, the maximum occurring in either January or February, and the minimum in either July or August. Combining the two series, the greatest monthly average change in twenty-four hours is 0.281 in January, and the least 0.121 in July, the annual mean being 0.198. The quarterly means are as follows :

	Winter.	Spring.	Summer.	Autumn.
1860 to 1862	0.273	0.197	0.137	0.192
1854 to 1859	0.269	0.211	0.121	0.189
1854 to 1862	0.270	0.206	0.126	0.190

In Table XXV. are given the quotients arising from the division of the algebraical sum of the changes that accompanied each resultant wind by the number of the resultant winds in the same month and direction. The numbers in the final column are derived from the sums and numbers of the changes corresponding to each wind, and will not therefore be generally equal to the arithmetic means of the numbers in the monthly columns. From both series it is seen that the barometer rises on the average of the year when the resultant wind is from N., N.W., and W., and that it falls with a resultant wind from other quarters. It is also found, both from Table XXV. and the analogous table in the volume for 1854-59, that these statements hold good in nearly every month taken separately. A comparison of the signs of Tables XXV. and XXIII. corresponding to the several winds, brings out the fact that the same winds that accompany a relatively *high* barometer are for the most part those that, as resultants, accompany a fall; and that the winds that correspond to a low barometer commonly accompany a rise.

TABLES RELATIVE TO THE PRESSURE OF DRY AIR—(TABLES XXVII. TO XXXVI).

The approximate normals of reference for the pressure of dry air are the hourly means in each month on the average of eighteen years.

Table XXVII. gives the monthly means of the pressure of dry air at each of the six observation hours in the years 1860, 1861, and 1862. The final columns of each monthly part are collected in Table XXVIII., and the lowest horizontal lines in Table XXIX.

Tables XXX. and XXXI. give, for each month and for each of the six observation hours, the mean abnormal variations of the pressure of dry air.

The following are the quarterly means of the abnormal variations of the pressure of dry air, together with those of the barometric pressure; both being derived from the same three years:

	Spring.	Summer.	Autumn.	Winter.	Year.
Dry Air,	0.215	0.184	0.211	0.258	0.217
Barometer,	0.191	0.128	0.175	0.233	0.183

The maxima and minima are as follows:

	Maxima.	Minima.
Dry Air,	0.282 in December;	0.167 in July;
Barometer,	0.257 in December;	0.144 in August.

From Table XXXI. a faint trace of a diurnal period is observable, better marked in winter than in summer, the mean digression on the average of the year being slightly less at 10 P.M. than at 2 P.M. and the morning hours. The following are the half-yearly and yearly means of the abnormal variations, without regard to sign, at the different hours.

	2	4	10	12	18	20
Winter	0.241	0.237	0.237	0.237	0.241	0.242
Summer	0.198	0.197	0.188	0.190	0.198	0.196
Year	0.220	0.217	0.213	0.213	0.219	0.219

In Tables XXXII. and XXXIII. the abnormal variations, with their proper signs, are arranged according to the simultaneous direction of the wind.

From Table XXXII. it is seen that the pressure of dry air, as in the case of the barometer, is above the normal with the wind from N. and N.E. in both half-years, and below the normal with a wind from S.W. or W.; it is also above the normal in summer and below it in winter with a south wind. Unlike the total barometric pressure however, the pressure of dry air is above the normal in both seasons with a N.W. wind; and while in both seasons, with S.E. and E. winds, the barometer is above the normal, the pressure of dry air, though above the normal on the average of the year, is below the normal in winter. The annual means have the same signs as those for the total pressure, excepting that the pressure of dry air with a N.W. wind is decidedly above the normal, and with a south wind slightly below it.

In Table XXXIV. are exhibited the monthly and annual means of the changes in the pressure of dry air that take place in the 24 hours, commencing and ending with 6 A.M. The average changes in the four quarters and year, together with the corresponding averages of the total pressure, are given herewith:

	Spring.	Summer.	Autumn.	Winter.	Year.
Dry Air,	0.228	0.207	0.237	0.302	0.243
Barometer,	0.197	0.137	0.192	0.273	0.199

Hence while the average change in the pressure of dry air, in the different seasons, follows the same order of progression as that of the barometer, its changes are considerably greater. The range, or ratio of the winter to the summer change, is greater for the barometer than for the dry air, the ratios being respectively 1.99 and 1.46.

From Table XXXV., in which the average changes of the pressure of dry air in twenty-four hours, with their proper signs, are arranged according to the daily resultant direction of the wind, we find, on the average of the year, that the pressure increases in twenty-four hours with a resultant wind from N., N.W., and W., and decreases with a resultant wind from any other quarter. It appears further, that with eight exceptions only out of the ninety-six cases, this holds good for each separate month. The greatest rise on the average of the year occurs with a N.W. resultant, and the greatest fall with one from S.E., the range amounting to 0.491.

TABLES RELATIVE TO THE PRESSURE OF VAPOUR—(XXXVII. TO XLVI.)

The approximate normals referred to in the tables that follow are the monthly means at the six observation hours, derived from the records of eighteen years.

In Table XXXVII. are given the monthly means of the pressure of vapour at each observation hour for each of the three years, 1860, 1861, and 1862. The monthly means, on the average of the six hours, are given for each year in Table XXXVIII., and the monthly means at each hour on the average of the three years in Table XXXIX.

Table XL. exhibits the means of the abnormal variations of the pressure of vapour, without regard to sign, for each month in each of the years, 1860, '61, and '62, as well as on the average of the same three years. The greatest monthly mean digression, 0.099, is in August, and the least, 0.040, occurs both in January and in March. The transition from month to month is not perfectly regular, but it follows in the main the annual variation of the pressure of vapour. The general similarity in the annual fluctuations of the average amount of the pressure of vapour, and of its abnormal variations, will be seen by the annexed table, from which also it appears that the irregular variation averages about one-fourth of the whole pressure of vapour.

	Spring.	Summer.	Autumn.	Winter.	Year.
Pressure of Vapour.....	0.200	0.441	0.285	0.119	0.261
Abnormal Variations	0.055	0.090	0.071	0.043	0.065

It may be noticed further, that the digressions in the amount of vapour, *as compared with the whole amount*, are least in summer when the whole amount is greatest, and greatest in winter when the whole amount is least.

Table XLI. shews the monthly and annual means of the abnormal variation of the pressure of vapour at each observation hour for the period 1860 to 1862. On the average of the year the abnormal digression is greatest at 2 p.m. and least at 6 a.m., which are respectively the hours of greatest and least pressure of vapour among the six hours of observation.

The mean extent of the abnormal variations at 2^h p.m. and 4^h p.m. will be found to be greater than at the other hours in every month but January and February, and in every quarter but the winter. By comparing the annual means of the abnormal variations at the six different hours with the annual means of the pressure of vapour at the same hours, as shewn together in the annexed table, it appears that the former are nearly one-fourth of the latter.

	2	4	10	12	18	20
Pressure of Vapour.....	0.280	0.275	0.256	0.250	0.247	0.261
Abnormal Variations	0.071	0.070	0.064	0.063	0.061	0.062

four hours, we find, on
and from N.,
that with
the greatest
from S.E.,

s at the six

ration hour
e six hours,
rage of the

regard to
same three
urs both in
follows in
fluctuations
een by the
urth of tho

h the whole
n the whole

e of vapour
nal digres-
st pressure

reater than
inter. By
ual means
rs that the

In Tables XLII. and XLIII. the mean abnormal variations of the pressure of vapour, with their proper signs, are arranged according to the simultaneous direction of the wind.

From Table XLII. it appears that the pressure of vapour is below the normal with a wind from N., N.W., and W., on the average of the year and for both half-years, and that this is true for each year taken singly. With a wind from any other quarter the pressure of vapour is above the normal on the average of the three years, as well as on the average of the three winters; this, excepting as regards the N.E. wind in 1862, being also true for each separate year. With winds from N.E., E., and S.E., the pressure of vapour during the summer half-year is sometimes above and sometimes below the normal. On the average of the year the pressure is greatest with a wind from S.W., and least with a wind from N.W. In both seasons the least vapour occurs with a wind from N.W., but in winter the largest amount of vapour is with a wind from the S.E., a quarter which in summer is accompanied by an amount of vapour below the average.

From Table XLIII. wherein the mean differences above or below the normal with different winds are given for every month, it is seen that the pressure of vapour is below the normal with a wind from N., N.W., and W., in every month, with one exception for the north wind and one exception for west. With winds from other points the pressure is above the normal in most months, seventeen exceptions occurring among the sixty monthly means for the said five winds.

The mean changes in the pressure of vapour between 6 A.M. and 6 P.M., on consecutive days, and without regard to sign, are given for each month and year in Table XLIV. The average change for the year is 0.060, the greatest change being 0.093 in September, and the least 0.038 in January. The quarterly means are :

Spring.	Summer.	Autumn.	Winter.
0.044	0.087	0.067	0.042

It may be seen from Table XLV., which gives the mean changes, with their proper signs arranged according to the resultant direction of the wind, that on the average of the year, as well as in each separate month, with one exception only out of the thirty-six cases, the vapour diminishes in twenty-four hours with a resultant wind from N., N.W., and N. With a resultant wind from any other quarter the vapour increases on the average of the year as well as in each separate month, with nine exceptions out of the sixty cases. The greatest increase +0.085 is with a S.E. resultant, and the greatest decrease -0.044 with a resultant from N.W.

RELATIVE HUMIDITY—(TABLES XLVII. TO LV.)

The approximate normals with which the humidity at observation is compared, are the monthly means at each observation hour derived from the records of eighteen years.

In Table XLVII. the monthly means are given for each hour and for each of the three years. The six-hour means for each of the three years are shewn in Table XLVIII., and the three-year means at each observation hour in Table XLIX.

The abnormal variations without regard to sign, as given in Table L., are greatest in June, on the average of the three years, and least in December and January. The quarterly averages given in the annexed table, whercin saturation is expressed by 100, shew however that the greatest irregularity occurs in spring and the least in winter.

Spring.	Summer.	Autumn.	Winter.	Year.
11.0	10.0	8.3	8.2	9.4

In the diurnal period the abnormal digressions, as given in Table LI., are greatest on the average of the year at 4 p.m. and least at 6 A.M. They are greater at 2 p.m. and 4 p.m., taken together, than at the other four hours, not only on the average of the year, but in each month, the monthly and quarterly averages being as follows :

	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
$\frac{h}{2}$ and $\frac{h}{4}$	10.0	11.0	12.8	14.0	13.0	13.9	11.5	11.0	10.8	10.4	10.2	10.2
$\frac{h}{10}, \frac{h}{12}, \frac{h}{18}, \frac{h}{20}, \dots$	7.0	7.5	8.5	9.7	11.1	11.5	8.4	6.8	7.1	7.3	7.1	6.8
	Spring.		Summer.		Autumn.		Winter.		Year.			
$\frac{h}{2}$ and $\frac{h}{4}$	13.3		12.2		10.5		10.5		11.6			
$\frac{h}{10}, \frac{h}{12}, \frac{h}{18}, \frac{h}{20}, \dots$	9.8		8.9		7.2		7.1		8.2			

In Tables LIII. and LIII. the mean abnormal variations, with their proper signs, are arranged according to the direction of the wind. In Table LIII., where the mean results are given for each half-year and each year, it sometimes occurs* that with half-yearly means of opposite signs, the yearly mean has the same sign as the half-year, of which the digression is numerically the smaller of the two. This is occasioned by the excess in the *number* of winds in the half-year for which the mean digression is numerically less, and by the fact that the annual mean is obtained by dividing the algebraical sum of all the abnormal variations accompanying the wind in question by the number of those winds. A similar explanation is applicable wherever cases of such apparent contradiction are found.

With winds from N.E., E., and S.E., the air is relatively damp, and with winds from W., N.W., and N., the air is relatively dry. The most damp wind is from E. and the most dry wind from N.W.; but the range is small, amounting only to 9.0. The dampness of the E. wind is greater in winter than in summer, and the dryness of the N.W. wind is greater in summer than in winter, but the range between these two winds remains nearly the same in both seasons.

On referring to table LIII. it is seen that the N.W. wind is relatively dry in every month, and that the east wind, with one exception, is relatively damp in every month. Winds from N., N.W., and W., are relatively dry in every month with six exceptions out of the thirty-six cases; winds from N.E., E., S.E., and S.W. are relatively damp, with ten exceptions out of forty-eight cases, while the S. winds, which are dry in the summer half-year and damp in winter, are dry in four of the six summer months and damp in four of the six winter months.

The average change in relative humidity in twenty-four hours, without regard to sign, is shewn by Table LIV. to be 8.9; the greatest monthly mean being 12.4 in June, and the least 7.3 in February. The quarterly means are 10.2 in spring, 9.6 in summer, 7.8 in autumn, and 7.7 in winter.

According to Table LV. the humidity on the average of the year increases in twenty-four hours when accompanied by a resultant wind from N.E., E., S.E., and S., and diminishes with a resultant from S.W., W., N.W., and N.; the greatest increase of humidity being with a resultant wind from E., and the greatest diminution with one from N.W. The range between the E. and N.W. winds however, is only 6.1.

It is just to remark with reference to Tables L. to LV. that the observations of only three years are materials too scanty to justify our regarding as conclusive the results that relate to the abnormal variations and diurnal changes of the relative humidity.

average of
than at the
quarterly

Dec.
10.2
6.8

arranged
each half-
yearly mean
two. This
is numeri-
cal of all the
similar expla-

N.W., and
N.W.; but
ter than in
ge between

h, and that
, and W.,
N.E., E.,
winds, which
s and damp

shewn by
uary. The

hours when
from S.W.,
the greatest
6.1.

years are
urnal varia-

EXTENT OF SKY CLOUDED—(TABLES LVIII. TO LXI.)

In Table LVIII. the monthly and annual means of the extent of sky clouded (the hemisphere being unity,) are given for each of the six observation hours, and for each of the three years. Table LIX. contains the final columns of the several monthly parts of Table LVIII., and Table LX. the lower lines of the same monthly parts. From Table LIX. we find, on the average of the last three, as well as on that of the preceding six years, that December and August are the most cloudy and the least cloudy months respectively. The mean amounts of cloudiness in each quarter are proportional to the following numbers, the annual mean being unity :

	Winter.	Spring.	Summer.	Autumn.
1854-'59	1.22	0.97	0.81	1.94
1860-'62	1.19	0.92	0.82	1.05

Table LX. shews that 2 P.M. is the most cloudy hour on the average of the year, in both series, and that 10 P.M., or midnight, are the hours most free from clouds. If the monthly means at the several hours given in this table, as well as those from the corresponding table in the preceding volume, be combined in quarterly averages, it is found that in every quarter 10 P.M. and midnight are nearly equally cloudy, and that they are less cloudy than other hours. In both series the amount of cloud at 2 P.M. in winter, is equalled or exceeded by that at 6 A.M. and 8 A.M.; but, excepting in the summer quarter for 1860-'62, when 2 P.M. is very decidedly in excess, the difference between the hours 6 A.M., 8 A.M., 2 P.M., and 4 P.M., with respect to the amount of cloud, never exceeds .04. The diurnal ranges in the four quarters are as follows :

	Winter.	Spring.	Summer.	Autumn.
1854-'59	11	12	16	12
1860-'62	10	11	18	8

The extent of sky clouded under different surface winds is shewn in Table LXI. On the average of the whole year it appears from both series that there are two maxima; the principal one between E. and N.E., and the second at or about S.W. There are also two minima; one at or about S., and the other between N. and N.W. The two minima are nearly equal in the earlier series, but in the last three years the lowest minimum is decidedly at N.W. As the number of times any given wind blows is not the same in different months, neither the annual nor quarterly means under different winds can be deduced with strictness from the numbers in this table. But if, in order to form an approximate estimate of the general character of the progression in the amount of cloud from point to point of the compass that occurs in the different seasons, the monthly means corresponding to the different winds be grouped in quarterly averages, it is found that the double progression disappears in the winter and autumn, the minimum in both of these seasons being at N.W., and the winter maximum at or to the south of E. In spring and summer the progression is double, the principal maximum, as before, is between E. and N.E.; but in spring the principal minimum is between S. and S.E., while in summer the principal minimum is decidedly at N.W.

ON THE COMPARATIVE DURATION OF THE SURFACE WINDS FROM THE SIXTEEN PRINCIPAL POINTS, AND
THE ANNUAL DISTRIBUTION OF EACH SEPARATE WIND WITH RESPECT TO ITS DURATION—(TABLES
LXII. TO LXVIII.).

These tables have been constructed in a manner similar to that employed for the years 1853 to 1859, in the Toronto Meteorological Results already published. The three years, 1860-'62, have been incorpo-

rated with the preceding seven years, and in order to shew the extent of the correspondence in the results of different years, in ten years, 1853-'62, have been discussed not only collectively but in two equal groups. In Table LXII. are given the absolute duration in hours of each of the sixteen winds and the calms, in each month and in the year, for the two partial periods as well as for the ten years collectively, the numbers in this table being derived from hourly records by Robinson's anemometer. The absolute durations for each quarter are given in Table LXIII. The numbers in Table LXIV. are the quotients arising from the division of the numbers in Table LXII. by the monthly and annual arithmetic means, ~~excluding the calms~~; Table LXV. being derived in a like manner from Table LXIII.

According to the results furnished by the ten years, winds from S.S.W. through W. to N. are above the average of the sixteen winds on the average of the year; but the N.W. and N.N.W. winds alone are above the average in each separate month. The E. and E.N.E. winds are above the average on the whole year, and the E. is above the average in every month but December and January. The N. wind is above the average in every month but March, June, and November. The S. wind is below the average on the average of the year, as well as in every month but June, July, August, and September. The principal maximum on the whole year is at N.N.W., and the principal minimum at S.E.; there is also a second maximum at E., and a second minimum about N.N.E. to N.E. The positions of the maxima and minima among the points of the compass do not remain permanently fixed through the year. Referring to Table LXV. wherein the nature of their movements will be best seen, we find that in winter the progression is single, the maximum being at W.S.W. and the minimum between S.E. and S. In spring the maximum of the winter has moved northward to N.W., but it is equalled or surpassed by the eastern maximum which has reappeared together with the second minimum at N.N.E. In summer the western maximum is broken, as it were, into two almost equal maxima, the principal one being restored to N.N.W. while the second has receded to S.S.W., the eastern becoming thereby a third maximum. The first and second minima occupy the same positions as before, while a third minimum has appeared at W.S.W. In autumn the two western maxima of the summer coalesce and jointly approach the winter position, being now at W.N.W. The eastern maximum continues but it is considerably softened down.

Table LXVI., giving the quarterly ratios for the period of ten years, where calms are *included* in the divisors, has been formed to aid in the construction of Tables XI. and XII., wherein extraordinarily high and low temperatures are classed under the several winds.

The ratios in Table LXVII. shew the annual distribution of each wind taken singly. The only instance of an uninterrupted annual period is that of the south wind, which passes continuously from its maximum in July, to its minimum in December. The winds from S.E., S.S.E., and S.S.W. follow the same order as that of the south wind in their progression, but not with equal regularity. Winds from E.S.E., E., and E.N.E. have their maxima in April or May. Proceeding from the S.S.W. towards the west, we find that the transition from month to month, in the frequency of S.W. winds, follows no definite rule; at W.S.W. and W. the maxima are found in the cold months and the minima in the warm months, while the winds from W.N.W. through N. to N.E. are very irregular. It may be remarked however, with reference to these latter, as well as to the S.W. winds, that although irregular as regards the differences between month and month, there is a fair correspondence in the different groups of years.

The general character of the annual march in the frequency of each wind taken singly, and the change in the position of the maxima in the annual period as different winds are considered, will be better seen from the quarterly averages of the ratios of Table LXVII., which are given in Table LXVIII. Com-

the results
in two equal
parts and the
collectively,
no absolute
quotients
of the arithmetic means,
are above
alone are
in the whole
and is above
range on the
the principal
so a second
and minimum
ing to Table
progression is
the maximum
maximum
maximum is
while the
and second
In autumn
ing now at

ded in the
narily high

The only
ously from
S.W. follow
y. Winds
W. towards
follows no
in the warm
marked how
regards the
years.

the change
better seen
II. Com-

mencing at E.N.E., the quarter of maximum frequency for E.N.E. and E. is spring; for E.S.E. and other points through S. to S.W. the greatest frequency is in summer. At S.W. a sudden breach of continuity takes place, the winter becoming the quarter of greatest frequency for winds from S.W., W.S.W., and W. With respect to winds from W.N.W. through N. to N.E. nothing of any definite character can be stated.

ON THE WINDS IN THE UPPER STRATA AS SHEWN BY THE MOTION OF THE CLOUDS.
(TABLES LXIX. TO LXXIV.).

At each of the six observation hours a note was always made of the direction from which the clouds were moving. When the motion was too slow to be detected by the eye, or when no clouds were visible, the sky in these tables is designated as being calm or clear. It is probable that in many instances, and especially at the night hours, the clouds have been recorded as motionless in consequence of the inability of the observer to perceive the motion, and hence in part may be explained the very large number of cases of calm sky, amounting to nearly one-third of all the observations.

In Table LXIX. are given the absolute number of each upper wind, for each month as well as for the two half-years and year, as furnished by six observations daily during the nine years 1844-62. The quotients resulting from the division of the numbers in Table LXIX. by the monthly, half yearly, and yearly means, *including* the cases of clear and calm sky, are given in Table LXX.

It is seen that while the winds from west are far more numerous than all other upper winds, they are exceeded in the six winter months separately and collectively by the calms, while in the summer the westerly upper winds exceed the calms in the six months collectively, and in each of the months, June, July, August, and September. The cases of clear sky are above the average of the observations, or more than one in ten, in every month but November and December.

The relative frequency of the different upper winds is seen better in Table LXXI., wherein the numbers of Table LXIX. are expressed in terms of the means of the eight winds, without including the cases of calm or clear sky.

Of the upper currents, that from the west greatly exceeds in frequency all others; not only on the average of the year, for which it is 3.80 times as frequent as the average frequency of all winds, but also in each month taken singly. Next in order of frequency in each month separately, is the wind from N.E., which has a more than average frequency in each single month, its relative frequency on the average of the year being 1.91. The S.W. wind is a little above the average of all winds in winter, and under the average in summer, while the winds from the other five points are greatly below the average in every month, the average of the ratios that express their relative frequency in the year being only 0.27. The least frequent upper wind is from the south, the next in order being the N.E. wind.

In Tables LXXII. and LXXIII. the half-yearly and yearly ratios in Tables LXX. and LXXI., derived from six daily observations, are compared with the ratios from the three hours, 8 A.M., 2 P.M., and 4 P.M., and with the ratios from the hours 10 P.M., midnight, and 6 A.M.

While the calms are more numerous than the west winds on the average of the year, when the six hours are reckoned together, it appears from LXXII., that for the three hours 8 A.M., 2 P.M., and 4 P.M., the west winds greatly exceed the calms on the average of the year, as well as in the winter and summer

separately, the excess being greatest in the summer. For the other three hours the calms are greatly in excess in both seasons, and chiefly so in the winter.

In Table LXXIII., where the winds alone are compared without reference to the calm or clear sky, it is found that the west wind maintains its preponerance materially to the same extent in each group of hours. The N.W. wind is the second, and the S.W. wind the third in order of frequency, in both seasons and in both groups of hours; but both in summer and in winter the excess of the N.W. over the S.W. wind is greatest in the night and least in the day.

The annual distribution of each upper wind is given in Table LXXIV., together with the ratios expressing the relative frequency of each wind in the winter as compared with the summer. The monthly numbers are derived from those of Table LXXII. by dividing these latter by the annual means corresponding to each wind. The three final columns are obtained by dividing the numbers in the columns of Table LXXII. headed "winter," by the numbers in the columns headed "summer."* From the three final columns it is seen that the N., N.W., and S.W. winds, on the average of the six hours, are nearly equally frequent in winter and in summer, the frequency in winter slightly exceeding that in summer during the day and falling short of it in the night; the greatest disparity in the day and night occurring with the S.W. wind. The S.E. and S. winds are in excess in winter as compared with summer, both during the day and during the night. The E., N.E., and W. winds are less frequent in winter than in summer at all hours, and particularly during the night. In fact, with respect to all upper winds, it may be stated that the ratios expressing their relative frequency in winter as compared with their frequency in the summer, are greater at the hours 8 A.M., 2 P.M., and 4 P.M., than at 10 P.M., midnight, and 6 A.M. The calms of winter greatly exceed in frequency those of summer, the preponderance being nearly the same at all hours. Cases of clear sky, on the contrary, are most frequent in summer, their preponderance in summer being the same for both groups of hours.

In Table LXXV. are collected the absolute number of each upper wind at each separate observation hour, together with the aggregate number in each of the two groups of hours, namely at 10 P.M., midnight, and 6 A.M., and at 8 A.M., 2 P.M., and 4 P.M.

Table LXXVI. shews for each hour separately, as well as for each of the two groups of hours, the relative frequency of each upper wind, together with the cases of calm and clear sky. It is derived from the preceding table by dividing the numbers therein contained by the arithmetic means given in the last line but one. Table LXXVII. has been computed in a similar manner from Table LXXV., the divisors being the arithmetic means in the lowest line of that table.

From Table LXXVI. we learn that while upper winds from the west greatly exceed all other winds at every hour, the calms are more numerous than the west winds at the hours 6 A.M., 10 P.M., and midnight, both separately and collectively, and less numerous at 8 A.M., 2 P.M., and 4 P.M. Cases of clear sky are above the average, or exceed one in ten of the observations at every hour but 2 P.M. Midnight is the hour at which a clear sky is most frequent; clear sky being generally more than twice as frequently found during the hours 6 A.M., 10 P.M., and midnight, as at the other hours.

From Table LXXVII., in which the different upper winds are compared, without reference to the cases of calm or clear sky, we find that the west is by far the most frequent of the upper winds, but that

* The numbers which in Table LXXII. are given to two decimal places, were carried out to three places when the divisions were performed; hence the want of perfect accordance between Table LXXII. and the three final columns of Table LXXIV.

its preponderance is nearly the same at all hours. The N.W. is the second in order of frequency, and the S.W. the third. The N.W. wind at all hours is above the average of all winds; but the excess in its frequency is greater in the night than in the day. The S.W. wind is equal to the average of all winds during the day, but considerably below it in the night. Winds from the remaining five points are greatly below the average at all hours.

The diurnal distribution of each upper wind, taken singly, is shewn in Table LXXVIII. The hourly numbers are derived from those of Table LXXV, by dividing them by the six-hour means in the final column of that table. The three final columns of Table LXXVIII. give the ratios of the absolute number of each upper wind at the day hours to the absolute number of the same wind at the hours of the night.

From the hourly numbers in this table it appears that every upper wind is above the average during the three hours 8 A.M., 2 P.M., and 4 P.M., and below the average at the other hours, an exception being made with respect to the S.E. wind at 6 A.M. The excess in the frequency of all upper winds at the three day hours, as compared with that of the other three hours, is shewn by the final columns to be greater in winter than in summer. It is greatest for the N.E. wind and least for the N.W. wind. Both calm sky and clear sky are less than half as frequent in the day as in the night, the disparity between the day and night being the same for both, and being also the same in winter as in summer.

Tables LXXIX., LXXX., LXXXI., are derived from the observations of three years only, namely, 1860, '61, and '62, and are designed to shew the relations between the upper currents and the simultaneous surface winds. The method employed in the computation is sufficiently explained by their titles.

It is obvious that if the decimal points be omitted in Table LXXX. the numbers under any point of the compass will shew how often, out of a thousand times that the corresponding surface wind occurs, it is accompanied by each of the several upper currents. Thus out of 1000 times that a surface north wind blows, calms aloft occur 375 times, a clear sky 273 times, an upper current from N. 38 times, and one from S. only 4 times. With a surface wind from E. the clouds move more frequently from W. than from E. With surface S. winds the clouds never move from the N., and they are about 27 times as numerous from the W. as they are from the S. Whatever be the surface wind, unless it be from S., S.W., or W., it is more frequently attended by a calm sky than by a motion of the clouds in any one specified direction; but when the lower wind is from S., S.W., or W., a motion of the clouds from W. is more frequent than a calm sky. A clear sky is above the average with surface winds from every point but the E.; in other words it occurs in the ratio of more than one for every ten times that each surface wind blows. Calms aloft and upper winds from W. are above the average during every surface wind. A calm sky is more frequently an accompaniment of an easterly surface wind than of a surface calm.

Whatever be the surface wind, it appears from Table LXXXI. that the westerly upper wind far exceeds all other upper winds in frequency. It is most frequent with a surface wind from W., and least frequent with one from E. The N.W. upper current which stands next in order of frequency is above the average of the upper currents with all surface winds but those from E. and N.E. The S.W. is above the average of upper currents when the surface wind is from any point from N.E. through S. to S.W., and below the average with a surface wind from N., N.W., and W. The upper current from E. is above the average when the surface wind is from N.E. and E., but below the average with all other surface winds. The S.E. upper wind is far below the average, excepting during an easterly surface wind, when it is slightly above. Upper winds from N., N.E., and S. are below the average whatever be the winds at the surface.

ON THE RELATIVE FREQUENCY OF THE DIFFERENT WINDS DURING DAYS OF RAIN OR SNOW, FROM THE
HOURLY RECORDS OF TEN YEARS (TABLES LXXXII. TO LXXXVI.)

The object of these tables is to compare the different winds with reference to the number of hours that they blow during days in any part of which a fall of rain or snow takes place. If all winds continued for an equal number of hours through the year or through the particular season under consideration, it would be sufficient to compare the *absolute* durations of the several winds on days of precipitation; but as there is a very great inequality in the frequency of winds from different points of the compass, (winds from N.W., for example, being more than three times as numerous as those from S.E.,) an undue prominence would be given to the winds of greater general frequency, if the comparison were to be made between the absolute durations. Hence it becomes requisite that the absolute durations of each wind during the days of rain included within a given period of time, should be divided by the whole duration of the same wind within the same period. The quotients form what may be termed the *relative* durations of the several winds, and constitute the proper quantities for intercomparison.

As winds of comparatively rare occurrence on days of heavy rain were found, according to the results in the preceding volume, to blow very frequently on days of light rain, the adoption of some classification of the rainy days became necessary. In these tables the days of rain have been arranged in three classes, which have been considered separately, as well as collectively in one group. Class I. includes days of light rain, in which the whole amount in the day did not exceed one-tenth of an inch. Class II. includes days of moderate rain, or over one-tenth and less than half an inch, while Class III. comprises days of heavy rain, wherein the fall in the day amounted to half an inch and upwards. The days in which snow fell are classified in a similar manner and with the same limits, one inch of snow being regarded as equivalent to one-tenth of an inch of rain. With a view of learning whether the relative duration or frequency of a wind during rain is affected by the season, the computations have been made separately for the winter half-year (October to March,) for the summer half-year (April to September,) and for the year as a whole. As the falls of snow after March are not sufficiently numerous to furnish materials for a separate discussion, no separation of the seasons has been made in the case of the snow.

Again, for the purpose of comparing the corresponding results in different years, the observations of the ten years, 1853 to 1862, have been discussed in two separate equal groups as well as in one.

In Table LXXXII. the durations of the different winds on days of rain during the six months, October to March, are examined. The three lines marked (1) in Class I. contain the absolute number of hours that each wind blew on days of light rain. The three lines marked (2) give the absolute number of hours that the same winds blew, with and without rain, during the same period. The three lines marked (3) are the relative durations of the several winds, being the quotients arising from the division of the numbers in (1) by the corresponding numbers in (2). It is clear, if the decimal points be disregarded, that any one of the numbers in (3) will indicate the number of hours comprised in days of light rain out of a thousand hours during which the corresponding wind blew within the same period; thus out of 1000 calm hours in the winters of the ten years, 1853 to 1862, 169 occurred during days of light rain. The ratios in the three lines marked (4) are obtained from the numbers in (3) by dividing each number by the arithmetic mean of the seventeen numbers in the same line. The computations for the other classes in Table LXXXII. as well as those of the corresponding tables for the rain in the summer half-year, for the rain in the whole year, and for the snow (Tables LXXXIII., LXXXIV., and LXXXV.,) have been made in a precisely similar manner.

Table LXXXVI. gives a synopsis of the final ratios in the four preceding tables.

Referring to the results of the ten years, it appears that in winter, whatever be the class of the rainy days, the progression in the relative frequency of the different winds is single, the maximum being at or near east, and the minimum at N.W. The amplitude however, is much greater for the heavier falls, the ratio of the maximum to the minimum being more than 20:1 in Class III., and less than 2:1 in Class I. The portion of the compass also for which the relative frequency is above the average extends from E.N.E. through S. to S.W. in Class I., but is limited in Class III. to the points N.E. to S.S.E. inclusive.

Between the summer and the winter half-years a marked contrast may be noticed, and particularly as regards the lighter rain, a double progression in the summer occurring in each of the classes.

For light rain the principal maximum is at W.S.W. and the principal minimum at or near N.N.E. the second maximum being at or near E., and a second minimum between S. and S.E. In Class II. the maximum at E. slightly exceeds the western maximum, and in Class III. the excess is very decided. Where rain is considered without reference to its amount, the eastern and western maxima are very nearly equal. Taking the whole year together, the distribution of the winds in Class I. is chiefly governed by the summer half-year, while Classes II. and III. resemble the same classes in the winter, although with less prominently marked features.

For the light snow there is a tolerably well defined maximum at or near W. From W., in both directions round the compass, there is a descent in the frequency of the winds as far as E. and S., at which points there are what may be termed two equal minima including a space, for which, on the average of the three included points, the winds are about 1.3 times as numerous as at either E. or S.

Classes II. and III., which possess the same general characters as regards the distribution of the winds, are in striking contrast with that which is found to accompany the lighter falls of snow; the most frequent wind on days of moderate or heavy snow being very decidedly at or near N.E., while the western maximum is nearly obliterated. The wind of most rare occurrence during such falls being from points at or near S. When no account is taken of the amount of snow that falls in the day, the most frequent wind relatively is from W., the N.E. wind forming a second maximum not very inferior to that at W. The least frequent wind is from S., and there is also a second depression between the maxima at W. and N.E., but which does not fall below the average as compared with all the points of the compass.

DEPTH AND FREQUENCY OF RAIN AND SNOW—(TABLES LXXXVII. AND LXXXVIII.)

The number of days of rain, together with its depth in inches, for each month of the years 1860, 1861, and 1862, are given in Table LXXXVII. Table LXXXVIII. contains the corresponding results for snow. The monthly and annual averages for the three years are accompanied by the corresponding averages derived from a series of years. The average frequency of rain and snow is derived, in each case, from the records of twenty-three years; but owing to breaks that occurred in the early part of the series, the average monthly and annual depths of rain are obtained from twenty-one, and those of snow from twenty years.

From these tables it appears that both rain and snow were much more *frequent* on the average of the three years, 1860-'62, than on that of the whole series, for the year taken collectively, as well as for almost every month; days of rain being about 20 per cent. and days of snow about 30 per cent. more

numerous in the partial than in the entire series. The *depth* of rain, on the contrary, was nearly 16 per cent. in defect on the average of the three years, as compared with the general average of the whole series. The average annual depth of snow was in excess, but not to such an extent as to compensate for the deficient rain; the general deficiency in the precipitation of the three years (allowing an inch of rain to be equivalent to ten inches of snow,) being more than 11 per cent.

The results derived from the hourly rain gauge, which was in operation from April to November in 1861, and from April to October in 1862, do not reveal any distinct diurnal period, either as regards the depth or the frequency of the rain. Arranging the day in six periods of four hours each, and dividing the depth of rain in each period and its frequency (or the number of hours in any part of which a fall occurred,) by the average depth and frequency in the six periods, the ratios are obtained which are given in the annexed table.

PERIODS.	6 A.M. to 10 A.M.	10 A.M. to 2 P.M.	2 P.M. to 6 P.M.	6 P.M. to 10 P.M.	10 P.M. to 2 A.M.	2 A.M. to 6 A.M.	Average of the six periods.
Depth, 1861....	0.55	0.76	1.37	1.41	1.07	0.84	1.00
" 1862....	1.18	0.93	1.00	0.92	0.97	1.00	1.00
Frequency, 1861....	0.84	0.84	1.02	1.06	1.08	1.16	1.00
" 1862....	1.18	0.97	0.92	0.99	0.88	1.06	1.00

MEAN ANNUAL VARIATIONS OF TEMPERATURE FROM THE OBSERVATIONS OF TWENTY-THREE YEARS.

Table LXXXIX. is an extension of a similar table published by General Sabine on page 163 of the Philosophical Transactions for the year 1853. The monthly and annual means of temperature for the twelve years, 1841-'52, are reprinted from the paper of General Sabine. For the eleven years, 1853-'63, inclusive, the monthly means derived from six observations each day are corrected for diurnal variation, by aid of the tables by General Sabine on pp. 145 and 146 of the same volume.

For the sake of comparison the means furnished by the twelve years, 1841-'52, have been introduced, together with the normal temperatures for the parallel of latitude $43^{\circ} 40' N.$ as computed by Dove.

The probable variabilities of the several months at the foot of the table are derived from the squares of the differences between the partial and general monthly means. The quarterly averages of these numbers, from the series of twenty-three years, and from the twelve years, are as follows:

	Winter.	Spring.	Summer.	Autumn.
1841-'52	2.6	2.2	1.4	1.8
1841-'63	2.8	2.0	1.5	1.6

The probable variability of a single year, as regards the annual mean, is $0^{\circ}.63$ for the twelve years, and $0^{\circ}.61$ for the twenty-three years.

early 16 per
whole series.
for the defi-
of rain to be

November in
regards the
and dividing
which a fall
eh are given

Average
of the
six periods.

1.00

1.00

1.00

1.00

EE YEARS.

e 163 of the
ture for the
rs, 1853-'63,
al variation,

introduced,
Dove.
the squares
these num-

elve years,

INTRODUCTION.

xxiii.

The following are the quarterly averages of the probable errors of the monthly means for the whole period :

	Winter.	Spring.	Summer.	Autumn.
	0.59	0.42	0.30	0.33

The probable error of the general mean temperature for the year is 0.18 for the twelve years, and 0°.13 for the twenty-three years.

If the assumption be accepted that the monthly means of temperature for the whole period are the temperatures proper to epochs separated by twelve equal intervals, the January mean being the mean temperature corresponding to the 15th of that month, regarded as the zero of time, then will the temperature T_n , corresponding to the time (n), (the unit of time being the twelfth part of a year) be represented by the expression.

$$T_n = 44.18 + 22.32 \sin(n \times 30^\circ + 261.2^\circ) + 0.74 \sin(2n \times 30^\circ + 78.25^\circ) \\ + 0.61 \sin(3n \times 30^\circ + 181.8^\circ) + 0.29 \sin(4n \times 30^\circ + 38.3^\circ) \\ + 0.74 \sin(5n \times 30^\circ + 51.7^\circ) + 0.30 \cos(6n \times 30^\circ). \quad (\text{I.})$$

The two assumptions, (1) that the mean temperature of a month is identical with the temperature of its middle point, and (2) that the months are all of equal length, are evidently not in strict accordance with fact. A nearer approach to accuracy will be made by admitting the second of the preceding assumptions only and by applying corrections for the errors introduced by the first. Making the necessary corrections* to the coefficients of the several terms of equation (I.), the mean diurnal temperature T'_n at any time (n), will be given more accurately by the following equation :

$$T'_n = 44.18 + 22.58 \sin(n \times 30^\circ + 261.2^\circ) + 0.77 \sin(2n \times 30^\circ + 78.25^\circ) \\ + 0.68 \sin(3n \times 30^\circ + 181.8^\circ) + 0.35 \sin(4n \times 30^\circ + 38.3^\circ) \\ + 1.00 \sin(5n \times 30^\circ + 51.7^\circ) + 0.47 \cos(6n \times 30^\circ). \quad (\text{II.})$$

If the values 0, 1, 2, &c., 11, be substituted for (n) in equations (I.) and (II.), the corresponding values of T_n in equation (I.) will be identical with the monthly means from the records of twenty-three years given in Table LXXXIX.; and the corresponding values of T'_n in equation (II.) will be the diurnal means proper to the middle points of the twelve months regarded as of equal length.

MEAN ANNUAL VARIATIONS OF BAROMETRIC PRESSURE FROM THE OBSERVATIONS OF TWENTY-THREE YEARS.

The mean barometric pressures for every month of each of the twenty-three years, 1841 to 1863, have been collected in Table XC. During the seven years, July 1842 to June 1848, the readings were made at each of the twenty-four hours. From this series tables were formed containing, for each month and hour, the differences between the monthly means for the hour and the monthly means on the average of twenty-four hours. These differences being regarded as corrections for diurnal variation, were employed in reducing the monthly means in the remainder of the series, during the greater part of which the observations were made six times only each day.

* See note on page xxiv.

The greatest probable variability of a monthly mean, .0562, is in February, and the least, .0292, is in July. The quarterly averages of the probable monthly variabilities and of the probable errors of the monthly means given by the whole series, are the following :

	Winter.	Spring.	Summer.	Autumn.
Variability0435	.0433	.0313	.0414
Probable error....	.0091	.0090	.0065	.0086

The probable variability of a single year with respect to the annual mean barometric pressure is .0125, and the probable error of the general annual mean for the whole series is .0026.

Making the same assumptions as in the case of temperature, B_n , the diurnal mean of barometric pressure at any time (n), may be computed from the following formula, derived from the twelve monthly means at the foot of Table XC.

$$\begin{aligned} B_n = & 29.6190 + .0350 \sin(n \times 30^\circ + 147^\circ 21') + .0126 \sin(2n \times 30^\circ + 11^\circ 29') \\ & + .0177 \sin(3n \times 30^\circ + 116^\circ 34') + .0067 \sin(4n \times 30^\circ + 133^\circ 11') \\ & + .0081 \sin(5n \times 30^\circ + 251^\circ 20') - .0018 \cos(6n \times 30^\circ). \end{aligned} \quad (\text{I.})$$

The diurnal means are given more accurately by the following equation, obtained from (I.) by applying the requisite corrections* to the co-efficients:

$$\begin{aligned} B'_n = & 29.6190 + .0354 \sin(n \times 30^\circ + 147^\circ 21') + .0132 \sin(2n \times 30^\circ + 11^\circ 29') \\ & + .0197 \sin(3n \times 30^\circ + 116^\circ 34') + .0081 \sin(4n \times 30^\circ + 133^\circ 11') \\ & + .0109 \sin(5n \times 30^\circ + 251^\circ 20') - .0028 \cos(6n \times 30^\circ). \end{aligned} \quad (\text{II.})$$

* To effect the corrections to which reference is made above, the coefficients of the terms in (I.) which involve $\frac{n}{\pi}$, $\frac{2n}{\pi}$, $\frac{3n}{\pi}$, &c., are multiplied respectively by the factors $\frac{12}{\sin \frac{\pi}{12}}$, $\frac{12}{\sin \frac{2\pi}{12}}$, $\frac{12}{\sin \frac{3\pi}{12}}$, &c. The products resulting from this multiplication are the coefficients of the corresponding terms in (II.).

least, .0292, is
e errors of the

etric pressure is

barometric pres-
monthly means

29).

11).

(I.)

n (I.) by apply-


29).

11).

(II.)

TORONTO

METEOROLOGICAL RESULTS.



GENERAL METEOROLOGICAL ABSTRACT.—JANUARY, 1860.

TORONTO METEOROLOGICAL RESULTS.

1

DATE.	DAILY MEANS.			WIND.			EXTREMES OR TEMPERATURE.			RAIN.			SNOW.			RAIN AND MELTED SNOW.		
	Temperature of the Air.	Vapour,	Relative Humidity.	Barometric Pressure.	Dry Air Pressure of Surface of Clouded Sky.	Relative Humidity.	Wind Velocity.	Maximum Velocity.	Minimum Velocity.	Depth in Inches.	Approximate Duration in hours.	Depth in Inches.	Approximate Duration in hours.	Depth in Inches.	Approximate Duration in hours.	Depth in Inches.	Approximate Duration in hours.	
1	4.40	0.045	83	30.104	30.058	0.72	N 86 W	4.45	6.54	9.6	-3.0	12.6	2.0	0.020	...	
2	16.42	0.080	87	29.772	29.682	1.00	S 66 W	12.54	12.98	22.2	-1.8	13.6	...	0.2	11.0	0.50	11.0	
3	9.32	0.060	83	30.111	30.048	0.50	S 58 W	11.43	10.8	20.2	9.4	6.8	...	0.2	1.5	0.20	1.5	
4	12.33	0.064	83	30.111	30.048	0.62	S 77 W	9.60	9.77	18.4	-6.8	25.2	
5	24.32	1.08	79	29.758	29.650	0.92	S 52 W	9.76	9.92	33.9	11.2	22.1	...	*	0.3	*.3	...	
6	7	35.18	193	93	3.392	1.00	S 54 W	3.68	3.78	37.0	28.1	8.9	0.200	8.2200	8.2	
7	9	33.82	171	87	3.392	1.00	S 68 W	7.03	8.00	38.6	33.4	5.2	*	0.5	...	*	0.5	
8	10	38.22	206	90	5.949	3.33	1.00	N 15 E	6.04	6.18	29.0	32.0	7.0	1.70	*	*.170	6.5	
9	11	19.60	69	37	7.443	1.00	N 25 W	11.07	11.27	42.5	33.8	8.7	3.40	8.2	0.1	0.3	3.50	
10	12	18.85	0.74	77	8.112	0.733	0.42	S 78 W	8.51	11.63	24.0	10.4	13.6	...	0.2	2.9	0.20	
11	13	12.32	68	88	9.50	0.882	0.70	N 43 E	5.00	5.33	19.2	9.5	10.6	...	0.5	
12	14	20.08	67	87	4.73	0.76	0.73	S 67 W	2.24	2.51	28.0	9.9	18.5	...	0.5	0.50	5.0	
13	15	32.98	151	79	3.60	2.09	0.78	N 65 W	13.89	14.10	37.0	22.4	14.6	
14	16	22.33	106	90	6.47	1.00	N 17 W	6.39	8.46	33.5	33.2	33.5	
15	17	20.20	87	81	4.91	4.04	1.00	N 17 W	4.57	4.81	35.2	19.4	3.8	...	0.1	2.0	0.10	
16	18	25.25	118	86	3.364	2.45	0.78	S 78 W	7.78	7.75	20.2	17.0	12.2	...	2.0	12.0	0.20	
17	19	24.33	105	80	5.10	4.05	0.75	S 73 W	7.15	10.18	28.5	12.5	12.5	...	*	0.5	0.5	
20	20	34.68	155	77	3.38	0.62	0.62	S 52 W	9.65	9.81	40.2	23.9	16.3	
21	21	33.98	152	77	4.42	0.290	0.23	N 79 W	4.16	4.58	42.4	31.7	10.7	
22	23	30.15	119	70	8.83	7.18	0.23	S 56 W	4.11	4.49	35.5	23.0	12.5	...	0.2	0.5	0.20	
24	24	41.48	203	77	4.42	2.19	0.98	S 70 W	12.64	13.49	46.4	30.8	15.6	
25	25	31.92	113	64	6.21	5.08	0.32	N 66 W	18.88	19.28	37.8	32.5	5.3	0.8	
26	26	20.68	082	75	7.51	6.69	0.93	N 59 W	1.51	1.51	23.0	19.5	3.5	
27	27	19.57	0.988	81	5.567	4.69	0.57	N 71 W	5.63	8.63	27.4	16.8	10.6	...	1.0	5.2	100	
28	28	17.13	0.683	84	4.402	4.10	0.50	N 3 W	4.50	7.60	24.4	10.8	13.6	...	2.5	5.0	250	
29	29	28.35	108	68	3.79	2.71	0.42	S 56 W	14.30	16.20	38.5	8.0	30.5	...	0.1	0.8	...	
30	30	2.35	0.637	74	8.72	0.834	0.67	N 9 W	9.19	9.36	7.9	4.3	3.6	...	0.1	1.5	0.10	
31	31	23.38	0.110	81	20.643	29.533	0.71	N 89 W	6.09	9.37	26.83	17.58	12.25	0.740	24.2	8.7	50.5	
																	1.610 74.7	

* Inappreciable.

GENERAL METEOROLOGICAL ABSTRACT.—FEBRUARY, 1860.

2

* Inappreciable.

GENERAL METEOROLOGICAL ABSTRACT.—MARCH, 1860.

Days.	DAILY MEANS.				WIND.				EXTREMES or TEMPERATURE.				RAIN.				SNOW.				RAIN AND MELTED SNOW.				
	Temperature of the Air. F.	Humidity. Percent.	Pressure of Vapors. Inches.	Pressure of Hyd. Air. Inches.	Wind. Velocity. Miles.	Wind. Direction. Degrees.	Wind. Velocity. Miles.	Wind. Direction. Degrees.	Maximun Temperature. Inches.	Minimun Temperature. Inches.	Difference in Hours.	Avg. Duration in Hours.	Maximun Velocity. Miles.	Wind. Direction. Degrees.	Maximun Velocity. Miles.	Wind. Direction. Degrees.	Approximate Depth, in Inches.	Approximate Duration in Hours.	Maximun Velocity. Inches.	Wind. Direction. Degrees.	Approximate Depth, in Inches.	Approximate Duration in Hours.	Approximate Depth, in Inches.	Approximate Duration in Hours.	
1	42.85	0.262	95	29.409	29.237	1.00	8.83	W	46.5	8.5	0.017	3.0	0.017	3.0	
2	40.15	.184	74	.785	.602	0.38	N 69° W	3.90	12.76	48.4	35.0	13.4015	4.0015	4.0	
3	41.07	.173	68	.314	.172	0.77	S 69° W	13.76	20.35	50.2	31.2	16.0	.015	4.0	0.2	2.0020	2.0	
4	
5	31.63	.137	76	.351	.214	0.28	S 68° E	4.60	7.41	37.0	20.5	7.5	
6	33.08	.153	81	.543	.390	0.67	N 78° E	12.35	12.96	37.2	23.5	13.7	.265	6.5275	7.0
7	45.42	.232	73	.193	.28.961	0.33	N 96° W	7.94	9.68	62.5	32.4	30.1	
8	40.35	.130	52	.459	.29.329	0.67	N 69° W	13.07	14.07	47.5	38.8	8.7	
9	26.57	.094	67	.411	.317	0.85	N 49° W	19.28	20.01	34.2	26.0	8.2	
10	23.77	.085	65	.541	.456	0.92	N 52° W	18.02	18.36	28.5	24.0	13.5	
11		
12	18.28	.075	74	.582	.507	0.42	S 31° W	19.81	20.46	22.0	19.2	2.8	
13	25.65	.109	78	.800	.690	0.63	S 55° W	4.06	5.22	12.8	17.6		
14	34.00	.145	72	.753	.608	0.13	S 82° W	2.70	3.91	43.5	23.4	20.1		
15	36.17	.149	70	.885	.736	0.00	S 25° W	1.92	3.82	46.0	26.6	19.4		
16	36.83	.162	73	.864	.762	0.00	S 74° E	3.07	3.50	46.5	28.0	18.5		
17	40.18	.179	71	.854	.765	0.02	N 75° E	3.86	4.32	47.2	28.0	19.2		
18		
19	40.78	.229	89	.531	.312	0.62	N 73° E	5.80	6.46	51.0	33.0	18.0		
20	34.60	.148	71	.400	.252	0.90	N 58° W	20.38	22.47	43.3	35.0	8.3	*	
21	24.78	.082	60	.691	.609	0.22	N 44° W	28.77	28.83	29.4	17.8	11.6		
22	31.07	.099	55	.483	.333	0.25	N 51° W	20.00	21.48	38.8	22.7	16.1		
23	30.33	.134	80	.324	.190	0.57	S 82° W	5.46	10.08	35.0	26.0	9.0		
24	25.68	.107	76	.268	.160	0.95	N 64° W	17.14	17.56	28.4	23.4	5.0		
25	25.35		
26	27	33.78	.131	.59	.680	.602	N 53° W	16.16	16.44	31.2	21.4	9.8		
28	31.30	.131	74	.398	.350	0.08	S 47° W	12.28	13.63	43.0	21.0	22.0		
29	39.07	.135	58	.303	.207	0.55	N 82° W	12.81	14.71	37.4	28.0	9.4		
30	45.03	.189	67	.222	.033	0.18	N 86° W	4.45	6.13	53.0	28.8	26.2		
31	53.20	.270	64	.108	.28.883	1.00	S 45° W	10.82	11.52	57.5	29.0	25.5		
	34.48	0.148	71	29.511	29.363	0.49	N 64° W	7.61	12.41	41.83	27.35	14.54	0.882	22.0	2.4	22.4	1.122	14.4							

* Immeasurable.

* Immeasurable.

GENERAL METEOROLOGICAL ABSTRACT.—APRIL, 1860.

TORONTO METEOROLOGICAL RESULTS.

GENERAL METEOROLOGICAL ABSTRACT.—APRIL, 1860.

DAILY MEANS.	EXTREMES OF TEMPERATURE.				RAIN.				SNOW.				RAIN AND MELTED SNOW.					
	WIND.		PRESSURE.		DEPTH IN INCHES.		DEPTH IN FEET.		DEPTH IN INCHES.		DEPTH IN FEET.		DEPTH IN INCHES.		DEPTH IN FEET.			
	Wind Velocity Miles per hour.	Wind Velocity Miles per hour.	Mean Velocity Miles per hour.	Mean Velocity Miles per hour.	Maximum Wind Velocity Miles per hour.	Maximum Wind Velocity Miles per hour.	Appreciation in Depth in inches.	Appreciation in Depth in feet.	Appreciation in Depth in inches.	Appreciation in Depth in feet.	Appreciation in Depth in inches.	Appreciation in Depth in feet.	Appreciation in Depth in inches.	Appreciation in Depth in feet.	Appreciation in Depth in inches.	Appreciation in Depth in feet.		
DATE.	Temperature of Air.	Pressure of Air.	Barometer.	Humidity.	Temperature of Air.	Pressure of Air.	Wind Velocity Miles per hour.	Wind Velocity Miles per hour.	Temperature of Air.	Pressure of Air.	Wind Velocity Miles per hour.	Wind Velocity Miles per hour.	Temperature of Air.	Pressure of Air.	Wind Velocity Miles per hour.	Wind Velocity Miles per hour.		
1	56.42	0.85	29.449	57	29.414	0.40	21.84	9.57	32.7	19.5	13.2	3.5	33.0	3.5	10	0.010	1.0	
2	56.42	0.85	29.449	57	29.414	0.40	31.51	7.63	52.4	26.8	25.6	5.0	51.6	5.0	10	0.010	1.0	
3	49.93	.191	29.449	57	28.889	.080	7.05	4.17	45.8	52.0	32.2	19.8	41.0	5.8	10	.410	.58	
4	46.70	.213	29.449	57	40.09	.009	6.6	6.6	53.1	6.94	48.5	36.0	12.5	0.2	10	.910	0.2	
5	39.43	.176	29.449	57	29.035	.211	7.2	6.6	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
6	39.43	.176	29.449	57	29.035	.211	7.2	6.6	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
7	41.42	.196	29.449	57	.608	.410	6.6	6.6	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
8	41.42	.196	29.449	57	.608	.410	6.6	6.6	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
9	42.08	.202	29.449	57	.613	.411	7.5	7.5	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
10	41.85	.242	29.449	57	.441	.199	1.00	1.00	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
11	39.35	.170	29.449	57	.611	.471	2.7	2.7	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
12	39.35	.199	29.449	57	.516	.316	5.52	5.52	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
13	40.72	.200	29.449	57	.532	.352	6.6	6.6	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
14	27.50	.111	29.449	57	.705	.594	7.5	7.5	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
15	29.449	57	.705	.594	7.5	7.5	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
16	39.00	.202	29.449	57	.84	.496	2.34	2.34	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
17	40.53	.175	29.449	57	.68	.742	3.56	3.56	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
18	33.90	.115	29.449	57	.61	.20191	0.676	0.676	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
19	43.73	.179	29.449	57	.63	.29.690	29.681	0.632	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
20	44.88	.285	29.449	57	.496	.490	2.33	2.33	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
21	40.42	.298	29.449	57	.84	.435	1.37	1.37	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
22	36.98	.151	29.449	57	.68	.616	6.6	6.6	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
23	36.98	.151	29.449	57	.68	.616	6.6	6.6	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
24	32.47	.143	29.449	57	.582	.439	0.50	0.50	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
25	33.85	.141	29.449	57	.723	.524	3.83	3.83	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
26	26.70	.151	29.449	57	.656	.605	0.58	0.58	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
27	40.87	.160	29.449	57	.64	.881	.721	.721	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
28	41.63	.193	29.449	57	.73	.838	.615	.615	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
29	30.72	.257	29.449	57	.70	.577	.320	.320	31.16	16.18	28.3	5.72	32.0	17.0	10	.052	.58	
30	39.55	.0185	29.578	74	29.393	0.50	N 35° W	41.10	10.30	47.04	32.19	14.84	12.82	40.6	0.3	7.2	1.312	47.8

* Inappreciable.

TORONTO METEOROLOGICAL RESULTS.

GENERAL METEOROLOGICAL ABSTRACT.—MAY, 1860.

DAy	DAILY MEANS.			WIND.			EXTREMES of TEMPERATURE.			RAIN.			SNOW.				
	Temperature of Air.	Pressure of Atmosphere.	Humidity.	Wind Velocity.	Direction.	Clouded Sky.	Min. Velocity.	Max. Velocity.	Difference.	Min. Temperature.	Max. Temperature.	Difference.	Min. Rain.	Max. Rain.	Depth in inches.		
1	41.30	0.201	.77	.39.710	.20.510	.98	N 36° W	4.40	.740	42°.5	52°	0.038	2.0	..	0.022	2.0	
2	46.90	0.260	.79	.836	.576	.45	S 63° W	1.19	3.50	57.0	32.5	.015	2.5	..	.015	2.5	
3	53.38	.258	.63	.714	.322	.012	S 37° W	2.20	3.90	69.0	47.8	.212	
4	56.92	.318	.69	.640	.322	.065	S 43° W	1.07	2.03	67.2	48.0	.19.2	
5	56.13	.289	.66	.628	.339	.073	N 80° E	4.39	5.34	68.8	44.2	.24.6	
6449	.79	.008	N 73° W	2.96	3.82	73.2	50.3	.080	1.0	..	.080	1.0	
7	62.02449	.79	.008	N 73° W	0.73	1.00	70.0	50.3	.22.9935	6.0	
8	59.78	.421	.82	.515	.605	.065	N 9° E	3.53	7.86	64.5	56.2	.8.3205	4.3	
9	57.02	.414	.89	.383	.28.969	1.00	S 68° E	6.40	9.41	63.2	54.2	.9.0	.205	4.3	
10	54.30381	.90	.562	N 61° E	1.00	1.181	4.98	3.77	.6.3055	5.8	
11	57.03	.391	.84	.725	.334	.093	S 49° E	8.89	9.11	62.5	50.8	.11.7	
12	58.45	.408	.83	.752	.344	.072	S 52° E	8.64	9.32	65.0	52.0	13.0	
13	S 36° E	5.65	5.57	68.8	53.2	15.6	
14	61.18366	.67	.226	0.15	5.62	7.6	
15	56.12	.285	.63	.662	.377	.000	S 82° E	1.15	4.70	70.0	52.4	17.6	
16	59.42	.295	.59	.608	.372	.000	N 75° E	2.75	4.80	63.2	52.2	11.0	
17	55.03	.296	.68	.720	.424	.043	N 65° E	5.46	7.32	67.0	50.6	.22.2	
18	54.05	.370	.88	.426	.56	.063	S 64° E	4.06	4.97	61.8	48.5	13.3052	1.8	
19	52.23	.308	.76	.160	.28.532	.072	S 72° E	15.78	16.52	60.1	49.0	11.1	.005	1.8	
20	S 36° W	4.56	10.54	53.0	37.4	16.0	
21	41.50224	.85	.419	29.195	1.00	N 55° E	11.49	11.77	44.6	38.8	.135	
22	51.13	.301	.78	.581	.280	.027	S 12° W	5.37	8.22	63.0	40.8	.29.2	
23	50.88	.276	.75	.773	.498	.005	S 27° E	1.14	3.37	61.8	41.0	20.8	
24	51.58297	.70	.686	N 55° E	3.89	4.80	62.0	44.4	17.6	
25	61.43	.328	.62	.530	.530	.202	S 53° E	7.23	7.76	60.2	52.0	.17.2	.013	..	.013	.12	
26	59.40	.416	.80	.301	.28.885	.042	N 1° W	4.59	8.83	71.0	49.8	21.2	.010	0.3	
27	N 55° W	4.78	6.17	65.6	50.2	15.4	
28	57.02	.332	.71	.540	.29.197	..	S 82° W	1.95	4.56	67.5	47.0	.20.5	
29	59.47	.352	.71	.572	.220	0.85	N 56° E	8.28	8.97	68.6	49.5	19.1	.005	0.3	..	.005	0.3
30	62.95	.451	.79	.319	.28.868	0.60	S 17° W	2.70	6.57	74.5	53.5	21.0	* 0.1	* 0.1	..
31	59.15	.430	.85	.357	.28.927	1.00	N 76° W	5.30	7.79	67.0	56.5	10.5	.020	2.7	..	.020	2.7
	53.53	0.338	.76	.29.366	29.298	0.57	N 26° E	2.66	7.17	63.97	47.79	16.18	1.815	36.1	..	1.815	36.1

* Inappreciable.

* Inappreciable.

TORONTO METEOROLOGICAL RESULTS.

GENERAL METEOROLOGICAL ABSTRACT.—JUNE, 1860.

Date.	DAILY MEANS.				WIND.				EXTREMES OF TEMPERATURE.				RAIN.				SNOW.				RATES AND MATERIALS.							
	Temperature of Air	Pressure of Vapor	Humidity	Relative Air	Pressure of Vapor	Humidity	Relative Air	Wind Velocity	Maximum	Minimum	Difference	Wind Velocity	Maximum	Minimum	Difference	Wind Velocity	Maximum	Minimum	Difference	Appreciable	Depth in inches.	Appreciable	Depth in inches.	Appreciable	Depth in hours.	Appreciable	Depth in hours.	Appreciable
1	60.83	.314	60	.588	20.228	0.48	N 36° W	12.98	71.2	53.4	17.8	**	***	***	***	***	***	***	***	***	***	***	***	***	***	***	***	
2	63.42	.294	50	.594	0.17	294	West	3.40	5.82	75.5	54.8	20.7	**	***	***	***	***	***	***	***	***	***	***	***	***	***	***	
3	57.93	.439	91	.002	28.562	1.00	S 30° W	5.37	5.66	71.8	50.2	21.6	**	***	***	***	***	***	***	***	***	***	***	***	***	***	***	
4	60.95	.423	78	.114	28.631	0.78	N 36° W	12.06	12.13	66.0	57.6	9.0	0.375	5.0	**	***	***	***	***	***	***	***	***	***	***	***	***	***
5	57.85	.358	74	.195	28.857	0.63	N 86° E	4.81	7.13	66.0	54.4	11.6	* 0.63	1.0	**	***	***	***	***	***	***	***	***	***	***	***	***	***
6	58.08	.365	76	.138	28.773	0.67	N 72° W	5.40	6.65	71.0	53.3	17.7	.045	2.5	**	***	***	***	***	***	***	***	***	***	***	***	***	***
7	56.18	.344	76	.197	28.853	0.85	N 74° W	15.95	16.84	62.5	49.2	13.3	.015	2.5	**	***	***	***	***	***	***	***	***	***	***	***	***	***
8	53.32	.277	59	.411	20.134	0.48	N 41° W	20.44	20.61	63.0	52.0	18.0	**	***	***	***	***	***	***	***	***	***	***	***	***	***	***	
9	67.15	.393	60	.538	145	0.25	N 39° W	14.56	14.71	68.0	49.2	18.8	**	***	***	***	***	***	***	***	***	***	***	***	***	***	***	
10	62.42	.371	65	.581	21.0	0.63	N 74° W	4.83	8.27	80.4	51.5	28.9	**	***	***	***	***	***	***	***	***	***	***	***	***	***	***	
11	63.85	.299	53	.607	.308	0.40	N 59° E	4.33	5.54	74.0	51.8	22.2	**	***	***	***	***	***	***	***	***	***	***	***	***	***	***	
12	65.63	.395	64	.569	.308	0.40	N 59° E	2.66	4.10	70.8	53.0	17.8	**	***	***	***	***	***	***	***	***	***	***	***	***	***	***	
13	64.95	.484	78	.589	.175	104	0.62	N 37° W	5.68	75.8	52.2	26.0	0.40	2.7	**	***	***	***	***	***	***	***	***	***	***	***	***	***
14	60.70	.400	75	.703	.303	0.70	N 10° W	7.79	7.86	66.5	54.0	12.5	**	***	***	***	***	***	***	***	***	***	***	***	***	***	***	
15	64.46	.463	76	.504	.041	0.77	S 79° E	3.91	4.35	72.8	56.4	19.6	**	***	***	***	***	***	***	***	***	***	***	***	***	***	***	
16	67.17	.447	75	.504	.319	0.58	N 15° W	5.23	6.32	74.0	57.2	16.8	.475	2.0	**	***	***	***	***	***	***	***	***	***	***	***	***	***
17	64.58	.463	76	.504	.319	0.58	N 15° W	5.38	6.40	76.0	55.0	21.8	* 0.5	0.5	**	***	***	***	***	***	***	***	***	***	***	***	***	
18	63.62	.482	81	.461	.403	0.28	N 25° W	5.01	6.30	76.8	55.0	21.8	* 0.5	0.5	**	***	***	***	***	***	***	***	***	***	***	***	***	
19	66.03	.545	88	.654	.403	0.29	N 50° E	7.72	7.98	61.0	50.0	2.0	* .005	3.1	**	***	***	***	***	***	***	***	***	***	***	***	***	
20	58.55	.434	64	.826	.493	0.40	N 79° E	4.35	5.24	69.2	55.5	13.7	**	***	***	***	***	***	***	***	***	***	***	***	***	***	***	
21	60.38	.333	64	.814	.982	0.23	S 13° W	1.90	4.23	75.0	49.5	25.5	**	***	***	***	***	***	***	***	***	***	***	***	***	***	***	
22	68.75	.432	73	.504	.305	0.87	S 87° E	3.56	4.40	79.8	53.0	26.8	**	***	***	***	***	***	***	***	***	***	***	***	***	***	***	
23	68.48	.512	74	.774	.262	0.97	S 14° W	3.98	4.86	78.2	58.0	20.2	**	***	***	***	***	***	***	***	***	***	***	***	***	***	***	
24	68.80	.541	73	.774	.184	0.77	N 24° W	3.76	4.70	76.8	65.4	11.4	.003	4.0	**	***	***	***	***	***	***	***	***	***	***	***	***	***
25	63.68	.381	65	.766	.385	0.13	S 84° E	5.29	5.55	69.8	60.2	9.6	**	***	***	***	***	***	***	***	***	***	***	***	***	***	***	
26	71.23	.657	73	.517	.296	0.57	S 16° W	7.02	7.96	81.6	54.0	27.6	**	***	***	***	***	***	***	***	***	***	***	***	***	***	***	
27	69.40	.624	86	.505	.286	0.90	S 91° W	7.70	8.18	68.4	60.6	10.6	.750	7.0	**	***	***	***	***	***	***	***	***	***	***	***	***	***
28	65.70	.371	58	.554	.201	0.63	N 72° W	6.63	8.56	78.8	60.0	18.8	**	***	***	***	***	***	***	***	***	***	***	***	***	***	***	
29	63.16	0.414	71	29.498	0.58	N 41° W	3.13	7.61	-7.58	55.33	12.21	2.136	31.8	**	***	***	***	***	***	***	***	***	***	***	***	***	***	
30	63.16	0.414	71	29.498	0.58	N 41° W	3.13	7.61	-7.58	55.33	12.21	2.136	31.8	**	***	***	***	***	***	***	***	***	***	***	***	***	***	

* Inappreciable.

* Inappreciable.

GENERAL METEOROLOGICAL ABSTRACT.—JULY, 1860.

TORONTO METEOROLOGICAL RESULTS.

Days.	DAILY MEANS.			WIND.			EXTREMES OF TEMPERATURE.			RAIN.			SNOW.			
	Temperature of the Air.	Humidity.	Pressure of Vapor.	Clouded Sky.	Pressure of Vapor.	Wind Velocity.	Maxim.	Minim.	Velocity.	Depth in inches.	Approximate duration in hours.	Depth in inches.	Approximate duration in hours.	Depth in inches.	Approximate duration in hours.	
1	65.27	0.456	.73	29.656	.29	.199	0.53	N 55° E	6.60	69.2	59°	10.0	
2	63.67	.498	.84	.495	.000	1.00	N 56° E	4.01	68.2	60.0	8.2	11.5	2.0	
3	61.98	.499	.90	.518	1.00	N 53° E	7.53	69.4	60.5	14.5	0.115	2.0	...	
4	61.57	.416	.77	.405	28.989	.80	N 66° E	5.03	7.52	72.0	54.7	17.3	.005	14.5
5	61.52	.348	.64	.610	29.292	.30	N 66° E	5.39	6.58	68.5	53.7	14.8
6	65.78	.408	.65	.611	.203	0.05	S 71° E	4.03	5.59	73.6	52.5	21.1
7	65.33	.499	.79	.364	28.865	.28	S 30° E	4.03	6.65	71.4	58.0	13.4	.480	7.0
8	65.67	.408	.79	.617	29.200	0.58	N 52° E	3.67	5.15	70.8	63.2	9.6
9	63.88	.314	.63	.793	.481	0.22	N 40° W	4.75	8.07	68.0	49.6	18.4
10	62.42	.348	.63	.818	.470	0.27	N 27° W	8.52	8.72	70.4	50.2	20.2
11	63.75	.441	.75	.747	.306	0.05	S 36° E	1.76	3.93	72.5	52.5	20.0
12	64.70	.446	.74	.673	.227	0.05	S 17° W	5.41	6.52	77.0	52.0	25.0
13	68.38	.482	.72	.493	.011	0.40	N 65° W	8.75	11.49	83.8	63.2	20.6
14	62.85	.417	.74	.654	.237	0.12	S 7° W	3.46	4.46	72.5	53.0	19.5	1.5
15	63.18	.521	.89	.525	.005	0.73	S 46° E	2.57	4.35	71.2	55.0	51.0	8.3
16	69.00	.565	.70	.486	28.901	0.47	S 80° W	7.89	8.93	88.0	62.7	25.3
17	69.58	.470	.66	.499	29.029	0.48	S 44° W	4.35	4.97	81.8	61.9	19.9
18	66.93	.442	.64	.324	28.884	0.32	N 53° W	14.70	15.47	76.0	62.2	13.8	.475	1.0
19	57.68	.323	.66	.335	.061	0.30	N 30° W	12.52	13.25	68.5	53.0	15.5	*	0.1
20	66.38	.500	.77	.460	.206	0.47	S 75° W	8.66	9.35	74.5	48.8	30.7
21	61.27	.346	.64	.442	.203	0.27	S 11° E	2.39	4.06	70.8	52.5	18.3
22	62.43	.343	.62	.679	.336	0.05	S 45° E	1.23	6.35	65.0	55.0	11.0	1.7
23	66.38	.323	.66	.335	.061	0.65	N 61° E	6.73	7.28	67.5	49.0	18.5	.655	5.0
24	63.25	.323	.57	.552	.206	0.43	S 11° E	2.39	4.06	78.6	60.0	12.8	.935	4.0
25	63.20	.417	.71	.614	.227	0.03	N 62° W	5.58	7.87	74.0	61.2	10.4	0.43	0.3
26	67.37	.535	.80	.538	.003	0.70	S 45° E	1.23	6.35	65.0	55.0	11.0	1.7
27	60.00	.330	.64	.735	.405	0.07	S 45° E	1.23	6.35	65.0	55.0	11.0	1.7
28	62.43	.343	.62	.679	.336	0.65	N 61° E	6.73	7.28	67.5	49.0	18.5	.655	5.0
29	66.38	.500	.77	.460	.206	0.53	N 39° W	4.07	7.27	74.0	61.2	12.8
30	61.27	.346	.64	.442	.203	0.18	S 57° W	1.90	2.93	74.0	54.8	19.2	0.43	0.3
31	63.92	.427	.72	.295	.564	0.43	N 60° W	2.15	7.29	72.99	55.85	7.15	4.336	4.84

* Inappreciable.

TORONTO METEOROLOGICAL RESULTS.

GENERAL METEOROLOGICAL ABSTRACT.—AUGUST, 1869.

Date	DAILY MEANS.			WIND.			EXTREMES OF TEMPERATURE.		RAIN.		SNOW.		RAIN AND MELTED SNOW			
	Atmospheric Pressure of Mercury.	Relative Humidity.	Temperature of the Air.	Clouded Sky.	Pressure of Dry Air.	Relative Humidity.	Mean Velocity Wind.	Velocity Wind.	Maximum.	Minimum.	Depth in inches.	Duration in hours.	Approximate depth in inches.	Duration in hours.	Approximate depth in inches.	Duration in hours.
1	35.98	0.353	66	20.738	29.405	0.30	S 10 W	68.2'	48.0	26.2
2	62.67	.461	81	.728	.267	0.25	N 81 E	1.47	4.16	71.0	40.8	21.2
3	65.08	.513	87	.529	28.986	0.87	S 25 E	4.58	68.5	60.0	8.5	0.125	2.5
4	71.65	.622	70	.633	29.011	0.33	N 74 W	7.79	8.85	33.2	62.3	20.4
5	67.83	.556	82	.606	0.46	0.13	N 81 E	4.01	4.62	73.8	50.8	14.0
6	67.83	.556	82	.606	0.46	0.13	S 88 E	2.25	2.36	75.4	58.5	16.9
7	73.82	.608	84	.464	28.706	0.80	S 51 W	2.74	4.39	94.6	63.2	19.4	*	0.6	0.6	0.6
8	72.08	.476	62	.456	28.986	0.30	N 83 W	3.56	6.16	87.9	67.0	20.0
9	67.87	.529	78	.485	28.956	0.40	S 43 W	3.56	4.27	76.8	62.1	14.7	.035	1.5
10	65.08	.463	72	.513	29.050	0.68	N 65 W	7.14	10.41	79.8	57.5	22.3
11	58.42	.359	49	.691	.351	0.05	S 22 E	1.55	6.09	73.4	49.0	24.4
12	57.32	.346	75	.596	.250	0.63	N 9 W	2.60	6.39	73.5	49.4	24.4	.675	10.0
13	56.47	.366	79	.805	.439	0.20	N 21 W	4.56	6.33	64.4	30.0	14.4
14	59.23	.384	76	.879	.267	0.27	S 18 E	3.14	4.40	67.6	48.0	19.6
15	60.07	.403	78	.780	.377	0.57	S 6 W	3.48	4.34	68.0	51.0	17.0
16	66.03	.461	73	.663	.454	0.45	N 82 W	0.69	3.63	74.5	53.5	21.0
17	68.73	.355	73	.409	28.964	0.26	S 47 W	3.15	5.33	78.0	61.2	15.1
18	68.73	.355	73	.409	29.025	0.75	S 70 W	2.45	3.04	75.4	59.2	16.2	.303	3.7
19	68.85	.551	79	.576	.29	0.25	S 55 W	1.99	4.20	80.5	64.8	15.6
20	67.47	.529	79	.651	.132	0.10	S 86 E	4.66	73.5	50.2	16.3
21	65.98	.433	71	.671	.218	0.62	N 71 E	3.33	4.66	73.0	56.5	16.2
22	71.02	.601	80	.533	.283	0.77	N 15 W	3.96	6.37	77.6	61.2	16.4	.890	2.3
23	66.08	.557	87	.422	.28	0.82	N 53 W	11.58	11.67	71.8	61.6	10.2	.350	6.2
24	64.80	.478	78	.409	28.931	0.50	S 50 W	3.24	4.55	73.0	53.8	19.2
25	64.80	.478	78	.409	29.195	.063	N 81 W	8.28	9.20	68.4	54.0	14.4	* .062	1.0
26	58.80	.365	74	.560	.297	0.23	S 33 W	2.96	4.13	67.0	48.8	20.2
27	56.67	.306	67	.603	.250	0.05	S 20 W	3.70	4.26	69.5	52.0	17.5
28	60.58	.358	68	.648	.401	0.42	S 49 W	4.08	5.82	73.4	49.5	23.9	.375	3.2
29	64.83	.489	79	.401	28.912	0.40	N 63 W	6.93	8.29	75.5	57.0	16.5	.013	1.8
30	62.78	.403	72	.399	28.496	0.40	N 70 W	1.83	5.80	73.73	56.26	17.46	3.405	40.4
31	64.46	0.463	76	.29582	2.119	0.43	N 70 W	1.83	5.80	73.73	56.26	17.46	3.405	40.4

* Inappreciable.

DAY	DAILY MEANS.				WIND.				EXTREMES of TEMPERATURE.				RAIN.				SNOW.				RAINS AND METEOR. SNO.				
	Relative Humidity.	Pressure of Vapour.	Temperature of Air.	Pressure of Dew-point.	Gloomy Sky.	Wind Velocity.	Wind Velocity.	Wind Velocity.	Max. Temp.	Min. Temp.	Difference.	Max. Wind.	Min. Wind.	Difference.	Max. Rain.	Min. Rain.	Difference.	Max. Depth.	Min. Depth.	Difference.	Max. Depth.	Min. Depth.	Difference.		
30	64.83	.363	73	.301	28.302	0.42	5.49	W	4.98	4.82	.13	73.4	49.3	23.9	.362	3.2379	3.2		
31	62.78	.403	72	.399	28.396	0.40	5.63	W	6.93	8.20	73.5	57.0	16.5	.013	1.8013	1.8			
	64.46	0.463	76	.29	582	2	.19	0.43	N 70° W	1.83	5.80	73.5	56.2	17.2	46	3.405	40.4	3.405	40.4	
1	54.43	0.255	62	29.650	29.395	0.05	N 36° W	7.86	8.06	61.2	50.8	13.4	
2	59.47	.386	76	.881	.405	0.00	S 27° W	1.73	2.64	65.8	45.2	20.6	
3	65.07	.532	84	.293	0.83	S 23° E	4.65	4.93	66.6	54.0	20.8	
4	69.75	.617	85	.674	.036	0.85	N 48° E	2.63	2.93	72.2	54.0	18.2	*	0.2	
5	66.15	.429	67	.725	.296	0.35	N 35° W	2.21	4.45	75.8	62.6	13.2	*	*	
6	76.57	.492	75	.592	.100	0.77	N 43° E	1.45	4.33	74.2	64.5	9.7	
7	54.13	.360	83	.683	.223	0.82	N 9° E	1.37	3.45	72.5	55.8	16.7	
8	58.93	.598	71	.654	.590	0.60	S 64° W	4.51	7.00	66.0	52.2	13.8	
9	49.60	.284	77	.522	.238	0.38	N 59° W	8.39	10.32	62.8	46.9	16.8	
10	46.37	.191	62	.753	.422	0.42	S 39° W	10.47	10.55	52.8	39.0	13.8	
11	54.00	.270	66	.861	.582	0.07	N 67° W	3.80	3.91	47.0	38.8	28.2	
12	59.62	.378	76	.867	.489	0.39	S 3° W	3.59	3.67	70.8	45.7	26.1	
13	63.92	.473	79	.712	.239	0.22	S 18° W	5.64	6.44	72.0	49.2	22.8	
14	57.85	.386	80	.616	.236	0.30	S 80° W	3.32	4.92	62.8	46.9	16.8	
15	55.98	.557	78	.654	.238	0.33	S 79° E	1.63	1.87	68.0	51.5	16.5	
16	55.07	.372	80	.703	.331	0.63	S 79° E	0.15	1.59	65.2	48.9	16.3	
17	60.98	.462	86	.579	.117	0.75	N 16° W	2.13	5.59	69.2	51.2	18.0	
18	53.42	.363	88	.394	.031	0.68	N 73° W	6.20	7.76	60.2	55.8	4.4	
19	47.80	.239	74	.526	.297	0.25	S 80° W	7.96	8.46	39.4	41.6	17.8	*	0.2
20	55.98	.321	73	.511	.190	0.55	S 60° W	7.36	8.77	66.8	40.6	26.2	
21	58.68	.384	78	.367	.28	0.95	N 81° E	4.21	4.57	57.8	39.2	18.6	
22	53.32	.273	69	.343	.20	0.64	S 86° W	5.72	6.20	63.0	50.0	13.0	
23	47.85	.204	66	.775	.671	0.53	S 76° W	11.82	12.29	64.0	50.8	13.2	
24	50.90	.244	79	.763	.525	0.80	N 74° W	3.03	4.70	52.0	41.4	10.6	
25	42.47	.175	64	.800	.715	0.43	N 44° W	10.49	10.68	52.0	40.2	11.8	
26	39.90	.151	61	.30	.961	.910	0.58	S 12° W	1.00	1.88	45.2	35.2	10.0	
27	30...	
28	55.34	0.342	74	29.673	29.332	0.48	N 71° W	2.63	5.75	3.12	4.29	15.83	1.950	36.2

* Imprecise.

GENERAL METEOROLOGICAL ABSTRACT.—SEPTEMBER, 1860.

TORONTO METEOROLOGICAL RESULTS.

GENERAL METEOROLOGICAL ABSTRACT.—OCTOBER, 1860.

Days	DAILY MEANS.			WIND.			Extremes of Temperature.		RAIN.		SNOW.		RAINS AND MELTED SNOW.		
	Temperature of Air,	Pressure of Air,	Humidity,	Pressure of Water,	Humidity,	Wind Velocity,	Clouded Days,	Minimum	Maximum	Depth in hours,	Approximate Depth in hours,	Depth in hours,	Approximate Depth in hours,	Depth in hours,	
1	47.20	0.301	92	29.672	.454	1.00	N 44 E	2.33	45.8	37.2	12.6	0.325	7.5		
2	50.25	.307	85	.766	.526	1.00	N 27 W	1.11	54.2	46.4	7.8	.55	.325		
3	53.93	.320	77	.846	.580	2.19	S 84 E	4.88	59.2	49.4	9.8	.040	6.5		
4	54.60	.380	59	.671	.291	1.00	S 63 E	1.06	1.14	59.0	50.2	.8.8	.005	3.5	
5	52.97	.306	76	.722	.416	0.43	N 29 W	8.42	9.08	64.0	52.0	12.0	.005	3.5	
6	37.77	.176	73	.870	.693	0.02	N 78 W	0.87	4.20	53.0	20.8	23.2			
7							N 81 E	1.31	2.78	53.2	33.8	19.4			
8	46.68	.227	70	.335	.301	.007	N 46 W	17.87	18.13	52.6	45.0	7.6			
9	42.97	.224	80	.301	.017	.97	N 79 W	5.42	5.51	48.0	38.0	10.0	*	.02	
10	51.80	.303	77	.144	.28	.841	S 39 W	10.02	11.28	59.2	40.2	19.0			
11	42.27	.178	65	.430	.20	.233	N 59 W	11.07	11.38	51.5	45.7	5.8			
12	37.45	.158	71	.711	.533	.042	N 86 W	7.01	7.27	45.8	28.4	17.4	*	.05	
13	39.33	.176	73	.821	.615	.042	N 39 W	5.57	5.37	46.2	30.5	15.7	*	.02	
14							N 24 E	4.23	6.27	48.5	38.0	10.5			
15	41.57	.198	75	.738	.560	.063	S 82 W	4.75	6.38	49.2	37.5	5.0			
16	47.98	.271	79	.686	.316	.063	S 75 W	3.94	10.29	55.2	35.0	20.2	.190	4.8	
17	43.42	.199	71	.858	.639	.068	N 7 E	10.69	10.83	49.0	40.2	8.8			
18	44.20	.229	79	.961	.732	.066	N 20 E	3.80	3.91	43.0	20.0	10.0			
19	47.08	.261	81	.891	.630	.062	N 20 E	4.29	6.23	63.0	39.0	14.0			
20	47.50	.246	75	.798	.532	1.00	N 47 E	11.20	11.37	52.5	44.8	7.7			
21							N 70 E	12.08	12.13	51.5	44.5	7.0	.220	8.5	
22	49.20	.339	97	.615	.276	1.00	N 85 E	6.41	6.45	51.4	48.0	3.4			
23	49.18	.335	96	.490	.165	1.00	S 77 E	1.83	3.12	52.8	48.0	4.8	.095	9.3	
24	48.45	.300	87	.540	.210	0.62	S 75 W	2.72	3.10	55.0	43.4	11.6	.105	3.1	
25	48.65	.294	86	.530	.296	0.73	N 70 W	1.98	5.33	55.1	38.8	16.3			
26	43.42	.223	80	.746	.523	.013	S 41 W	6.90	7.28	51.0	41.0	10.0			
27	42.70	.233	83	.910	.676	.077	S 74 E	11.10	11.66	48.2	33.8	14.4			
28							S 69 E	5.45	9.21	55.5	44.0	11.5	.225	7.5	
29	52.13	.341	47	.799	.458	.050	S 84 W	3.78	6.12	60.0	51.5	8.5			
30	55.55	.393	89	.743	.350	0.60	N 88 E	5.21	5.81	62.4	44.0	18.4	.010	0.5	
31	58.03	.415	87	.706	.291	1.00	N 33 W	0.63	4.84	68.0	52.0	16.0	*	.02	
41.25	0.272	81	26.671	29.399	0.70	N 9 W	2.00	6.93	53.65	41.58	12.06	1.618	66.6	*	0.2
															1.618 36.8

Inappreciable.

TORONTO METEOROLOGICAL RESULTS.

GENERAL METEOROLOGICAL ABSTRACT.—NOVEMBER, 1860.

* Inapplicable.

TORONTO METEOROLOGICAL RESULTS.

GENERAL METEOROLOGICAL ABSTRACT.—DECEMBER, 1860.

Date	DAILY MEANS.			WIND.			EXTREMES OF TEMPERATURE.		RAIN.			SNOW.			RAIN AND MELTED SNOW.	
	Temperature of Air	Humidity	Barometer	Pressure of Vapor	Clouded Sky	Wind Velocity	Minimum	Maximum	Difference	Depth in inches	Approximate Depth in inches	Depth in inches	Approximate Depth in inches	Approximate Duration in hours	Approximate Duration in hours	
1	23.28	0.01	81	29.242	29.140	1.00	N 53° W	W 20.47	25.2	21.2	4.0	0.1	3.0	0.010
2	31.22	.155	88	.575	.420	1.00	N 75° W	W 14.65	14.75	30.8	21.8	9.0	...	*	0.2	
3	28.38	.135	87	.488	.363	1.00	N 78° W	W 5.24	6.76	33.0	28.8	4.2	...	*	3.0	
4	5	23.70	90	.492	.287	0.88	N 53° W	W 7.69	7.89	31.8	28.0	3.8	...	*	7.0	
5	26.25	.129	90	.307	.178	0.82	S 44° W	W 6.04	6.36	30.4	23.0	7.4	...	*	0.2	
6	7	30.32	83	.574	.435	0.52	S 31° W	W 4.96	5.53	36.0	23.7	12.3	...	*	8.8	
8	24.92	.117	87	.852	.735	0.83	S 1° W	W 3.60	5.83	30.0	24.0	6.0	
9	10	31.20	84	.165	.94	.239	S 68° W	W 2.95	5.56	35.2	24.2	11.0	...	1.5	2.0	
11	20.75	.084	74	.501	.417	0.70	S 82° W	W 11.37	14.18	29.0	19.5	9.5	...	4.5	12.0	
12	25.20	.108	79	.174	.096	0.82	S 69° W	W 18.75	19.04	25.0	13.0	15.0	...	*	2.0	
13	11.25	.063	84	.580	.317	0.12	S 68° W	W 11.73	12.75	17.6	12.5	4.1	...	0.1	0.5	
14	1.08	.037	79	.30.189	.30.152	0.18	S 15° W	W 5.30	5.63	9.0	7.0	16.0	...	0.2	3.0	
15	13.20	.061	78	.29.989	.29.928	0.60	S 56° W	W 1.45	6.18	24.0	1.3	22.7	
16	21.38	.105	81	.30.049	.34.4	1.00	S 60° W	W 8.47	8.52	34.5	11.0	23.5	
17	24.92	.119	87	.30.029	.31.0	1.00	S 28° E	W 6.89	7.89	26.0	21.4	4.6	...	*	1.0	
18	19	35.45	84	.196	.29.434	.238	S 84° E	W 8.78	9.88	37.8	23.5	14.3	1.265	* 1.0	1.0	
20	37.05	.178	80	.28.997	.28.820	0.66	S 89° W	W 11.06	13.23	39.0	33.2	3.8	...	0.47	3.5	
21	28.27	.136	87	.20.236	.29.100	0.88	S 9° W	W 8.34	10.17	31.2	27.5	3.7	...	5.0	.500	
22	21.15	.094	82	.490	.396	0.93	S 86° W	W 10.02	10.48	25.8	21.4	4.4	...	*	1.0	
23	18.30	.084	84	.56	.576	0.75	S 82° W	W 5.58	5.59	22.5	10.4	12.1	
24	25	11.0	85	.30.058	.30.068	1.00	S 61° W	W 6.49	7.12	24.4	13.8	10.6	...	*	1.0	
26	23.88	.110	86	.30.189	.30.068	1.00	S 42° W	W 10.23	10.27	28.0	19.0	9.0	...	*	1.0	
27	25.80	.121	86	.30.170	.30.069	0.95	S 66° E	W 9.16	10.64	28.8	23.5	5.3	
28	21.70	.101	86	.29.880	.29.730	1.00	S 20° E	W 9.21	13.64	31.2	18.2	16.0	...	0.50	4.0	
29	31.03	.150	86	.30.062	.30.062	0.83	S 13° W	W 10.59	11.01	25.0	16.0	9.0	...	0.2	0.5	
30	31	17.82	.076	.79	.30.062	.30.062	S 66° W	W 10.91	11.05	25.5	15.0	10.5	...	*	0.1	
31	24.03	0.115	84	29.667	29.551	0.83	N 62° W	W 4.66	10.14	28.79	19.25	9.54	1.362	24.0	13.5	
														68.4	2.712	
															92.4	

* Inappreciable.

TORONTO METEOROLOGICAL RESULTS.

GENERAL METEOROLOGICAL ABSTRACT.—JANUARY, 1861.

Days	DAILY MEANS.		WIND.				Extremes of Temperature.		RAIN.		SNOW.		Rain and Melted Snow.	
	Barometeric Pressure of Atmosphere,	Relative Humidity,	Pressure Atm. of Clouded Sky,	Velocity of Wind,	Difference in Mean Velocity,	Maximum Wind.	Difference in Minimum Wind.	Hours of Apprehension in hours,	Depth in inches,	Hours of Apprehension in hours,	Depth in inches,	Hours of Apprehension in hours,	Depth in inches,	Hours of Apprehension in hours,
30	21.32	.076	.79	30.062	.986	.083	S 73 W 10.79 11.01	25.0	16.0	9.0	* .3	3.0
31	17.82	.140	.88	29.530	.96	.083	S 66 W 10.91 11.05	25.5	15.0	10.5010	3.0
	24.01	0.115	.84	29.667	29.551	.083	N 62° W 4.66 10.14	28.79	19.25	9.54	1.362	24.0	13.5	68.4
														2.712 92.4
1	27.60	0.160	.72	29.781	.97	S 62° W	8.80 Miles.	36.2	15.0	12.2
2	28.75	.140	.88	29.708	.97	S 29 W	7.03 Miles.	23.0	7.0
3	15.12	.075	.86	29.635	.908	S 17 W	8.15 Miles.	33.8	18.0	12.0
4	17.17	.081	.84	29.703	.622	S 71 W	9.91 Miles.	11.3	22.2	9.3
5	19.98	.089	.85	29.854	.765	S 66 W	9.98 Miles.	9.35	24.0	14.0	10.0
6	7	23.00	...	118 94	...	S 81 E	3.74 Miles.	5.35	32.0	18.0	11.9
8	21.42	103	90	716	.417	S 21 E	5.66 Miles.	7.60	23.0	6.4	0.650	2.0	3.0	...
9	22.43	107	89	419	.312	S 95 N 12 W	3.03 Miles.	3.20	24.4	17.0	7.5
10	11.63	.071	.97	4.55	.386	S 47 W	12.42 Miles.	13.83	23.2	15.6	7.6
11	0.12	.043	.54	601	.558	1.00 N 18 E	11.87 Miles.	12.71	24.8	-4.8	8.8
12	-4.15	.032	.80	.904	.872	0.28 N 16 W	7.16 Miles.	7.25	-1.2	-7.0	5.8
13	S 64 E	1.93 Miles.	14.78	14.0	-11.2	26.2
14	22.67	113	90	718	.606	S 92 N 16 W	5.86 Miles.	6.55	27.3	8.0	19.3
15	27.38	140	94	551	.411	1.00 N 63 E	13.75 Miles.	13.86	20.8	20.5	9.3	140	7.0	...
16	32.53	181	98	116	.293	1.00 N 10 E	1.25 Miles.	19	35.5	24.8	10.7	47.5
17	26.60	128	88	703	.29.575	1.00 N 49 W	6.29 Miles.	7.53	30.5	26.2	3.8
18	28.02	145	94	415	.271	1.00 N 39 E	4.41 Miles.	12.37	37.0	24.0	13.0
19	29.88	147	87	360	.213	1.00 N 16 W	16.31 Miles.	16.42	32.2	5.4
20	S 85 W	7.35 Miles.	7.62	23.5	20.4	3.1
21	12.45	S 73 W	5.21 Miles.	5.44	22.0	6.0	16.0
22	7.47	.056	.85	956	.890	0.02 S 21 E	2.60 Miles.	2.73	18.0	-2.6	20.6
23	21.32	.100	.83	30.176	.096	S 0.0 W	14.40 Miles.	30.0	6.7	23.3
24	30.38	.161	.95	29.530	.29.369	S 5 W	5.06 Miles.	15.37	33.8	24.2	9.6	0.20	1.5	...
25	15.03	.067	.75	699	.632	S 30 W	9.81 Miles.	9.88	24.2	14.7	9.5
26	11.25	.065	.86	7.796	.730	S 68 W	2.95 Miles.	3.25	21.0	1.7	19.3
27	25.37	112	82	584	...	S 32 W	6.77 Miles.	7.66	30.0	8.5	21.5
28	28.45	146	91	260	.115	S 77 W	9.07 Miles.	10.45	30.2	21.0	9.2
29	19.35	.093	.87	614	.522	S 67 W	13.97 Miles.	15.21	33.0	24.5	10.5
30	15.75	.078	.88	653	.575	S 60 W	7.80 Miles.	7.83	17.6	15.0	2.6
						N 86 W	29.632 Miles.	29.549	0.76	9.30	23.44 Miles.	12.21	0.685	14.5
														20.6 33.6
														2.745 108.1

* Inappreciable.

TORONTO METEOROLOGICAL RESULTS.

GENERAL METEOROLOGICAL ABSTRACT—FEBRUARY, 1861.

Days	DAILY MEANS.				WIND.				EXPOSED OF TEMPERATURE.				RAIN.				SNOW.				
	Relative Humidity		Pressure of Atmosphere		Relative Humidity		Velocity Wind		Maximun Temperature		Minimun Temperature		Difference in Hours.		Approximate Duration in hours.		Rain and Melted Snow.		Depth in Inches.		
	Response of Thermometer		Response of Barometer		Clouded Sky		Sunny		Maximun Velocity		Minimun Velocity		Depth in Inches.		Approximate Duration in hours.		Depth in Inches.		Approximate Duration in hours.		
1	20.50	0.111	93	29.462	29.351	0.90	N 76° E	8.00	35° 2	5° 0	2° 2	22.7	7.0	10.5	0.700	10.5		
2	26.55	.135	92	.372	.257	1.00	N 52° W	11.63	11.86	11.39	11.39	22.4	9.3	13.1	1.4		
3	26.83	.126	131	.89	.837	.066	S 85° W	5.30	7.19	12.1	12.1	19.3	
4	26.67	.126	664	.87	.837	.538	S 59° W	7.24	7.63	31.4	31.4	4.6	* 0.1	* 2.5	* 0.10	* 2.5		
5	24.55	.105	77	.240	.135	0.88	S 88° W	9.27	16.16	35.0	35.0	22.8	12.2	2.0	3.8	.200	3.8		
6	24.55	.105	77	.529	.409	0.75	N 44° W	19.07	19.76	9.0	9.0	18.8	8.0	5.1	.800	5.1		
7	-7.72	.030	81	.529	.306	0.97	N 92° W	3.91	6.39	8.2	8.2	-29.8	0.5	4.0	.650	4.0		
8	-5.23	.033	90	.30160	.20.067	1.00	N 60° W	3.91	6.39	8.2	8.2	-29.8	* 3.0	* 3.0	* 3.0	* 3.0		
9	18.10	.040	89	.21.960	.20.860	1.00	N 70° E	6.96	7.62	29.8	29.8	-2.6	32.4	
10	41.52	.250	95	.395	.145	.97	N 50° E	3.05	4.08	44.6	44.6	37.6	7.0	.300	7.0	
11	33.70	.168	86	.266	.069	0.88	S 57° W	12.19	13.39	39.0	39.0	34.0	5.0	
12	30.28	.145	86	.753	.607	0.55	N 55° W	9.38	12.37	34.2	34.2	29.6	4.6	
13	28.85	.131	82	.662	.531	1.00	N 63° E	19.85	26.06	32.0	32.0	25.0	7.0	.045	4.5	
14	32.65	.176	97	.108	.28.932	1.00	S 76° E	1.62	10.34	33.4	33.4	28.2	5.2	
15	31.27	.140	79	.214	.29.074	0.57	S 58° W	6.89	8.22	34.8	34.8	31.0	3.8	
16	27.25	.119	79	.505	.386	1.00	S 72° W	10.89	10.78	32.2	32.2	25.0	7.2	
17	27.25	.119	79	.505	.261	0.87	S 19° E	7.36	9.32	32.6	32.6	19.2	3.2	
18	27.50	.138	90	.399	.018	0.83	S 85° E	12.80	13.34	36.0	36.0	19.2	13.4	
19	21.40	.145	81	.163	.619	0.88	S 76° E	10.14	11.18	36.4	36.4	26.0	13.2	
20	21.23	.089	75	.553	.464	0.52	N 60° W	20.06	20.20	22.5	22.5	6.7		
21	21.02	.697	85	.716	.619	0.88	S 76° E	10.14	11.18	36.4	36.4	26.0	13.2	
22	29.60	.139	86	.176	.637	1.00	S 57° W	3.31	16.43	14.57	14.57	19.7	8.0	.430	7.1	* 0.5	* 0.5	.430	7.6		
23	29.67	.128	76	.734	.606	0.83	S 45° W	13.01	12.31	12.58	12.58	39.2	8.1	
24	36.15	.166	78	.847	.680	0.45	S 77° W	3.76	5.12	14.48	14.48	27.8	17.0	
25	37.32	.187	84	.655	.468	0.97	N 54° W	3.04	6.18	46.0	46.0	32.8	13.2	.010	2.5040	2.5
26	29.44	.29.544	84	0.83	N 77° W	3.86	10.58	32.37	18.54	13.83	13.83	8.15	21.1	29.7	93.6	3.785	114.7	..	Inappreciable.		

GENERAL METEOROLOGICAL ABSTRACT—MARCH, 1861.

DATA.	DAILY MEANS.				EXTREME OF TEMPERATURE.				RAIN.				SNOW.			
	Temperature of Air.		Humidity.		Wind.		Velocity of Wind.		Rain.		Depth in inches.		Snow.		Rain and Melted Snow.	
	Pressure of Air.	Vapour.	Relative of Humidity.	Temperature of Sky.	Clouded Sky.	Clear Sky.	Velocity.	Direction.	Quantity.	Velocity.	Quantity.	Hours.	Approximate Depth in hours.	Quantity.	Hours.	Approximate Depth in hours.
1 26.72	0.206	85	29.545	29.339	0.98	N 5° W	2.60	44.2	.34.4	9.8	0.007	2.0	...	0.007	2.0	
2 37.42	.209	93	32.9	1.06	N 35° W	0.48	1.81	43.2	9.0	0.80	2.0	...	0.80	2.0		
3 29.35	..	109	66	..	S 85° W	5.47	6.85	35.0	12.4	1.70	2.5	...	1.70	3.5		
4 19.53	.080	76	632	0.45	S 85° W	13.25	17.25	38.8	31.8	7.0	...	2.5	8.7	2.5		
5 17.47	.068	84	582	0.63	S 80° W	16.81	17.83	24.6	11.8	12.8	...	0.1	0.5	.010		
6 8.20	.055	84	30.055	0.28	N 55° W	21.27	22.62	25.0	18.2	6.8	...	0.2	5.0	.020		
7 33.12	1.49	77	29.482	0.00	S 74° E	1.72	6.65	24.4	-3	21.9	0.50	2.5	3.0	5.5		
8 34.63	1.62	81	31.3	0.97	N 44° E	2.45	9.79	42.6	9.3	33.3	0.50	2.5	3.0	5.5		
9 10	..	116	81	..	N 87° W	8.45	9.69	39.8	32.2	7.6	...	0.1	0.3	.010		
11 24.90	1.11	75	75	..	N 33° W	14.30	14.54	55.4	19.1	6.3		
12 27.40	1.11	74	639	.679	S 55° W	13.36	14.23	33.6	9.3	24.3	...	0.5	0.5	0.5		
13 21.35	.091	80	779	.528	S 10° E	8.38	11.61	32.0	24.8	7.2	...	1.0	5.0	1.0		
14 20.38	.088	79	825	.688	N 10° W	9.24	10.22	28.2	18.8	9.4	...	0.2	5.0	.020		
15 26.77	1.19	81	655	.738	N 45° E	2.96	5.18	26.0	17.0	9.0	...	0.2	1.5	...		
16 28.12	1.12	74	319	.539	S 11° W	3.37	3.88	33.8	14.8	9.0		
17	..	116	81	.197	N 66° W	13.30	15.81	44.0	22.2	21.8		
18	3.48	0.41	77	30.100	S 10° W	13.65	13.75	9.2	0.2	9.0		
19	12.77	0.69	75	29.555	S 35° E	6.76	6.76	9.8	-5.2	1.5		
20	22.20	0.98	81	662	S 22° N	4.49	4.87	5.29	19.4	3.5		
21	11.13	83	669	.563	S 83° E	6.62	7.96	11.16	27.0	12.6	...	2.0	11.0	...		
22	30.37	131	77	.609	S 53° N	3.34	9.47	10.02	32.2	20.5	11.7	...	0.5	.30		
23	34.47	1.78	88	.722	S 03° N	6.11	3.61	38.9	22.5	16.4		
24	..	100	88	.265	S 30° W	2.11	3.61	38.9	22.5	16.4		
25	36.18	..	85	..	N 75° W	14.07	15.55	31.5	23.4	8.1	...	1.0	21.0	8.7		
26	38.97	..	95	.658	N 45° E	4.22	5.65	39.9	25.8	14.1	...	4.0	4.0	4.0		
27	37.22	.227	84	.294	N 00° N	1.00	N 81° E	6.77	8.56	42.8	34.2	12.5468		
28	32.02	.190	84	.314	N 74° E	1.24	0.88	S 82° W	17.29	18.71	45.0	35.8	13.5865	
29	31.63	..	78	.544	N 47° E	0.95	N 77° W	3.24	8.73	38.4	25.4	13.0	...	0.2		
30	..	138	78	.220	N 41° W	5.62	11.12	43.8	32.4	11.4	27.5	11.5275		
31581	N 89° E	10.29	10.45	33.2	25.0	8.2	0.2	0.8	0.2	0.8		
	26.92	0.127	80	29.021	29.493	0.62	N 54° W	4.33	10.56	33.53	20.71	12.82	2.125	55.2	7.1	

* Inappreciable.

* Inappreciable.

TORONTO METEOROLOGICAL RESULTS.

GENERAL METEOROLOGICAL ABSTRACT.—APRIL, 1861.

Days.	DAILY MEANS.				WIND.		EXTREMES or TEMPERATURE.		RAIN.		SNOW.		RAIN AND MELTED SNOW.			
	Temperature. of Air.	Pressure of Vapour.	Humidity.	Barometer.	Wind Velocity, Miles.	Wind Velocity, Fathoms.	Maximum Temperature.	Minimum Temperature.	Approximate duration in hours.	Depth in inches.	Approximate duration in hours.	Depth in inches.	Approximate duration in hours.	Approximate duration in hours.		
1	32.38	0.162	.89	29.611	29.449	1.00	\$ 89 N 63 W	3.20	12.22	16.38	.53	*	0.5	14.5	.606 15.0	
2	32.65	0.150	.81	27.757	60.17	0.70	S 89 N 63 W	3.20	3.15	36.4	27.13	
3	32.06	0.148	.81	29.948	80.0	0.50	N 70 E	0.67	1.28	39.0	23.8	15.2	
4	34.28	0.156	.79	30.041	88.6	0.15	N 84 E	1.83	2.24	38.8	20.1	9.7	
5	37.80	0.126	.56	29.947	82.1	0.45	N 57 E	2.27	3.23	45.0	28.9	16.1	
6	41.47	0.151	.58	28.84	.733	0.45	N 80 E	12.95	12.98	46.8	34.8	12.0	
7	40.30	0.191	.75	27.75	...	0.72	N 83 E	13.02	13.20	43.5	36.0	7.5	
8	43.67	0.183	.56	63.69	.478	0.72	N 86 E	16.04	16.06	45.0	34.8	10.2	.015	1.0	.015 1.0	
9	44.13	0.147	.53	.739	.457	0.53	N 79 E	14.63	14.70	50.2	36.4	13.8	.015	0.5	.015 0.5	
10	47.88	0.201	.59	.633	.392	0.00	N 67 E	2.31	4.47	53.8	36.8	17.0	.072	4.5	.072 4.5	
11	47.83	0.287	.87	.436	.432	0.23	N 87 E	7.52	7.78	56.8	35.2	21.6	.45	
12	48.62	0.301	.88	.215	29.914	1.49	1.00	S 78 E	4.30	4.60	58.0	41.0	17.0	.515	6.5	.515 6.5
13	35.95	0.170	.80	26.461	...	0.95	N 76 W	3.09	5.78	60.8	42.5	18.7	*	0.5	...	
14	35.95	0.146	.62	.492	3.46	1.00	N 42 E	1.38	2.88	39.0	34.2	4.8	
15	38.50	0.125	.60	.260	1.35	0.32	N 38 E	15.99	16.76	44.5	32.0	11.9	
16	36.03	0.158	.86	.218	.660	0.10	N 21 E	4.54	5.40	35.2	27.6	7.6	
17	32.45	0.160	.74	.505	3.46	0.18	N 34 W	12.12	12.33	48.0	26.0	18.4	
18	36.27	0.191	.72	.587	.396	0.52	S 52 W	6.63	7.40	50.0	27.0	23.0	
19	35.95	0.170	.70	27.75	...	0.95	N 42 E	1.38	2.88	39.0	34.2	4.8	
20	41.30	0.207	.70	.631	26.461	0.95	N 42 E	12.54	13.57	43.9	32.0	11.9	
21	54.77	0.307	.70	.339	...	0.32	S 73 W	5.96	7.68	55.2	33.8	21.4	.025	1.0	.025 1.0	
22	49.02	0.304	.87	.490	1.86	0.87	N 68 E	2.10	4.32	51.8	41.3	25.2	0.4	
23	50.02	0.245	.71	.328	.083	0.58	S 63 W	11.99	12.10	62.3	41.8	20.5	.037	1.2	.037 1.2	
24	43.38	0.176	.61	.579	.403	0.42	N 69 W	10.19	10.83	52.2	30.0	12.2	0.2	
25	43.27	0.194	.69	.644	.450	0.52	S 89 E	9.19	9.50	49.4	32.0	17.4	
26	48.67	0.286	.82	.217	28.931	0.83	N 82 E	6.12	9.21	58.3	40.2	18.1	.615	6.2	.615 6.2	
27	28	51.03	0.266	.70	...	29.152	0.57	S 32 W	10.13	11.62	59.2	48.2	16.0
28	30	48.12	0.236	.70	.434	.198	S 40 W	3.41	5.56	66.2	37.4	28.8	
29	42.02	0.199	.73	29.564	29.365	0.61	N 35 E	2.31	8.90	49.71	33.33	14.36	1.619	29.3	6.9 22.0	
30	42.02	0.199	.73	29.564	29.365	0.61	N 35 E	2.31	8.90	49.71	33.33	14.36	1.619	29.3	6.9 22.0	
															51.3	

Inappreciable.

TORONTO METEOROLOGICAL RESULTS.

GENERAL METEOROLOGICAL ABSTRACT.—MAY, 1861.

• Inappreciable.

42.02	0.199	73	29.564	29.365	0.61	N 3° E	2.31	8.90	45.71	33° 35'	14° 36'	1.619	29.3	6.9	22.0	2.309	51.3
-------	-------	----	--------	--------	------	--------	------	------	-------	---------	---------	-------	------	-----	------	-------	------

Date.	DAILY MEANS.				WIND.				EXTREMES OF TEMPERATURE.				RAIN.		SNOW.		RAIN AND MELTED SNOW.	
	Temperature of Air.	Humidity.	Pressure of Atmosphere.	Wind Velocity.	Direction.	Velocity.	Direction.	Velocity.	Maximum Temperature.	Minimum Temperature.	Difference.	Depth of Rain.	Duration in hours.	Depth of Snow.	Duration in hours.	Depth of Melted Snow.	Duration in hours.	
1	34° 53'	0.143	70	29.668	29.525	0.37	S 37° W	5.20	21.32	41.0	32°.5	.5	0.5	1.0	0.050	1.0	
2	36.03	.116	69	.817	.671	0.37	N 43° W	5.20	7.33	45.4	28.0	17.4	
3	37.72	.125	.56	.661	.536	0.42	N 70° E	5.27	6.32	44.5	32.0	12.5	
4	42.57	.136	.51	.652	.516	0.00	N 37° W	4.44	4.44	40.8	32.0	17.8	
5	47.75	..	89	28.679	1.00	..	N 89° E	11.33	11.46	54.8	31.8	23.0	0.260	2.2	2.0	2.2	
6	49.52	..	75	28.872	28.607	0.98	S 22° E	4.04	15.96	53.2	43.0	10.2	1.845	10.3	1.815	10.3	
7	44.17	.206	.72	29.322	29.116	0.63	S 40° W	8.74	9.06	50.5	40.4	10.1	* 1.00	5.4100	5.5	
8	44.93	.224	.74	.585	.301	0.12	S 26° E	2.08	2.59	54.0	32.8	21.2	.. 1.0	* 1.0	
9	45.05	.231	.76	.369	.138	0.87	N 69° E	6.71	7.17	48.6	31.8	11.8	1.65	6.2165	6.2	
10	48.58	.210	.79	.416	.146	0.58	S 43° W	6.69	2.92	57.4	43.5	13.9	
11	47.82	..	89	29.6	S 61° E	1.70	3.07	60.2	42.8	15.4	
12	50.45	.250	.71	.407	.148	0.76	S 79° W	11.06	11.27	54.8	44.4	10.4	
13	48.90	.210	.73	.457	.148	0.38	N 75° W	3.72	7.16	60.2	37.8	22.4	1.70	3.0170	3.0	
14	44.33	.210	.73	.518	.310	0.00	S 54° W	1.14	2.47	60.9	41.3	19.6	
15	44.13	.208	.71	.372	.372	0.28	N 37° W	14.23	14.6	50.4	41.0	9.4	
16	50.53	..	69	S 52° E	7.21	7.54	48.4	40.0	8.4	* 0.9	* 0.9	
17	52.70	.188	.48	.694	.506	0.07	S 22° W	11.65	11.83	61.0	45.2	15.8	
18	52.45	.236	.59	.310	.575	0.00	S 54° W	1.14	2.47	60.9	41.3	19.6	
19	50.90	.249	.67	.779	.613	0.28	N 37° W	14.23	14.6	50.4	41.0	9.4	
20	56.82	.318	.68	.497	.179	0.86	S 77° E	4.10	4.44	59.8	37.8	22.0	
21	62.25	.419	.74	.389	.28.971	0.20	S 60° W	10.78	11.08	73.0	53.5	19.5	
22	47.33	..	76	S 87° E	1.79	7.10	65.2	51.0	14.2	.. 1.15	
23	48.00	.197	.60	.563	.29.855	0.60	N 45° W	10.28	11.50	60.5	40.4	20.1	* .080	4.7	* .080	4.7	
24	45.32	.215	.69	.777	.562	0.23	S 47° W	4.81	6.07	55.0	42.2	12.8	
25	51.62	.233	.69	.910	.677	0.00	S 33° W	4.22	4.61	61.4	33.0	28.4	
26	54.80	.286	.67	.868	.583	0.00	S 73° E	2.82	3.12	63.2	40.2	25.0	
27	47.50	0.232	69	29.545	29.313	0.49	N 47° W	3.60	9.17	55.63	40.04	15.65	3.380	47.9	0.5	1.0	3.430	48.9

• Inappreciable.

GENERAL METEOROLOGICAL ABSTRACT.—JUNE, 1861.

TORONTO METEOROLOGICAL RESULTS

Invaluable.

61° 29'	0.377	69	29.570	29.192	0.45	N 36° W	2.29	6.11	70.36	51.26	10.11	2.320	32.3	2.320	32.3
---------	-------	----	--------	--------	------	---------	------	------	-------	-------	-------	-------	------	-----	-----	-------	------

* Inappreciable.

GENERAL METEOROLOGICAL ABSTRACT.—JULY, 1861.

DAYS.	DAILY MEAN.			WIND.			EXTREMES OF TEMPERATURE.			RAIN.			SNOW.			RAIN AND MELTED SNOW.				
	Temperature of Air.	Humidity.	Pressure of Atmosphere.	Wind Velocity.	Wind Direction.	Wind Velocity.	Wind Velocity.	Wind Direction.	Wind Velocity.	Wind Velocity.	Wind Direction.	Wind Velocity.	Wind Velocity.	Wind Direction.	Wind Velocity.	Wind Velocity.	Wind Velocity.			
	Temperature of Water.	Humidity.	Pressure of Atmosphere.	Wind Velocity.	Wind Direction.	Wind Velocity.	Wind Velocity.	Wind Direction.	Wind Velocity.	Wind Velocity.	Wind Direction.	Wind Velocity.	Wind Velocity.	Wind Direction.	Wind Velocity.	Wind Velocity.	Wind Velocity.			
1	58° 23'	0.337	67	29.362	29.225	0.73	N 21° W	6.17	63.5	64.6	53.5	11.1			
2	54.47	.277	66	.657	.380	0.62	N 44° W	9.25	67.2	47.0	20.1			
3	65.90	.378	58	.690	.312	0.25	N 37° W	1.99	3.40	78.5	49.4			
4	67.53	.480	70	.682	.202	0.08	S 1° E	2.08	2.37	80.0	51.2	28.8			
5	69.77	.477	65	.589	.064	0.64	S 0.6° S	3.00	3.22	81.4	53.0	26.4			
6	69.93	.500	69	.564	.389	0.12	S 0.6° S	1.89	1.90	80.8	63.6	21.2			
7	73.85	.638	79	.427	.247	28	N 60° W	2.11	2.17	84.5	61.4	23.1	0.2	0.12			
8	72.05	.596	75	.319	.28	72.22	S 34° W	4.76	4.91	84.2	63.8	18.4	0.3	0.15		
9	74.35	.478	77	.371	.28	9.13	S 51° W	4.86	5.20	83.0	65.0	18.0	0.7	0.1		
10	74.47	.396	77	.421	.29	0.63	0.72	0.56	S 12° E	5.91	8.30	75.2	55.8	19.4	87.5	11.2		
11	57.98	.532	73	.589	.237	0.53	S 42° W	8.07	8.35	61.2	33.2	8.0	0.13913	2.0		
12	59.92	.390	76	.779	.389	0.72	S 19° W	0.45	1.94	69.6	54.0	17.6		
13	60.49	...	76	S 16° E	0.69	1.35	71.4	50.8	20.6		
14	60.75	...	76	S 38° W	0.63	0.96	65.2	51.6	13.6	4.5		
15	64.30	...	71	S 69° W	0.35	7.40	57.2	46.8	24.5	1.5		
16	65.87	...	71	0.17	S 21° E	1.95	2.14	72.4	47.0	25.4		
17	64.8	4.66	73	.563	.154	0.38	S 21° E	0.30	0.35	61.2	33.2	22.0	0.07	0.07	0.2		
18	67.28	.322	78	.306	28	0.04	N 73° S	2.58	3.90	4.58	77.2	53.2		
19	66.63	.521	80	.369	.75	0.75	S 19° W	1.58	3.60	76.0	60.4	15.6	4.00	5.5	5.5		
20	63.38	.429	72	.411	28	4.981	0.17	N 44° W	10.01	10.08	72.0	57.5	14.5	
21	60.77	...	75	S 30° W	3.65	4.50	69.8	52.0	17.8	0.10	0.5		
22	60.77	...	73	S 36° W	2.72	4.87	68.5	51.8	16.7	*	0.2	*	0.2		
23	65.87	3.54	57	.675	.321	0.52	N 1° E	4.03	4.79	75.8	54.0	21.8		
24	65.12	.511	80	.729	.218	0.26	S 31° E	1.63	1.88	73.2	55.5	17.7		
25	67.83	.433	67	.636	.243	0.78	S 7° E	3.92	4.15	77.0	52.4	24.6		
26	70.30	.577	78	.557	.28	9.80	1.00	S 17° W	3.66	4.12	78.0	63.4	24.6	2.5	
27	68.68	.558	80	.563	.29	0.65	0.90	S 83° E	4.39	4.57	79.0	65.0	14.0	0.08	2.5	0.85	2.5	
28	66.57	5.37	82	.450	28	0.55	...	S 13° E	7.10	9.35	74.7	64.5	10.2	.500	3.5	
29	66.57	5.58	73	.558	29.000	0.22	S 15° W	3.30	3.82	80.8	58.2	22.6	1.5	0.3	0.3	0.3	
30	71.20	.653	82	.506	28	8.83	0.65	S 26° W	1.85	3.51	81.4	60.8	14.6	.010	0.5	0.10	0.5
31	73.02	...	73	S 74° W	1.43	4.06	74.67	58.23	18.44	2.635	34.7	2.635	34.7
	63.37	0.407	73	29.551	29.084	0.56	

* Inappreciable.

TORONTO METEOROLOGICAL RESULTS.

GENERAL METEOROLOGICAL ABSTRACT.—AUGUST, 1861.

DAYS	DAILY MEANS.				WIND.				EXTREMES of TEMPERATURE.		RAIN.		SNOW.				
	Temperature of the Air.	Humidity.	Pressure of Vapour.	Pressure of Mercurial.	Clouded Sky.		Clear Sky.		Minimum.	Maximum.	Difference.	Depth in inches.	Approximate duration in hours.	Depth in inches.	Approximate duration in hours.		
					Pressure of Mercurial.	Humidity.	Velocity.	Velocity.									
1	71° 63	83	29.552	28.911	0.22	S 5° E	1.41	2.23	78.2	66.4	11.8		
2	72.46	82	.474	.28.825	0.08	N 34° E	1.70	4.37	81.8	62.4	19.4		
3	74.20	650	.78	.603	28.053	...	6.07	6.4	80.0	66.2	13.8	0.005	1.5		
4	73.17	646	.78	.484	28.828	0.35	N 84° W	0.87	3.26	82.0	67.0	15.0	
5	66.92	560	.78	.562	29.002	0.60	S 80° E	1.25	1.84	76.8	61.8	15.0	0.13	0.3	
6	62.65	516	.90	.623	1.07	1.00	N 56° E	4.78	4.95	66.2	61.6	4.6	.465	10.7	
7	65.18	476	.77	.612	1.35	1.00	N 74° E	4.28	4.47	69.5	61.0	8.5	0.13	0.5	
8	66.57	580	.89	.488	28.908	0.83	N 68° E	1.91	3.29	71.8	64.0	7.8	0.15	4.0	
9	69.15	525	.72	.434	28.909	0.68	S 29° W	2.84	6.87	83.4	59.2	24.2	
10	69.15	525	.72	.530	1.00	1.00	N 43° E	0.59	1.01	62.4	57.8	4.6	.760	12.5	
11	62.92	508	.84	.590	29.112	0.65	N 54° E	4.06	5.09	69.0	55.0	14.0	
12	66.77	425	.80	.689	.264	0.50	S 7° W	0.91	1.39	70.6	53.0	17.2	
13	69.40	386	.72	.854	.489	0.51	S 25° W	0.68	1.02	72.0	47.0	25.0	
14	61.95	391	.72	.842	.451	0.03	S 25° W	0.68	1.02	72.0	47.0	25.0	
15	64.70	450	.74	.727	.247	0.60	S 6° E	2.93	3.21	74.2	53.4	20.8	
16	66.18	507	.78	.721	.214	0.58	S 65° W	0.36	1.88	77.0	54.0	23.0	*	0.5	
17	66.18	507	.78	.721	.214	0.58	S 80° W	0.12	2.68	75.3	53.0	22.3	0.5	0.5	
18	64.13	469	.78	.730	.251	0.62	N 78° E	2.17	2.60	70.6	57.2	13.4	
19	64.13	469	.78	.730	.251	0.62	N 78° E	2.17	2.60	70.6	57.2	13.4	
20	64.23	424	.71	.855	.431	0.67	S 89° W	3.19	7.24	70.2	58.4	11.8	.575	4.0	
21	70.48	621	.83	.620	28.909	0.97	S 49° W	3.86	7.08	80.0	58.5	21.5	
22	61.70	451	.81	.571	.29.120	0.42	N 44° W	9.73	9.92	70.8	60.5	10.3	*	0.3	
23	63.15	371	.66	.741	.370	0.12	N 75° W	4.45	5.21	75.4	51.2	24.2	
24	65.18	455	.74	.788	.333	0.22	S 41° W	0.88	1.85	75.5	52.4	23.1	
25	67.92	560	.82	.699	.158	0.56	S 50° E	1.10	1.60	54.5	54.5	1.0	1.15	1.5	
26	68.03	.557	.81	.631	.031	0.68	S 21° W	1.82	1.98	70.6	62.0	14.6	.605	2.2	
27	66.07	.515	.85	.631	.086	0.55	S 23° W	1.87	2.33	74.8	58.4	16.4	1.16	0.8	
28	63.27	.473	.81	.607	.131	0.43	N 75° W	1.20	2.38	74.2	59.8	14.4	.035	0.2	
29	60.45	.354	.67	.659	.305	0.15	N 49° W	10.71	10.92	70.0	51.0	19.0	
30	59.40	.315	.63	.784	.469	0.38	N 31° W	1.85	3.34	69.8	40.8	20.0	
31	59.40	.315	.63	.784	.469	0.38	N 31° W	1.85	3.34	69.8	40.8	20.0	
32	63.48	0.495	.78	29.653	29.158	0.54	N S E	0.46	4.21	74.30	58.15	16.2	.953	43.3

* Inappreciable.

	65.48	0.495	78	29.653	29.158	0.54	N	S E	0.46	4.21	74.30	58.15	16.16	2.953	43.3	2.353	43.3
--	-------	-------	----	--------	--------	------	---	-----	------	------	-------	-------	-------	-------	------	-----	-----	-------	------

* Imprecise.

GENERAL METEOROLOGICAL ABSTRACT—SEPTEMBER, 1861.

DAY	DAILY MEANS.			WIND.			EXTREMES OF TEMPERATURE.			RAIN.			SNOW.			RAIN AND MELTED SNOW.		
	Temperature Atmer.	Vapor pressure of air.	Relative humidity.	Pressure of atmosphere.	Wind direction.	Wind velocity.	Wind effect.	Mean Velocity.	Maximum.	Minimum.	Difference.	Hours.	Depth in inches.	Approximate duration in hours.	Hours.	Depth in inches.	Approximate duration in hours.	Hours.
1	69.33	0.588	81	29.566	28.078	0.77	S 88 E	4.45	4.70	4.17	.52.2	11.0	*	0.2	...	*	0.2	
2	66.47	.472	70	.611	.29.139	0.25	S 41 W	6.38	8.38	78.0	.61.2	16.8	.002	0.2	...	0.2	0.2	
3	58.45	.346	71	.894	.458	0.47	N 64 W	8.08	10.21	78.8	.61.5	17.3	.070	0.2070	0.2	
4	61.28	.415	77	.588	.173	0.57	S 18 E	2.76	3.02	69.2	.52.2	13.6
5	63.87	.410	69	.418	.008	0.28	S 87 W	6.56	7.20	73.0	.50.2	19.0
6	55.47	.327	73	.788	.461	0.67	N 20 W	2.30	2.88	74.4	.50.0	14.4	*	0.1	...	*	0.1	0.1
7	57.63	.366	77	.638	.533	0.58	S 66 E	2.01	2.58	64.8	.47.2	17.6	*	0.1	...	*	0.1	0.1
8	59.58	.417	81	.782	.345	0.97	N 80 E	3.98	4.17	64.0	.33.2	10.8	.345	0.7	...	*	0.7	0.7
9	61.17	.491	90	.407	.28.316	0.77	N 9 W	2.47	3.55	63.4	.38.0	5.4	.030	1.0	...	*	0.30	1.0
10	60.73	.382	74	.560	.179	0.47	N 66 W	3.12	3.12	70.4	.36.8	13.6
11	63.33	.444	75	.683	.239	0.40	S 18 W	2.74	2.77	73.0	.49.0	24.0
12	57.47	.361	80	.608	.047	0.70	S 41 W	1.23	2.05	68.0	.38.5	9.5	.115	5.5	...	*	5.5	5.5
13	52.03	.327	84	.529	.415	0.52	S 27 W	2.55	2.92	75.0	.38.2	16.8
14	56.33	.332	73	.747	.415	0.82	N 78 E	1.87	2.30	63.0	.30.0	13.0
15	52.33	.328	74	.571	.244	0.62	N 15 E	3.49	4.29	63.0	.33.6	9.4	*	0.5	...	*	0.5	0.5
16	56.33	.332	73	.747	.415	0.82	N 78 E	1.87	2.30	63.0	.30.0	13.0
17	56.33	.328	74	.571	.244	0.62	N 15 E	3.49	4.29	63.0	.33.6	9.4	*	0.5	...	*	0.5	0.5
18	60.43	.421	81	.567	.090	0.16	S 19 W	4.16	5.85	68.0	.49.5	18.5
19	66.93	.459	69	.486	.027	0.82	S 62 W	4.20	7.23	74.8	.52.0	21.8	*	0.5	...	*	0.5	0.5
20	57.73	.421	88	.467	.046	1.00	N 44 E	7.32	8.20	61.1	.66.0	5.2	.895	12.0	...	*	12.0	12.0
21	52.03	.327	84	.529	.201	0.83	N 4 W	8.20	8.44	55.2	.50.6	4.6	0.43	5.0	...	*	0.43	5.0
22	56.33	.332	73	.747	.415	0.82	N 78 E	1.87	2.30	63.0	.30.0	13.0
23	56.03	.361	76	.568	.207	0.10	S 67 W	8.01	8.24	68.8	.47.2	21.6
24	56.00	.324	76	.525	.401	0.15	S 1 W	1.25	1.94	64.8	.45.5	19.3
25	61.15	.433	81	.603	.170	0.38	S 25 E	1.99	2.17	69.8	.47.8	22.0
26	62.72	.473	82	.378	.28.905	0.82	S 36 W	3.88	5.13	72.0	.59.8	12.2	.175	3.5	...	*	3.5	3.5
27	52.53	.348	88	.331	.28.883	1.00	N 44 W	5.07	7.06	59.2	.33.0	6.2	1.865	11.0	...	*	1.865	11.0
28	47.58	.263	80	.488	.29.225	0.60	S 85 W	7.48	7.75	56.5	.46.4	10.1	.020	0.6	...	*	0.20	0.6
29	48.95	.281	81	.30.085	.802	0.85	S 82 W	2.76	3.29	54.3	.37.1	9.5	*	0.37	9.5
30	48.95	.281	81	.30.085	.802	0.85	N 46 E	1.18	2.03	55.0	.46.4	8.6	*	0.37	8.6
	56.07	0.400	79	29.608	29.200	0.60	N 71 W	1.39	4.81	68.38	51.80	14.58	3.607	60.1	...	*	3.607	60.1

* Imprecise.

GENERAL METEOROLOGICAL ABSTRACT.—OCTOBER, 1861.

Days	DAILY MEANS.				WIND.				EXTREMES OF TEMPERATURE.				RAIN.				SNOW.				
	Temperature of the Air.	Vapour of the Air.	Humidity.	Pressure.	Wind.	Velocity.	Direction.	Wind.	Min.	Max.	Difference.	Wind.	Velocity.	Direction.	Wind.	Min.	Max.	Depth in inches.	Duration in hours.	Depth in inches.	Duration in hours.
1	58.25	0.341	84	29.587	29.546	0.83	N 80 E	58.4	44.0	14.2	
2	59.12	.442	88	54.0	0.68	0.65	S 42 W	2.53	2.63	1.1	
3	56.07	303	70	.702	0.38	0.52	N 52 W	5.23	7.16	65.2	11.2	0.018	1.5	
4	51.52	.559	94	.626	1.06	1.00	S 64 E	8.43	9.10	53.0	5.1	0.25	15.0	
5	55.25	.419	96	.434	0.15	1.00	S 40 E	2.49	3.00	59.4	12.4	0.12	0.5	
6	47.40	284	86	.755	0.57	0.92	S 23 W	5.94	7.32	55.5	51.1	4.4	4.15	7.5
7	50.28	.301	82	.861	.560	0.02	S 21 E	1.66	3.48	56.5	38.8	14.7	
8	52.22	.291	75	.927	0.17	0.92	S 86 E	1.57	4.37	58.0	43.5	14.5	
9	52.33	.327	84	.775	0.51	0.37	S 69 E	4.32	5.06	59.0	44.5	14.5	
10	52.75	.360	90	.453	0.63	0.43	S 72 W	2.15	3.76	60.6	41.2	16.4	17.5	
11	52.25	.235	77	.171	2.437	0.92	S 59 W	2.02	2.99	63.8	47.4	16.4	0.58	2.5	
12	45.25	13	73	.555	2.282	0.07	S 61 W	10.86	11.35	54.0	50.8	12.3	0.065	1.0	
13	50.12	.573	73	.555	2.75	0.07	S 51 W	5.74	6.73	67.4	55.5	18.5	0.05	0.3	
14	46.23	.229	74	.737	.528	0.07	S 69 E	2.54	4.02	55.2	31.9	20.2	
15	52.73	.362	90	.638	.275	1.00	S 84 E	2.23	2.49	57.3	45.5	11.8	
16	56.83	.381	88	.571	.160	0.88	N 50 E	2.97	4.07	62.0	50.0	12.0	0.80	2.1	
17	56.83	.407	88	.420	.065	1.00	N 63 E	3.49	4.29	62.0	7.0	0.46	
18	51.63	.329	82	.926	.420	.100	N 79 W	6.04	8.02	57.2	52.0	5.2	0.72	5.0	
19	51.63	27	73	.836	.639	0.58	N 55 W	6.39	7.07	52.4	40.5	11.9	
20	44.40	222	73	.847	.625	0.12	S 45 E	8.97	9.43	52.0	33.5	18.5	
21	51.03	.530	87	.476	.146	0.73	S 8 W	2.80	4.47	57.2	38.8	18.4	4.65	5.9	
22	40.75	198	77	.417	.219	0.58	S 80 W	6.63	10.29	47.4	43.8	3.6	0.43	1.0	
23	34.27	155	78	.899	.744	0.32	N 55 W	5.90	6.58	42.0	30.5	11.5	
24	45.77	.241	78	.926	.085	0.58	S 17 W	6.04	6.32	51.8	29.0	22.8	
25	44.22	.247	83	.836	.639	0.58	S 72 W	4.76	6.50	53.2	43.8	9.4	*	1.0	
26	38.52	176	74	.760	.614	0.02	S 34 E	1.56	4.88	46.2	31.4	14.8	
27	42.82	.227	81	.289	.002	0.20	S 75 E	2.20	3.71	47.0	31.0	16.0	16.8	10.5	
28	40.82	.229	89	.001	2.802	0.63	S 83 E	3.64	5.17	48.8	32.0	4.5	0.70	1.7	
29	43.48	.236	84	.608	.29372	0.87	N 55 W	3.52	4.08	49.0	35.4	13.6	
30	48.74	0.292	82	29.619	26.327	0.61	N 61 W	1.06	5.96	55.34	41.62	13.73	1.933	53.6	*	1.0	1.393	54.6			

Inappreciable.

48.74	0.292	82	29.619	29.327	0.61	N 61 W	1.06	5.96	5.34	41.62	13.73	1.983	53.6	*	1.0	1.983	54.6
-------	-------	----	--------	--------	------	--------	------	------	------	-------	-------	-------	------	---	-----	-------	------

* Imprecise.

GENERAL METEOROLOGICAL ABSTRACT—NOVEMBER, 1861.

TORONTO METEOROLOGICAL RESULTS.

23

Days.	Relative Air Pressure of Thermometer	Pressure of Atmosphere	Pressure of Clouded Sky	Pressure of Dry Air	DAILY MEANS.		WIND.		EXTREMES OF TEMPERATURE.		RAIN.		SNOW.		RAYS AND MELTED SNOW.		
					Presented	Maxim.	Min.	Wind.	Wind.	Difference.	Min.	Maxim.	Depth in inches.	Approximate in inches.	Depth in inches.	Approximate in inches.	
1	42.45	0.222	81	29.877	29.655	1.00	N 5 E	10.77	10.94	.44.2	36.0	5.2	0.415	5.5	...	415 5.5	
2	40.33	.235	94	.325	1.00	51 E	18.30	20.73	44.0	38.5	5.5	3.132	19.0	...	3.132 19.0		
3	879 W	7.83	8.25	44.5	37.0	7.5	3.195	6.0	...	3.195 6.0		
4	40.65	.206	83	.302	.996	0.95	875 W	6.85	6.85	46.5	36.0	10.5	.005	0.2	*	0.2	
5	46.07	.256	82	.228	28.973	0.78	843 W	2.46	3.82	52.4	38.6	13.8	
6	43.80	.219	77	.305	29.086	0.53	N 36 W	6.23	6.85	51.6	41.0	10.6	*	0.5	*	0.5	
7	38.25	1.64	73	.543	.379	0.38	N 39 W	2.89	3.13	46.0	33.6	12.4	
8	40.65	.269	82	.431	.222	1.00	N 57 W	0.66	1.42	45.2	31.4	13.8	*	0.4	*	0.4	
9	35.92	1.52	73	.570	.418	0.32	N 22 W	2.26	2.42	44.0	31.0	13.0	
10	S 51 E	7.73	9.14	48.8	38.4	20.4	.060	2.0	
11	42.82	.210	75	.540	.331	0.47	N 80 W	15.20	15.40	49.6	43.0	6.6	.020	1.0	...	0.20 1.0	
12	37.83	.179	79	.911	.732	0.72	S 45 E	1.37	8.56	46.2	29.5	16.7	.025	0.5	...	0.25 0.5	
13	41.73	.201	75	.648	.447	0.88	N 37 W	1.11	3.44	45.0	34.0	11.0	
14	37.38	1.58	69	.610	.452	0.95	N 24 W	5.34	6.43	45.0	36.0	9.0	
15	31.70	1.18	66	.473	.355	0.82	N 36 W	11.14	11.82	35.8	29.2	6.6	...	0.1	1.6	0.10 1.6	
16	32.37	1.43	78	.587	.444	0.82	N 37 W	18.92	19.10	35.0	27.6	7.4	...	0.1	3.0	0.10 3.0	
17	N 43 W	3.7	3.7	28.0	28.0	12.2	
18	33.25	1.44	75	.761	.617	0.02	N 17 W	9.11	9.38	41.0	23.0	18.0	
19	35.47	1.38	71	.890	.752	0.12	S 79 E	4.12	4.89	30.4	25.0	13.8	
20	38.18	.182	77	.619	.497	0.98	S 82 E	5.18	5.30	42.2	30.0	12.2	.127	4.8	...	1.27 4.8	
21	38.50	2.19	93	.805	.587	1.00	S 77 E	0.45	0.60	43.8	37.0	6.8	*	1.5	
22	39.85	.204	83	.453	.249	1.00	S 72 E	0.85	1.42	44.0	34.4	9.6	*.230	3.2	...	230 3.2	
23	35.20	.167	81	.158	.28.391	1.00	S 39 W	6.57	7.35	37.4	35.5	3.9	...	0.3	3.1	0.30 3.1	
24	S 58 W	1.47	1.64	34.0	30.0	4.0	...	*	0.5	0.5	
25	30.67	1.48	86	.344	.29.196	0.55	N 16 W	0.47	0.48	37.0	23.3	13.7	
26	32.18	.143	79	.685	.541	0.20	S 57 W	1.84	2.31	40.8	35.0	15.8	
27	35.02	.166	81	.475	.309	0.98	S 32 W	7.32	11.33	38.2	28.6	9.6	.055	3.0	*	1.27 4.2	
28	30.43	1.39	81	.602	.464	0.95	S 30 E	2.41	5.52	37.0	27.8	9.2	...	2.0	5.0	.000 5.0	
29	35.73	.177	84	.334	.157	0.93	S 78 W	6.84	7.10	39.0	39.2	9.8	*	0.5	*	0.5	
30	31.33	.137	77	.486	.330	0.87	S 81 W	8.34	8.42	34.0	30.5	3.5	...	0.5	2.8	.050 2.8	
31	37.14	0.178	79	29.537	29.359	0.74	N 46 W	1.94	7.44	42.39	31.99	10.40	4.294	48.1	3.2	17.4	4.614 65.5

* Imprecise.

GENERAL METEOROLOGICAL ABSTRACT.—DECEMBER, 1861.

TORONTO METEOROLOGICAL RESULTS.

Days	DAILY MEANS.				WIND.				EXTREMES OF TEMPERATURE.				RAIN.				SNOW.				RAYS AND MELTED SNOW.						
	Temperature of the Air	Humidity	Pressure of Atmosphere	Wind Velocity	Pressure of Atmosphere	Velocity of Wind	Wind Direction	Wind Velocity	Maximum	Minimum	Mean Velocity	Pressure of Atmosphere	Velocity of Wind	Wind Direction	Wind Velocity	Depth in inches	Duration in hours	Apparatus in hours	Depth in inches	Duration in hours	Apparatus in hours	Depth in inches	Duration in hours	Apparatus in hours	Depth in inches	Duration in hours	Apparatus in hours
	°	%	mm.	miles	mm.	miles		miles	°F.	°F.	°F.	mm.	mm.		miles		hours	hours	hours	hours	hours	hours	hours	hours	hours	hours	
1	19.98	0.090	33	29.488	0.25	N 75 W	1.87	26.5	34.7	3.0	10.0	0.300	10.0														
2	14.25	0.068	83	.827	.758	N 23 W	5.47	6.41	3.84	14.2	14.2	* * *	0.5	0.5													
3	28.63	.133	85	.714	.581	S 42 W	9.98	10.23	34.4	8.0	26.4																
4	34.63	.156	77	.861	.705	S 72 E	2.31	2.27	40.8	26.3	14.5	*	1.5														
5	41.92	.297	80	.697	.400	S 25 E	0.90	1.76	46.4	34.4	12.0																
6	47.92	.269	89	.674	.375	S 38 W	6.24	6.37	52.0	39.5	12.5																
7	35.5	.279	94	.353	.256	S 66 E	0.42	0.77	55.1	45.0	10.1																
8	44.60	.329	95	.331	.092	1.00	N 73 W	10.30	14.03	55.2	42.4	12.8															
9	48.45	.329	96	.692	.64	N 49 W	9.70	10.33	32.2	27.3	4.9																
10	26.23	.092	64	.980	.888	N 49 W	6.41	6.50	35.6	18.8	16.8																
11	30.68	.133	78	.501	.301	S 54 W	0.50	0.55	4.29	4.35	41.6	30.0	11.6														
12	35.15	.147	71	.501	.208	S 65 W	0.60	0.76	5.79	7.04	43.8	32.6	11.2														
13	37.75	.156	70	.29.761	.605	S 76 W	0.60	0.76	5.79	7.04	43.8	32.6	11.2														
14	42.45	.195	71	.391	.396	S 66 E	5.98	6.40	38.0	23.0	9.8																
15	35.78	.160	75	.702	.624	S 12 E	2.80	2.89	7.14	41.0	29.4	11.6															
16	35.67	.141	69	.765	.617	S 34 W	3.05	3.65	43.2	31.8	11.4																
17	41.78	.217	81	.321	.304	1.00	N 84 W	8.56	10.83	48.0	30.2	17.8	*	0.2													
18	29.21	.092	73	.30.072	.625	S 82 W	0.82	0.88	4.44 W	13.27	15.04	29.8	20.2	9.6													
19	16.48	.068	74	.30.072	.30.074	S 86 W	5.33	6.26	23.0	12.0	11.0																
20	23.73	.107	87	.29.323	.29.226	N 50 E	7.58	8.52	30.5	13.2	17.3																
21	11.68	.068	91	.741	.673	N 22 W	7.01	7.61	20.0	8.0	12.0																
22	28.06	.173	91	.467	.294	S 15 W	5.41	14.29	41.5	18.2	23.3																
23	24.02	.090	70	.874	.784	N 64 W	18.23	20.27	30.0	24.3	5.7																
24	18.22	.080	79	.30.072	.992	N 88 E	4.75	6.63	27.3	9.5	17.8																
25	27.22	.105	71	.29.302	.497	N 66 W	3.72	6.15	32.2	18.7	13.5																
26	36.60	.175	80	.408	.234	S 76 W	3.38	5.24	33.2	27.3	5.7																
27	31.13	0.151	79	29.546	29.595	N 72 W	3.50	7.96	37.03	24.23	12.80	0.560	18.6	6.8	30.0	1.240	48.6										

* Inappreciable.

31.13	0.151	79	29.746	29.595	0.62	N 72 W	3.50	7.96	37.03	24.23	12.80	0.560	18.6	6.8	30.0	1.240	48.6
-------	-------	----	--------	--------	------	--------	------	------	-------	-------	-------	-------	------	-----	------	-------	------

* Inappreciable.

GENERAL METEOROLOGICAL ABSTRACT—JANUARY, 1862.

DAYS.	DAILY MEANS.				WIND.				PRESSURE OF TEMPERATURE.				RAIN.				SNOW.			
	Temperature of Air.	Pressure of Vapour.	Relative Humidity.	Barometric Pressure of Air.	Wind direction.	Wind velocity.	Depth of snow.	Wind direction.	Mean Velocity.	Maximum.	Minimum.	Depth of rain.	Dependence in sources.	Dependence in modes.	Dependence in time.	Dependence in hours.	Dependence in days.	Dependence in months.	Dependence in years.	Rate of Melted Snow.
1	30.68	0.135	.70	29.248	29.113	6.70	N 70 W	24.91	26.31	44.5	32.0	12.5	*	0.2	0.1	1.0	.016	1.2		
2	12.45	.154	.69	.919	.865	0.12	N 31 W	14.43	14.50	19.0	13.9	5.1		
3	2.42	.036	.73	.993	.872	0.68	N 3 E	8.87	9.07	8.2	5.6	10.8		
4	6.25	.037	.63	.788	.751	0.18	N 5 W	4.09	4.30	11.0	-1.8	12.8		
5	17.32	.081	.85	.471	.390	...	N 56 E	7.18	7.52	19.8	0.5	19.3	1.0	0.4	1.0		
6	7	22.60	.094	.78	.941	.847	N 75 W	3.96	4.33	21.8	11.0	10.8	0.3	3.0	.030	3.0		
8	30.20	1.44	.84	.745	.600	1.00	S 7 W	3.97	3.97	30.2	13.2	17.0		
9	35.32	1.75	.85	.626	.451	1.00	S 36 W	6.43	7.77	38.0	17.1	20.9	0.5	3.5	.050	3.5		
10	34.40	1.61	.76	.409	.298	0.75	N 68 W	4.18	5.07	37.0	20.0	8.0	0.5	0.5	0.05	0.5		
11	15.85	.078	.87	.583	.505	0.98	S 85 E	10.72	15.23	43.2	34.5	8.7	.060	2.5	0.60	2.5		
12	4.35	.042	.78	.030	.026	30.668	N 75 W	11.85	12.40	32.4	13.2	19.2	.050	2.0	0.2	1.5	.070	3.5		
13	12.00	.061	.76	.151	.091	0.60	N 83 E	6.10	8.40	7.21	1.5	5.6		
14	27.60	1.36	.90	.20	.329	29.194	S 64 W	10.57	14.94	32.7	12.2	20.5	2.5	5.8	.250	5.8		
15	14.12	1.59	.71	.639	.579	0.20	S 67 W	10.38	10.65	20.0	15.7	4.3		
16	18.78	.091	.87	.974	.883	1.00	N 13 E	4.71	5.20	24.0	2.0	22.0	1.5	1.5	1.5	1.5		
18	24.43	.119	.91	.680	.561	1.00	N 50 E	6.76	6.82	26.2	20.5	5.7	3.0	8.0	.300	8.0		
19	22.37	.066	.80	.566	.470	0.08	N 42 E	6.53	6.56	26.2	19.5	6.7	1.5	5.0	.150	5.0		
21	23.83	.110	.85	.737	.627	1.00	N 35 E	7.50	8.23	28.6	15.0	13.7	0.5	3.0	.050	3.0		
22	28.87	.139	.87	.687	.548	1.00	N 79 W	2.01	2.90	32.6	23.7	8.9	0.2	3.0	.020	3.0		
23	28.27	.133	.86	.637	.570	0.97	N 63 E	6.76	7.62	32.9	25.9	7.0	0.1	2.0	0.10	2.0		
24	29.35	.136	.83	.719	.583	0.97	N 84 E	10.00	10.34	31.5	26.3	5.2	1.5	3.0	.150	3.0		
25	27.80	.122	.81	.331	.205	0.98	S 36 W	4.67	10.07	32.0	22.0	10.0	5.0	13.7	.060	13.7		
26	16.37	.081	.85	.30	.136	30.055	N 84 W	12.98	13.01	25.8	23.0	2.8		
27	24.00	.119	.91	.26748	.29.629	1.00	S 66 E	2.46	3.46	27.2	5.0	22.2		
28	22.23	.154	.84	.508	.349	0.90	S 84 E	14.09	14.17	29.2	15.4	13.8	*	2.0	5.5		
29	23.38	.096	.76	.757	.661	0.52	N 5 W	5.56	7.65	29.8	20.3	9.5	0.2	*	0.2	0.2		
31	21.30	.097	.82	.907	.811	0.53	S 76 E	7.28	7.78	30.2	9.0	21.2		
	21.71	0.103	.81	29.727	29.624	0.73	N 26 W	2.69	8.82	27.58	13.03	12.55	0.115	7.2	27.4	88.9	2.855	96.1		

* Inappreciable.

GENERAL METEOROLOGICAL ABSTRACT.—FEBRUARY, 1862.

Days	DAILY MEANS.			WIND.			EXTREMS. OF TEMPERATURE.			RAIN.			SNOW.			RIS AND MELTED SNOW		
	Temperature of Air, of the Atmosphere	Vapour of Humidity	Relative Humidity	Clouded Sky	Pressure of Atmosphere	Velocity Wind	Maximum Velocity	Mean Velocity	Minimum	Difference	Depth in hours.	Deposition in hours.						
1	26.33	.109	.76	.29	.704	.29	.595	.58	N 86 W	3.01	33.5	.21	9.3	*	0.6	
2	
3	19.08	.660	.86	.754	.634	1.00	N 72 W	4.41	5.62	24.2	14.2	14.2	1.2	1.2	120	8.0	..	
4	20.53	.696	.86	.861	.765	0.88	N 63 W	3.35	3.47	23.4	10.8	8.6	3.5	3.5	100	2.5	..	
5	24.20	.116	.84	.854	.738	0.72	S 72 E	3.95	4.18	32.8	10.2	22.6	1.8	1.8	125	3.1	..	
6	34.53	.170	.85	.309	.139	1.00	N 76 W	9.18	11.39	37.8	26.0	8.8	1.3	1.3	175	3.1	..	
7	24.47	.109	.81	.726	.617	0.38	S 49 W	6.61	8.08	30.8	27.4	3.4	1.0	1.0	100	6.5	..	
8	14.90	.077	.90	.708	.631	1.00	N 36 E	1.87	2.91	21.0	12.0	9.0	0.4	0.4	5.5	.040	5.5	
9	N 80 W	8.98	9.21	21.0	13.7	7.8	
10	19.62	.666	.80	.517	.459	0.83	S 18 W	3.90	4.22	25.6	10.5	15.1	
11	28.63	.442	.89	.411	.387	1.00	S 16 E	7.39	10.23	22.4	20.3	12.1	
12	29.42	.133	.82	.418	.293	0.95	S 86 W	5.27	6.92	32.5	5.5	5.5	
13	26.70	.131	.90	.516	.384	1.00	S 38 E	2.89	6.23	30.0	21.7	8.3	
14	12.03	.061	.77	.669	.52	N 57 W	8.96	9.51	19.0	15.0	4.0		
15	9.58	.051	.75	.650	.508	0.48	S 67 W	10.93	11.49	17.2	-5.2	22.4	
16	N 21 W	1.41	3.04	17.8	2.0	15.8	
17	25.70	.132	.92	.729	.507	0.87	N 78 E	6.45	8.6	7.6	7.6	7.6	0.0	3.7	
18	32.83	.149	.79	.726	.577	1.00	S 88 W	10.60	11.25	27.0	26.0	11.0	
19	26.52	.127	.98	.610	.483	1.00	S 38 E	11.32	13.06	31.2	22.6	8.6	
20	21.07	.093	.83	.746	.653	0.63	S 57 W	10.66	11.29	25.8	20.2	5.6	
21	21.42	.094	.81	.738	.645	0.95	S 45 W	1.23	2.38	30.0	0.2	20.8	
22	20.72	.144	.87	.409	.265	1.00	S 38 E	0.47	1.52	34.0	18.0	16.0	
23	N 59 E	3.48	4.19	36.2	26.3	9.9	
24	18.85	.097	.84	.482	.385	0.37	N 42 W	18.29	18.72	32.0	13.6	18.4	
25	11.23	.067	.88	.982	.913	0.28	N 88 E	2.84	4.36	25.5	+4.5	30.0	
26	28.13	.112	.92	.400	.258	1.00	N 55 E	4.04	8.19	31.0	12.5	18.5	8.5	8.5	
27	20.30	.085	.77	.409	.324	0.78	N 68 W	19.76	19.83	26.0	22.0	4.0	*	
28	14.18	.066	.81	.528	.462	0.52	N 48 W	23.99	24.13	17.2	11.1	6.1	
	22.50	0.107	.84	20.608	20.501	0.78	N 55 W	3.93	8.52	28°.25	13.41	1.5	1.0	1.0	23.1	73.4	2490	
																	87.4	

Inappreciable.

22.50	0.407	84	29.608	29.501	0.748	N 55 W	3.93	8.32	28.25	1.0	41.1284	0.180	14.0	23.1	1.34	2.400	51.4
-------	-------	----	--------	--------	-------	--------	------	------	-------	-----	---------	-------	------	------	------	-------	------

* Immeasurable.

GENERAL METEOROLOGICAL ABSTRACT.—MARCH, 1862.

DATE.	DAILY MEANS.				WIND.				EXTREMES OF TEMPERATURE.				RAIN.				SNOW.			
	Temperature of Air.	Pressure of Vapor.	Relative Humidity.	Temperature of Dew-point.	Clouded Sky.		Clear Sky.		Maximum Velocity.	Minimum Velocity.	Wind.	Extremes of Temperature.	Difference in Height.	Rain.	Snow.	Rain and Melted Snow.				
					Hours.	Minutes.	Hours.	Minutes.												
1	17.55	0.075	76	29.638	29.564	0.05	N 52 W	10.56	9.78	11.17	28.0	1.6	16.4	2.0	.050	2.0				
2	16.88	0.075	76	29.564	29.490	0.05	N 58 E	9.92	10.56	28.0	8.0	20.0	19.0	5.0	1.000	17.5				
3	30.70	0.075	94	28.896	1.00	0.00	S 79 E	7.90	10.67	34.9	2.0	15.9	7.00	12.5	3.0	1.000				
4	23.78	0.107	83	1.37	29.030	0.98	S 67 W	8.36	10.81	26.2	2.5	4.7	0.2	0.2				
5	23.07	0.103	82	1.291	1.188	0.72	S 87 W	7.83	8.10	29.5	16.5	14.0	...*	...*	0.2	0.2				
6	24.68	0.111	83	1.347	23.36	0.90	S 38 W	7.72	8.10	39.0	20.3	9.2	...*	...*	0.2	0.2				
7	32.83	0.147	78	1.497	3.46	0.80	N 65 W	5.22	5.61	39.5	23.8	15.7	...*	...*	0.2	0.2				
8	34.52	0.153	77	1.696	5.43	1.00	N 76 W	2.50	4.19	37.2	22.0	5.2	...*	...*	0.2	0.2				
9	10	35.95	1.190	88	3.53	1.16	S 72 E	7.36	7.91	39.0	32.0	7.0	...*	...*	0.2	0.2				
11	11	30.38	1.37	81	6.28	1.91	S 97 W	9.77	11.50	39.0	33.0	6.0	16.0	4.0	...*	0.2	0.2			
12	12	27.02	1.07	74	7.02	1.65	S 60 W	6.93	7.55	37.8	24.8	13.0	...*	...*	0.2	0.2				
13	13	25.25	1.11	81	7.61	1.65	N 75 E	12.28	12.35	28.0	20.3	7.4	...*	...*	0.2	0.2				
14	14	29.25	1.39	86	6.28	1.89	1.00	N 61 E	8.04	8.59	32.0	25.0	7.0	5.25	13.0	...*	0.2			
15	15	31.63	1.71	96	2.54	0.82	1.00	N 48 E	10.33	10.44	33.2	29.8	3.4	7.45	14.0	3.0	9.0			
16	16	29.58	1.44	87	4.75	1.33	S 28 E	3.77	4.60	32.0	27.0	0	1.0	1.0	1.00	4.0				
17	17	28.75	1.21	78	7.64	1.91	S 33 W	4.68	5.25	35.0	20.7	14.3	...*	...*	0.2	0.2				
18	18	30.97	1.31	75	7.43	1.43	S 36 E	3.00	4.51	34.5	23.8	12.2	...*	...*	0.2	0.2				
19	19	31.57	1.70	96	1.74	1.04	N 80 W	16.89	17.11	34.5	25.8	8.7	...*	...*	0.2	0.2				
20	20	31.97	1.71	96	1.74	1.04	N 63 E	9.14	11.57	33.1	20.2	3.9	...*	...*	0.2	0.2				
21	21	31.90	1.65	91	28.916	0.75	N 65 W	3.67	4.59	37.8	20.0	8.8	0.20	2.5	0.2	0.2				
22	22	26.60	1.17	82	3.91	29.274	S 34 W	12.26	12.50	40.5	26.6	13.9	...*	...*	0.2	0.2				
23	23	23.80	1.01	78	5.57	0.62	N 50 W	7.01	7.25	34.0	15.4	8.6	...*	...*	0.2	0.2				
24	24	26.30	1.11	77	6.54	0.10	S 82 W	3.21	4.01	35.6	12.0	23.6	...*	...*	0.2	0.2				
25	25	32.32	1.20	67	6.71	0.47	N 68 W	8.18	8.87	43.2	20.7	22.5	...*	...*	0.2	0.2				
26	26	32.77	1.05	72	7.31	0.18	N 2 E	7.90	8.94	35.2	23.8	11.4	...*	...*	0.2	0.2				
27	27	27.98	1.19	77	7.30	0.61	N 89 E	10.74	11.00	32.4	26.5	12.1	...*	...*	0.2	0.2				
28	28	36.43	1.82	83	5.87	4.05	S 72 E	7.53	11.32	37.5	20.8	7.7	0.40	3.5	0.40	3.5				
29	29	31.90	1.82	83	5.87	4.05	S 78 W	14.31	15.12	40.0	33.5	6.5	...*	...*	0.2	0.2				
30	30	29.504	0.132	82	29.504	29.372	0.63	N 12 W	2.50	9.38	34.64	23.12	11.52	2.560	54.0	18.5	52.8			
31	31	28.79	0.132	82	29.504	29.372	0.63	N 12 W	2.50	9.38	34.64	23.12	11.52	2.560	54.0	18.5	52.8			

* Immeasurable.

GENERAL METEOROLOGICAL ABSTRACT—APRIL, 1862.

TORONTO METEOROLOGICAL RESULTS.

Date	DAILY MEANS.				WIND.				EXTREMES OF TEMPERATURE.				RAIN.				SNOW.			
	Temperature of Air	Pressure of Air	Relative Humidity	Barometer	Wind Velocity	Wind Direction	Wind Velocity	Wind Direction	Minimum Temperature	Maximum Temperature	Wind Velocity	Wind Direction	Wind Velocity	Wind Direction	Depth of Rain	Amount of Rain	Depth of Snow	Amount of Snow		
					Miles	Miles	Miles	Miles	Miles	Miles	Miles	Miles	Miles	Miles	Inches	inches	Inches	inches		
1	34.23	0.104	83	29.889	29.725	0.42	N 50° E	39.6	30.4	9.2010	1.0		
2	39.27	.192	80	.434	.241	0.87	S 62° E	3.71	15.5	45.7	29.8	15.9	.010	1.0010	1.5		
3	40.25	.179	72	.745	.566	0.77	N 77° W	8.07	10.24	45.4	31.2	11.2010	1.5		
4	32.50	.149	81	.784	.335	0.85	S 80° E	15.47	15.56	45.4	31.6	4.4	.555	5.0	0.1	1.5	.365	5.8		
5	36.40	.168	79	.378	.210	0.85	S 87° W	6.45	12.51	41.8	26.4	11.4	* *		
6	N 69° W	11.85	12.03	35.0	29.8	6.2		
7	26.17	.117	82	.874	.757	0.72	S 72° E	8.41	9.4	31.2	14.5	16.7		
8	29.30	.182	79	.647	.514	1.00	N 80° E	19.47	19.39	32.2	26.2	6.0		
9	33.22	.131	69	.576	.446	0.95	N 72° E	11.17	11.83	36.8	29.0	7.8		
10	35.67	.116	70	.901	.755	0.90	N 32° E	2.71	5.00	43.0	29.1	14.9		
11	37.08	.145	66	.30.002	.948	0.10	N 20° E	1.70	4.79	46.4	30.0	16.4		
12	39.73	.128	53	.017	.889	0.35	N 85° E	10.42	10.43	45.2	30.0	15.2		
13	N 77° E	6.74	6.49	45.8	36.0	9.8	.020	1.0020	1.0		
14	46.37	.247	79	.792	.545	0.93	N 88° E	3.84	4.18	57.0	39.8	17.2		
15	49.26	.288	82	.880	.631	0.63	S 81° E	6.11	6.17	55.4	40.5	14.0	*	..	* * 1.5	..	* 0.6	0.5		
16	54.32	.338	79	.721	.383	0.85	S 67° E	2.58	3.34	66.8	44.5	22.3	.005	0.5		
17	56.85	.334	82	.747	.363	0.98	S 80° W	4.27	6.92	68.0	51.4	16.6	.100	5.5		
18	S 74° W	5.25	8.98	57.9	45.2	11.8	.115	5.0		
19	41.30	.177	68	.772	.535	0.82	N 49° W	4.54	7.82	47.8	37.0	10.8		
20	S 87° E	5.44	7.48	44.8	35.0	9.8		
21	37.40	.159	89	.491	.232	1.00	S 92° E	18.93	18.95	38.8	34.6	4.2	.155	21.2		
22	39.88	.217	87	.107	.28.050	0.97	N 64° W	11.08	14.04	40.0	36.6	11.4	.020	2.0020	2.0		
23	31.12	.147	84	.565	.29.448	0.53	N 68° W	17.30	18.11	33.8	28.0	5.8	* 1.5	..	* 1.5	..		
24	35.92	.126	59	.801	.676	0.07	S 55° W	3.97	9.84	46.0	29.2	16.8		
25	33.00	.111	58	.888	.777	0.50	S 83° E	4.42	5.37	43.2	27.8	15.4		
26	39.87	.138	56	.977	.839	0.40	S 87° E	3.99	6.40	48.0	25.9	20.1		
27	27.27	.157	57	.614	.379	0.92	S 81° W	5.35	10.76	56.4	41.5	14.9	.025	1.0025	1.0		
28	48.10	.235	79	.763	.006	0.35	S 38° W	9.30	9.54	52.2	30.4	12.8		
29	43.97	.157	61	.702	.515	0.28	N 87° E	6.30	7.47	54.0	30.5	25.5		
30	45.45	.188	61		
	30.56	0.184	73	29.726	29.542	0.65	N 50° E	2.48	9.77	46.3	33.43	12.01	2.255	43.7	0.2	3.8	2.255	47.5		

* Improbable.

39.56	0.184	73	29.526	0.63	29.542	0.63	3.74	40.34	33.45	12.31	2.26	49.4	0.2	3.8	-2.50
-------	-------	----	--------	------	--------	------	------	-------	-------	-------	------	------	-----	-----	-------

* Impreciseable.

GENERAL METEOROLOGICAL ABSTRACT.—MAY, 1862.

DATE	DAILY MEANS.			WIND.			EXTREMES OF TEMPERATURE.			RAIN.			KNOW.			PAS AND MELTED SNOW.		
	Barometer. of die Air.	Barometer. middle.	Pressure of Vapour.	Wind direction.	Wind pressure.	Wind velocity.	Max. Velocity.	Min. Velocity.	Difference.	Depth in inches.	Approximate duration in hours.	Depth in inches.	Approximate duration in hours.	Depth in inches.	Approximate duration in hours.	Depth in inches.	Approximate duration in hours.	
1	42.27	0.235	.87	29.514	29.279	1.00	N 82° E	14.34	49.0	4.6	7.4	0.255	8.5	0.255	8.5	
2	44.37	0.257	.88	43.3	41.76	0.73	S 30 W	4.34	6.77	53.5	39.8	13.7	3.0	1.35	3.0	
3	46.43	0.239	.76	.523	.524	0.30	S 62 W	6.08	8.03	53.2	39.2	14.0	0.07	0.5007	
4	47.11	0.222	.55	46.45	45.45	1.17	N 45 W	1.36	5.94	55.0	36.5	18.5	0.5	
5	46.45	0.230	.73	.419	.519	0.88	N 39 W	11.10	12.27	54.6	40.0	14.6	
6	52.35	0.218	.56	.410	.492	0.77	N 53 W	16.38	17.14	63.8	48.0	20.8	
7	45.90	0.145	.48	.577	.432	0.15	N 30 W	10.34	10.43	52.6	49.2	12.4	
8	51.97	0.237	.61	.722	.485	0.20	N 80 W	6.22	8.19	68.0	32.4	35.6	
9	65.18	0.286	.45	.500	.504	0.17	N 57 W	16.33	16.33	78.5	41.5	37.0	
10	60.83	0.271	.32	.627	.556	0.00	N 26 W	15.69	15.81	70.5	54.6	15.9	
11	58.03	0.304	.63	.695	.591	0.35	S 47 W	1.64	3.24	59.6	40.2	19.4	
12	50.20	0.324	.68	.582	.507	0.73	N 35 E	3.14	8.86	4.6	6.5	67.5	39.4	28.1	*	*	* 1.0	
13	52.48	0.234	.60	.734	.500	0.00	N 69 E	5.37	6.85	62.2	42.6	19.6	
14	56.20	0.253	.57	.778	.525	0.17	N 73 E	1.53	2.73	65.5	44.4	21.1	
15	56.93	0.375	.72	.731	.556	0.00	S 42 E	0.78	1.17	74.0	43.6	30.4	
16	60.75	0.391	.73	.573	.573	1.84	S 08 E	0.62	1.73	70.6	71.8	49.0	23.4	
17	60.75	0.391	.73	.573	.573	1.84	S 08 E	0.62	1.73	70.6	71.8	51.6	20.2	
18	41.87	0.177	.182	.69	.507	.325	1.00	N 68 W	8.35	8.55	45.6	40.8	4.8	
19	44.08	0.204	.70	.664	.664	0.70	S 73 E	4.80	6.53	51.0	32.6	18.4	
20	49.26	0.287	.81	.452	.165	0.78	S 58 W	4.62	12.67	51.0	42.0	14.0	0.85	0.3	
21	54.37	0.291	.68	.597	.307	0.50	S 69 W	10.97	12.92	64.4	47.8	16.6	
22	54.37	0.291	.68	.597	.307	0.50	S 69 W	10.97	12.92	64.4	47.8	16.6	
23	50.15	0.181	.52	.780	.698	0.10	N 53 W	12.64	13.82	59.8	45.1	12.7	
24	46.07	0.183	.58	.833	.670	0.12	S 7 E	1.43	4.31	33.5	33.8	19.7	
25	52.8	0.227	.68	.521	.553	0.72	S 15 W	4.63	4.96	58.5	37.0	21.5	
26	58.28	0.268	.67	.363	.636	0.70	S 80 E	3.29	3.42	59.2	42.5	16.7	0.015	0.8	*	0.1	*	
27	51.25	0.193	.33	.603	.410	0.03	N 30 W	6.65	6.78	63.0	49.6	21.8	
28	55.15	0.227	.62	.587	.360	0.20	S 58 W	1.36	3.22	64.1	36.0	28.1	
29	55.08	0.238	.54	.508	.570	0.67	S 66 E	1.79	2.45	60.0	44.8	15.2	
30	51.40	0.242	.32	.521	.579	0.65	N 56 E	3.01	3.51	66.2	41.4	24.8	
31	52.17	0.253	.65	29.590	29.337	0.45	N 52 W	2.80	7.87	61.43	42.01	16.33	1.427	23.2	1.427	23.2	1.427	

* Impreciseable.

TORONTO METEOROLOGICAL RESULTS.

GENERAL METEOROLOGICAL ABSTRACT.—JUNE, 1862.

	DAILY MEANS.		WIND.		EXTRACTS OF THERMOMETER.		RAIN.		SNOW.		RAIN AND MELTED SNOW.	
	Days.	Hours.	Pressure of Barometer.	Temperature of Water.	Wind.	Velocity of Wind.	Mean Velocity of Wind.	Difference of Temperature.	Depth in Inches.	Depths in Inches.	Apparatus in borders.	Apparatus in borders.
1	56.28	0.352	77	20.433	N 82 E	3.04	3.10	.63	49.6	13.4
2	56.28	0.343	76	20.681	N 87	2.27	3.29	.63	46.0	17.0
3	56.33	1.00	55.5	20.679	N 35 E	4.23	4.81	.62	53.0	9.5
4	56.15	.528	57	56.6	N 41	7.00	64.5	.54	44.0	10.1
5	60.95	.293	53	57.55	N 47 S 81 E	1.07	2.16	.71	44.0	48.8	21.2	...
6	60.92	.335	62	54.9	N 47 S 81 E	1.18	1.86	.70	44.0	27.4
7	57.47	.322	67	55.6	N 62 N 16 W	1.62	1.7	1.61	53.0	9.2	.065	.10
8	59.18	...	52	57.8	N 7 W	2.31	3.14	.68	44.0	13.6
9	59.18	2.90	58	57.78	N 22 S 34 W	1.48	1.66	.76	47.0	29.0
10	61.58	0.316	58	56.52	S 33 S 1 E	0.93	1.1	1.44	50.8	20.2
11	61.17	3.79	71	59.305	S 34 W	0.75	1.1	2.87	51.0	10.0
12	68.25	.461	67	51.14	S 28 S 53	0.53	0.85	1.75	80.8	53.4
13	61.58	.365	64	45.8	S 57 N 54 W	0.70	0.70	1.42	69.0	57.4
14	58.33	.310	63	.677	S 26 N 41 E	0.85	1.41	1.87	53.0	57.8	.025	.10
15	58.05	.296	62	.659	S 6 E	0.77	0.9	1.82	65.0	56.8	1.0	...
16	53.28	.321	58	.681	S 25 S 88 E	0.25	0.39	0.04	60.6	43.8
17	62.85	.421	73	.463	S 55 W	0.62	0.62	0.35	59.2	32.0
18	59.90	.458	88	.214	S 75 N 56 W	1.00	1.00	1.79	8.78	67.2	56.0	...
19	52.10	.286	72	.394	S 10 N 10 E	0.62	0.38	0.26	7.10	60.4	49.3	...
20	54.57	.316	74	.512	S 63 S 25 W	0.63	0.29	4.18	5.33	41.6	21.9	...
21	58.05	.296	62	.659	S 20 N 35 W	0.20	0.35	8.74	10.69	71.0	46.8	...
22	57.18	.337	53	.688	S 35 W	0.97	1.41	4.83	65.2	42.0	23.2	...
23	57.18	.337	53	.688	S 68 E	0.91	1.46	7.06	61.8	52.4	9.4	...
24	56.70	.336	84	.688	S 97 N 57 E	0.91	1.83	5.56	61.6	55.4	.035	.06
25	64.05	.371	63	.646	S 46 W	2.75	2.20	5.36	9.17	74.0	51.2	...
26	73.32	.304	38	.623	S 38 W	0.57	0.57	7.46	7.86	85.4	64.0	...
27	70.52	.398	53	.567	S 33 W	0.30	0.30	4.17	6.57	70.6	55.2	...
28	69.57	.415	59	.606	S 71 E	0.53	1.45	3.21	91.0	58.0	23.6	...
29	62.85	.363	67	.347	S 19 W	0.65	0.34	3.36	77.0	58.5	18.5	...
30	62.85	.364	66	.29219	S 26 W	0.60	0.60	1.77	5.98	69.12	50.97	1.007
	60.52	0.346	66	.29219	S 26 W	0.60	0.60	1.77	5.98	69.12	50.97	1.007

* Imprecise.

GENERAL METEOROLOGICAL ABSTRACT.—JULY, 1862.

DAYS	DAILY MEANS			WIND.			EXTREMES OF TEMPERATURE.			RAIN.			SNOW.			RAIN AND MELTED SNOW.		
	Temperature of Air	Pressure of Atmosphere	Relative Humidity	Wind Velocity	Direction	Speed	Cloud Altitude	Velocity	Direction	Altitude	Depth in inches	Duration	Altitude in feet	Depth in inches	Duration	Altitude in feet	Depth in inches	Altitude in feet
1	60° 53'	30.409	29.100	0.22	S 10° W	4.65	5.66	70.2	49.8	20.4
2	61.47	.326	60	.498	S 76 E	2.97	3.33	70.8	50.2	20.6
3	65.60	.461	74	.775	S 114	0.60	S 51 E	1.86	2.83	77.0	49.8	27.2
4	72.22	.548	70	.875	S 27	0.63	S 12 W	3.53	3.97	83.8	55.0	28.8
5	77.58	.698	64	.747	S 41 W	6.42	6.88	90.2	63.9	27.7
6	71.42	.447	58	.431	S 62 W	6.51	9.43	95.5	67.8	27.7
7	73.67	.557	68	.342	S 77 W	2.35	3.02	82.5	61.2	21.3
8	69.63	.488	69	.319	S 28 W	8.23	8.45	79.5	64.6	14.9	0.8
9	65.55	.800	51	.635	S 19 W	7.75	8.63	76.0	54.2	21.8
10	63.32	.390	68	.670	S 17 W	1.68	1.77	83.0	77.5	48.2
11	70.47	.473	64	.433	S 59 W	0.98	0.67	67.7	6.62	7.5	31.9
12	66.70	.565	86	.342	S 25 W	5.66	7.25	83.9	52.0	31.9	*	0.1	0.1
13	64.78	.512	83	.443	S 90 E	7.65	8.38	80.4	62.6	17.8	.625	8.0625
14	64.28	.553	92	.466	S 91 W	0.68	0.59	54.6	74.2	6.0	12.2	*	0.1	0.1
15	62.32	.424	73	.599	S 17 W	2.80	3.32	73.0	50.8	13.2	1.40	2.0	2.0
16	61.52	.389	71	.698	S 36 W	0.65	0.66	7.13	8.30	70.6	61.0
17	62.82	.429	72	.713	S 26 W	0.47	0.46	2.71	6.10	68.8	50.4	18.4
18	66.70	.565	86	.333	S 96 W	0.92	0.80	4.73	6.29	72.0	62.3	9.7	.335	4.5
19	64.23	.545	91	.499	S 77 W	0.69	0.81	8.13	78.8	63.8	15.0	0.290	1.6	1.6
20	67.35	.569	85	.545	S 44 E	0.77	0.77	6.0	0.33	1.37	61.4	5.8	.005	1.0005
21	70.23	.510	71	.310	S 80 W	1.00	1.42	4.18	5.07	72.8	62.8	10.0	.352	3.0352
22	62.07	.384	68	.498	S 114	0.65	0.65	5.72	5.56	81.0	65.2	14.8	.880	2.5880
23	64.57	.411	65	.567	S 116	0.07	S 71 W	4.40	6.60	72.0	51.8	20.2	.005	1.0005
24	63.27	.473	81	.409	S 83 W	4.94	5.95	70.8	54.0	16.8	.090	1.0	1.0	
25	62.07	.384	68	.498	S 114	0.65	S 71 W	4.40	6.60	72.0	51.8	20.2	.005	1.0005
26	66.57	.411	65	.567	S 116	0.23	S 72 W	3.98	6.86	76.8	56.0	20.8
27	64.75	.532	87	.528	S 7 W	3.76	3.75	75.0	53.4	21.6	
28	66.93	.551	83	.502	S 34 W	4.00	5.18	72.0	60.0	12.0	.815	6.3815	
29	68.48	.489	71	.628	S 17 W	0.67	S 34 W	4.17	N 38 W	7.52	8.18	75.0	63.0	12.0440
30	70.25	.534	72	.701	.167	0.03	S 87 W	0.91	4.43	76.3	60.5	16.0
31	69.70	.473	71	29.547	29.075	0.56	S 80 W	1.42	5.80	76.42	58.5	14.18	28.5	344	35.4	5.344

• Imprecise.

TORONTO METEOROLOGICAL RESULTS.

GENERAL METEOROLOGICAL ABSTRACT - AUGUST, 1892.

Days	DAILY MEANS			WIND			EXPOSURE			RAINFALL			SNOW			RAIN AND METEOROLOGICAL RESULTS.			
	Temperature of Outside Air	Pressure of Vapour over Water	Relative Humidity	Direction of Wind	Speed of Wind	Depth of Wind	Maximum Wind Velocity	Distance from Water	Maximum Wind Velocity	Distance from Water	Depth in Inches	Apparatus in Inches	Depth in Inches	Apparatus in Inches	Depth in Inches	Apparatus in Inches	RAIN AND METEOROLOGICAL RESULTS.		
1	72.72	0.507	71	29.651	29	.685	0.37	S 63 W	3.11	61.4	81.1	60.0	21.1	
2	71.82	0.565	73	64.1	0.45	N 69 E	2.60	3.71	79.8	68.4	13.4	60.6	
3	71.22	0.517	75	53.9	28	.507	0.60	S 24 W	8.60	8.93	84.0	68.8	15.2	
4	70.17	0.553	79	58.8	29	.016	0.65	N 61 W	6.57	7.60	79.8	76.2	9.6	* 1.4	
5	69.69	0.460	70	47.2	0.26	S 73 W	1.27	5.03	73.5	53.5	22.0	0.45	1.3	
6	69.65	0.588	81	67.3	0.65	S 39 E	2.15	4.64	76.5	63.0	13.5	3.12	2.0	
7	70.08	0.441	79	47.8	0.40	S 64 W	6.55	7.20	89.5	70.0	19.5	0.2	0.2	
8	73.73	0.650	76	57.8	25	.533	N 70 W	9.28	10.05	84.6	71.5	13.1	.636	0.5
9	71.33	0.486	75	47.9	28	.576	0.42	S 69 W	3.60	4.46	76.2	66.3	15.7
10	67.88	0.363	59	40.7	29	.284	0.12	S 49 W	9.44	10.00	79.8	58.6	21.2	.651	1.0
11	63.15	0.353	68	47.1	3.88	0.40	5 E	1.57	3.99	79.4	53.3	24.1
12	62.17	0.402	68	51.6	0.24	S 77 W	4.06	6.19	69.2	56.8	12.4	.618	10.9	
13	61.10	0.323	61	66.7	3.64	0.52	S 10 W	8.51	8.92	60.5	52.8	16.7
14	58.89	0.368	73	48.6	0.18	S 65 W	2.15	4.52	68.8	52.5	16.3	
15	57.59	0.358	75	47.6	22	.555	0.18	S 42 W	3.42	3.92	70.8	46.0	24.8
16	57.59	0.358	75	47.6	22	.555	0.18	S 8 E	2.27	3.76	74.0	53.5	20.5	.429	0.8
17	57.59	0.358	75	47.6	22	.555	0.18	S 43 E	1.79	3.28	75.0	54.2	20.8
18	57.59	0.358	75	47.6	22	.555	0.18	S 30 W	0.70	4.64	73.0	55.3	23.5
19	57.59	0.358	75	47.6	22	.555	0.18	S 75 E	5.11	5.60	74.8	56.9	19.8	.531	3.0
20	58.55	0.582	74	63.2	0.671	0.75	S 81 E	7.55	79.2	65.8	13.4	4.5	
21	58.55	0.582	74	63.2	0.671	0.75	S 79 W	5.68	7.38	9.49	5.58	29.0	
22	68.50	0.408	51	58.8	28	.828	0.65	S 65 W	0.13	0.25	7.38	5.38	29.0
23	61.42	0.459	76	65.8	0.20	.330	0.687	S 33 W	5.79	6.79	62.0	48.4	13.6
24	61.42	0.459	76	65.8	0.20	.330	0.687	S 65 E	4.97	5.79	62.0	48.4	13.6
25	66.07	0.463	70	67.8	21	.578	..	S 48 W	6.48	8.42	2.98	3.62	75.8
26	72.68	0.582	74	60.4	28.022	0.47	2.72	S 70 W	2.58	5.28	84.0	63.6	20.4	2.42	3.0
27	67.28	0.596	80	48.2	28	.886	1.00	S 67 E	1.51	4.05	71.5	64.2	7.3	.293	6.0
28	69.45	0.571	61	40.0	28	.829	0.42	S 60 W	8.33	9.57	78.0	64.0	14.0
29	61.00	0.355	64	68.1	20	.326	0.08	S 27 W	7.29	7.44	71.0	56.2	14.8
30	57.43	0.297	63	8.85	S 36 E	2.58	5.46	69.2	46.2	23.0	.636	8.5
31	
	67.60	0.510	74	29.616	0.45	N 78 W	1.67	5.96	76.11	58.22	1 ²	3.483	27.1

Inapplicable.

67.60 0.510 74 28.616 0.45 N 21.956 3.01 0.611 5.522 11.53 0.71 0.71

Inappreciable.

GENERAL METEOROLOGICAL ABSTRACT.—SEPTEMBER, 1862.

TORONTO METEOROLOGICAL RESULTS.

33

Date.	DAILY MEANS.			WIND.			EXTREMES OF TEMPERATURE.			RAIN.			SNOW.			RAIN AND MELTED SNOW.		
	Pressure of Dry Air.	Humidity.	Vapour of Water.	Pressure of Dry Air.	Humidity.	Vapour of Water.	Maximum.	Minimum.	Avgd. Velocity.	Wind.	Velocity.	Wind.	Depth in Rain.	App'd. in Rain.	Depth in Snow.	App'd. in Snow.	Depth in Melted Snow.	
1	61.38	0.484	.84	29.197	28.713	0.75	N 54 W	12.26	1.46	7.1	9	63.6	2.4		
2	47.45	.221	.68	.617	.29	.306	0.20	0.27	W	8.67		
3	55.50	.319	.71	.742	.422	0.23	S 27	W	2.84	3.10	61.8	29.0	17.0		
4	64.45	.483	.80	.636	.213	0.17	S 28	W	5.65	5.71	73.2	52.8	22.4		
5	69.32	.557	.78	.624	.067	0.78	S 27	W	1.68	2.41	77.0	64.2	12.8		
6	69.63	.654	.92	.555	.28.901	1.00	S 22	W	7.52	7.67	73.5	58.0	10.7	0.040	0.40	0.33		
7	S 57	W	8.76	9.87	75.5	65.0	10.5		
8	67.27	.450	.68	.568	.29.118	0.481	S 63	N 64	8	1.14	2.09	65.6	49.4	15.6	
9	57.97	.388	.82	.879	.481	0.68	S 79	W	8.76	9.87	75.5	65.0	10.5		
10	61.65	.403	.74	.837	.464	0.60	S 77	E	2.42	2.87	69.9	49.8	10.2		
11	66.22	.508	.78	.619	.161	0.28	S 60	E	2.12	2.57	74.0	54.1	19.6	0.007	0.07	0.10		
12	61.55	.430	.47	.696	.266	0.73	S 27	W	10.97	11.64	68.8	63.2	5.6	0.005	0.05	0.05		
13	48.70	.280	.81	.980	.700	0.53	S 68	E	2.67	3.75	55.0	44.8	10.2		
14	S 77	E	6.82	2.25	62.0	45.0	9.0		
15	57.48	.337	.91	.786	.349	0.77	S 22	E	2.72	4.06	65.0	48.6	16.4	0.037	0.37	1.5		
16	56.97	.206	.79	.828	.402	0.59	S 83	E	3.59	3.98	62.2	54.6	7.5		
17	62.22	.514	.59	.573	.059	0.55	S 56	E	5.81	6.24	51.8	52.8	15.2		
18	66.45	.642	.83	.458	.28.917	0.70	S 92	W	5.72	6.17	74.0	64.0	10.0	0.165	1.65	2.22		
19	58.30	.381	.77	.765	.29.384	0.50	S 32	W	2.81	4.86	66.8	55.4	11.4		
20	58.22	.396	.92	.751	.355	0.30	S 32	W	2.69	3.85	57.8	49.8	10.8		
21	S 53	E	2.50	3.00	65.0	54.2	10.8		
22	62.07	.473	.85	.757	.254	0.23	S 10	W	2.28	2.62	73.8	48.4	25.4	0.415	4.15	3.85		
23	64.75	.488	.90	.557	.018	0.65	S 59	W	2.63	7.29	75.0	53.2	21.2		
24	53.40	.270	.68	.681	.411	0.35	S 17	W	6.73	6.92	60.2	52.8	7.4		
25	54.58	.393	.77	.730	.397	0.05	S 26	W	3.35	4.38	64.0	45.0	10.0		
26	57.48	.378	.78	.720	.342	0.00	S 28	W	2.63	2.99	67.8	45.0	22.8		
27	57.42	.395	.84	.682	.287	0.53	S 34	E	0.61	1.04	66.2	45.2	14.2		
28	58.88	.415	.82	.611	.196	0.62	S 8	E	0.66	1.50	70.6	52	18.6		
29	58.22	.293	.82	.790	.506	1.00	S 64	E	5.08	5.52	...	46.0	31.6	8.3	...	3.15	8.3	
30	49.63	0.418	.80	.29	.683	29.265	0.47	S 59	W	1.07	5.11	68.43	52.77	1.66	2.344	20.7	...	
	50.59	2.344	20.7	

TORONTO METEOROLOGICAL RESULTS.

GENERAL METEOROLOGICAL ABSTRACT—OCTOBER, 1862

DAYS	DAILY MEANS		WIND		EXTREMES OF TEMPERATURE.		RAIN.		SNOW.		RAIN AND MELTED SNOW.	
	Temperature of Air	Pressure of Air	Clouded Sky	Clouded Air	Mean Velocity	Difference	Depth in inches	Approximate in hours	Depth in inches	Approximate in hours	Depth in inches	Approximate in hours
1	50° 27'	0.338	92	29.747	29.400	1.00	N 60° E 5.50	5.2	45° 0	12.0	0.085	3.5
2	58.47	0.473	96	0.617	.104	1.00	S 33° E 0.65	51.4	11.8	.085	5.0	
3	60.80	.521	97	.645	.124	1.00	N 78° E 3.48	62.5	53.9	.565	10.0	
4	61.03	.400	72	.499	.069	0.67	N 86° W 15.05	17.21	73.8	.072	0.9	
5	54.63	.370	55	.555	.115	.05	S 23° W 2.70	5.88	60.0	.104	10.6	
6	67.55	.503	88	.563	.292	28.796	0.40	4.1	47.6	.185	6.7	
7	67.55	.503	88	.563	.292	0.40	S 41° W 6.31	6.32	74.0	1.0	1.0	
8	59.38	.554	77	.594	.284	0.55	S 52° W 8.27	16.89	76.6	.126	2.5	
9	54.95	.329	75	.476	.207	20.697	1.00	N 17° W 3.08	3.29	58.8	.030	4.7
10	48.42	.292	56	.624	.342	1.00	N 1° E 8.95	9.53	52.8	.270	8.7	
11	46.15	.212	76	.530	.508	0.47	N 20° W 5.40	5.58	55.6	.050	1.5	
12	51.10	.310	55	.576	.266	.00	N 73° E 3.00	3.81	52.6	.250	1.5	
13	46.02	.310	53	.601	.432	0.84	N 25° W 8.01	8.34	58.2	.100	2.5	
14	38.92	.191	61	.521	.630	0.96	N 68° E 2.66	3.43	42.0	.114	1.5	
15	47.03	.260	99	.482	.192	0.93	S 44° W 4.70	4.92	49.4	.055	7.9	
16	44.98	.219	74	.504	.485	0.32	N 65° W 5.08	5.35	53.5	.040	1.5	
17	42.17	.206	76	.578	.523	0.18	S 25° W 13.03	13.76	61.2	.230	2.5	
18	47.70	.265	79	.578	.402	.00	S 61° W 8.79	9.09	55.0	.020	2.5	
19	42.18	.213	78	.615	.402	0.80	S 21° W 6.61	7.40	29.5	.223	3.0	
20	48.17	.202	57	.117	28.815	0.95	S 56° W 16.22	16.57	51.8	.035	4.4	
21	48.17	.202	57	.117	29.197	0.88	S 67° W 16.22	16.57	48.2	.400	0.2	
22	42.17	.206	76	.543	.633	0.53	S 88° W 2.32	4.80	45.0	.032	1.5	
23	38.23	.192	53	.558	.656	0.40	S 79° W 8.25	11.67	56.0	.140	1.5	
24	47.45	.253	77	.663	.410	1.00	S 79° W 5.61	6.50	39.0	.100	4.0	
25	35.37	.173	55	.974	.709	0.95	N 11° W 5.11	5.57	31.8	3.0	1.0	
26	34.47	.167	53	.540	.555	.05	N 14° W 3.73	3.69	3.82	2.8	.50	
27	34.47	.167	53	.540	.555	.065	S 47° W 4.32	4.95	43.0	.262	3.0	
28	40.43	.216	56	.648	.432	0.95	S 14° E 3.01	3.20	45.4	.315	6.5	
29	42.13	.223	53	.702	.479	0.60	S 88° W 7.20	7.33	49.9	.400	1.5	
30	44.75	.235	79	.647	.412	0.33	S 24° W 2.09	3.70	53.5	.207	4.0	
31	51.17	.213	71	.519	.250	0.10	S 42° W 6.19	6.42	59.0	.408	1.5	
	48.70	0.300	82	29.619	29.319	0.72	N 78° W 5.53	54.80	41.43	13.36	2.684	71.7
												4.0
												2.734
												75.7

* Improbable.

GENERAL METEOROLOGICAL ABSTRACT—NOVEMBER, 1862.

TORONTO METEOROLOGICAL RESULTS.

35

Days	DAILY MEANS.				WIND.				EXTREMES. OF TEMPERATURE.				RAIN.				SNOW.			
	Temperature of Air.	Humidity.	Pressure of Atmosphere.	Wind Velocity.	Direction.	Pressure of Atmosphere.	Wind Velocity.	Direction.	Minimum.	Maximum.	Direction.	Pressure of Atmosphere.	Wind Velocity.	Direction.	Depth in inches.	Approximate duration in hours.	Rain and Melted Snow.			
1	50.35	0.265	72	29.536	29.271	0.28	3.34 W	2.83 miles	58°	39°.0	19.0	0.680	5.5680	5.5	...				
2	37.65	..160	70	.578	.418	0.45	N 45 W	10.40	12.26	53.0	44.4	8.6	0.2	... * 0.2	...					
3	36.50	179	81	.796	.473	0.45	N 52 W	10.41	11.55	45.8	36.8	9.0	* .050	2.5050	2.5	...			
4	38.37	185	78	.550	.365	0.36	N 33 W	8.84	8.99	49.3	31.0	18.3	21.5					
5	28.62	165	68	.687	.582	0.45	N 25 E	5.31	6.22	33.6	24.4	9.2050	1.0050	1.0	...			
6	27.32	102	69	.775	.673	0.47	N 38 E	7.88	8.13	31.4	22.6	8.8	0.2	1.0020	1.0	...			
7	26.65	128	89	.736	.607	1.00	N 10 E	5.46	5.99	28.8	23.4	5.4	0.5	5.5050	5.5	...			
8	10	37.95	73	.733	.633	0.48	N 66 W	6.40	6.72	42.0	27.0	15.0050	1.5045	1.5	...			
9	11	44.80	236	.79	.494	.258	N 75 S 51 W	5.42	8.35	50.2	31.0	19.2	1.5050	1.5045	1.5	...		
12	12	39.33	182	.76	.678	0.46	N 69 W	5.25	5.33	44.0	36.0	5.0050	1.5045	1.5	...			
13	13	39.38	185	.76	.603	0.93	N 55 W	5.60	5.83	42.8	35.0	7.8050	1.5045	1.5	...			
14	14	35.60	157	.75	.933	.776	N 41 W	9.53	11.97	43.2	35.8	7.4	0.2	0.5050	0.7	...			
15	15	22.80	.069	.81	.30	.427	20.329	.222	1.75	27.6	16.2	11.4	.225	2.9225	2.9	...			
16	16	22.56	.166	.73	.799	.633	N 33 W	2.80	4.35	46.5	35.5	16.1	1.5050	1.5045	1.5	...		
17	17	43.13	.236	.79	.494	.635	N 75 S 51 W	5.42	8.35	50.2	31.0	19.2	1.5050	1.5045	1.5	...		
18	18	39.57	.210	.86	.895	.635	N 100 E	2.85	3.37	45.0	35.5	7.2	1.5050	1.5045	1.5	...		
19	19	42.58	.262	.96	.439	.178	N 10 E	5.75	5.82	44.2	39.0	5.2	.660	23.0660	23.0	...			
20	20	34.87	.173	.85	.350	.176	N 97 N	7.45	8.07	38.5	32.8	4.7	.285	9.0285	9.0	...			
21	21	29.68	.145	.87	.508	.363	N 70 W	1.42	1.57	33.4	25.5	7.9050	1.5050	1.5	...			
22	22	31.45	.146	.83	.505	.358	N 75 S 30 W	10.54	10.82	37.2	25.5	11.7050	1.5050	1.5	...			
23	23	29.955	.29	.719	1.00	N 117 W	3.71	4.40	49.6	35.0	14.6	1.5050	1.5045	1.5	...			
24	24	35.73	.165	.79	.592	.427	S 33 W	2.85	3.37	45.0	35.5	7.2	1.5050	1.5045	1.5	...		
25	25	39.40	.199	.82	.472	.273	S 98 E	3.21	3.92	44.0	34.5	9.5	0.5050	0.5050	0.5	...		
26	26	34.57	.154	.77	.469	.315	S 66 W	6.54	6.80	38.5	32.5	6.0050	1.5050	1.5	...			
27	27	31.82	.147	.81	.348	.201	N 34 W	5.32	6.41	34.2	32.0	2.2050	1.5050	1.5	...			
28	28	30.50	.142	.83	.191	.049	N 73 W	4.25	4.52	33.0	29.1	5.9050	1.5050	1.5	...			
29	29	30.85	.154	.89	.397	.243	N 43 W	0.74	1.81	36.0	25.0	11.0050	1.5050	1.5	...			
30	30	N 36 E	1.37	2.90	25.0	20.7	5.3050	1.5050	1.5	...			
	35.58	0.171	80	29.636	29.465	0.79	N 46 W	3.00	6.60	40.50	30.50	10.09	2.205	51.5	5.3	51.0	2.735	102.5		

*, Inappreciable.

TORONTO METEOROLOGICAL RESULTS.

GENERAL METEOROLOGICAL ABSTRACT—DECEMBER, 1862.

Date	DAILY MEANS.			WIND.			EXTREMES OF TEMPERATURE.			RAIN.			SNOW.			RUM AND MELTED SNOW.		
	Temperature	Humidity	Pressure of Air	Clouded Sky	Visibility	Wind	Minimum	Maximum	Difference	Minimum	Maximum	Difference	Depth in inches	Approximate hours	Depths in inches	Approximate hours	Depths in inches	Approximate hours
1	33.07	0.162	.55	29.665	29.503	N 44° W	5.14	5.51	.358	32.0	3.8	28.2	3.0	6.5	0.300	6.5		
2	24.45	.110	.83	.861	.751	N 63° W	3.50	4.03	.398	18.2	11.6	23.5	8.1	0.1	*	0.1		
3	28.48	.131	.84	.602	.471	S 12° W	3.10	4.92	.316	23.5	8.1	23.5	8.1	3.0	7.5	.300	7.5	
4	29.95	.130	.78	.515	.415	S 81° W	5.33	5.45	.340	19.4	14.6	20.5	9.7	
5	26.48	.110	.75	.552	.242	S 73° W	7.56	8.87	.302	20.5	9.7	20.5	9.7	
6	8.97	.060	.76	.421	.371	N 52° W	21.30	21.52	13.6	10.6	3.0	10.5	1.5	12.0	0.1	11.1	.350	11.0
7	19.38	.102	.94	.797	.695	S 34° E	4.83	6.92	10.5	7.30	2.0	23.8	8.1	2.5	11.1			
8	25.53	.122	.88	.812	.690	S 70° W	4.72	7.01	30.6	21.3	9.3	25.9	12.1	
9	34.72	.166	.82	.670	.504	S 50° W	9.94	10.07	38.0	21.8	10.5	32.5	12.5	
10	41.60	.209	.79	.549	.340	S 51° W	11.08	11.08	47.0	34.5	12.5	34.5	12.5	
11	38.37	.177	.76	.766	.560	S 73° E	5.10	6.10	42.0	31.0	8.0	42.0	17.0	
12	40.50	.231	.91	.647	.416	S 75° E	7.5	8.52	5.08	45.4	37.0	8.4	42.0	17.0	
13	44.27	.255	.67	.710	.510	S 22° E	2.14	2.52	5.01	32.9	10.3	21.0	2.5	
14	29.63	.69	.210	.925	.100	S 88° W	4.19	6.93	50.0	44.5	5.5	1.250	15.0	0.12	1.0	1.250	16.0	
15	31.42	.135	.75	.263	.20	S 71° W	14.64	15.69	35.8	32.5	3.3	0.5	1.3	0.650	1.3			
16	17.98	.079	.91	.30.631	.363	S 23° W	7.38	7.77	21.8	16.0	5.8	0.5	*	0.12	*	0.2		
17	27.07	.116	.78	.29.913	.797	S 20° W	14.63	15.17	33.0	15.6	17.4	
18	16.73	.086	.89	.30.694	.918	S 20° W	18.17	18.83	34.9	17.5	16.5	
19	5.30	.041	.74	.30.370	.30.329	S 55° W	2.32	3.35	16.5	23.4	19.9	
20	33.08	.160	.85	.29.655	.29.494	S 40° E	5.51	5.82	27.0	5.3	21.7	...	0.2	2.5	0.220	2.5		
21	31.62	.148	.82	.872	.724	S 72° W	4.05	4.58	39.0	20.7	18.3	
22	32.48	.168	.87	.778	.610	S 17° E	4.66	6.53	36.0	31.5	4.5	
23	36.30	.190	.93	.265	.096	S 52° W	2.47	2.90	46.6	32.6	14.0	41.0	16.2	*	0.7			
24	34.98	.178	.88	.444	.266	S 23° E	4.14	5.74	38.0	35.2	2.8	0.345	16.0	
25	37.05	.177	.80	.546	.369	S 44° W	6.43	6.57	39.8	32.8	3.8	
26	26.68	.108	.74	.661	.553	S 61° W	5.72	7.37	41.2	33.2	8.0	
27	21.00	.097	.85	.883	.786	S 13° W	0.94	1.98	27.8	15.0	12.8	
28	28.78	0.142	.82	.29.678	.29.536	N 73° W	3.17	7.58	34.1	28.5	5.7	54.1	9.945	41.2	10.4	30.1	2.985	71.3

* Inapplicable.

TABLE I.
Monthly Means of the Temperature of the Air, at each of the observation hours, from 1860 to 1862 inclusive.

Toronto Astronomical Time.	2 ^h		4 ^h		10 ^h		12 ^h		18 ^h		20 ^h		Mean.		2 ^h		4 ^h		10 ^h		12 ^h		18 ^h		20 ^h		
	2 ^h	4 ^h	10 ^h	12 ^h	18 ^h	20 ^h	2 ^h	4 ^h	10 ^h	12 ^h	18 ^h	20 ^h	2 ^h	4 ^h	10 ^h	12 ^h	18 ^h	20 ^h	2 ^h	4 ^h	10 ^h	12 ^h	18 ^h	20 ^h	Mean.		
January	1860	26.31	25.67	25.12	21.68	21.51	21.51	21.62	21.60	23.38	23.38	23.38	23.38	1860	70.30	70.07	69.45	58.94	59.33	64.45	63.92	63.92	63.92	63.92	63.92	63.92	
	"	1861	22.86	22.67	18.51	17.78	17.91	19.86	19.86	19.86	19.86	19.86	19.86	"	1861	71.49	71.38	62.41	60.76	60.56	65.40	65.37	65.37	65.37	65.37	65.37	65.37
	"	1862	25.01	24.60	21.53	20.44	19.25	19.45	21.71	21.71	21.71	21.71	21.71	"	1862	73.08	73.21	63.46	62.52	61.33	66.60	66.70	66.70	66.70	66.70	66.70	66.70
	Means	24.73	24.31	21.36	20.21	19.51	19.79	21.65	21.65	21.65	21.65	21.65	21.65	Means	71.62	71.62	62.11	60.74	60.41	65.48	65.33	65.33	65.33	65.33	65.33	65.33	
February	1860	26.57	26.45	21.66	20.62	20.48	21.17	22.83	22.83	22.83	22.83	22.83	22.83	1860	71.18	70.35	61.68	60.24	58.60	64.21	64.46	64.46	64.46	64.46	64.46	64.46	
	"	1861	28.82	28.45	26.05	25.32	23.49	24.37	25.06	25.06	25.06	25.06	25.06	"	1861	71.75	70.91	62.67	61.45	60.48	65.50	65.48	65.48	65.48	65.48	65.48	65.48
	"	1862	25.34	25.16	21.87	21.67	20.65	20.91	22.50	22.50	22.50	22.50	22.50	"	1862	74.06	74.10	64.58	63.27	62.36	67.25	67.25	67.25	67.25	67.25	67.25	67.25
	Means	26.91	26.69	23.19	22.54	21.34	21.34	22.10	23.80	23.80	23.80	23.80	23.80	Means	72.33	71.35	62.98	61.65	60.48	65.68	65.55	65.55	65.55	65.55	65.55	65.55	
March	1860	30.92	40.21	32.29	32.26	29.39	31.81	34.48	34.48	34.48	34.48	34.48	34.48	1860	61.33	60.44	52.60	51.56	50.17	55.71	55.34	55.34	55.34	55.34	55.34	55.34	
	"	1861	30.87	31.08	26.72	25.14	23.17	24.79	26.92	26.92	26.92	26.92	26.92	"	1861	64.72	64.46	56.80	55.61	54.38	58.44	58.44	58.44	58.44	58.44	58.44	58.44
	"	1862	33.14	32.83	28.37	26.88	24.67	26.89	28.79	28.79	28.79	28.79	28.79	"	1862	66.26	65.73	57.01	55.40	54.34	58.77	58.77	58.77	58.77	58.77	58.77	58.77
	Means	34.64	34.71	29.39	28.09	25.74	27.79	30.06	30.06	30.06	30.06	30.06	30.06	Means	64.17	63.54	55.48	54.10	52.36	57.64	58.00	58.00	58.00	58.00	58.00	58.00	
April	1860	40.80	44.35	37.09	37.06	34.73	38.97	38.55	38.55	38.55	38.55	38.55	38.55	1860	52.06	51.19	45.70	45.13	43.74	45.71	47.25	47.25	47.25	47.25	47.25	47.25	
	"	1861	46.92	47.35	41.29	39.87	36.61	40.09	42.02	42.02	42.02	42.02	42.02	"	1861	53.94	52.96	47.50	46.50	43.81	47.71	48.74	48.74	48.74	48.74	48.74	48.74
	"	1862	44.09	43.79	27.77	37.14	36.02	38.55	38.56	38.56	38.56	38.56	38.56	"	1862	53.79	52.23	47.94	47.15	44.53	46.60	48.70	48.70	48.70	48.70	48.70	48.70
	Means	45.36	45.16	39.02	38.01	35.79	39.60	40.38	40.38	40.38	40.38	40.38	40.38	Means	53.26	52.13	47.05	46.26	44.93	46.67	48.23	48.23	48.23	48.23	48.23	48.23	
May	1860	61.39	61.14	63.42	51.33	50.77	55.11	55.53	55.53	55.53	55.53	55.53	55.53	1860	41.80	40.80	37.00	36.19	35.39	36.32	37.95	37.95	37.95	37.95	37.95	37.95	
	"	1861	53.27	52.93	44.94	43.19	43.08	47.56	47.50	47.50	47.50	47.50	47.50	"	1861	40.83	39.68	36.21	35.45	35.05	35.03	37.14	37.14	37.14	37.14	37.14	37.14
	"	1862	58.26	58.52	49.41	47.78	47.36	51.71	52.17	52.17	52.17	52.17	52.17	"	1862	39.59	38.05	34.34	33.46	33.46	34.31	35.58	35.58	35.58	35.58	35.58	35.58
	Means	57.64	57.53	49.26	47.43	47.07	51.46	51.73	51.73	51.73	51.73	51.73	51.73	Means	40.74	39.51	35.85	35.11	34.70	35.42	36.84	36.84	36.84	36.84	36.84	36.84	
June	1860	68.65	68.98	60.25	58.95	59.17	62.97	63.16	63.16	63.16	63.16	63.16	63.16	1860	26.44	25.18	23.02	23.30	23.31	22.77	24.00	24.00	24.00	24.00	24.00	24.00	
	"	1861	67.70	67.08	58.68	56.70	56.42	61.17	61.29	61.29	61.29	61.29	61.29	"	1861	34.84	33.94	30.52	29.58	29.14	28.76	31.13	31.13	31.13	31.13	31.13	31.13
	"	1862	68.57	68.15	57.33	56.45	55.44	61.20	60.52	60.52	60.52	60.52	60.52	"	1862	31.57	30.57	28.03	27.15	27.73	27.73	28.78	28.78	28.78	28.78	28.78	28.78
	Means	67.64	67.40	58.75	57.37	57.01	61.78	61.66	61.66	61.66	61.66	61.66	61.66	Means	30.45	29.90	27.19	26.68	26.73	26.37	27.97	27.97	27.97	27.97	27.97	27.97	

TORONTO METEOROLOGICAL RESULTS.

TABLE II.

Monthly and Annual Means of the Temperature of the Air, furnished by six daily observations, from 1860 to 1862 inclusive.

	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Year.
1860	23°.38	22°.83	31°.48	35°.55	55°.53	63°.16	63°.92	61°.46	55°.34	47°.25	37°.95	27°.00	44.32
1861	19°.86	26.06	26.92	42.02	47.50	61.29	65.37	65.48	59.07	48.74	37.14	31.13	44.22
1862	21.71	22.50	28.70	33.66	52.17	60.62	66.70	67.60	59.59	48.70	35.58	28.78	44.55
Mean 1860 to 1862 ..	21.65	22.80	30.06	40.38	51.73	61.66	65.33	65.85	58.00	48.23	36.89	27.97	44.30
Normal Means	24.85	23.69	30.23	41.22	51.55	61.09	66.33	65.72	57.42	44.99	36.14	27.06	44.19
Difference from Normal..... ..	-3.20	+0.11	-0.17	-0.84	+0.18	+0.57	-1.00	+0.13	+0.58	+3.24	+0.75	+0.91	+0.11
Mean 1854 to 1859 ..	22.46	20.62	29.13	40.35	51.45	61.26	68.80	65.87	58.43	46.24	36.57	24.79	43.83

TABLE III.

Monthly Means of the Temperature of the Air at each of the six observation hours, for the period 1860 to 1862 inclusive.

Toronto Astronomical Time.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Year.
2h	21°.73	26.91	34°.64	45°.30	62°.64	62°.64	71°.62	72°.33	64°.17	53°.26	40°.74	30°.95	49.16
4h	24.31	26.69	34.71	45.16	57.53	67.40	71.62	71.95	63.54	52.13	39.51	29.90	48.71
10h	21.36	23.19	29.39	39.02	49.26	58.75	62.11	62.98	55.48	47.05	35.85	27.19	42.64
12h	20.21	22.54	28.60	38.06	47.43	57.37	60.74	61.65	54.10	46.26	35.11	26.68	41.52
18h	19.51	21.34	25.74	35.79	47.07	57.01	60.41	60.48	52.96	44.03	34.70	26.73	40.48
20h	19.79	22.10	27.70	39.00	51.46	61.78	65.48	65.68	57.64	46.67	35.42	26.37	43.26

TABLE IV.

Monthly Mean Abnormal Variations of Temperature, without regard to sign, or mean differences without regard to sign between the normal temperature of the day and hour and the observed temperature of the same day and hour, for each month of the years 1860 to 1862.

	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Year.
1860	9.2	10.4	8.3	6.6	6.1	4.0	4.4	4.7	6.1	8.0	5.8	6.2	6.4
1861	8.1	8.9	8.1	4.9	6.3	5.2	4.8	4.3	4.6	5.7	3.4	10.1	6.1
1862	7.9	7.0	4.4	6.0	6.6	4.8	4.4	4.9	5.8	6.7	5.3	8.5	6.9
Mean 1860 to 1862 ..	8.4	8.8	6.9	6.6	6.7	4.7	4.6	4.6	6.6	6.1	4.8	8.2	6.1
Mean 1854 to 1859 ..	9.4	10.1	7.9	6.5	5.1	5.7	5.6	4.5	5.9	6.0	5.8	8.8	6.7
Mean 1854 to 1862 ..	9.1	9.7	7.6	6.5	5.3	5.4	5.2	4.5	5.8	6.0	5.5	8.6	6.5

TABLE V.

Monthly Mean Abnormal Variations of Temperature, without regard to sign, at each observation hour, for the period 1860-1862.

Year.	Toronto Astronomical Time.	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.	
														1860 to 1862	1854 to 1859
44.32	2h	2.6	2.9	2.0	2.2	2.4	2.6	2.9	2.6	2.7	2.6	2.5	2.9	2.2	2.8
44.22	4	7.6	7.7	6.9	6.1	6.3	5.6	6.0	4.6	5.0	5.3	4.3	8.0	6.2	6.6
44.35	10	8.5	9.3	6.4	5.5	5.5	4.3	3.7	4.7	5.8	6.0	4.5	8.5	6.1	6.5
44.30	12	8.7	9.0	6.5	5.4	5.4	4.3	4.3	4.9	5.8	6.8	4.8	8.4	6.2	6.7
44.39	18	9.0	9.6	7.6	4.7	5.4	4.1	3.7	5.0	5.4	6.6	6.6	8.4	6.3	6.9
44.19	20	8.8	9.1	7.4	5.0	5.1	4.2	3.7	4.1	4.7	5.8	5.3	8.2	6.0	6.7

TABLE VI.

Half-yearly and Yearly Mean Abnormal Variations of Temperature, without regard to sign, for the six observation hours.

Year.	Toronto Astronomical Time.	2h	4h	10h	12h	18h	26h	2h and 4h	10h, 12h, 18h, and 26h.			All hours.
									Winter	Summer	Year	
1854 to 1859	Winter	2.6	2.4	2.9	2.1	2.6	2.4	2.48	2.25	2.00	2.37	2.00
	Summer	5.9	5.8	5.1	5.2	5.1	5.1	5.90	5.13	5.37	5.69	5.18
	Year	6.8	6.6	6.5	6.7	6.8	6.7	6.69	6.69	6.69	6.69	6.68
1860 to 1862	Winter	6.7	6.6	7.3	7.4	7.8	7.4	6.68	7.48	7.20	7.48	7.20
	Summer	5.7	5.7	4.9	5.0	4.7	4.5	5.72	4.78	5.08	5.72	5.08
	Year	6.2	6.2	6.1	6.2	6.2	5.9	6.20	6.13	6.13	6.13	6.13
1854 to 1862	Winter	7.3	7.1	7.7	7.9	8.4	8.1	7.22	7.09	7.75	7.09	7.75
	Summer	5.9	5.8	5.0	5.2	5.0	4.9	5.81	5.01	5.30	5.01	5.30
	Year	6.6	6.5	6.4	6.5	6.7	6.5	6.53	6.50	6.53	6.50	6.53

TABLE VII.

Probable Variability of the Monthly Means of Temperature at each of the six observation hours, in a single year, together with their half-yearly and yearly averages, from the years 1854 to 1862 inclusive.

Year.	Means.	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.		
														Winter.	Summer	Year.
44.32	2h	3.49	2.47	2.43	1.94	2.38	2.20	2.26	1.66	1.95	1.69	1.45	3.12	2.44	2.08	2.26
44.22	4	3.31	2.54	2.59	1.76	2.36	2.13	2.09	1.37	1.71	1.46	1.30	3.10	2.38	1.90	2.14
44.35	10	3.53	3.52	2.69	1.66	1.82	1.76	1.36	1.10	1.21	1.54	1.26	3.02	2.59	1.47	2.03
44.30	12	3.67	3.85	2.76	1.52	1.76	1.88	1.33	1.14	1.07	1.56	1.30	3.04	2.70	1.45	2.07
44.39	18	3.90	3.65	2.98	1.32	1.72	1.85	1.59	1.09	1.25	1.48	1.24	3.20	2.74	1.47	2.11
44.19	20	3.89	3.57	2.85	1.38	1.95	1.99	1.67	1.01	1.26	1.59	1.28	3.12	2.71	1.54	2.13
44.1	2-1	3.40	2.50	2.51	1.85	2.37	2.16	2.22	1.52	1.83	1.58	1.37	3.11	2.41	1.99	2.20
44.7	10-20	3.75	3.65	2.82	1.45	1.81	1.87	1.49	1.08	1.20	1.54	1.26	3.10	2.69	1.48	2.08
44.5	All hours	3.63	3.27	2.72	1.58	2.00	1.97	1.73	1.23	1.11	1.55	1.30	3.10	2.59	1.65	2.12

TORONTO METEOROLOGICAL RESULTS.

TABLE VIII.

Mean Abnormal Variations of Temperature, with their proper signs, arranged according to the direction of the wind at the hours of observation, for each year and for the winter and summer half-years, from 1860 to 1862, the summer being understood to begin April 1st and end September 30th.

	1860.			1861.			1862.			1860 to 1862.		
	Winter.	Summer.	Year.	Winter.	Summer.	Year.	Winter.	Summer.	Year.	Winter.	Summer.	Year.
N.	-3.77	+0.78	-1.57	-6.21	-1.37	-3.69	-5.57	-0.57	-3.18	-5.52	-0.30	-2.78
N. E.	+2.15	+1.49	+1.73	+1.44	-1.60	+0.08	-2.42	-0.94	-1.71	+0.19	-0.14	+0.02
E.	+6.22	-0.06	+3.01	+2.67	-0.73	+1.01	+1.28	-1.23	-0.16	+3.24	-0.75	+1.13
S. E.	+4.87	-2.48	+1.09	+3.16	-0.43	+0.98	+5.59	-0.05	+2.06	+3.43	-0.83	+1.35
S.	+5.60	+0.74	+1.58	+3.77	+0.17	+1.34	+3.08	+1.17	+1.74	+3.90	+0.73	+1.56
S. W.	+4.34	+1.89	+3.53	+5.00	+2.34	+4.07	+6.18	+3.61	+5.05	+5.11	+2.70	+4.21
W.	-0.27	-0.38	-0.31	-0.54	+0.22	-0.30	+0.57	+1.91	+1.07	-0.69	+0.56	+0.14
N. W.	-2.91	-2.34	-2.63	-3.25	-1.66	-2.42	-3.55	+0.03	-1.73	-3.22	-1.36	-2.27
Calms	-2.04	-0.96	-1.65	+0.67	-0.04	+0.16	+0.63	+0.69	+0.66	-0.23	-0.02	-0.10

TABLE IX.

Mean Abnormal Variations of Temperature, with their proper signs, arranged according to the direction of the wind at the hours of observation, in each month and in the year, for the period 1860 to 1862.

	Jan.	Feb.	March	April	May	June	July	Aug.	Sept	Oct.	Nov.	Dec.	Year.	
	1860 to 1862 1864 to 1869													
N.	-14.07	-7.48	-8.00	-3.12	+5.01	+1.36	-3.35	-0.10	-0.96	+0.77	-1.81	-7.65	-2.78	-2.80
N. E.	-3.73	+4.69	-3.15	-1.86	+1.76	-0.07	-2.15	-0.69	-0.51	+2.97	+0.01	+1.03	+0.02	-1.81
E.	+2.36	+2.51	+0.37	+3.37	-0.87	-3.04	-2.60	-0.12	+0.83	+5.10	+4.04	+6.01	+1.13	+1.73
S. E.	+3.54	+0.65	+2.14	-0.23	+0.20	-3.20	-2.02	-0.55	+0.09	+6.69	+4.48	+6.79	+1.35	+1.79
S.	+4.56	+5.36	+2.10	-2.50	+1.78	+1.26	-0.79	0.00	+3.39	+4.15	+3.98	+2.31	+1.56	+2.89
S. W.	+2.49	+6.19	+3.82	+2.54	+0.91	+1.67	+2.17	+2.75	+5.27	+8.37	+1.13	+8.74	+4.21	+3.45
W.	-2.82	+0.90	+1.78	-1.83	+0.15	+1.42	+0.65	+2.60	+0.97	+1.41	-0.65	-0.85	+0.14	-2.18
N. W.	-6.49	-6.37	-3.33	-3.22	-0.85	+2.44	-1.94	-1.99	-3.41	-0.16	-0.25	-2.71	-2.27	-3.51
Calms	-4.81	-1.13	+2.03	-1.38	-0.96	-0.67	+1.49	+0.23	-0.18	+2.05	+0.08	+0.81	-0.10	+1.33

TABLE X.

Showing, for the period 1853 to 1862, the number of times in each month in which the temperature at the hour of observation differed from the normal to the extent of 15° and upwards, with the average value of the corresponding deviation.

Relatively high tem- peratures.	Jan.	Feb.	March	April	May	June	July	Aug.	Sept	Oct.	Nov.	Dec.	
	Number	47	83	55	15	21	35	21	12	30	53	37	46
	Ratio to mean	1.21	2.19	1.15	0.10	0.55	0.92	0.55	0.32	0.79	1.40	0.98	1.21
Relatively low tem- peratures.	Average excess	17.5	18.3	17.6	16.9	17.9	15.5	16.5	17.4	16.9	17.7	17.1	18.8
	Number	210	227	134	24	26	15	9	5	16	8	28	178
	Ratio to mean	3.16	2.99	1.77	0.32	0.34	0.20	0.12	0.07	0.21	0.11	0.37	2.35
Relatively low tem- peratures.	Average defect	22.9	21.5	20.4	17.3	17.7	18.0	15.5	16.6	16.6	16.3	18.8	20.9

TABLE XI.
Relative frequency of the several Winds during extraordinarily high temperatures, when the temperature at the hour of observation exceeded the norm by 15° and upwards, with the average excess of temperature for each wind, deduced from observations in the ten years 1853 to 1862.

		Duration of each wind without regard to temperature in terms of the average duration of all winds, from Table LXVI.		Relative frequency of each wind during high temperatures, or ratios of (1) to (2).		Ratios of the numbers in the four preceding columns to their respective means for all winds.				Average excess for each wind.	
(1)		(2)		(3)		(4)		(5)			
Wind	Number	Wind	Number	Wind	Number	Wind	Number	Wind	Number	Wind	Number
N.	5	0	5	1.18	1.03	1.01	1.07	4.2	0.0	2.0	4.7
N.N.E.	3	2	3	0.88	0.63	0.67	0.77	3.4	3.2	3.0	3.9
N.E.	8	0	5	0.87	0.76	0.62	0.77	9.2	0.0	6.5	6.5
E.N.E.	16	8	1	7	0.75	1.45	0.98	1.02	21.3	5.5	1.0
E.	16	3	0	18	0.85	1.73	1.34	1.13	18.9	1.7	0.0
E.S.E.	7	0	1	2	0.40	0.75	0.78	0.56	17.3	0.0	1.3
S.E.	4	1	1	6	0.23	0.47	0.58	0.42	17.6	2.1	1.7
S.S.E.	8	3	2	1	0.24	0.42	0.72	0.48	33.9	7.2	2.8
S.	9	3	11	5	0.24	0.66	1.20	0.74	37.5	4.6	8.5
S.S.W.	16	12	13	17	0.71	1.01	1.53	1.14	22.5	11.9	8.5
S.W.	37	10	7	18	1.46	0.88	0.97	1.15	25.3	11.3	7.2
W.S.W.	19	8	5	8	2.29	0.79	0.57	1.23	8.3	10.1	8.8
W.	8	12	9	6	1.97	1.13	0.79	1.26	4.1	10.6	11.4
W.N.W.	3	11	2	5	1.31	1.45	1.06	1.37	2.3	7.6	1.9
N.W.	5	10	6	1	1.23	1.54	1.41	1.21	4.1	6.5	4.3
N.N.W.	0	2	3	6	1.40	1.51	1.57	1.25	0.0	1.3	1.9
Calms.	12	6	3	7	0.99	0.79	1.11	1.43	12.1	7.6	2.7

TORONTO METEOROLOGICAL RESULTS.

TABLE XII.

Relative frequency of the several winds during extraordinarily low temperatures, when the temperature at the hour of observation was below the normal by 1.5° and upwards, with the average effect of temperature for each wind, deduced from observations in the ten years 1853 to 1862.

	Duration of each wind with out regard to temperature, in terms of the average dura- tion of all winds, from Table LXVI.		Relative frequency of each wind during low tem- peratures, or ratios of (1) to (2).				Ratio of the numbers in the four preceding columns to their respective means for all winds.				Average effect for each wind.	
	(1)	(2)	Wind. Number.	Wind. Number.	Wind. Number.	Wind. Number.	Wind. Number.	Wind. Number.	Wind. Number.	Wind. Number.	Average.	
N.	116	9	1	1.18	1.03	1.01	1.07	98.0	8.8	1.0	11.2	
S.N.E.	56	10	1	0.88	0.63	0.67	0.77	63.8	15.9	1.5	6.5	
N.E.	42	5	0	0.87	0.76	0.62	0.77	48.3	6.6	4.8	0	
E.N.E.	3	8	3	2	0.75	1.45	0.98	1.02	4.0	5.5	3.1	
E.	2	5	3	3	0.85	1.73	1.34	1.13	2.4	2.9	2.2	
E.S.E.	0	0	2	0.40	0.75	0.78	0.56	0.0	0.0	3.6	0.0	
S.E.	2	0	0	0.23	0.47	0.58	0.42	8.8	0.0	0.0	0.29	
S.S.E.	0	0	0	0.24	0.42	0.72	0.48	0.0	0.0	0.0	0.0	
S.	0	2	0	0.24	0.66	1.29	0.74	0.0	1.5	0.0	0.0	
S.S.W.	0	3	0	1	0.71	1.01	1.53	1.14	0.0	3.0	0.0	
S.W.	13	0	0	1.46	0.88	0.97	1.15	8.9	0.0	0.0	0.29	
W.S.W.	60	8	1	2.29	0.79	0.57	1.23	26.2	10.1	1.7	9.8	
W.	84	13	1	0	1.97	1.13	0.79	1.26	42.7	11.5	1.3	
W.N.W.	59	38	2	5	1.31	1.45	1.06	1.37	45.0	26.2	1.9	
N.W.	62	42	6	5	1.23	1.54	1.41	1.21	50.3	27.3	4.3	
N.N.W.	112	35	5	5	1.40	1.51	1.58	1.25	80.2	23.1	3.2	
Calms	33	8	1	0	0.99	0.79	1.11	1.43	33.3	10.1	0.9	

TABLE XIII.

Mean differences without regard to sign between the temperatures of the air at 6 a.m. on consecutive days, for each month in the years 1860 to 1862, the effect of annual variation being eliminated.

	Jan.	Feb.	March.	April.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
1860	8.4	9.6	5.3	5.2	4.0	3.9	4.2	4.3	6.3	5.1	5.8	6.4	5.7
1861	8.1	9.0	9.3	3.9	3.7	4.8	3.9	3.8	6.5	5.5	4.9	7.6	5.9
1862	9.4	10.2	4.7	4.1	6.3	4.7	4.1	5.4	6.4	8.2	5.9	8.9	6.5
Means	8.6	9.6	6.4	4.4	4.7	4.5	4.1	4.5	6.4	6.3	5.5	7.6	6.1
1860 to 1862	10.0	10.0	6.7	4.9	4.2	3.9	3.7	4.2	6.2	6.8	7.2	8.3	6.3
1854 to 1859	10.0	10.0	6.7	4.9	4.2	3.9	3.7	4.2	6.2	6.8	7.2	8.3	6.3

TABLE XIV.

Mean change of temperature with its proper sign, from 6 a.m. to 6 a.m., for the period 1860 to 1862, arranged according to the daily resultant direction of the wind, the effect of annual variation being eliminated.

	Jan.	Feb.	March.	April.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.	
	1860 to 1862												1854 to 1859	
N.	-3.0	-2.2	-3.4	-2.3	-0.1	-2.0	-2.3	-3.0	-3.7	-2.0	-0.8	+1.4	-1.9	-3.3
N.E.	+6.8	+4.1	+2.7	+3.0	+0.4	+0.3	+0.6	-1.2	+0.7	+1.3	+2.3	+4.8	+2.4	+1.5
E.	+10.6	+12.9	+5.1	+2.2	+1.1	-0.3	+1.9	+1.7	+7.3	+4.8	+1.4	+8.1	+4.0	+3.5
S.E.	*	+5.1	-4.8	+5.5	+2.0	+6.6	+2.2	+8.1	+9.0	+3.2	+11.6	+10.9	+6.3	+4.6
S.	-3.8	+7.8	+8.4	+5.8	+1.4	+4.7	+2.7	+1.1	+6.1	+14.1	+6.0	+0.6	+3.9	+3.9
S.W.	+5.3	+3.4	+5.7	+2.5	+3.5	+4.0	+2.5	+1.8	-0.2	+1.3	-0.3	+5.2	+2.7	+2.2
W.	-3.5	-2.7	-2.2	-4.4	-2.6	-2.2	-3.2	-3.5	-3.7	-2.5	-5.0	-3.2	-2.9	-2.9
N.W.	-12.4	-7.4	-4.5	-4.2	-2.6	-2.1	-2.3	-4.3	-6.6	-5.1	-3.5	-4.3	-4.5	-4.5

* No case of a S.E. resultant occurred in January.

TABLE XV.

Highest and lowest temperatures in each month, for the years 1860, 1861, and 1862.

MAXIMA.

	Jan.	Feb.	March.	April.	May.	June.	July.	Aug.	Sept.	Ott.	Nov.	Dec.	Year.
	Tempert.												Date.
1860	46.4	50.2	67.0	61.8	74.5	81.6	88.0	87.0	75.8	68.0	64.5	39.0	88.0 July 19.
1861	37.0	46.0	47.4	67.0	73.0	87.8	84.5	85.2	78.8	71.0	52.4	35.2	87.8 June 9.
1862	44.5	37.8	43.2	68.0	78.5	85.4	95.5	89.5	79.4	76.6	58.0	50.1	95.5 July 6.
Means	42.6	44.7	52.5	65.6	75.3	84.9	89.3	87.2	78.0	71.9	58.3	48.1	90.4
1860 to 1862	42.6	44.7	52.5	65.6	75.3	84.9	89.3	87.2	78.0	71.9	58.3	48.1	90.4
1854 to 1859	42.9	43.4	52.2	64.7	75.9	87.6	91.2	86.6	82.2	70.8	57.5	46.7	92.5

MINIMA.

	Jan.	Feb.	March.	April.	May.	June.	July.	Aug.	Sept.	Ott.	Nov.	Dec.	Tempert.
	Date.												
1860	-6.8	-8.5	12.8	10.5	32.5	49.2	45.8	46.8	28.7	28.4	13.2	-7.0	-8.5 Feb. 1.
1861	-11.2	-20.8	-5.2	23.8	28.0	41.6	47.0	47.0	37.1	29.0	23.0	5.5	-20.8 Feb. 8.
1862	-2.6	-5.2	8.0	14.5	32.4	39.4	48.2	42.8	39.0	26.2	16.2	3.4	-5.2 Feb. 15.
Means	-6.9	11.5	5.2	19.3	31.0	43.4	46.3	45.5	34.9	27.9	17.5	-1.6	-11.5
1860 to 1862	-6.9	11.5	5.2	19.3	31.0	43.4	46.3	45.5	34.9	27.9	17.5	-1.6	-11.5
1854 to 1859	-10.5	11.0	-1.8	15.9	31.0	37.2	47.5	43.8	34.9	25.4	13.6	-3.1	18.1

TORONTO METEOROLOGICAL RESULTS.

TABLE XVI.
Monthly Means of the Barometric Pressure, at each of the six observation hours, from 1850 to 1862 inclusive.
 Barometer at $35^{\circ} = 27$ inches + the numbers in the table.

Harmonized at $\omega_0 = 20$ nodes											
	Toronto		Astronomical Time		Astronomical Time		Toronto		Astronomical Time		
	2 ^a	4 ^b	10 ^b	12 ^b	1 ^b	2 ^b	4 ^b	10 ^b	12 ^b	1 ^b	2 ^b
January	1860	2,624	2,646	2,641	2,654	2,672	2,643	2,656	2,676	2,657	2,651
"	1861	2,630	2,663	2,661	2,652	2,664	2,652	2,655	2,654	2,667	2,664
"	1862	2,639	2,713	2,746	2,746	2,758	2,727	2,730	2,749	2,752	2,751
Means	2,650	2,659	2,685	2,683	2,677	2,691	2,674	2,684	2,694	2,687	2,686
July	1860	2,624	2,638	2,653	2,658	2,672	2,643	2,656	2,676	2,657	2,651
"	1861	2,633	2,652	2,658	2,656	2,674	2,652	2,655	2,674	2,657	2,651
"	1862	2,652	2,697	2,739	2,620	2,729	2,678	2,727	2,740	2,757	2,756
Means	2,675	2,675	2,697	2,697	2,688	2,701	2,674	2,695	2,704	2,697	2,696
August	1860	2,624	2,638	2,653	2,658	2,672	2,643	2,656	2,676	2,657	2,651
"	1861	2,633	2,652	2,658	2,656	2,674	2,652	2,655	2,674	2,657	2,651
"	1862	2,652	2,697	2,739	2,620	2,729	2,678	2,727	2,740	2,757	2,756
Means	2,675	2,675	2,697	2,697	2,688	2,701	2,674	2,695	2,704	2,697	2,696
September	1860	2,624	2,638	2,653	2,658	2,672	2,643	2,656	2,676	2,657	2,651
"	1861	2,633	2,652	2,658	2,656	2,674	2,652	2,655	2,674	2,657	2,651
"	1862	2,652	2,697	2,739	2,620	2,729	2,678	2,727	2,740	2,757	2,756
Means	2,675	2,675	2,697	2,697	2,688	2,701	2,674	2,695	2,704	2,697	2,696
October	1860	2,624	2,638	2,653	2,658	2,672	2,643	2,656	2,676	2,657	2,651
"	1861	2,633	2,652	2,658	2,656	2,674	2,652	2,655	2,674	2,657	2,651
"	1862	2,652	2,697	2,739	2,620	2,729	2,678	2,727	2,740	2,757	2,756
Means	2,675	2,675	2,697	2,697	2,688	2,701	2,674	2,695	2,704	2,697	2,696
November	1860	2,624	2,638	2,653	2,658	2,672	2,643	2,656	2,676	2,657	2,651
"	1861	2,633	2,652	2,658	2,656	2,674	2,652	2,655	2,674	2,657	2,651
"	1862	2,652	2,697	2,739	2,620	2,729	2,678	2,727	2,740	2,757	2,756
Means	2,675	2,675	2,697	2,697	2,688	2,701	2,674	2,695	2,704	2,697	2,696
December	1860	2,624	2,638	2,653	2,658	2,672	2,643	2,656	2,676	2,657	2,651
"	1861	2,633	2,652	2,658	2,656	2,674	2,652	2,655	2,674	2,657	2,651
"	1862	2,652	2,697	2,739	2,620	2,729	2,678	2,727	2,740	2,757	2,756
Means	2,675	2,675	2,697	2,697	2,688	2,701	2,674	2,695	2,704	2,697	2,696

TABLE XVII.

*Monthly and annual means of the Barometric Pressure furnished by six daily observations, from 1860 to 1862.*Barometer at temperature $32^{\circ} = 27$ inches + the numbers in the table.

	Jan.	Feb.	March.	April.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
1860	2.643	2.632	2.511	2.578	2.566	2.498	2.564	2.582	2.673	2.671	2.523	2.667	2.592
1861	2.652	2.544	2.621	2.561	2.545	2.570	2.551	2.652	2.608	2.619	2.537	2.746	2.601
1862	2.727	2.608	2.504	2.726	2.590	2.564	2.547	2.610	2.68	2.619	2.636	2.678	2.625
Means 1860 to 1862	2.674	2.593	2.545	2.623	2.567	2.544	2.554	2.617	2.655	2.636	2.565	2.697	2.606
Means 1854 to 1859	2.667	2.639	2.538	2.572	2.596	2.544	2.614	2.606	2.675	2.653	2.595	2.670	2.614

TABLE XVIII.

*Monthly means of the Barometric Pressure at each of the six observation hours, for the period 1860 to 1862.*Barometer at temperature $32^{\circ} = 27$ inches + the numbers in the table.

Toronto Astronomical Time.	Jan.	Feb.	March.	April.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
2 ^h	2.650	2.575	2.525	2.611	2.560	2.536	2.544	2.607	2.640	2.616	2.58	2.682	2.591
4	2.639	2.575	2.523	2.599	2.548	2.523	2.535	2.593	2.633	2.617	2.556	2.693	2.588
10	2.685	2.597	2.557	2.620	2.568	2.541	2.561	2.618	2.664	2.640	2.574	2.704	2.611
12	2.683	2.597	2.551	2.621	2.567	2.540	2.562	2.620	2.666	2.630	2.568	2.701	2.610
18	2.677	2.608	2.551	2.636	2.573	2.557	2.554	2.627	2.658	2.646	2.566	2.692	2.610
20	2.691	2.617	2.564	2.648	2.587	2.567	2.567	2.637	2.669	2.660	2.580	2.708	2.625

TABLE XIX.

Monthly mean abnormal variations of the Barometer without regard to sign, for each month of the years 1860 to 1862.

	Jan.	Feb.	March.	April.	May.	June.	July.	August.	Sept.	Oct.	Nov.	Dec.	Year.
1860	0.207	0.238	0.215	0.207	0.139	0.185	0.124	0.122	0.143	0.167	0.231	0.334	0.193
1861	.206	.262	.222	.194	.214	.118	.109	.103	.163	.197	.195	.227	0.184
1862	.229	.191	.192	.229	.112	.139	.155	.118	.125	.153	.204	.212	0.170
Means 1860 to 1862	0.211	0.230	0.210	0.209	0	0.147	0.123	0.114	0.144	0.172	0.210	0.257	0.182
Means 1854 to 1859	0.219	0.215	0.223	0.196	0.151	0	0	0.121	0.119	0.162	0.183	0.228	0.225
Means 1854 to 1862	0.237	0.220	0.219	0.200	0.132	0.135	0.122	0.117	0.156	0.179	0.222	0.236	0.183

TORONTO METEOROLOGICAL RESULTS.

TABLE XX.

Monthly Mean Abnormal Variations of the Barometer, without regard to sign, at each of the six observation hours, for the period 1860 to 1862.

Toronto Astronomical Time.	Jan.	Feb.	March.	April.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.	
													1860 to 1862.	1854 to 1859.
2h	.0220	.0228	.0215	.0210	.0155	.0149	.0125	.0113	.0146	.0177	.0207	.0265	.0184	.0187
4	.217	.224	.210	.211	.153	.143	.117	.110	.140	.173	.201	.260	.180	.182
10	.210	.221	.206	.210	.152	.135	.110	.112	.134	.170	.206	.265	.178	.175
12	.205	.225	.207	.212	.154	.138	.112	.115	.137	.172	.212	.260	.179	.177
18	.215	.240	.206	.205	.159	.160	.135	.118	.151	.168	.217	.245	.185	.188
20	.218	.241	.213	.205	.159	.159	.137	.117	.155	.173	.218	.248	.187	.191

TABLE XXI.

Probable Variability of the Monthly Means of the Barometer at each hour, in a single year, derived from observations in the nine years, 1854 to 1862.

Toronto Astronomical Time.	January.	February	March.	April.	May.	June.	July.	August.	September	October.	November	December	Year.	
													1860	1854
2h	.0267	.0550	.0453	.0492	.0301	.0393	.0245	.0294	.0300	.0343	.0540	.0330	.0376	
4	.0251	.0521	.0452	.0487	.0287	.0388	.0229	.0280	.0280	.0330	.0533	.0325	.0364	
10	.0271	.0574	.0455	.0471	.0246	.0391	.0214	.0278	.0271	.0356	.0539	.0296	.0355	
12	.0289	.0456	.0455	.0480	.0254	.0388	.0214	.0279	.0278	.0372	.0553	.0299	.0360	
18	.0337	.0514	.0453	.0490	.0336	.0408	.0307	.0296	.0316	.0315	.0591	.0374	.0395	
20	.0325	.0581	.0461	.0499	.0353	.0420	.0283	.0294	.0315	.0323	.0568	.0389	.0395	
Means0291	.0508	.0455	.0486	.0294	.0398	.0219	.0287	.0293	.0340	.0554	.0355	.0374	

TABLE XXII.

Mean Abnormal Variations of the Barometer, with their proper signs, arranged according to the direction of the wind, for each year and for the winter and summer half years, from 1860 to 1862, the summer being considered to begin April 1st, and end September 30th.

	1860.			1861.			1862.			Means 1860-1862.		
	Winter.	Summer.	Year.	Winter.	Summer.	Year.	Winter.	Summer.	Year.	Winter.	Summer.	Year.
N.	+0.155	+0.008	+0.062	+0.063	+0.024	+0.043	+0.136	+0.066	+0.103	+0.114	+0.032	+0.071
N.E.	+.034	-.041	-.013	-.005	-.031	-.016	+.075	+.068	+.086	+.093	+.006	+.020
E.	+.029	-.012	+.008	-.02	+.004	+.001	+.034	+.041	+.038	+.019	+.015	+.017
S.E.	+.030	+.055	+.043	+.042	+.027	+.033	+.067	+.090	+.053	+.010	+.055	+.037
S.	-.055	+.026	+.012	-.020	+.035	+.014	-.033	+.018	+.003	-.037	+.026	+.009
S.W.	-.171	-.158	-.167	-.063	-.092	-.073	-.128	-.075	-.105	-.119	-.104	-.114
W.	-.069	-.131	-.083	-.037	-.175	-.082	-.056	-.042	-.051	-.055	-.115	-.076
N.W.	-.009	-.017	-.013	-.031	-.019	-.011	-.027	-.010	-.009	-.011	-.024	-.005
Calms.	+.012	+.018	+.033	+.015	+.030	+.018	+.062	+.049	+.056	+.027	+.032	+.030

TABLE XXIII.
Mean Abnormal Variations of the Barometer, with their proper signs, arranged according to the direction of the wind, in each month and in the year, for the period 1860-1862.

	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.	
	1860	1862	1859	1860	1862	1859	1860	1862	1859	1860	1862	1859	1860 to 1862	1853 to 1859
N.	+ 0.152	+ 0.105	+ 0.018	+ 0.031	+ 0.054	+ 0.021	- 0.021	+ 0.048	+ 0.045	+ 0.192	+ 0.021	+ 0.214	+ 0.071	+ 0.069
N.E.	+ .031	- .109	+ .024	+ .010	- .005	- .002	- .036	- .003	+ .000	+ .000	+ .110	+ .020	+ .052	+ .015
E.	+ .148	- .084	+ .010	+ .198	+ .047	+ .084	- .045	+ .004	+ .005	+ .009	+ .057	+ .140	+ .017	+ .015
E.E.	- .126	+ .107	+ .135	- .022	+ .070	+ .042	+ .061	+ .010	+ .027	+ .054	+ .151	+ .150	+ .037	+ .041
S.E.	- .045	- .207	- .118	- .042	- .120	- .111	- .082	- .129	- .108	- .203	- .028	- .114	- .115	- .115
S.	+ .076	+ .007	+ .144	- .005	- .181	- .174	- .114	- .162	- .089	- .163	- .097	- .044	+ .010	+ .010
S.W.	+ .084	+ .027	+ .028	+ .028	+ .018	+ .007	+ .000	+ .025	+ .003	+ .002	+ .015	+ .005	+ .017	+ .017
W.	+ .073	+ .140	+ .012	+ .011	+ .041	+ .021	+ .017	+ .021	+ .030	+ .020	+ .109	+ .138	+ .030	+ .030
Calms														

TABLE XXIV.
Mean Differences without regard to sign between the barometric pressures at 6 a.m. on consecutive days, for each month in the years 1860 to 1862.

	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.	
	1860	1862	1859	1860	1862	1859	1860	1862	1859	1860	1862	1859	1860 to 1862	1853 to 1859
1860	0.255	0.248	0.174	0.223	0.123	0.135	0.188	0.114	0.184	0.144	0.258	0.247	0.189	0.208
1861	- .254	- .303	- .161	- .161	- .161	- .161	- .161	- .161	- .161	- .161	- .161	- .161	- .255	- .255
1862	.351	.289	.360	.291	.351	.356	.162	.142	.166	.166	.211	.184	.255	.201
Means 1860 to 1862	0.282	0.280	0.235	0.195	0.160	0.148	0.144	0.120	0.163	0.192	0.220	0.249	0.199	
Means 1854 to 1856	0.280	0.253	0.257	0.217	0.160	0.126	0.110	0.128	0.151	0.168	0.248	0.273	0.198	
Means 1854 to 1862	0.281	0.265	0.250	0.210	0.160	0.133	0.121	0.125	0.155	0.176	0.239	0.265	0.198	

TABLE XXV.
Mean change in the Barometer with its proper sign, from 6 a.m. to 6 a.m., for the period 1860 to 1862, arranged according to the daily resultant direction of the wind.

	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.	
	1860	1862	1859	1860	1862	1859	1860	1862	1859	1860	1862	1859	1860 to 1862	1854 to 1859
N.	+ 0.107	+ 0.093	+ 0.194	+ 0.105	- 0.116	+ 0.137	+ 0.106	- 0.031	+ 0.094	- 0.000	+ 0.086	+ 0.067	+ 0.076	+ 0.097
N.E.	- .237	- .347	- .170	- .120	- .074	+ .061	- .081	+ .016	- .055	- .060	- .172	- .233	- .137	- .162
E.	- .548	- .352	- .267	- .151	- .125	- .135	- .097	- .071	- .237	- .107	- .254	- .246	- .196	- .170
E.E.	- .352	- .312	- .312	- .318	- .303	- .303	- .303	- .303	- .303	- .303	- .303	- .303	- .303	- .303
S.E.	- .674	- .265	- .312	- .114	- .097	- .072	- .100	- .117	- .045	- .065	- .246	- .170	- .174	- .174
S.	+ .094	+ .106	+ .120	+ .120	+ .097	+ .148	+ .099	+ .150	+ .097	+ .154	+ .174	+ .170	- .130	- .069
F.W.	+ .094	+ .106	+ .120	+ .120	+ .097	+ .148	+ .099	+ .150	+ .097	+ .154	+ .174	+ .170	- .130	- .069
W.	+ .192	+ .170	+ .112	+ .213	+ .156	+ .050	+ .101	+ .048	+ .180	+ .119	+ .228	+ .134	+ .142	+ .144
N.W.	+ .234	+ .330	+ .183	+ .148	+ .154	+ .139	+ .133	+ .192	+ .156	+ .248	+ .163	+ .233	+ .183	+ .170

* No case of a S.E. resultant occurred in January.

TABLE XXVI.
Highest and lowest Readings of the Barometer in each month, for the years 1860, 1861, and 1862.
Barometer at 32° = 27 inches + the numbers in the Table.

HIGHEST.

Year.	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Barometer.	Date.
	1860	1862	1859	1860	1862	1859	1860	1862	1859	1860	1862	1859	1860 to 1862	1854 to 1859
1860	5.142	3.150	2.034	3.265	2.846	2.859	2.839	2.043	3.170	2.982	2.959	3.267	December 14.	
1861	3.330	3.144	3.200	3.129	2.955	2.810	2.830	2.912	3.104	3.654	3.000	3.330	January 22.	
1862	3.300	3.138	2.824	3.117	2.942	3.109	2.957	2.977	3.031	3.039	3.469	3.453	November 15.	
Means.....	3.257	3.139	2.987	3.167	2.928	2.920	2.873	2.977	3.102	3.025	3.143	3.301	3.355	

LOWEST.

Year.	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Barometer.	Date.
	1860	1862	1859	1860	1862	1859	1860	1862	1859	1860	1862	1859	1860 to 1862	1854 to 1859
1860	2.153	1.920	2.044	1.890	2.088	1.993	2.157	2.211	2.233	2.019	1.844	1.838	1.838	December 20.
1861	2.006	1.979	2.034	2.055	1.914	2.170	2.200	2.382	2.070	1.998	2.095	2.171	1.644	May 6.
1862	1.965	2.011	1.845	2.076	2.238	2.163	2.193	2.326	2.107	2.047	2.132	2.103	1.805	March 3.
Means.....	2.042	1.970	1.961	2.009	1.990	2.084	2.207	2.306	2.130	2.021	1.991	2.038	1.762	

TORONTO METEOROLOGICAL RESULTS.

TABLE XXVII.
Monthly means of the Pressure of Dry Air, at each of the six observation hours from 1860 to 1862 inclusive.

Pressure of Dry Air at $32^{\circ} = 27$ inches + the numbers in the table.

Toronto Astronomical Time		2 ^h	4 ^h	6 ^h	10 ^h	12 ^h	14 ^h	20 ^h	Means	Toronto Astronomical Time		2 ^h	4 ^h	6 ^h	10 ^h	12 ^h	18 ^h	20 ^h	Means
January .. 1860	2.506	2.510	2.522	2.532	2.543	2.549	2.555	2.533	2.514	July..... 1860	2.114	2.130	2.155	2.167	2.137	2.117	2.137	2.137	
" 1861	2.521	2.531	2.562	2.564	2.547	2.547	2.549	2.562	2.561	" 1861	2.025	2.027	2.110	2.123	2.068	2.065	2.084	2.084	
" 1862	2.548	2.604	2.645	2.647	2.624	2.637	2.624	2.624	2.624	" 1862	2.039	2.037	2.058	2.101	2.098	2.076	2.075	2.075	
Means	2.558	2.548	2.579	2.580	2.575	2.591	2.569	2.569	2.569	Means	2.059	2.065	2.121	2.129	2.119	2.097	2.069	2.069	
February .. 1860	2.488	2.484	2.541	2.543	2.528	2.536	2.520	2.480	2.480	August... 1860	2.069	2.075	2.130	2.140	2.165	2.127	2.119	2.119	
" 1861	2.484	2.488	2.598	2.598	2.496	2.444	2.453	2.414	2.481	" 1861	2.062	2.095	2.185	2.200	2.160	2.169	2.158	2.158	
" 1862	2.483	2.483	2.561	2.561	2.492	2.515	2.525	2.501	2.501	" 1862	2.054	2.042	2.135	2.153	2.149	2.129	2.106	2.106	
Means	2.455	2.454	2.480	2.481	2.496	2.505	2.478	2.478	2.478	Means	2.072	2.071	2.150	2.158	2.173	2.142	2.128	2.128	
March ... 1860	2.329	2.326	2.384	2.382	2.380	2.385	2.363	2.363	2.363	September.. 1860	2.281	2.291	2.367	2.362	2.323	2.332	2.332	2.332	
" 1861	2.472	2.465	2.492	2.492	2.490	2.513	2.528	2.493	2.493	" 1861	2.162	2.162	2.232	2.234	2.226	2.242	2.296	2.296	
" 1862	2.343	2.346	2.387	2.387	2.359	2.381	2.366	2.372	2.372	" 1862	2.217	2.218	2.286	2.304	2.291	2.272	2.295	2.295	
Means	2.378	2.379	2.421	2.420	2.425	2.433	2.409	2.409	2.409	Means	2.220	2.222	2.291	2.302	2.298	2.277	2.290	2.290	
April .. 1860	2.365	2.359	2.388	2.390	2.419	2.427	2.395	2.395	2.395	October .. 1860	2.266	2.271	2.327	2.327	2.325	2.308	2.415	2.399	
" 1861	2.397	2.392	2.350	2.351	2.411	2.410	2.365	2.365	2.365	" 1861	2.292	2.300	2.314	2.314	2.381	2.365	2.327	2.327	
" 1862	2.519	2.518	2.561	2.560	2.552	2.560	2.512	2.512	2.512	" 1862	2.264	2.276	2.319	2.330	2.358	2.365	2.319	2.319	
Means	2.407	2.403	2.430	2.433	2.461	2.466	2.433	2.433	2.433	Means	2.307	2.316	2.350	2.356	2.379	2.332	2.349	2.349	
May .. 1860	2.189	2.180	2.230	2.247	2.265	2.258	2.228	2.228	2.228	November.. 1860	2.201	2.201	2.313	2.329	2.329	2.349	2.327	2.327	
" 1861	2.297	2.292	2.333	2.334	2.312	2.313	2.313	2.313	2.313	" 1861	2.346	2.344	2.363	2.363	2.352	2.359	2.359	2.359	
" 1862	2.302	2.303	2.338	2.336	2.336	2.336	2.337	2.337	2.337	" 1862	2.420	2.446	2.484	2.486	2.464	2.481	2.465	2.465	
Means	2.262	2.258	2.309	2.314	2.312	2.293	2.293	2.293	2.293	Means	2.359	2.371	2.392	2.385	2.369	2.384	2.384	2.384	
June .. 1860	2.053	2.036	2.169	2.120	2.103	2.055	2.084	2.084	2.084	December.. 1860	2.540	2.556	2.663	2.559	2.557	2.553	2.551	2.551	
" 1861	2.157	2.162	2.192	2.192	2.205	2.234	2.216	2.192	2.192	" 1861	2.570	2.579	2.589	2.596	2.605	2.629	2.595	2.595	
" 1862	2.178	2.184	2.227	2.225	2.235	2.263	2.235	2.219	2.219	" 1862	2.513	2.524	2.550	2.547	2.528	2.554	2.536	2.536	
Means	2.129	2.121	2.176	2.187	2.200	2.179	2.165	2.165	2.165	Means	2.541	2.553	2.567	2.557	2.579	2.560	2.560	2.560	

TABLE XXVIII.

*Monthly and annual means of the Pressure of Dry Air, furnished by six daily observations, from 1860 to 1862.*Pressure of Dry Air at $32^{\circ} = 27$ inches + the numbers in the table.

	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
1860	2.533	2.520	2.363	2.393	2.228	2.084	2.137	2.119	2.332	2.309	2.327	2.551	2.334
1861	2.540	2.414	2.493	2.365	2.313	2.102	2.084	2.158	2.209	2.327	2.359	2.595	2.338
1862	2.624	2.501	2.372	2.542	2.337	2.219	2.075	2.108	2.265	2.319	2.465	2.536	2.363
Means 1860 to 1862	2.569	2.478	2.409	2.433	2.293	2.165	2.099	2.128	2.260	2.349	2.384	2.500	2.345
Means 1854 to 1859	2.536	2.536	2.405	2.388	2.331	2.137	2.107	2.148	2.202	2.406	2.419	2.551	2.356

TABLE XXIX.

*Monthly means of the Pressure of Dry Air at each of the six observation hours, for the period 1860 to 1862.*Pressure of Dry Air at $32^{\circ} = 27$ inches + the numbers in the table.

Toronto Astronomical Time.	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
2 ^h	2.538	2.455	2.378	2.407	2.202	2.129	2.050	2.072	2.220	2.307	2.359	2.541	2.311
4	2.548	2.454	2.379	2.403	2.258	2.121	2.065	2.071	2.222	2.316	2.371	2.553	2.313
10	2.579	2.480	2.421	2.430	2.300	2.170	2.124	2.150	2.291	2.350	2.395	2.567	2.355
12	2.580	2.481	2.420	2.433	2.309	2.187	2.129	2.158	2.302	2.356	2.392	2.567	2.350
18	2.575	2.496	2.425	2.461	2.314	2.200	2.119	2.173	2.298	2.379	2.385	2.557	2.305
20	2.591	2.505	2.433	2.466	2.312	2.179	2.097	2.142	2.277	2.382	2.399	2.579	2.364

TABLE XXX.

Monthly mean abnormal variations of the Pressure of Dry Air without regard to sign, for each month of the years 1860 to 1862.

	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
1860	0.238	0.258	0.224	0.229	0.178	0.207	0.182	0.183	0.201	0.194	0.235	0.351	0.223
1861	.226	.283	.257	.227	.251	.180	.164	.174	.210	.235	.200	.259	.222
1862	.256	.211	.205	.231	.129	.192	.155	.215	.193	.216	.218	.237	.205
Means 1860 to 1862	0.240	0.251	0.229	0.229	0.186	0.193	0.167	0.191	0.201	0.215	0.218	0.282	0.217

TORONTO METEOROLOGICAL RESULTS.

TABLE XXXI.

Monthly Mean Abnormal Variations of the Pressure of Dry Air, without regard to sign, at each of the six observation hours, for the period 1860 to 1862.

Toronto Astronomical Time.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Year.
28	0.246	0.210	0.236	0.228	0.194	0.203	0.164	0.194	0.205	0.222	0.212	0.291	0.220
4	.242	.240	.232	.241	.193	.196	.161	.192	.200	.214	.207	.289	.217
10	.232	.215	.220	.232	.186	.173	.155	.194	.191	.214	.214	.294	.213
12	.229	.250	.221	.230	.187	.177	.160	.193	.190	.218	.217	.287	.213
18	.244	.267	.230	.221	.182	.204	.180	.186	.210	.212	.227	.264	.219
20	.247	.263	.232	.221	.175	.203	.179	.183	.212	.212	.229	.269	.219

TABLE XXXII.

Mean Abnormal Variations of the Pressure of Dry Air, with their proper signs, arranged according to the direction of the wind, for each year and for the winter and summer half years, from 1860 to 1862, the summer being considered to begin April 1st, and end September 30th.

	1860.			1861.			1862.			Mean 1860-1862.			
	Winter.	Summer.	Year.	Winter.	Summer.	Year.	Winter.	Summer.	Year.	Winter.	Summer.	Year.	
	N.	+0.157	+0.128	+0.083	+0.184	+0.064	+0.073	+0.161	+0.118	+0.141	+0.140	+0.068	+0.101
N.E.	+.009	-.068	-.035	-.029	-.020	-.089	+.025	+.089	+.128	+.108	+.025	+.008	+.016
E.	-.018	-.004	-.011	-.028	-.012	-.008	+.009	+.009	+.014	+.029	-.012	+.021	+.005
S.E.	-.017	+.086	+.036	+.017	+.029	+.024	-.103	+.089	+.017	-.028	+.006	+.026	
S.	-.080	+.021	+.004	-.010	+.015	+.002	-.047	+.001	-.014	-.051	+.015	-.003	
S.W.	-.182	-.180	-.183	-.094	-.131	-.107	-.181	-.143	-.164	-.149	-.152	-.150	
W.	-.053	-.107	-.072	-.025	-.165	-.071	-.049	-.068	-.045	-.013	-.100	-.063	
N.W.	+.025	+.055	+.040	+.059	-.006	+.025	+.018	+.036	+.042	+.042	+.032	+.036	
Calms.	+.014	+.030	+.039	-.026	+.045	+.026	+.047	+.058	+.052	+.019	+.016	+.035	

TABLE XXXIII.

Mean Abnormal Variations of the Pressure of Dry Air, with their proper signs, arranged according to the direction of the wind in each month and in the year, for the period 1860 to 1862.

	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Year.
N.	+0.157	+0.124	+0.042	+0.061	+0.070	+0.085	+0.043	+0.072	+0.075	+0.129	+0.037	+0.247	+0.101
N.E.	+.018	-.145	+.016	+.036	-.048	+.040	-.031	-.029	+.009	+.068	+.045	+.101	+.016
E.	+.012	-.045	-.009	+.045	-.037	+.009	-.025	-.003	+.007	-.078	+.060	+.032	+.005
S.E.	-.176	-.102	+.027	+.190	+.053	+.163	-.039	-.084	+.088	-.073	+.004	+.106	+.026
S.	-.164	-.221	+.152	+.042	+.058	+.089	+.006	-.005	-.037	+.010	-.158	+.111	-.003
S.W.	-.069	-.145	-.206	-.110	-.052	-.165	-.157	-.144	-.226	-.265	-.181	-.083	-.150
W.	+.087	+.004	-.121	-.043	-.178	-.068	-.107	-.422	-.054	-.160	-.070	-.034	-.063
N.W.	+.115	+.056	+.021	+.071	-.017	-.021	+.030	+.041	+.070	+.023	-.041	+.082	+.036
Calms.	+.087	+.117	+.019	+.048	+.067	+.065	-.017	+.050	+.068	+.067	-.107	+.135	+.035

TORONTO METEOROLOGICAL RESULTS.

TABLE XXXIV.
Mean Differences, without regard to sign, between the Pressures of Dry Air at 6 a.m. on consecutive days, for each month in the years 1860 to 1862.

	Jan.	Feb.	March.	April.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
1860	0.283	0.283	0.192	0.261	0.161	0.211	0.285	0.186	0.227	0.178	0.266	0.263	0.233
1861	.275	.354	.366	.183	.251	.218	.149	.135	.236	.257	.240	.302	.247
1862	.371	.326	.216	.227	.190	.227	.192	.262	.240	.276	.215	.261	.250
Means 1860 to 1862	0.310	0.321	0.258	0.221	0.201	0.219	0.200	0.194	0.234	0.237	0.240	0.275	0.243

TABLE XXXV.
Mean Changes in the Pressure of Dry Air, with their proper signs, from 6 a.m. to 6 a.m., for the period 1860 to 1862, arranged according to the daily resultant direction of the wind.

	Jan.	Feb.	March.	April.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
N.	+0.119	+0.100	+0.215	+0.132	-0.112	+0.111	+0.122	-0.033	+0.161	+0.023	+0.166	+0.072	+0.096
N.E.	-.263	-.368	-.190	-.115	-.093	+.070	-.089	+.045	-.058	-.111	-.194	-.258	-.152
E.	-.598	-.413	-.301	-.178	-.142	-.155	-.149	-.115	-.345	-.253	-.276	-.305	-.240
S.E.	*	-.228	-.284	-.154	-.122	-.410	-.171	-.330	-.288	+.008	-.495	-.261	-.264
S.	+.032	-.404	-.478	-.171	+.008	-.279	-.160	-.110	-.178	-.450	-.364	-.170	-.189
S.W.	-.122	-.110	-.151	-.018	-.116	-.222	-.145	-.112	-.021	-.019	-.065	-.228	-.109
W.	+.214	+.189	+.133	+.245	+.201	+.098	+.224	+.087	+.219	+.169	+.251	+.166	+.180
N.W.	+.282	+.354	+.195	+.175	+.181	+.202	+.195	+.284	+.267	+.293	+.135	+.252	+.227

* No case of a S.E. resultant occurred in January.

TABLE XXXVI.
Highest and Lowest Pressures of Dry Air in each month, for 1860, 1861, and 1862.
Pressure of Dry Air at 32° = 27 inches + the numbers in the table.

HIGHEST.

Year.	Jan.	Feb.	March.	April.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
1860	3.110	3.070	2.787	3.160	2.685	2.550	2.551	2.561	3.025	2.788	2.886	3.222	3.222 Dec. 14.
1861	3.265	3.103	3.160	2.963	2.772	2.500	2.472	2.600	2.819	2.886	2.861	3.127	3.265 Jan. 23.
1862	3.217	3.090	2.755	3.014	2.735	2.878	2.483	2.700	2.786	2.873	3.356	3.371	3.371 Dec. 20.
Means...	3.197	3.088	2.904	3.046	2.731	2.673	2.502	2.622	2.877	2.819	3.031	3.240	3.286

LOWEST.

1860	1.985	1.783	1.617	1.689	1.665	1.469	1.490	1.600	1.865	1.684	1.621	1.643	1.469 June 4.
1861	1.832	1.826	1.833	1.710	1.333	1.500	1.611	1.611	1.740	1.734	1.773	1.825	1.333 May 6.
1862	1.721	1.841	1.616	1.835	1.805	1.637	1.550	1.601	1.441	1.719	1.982	1.858	1.441 Sept. 1.
Means...	1.816	1.817	1.703	1.715	1.601	1.535	1.550	1.605	1.682	1.712	1.792	1.709	1.414

TORONTO METEOROLOGICAL RESULTS.

TABLE XXXVII.
Monthly Means of the Pressure of Vapour, at each of the six observation hours, from 1860 to 1862 inclusive.

Toronto Astronomical Time	2 ^h		4 ^h		10 ^h		12 ^h		18 ^h		20 ^h		Mean.
	January	February	March	April	May	June	July	August	September	October	November	December	
January . . . 1860	0.115	0.114	0.114	0.109	0.109	0.106	0.104	0.110	0.136	0.126	0.116	0.114	0.115
" " 1861	.109	.109	.102	.109	.109	.102	.102	.102	.125	.122	.112	.110	.110
" " 1862	.110	.109	.101	.109	.109	.100	.100	.100	.128	.125	.112	.110	.110
Means	0.111	0.111	0.106	0.102	0.101	0.100	0.105	0.112	0.136	0.125	0.112	0.110	0.110
February . . . 1860	0.116	0.119	0.110	0.109	0.110	0.110	0.110	0.112	0.136	0.125	0.116	0.114	0.115
" " 1861	.136	.135	.133	.132	.132	.122	.122	.125	.150	.146	.136	.134	.134
" " 1862	.109	.109	.107	.107	.108	.105	.104	.107	.128	.125	.115	.113	.113
Means	0.120	0.121	0.117	0.116	0.117	0.112	0.112	0.116	0.136	0.125	0.116	0.114	0.115
March 1860	0.168	0.162	0.144	0.144	0.138	0.136	0.143	0.148	0.176	0.166	0.156	0.154	0.155
" " 1861	.131	.131	.130	.130	.126	.123	.123	.125	.157	.146	.136	.134	.134
" " 1862	.142	.139	.134	.134	.128	.120	.120	.127	.152	.142	.132	.130	.130
Means	0.147	0.144	0.136	0.131	0.131	0.126	0.126	0.131	0.166	0.156	0.146	0.144	0.145
April 1860	0.198	0.193	0.191	0.191	0.182	0.170	0.175	0.185	0.224	0.216	0.206	0.205	0.205
" " 1861	.216	.205	.204	.204	.197	.179	.179	.190	.240	.232	.216	.214	.214
" " 1862	.199	.190	.177	.177	.177	.163	.163	.184	.239	.232	.213	.212	.212
Means	0.204	0.196	0.191	0.187	0.175	0.159	0.159	0.189	0.259	0.255	0.234	0.234	0.234
May 1860	0.371	0.365	0.331	0.314	0.315	0.321	0.338	0.348	0.414	0.428	0.426	0.426	0.426
" " 1861	.243	.241	.225	.225	.221	.222	.222	.222	.286	.284	.282	.282	.282
" " 1862	.279	.261	.246	.238	.239	.230	.230	.233	.293	.282	.275	.275	.275
Means	0.298	0.280	0.267	0.258	0.258	0.259	0.259	0.274	0.358	0.354	0.349	0.349	0.349
June 1860	0.433	0.441	0.397	0.384	0.390	0.428	0.414	0.414	0.518	0.516	0.493	0.493	0.493
" " 1861	.409	.388	.370	.356	.354	.388	.377	.377	.537	.536	.498	.498	.498
" " 1862	.378	.377	.328	.321	.320	.319	.319	.316	.501	.492	.467	.467	.467
Means	0.407	0.402	0.365	0.354	0.358	0.388	0.379	0.379	0.535	0.525	0.479	0.479	0.479
July 1860	0.440	0.441	0.409	0.394	0.401	0.442	0.429	0.419	0.518	0.516	0.488	0.488	0.488
" " 1861	.461	.461	.411	.411	.411	.451	.439	.439	.538	.538	.493	.493	.493
" " 1862	.462	.462	.401	.401	.401	.452	.452	.452	.538	.538	.498	.498	.498
Means	0.455	0.471	0.440	0.433	0.435	0.462	0.453	0.453	0.534	0.534	0.496	0.496	0.496
August 1860	0.502	0.485	0.456	0.456	0.441	0.450	0.450	0.450	0.560	0.560	0.470	0.470	0.470
" " 1861	.552	.538	.472	.472	.461	.450	.450	.450	.590	.590	.495	.495	.495
" " 1862	.552	.538	.482	.482	.483	.479	.479	.479	.515	.515	.510	.510	.510
Means	0.535	0.524	0.468	0.468	0.453	0.453	0.453	0.453	0.560	0.560	0.489	0.489	0.489
September 1860	0.369	0.335	0.336	0.336	0.327	0.327	0.327	0.327	0.350	0.350	0.342	0.342	0.342
" " 1861	.436	.436	.379	.379	.376	.376	.376	.376	.401	.401	.390	.390	.390
" " 1862	.436	.436	.404	.404	.389	.389	.389	.389	.423	.423	.418	.418	.418
Means	0.420	0.411	0.373	0.364	0.360	0.360	0.360	0.360	0.391	0.391	0.357	0.357	0.357
October 1860	0.296	0.285	0.269	0.269	0.265	0.265	0.265	0.265	0.298	0.298	0.265	0.265	0.265
" " 1861	.312	.302	.298	.298	.298	.298	.298	.298	.329	.329	.288	.288	.288
" " 1862	.318	.318	.302	.302	.302	.302	.302	.302	.330	.330	.292	.292	.292
Means	0.309	0.302	0.290	0.283	0.283	0.283	0.283	0.283	0.307	0.307	0.278	0.278	0.278
November 1860	0.193	0.187	0.175	0.175	0.175	0.175	0.175	0.175	0.192	0.192	0.193	0.193	0.193
" " 1861	.180	.179	.175	.175	.175	.175	.175	.175	.192	.192	.181	.181	.181
" " 1862	.182	.175	.169	.169	.167	.167	.167	.167	.187	.187	.166	.166	.166
Means	0.189	0.185	0.179	0.175	0.175	0.175	0.175	0.175	0.190	0.190	0.180	0.180	0.180
December 1860	0.206	0.206	0.192	0.192	0.192	0.192	0.192	0.192	0.213	0.213	0.195	0.195	0.195
" " 1861	.157	.157	.149	.149	.149	.149	.149	.149	.172	.172	.151	.151	.151
" " 1862	.152	.152	.144	.144	.144	.144	.144	.144	.167	.167	.142	.142	.142
Means	0.141	0.140	0.137	0.133	0.133	0.133	0.133	0.133	0.153	0.153	0.136	0.136	0.136

TABLE XXXVIII.

Monthly and annual means of the Pressure of Vapour furnished by six daily observations, from 1860 to 1862.

	Jan.	Feb.	March.	April.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
1860	0.110	0.112	0.148	0.185	0.338	0.414	0.427	0.463	0.342	0.272	0.195	0.115	0.260
1861	.102	.130	.127	.199	.232	.377	.407	.495	.400	.292	.178	.151	.262
1862	.103	.107	.132	.184	.253	.346	.473	.510	.418	.300	.171	.142	.262
Means 1860 to 1862	0.105	0.116	0.136	0.189	0.274	0.379	0.456	0.489	0.387	0.288	0.181	0.136	0.261
Means 1854 to 1859	0.112	0.104	0.133	0.184	0.266	0.407	0.507	0.458	0.384	0.247	0.176	0.120	0.258

TABLE XXXIX.

Monthly means of the Pressure of Vapour at each of the six observation hours, for the period 1860 to 1862.

Toronto Astronomical Time.	Jan.	Feb.	March.	April.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
2 ^h	0.111	0.120	0.147	0.204	0.298	0.407	0.485	0.535	0.420	0.309	0.180	0.141	0.280
4	.111	.121	.144	.196	.289	.402	.471	.524	.411	.302	.185	.140	.275
10	.106	.117	.136	.191	.267	.365	.440	.468	.373	.290	.179	.137	.256
12	.102	.116	.131	.187	.258	.354	.433	.462	.364	.283	.175	.133	.250
18	.101	.112	.126	.175	.259	.358	.435	.453	.360	.267	.180	.135	.247
20	.100	.112	.131	.182	.275	.388	.469	.495	.391	.278	.180	.129	.261

TABLE XL.

Monthly mean abnormal variations of the Pressure of Vapour, without regard to sign, for each month of the years 1860 to 1862.

	Jan.	Feb.	March.	April.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
1860	0.043	0.051	0.046	0.048	0.081	0.078	0.091	0.094	0.111	0.067	0.050	0.035	0.067
1861	.037	.041	.047	.054	.072	.087	.091	.090	.078	.076	.037	.000	.064
1862	.039	.033	.028	.050	.062	.033	.079	.113	.090	.092	.041	.046	.084
Means 1860 to 1862	0.019	0.042	0.040	0.053	0.072	0.083	0.087	0.099	0.095	0.073	0.046	0.047	0.065

TABLE XLII.

Monthly mean abnormal variations of the Pressure of Vapour, without regard to sign, at each of the six observation hours, for the period 1860 to 1862.

Toronto Astronomical Time.	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
2h	.010	.040	.043	.062	.085	.093	.092	.106	.110	.082	.017	.049	.071
4	.038	.011	.043	.061	.087	.092	.097	.108	.106	.078	.045	.048	.070
10	.040	.014	.036	.056	.065	.076	.081	.096	.094	.081	.016	.049	.064
12	.041	.045	.055	.049	.066	.073	.083	.098	.089	.081	.045	.047	.063
18	.041	.043	.042	.018	.063	.078	.084	.091	.079	.074	.045	.047	.061
20	.039	.041	.042	.049	.065	.083	.083	.091	.093	.073	.044	.043	.062

TABLE XLIII.

Mean abnormal variations of the Pressure of Vapour, with their proper signs, arranged according to the direction of the wind, for each year and for the winter and summer half years, from 1860 to 1862, the summer being considered to begin April 1st, and end September 30th.

	1860.			1861.			1862.			Mean 1860 to 1862.		
	Winter.	Summer.	Year.	Winter.	Summer.	Year.	Winter.	Summer.	Year.	Winter.	Summer.	Year.
N.	-0.020	-0.011	-0.015	-0.019	-0.042	-0.031	-0.025	-0.052	-0.038	-0.022	-0.033	-0.028
N.E.	+.031	+.031	+.031	+.028	-.010	+.011	-.013	-.030	-.021	+.013	-.000	+.006
E.	+.056	+.001	+.027	+.032	-.008	+.012	+.025	-.004	+.008	+.036	-.004	+.015
S.E.	+.059	-.022	+.017	+.031	-.001	+.012	+.047	+.001	+.018	+.045	-.006	+.015
S.	+.032	+.007	+.012	+.018	+.008	+.011	+.014	+.017	+.016	+.020	+.011	+.013
S.W.	+.016	+.038	+.023	+.031	+.040	+.034	+.052	+.068	+.059	+.032	+.050	+.039
W.	-.013	-.022	-.016	-.009	-.009	-.009	-.006	-.004	-.006	-.010	-.012	-.010
N.W.	-.021	-.065	-.045	-.024	-.041	-.033	-.020	-.045	-.033	-.023	-.052	-.037
Calm.	-.001	-.018	-.008	+.013	-.015	-.008	+.016	-.009	+.004	+.009	-.014	-.005

TABLE XLIII.

Mean abnormal variations of the Pressure of Vapour, with their proper signs, arranged according to the direction of the wind, in each month and in the year, for the period 1860 to 1862.

	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
N.	-0.051	-0.024	-0.009	-0.005	-0.022	-0.058	-0.067	-0.019	-0.034	+0.008	-0.014	-0.036	-0.028
N.E.	-.012	+.036	-.003	-.003	+.031	-.035	-.030	+.021	-.013	+.035	+.018	+.005	+.006
E.	+.015	+.023	+.011	+.006	-.000	-.062	-.018	+.023	+.033	+.086	+.046	+.035	+.015
S.E.	+.023	+.017	+.013	-.007	-.008	-.077	-.009	+.053	-.008	+.086	+.053	+.033	+.015
S.	+.031	+.026	+.001	-.019	+.012	-.031	-.009	+.026	+.058	+.027	+.009	+.015	+.013
S.W.	+.013	+.029	+.021	+.021	+.007	+.046	+.045	+.065	+.092	+.101	-.013	+.054	+.039
W.	-.017	+.002	-.001	-.003	-.008	-.029	-.012	+.017	-.012	-.001	-.026	-.014	-.010
N.W.	-.033	-.032	-.025	-.016	-.038	-.040	-.074	-.061	-.076	-.011	-.012	-.021	-.037
Calm.	-.021	-.005	+.016	-.002	-.027	-.044	-.000	+.002	-.014	+.019	-.002	+.002	-.005

TABLE XLIV.

Mean Differences, without regard to sign, between the Pressures of Vapour at 6 a.m. on consecutive days, for each month in the years 1860 to 1862.

	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
1860	.010	.034	.041	.045	.046	.101	.107	.082	.089	.053	.047	.034	.061
1861	.034	.041	.047	.041	.044	.083	.070	.073	.089	.059	.043	.050	.050
1862	.041	.043	.029	.057	.064	.088	.070	.110	.102	.082	.013	.049	.063
Means 1860 to 1862038	.043	.039	.041	.051	.091	.144	.088	.093	.065	.044	.044	.060

TABLE XLV.

Mean changes in the Pressures of Vapour with their proper signs, from 6 a.m. to 6 a.m., for the period 1860 to 1862, arranged according to the daily resultant direction of the wind.

	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
N.	-0.011	-0.007	-0.021	-0.027	-0.004	-0.014	-0.010	+0.001	-0.008	-0.020	+0.020	-0.005	-0.020
N.E.	+ .020	+ .022	+ .011	+ .019	+ .018	- .007	+ .010	+ .020	+ .014	+ .021	+ .022	+ .021	+ .013
E.	+ .019	+ .001	+ .034	+ .027	+ .019	+ .020	+ .032	+ .013	+ .198	+ .058	+ .022	+ .050	+ .043
S.E.	+ .023	+ .028	+ .025	+ .005	+ .031	+ .128	+ .090	+ .131	+ .109	+ .034	+ .091	+ .066	+ .085
S.	+ .016	+ .042	+ .035	+ .059	+ .009	+ .119	+ .074	+ .012	+ .082	+ .136	+ .017	+ .001	+ .057
S.W.	+ .028	+ .041	+ .031	+ .019	+ .010	+ .035	+ .015	+ .028	+ .001	+ .015	+ .011	+ .027	+ .021
W.	+ .022	+ .019	+ .022	+ .032	+ .048	+ .039	+ .004	+ .010	+ .001	+ .014	+ .023	+ .032	+ .030
N.W.	+ .018	+ .024	+ .012	+ .027	+ .027	+ .063	+ .062	+ .032	+ .111	+ .045	+ .029	+ .010	+ .014

* No case of a S.E. resultant occurred in January.

TABLE XLVI.

Highest and lowest Pressures of Vapour in each month, for the years 1860, 1861, and 1862.

HIGHEST.

Year.	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.	
1860	.293	.292	.431	.388	.076	.714	.710	.822	.071	.450	.481	.322	.922	August 7th.
1861	.160	.274	.278	.410	.615	.676	.704	.828	.072	.494	.235	.390	.823	August 10th.
1862	.244	.199	.220	.456	.484	.643	.734	.928	.718	.671	.265	.343	.828	August 8th.
Means.....	.234	.235	.313	.419	.058	.678	.710	.824	.080	.540	.301	.322	.824	

LOWEST.

1860	0.023	0.022	0.015	0.050	0.163	0.142	0.195	0.269	0.104	0.131	0.013	0.029	0.022	February 1.
1861	0.20	0.12	0.28	0.09	0.102	0.150	0.250	0.275	0.182	0.118	0.09	0.03	0.012	February 7.
1862	0.25	0.11	0.02	0.070	0.110	0.090	0.221	0.27	0.172	0.130	0.083	0.03	0.025	January 4.
Means.....	0.023	0.023	0.015	0.070	0.145	0.132	0.211	0.250	0.133	0.162	0.074	0.020	0.020	

TORONTO METEOROLOGICAL RESULTS.

TABLE XLVII.
Monthly means of the Relative Humidity, at each of the six observation hours, from 1860 to 1862 inclusive.

Toronto Astronomical Time.	2 ^h			4 ^h			6 ^h			8 ^h			10 ^h			12 ^h			14 ^h			16 ^h			18 ^h			20 ^h			Means.		
	2 ^h	4 ^h	6 ^h	8 ^h	10 ^h	12 ^h	14 ^h	16 ^h	18 ^h	20 ^h	Astronomical Time.	2 ^h	4 ^h	6 ^h	8 ^h	10 ^h	12 ^h	14 ^h	16 ^h	18 ^h	20 ^h	Means.											
January ... 1860	73	76	82	84	85	82	81	81	81	81	July ... 1860	59	56	78	80	81	74	72	72	72	72	72											
" 1861	84	84	89	90	90	88	88	88	88	88	" 1861	65	64	78	80	80	71	73	73	73	73	73											
" 1862	75	77	80	82	86	84	84	84	84	84	" 1862	62	60	76	79	82	74	72	72	72	72	72											
Means	78	79	84	85	87	85	83	83	83	83	Means	62	60	77	80	81	74	72	72	72	72	72											
February	72	74	82	85	87	84	81	81	81	81	August ... 1860	65	64	80	82	85	77	75	75	75	75	75											
" 1861	80	79	88	87	86	86	84	84	84	84	" 1861	70	69	81	83	84	78	78	78	78	78	78											
" 1862	77	78	86	87	89	87	84	84	84	84	" 1862	65	64	77	80	82	75	74	74	74	74	74											
Means	76	77	85	86	88	86	83	83	83	83	Means	67	66	79	82	84	77	76	76	76	76	76											
March ... 1860	65	61	73	73	74	74	71	71	71	71	September ... 1860	64	64	79	81	83	75	75	75	75	75	75											
" 1861	70	70	84	83	87	83	80	80	80	80	" 1861	70	69	79	81	83	81	81	81	81	81	81											
" 1862	75	74	84	85	86	87	84	82	82	82	" 1862	69	69	84	86	88	82	82	80	80	80	80											
Means	70	68	80	81	84	80	78	78	78	78	Means	68	68	81	83	86	79	78	78	78	78	78											
April ... 1860	64	64	80	80	81	73	74	74	74	74	October ... 1860	71	74	85	84	88	83	81	81	81	81	81											
" 1861	65	62	75	75	81	74	73	73	73	73	" 1861	73	73	87	88	89	84	84	84	84	84	84											
" 1862	66	64	75	78	80	75	73	73	73	73	" 1862	75	77	85	86	86	83	84	82	82	82	82											
Means	65	63	77	79	81	74	73	73	73	73	Means	73	75	86	86	88	84	82	82	82	82	82											
May ... 1860	67	67	79	81	83	76	76	76	76	76	November ... 1860	74	75	81	82	86	84	84	84	84	84	84											
" 1861	58	58	73	77	78	71	72	66	69	69	" 1861	69	72	81	81	81	84	84	84	84	84	84											
" 1862	56	53	70	71	72	66	65	66	66	66	" 1862	73	74	83	84	85	82	82	80	80	80	80											
Means	60	59	74	76	78	71	70	70	70	70	Means	72	74	82	82	86	83	83	83	83	83	83											
June ... 1860	62	62	62	75	76	78	74	71	71	71	December ... 1860	78	81	87	86	88	87	87	87	87	87	87											
" 1861	59	59	69	74	76	75	70	69	69	69	" 1861	72	73	82	82	85	81	81	81	81	81	81											
" 1862	59	60	69	70	70	72	64	66	66	66	" 1862	78	79	84	85	87	83	83	82	82	82	82											
Means	60	60	73	74	75	73	69	69	69	69	Means	76	78	84	84	87	84	84	84	84	84	84											

TABLE XLVIII.

Monthly and annual means of the Relative Humidity furnished by six daily observations, from 1860 to 1862.

	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
1860	81	81	71	74	76	71	72	76	74	81	80	84	76
1861	88	84	80	73	69	69	73	78	79	82	79	79	78
1862	81	84	82	73	65	66	72	74	80	82	80	83	77
Means 1860 to 1862	83	83	78	73	70	69	72	76	78	82	80	82	77
Means 1854 to 1859	82	80	77	72	70	74	73	73	77	76	78	81	76

TABLE XLIX.

Monthly means of the Relative Humidity at each of the six observation hours, for the period 1860 to 1862.

Toronto Astronomical Time.	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
2h	78	76	70	65	60	60	62	67	68	73	72	76	69
4	79	77	68	63	59	60	60	66	68	75	74	78	69
10	84	85	80	77	74	73	77	79	81	86	82	84	80
12	85	86	81	79	76	74	80	82	83	86	82	84	81
18	87	88	84	81	78	75	81	84	86	88	86	87	84
20	85	86	80	74	71	69	74	77	79	84	83	84	79

TABLE L.

Monthly mean abnormal variations of the Relative Humidity without regard to sign, for each month of the years 1860 to 1862.

	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
1860	8.2	8.1	13.3	10.5	11.2	11.1	9.9	7.7	10.3	8.4	8.6	7.1	9.5
1861	7.5	9.5	12.0	10.3	10.9	7.6	8.1	7.3	8.1	8.3	9.8	9.0	
1862	8.4	8.4	8.5	10.9	13.8	15.0	11.0	8.8	7.4	8.8	7.5	7.2	9.6
Means 1860 to 1862	8.0	8.7	10.0	11.1	11.8	12.3	9.5	8.2	8.3	8.4	8.1	8.0	9.4

TABLE LI.

Monthly mean abnormal variations of the Relative Humidity without regard to sign, at each of the six observation hours, for the period 1860 to 1862.

Toronto Astronomical Time.	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
2h	9.9	11.2	12.2	12.9	12.4	14.0	11.2	10.3	10.9	10.0	10.6	10.5	11.3
4	10.2	10.9	13.5	13.1	13.7	13.8	11.9	11.7	10.8	10.9	9.8	10.0	11.9
10	7.7	8.4	9.1	9.9	11.8	11.2	9.2	7.4	7.3	7.2	7.3	7.0	8.6
12	6.6	8.3	8.6	8.7	11.0	11.2	8.2	6.5	6.6	7.4	7.6	7.3	8.2
18	6.2	6.6	8.1	8.2	10.1	11.1	7.3	6.2	6.0	6.5	6.4	6.2	7.4
20	7.6	6.6	8.3	12.1	11.6	12.7	9.1	7.2	8.4	8.3	7.1	6.9	8.8

TORONTO METEOROLOGICAL RESULTS.

TABLE LII.

Mean abnormal variations of the Relative Humidity, with their proper signs, arranged according to the direction of the wind, for each year and for the winter and summer half years, from 1860 to 1862, the summer being considered to begin April 1st, and end September 30th.

	1860.			1861.			1862.			Mean 1860 to 1862.		
	Winter.	Summer.	Year.	Winter.	Summer.	Year.	Winter.	Summer.	Year.	Winter.	Summer.	Year.
N.	0.0	-3.6	-2.1	+1.9	-4.7	-1.6	-0.2	-9.2	-4.5	+0.5	-5.8	-2.8
N.E.	+1.0	+1.4	+4.6	+5.6	+0.6	+3.3	+2.4	-2.5	0.0	+3.2	+1.1	+2.6
E.	+6.0	+0.9	+3.9	+4.8	+1.6	+3.2	+7.0	+2.5	+4.4	+6.2	+1.8	+3.9
S.E.	+3.1	-0.7	+1.1	+4.4	-0.9	+1.2	+5.0	-2.6	+0.3	+3.2	-1.4	+0.9
S.	+2.1	-0.7	-0.2	+3.2	-1.6	0.0	-0.1	-2.3	-1.6	+1.6	-1.5	-0.7
S.W.	-2.4	+1.7	-1.1	+0.8	-0.3	+0.4	+2.1	+0.9	+1.6	0.0	+0.8	+0.3
W.	-5.2	-3.4	-1.5	-1.3	-2.5	-1.7	-1.5	-6.0	-8.2	-2.8	-4.0	-3.2
N.W.	-4.9	-5.6	-5.3	-2.1	-5.1	-3.9	-2.2	-9.7	-6.0	-3.3	-6.8	-5.1
Calms.	+1.2	+1.1	+1.2	+2.5	-1.7	-0.6	+4.8	-1.5	+1.8	+2.8	-1.3	+0.4

TABLE LIII.

Mean abnormal variations of the Relative Humidity, with their proper signs, arranged according to the direction of the wind, in each month and in the year, for the period 1860 to 1862.

	1860 to 1862.												
	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Year.
N.	-1.6	+3.3	+3.8	-0.5	-8.9	-12.6	-4.4	-5.6	-2.3	+0.1	-1.7	+1.2	-2.8
N.E.	+5.1	+9.4	+5.6	+1.8	+3.0	-5.7	+2.5	+2.6	-0.3	+1.9	+1.2	+2.1	+2.6
E.	+5.6	+9.6	+4.7	+2.0	+2.0	-2.9	+3.6	+3.5	+1.1	+0.1	+0.6	+5.3	+3.9
S.E.	+5.5	+8.5	+2.2	-5.6	-5.5	-3.5	+2.5	+3.7	+0.4	+6.3	+4.4	-2.4	+0.9
S.	+6.5	-0.3	-1.2	+2.1	-1.1	-9.6	-0.2	+0.7	-0.4	+0.2	+0.8	+5.3	-0.7
S.W.	+9.1	+1.1	-1.3	-2.1	-1.1	+1.2	+1.4	+1.1	+2.7	+2.1	-1.6	+0.6	+0.3
W.	-3.2	+0.1	-1.3	+0.9	-2.0	-9.0	-3.1	-5.1	-7.5	-1.2	-6.1	-3.0	-3.2
N.W.	-2.5	-2.1	-3.8	-1.1	-6.3	-12.5	-0.1	-5.4	-8.7	-3.4	-4.5	-2.0	-5.1
Calms.	+0.1	+1.2	+1.6	+6.6	-1.7	-6.8	-1.8	-0.8	+0.8	+5.9	+2.5	-0.1	+0.4

TABLE LIV.

Mean difference, without regard to sign, between the Relative Humidity at 6 a.m. on consecutive days, for each month in the years 1860 to 1862.

	1860 to 1862.												
	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Year.
1860	9.5	6.7	12.9	10.0	10.1	12.5	10.2	7.7	11.4	7.5	10.8	6.6	9.7
1861	4.8	9.0	7.8	14.2	11.0	11.4	7.5	6.3	5.7	8.9	7.3	10.0	8.7
1863	9.4	6.1	6.1	8.2	11.0	13.4	8.6	9.1	6.0	7.1	6.7	7.5	8.3
Means 1860 to 1862	7.9	7.3	8.9	10.8	11.0	12.4	8.8	7.7	7.7	7.8	7.9	8.0	8.9

TABLE LV.

Mean changes in the Relative Humidity, with the proper signs, from 6 a.m. to 6 p.m., for the period 1860 to 1862, arranged according to the daily resultant direction of the wind.

	Jan.	Feb.	March.	April.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
Year.													
N.	+0.4	+0.8	-7.5	-2.8	-6.3	+2.6	-3.9	-3.2	-5.1	-3.7	-1.4		
N.E.	+1.4	+1.1	-0.7	+2.9	+3.3	-0.2	-3.0	+1.2	+0.8	+4.2	+5.5	+0.8	+10.5
E.	+3.5	+2.3	-3.4	+3.5	+2.6	+4.9	+4.8	-1.1	+5.3	+6.2	+1.1	-0.8	+3.9
S.E.	*	+0.5	-0.0	+2.0	-1.0	+5.3	-8.7	+0.0	+7.0	-2.6	+5.8	+1.3	+3.3
S.	+1.5	-1.0	-0.0	+21.5	-2.8	+9.5	-5.1	+6.6	+0.7	-1.0	-2.0	-6.0	+2.8
S.W.	+0.5	+1.1	-0.0	-7.5	+4.2	+5.4	+0.9	-0.2	+2.6	-6.2	-6.3	-0.6	-6.4
W.	-1.8	-0.3	-2.8	-0.0	-1.0	-5.3	-0.2	+0.1	-0.7	-2.9	-0.5	-1.3	-1.9
N.W.	+0.4	+2.4	+0.8	-1.0	-2.8	-6.3	-6.6	-0.0	-4.8	+1.3	-2.2	+0.7	-2.2

* No case of a S.E. resultant occurred in January.

TABLE LVI

Comparative view of the Annual variations of certain Meteorological elements derived from the series 1842-48, 1854-59, and 1860-1862.

Year.	Months.	Temperature.				Barometer.				Pressure of Dry Air.				Pressure of Vapour.				Relative Humidity.			
		1842 1818	1854 1859	1860 1862	1848	1842 1848	1854 1859	1860 1862	1842 1848	1854 1859	1860 1862	1842 1848	1854 1859	1860 1862	1842 1848	1854 1859	1860 1862	1842 1848	1854 1859	1860 1862	
-2.8	January	-10.00	-21.44	-22.71	-0.003	+0.003	+0.006	+0.159	+0.200	+0.221	-142	-146	-166	+5	+6	+6					
-2.6	February	-21.00	-21.09	-20.80	-0.007	+0.015	+0.011	+0.149	+0.153	+0.156	-154	-154	-145	-2	+4	+6					
-3.0	March	-14.51	-14.58	-14.14	+0.001	+0.076	+0.001	+0.131	+0.049	+0.064	-129	-129	-125	-3	+1	+1					
+0.9	April	-1.04	-3.47	-3.91	+0.09	-0.042	+0.017	+0.093	+0.062	+0.088	-95	-95	-64	-1	-1	-1					
-0.7	May	+8.59	+7.62	+7.33	-0.056	-0.059	-0.059	-0.029	-0.025	-0.032	-152	-153	-153	+0.008	+0.013	+0.013	-6	-7	-7		
+0.4	June	+1.47	+1.45	+17.28	-0.054	-0.054	-0.002	-0.182	-0.219	-0.180	-153	-153	-149	+0.118	-2	-2	-8				
-0.7	July	+21.57	+21.55	+21.50	-0.052	-0.052	-0.002	-0.157	-0.209	-0.159	-209	-209	-214	+0.249	+0.95	-4	-3	-5			
+0.3	August	+21.42	+22.08	+21.69	+0.017	-0.008	+0.011	+0.213	+0.206	+0.207	-157	-157	-159	+0.098	+0.17	+0.17	+1	+1	+1		
-3.2	September	+13.27	+14.05	+13.75	+0.026	+0.011	+0.044	+0.092	+0.064	+0.076	+118	+118	+120	-2	-3	-3					
-5.1	October	-0.12	+2.42	+3.04	+0.042	+0.036	+0.050	+0.064	+0.050	+0.064	-122	-122	-111	-0.011	+0.027	+0.04	-0	-5	-5		
+0.4	November	-8.08	-7.59	-7.41	+0.035	+0.010	-0.041	+0.083	+0.061	+0.059	-17.8	-17.8	-18.2	-0.082	-0.080	+0.06	+2	+3	+3		
	December	-10.89	-19.03	-10.32	+0.022	+0.056	+0.001	+0.155	+0.195	+0.215	-125	-128	-125	+3	+5	+5					

TABLE LVII.

Comparative view of the Annual means of the diurnal variations at the six observation hours for the same three series.

Year.	Hours.	Temperature.				Barometer.				Pressure of Dry Air.				Pressure of Vapour.				Relative Humidity.			
		1842 1848	1854 1859	1860 1862	1842 1848	1854 1859	1860 1862	1842 1848	1854 1859	1860 1862	1842 1848	1854 1859	1860 1862	1842 1848	1854 1859	1860 1862	1842 1848	1854 1859	1860 1862		
0.7	2	+0.90	+0.74	+0.84	-0.013	-0.015	-0.016	-0.040	-0.032	-0.034	+0.028	+0.017	+0.010	-9	-9	-8					
8.7	4	+5.06	+4.33	+4.39	-0.08	-0.019	-0.018	-0.041	-0.011	-0.034	+0.021	+0.012	+0.014	-9	-8	-8					
8.3	10	-2.39	-1.85	-1.68	-0.011	+0.035	+0.005	+0.011	+0.009	+0.010	-0.012	-0.005	-0.005	+3	+4	+3					
8.9	12	-3.42	-3.09	-2.80	-0.005	+0.002	+0.001	+0.012	+0.012	+0.014	-0.017	-0.010	-0.011	+5	+5	+4					
8.9	14	-4.50	-3.93	-3.81	+0.010	+0.005	+0.004	+0.032	+0.025	+0.029	-0.022	-0.014	-0.011	+7	+7	+7					
8.9	18	-4.50	-3.93	-3.81	+0.010	+0.005	+0.004	+0.032	+0.025	+0.029	-0.022	-0.014	-0.011	+7	+7	+7					
8.9	20	-1.21	-1.22	-1.00	+0.025	+0.020	+0.019	+0.025	+0.020	+0.019	.000	.000	.000	+2	+2	+2					



TORONTO METEOROLOGICAL RESULTS.

TABLE LVIII.
Monthly means of the extent of Sky Clouded, at each of the six observation hours, for the years 1860 to 1862.

Astronomical Time.	Toronto			Astronomical Time.			Toronto			Astronomical Time.		
	2 ^h	4 ^h	10 ^h	2 ^h	4 ^h	10 ^h	2 ^h	4 ^h	10 ^h	12 ^h	18 ^h	20 ^h
January .. 1860	0.81	0.77	0.54	0.47	0.82	0.86	0.71					
" " 1861	.77	.76	.68	.70	.80	.85	.76					
" " 1862	.81	.82	.62	.68	.70	.73	.73					
Means	0.80	0.78	0.61	0.62	0.77	0.81	0.73					
February .. 1860	0.64	0.70	0.60	0.62	0.74	0.74	0.67					
" " 1861	.69	.76	.79	.78	.78	.80	.83					
" " 1862	.82	.72	.67	.72	.72	.74	.78					
Means	0.78	0.76	0.69	0.71	0.77	0.79	0.83					
March .. 1860	0.47	0.49	0.41	0.46	0.56	0.51	0.49					
" " 1861	.59	.61	.62	.61	.66	.63	.62					
" " 1862	.59	.71	.60	.55	.66	.69	.63					
Means	0.55	0.60	0.54	0.54	0.54	0.63	0.61	0.58				
April .. 1860	0.67	0.65	0.49	0.49	0.59	0.66	0.59					
" " 1861	.65	.63	.54	.55	.67	.64	.61					
" " 1862	.68	.65	.53	.60	.68	.74	.65					
Means	0.67	0.64	0.52	0.55	0.65	0.68	0.62					
May .. 1860	0.64	0.61	0.50	0.51	0.63	0.56	0.57					
" " 1861	.54	.50	.36	.45	.53	.56	.49					
" " 1862	.46	.44	.45	.36	.50	.47	.45					
Means	0.55	0.52	0.44	0.44	0.55	0.53	0.50					
June .. 1860	0.66	0.65	0.47	0.44	0.60	0.65	0.58					
" " 1861	.52	.53	.38	.42	.37	.46	.45					
" " 1862	.74	.70	.42	.46	.69	.68	.60					
Means	0.64	0.63	0.42	0.44	0.52	0.60	0.54					
July .. 1860	0.50	0.47	0.40	0.40	0.43	0.43	0.35					
" " 1861	.66	.67	.49	.42	.42	.42	.54					
" " 1862	.72	.59	.29	.36	.45	.45	.45					
Means	0.63	0.58	0.44	0.44	0.49	0.54	0.52					
August .. 1860	0.50	0.47	0.46	0.46	0.39	0.30	0.43					
" " 1861	.64	.64	.42	.58	.54	.54	.54					
" " 1862	.57	.52	.29	.45	.49	.49	.45					
Means	0.57	0.54	0.40	0.47	0.44	0.47	0.47					
September .. 1860	0.57	0.60	0.35	0.36	0.53	0.53	0.46					
" " 1861	.72	.62	.58	.50	.58	.58	.64					
" " 1862	.42	.38	.44	.53	.56	.56	.50					
Means	0.57	0.53	0.46	0.46	0.56	0.53	0.52					
October .. 1860	0.69	0.69	0.64	0.64	0.77	0.74	0.70					
" " 1861	.69	.65	.51	.51	.66	.66	.64					
" " 1862	.76	.77	.70	.70	.76	.76	.73					
Means	0.71	0.70	0.62	0.64	0.67	0.73	0.68					
November .. 1860	0.77	0.78	0.66	0.63	0.68	0.71	0.70					
" " 1861	.72	.74	.73	.73	.75	.75	.78					
" " 1862	.82	.79	.74	.75	.79	.79	.82					
Means	0.77	0.76	0.71	0.70	0.74	0.77	0.74					
December .. 1860	0.80	0.81	0.75	0.84	0.86	0.89	0.83					
" " 1861	.64	.64	.57	.55	.61	.64	.62					
" " 1862	.70	.73	.78	.86	.77	.77	.75					
Means	0.72	0.73	0.70	0.75	0.75	0.74	0.73					

TABLE LIX.

Monthly and Annual means of the extent of sky clouded, from six daily observations, for 1860 to 1862.

	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
1860	.71	.67	.49	.59	.57	.58	.43	.43	.48	.70	.70	.83	.60
1861	.70	.83	.62	.61	.49	.45	.56	.54	.60	.61	.74	.62	.62
1862	.73	.78	.63	.65	.45	.60	.56	.45	.47	.72	.79	.75	.63
Means 1860 to 1862	.73	.76	.58	.62	.50	.54	.52	.47	.52	.68	.74	.73	.62
Means 1854 to 1859	.71	.69	.60	.59	.52	.55	.46	.44	.48	.60	.74	.75	.59

TABLE LX.

Monthly means of the extent of sky clouded at each observation hour, for the period 1860 to 1862.

Toronto Astronomical Time.	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.	1860 to 1862	1854 to 1859.
2 ^h	0.80	.78	.55	.67	.55	.64	.63	.57	.57	.71	.77	.72	.66	.65	
4	.78	.76	.60	.64	.52	.63	.58	.54	.53	.70	.76	.73	.65	.63	
10	.61	.69	.54	.52	.44	.42	.44	.40	.46	.62	.71	.70	.55	.52	
12	.62	.71	.54	.55	.44	.44	.44	.41	.46	.64	.70	.75	.56	.52	
18	.77	.80	.63	.65	.55	.52	.49	.47	.56	.67	.74	.75	.63	.62	
20	.81	.83	.61	.68	.53	.60	.54	.44	.53	.71	.77	.74	.65	.62	

TABLE LXI.

Mean clouded sky, arranged according to the direction of the wind at the hour of observation, in each month and in the year, for the period 1860 to 1862.

	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year.	1860 to 1862	1854 to 1859.
N.	0.70	0.62	0.58	0.52	0.30	0.38	0.47	0.36	0.56	0.67	0.76	0.58	0.54	0.50	
N.E.	.87	.93	.71	.77	.65	.76	.62	.70	.71	.75	.84	.89	.76	.74	
E.	.97	.97	.77	.70	.66	.68	.73	.62	.58	.80	.79	.97	.77	.70	
S.E.	1.00	.98	.46	.46	.36	.75	.49	.55	.60	.71	.93	.89	.63	.57	
S.	0.96	.65	.42	.54	.38	.55	.55	.48	.47	.65	.84	.84	.56	.48	
S.W.	.77	.74	.59	.65	.54	.62	.66	.62	.64	.57	.78	.76	.68	.67	
W.	.59	.74	.58	.68	.72	.52	.59	.36	.41	.68	.71	.73	.63	.57	
N.W.	.57	.59	.49	.52	.42	.40	.36	.32	.42	.56	.62	.65	.49	.53	
Calms.	.72	.76	.43	.59	.34	.43	.47	.46	.48	.69	.75	.59	.55	.55	

TABLE LXII.
Number of hours that each wind blew in each month and in the year, for the periods 1853-1857, 1858-1862, and 1863-1867.

1853-1857.

	N.	N.N.E.	N.E.	E.N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.S.W.	S.W.	W.S.W.	W.	W.N.W.	N.W.	N.N.W.	Calm.	Suns.
January	290	189	211	126	143	98	52	42	51	130	296	598	407	228	250	313	294	3720
February	246	191	142	128	209	91	36	52	81	158	245	370	439	320	234	305	137	3384
March	132	77	77	157	67	99	45	127	198	361	299	428	495	472	336	136	3715	
April	320	233	197	234	331	201	108	166	150	271	179	119	122	265	314	265	3600	
May	317	116	153	296	339	263	131	120	236	328	187	136	205	202	208	174	3456	
June	180	99	172	286	356	182	123	108	251	328	264	114	122	173	193	300	300	3720
July	230	173	144	202	303	194	140	263	313	357	179	110	207	292	296	346	281	3720
August	293	209	145	149	218	114	170	157	240	297	206	141	126	196	263	314	330	3544
September	269	218	164	195	237	143	150	157	208	253	280	183	274	294	283	295	397	3800
October	210	151	209	214	265	111	56	104	219	221	174	221	174	265	365	187	207	196
November	148	101	117	276	273	97	108	104	105	220	366	686	396	248	235	302	240	3720
December	284	241	196	160	135	100	53	49	41	118	347	118	465	365	280	301	314	43549
Year.....	2899	1998	1632	2413	2937	1681	1228	1307	2022	2900	*3034	3215	3198	3052	3095	3663	2975	

1858-1862.

	N.	N.N.E.	N.E.	E.N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.S.W.	S.W.	W.S.W.	W.	W.N.W.	N.W.	N.N.W.	Calm.	Suns.
January	220	17	9	163	206	70	38	60	57	181	389	533	388	278	283	247	247	3720
February	207	72	0	186	198	55	59	54	47	197	274	372	401	314	283	247	200	3384
March	162	58	58	331	348	102	77	63	59	150	197	240	382	247	265	147	3720	
April	204	142	186	410	547	170	107	89	103	117	132	172	265	337	237	303	79	3600
May	196	160	216	450	439	149	89	120	175	245	92	99	165	257	312	319	177	3720
June	237	101	104	251	319	154	62	114	270	341	227	111	167	172	388	382	190	3600
July	204	122	85	218	250	183	142	163	322	330	189	135	145	242	342	389	259	3720
August	182	163	153	158	278	176	108	125	273	300	185	129	174	295	392	404	225	3720
September	261	137	149	194	218	100	103	113	209	388	180	145	237	231	233	261	421	3600
October	251	186	159	244	269	147	67	74	142	239	278	183	274	429	270	241	267	3720
November	226	185	190	183	245	115	50	58	65	131	238	441	336	310	279	220	3552	
December	261	202	188	204	189	100	49	44	28	122	313	456	473	280	301	314	146	3720
Year....	2611	1750	1955	2932	3366	1521	951	1077	1750	2441	2694	3032	3407	3612	3840	3710	2567	43806

TORONTO METEOROLOGICAL RESULTS.

TABLE LXII.—*continue¹.*
1853-1862.

	N.	N.N.E.	N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.S.W.	S.W.	W.S.W.	W.	W.W.W.	N.W.	N.N.W.	Calms.	Sums.	
January	510	359	420	289	349	168	92	102	108	311	685	1131	795	506	516	609	490	7440
February	453	315	302	314	407	146	95	106	128	335	519	748	840	634	517	532	237	6768
March	294	165	233	488	536	189	176	108	186	349	558	539	810	942	978	601	283	7435
April	524	375	383	644	876	371	215	196	238	388	311	291	387	522	502	617	344	7200
May	513	276	374	746	338	412	220	340	411	573	279	194	272	421	521	748	402	7440
June	397	200	276	537	675	186	222	521	669	491	247	372	374	596	694	364	1056	7440
July	434	295	328	420	553	577	282	426	656	707	308	249	267	415	535	689	559	7440
August	475	372	298	307	490	290	278	282	513	597	391	239	381	577	688	750	506	7440
September	530	355	313	389	455	233	253	270	417	641	410	286	363	447	496	575	751	7194
October	461	337	368	458	474	255	123	178	261	460	452	376	548	723	553	536	654	7120
November	374	286	307	459	518	212	158	182	170	351	604	906	701	575	497	486	416	7182
December	545	443	384	354	324	200	102	93	69	240	680	1041	869	628	536	616	436	7440
Year	5510	3778	3887	5405	6303	3202	2179	2384	3772	5641	5728	6247	6695	6634	6935	7373	5512	87355

TABLE LXIII.
Number of hours that each wind blew in each quarter, including December of the same year, for 1853-57, 1858-62, and 1853-1862.

	N.	N.N.E.	N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.S.W.	S.W.	W.S.W.	W.	W.W.W.	N.W.	N.N.W.	Calms.	Sums.	
1853-57	820	621	549	404	447	289	143	143	173	406	888	1553	1242	796	719	920	671	10824
1854-52	688	496	557	553	563	225	146	158	132	500	976	1367	1262	872	850	867	592	10824
1853-52	1508	1117	1106	957	1080	514	289	301	305	906	1864	2820	2504	1668	1589	1477	1263	21648
1853-57	769	420	432	687	858	551	338	271	513	798	727	513	657	844	946	1079	626	11035
1853-52	562	390	558	1191	1394	421	273	272	337	512	421	511	812	1041	1055	887	403	11040
1853-52	1331	816	990	1878	2252	972	611	543	850	1310	1148	1024	1469	1885	2001	1966	1029	22075
1853-57	683	481	461	637	877	490	433	528	804	1002	649	360	534	657	697	845	755	10896
1858-62	623	386	342	627	847	513	312	402	865	971	691	375	486	709	1122	1185	674	11040
1853-62	1306	867	803	1264	1724	1003	745	930	1669	1973	1250	745	1020	1366	1819	2033	1429	21936
AVERAGE	627	470	490	685	715	351	314	365	532	694	70	789	765	733	816	923	10794	
1858-62	738	508	498	621	732	362	220	245	416	758	696	779	847	930	813	781	1597	1821
1853-62	1365	978	988	1306	1447	713	534	610	948	1452	1466	1568	1612	1745	1546	1802	21646	

TORONTO METEOROLOGICAL RESULTS.

TABLE LXIV.
Ratios showing for each month separately the comparative duration of the different winds, being the numbers in Table L.XVII. expressed in terms of their respective means for all winds, EXCLUDING clouds.
1853-1857.

	N.	N. N. E.	N. E.	E.	E. S. E.	S. E.	S. S. E.	S.	S. S. W.	S. W.	W. W. S. W.	W.	W. N. W.	N. W.	N. N. W.	
January.....	1.35	0.88	0.99	0.59	0.67	0.46	0.25	0.20	0.24	0.61	1.38	2.79	1.90	1.06	1.17	1.46
February.....	1.21	0.94	0.70	0.63	1.03	0.45	0.18	0.26	0.40	0.78	1.21	1.82	2.16	1.58	1.15	1.50
March.....	0.59	0.34	0.70	0.84	0.39	0.44	0.20	0.57	0.89	1.61	1.34	1.91	2.21	2.11	1.50	
April.....	1.54	1.12	0.94	1.12	1.59	0.96	0.52	0.51	0.72	1.30	0.86	0.57	0.58	0.89	1.27	1.51
May.....	1.45	0.53	0.72	1.36	1.55	1.20	0.60	0.55	1.08	1.50	0.86	0.44	0.49	0.75	0.96	1.96
June.....	0.78	0.48	0.84	0.39	1.74	0.89	0.60	0.53	1.22	1.60	1.29	0.66	1.00	0.98	1.01	0.99
July.....	1.08	0.81	0.67	0.65	1.42	0.91	0.66	1.23	1.46	1.76	0.84	0.53	0.57	0.81	0.90	1.40
August.....	1.33	0.97	0.68	0.69	1.01	0.53	0.79	0.73	1.12	1.38	0.96	0.51	0.96	1.31	1.38	1.61
September.....	1.32	1.07	0.80	0.96	1.16	0.70	0.73	0.77	1.02	1.24	1.13	0.69	0.62	0.96	1.29	1.54
October.....	1.05	0.75	1.04	1.07	1.02	0.55	0.28	0.52	1.09	1.10	0.87	0.91	1.37	1.47	1.41	1.47
November.....	0.70	0.47	0.55	1.30	1.28	0.46	0.51	0.49	0.49	1.03	1.72	2.19	1.72	1.25	0.88	0.97
December.....	1.31	1.11	0.90	0.69	0.62	0.46	0.24	0.22	0.19	0.54	1.60	2.69	1.82	1.14	1.08	1.39
Year.....	1.14	0.79	0.76	0.95	1.16	0.66	0.48	0.52	0.80	1.14	1.20	1.27	1.26	1.20	1.22	1.44
Arithmetic Means	{ 1.14	0.79	0.77	0.95	1.16	0.66	0.48	0.52	0.80	1.15	1.19	1.26	1.26	1.20	1.22	1.44

1858-1862.

	N.	N. N. E.	N. E.	E. N. E.	E.	E. S. E.	S. E.	S. S. E.	S.	S. S. W.	S. W.	W. S. W.	W.	W. N. W.	N. W.	N. N. W.
January.....	1.00	0.77	0.95	0.74	0.94	0.32	0.17	0.27	0.26	0.82	1.77	2.42	1.76	1.26	1.21	1.34
February.....	1.04	0.62	0.80	0.94	0.99	0.28	0.30	0.27	0.24	0.99	1.38	1.08	1.58	1.42	1.24	
March.....	0.73	0.49	0.70	1.48	1.56	0.46	0.34	0.28	0.26	0.67	0.88	1.08	1.71	2.01	2.27	1.19
April.....	0.93	0.64	0.85	0.85	1.86	2.49	0.77	0.49	0.40	0.47	0.53	0.60	0.78	1.20	1.53	1.08
May.....	0.88	0.72	0.98	2.03	2.25	0.67	0.40	0.54	0.79	1.11	0.42	0.45	0.75	1.16	1.41	1.44
June.....	1.11	0.47	0.49	1.18	1.50	0.72	0.29	0.53	1.27	1.60	1.07	0.52	0.78	0.81	1.82	1.84
July.....	0.94	0.56	0.39	1.01	1.16	0.65	0.66	0.75	1.49	1.53	0.87	0.62	0.67	1.12	1.58	
August.....	0.83	0.75	0.70	0.72	1.27	0.81	0.49	0.57	1.25	1.37	0.85	0.59	0.80	1.35	1.80	1.85
September.....	1.31	0.69	0.75	0.98	1.10	0.50	0.32	0.57	1.05	1.95	0.91	0.73	1.19	1.26	1.17	1.31
October.....	1.16	0.86	0.74	1.13	1.24	0.68	0.31	0.34	0.66	1.10	1.28	0.89	1.27	1.98	1.25	1.11
November.....	1.08	0.88	0.90	0.87	1.17	0.55	0.24	0.28	0.31	0.62	1.13	2.10	1.60	1.48	1.48	1.33
December.....	1.18	0.92	0.85	0.93	0.86	0.45	0.22	0.20	0.13	0.55	1.42	2.07	2.15	1.27	1.37	1.43
Year.....	1.01	0.69	0.76	1.16	1.38	0.59	0.37	0.42	0.68	1.06	1.05	1.18	1.32	1.40	1.49	1.44
Arithmetic Means	{ 1.02	0.69	0.76	1.15	1.38	0.59	0.37	0.42	0.68	1.07	1.05	1.18	1.32	1.40	1.49	1.44

TABLE LXIV.—*Continued.*
1853-1862.

	N.	N. N. E.	E. N. E.	E.	E. S. E.	S. E.	S. S. E.	S.	S. S. W.	S. W.	W. S. W.	W.	W. N. W.	X. W.	X. N. W.	
January.....	1.17	0.83	0.97	0.67	0.80	0.39	0.21	0.23	0.25	1.72	1.58	2.60	1.83	1.16	1.19	1.49
February.....	1.13	0.78	0.75	0.78	1.01	0.36	0.24	0.26	0.32	0.88	1.29	2.66	1.58	1.29	1.37	
March.....	0.66	0.37	0.52	1.09	1.20	0.42	0.39	0.24	0.42	0.78	1.25	1.21	1.81	2.19	1.34	
April.....	1.22	0.87	0.89	1.50	2.05	0.87	0.50	0.46	0.59	0.91	0.73	0.68	0.90	1.22	1.17	1.44
May.....	1.17	0.63	0.85	1.50	1.90	0.94	0.50	0.55	0.93	1.20	0.63	0.44	0.62	0.96	1.18	1.70
June.....	0.45	0.43	0.66	1.38	1.61	0.80	0.44	0.53	1.25	1.60	1.17	0.59	0.89	1.43	1.42	
July.....	1.01	0.69	0.53	0.98	1.29	0.88	0.66	0.99	1.48	1.64	0.86	0.58	0.62	0.96	1.24	1.60
August.....	1.10	0.86	0.69	0.71	1.14	0.67	0.64	0.65	1.18	1.38	0.90	0.55	0.88	1.33	1.59	1.75
September.....	1.32	0.88	0.78	0.97	1.13	0.60	0.63	0.67	1.04	1.59	1.02	0.71	0.90	1.11	1.23	1.43
October.....	1.11	0.81	0.88	1.10	1.14	0.62	0.29	0.43	0.87	1.10	1.09	0.90	1.31	1.73	1.33	1.29
November.....	0.88	0.68	0.73	1.09	1.22	0.50	0.37	0.38	0.40	0.83	1.43	2.14	1.66	1.36	1.18	1.15
December.....	1.24	1.01	0.88	0.81	0.74	0.46	0.23	0.21	0.16	0.55	1.51	2.38	1.98	1.21	1.22	1.41
Year.....	1.08	0.74	0.76	1.06	1.27	0.63	0.43	0.47	0.74	1.10	1.12	1.22	1.20	1.30	1.36	1.44
Arithmetic Means }	1.08	0.74	0.76	1.05	1.27	0.63	0.43	0.47	0.74	1.11	1.12	1.22	1.20	1.30	1.35	1.44

TABLE LXV.
Ratios showing for each quarter the comparative duration of the different winds, being the numbers in Table LXIII, expressed in terms of their respective means for all winds, EXCLUDING calms

	N.	N. N. E.	E. N. E.	E.	E. S. E.	S. E.	S. S. E.	S.	S. S. W.	S. W.	W. S. W.	W.	W. N. W.	X. W.	X. N. W.		
1853-57	1.29	0.98	0.86	0.64	0.77	0.45	0.23	0.22	0.27	0.64	1.40	2.45	1.96	1.25	1.13	1.45	
1858-62	1.08	0.78	0.87	0.86	0.93	0.35	0.23	0.25	0.21	0.78	1.53	2.14	1.97	1.36	1.33	1.34	
1853-62	1.18	0.88	0.87	0.75	0.85	0.10	0.23	0.24	0.24	0.71	1.46	2.29	1.96	1.31	1.23	1.40	
MATTHEW	1.853-57	1.18	0.65	0.66	1.06	1.32	0.85	0.52	0.42	0.79	1.23	1.12	0.79	1.01	1.30	1.45	1.65
APRIL	1.858-62	0.84	0.59	0.84	2.10	0.63	0.41	0.41	0.51	0.77	0.63	0.77	1.22	1.57	1.59	1.33	
MAY	1.853-62	1.01	0.62	0.75	1.43	1.71	0.74	0.46	0.41	0.65	1.00	0.87	0.78	1.12	1.43	1.52	1.50
JUNE	1.853-57	1.08	0.76	0.73	1.01	1.38	0.77	0.68	0.83	1.27	1.58	1.02	0.57	0.84	1.04	1.10	1.34
JULY	1.858-62	0.96	0.60	0.53	0.97	1.31	0.79	0.48	0.62	1.33	1.50	0.93	0.58	0.75	1.09	1.73	1.83
AUGUST	1.853-62	1.02	0.68	0.63	0.99	1.34	0.75	0.58	0.73	1.30	1.54	0.97	0.57	0.80	1.07	1.42	1.59
SEPTEMBER	1.853-57	1.02	0.76	0.79	1.11	1.46	0.57	0.51	0.59	0.86	1.13	1.25	1.28	1.24	1.32	1.19	1.32
OCTOBER	1.858-62	1.18	0.81	0.80	0.99	1.17	0.57	0.43	0.49	0.76	1.17	1.18	1.26	1.36	1.58	1.30	1.25
NOVEMBER	1.853-62	1.10	0.79	0.80	1.05	1.17									1.40	1.24	1.29

* * For Table LXVI, see Introduction, foot of page vii.

TORONTO METEOROLOGICAL RESULTS.

TABLE LXVII.
Ratios showing the annual distribution of each wind, being the numbers in Table LXIV, expressed in terms of their respective yearly Arithmetic means.

1853-1857.

	N.	N.N.E.	N.E.	E.N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.S.W.	S.W.	W.S.W.	W.	W.N.W.	N.W.	N.N.W.
January.....	1.18	1.12	1.29	0.62	0.58	0.69	0.52	0.38	0.30	0.53	1.16	2.21	1.51	0.89	0.96	1.01
February.....	1.06	1.19	0.91	0.66	0.59	0.68	0.37	0.49	0.50	0.68	1.01	1.45	1.72	1.31	0.95	1.04
March.....	0.52	0.44	0.45	0.74	0.72	0.59	0.92	0.39	0.71	0.78	1.35	1.06	1.52	1.84	1.73	1.04
April.....	1.34	1.41	1.24	1.18	1.37	1.35	1.07	0.90	0.90	1.14	0.72	0.45	0.46	0.74	1.04	1.04
May.....	1.27	0.67	0.94	1.42	1.34	1.81	1.24	1.06	1.35	1.31	0.72	0.35	0.39	0.63	0.79	1.36
June.....	0.68	0.61	1.10	0.99	1.50	1.34	1.24	1.02	1.53	1.40	1.08	0.52	0.79	0.82	0.83	0.68
July.....	0.94	1.02	0.88	1.46	1.22	1.37	1.35	2.38	1.83	1.54	0.70	0.42	0.45	0.67	0.74	0.97
August.....	1.19	1.23	0.88	0.73	0.87	0.80	1.64	1.41	1.39	1.21	0.80	0.41	0.77	1.09	1.13	1.12
September....	1.15	1.35	1.05	1.00	1.06	1.52	1.49	1.27	1.08	0.95	0.55	0.49	0.80	1.06	1.07	1.07
October.....	0.92	0.95	1.36	1.12	0.88	0.83	0.58	1.00	1.37	0.96	0.73	0.72	1.09	1.22	1.16	1.02
November....	0.61	0.60	0.72	1.36	1.10	0.69	1.05	0.95	0.62	0.90	1.44	1.73	1.36	1.04	0.72	0.67
December....	1.14	1.40	1.18	0.72	0.53	0.69	0.50	0.44	0.24	0.47	1.34	2.13	1.45	0.95	0.39	0.96

1858-1862.

	N.	N.N.E.	N.E.	E.N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.S.W.	S.W.	W.S.W.	W.	W.N.W.	N.W.	N.N.W.
January.....	0.98	1.12	1.25	0.64	0.68	0.54	0.47	0.65	0.38	0.77	1.69	2.05	1.33	0.90	0.81	0.94
February.....	1.02	0.90	1.06	0.81	0.72	0.47	0.80	0.65	0.35	0.92	1.31	1.61	1.32	1.13	0.55	0.86
March.....	0.71	0.57	0.92	1.28	1.13	0.78	0.93	0.67	0.39	0.63	0.84	0.91	1.29	1.43	1.52	0.83
April.....	0.91	0.93	1.12	1.61	1.81	1.31	1.32	0.97	0.69	0.50	0.57	0.66	0.91	1.09	0.72	0.96
May.....	0.87	1.05	1.29	1.76	1.64	1.14	1.09	1.16	1.03	1.16	1.03	0.40	0.38	0.56	0.83	1.00
June.....	1.09	0.69	0.64	1.02	1.09	1.23	0.79	1.25	1.86	1.49	1.02	0.44	0.59	0.58	1.22	1.28
July.....	0.93	0.82	0.52	0.87	0.84	1.44	1.78	1.80	2.13	1.42	0.83	0.53	0.51	0.80	1.06	1.25
August.....	0.82	1.08	0.92	0.63	0.92	1.37	1.34	1.37	1.84	1.28	0.81	0.50	0.60	0.96	1.21	1.29
September....	1.29	1.00	0.99	0.85	0.80	0.68	1.40	1.36	1.55	1.82	0.86	0.62	0.90	0.79	0.91	0.77
October.....	1.14	1.24	0.97	0.98	0.90	1.16	0.84	0.82	0.96	1.03	1.23	0.76	0.96	1.42	0.84	0.77
November....	1.06	1.27	1.19	0.75	0.85	0.93	0.64	0.66	0.45	0.58	1.98	1.78	1.21	1.05	0.99	0.92
December....	1.17	1.33	1.13	0.80	0.62	0.77	0.60	0.48	0.19	0.52	1.36	1.76	1.62	0.91	0.92	0.99

TORONTO METEOROLOGICAL RESULTS.

67

TABLE LXVII.—*Continued.*
1853–1862.

	N.	N.N.E.	N.E.	E.N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.S.W.	S.W.	W.S.W.	W.	W.N.W.	N.W.	N.N.W.
November	1.06	1.27	1.19	0.75	0.85	0.64	0.66	0.45	0.58	1.08	1.18	1.24	1.00	0.99	0.97	0.95
December	1.17	1.33	1.13	0.80	0.62	0.77	0.60	0.48	0.19	0.52	1.36	1.76	1.62	0.91	0.92	0.99

TABLE LXVIII.
Quarterly averages of the ratios in Table LXVII.

	N.	N.N.E.	N.E.	E.N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.S.W.	S.W.	W.S.W.	W.	W.N.W.	N.W.	N.N.W.
Winter	1.13	1.24	1.13	0.67	0.69	0.46	0.44	0.35	0.56	1.17	1.93	1.56	1.05	0.93	1.00	
Spring	1.04	0.84	0.88	1.11	1.14	1.28	1.08	0.81	0.99	1.08	0.93	0.62	0.70	1.07	1.15	
Summer	0.94	0.95	0.95	1.06	1.20	1.17	1.41	1.60	1.58	1.38	0.86	0.45	0.67	0.86	0.90	0.92
Autumn	0.89	0.97	1.04	1.16	0.99	0.86	1.05	1.15	1.09	0.98	1.04	1.00	0.98	1.02	0.98	0.92
<hr/>																
1853 to 1857	Winter	1.06	1.12	1.15	0.75	0.67	0.59	0.62	0.59	0.31	0.74	1.45	1.81	1.49	0.93	0.93
	Spring	0.83	0.85	0.86	1.11	1.53	1.08	1.11	0.98	0.75	0.72	0.60	0.65	0.92	1.12	1.06
	Summer	0.95	0.86	0.69	0.84	6.95	1.35	1.30	1.48	1.96	1.40	0.89	0.49	0.57	0.78	1.16
	Autumn	1.16	1.17	1.05	0.86	0.85	0.98	0.96	0.95	0.99	1.14	1.06	1.05	1.02	1.12	0.87
<hr/>																
1858 to 1862	Winter	1.09	1.18	1.14	0.71	0.67	0.64	0.53	0.51	0.33	0.65	1.30	1.87	1.53	1.01	0.97
	Spring	0.94	0.84	0.99	1.35	1.35	1.19	1.09	0.89	0.87	0.90	0.78	0.64	0.86	1.10	1.04
	Summer	0.94	0.91	0.83	0.94	1.06	1.25	1.36	1.55	1.76	1.39	0.87	0.47	0.62	0.82	1.10
	Autumn	1.02	1.06	1.04	0.99	0.92	1.01	1.05	1.04	1.06	1.05	1.03	1.00	1.07	0.92	0.89

TORONTO METEOROLOGICAL RESULTS.

WINDS in the upper strata as shown by the motion of the clouds, from six daily observations during the years 1851 to 1862 inclusive.

TABLE LXIX.

Absolute number of the several upper currents in each month, in the two half-years, and in the year; the summer half-year being considered to extend from April to September inclusive.

	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Winter.	Summer.	Year.
N.	18	12	12	25	26	24	27	28	26	45	33	38	159	150	314
N.E.	19	9	8	17	33	21	21	13	17	28	17	10	91	122	213
E.	24	32	38	49	46	41	54	34	35	43	50	28	215	262	477
S.E.	23	18	23	13	38	19	21	16	18	27	35	15	141	128	269
S.	10	6	15	11	30	10	4	9	15	21	23	20	95	79	174
S.W.	71	80	70	92	84	76	77	80	63	116	94	66	497	502	999
W.	273	259	292	361	282	413	481	415	395	305	319	253	1701	2347	4048
N.W.	137	123	164	140	155	145	183	235	156	200	214	181	1019	1014	2033
Calm Sky (a)	677	607	505	377	379	363	281	269	305	412	478	616	3325	1974	5299
Clear Sky (a)	185	167	301	272	364	283	312	362	312	263	135	132	1184	1935	3118
Means including (a)	144	131	143	136	144	140	146	146	140	146	140	139	812	852	1694
Means excluding (a)	72	67	78	88	87	94	109	104	94	98	98	76	490	576	1066

TABLE LXX.

Ratios shewing for each month separately as well as for the winter, summer, and year, the relative frequency of the several upper currents, including the cases of calm sky and clear sky, being the numbers in Table LXIX, expressed in terms of the means in the lowest line but one in the same table.

	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Winter.	Summer.	Year.
N.	0.13	0.09	0.08	0.18	0.18	0.17	0.18	0.19	0.18	0.31	0.24	0.27	0.19	0.18	0.19
N.E.	0.13	0.07	0.06	0.13	0.23	0.15	0.14	0.09	0.12	0.19	0.12	0.07	0.11	0.14	0.13
E.	0.17	0.24	0.27	0.30	0.32	0.32	0.37	0.23	0.25	0.29	0.36	0.20	0.26	0.31	0.28
S.E.	0.16	0.11	0.16	0.10	0.26	0.14	0.16	0.11	0.13	0.18	0.25	0.11	0.17	0.15	0.16
S.	0.07	0.05	0.10	0.08	0.21	0.07	0.03	0.06	0.11	0.11	0.17	0.14	0.11	0.09	0.10
S.W.	0.49	0.61	0.49	0.68	0.59	0.54	0.53	0.55	0.66	0.79	0.67	0.48	0.59	0.59	0.59
W.	1.90	1.97	2.01	2.60	1.96	2.95	3.29	2.81	2.82	2.09	2.28	1.83	2.02	2.75	2.39
N.W.	0.95	0.91	1.15	1.03	1.08	1.04	1.25	1.61	1.11	1.37	1.53	1.30	1.21	1.19	1.20
Calm Sky.	4.71	4.62	3.54	2.78	2.61	2.60	1.92	1.84	2.17	2.82	3.42	4.65	3.95	2.32	3.13
Clear Sky.	1.29	1.27	2.11	2.06	2.63	2.02	2.13	2.48	2.44	1.89	0.97	0.95	1.40	2.27	1.84

TABLE LXXI.

Ratios shewing for each month separately, as well as for the two half-years and year, the relative frequency of the several upper currents, excluding the cases of calm sky and clear sky, being the numbers in Table LXIX, expressed in terms of the means in the lowest line of that table.

	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Winter.	Summer.	Year.
N.	0.25	0.18	0.15	0.28	0.30	0.26	0.25	0.27	0.27	0.46	0.34	0.50	0.32	0.27	0.29
N.E.	0.26	0.13	0.10	0.19	0.38	0.22	0.19	0.12	0.18	0.28	0.17	0.13	0.19	0.21	0.26
E.	0.33	0.47	0.49	0.55	0.53	0.47	0.50	0.33	0.37	0.44	0.51	0.37	0.44	0.46	0.45
S.E.	0.32	0.27	0.30	0.15	0.44	0.20	0.22	0.15	0.19	0.27	0.36	0.23	0.29	0.22	0.25
S.	0.14	0.09	0.19	0.12	0.35	0.11	0.04	0.09	0.16	0.21	0.23	0.26	0.19	0.14	0.16
S.W.	0.99	1.19	0.90	1.01	0.97	0.81	0.71	0.77	0.98	1.18	0.96	0.86	1.02	0.87	0.94
W.	3.80	3.81	3.76	4.08	3.25	4.39	4.42	4.00	4.18	3.11	3.25	3.31	3.47	4.07	3.80
N.W.	1.91	1.82	2.11	1.58	1.79	1.54	1.68	2.27	1.65	2.04	2.18	2.37	2.08	1.76	1.91

use.
mer
Year.
314
218
477
174
269
990
4048
1033
5299
3118
1694

of the
LXXX.

Year.
0.19
0.13
0.28
0.16
0.10
0.59
2.39
1.20
3.13
1.84

y of the
LXXX.

Comparison of the half-yearly and yearly ratios in Tables LXXV and LXXVI, with the ratios derived in a similar manner from the observations made at 8 a.m., 2 p.m., and 4 p.m., as well as those derived from the observations at 10 p.m., midnight, and 6 a.m.

TABLE LXXII.

INCLUDING CALM SKY AND CLEAR SKY.

		10 P.M., Mid., and 6 A.M.		Six Hours.		EXCLUDING CALM SKY AND CLEAR SKY.		S.A.M., 2 P.M., and 4 P.M.		Six Hours.	
		S.A.M., 2 P.M., and 4 P.M.		Winter, Summer		Year.		S.A.M., Mid., and 6 A.M.		Six Hours.	
		Winter.	Summer.	Winter.	Summer.	Year.	Winter.	Summer.	Year.	Winter.	Summer.
N.	0.25	0.23	0.24	0.12	0.13	0.19	0.18	0.19	N.	0.30	0.27
N.E.	0.17	0.21	0.19	0.04	0.05	0.11	0.14	0.13	N.E.	0.21	0.23
E.	0.41	0.43	0.42	0.10	0.18	0.14	0.26	0.28	E.	0.49	0.48
S.E.	0.24	0.21	0.23	0.09	0.09	0.17	0.15	0.16	S.E.	0.29	0.24
S.	0.16	0.12	0.14	0.06	0.06	0.11	0.09	0.10	S.	0.19	0.14
S.W.	0.91	0.83	0.87	0.27	0.34	0.59	0.59	0.59	S.W.	1.09	0.93
W.	2.92	3.63	3.28	1.11	1.86	1.49	2.02	2.75	W.	3.52	4.06
N.W.	1.57	1.40	1.53	0.84	0.89	0.87	1.21	1.19	N.W.	1.90	1.66
Calm sky	2.50	1.43	1.96	5.41	3.22	4.31	3.95	2.32	Calm sky	3.13	2.32
Clear sky	0.87	1.40	1.14	1.94	3.16	2.55	1.40	2.27	Clear sky	1.84	1.75

TABLE LXXIII.

EXCLUDING CALM SKY AND CLEAR SKY.

		10 P.M., Mid., and 6 A.M.		Six Hours.		S.A.M., 2 P.M., and 4 P.M.		Six Hours.		S.A.M., 2 P.M., and 4 P.M.	
		Winter, Summer		Year.		Winter, Summer		Year.		Winter, Summer	
		Winter.	Summer.	Year.	Winter.	Summer.	Year.	Winter.	Summer.	Year.	Winter.
N.	0.25	0.23	0.24	0.12	0.13	0.19	0.18	0.19	N.	0.30	0.27
N.E.	0.17	0.21	0.19	0.04	0.05	0.11	0.14	0.13	N.E.	0.21	0.23
E.	0.41	0.43	0.42	0.10	0.18	0.14	0.26	0.28	E.	0.49	0.48
S.E.	0.24	0.21	0.23	0.09	0.09	0.17	0.15	0.16	S.E.	0.29	0.24
S.	0.16	0.12	0.14	0.06	0.06	0.11	0.09	0.10	S.	0.19	0.14
S.W.	0.91	0.83	0.87	0.27	0.34	0.59	0.59	0.59	S.W.	1.09	0.93
W.	2.92	3.63	3.28	1.11	1.86	1.49	2.02	2.75	W.	3.52	4.06
N.W.	1.57	1.40	1.53	0.84	0.89	0.87	1.21	1.19	N.W.	1.90	1.66

TABLE LXXIV.

Ratios showing the annual distribution of each upper wind, and comparing the frequency of each wind in the six winter months, with its frequency in the six summer months.

		January		February		March		April		May		June		July		August		Sept.		October		Nov.		Dec.		Ratios comparing Winter with Summer	
		January	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	2b. 4b & 20b. 10b. 12b & 18b. six hours.
N.	0.68	0.49	0.46	1.00	0.98	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	
N.E.	1.06	0.55	0.45	1.00	1.84	1.20	1.14	1.12	1.31	0.83	0.71	0.97	1.54	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	
E.	0.59	0.87	0.94	1.28	1.14	1.12	1.12	1.12	1.12	1.04	0.69	0.81	1.17	1.58	0.68	1.13	1.13	1.13	1.13	1.13	1.13	1.13	1.13	1.13	1.13	1.13	
S.E.	1.01	0.87	1.02	0.79	1.67	0.61	1.67	0.66	0.66	0.66	0.66	0.66	0.66	0.66	0.66	0.66	0.66	0.66	0.66	0.66	0.66	0.66	0.66	0.66	0.66	0.66	
S.	0.68	0.45	1.02	0.79	2.04	0.70	0.26	0.00	1.04	1.40	1.61	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	
S.W.	0.84	1.03	0.83	1.15	0.99	0.92	0.89	0.93	1.38	1.19	1.18	0.88	0.86	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	
W.	0.80	0.78	0.96	0.86	1.11	0.82	1.24	1.38	1.19	1.18	1.18	0.88	1.14	1.28	1.09	1.48	1.48	1.48	1.48	1.48	1.48	1.48	1.48	1.48	1.48	1.48	
N.W.	1.50	1.47	1.13	0.88	0.84	0.83	0.61	0.39	1.04	1.16	1.55	1.33	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	
Calm sky.	0.70	0.69	1.15	1.09	1.38	1.10	1.16	1.55	1.33	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00		
Clear sky.	0.70	0.69	1.15	1.09	1.38	1.10	1.16	1.55	1.33	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00		

TORONTO METEOROLOGICAL RESULTS.

WINDS IN THE UPPER STRATA.

TABLE LXXV.
Absolute number of the several upper currents at each of the six observation hours.

	6 A.M.	S.A.M.	2 P.M.	4 P.M.	10 P.M.	Mid.	6 A.M.	8 A.M.	2 P.M.	4 P.M.	10 P.M.	Mid.	6 A.M.	8 A.M.	2 P.M.	4 P.M.
	Mid.	2 P.M.	4 P.M.	Mid.	2 P.M.	4 P.M.	Mid.	2 P.M.	4 P.M.	Mid.	2 P.M.	4 P.M.	Mid.	2 P.M.	4 P.M.	Mid.
N.	45	69	68	71	31	30	106	208	52	N.	0.31	0.26	0.28	0.33	0.32	0.28
N.E.	24	59	61	52	12	14	50	163	35	N. E.	0.17	0.23	0.20	0.13	0.16	0.15
E.	77	129	120	107	20	24	121	356	79	E.	0.53	0.59	0.46	0.42	0.21	0.37
S.E.	47	72	63	57	10	20	77	192	45	S.E.	0.32	0.24	0.22	0.10	0.22	0.23
S.	27	42	45	34	12	14	53	121	29	S.	0.19	0.17	0.13	0.13	0.16	0.16
S.W.	137	229	274	236	63	58	260	739	166	S.W.	0.94	1.04	0.92	0.68	0.64	1.01
W.	532	787	974	1033	372	350	1254	2794	675	W.	3.66	3.38	4.04	3.87	3.89	3.80
N.W.	273	381	470	433	246	210	729	1304	359	N.W.	1.88	1.73	1.81	1.77	2.33	2.29
Calm sky (a)	1167	668	509	497	1216	247	3650	1069	883							
Clear sky (a)	497	412	265	294	818	832	2147	971	520							
Means including (a)	283	283	286	283	280	280	843	852	282							
Means excluding (a)	145	220	259	255	96	90	331	735	178							

TABLE LXXVI.
Ratios showing for each hour separately the relative frequency of the different upper winds, including the cases of calm and clear sky.

	6 A.M.	S.A.M.	2 P.M.	4 P.M.	10 P.M.	Mid.	6 A.M.	8 A.M.	2 P.M.	4 P.M.	10 P.M.	Mid.	6 A.M.	8 A.M.	2 P.M.	4 P.M.
	Mid.	2 P.M.	4 P.M.	Mid.	2 P.M.	4 P.M.	Mid.	2 P.M.	4 P.M.	Mid.	2 P.M.	4 P.M.	Mid.	2 P.M.	4 P.M.	Mid.
N.	0.16	0.24	0.24	0.25	0.11	0.13	0.24	N.	0.86	1.32	1.30	1.36	0.56	0.57	2.02	1.86
N.E.	0.09	0.18	0.21	0.18	0.04	0.05	0.06	N.E.	0.68	1.41	1.72	1.47	0.34	0.39	4.00	2.76
E.	0.27	0.46	0.42	0.38	0.07	0.09	0.14	E.	0.97	1.62	1.51	1.35	0.25	0.30	3.95	2.33
S.E.	0.17	0.25	0.22	0.20	0.04	0.07	0.09	S.E.	1.05	1.61	1.41	1.27	0.22	0.45	2.52	2.44
S.	0.10	0.15	0.16	0.12	0.04	0.05	0.06	S.	0.93	1.45	1.35	1.17	0.41	0.48	2.52	2.25
S.W.	0.48	0.81	0.96	0.83	0.23	0.21	0.31	S.W.	0.82	1.38	1.65	1.42	0.39	0.35	3.33	2.41
W.	1.88	2.78	3.42	3.64	1.33	1.25	1.49	W.	0.79	1.17	1.44	1.53	0.52	0.55	2.63	1.96
N.W.	0.97	1.34	1.65	1.60	0.88	0.75	0.87	N.W.	0.81	1.12	1.38	1.34	0.73	0.62	1.87	1.68
Calm sky.	4.13	2.34	1.79	1.75	4.34	4.46	4.31	Calm sky.	1.32	0.75	0.58	1.38	1.41	1.46	0.45	0.45
Clear sky.	1.76	1.45	0.93	1.4	2.92	2.97	2.55	Clear sky	0.96	0.79	0.51	0.57	1.57	1.60	0.45	0.45

TABLE LXXVII.

Ratios showing for each hour separately the relative frequency of the different upper winds, excluding the cases when the sky was clear and when it was clear.

	6 A.M.	S.A.M.	2 P.M.	4 P.M.	10 P.M.	Mid.	6 A.M.	8 A.M.	2 P.M.	4 P.M.	10 P.M.	Mid.	6 A.M.	8 A.M.	2 P.M.	4 P.M.
	Mid.	2 P.M.	4 P.M.	Mid.	2 P.M.	4 P.M.	Mid.	2 P.M.	4 P.M.	Mid.	2 P.M.	4 P.M.	Mid.	2 P.M.	4 P.M.	Mid.
N.	0.16	0.24	0.24	0.25	0.11	0.13	0.24	N.	0.86	1.32	1.30	1.36	0.56	0.57	2.02	1.86
N.E.	0.09	0.18	0.21	0.18	0.04	0.05	0.06	N.E.	0.68	1.41	1.72	1.47	0.34	0.39	4.00	2.76
E.	0.27	0.46	0.42	0.38	0.07	0.09	0.14	E.	0.97	1.62	1.51	1.35	0.25	0.30	3.95	2.90
S.E.	0.17	0.25	0.22	0.20	0.04	0.07	0.09	S.E.	1.05	1.61	1.41	1.27	0.22	0.45	2.52	2.44
S.	0.10	0.15	0.16	0.12	0.04	0.05	0.06	S.	0.93	1.45	1.35	1.17	0.41	0.48	2.52	2.25
S.W.	0.48	0.81	0.96	0.83	0.23	0.21	0.31	S.W.	0.82	1.38	1.65	1.42	0.39	0.35	3.33	2.41
W.	1.88	2.78	3.42	3.64	1.33	1.25	1.49	W.	0.79	1.17	1.44	1.53	0.52	0.55	2.63	1.96
N.W.	0.97	1.34	1.65	1.60	0.88	0.75	0.87	N.W.	0.81	1.12	1.38	1.34	0.73	0.62	1.87	1.68
Calm sky.	4.13	2.34	1.79	1.75	4.34	4.46	4.31	Calm sky.	1.32	0.75	0.58	1.38	1.41	1.46	0.45	0.45
Clear sky.	1.76	1.45	0.93	1.4	2.92	2.97	2.55	Clear sky	0.96	0.79	0.51	0.57	1.57	1.60	0.45	0.45

TORONTO METEOROLOGICAL RESULTS.

Winds in the Upper Strata, from six daily observations in the years 1860 to 1862, considered with reference to the Simultaneous Surface Winds.

TABLE LXXXIX.
Absolute Number of the several Upper Currents accompanying each Surface Wind.

the directions were moving	SURFACE WINDS.								Calm.
	N.	N.E.	E.	S.E.	S.	S.W.	W.	N.W.	
N.	29	1	2	0	5	3	3	51	2
N.E.	27	14	5	0	1	0	7	3	3
E.	16	44	86	6	2	1	6	1	1
S.E.	6	12	47	2	2	6	2	1	1
S.	3	9	12	3	6	3	4	2	4
S.W.	10	30	66	24	42	104	51	229	22
W.	106	71	116	61	159	258	313	99	99
N.W.	74	24	27	24	60	66	138	208	52
Calm.	288	253	320	75	116	228	225	301	180
Clear.	210	68	65	35	66	102	126	311	120
Total No. of each surface wind	769	526	746	237	454	773	861	1148	485
Mean*	77	53	75	24	45	77	86	115	48
Total No., excluding calm and clear sky	271	205	361	127	272	443	510	536	185
Mean†	34	26	45	16	34	55	64	67	23

TABLE LXXX.

Relative frequency of the different Upper Currents, including the cases when the sky was calm and when it was clear, being the numbers in Table LXXXIX, expressed in terms of the means of each corresponding surface wind given in the line marked *.

SURFACE WINDS.

the directions were moving	SURFACE WINDS.								Calm.
	N.	N.E.	E.	S.E.	S.	S.W.	W.	N.W.	
N.	0.38	0.02	0.03	0.08	0.00	0.06	0.03	0.44	0.04
N.E.	0.35	0.27	0.07	0.21	0.02	0.00	0.00	0.06	0.05
E.	0.21	0.84	1.15	0.35	0.04	0.01	0.01	0.65	0.04
S.E.	0.08	0.23	0.63	0.08	0.04	0.08	0.00	0.02	0.13
S.	0.04	0.17	0.6	0.13	0.13	0.04	0.05	0.08	0.09
S.W.	0.13	0.57	0.88	1.01	0.93	1.35	0.45	0.35	0.04
W.	1.38	1.25	1.55	2.57	3.50	3.34	3.61	1.99	0.17
N.W.	0.36	0.46	0.36	1.01	1.32	0.85	1.60	1.81	0.17
Calm.	3.75	4.81	4.29	3.16	2.96	2.95	2.61	2.62	3.71
Clear.	2.73	2.9	0.87	1.48	1.45	1.32	1.46	2.47	2.47

TABLE LXXXI.

Relative frequency of the different Upper Currents, excluding the cases when the sky was calm and clear, being the numbers in Table LXXXIX, expressed in terms of the means of each corresponding surface wind given in the line marked †.

SURFACE WINDS.

the directions were moving	SURFACE WINDS.								Calm.
	N.	N.E.	E.	S.E.	S.	S.W.	W.	N.W.	
N.	0.86	0.04	0.04	0.13	0.00	0.09	0.00	0.05	0.09
N.E.	0.80	0.55	0.1	0.31	0.03	0.00	0.10	0.13	0.13
E.	0.47	1.72	1.91	0.38	0.06	0.02	0.02	0.09	0.09
S.E.	0.18	0.47	1.04	0.13	0.06	0.11	0.00	0.03	0.04
S.	0.09	0.35	0.27	0.19	0.18	0.05	0.06	0.03	0.17
S.W.	3.29	1.17	1.46	1.51	1.24	1.88	0.86	0.46	0.95
W.	3.13	2.77	3.84	4.68	4.66	4.91	3.42	4.29	4.29
N.W.	2.18	0.94	0.60	1.51	1.76	1.19	2.16	3.10	2.25

TABLE LXXXII.

Comparative frequency of the several winds on the days in any part of the winter half-year, for the years 1850-51, 1855-56, and 1853-1862.

CLASS I. Light rain not exceeding 0.100 of an inch.

	N.	N.E.	E.N.E.	E.	E.S.E.	S.E.	S.	S.W.	S.W.S.	W.	W.W.W.	N.W.	S.N.W.	Calm.
On days of light rain (1)	1853-1857	170	100	138	207	185	114	71	93	126	171	209	246	245
	1858-1862	132	75	136	244	283	117	82	57	75	202	200	350	297
	1853-1862	362	175	274	451	468	231	153	150	201	381	559	696	196
												655	570	386
On days with and without rain (2)	1853-1857	1310	950	1052	1051	1153	584	406	396	624	1016	1789	2500	245
	1858-1862	1327	955	1062	1311	1455	589	340	353	398	1020	1680	2241	2309
	1853-1862	2637	1903	2014	2362	2008	1173	846	749	1022	2066	3478	4741	2508
												1463	3908	3357
Absolute value of each hour's mean wind speed (3)	1853-1857	130	105	145	197	160	145	175	235	202	171	167	138	129
	1858-1862	69	57	128	186	195	199	241	161	188	154	156	158	191
	1853-1862	114	99	136	191	174	197	265	200	197	184	161	146	146
Relative duration of each wind, or ratio of (1) to (2)	1853-1857	0.67	0.92	1.26	1.02	1.24	1.12	1.51	1.01	1.18	1.24	1.09	1.07	1.12
	1858-1862	0.62	0.89	1.16	0.98	1.24	1.24	1.24	1.24	1.24	1.24	1.24	1.02	0.92
	1853-1862	0.72	0.58	1.02	0.86	1.21	1.13	1.24	1.29	1.26	1.24	1.16	0.90	0.92
Ratio of numbers in (3) to their respective means for all	1853-1857	0.83	0.67	0.92	1.26	1.02	1.24	1.12	1.01	1.18	1.24	1.09	1.07	1.12
	1858-1862	0.62	0.49	0.86	1.16	1.24	1.24	1.24	1.24	1.24	1.24	1.24	1.02	0.92
	1853-1862	0.72	0.58	1.02	0.86	1.21	1.13	1.24	1.29	1.26	1.24	1.16	0.90	0.92

(Class II). Moderate rain (more than 0.100 and less than 0.500 of an inch).

	S.	N.E.	N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.W.	W.	W.N.W.	N.W.	S.W.W.	W.	W.N.W.	S.W.W.	Calm.
On days of moderate rain (1)	1853-1857	40	75	94	241	320	98	78	72	99	107	126	121	97	91	71	106
	1858-1862	89	76	106	311	363	135	65	72	65	157	186	171	78	42	82	70
	1853-1862	129	151	200	552	683	233	144	144	95	264	312	292	175	133	153	119
Absolute duration in hours of each wind (2) or ratio of (1) to (2)	1853-1857	1310	950	952	1051	1153	584	406	396	624	1046	1789	2500	2309	1850	1611	1758
	1858-1862	1327	955	1062	1311	1455	589	340	353	398	1020	1689	2241	2254	2058	1936	1642
	1853-1862	2637	1005	2014	2392	2608	1173	746	749	1022	2606	3478	1741	4563	3908	3537	3401
Relative duration of each wind (3)	1853-1857	0.31	0.79	0.99	0.229	0.278	0.168	0.192	0.182	0.158	1.02	0.70	0.48	0.42	0.49	0.43	0.28
	1858-1862	0.67	0.80	1.00	0.237	0.249	0.194	0.204	0.194	0.166	1.54	1.01	0.76	0.35	0.20	0.42	0.43
	1853-1862	0.49	0.79	0.99	0.234	0.262	0.199	0.163	0.192	0.128	0.99	0.62	0.38	0.34	0.45	0.35	0.94
Ratio of numbers in (2) to their respective means for all cases (4)	1853-1857	0.28	0.72	0.90	2.68	2.52	1.52	1.74	1.65	1.43	0.92	0.64	0.43	0.38	0.44	0.39	0.25
	1853-1862	0.62	1.54	0.64	0.85	0.80	1.90	2.00	2.24	1.70	1.55	1.64	1.37	1.09	0.77	0.53	0.69
	1853-1862	0.42	1.02	0.42	0.85	0.80	1.90	2.00	2.24	1.70	1.55	1.64	1.37	1.09	0.77	0.53	0.80

TORONTO METEOROLOGICAL RESULTS.

CLASS III. Rain 0.500 of an inch and upwards.

	N.	N.N.E.	N.E.	E.	E.N.E.	E.S.E.	S.E.	S.S.E.	S.	S.S.W.	S.W.	W.S.W.	W.	W.N.W.	N.W.	N.N.W.	Calms.	
On days of heavy rain.	1853-1857	3	11	19	96	136	54	21	16	26	35	27	14	11	5	5	25	
min. (1)	1858-1862	37	38	46	154	151	29	7	11	7	13	30	23	12	17	9	25	
1853-1862	40	49	65	250	287	83	28	27	19	39	50	57	37	23	17	22	34	
On days with and without rain.	1853-1857	1310	950	952	16	153	584	406	396	324	1046	1789	2500	2309	1850	1661	1758	1400
(2)	1858-1862	227	955	1002	1311	1455	589	310	353	308	1020	1689	2241	2254	2058	1936	1642	1216
On days with and without rain.	1853-1862	2637	1905	2014	2302	2608	1173	746	749	1022	2066	3478	4741	4563	3908	3357	3400	2616
Absolute duration in hours.	Relative duration of each wind, or ratios of (1) to (2).	002	012	020	001	118	062	032	040	019	025	020	.011	.006	.003	.003	.018	
(3)	1853-1862	.028	.040	.043	.117	.104	.049	.021	.031	.018	.013	.009	.013	.010	.006	.010	.007	
Ratios of the numbers in (3) to their respective means for all winds.	1853-1857	.015	.026	.032	.106	.110	.071	.038	.036	.019	.019	.014	.012	.008	.006	.006	.013	
(4)	1858-1862	.06	.38	.63	.287	.373	.291	.144	.126	.60	.79	.63	.35	.19	.19	.10	.09	
Relative duration of each wind, or ratios of (1) to (2).	1853-1862	0.91	1.30	1.35	3.79	3.57	1.39	0.68	1.00	0.58	0.42	0.29	0.42	0.32	0.19	0.32	0.25	
(3)	1853-1862	0.48	0.82	1.01	3.35	3.48	2.25	1.20	1.14	0.60	0.60	0.44	0.38	0.29	0.19	0.16	0.19	

Rain without reference to amount.

	N.	N.N.E.	N.E.	E.	E.N.E.	E.S.E.	S.E.	S.S.E.	S.	S.S.W.	S.W.	W.S.W.	W.	W.N.W.	N.W.	N.N.W.	Calms.	
On days of rain.	1853-1857	213	186	251	544	641	266	170	181	237	312	460	494	409	278	259	291	376
min. (1)	1858-1862	258	189	288	709	797	281	155	140	118	372	461	551	458	448	287	345	345
1853-1862	471	375	539	1253	1438	547	325	321	335	634	921	1045	867	726	556	576	721	
On days with and without rain.	1853-1857	1310	950	952	1051	1153	584	406	396	624	1046	1789	2500	2309	1850	1661	1758	1400
(2)	1858-1862	1327	955	1062	1311	1455	589	340	333	398	1020	1689	2241	2254	2058	1936	1642	1216
1853-1862	2637	1905	2014	2302	2608	1173	746	749	1022	2066	3478	4741	4563	3908	3357	3400	2616	
Absolute duration in hours.	Relative duration of each wind, or ratios of (1) to (2).	002	012	020	001	118	062	032	040	019	025	020	.011	.006	.003	.003	.018	
(3)	1853-1862	.028	.040	.043	.117	.104	.049	.021	.031	.018	.013	.009	.013	.010	.006	.010	.007	
Ratios of the numbers in (3) to their respective means for all winds.	1853-1857	.06	.38	.63	.287	.373	.291	.144	.126	.60	.79	.63	.35	.19	.19	.10	.09	
(4)	1858-1862	0.48	0.82	1.01	3.35	3.48	2.25	1.20	1.14	0.60	0.60	0.44	0.38	0.29	0.19	0.16	0.19	

TORONTO METEOROLOGICAL RESULTS.

TABLE LXXXIII.
*Comparative frequency of the several winds on the days in any part of which rain fell during the summer half-year, for the years 1853-57,
 1858-62, and 1863-67.*

Class I. Light rain not exceeding 0.100 of an inch.

	N.	N.N.E.	S.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.W.	W.S.W.	W.	W.N.W.	N.W.	N.N.W.	Calms			
On days of light rain.	1853-1857	198	325	396	214	139	230	391	290	204	214	204	262	212	350			
	1858-1862	280	110	202	317	182	155	160	115	205	421	316	324	428	236			
	1863-1867	515	400	612	878	369	254	444	812	600	538	616	680	812	615			
On days with and without rain.	1853-1857	1580	1048	980	1362	1784	1097	822	911	1398	1854	1245	715	880	1202	1434	1905	1575
	1858-1862	1284	825	815	1681	2111	932	611	724	1352	1721	1065	791	1153	1554	1904	2068	1351
	1863-1867	1873	1873	1873	2043	3865	2029	1433	1635	2750	3575	2250	1506	2042	2746	2338	3973	2926
Relative duration of each wind, or ratios of (1) to (2), or ratios of (1) to (3).	1853-1857	148	149	202	259	222	195	168	153	171	211	233	285	308	229	183	164	228
	1858-1862	218	133	226	206	225	166	161	150	152	245	308	422	323	218	288	207	189
	1863-1867	179	142	214	221	182	166	155	161	227	267	357	316	247	243	186	210	
Ratios of the numbers in (3) to their respective means for all winds.	1853-1857	0.72	0.73	0.49	1.17	1.98	0.95	0.82	0.75	0.84	1.03	1.14	1.39	1.51	1.07	0.89	0.80	1.11
	1858-1862	0.95	0.58	0.49	0.90	0.98	0.72	0.71	0.69	0.66	1.07	1.34	1.84	1.41	1.17	1.25	0.90	0.82
	1863-1867	0.82	0.65	0.48	1.02	1.03	0.84	0.76	0.71	0.74	1.04	1.23	1.64	1.45	1.14	1.12	0.85	0.66

Class II. Moderate rain (more than 0.100 and less than 0.500 of an inch).

	N.	N.N.E.	S.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.W.	W.S.W.	W.	W.N.W.	N.W.	N.N.W.	Calms			
On days of moderate rain.	1853-1857	163	157	261	207	102	125	130	220	292	220	135	125	166	191	195	150	
	1858-1862	138	81	153	321	306	146	63	78	118	192	172	124	138	158	179	130	
	1863-1867	301	238	236	582	574	308	185	298	338	484	392	259	263	324	376	374	
On days with and without rain.	1853-1857	1389	1048	980	1362	1784	1097	822	911	1398	1854	1245	715	880	1202	1434	1905	1575
	1858-1862	1881	825	893	1681	2111	932	611	724	1352	1721	1065	791	1153	1554	1904	2068	1351
	1863-1867	2873	1873	1873	3043	3865	2029	1433	1635	2750	3575	2250	1506	2042	2746	2338	3973	2926
Relative duration of each wind, or ratios of (1) to (2), or ratios of (1) to (3).	1853-1857	102	150	187	192	156	148	152	113	135	157	177	189	141	138	133	102	905
	1858-1862	108	998	171	191	145	157	103	108	987	112	171	157	120	192	.097	.087	.103
	1863-1867	105	127	179	191	147	152	131	127	125	185	174	172	129	118	113	.094	.099
Ratios of the numbers in (3) to their respective means for all winds.	1853-1857	0.69	1.01	1.27	1.30	1.01	1.00	1.03	0.97	1.06	1.20	1.28	0.95	0.93	0.90	0.69	0.64	
	1858-1862	0.87	0.73	1.37	1.53	1.16	1.26	0.83	0.87	0.70	0.90	1.37	1.26	0.96	0.82	0.78	0.70	0.82
	1863-1867	0.77	0.93	1.31	1.40	1.08	1.12	0.96	0.93	0.90	0.99	1.28	1.26	0.95	0.87	0.83	0.69	0.73

CLASS III. Rain 0.500 of an inch and upwards.

CLASS III. Rain 0-500 of an inch and upwards.											Calms.			
	N.	S.N.E.	E.N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.W.	W.S.W.	W.	W.N.W.	N.W.	N.N.W.
On days of heavy rain. (1)	1853-1857	99	48	112	163	218	162	47	72	59	107	89	19	40
	1858-1862	62	60	83	284	220	30	50	22	49	111	115	46	52
On days with and without rain. (2)	1853-1857	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862
	1853-1857	1589	1048	986	1362	1784	1097	822	911	1398	1854	715	886	1202
Absolute difference in hours of quiet wind from ratios of (1) to (2).	1853-1857	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862
	1853-1857	557	446	114	121	122	93	67	79	142	105	71	0.27	0.41
Relative duration of each wind, or ratios of (1) to (2). (3)	1853-1857	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862
	1853-1857	0.87	0.76	1.74	1.85	1.86	1.42	0.87	1.21	0.64	0.86	0.41	0.69	0.87
Ratios of the numbers in (3) to their respective means for all winds. (4)	1853-1857	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862	1853-1862
	1853-1857	0.75	1.13	1.44	2.62	1.61	0.98	1.27	0.47	0.56	0.90	1.77	0.50	0.70
	1853-1862	0.82	0.89	1.60	2.28	1.72	1.21	1.03	0.89	0.60	0.94	1.40	0.66	0.69

Rain without reference to amount.

	N.	N.N.E.	N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.W.	W.S.W.	W.	W.S.W.	N.W.	N.N.W.	Calms.			
On days of rain.	1853-1857	488	361	493	751	881	478	310	341	518	740	309	258	439	499	518	580	573
(1)	1854-1862	480	251	438	952	1009	360	213	215	372	724	307	504	562	623	781	676	450
On days with and without rain.	1853-1862	968	612	931	1703	1890	838	523	556	890	1514	1196	862	1001	1122	1260	1256	1063
(2)	1854-1857	1859	1048	980	1362	1784	1097	822	911	1398	1854	1245	715	859	1202	1434	1905	1575
Relative duration of each wind, hours, or ratios of (1) to (2).	1853-1862	1284	825	893	1681	2111	932	611	724	1382	1721	1005	791	1351	1554	1904	2008	1351
Ratios of the numbers in (3) to their respective means for all winds.	1853-1857	374	304	490	566	478	386	319	297	421	594	437	371	481	415	361	394	364
(3)	1853-1862	337	327	497	560	485	413	365	340	324	425	532	572	490	407	389	316	313
1853-1857	0.73	0.82	1.20	1.32	1.18	1.04	0.90	0.89	0.89	1.02	1.15	1.20	1.18	1.09	0.86	0.73	0.87	
1853-1862	0.89	0.73	1.17	1.35	1.14	1.02	0.83	0.71	0.66	1.01	1.42	1.52	1.16	0.96	0.98	0.78	0.76	
1853-1857	0.74	0.82	1.20	1.32	1.18	1.04	0.90	0.89	0.89	1.02	1.15	1.20	1.18	1.09	0.86	0.73	0.87	
1853-1862	0.81	0.74	1.17	1.34	1.16	1.00	0.87	0.81	0.77	1.01	1.27	1.37	1.17	0.97	0.93	0.75	0.82	

TORONTO METEOROLOGICAL RESULTS.

TABLE LXXXIV.
Comparative frequency of the several winds on the days in any part of which rain fell during the year, for the years 1853-1857, 1858-62,
and 1853-1862.

CLASS I. Light rain not exceeding 0.100 of an inch.

	N.	N.N.E.	S.E.	E.S.E.	E.	E.S.W.	S.E.	S.S.E.	S.	S.S.W.	S.W.	W.S.W.	W.	W.N.W.	N.W.	N.N.W.	Calm.	
On days of light rain.	256	336	581	328	209	232	315	570	589	550	572	440	446	549	604			
1853-1857	405	591	765	272	182	172	280	623	634	729	810	732	626	452				
1858-1862	817	441	674	1123	1346	600	391	404	65	1193	1159	1234	1301	1250	1198	1175	1056	
On days with and without rain.	1853-1857	2899	1998	1932	2413	2937	1681	1228	1307	2022	2900	3034	3215	3198	3052	3095	2963	2975
1858-1862	2611	1780	1955	2892	3366	1521	951	1077	1750	2741	2694	3032	3407	3812	3840	3710	2567	
Relative duration of each wind, or ratios of (1) to (2).	1853-1862	5610	3778	3887	5403	6303	4202	2779	2884	5772	5641	5728	6247	6805	6694	6935	7373	5542
Ratios of the numbers in (3) to their respective means for all winds.	1853-1857	140	128	174	220	198	195	170	178	181	197	191	171	179	144	150	.203	
(3)	1858-1862	158	104	173	198	215	179	191	160	160	227	212	226	214	224	196	169	.176
1853-1862	148	117	173	298	267	187	179	169	171	211	202	198	197	188	173	159	.191	
1853-1857	0.50	0.73	1.00	1.26	1.14	1.12	0.97	1.02	1.04	1.13	1.11	0.98	1.03	0.83	0.82	0.86	1.16	
1858-1862	0.84	0.56	0.93	1.05	1.15	0.96	1.02	0.85	0.86	1.21	1.18	1.21	1.14	1.20	1.05	0.90	0.94	
1853-1862	0.82	0.65	0.96	1.15	1.14	1.03	0.99	0.93	0.94	1.17	1.12	1.09	1.03	1.04	0.35	0.88	1.05	

CLASS II. Moderate rain (more than 0.100 and less than 0.500 of an inch.)

	N.	N.N.E.	N.E.	E.N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.S.W.	S.W.	W.S.W.	W.	W.N.W.	N.W.	N.N.W.	Calm.	
On days of moderate rain.	1853-1857	203	222	277	502	587	260	203	202	319	399	346	256	222	257	262	244	256
1858-1862	227	157	250	632	670	281	129	150	184	349	358	295	216	200	267	249	279	
1853-1862	430	380	536	1134	1257	541	332	352	503	748	704	531	438	457	520	493	535	
On days with and without rain.	1853-1857	2899	1998	1932	2413	2937	1681	1228	1307	2022	2900	3034	3215	3198	3052	3095	2963	2975
1858-1862	2611	1780	1955	2892	3366	1521	951	1077	1750	2741	2694	3032	3407	3812	3840	3710	2567	
1853-1862	5510	3778	3887	5403	6303	4202	2779	2884	5772	5641	5728	6247	6805	6694	6935	7373	5542	
Relative duration of each wind, or ratios of (1) to (2).	1853-1857	0.70	1.16	1.43	2.08	2.00	1.55	1.65	1.54	1.58	1.38	1.14	0.80	0.69	0.84	0.85	0.67	
(3)	1858-1862	0.87	0.78	1.03	1.38	2.10	1.93	1.63	1.52	1.48	1.33	1.23	0.88	0.66	0.69	0.70	0.67	
1853-1862	0.57	0.94	1.16	1.09	1.62	1.34	1.26	1.28	1.12	0.93	0.65	0.56	0.48	0.54	0.54	0.70		
Ratios of the numbers in (3) to their respective means for all winds.	1853-1857	0.75	1.13	1.80	1.60	1.58	1.16	1.19	1.08	1.01	1.14	0.83	0.54	0.47	0.60	0.57	0.38	
1858-1862	0.65	0.86	1.15	1.73	1.61	1.41	1.26	1.23	1.11	1.11	1.02	0.73	0.55	0.57	0.63	0.56	0.81	
1853-1862	0.65	0.95	1.15	1.73	1.61	1.41	1.26	1.23	1.11	1.11	1.02	0.73	0.55	0.57	0.63	0.56	0.81	

TORONTO METEOROLOGICAL RESULTS.

CLASS III. Rain 0.500 of an inch and upwards.

	N.	N.E.	N.	E.N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.W.	W.S.W.	W.	W.N.W.	N.W.	N.N.W.	Cards	
On days of heavy rain.	93	59	131	261	354	156	68	88	71	133	124	46	54	80	69	89	
(1)	1858-1862	99	121	458	371	88	57	33	56	124	130	76	61	59	86	44	
On days with and without rain.	1853-1862	192	157	260	639	75	244	125	121	127	257	254	122	129	141	128	133
(2)	1858-1862	5289	1982	2413	2337	1681	1228	1307	2027	2906	3034	3215	3498	3452	3005	3663	2475
Absolute difference in of each wind of each number in (1) to (2).	1853-1862	3611	1780	1955	2992	3566	1521	951	107	1750	2741	3694	3032	2407	3612	2840	2370
(3)	1853-1862	5510	3778	3887	5405	6503	3202	2179	2984	3772	5644	5728	6247	6005	6664	6935	7373
Relative duration of each wind, or ratios of (1) to (2).	1853-1857	0.32	0.80	0.68	1.08	1.21	0.93	0.55	0.67	0.35	0.46	0.41	0.14	0.17	0.26	0.22	0.30
(3)	1858-1862	0.33	0.65	0.66	1.46	1.04	0.58	0.60	0.31	0.31	0.66	0.44	0.19	0.20	0.17	0.18	0.24
Ratios of the numbers in (3) to their respective means for all winds.	1853-1857	0.66	1.40	2.22	2.49	1.91	1.13	1.38	0.72	0.46	0.84	0.29	0.35	0.54	0.45	0.43	0.62
(3)	1858-1862	0.81	1.17	1.40	3.09	2.20	1.23	1.27	0.66	0.68	0.95	1.02	0.53	0.47	0.36	0.32	0.49
(4)	1853-1862	0.73	0.87	1.39	2.68	2.33	1.58	1.19	1.06	0.71	0.96	0.92	0.40	0.42	0.44	0.37	0.50

Pain without reference to amount.

	N.	N.E.	E.N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.W.	W.	W.S.W.	W.	W.W.W.	N.W.	N.N.W.	Colins.	
On days of rain.	1855-1857	701	547	744	1296	744	480	522	755	1102	1050	852	848	777	777	949	
(1)	1856-1862	728	440	726	1661	800	641	368	955	520	1066	1058	1055	1020	1071	1075	961
On days with and without rain.	1855-1862	1439	987	1470	2956	3328	1385	848	877	1215	2148	2117	1907	1698	1848	1855	1832
(2)	1855-1862	5510	3778	3887	5405	6503	3202	2179	1228	1307	2022	2900	3034	3215	3198	3052	2975
Absolute duration in hours of each wind.	Relative duration of each wind, or ratios of (1) to (2).	1855-1857	2899	1998	2413	2937	1681	951	1077	1750	2741	2894	3482	3407	3612	3440	3710
(3)	1856-1862	2611	1780	1956	2492	3566	1521	951	1077	1750	2741	2894	3482	3405	3612	3440	3710
Ratios of the numbers in (3) to their respective means for all winds.	1855-1857	0.70	0.79	1.11	1.55	1.50	1.28	1.13	1.15	1.08	1.10	1.01	1.07	0.77	0.74	0.69	0.92
(4)	1856-1862	0.81	0.81	1.06	1.58	1.44	1.29	1.10	0.94	1.14	1.12	0.99	0.85	0.84	0.80	0.74	0.86

TORONTO METEOROLOGICAL RESULTS.

TABLE LXXXV.
Comparative frequency of the several winds on the days in any part of which snow fell during the year, for the years 1853-57,
1858-62, and 1853-62.

CLASS I. Light snow not exceeding one inch.

	S.	N.N.E.	N.E.	E.N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.W.	W.S.W.	W.	W.W.	N.W.	S.W.	S.S.W.	Calm.
On days of light snow.	1853-1857	464	298	274	151	168	169	105	84	120	168	486	806	835	614	586	622
(1)	1853-1862	471	275	183	227	230	163	69	105	111	211	388	862	939	917	724	644
1853-1862	635	574	457	378	398	272	174	189	231	379	874	1668	1794	1561	1319	1266	
On days with and without snow.	1853-1862	2849	1998	1322	2413	2037	1681	1228	1307	2022	2060	2034	3215	3108	3052	3095	3063
(2)	1853-1862	5310	3778	3887	5405	6503	3204	2173	2384	3772	5611	5728	6247	6635	6654	6335	7475
Relative duration of each wind, or ratios of (1) to (2).	1853-1857	160	150	142	103	657	101	686	684	59	658	160	251	261	201	170	107
1853-1862	180	154	104	976	694	698	673	697	663	144	144	284	281	286	189	174	108
1853-1862	170	152	118	970	661	685	680	679	661	167	153	266	272	234	189	172	668
Ratios of the numbers in (3) to their respective means for all winds.	1853-1857	1.20	1.12	1.06	0.47	0.43	0.76	0.64	0.48	0.44	0.43	1.20	1.88	1.96	1.51	1.42	1.27
(4)	1853-1862	1.29	1.10	0.67	0.54	0.46	0.49	0.52	0.69	0.45	0.55	1.03	2.03	2.01	1.87	1.35	1.24
1853-1862	1.24	1.11	0.86	0.51	0.45	0.62	0.58	0.45	0.49	1.12	1.94	1.49	1.71	1.38	1.26	1.22	

CLASS II. Moderate snow, more than one inch and less than five inches.

	N.	N.N.E.	N.E.	E.N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.W.	W.S.W.	W.	W.W.	N.W.	S.W.	S.S.W.	Calm.
On days of moderate snow.	1853-1857	113	170	203	121	126	77	74	34	56	87	125	125	145	112	103	214
(1)	1853-1862	124	150	206	212	198	58	27	50	36	52	70	104	152	76	123	138
1853-1862	207	220	409	323	324	135	101	84	92	139	205	239	297	188	226	332	
On days with and without snow.	1853-1857	1859	1998	1932	2413	2037	1681	1228	1307	2022	2060	2034	3215	3198	3052	3095	3363
(2)	1853-1862	2511	1780	1655	2992	3565	2912	3516	1231	951	1077	150	2141	2694	3032	3467	3612
1853-1862	5510	3778	3887	5405	6503	3202	2170	2384	3772	5641	5728	6247	6605	6664	6335	7375	5542
Relative duration of each wind, or ratios of (1) to (2).	1853-1857	0.93	0.85	0.84	0.84	0.84	0.84	0.84	0.84	0.84	0.84	0.84	0.84	0.84	0.84	0.84	0.84
1853-1862	0.98	0.48	0.85	0.85	0.85	0.85	0.85	0.85	0.85	0.85	0.85	0.85	0.85	0.85	0.85	0.85	0.85
1853-1862	0.98	0.48	0.85	0.85	0.85	0.85	0.85	0.85	0.85	0.85	0.85	0.85	0.85	0.85	0.85	0.85	0.85
Ratios of the numbers in (3) to their respective means for all winds.	1853-1857	1.04	1.79	2.22	0.96	0.91	0.97	1.27	0.59	0.59	0.59	0.89	0.95	0.78	0.70	1.22	0.49
(4)	1853-1862	1.04	1.82	2.26	1.53	1.21	1.82	0.60	0.99	0.45	0.41	0.56	0.73	0.97	0.45	0.69	0.0
1853-1862	1.02	1.81	2.12	1.52	1.06	0.89	0.98	0.75	0.51	0.53	0.77	0.81	0.96	0.60	0.70	1.02	0.55

ratios of the numbers in (3) to their respective means for all winds.

(4)

CLASS III. Heavy Snow, five inches and upwards.

	N.	S.	E.N.E.	E.	E.S.E.	E.W.	S.E.	S.S.E.	S.	S.W.	W.	W.W.	W.	W.W.W.	S.W.	S.W.W.	Gains.	
On days of heavy snow.	1853-1857	38	44	42	63	19	2	1	3	5	6	14	10	12	13	22	7	
(1)	1858-1862	21	21	37	52	62	6	3	0	6	17	15	15	8	18	19	9	
1853-1862	39	65	81	94	125	25	5	4	3	11	23	29	25	20	31	41	16	
On days with and without snow.	1853-1857	2869	1998	1932	2413	2037	1681	1228	1307	2022	2900	3034	3215	3198	3052	3095	3163	2975
(2)	1858-1862	2611	1780	1935	2092	3566	1521	951	1077	1750	2741	2694	31032	3407	3612	3849	3710	2567
Relative duration of each wind, or ratios of (1) to (2).	1853-1862	0.08	0.12	0.19	0.17	0.17	0.04	0.03	0.02	0.00	0.02	0.02	0.04	0.04	0.04	0.06	0.02	
(3)	1853-1862	0.11	0.17	0.21	0.17	0.19	0.08	0.02	0.02	0.01	0.02	0.04	0.05	0.04	0.03	0.05	0.04	
Ratios of the numbers in (3) to their respective means for all winds.	1853-1857	1.59	2.69	2.81	2.08	2.57	1.34	0.25	0.12	0.24	0.25	0.49	0.49	0.49	0.49	0.73	0.24	
(4)	1853-1862	1.17	1.76	2.78	2.49	0.59	0.44	0.44	0.00	0.29	0.88	0.73	0.59	0.29	0.73	0.59	0.39	
Relative duration of each wind, or ratios of (1) to (2).	1853-1862	1.44	2.22	2.75	2.22	2.49	1.05	0.26	0.13	0.26	0.52	0.65	0.52	0.39	0.65	0.78	0.39	

Snow without reference to amount.

	N.	S.	E.N.E.	E.	E.S.E.	E.W.	S.E.	S.S.E.	S.	S.W.	W.	W.W.	W.	W.W.W.	S.W.	S.W.W.	Gains.	
On days of snow.	1853-1857	64.5	51.3	52.1	31.4	35.7	22.5	18.1	11.9	17.9	23.0	62.7	93.5	89.0	73.8	70.2	85.8	36.5
(1)	1858-1862	61.6	44.6	51.6	4.91	4.75	9.0	1.58	1.47	2.89	4.75	9.81	11.26	10.31	8.01	8.01	3.38	7.63
On days with and without snow.	1853-1857	2869	1998	1932	2413	2037	1681	1228	1307	2022	2900	3034	3215	3198	3052	3095	3163	2975
(2)	1858-1862	2611	1780	1935	2092	3566	1521	951	1077	1750	2741	2694	31032	3407	3612	3849	3710	2567
Relative duration of each wind, or ratios of (1) to (2).	1853-1862	0.08	0.12	0.19	0.17	0.19	0.04	0.03	0.02	0.01	0.02	0.04	0.05	0.04	0.03	0.05	0.04	
(3)	1853-1862	0.11	0.17	0.21	0.17	0.19	0.08	0.02	0.02	0.01	0.02	0.04	0.05	0.04	0.03	0.05	0.04	
Ratios of the numbers in (3) to their respective means for all winds.	1853-1857	1.17	1.56	1.43	0.69	0.64	0.84	0.78	0.48	0.47	0.48	1.00	1.57	1.64	1.28	1.20	0.65	
(4)	1858-1862	1.22	1.30	1.37	0.85	0.71	0.57	0.54	0.76	0.43	0.51	0.91	1.08	1.21	1.48	1.17	1.12	
Relative duration of each wind, or ratios of (1) to (2).	1853-1862	1.20	1.33	1.40	0.78	0.68	0.71	0.67	0.61	0.45	0.49	1.00	1.62	1.67	1.38	1.18	0.66	

TORONTO METEOROLOGICAL RESULTS.

TABLE LXXXVI.
Synopsis of the Final Ratios given in Tables LXXXVII. to LXXXVII., corresponding to the different classes of Rain and Snow.

Rain in the Winter Half-year.																		
	N.	S.N.E.	S.E.	E.N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.S.W.	W.	W.S.W.	W.	W.N.W.	N.W.	N.N.W.	Calm.	
1853-57.	Class I.....	0.83	0.67	0.92	1.26	1.02	1.24	1.12	1.50	1.29	1.00	1.07	0.88	0.82	0.61	0.70	0.86	1.12
	" II.....	0.38	0.72	0.90	2.08	2.52	1.52	1.74	1.65	1.43	0.93	0.63	0.44	0.38	0.44	0.39	0.25	0.69
	" III.....	0.06	0.38	0.63	2.88	3.73	2.91	1.64	1.26	0.60	0.79	0.63	0.35	0.19	0.19	0.10	0.09	0.57
	Rain generally.....	0.55	0.66	0.88	1.73	1.86	1.62	1.40	1.53	1.27	1.00	0.86	0.66	0.59	0.50	0.52	0.56	0.90
1858-62.	Class I.....	0.62	0.49	0.80	1.16	1.22	1.24	1.51	1.01	1.17	1.24	0.96	0.98	0.99	1.19	0.66	0.76	1.01
	" II.....	0.54	0.64	0.80	1.90	2.00	1.84	1.55	1.63	1.33	1.23	0.68	0.61	0.28	0.31	0.34	0.34	0.92
	" III.....	0.91	1.30	1.39	3.70	3.37	1.59	0.68	1.00	0.58	0.42	0.29	0.42	0.32	0.19	0.19	0.32	0.23
	Rain generally.....	0.61	0.63	0.86	1.71	1.74	1.51	1.44	1.26	1.18	1.15	0.86	0.78	0.64	0.48	0.55	0.55	0.90
1863-67.	Class I.....	0.72	0.58	0.86	1.20	1.13	1.24	1.29	1.26	1.24	1.16	1.02	0.92	0.90	0.92	0.67	0.81	1.07
	" II.....	0.42	0.67	0.84	2.00	2.24	1.70	1.65	1.64	1.37	1.09	0.77	0.53	0.32	0.29	0.37	0.30	0.80
	" III.....	0.48	0.82	1.01	3.36	3.48	2.25	1.30	1.14	0.60	0.60	0.44	0.38	0.29	0.19	0.16	0.19	0.41
	Rain generally.....	0.58	0.64	0.87	1.72	1.79	1.52	1.42	1.40	1.23	1.08	0.86	0.72	0.62	0.61	0.50	0.55	0.90
Rain in the Summer Half-year.																		
	N.	S.N.E.	S.E.	E.N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.S.W.	W.	W.S.W.	W.	W.N.W.	N.W.	N.N.W.	Calm.	
1853-57.	Class I.....	0.52	0.73	0.99	1.17	1.08	0.95	0.82	0.75	0.84	1.03	1.14	1.39	1.50	1.08	0.89	0.80	1.11
	" II.....	0.69	1.01	1.27	1.30	1.01	1.00	1.03	0.97	1.06	1.06	1.20	1.38	0.95	0.93	0.90	0.69	0.64
	" III.....	0.57	0.70	1.74	1.85	1.86	1.42	0.87	1.21	0.64	0.39	1.08	0.41	0.69	0.57	0.69	0.58	0.63
	Rain generally.....	0.73	0.82	1.20	1.32	1.18	1.04	0.90	0.90	0.89	1.02	1.15	1.20	1.18	0.99	0.86	0.73	0.87
1868-69.	Class I.....	0.95	0.58	0.98	0.90	0.99	0.72	0.71	0.69	0.66	1.97	1.34	1.84	1.41	1.17	1.25	0.90	0.82
	" II.....	0.87	0.79	1.37	1.53	1.16	1.26	0.83	0.87	0.70	0.90	1.37	1.26	0.96	0.82	0.78	0.70	0.83
	" III.....	0.5	1.13	1.44	2.62	1.61	0.98	1.27	0.47	0.56	0.39	1.77	0.90	0.70	0.50	0.39	0.51	0.40
	Rain generally.....	0.89	0.73	1.17	1.35	1.14	0.92	0.83	0.71	0.66	1.01	1.42	1.32	1.16	0.96	0.98	0.78	0.76
1863-69.	Class I.....	0.82	0.65	0.95	1.02	1.03	0.84	0.76	0.71	0.74	1.04	1.23	1.64	1.45	1.13	1.12	0.86	0.96
	" II.....	0.77	0.93	1.31	1.40	1.08	1.12	0.96	0.93	0.90	0.99	1.28	1.26	0.95	0.87	0.83	0.69	0.73
	" III.....	0.82	0.89	1.60	2.28	1.72	1.21	1.05	0.89	0.60	0.94	1.40	0.66	0.69	0.51	0.55	0.52	0.82
	Rain generally.....	0.81	0.78	1.19	1.34	1.16	0.99	0.87	0.81	0.67	1.01	1.27	1.37	1.17	0.97	0.93	0.75	0.82

1853-57.	Rain generally	0.81	0.78	1.19	1.34	1.16	0.99	0.87	0.81	0.67	1.01	1.27	1.37	1.17	0.93	0.73	0.52
----------	----------------------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

Rain in the whole year.

	N.	S.N.E.	N.E.	E.N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.W.	W.S.W.	W.	W.N.W.	N.W.	N.N.W.	Calms.	
Class I.....	0.80	0.73	1.00	1.26	1.14	1.12	0.97	1.02	1.04	1.13	1.11	0.98	1.03	0.82	0.83	1.16	
" II.....	0.57	0.94	1.16	1.09	1.63	1.26	1.34	1.25	1.28	1.12	0.93	0.65	0.56	0.68	0.69	0.54	
" III.....	0.66	0.62	1.40	2.22	2.49	1.91	1.13	1.38	0.72	0.95	0.84	0.29	0.35	0.54	0.45	0.43	
Rain generally	0.70	1.11	1.55	1.50	1.28	1.13	1.15	1.08	1.10	1.01	0.77	0.77	0.74	0.72	0.69	0.92	
1858-62.	Class I.....	0.84	0.56	0.93	1.05	1.15	0.96	1.02	0.85	0.86	1.21	1.13	1.21	1.14	1.20	1.05	0.90
" II.....	0.74	1.13	1.58	1.80	1.60	1.58	1.16	1.19	0.90	1.08	1.14	0.83	0.54	0.47	0.60	0.57	
" III.....	0.81	1.17	1.40	3.09	2.20	1.23	1.27	1.66	0.68	0.95	1.02	0.53	0.47	0.36	0.32	0.49	
Rain generally	0.81	1.06	1.58	1.44	1.20	1.10	0.94	0.84	1.14	1.12	0.99	0.85	0.84	0.80	0.74	0.56	
1858-62.	Class I.....	0.82	0.65	0.96	1.15	1.14	1.03	0.99	0.93	0.94	1.17	1.12	1.09	1.09	1.04	0.95	0.88
" II.....	0.65	0.86	1.15	1.75	1.61	1.41	1.27	1.23	1.11	1.11	1.02	0.73	0.55	0.57	0.63	0.56	
" III.....	0.72	0.87	1.39	2.68	2.33	1.58	1.19	1.06	0.71	0.93	0.92	0.40	0.42	0.44	0.37	0.46	
Rain generally	0.75	1.08	1.57	1.47	1.24	1.11	1.05	0.97	1.12	1.06	0.87	0.81	0.79	0.76	0.71	0.89	

Snow.

	N.	S.N.E.	N.E.	E.N.E.	E.	E.S.E.	S.E.	S.S.E.	S.	S.W.	W.S.W.	W.	W.N.W.	N.W.	N.N.W.	Calms.	
Class I.....	1.20	1.12	1.06	0.47	0.43	0.76	0.64	0.48	0.44	0.44	0.44	0.63	0.35	0.89	0.95	0.78	1.27
" II.....	1.04	1.79	2.22	1.05	0.91	0.97	1.27	0.55	0.59	0.12	0.24	0.25	0.49	0.49	0.49	0.73	0.49
" III.....	1.59	2.69	2.81	2.05	2.57	1.34	0.25	0.83	0.78	0.48	0.47	0.48	1.09	1.58	1.64	1.28	0.25
Snow generally	1.17	1.36	1.43	0.69	0.65	0.83	0.83	0.83	0.83	0.83	0.83	0.83	0.83	0.83	0.83	0.83	0.65
1858-62.	Class I.....	1.29	1.10	0.67	0.54	0.46	0.49	0.52	0.69	0.45	0.55	1.03	2.03	2.01	1.87	1.35	1.24
" II.....	1.04	1.82	3.26	1.53	1.21	0.82	0.60	0.99	0.45	0.41	0.56	0.73	0.97	0.45	0.69	0.80	0.65
" III.....	1.17	1.76	2.78	2.49	2.49	0.59	0.44	0.44	0.00	0.29	0.88	0.73	0.59	0.29	0.73	0.59	0.59
Snow generally	1.22	1.30	1.37	0.85	0.71	0.57	0.54	0.76	0.44	0.51	0.91	1.68	1.71	1.48	1.16	1.12	0.68
1858-62.	Class I.....	1.24	1.11	0.86	0.51	0.45	0.62	0.58	0.58	0.45	0.49	1.12	1.94	1.99	1.71	1.38	1.26
" II.....	1.02	1.81	2.72	1.32	1.06	0.89	0.75	0.51	0.53	0.77	0.81	0.96	0.60	0.70	1.02	0.55	0.72
" III.....	1.44	2.22	2.75	2.22	2.48	1.05	0.26	0.13	0.26	0.52	0.65	0.39	0.65	0.78	0.39	0.55	0.55
Snow generally	1.20	1.33	1.40	0.75	0.68	0.71	0.67	0.61	0.45	0.49	1.00	1.62	1.67	1.39	1.18	1.18	0.66

TORONTO METEOROLOGICAL RESULTS.

TABLE LXXXVII.

Number of days in which rain fell with its depth in inches, for each month of the years 1860, 1861, and 1862.

Number of days.

	Jan.	Feb.	March.	April.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
1860	6	7	5	11	16	14	13	14	14	15	12	3	130
1861	4	4	8	12	12	13	16	15	17	15	14	6	136
1862	5	3	8	10	8	10	15	15	9	19	11	5	118
Means for the 3 years.	5	5	7	11	12	12	15	15	13	16	12	5	128
Means for 23 years...	5	4	6	9	11	12	10	10	11	12	10	5	106

Depth in inches.

1860	0.740	1.330	0.882	1.282	1.815	2.136	4.336	3.405	1.950	1.618	2.560	1.362	23.484
1861	0.685	0.815	2.125	1.619	3.380	2.329	2.635	2.953	3.607	1.933	4.294	0.560	26.995
1862	0.115	0.180	2.560	2.235	1.427	1.007	5.344	3.483	2.344	2.684	2.205	1.945	25.529
Means for the 3 years.	0.513	0.775	1.850	1.712	2.207	1.824	4.105	3.280	2.637	2.098	3.023	1.289	25.319
Means for 21 years...	1.260	1.034	1.553	2.390	3.084	2.961	3.672	3.034	3.788	2.529	3.192	1.637	30.143

TABLE LXXXVIII.

Number of days in which snow fell with its depth in inches, for each month of the years 1860, 1861, and 1862.

Number of days.

	Jan.	Feb.	March.	April.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
1860	16	13	11	5	1	8	21	75
1861	23	17	14	4	1	1	8	8	76
1862	19	17	11	4	2	11	8	72
Means for the 3 years.	19	16	12	4	0	0	0	0	0	1	9	12	74
Means for 23 years ...	12	12	9	3	0	0	0	0	0	2	6	13	58

Depth in inches.

1860	8.7	18.8	2.4	0.3	*	1.9	13.5	45.0
1861	20.6	29.7	7.1	6.9	0.5	*	3.2	6.8	74.8
1862	27.4	23.1	18.5	0.2	0.5	5.3	10.4	85.4
Means for the 3 years.	18.9	23.9	9.3	2.5	0.2	0.2	3.5	10.0	68.6
Means for 20 years...	14.3	18.3	9.3	2.4	0.1	0.8	3.2	14.5	62.8

* Inappreciable.

*

TABLE LXXXIX.

Mean monthly and annual temperatures in each of the twenty-three years, from 1841 to 1863 inclusive, with the average monthly and annual temperatures in the whole period. Also the probable variability of a single year, and of each month in a single year, together with the probable errors of the monthly and annual averages for the whole period.

	Jan.	Feb.	March.	April.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Mean Annual Temperatures.	Differences from Average.
1841	25°.6	22°.4	27°.7	39°.2	50°.5	65°.6	65°.0	64°.4	61°.3	41°.6	35°.0	28°.7	43.92	-0.26
1842	27.0	26.9	35.8	43.1	49.1	55.6	64.7	65.7	55.7	45.1	33.3	24.7	43.96	-0.22
1843	28.7	14.5	21.3	40.9	49.1	58.4	64.5	66.4	59.1	41.8	33.5	30.0	42.35	-1.83
1844	20.2	26.0	31.3	47.5	53.6	59.9	66.0	64.3	58.6	43.3	34.9	28.2	44.48	+0.30
1845	26.5	26.0	35.4	42.1	49.6	61.0	66.2	67.9	66.0	46.4	36.8	21.1	44.58	+0.40
1846	25.7	20.4	33.1	44.0	55.5	63.3	68.0	68.4	63.6	44.6	41.3	27.5	46.36	+2.18
1847	23.3	21.5	26.2	39.2	54.4	58.4	68.0	65.1	55.6	44.0	38.6	30.1	43.70	-0.48
1848	28.7	26.6	28.6	41.3	54.1	62.9	65.5	69.2	54.2	46.3	34.5	29.1	45.08	+0.90
1849	18.5	19.5	33.5	39.0	48.0	63.2	68.4	66.3	58.2	45.3	42.6	26.5	44.00	-0.09
1850	29.7	26.0	29.8	37.9	47.6	64.3	68.9	66.8	56.5	45.4	38.8	21.7	44.45	+0.27
1851	25.5	27.6	32.4	41.3	51.3	59.2	65.0	63.6	60.0	47.4	32.9	21.5	43.98	-0.20
1852	18.4	23.4	27.7	38.2	51.4	60.8	66.8	65.9	57.5	48.0	36.0	31.9	43.84	-0.34
1853	22.9	24.2	30.8	41.9	50.8	65.4	65.5	68.7	58.9	44.5	38.7	25.4	44.80	+0.62
1854	23.5	21.2	30.8	41.1	52.1	64.1	72.4	68.1	61.1	49.5	36.9	21.9	45.23	+1.05
1855	25.9	15.6	28.6	42.5	53.0	59.9	67.9	64.1	59.6	45.4	38.6	26.9	43.98	-0.20
1856	16.0	15.8	23.2	42.3	50.4	62.1	69.8	63.6	57.2	45.4	37.4	22.9	42.18	-2.00
1857	12.7	28.7	28.0	35.4	48.8	50.9	67.7	65.4	58.7	45.5	33.6	31.0	42.75	-1.43
1858	30.0	17.1	28.6	41.5	48.8	66.1	67.8	67.7	59.2	48.8	34.2	27.4	44.76	+0.58
1859	26.4	26.2	36.5	39.6	55.1	58.2	66.8	66.7	55.2	43.0	38.9	17.9	44.21	+0.03
1860	23.3	23.0	34.6	39.6	55.5	63.1	63.8	64.5	55.4	47.3	38.0	24.0	44.34	+0.16
1861	19.8	22.6	27.1	42.1	47.4	61.2	65.3	65.5	59.1	48.8	37.2	31.2	44.24	+0.06
1862	21.7	22.6	28.0	39.6	52.1	60.5	66.6	67.7	59.7	48.7	35.6	28.8	44.37	+0.19
1863	28.0	22.6	26.0	42.1	54.2	60.1	67.5	66.6	55.9	46.0	39.1	27.0	44.59	+0.41
Means 1841-1863	23°.91	22°.78	29°.82	40°.93	51°.41	61°.31	66°.87	66°.20	58°.10	45°.74	36°.80	26°.35	44.18	
Normal temperatures in lat 43° 40' N.	32°.8	34°.7	40°.1	50°.2	55°.1	64°.0	68°.7	68°.5	61°.5	53°.8	43°.2	36°.0	51°.0	
Probable variabilities in a single year.	3.10	2.76	2.65	1.64	1.77	1.92	1.33	1.14	1.56	1.48	1.76	2.59	0.61	
Probable errors of general averages.	0.65	0.58	0.55	0.34	0.37	0.40	0.28	0.24	0.33	0.31	0.37	0.54	0.13	
Means 1841-1852	21°.97	23°.40	30°.23	41°.14	51°.18	61°.05	66°.41	66°.16	58°.02	41°.93	36°.51	26°.75	44.23	

TORONTO METEOROLOGICAL RESULTS.

TABLE XC.

Mean monthly and annual barometric pressures in each of the twenty-three years from 1841 to 1863 inclusive, with the average monthly and annual pressures in the whole period. Also the probable variability in a single year, and of each month in a single year, together with the probable errors of the monthly and annual averages for the whole period.

	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Year.
1841	.6036	29.4918	29.6540	29.6234	29.5452	29.5428	29.6171	29.6977	29.6085	29.6440	29.6078	29.6070	29.6040
1842	.6091	.6477	.6343	.6551	.6888	.6850	.6567	.7100	.6594	.6397	.6094	.6504	.6122
1843	.6061	.5566	.6580	.6084	.6149	.6520	.6203	.6799	.6911	.6425	.6682	.6659	.6120
1844	.6118	.6004	.6524	.7374	.5496	.6002	.5368	.6289	.7307	.6356	.6110	.5486	.6177
1845	.6244	.6748	.5922	.5901	.6355	.5976	.6082	.6320	.6011	.7941	.5084	.6918	.6090
1846	.6172	.6620	.6026	.7017	.5067	.6030	.6847	.6882	.6237	.6960	.6714	.6424	.6284
1847	.5964	.6202	.6756	.6731	.6837	.6619	.6313	.6360	.6099	.6753	.6851	.6567	.6253
1848	.6667	.6084	.6470	.7310	.4961	.6435	.6715	.6412	.5874	.5999	.6555	.6776	.6188
1849	.8046	.7529	.7133	.5814	.6717	.6288	.6826	.6194	.6820	.6038	.6886	.6799	.6070
1850	.6869	.4945	.6004	.6653	.6551	.6409	.6878	.6006	.6233	.5977	.6565	.6756	.6164
1851	.6106	.7562	.6570	.5991	.0288	.6003	.5587	.6691	.7586	.5908	.6347	.6668	.6446
1852	.6756	.6281	.5916	.4147	.6187	.5208	.6104	.6658	.7013	.6613	.5761	.5995	.5886
1853	.7169	.6856	.6551	.5672	.5966	.6155	.6538	.5894	.6419	.6500	.7948	.6001	.6306
1854	.6106	.6979	.5264	.6362	.6047	.5494	.6389	.6464	.7000	.6979	.4419	.5890	.6083
1855	.6433	.6280	.5147	.6522	.6400	.5114	.6007	.6517	.7200	.5588	.6670	.7037	.6255
1856	.6734	.4915	.5610	.5773	.5808	.6464	.5808	.6195	.5999	.7003	.7648	.7130	.6006
1857	.7400	.7393	.5975	.5283	.6339	.4247	.5869	.5930	.7118	.6695	.5265	.6205	.6060
1858	.6791	.6835	.6215	.4970	.5824	.6037	.6038	.6181	.6497	.6837	.6294	.6960	.6273
1859	.6808	.6353	.4143	.6333	.6584	.6176	.6409	.5977	.6684	.6170	.6773	.7109	.6215
1860	.6467	.6356	.5129	.5758	.5646	.4958	.5628	.5812	.6731	.6735	.5253	.6688	.5929
1861	.6555	.5473	.6224	.5623	.5440	.5678	.6491	.0517	.6082	.6215	.5398	.7478	.6015
1862	.7312	.6109	.5054	.7241	.5881	.5622	.5400	.6148	.6828	.6212	.6301	.6795	.6264
1863	.6504	.7954	.6667	.6436	.6156	.5503	.5949	.6440	.7322	.6996	.5584	.6992	.6542
Means 1841-1863.	.6516	.6254	.5947	.5990	.6858	.6659	.6075	.6272	.6018	.6473	.6120	.6600	.6190
Probable varieties in a single year.	0.0418	0.0502	0.0406	0.0520	0.0314	0.0831	0.0292	0.0316	0.0358	0.0374	0.0510	0.0325	0.0125
Probable errors of general averages.	0.0087	0.0117	0.0097	0.0108	0.0066	0.0069	0.0061	0.0066	0.0076	0.0078	0.0106	0.0065	0.0026

with
year,
s for

ear.

.6040
.6122
.6120
.6177
.6099
.6284
.0263
.6188
.6670
.6154
.6446
.5886
.6306
.6083
.0255
.6006
.6060
.6273
.6216
.5929
.6015
.0254
.6542

.6190

0.0125

0 0026

