



IMAGE EVALUATION TEST TARGET (MT-3)



Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

43 26
32 25
36 22
20 18

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques



1982

11
10
9
8
7
6
5
4
3
2
1

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

- Coloured covers/
Couverture de couleur
- Covers damaged/
Couverture endommagée
- Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée
- Cover title missing/
Le titre de couverture manque
- Coloured maps/
Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur
- Bound with other material/
Relié avec d'autres documents
- Tight binding may cause shadows or distortion
along interior margin/
La reliure serrée peut causer de l'ombre ou de la
distortion le long de la marge intérieure
- Blank leaves added during restoration may
appear within the text. Whenever possible, these
have been omitted from filming/
Il se peut que certaines pages blanches ajoutées
lors d'une restauration apparaissent dans le texte,
mais, lorsque cela était possible, ces pages n'ont
pas été filmées.

Additional comments:/
Commentaires supplémentaires: Pagination continued from Vol III. Irregular pagination: 3137 - 3306, 3311-
[3314], 3307 - 3310 p.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured pages/
Pages de couleur
- Pages damaged/
Pages endommagées
- Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- Pages detached/
Pages détachées
- Showthrough/
Transparence
- Quality of print varies/
Qualité inégale de l'impression
- Includes supplementary material/
Comprend du matériel supplémentaire
- Only edition available/
Seule édition disponible
- Pages wholly or partially obscured by errata
slips, tissues, etc., have been refilmed to
ensure the best possible image/
Les pages totalement ou partiellement
obscurcies par un feuillett d'errata, une pelure,
etc., ont été filmées à nouveau de façon à
obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
12X	16X	20X	24X	28X	32X

The copy filmed here has been reproduced thanks to the generosity of:

Library.
Department of Fisheries and Oceans

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol → (meaning "CONTINUED"), or the symbol ▽ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

L'exemplaire filmé fut reproduit grâce à la générosité de:



Bibliothèque.
Ministère des pêches et océans

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole → signifie "A SUIVRE", le symbole ▽ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.



X
UNIT

DESCRIPTION
THE W

SMITHSONIAN INSTITUTION.
UNITED STATES NATIONAL MUSEUM.

BULLETIN

OF THE

UNITED STATES NATIONAL MUSEUM.

No. 47.

THE FISHES OF NORTH AND MIDDLE AMERICA:

DESCRIPTIVE CATALOGUE OF THE SPECIES OF FISH-LIKE VERTEBRATES FOUND IN
THE WATERS OF NORTH AMERICA, NORTH OF THE IsthMUS OF PANAMA.

BY

DAVID STARR JORDAN, Ph. D.,

PRESIDENT OF THE LELAND STANFORD JUNIOR UNIVERSITY AND OF THE
CALIFORNIA ACADEMY OF SCIENCES,


AND

BARTON WARREN EVERMANN, Ph. D.,

ICHTHYOLOGIST OF THE UNITED STATES FISH COMMISSION.

PART IV.

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1900.



SMITHSONIAN INSTITUTION.

UNITED STATES NATIONAL MUSEUM.

THE FISHES

OF

NORTH AND MIDDLE AMERICA:

A DESCRIPTIVE CATALOGUE

OF THE

SPECIES OF FISH-LIKE VERTEBRATES FOUND IN THE
WATERS OF NORTH AMERICA, NORTH OF
THE IsthMUS OF PANAMA.

BY

DAVID STARR JORDAN, PH. D.,

PRESIDENT OF THE LELAND STANFORD JUNIOR UNIVERSITY AND OF THE
CALIFORNIA ACADEMY OF SCIENCES,

AND

BARTON WARREN EVERMANN, PH. D.,

ICHTHYOLOGIST OF THE UNITED STATES FISH COMMISSION.

PART IV.

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1900.

② L
6-8
J 82

P R E F A C E.

This volume is fourth and last of a descriptive catalogue of the fishes and fish-like vertebrates of North and Middle America. Part I, *Branchiosomatidae* to *Priacanthidae* inclusive (pp. 1 to 1240), was published October 3, 1896; Part II, *Lutianidae* to *Cephalicanthidae* inclusive (pp. 1241 to 2183), was published October 3, 1898; Part III, *Callionymidae* to *Ogcoccephalidae* inclusive, including also "Addenda," an "Artificial Key to the Families of the True Fishes," a "Glossary of Technical Terms," and an "Index" complete for Parts I, II, and III (pp. 2183*a* to 3136), was published November 28, 1898; and Part IV, the present volume, appears on June 26, 1900.

Since the publication of Part III, investigations by Dr. Jordan, in Mexico, and by Dr. Evermann, in Puerto Rico, have added greatly to our knowledge of the fishes of those regions, and have made it desirable to incorporate in the present volume still further addenda.

Recent studies by Dr. Günther, of certain Linnean types, show that some changes in nomenclature are necessary, which are also included.

DAVID STARR JORDAN.

BARTON WARREN EVERMANN.

LELAND STANFORD JUNIOR UNIVERSITY,
PALO ALTO, SANTA CLARA COUNTY, CALIFORNIA.

March 15, 1900.

(iii)

10024
88
37
Y.

A SYSTEMATIC ARRANGEMENT OF THE FISHES OF NORTH AND MIDDLE AMERICA.

In these pages the authors exhibit, as clearly as the method will permit, their present views as to the genetic relations of the fishes and fish-like vertebrates of North and Middle America. This list is, in fact, a Table of Contents complete for the four volumes and corrected to include the Addenda (pp. 2745-2873 of Part III and pp. 3137-3197 of Part IV). From this Systematic Arrangement it is seen that the fish fauna of North and Middle America, as now understood by the present authors, embraces 3 classes, 30 orders, 225 families, 1113 genera, 325 subgenera, 3263 species, and 133 subspecies.

	Page.
CLASS LEPTOCARDII	2
ORDER AMPHOXI	2
<i>Family Branchiostomidae</i>	2
Genus <i>Branchiostoma</i> , Costa	3
<i>lanceolatum</i> (Pallas)	3
<i>caribaeum</i> , Sundevall	3
<i>californiense</i> , Gill	4
Genus <i>Asymmetron</i> , Andrews	4
<i>lucayanum</i> , Andrews	4
CLASS MARSIPOBANCHII	4
ORDER HYPEROTRETI	5
<i>Family Heptatremidae</i>	5
Genus <i>Polistotrema</i> , Gill	6
<i>stouti</i> (Lockington)	6
<i>Family Myxinidae</i>	7
Genus <i>Myxine</i> , Linnaeus	7
<i>glutinosa</i> , Linnaeus	7
ORDER HYPEROARTHRI	8
<i>Family Petromyzonidae</i>	8
Genus <i>Bathymyzon</i> , Gill	9
<i>bairdii</i> (Gill)	9
Genus <i>Petromyzon</i> (Arvedi) Linnaeus	9
<i>marinus</i> , Linnaeus	10
<i>unicolor</i> (De Kay)	10
Genus <i>Ichthyomyzon</i> , Girard	10
<i>concolor</i> (Kirtland)	11
<i>castaneus</i> , Girard	11
Genus <i>Entosphenus</i> , Gill	11
<i>tridentatus</i> (Girardner)	12
<i>cantschatius</i> (Tilesius)	2745
Genus <i>Lampetra</i> , Gray	12
<i>aurea</i> (Bean)	13
<i>spadicea</i> , Bean	13
<i>eibaria</i> (Girard)	13
<i>wilderi</i> , Gage	13
CLASS PISCES	14
SUBCLASS SELACHII	15
ORDER DIPLOSPONDYLI	16
<i>Family Chlamydoselachidae</i>	16
Genus <i>Chlamydoselachus</i> , Garman	16
<i>anguineus</i> , Garman	16

CLASS PISCES—Continued.	Page.
SUBCLASS SELACHII—Continued.	
ORDER DUPLOSPONDYLI—Continued.	
<i>Family Hexanchidae</i>	17
Genus Notorhynchus, Ayres.....	17
maculatus, Ayres	17
Genus Hexanchus, Rafinesque.....	18
coronus, Jordan & Gilbert.....	18
griseus (Gmelin).....	19
ORDER ASTEROSPONDYLI	19
SUBORDER PROARTHRII	
<i>Family Heterodontidae</i>	19
Genus Gyroplectodus, Gill	20
francisci (Girard)	20
quoyi (Fréminville)	21
SUBORDER GALEI	21
<i>Family Scylliorhinidae</i>	22
Genus Scylliorhinus, Blainville	22
profundorum, Goode & Bean	22
Genus Catalus, Smith	23
Subgenus Catalus.....	24
xanthurus, Gilbert	24
brunneus, Gilbert	24
cephalus, Gilbert	24
retifer (Garman)	25
Subgenus Cephaloscyllium, Gill.....	25
uter, Jordan & Gilbert	25
<i>Family Ginglymostomidae</i>	25
Genus Ginglymostoma, Müller & Henle	26
cirratum (Gmelin)	26
<i>Family Pseudotriakidae</i>	26
Genus Pseudotriakis, Capello	27
microdon, Capello	27
<i>Family Galeidae</i>	27
Genus Mustelus, Cuvier	28
lumulatus, Jordan & Gilbert	28
canis (Mitchill)	29
Genus Galeus, Rafinesque	29
dorsalis (Gill)	30
californicus (Gill)	30
Genus Rhinotriacus, Gill	30
henlei, Gill	31
Genus Triakis, Müller & Henle	31
semifasciatum, Girard	31
Genus Galeorhinus, Blainville	31
zyopterus, Jordan & Gilbert	32
Genus Galeocerdo, Müller & Henle	32
tigrinus, Müller & Henle	32
Genus Prionace, Cantor	33
glauca (Linnaeus)	33
Genus Carcharhinus, Blainville	33
Subgenus Platypodon, Gill	35
obseurus (Le Sueur)	35
platyrhynchos (Gilbert)	36
falciformis (Bibron)	36
aeronotus (Poey)	36
perezi (Poey)	36

	Page.
CLASS PISCES—Continued.	
SUBCLASS SELACHII—Continued.	
ORDER ASTEROSPONDYLI—Continued.	
SUBORDER GALEI—Continued.	
<i>Family Cetorhinidae</i> —Continued.	
Genus <i>Carcharhinus</i> , Blainville—Continued.	
Subgenus <i>Platypodon</i> , Gill—Continued.	
<i>remotus</i> (Vauclusienus)	37
<i>henlei</i> (Valenciennes)	37
<i>cerdale</i> , Gilbert	2746
Subgenus <i>Carcharhinus</i>	37
<i>millberti</i> (Müller & Henle).....	37
<i>lamelia</i> (Jordan & Gilbert)	37
<i>lamia</i> (Rafinesque)	38
<i>platyodon</i> (Poey)	39
<i>fronto</i> (Jordan & Gilbert)	39
<i>nicaraguensis</i> (Gill & Bransford)	39
Subgenus <i>Isohypophodon</i> , Gill	40
<i>athalurus</i> (Jordan & Gilbert)	40
<i>limbatus</i> (Müller & Henle)	40
<i>oxyrhynchus</i> (Müller & Henle)	40
<i>velox</i> , Gilbert	2747
Genus <i>Hypoprion</i> , Müller & Henle	41
<i>brevirostris</i> , Poey	41
<i>signatus</i> , Poey	41
Genus <i>Aprionodon</i> , Gill	42
<i>isodon</i> (Müller & Henle)	42
Genus <i>Scyliodon</i> , Müller & Henle	42
<i>longurio</i> (Jordan & Gilbert)	42
<i>terra-novae</i> (Richardson)	43
<i>Family Sphyrnidae</i>	43
Genus <i>Sphyrna</i> , Rafinesque	43
Subgenus <i>Reniceps</i> , Gill	44
<i>tiburo</i> (Linnaeus)	44
Subgenus <i>Platysqualus</i> , Swamson	44
<i>tudes</i> (Cuvier)	44
Subgenus <i>Sphyrna</i>	45
<i>zygana</i> (Linnaeus)	45
<i>Family Alopiidae</i>	45
Genus <i>Alopias</i> , Rafinesque	45
<i>vulpes</i> (Gmelin)	45
<i>Family Carchariidae</i>	46
Genus <i>Carcharias</i> , Rafinesque	46
Subgenus <i>Engomphodus</i> , Gill	46
<i>litoralis</i> (Mitchill)	46
<i>Family Lamnidae</i>	47
Genus <i>Isurus</i> , Rafinesque	47
Subgenus <i>Isuropsis</i> , Gill	48
<i>dekeyi</i> (Gill)	48
Subgenus <i>Isurus</i>	48
<i>oxyrhynchus</i> , Rafinesque	48
Genus <i>Lamna</i> , Cuvier	49
<i>cornubica</i> (Gmelin)	49
Genus <i>Carcharodon</i> , Smith	50
<i>carcharias</i> (Linnaeus)	50
<i>Family Cetorhinidae</i>	50
Genus <i>Cetorhinus</i> , Blainville	51
<i>maximus</i> (Gunner)	51

	Page.
CLASS PISCES—Continued.	
SUBCLASS SELACHII—Continued.	
ORDER ASTEROSPONDYLI—Continued.	
SUBORDER GALEI—Continued.	
<i>Family Rhinodontidae</i>	52
Genus <i>Miceristedus</i> Gill.....	52
<i>punctatus</i> , Gill.....	52
<i>ORDER CYCLOSPONDYLI</i>	52
SUBORDER CYCLOSPONDYLI	53
<i>T. nily Squalid'e</i>	53
Genus <i>Squalus</i> (Artedi) Linnaeus.....	53
<i>acanthias</i> , Linnaeus	54
<i>sucklini</i> (Girard)	54
Genus <i>Centroscymnus</i> , Bocage & Capello.....	54
<i>celeopis</i> , Bocage & Capello	55
Genus <i>Etmopterus</i> , Rafinesque	55
<i>pusillus</i> (Lowe)	55
Genus <i>Centroscyllium</i> , Müller & Henle.....	56
<i>fabricii</i> (Reinhardt)	56
<i>Family Dalatiidae</i>	56
Genus <i>Somniosus</i> , Le Sueur	56
<i>microcephalus</i> (Bloch)	57
<i>Family Echinorhinidae</i>	57
Genus <i>Echinorhinus</i> , Blainville	57
<i>spinosus</i> (Gmelin)	58
SUBORDER TECTOSPONDYLI.....	58
<i>Family Squatinidae</i>	58
Genus <i>Squatina</i> , Duméril	58
<i>squatina</i> (Linnaeus)	58
<i>ORDER BATOIDÆ</i>	59
SUBORDER SARCURA.....	60
<i>Family Pristidae</i>	60
Genus <i>Pristis</i> , Latham	60
<i>zephyreus</i> , Jordan & Starks	60; 2749
<i>pectinatus</i> , Latham	60
<i>Family Rhinobatidae</i>	61
Genus <i>Rhinobatos</i> , Blech & Schneider	61
<i>lentiginosus</i> , Garman	62
<i>stellio</i> , Jordan & Rutter	2750
<i>glaucostigma</i> , Jordan & Gilbert	62
<i>leucorrhynchus</i> , Günther	62
<i>productus</i> , Ayres	63
<i>percellens</i> (Walbaum)	63
<i>spinifer</i> , Günther	63
<i>planiceps</i> , Garman	64
Genus <i>Zapteryx</i> , Jordan & Gilbert	64
<i>exasperatus</i> (Jordan & Gilbert)	64
<i>xyster</i> , Jordan & Evermann	65
Genus <i>Platyrrhinoidis</i> , Garman	65
<i>triseriatus</i> (Jordan & Gilbert)	65
<i>Family Rajidae</i>	66
Genus <i>Raja</i> (Arteilli) Linnaeus	66
<i>erinaceus</i> , Mitchell	68
<i>ocellata</i> , Mitchell	68
<i>fylle</i> , Lütken	69
<i>radiata</i> , Donovan	69
<i>plutonia</i> , Garman	69
<i>ackleyi</i> , Garman	70
<i>ornata</i> , Garman	70

	Page.
CLASS PISCES—Continued.	
SUBCLASS SELACHII—Continued.	
ORDER BATOIDEI—Continued.	
SUBORDER SARCURA—Continued.	
<i>Family Rajidae—Continued.</i>	
Genus <i>Raja</i> (Artedi) Linnaeus—Continued.	
<i>eglæteria</i> , Bosc	71
<i>senta</i> , Garman	71
<i>levis</i> (Mitchill)	71
<i>rhina</i> , Jordan & Gilbert	72
<i>binoculata</i> , Girard	72
<i>inornata</i> , Jordan & Gilbert	73
<i>equatorialis</i> , Jordan & Bollman	74
<i>rossispinis</i> , Gill & Townsend	2751
<i>interrupta</i> , Gill & Townsend	2751
<i>parmifera</i> , Bean	74
<i>stellulata</i> , Jordan & Gilbert	75
<i>aleutica</i> , Gilbert	75
<i>trachura</i> , Gilbert	75
<i>abyssicola</i> , Gilbert	76
<i>Family Narcobatidae</i>	76
Genus <i>Tetronarce</i> , Gill	77
<i>œcidentalis</i> (Storer)	77
<i>californica</i> (Ay)	77
Genus <i>Narcine</i> , Henle	78
<i>brasiliensis</i> (Ölfers)	78
<i>entemedor</i> , Jordan & Starks	2752
Genus <i>Discopyge</i> , Tschudi	78
<i>omnata</i> , Jordan & Gilbert	78
<i>SUBORDER MASTICURA</i>	79
<i>Family Dasyatidae</i>	79
Genus <i>Urolophus</i> , Müller & Henle	79
<i>halleri</i> , Cooper	80
<i>nebulosus</i> , Garman	80
<i>umbrifer</i> , Jordan & Starks	2752
<i>jamaicensis</i> (Cuvier)	81
<i>mundus</i> (Gill)	81; 2752
<i>goodri</i> , Jordan & Bollman	81
<i>aspidurus</i> , Jordan & Gilbert	81
<i>rogersi</i> , Jordan & Starks	2752
Genus <i>Dasyatis</i> , Rafinesque	82
Subgenus <i>Hepterygon</i> , Müller & Henle	83
<i>centrura</i> (Mitchill)	83
Subgenus <i>Dasyatis</i>	83
<i>hastata</i> (De Kay)	83
<i>gymnura</i> (Müller)	84
<i>sabina</i> (Le Sueur)	84
<i>longa</i> , Garman	85
<i>dipterura</i> , Jordan & Gilbert	85
<i>say</i> (Le Sueur)	86
Genus <i>Pteroplatea</i> , Müller & Henle	86
<i>maelura</i> (Le Sueur)	86
<i>crebripunctata</i> , Peters	87
<i>marmorata</i> , Cooper	87
<i>rava</i> , Jordan & Starks	2754
<i>Family Myliobatidae</i>	87
Genus <i>Aetobatus</i> , Blainville	88
<i>narinari</i> (Enphrasen)	88

	Page.
CLASS PISCES—Continued.	
SUBCLASS SELACHII—Continued.	
ORDER BATOIDEI—Continued.	
SUPERORDER MASTICURA—Continued.	
<i>Family Myliobatidae—Continued.</i>	
Genus <i>Myliobatis</i> , Duméril	89
Subgenus <i>Myliobatis</i>	89
<i>fremivillei</i> , Le Sueur	89
Subgenus <i>Holorhinus</i> , Gill	89
<i>californicus</i> , Gill	89
<i>asperrimus</i> , Gilbert	2754
<i>goodei</i> , Garman	2755
Genus <i>Rhinoptera</i> , Kuhl.....	90
Subgenus <i>Rhinoptera</i>	90
<i>bonasus</i> (Mitchill)	90
<i>steindachneri</i> , Evermann & Jenkins	91
Subgenus <i>Micromesistius</i> , Gill	91
<i>ensenatus</i> , Rosa Smith	91
<i>Family Aodontidae</i>	91; 2756
Genus <i>Aodon</i> , Lacépède	91
<i>hypostomus</i> (Bancroft)	92
Genus <i>Ceratobatis</i> , Boulenger	2756
<i>robertsi</i> , Boulenger	2756
Genus <i>Manta</i> , Bancroft	92
<i>birostris</i> (Walbaum)	92
SUBCLASS HOLOCEPHALI.....	93
ORDER CHIMAEROIDEL.....	93
<i>Family Chimaeridae</i>	93
Genus <i>Chimæra</i> , Linnaeus	94
<i>monstrosa</i> , Linnaeus	94
<i>affinis</i> , Capello	95
Genus <i>Hydrolagus</i> , Gill	95
<i>colliei</i> (Lay & Bennett)	95
Genus <i>Harriotta</i> , Goode & Bean	96
<i>raleighana</i> , Goode & Bean	96
SUBCLASS TELEOSOMI.....	97
Series <i>Ganoidei</i>	100
Chondroganoidea.....	100
ORDER SELACHOSTOMI.....	100
<i>Family Polyodontidae</i>	101
Genus <i>Polyodon</i> , Lacépède	101
<i>spathula</i> (Walbaum)	101
ORDER CHONDROSTEI.....	102
<i>Family Acipenseridae</i>	102
Genus <i>Acipenser</i> , Linnaeus	103
<i>transmontanus</i> , Richardson	104
<i>mediostriatus</i> , Ayres	104
<i>sturio</i> , Linnaeus	105
<i>ruberundus</i> , Le Sueur	106
<i>brevirostrum</i> , Le Sueur	106
Genus <i>Scaphirhynchus</i> , Heckel	107
<i>platorynchus</i> (Rafinesque)	107
Holostei	107
ORDER RHOMBOGANOIDÆ.....	108
<i>Family Lepisosteidae</i>	108
Genus <i>Lepisosteus</i> , Lacépède	109
Subgenus <i>Lepisosteus</i>	109
<i>osseus</i> (Linnaeus)	109

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER RHOMBOGANOIDEA—Continued.	
Family <i>Lepisosteidae</i> —Continued.	
Genus <i>Lepisosteus</i> , Lacépède—Continued.	
Subgenus <i>Cylindrostenus</i> , Rafinesque.....	110
<i>platostomus</i> , Rafinesque	110
Subgenus <i>Atractosteus</i> , Rafinesque.....	111
<i>tristechus</i> (Bloch & Schneider)	111
<i>tropius</i> (Gill)	111
ORDER CYCLOGANOIDEA.....	111
Family <i>Amiidae</i>	112
Genus <i>Amia</i> , Linnaeus	112
<i>calva</i> , Linnaeus.....	113
Series Teleostei	113
Ostariophysai	114
ORDER NEMATOGNATHI.....	114
Family <i>Siluridae</i>	115
Genus <i>Felichthys</i> , Swainson	116
<i>panamensis</i> (Gill)	117
<i>bägre</i> (Linnaeus)	117
<i>plumimaculatus</i> (Steindachner).....	117
<i>eydouxii</i> (Cuvier & Valenciennes)	118
<i>filamentosus</i> , Swainson	118
<i>felis</i> (Linnaeus).....	118; 3197
<i>bahiensis</i> (Castelnau)	118
Genus <i>Galeichthys</i> , Cuvier & Valenciennes.....	122; 2770
Subgenus <i>Galeichthys</i>	2771
<i>lentiginosus</i> (Eigenmann & Eigenmann)	122; 2771
<i>peruanus</i> , Lütken	122; 2771
Subgenus <i>Hexanematicthys</i> , Bleeker	128; 2772
<i>milberti</i> (Cuvier & Valenciennes).....	128; 3197
<i>seemannii</i> (Günther)	128; 2772
<i>gilberti</i> , Jordan & Williams.....	2773
<i>jordani</i> (Eigenmann & Eigenmann)	128; 2774
<i>azureus</i> , Jordan & Williams	2775
<i>cerulescens</i> (Günther)	129; 2776
<i>xenuchen</i> , Gilbert.....	2777
<i>guatemalensis</i> (Günther).....	129; 2778
<i>surinamensis</i> (Bleeker).....	129; 2780
<i>dasycephalus</i> (Günther)	130; 2780
<i>longicephalus</i> (Eigenmann & Eigenmann)	130; 2781
<i>rugispinis</i> (Cuvier & Valenciennes)	130; 2781
<i>labiatus</i> (Boulenger).....	3137
<i>phrygiatus</i> (Cuvier & Valenciennes)	130; 2782
Genus <i>Sciaideichthys</i> , Bleeker.....	122
<i>etroscheli</i> (Gill)	122
<i>emphysetus</i> (Müller & Troschel).....	122
<i>temminckianus</i> (Cuvier & Valenciennes).....	122
<i>flavescens</i> (Cuvier & Valenciennes)	123
<i>mesops</i> (Cuvier & Valenciennes)	123
<i>proops</i> (Cuvier & Valenciennes)	123
<i>passany</i> (Cuvier & Valenciennes)	124
<i>albicans</i> (Cuvier & Valenciennes)	124
Genus <i>Aspistor</i> , Jordan & Evermann.....	2763
<i>luniseutis</i> (Cuvier & Valenciennes)	125; 2763
Genus <i>Selenaspis</i> , Bleeker	124
<i>herzbergii</i> (Bloch).....	124

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER NEMATOGNATHI—Continued.	
Family Siluridae—Continued.	
Genus <i>Selenaspis</i> , Bleeker—Continued.	
<i>dowi</i> (Gill)	125
<i>parkeri</i> (Traill).....	125; 2764
Genus <i>Netuma</i> , Bleeker.....	126; 2764
Subgenus <i>Notarius</i> , Gill	2764
<i>grandicassis</i> (Cuvier & Valenciennes).....	126; 2764
<i>stricticassis</i> (Cuvier & Valenciennes)	126; 2765
Subgenus <i>Netuma</i>	2765
<i>dubia</i> (Bleeker)	126; 2765
<i>kessleri</i> (Steindachner).....	127; 2765
<i>insculpta</i> (Jordan & Gilbert).....	127; 2765
<i>planiceps</i> (Steindachner).....	127; 2766
<i>platypteron</i> (Günther).....	127; 2767
<i>oscula</i> (Jordan & Gilbert).....	127; 2768
<i>elattura</i> (Jordan & Gilbert)	128; 2769
<i>insularum</i> , Flora Hartley Greene.....	2770
Genus <i>Tachysurus</i> , Lacépède.....	131; 2782
<i>nuchalis</i> (Günther).....	131; 2782
<i>fissus</i> (Cuvier & Valenciennes)	131; 2782
<i>spixii</i> (Agassiz)	131; 2783
<i>melanopus</i> (Günther).....	132; 2784
<i>festa</i> , Boulenger	3138
<i>liropus</i> , Susan B. Bristol	2784
<i>cuneolane</i> , Gilbert	2785
<i>furthii</i> (Steindachner).....	132; 2787
<i>variolosus</i> (Cuvier & Valenciennes)	132; 2788
<i>multiradiatus</i> (Günther)	132; 2788
Genus <i>Cathorops</i> , Jordan & Gilbert	133; 2788
<i>hypophthalmus</i> (Steindachner)	133; 2788
<i>gulosus</i> (Eigenmann & Eigenmann)	133
Genus <i>Ictalurus</i> , Rafinesque.....	133; 2788
<i>furcatus</i> (Le Sueur)	134; 2788
<i>anguilla</i> , Evermann & Kendall	2788
<i>punctatus</i> (Rafinesque)	134; 2788
<i>meridionalis</i> (Günther).....	135
Genus <i>Ictalurus</i> , Jordan & Snyder.....	3138
<i>balsanus</i> , Jordan & Snyder	3139
Genus <i>Ameiurus</i> , Rafinesque.....	135
Subgenus <i>Haustor</i> , Jordan & Evermann	137; 3197
<i>lacustris</i> (Walbaum)	137
<i>lucus</i> (Girard)	137
<i>eatus</i> (Linnæus)	138
<i>pricei</i> (Rutter)	2790
<i>dugesii</i> Bean	138; 2789
<i>okeechobensis</i> (Heilprin)	138
Subgenus <i>Ameiurus</i>	139
<i>erebennus</i> , Jordan	139
<i>natalis</i> (Le Sueur)	139
<i>vulgaris</i> (Thompson).....	140
<i>nebulosus</i> (Le Sueur)	140
<i>catulus</i> (Girard)	141
<i>marmoratus</i> (Holbrook).....	141
<i>melas</i> (Rafinesque)	141
<i>platycephalus</i> (Girard)	142

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER NEMATOGNATHI—Continued.	
<i>Family Siluridae</i> —Continued.	
Genus <i>Ameiurus</i> , Rafinesque—Continued.	
Subgenus <i>Gronias</i> , Cope	142
<i>nlgrilabris</i> (Cope)	142
Genus <i>Leptops</i> , Rafinesque	142
<i>olivaris</i> (Rafinesque).....	143
Genus <i>Noturus</i> , Rafinesque.....	143
<i>flavus</i> , Rafinesque.....	144
Genus <i>Schilbeodes</i> , Bleeker.....	144
Subgenus <i>Schilbeodes</i>	146
<i>gyrinus</i> (Mitchill).....	146
<i>leptacanthus</i> (Jordan)	146
Subgenus <i>Rabida</i> , Jordan & Evermann	146
<i>nocturnus</i> (Jordan & Gilbert).....	146
<i>tenebris</i> (Gilbert & Swain)	147
<i>exillis</i> (Nelson)	147
<i>insignis</i> (Richardson)	147
<i>gillerti</i> (Jordan & Evermann)	148
<i>eleutherns</i> (Jordan)	148
<i>miurus</i> (Jordan)	148
<i>furlosus</i> (Jordan & Meek)	149
Genus <i>Rhamdia</i> , Bleeker.....	149
Subgenus <i>Rhamdia</i>	150
<i>wagneri</i> (Günther)	150
Subgenus <i>Rhamdella</i> , Eigenmann & Eigenmann	151
<i>baronis-mulleri</i> (Troschel)	151
<i>motaguensis</i> (Günther)	151
<i>brachyptera</i> (Cope).....	151
<i>salvini</i> (Günther)	152
<i>hypselurus</i> (Günther)	152
<i>laticauda</i> (Heckel)	152
<i>godmani</i> (Günther)	152
<i>guatemalensis</i> (Günther)	152
<i>nicaraguensis</i> (Günther)	152
<i>microptera</i> (Günther)	153
<i>managuensis</i> (Günther)	153
<i>polycalus</i> (Günther)	153
<i>petenensis</i> (Günther)	153
<i>parryi</i> (Eigenmann)	153
Genus <i>Pimelodella</i> , Eigenmann & Eigenmann	153
<i>chagresi</i> (Steindachner)	154
<i>modesta</i> (Günther)	154
Genus <i>Pimelodus</i> , Lacépède	154
<i>maeulatus</i> , Lacépède	155
<i>Family Loricariidae</i>	155
Genus <i>Loricaria</i> , Linnaeus	156
Subgenus <i>Hemiodon</i> , Kner	157
<i>panamensis</i> , Eigenmann & Eigenmann	157
Subgenus <i>Sturisoma</i> , Swainson	157
<i>rostrata</i> , Spix	157
Subgenus <i>Rineloricaria</i> , Bleeker	158
<i>lima</i> , Kner	158
<i>bransfordi</i> , Gill	158
Subgenus <i>Parahemiodon</i> , Bleeker.....	153
<i>uracantha</i> , Kner & Steindachner	158

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER NEMATOGNATHI—Continued.	
<i>Family Loricariidae</i> —Continued.	
Genus <i>Loricaria</i> , Linnaeus—Continued.	
Subgenus <i>Loricaria</i>	159
variegata, Steindachner	159
Genus <i>Hemiancistrus</i> , Bleeker	159
guacharote (Cuvier & Valenciennes)	159
aspidelepis (Günther)	159
Genus <i>Clariostomus</i> , Kner	160
fischeri, Steindachner	160
Genus <i>Ancistrus</i> , Kner	160
chagresi, Eigenmann & Eigenmann	160
ORDER PLECTOSPONDYLI	160
SUBORDER EVENTOGNATHI	191
<i>Family Catostomidae</i>	191
Genus <i>Ictiobus</i> , Rafinesque	163
Subgenus <i>Sclerognathus</i> , Cuvier & Valenciennes	163
cyprinella (Cuvier & Valenciennes)	163
Subgenus <i>Ictiobus</i>	164
urus (Agassiz)	164
meridionalis (Günther)	164
bubalus (Rafinesque)	164
Genus <i>Carpoides</i> , Rafinesque	165
carpio (Rafinesque)	166
diformis, Cope	166
thompsoni, Agassiz	167
velifer (Rafinesque)	167
cyprinus (Le Sueur)	168
Genus <i>Cyclopterus</i> , Rafinesque	168
eloungatus (Le Sueur)	168
Genus <i>Pantosteus</i> , Cope	169
arizone, Gilbert	170
generosus (Girard)	170
plebeius (Baird & Girard)	170
delphinus (Cope)	171
guzmaniensis (Girard)	171
jordani, Evermann	171
araopis (Jordan)	172
clarki (Baird & Girard)	172
Genus <i>Catostomus</i> , Le Sueur	173
Subgenus <i>Catostomus</i>	174
latipinnis, Baird & Girard	174; 2790
discobolus, Cope	175; 2791
griseus (Girard)	175
retropinnis, Jordan	2791
pocatello, Gilbert & Evermann	175
catostomus (Forster)	176
rimiculus, Gilbert & Snyder	2792
snyderi, Gilbert	2792
tahoensis, Gill & Jordan	177
Subgenus <i>Decactylus</i> , Rafinesque	177
occidentalis, Ayres	178; 2792
tsiltcoosensis, Evermann & Meek	2793
bernardini, Girard	178
macrochellus, Girard	178
commersonii, Lacépède	178
ardens, Jordan & Gilbert	179

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOSPONDYLI—Continued.	
SUBORDER EVENTOGNATHI—Continued.	
<i>Family Catostomidae</i> —Continued.	
Genus <i>Catostomus</i> , Le Sueur—Continued.	
Subgenus <i>Decactylus</i> , Rafinesque—Continued.	
<i>gila</i> , Kirsch	180
<i>insignis</i> , Baird & Girard	180
Subgenus <i>Hypentelium</i> , Rafinesque.....	181
<i>nigricans</i> , Le Sueur	181
<i>rhothoecus</i> , Thoburn	181
Genus <i>Deltistes</i> , Seale.....	2794
<i>luxatus</i> (Cope)	183; 2794
Genus <i>Chasmistes</i>	182
<i>fecundus</i> (Cope & Yarrow).....	180; 2794
<i>Ilorus</i> , Jordan	183
<i>cujns</i> , Cope.....	183
<i>brevirostris</i> , Cope	183
<i>stomias</i> , Gilbert	2794
<i>copel</i> , Evermann & Meek	2795
Genus <i>Xyrauchen</i> , Eigenmann & Kirsch.....	184
<i>cypho</i> (Lockington)	184
<i>incompahgno</i> , Jordan & Evermann	184
Genus <i>Erimyzon</i> , Jordan	185
<i>sucetta</i> (Lacépède)	185
<i>oblongus</i> (Mitchill)	186
Genus <i>Minytrema</i> , Jordan	186
<i>melanops</i> (Rafinesque)	187
Genus <i>Moxostoma</i> , Rafinesque	187
<i>papillosum</i> (Cope)	189
<i>anisurum</i> (Rafinesque)	190
<i>collapsum</i> (Cope)	190
<i>bucco</i> (Cope)	190
<i>pidiense</i> (Cope)	191
<i>coregonus</i> (Cope)	191
<i>album</i> (Cope)	191
<i>thalassinum</i> (Cope)	191
<i>congestum</i> (Baird & Girard)	192
<i>austrinum</i> , Bean	192
<i>aureolum</i> (Le Sueur)	192
<i>robustum</i> (Cope)	193
<i>macrolepidotum</i> (Le Sueur)	193
<i>crassilabre</i> (Cope)	194
<i>lesueuri</i> (Richardson)	194
<i>breviceps</i> (Cope)	196
<i>conus</i> (Cope)	196
<i>peccilurum</i> (Jordan)	196
<i>rupisartes</i> , Jordan & Jenkins	196
<i>cervinum</i> (Cope)	197
Genus <i>Placopbarynx</i> , Cope	197
<i>duquesnii</i> (Le Sueur)	198
Genus <i>Lagochila</i> , Jordan & Brayton	198
<i>lacera</i> , Jordan & Brayton	199
<i>Family Cyprinidae</i>	199
Genus <i>Campostoma</i> , Agassiz	204
<i>ornatum</i> , Girard	205; 2796
<i>anomalum</i> (Rafinesque)	205
<i>formosulum</i> , Girard	206

	Page.
CLASS PISCES--Continued.	
SUBCLASS TELEOSTOMI--Continued.	
ORDER PLECTOSPONDYLI--Continued.	
SUBORDER EVENTOGNATHI--Continued.	
<i>Family Cyprinidae</i> --Continued.	
Genus <i>Orthodon</i> , Girard	206
<i>microlepidotus</i> (Ayres)	207
Genus <i>Oxygeneum</i> , Forbes	207
<i>pulverulentum</i> , Forbes	207
Genus <i>Acrocheilus</i> , Agassiz	207
<i>alutaceus</i> , Agassiz & Pickering	208
Genus <i>Lavinia</i> , Girard	208
<i>exiliacula</i> , Baird & Girard	209
Genus <i>Chrosomus</i> , Rafinesque	209
<i>erythrogaster</i> , Rafinesque	209
<i>eos</i> (Cope)	210
<i>dakotensis</i> , Evermann & Cox	210
<i>oreas</i> , Cope	211
Genus <i>Algансea</i> , Girard	211
<i>tinella</i> (Cuvier & Valenciennes)	211
<i>laeustris</i> , Steindachner	2140
<i>taraseorum</i> , Steindachner	2796
<i>dugesii</i> , Bean	211
<i>sallaei</i> (Günther)	212
Genus <i>Hybognathus</i> , Agassiz	212
Subgenus <i>Hybognathus</i>	213
<i>nuchalis</i> , Agassiz	213
<i>argyritis</i> , Girard	214
<i>hayi</i> , Jordan	214
Subgenus <i>Dionda</i> , Girard	214
<i>serena</i> (Girard)	214
<i>episeopu</i> (Girard)	214
<i>nubila</i> (Forbes)	215
<i>amara</i> (Girard)	215
<i>melanops</i> (Girard)	216
<i>plumbea</i> (Girard)	216
Genus <i>Pimephales</i> , Rafinesque	216
<i>promelas</i> , Rafinesque	217
<i>maculosus</i> (Girard)	217
<i>confertus</i> (Girard)	217
<i>notatus</i> (Rafinesque)	218
Genus <i>Mylopharodon</i> , Ayres	218
<i>conocephalus</i> (Baird & Girard)	219
Genus <i>Mylocheilus</i> , Agassiz	219
<i>caurinus</i> (Richardson)	219
Genus <i>Stypondon</i> , Garman	220
<i>signifer</i> , Garman	220
Genus <i>Semotilus</i> , Rafinesque	220
Subgenus <i>Leucosomus</i> , Heckel	221
<i>corporalis</i> (Mitchill)	221
Subgenus <i>Semotilus</i>	222
<i>atromaculatus</i> (Mitchell)	222
<i>thoreauianus</i> (Jordan)	223
Genus <i>Pogonichthys</i> , Girard	223
<i>maerolepidotus</i> (Ayres)	223
Genus <i>Ptychocheilus</i> , Agassiz	224
<i>oregonensis</i> (Richardson)	224
<i>grandis</i> (Ayres)	225; 2796
<i>lucius</i> , Girard	225

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOSPONDYLI—Continued.	
SUBORDER EVENTOGNATHI—Continued.	
Family Cyprinidae—Continued.	
Genus <i>Gila</i> , Baird & Girard.....	226
<i>elegans</i> , Baird & Girard	226
<i>robusta</i> , Baird & Girard	227
<i>semilunata</i> , Cope & Yarrow	228
Genus <i>Leuciscus</i> , Cuvier.....	228
Subgenus <i>Sibona</i> , Girard	231
<i>crassicauda</i> (Baird & Girard)	231
Subgenus <i>Tigoma</i> , Girard	231
<i>conformis</i> (Baird & Girard)	231
<i>bicolor</i> (Girard)	232
<i>lineatus</i> (Girard)	232
<i>nigrescens</i> (Girard)	233
<i>purpureus</i> (Girard)	234
<i>intermedius</i> (Girard)	235
<i>niger</i> (Cope)	235
<i>alicia</i> (Jouy)	236
Subgenus <i>Cheonda</i> , Girard.....	236
<i>cooperi</i> (Girard)	236
<i>humboldti</i> (Girard)	236
<i>egregius</i> (Girard)	237
<i>hydropbiox</i> (Cope)	238
Subgenus <i>Richardsonius</i> , Girard	238
<i>balteatus</i> (Richardson)	238
<i>iuslawi</i> , Evermann & Meek	2797
Subgenus <i>Clinostomus</i> , Girard	239
<i>vandoisniius</i> , Cuvier & Valenciennes	239
<i>elongatus</i> (Kirtland)	240
<i>nachtriebi</i> , Cox	2798
Subgenus <i>Phoxinus</i> , Rafinesque	240
<i>neogaeus</i> (Cope)	240
<i>margarita</i> (Cope)	241
<i>orculti</i> (Eigenmann & Eigenmann)	241
Subgenus <i>Hemitremia</i> , Cope	242
<i>milnerianus</i> (Cope)	242
<i>flammens</i> (Jordan & Gilbert)	242
Subgenus <i>Iotichthys</i> , Jordan & Evermann	243
<i>phlegethonitis</i> (Cope)	243
Genus <i>Rutilus</i> , Rafinesque	243
Subgenus <i>Opsopodus</i>	248
<i>osculus</i> , Evermann	248
<i>emilia</i> , Hay	248
<i>megalops</i> (Forbes)	248
Subgenus <i>Opsopoea</i> , Jordan & Evermann	249
<i>bolmani</i> , Gilbert	249
Genus <i>Abramia</i> , Cuvier	249
Subgenus <i>Notemigonus</i> , Rafinesque	250
<i>crysoleneas</i> (Mitchill)	250
<i>bosci</i> (Cuvier & Valenciennes)	251
<i>gardonensis</i> (Cuvier & Valenciennes)	251

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOSPONDYLI—Continued.	
SUBORDER EVENTOOGNATHI—Continued.	
<i>Family Cyprinidae—Continued.</i>	
Genus <i>Cochlognathus</i> , Baird & Girard	251
<i>ornata</i> , Baird & Girard.....	252
<i>bifurcata</i> , Cope.....	252
Genus <i>Cliola</i> , Girard	252
<i>vigilax</i> (Baird & Girard).....	253
<i>smithii</i> , Evermann & Cox	253
Genus <i>Notropis</i> , Rafinesque	254
Subgenus <i>Aztecula</i> , Jordan & Evermann.....	258; 2790
<i>aztecus</i> , Woolman.....	528
Subgenus <i>Chriope</i> , Jordan	258
<i>bifrenatus</i> (Cope)	258
<i>jordani</i> , Eigenmann & Eigenmann	259
<i>maeulatus</i> (Hay)	259
<i>anogenus</i> , Forbes	259
<i>cayuga</i> , Meek	260
<i>atrocaudalis</i> , Evermann	260
<i>heterodon</i> (Cope)	261
<i>callentis</i> , Jordan & Snyder	3197
<i>muskoka</i> , Meek	3141
<i>welaka</i> , Evermann & Kendall	2799
Subgenus <i>Alburnops</i> , Girard	261
<i>fretensis</i> (Cope)	261
<i>blennius</i> (Girard)	261
<i>buchananii</i> , Meek	2800
<i>sabine</i> , Jordan & Gilbert	262
<i>volucellus</i> (Cope)	263
<i>seyilla</i> (Cope)	263
<i>procne</i> (Cope)	264
<i>rasconis</i> , Jordan & Snyder	3141
<i>nigrotentaculus</i> (Günther)	264
<i>kanawha</i> , Jordan & Jenkins	264
<i>braytoni</i> , Jordan & Evermann	264
<i>spectrunculus</i> (Cope)	265
<i>ozarcanus</i> , Meek	265
<i>chihualma</i> , Woolman	265
<i>topeka</i> , Gilbert	266
Subgenus <i>Hudsonius</i> , Girard	266
<i>gilberti</i> , Jordan & Meek	266
<i>piptolepis</i> (Cope)	266
<i>simus</i> (Cope)	267
<i>longirostris</i> (Hay)	267
<i>nux</i> , Evermann	267
<i>noconis</i> , Evermann	268
<i>shumardi</i> (Girard)	268
<i>illecebrosus</i> (Girard)	268
<i>hudsonius</i> (De Witt Clinton)	269
<i>seiene</i> (Jordan)	269
<i>amarus</i> (Girard)	270
<i>saludanus</i> (Jordan & Brayton)	270
Subgenus <i>Codoma</i> , Girard	270
<i>ornatus</i> (Girard)	270
Subgenus <i>Moniana</i> , Girard	271
<i>formosus</i> (Girard)	271
<i>frigidus</i> (Girard)	271

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOSPONDYLI—Continued.	
SUBORDER EVENTOGNATHI—Continued.	
<i>Family Cyprinidae</i> —Continued.	
Genus <i>Notropis</i> , Rafinesque—Continued.	
Subgenus <i>Moulana</i> , Girard—Continued.	
<i>lutrensis</i> (Baird & Girard)	271
<i>proserpina</i> (Girard)	272
<i>callisema</i> (Jordan)	272
Subgenus <i>Cyprinella</i> , Girard	273
<i>bimaculatus</i> (Baird & Girard)	273
<i>ludibundus</i> (Girard)	273
<i>macrostomus</i> (Girard)	274
<i>texanus</i> (Girard)	274
<i>notatus</i> (Girard)	274
<i>venustus</i> (Girard)	274
<i>cereostigma</i> (Cope)	275
<i>stigmaturus</i> (Jordan)	275
<i>trichroistius</i> (Jordan & Gilbert)	275
<i>callistius</i> (Jordan)	276
<i>eurystomus</i> (Jordan)	276
<i>cernleus</i> (Jordan)	277
<i>nivens</i> (Cope)	277
<i>chloristius</i> (Jordan & Brayton)	278
<i>whipplei</i> (Girard)	278
<i>analostomus</i> (Girard)	279
<i>galacturus</i> (Cope)	279
<i>camurus</i> (Jordan & Meek)	279
<i>xenurus</i> (Jordan)	280
<i>hypselopterus</i> (Günther)	280
<i>pyrrhomelas</i> (Cope)	280
<i>garmani</i> , Jordan	281
Subgenus <i>Luxilus</i> , Rafinesque	281
<i>cornutus</i> (Mitchill)	281
<i>frontalis</i> (Agassiz)	283
<i>cyanus</i> (Cope)	283
<i>erasinus</i> (Cope)	283
<i>albeolus</i> (Jordan)	283
<i>lacertosus</i> (Cope)	284
Subgenus <i>Hydrophilox</i> , Jordan	284
<i>macleodaldi</i> , Jordan & Jenkins	284
<i>coccogenis</i> (Cope)	284
<i>zonatus</i> (Agassiz)	285
<i>zonistius</i> (Jordan)	285
<i>rubericrocens</i> (Cope)	286
<i>chlorocephalus</i> (Cope)	286
<i>lutipinnis</i> (Jordan & Brayton)	286
<i>chamberlaini</i> , Evermann	2800
<i>chiliticus</i> (Cope)	287
<i>altipinnis</i> (Cope)	287
<i>rosens</i> (Jordan)	287
<i>chalybeus</i> (Cope)	288
<i>chrosomus</i> (Jordan)	288
<i>xenocephalus</i> (Jordan)	289
Subgenus <i>Oreula</i> , Jordan & Evermann	289; 3140
<i>oreca</i> , Woolman	289
Subgenus <i>Notropis</i>	290
<i>ariommus</i> (Cope)	290

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOSPONDYLI—Continued.	
SCHROEDER EVENTOGNATHI—Continued.	
Family Cyprinidae—Continued.	
Genus <i>Notropis</i> , Rafinesque—Continued.	
Subgenus <i>Notropis</i> —Continued.	
<i>senibriceps</i> (Cope)	290
<i>jejunus</i> (Forbes)	290
<i>swalni</i> , Jordan	290
<i>amabilis</i> (Girard)	291
<i>lencodinus</i> (Cope)	291
<i>telescopus</i> (Cope)	
<i>arcansum</i> , Meek	292
<i>socius</i> (Girard)	292
<i>notemigonoides</i> , Evermann	292
<i>stillbins</i> , Jordan	293
<i>atherinoides</i> , Rafinesque	293
<i>urge</i> (Cope)	294
<i>dilecta</i> (Girard)	294
<i>louisiana</i> , Evermann	294
<i>fineus</i> , Evermann	294
<i>rubrifrons</i> (Cope)	295
<i>photogenis</i> (Cope)	295
<i>anuenus</i> (Abbott)	296
<i>sceptiens</i> (Jordan & Gilbert)	296
<i>micropteryx</i> (Cope)	296
<i>metallifer</i> , Jordan & Meek	297
Subgenus <i>Lythrurus</i> , Jordan	297
<i>bellus</i> (Hay)	297
<i>lirus</i> (Jordan)	297
<i>roseipinnis</i> , Hay	298
<i>umbratilis</i> (Girard)	298
<i>umbratilis</i> (Girard)	299
<i>atripes</i> (Jordan)	300
<i>lythrurus</i> (Jordan)	300
<i>cyanoccephalus</i> (Copeland)	300
<i>ardens</i> (Cope)	301
<i>fasciolaris</i> , Gilbert	301
<i>matutinus</i> (Cope)	301
<i>punctulatus</i> (Hay)	301
Genus <i>Xystrosus</i> , Jordan & Snyder	3142
<i>popoche</i> , Jordan & Snyder	3142
Genus <i>Ericymba</i> , Cope	302
<i>buccata</i> , Cope	302
Genus <i>Phenacobius</i> , Cope	302
<i>tertulus</i> , Cope	303
<i>miraibilis</i> (Girard)	303
<i>scopifer</i> (Cope)	303
<i>uranops</i> , Cope	304
<i>catostomus</i> Jordan	304
Genus <i>Evarra</i> , Woolman	304
<i>eigenmanni</i> , Woolman	304
Genus <i>Falcila</i> , Jordan & Snyder	3143
<i>chapalae</i> , Jordan & Snyder	3143
Genus <i>Tiaroga</i> , Girard	305
<i>cobitis</i> , Girard	305
Genus <i>Rhinichthys</i> , Agassiz	305
<i>earractae</i> (Cuvier & Valenciennes)	306
<i>duleis</i> (Girard)	306

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOSPONDYLI—Continued.	
SUBORDER EVENTOGNATHI—Continued.	
Family Cyprinidae—Continued.	
Genus <i>Rhinichthys</i> , Agassiz—Continued.	
<i>sinus</i> , Garman.....	307
<i>atronasus</i> (Mitchill)	307
<i>crocens</i> (Storer).....	308
<i>lunatus</i> (Cope)	308
<i>meleagris</i> (Agassiz).....	308
Genus <i>Agosia</i> , Girard	308
Subgenus <i>Apocope</i> , Cope.....	309
<i>oseula</i> (Girard)	309
<i>yarrowi</i> , Jordan & Evermann.....	309
<i>comesii</i> (Yarrow).....	310
<i>adobe</i> , Jordan & Evermann	310
<i>nevadensis</i> (Gilbert).....	310
<i>nubila</i> (Girard)	311
<i>carringtonii</i> (Cope)	311
<i>klamathensis</i> , Evermann & Meek	314
<i>velifera</i> (Gilbert).....	312
<i>umatilla</i> , Gilbert & Evermann	313
<i>falcata</i> , Eigenmann & Eigenmann	313
Subgenus <i>Agosia</i>	313
<i>chrysogaster</i> , Girard	313
Genus <i>Hybopsis</i> , Agassiz	314
Subgenus <i>Erimystax</i> , Jordan	315
<i>tetranemus</i> , Gilbert	315
<i>astivallis</i> (Girard)	316
<i>marconis</i> , Jordan & Gilbert	316
<i>hyostomus</i> (Gilbert)	316
<i>gelidus</i> (Girard).....	316
<i>meeki</i> , Jordan & Evermann	317
<i>montanus</i> , Meek	317
<i>cumingii</i> (Günther)	318
<i>monacanth</i> (Cope)	318
<i>dissimilis</i> (Kirtland)	318
<i>watauga</i> , Jordan & Evermann	319
Subgenus <i>Hybopsis</i>	319
<i>labrosus</i> (Cope)	319
<i>hypsinotus</i> (Cope)	320
<i>rubrifrons</i> (Jordan)	320
<i>amblops</i> (Rafinesque)	320
<i>storerianus</i> (Kirtland)	321
Subgenus <i>Yuriria</i> , Jordan & Evermann	321
<i>altus</i> (Jordan)	321
Subgenus <i>Noconis</i> , Girard	322
<i>kentuckiensis</i> (Rafinesque)	322
Genus <i>Couesius</i> , Jordan	323
<i>squamiferinus</i> (Cope).....	323
<i>plumbeus</i> (Agassiz).....	323
<i>dissimilis</i> (Girard).....	324
<i>greeni</i> , Jordan	324
<i>adustus</i> , Woolman	325
Genus <i>Platygobio</i> , Gill	325
<i>physignathus</i> (Cope)	325
<i>gracilis</i> (Richardson)	326
<i>pallidus</i> , Forbes.....	326

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOSPONDYLI—Continued.	
SUBORDER EVENTOGNATHI—Continued.	
Family Cyprinidae—Continued.	
Genus Exoglossum, Rafinesque	327
maxillingua (Le Sueur)	327
Genus Lepidomeda, Cope	328
vittata, Cope	328
jarrovi, Cope	328
Genus Meda, Girard	328
fulgidu, Girard	329
Genus Plagopterus, Cope	329
argentissimus, Cope	329
SUBORDER HETEROGNATHI.....	329
Family Erythrinidae.....	330
Genus Macrodon, Müller	330
microlepis, Günther	330
Family Characinae.....	331
Genus Curimata (Cuvier) Cloquet	332
magdalene, Steindachner	332
Genus Piabucina, Cuvier & Valenciennes	332
panamensis, Gill	332
festae, Boulenger	3145
Genus Tetragonopterus, Cuvier	333
Subgenus Astyanax, Baird & Girard	333
aneus, Günther	333
rutilus, Jenyns	334
panamensis, Günther	334
microphthalmus, Günther	334
ørstedii, Kröver	334
petenensis, Günther	335
seabripiennis, Jenyns	335
humilis, Günther	335
brevimanus, Günther	335
mexicanus, Filippi	335
argentatus (Baird & Girard)	336
Genus Brycon, Müller & Troschel	337
Subgenus Chalcinopsis, Kner	337
dentex, Günther	337
striatulus (Kner)	337
Genus Gasteropelecus (Gronow) Pallas	337
maculatus, Steindachner	338
Genus Reboides, Günther	338
guatemalensis (Günther)	338
Genus Brachycharax, Gill	338
bransfordi, Gill	339
Genus Luciocharax, Steindachner	339
inseulatus, Steindachner	339
SUBORDER GYMONOTI	339
Family Gymnotidae	340
Genus Giton, Kaup	340
fasciatus (Pallas)	340
Genus Eigenmannia, Jordan & Evermann	341
humboldti (Steindachner)	341
ORDER SYMBRANCHIA	341
Family Symbranchidae	342
Genus Symbranchus, Bloch	342
marmoratus, Bloch	342

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER CARENCHELVI.	343
<i>Family Derichthyidae</i>	343
Genus <i>Derichthys</i> , Gill	343
<i>serpentinus</i> , Gill	343
ORDER APODES	344
SCHORDER ENCHELYCEPHALI	346
<i>Family Anguillidae</i>	346
Genus <i>Anguilla</i> , Shaw	347
<i>chrysypa</i> , Rafinesque	348
<i>Family Simenichelyidae</i>	348
Genus <i>Simenichelys</i> , Gill	349
<i>parasiticus</i> , Gill	349
<i>Family Iyophidae</i>	349
Genus <i>Iyophis</i> , Gilbert	349
<i>brunnneus</i> , Gilbert	350
<i>Family Synaphobranchidae</i>	350
Genus <i>Synaphobranchus</i> , Johnson	351
<i>pinnatus</i> (Gronow)	351
Genus <i>Histiobranchus</i> , Gill	351
<i>bathybius</i> , Günther	352
<i>infernalis</i> , Gill	352
<i>Family Leptocephalidae</i>	352
Genus <i>Leptocephalus</i> (Gronow) Seopoli	353
<i>conger</i> (Linnaeus)	354
<i>candilimbatus</i> (Poey)	355
Genus <i>Congrelbus</i> , Ogilby	355; 2801
<i>balearicus</i> (De la Roche)	356
<i>maerurus</i> (Gilbert)	356
<i>prorigera</i> (Gilbert)	357
<i>nitens</i> (Jordan & Bollman)	357
<i>flavus</i> (Goode & Bean)	357
Genus <i>Uroconger</i> , Kaup	358
<i>vicinus</i> , Vaillant	358
<i>Family Muraenesocidae</i>	358
Genus <i>Muraenesox</i> , McClelland	359
Subgenus <i>Muraenesox</i>	359
<i>coniceps</i> , Jordan & Gilbert	359
<i>savanne</i> (Cuvier)	360
Genus <i>Xenomystax</i> , Gilbert	360
<i>atrarius</i> , Gilbert	361
Genus <i>Hoplunnis</i> , Kaup	361
<i>schmidtii</i> , Kaup	361
<i>diomedianus</i> , Goode & Bean	361
Genus <i>Neconger</i> , Girard	362
<i>muconatus</i> , Girard	362
<i>vermiformis</i> , Gilbert	362
Genus <i>Leptocenger</i> , Poey	362
<i>perlongus</i> (Poey)	363
Genus <i>Stibliscens</i> , Jordan & Bollman	363
<i>edwardsi</i> , Jordan & Bollman	363
Genus <i>Gordlichthys</i> , Jordan & Davis	363
<i>irretitus</i> , Jordan & Davis	363
<i>Family Nettastomatidae</i>	364
Genus <i>Chiopis</i> , Rafinesque	364
<i>equatorialis</i> , Gilbert	364
Genus <i>Venechia</i> , Jordan & Davis	365
<i>procera</i> , Goode & Bean	365

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER APODES—Continued.	
SUBORDER ENCHELYCEPHALI—Continued.	
<i>Family Nemichthyidae</i>	366
Genus Serrivomer, Gill & Ryder	367
beanii, Gill & Ryder	367
Genus Spinivomer, Gill & Ryder	367
goodei, Gill & Ryder	367
Genus Avocettina, Jordan & Davis	367
infans (Günther)	367
gilli (Bean)	368; 2801
elongata (Gill & Ryder)	369; 2802
Genus Labichthys, Gill & Ryder	368
carinatus, Gill & Ryder	368
Genus Neumichthys, Richardson	369
scolopaceus, Richardson	369
avocetta, Jordan & Gilbert	369
<i>Family Myriidae</i>	370
Genus Ahlia, Jordan & Davis	370
egmontis (Jordan)	370
Genus Myrophis, Lütken	371
punctatus, Lütken	371
vafer, Jordan & Gilbert	372
Genus Chilorhinus, Lütken	372
suensonii, Lütken	372
<i>Family Ophichthyidae</i>	372
Genus Sphagebranchus, Bloch	373
anguiformis (Peters)	374
selachops (Jordan & Gilbert)	374
Genus Verma, Jordan & Evermann	374
kendalli (Gilbert)	375
Genus Letharchus, Goode & Bean	375
velifer, Goode & Bean	375
Genus Myrichthys, Girard	375
tigrinus, Girard	376
xysturus, Jordan & Gilbert	2802
pantostigmus, Jordan & McGregor	2802
oenatus (Kaup)	376
acuminatus (Gronow)	376
Genus Pisodonophis, Kaup	377
ernentifer, Goode & Bean	377
daspilotus, Gilbert	2803
Genus Callechelys, Kaup	378
murena, Jordan & Evermann	378
Genus Baseanichthys, Jordan & Davis	378
s. uticaris (Goode & Bean)	378
peninsulae (Gilbert)	379
baseanium (Jordan)	379
Genus Quasistremus, Jordan & Davis	380
nothochir (Gilbert)	380
evionthas (Jordan & Bollman)	380
Genus Ophichthus, Thunborg & Ahl	381
Subgenus Cryptopterus, Kaup	382
puncticeps (Kaup)	382
Subgenus Ophichthus	382
ophis (Linnæus)	382; 2804
retropinnis (Eigenmann)	383

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER APODES—Continued.	
SUBORDER ENCHELYCEPHALI—Continued.	
<i>Family Ophichthyidae</i> —Continued.	
Genus <i>Ophichthus</i> , Thunberg & Ahl—Continued.	
Subgenus <i>Muraenopsis</i> , Kaup	383
guttifer (Bean & Dresel).....	383
ocellatus (Le Sueur)	383
triserialis (Kaup)	384
Subgenus <i>Seytalophis</i> , Kaup.....	384
gomesii (Castelnau).....	384
zophochir (Jordan & Gilbert).....	385
magnioculis (Kaup).....	385
parilis (Richardson).....	386
Genus <i>Mystriophis</i> , Kaup	386
Subgenus <i>Echiopsis</i> , Kaup.....	386
intertinctus (Richardson).....	386
Genus <i>Seytalichthys</i> , Jordan & Davis	387
mifurus (Jordan & Gilbert).....	387
Genus <i>Brachysomophis</i> , Kaup	387
errocodilians (Bennett)	388
SUBORDER COLOCEPHALI	
<i>Family Muranidae</i>	
Genus <i>Enchelycore</i> , Kaup	389
nigricans (Bonnaterre).....	389
Genus <i>Pythonichthys</i> , Poey	390
sanguineus, Poey	390
Genus <i>Rabula</i> , Jordan & Davis	390
aque-dulcis (Cope)	390
marmorea (Valenciennes).....	391
panamensis (Steindachner).....	391
longicauda (Peters)	391
Genus <i>Lycodontis</i> , McClelland	392
Subgenus <i>Lycodontis</i>	393
verrilli (Jordan & Gilbert)	393
vicinus (Castelnau)	394
virescens (Poey)	394
polygonus (Poey)	394
moringa (Cuvier).....	395
mordax (Ayres).....	395
funebris (Ranzani)	396
sancta-helene (Günther)	397
castaneus (Jordan & Gilbert)	396; 2804
pietus (Ahl)	2805
dovii (Günther)	397
conspersus (Poey)	397
miliaris (Kaup)	397
elaboratus (Poey).....	398
obsecuratus (Poey).....	398
jordani, Evermann & Marsh	3145
chlevastes (Jordan & Gilbert).....	398
Subgenus <i>Priodonophis</i> , Kaup	399
ocellatus (Agassiz).....	399
saxicola (Jordan & Davis).....	399
nigromarginatus (Grandidier)	399
Genus <i>Muraena</i> (Artedi). Linnaeus.....	400
insularum, Jordan & Davis	400

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER APODES—Continued.	
SUPERORDER COLOCEPHALI—Continued.	
<i>Family Muraenidae—Continued.</i>	
Genus <i>Muraena</i> (Artedi) Linnaeus—Continued.	
<i>argus</i> (Steindachner).....	401
<i>clepsydra</i> , Gilbert.....	2805
<i>retifera</i> , Goode & Bean	401
<i>melanotis</i> (Kaup).....	401
<i>lentiginosa</i> , Jenyns.....	402
Genus <i>Echidna</i> , Forster.....	402
<i>nocturna</i> (Cope).....	402
<i>catenata</i> (Bloch).....	403
Genus <i>Uropterygius</i> , Rüppell.....	403
Subgenus <i>Scutica</i> , Jordan & Evermann	404
<i>necturus</i> (Jordan & Gilbert)	404
Genus <i>Channichthys</i> , Richardson	404
<i>vittata</i> (Richardson)	404
ORDER LYOMERI.	404
<i>Family Saccopharyngidae</i>	405
Genus <i>Saccopharynx</i> , Mitchell.....	405
<i>ampullaceus</i> (Harwood).....	406
<i>Family Eurypharyngidae</i>	406
Genus <i>Gastrostomus</i> , Gill & Ryder	406
<i>hairii</i> , Gill & Ryder	406
ORDER ISOSPONDYLI.	407
<i>Family Elopidae</i>	408
Genus <i>Tarpon</i> , Jordan & Evermann	409
<i>atlanticus</i> (Cuvier & Valenciennes)	409
Genus <i>Elops</i> , Linnaeus	409
<i>saurns</i> , Linnaeus	410
<i>Family Albulidae</i>	410
Genus <i>Albulia</i> (Gronow) Scopoli	411; 2807
<i>vulpes</i> (Linnaeus)	411
<i>Family Hiodontidae</i>	412
Genus <i>Hiodon</i> , Le Sueur	412
Subgenus <i>Amphiodon</i> , Rafinesque	413
<i>alosoides</i> (Rafinesque)	413
Subgenus <i>Hiodon</i>	413
<i>tergisus</i> , Le Sueur	413
<i>selenops</i> , Jordan & Bean	414
<i>Family Chanidae</i>	414
Genus <i>Chanos</i> , Lacépède	414
<i>chanos</i> (Forskål)	414
<i>Family Dorosomatidae</i>	415
Genus <i>Dorosoma</i> , Rafinesque	415
<i>cepedianum</i> (Le Sueur)	416
<i>exile</i> , Jordan & Gilbert	416
<i>mexicanum</i> (Günther)	416
<i>petenense</i> (Günther)	417
Genus <i>Signalosa</i> , Evermann & Kendall	2809
<i>achaftalaye</i> , Evermann & Kendall	2809
<i>Family Clariidae</i>	417
Genus <i>Jenkinsia</i> , Jordan & Evermann	418
<i>acuminata</i> (Gilbert)	419
<i>lamprotentis</i> (Gosse)	419
<i>stolifera</i> (Jordan & Gilbert)	419

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ISOSPONDYLI—Continued.	
<i>Family Clupeidae—Continued.</i>	
Genus <i>Etrumeus</i> , Bleeker.....	419
<i>sadina</i> (Mitchill)	420
Genus <i>Perklusia</i> , Rosa Smith Eigenmann	420
<i>othonops</i> , R. S. Eigenmann.....	420
Genus <i>Clupea</i> (Arted) Linnaeus	421
<i>harengus</i> , Linnaeus.....	421
<i>pallasii</i> , Cuvier & Valenciennes	422
<i>caruleus</i> (Girard)	423
<i>pseudohispanicus</i> (Poey).....	423
Genus <i>Pomolobus</i> , Rafinesque	424
<i>chrysichleris</i> , Rafinesque	425
<i>mediooculis</i> , Mitchill	425
<i>pseudoharengus</i> (Wilson)	426
<i>astivalis</i> (Mitchill)	426
Genus <i>Alosa</i> , Cuvier.....	427
<i>sapidissima</i> (Wilson)	427
<i>alabama</i> , Jordan & Evermann.....	2810
Genus <i>Sardinella</i> , Cuvier & Valenciennes.....	428
Subgenus <i>Sardinella</i>	429
<i>anchovia</i> (Cuvier & Valenciennes)	429
<i>clupeola</i> (Cuvier & Valenciennes)	429
<i>apicalis</i> (Müller & Troschel)	429
<i>bishopi</i> (Müller & Troschel)	430
Subgenus <i>Harengula</i> , Cuvier & Valenciennes.....	430
<i>sardina</i> (Poey)	430
<i>macrophthalmus</i> (Rauzani)	430
<i>tbrissina</i> (Jordan & Gilbert)	430
<i>humeralis</i> (Cuvier & Valenciennes)	431
Subgenus <i>Lile</i> , Jordan & Evermann.....	431
<i>stolifera</i> (Jordan & Gilbert)	431
Genus <i>Opisthonema</i> , Gill.....	432
<i>ogibum</i> (Le Sueur)	432
<i>libertate</i> (Günther)	433
Genus <i>Brevoortia</i> , Gill.....	433
<i>tyrannus</i> (Latrobe)	433
<i>aurea</i> (Agassiz)	434
<i>brevicaudata</i> , Goode	434
<i>patronus</i> , Goode.....	434
Genus <i>Chirocentrodon</i> , Günther	435
<i>teniatus</i> , Günther	435
Genus <i>Ilisha</i> , Gray.....	435
<i>flavipinnis</i> (Valenciennes)	435
<i>bleekeriana</i> (Poey)	436
<i>furthi</i> (Steindachner)	436
Genus <i>Oplisopterus</i> , Gill	436
<i>latipinnis</i> (Jordan & Gilbert)	437
<i>dovii</i> (Günther)	437
<i>macrops</i> (Günther)	437
Genus <i>Odontognathus</i> , Lacépède	437
<i>mucronata</i> , Lacépède	438
<i>panamensis</i> (Steindachner)	438
Genus <i>Pristigaster</i> , Cuvier	438
<i>cayamus</i> , Cuvier	438

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER ISOSPONDYLI—Continued.	
<i>Family Engraulidae</i>	439
Genus <i>Stolephorus</i> , Lacépède	439
<i>miarchus</i> , Jordan & Gilbert	441
<i>perfasciatus</i> (Poey)	441
<i>exiguum</i> , Jordan & Gilbert	442
<i>cabanus</i> (Poey)	442
<i>perthecatus</i> , Goode & Bean.....	442
<i>ischanus</i> , Jordan & Gilbert.....	442
<i>brownii</i> (Gmelin).....	443
<i>culturatus</i> , Gilbert	443
<i>delicatissimus</i> (Girard)	444
<i>cherostomus</i> (Goode)	444
<i>argyrophanus</i> (Cuvier & Valenciennes)	444
<i>curtus</i> , Jordan & Gilbert	445
<i>opercularis</i> , Jordan & Gilbert	445
<i>mitchilli</i> (Cuvier & Valenciennes)	446
<i>lucidus</i> , Jordan & Gilbert	446
<i>rastralis</i> , Gilbert & Pierson	2811
<i>mundulus</i> , Gilbert & Pierson	2812
<i>naso</i> , Gilbert & Pierson	2813
<i>starksii</i> , Gilbert & Pierson	2813
<i>clupeoides</i> (Swainson).....	447
<i>productus</i> (Poey).....	447
<i>gilberti</i> , Evermann & Marsh	3146
<i>garmani</i> , Evermann & Marsh	3146
<i>compressus</i> (Girard)	447
<i>panamensis</i> (Steindachner).....	448
<i>spinifer</i> (Cuvier & Valenciennes)	448
<i>seofieldi</i> , Jordan & Culver.....	2814
<i>astilbe</i> , Jordan & Rutter	2815
<i>robertsi</i> , Jordan & Rutter	2815
<i>macrolepidotus</i> (Kner & Steindachner)	449: 2815
Genus <i>Engraulis</i> , Cuvier.....	448
<i>mordax</i> , Girard.....	448
Genus <i>Cetengraulis</i> , Günther.....	450
<i>mysticetus</i> (Günther)	450
<i>edentulus</i> (Cuvier)	450
<i>engyumen</i> , Gilbert & Pierson.....	2815
Genus <i>Pterengraulis</i> , Günther	450
<i>atherinooides</i> (Linnaeus)	450
Genus <i>Lycengraulis</i> , Günther	451
<i>grossidens</i> (Cuvier)	451
<i>poeyi</i> (Kner & Steindachner).....	445: 2811
<i>Family Alepocephalidae</i>	451
Genus <i>Alepocephalus</i> , Risso.....	452
<i>productus</i> , Gill	452
<i>agassizii</i> , Goode & Bean.....	453
<i>tenebrosus</i> , Gilbert.....	453
Genus <i>Mitchillina</i> , Jordan & Evermann	453
<i>bairdii</i> (Goode & Bean).....	454
Genus <i>Bathytroctes</i> , Günther	454
<i>stomias</i> , Gilbert	454
Genus <i>Talismania</i> , Goode & Bean	455
<i>antillarum</i> , Goode & Bean	455
<i>equatorialis</i> , Goode & Bean	456

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER ISOSPONDYLI—Continued.	
Family Alepocephalidae—Continued.	
Genus Conocara, Goode & Bean.....	456
macdonaldi, Goode & Bean.....	457
macroptera (Vaillant)	457
Genus Platypterus, Günther	458
apus, Günther.....	458
Genus Ericara, Gill & Townsend	2816
salmoncea, Gill & Townsend	2816
Genus Aleposomus, Gill	459
copei, Gill.....	459
Family Salmonidae	460
Genus Coregonus (Artedi) Linnaeus.....	461
Subgenus Prosopium, Milner	462
coulterii, Eigenmann & Eigenmann	462
williamsoni, Girard	463
cismontanus, Jordan	463
kennicotti, Milner.....	464
richardsonii, Günther.....	465
quadrilateralis, Richardson	465
Subgenus Coregonus.....	465
clupeiformis (Mitchill)	465
nelsonii, Bean	466
labradoricus, Richardson	466
Genus Argyrosomus, Agassiz.....	467
Subgenus Argyrosomus.....	468
osmeriformis (H. M. Smith)	468
artedi (Le Sueur)	468
sisco, Jordan	469
boyi, Gill.....	469
pusillus (Bean)	470
lucidus (Richardson)	470
laurettae (Bean)	471
alascanus, Scofield.....	2817
prognathus (H. M. Smith)	471
nigripinnis, Gill.....	472
Subgenus Allosomus, Jordan	473
tullibee (Richardson)	473
bisselli, Böhlman	473
Genus Stenodus, Richardson.....	473
mackenzii (Richardson)	474
Genus Oncorhynchus, Seckley.....	474
Subgenus Oncorhynchus.....	478
gorbuscha (Walbaum)	478
keta (Walbaum)	478
tschawytscha (Walbaum)	479
kisutch (Walbaum)	480
Subgenus Hypsifario, Gill	481
nerka (Walbaum)	481
Genus Salmo (Artedi) Linnaeus.....	483
Subgenus Salmo.....	486
salar, Linnaeus	486
sebago (Girard)	487
ouananiche, McCarthy	487
Subgenus Trutta, Linnaeus.....	487
mykiss, Walbaum	487; 492; 2818

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER ISOSPONDYLI—Continued.	
<i>Family Salmonidae</i> —Continued.	
Genus <i>Salmo</i> (Artedi) Linnaeus—Continued.	
Subgenus <i>Trutta</i> , Linnaeus—Continued.	
<i>clarkii</i> (Richardson).....	492; 2819
<i>lewisi</i> (Girard)	493; 2819
<i>gibbsii</i> (Suckley).....	493; 2819
<i>henshawi</i> (Gill & Jordan)	493; 2819
<i>tahoensis</i> , Jordan & Evermann.....	2870
<i>virginicus</i> (Girard)	495; 2819
<i>spilurus</i> (Cope).....	495; 2819
<i>pleuriticus</i> (Cope)	496; 2819
<i>bouvieri</i> (Bendire)	496; 2819
<i>stomias</i> (Cope)	497; 2819
<i>macleodii</i> , Jordan & Evermann	497; 2819
<i>decellvifrons</i> , Meek	3147
<i>jordani</i> , Meek	3148
<i>gairdneri</i> , Richardson	497; 498
<i>kamloops</i> (Jordan)	499
<i>beardsleei</i> , Jordan & Seale.....	2819
<i>crescentis</i> , Jordan & Beardslee.....	2821
<i>bathaeator</i> , Meek	3149
<i>ridens</i> , Gibbons.....	500
<i>masoni</i> (Suckley)	501
<i>shasta</i> (Jordan).....	502
<i>gilberti</i> (Jordan)	502
<i>stonei</i> (Jordan).....	503
<i>agu-a-bonita</i> (Jordan)	503
Genus <i>Cristivomer</i> , Gill & Jordan	504
<i>nainayewash</i> (Walbaum).....	504
<i>siscowet</i> (Agassiz)	505
Genus <i>Salvelinus</i> (Nilsson) Richardson	506
<i>fontinalis</i> (Mitchill).....	506
<i>agassizii</i> (Garman).....	507
<i>malma</i> (Walbaum)	507; 2823
<i>parkei</i> (Suckley)	507; 2823
<i>kundschach</i> (Pallas).....	2822
<i>alpinus</i> (Linnaeus).....	508
<i>alipes</i> (Richardson)	509
<i>stagnalis</i> (Fabricius).....	510
<i>arcturus</i> (Günther)	510
<i>aureolus</i> (Bean)	511
<i>oquassa</i> (Girard)	514
<i>naresi</i> (Günther)	515
<i>marstoni</i> , Garman.....	515
<i>Family Thymallidae</i>	517
Genus <i>Thymallus</i> , Cuvier	517
<i>signifer</i> (Richardson)	517
<i>tricolor</i> , Cope	518; 2871
<i>montanus</i> (Milner)	519; 2871
<i>Family Argentiniidae</i>	519
Genus <i>Mallotus</i> , Cuvier.....	520
<i>vilosus</i> (Müller)	520
Genus <i>Thaleichthys</i> , Girard	521
<i>pacificus</i> (Richardson)	521

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ISOSPONDYLI—Continued.	
<i>Family Argentinidae—Continued.</i>	
Genus <i>Osmerus</i> (Artedi) Linnaeus	522
Subgenus <i>Spirinchus</i> , Jordan & Evermann	522
<i>thaleichthys</i> , Ayres	522
<i>attenuatus</i> , Lockington	523
Subgenus <i>Osmerus</i>	523
<i>mordax</i> (Mitchill)	523
<i>spectrum</i> (Cope)	523
<i>abbotti</i> (Cope)	524
<i>dentex</i> , Steindachner	524
<i>albatrossis</i> , Jordan & Gilbert	2823
Genus <i>Mesopna</i> , Gill	524
<i>pretiosus</i> (Girard)	525
<i>olidus</i> (Pallas)	525
Genus <i>Argentina</i> (Artedi) Linnaeus	525
<i>silus</i> , Ascanius	526
<i>sialis</i> , Gilbert	526
<i>striata</i> , Goode & Bean	526
Genus <i>Leucoglossus</i> , Gilbert	527
<i>stilbius</i> , Gilbert	527
<i>Family Microstomidae</i>	527
Genus <i>Nansenia</i> , Jordan & Evermann	528
<i>grøenlandica</i> (Reinhardt)	528
Genus <i>Bathylagus</i> , Günther	528
<i>benedicti</i> , Goode & Bean	529
<i>uryops</i> , Goode & Bean	529
<i>pacificus</i> Gilbert	530
<i>borealis</i> , Gilbert	2824
<i>milleri</i> , Jordan & Gilbert	2825
ORDER INOMI.....	530
<i>Family Synodontidae</i>	532
Genus <i>Trachinocéphalus</i> , Gill	533
<i>nyops</i> (Forster)	533
Genus <i>Synodus</i> (Gronow) Scopoli	533; 2807
<i>intermedius</i> (Agassiz)	533
<i>evermanni</i> , Jordan & Bollman	535
<i>poeyi</i> , Jordan	536
<i>synodus</i> (Linnaeus)	536
<i>lacertinus</i> , Gilbert	536
<i>saurus</i> (Linnaeus)	537
<i>scituliceps</i> , Jordan & Gilbert	537
<i>jenkinsi</i> , Jordan & Bollman	537
<i>festens</i> (Linnaeus)	538
<i>lucioceps</i> (Ayres)	539
Genus <i>Bathyaurus</i> , Günther	539
<i>ferox</i> , Günther	539
Genus <i>Bathyaco</i> , Goode & Bean	540
<i>nigricans</i> , Goode & Bean	540
<i>Family Autopidae</i>	541
Genus <i>Chlorophthalmus</i> , Bonaparte	541
<i>agassizii</i> , Bonaparte	541
<i>chalybeinus</i> (Goode)	542
<i>trunculentus</i> , Goode & Bean	542
<i>Family Benthosauridae</i>	543
Genus <i>Benthosaurus</i> , Goode & Bean	543
<i>grallator</i> , Goode & Bean	543

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER INIOMI—Continued.	
<i>Family Bathypteroide</i>	544
Genus <i>Bathypterois</i> , Günther.....	544
Subgenus <i>Synapteretinus</i> , Goode & Bean	545
<i>quadrifilis</i> , Günther.....	545
<i>longipes</i> , Günther.....	546
<i>Family Ipnopidae</i>	546
Genus <i>Ipnops</i> , Günther.....	546
<i>murrayi</i> , Günther	547
<i>Family Rondeletiidae</i>	547
Genus <i>Rondeletia</i> , Goode & Bean	548
<i>bicolor</i> , Goode & Bean	548
<i>Family Cetomimidae</i>	548
Genus <i>Cetomimus</i> , Goode & Bean	549
<i>gillii</i> , Goode & Bean.....	549
<i>storeeri</i> , Goode & Bean	550
<i>Family Myctophidae</i>	550
Genus <i>Macrostoerna</i> , Risso	554
<i>quercentium</i> (Goode & Bean).....	554
<i>margaritiformis</i> (Goode & Bean)	555
<i>angustidens</i> , Risso	555
<i>castaneum</i> (Goode & Bean)	556
<i>caudispinosum</i> (Johnson)	556
Genus <i>Ceratoscopelus</i> , Günther	557
<i>madeirensis</i> (Lowe)	557
Genus <i>Lampanyetus</i> , Bonaparte	557
<i>erectodilus</i> (Risso)	558
<i>townsendi</i> (Eigenmann & Eigenmann)	558
<i>alatus</i> , Goode & Bean	559
<i>guntheri</i> , Goode & Bean	559
<i>gemmifer</i> , Goode & Bean	559
<i>laeerta</i> (Goode & Bean)	560
Genus <i>Lampadena</i> , Goode & Bean	560
<i>speculigera</i> , Goode & Bean	561
Genus <i>Nannobrachium</i> , Günther	561
<i>leucopsarum</i> (Eigenmann & Eigenmann)	562
<i>nannocheir</i> (Gilbert)	562
<i>mexicanum</i> (Gilbert)	563
<i>regale</i> (Gilbert)	563
<i>maedonaldi</i> , (Goode & Bean)	563
Genus <i>Diaphus</i> , Eigenmann & Eigenmann	564
<i>theta</i> , Eigenmann & Eigenmann	564
Genus <i>Aethoprora</i> , Goode & Bean	565
<i>lucida</i> , Goode & Bean	565
<i>effulgens</i> , Goode & Bean	566
Genus <i>Colletta</i> , Goode & Bean	567
<i>rufinesquei</i> (Cocco)	567
<i>nocturna</i> (Poey)	567
Genus <i>Rhinoseopelus</i> , Lütken	568
<i>coccoi</i> (Cocco)	568
<i>andreae</i> (Lütken)	569
<i>rarus</i> (Lütken)	569
Genus <i>Myctophum</i> , Rafinesque	569
<i>punctatum</i> , Rafinesque	570
<i>affine</i> (Lütken)	570
<i>opalinum</i> , Goode & Bean	571
<i>humboldti</i> (Risso)	571

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER INIOMI—Continued.	
<i>Family Myctophidae</i> —Continued.	
Genus <i>Myctophum</i> , Rafinesque—Continued.	
<i>californiense</i> , Eigenmann & Eigenmann.....	572
<i>gracile</i> (Lütken).....	572; 3150
<i>benoiti</i> (Cocco).....	573
<i>hygomii</i> (Lütken).....	573
Genus <i>Benthosema</i> , Goode & Bean	573
<i>mulleri</i> (Gmelin).....	574
<i>arcticum</i> (Lütken).....	574
Genus <i>Dasy scopelus</i> , Günther	574
<i>spinosus</i> (Steindachner).....	575
Genus <i>Tarletonbeania</i> , Eigenmann & Eigenmann.....	575
<i>crenularis</i> (Jordan & Gilbert).....	575
<i>tenua</i> , Eigenmann & Eigenmann	575
<i>Family Maurolicidae</i>	576
Genus <i>Maurolicus</i> , Cocco	576
<i>penicanti</i> (Walbaum).....	577
Genus <i>Vineiguerria</i> , Jordan & Evermann	577
<i>attenuata</i> (Cocco).....	577
Genus <i>Valenciennellus</i> , Jordan & Evermann	577
<i>tripunctatus</i> (Esmark).....	578
<i>Family Chauliodontidae</i>	578
Genus <i>Gonostoma</i> , Rafinesque	578
<i>denudatum</i> , Rafinesque	579
<i>brevidens</i> , Kner & Steindachner	579
Genus <i>Zaphotias</i> , Goode & Bean	580; 2826
<i>pedaliotus</i> (Goode & Bean).....	580; 2826
Genus <i>Cyclothona</i> , Goode & Bean	581
Subgenus <i>Cyclothona</i>	582
<i>microdon</i> (Günther).....	582
<i>bathyphila</i> (Vaillant).....	582
Subgenus <i>Sigmops</i> , Gill	583
<i>elongata</i> (Günther).....	583
<i>megalops</i> Lütken	3150
Genus <i>Yarrella</i> , Goode & Bean	583
<i>blackfordi</i> , Goode & Bean	584
Genus <i>Chauliodus</i> , Bloch & Schneider	584
<i>sloanei</i> , Bloch & Schneider	585
<i>macouni</i> , Bean	585
<i>Family Astronesthidae</i>	586
Genus <i>Astronesthes</i> , Richardson	586
<i>niger</i> , Richardson	586
<i>gemmifer</i> , Goode & Bean	586
<i>richardsoni</i> (Poey)	587
<i>Family Stomiatidae</i>	587
Genus <i>Stomias</i> , Cuvier	588
<i>ferox</i> , Reinhardt	588
<i>affinis</i> , Günther	588
Genus <i>Echiostoma</i> , Lowe	589
<i>barbatum</i> , Lowe	589
<i>margarita</i> , Goode & Bean	589
Genus <i>Grammatostomias</i> , Goode & Bean	590
<i>dentatus</i> , Goode & Bean	590
Genus <i>Photonectes</i> , Günther	591
<i>gracilis</i> , Goode & Bean	591

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER INOMI—Continued.	
<i>Family Malacosteidae</i>	592
Genus <i>Malacosteus</i> , Ayres	592
<i>ulger</i> , Ayres	593
<i>Family Plagyodontidae</i>	593; 2826
Genus <i>Plagyodus</i> (Steller)	594; 2826
Subgenus <i>Plagyodus</i>	595
<i>ferox</i> (Lowe)	595
<i>ascenlapinus</i> (Bean)	595
Subgenus <i>Canlopus</i> , Gill	596
<i>alitivellis</i> (Poey)	596
<i>borealis</i> (Gill)	596
<i>serra</i> (Gill)	597
<i>Family Odontostomidae</i>	597
Genus <i>Omosudis</i> , Günther	598
<i>lowii</i> , Günther	598
<i>Family Paralepididae</i>	599
Genus <i>Sudis</i> , Ratiñesque	599
<i>intermedius</i> (Poey)	600
<i>ringens</i> , Jordan & Gilbert	600
Genus <i>Arctozenus</i> , Gill	601
<i>borealis</i> (Reinhardt)	601
<i>cornutus</i> (Jordan & Gilbert)	601
Genus <i>Paralepis</i> , Risso	602
<i>eoregonoides</i> , Risso	602
<i>Family Sternopychidae</i>	603
Genus <i>Sternopyx</i> , Hermann	603
<i>diaphana</i> , Hormann	603
Genus <i>Argyropelecus</i> , Cocco	603
<i>hemigymnus</i> , Cocco	604
<i>olfersi</i> (Cuvier)	604
<i>Family Idiacanthidae</i>	604
Genus <i>Idiacanthus</i> , Peters	605
<i>ferox</i> (Günther)	605
<i>antrostomus</i> , Gilbert	605
ORDER LYOPOMI	606
<i>Family Halosauridae</i>	606
Genus <i>Halesaurus</i> , Johnson	607
<i>oweni</i> , Johnson	607
<i>guntheri</i> , Goode & Bean	608
Genus <i>Halosaropsis</i> , Collett	608; 2826
<i>rostrata</i> (Günther)	609
<i>macrochir</i> (Günther)	609
<i>goodei</i> (Gill)	610
<i>gracilis</i> (Goode & Bean)	610
<i>pallida</i> (Goode & Bean)	611
ORDER HETEROMI	612
<i>Family Notacanthidae</i>	613
Genus <i>Notacanthus</i> , Bloch	614
<i>chemnitzi</i> , Bloch	614
<i>analis</i> , Gill	615
<i>phasganorus</i> , Goode	616
Genus <i>Macdonaldia</i> , Goode & Bean	616
<i>rostrata</i> (Collett)	617
<i>challengeri</i> (Vaillant)	617
<i>alta</i> , Gill & Townsend	2826
<i>longa</i> , Gill & Townsend	2826

	Page.
CLASS OSTEES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER HETEROCHILOMORPHI—Continued.	
Family Lipogenyidae	619
Genus Lipogenys, Goode & Bean	619
gilli, Goode & Bean	619
ORDER XENOMORPHI.....	620
Family Dallidae	620
Genus Dallia, Bean	621
pectoralis, Bean	621
ORDER HAPLOMORPHI	622
Family Umbridae	622
Genus Umbra (Krümer) Scopoli	623; 2407
Subgenus Melanura, Agassiz	623
linsi (Kirtland)	623
pygmaea (De Kay)	624
Family Luciidae	624
Genus Lucius, Radresque	625
Subgenus Kenoza, Jordan & Evermann	626
americanus (Gmelin)	626
vermiculatus (Le Sueur)	627
reflexatus (Le Sueur)	627
Subgenus Lucius	628
lucius (Linnaeus)	628
Subgenus Mascalongus, Jordan	629
masquinougy (Mitchill)	629
ohiensis (Kirtland)	629
immaculatus (Garrard)	630
Family Poeciliidae	630
Genus Fundulus, Lacépède	632
Subgenus Fundulus	637
punctatus, Günther	637
vinetus, Jordan & Gilbert	637
pallidus, Evermann	638
similis (Baird & Girard)	638
majalis (Walbaum)	639
parvipinnis, Girard	640
heterocellus (Linnaeus)	640
macrolepidotus (Walbaum)	641
badins, Garman	2827
grandis (Baird & Girard)	641; 2827
fonticola, Cuvier & Valenciennes	643
bermudae, Günther	643
robustus, Bean	644
labialis, Günther	644
Subgenus Fontinus, Jordau & Evermann	645
adinia, Jordan & Gilbert	645
diaphanus (Le Sueur)	645
menona (Jordan & Copeland)	645
extensus, Jordan & Gilbert	646
seminolis, Girard	647; 2828
Subgenus Planterus, Garman	2828
zebrinus, Jordan & Gilbert	646; 2828
Subgenus Xenisma, Jordan	648
catenatus (Storer)	648
stellifer (Jordan)	648
Subgenus Gambusinus, Jordan & Evermann	649
rathbuni, Jordan & Meek	649

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER HALOPOMI—Continued.	
<i>Family Poeciliidae</i> —Continued.	
Genus <i>Fundulus</i> , Lacépède—Continued.	
Subgenus <i>Gaibiusinus</i> , Jordan & Evermann—Continued.	
<i>albolineatus</i> , Gilbert	649
<i>confinitus</i> , Goode & Bean.....	642; 650; 2828
<i>funduloides</i> (Evermann)	650
Subgenus <i>Zygonectes</i> , Agassiz.....	650
<i>dovii</i> (Günther)	650
<i>floripinnis</i> (Cope)	651
<i>jenkinsi</i> (Evermann)	651
<i>pulvereus</i> (Evermann)	652
<i>scartes</i> , Meek.....	654
<i>seciadens</i> , Cope.....	649; 654; 2828
<i>leucie</i> (Baird)	654
<i>chrysotus</i> , Holbrook	655; 2828
<i>cingulatus</i> , Cuvier & Valenciennes.....	656; 2829
<i>nottii</i> (Agassiz)	656; 2830
<i>dispar</i> (Agassiz)	658
<i>notatus</i> (Rafinesque).....	659
<i>goodei</i> (Jordan)	664; 2831
Genus <i>Aplochelius</i> , McClelland.....	2830
<i>dovii</i> (Günther)	650; 2830
Genus <i>Adinia</i> , Girard	660
<i>guatemalensis</i> (Günther).....	660
<i>pachycephala</i> (Günther).....	660
<i>dugesii</i> (Beau).....	661
<i>multifasciata</i> , Girard.....	661
Genus <i>Rivulus</i> , Poey	662
<i>cylindraceus</i> , Poey	662; 2830
<i>isthmensis</i> , Garman.....	2830
Genus <i>Lucania</i> , Girard	663
<i>ommuta</i> (Jordan)	663
<i>venusta</i> , Girard	665
<i>parva</i> (Baird & Girard).....	665
Genus <i>Girardinichthys</i> , Bleeker	666
<i>innominatus</i> , Bleeker.....	666
Genus <i>Empetrichthys</i> , Gilbert	666
<i>merriami</i> , Gilbert	667
Genus <i>Characodon</i> , Günther	667
<i>lateralis</i> , Günther	668
<i>bilineatus</i> , Bean.....	668
<i>variatus</i> , Bean.....	669
<i>eucanthus</i> , Jordan & Snyder	3150
<i>furcidens</i> , Jordan & Gilbert.....	669
<i>eiseni</i> , Rutter	2831
<i>garmani</i> , Jordan & Evermann	2831
Genus <i>Cyprinodon</i> , Lacépède	670
<i>variegatus</i> , Lacépède	671
<i>rivendredi</i> (Poey)	673; 2832
<i>eximus</i> , Girard	673; 2832
<i>bovinus</i> , Baird & Girard.....	673
<i>macularius</i> , Baird & Girard.....	674
<i>balleyi</i> (Gilbert)	675
<i>marte</i> , Steindachner	675
<i>carpio</i> , Günther	675

CLASS PISCES—Continued.

Page.

SUBCLASS TELEOSTOMI—Continued.

ORDER HAPLOMI—Continued.

Family *Paciliidae*—Continued.

Genus <i>Cyprinodon</i> , Lacépède—Continued.	
latifasciatus, Garman	676
Genus <i>Jordanella</i> , Goode & Bean	677
florida, Goode & Bean	677
Genus <i>Pseudoxiphophorus</i> , Bleeker	678
bimaculatus (Heckel)	678
Genus <i>Gambusia</i> , Poey	678
punctata, Poey	679
puncticulata, Poey	680
affinis (Baird & Girard)	680
tridentiger, Garman	2833
nobilis (Baird & Girard)	682
nicaraguensis, Günther	682
gracilis, Heckel	683; 2832
episcopi, Steindachner	683
melapleura (Gosse)	659; 2830
Genus <i>Belonesox</i> , Kner	684
belizanus, Kner	684
Genus <i>Anableps</i> (Artedi) Söpoli	684; 2807
dovii, Gill	685
Genus <i>Goodea</i> , Jordan	685
atripinnis, Jordan	685
Genus <i>Xenendum</i> , Jordan & Snyder	3151
caliente, Jordan & Snyder	3152
xaliscome, Jordan & Snyder	3153
luitpoldii, Steindachner	2832; 3152
Genus <i>Platypoecilus</i> , Günther	685
maculatus, Günther	686
quitzeoensis, B. A. Bean	2873
mentalis, Gill	686
Genus <i>Heterandria</i> , Agassiz	686
uninotata (Poey)	687
metallica (Poey)	687
formosa, Agassiz	687
pleurospilus (Günther)	688
Genus <i>Lebiasina</i> , De Filippi	689
pacificus, De Filippi	689
Genus <i>Acropocilia</i> , Hilgendorf	690
tridens (Hilgendorf)	690
Genus <i>Pecilia</i> , Bloch & Schneider	690
vivipara, Bloch & Schneider	691
butleri, Jordan	691
gillii (Kner & Steindachner)	692
vittata, Guicheon	692; 2833
mexicana, Steindachner	692
thermalis, Steindachner	693
chisoyensis, Günther	693
petenensis, Günther	694
sphenops, Cuvier & Valenciennes	694
dovii, Günther	695
couchiana (Girard)	695
bonaparti, Steindachner	695
reticulata, Peters	696; 2833
arubensis, Van Lidth de Jeude	696

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER HAPLOMI—Continued.	
<i>Family Poeciliidae—Continued.</i>	
Genus <i>Poecilia</i> , Bloch & Schneider—Continued.	
<i>dominicensis</i> , Cuvier & Valenciennes	696; 2834
<i>spilurus</i> , Günther	697
<i>elongatus</i> , Günther	697
<i>presidionis</i> , Jordan & Culver	697
<i>versicolor</i> (Günther)	688; 2833
<i>occidentalis</i> (Baird & Girard)	689; 2833
<i>enneata</i> , Garman	2834
<i>limantouri</i> , Jordan & Snyder	3153
Genus <i>Mollenkia</i> , Le Sueur	698
<i>jonesi</i> , Günther	698
<i>formosa</i> (Girard)	699
<i>latipinnna</i> , Le Sueur	699
<i>petenensis</i> , Günther	700
Genus <i>Xiphophorus</i> , Heckel	701
<i>helleri</i> , Heckel	701
<i>montezumae</i> , Jordan & Snyder	3154
<i>guntheri</i> , Jordan & Evermann	702
<i>Family Amblyopidae</i>	702
Genus <i>Chologaster</i> , Agassiz	703
<i>cornutus</i> , Agassiz	703
<i>agassizii</i> , Putnam	704
<i>papilliferus</i> , Forbes	704
Genus <i>Typhlichthys</i> , Girard	704
<i>subterraneus</i> , Girard	704
Genus <i>Troglichthys</i> , Eigenmann	3156
<i>rose</i> , Eigenmann	2835; 3156
Genus <i>Amblyopsis</i> , De Kay	706
<i>speleana</i> , De Kay	706
ORDER SYNENTOGNATHI	707
<i>Family Esocidae</i>	708
Genus <i>Tylosurus</i> , Cocco	708
<i>notatus</i> (Poey)	710
<i>scapularis</i> , Jordan & Gilbert	711
<i>timucu</i> (Walbaum)	711
<i>enryops</i> , Bean & Dresel	711
<i>diplostenia</i> (Cope)	712
<i>micros</i> (Günther)	712
<i>angusticeps</i> (Günther)	712
<i>ardeola</i> (Cuvier & Valenciennes)	713
<i>stolzmanni</i> (Steindachner)	713
<i>exilis</i> (Girard)	714
<i>marinus</i> (Walbaum)	714
<i>almeldae</i> (Quoy & Gaimard)	715
<i>fedlator</i> , Jordan & Gilbert	715
<i>raphidoma</i> (Ranzau)	715
<i>galeatus</i> (Cuvier & Valenciennes)	716
<i>pacificus</i> (Steindachner)	716
<i>acns</i> (Lacépède)	716
<i>caribbeus</i> (Le Sueur)	717
Genus <i>Athlennes</i> , Jordan & Fordice	717
<i>hlana</i> (Cuvier & Valenciennes)	718
<i>Family Hemiramphidae</i>	718
Genus <i>Chirodorbus</i> , Goode & Bean	719
<i>atherinoides</i> , Goode & Bean	719

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER SYNENTOGNATHI—Continued.	
<i>Family Hemiramphidae</i> —Continued.	
Genus <i>Hyporhamphus</i> , Gill.....	719
<i>unifasciatus</i> (Ranzani).....	720
<i>roberti</i> (Cuvier & Valenciennes).....	721
<i>rosea</i> (Jordan & Gilbert).....	721
Genus <i>Heuriacanthus</i> , Cuvier.....	722
<i>brasiliensis</i> (Linnaeus).....	722
<i>balao</i> , Le Sueur.....	723
Genus <i>Euleptorhamphus</i> , Gill.....	723
<i>velox</i> , Poey.....	724
<i>Family Scombracidae</i>	724
Genus <i>Scombrusox</i> , Lacépède	725
<i>saurus</i> (Walbaum).....	725
Genus <i>Cololabis</i> , Gill	726
<i>brevirostris</i> (Peters)	726
<i>Family Exocatidae</i>	726
Genus <i>Fodiator</i> , Jordan & Meek.....	727
<i>acutus</i> (Cuvier & Valenciennes)	728
Genus <i>Parexocetus</i> , Bleeker.....	728
<i>mesogaster</i> (Bloch).....	728
Genus <i>Exocetus</i> , Linnaeus.....	730; 2835
<i>volitans</i> , Linnaeus.....	729; 734; 2835
Genus <i>Cypsilurus</i> , Swainson	2835
<i>heterurus</i> (Rafinesque).....	735; 2836
<i>lutkeni</i> (Jordan & Evermann).....	736; 2836
<i>fmicatus</i> (Mitchill).....	737; 2836
<i>nigricans</i> (Bennett)	737; 2836
<i>xenopterus</i> (Gilbert).....	738; 2836
<i>lineatus</i> (Cuvier & Valenciennes).....	739; 2836
<i>cyanopterus</i> (Cuvier & Valenciennes).....	739; 2836
<i>bahiensis</i> (Ranzani).....	739; 2836
<i>californicus</i> (Cooper).....	740; 2836
<i>callipterus</i> (Günther).....	740; 2836
<i>gibbifrons</i> (Cuvier & Valenciennes).....	741; 2836
Genus <i>Exonantes</i> , Jordan & Evermann	2835
<i>exsiliens</i> (P. L. S. Müller).....	732; 2836
<i>rondeletii</i> (Cuvier & Valenciennes)	733; 2836
<i>vinegarae</i> (Jordan & Meek).....	734; 2836
<i>rufipinnis</i> (Cuvier & Valenciennes)	735; 2836
<i>affinis</i> (Günther).....	2836
ORDER HEMIBRANCHII.	741
<i>Family Gasterosteidae</i>	742
Genus <i>Encalia</i> , Jordan	743
<i>inconstans</i> (Kirtland)	744; 3157
<i>cayuga</i> , Jordan.....	744
<i>pygmaea</i> (Agassiz)	744
Genus <i>Pygosteus</i> , Brevoort	745
<i>pungitius</i> (Linnaeus)	745
<i>brachypoda</i> (Bean)	746
Genus <i>Gasterosteus</i> (Artedi) Linnaeus	746
<i>aculeatus</i> , Linnaeus	747
<i>bispinosus</i> , Walbaum	748
<i>atkinsii</i> (Bean)	748
<i>cuvieri</i> (Girard)	749
<i>gladiunculus</i> , Kendall.....	2836
<i>cataphractus</i> (Pallas)	749

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER HEMIBRANCHII—Continued.	
Family Gasterosteidae—Continued.	
Genus <i>Gasterostes</i> (Artedi) Linnaeus—Continued.	
williamsoni, Girard	750
microcephalus (Girard)	751
<i>Apeltes</i> , De Kay	752
quadraeans (Mitchill)	752
Family <i>Aulorhynchidae</i>	752
Genus <i>Aulorhynchus</i> , Gill	753
<i>flavidus</i> , Gill	754
Family <i>Aulostomidae</i>	754
Genus <i>Aulostomus</i> , Lacépède.....	754
<i>maculatus</i> , Valenciennes	754
<i>einereus</i> , Poey	755
Family <i>Fistulariidae</i>	755
Genus <i>Fistularia</i> , Linnaeus.....	756
<i>tabacaria</i> , Linnaeus	757
<i>depressa</i> , Günther	757
<i>petimba</i> , Lacépède	758
Family <i>Macrorhamphosidae</i>	758
Genus <i>Macrorhamphosus</i> , Lacépède	759
<i>scolopax</i> (Linnaeus)	759
ORDER LOPHOBRANCHII.....	759
SUBORDER SYNGNATHI.....	760
Family <i>Syngnathidae</i>	760
Genus <i>Siphonoma</i> , Radinesque	761
Subgenus <i>Dermatostethus</i> , Gill	763
<i>punctipinnis</i> (Gill)	763
Subgenus <i>Siphonoma</i>	763
<i>carinatum</i> , Gilbert	763
<i>californiense</i> (Storer)	764
<i>griseolineatum</i> (Ayres)	764
<i>leptorhynchum</i> (Girard)	764
<i>fistulatum</i> (Peters)	765
<i>barbare</i> , Swain	765
<i>mackayi</i> Swain & Meek	766
<i>floridae</i> , Jordan & Gilbert	766
<i>poeyi</i> , Jordan & Evermann	766
<i>aulicus</i> , Swain	767
<i>pelagicum</i> (Osbeck)	767
<i>rousseau</i> (Kaup)	767
<i>elucens</i> (Poey)	768
<i>jonesi</i> (Günther)	768
<i>robertsi</i> , Jordan & Rutter	2837
<i>sinaloae</i> , Jordan & Starks	2838
<i>brachycephalum</i> (Poey)	769
<i>affine</i> (Günther)	769
<i>scovellii</i> , Evermann & Kendall	769
<i>bairdianum</i> (Duméril)	770
<i>louisianae</i> (Günther)	770
<i>fuscum</i> (Storer)	770
<i>starksii</i> , Jordan & Cuvier	771; 2838
<i>arctum</i> , Jenkins & Evermann	771
<i>erinigerum</i> , Bean & Dresel	771
Genus <i>Corythoichthys</i> , Kaup	772; 2838
<i>albirostris</i> , Heckel	772; 2838
<i>cayennensis</i> (Sauvage)	772; 2838

	Page.
CLASS PISCES—Continued.	
SUCLASS TELEOSTOMI—Continued.	
ORDER LOPHIORHANCHII—Continued.	
SUBORDER SYGNATHI—Continued.	
<i>Family Syngnathidae</i> —Continued.	
Genus <i>Corythoichthys</i> , Kaup—Continued.	
cayorum, Evermann & Kendall	2838
Genus <i>Doryrhamphus</i> , Kaup	773
Subgenus <i>Dorylethys</i> , Kaup.....	773
<i>lineatus</i> (Valenciennes).....	773
Subgenus <i>Doryrhamphus</i>	773
<i>californiensis</i> , Gill	773
Genus <i>Syngnathus</i> , Linnaeus	774
Subgenus <i>Syngnathus</i>	774
<i>heckeli</i> (Kaup).....	774; 2839
Genus <i>Ostholox</i> , Cope	775
<i>pellucidus</i> , Cope	775
Genus <i>Hippocampus</i> , Rafinesque	775
<i>ingens</i> , Girard	776
<i>hudsonius</i> , De Kay	777
<i>punctulatus</i> , Guichenot	777
<i>stylifer</i> , Jordan & Gilbert	778
<i>zosteræ</i> , Jordan & Gilbert	778
ORDER ACANTHOPTERI	779
SUBORDER SALMOPERCÆ	782
<i>Family Percopsidae</i>	783
Genus <i>Percopsis</i> , Agassiz	783
<i>guttatus</i> , Agassiz	784
Genus <i>Columbia</i> , Eigenmann & Eigenmann	784
<i>transmontana</i> , Eigenmann & Eigenmann	784
SUBORDER XENARCHI	785
<i>Family Aphredoderidae</i>	785
Genus <i>Aphredoderus</i> , Le Sueur	786
<i>sayanus</i> (Gilliams).....	786
SUBORDER PERCESSES	787
<i>Family Atherinidae</i>	788
Genus <i>Atherina</i> (Artedi) Linnaeus	789
<i>stipes</i> , Müller & Troschel	790
<i>laticeps</i> , Poey	790
<i>area</i> , Jordan & Gilbert	790
<i>barringtonensis</i> , Goode	791
<i>carolina</i> , Cuvier & Valenciennes	791
<i>microps</i> , Poey	791
Genus <i>Chiromystus</i> , Swainson	792; 2839
<i>chapalu</i> , Jordan & Snyder	3159
<i>promelas</i> , Jordan & Snyder	3160
<i>diazii</i> , Jordan & Snyder	3161
<i>crystallinum</i> , Jordan & Snyder	3162
<i>lernae</i> , Jordan & Snyder	3163
<i>ocotlane</i> , Jordan & Snyder	3163
<i>estor</i> , Jordan	792; 2839; 3165
<i>album</i> (Steindachner)	3165
<i>humboldtianum</i> (Cuvier & Valenciennes)	793; 2839
<i>grandocule</i> (Steindachner)	2839
Genus <i>Elopsarum</i> , Jordan & Evermann	2840
<i>bartoni</i> , (Jordan & Evermann)	792; 2840
<i>jordani</i> , (Woolman)	793; 2840; 3159
<i>arge</i> , Jordan & Snyder	3158

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER Acanthopteri—Continued.	
SUBORDER PERCESOES—Continued.	
Family Atherinidae—Continued.	
Genus Kirtlandia, Jordan & Evermann.....	794
vagrans (Goode & Bean)	794
laciniiata (Swain)	795
martinica (Cuvier & Valenciennes)	795
Genus Menidia (Bonaparte) Jordan & Gilbert	796
peninsule (Goode & Bean)	797
gracilis (Günther)	797
beryllina (Cope)	797
andens, Hay	798
gilberti, Jordan & Böllman	798
sardina (Jenkins & Evermann)	799
menidia (Linnaeus)	800
notata (Mitchill)	800; 2840
chura, Evermann & Jenkins.....	801
Genus Leuresthes, Jordan & Gilbert.....	801
crameri, Jordan & Evermann	802
tennis (Ayres)	802
Genus Eurystole, Jordan & Evermann.....	802
erriacha (Jordan & Gilbert)	803
Genus Thyrina, Jordan & Culver.....	803
evermanni, Jordan & Culver	804
crystallina, Jordan & Culver	804
guatemalensis (Günther)	801; 2840
pachylepis (Günther)	801; 2840
Genus Atherinella, Steindachner	805
panamensis, Steindachner	805
Genus Labidesthes, Cope	805
sicculus (Cope)	805
Genus Atherinopsis, Girard	806
californiensis, Girard	806
Genus Atherinops, Steindachner	807
insularum, Gilbert	807
affinis (Ayres)	807
regis, Jenkins & Evermann	808
Family Mugilidae.....	808
Genus Mugil (Arteci) Linnaeus.....	809
brasiliensis, Agassiz	810
cephalus, Linnaeus	811
incilis, Hancock	812
thoburni, Jordan & Starks	812
enrema, Cuvier & Valenciennes	813
hospe, Jordan & Culver	814
gainardianus, Desmarest	814
setosus, Gilbert	815
trichodon, Poey	816
Genus Chenomugil, Gill.....	816
proboscideus (Günther)	816
Genus Querimana, Jordan & Gilbert	817
harengus (Günther)	817
gyrans, Jordan & Gilbert	818
Genus Agonostomus, Bennett	818
Subgenus Dajaus, Cuvier & Valenciennes	819
percoides, Günther	819

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER PERCESOCES—Continued.	
<i>Family Mugilidae</i> —Continued.	
Genus <i>Agonostomus</i> , Bennett—Continued.	
Subgenus <i>Dajaus</i> , Cuvier & Valenciennes—Continued.	
<i>monticola</i> (Bancroft)	819
<i>nasutus</i> , Günther	819
<i>microps</i> , Günther	820
Genus <i>Joturus</i> , Poey	820
<i>pichardi</i> , Poey	821
Genus <i>Neomugil</i> , Vaillant	3165
<i>digiueti</i> , Vaillant	3165
<i>Family Sphyraenidae</i>	822
Genus <i>Sphyraena</i> (Arvedi) Bloch & Schneider	822
<i>barracuda</i> (Walbaum)	823; 2841
<i>ensis</i> , Jordan & Gilbert	824
<i>guachancho</i> , Cuvier & Valenciennes	824
<i>piendilla</i> , Poey	824
<i>borealis</i> , De Kay	825
<i>argentea</i> , Girard	826
<i>sphyraena</i> (Linnaeus)	826
SUBORDER RHEGNOPTERI	827
<i>Family Polynemidae</i>	827
Genus <i>Polynemus</i> (Gronow) Linnaeus	828
<i>quinquarius</i> , Linnaeus	828
Genus <i>Polydactylus</i> , Lacépède	828
<i>approximans</i> (Lay & Bennett)	829
<i>virginicus</i> (Linnaeus)	829
<i>octonemus</i> (Girard)	830
<i>opercularis</i> (Gill)	830
Group <i>Ammodytoidae</i>	831
<i>Family Ammodytidae</i>	831
Genus <i>Ammodytes</i> (Arvedi) Linnaeus	832
<i>dubius</i> , Reinhardt	832
<i>americanus</i> , De Kay	833
<i>personatus</i> , Girard	833
Group <i>Berycoidei</i>	833
Genus <i>Rhynchias</i> , Gill	2841
<i>setipinnis</i> (Pallas)	2841
Group <i>Berycoidei</i>	833
<i>Family Bathylupeidae</i>	834
Genus <i>Bathylaea</i> , Alcock	834
<i>argentea</i> , Goode & Bean	835
<i>Family Stephanoberycidae</i>	835
Genus <i>Stephanobryx</i> , Gill	836
<i>mone</i> , Gill	836
<i>gillii</i> , Goode & Bean	836
<i>Family Trachichthyidae</i>	836
Genus <i>Hoplostethus</i> , Cuvier & Valenciennes	837
<i>mediterraneus</i> , Cuvier & Valenciennes	837
<i>Family Berycidae</i>	837
Genus <i>Caulolepis</i> , Gill	838
<i>longidens</i> , Gill	839
Genus <i>Anoplogaster</i> , Günther	839
<i>cornutus</i> (Cuvier & Valenciennes)	840
Genus <i>Poromitra</i> , Goode & Bean	840
<i>capito</i> , Goode & Bean	840

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER RHEGNOPTERI—Continued.	
Family <i>Berycidae</i> —Continued.	
Genus <i>Plectromus</i> , Gill	840
<i>sueborbitalis</i> , Gill	841
<i>lugubris</i> (Gilbert)	842
<i>beanii</i> (Günther)	842
<i>crassiceps</i> , Günther	843
<i>eristiciceps</i> (Gilbert)	843
Genus <i>Beryx</i> , Cuvier	844
<i>decadactylus</i> , Cuvier & Valenciennes	844
<i>splendens</i> , Lowe	844
Family <i>Holocentridae</i>	845
Genus <i>Myripristis</i> , Cuvier	846
Subgenus <i>Ostichthys</i> (Langsdorf)	846
<i>trachypoma</i> , Günther	846
Subgenus <i>Myripristis</i>	846
<i>jacobus</i> , Cuvier & Valenciennes	846
<i>occidentalis</i> , Gill	847
<i>paeциlopus</i> (Gill)	847
<i>clarionensis</i> , Gilbert	2842
Genus <i>Holocentrus</i> (Gronow) Scopoli	847
<i>ascensionis</i> (Osbeck)	848
<i>rufus</i> (Walbaum)	849
<i>slecker</i> , Cope	849
<i>suborbitalis</i> , Gill	850
<i>cornutus</i> , Poey	851
<i>brachypterus</i> , Poey	852
<i>vexillarius</i> , Poey	852
<i>osculus</i> , Poey	853
<i>sancti-pauli</i> , Günther	853
Genus <i>Flammeo</i> , Jordan & Evermann	852; 2871
<i>marianus</i> (Cuvier & Valenciennes)	852; 2871
Genus <i>Plectrypops</i> , Cill	853
<i>retrospinis</i> (Guichenot)	853
Family <i>Polymixiidae</i>	854
Genus <i>Polymixia</i> , Lowe	854
<i>lowei</i> , Günther	854
Family <i>Mullidae</i>	855
Genus <i>Mullus</i> , Linnaeus	856
<i>auratus</i> , Jordan & Gilbert	856
Genus <i>Upeneus</i> , Cuvier	857; 2843
<i>rathbuni</i> , Evermann & Jenkins	857
<i>maculatus</i> (Bloch)	858
<i>dentatus</i> , Gill	859
<i>parvus</i> , Poey	859
<i>martiniens</i> , Cuvier & Valenciennes	859
<i>xanthogrammus</i> , Gilbert	860
<i>grandisquamis</i> , Gill	860
Group <i>Scombroidei</i>	860
Family <i>Scombridae</i>	863
Genus <i>Scomber</i> (Arvedi) Linnaeus	865
Subgenus <i>Scomber</i>	865
<i>scombrus</i> , Linnaeus	865
Subgenus <i>Pneumatophorus</i> , Jordan & Gilbert	866
<i>coilas</i> , Gmelin	866

	Page.
840	
841	
842	
842	
843	
843	
844	
844	
844	
845	
845	
846	
846	
846	
846	
846	
846	
846	
847	
847	
847	
847	
847	
847	
847	
847	
848	
849	
849	
850	
851	
852	
852	
853	
853	
2871	
2871	
853	
853	
854	
854	
855	
856	
856	
2843	
857	
858	
859	
859	
859	
860	
860	
860	
860	
860	
861	
862	
862	
862	
862	
863	
863	
865	
865	
865	
866	
866	
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER RHEGNOPTERI—Continued.	
<i>Family Scombridae</i> —Continued.	
Genus <i>Auxis</i> , Cuvier.....	867
<i>thazard</i> (Lacépède).....	867
Genus <i>Gymnosarda</i> , Gill	868
<i>pelanis</i> (Linnaeus)	868
<i>alleterata</i> (Rafinesque).....	869
Genus <i>Thunnus</i> , South.....	869
<i>thynnus</i> (Linnaeus).....	870
Genus <i>Gerimo</i> , Jordan.....	870
<i>alalunga</i> (Gmelin)	871
Genus <i>Sarda</i> , Cuvier.....	871
<i>sarda</i> (Bloch).....	872
<i>chilensis</i> , Cuvier & Valenciennes.....	872
Genus <i>Scomberomorus</i> , Lacépède	873
<i>concolor</i> (Lockington).....	873
<i>maculatus</i> (Mitchilli)	874
<i>sierra</i> , Jordan & Starks.....	874
<i>regalis</i> (Bloch)	875
<i>cavalla</i> (Cuvier & Valenciennes)	875
Genus <i>Acanthocybium</i> , Gill.....	876
<i>solandri</i> (Cuvier & Valenciennes).....	876
<i>Family Gempylidae</i>	877
Genus <i>Escarlar</i> , Jordan & Evermann	878; 2843
<i>violaceus</i> (Bean).....	878; 2843
Genus <i>Ruvettus</i> , Cocco	879
<i>pretiosus</i> , Cocco	879
Genus <i>Epinnula</i> , Poey	880
<i>magistralis</i> , Poey	880
Genus <i>Neolotus</i> , Johnson	881
<i>tripes</i> , Johnson	881
Genus <i>Promethichthys</i> , Gill	882
<i>prometheus</i> , Cuvier & Valenciennes	882
<i>parvipinnis</i> (Goode & Bean).....	883
Genus <i>Genypterus</i> , Cuvier & Valenciennes	883
<i>serpens</i> , Cuvier & Valenciennes	884
<i>Family Lepidopidae</i>	884
Genus <i>Aphanopus</i> , Lowe	885
<i>mlnor</i> , Collett	885
Genus <i>Evoxymetopon</i> , Poey	885
<i>teniatus</i> , Poey	886
Genus <i>Lepidopus</i> , Gouan	886
<i>xantusi</i> , Goode & Bean	886; 2843
Genus <i>Benthodesmus</i> , Goode & Bean	887
<i>atlanticus</i> , Goode & Bean	887
<i>Family Trichiuridae</i>	888
Genus <i>Trichiurus</i> , Linnaeus	889
<i>lepturus</i> , Linnaeus	889
<i>Family Istiophoridae</i>	890
Genus <i>Istiophorus</i> , Lacépède	890
<i>nigricans</i> (Lacépède).....	891
Genus <i>Tetrapturus</i> , Rafinesque	891
<i>imperator</i> (Büch & Schneider).....	892
<i>ampius</i> , Poey	892

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER RHEGNOPTERI—Continued.	
<i>Family Xiphidae</i>	893
Genus <i>Xiphias</i> , Linneus.....	893
<i>gladius</i> , Linneus.....	894
<i>Family Nematiidae</i>	894
Genus <i>Nematus</i> , Gill	895
<i>pectoralis</i> , Gill.....	895
<i>Family Carangidae</i>	895
Genus <i>Oligoplites</i> , Gill.....	898
<i>saurus</i> (Bloch & Schneider)	898
<i>saliens</i> (Bloch).....	899
<i>palometa</i> (Cuvier & Valenciennes)	899
<i>cetus</i> (Günther).....	899
<i>mundus</i> , Jordan & Starks	2844
Genus <i>Nauferus</i> , Rafinesque	900
<i>ductor</i> , Linneus	900
Genus <i>Seriola</i> , Cuvier.....	901
<i>dorsalis</i> (Gill).....	902
<i>zonata</i> (Mitchill).....	902
<i>carolinensis</i> , Holbrook.....	902
<i>lalandi</i> , Cuvier & Valenciennes	903
<i>dumerilli</i> (Risso)	903
Subgenus <i>Zonichthys</i> , Swainson.....	904
<i>mazatlana</i> , Steindachner.....	904
<i>fasciata</i> (Bloch).....	904
<i>rivoliana</i> Cuvier & Valenciennes.....	904
<i>falcata</i> , Cuvier & Valenciennes	905
Genus <i>Elagatis</i> , Bennett.....	906
<i>bipinnulatus</i> (Quoy & Gaimard)	906
Genus <i>Decapterus</i> , Bleeker	907
<i>punctatus</i> (Agassiz)	907
<i>scombrinus</i> (Valenciennes).....	908
<i>sanctae-helene</i> (Cuvier & Valenciennes).....	908
<i>hypodus</i> , Gill	908
<i>macarellus</i> (Cuvier & Valenciennes)	909
Genus <i>Trachurus</i> , Rafinesque	909
<i>symmetricus</i> (Ayres).....	909; 2844
<i>trachurus</i> (Linneus).....	910
Genus <i>Trachurops</i> , Gill.....	911
<i>crumenophthalmus</i> (Bloch)	911
Genus <i>Hemicarax</i> , Bleeker	912
<i>amblyrhynchus</i> (Cuvier & Valenciennes).....	912
<i>falcatus</i> (Holbrook)	912; 2845
<i>zelotes</i> , Gilbert	2845
<i>atrimanus</i> (Jordan & Gilbert)	913
<i>secundus</i> (Poey)	914
<i>furthii</i> (Steindachner)	914
<i>leucurus</i> (Günther)	914
Genus <i>Caranx</i> , Lacépède	915
Subgenus <i>Selar</i> , Bleeker	918
<i>vinctus</i> , Jordan & Ghoert	918
Subgenus <i>Caranx</i>	919
<i>ruber</i> (Bloch)	919
<i>bartholomaei</i> , Cuvier & Valenciennes	919
Subgenus <i>Tricropodus</i> Rafinesque.....	920
<i>hippos</i> (Linnaeus)	920

Page.
.. 893
.. 893
.. 894
.. 894
.. 895
.. 895
.. 895
.. 895
.. 896
.. 896
.. 897
.. 897
.. 898
.. 898
.. 899
.. 899
.. 899
.. 899
.. 899
.. 899
.. 900
.. 900
.. 900
.. 901
.. 901
.. 902
.. 902
.. 902
.. 903
.. 903
.. 904
.. 904
.. 904
.. 904
.. 905
.. 905
.. 906
.. 906
.. 907
.. 907
.. 908
.. 908
.. 908
.. 909
.. 909
.. 909
.. 910
.. 911
.. 911
.. 912
.. 912
.. 912
.. 913
.. 913
.. 914
.. 914
.. 914
.. 915
.. 915
.. 918
.. 918
.. 919
.. 919
.. 919
.. 920
.. 920
.. 920
.. 920

CLASS PISCES—Continued.

SUBCLASS TELEOSTOMI—Continued.

ORDER ACANTHOPTERI—Continued.

SUPERORDER RHEGNOPTERI—Continued.

Family Carangidae—Continued.

Genus Caranx, Lacépède—Continued.

Subgenus Paratracetus, Gill	921
cryson (Mitchill)	921
pisquetus, Cuvier & Valenciennes	921; 2846
caballus (Günther)	921
Subgenus Carangichthys, Bleeker	922
marginatus, Gill	922
latus, Agassiz	923
medusicola, Jordan & Starks	924
lugubris, Poey	924
melampygus, Cuvier & Valenciennes	925
Subgenus Uraspis, Bleeker	926
guara (Bonnaterre)	926
Genus Gnathanedon, Bleeker	927
speciosus (Forskal)	928
Genus Carangooides, Bleeker	928
orthogrammus (Jordan & Gilbert)	928
Genus Citula, Cuvier	929
dorsalis (Gill)	930
Genus Alectis, Rafinesque	931
ciliaris (Bloch)	931
Genus Hynnis, Cuvier	932
eubensis (Poey)	932
hopkinsi, Jordan & Starks	933
Genus Vomer, Cuvier & Valenciennes	933
dorsalis, Gill	934
setipinnis (Müller)	934
spixii (Swainson)	2846
gabonensis, Guichenot	934
Genus Selene, Lacépède	935
terstedii, Lütken	935
vomer (Linnaeus)	936
Genus Chloroscombrus, Girard	937
orqueia, Jordan & Gilbert	937
chrysurus (Linnaeus)	938
ectenurus, Jordan & Osgood	2847
Genus Trachinotus, Lacépède	939
glauces (Bloch)	940
rhodopus, Gill	941
falcatus (Linnaeus)	941
rhomboides (Bloch)	2847
culveri, Jordan & Starks	942
kennedyi, Steindachner	942
goodei, Jordan & Evermann	943
argenteus, Cuvier & Valenciennes	944
carolinus (Linnaeus)	944
paloma, Jordan & Starks	945
cayennensis, Cuvier & Valenciennes	945
Genus Zalocys, Jordan & McGregor	2848
stilbe, Jordan & McGregor	2848
Family Pomatomidae	945
Genus Pomatomus, Lacépède	946
saltatrix (Linnaeus)	946

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUPERORDER RHEOOPTERI—Continued.	
<i>Family Rachycentridae</i>	947
Genus <i>Rachycentron</i> , Kaup	948
<i>canadus</i> (Linnaeus)	948
<i>Family Nomeidae</i>	948
Genus <i>Nomens</i> , Cuvier	949
<i>gronovii</i> (Gmelin)	949
Genus <i>Psenes</i> , Cuvier & Valenciennes	950
<i>pellucidus</i> , Littken	950
<i>cyanophrys</i> , Cuvier & Valenciennes	950
<i>maculatus</i> , Littken	951
<i>regulus</i> , Poey	951
<i>Family Coryphaenidae</i>	951
Genus <i>Coryphaena</i> , Linnaeus	952
<i>hippurus</i> , Linnaeus	952
<i>equisetis</i> , Linnaeus	953
<i>Family Lampridae</i>	953
Genus <i>Lampris</i> , Retzius	954
<i>luna</i> (Gmelin)	954
<i>Family Pteraclidae</i>	955
Genus <i>Pteraclis</i> , Gronow	955
<i>carolinus</i> , Cuvier & Valenciennes	956
<i>Family Bramidae</i>	956
Genus <i>Thraetes</i> , Lowe	957
<i>sauvagii</i> (Lunel)	957
Genus <i>Brama</i> , Bloch & Schneider	958
<i>agassizii</i> , Poey	959
<i>brevirostris</i> , Poey	959
<i>ratti</i> (Bloch)	959
<i>Family Steinegeriidae</i>	960
Genus <i>Steinegeria</i> , Jordan & Evermann	960
<i>rubescens</i> , Jordan & Evermann	961
<i>Family Stromateidae</i>	964
Genus <i>Centrolophus</i> , Lacépède	962; 3166
<i>niger</i> (Gmelin)	963
Genus <i>Palinurichthys</i> , Bleeker	963
<i>perciformis</i> (Mitchill)	964
Genus <i>Peprilus</i> , Cuvier	965; 3197
<i>parr</i> (Linnaeus)	965
<i>xanthurus</i> (Quoy & Gaimard)	966
Genus <i>Palometa</i> , Jordan & Evermann	966; 2849
<i>palometa</i> (Jordan & Bollman)	966
<i>medius</i> (Peters)	967
<i>simillimus</i> (Ayres)	967
Genus <i>Torontos</i> , Gill	967; 2849
<i>tr. acanthus</i> (Peck)	967; 2849
<i>Family Icosteidae</i>	968
Genus <i>Ieichthys</i> , Jordan & Gilbert	969
<i>lockingtoni</i> , Jordan & Gilbert	969
Genus <i>Schedophilus</i> , Cocco	970
<i>medusophagus</i> , Cocco	970
Genus <i>Icostenus</i> , Lockington	972
<i>enigmaticus</i> , Lockington	972

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER RHEGNOPTERI—Continued.	
<i>Family Aerotidae</i>	973; 2849
Genus <i>Aerotus</i> , Beau	973
willoughbyi, Beau	973
<i>Family Zaproridae</i>	2849
Genus <i>Zaprorus</i> , Jordan	2850
silenus, Jordan	2850
<i>Family Grammicolepididae</i>	973
Genus <i>Grammicolepis</i> , Poey	974
brachi, senilis, Poey	974
<i>Family Tetragonuridae</i>	975
Genus <i>Tetragonurus</i> , Risso	975
cuvieri, Risso	976
<i>Family Pempheridae</i>	977
Genus <i>Pempheris</i> , Cuvier & Valenciennes	977
mexicanus, Cuvier & Valenciennes	978
schomburgkii, Müller & Troschel	978
mulleri, Poey	978
poeyi, Beau	979
Group <i>Percoldea</i>	979
<i>Family Elassomidae</i>	981
Genus <i>Elassoma</i> , Jordan	982
zonatum, Jordan	982
evergladel, Jordan	982
<i>Family Centrarchidae</i>	984
Genus <i>Pomoxis</i> , Rafinesque	986
annularis, Rafinesque	987
sparoides (Lacépède)	987
Genus <i>Centrarchus</i> , Cuvier & Valenciennes	988
macropterus (Lacépède)	988
Genus <i>Acantharchus</i> , Gill	989
pomotis (Baird)	989
Genus <i>Ambloplites</i> , Rafinesque	989
rupestris (Rafinesque)	990
cauvifrons, Cope	990
Genus <i>Archoplites</i> , Gill	990
interruptus (Girard)	991
Genus <i>Chienobryttus</i> , Gill	991
gulosus (Cuvier & Valenciennes)	992
Genus <i>Enneacanthus</i> , Gill	992
obesus (Baird)	993
gloriosus (Holbrook)	993
Genus <i>Mesogonistius</i> , Gill	994
chaetodon (Baird)	995
Genus <i>Apomotis</i> , Rafinesque	995
cyanellus (Rafinesque)	996
ischyrus (Jordan & Nelson)	997
phenax (Cope & Jordan)	997
punctatus (Cuvier & Valenciennes)	997
symmetricus (Forbes)	998
Genus <i>Lepomis</i> , Rafinesque	999
Subgenus <i>Lepomis</i>	1001
auritus (Linnaeus)	1001
solis (Cuvier & Valenciennes)	1001
miniatus, Jordan	1002

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER RHEGNOPTERI—Continued.	
<i>Family Centrarchidae—Continued.</i>	
Genus <i>Lepomis</i> , Rafinesque—Continued.	
Subgenus <i>Xenotis</i> , Jordan.....	1002
<i>garmani</i> , Forbes	1002
<i>megalotis</i> (Rafinesque).....	1002
Subgenus <i>Helioperca</i> , Jordan.....	1004
<i>humilis</i> (Girard)	1004
<i>haplognathus</i> , Cope	1004
<i>macrochirus</i> , Rafinesque	1005
<i>pallidus</i> (Mitchill)	1005
Genus <i>Eupomotis</i> , Gill & Jordan.....	1006
Subgenus <i>Xystropilus</i> , Jordan.....	1006
<i>pallidus</i> (Agassiz).....	1006
Subgenus <i>Eupomotis</i>	1007
<i>heros</i> (Baird & Girard)	1007
<i>holbrooki</i> (Cuvier & Valenciennes)	1008
<i>euryorus</i> (McKay)	1008
<i>gibbosus</i> (Linnaeus).....	1009
Genus <i>Micropterus</i> , Lacépède	1010
<i>dolomieu</i> , Lacépède	1011
<i>salmoïdes</i> (Lacépède)	1012
<i>Family Kuhliidae</i>	1013
Genus <i>Kuhlia</i> , Gill.....	1013
<i>arge</i> , Jordan & Bollman	1014
<i>xenura</i> (Jordan & Gilbert)	1015
<i>Family Percidae</i>	1015
Genus <i>Stizostedion</i> , Rafinesque	1020
Subgenus <i>Stizostedion</i>	1021
<i>vitreum</i> (Mitchill)	1021
Subgenus <i>Cynopercaria</i> , Gill & Jordan.....	1022
<i>canadense</i> (Smith)	1022
<i>griseum</i> (De Kay).....	1022
<i>boreum</i> (Girard)	1022
Genus <i>Perea</i> (Artedi) Linnaeus	1023
<i>flavescens</i> (Mitchill)	1023
Genus <i>Percina</i> , Haldeman	1024
<i>rex</i> , Jordan & Evermann	1025
<i>caprodes</i> (Rafinesque)	1026
<i>zebra</i> (Agassiz)	1027
Genus <i>Hydropterus</i> , Agassiz	1028
Subgenus <i>Alvordius</i> , Girard	1030
<i>phoxocephalus</i> (Nelson).....	1030
<i>macrocephalus</i> (Cope).....	1031
<i>maculatus</i> (Girard)	1031
<i>aspro</i> (Cope & Jordan)	1032
<i>guntheri</i> (Eigenmann & Eigenmann)	1033
<i>pelatus</i> (Stauffer)	1034
<i>onachitus</i> (Jordan & Gilbert)	1035
<i>roanoka</i> (Jordan & Jenkins)	1036
Subgenus <i>Ericosma</i> , Jordan & Copeland	1030
<i>eviles</i> (Jordan & Copeland)	1030
Subgenus <i>Serraria</i> , Gilbert	1037
<i>scierus</i> , Swain	1037
<i>serrula</i> , Jordan & Gilbert	1038
<i>maxinkuckiensis</i> , Evermann	3166

Systematic Arrangement.

LI

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER RHEGNOPTERI—Continued.	
Family <i>Percidae</i> —Continued.	
Genus <i>Hadropterus</i> , Agassiz—Continued.	
Subgenus <i>Hadropterus</i>	1038
<i>nigrofasciatus</i> , Agassiz	1038
<i>Geius Hypophomus</i> , Cope	1039
Subgenus <i>Swainia</i> , Jordan & Evermann.....	1040
<i>squamatus</i> (Gilbert & Swain).....	1040
Subgenus <i>Hypophomus</i>	1040
<i>aurantiacus</i> (Cope)	1040
<i>cymatotenia</i> (Gilbert & Meek)	1041
<i>niangue</i> (Gilbert & Meek)	1042
<i>spilotus</i> (Gilbert)	1043
<i>Genus Cottogaster</i> , Putnam	1044
Subgenus <i>Cottogaster</i>	1044
<i>uranidea</i> (Jordan & Gilbert)	1044
<i>copelandi</i> (Jordan)	1045
Subgenus <i>Inostoma</i> , Jordan	1046
<i>shumardi</i> (Girard)	1046
<i>cheneyi</i> , Evermann & Kendall	2851
<i>Genus Ulocentrus</i> , Jordan	1047
<i>stigmata</i> (Jordan)	1047
<i>gilberti</i> , Evermann & Thoburn	1049
<i>meadie</i> , Jordan & Evermann	2852
<i>verecunda</i> (Jordan & Evermann)	1049
<i>histrio</i> (Jordan & Gilbert)	1050
<i>simotera</i> (Cope)	1051
<i>phlox</i> (Cope)	1052
<i>Genus Diplesion</i> , Rafinesque	1052
<i>blennioides</i> (Rafinesque)	1053
<i>Genus Boleosoma</i> , De Kay	1054
<i>longimanus</i> (Jordan)	1054
<i>podostemone</i> (Jordan & Jenkins)	1055
<i>nigrum</i> (Rafinesque)	1056
<i>olmstedi</i> (Storer)	1057
<i>effulgens</i> (Girard)	1058
<i>vexillare</i> (Jordan)	1058
<i>maculaticeps</i> (Cope)	1058
<i>mesatum</i> (Cope)	1059
<i>susane</i> (Jordan & Swain)	1059
Subgenus <i>Vaillantia</i> , Jordan	1060
<i>camurum</i> , Forbes	1060
<i>Genus Crystallaria</i> , Jordan & Gilbert	1060
<i>asprella</i> (Jordan)	1061
<i>Genus Ammocrypta</i> , Jordan	1061
<i>pellucida</i> (Baird)	1062
<i>clara</i> (Jordan & Meek)	1063
<i>vivax</i> (Hay)	1063
<i>beamii</i> , Jordan	1064
<i>Genus Ioa</i> , Jordan & Brayton	1064
<i>vitrea</i> (Cope)	1064
<i>vlgil</i> , Hay	1065
<i>Genus Etheostoma</i> , Rafinesque	1066
Subgenus <i>Poecilichthys</i> , Agassiz	1069
<i>variatum</i> , Kirtland	1069

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUPERORDER RHEGNOPTERI—Continued.	
<i>Family Percidae</i> —Continued.	
Genus Etheostoma, Rafinesque—Continued.	
Subgenus <i>Nanostoma</i> , Putnam	1070
swannanum, Jordan & Evermann	1070
thalassinum (Jordan & Brayton)	1071
inscriptum (Jordan & Brayton)	1072
blennius, Gilbert & Swain	1072
rupestre, Gilbert & Swain	1073
elegans (Hay)	1074
zonale (Cope)	1075
arcansum, Jordan & Gilbert	1075
Subgenus <i>Nothonotus</i> , Agassiz	1076
camurum (Cope)	1076
vulgaratum (Cope)	1077
maculatum, Kirkland	1077
cinerum, Storer	1078
tessellatum, Storer	1078
rufulineatum (Cope)	1079
jordani, Gilbert	1079
Subgenus <i>Torrentaria</i> , Jordan & Evermann	1080
sagitta, Jordan & Swain	1080
australe, Jordan	1081
Subgenus <i>Nvicola</i> , Jordan & Evermann	1082
boreale (Jordan)	1082
Subgenus <i>Rafinesquius</i> , Jordan & Evermann	1082
pottsi (Girard)	1082
Subgenus <i>Oligocephalus</i> , Girard	1083
lowei, Jordan & Meek	1083
aubeeanbei, Evermann	1084
jessiae (Jordan & Brayton)	1084
interiorinatum, Gilbert & Swain	1086
lepidogenys, Evermann & Kendall	1087
carolinum, Storer	1088
spectabile (Agassiz)	1089
lepidum (Balrd & Girard)	1089
tippecanoe, Jordan & Evermann	1090
punctulatum (Agassiz)	1090
cragini, Gilbert	1091
obeyense, Kirsch	1092
pagel, Meek	1092
virgatum (Jordan)	1093
Subgenus <i>Claricola</i> , Jordan & Evermann	1093
julie, Meek	1093
astesiae (Hay)	1094
alabamae (Gilbert & Swain)	1095
whipplei (Girard)	1095
squamiceps, Jordan	1096
Subgenus <i>Etheostoma</i>	1097
flabellare, Rafinesque	1097
cumberlandicum, Jordan & Swain	1098
lineolatum (Agassiz)	1098
Genus <i>Alvarius</i> , Girard	1099
lateralis, Girard	1099
Genus <i>Psychromaster</i> , Jordan & Evermann	1099
tuscumbia (Gilbert & Swain)	1100

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER RHEGNOPTERI—Continued.	
Family <i>Percidae</i> —Continued.	
Genus <i>Copelandellus</i> , Jordan & Evermann.....	1100
<i>quiescens</i> (Jordan).....	1100
Genus <i>Boleichthys</i> , Girard.....	1101
<i>fusiformis</i> (Girard)	1101
<i>exilis</i> , Girard.....	1103
Genus <i>Micropercia</i> , Putnam	1103
<i>praelaris</i> , Hay	1103
<i>punctulata</i> , Putnam.....	1104
<i>fonticola</i> (Jordan & Gilbert)	1104
Family <i>Cheilodipteridae</i>	1105
Genus <i>Apogon</i> Lacépède.....	1106
<i>imberbis</i> (Linneus).....	1107
<i>dovii</i> , Günther	1108
<i>retroscilla</i> (Gill)	1108
<i>maculatus</i> (Poey)	1109
<i>binotatus</i> (Poey)	1109
<i>pigmentarius</i> (Poey).....	1109
<i>atricaudus</i> , Jordan & McGregor.....	2853
Genus <i>Apogonichthys</i> , Bleeker	1110
<i>alutatus</i> (Jordan & Gilbert)	1110
<i>stellatus</i> , Cope	1110
<i>puncticulatus</i> , Poey.....	1111
Genus <i>Glossamia</i> , Gill	1111
<i>pandionis</i> (Goode & Bean)	1111
Genus <i>Epigonus</i> , Rafinesque	1111
<i>occidentalis</i> , Goode & Bean	1112
Genus <i>Cheilodipterus</i> , Lacépède	1112
<i>affinis</i> , Poey	1113
Genus <i>Amia</i> Ichthys, Poey	1113
<i>dapterus</i> (Poey).....	1113
Genus <i>Sphyraenops</i> , Gill	1114
<i>bairdianus</i> , Poey	1114
Genus <i>Scombrops</i> , Temminck & Schlegel	1114
Subgenus <i>Latebrus</i> , Poey	1114
<i>oculatus</i> (Poey).....	1114
Genus <i>Hypoclydonia</i> , Goode & Bean	1115
<i>bella</i> , Goode & Bean	1115
Family <i>Centropomidae</i>	1116
Genus <i>Centropomus</i> , Lacépède.....	1117
<i>viridis</i> , Lockington	1118
<i>undecimalis</i> (Bloch)	1118
<i>nigrescens</i> , Günther	1119
<i>pedimacula</i> , Poey.....	1119
<i>grandoculatus</i> , Jenkius & Evermann.....	1120
<i>cuvieri</i> , Bocourt	1121
<i>mexicanus</i> , Bocourt	1121
<i>parallelus</i> , Poey	1122
<i>pectinatus</i> , Poey	1122
<i>unionensis</i> , Bocourt	1122
<i>armatus</i> , Gill	1123
<i>robalito</i> , Jordan & Gilbert	1123
<i>ensiferus</i> , Poey.....	1123; 2853

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER XENOPTERI—Continued.	
<i>Family Serranidae</i>	1126
Genus <i>Roccus</i> , Mitchell	1131
Subgenus <i>Lepibrama</i> , Rafinesque	1132
<i>chrysops</i> (Rafinesque).....	1132
Subgenus <i>Roccus</i>	1132
<i>lineatus</i> (Bloch)	1132
Genus <i>Morone</i> , Mitchell.....	1133
<i>interrupta</i> , Gill	1134
<i>americana</i> (Gmelin).....	1134
Genus <i>Liopropoma</i> , Gill	1135
<i>aberrans</i> (Poey).....	1136
Genus <i>Chorististium</i> , Gill.....	1136
<i>ruberum</i> (Poey).....	1136
Genus <i>Stereolepis</i> , Ayres	1137
<i>gigas</i> , Ayres	1137
Genus <i>Polypyron</i> , Cuvier.....	1138
<i>americanus</i> (Bloch & Schneider).....	1139
Genus <i>Gonioplectrus</i> , Gill.....	1139
<i>hispanus</i> (Cuvier & Valenciennes).....	1140
Genus <i>Petrometopon</i> , Gill.....	1140
<i>panamensis</i> (Steindachner)	1141
<i>cruentatus</i> (Laëpède)	1141
<i>coronatus</i> (Cuvier & Valenciennes)	1142
Genus <i>Bodianus</i> , Bloch	1143
<i>teniops</i> (Cuvier & Valenciennes)	1144
<i>fulvus</i> (Linnaeus)	1144
<i>ruber</i> (Bloch & Schneider)	1145
<i>punctatus</i> (Linnaeus)	1146
Subgenus <i>Menophorus</i> , Poey	1146
<i>dubius</i> (Poey).....	1146
<i>punctiferus</i> (Poey).....	1147
Subgenus <i>Euneistus</i> , Jordan & Evermann	1147
<i>acanthistius</i> (Gilbert).....	1147
Genus <i>Epinephelus</i> , Bloch	1148
Subgenus <i>Schistorus</i> , Gill.....	1151
<i>mystacinus</i> (Poey)	1151
Subgenus <i>Epinophelus</i>	1152
<i>analogus</i> , Gill	1152
<i>adscensionis</i> (Osbeck)	1152
<i>guazza</i> (Linnaeus)	1154
<i>labriformis</i> (Jenyns)	1155
<i>flavolimbatus</i> , Poey	1155
<i>niphobles</i> , Gilbert & Starks.....	2853
<i>niveatus</i> (Cuvier & Valenciennes)	1156
<i>striatus</i> (Bloch).....	1157
<i>guttatus</i> (Linnaeus)	1158; 3197
<i>drummond-hayi</i> , Goode & Bean	1159
<i>morio</i> (Cuvier & Valenciennes)	1160
Genus <i>Garrupa</i> , Jordan	1161
<i>nigrita</i> (Holbrook)	1161
Genus <i>Promicrops</i> (Gill) Poey	1162
<i>itaiara</i> (Lichtenstein)	1162; 3197
Genus <i>Alphestes</i> , Bloch & Schneider.....	1164
<i>chloropterus</i> (Cuvier & Valenciennes)	1164; 2854
<i>multiguttatus</i> (Günther)	1165

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER RHEGNOPTERI—Continued.	
Family Serranidae—Continued.	
Genus Dermatolepis, Gill	1166
Subgenus Lioperca, Gill.....	1167
<i>inermia</i> (Cuvier & Valenciennes)	1167
<i>zanculus</i> , Evermann & Kendall	2854
Subgenus Dermatolepis	1168
<i>punctatus</i> , Gill.....	1168
Genus Mycteroperca, Gill.....	1169
Subgenus Archopercia, Jordan & Evermann.....	1171
<i>boulengeri</i> , Jorlax & Starks	1171
Subgenus Triostropis, Gill.....	1172
<i>venenosa</i> (Linnaeus)	1172
<i>apua</i> (Bloch)	1173
<i>bonaei</i> (Poey)	1174
<i>xanthosticta</i> , Jordan & Swain.....	1176
<i>jordani</i> (Jenkins & Evermann)	1176
<i>microlepis</i> (Goode & Bean)	1177
<i>interstitialis</i> (Poey).....	1178
<i>dimidiata</i> (Poey).....	1179
<i>xenarha</i> , Jordan.....	1180
Subgenus Parepinephelua, Bleeker.....	1180
<i>rubra</i> (Bloch)	1180
Subgenus Xystroperca, Jordan & Evermann.....	1181
<i>pardalis</i> , Gilbert	1181
Subgenus Mycteroperca	1183
<i>olfax</i> (Jenyns)	1183
<i>ruberrima</i> , Jordan & Bollman	1183
<i>rosacea</i> (Streets)	1184
<i>falcata</i> (Poey).....	1184
<i>phenax</i> , Jordan & Swain	1185
<i>venadorum</i> , Jordan & Starks.....	1186
<i>callifura</i> , Poey	1186
<i>hopkinsi</i> , Jordan & Rutter	2855
<i>boulengeri</i> , Jordan & Starks.....	2856
<i>tigris</i> (Cuvier & Valenciennes)	1187
<i>amelopardalis</i> (Poey)	1187
Genus Cratinus, Steindachner	1188
<i>agassizii</i> , Steindachner.....	1188
Genus Hypoplectrus, Gill.....	1189
<i>lamprurus</i> (Jordan & Gilbert)	1190
<i>nunicolor</i> (Walbaum).....	1190
<i>puella</i> (Cuvier & Valenciennes)	1192
<i>vitulina</i> (Poey)	1192
<i>plinnivarius</i> (Poey).....	1192
<i>guttavarius</i> (Poey)	1192
<i>gummigutta</i> (Poey).....	1192
<i>erocotus</i> (Cope)	1192
<i>aberrans</i> (Poey).....	1193
<i>accensus</i> (Poey).....	1193
<i>affinis</i> (Poey).....	1193
<i>chlorurus</i> (Cuvier & Valenciennes)	1193
<i>nlgricana</i> (Poey)	1193
<i>indigo</i> (Poey)	1193
<i>bovinus</i> (Poey).....	1193
<i>gemma</i> , Goode & Bean.....	1193

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER RHEGNOPTERI—Continued.	
Family Serranidae—Continued.	
Genus <i>Paralabrax</i> , Girard.....	1194
<i>nebulifer</i> (Girard).....	1195
<i>maculatofasciatus</i> (Steindachner).....	1196
<i>humeralis</i> (Cuvier & Valenciennes).....	1196
<i>clathratus</i> (Girard)	1197
Genus <i>Centropristes</i> , Cuvier	1198
Subgenus <i>Centropristes</i>	1199
<i>rufus</i> , Cuvier & Valenciennes.....	1199
<i>striatus</i> (Linnaeus)	1199
<i>ocyrurus</i> (Jordan & Evermann).....	1200
Subgenus <i>Triloburus</i> , Gill.....	1201
<i>philadelphicus</i> (Linnaeus).....	1201
Genus <i>Diplectrum</i> , Holbrook	1203
Subgenus <i>Haliopercus</i> , Gill	1204
<i>sciurus</i> , Gilbert.....	1204
<i>radiale</i> (Quoy & Gaimard)	1204
<i>maeropoma</i> (Günther)	1205
<i>enrypteatum</i> , Jordan & Bollman	1206
Subgenus <i>Diplectrum</i>	1207
<i>formosum</i> (Linnaeus).....	1207
Genus <i>Prionodes</i> , Jenyns	1208
Subgenus <i>Prionodes</i>	1210
<i>equidens</i> (Gilbert).....	1210
<i>tusculus</i> (Poey).....	1211
<i>phebe</i> (Poey)	1211
<i>baldwini</i> , Evermann & Marsh.....	3168
<i>fasciatus</i> , Jenyns	1212
<i>bulleri</i> (Boulenger).....	1213
Subgenus <i>Mentipercus</i> , Gill	1214
<i>tigrinus</i> (Bloch).....	1214
<i>tabacarius</i> (Cuvier & Valenciennes)	1215
<i>flavescens</i> (Cuvier & Valenciennes).....	1215
<i>luciopercanus</i> (Poey)	1216
<i>stilostigma</i> , Jordan & Bollman.....	1216
Genus <i>Dules</i> , Cuvier.....	1217
<i>subligatus</i> (Cope)	1218
<i>displurus</i> (Günther).....	1219
<i>auriga</i> , Cuvier & Valenciennes.....	1220
Genus <i>Paranthias</i> , Guichenot	1221
<i>foreifer</i> (Cuvier & Valenciennes)	1221
Genus <i>Hemianthias</i> , Steindachner	1222
<i>peruanus</i> , Steindachner.....	1222
<i>vivanus</i> (Jordan & Swain)	1223
Genus <i>Pronogrammus</i> , Gill	1224
<i>eos</i> , Gilbert	1224
<i>multifasciatus</i> , Gill	1226
Genus <i>Anthias</i> , Bloch.....	1226
<i>asperillnguis</i> , Günther	1227
Genus <i>Ocyanthias</i> , Jordan & Evermann	1227
<i>martiniensis</i> (Guichenot)	1228
Genus <i>Gramma</i> , Poey	1228
<i>loreto</i> , Poey	1229
Genus <i>Khegma</i> , Gilbert.....	3169
<i>thaumastium</i> , Gilbert.....	3170

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER RHEONOPTERI—Continued.	
<i>Family Serranidae</i> —Continued.	
Genus <i>Ryptileus</i> , Cuvier.....	1229
xanti, Gill.....	1231
bicolor (Valenciennes)	1231
saponaceus (Bloch & Schneider).....	1232
arenatus, Cuvier & Valenciennes	1232
coriaceus (Cope)	1233
Subgenus <i>Promicropodus</i> , Gill.....	1233
<bistrispinus< b=""> (Mitchill).....</bistrispinus<>	1233
<bigrispinus< b="">, Gill.....</bigrispinus<>	1234
<i>Family Lobotidae</i>	1235
Genus <i>Lobotes</i> , Cuvier.....	1235
surinamensis (Bloch)	1235
pacificus, Gilbert.....	2857
<i>Family Priacanthidae</i>	1236
Genus <i>Priacanthus</i> , Cuvier	1237
arenatus, Cuvier & Valenciennes	1237
ermentatus (Lacépède).....	1238
carolinus, Lesson	2858
Genus <i>Pseudopriacanthus</i> , Bleeker	1239
serrula, Gilbert	1239
altus (Gill)	1239
<i>Family Lutianidae</i>	1241
Genus <i>Hoplopagrus</i> , Gill.....	1244
guntheri, Gill	1244
Genus <i>Evophites</i> , Gill	1245
viridis (Valenciennes).....	1246
Genus <i>Neomensis</i> , Girard	1247
Subgenus <i>Neomensis</i>	1251
jordani, Gilbert	1251
novemfasciatus (Gill)	1252
cyanopterus (Cuvier & Valenciennes)	1254
griseus (Linnaeus)	1255
jocu (Bloch & Schneider)	1257
apodus (Walbaum)	1258
argentiventris (Peters)	1260
lutjanoides (Poey)	1261
buccanella (Cuvier & Valenciennes)	1261
vivanus (Cuvier & Valenciennes)	1262
hastingsi, Bean	2858
aya (Bloch)	1264
analis (Cuvier & Valenciennes)	1265
colorado (Jordan & Gilbert)	1267
brachypterus (Cope)	1268
guttatus (Steindachner)	1269
synagris (Linnaeus)	1270
ambigua (Poey)	1271
mahogoni (Cuvier & Valenciennes)	1272
Subgenus <i>Raizeria</i> , Jordan & Fesler	1273
aratus (Günther)	1273
Genus <i>Rabirubia</i> , Jordan & Fesler	1274
lhermis (Peters)	1274
Genus <i>Ocyurus</i> , Gill	1275
chrysurus (Bloch)	1275

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI - Continued.	
SUBORDER RHEGNOPTERI—Continued.	
<i>Family Lutianidae</i> —Continued.	
Genus Rhomboplites (Gill)	1276
aurorubens (Cuvier & Valenciennes)	1277
Genus Apsilus, Cuvier & Valenciennes	1278
Subgenus Tropidinius, Gill.....	1278
dentatus, Gulchenot.....	1278
Genus Apriou, Cuvier & Valenciennes.....	1279
Subgenus Platynius, Gill	1280
macropthalmus (Müller & Troschel)	1280
Genus Etelis, Cuvier & Valenciennes.....	1281
oculatus (Cuvier & Valenciennes).....	1282
aquilonaris (Goode & Bean).....	1283
Genus Verilus, Poey	1283
sordidus, Poey	1284
Genus Xenocys, Jordan & Bollman.....	1285
jessiae, Jordan & Bollman	1285
Genus Xenistius, Jordan & Gilbert.....	1286
californiensis (Steindachner).....	1286
Genus Xenichthys, Gill.....	1287
tgassizii, Steindachner	1287
xanti, Gill.....	1287
Genus Nemipterus, Swainson	1288
macronemus (Günther)	1289
<i>Family Haemulidae</i>	1289
Genus Haemulon, Cuvier.....	1291
sexfasciatum, Gill.....	1294
album, Cuvier & Valenciennes	1295
macrostomum, Günther	1296
bonariense, Cuvier & Valenciennes	1297
parra (Desmarest).....	1297
scudderii, Gill.....	1299
helenae, Boulenger	3171
carbonarium, Poey	1300
steindachneri (Jordan & Gilbert)	1301
melanurum (Linnaeus)	1302
scirurus (Shaw).....	1303
plumieri (Lacépède)	1304
flavolineatum (Desmarest)	1306
Genus Brachygensis, Steudler	1307
chrysargyreus (Günther)	1307
Genus Bathystoma, Steudler	1308
rimator (Jordan & Swain)	1308
aurolineatum (Cuvier & Valenciennes)	1310
striatum (Linnaeus)	1310
Genus Lythrypnus, Jordan & Swain	1311
flaviguttatum (Gill).....	1312
opalescens, Jordan & Starks	1312
Genus Orthostichus, Gill	1313
maculicauda, Gill	1313
Genus Anisotremus, Gill	1314
Subgenus Paraconodon, Bleeker.....	1316
pacifici (Günther)	1316
caesius (Jordan & Gilbert)	1313
dovii (Günther)	1317

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER RHEGNOPTERI—Continued.	
<i>Family Haemulidae</i> —Continued.	
Genus <i>Anisotremus</i> , Gill—Continued.	
Subgenus <i>Anisotremus</i>	1318
<i>surinamensis</i> (Bloch).....	1318
<i>interruptus</i> (Gill)	1319
<i>bicolor</i> (Castelnau)	1319
<i>scapularis</i> (Tschudi)	1320
<i>do idsonii</i> (Steindachner)	1321
<i>spleniatus</i> (Poey)	1321
<i>tenuatus</i> , Gill	1322
<i>virginicus</i> (Linnaeus)	1322
<i>serrula</i> (Cuvier & Valenciennes)	1323
Genus <i>Conodon</i> , Cuvier & Valenciennes	1324
<i>nobilis</i> (Linnaeus)	1324
<i>serrifer</i> , Jordan & Gilbert	1324
Genus <i>Brachydeuterus</i> , Gill	1325
<i>nitidus</i> (Steindachner)	1326
<i>corvinaformis</i> (Steindachner)	1326
<i>leuciscus</i> (Günther)	1327
<i>axillaris</i> (Steindachner)	1328
Genus <i>Pomadasys</i> , Lacépède	1329
Subgenus <i>Rhenicus</i> , Jordan & Evermann	1331
<i>panamensis</i> (Steindachner)	1331
Subgenus <i>Pristipoma</i> , Cuvier	1331
<i>bayanus</i> , Jordan & Evermann	1331
<i>productus</i> (Poey)	1332
<i>macracanthus</i> (Günther)	1332
<i>andrei</i> (Sauvage)	1332
Subgenus <i>Rhencous</i> , Jordan & Evermann	1333
<i>crocro</i> (Cuvier & Valenciennes)	1333
<i>branicki</i> (Steindachner)	1333
<i>ramosus</i> (Poey)	1334
<i>labraciforme</i> (Boulenger)	1372
Genus <i>Orthopristis</i> , Girard	1334
Subgenus <i>Orthopristis</i>	1336
<i>forbesi</i> , Jordan & Starks	1336
<i>reddingi</i> , Jordan & Richardson	1336
<i>chalceus</i> (Günther)	1337
<i>chrysopterus</i> (Linnaeus)	1338
<i>poeyi</i> , Seudder	1339
<i>cantharinus</i> (Jenyns)	1339
Subgenus <i>Evapristis</i> , Jordan & Evermann	1340
<i>lethopristis</i> , Jordan & Fesler	1340
Genus <i>Isaciella</i> , Jordan & Fesler	1340
<i>breviplinis</i> (Steindachner)	1341
Genus <i>Microlepidotus</i> , Gill	1341
<i>inornatus</i> , Gill	1341
Genus <i>Genyatremus</i> , Gill	1342
<i>luteus</i> (Bloch)	1342
<i>Family Sparidae</i>	1343
Genus <i>Otrynter</i> , Jordan & Evermann	1344
<i>caprinus</i> (Bean)	1345
Genus <i>Stenotomus</i> , Gill	1345
<i>chrysops</i> (Linnaeus)	1346
<i>aculeatus</i> (Cuvier & Valenciennes)	1346

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER RHEGOPTERI—Continued.	
<i>Family Sparidae</i> —Continued.	
Genus <i>Calamus</i> , Swainson	1347
Subgenus <i>Calamus</i>	1349
<i>calamus</i> (Cuvier & Valenciennes).....	1349
<i>proridens</i> , Jordan & Gilbert.....	1350
<i>kendalli</i> , Evermann & Marsh	3172
<i>pennatula</i> , Gulebenot	1351
<i>bajonade</i> (Bloch & Schneider).....	1352
Subgenus <i>Grammateus</i> , Poey	1353
<i>brachysomus</i> (Lockington).....	1353
<i>leucosteus</i> , Jordan & Gilbert.....	1353
<i>macrops</i> , Poey	1354
<i>taurinus</i> (Jenyns).....	1354
<i>benna</i> (Cuvier & Valenciennes)	1354
<i>arctifrons</i> , Goode & Bean.....	1355
<i>medius</i> (Poey).....	1356
Genus <i>Pagrus</i> , Cuvier	1356
<i>Pagrus</i> (Linnaeus).....	1356
Genus <i>Lagodon</i> , Holbrook	1357
<i>rhomboides</i> (Linnaeus).....	1358
Genus <i>Archosargus</i> , Gill.....	1358
Subgenus <i>Salema</i> , Jordan & Evermann.....	1359
<i>unimaculatus</i> (Bloch)	1359
<i>pontica</i> (Steindachner)	1360
<i>tridens</i> (Poey).....	1360
Subgenus <i>Archosargus</i>	1361
<i>probatocephalus</i> (Walbaum)	1361
<i>aries</i> (Cuvier & Valenciennes)	1361
Genus <i>Diplodus</i> , Rafinesque	1362
<i>holbrookii</i> (Bean)	1362
<i>argenteus</i> (Cuvier & Valenciennes)	1363
<i>sargus</i> (Linnaeus)	1363
<i>Family Menidae</i>	1364
Genus <i>Spicara</i> , Rafinesque	1364
<i>martinica</i> (Cuvier & Valenciennes)	1364
Genus <i>Emmelichthys</i> , Richardson	1365
Subgenus <i>Inermia</i> , Poey	1365
<i>vittatus</i> (Poey)	1365
<i>Family Gerride</i>	1366
Genus <i>Eucinostomus</i> , Baird & Girard	1367
<i>dowi</i> (Gill)	1367
<i>pseudogula</i> , Poey	1368
<i>harengulus</i> , Goode & Bean	1368
<i>californiensis</i> (Gill)	1369
<i>gula</i> (Cuvier & Valenciennes)	1370
Genus <i>Ulema</i> , Jordan & Evermann	1371
<i>lefroyi</i> (Goode)	1371
Genus <i>Xystiema</i> , Jordan & Evermann	1372
<i>cinerereum</i> (Walbaum)	1372
Genus <i>Gerres</i> , Cuvier	1373
Subgenus <i>Moharra</i> , Poey	1374
<i>rhombus</i> , Cuvier & Valenciennes	1374
Subgenus <i>Diapterus</i> , Ranzani	1375
<i>aureolus</i> , Jordan & Gilbert	1375

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER RHEGNOPTERI—Continued.	
<i>Family Gerridae</i> —Continued.	
Genus <i>Gerris</i> , Cuvier—Continued.	
Subgenus <i>Diplapterus</i> , Ranzani—Continued.	
<i>peruvianus</i> , Cuvier & Valenciennes	1376
<i>olithostomus</i> , Goode & Bean	1376
Subgenus <i>Gerres</i>	1377
<i>brevimanus</i> , Günther	1377
<i>lineatus</i> (Humboldt)	1377
<i>brasiliensis</i> , Cuvier & Valenciennes	1378
<i>embryx</i> , Jordan & Starks	1379
<i>plumieri</i> , Cuvier & Valenciennes	1379
<i>mexicanus</i> , Steindachner.....	1380
<i>Family Kyphoside</i>	1380
Genus <i>Girella</i> , Gray	1381
<i>nigricans</i> (Ayres)	1382
Genus <i>Doydilodon</i> , Valenciennes	1382
<i>fremivillei</i> , Valenciennes	1382
Genus <i>Hermosilla</i> , Jenkins & Evermann	1383
<i>azurea</i> , Jenkins & Evermann	1383
Genus <i>Kyphosus</i> , Lacépède	1384
<i>analogus</i> (Gill)	1385
<i>incisor</i> (Cuvier & Valenciennes)	1386
<i>elegans</i> (Peters)	1387
<i>sextatrix</i> (Linnaeus)	1387
<i>lutescens</i> (Jordan & Gilbert)	1388
Genus <i>Sectator</i> , Jordan & Fesler	1389
<i>ocyrurus</i> (Jordan & Gilbert)	1389
Genus <i>Medialuna</i> , Jordan & Fesler	1390
<i>californiensis</i> (Steindachner)	1391
<i>Family Sciaenide</i>	1392
Genus <i>Seriphus</i> , Ayres	1397
<i>politus</i> , Ayres	1397
Genus <i>Isopisthus</i> , Gill	1398
<i>remifer</i> , Jordan & Gilbert	1398
<i>parvipinnis</i> (Cuvier & Valenciennes)	1399
Genus <i>Buccome</i> , Jordan & Evermann	1400
<i>predatoria</i> (Jordan & Gilbert)	1400
Genus <i>Cynoscion</i> , Gill	1400
Subgenus <i>Cynoscion</i>	1403
<i>acoupa</i> (Lacépède)	1403
<i>squamipinnis</i> (Günther)	1404
<i>othopterus</i> , Jordan & Gilbert	1404
<i>obliquatus</i> (Valenciennes)	1405
<i>jamaicensis</i> (Vaillant & Bocourt)	1406
<i>nothus</i> (Holbrook)	1406
<i>regalis</i> (Bloch & Schneider)	1407
<i>thalassinus</i> (Holbrook)	1407
<i>reticulatus</i> (Günther)	1408
<i>nebulosus</i> (Cuvier & Valenciennes)	1409
<i>parvipinnis</i> , Ayres	1410
<i>xanthulus</i> , Jordan & Gilbert	1410
<i>albus</i> (Günther)	1411
<i>macdonaldi</i> , Gilbert	1411
<i>stolzmanni</i> (Steindachner)	1412

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER RHEGNOPTERI—Continued.	
Family Sciaenidae—Continued.	
Genus <i>Cynoscion</i> , Gill—Continued.	
Subgenus <i>Atractoscion</i> , Gill.....	1413
<i>nobilis</i> (Ayres).....	1413
<i>phoxocephalus</i> , Jordan & Gilbert	1413
<i>leiarchus</i> (Cuvier & Valenciennes).....	1414
<i>virecens</i> (Cuvier & Valenciennes)	1415
<i>microlepidotus</i> (Cuvier & Valenciennes)	1415
Genus <i>Sagenichthys</i> , Berg.....	1416
<i>ancylodon</i> (Bloch & Schneider).....	1416
Genus <i>Nebrius</i> , Cuvier & Valenciennes.....	1416
<i>intropis</i> , Cuvier & Valenciennes.....	1417
<i>occidentalis</i> , Vaillant.....	3173
Genus <i>Plagioscion</i> , Gill	1418
<i>squamossissimus</i> (Heckel)	1418
<i>heterolepis</i> (Bleeker)	1419
<i>surinameensis</i> (Bleeker)	1419
Genus <i>Larimus</i> , Cuvier & Valenciennes	1420
Subgenus <i>Amblyscion</i> , Gill.....	1421
<i>argenteus</i> (Gill)	1421
Subgenus <i>Larimus</i>	1421
<i>effulgens</i> , Gilbert	1421
<i>acclivis</i> , Jordan & Bristol	1422
<i>breviceps</i> , Cuvier & Valenciennes	1423
<i>pacificus</i> , Jordan & Bollman	1424
<i>fasciatus</i> , Holbrook	1424
Genus <i>Odontoscion</i> , Gill	1425
<i>dentex</i> (Cuvier & Valenciennes)	1425
<i>xanthops</i> , Gilbert	1426
Genus <i>Corvula</i> , Jordan & Eigenmann	1427
<i>macrops</i> , (Steindachner)	1427
<i>sialis</i> , Jordan & Eigenmann	1428
<i>subaequalis</i> (Poey)	1429
<i>sanetæ-luciae</i> , Jordan	1429
<i>batabana</i> (Poey)	1430
Genus <i>Elattarchus</i> , Jordan & Evermann	1431
<i>archidium</i> (Jordan & Gilbert)	1431
Genus <i>Bairdiella</i> , Gill	1432
Subgenus <i>Bairdiella</i>	1433
<i>chrysura</i> (Lacépède)	1433
<i>ensifera</i> (Jordan & Gilbert)	1434
<i>icistia</i> (Jordan & Gilbert)	1435
<i>roachus</i> (Cuvier & Valenciennes)	1436
Subgenus <i>Nector</i> , Jordan & Evermann	1436
<i>armata</i> , Gill.....	1436
<i>aluta</i> , Jordan & Gilbert	1437
<i>chrysoleuca</i> (Günther)	1438
<i>niacantha</i> (Boulenger)	3173
Genus <i>Stellifer</i> (Cuvier) Oken	1439
Subgenus <i>Zestis</i> , Gilbert	1440
<i>escitans</i> (Jordan & Gilbert)	1440
<i>furthi</i> (Steindachner)	1441
Subgenus <i>Zestidium</i> , Gilbert	1442
<i>illecebrosus</i> , Gilbert	1442

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER RHEO(N)OPTERI—Continued.	
<i>Family Sciaenidae</i> —Continued.	
Genus <i>Stellifer</i> (Cuvier) Oken—Continued.	
Subgenus <i>Stellifer</i>	1443
<i>stellifer</i> (Bloch)	1443
<i>lanceolatus</i> (Holbrook).....	1443
<i>erleymba</i> (Jordan & Gilbert)	1444
<i>microps</i> (Steindachner)	1445
Subgenus <i>Stelicarens</i> , Gilbert.....	1445
<i>zestocarne</i> , Gilbert	1445
Genus <i>Ophioscion</i> , Gill	1446
Subgenus <i>Ophioscion</i>	1447
<i>adustus</i> (Agassiz)	1447
<i>typicus</i> , Gill	1448
<i>trabo</i> , Gilbert	1448
<i>simulus</i> , Gilbert	1449
<i>iniceps</i> (Jordan & Gilbert)	1451
<i>sclerurus</i> (Jordan & Gilbert).....	1452
Subgenus <i>Sigmarus</i> , Gilbert.....	1452
<i>vermicularis</i> (Günther)	1452
Genus <i>Schenops</i> , Gill	1453
<i>ocellatus</i> (Linneus)	1453
Genus <i>Sciaena</i> (Arvedi) Linnaeus	1454
Subgenus <i>Callans</i> , Jordan	1455
<i>delicosa</i> (Tschudi).....	1455
Subgenus <i>Cheilotrema</i> , Tschudi	1456
<i>saturna</i> (Girard)	1456
Genus <i>Roneador</i> , Jordan & Gilbert	1457
<i>stearnsi</i> (Steindachner)	1457
Genus <i>Lefostomus</i> , Lacépède	1458
<i>xanthurus</i> , Lacépède.....	1458
Genus <i>Pachypops</i> , Gill.....	1459
<i>furcatus</i> (Lacépède)	1459
Genus <i>Genyonemus</i> , Gill	1460
<i>lineatus</i> (Ayres)	1460
Genus <i>Micropogon</i> , Cuvier & Valenciennes	1461
<i>undulatus</i> (Linneus)	1461
<i>furnieri</i> (Desmarest)	1462
<i>megalops</i> , Gilbert	1463
<i>etenes</i> , Jordan & Gilbert.....	1463
<i>altipinnis</i> , Günther.....	1464
Genus <i>Umbrina</i> , Cuvier	1465
<i>broussoneti</i> , Cuvier & Valenciennes	1466
<i>coroides</i> , Cuvier & Valenciennes	1466
<i>roneador</i> , Jordan & Gilbert	1467
<i>xanti</i> , Gill	1467
<i>sinaloae</i> , Scofield	1468
<i>galapagorum</i> , Steindachner	1468
<i>dorsalis</i> , Gill	1469
Genus <i>Menticirrhus</i> , Gill	1469
<i>sinus</i> , Jordan & Eigenmann	1472
<i>nasus</i> (Günther)	1473
<i>panamensis</i> (Steindachner)	1473
<i>martinicensis</i> (Cuvier & Valenciennes)	1473
<i>americanus</i> (Linnaeus)	1474

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER RHEGNOPTERI—Continued.	
<i>Family Sciaenidae</i> —Continued.	
Genus <i>Menticirrhus</i> , Gill—Continued.	
<i>saxatilis</i> (Bloch & Schneider)	1475
<i>undulatus</i> (Girard)	1476
Subgenus <i>Umbrula</i> , Jordan & Eigenmann	1476
<i>elongatus</i> (Giüthler)	1476
<i>ittoralis</i> (Holbrook)	1477
Genus <i>Paralonchurus</i> , Bocourt	1477
Subgenus <i>Polyplemus</i> , Berg	1478
<i>dinnerili</i> (Bocourt)	1478
Subgenus <i>Zcnoseion</i> , Jordan & Evermann	1479
<i>rathbuni</i> (Jordan & Bollman)	1479
Subgenus <i>Zaclemus</i> , Gilbert	1480
<i>goodei</i> , Gilbert	1480
Subgenus <i>Paralonchurus</i>	1481
<i>petersi</i> , Bocourt	1481
Genus <i>Lonchirus</i> , Bloch	1481
<i>lanceolatus</i> (Bloch)	1481
Genus <i>Pogonias</i> , Lacépède	1482
<i>cromis</i> (Linnaeus)	1482
<i>courbina</i> (Lacépède)	1483
Genus <i>Aplodinotus</i> , Rafinesque	1483
<i>gurniens</i> , Rafinesque	1484
Genus <i>Eques</i> , Bloch	1485
Subgenus <i>Pareques</i> , Gill	1486
<i>viola</i> , Gilbert	1486
<i>acutainatus</i> (Bloch & Schneider)	1487
<i>umbrosus</i> , Jordan & Eigenmann	1487
<i>pnactatus</i> , Bloch & Schneider	1488
<i>pulcher</i> , Steindachner	1489
Subgenus <i>Eques</i>	1489
<i>lanceolatus</i> (Linnaeus)	1489
Group <i>Cirrhitidei</i>	1490
<i>Family Cirrhitidae</i>	1490
Genus <i>Cirrhites</i> , Lacépède	1491
<i>rivulatus</i> , Valenciennes	1491
<i>betaurus</i> , Gill	1492
SUBORDER HOLCONOTI	1493
<i>Family Embiotocidae</i>	1493
Genus <i>Hystericarpus</i> , Gibbons	1495
<i>traski</i> , Gibbons	1496
Genus <i>Abeona</i> , Girard	1496
<i>minima</i> (Gibbons)	1497
<i>aurora</i> , Jordan & Gilbert	1497
Genus <i>Cymatogaster</i> , Gibbons	1498
<i>aggregatus</i> , Gibbons	1498
Genus <i>Brachylistius</i> , Gill	1499
<i>frenatus</i> , Gill	1499
Genus <i>Zalembius</i> , Jordan & Evermann	1499
<i>rosaceus</i> (Jordan & Gilbert)	1500
Genus <i>Hypocritichthys</i> , Gill	1500
<i>analis</i> (Alexander Agassiz)	1500

	Page.	
CLASS PISCES—Continued.		
SUBCLASS TELEOSTOMI—Continued.		
ORDER ACANTHOPTERI—Continued.		
SUBORDER HOLCONOTI—Continued.		
Family Embiotocidae—Continued.		
475	Genus Hyperprosopon, Gibbons.....	1501
476	argentens, Gibbons.....	1501
476	agassizii, Gill.....	1502
476	Genus Holconotus, Agassiz.....	1502
476	rhodoterns, Agassiz.....	1502
477	Genus Amphisticus, Agassiz.....	1503
477	argentens, Agassiz.....	1503
478	Genus Embiotoca, Agassiz.....	1504
478	jacksoni, Agassiz.....	1504
479	Genus Taeniotoca, Alexander Agassiz.....	1505
479	lateralis (Agassiz).....	1505
480	Genus Phancredon, Girard.....	1506
480	fureatus, Girard.....	1506
481	artipes (Jordan & Gilbert).....	1507
481	Genus Rhacochilus, Agassiz.....	1507
481	toxotes, Agassiz.....	1507
482	Genus Hypsurus, Alexander Agassiz.....	1508
482	caryi (Agassiz).....	1508
482	Genus Damalichthys, Girard.....	1509
483	argyrosomus (Girard).....	1509
483	SUBORDER CHROMIDES.....	1511
484	Family Cichlidæ.....	1512
485	Genus Petenia, Günther.....	1513
486	splendida, Günther.....	1513
486	Genus Evidens, Eigenmann & Bray.....	1513
487	coruleopunctatus (Kner & Steindachner).....	1514
487	Genus Cichlasoma, Swainson.....	1514
488	Subgenus Cichlasoma.....	1515
489	rectangulare (Steindachner).....	1515
489	bartoni (Bean).....	1515
489	godmani (Günther).....	1516
490	sieboldii (Kner & Steindachner).....	1516
490	intermedium (Günther).....	1517
491	anguliferum (Günther).....	1517
491	fenestratum (Günther).....	1518
492	montezuma (Heckel).....	1518
493	macracanthum (Günther).....	1518
493	parva (Günther).....	1519
495	margaritiferum (Günther).....	1519
496	spilurum (Günther).....	1520
496	longimanus (Günther).....	1520
497	bifasciatum (Steindachner).....	1521
497	helleri (Steindachner).....	1521
498	balteatum (Gill & Bransford).....	1521
498	rostratum (Gill & Bransford).....	1522
499	melanopogon (Steindachner).....	1523
499	melanurum (Günther).....	1523
500	nebulosum (Günther).....	1524
500	lentiginosum (Steindachner).....	1524
500	deppii (Heckel).....	1524
500	Subgenus Archocentrus, Gill.....	1525
	nigrafasciatum (Günther).....	1525

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUPERFAMILY CHROMIDES—Continued.	
<i>Family Cichlidae—Continued.</i>	
Genus <i>Cichlasoma</i> , Swainson—Continued.	
Subgenus <i>Archocentrus</i> , Gill—Continued.	
<i>multispinosum</i> (Günther)	1525
<i>centrarchus</i> (Gill & Bransford)	1526
<i>steindachneri</i> , Jordan & Snyder.....	3173
Genus <i>Heros</i> , Heckel.....	1526
<i>friedrichsthalii</i> , Heckel.....	1528
<i>salvini</i> , Günther.....	1528
<i>affinis</i> , Günther.....	1529
<i>maculipinnis</i> , Steindachner.....	1529
<i>trimaculatus</i> , Günther.....	1529
<i>labiatus</i> , Günther	1530
<i>lobochilus</i> , Günther.....	1530
<i>erythraeus</i> , Günther.....	1531
<i>basilaris</i> , Gill & Bransford.....	1532
<i>nicaraguensis</i> , Günther	1532
<i>managuensis</i> , Günther	1533
<i>aureus</i> , Günther	1533
<i>citrinellus</i> , Günther.....	1534
<i>motaguensis</i> , Günther.....	1534
<i>oblongus</i> , Günther.....	1535
<i>dovii</i> , Günther.....	1535
<i>gibbleps</i> , Steindachner.....	1536
<i>microphthalmus</i> , Günther.....	1536
<i>urophthalmus</i> , Günther.....	1537
<i>troschell</i> , Steindachner.....	1537
<i>cyanoguttatus</i> (Baird & Girard).....	1537
<i>pavonaceus</i> , Garman.....	1538
<i>altifrons</i> , Kner & Steindachner.....	1538
<i>beamii</i> , Jordan.....	1538
<i>tetraacanthus</i> (Cuvier & Valenciennes).....	1539
<i>istianus</i> , Jordan & Snyder.....	3174
Genus <i>Theraps</i> , Günther.....	1540
<i>irregularis</i> , Günther	1540
Genus <i>Neetroplus</i> , Günther.....	1541
<i>nomatopus</i> , Günther	1541
<i>nicaraguensis</i> , Gill & Bransford.....	1542
<i>carpintis</i> , Jordan & Snyder.....	3175
Genus <i>Satanoperca</i> , Günther.....	1542
<i>crassilabris</i> (Steindachner).....	1542
<i>Family Pomacentridae</i>	1543
Genus <i>Azurina</i> , Jordan & McGregor.....	1544
<i>hlrunido</i> , Jordan & McGregor.....	1544
Genus <i>Chromis</i> , Cuvier.....	1545
Subgenus <i>Furcaria</i> , Poey.....	1546
<i>atrilobatus</i> , Gill.....	1546
<i>cyaneus</i> (Poey).....	1547
<i>multilineatus</i> (Guichenot).....	1547
Subgenus <i>Ayresia</i> , Cooper.....	1548
<i>punctiphonis</i> (Cooper).....	1548
Subgenus <i>Heliastes</i> , Cuvier & Valenciennes.....	1548
<i>insolatus</i> (Cuvier & Valenciennes).....	1548
<i>enchrysurus</i> , Jordan & Gilbert.....	1548

CLASS FISHES—Continued.

SUBCLASS TELEOSTOMI—Continued.

ORDER ACANTHOPTERI—Continued.

SUBORDER CHROMIDES—Continued.

Family Pomacentridæ—Continued.

Page.		Page.
1525	Genus Eupomacentrus, Bleeker	1549
1526	Subgenus Eupomacentrus	1551
3173	leucorus (Gilbert)	1551
1526	adustus (Troschel)	1551
1528	fuscus (Cuvier & Valenciennes)	1552
1528	diacanthus, Jordan & Rutter	1552
1528	rectifranum (Gill)	1553; 3176
1528	analis (Poey)	1554
1529	otophorus (Poey)	1555
1529	leucostictus (Müller & Troschel)	1555
1529	flaviventer (Troschel)	1557
1530	partitus (Poey)	1558
1530	planifrons (Cuvier & Valenciennes)	1559
1531	Genus Nexilarius, Gilbert	1559
1532	concolor (Gill)	1559
1532	Genus Abudefduf, Forskål	1560
1533	Subgenus Glyphtodon, Lacépède	1561
1533	saxatilis (Linnaeus)	1561
1534	Subgenus Euschistodus, Gill	1562
1534	declivifrons (Gill)	1562
1535	analogus (Gill)	1563
1535	taurus (Müller & Troschel)	1563
1536	rndis (Poey)	1563
1536	Genus Hypsopops, Gill	1564
1537	rubicundus (Girard)	1564
1537	Genus Microspathodon, Günther	1565
1537	bairdii (Gill)	1566
1538	chrysurus (Cuvier & Valenciennes)	1567
1538	niveatus (Poey)	1567
1538	dorsalis (Gill)	1568
1539	SUBORDER PHARYNGOGNATHI	1571
3174	Family Labridæ	1571
1540	Genus Centrolabrus, Günther	1575
1540	exoletus (Linnaeus)	1576
1541	Genus Tautogolabrus, Günther	1576
1541	adspersus (Walbaum)	1577
1542	Genus Tautoga, Mitchell	1577
3175	onitis (Linnaeus)	1578
1542	Genus Lachnolaimus, Cuvier & Valenciennes	1579
1542	maximus (Walbaum)	1579
1513	Genus Harpe, Lacépède	1581
1544	diplotenia, (Gill)	1582
1544	rufa (Linnaeus)	1583
1545	eclancheri (Valenciennes)	1583
1546	pulchella (Poey)	1584
1546	Genus Decodon, Günther	1584
1547	puellaris (Poey)	1584
1547	Genus Plimelometopon, Gill	1585
1548	pulcher (Ayres)	1585
1548	darwini (Jenyns)	1586
1548	Genus Clepticus, Cuvier	1586
1548	parra (Bloch & Schneider)	1586
... 1548		

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER PHARYNGOGNATHI—Continued.	
Family Labridæ—Continued.	
Genus Iridio, Jordan & Evermann	1587
radiatus (Linnaeus)	1590
nicholsi (Jordan & Gilbert)	1591
sellifer (Gilbert)	1592
semicinctus (Ayres)	1592
garnoti (Cuvier & Valenciennes)	1593
cyanocephalus (Bloch)	1594
maculipinna (Müller & Troschel)	1594
bivittatus (Bloch)	1595
dispidus (Günther)	1597
kirschi, Jordan & Evermann	1598
poeyi (Steindachner)	1599
caudalis (Poey)	1599
pictus (Poey)	1599
Genus Oxyjulis, Gill	1601
californicus (Günther)	1601
Genus Eumeeekia, Jordan & Evermann	1601
venusta (Jenkins & Evermann)	1602
Genus Julidio, Jordan & Evermann	1602
adustus (Gilbert)	1602
notospilus (Günther)	1603
Genus Pseudojulis, Bleeker	1604
inornatus, Gilbert	1604
melanotis, Gilbert	1605
Genus Thalassoma, Swainson	1605; 2859
lucasianum (Gill)	1607; 2859
socorroense, Gilbert	1607; 2859
nitidum (Günther)	1608; 2859
nitidissimum (Goode)	1608; 2859
steindachneri (Jordan)	1609; 2859
bifasciatum (Bloch)	1609; 2859
grammaticum, Gilbert	1610; 2859
virens, Gilbert	1610; 2859
Genus Doratonotus, Günther	1611
megalepis, Günther	1611
decoris, Evermann & Marsh	3177
Genus Xyrula, Jordan	1612
jessiae (Jordan)	1612
Genus Novaculichthys, Bleeker	1613
rosipes (Jordan & Gilbert)	1614
ventralis (Bean)	1615
infirmus (Bean)	1616
marticensis (Cuvier & Valenciennes)	1616
Genus Xyrichthys, Cuvier	1617
mundiceps, Gill	1618
psittacus (Linnaeus)	1618
modestus, Poey	1619
Genus Inlistius, Gill	1619
mundicorpus, Gill	1620
Family Scaridae.....	1620
Genus Cryptotomus, Cope	1621
dentiens (Poey)	1623
retractus (Poey)	1623
ustus (Cuvier & Valenciennes)	1624

CLASS PISCES—Continued.

Page.

SUBCLASS TELEOSTOMI—Continued.

ORDER ACANTHOPTERI—Continued.

SUBORDER PHARYNGOGNATHI—Continued.

Family Scaridae—Continued.

Page.

1587	Genus Cryptotomus, Cope—Continued.	
1590	auropunctatus (Cuvier & Valenciennes)	1624
1591	beryllinus, Jordan & Swain.....	1625
1592	rosenii, Cope	1626
1592	Genus Calotomus, Gilbert	1626
1593	xenodon, Gilbert.....	1626
1594	Genus Sparisoma, Swainson	1627
1594	Subgenus Sparisoma	1630
1595	xystrodon, Jordan & Swain	1630
1597	atomarium (Poey).....	1631
1598	radicans (Cuvier & Valenciennes)	1631
1599	hoplomystax (Cope).....	1632
1599	niphobles, Jordan & Bollman	1633
1599	aurofrenatum (Cuvier & Valenciennes).....	1634
1601	oxybrachium (Poey)	1634
1601	abildgaardii (Bloch).....	1635
1601	distinctum (Poey)	1635
1602	chrysopterum (Bloch & Schneider).....	1636
1602	lorito, Jordan & Swain.....	1637
1602	viride (Bonnaterre)	1638
1603	Subgenus Euscarus, Jordan & Evermann	1639
1604	strigatum (Günther)	1639
1604	flavescens (Bloch & Schneider)	1639
1605	rubripinne (Cuvier & Valenciennes)	1640
5; 2850	brachiale (Poey)	1641
7; 2850	maschalepilos (Bleeker)	1641
7; 2850	frondosum (Cuvier)	1642
8; 2850	Genus Scarus, Forskål	1642
8; 2850	Subgenus Scarus	1645
9; 2850	punctulatus (Cuvier & Valenciennes).....	1645
9; 2850	bollmani, Jordan & Evermann	1646
0; 2850	tenuipterus, Desmarest	1646
0; 2850	aracanga (Günther)	1647
.. 1611	trispinosus, Cuvier & Valenciennes	1648
.. 1611	cuzauila, Bean.....	1648
.. 3177	vetula, Bloch & Schneider.....	1649
.. 1612	gnathodus, Poey	1650
.. 1612	Subgenus Calliodon (Gronow) Schneider.....	1650
.. 1613	croicensis (Bloch)	1650
.. 1614	evermanni, Jordan	1651
.. 1615	flavomarginatus, Cuvier & Valenciennes	1652
.. 1616	acus, Poey	1652
.. 1616	caeruleus (Bloch).....	1652
.. 1617	emblematus, Jordan & Rutter	1654
.. 1618	Genus Pseudoscarus, Bleeker	1655
.. 1618	Subgenus Pseudoscarus	1655
.. 1619	celestinus (Cuvier & Valenciennes).....	1655
.. 1619	simplex, Poey	1656
.. 1620	pleianus (Poey)	1656
.. 1620	Subgenus Loro, Jordan & Evermann	1657
.. 1621	guacamala (Cuvier)	1657
.. 1621	perile (Jordan & Gilbert)	1658
.. 1623	Group Zeoldea	1659
.. 1624		

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER ACANTHOPTERI—Continued.	
SUBORDER PHARYNGOGNATHI—Continued.	
<i>Family Zeidae</i>	1659
Genus <i>Zenopsis</i> , Gill.....	1660
<i>ocellatus</i> (Storer)	1660
Genus <i>Zenion</i> , Jordan & Evermann	1661
<i>hololepis</i> (Goode & Bean)	1661
Genus <i>Oreosoma</i> , Cuvier & Valenciennes	1662
<i>atlanticum</i> , Cuvier & Valenciennes	1662
Group <i>Caproidea</i>	1663
<i>Family Caproidae</i>	1663
Genus <i>Antigonia</i> , Lowe.....	1664
<i>capros</i> , Lowe.....	1665
SUBORDER SQUAMIPINNES	1665
<i>Family Ephippidae</i>	1666
Genus <i>Chetodipterus</i> , Lacépède.....	1667
<i>faber</i> (Broussonet)	1668
<i>zonatus</i> (Girard)	1668
Genus <i>Parapsettus</i> , Steindachner	1669
<i>panamensis</i> , Steindachner	1669
<i>Family Chaetodontidae</i>	1669
Genus <i>Prognathodes</i> , Gill.....	1671
<i>aculeatus</i> (Poey)	1671
Genus <i>Forcipiger</i> , Jordan & McGregor	1671
<i>flavissimus</i> , Jordan & McGregor	1671
Genus <i>Chaetodon</i> (Arvedi) Linnaeus	1672
Subgenus <i>Chatodontops</i> , Bleeker.....	1673
<i>nigrirostris</i> (Gill)	1673
<i>ocellatus</i> , Bloch	1674
<i>humeralis</i> , Günther	1674
<i>sedentarius</i> , Poey	1675
<i>aya</i> , Jordan	1675
<i>ateniatus</i> (Poey)	1676
<i>striatus</i> , Linnaeus	1677
Subgenus <i>Chatodon</i>	1677
<i>capistratus</i> , Linnaeus	1677
<i>bricei</i> , Smith	1678
Genus <i>Pomacanthus</i> , Lacépède	1679
Subgenus <i>Pomacanthus</i>	1679
<i>arcuatus</i> (Linnaeus)	1679
<i>paru</i> (Bloch)	1680
Subgenus <i>Pomacanthodes</i> , Gill	1681
<i>zonopectus</i> (Gill)	1681
Genus <i>Holacanthus</i> , Lacépède	1682
<i>passer</i> , Valenciennes	1682
<i>clarionensis</i> , Gilbert	1683
<i>tricolor</i> (Bloch)	1684
Genus <i>Angelichthys</i> , Jordan & Evermann	1684
<i>ciliaris</i> (Linnaeus)	1684
<i>isabelita</i> , Jordan & Rutter	1685
<i>iodocus</i> , Jordan & Rutter	1686
<i>Family Zanclidae</i>	1687
Genus <i>Zanclus</i> , Cuvier & Valenciennes	1687
<i>cornutus</i> (Linnaeus)	1687
<i>Family Teuthididae</i>	1688
Genus <i>Teuthis</i> , Linnaeus	1689
<i>triostegus</i> (Linnaeus)	1690

CLASS
SUBC
ORD
SUSORD
SE

P

age.	CLASS PISCES—Continued.	Page.
	SUBCLASS TELEOSTOMI—Continued.	
	ORDER Acanthopteri—Continued.	
	SUBORDER SQUAMIPINNES—Continued.	
	<i>Family Teuthidae</i> —Continued.	
1659	Genus <i>Teuthis</i> , Linnaeus—Continued.	
1660	ceruleus (Bloch & Schneider)	1601
1661	hepatus, Linnaeus	1601
1661	crestonius, Jordan & Starks	1692
1662	bahianus (Castelnau)	1693
1662	aliala (Lesson)	1693
1663	Genus <i>Xesurus</i> , Jordan & Evermann	1604
1663	punctatus (Gill)	1604
1664	clarionis, Gilbert & Starks	1605
1665	latilavivus (Valenciennes)	1605
1665	ORDER PLECTOGNATHI.	1696
1666	SUBORDER SCLERODERMI	1697
1667	<i>Family Triacanthidae</i>	1697
1668	Genus <i>Hollandia</i> , Poey	1697
1668	hollandi, Poey	1698
1669	<i>Family Balistidae</i>	1698
1669	Genus <i>Balistes</i> (Artedi) Linnaeus	1609
1669	Subgenus <i>Caprifascus</i> , Rafinesque	1700
1671	polylepis, Steindachner	1700
1671	nauprarium, Jordan & Starks	1700
1671	carolinensis, Gmelin	1701
1671	forcipatus, Gmelin	1702
1672	Subgenus <i>Balistes</i>	1703
1673	vetula, Linnaeus	1703
1673	Genus <i>Pachynathus</i> , Swainson	1703
1674	epistratus (Shaw)	1704
1674	Genus <i>Canthidermis</i> , Swainson	1705
1675	sobaeo, Poey	1705
1675	sufflamen (Mitchill)	1706
1676	maculatus (Bloch)	1706
1677	willughbeii (Lay & Bennett)	1707
1677	Genus <i>Xanthichthys</i> , Kaup	1708
1677	ringens (Linnaeus)	1709
1678	mento (Jordan & Gilbert)	1710
1679	Genus <i>Melichthys</i> , Swainson	1711
1679	piceus (Poey)	1711
1679	bispinosus, Gilbert	1711
1680	<i>Family Monacanthidae</i>	1712
1681	Genus <i>Cantherines</i> , Swainson	1713
1681	pullus (Ranzani)	1713
1682	carolinus, Jordan & McGregor	1713
1682	Genus <i>Monacanthus</i> , Cuvier	1714
1683	ciliatus (Mitchill)	1714
1684	hispidus (Linnaeus)	1715
1684	spilonotus, Cope	1716
1684	oppositus, Poey	1716
1685	Genus <i>Pseudomonacanthus</i> , Bleeker	1717
1686	amphioxys (Cope)	1717
1687	Genus <i>Ceratacanthus</i> , Gill	1718; 2860
1687	schoepfii (Walbaum)	1718; 2860
1687	punctatus (Agassiz)	1718; 2860
1688	Subgenus <i>Osbeckia</i> , Jordan & Evermann	1719
1688	scriptus (Osbeck)	1719; 2860
1689	Genus <i>Aluterus</i> , Cuvier	1717; 2860
1690	monoceros (Osbeck)	1720; 2860; 3178

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER OSTRACODERMI	1720
<i>Family Ostraciidae</i>	1721
Genus <i>Lactophrys</i> , Swainson	1721
Subgenus <i>Rhinesomus</i> , Swainson	1722
<i>triquerter</i> (Linnaeus)	1722
Subgenus <i>Chapinus</i> , Jordan & Evermann	1723
<i>bicaudalis</i> (Linnaeus)	1723
Subgenus <i>Lactophrys</i>	1723
<i>trigonus</i> (Linnaeus)	1723
Subgenus <i>Acanthostracion</i> , Bleeker	1724
<i>tricornis</i> (Linnaeus)	1724
SUBORDER GYMNODONTES	1726
<i>Family Tetraodontidae</i>	1726
Genus <i>Lagocephalus</i> , Swainson	1727
<i>levigatus</i> (Linnaeus)	1728
<i>pachycephalus</i> (Ranzani)	1728
Genus <i>Spherooides</i> , Lacépède	1729
Subgenus <i>Spherooides</i>	1731
<i>angusticeps</i> (Jenyns)	1731
<i>lobatus</i> (Steindachner)	1731
<i>spengleri</i> (Bloch)	1732
<i>maculatus</i> (Bloch & Schneider)	1733
<i>nephelus</i> (Goode & Bean)	1733; 3178
<i>marmoratus</i> (Ranzani)	1733
Subgenus <i>Cheilichthys</i> , Müller	1734
<i>testudineus</i> (Linnaeus)	1734
<i>annulatus</i> (Jenyns)	1735
<i>politus</i> (Girard)	1736
<i>formosus</i> (Günther)	1736
<i>furthi</i> (Steindachner)	1737
<i>trichocephalus</i> (Cope)	1737
<i>pachygaster</i> (Müller & Troschel)	1738
Genus <i>Ovoides</i> , Lacépède	1738
<i>orethizon</i> (Jordan & Gilbert)	1739
<i>setosus</i> (Rosa Smith)	1739
Genus <i>Colomesus</i> , Gill	1740
<i>psittacus</i> (Bloch & Schneider)	1740
<i>Family Canthigasteridae</i>	1740
Genus <i>Canthigaster</i> , Swainson	1741
<i>punctatissimus</i> (Günther)	1741
<i>rostratus</i> (Bloch)	1741
<i>Family Diodontidae</i>	1742
Genus <i>Trichodiodon</i> , Bleeker	1743
<i>pilosus</i> (Mitchill)	1743
Genus <i>Diodon</i> , Linnaeus	1744
<i>hystrix</i> , Linnaeus	1745
<i>holacanthus</i> , Linnaeus	1746
<i>maculifer</i> , Kaup	1747
Genus <i>Chilomycterus</i> , Bibron	1747
Subgenus <i>Cyclichthys</i> , Kaup	1748
<i>schöpfii</i> (Walbaum)	1748
<i>spinulosus</i> , Linnaeus	1749
<i>antillarum</i> , Jordan & Rutter	1749
<i>antennatus</i> (Cuvier)	1750
Subgenus <i>Chilomycterus</i>	1750
<i>atinga</i> (Linnaeus)	1750
<i>californiensis</i> , Elgemann	1751

CLASS PISCES—Continued.

SUBCLASS TELEOSTOMI—Continued.

ORDER PLECTOGNATHI—Continued.

SUBORDER GYMNOdontes—Continued.

Family *Diodontidae*—Continued.

Genus <i>Lyosphera</i> , Evermann & Kendall.....	1751
<i>globosa</i> , Evermann & Kendall.....	1751
Family <i>Molidae</i>	1752
Genus <i>Mola</i> , Cuvier	1753
<i>mola</i> (Linnaeus)	1753
Genus <i>Ranzania</i> , Nardo	1755
<i>truncata</i> (Retzius)	1755
Suborder LORICATI	1756
Family <i>Scorpaenidae</i>	1758
Genus <i>Sebastes</i> , Cuvier	1760
<i>marinus</i> (Linnaeus)	1760
<i>Sebastolobus</i> , Gill	1761
<i>alascanus</i> , Bean	1761
<i>altivelis</i> , Gilbert	1763
Genus <i>Sebastodes</i> , Gill	1765
Subgenus <i>Emmelas</i> , Jordan & Evermann	1777
<i>glaucescens</i> (Hilgendorf)	1777
Subgenus <i>Sebastodes</i>	1778
<i>jordani</i> , Gilbert	1778
<i>goodei</i> , Eigenmann & Eigenmann	1779
<i>pancispinis</i> (Ayres)	1780
Subgenus <i>Sebastosomus</i> , Gill	1781
<i>flavidus</i> (Ayres)	1781
<i>serranoides</i> , Eigenmann & Eigenmann	1782
<i>melanops</i> (Girard)	1782
Subgenus <i>Primospina</i> , Eigenmann & Beeson	1783
<i>ciliatus</i> (Tilesius)	1783
<i>mystinus</i> , Jordan & Gilbert	1784
Subgenus <i>Acutomentum</i> , Eigenmann & Beeson	1785
<i>entomelas</i> (Jordan & Gilbert)	1785
<i>rufus</i> , Eigenmann & Eigenmann	1786
<i>maedonaldi</i> (Eigenmann & Beeson)	1786
<i>brevispinis</i> (Bean)	1787
<i>ovalis</i> (Ayres)	1788
<i>eigenmanni</i> , Cramer	1789
<i>hopkinsii</i> , Cramer	1789
<i>alutus</i> (Gilbert)	1790
<i>proriger</i> (Jordan & Gilbert)	1792
Subgenus <i>Rosicola</i> , Jordan & Evermann	1793
<i>pinniger</i> (Gill)	1793
<i>miniatum</i> (Jordan & Gilbert)	1794
<i>atrorubens</i> (Gilbert)	1796
<i>atrovirens</i> (Jordan & Gilbert)	1797
Subgenus <i>Zalopyr</i> , Jordan & Evermann	1795; 2960
<i>aleutianus</i> , Jordan & Evermann	1795; 2860
Subgenus <i>Eosebastes</i> , Jordan & Evermann	1798
<i>saxicola</i> (Gilbert)	1798
<i>crameri</i> , Jordan	1799
<i>seunicinctus</i> , Gilbert	1800
<i>diploproa</i> (Gilbert)	1801
<i>aurora</i> (Gilbert)	1802
<i>melanostomus</i> , Eigenmann & Eigenmann	1803
<i>introniger</i> (Gilbert)	1803

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER Loricati—Continued.	
Family <i>Scorpaenidae</i> —Continued.	
Genus <i>Sebastodes</i> , Gill—Continued.	
Subgenus <i>Sebastomus</i> , Gill	1805
<i>ruberrimus</i> , Cramer.....	1805
<i>constellatus</i> (Jordan & Gilbert)	1806
<i>umbrosus</i> (Jordan & Gilbert)	1807
<i>rosaceus</i> (Girard)	1808
<i>ayre</i> . Gilbert & Cramer	1808
<i>rhodochloris</i> (Jordan & Gilbert)	1809
<i>eos</i> , Eigenmann & Eigemann.....	1810
<i>gilli</i> , Eigenmann & Eigenmann	1811
<i>chlorostictus</i> (Jordan & Gilbert).....	1811
<i>rupestris</i> (Gilbert)	1812
Subgenus <i>Hispanicus</i> , Cramer	1813
<i>siensis</i> (Gilbert).....	1813
<i>zaeocentrus</i> (Gilbert).....	1814
<i>elongatus</i> (Ayles).....	1815
<i>levis</i> (Eigenmann & Eigemann).....	1816
<i>ruberivinctus</i> (Jordan & Gilbert)	1817
Subgenus <i>Autospina</i> , Eigenmann & Peeson	1817
<i>auriculatus</i> (Girard).....	1817
<i>dallii</i> (Eigenmann & Beeson).....	1818
Subgenus <i>Pteropodus</i> , Eigenmann & Beeson	1819
<i>rastrelliger</i> (Jordan & Gilbert)	1819
<i>caurinus</i> (Richardson)	1820
<i>vexillaris</i> (Jordan & Gilbert)	1821
<i>maliger</i> (Jordan & Gilbert)	1822
<i>gilberti</i> , Cramer	1823
<i>carnatus</i> (Jordan & Gilbert)	1824
<i>chrysomelas</i> (Jordan & Gilbert)	1825
<i>nebulosus</i> (Ayles)	1826
Subgenus <i>Sebastichthys</i> , Gill	1827
<i>sericeps</i> (Jordan & Gilbert)	1827
<i>nigrocinetus</i> (Ayles)	1827
Subgenus <i>Sebastosomus</i> , Gill	1829; 2860
<i>taczanowskii</i> (Steindachner)	1831; 2860
Genus <i>Sebastopsis</i> , Gill.....	1835
<i>xyris</i> , Jordan & Gilbert.....	1835
Genus <i>Helicolenus</i> , Goode & Bean	1836
<i>dactylopterus</i> (De la Roche)	1837
<i>maderensis</i> , Goode & Bean	1837
Genus <i>Scorpaena</i> (Artedi) Linnaeus	1839
<i>agassizii</i> , Goode & Bean	1840
<i>eristulata</i> , Goode & Bean	1841
<i>brasiliensis</i> , Cuvier & Valenciennes	1842
<i>histrio</i> , Jenyns	1843
<i>pannosa</i> , Cramer	1845
<i>guttata</i> , Girard	1847
<i>plumieri</i> , Bloch	1848
<i>mystes</i> , Jordan & Starks	1849
<i>grandicornis</i> , Cuvier & Valenciennes	1850
<i>russula</i> , Jordan & Bollman	1851
<i>sonore</i> , Jenkins & Evermann	1852
<i>inermis</i> , Cuvier & Valenciennes	1853
<i>nematophthalmus</i> , (Günther)	2861

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER LORICATI—Continued.	
<i>Family Scorpaenidae</i> —Continued.	
Genus <i>Pontinus</i> , Poey	1854
<i>macrolepis</i> , Goode & Bean	1855
<i>castor</i> , Poey	1856
<i>pollux</i> , Poey	1857
<i>rathbuni</i> , Goode & Bean	1857
<i>longispinis</i> , Goode & Bean	1858
<i>sierra</i> (Gilbert)	1859
Genus <i>Setarches</i> , Johnson	1860
<i>parmatus</i> , Goode	1860
<i>Family Anoplopomatidae</i>	1861
Genus <i>Anoplopoma</i> , Ayres	1861
<i>fimbria</i> (Pallas)	1862
Genus <i>Eriilepis</i> , Gill	1862
<i>zonifer</i> (Lockington)	1863
<i>Family Hexagrammidae</i>	1863
Genus <i>Pleurrogrammus</i> , Gill	1864
<i>monoptygius</i> (Pallas)	1864
Genus <i>Hexagrammos</i> (Steller) <i>Tilesius</i>	1866
<i>decagrammus</i> (Pallas)	1867
<i>octogrammus</i> (Pallas)	1869
<i>stelleri</i> , <i>Tilesius</i>	1871
<i>supercliosus</i> (Pallas)	1872
<i>lagocephalus</i> (Pallas)	1873
Genus <i>Ophiodon</i> , Girard	1875
<i>elongatus</i> , Girard	1875
Genus <i>Zaniolepis</i> , Girard	1876
<i>latipinnis</i> , Girard	1876
<i>frenatus</i> , Eigenmann	1877
Genus <i>Oxylebius</i> , Gill	1878
<i>pictus</i> , Gill	1878
<i>Family Cottidae</i>	1879
Genus <i>Jordania</i> , Starks	1884
<i>zonope</i> , Starks	1884
Genus <i>Paricelinus</i> , Eigenmann & Eigenmann	1885
<i>hopliticus</i> , Eigenmann & Eigenmann	1886
Genus <i>Alcidea</i> , Jordan & Evermann	1886
<i>tibourbi</i> (Gilbert)	1887
Genus <i>Scorpenichthys</i> , Girard	1889
<i>marmoratus</i> (Ayres)	1889
Genus <i>Chitonotus</i> , Lockington	1889
<i>pudgetensis</i> (Steindachner)	1890
Genus <i>Tarandichthys</i> , Jordan & Evermann	1891
<i>cavifrons</i> (Gilbert)	1891
<i>filamentosus</i> (Gilbert)	1892
<i>tenula</i> (Gilbert)	1893
Genus <i>Icelinus</i> , Jordan	1894
<i>fimbriatus</i> , Gilbert	1894
<i>oculatus</i> , Gilbert	1895
<i>borealis</i> , Gilbert	1896
<i>quadriseriatus</i> (Lockington)	1897
<i>strabo</i> , Starks	1897
Genus <i>Astrolytes</i> , Jordan & Starks	1898
<i>notospilotus</i> (Girard)	1899
<i>fenestralis</i> (Jordan & Gilbert)	1899

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER LORICATI—Continued.	
<i>Family Cottidae</i> —Continued.	
Genus <i>Archistes</i> , Jordan & Gilbert	1900
<i>plumarius</i> , Jordan & Gilbert	1900
Genus <i>Artedius</i> , Girard	1902
<i>lateralis</i> (Girard)	1902
<i>asperulus</i> , Starks	1903
Genus <i>Axyrias</i> , Starks	1903
<i>harringtoni</i> , Starks	1904
Genus <i>Arteediellus</i> , Jordan	1905
<i>uncinatus</i> (Reinhardt)	1905
<i>atlanticus</i> , Jordan & Evermann	1906
<i>pacificus</i> , Gilbert	1906
Genus <i>Ruscarius</i> , Jordan & Starks	1908
<i>meanyi</i> , Jordan & Starks	1908
Genus <i>Rastrinus</i> , Jordan & Evermann	1909
<i>scutiger</i> (Bean)	1909
Genus <i>Icelus</i> , Kröyer	1911
<i>bicornis</i> (Reinhardt)	1911
<i>spiniger</i> , Gilbert	1914
<i>euryops</i> , Beau	1915
<i>vicinalis</i> , Gilbert	1916
<i>canaliculatus</i> , Gilbert	1917
<i>australis</i> , Eigenmann & Eigenmann	1918
Genus <i>Radulinus</i> , Gilbert	1919
<i>beleoides</i> , Gilbert	1919
<i>asprellus</i> , Gilbert	1920
Genus <i>Stelgistrum</i> , Jordan & Gilbert	1921
<i>stejnegeri</i> , Jordan & Gilbert	1921
Genus <i>Triglops</i> , Reinhardt	1923
<i>pingeli</i> , Reinhardt	1923
<i>beani</i> , Gilbert	1924
<i>scepticus</i> , Gilbert	1925
Genus <i>Sternias</i> , Jordan & Evermann	1926
<i>xenostethus</i> (Gilbert)	1927
Genus <i>Prionistius</i> , Bean	1927
<i>macellus</i> , Bean	1928
Genus <i>Elannura</i> , Gilbert	1930
<i>forficata</i> , Gilbert	1930
Genus <i>Melletes</i> , Bean	1932
<i>papilio</i> , Bean	1932
Genus <i>Hemilepidotus</i> , Cuvier	1934
<i>jordani</i> , Bean	1934
<i>hemilepidotus</i> (Tilesius)	1935
Genus <i>Calyceilepidotus</i> , Ayres	1936
<i>sphinosus</i> , Ayres	1937
Genus <i>Enophrys</i> , Swainson	1937
Subgenus <i>Aspicottus</i> , Girard	1938
<i>bison</i> (Girard)	1938
Subgenus <i>Enophrys</i>	1938
<i>claviger</i> (Cuvier & Valenciennes)	1938
Genus <i>Ceratocottus</i> , Gill	1939
<i>lucasi</i> , Jordan & Gilbert	1940
<i>dicerous</i> (Pallas)	1940

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SCHORDER LORICATI—Continued.	
Family Cottidae—Continued.	
Genus <i>Cottus</i> (Arteell) Linnaeus	1941
Subgenus <i>Pegedelctis</i> , Rafinesque	1944
asper, Richardson	1944
gulosus (Girard)	1944
evermanni, Gilbert	1945
rhothens, Rosa Smith	1946
shasta, Jordan & Starks	1947
punctulatus (Gill)	1948
semiscaber (Cope)	1949
ictulops (Rafinesque)	1950
Subgenus <i>Tauridea</i> , Jordan & Rice	1952
ricel, Nelson	1952
Subgenus <i>Cottus</i>	1953
onychus, Eigenmann & Eigemann	1953
polllearis (Jordan & Gilbert)	1953
cognatus, Richardson	1954
perplexus, Gilbert & Evermann	1955
klamathensis, Gilbert	1955
aleuticus, Gilbert	1957
minutus, Pallas	1958
beldingii, Eigenmann & Eigemann	1958
philomus, Eigenmann & Eigenmann	1959
anna, Jordan & Starks	1960
spilotus (Cope)	1961
lelopomus, Gilbert & Evermann	1962
princeps, Gilbert	1962
Genus <i>Uranidea</i> , De Kay	1963
bendirei (Bean)	1964
greeni, Gilbert & Culver	1965
marginalis, Bean	1965
tenuis, Evermann & Seale	1966
franklini (Agassiz)	1967
kumlienii, Hoy	1967
gracilis (Heckel)	1968
formosa (Girard)	1969
hoyi, Putnam	1969
Genus <i>Myoxocephalus</i> (Steller) Tilesius	1970
Subgenus <i>Acanthocottus</i> , Girard	1971
bubalis (Euphrasen)	1971
reneus (Mitchill)	1972
scorpioides (Fabricius)	1973
scorpius (Linnaeus)	1974
greenlandicus (Cuvier & Valenciennes)	1974
octodecimspinosus (Mitchill)	1976
Subgenus <i>Myoxocephalus</i>	1976
polyacanthocephalus, Pallas	1976
jaok (Cuvier & Valenciennes)	1977
verrucosus (Bean)	1979
axillaris (Gill)	1980
stelleri, Tilesius	1981
medinicus, B. A. Bean	1983
nivosus (Herzenstein)	1984
niger (Bean)	1985

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER LORICATI—Continued.	
Family <i>Cottidae</i> —Continued.	
Genus <i>Megalocottus</i> , Gill	1987
<i>platycephalus</i> (Pallas)	1987
<i>laticeps</i> (Gilbert)	1988
Genus <i>Zesticelus</i> , Jordan & Evermann	1990
<i>profundorum</i> (Gilbert)	1990
Genus <i>Dasy cottus</i> , Bean	1991
<i>setiger</i> , Bean	1991
Genus <i>Cottunculus</i> , Collett	1992
<i>micros</i> , Collett	1992
<i>thomsonii</i> (Günther)	1993
Genus <i>Malacocottus</i> , Bean	1994
<i>zonurus</i> , Bean	1994
Genus <i>Argyrocottus</i> , Herzenstein	1995
<i>zanderi</i> , Herzenstein	1995
Genus <i>Porocottus</i> , Gill	1996
<i>sellaris</i> (Gilbert)	1996
<i>quadratus</i> , B. A. Bean	1998
<i>polaris</i> (Sabine)	1998
<i>quadrifilis</i> , Gill	1999
<i>tentaculatus</i> (Kner)	2000
<i>bradfordi</i> , Rutter	2862
Genus <i>Oncocottus</i> , Gill	2000
<i>quadricornis</i> (Linnaeus)	2001
<i>hexacornis</i> (Richardson)	2002
Genus <i>Triglopsis</i> , Girard	2005
<i>thompsoni</i> , Girard	2005
Genus <i>Gymno canthus</i> , Swainson	2006
<i>distilliger</i> (Pallas)	2006
<i>tricuspidis</i> (Reinhardt)	2008
<i>galeatus</i> (Bean)	2010
Genus <i>Leiocottus</i> , Girard	2010
<i>birnau</i> , Girard	2011
Genus <i>Leptocottus</i> , Girard	2011
<i>ermatus</i> , Girard	2012
Genus <i>Clinocottus</i> , Gill	2012
<i>analis</i> (Girard)	2012
Genus <i>Oligocottus</i> , Girard	2013
<i>maculosus</i> , Girard	2013
<i>borealis</i> , Jordan & Snyder	2014
<i>snyderi</i> , Greeley	2871
Genus <i>Signistes</i> , Rutter	2863
<i>caulias</i> , Rutter	2863
Genus <i>Blennicottus</i> , Gill	2017; 2864
Subgenus <i>Oxycottus</i> , Jordan & Evermann	2015; 2864
<i>enticeps</i> (Gilbert)	2015; 2864
<i>embryum</i> (Jordan & Starks)	2016; 2864
Subgenus <i>Blennicottus</i>	2016; 2864
<i>recalvus</i> , Greeley	3178
<i>globiceps</i> (Girard)	2017
<i>bryosus</i> , Jordan & Starks	2017
Genus <i>Ruschulus</i> , Greeley	3179
<i>rimensis</i> , Greeley	3179
Genus <i>Dialarchus</i> , Greeley	3180
<i>snyderi</i> , Greeley	3181

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER LORICATI—Continued.	
<i>Family Cottidae</i> —Continued.	
Genus <i>Eximia</i> , Greeley	3182
<i>rubellio</i> , Greeley	3182
Genus <i>Histiocottus</i> , Gill	2018
<i>bilobus</i> (<i>Cuvier & Valenciennes</i>).....	2018
Genus <i>Blepsias</i> , Cuvier	2018
<i>cirrhosus</i> (<i>Pallas</i>).....	2018
Genus <i>Nauticus</i> , Jordan & Evermann	2019
<i>pribiloviensis</i> , Jordan & Gilbert	2019
Genus <i>Nautichthys</i> , Girard	2020
<i>oenofasciatus</i> (<i>Girard</i>).....	2021
Genus <i>Ulca</i> , Jordan & Evermann	2021
<i>marmorata</i> (<i>Bean</i>)	2021
Genus <i>Hemitripterus</i> , Cuvier	2022
<i>americanus</i> (<i>Gmelin</i>).....	2023
<i>cavifrons</i> , Lockington	2023
Genus <i>Synchirus</i> , Bean	2023
<i>gilli</i> , Bean.....	2024
Genus <i>Ascelichthys</i> , Jordan & Gilbert	2024
<i>rhodorus</i> , Jordan & Gilbert	2025
Genus <i>Psychrolutes</i> , Günther	2025
<i>paradoxus</i> , Günther	2026
Genus <i>Gilbertidia</i> , Berg	2027; 3183
<i>sigillata</i> , Jordan & Starks.....	2028
<i>Family Ramphocottidae</i>	2029
Genus <i>Ramphocottus</i> , Günther	2030
<i>richardsoni</i> , Günther	2030
<i>Family Agonidae</i>	2031
Genus <i>Percis</i> , Scopoli.....	2033
<i>japonicus</i> (<i>Pallas</i>)	2034
Genus <i>Agonomalus</i> , Guichenot	2036
<i>proboscidalis</i> (<i>Valenciennes</i>)	2037
Genus <i>Hypsagonus</i> , Gill	2038
<i>quadricornis</i> (<i>Cuvier & Valenciennes</i>)	2038
Genus <i>Stellerina</i> , Cramer	2041
<i>xyosterna</i> (<i>Jordan & Gilbert</i>)	2042
Genus <i>Ocea</i> , Jordan & Evermann	2043
<i>verrucosa</i> (<i>Lockington</i>)	2043
<i>dodecaedron</i> (<i>Tilesius</i>)	2044
Genus <i>Brachyopsis</i> , Gill	2046
<i>rostratus</i> (<i>Tilesius</i>)	2046
<i>segallensis</i> (<i>Tilesius</i>)	2048
Genus <i>Pallasina</i> , Cramer	2048
<i>barbata</i> (<i>Steindachner</i>).....	2049
<i>aix</i> , Starks.....	2050
Genus <i>Leptagonus</i> , Gill.....	2052
<i>deagonus</i> (<i>Bloch & Schneider</i>)	2052
Genus <i>Podothecus</i> , Gill	2054
<i>aceipiter</i> , Jordan & Starks.....	2055
<i>hamiltoni</i> , Jordan & Gilbert	2056
<i>gilberti</i> (<i>Colletti</i>)	2058
<i>thompsoni</i> , Jordan & Gilbert	2060
<i>acipenserinus</i> (<i>Tilesius</i>).....	2061
<i>veterinus</i> , Jordan & Starks	2063

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER LORICATI—Continued.	
<i>Family Agonidae</i> —Continued.	
Genus <i>Agonus</i> , Bloch & Schneider.....	2064
<i>cataphractus</i> , Linnaeus	2065
Genus <i>Stolegis</i> , Cramer	2067
<i>vulsa</i> (Jordani & Gilbert)	2067
Genus <i>Avertunculus</i> , Jordani & Starks	2069
<i>emmeline</i> , Jordan & Starks	2069
<i>sterletus</i> , Gilbert	2071
Genus <i>Sarrtori</i> , Cramer	2072
<i>frenatus</i> , Gilbert	2073
<i>leptorhynchus</i> (Gilbert)	2075
Genus <i>Xystes</i> , Jordan & Starks	2076
<i>axinophrys</i> , Jord. & Starks	2076
Genus <i>Bathyagonus</i> , Gilbert	2077
<i>nigripinnis</i> , Gilbert	2078
Genus <i>Xenochirrus</i> , Gilbert	2079
<i>pentacanthus</i> , Gilbert	2080
<i>alascannus</i> , Gilbert	2081
<i>latifrons</i> , Gilbert	2082
<i>triacanthus</i> , Gilbert	2084
Genus <i>Odontopyxis</i> , Lockington	2085
<i>triapinosus</i> , Lockington	2085
Genus <i>Bothragonus</i> , Gill	2086
<i>swaini</i> (Steindachner)	2086
Genus <i>Aspidophoroides</i> , Lacépède	2086
Subgenus <i>Ulcina</i> , Cramer	2089
<i>olrikii</i> , Lütken	2089
<i>guntheri</i> , Bean	2090
<i>monopterygius</i> (Bloch)	2091
<i>bartoni</i> , Gilbert	2092
Subgenus <i>Anoplagonus</i> , Gill	2093
<i>inermis</i> , Günther	2093
<i>Family Cyclopteridae</i>	2094
Genus <i>Cyclopterus</i> (Artedi) Linnaeus	2096
<i>bumpus</i> , Linnaeus	2096
Genus <i>Eumicrotremus</i> , Gill	2097
<i>spinosus</i> (Müller)	2098
<i>orbis</i> (Günther)	2099
Genus <i>Lethotremus</i> , Gilbert	2100
<i>mutiens</i> , Gillett	2101
<i>vinolentus</i> , Jordan & Starks	2101
Genus <i>Cyclopteroides</i> , Garman	2102
<i>gyrinops</i> , Garman	2102
Genus <i>Cyclopterichthys</i> , Steindachner	2103
<i>ventricosus</i> (Pallas)	2104
Genus <i>Liparops</i> , Gilman	2104
<i>stelleri</i> (Pallas)	2104
<i>Family Liparidae</i>	2105
Genus <i>Neoliparis</i> , Steindachner	2106
<i>atlanticus</i> , Jordan & Evermann	2107
<i>rutteri</i> , Gilbert & Snyder	2108
<i>callyodon</i> (Pallas)	2110
<i>mucosus</i> (Ayres)	2111
<i>dorsalis</i> , Jordan & Starks	2111
<i>greeni</i> , Jordan & Starks	2112

LAS
SUB
OR
S

G

Bul

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER LORICATI—Continued.	
Family <i>Liparidae</i> —Continued.	
Genus <i>Neoliparis</i> , Steindachner—Continued.	
<i>fissuratus</i> , Starks	2113
Genus <i>Liparis</i> (Arvedi) Scopoli	2114
Subgenus <i>Liparis</i>	2116
<i>liparis</i> (Linnæus)	2116
<i>cyclopus</i> , Günther	2118
<i>fucensis</i> , Gilbert	2119
<i>tunicatus</i> , Reinhardt	2120
<i>agassizii</i> , Putnam	2121
<i>herculeanus</i> , Scofield	2123
<i>dennysi</i> , Jordan & Starks	2124
<i>cyclostigma</i> , Gilbert	2125
Subgenus <i>Lyoliparis</i> , Jordan & Evermann	2126
<i>pulchellus</i> , Ayres	2126
Subgenus <i>Actinochir</i> , Gill	2127
<i>major</i> (Gill)	2127
Genus <i>Crystallichthys</i> , Jordan & Gilbert	2864
<i>mirabilis</i> , Jordan & Gilbert	2865
Genus <i>Bathyphasma</i> , Gilbert	2128
<i>ovigerum</i> , Gilbert	2128
Genus <i>Careproctus</i> , Krüger	2129
Subgenus <i>Cataphracta</i> , Jordan & Evermann	2131
<i>sinus</i> , Gilbert	2131
Subgenus <i>Careproctus</i>	2131
<i>colletti</i> , Gilbert	2131
<i>phasiana</i> , Gilbert	2132
<i>spectrum</i> , Bean	2133
<i>reinhardti</i> (Krüger)	2133
<i>rampilla</i> (Goode & Bean)	2134
<i>ostentum</i> , Gilbert	2134
<i>gelatinosus</i> (Pallas)	2134
Subgenus <i>Allocirr</i> , Jordan & Evermann	2135
<i>melanurus</i> , Gilbert	2135
Subgenus <i>Allineetes</i> , Jordan & Evermann	2136; 2866
<i>ectenes</i> , Gilbert	2136
Genus <i>Prognurus</i> , Jordan & Evermann	2866
<i>cypselurus</i> , Jordan & Gilbert	2866
Genus <i>Gyrinichthys</i> , Gilbert	2137
<i>mintyrenus</i> , Gilbert	2137
Genus <i>Amitra</i> , Goode	2138
<i>liparina</i> , Goode	2138
Genus <i>Paraliparis</i> , Collett	2139
Subgenus <i>Paraliparis</i>	2140
<i>böholensis</i> , Gilbert	2140
Subgenus <i>Amitrichthys</i> , Jordan & Evermann	2141
<i>cephalus</i> , Gilbert	2141
<i>rosaceus</i> , Gilbert	2142
<i>mento</i> , Gilbert	2142
<i>copei</i> , Goode & Bean	2143
<i>dactylopus</i> , Gilbert	2144
Subgenus <i>Hilgendorffia</i> , Goode & Bean	2144
<i>moehri</i> , Gilbert	2144
Genus <i>Rhinoliparis</i> , Gilbert	2145
<i>barbifilifer</i> , Gilbert	2145

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER CRANIOMI.	2146
<i>Family Triglidae</i>	2147
Genus <i>Prionotus</i> , Lacépède	2148
Subgenus <i>Gurnardus</i> , Jordan & Evermann	2152
<i>birostratus</i> , Richardson	2152
<i>gymnostethus</i> , Gilbert	2153
<i>xenisma</i> , Jordan and Bollman	2154
<i>loxias</i> , Jordan	2155
Subgenus <i>Merulinus</i> , Jordan & Evermann	2156
<i>carolinus</i> (Linnaeus)	2156
<i>scitulus</i> , Jordan	2157
<i>roseus</i> , Jordan & Evermann	2158
<i>alatus</i> , Goode & Bean	2159
Subgenus <i>Prionotus</i>	2160
<i>miles</i> , Jenyns	2160
<i>stephanophrys</i> , Lockington	2161
<i>quiescens</i> , Jordan & Bollman	2161
<i>albirostris</i> , Jordan & Bollman	2163
<i>rubio</i> , Jordan	2164
<i>ophryas</i> , Jordan & Swain	2164
<i>stearnsi</i> , Jordan & Swain	2166
<i>strigatus</i> (Cuvier & Valenciennes)	2167
<i>evolans</i> (Linnaeus)	2168
<i>punctatus</i> (Bloch)	2169
<i>beanii</i> , Goode	2170
<i>tribulus</i> (Cuvier)	2171
<i>horrens</i> , Richardson	2172
Genus <i>Bellator</i> , Jordan & Evermann	2173
<i>militaris</i> (Goode & Bean)	2173
<i>egretta</i> (Goode & Bean)	2174
Genus <i>Trigla</i> (Artedi) Linnaeus	2176
<i>cuculus</i> , Linnaeus	2177
<i>Family Peristediidae</i>	2177
Genus <i>Peristedion</i> , Lacépède	2178
<i>miniatum</i> , Goode	2178
<i>longispathum</i> , Goode & Bean	2178
<i>gracile</i> , Goode & Bean	2179
<i>platycephalum</i> , Goode & Bean	2180
Genus <i>Vulsiculus</i> , Jordan & Evermann	2181
<i>imberbis</i> (Poey)	2181
<i>Family Cephalacanthidae</i>	2182
Genus <i>Cephalacanthus</i> , Lacépède	2183
<i>volitans</i> (Linnaeus)	2183
Group Gobloidea	2184
<i>Family Callionymidae</i>	2184
Genus <i>Callionymus</i> , Linnaeus	2185
<i>bairdi</i> , Jordan	2185
<i>ngassizil</i> , Goode & Bean	2186
<i>calliurus</i> , Eigenmann & Eigenmann	2187
<i>pauciradiatus</i> , Gill	2188
<i>Family Gobiidae</i>	2188
Genus <i>Ioglossus</i> , Beau	2192
<i>caillurus</i> , Beau	2193
Genus <i>Philyampus</i> , Cuvier & Valenciennes	2194
<i>dormitor</i> (Lacépède)	2194
<i>lateralis</i> , Gill	2195

CLASS PISCES—Continued.

SUBCLASS TELEOSTOMI—Continued.

ORDER PLECTOGNATHI—Continued.

SUBORDER CRANIOMI—Continued.

Family Gobiidae—Continued.

	Page.
46	
47	
48	
152	
152	
153	
154	
155	
156	
156	
157	
158	
159	
2160	
2160	
2161	
2161	
2163	
2164	
2164	
2166	
2167	
2168	
2169	
2170	
2171	
2172	
2173	
2173	
2174	
2174	
2175	
2176	
2177	
2177	
2178	
2178	
2179	
2179	
2180	
2181	
2181	
2182	
2183	
2183	
2184	
2184	
2185	
2185	
2186	
2187	
2188	
2188	
2189	
2189	
2190	
2191	
2191	
2192	
2193	
2194	
2194	
2195	
Genus <i>Dormitator</i> , Gill.....	2195
<i>maenulatus</i> (Bloch).....	2196
Genus <i>Guavina</i> , Bleeker.....	2198
<i>guavina</i> (Cuvier & Valenciennes).....	2198
Genus <i>Electris</i> (Gronow) Bloch & Schneider.....	2199
<i>amblyopsis</i> Cope	2199
<i>abaeurus</i> , Jordan & Gilbert.....	2200
<i>pisonis</i> (Gmelin).....	2200
<i>perniger</i> (Cope).....	2201
<i>pictus</i> (Kner & Steindachner)	2201
Genus <i>Alexurus</i> , Jordan	2202
<i>armiger</i> , Jordan & Richardson.....	2203
Genus <i>Erotelis</i> , Poey	2203
<i>smaragdus</i> (Cuvier & Valenciennes).....	2204
Genus <i>Gymnoleotris</i> , Bleeker.....	2204
<i>semimundus</i> (Günther).....	2204
Genus <i>Chriolepis</i> , Gilbert.....	2205
<i>minutillus</i> , Gilbert	2205
Genus <i>Sicydium</i> , Cuvier & Valenciennes.....	2205
<i>plumieri</i> (Bloch).....	2206
<i>antillarum</i> , Ogilvie-Grant	2206
<i>vineente</i> , Jordan & Evermann.....	2207
<i>punctatum</i> , Perugia.....	2207
<i>caguitae</i> , Evermann & Marsh.....	3183
Genus <i>Cotylopus</i> , Guichenot	2207
Subgenus <i>Sicyosus</i> , Jordan & Evermann.....	2207; 2867
<i>gymnogaster</i> (Ogilvie-Grant).....	2207
<i>salvini</i> (Ogilvie-Grant).....	2208
Genus <i>Evorthodus</i> , Gill.....	2208
<i>breviceps</i> , Gill	2208
Genus <i>Lophogobius</i> , Gill.....	2209
<i>cyprinoides</i> (Pallas).....	2209
Genus <i>Gobius</i> (Artedi) Linnaeus.....	2210
Subgenus <i>Gobius</i>	2216
<i>soporator</i> , Cuvier & Valenciennes.....	2216
Subgenus <i>Ctenogobius</i> , Gill	2218
<i>nicholsii</i> , Bean	2218
<i>eigenmanni</i> , Garman	2218
<i>glaucofrenum</i> (Gill).....	2219
<i>manglicola</i> , Jordan & Starks.....	2220
<i>stigmaturus</i> , Goode & Bean.....	2220
<i>quadriforus</i> , Cuvier & Valenciennes.....	2221
<i>shufeldti</i> , Jordan & Eigenmann.....	2221
<i>boleosoma</i> , Jordan & Gilbert.....	2221
<i>fasciatus</i> (Gill)	2222
<i>encionius</i> , Jordan & Gilbert.....	2223
<i>stigmaticus</i> (Poey)	2224
<i>lyricus</i> , Girard	2224
<i>garmani</i> , Eigenmann & Eigenmann	2225
<i>zebra</i> , Gilbert	2226; 2867
Subgenus <i>Euctenogobius</i> , Gill	2226
<i>pocyt</i> , Steindachner	2226
<i>badius</i> (Gill)	2227

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER CRANIOMI—Continued.	
<i>Family Gobiidae—Continued.</i>	
Genus <i>Gobius</i> (Artedi) Linnaeus—Continued.	
Subgenus <i>Gobionellus</i> , Girard	2227
<i>mierodon</i> , Gilbert	2227
<i>smaragdus</i> , Cuvier & Valenciennes.....	2227
<i>strigatus</i> , O'Shangnessy	2228
<i>sagittula</i> (Günther)	2228
<i>hastatus</i> (Girard)	2229
<i>oceaniensis</i> , Pallas.....	2230
<i>bayamonensis</i> , Evermann & Marsh.....	3184
Subgenus <i>Lythrypnus</i> , Jordan & Evermann.....	2230
<i>dalli</i> , Gilbert	2230
Genus <i>Garmannia</i> , Jordan & Evermann.....	2231
Subgenus <i>Garmannia</i>	2232
<i>paradoxa</i> (Günther)	2232
<i>hemigymna</i> (Eigenmann & Eigenmann).....	2233
Subgenus <i>Enypnias</i> , Jordan & Evermann.....	2233
<i>seminuda</i> (Günther)	2233
Genus <i>Awaous</i> , Steindachner	2234
<i>flavus</i> (Cuvier & Valenciennes)	2235
<i>nelsoni</i> , Evermann	2235
<i>taiasica</i> (Lichtenstein)	2236
<i>mexicanus</i> (Günther)	2237
Genus <i>Bolmanna</i> , Jordan	2237
<i>ocellata</i> , Gilbert	2238
<i>chlamydes</i> , Jordan	2238
<i>maeropoma</i> , Gilbert	2239
<i>stigmatura</i> , Gilbert	2239
<i>boqueronensis</i> , Evermann & Marsh.....	3185
Genus <i>Aboma</i> , Jordan & Starks	2240
<i>etheostoma</i> , Jordan & Starks	2240
<i>luretiae</i> (Eigenmann & Eigenmann)	2241
<i>chiquita</i> (Jenkins & Evermann)	2241
Genus <i>Microgobius</i> , Poey	2242
<i>gulosus</i> (Girard)	2243
<i>eulepis</i> , Eigenmann & Eigenmann	2244
<i>meeki</i> , Evermann & Marsh	3185
<i>thalassinus</i> , Jordan & Gilbert	2245
<i>signatus</i> , Poey	2246
Genus <i>Zalypnus</i> , Jordan & Evermann	2246
<i>cyclolepis</i> (Gilbert)	2246
<i>emblematius</i> (Jordan & Gilbert)	2247
Genus <i>Eucyclogobius</i> , Gill	2248
<i>newberryi</i> (Girard)	2248
Genus <i>Lepidogobius</i> , Gill	2249
<i>lepidus</i> (Girard)	2249
Genus <i>Gilllichthys</i> , Cooper	2249
<i>mirabilis</i> , Cooper	2250
<i>detrusus</i> , Gilbert & Sebold	2251
Genus <i>Quietula</i> , Jordan & Evermann	2251
<i>y-cauda</i> (Jenkins & Evermann)	2251
Genus <i>Ilypnus</i> , Jordan & Evermann	2253
<i>gilberti</i> (Eigenmann & Eigenmann)	2253


CLASS PISCES—Continued.	Page.	
SUBCLASS TELEOSTOMI—Continued.		
ORDER PLECTOGNATHI—Continued.		
SUBORDER CRANIOMI—Continued.		
Family Gobiidae—Continued.		
227	Genus Clevelandia, Eigenmann & Eigenmann.....	2254
227	<i>ios</i> (Jordan & Gilbert).....	2254
227	<i>rosea</i> , Jordan & Evermann.....	2255
227	Genus Evermannia, Jordan.....	2256
228	<i>longipinnis</i> (Steindachner).....	2256
228	<i>zosterura</i> , (Jordan & Gilbert).....	2256
229	Genus Gobiosoma, Girard.....	2257
229	<i>histrio</i> , Jordan	2258
229	<i>molestem</i> , Girard	2258
229	<i>bosci</i> (Lacépède)	2259
229	<i>crescentale</i> , Gilbert	2259
229	<i>multifasciatum</i> , Steindachner.....	2260
229	Genus Barbulifer, Eigenmann & Eigenmann.....	2260
229	<i>ceutheecus</i> (Jordan & Gilbert).....	2260
229	Genus Typhlogobius, Steindachner	2261
229	<i>californiensis</i> , Stelndachner	2262
229	Genus Tyntlastes, Günther	2262
229	<i>brevis</i> (Günther)	2262
229	<i>sagitta</i> (Günther)	2263
229	Genus Gobioloides, Lacépède.....	2263; 2868
229	<i>broussonnetii</i> , Lacépède.....	2263
229	<i>peruanus</i> (Steindachner)	2264
229	Genus Cayennia, Sauvage.....	2265
229	<i>guichenoti</i> , Sauvage.....	2265
229	SUBORDER DISCOCEPHALI	2265
229	Family Echeneidae	2265
229	Genus Phtheirichthys, Gill	2268
229	<i>lineatus</i> (Menzies).....	2268
229	Genus Echeneis (Artedi) Linnaeus.....	2268
229	<i>naucrates</i> , Linnaeus	2269
229	<i>naucratoeides</i> , Zaiew	2270
229	Genus Remilegla, Gill.....	2270
229	<i>australis</i> (Bennett).....	2270
229	Genus Remora, Gill	2271
229	Subgenus Remora.....	2271
229	<i>remora</i> (Linnaeus).....	2271
229	Subgenus Remorina, Jordan & Evermann	2272
229	<i>albescens</i> (Temminck & Schlegel).....	2272
229	Subgenus Remoropsis, Gill	2272
229	<i>brachyptera</i> (Lowe)	2272
229	Genus Rhombochirrus, Gill.....	2273
229	<i>osteochir</i> (Cuvier).....	2273
229	Group Trachinidae	2273
229	Family Malacanthidae	2274
229	Genus Malacanthus, Cuvier	2275
229	<i>plumieri</i> (Bloch)	2275
229	Genus Caulolatilus, Gill	2276
229	<i>princeps</i> (Jenyns).....	2276
229	<i>microps</i> , Goode & Bean	2277
229	<i>cyanops</i> , Poey	2278
229	Genus Lopholatilus, Goode & Bean	2278
229	<i>chamaeleonticeps</i> , Goode & Bean	2278

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER DISCOCEPHALI—Continued.	
Family <i>Opistognathidae</i>	2279
Genus <i>Opistognathus</i> , Cuvier	2280
<i>lonchurum</i> , Jordan & Gilbert	2281
<i>punctatum</i> , Peters	2281
<i>macrognathum</i> , Poey	2281
<i>ommatum</i> , Jenkins & Evermann	2282
Genus <i>Gnathoprops</i> , Gill	2283
<i>scops</i> , Jenkins & Evermann	2283
<i>maxillosa</i> (Poey)	2284
<i>macrorops</i> (Poey)	2284
<i>rhomalea</i> (Jordan & Gilbert)	2285
<i>snyderi</i> , Jordan & Evermann	2285
<i>mystacina</i> , Jordan	2286
Genus <i>Lonchopisthus</i> , Gill	2286
<i>micrognathus</i> (Poey)	2287
Family <i>Bathynasteridae</i>	2287
Genus <i>Bathymaster</i> , Cope	2288
<i>signatus</i> , Cope	2288
Genus <i>Ronquilus</i> , Jordan & Starks	2289
<i>jordani</i> (Gilbert)	2289
Genus <i>Rathbunella</i> , Jordan & Evermann	2289
<i>hypolecta</i> (Gilbert)	2290
Family <i>Chiasmodontidae</i>	2291
Genus <i>Chiasmodon</i> , Johnson	2291
<i>niger</i> , Johnson	2291
Genus <i>Pseudoscopelus</i> , Lütken	2292
<i>scriptus</i> , Lütken	2292
Family <i>Chænichthyidae</i>	2293
Genus <i>Hypsicometes</i> , Goode	2293
<i>gobioides</i> , Goode	2294
Family <i>Trichodontidae</i>	2295
Genus <i>Trichodon</i> (Steller) Cuvier	2295
<i>trichodon</i> (Tilesius)	2295
Genus <i>Arctoscopus</i> , Jordan & Evermann	2297
<i>japonicus</i> (Steindachner)	2297
Family <i>Dactyloscopidae</i>	2297
Genus <i>Gillellus</i> , Gilbert	2298
<i>semicinctus</i> , Gilbert	2298
<i>arenicola</i> , Gilbert	2299
<i>ornatus</i> , Gilbert	2299
Genus <i>Dactyloscopus</i> , Gill	2300
Subgenus <i>Dactyloscopus</i>	2301
<i>pectoralis</i> , Gill	2301
<i>tridigitatus</i> , Gill	2301
<i>poeyi</i> , Gill	2302
<i>luuaticus</i> , Gilbert	2302
Subgenus <i>Esoscopus</i> , Jordan & Evermann	2303
<i>zelotes</i> , Jordan & Gilbert	2303
Genus <i>Dactylagnus</i> , Gill	2304
<i>mundus</i> , Gill	2304
Genus <i>Myxodagnus</i> , Gill	2305
<i>opercularis</i> , Gill	2305

	Page.	
CLASS PISCES—Continued.		
SUBCLASS TELEOSTOMI—Continued.		
ORDER PLECTOGNATHI—Continued.		
SUBORDER DISCOCEPHALI—Continued.		
Family <i>Uranoscopidae</i>		
279	Genus <i>Astroscopus</i> , Brevoort	2305
280	<i>y-gracum</i> (Cuvier & Valenciennes)	2306
281	<i>zephyreus</i> , Gilbert & Starks	2307
281	<i>guttatus</i> (Abbott)	2309
282	Genus <i>Kathetostoma</i> , Günther	2311
283	<i>averruncus</i> , Jordan & Bollman	2311
283	<i>albifigatum</i> , Bean	2312
284	SUBORDER HAPLODOCII	2313
284	Family <i>Batrachoididae</i>	2313
2285	Genus <i>Batrachoides</i> , Lacépède	2314; 2668
2285	<i>surinamensis</i> (Bloch & Schneider)	2314
2286	<i>pacifici</i> (Günther)	2314
2286	Genus <i>Opsanus</i> , Rafinesque	2315
2287	<i>tau</i> (<i>Liunicus</i>)	2315
2287	<i>pardus</i> (Goode & Bean)	2316
2288	Genus <i>Porichthys</i> , Girard	2317
2288	<i>porosissimus</i> (Cuvier & Valenciennes)	2319
2289	<i>notatus</i> , Girard	2321
2289	<i>margaritatus</i> (Richardson)	2322
2289	Genus <i>Thallassophryne</i> , Günther	2323
2290	<i>maculosa</i> , Günther	2324
2291	<i>reticulata</i> , Günther	2325
2291	Genus <i>Dæctor</i> , Jordan & Evermann	2325
2291	<i>dowi</i> (Jordan & Gilbert)	2325
2292	SUBORDER XENOPTERYGHII	2326
2292	Family <i>Gobiesocidae</i>	2326
2293	Genus <i>Caularchus</i> , Gill	2327
2293	<i>meandricus</i> (Girard)	2328
2294	Genus <i>Brysseteres</i> , Jordan & Evermann	2328
2295	<i>plumiger</i> (Gilbert)	2328
2295	Genus <i>Gobiesox</i> , Lacépède	2329
2295	Subgenus <i>Bryssophilus</i> , Jordan & Evermann	2330
2297	<i>papillifer</i> , Gilbert	2330
2297	Subgenus <i>Gobiesox</i>	2331
2297	<i>gyrinus</i> , Jordan & Evermann	2331
2298	<i>nigripinnis</i> (Peters)	2331
2298	<i>cephalus</i> , Lacépède	2332
2299	<i>tudes</i> , Richardson	2333
2299	<i>strumosus</i> , Cope	2333
2300	<i>virgatulus</i> , Jordan & Gilbert	2333
2301	<i>adustus</i> , Jordan & Gilbert	2334
2301	<i>funebris</i> , Gilbert	2334
2301	<i>pecilophthalmus</i> , Jenyns	2335
2302	<i>rhodospilus</i> , Günther	2335
2302	<i>macrophthalmus</i> , Günther	2335
2303	<i>cerasinus</i> , Cope	2336
2303	Subgenus <i>Sicyases</i> , Müller & Troschel	2336
2304	<i>erythrops</i> , Jordan & Gilbert	2336
2304	<i>rubiginosus</i> (Poey)	2337
2305	<i>carninus</i> (Poey)	2337
2305	<i>haeres</i> , Jordan & Bollman	2337
2306	<i>punctulatus</i> (Poey)	2338
2306	<i>fasciatus</i> (Peters)	2338

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER XENOPTERYGII—Continued.	
<i>Family Gobiesocidae</i> —Continued.	
Genus Rimicola, Jordan & Evermann.....	2338
muscicarum (Meek & Pierson).....	2338
elgenmanni (Gilbert)	2339
Genus Arbaciosa, Jordan & Evermann.....	2340
rhessodon (Rosa Smith).....	2340
humeralis (Gilbert)	2341
rupestris (Poey)	2341
zebra (Jordan & Gilbert).....	2341
eos (Jordan & Gilbert)	2343
Group Blennioidea	2343
<i>Family Blenniidae</i>	2344
Genus Enneanectes, Jordan & Evermann	2349
carminalis (Jordan & Gilbert).....	2350
Genus Gilliæ, Evermann & Marsh.....	3186
jordani, Evermann & Marsh	3186
Genus Dialommus, Gilbert.....	2868
fuscus, Gilbert	2868
Genus Heterostichus, Girard	2350
rostratus, Girard.....	2351
Genus Gibbonsia, Cooper	2351
elegans (Cooper)	2353
evides (Jordan & Gilbert).....	2352; 2869
Genus Neoclinus, Girard	2354
Subgenus Neoclinus	2354
blanchardi, Girard.....	2354
Subgenus Pterognathus, Girard	2355
satirdens, Girard	2355
Genus Malacoctenus, Gill	2356
ocellatus (Steindachner)	2356; 2869
varius (Poey).....	2357
macropus (Poey).....	2357
lugubris (Poey)	2357
culebræ, Evermann & Marsh	3187
gillii (Steindachner)	2358
puertoricensis, Evermann & Marsh	3189
bimaculatus (Steindachner)	2358
moorei, Evermann & Marsh.....	3188
delalandi (Cuvier & Valenciennes)	2358
versicolor (Poey)	2359
biguttatus (Cope)	2360
Genus Labrisomus, Swainson	2360
herminieri (Le Sueur)	2361
nuchipinnis (Quoy & Gaimard)	2362
xanti, Gill	2362
bucciferus, Poey.....	2363
microlepidotus, Poey	2363
Genus Miigerpes, Jordan & Evermann	2364
macrocephalus (Günther)	2364
Genus Gobioclinus, Gill	2364
gobio (Cuvier & Valenciennes)	2365
Genus Starksia, Jordan & Evermann	2365
crenophobates (Gilbert).....	2365
Genus Cryptotrema, Gilbert	2366
corallinum, Gilbert.....	2366

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER XENOPTERYGINI—Continued.	
<i>Family Blennidae</i> —Continued.	
Genus <i>Exernes</i> , Jordan & Evermann	2367
<i>asper</i> (Jenkins & Evermann)	2367
Genus <i>Anchenistius</i> , Evermann & Marsh	3190
<i>stahlii</i> , Evermann & Marsh.....	3190
Genus <i>Anchenopterus</i> , Günther	2369
Subgenus <i>Corallicola</i> , Jordan & Evermann.....	2369
<i>nigripinnis</i> (Steindachner)	2369
<i>altivelis</i> (Lockington)	2370
<i>marmoratus</i> (Steindachner)	2371
Subgenus <i>Anchenopterus</i>	2371
<i>affinis</i> (Steindachner)	2371
<i>monopthalmus</i> , Günther.....	2372
<i>integripinnis</i> (Rosa Smith)	2372
<i>albicaudus</i> , Evermann & Marsh.....	3191
<i>rubescens</i> , Evermann & Marsh.....	3191
<i>cingulatus</i> , Evermann & Marsh.....	3192
<i>fajardo</i> , Evermann & Marsh.....	3193
<i>fasciatus</i> (Steindachner)	2373
<i>nox</i> (Jordan & Gilbert)	2373
Genus <i>Paraclinus</i> , Mocquard.....	2374
<i>chaperi</i> , Mocquard	2374
Genus <i>Emmuron</i> , Jordan	2375
<i>bristoli</i> , Jordan	2375
Genus <i>Atopoclinus</i> , Vaillant	2376
<i>ringens</i> , Vaillant	2376
Genus <i>Runnula</i> , Jordan & Bollman	2377
<i>azalea</i> , Jordan & Bollman	2377
Genus <i>Blennius</i> (Artedi) Linnaeus	2377
Subgenus <i>Lipophrys</i> , Gill.....	2378
<i>carolinus</i> (Cuvier & Valenciennes)	2378
<i>fucorum</i> , Cuvier & Valenciennes	2379
<i>stearnsi</i> , Jordan & Gilbert	2379
<i>favorus</i> , Goode & Bean	2380
<i>pllicornis</i> , Cuvier & Valenciennes	2380
<i>marmoratus</i> , Poey	2381
<i>truncatus</i> (Poey).....	2381
<i>vinctus</i> , Poey	2382
<i>cristatus</i> , Linnaeus	2382
Genus <i>Scartella</i> , Jordan	2384
<i>microstoma</i> (Poey)	2384
Genus <i>Hyleurochilus</i> , Gill	2385
<i>geminatus</i> (Wood)	2385
Genus <i>Hypsoblennius</i> , Gill	2386
Subgenus <i>Hypsoblennius</i>	2386
<i>gilberti</i> (Jordan)	2386
<i>gentilis</i> (Girard)	2387
<i>striatus</i> (Steindachner)	2388
<i>ionthas</i> (Jordan & Gilbert)	2388
<i>hentz</i> (Le Sueur).....	2390
Subgenus <i>Blenniolus</i> , Jordan & Evermann.....	2390
<i>brevipinnis</i> (Günther).....	2390
Genus <i>Chasmodes</i> , Cuvier & Valenciennes	2391
<i>jenkinsi</i> (Jordan & Evermann)	2391
<i>quadrifasciatus</i> (Wood)	2392









IMAGE EVALUATION TEST TARGET (MT-3)



6"



Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

EE
128
132
125
22
20
18

EE
11
10
EE

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER XENOPTERYGINI—Continued.	
<i>Family Blennidae</i> —Continued.	
Genus <i>Chasmodes</i> , Cuvier & Valenciennes—Continued.	
<i>saburrae</i> , Jordan & Gilbert	2392
<i>novemlineatus</i> (Wood)	2393
<i>bosquianus</i> (Lacépède)	2394
Genus <i>Homesthes</i> , Gilbert	2394
<i>caulopus</i> , Gilbert	2394
Genus <i>Scartichthys</i> , Jordan & Evermann	2395
<i>rhabropunctatus</i> (Cuvier & Valenciennes)	2396
Genus <i>Rupiacartes</i> , Swainson	2396
<i>atlanticus</i> (Cuvier & Valenciennes)	2397
Genus <i>Entomacrodus</i> , Gill	2397
<i>chiostictus</i> (Jordan & Gilbert)	2398
<i>margaritaceus</i> (Poey)	2398
<i>decoratus</i> , Poey	2399
<i>nigricans</i> , Gill	2399
Genus <i>Salarichthys</i> , Guichenot	2400
<i>textilis</i> (Quoy & Gaimard)	2400
Genus <i>Coralliozetus</i> , Evermann & Marsh	3194
<i>cardone</i> , Evermann & Marsh	3194
Genus <i>Ophiolemnus</i> , Gill	2400
<i>webbi</i> (Valenciennes)	2401
<i>steindachneri</i> , Jordan & Evermann	2401
Genus <i>Emblemaria</i> , Jordan & Gilbert	2401
<i>atlantica</i> , Jordan & Evermann	2402
<i>nivipes</i> , Jordan & Gilbert	2402
<i>oculocirrhis</i> , Jordan	2403
Genus <i>Chenopsis</i> , Gill	2403
<i>ocellatus</i> , Poey	2403
Genus <i>Lucioblemmus</i> , Gilbert	2404
<i>alepidotus</i> , Gilbert	2404
Genus <i>Pholidichthys</i> , Bleeker	2405
<i>anguilliformis</i> , Lockington	2405
Genus <i>Pseudoblemmus</i> , Jenkins & Evermann	2406
<i>hypacanthus</i> , Jenkins & Evermann	2406
Genus <i>Stathmonotus</i> , Bean	2407
<i>hemphillii</i> , Bean	2407
Genus <i>Bryostemma</i> , Jordan & Starks	2408
<i>polyactocephalum</i> (Pallas)	2408
<i>nugator</i> , Jordan & Williams	2410
Genus <i>Apodichthys</i> , Girard	2411
<i>davidus</i> , Girard	2411
<i>univittatus</i> , Lockington	2412
Genus <i>Xeropis</i> , Jordan & Gilbert	2413
<i>fucorum</i> (Jordan & Gilbert)	2413
Genus <i>Ulviola</i> , Gilbert	2413
<i>sancte-rose</i> , Gilbert	2413
Genus <i>Pho'tis</i> (Gronow) Scopoli	2414
Subgenus <i>Jrocentrus</i> , Kner	2415
<i>pictus</i> (Kner)	2415
Subgenus <i>Rhodomenichthys</i> , Jordan & Evermann	2416
<i>dolichogaster</i> (Pallas)	2416
Subgenus <i>Pholis</i>	2417
<i>fasciatus</i> (Bloch & Schneider)	2417

CLASS I
SUBCL
ORDE
SUE
F

Fam

C

G

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER XENOPTERYGII—Continued.	
<i>Family Blenniidae</i> —Continued.	
Genus <i>Pholis</i> (Gronow) Scopoli—Continued.	
Subgenus <i>Pholis</i> —Continued.	
<i>gunnellus</i> (Linnaeus).....	2419
<i>ornatus</i> (Girard)	2419
Genus <i>Gunnellops</i> , Bleeker.....	2420
<i>roseus</i> (Pallas)	2420
Genus <i>Asternopteryx</i> , Rüppell.....	2420
<i>gunnelliformis</i> , Rüppell	2420
Genus <i>Anoplarchus</i> , Gill.....	2421
<i>atropurpureus</i> (Kittlitz)	2422; 2869
Genus <i>Alectrias</i> , Jordan & Evermann.....	2869
<i>alectrolophus</i> (Pallas)	2421; 2869
Genus <i>Xiphistes</i> , Jordan & Starks.....	2423
<i>ulvae</i> , Jordan & Starks	2423
<i>chirurus</i> (Jordan & Gilbert)	2424
Genus <i>Xiphidion</i> , Girard.....	2424
<i>mucosum</i> , Girard	2425
<i>rupicolle</i> (Jordan & Gilbert)	2426
Genus <i>Cebidichthys</i> , Ayres.....	2426
<i>violaceus</i> (Ayres)	2427
Genus <i>Plagiotogrammus</i> , Bean.....	2427
<i>hopkinsi</i> , Bean	2428
Genus <i>Opisthocentrus</i> , Kner.....	2428
<i>ocellatus</i> (Tilesius)	2429
Genus <i>Pholidapus</i> , Bean & Bent.....	2430
<i>dybowskii</i> (Steindachner)	2430
Genus <i>Plectobranchus</i> , Gilbert	2431
<i>evides</i> , Gilbert	2432
Genus <i>Leptoclinus</i> , Gill.....	2432
<i>maculatus</i> (Fries)	2433
Genus <i>Poroclinus</i> , Bean	2433
<i>rothrocki</i> , Bean	2434
Genus <i>Lumpenus</i> , Reinhardt	2435
Subgenus <i>Anisarchus</i> , Gill.....	2435
<i>medius</i> (Reinhardt)	2435
Subgenus <i>Lumpenus</i>	2436
<i>anguillaris</i> (Pallas)	2436
<i>mackayi</i> (Gilbert)	2436
<i>fabricii</i> (Cuvier & Valenciennes)	2437
<i>lampetraformis</i> (Walbaum)	2438
Genus <i>Stichæns</i> , Reinhardt.....	2439
<i>punctatus</i> (Fabricius)	2439
Genus <i>Ulvaria</i> , Jordan & Evermann	2440
<i>subbifurcata</i> (Storer)	2440
Genus <i>Eumesogrammus</i> , Gill	2441
<i>præcilius</i> (Krüyer)	2441
<i>Family Cryptacanthodidae</i>	2442
Genus <i>Delolepis</i> , Bean	2442
<i>virgatus</i> , Bean	2442
Genus <i>Cryptacanthodes</i> , Storer	2443
<i>maculatus</i> , Storer	2443
Genus <i>Lyconectes</i> , Gilbert	2444
<i>aleutensis</i> , Gilbert	2444

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTO? II—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER XENOPTERYGII—Continued.	
<i>Family Anarhichadidae</i>	2445
Genus <i>Anarhichas</i> (Arvedii) Linneus	2445
latifrons, Steenstrup & Hallgrímsson	2446
minor, Olafsen	2446
lupus, Linnaeus	2446
lepturus, Bean	2447
orientalis, Pallas	2447
Genus <i>Anarrhichthys</i> , Ayres	2447
ocellatus, Ayres	2448
<i>Family Cerdalidae</i>	2448
Genus <i>Cerdale</i> , Jordan & Gilbert	2448
ionthas, Jordan & Gilbert	2449
Genus <i>Microdesmus</i> , Günther	2450
dipus, Günther	2450
retropinnis, Jordan & Gilbert	2450
<i>Family Ptileichthyidae</i>	2451
Genus <i>Ptileichthys</i> , Bean	2452
goodei, Bean	2452
Group Ophidioidea	2453
<i>Family Scyhalinidae</i>	2453
Genus <i>Scyhalina</i> , Jordan & Gilbert	2454
cerdale, Jordan & Gilbert	2454
<i>Family Zoarcidae</i>	2455
Genus <i>Zoarcæs</i> , Gill	2456
Subgenus <i>Tacrozoarcæs</i> , Gill	2457
anguillaris (Peck)	2457
Genus <i>Embryx</i> , Jordan & Evermann	2458
crassilabris (Gilbert)	2458
crotalinus (Gilbert)	2458
Genus <i>Lycodopsis</i> , Collett	2460
pacificus (Collett)	2460
Genus <i>Aprodon</i> , Gilbert	2460
cortezianus, Gilbert	2461
Genus <i>Lycodes</i> , Reinhardt	2461
Subgenus <i>Lycodes</i>	2463
esmarkii, Collett	2463
vahlii, Reinhardt	2463
concolor, Gill & Townsend	2463
zoarchus, Goode & Bean	2464
reticulatus, Reinhardt	2465
perspicillum, Krüyer	2465
frigidus, Collett	2465
terre-novæ, Collett	2466
digitatus, Gill & Townsend	2466
palearis, Gilbert	2466
brevipes, Bean	2467
Subgenus <i>Lycius</i> , Jordan & Evermann	2468
nebulosus, Krüyer	2468
semihudius, Reinhardt	2468
Genus <i>Lycodalepis</i> , Bleeker	2468
polaris (Sabine)	2468
mucosus (Richardson)	2470
Genus <i>Lyconchelys</i> , Gill	2470; 2860
verrillii (Goode & Bean)	2470

LASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER XENOPTERYGI—Continued.	
<i>Family Zoarcidae</i> —Continued.	
Genus <i>Lyconchelys</i> , Gill—Continued.	
<i>paxillus</i> (Goode & Bean)	2471
<i>muraena</i> (Collett).....	3195
<i>porifer</i> (Gilbert).....	2471
Genus <i>Furcimanus</i> , Jordan & Evermann.....	2472
<i>diapterus</i> (Gilbert).....	2472
Genus <i>Lycodonus</i> , Goode & Bean	2473
<i>mirabilis</i> , Goode & Bean.....	2474
Genus <i>Lyconema</i> , Gilbert	2474
<i>barbatum</i> , Gilbert	2474
Genus <i>Bothrocara</i> , Bean.....	2475
<i>pusilla</i> (Bean)	2476
<i>mollis</i> , Bean	2476
Genus <i>Gymnelis</i> , Reinhardt.....	2477
<i>viridis</i> (Fabricius)	2477
<i>stigma</i> (Lay & Bennett).	2477
Genus <i>Lycocara</i> , Gill	2478
<i>parrii</i> (Ross)	2478
Genus <i>Melanostigma</i> , Günther.....	2478
<i>gelatinosum</i> , Günther	2479
<i>pammelas</i> , Gilbert	2479
<i>Family Derepodichthyidae</i>	2480
Genus <i>Derepodichthys</i> , Gilbert	2480
<i>alepidotus</i> , Gilbert	2480
<i>Family Ophidiidae</i>	2481
Genus <i>Lepophidium</i> , Gill.....	2482
<i>marmoratum</i> (Goode & Bean)	2482
<i>emmelas</i> (Gilbert)	2483
<i>stigmatistium</i> (Gilbert)	2483
<i>profundorum</i> (Gill)	2484
<i>cervinum</i> (Goode & Bean)	2484
<i>prorates</i> (Jordan & Bollman)	2485
<i>brevibarbe</i> (Cuvier)	2485
<i>pardale</i> (Gilbert)	2486
<i>microlepis</i> (Gilbert)	2486
Genus <i>Ophidion</i> (Artefi) Linnaeus	2487
<i>beani</i> , Jordan & Gilbert	2487
<i>holbrookii</i> (Putnam)	2487
<i>grelli</i> , Poey	2488
Genus <i>Chiaria</i> , Jordan & Evermann	2488
<i>taylori</i> (Girard)	2489
Genus <i>Rissoala</i> , Jordan & Evermann	2489
<i>marginata</i> (De Kay)	2489
Genus <i>Otophidium</i> , Gill	2490
<i>omostigmum</i> (Jordan & Gilbert)	2490
<i>indefatigabile</i> , Jordan & Bollman	2490
<i>galeoides</i> (Gilbert)	2491
<i>Family Lycodopodidae</i>	2491
Genus <i>Lycodapus</i> , Gilbert	2492
<i>dermatinus</i> , Gilbert	2492
<i>fierasfer</i> , Gilbert	2493
<i>parviceps</i> , Gilbert	2493
<i>extensus</i> (Gilbert)	2494

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER XENOPTERYGII—Continued.	
<i>Family Fierasferidae</i>	2494
Genus Fierasfer, Cuvier	2495
affinis (Günther)	2495
arenicola, Jordan & Gilbert	2496
bermudensis (Jones)	2497
<i>Family Brotulidae</i>	2498
Genus Brotula, Cuvier	2500
barbata (Bloch & Schneider)	2500
Genus Stygicola, Gill	2500
dentatus (Poey)	2500
Genus Lucifuga, Poey	2501
subterraneus, Poey	2501
Genus Brosmophycis, Gill	2502
marginatus (Ayres)	2502
Genus Ogilbia, Jordan & Evermann	2502
ventralis (Gill)	2503
cayorum, Evermann & Kendall	2503
Genus Bythites, Reinhardt	2504
fuscus, Reinhardt	2504
Genus Catatyx Günther	2504
rubirostris, Gilbert	2505
Genus Dicromista, Goode & Bean	2506
agassizii, Goode & Bean	2506
Genus Bassozetus, Gill	2507
normalis, Gill	2507
compressus (Günther)	2508
catena, Goode & Bean	2509
tentia (Günther)	2510
Genus Mcleia, Goode & Bean	2510
promelas (Gilbert)	2511
Genus Neobrythites, Goode & Bean	2512
gilli, Goode & Bean	2512
marginatus, Goode & Bean	2513
Genus Benthocometes, Goode & Bean	2514
robustus, Goode & Bean	2514
Genus Bassogigas, Gill	2515
gilli, Goode & Bean	2515
stelliferoides (Gilbert)	2516
Genus Barathrodemus, Goode & Bean	2517
manatinus, Goode & Bean	2517
Genus Nematus, Günther	2518
pectoralis, Goode & Bean	2518
Genus Porogadus, Goode & Bean	2519
miles, Goode & Bean	2520
Genus Penopus, Goode & Bean	2520
maedonaldi, Goode & Bean	2521
Genus Dicroleone, Goode & Bean	2522
intronigra, Goode & Bean	2522
Genus Mixonix, Günther	2523
laticeps (Günther)	2523
Genus Barathronus, Goode & Bean	2524
bicolor, Goode & Bean	2524
Genus Aphonus, Günther	2525
mollis, Goode & Bean	2525

CLASS PISCES—Continued.	Page.
SUB-LASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER XENOPTERYGINI—Continued.	
<i>Family Bregmacerotidae</i>	2525
Genus <i>Bregmaceros</i> , Thompson	2526
<i>maclellandii</i> , Thompson	2526
<i>atlanticus</i> , Goode & Bean	2527
<i>NB SUBORDER ANACANTHINI</i>	2528
<i>Family Merluccidae</i>	2529
Genus <i>Merluccius</i> , Rafinesque	2529
<i>merluccius</i> (Linnaeus)	2530
<i>bilinearis</i> (Mitchill)	2530
<i>productus</i> (Ayres)	2531
<i>Family Gadidae</i>	2531
Genus <i>Boreogadus</i> , Günther	2533
<i>saida</i> (Lepechin)	2533
Genus <i>Pollachius</i> , Nilsson	2534
<i>virens</i> (Linnaeus)	2534
Genus <i>Theragra</i> , Lucas	2535
<i>chaleogramma</i> (Pallas)	2535
<i>fucensis</i> (Jordan & Gilbert)	2536
Genus <i>Eleginops</i> , Fischer	2537
<i>navaga</i> (Kolreuter)	2537
Genus <i>Microgadus</i> , Gill	2538
<i>proximus</i> (Girard)	2539
<i>tomcod</i> (Walbaum)	2540
Genus <i>Gadus</i> (Artedi) Linnaeus	2540
<i>callarias</i> , Linnaeus	2541
<i>macrocephalus</i> , Tileanus	2541
<i>ogac</i> , Richardson	2542
Genus <i>Melanogrammus</i> , Gill	2542
<i>ieglefinus</i> , Linnaeus	2542
Genus <i>Lepidion</i> , Swainson	2543
<i>vereendum</i> , Jordan & Cramer	2543
Genus <i>Antimora</i> , Günther	2544
<i>viola</i> (Goode & Bean)	2544
<i>microlepis</i> , Bean	2545
Genus <i>Uraleptus</i> , Costa	2545
<i>maraldi</i> (Risso)	2545
Genus <i>Lotella</i> , Kaup	2546
<i>maxilaris</i> , Bean	2546
Genus <i>Physiculus</i> , Kaup	2547
<i>fulvus</i> , Bean	2547
<i>nematopus</i> , Gilbert	2548
<i>kaupi</i> , Poey	2548
<i>rastrelliger</i> , Gilbert	2549
Genus <i>Lota</i> (Cuvier) Oken	2550
<i>maenosa</i> (Le Sueur)	2550
Genus <i>Molva</i> , Fleming	2551
<i>molva</i> (Linnaeus)	2551
Genus <i>Urophycis</i> , Gill	2552
Subgenus <i>Urophycis</i>	2553
<i>regius</i> (Walbaum)	2553
<i>cirratus</i> (Goode & Bean)	2553
<i>floridanus</i> (Bean & Dresel)	2554
<i>earll</i> (Bean)	2554
Subgenus <i>Emphyicus</i> , Jordan & Evermann	2554
<i>tenuis</i> (Mitchill)	2555

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER ANACANTHINI—Continued.	
<i>Family Gadidae—Continued.</i>	
Genus <i>Urophycis</i> , Cuvier—Continued.	
Subgenus <i>Emphyceus</i> , Jordan & Evermann—Continued.	
<i>chuss</i> (Walbaum)	2555
<i>chesteri</i> (Goode & Bean)	2556
Genus <i>Lamponema</i> , Günther	2556
<i>barbatulum</i> , Goode & Bean	2556
<i>melanurrum</i> , Goode & Bean	2557
Genus <i>Gadropsar</i> s, Rafinesque	2557
<i>ensis</i> (Reinhardt)	2558
<i>argentatus</i> (Reinhardt)	2559
<i>septentrionalis</i> (Collett)	2559
Genus <i>Enchelyopus</i> , Bloch & Schneider	2560
<i>cimbricus</i> (Linnaeus)	2560
Genus <i>Brosme</i> (Cuvier) Oken	2561
<i>brosme</i> (Müller)	2561
<i>Family Macrouridae</i>	2561
Genus <i>Bathygadus</i> , Günther	2563
<i>arcuatus</i> , Goode & Bean	2564
<i>favosus</i> , Goode & Bean	2565
<i>maerops</i> , Goode & Bean	2566
<i>longifilis</i> , Goode & Bean	2566
Genus <i>Steindachnerina</i> , Goode & Bean	2567
<i>argentea</i> , Goode & Beau	2568
Genus <i>Trachyrhynchus</i> , Giorna	2568
<i>helolepis</i> , Gilbert	2569
Genus <i>Malacocephalus</i> , Günther	2569
<i>occidentalis</i> , Goode & Bean	2570
Genus <i>Moseleya</i> , Goode & Bean	2570
<i>cyclolepis</i> (Gilbert)	2570
Genus <i>Nematourus</i> , Günther	2571
<i>goodei</i> (Günther)	2571
<i>suborbitalis</i> (Gill & Townsend)	2572
Genus <i>Albatrossia</i> , Jordan & Evermann	2573
<i>peectoralis</i> (Gilbert)	2573
Genus <i>Bogoslovi</i> s, Jordan & Evermann	2574
<i>clarkii</i> , Jordan & Gilbert	2575
<i>firmisquamis</i> (Gill & Townsend)	2575
Genus <i>Challinura</i> , Goode & Bean	2576
<i>serrula</i> , Bean	2576
<i>filifera</i> , Gilbert	2577
<i>simula</i> , Goode & Bean	2578
Genus <i>Coryphaenoides</i> , Gunner	2578
<i>rupestris</i> , Gunner	2579
<i>carapinus</i> , Goode & Bean	2579
Genus <i>Hymenocephalus</i> , Giglioli	2580
<i>cavernosus</i> (Goode & Bean)	2580
Genus <i>Macrourus</i> , Bloch	2581
<i>berglax</i> , Laëpède	2582
<i>holotrachys</i> , Günther	2582
<i>bairdii</i> , Goode & Bean	2583
<i>lepturus</i> , Gill & Townsend	2584
<i>acrolepis</i> , Bean	2585
<i>stelgidolepis</i> , Gilbert	2585

	Page.
CLASS PISCES—Continued.	
SUBCLASS TELEOSTOMI—Continued,	
ORDER PLECTOGNATHI—Continued.	
SUBORDER ANACANTHINI—Continued.	
Family Macrouridae—Continued.	
Genus Macrourus, Bloch	—Continued.
cinerens, Gilbert	2586
ingolfi, Lütken	3105
Genus Celorhynchus, Giorna	2587
ocea (Goode & Bean)	2588
carminatus (Goode)	2588
caribbeus (Goode & Bean)	2589
scaphopsis (Gilbert)	2590
Genus Trachonurus, Günther	2591
sulcatus (Goode & Bean)	2591
Genus Lionurus, Günther	2592
flicanda (Günther)	2592
liolepis, Gilbert	2593
SUBORDER TENIOSOMI	2594
Family Regalecidae	2595
Genus Regalecus, Brünnich	2595
glesne (Ascanius)	2596
Family Trachypteridae	2597
Genus Trachypterus, Gouan	2599
rex-salmonorum, Jordan & Gilbert	2599
trachyurus, Poey	2600
Family Stylephoridae	2601
Genus Stylephorus, Shaw	2601
chordatus, Shaw	2601
SUBORDER HETEROSOMATA	2602
Family Pleuronectidae	2602
Genus Atheresthes, Jordan & Gilbert	2609
stomias (Jordan & Gilbert)	2609
Genus Reinhardtius, Gill	2610
hippoglossoides (Walbaum)	2611
Genus Hippoglossus, Cuvier	2611
hippoglossus (Linnaeus)	2611
Genus Lyopsetta, Jordan & Goss	2612
exilis (Jordan & Gilbert)	2612
Genus Eopsetta, Jordan & Goss	2613
jordani (Lockington)	2613
Genus Hippoglossoides, Gottsche	2614
platessoides (Fabricius)	2614
elassodon, Jordan & Gilbert	2615
robustus, Gill & Townsend	2616
hamiltoni, Jordan & Gilbert	2616
Genus Psettichthys, Girard	2617
melanostictus, Girard	2618
Genus Verasper, Jordan & Gilbert	2618
moseri, Jordan & Gilbert	2619
Genus Hippoglossina, Steindachner	2620
stomata, Eigenmann & Eigenmann	2620
macrops, Steindachner	2621
bollmani, Gilbert	2621
subanensis, Boulenger	3196
Genus Lioglossina, Gilbert	2622
tetrophthalmia, Gilbert	2622
Bull. No. 47, pt. 4—VII	

CLASS PISCES—Continued.	Page.
SUBCLASS TELEOSTOMI—Continued.	
ORDER PLECTOGNATHI—Continued.	
SUBORDER HETEROSOMATA—Continued.	
Family <i>Pleuronectidae</i> —Continued.	
Genus <i>Xystreurus</i> , Jordan & Gilbert.....	2623
<i>holepis</i> , Jordan & Gilbert	2623
Genus <i>Paralichthys</i> , Girard.....	2624
<i>californicus</i> (Ayres).....	2625
<i>magdalene</i> , Abbott.....	2871
<i>estuarinus</i> , Gilbert & Sefton	2626
<i>brasiliensis</i> (Ranzani).....	2626
<i>sinaloae</i> , Jordan & Abbott.....	2627; 2872
<i>woolmani</i> , Jordan & Williams.....	2628
<i>dentatus</i> (Linnaeus).....	2629
<i>lethostigma</i> , Jordan & Gilbert.....	2630
<i>squamiferus</i> , Jordan & Gilbert.....	2631
<i>albifrons</i> , Jordan & Gilbert.....	2631
<i>oblongus</i> (Mitchill).....	2632
Genus <i>Ramularia</i> , Jordan & Evermann.....	2633
<i>dendriticis</i> (Gilbert).....	2633
Genus <i>Aneylopsetta</i> , Gill.....	2634
<i>quadrocellata</i> , Gill	2634
Genus <i>Notosema</i> , Goode & Bean	2635
<i>dilectum</i> , Goode & Bean.....	2635
Genus <i>Gastropsetta</i> , B. A. Bean.....	2636
<i>frontalis</i> , B. A. Bean.....	2636
Genus <i>Pleuronichthys</i> , Girard	2637
<i>decurvens</i> , Jordan & Gilbert	2637
<i>verticalis</i> , Jordan & Gilbert.....	2638
<i>cenosus</i> , Girard.....	2638
Genus <i>Hypsopsetta</i> , Gill.....	2639
<i>guttulata</i> (Girard).....	2639
Genus <i>Parophrys</i> , Girard	2640
<i>vetulus</i> , Girard.....	2640
Genus <i>Inopsetta</i> , Jordan & Goss	2641
<i>ischyra</i> (Jordan & Gilbert).....	2641
Genus <i>Isopsetta</i> , Lockington	2642
<i>isolepis</i> (Lockington).....	2642
Genus <i>Lepidopsetta</i> , Gill	2642
<i>bilineata</i> (Ayres).....	2643
Genus <i>Limanda</i> , Gottsche	2644
<i>ferruginea</i> (Storer).....	2644
<i>aspera</i> (Pallas)	2645
<i>proboscidea</i> , Gilbert.....	2645
<i>beanii</i> , Goode.....	2646
Genus <i>Pseudopleuronectes</i> , Bleeker	2646
<i>americanus</i> (Walbaum).....	2647
<i>pinnifasciatus</i> (Kner)	2647
Genus <i>Pleuronectes</i> (Artedi) Linnaeus	2648
<i>quadriruberulatus</i> , Pallas.....	2648
Genus <i>Liopsetta</i> , Gill	2649
<i>glacialis</i> (Pallas)	2649
<i>putnami</i> (Gill)	2650
<i>obscura</i> (Herzenstein)	2651
Genus <i>Platichthys</i> , Girard	2651
<i>stellatus</i> (Pallas).....	2652

CLASS
SUBC
ORD
SU

	Page.	
CLASS PISCES—Continued.		
SUBCLASS TELEOSTOMI—Continued.		
ORDER PLECTOGNATHI—Continued.		
SUPERORDER HETEROCHTHONIA—Continued.		
Family Pleuronectidae—Continued.		
623	Genus <i>Microstomus</i> , Gottsche	2653
623	<i>kitt</i> (Walbaum)	2654
624	<i>pacificus</i> (Lockington)	2655
625	Genus <i>Embassichthys</i> , Jordan & Evermann	2655
871	<i>bathybium</i> (Gilbert)	2655
626	Genus <i>Glyptocephalus</i> , Gottsche	2656
626	<i>cynoglossus</i> (Linnaeus)	2657
852	<i>zachirus</i> , Lockington	2658
628	Genus <i>Lophopsetta</i> , Gill	2659
629	<i>maculata</i> (Mitchill)	2660
630	Genus <i>Platophrys</i> , Swainson	2660
631	<i>spinosus</i> (Poey)	2662
631	<i>constellatus</i> , Jordan	2663
632	<i>ocellatus</i> (Agassiz)	2663
633	<i>maculifer</i> (Poey)	2664
633	<i>ellipticus</i> (Poey)	2665
634	<i>lunatus</i> (Linnaeus)	2665
634	<i>leopardinus</i> (Günther)	2666
635	Genus <i>Perissias</i> , Jordan & Evermann	2667
635	<i>taenopterus</i> (Gilbert)	2667
636	Genus <i>Engyophrys</i> , Jordan & Bollman	2668
636	<i>saneti-laurentii</i> , Jordan & Bollman	2668
637	Genus <i>Trichopsetta</i> , Gill	2669
637	<i>ventralis</i> (Goode & Bean)	2669
638	Genus <i>Syacium</i> , Ranzani	2670
638	<i>papillosum</i> (Linnaeus)	2671
639	<i>micrurum</i> , Ranzani	2672
639	<i>latifrons</i> (Jordan & Gilbert)	2673
640	<i>ovale</i> (Günther)	2674
640	Genus <i>Cyclopsetta</i> , Gill	2675
641	<i>querma</i> (Jordan & Bollman)	2675
642	<i>chittendeni</i> , B. A. Bean	2676
642	<i>fimbriata</i> (Goode & Bean)	2676
642	Genus <i>Azevia</i> , Jordan	2677
642	<i>panamensis</i> (Steindachner)	2677
643	Genus <i>Citharichthys</i> , Bleeker	2678
644	Subgenus <i>Orthopsetta</i> , Gill	2679
644	<i>sordidus</i> (Girard)	2679
645	<i>fragilis</i> , Gilbert	2680
645	<i>xanthostigmus</i> , Gilbert	2680
646	<i>stigmaeus</i> , Jordan & Gilbert	2681
646	Subgenus <i>Citharichthys</i>	2682
647	<i>dinoceros</i> , Goode & Bean	2682
647	<i>platophrys</i> , Gilbert	2683
648	<i>arctifrons</i> , Goode	2685
648	<i>unicornis</i> , Goode	2683
649	<i>uhleri</i> , Jordan	2684
649	<i>macrops</i> , Dressel	2684
650	<i>epipopterus</i> , Günther	2685
651	<i>gilberti</i> , Jenkins & Evermann	2686
651	Genus <i>Etropus</i> , Jordan & Gilbert	2687
652	<i>microstomus</i> (Gill)	2687

CLASS PISCES—Continued.	Page.	
SUBCLASS TELEOSTOMI—Continued.		
ORDER PLECTOGNATHI—Continued.		
SUBORDER HETEROSOMATA—Continued.		
<i>Family Pleuronectidae</i> —Continued.		
Genus <i>Etronus</i> , Jordan & Gilbert—Continued.		
<i>rimosus</i> , Goode & Bean	2688	
<i>crossotus</i> , Jordan & Gilbert	2689	
Genus <i>Monolepis</i> , Goode.....	2690	
<i>sessilecauda</i> , Goode	2691	
<i>atrimana</i> , Goode & Bean	2692	
<i>Family Soleidae</i>	2692	
Genus <i>Achirus</i> , Lacépède	2693	
Subgenus <i>Balostoma</i> , Bean	2695	
<i>uehirus</i> (<i>Linnæus</i>)	2695	
<i>inscriptus</i> , Gosse	2696	
<i>kunzingeri</i> (Steindachner)	2697	
<i>lineatus</i> (<i>Linnæus</i>)	2697	
<i>mazatlanus</i> (Steindachner)	2698	
<i>fonsecensis</i> (Günther)	2699	
<i>fischeri</i> (Steindachner)	2699	
<i>scutum</i> (Günther)	2700	
Subgenus <i>Achirus</i>	2700	
<i>rhombifer</i> (Günther)	2700	
<i>fasciatus</i> , Lacépède	2700	
<i>panamensis</i> (Steindachner)	2702	
Genus <i>Aplonichthys</i> , Kaup	2702	
<i>unicolor</i> (Günther)	2702	
Genus <i>Gymnachirus</i> , Kaup	2703	
<i>laevigatus</i> , Günther	2703	
Genus <i>Syphurus</i> , Rafinesque	2704	
Subgenus <i>Syphurus</i>	2705	
<i>piger</i> (Goode & Bean)	2705	
<i>marginatus</i> (Goode & Bean)	2706	
<i>atramentatus</i> , Jordan & Bollman	2706	
<i>fasciolaris</i> , Gilbert	2707	
<i>elongatus</i> (Günther)	2707	
<i>atricaudatus</i> (Jordan & Gilbert)	2707	
<i>leei</i> , Jordan & Bollman	2708	
<i>plagiusa</i> (Bloch & Schneider)	2709	
<i>plagiusa</i> (<i>Linnæus</i>)	2710	
<i>pusillus</i> (Goode & Bean)	2710	
<i>diomedeanus</i> (Goode & Bean)	2711	
<i>williamsi</i> , Jordan & Culver	2711	
Subgenus <i>Acedia</i> , Jordan	2712	
<i>nebulosus</i> (Goode & Bean)	2712	
ORDER PEDICULATI	2712	
<i>Family Lophiidae</i>		
Genus <i>Lophius</i> (Artedi) <i>Linnæus</i>	2713	
<i>piscatorius</i> , <i>Linnæus</i>	2713	
Genus <i>Lophiomus</i> , Gill	2714	
<i>setigerus</i> (Vahl)	2714	
<i>Family Antennariidae</i>		2715
Genus <i>Pterophryne</i> , Gill	2715	
<i>histrio</i> (<i>Linnæus</i>)	2716	
<i>glbba</i> (Mitchill)	2717	
Genus <i>Antennarius</i> , Lacépède	2717	
<i>inops</i> , Poey	2718	

CLASS PISCES—Continued.

Page.

SUBCLASS TELEOSTOMI—Continued.

ORDER PEDICULATI—Continued.

Family Antennariidae—Continued.Genus *Antennarius*—Continued.

principis (Cuvier & Valenciennes)	2710
tenebrosus (Poey)	2719
reticularis, Gilbert	2719
strigatus, Gill	2720
sanguineus, Gill	2721
ocellatus (Bloch & Schneider)	2721
seaber (Cuvier)	2722
tigris, Poey	2723
nuttingii, Garman	2723
multicollatus (Cuvier & Valenciennes)	2724
radiosus, Garman	2725
Genus <i>Chimaera</i> , Lowe	2726
pictus, Lowe	2726
nuttingii, Garman	2727

Family Ceratiidae

Genus <i>Ceratias</i> , Krüyer	2727
holbolli, Krüyer	2729
Genus <i>Manculias</i> , Gill	2729
uranoscopus (Murray)	2729
shufeldti (Gill)	2730
Genus <i>Cryptopsaras</i> , Gill	2731
conesil, Gill	2731
Genus <i>Oncirodes</i> , Lütken	2732
eschrichtii, Lütken	2732
Genus <i>Himantolophus</i> , Reinhardt	2732
greenlandicus, Reinhardt	2733
Genus <i>Corynophorus</i> , Gill	2733
reinhardti (Lütken)	2733
Genus <i>Lioctetus</i> , Günther	2733
murrayi (Günther)	2733
Genus <i>Linophryne</i> , Collett	2734
lucifer, Collett	2734
Genus <i>Caulophryne</i> , Goode & Bean	2734
jordani, Goode & Bean	2735

Family Oyocephalidae

Genus <i>Oyocephalus</i> , Fischer	2735
vespertilio (Linnaeus)	2736
nasutus (Cuvier & Valenciennes)	2737
radiatus (Mitchill)	2737
Genus <i>Zalientes</i> , Jordan & Evermann	2738
elater (Jordan & Gilbert)	2738
Genus <i>Halieutichthys</i> , Poey	2739
aculeatus (Mitchill)	2739
caribbeus, Garman	2741
Genus <i>Halieutaea</i> , Cuvier & Valenciennes	2741
spongiosa, Gilbert	2741
Genus <i>Halieutella</i> , Goode & Bean	2742
lappa, Goode & Bean	2742
Genus <i>Dibranchus</i> , Peters	2743
atlanticus, Peters	2743

Since
new gene
localities
relations
been enla
the follow

Page 130
insert the fo

Head $3\frac{1}{2}$;
D, I, 7; A, 1.
width half
much smaller
premaxillary
head, 4 in in
this last a
narrow, not
occipital pro
G. rugispinus
maxillary 1 $\frac{1}{2}$
times as long
edge; dorsal
a little less than
twice as far
pectoral spine
shorter than the
deeply forked
deep; no maxilla
blackish. Length
Rio Peripa, E.

Arius labiatus, 1
Torino, Vol.
(Festa.)

ADDITIONAL ADDENDA.

Since the publication of Part III of this work a number of new genera and species of fishes have been described from localities within our limits, and our knowledge regarding the relations and distribution of previously known species has been enlarged. These additions and corrections are given in the following pages:

Page 130. After *Heramenatichthys rugispinus* (Cuvier & Valenciennes), insert the following:

195 (n). **GALEICHTHYS LABIATUS** (Boulenger).

Head $3\frac{1}{2}$; width of head $1\frac{1}{2}$ in its length; depth 5; eye 9 in head. D. I, 7; A. 17. Band of premaxillary teeth 4 times longer than wide, the width half the eye; palatine teeth conical, in 2 small oblique groups, much smaller than eye, and separated by a space equal to $\frac{1}{4}$ the length of premaxillary band; eye situated above level of mouth, in anterior half of head, 4 in interocular space; occiput and occipital prolongation granular, this last a little longer than wide, with an obtuse spine; foranelle narrow, not distinct, followed by a groove which does not reach the occipital prolongation; lips thick, the upper overhanging the lower, as in *G. rugispinus*, to which this species is closely related; barbels flattened; maxillary $1\frac{1}{2}$ in head, not reaching edge of opercle; external mandible $1\frac{1}{2}$ times as long as internal, 2 in head; gill membrane with a free, unbroken edge; dorsal spine with indistinct serrations, covered with skin, its length a little less than $\frac{1}{2}$ head; adipose dorsal as long at base as rayed dorsal, twice as far from it as from the caudal; humeral prolongation smooth; pectoral spine $2\frac{1}{2}$ in head, enveloped in skin like the dorsal spine, much shorter than the soft rays; ventral shorter, reaching origin of anal; caudal deeply forked, with pointed lobes; caudal peduncle $2\frac{1}{2}$ times longer than deep; no maxillary pore. Blackish gray above, silvery below; all the fins blackish. Length 470 mm. Known only from a single specimen from Rio Peripa, Ecuador. (Boulenger.) (*labiatus*, having large lips.)

Arius labiatus, BOULENGER, Bollettino dei Mus. di Zool. ed. Anat. Comp. della Univ. di Torino, Vol. XIII, No. 329, 6, Dec. 2, 1898, Rio Peripa, Ecuador. (Coll. Dr. Enrico Festa.)

Page 132. After *Tachysurus melanopus* (Günther), insert the following:

260 (a). TACHYSURES FESTE (Boulenger).

Head 4; depth 4; eye $6\frac{1}{2}$, above level of mouth, in anterior part of head, D. 17; A. 23. Premaxillary band of teeth 4 times longer than broad, its width $2\frac{1}{2}$ in eye, or equal to width of groups of vomer and palatine teeth, these last granular, forming 2 large, oblique groups, separated on the median line by a space equal to $\frac{1}{2}$ diameter of eye; width of head $1\frac{1}{2}$ in its length; occiput and occipital prolongation granular, the last as long as wide, and with blunt spine; fontanelle narrow, commencing in a groove between the anterior nostrils and reaching occiput, not reaching occipital prolongation; barbels flattened; maxillary as long as head, reaching beyond base of pectoral spine; external mandible a little shorter, as long as the internal; gill-membrane with a free border broken at the isthmus; anal spine striated, $1\frac{1}{2}$ in head, its anterior edge feebly toothed except at extremity, which has, like the posterior edge, stronger and more retrorse teeth; adipose dorsal twice as far from rayed dorsal as from caudal, its base $1\frac{1}{2}$ in base of rayed dorsal; pectoral spine $1\frac{1}{2}$ in head, its external edge furnished with a series of soft, small tubercles on its basal half, anterior edge armed with feeble retrorse teeth; ventral shorter, reaching origin of anal; caudal deeply forked; caudal peduncle twice as long as deep; axillary pore present, much smaller than the nostril. Color steel blue above, silvery below. This species closely resembles *Tachysurus liopnus*, Bristol. Length 210 mm. One specimen known, from Narrangal, Ecuador. (Boulenger.) (Named for the collector of the type, Dr. Enrico Festa.)

Arius festae, BOULENGER, Bollettino dei Mus. di Zool. ed. Anat. Comp. della Univ. di Torino, Vol. xii, No. 329, 5, Dec. 2, 1898, Narrangal, Ecuador.

Page 135. After *Ictalurus meridionalis* (Günther), insert:

76 (a). ISTLARIUS, Jordan & Snyder.

Istlaricus, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 118 (*balsanus*).

Body rather deep and compressed; head not widened, nor greatly depressed; eye large; lower jaw included; teeth in villiform bands on premaxillaries and dentaries; the premaxillary band convex anteriorly, with a short angular posterior extension on each side, no division of the band at symphysis; dentary band broad anteriorly, growing narrow and pointed posteriorly, with a distinct median division; no teeth on vomer or palatines; villiform teeth on upper and lower pharyngeals; gill-rakers on first arch 17, long and slender; branchiostegals 8; air bladder very large, extending almost to posterior end of body cavity, divided by a transverse constriction into two parts of nearly equal length, the anterior heart-shaped, posterior part oval; supraoccipital bone widely separated from interspinal; humeral process short, almost hidden by the skin; lateral line extending from below insertion of dorsal to caudal; skin covered with minute, hairlike villi; skin of head completely concealing bones of skull; barbels 8. Spines with

distal part of groove.

Istlaricus dentition more close than genu. (Named

Head 4; $2\frac{1}{2}$; distal dorsal 3; $1\frac{1}{2}$; ventral pressed, does not greatl edge of op convex; w jecting a d ones the m diameter of the edges o of gill open villiform ba convex ante no apparent anteriorly, no teeth on bands; low on first arc tubular, the barbel to the head comple in contact v diameter of from a perpe with minute ced by a s others gradu of the poste ray longest, lower lobe in two-thirds o basal part gr reaching orig few smal inferior barbules wit

distal parts soft, not branched, continuous with the hard parts; basal part of pectoral spine grooved posteriorly, weakly serrate above the groove.

Istlarinus has some of the characters of the genus *Leptops*, notably the dentition of the upper jaw and the weakness of the fin-spines; but it more closely resembles *Ictalurus*, and its relationship is probably with that genus.

(Named for the Rio Ixtla, the type locality of the type species.)

208 (a). **ISTLARIUS BALSANUS**, Jordan & Snyder.

Head 4; depth $4\frac{1}{2}$; depth of caudal peduncle $2\frac{1}{2}$ in head; eye $5\frac{1}{2}$; snout $2\frac{3}{4}$; distance between eyes $2\frac{1}{2}$; height of dorsal $1\frac{1}{2}$; length of base of dorsal 3; height of anal $1\frac{1}{2}$; length of base of anal 1; length of pectoral $1\frac{1}{2}$; ventrals $1\frac{1}{2}$; caudal 1; D. I, 6; A. 24. Body deep and somewhat compressed, deepest above ventrals, widest between pectorals; head narrow, not greatly depressed; eye large, nearer tip of snout than to posterior edge of opercle a distance equal to diameter of eye; interorbital space convex; width of mouth $2\frac{1}{2}$ in head, lower jaw included; upper jaw projecting a distance equal to diameter of pupil; barbels 8; of the inferior ones the median pair shorter; distance between their bases equal to the diameter of pupil; the outer pair when extended directly backward reach the edges of gill covers; maxillary barbels longest, reaching upper angle of gill opening; nostril barbels reaching middle of pupil. Teeth in broad villiform bands on premaxillaries and dentaries, the band on upper jaw convex anteriorly, with a short, angular posterior extension on each side, no apparent division of the band at symphysis; band on lower jaw broad anteriorly, narrow and pointed posteriorly, a distinct median division; no teeth on vomer or palatines; upper pharyngeal teeth in oval villiform bands; lower pharyngeal teeth in 2 narrow oblong bands; gill-rakers on first arch 17, long and slender; branchiostegals 8; anterior nostril tubular, the posterior with a raised rim extending on each side from the barbel to the posterior edge of the opening, where it is divided; skin of head completely concealing the bones of skull; supraoccipital bone not in contact with interspinal; humeral process about as long as vertical diameter of eye, almost hidden by the thick skin; lateral line extending from a perpendicular through insertion of dorsal to caudal; skin covered with minute, hair-like villi; dorsal spine with its distal third soft, preceded by a small, angular, immature spine; first branched ray longest, the others gradually shorter; adipose fin large, above middle of anal; length of the posterior free edge 3 times diameter of pupil; third or fourth anal ray longest, edge of fin rounded posteriorly; caudal deeply forked, the lower lobe rounded, the upper rather pointed; pectoral rays 1, 11, distal two-thirds of spine soft, not branched, continuous with the hard part; basal part grooved posteriorly, weakly serrate above the groove; ventrals reaching origin of anal. Color bluish slate above, light silvery below; a few small dark spots on the head and body; fins with dusky coloring; inferior barbels light; maxillary barbets with upper half dark; nasal barbules with light edges. Some specimens have many well-defined color

spots, while others have few or none. The young have no spots. Our specimens vary in length from 10 to 60 centimeters.

Istlaris balsanus has a large and rather complex air bladder. It lies close to the spinal column and extends almost to the posterior end of the body cavity. It is divided by a deep transverse constriction into two halves. The anterior part is heart-shaped, constricted dorso-ventrally. It is separated by a T-shaped partition into 3 chambers. Of these the anterior, transverse chamber is partly divided on the median line by a fold of the dorsal wall of the bladder over the vertebral column. The transverse chamber is connected on each side with the 2 posterior lateral chambers by large openings in the ends of the transverse wall. The posterior half of the bladder is without partitions. It is connected with the left lateral chamber only by a small opening. (Jordan & Snyder.)

Known only from Rio Ixtla at Puente de Ixtla, Morelos, Mexico. (Type, No. 6149, L. S. Jr. Univ. Mus. Coll. Jordan & Snyder.)

Istlanus balsanus, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 118, Rio Ixtla, Morelos, Mexico.

Page 211. After *Algansea tincella* (Cuvier & Valenciennes), insert:

337 (b). *ALGANSEA LACISTRIS*, Steindachner.

Head 3.67; depth 5; eye 4.75; snout 3.6; interorbital 3. D. III, 7; A. iii, 6; P. 17; V. 9; scales 76-13. Body elongate, strongly compressed behind; caudal peduncle long and slender, its length 5 in head, its depth 10; head pointed, rounded above, greatly increasing in width behind the eyes where the width is half the head's length; mouth slightly oblique, jaws equal; tip of chin slightly thickened; maxillary nearly reaching eye; gape 4 in head; preorbital boat-shaped, concave above, convex below, equal to eye, or 1.6 longer than wide; teeth 4-4, hooked, with compressed crowns. Origin of dorsal midway between anterior border of eye and base of caudal, a little in front of ventrals, whose origin is equally distant between center of eye and base of caudal; height of dorsal 1.75 in head, its base half its height; height of anal 2.14 in head, or twice its base; caudal forked, 1.2 in head, the middle rays not quite half length of longest; ventrals short, 2.17 in head, their tips not reaching vent; pectoral 1.6 in head. Scales firm, half-oval in shape, larger posteriorly; lateral line on middle of side, descending somewhat anteriorly. Color dark brownish gray above, a darker band along middle of side; color below lateral line becoming abruptly lighter gray, merging into silvery on under parts. One example 20 cm. long from Lake Patzcuaro, Mexico. (Steindachner.)

Algansea lacustris, STEINDACHNER, Einige Fischarten Mex., 10, pl. III, figs. 1-1b, 1895, Lake Patzcuaro, Mexico. (Coll. Princess Theresa von Bayern.)

Page 254. *Orcella*, proposed by us for a subgenus in *Notropis*, is preoccupied by *Orcella*, Gray, 1866, a genus of Cetacea. For our use of it we substitute *Orcula*.

Orcula, JORDAN & EVERMANN, new subgenus in *Notropis* (*oreca*).

Page

Head from dorsal curvilinear. Body slightly of maxillary scales becoming beyond the midwidth the long posterior tips of the very pale on side to upper jaw is a light also a similar lateral line 12 scales, in this species.

Close to before the larger and is also a la-

Known specimens length from

Notropis mu- and Gull No. 17, 3 Museum

Page 26

Head 4; of caudal pectoral 5 between dorsal outline of slightly rounded to

Page 260. After *Notropis anogenus*, Forbes, insert:

404 (a). **NOTROPIS MUSKOKA**, Meek.

Head 4; depth 5.5; eye 3.25; snout 3.75; D. 8; scales 36, 10 in series from dorsal to ventral fin; longest dorsal ray 1.34 in head; pectoral 1.5; ventral 2, reaching anal. Teeth 4-4, slightly hooked, grinding surface narrow.

Body rather terete and slender; snout bluntnish; mouth small and slightly oblique; lower jaw the shorter, slightly inclined; posterior end of maxillary scarcely reaching vertical from the front of orbit. Eighteen scales between nape and dorsal fin, the scales in this region being smaller and more crowded than on sides and posterior dorsal region, much resembling *Pimephales notatus* in this respect; lateral line incomplete, not extending beyond the fourth scale on each side. Origin of the first ray of dorsal fin midway between the base of the caudal fin and the tip of the snout; the longest (anterior) dorsal rays nearly three times the length of the posterior ones, the tips of the former extending considerably beyond the tips of the latter when the fin is deflexed. Color olivaceous, darker above, very pale below; a dark band about $\frac{1}{2}$ diameter of eye around snout and on side to base of caudal fin; on the snout this band is confined to the upper jaw; between this band and the darker color on the dorsal region is a lighter band of about the same width; a dark vertebral line present, also a similar one from base of anal to caudal fin. In some specimens the lateral line is absent on the first scales and appears at intervals on about 12 scales, but it is occasionally absent altogether. It is less developed in this species than in any other of the genus.

Close to *N. cayuga*, from which it differs in the reduced size of the scales before the dorsal, the more slender body, the sharper snout, the slightly larger and more oblique mouth, and the more incomplete lateral line. It is also a larger fish than *N. cayuga*.

Known only from Gull Lake, near Muskoka Lake, Ontario, where 24 specimens were obtained by Dr. Meek in September, 1899. They vary in length from 1.31 to 2.91 inches. (Named for Muskoka Lake.)

Notropis muskoka, MEEK. Notes on a Collection of Fishes and Amphibians from Muskoka and Gull Lakes: Publication 41, Field Columbian Museum, Zoological Series, Vol. I, No. 17, 308, November, 1899, Gull Lake, Ontario. (Type, No. 2964, Field Columbian Museum. Coll. Dr. Meek.)

Page 264. After *Notropis proeue* (Cope), insert:

412 (a). **NOTROPIS RASCONIS**, Jordan & Snyder.

Head 4; depth $3\frac{1}{2}$; eye $2\frac{1}{2}$ in head; snout $3\frac{1}{2}$; interorbital space 3; depth of caudal peduncle $2\frac{1}{2}$; height of dorsal $4\frac{1}{2}$ in length; anal 5; length of pectoral $5\frac{1}{2}$; ventral $5\frac{1}{2}$; caudal $3\frac{1}{2}$; D. I, 8; A. I, 8; P. 12; scales 33, 15 between dorsal and occiput, 10 between dorsal and ventral fins. Dorsal outline of body evenly rounded from snout to insertion of dorsal fin, slightly concave from latter point to base of caudal; ventral outline evenly rounded to end of anal base; deepest part of body just anterior to insertion of dorsal; width of body $\frac{1}{2}$ of its length. Snout pointed; mouth

oblique, lower jaw included; maxillary not quite reaching vertical from anterior margin of orbit. Teeth 4-4, frail and easily detached from the arch, slightly hooked, no evident grinding surface; gill-rakers short and blunt, reduced to slight elevations on lower part of first arch. First ray of dorsal fin longest, last ray much shorter; when the fin is elevated, the posterior margin is straight, when depressed, it is somewhat falcate; anal fin similar in shape; pectoral pointed; tips of ventrals reaching anal; caudal deeply forked, the lobes pointed. Color silvery, a dark lateral band, the width of a scale, extending from tip of snout through eye to base of caudal, where it ends in an elongate, black spot; body above lateral band stippled with black, the dots grouped closely together on top of head, and in a narrow dorsal band extending from occiput to insertion of dorsal, in a sharply defined black line along base of dorsal fin, and also on the edges of the dorsal scales; body below dark band immaculate, except a dusky line along base of anal fin; all the fin rays dusky, especially the partly developed caudal rays, which are noticeably darker than the others.

Notropis rasconis is closely related to *N. nigrotentacularis*, from which it differs in having a more compressed body, a longer snout, larger eye, longer and more pointed fins, and in having the color-band narrower on the body and more marked on the snout. (Jordan & Snyder.)

Known only from the Rio Verde near Rascon, San Luis Potosi, Mexico, where the type (No. 6153, L. S. Jr. Univ. Mus.) was collected by J. O. Snyder.

(Named for Rascon, near which place the type was obtained.)

Notropis rasconis, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 121, Rio Verde, near Rascon, Mexico.

Page 302. Before *Erieymba*, Cope, insert:

123 (a). XYSTROSUS, Jordan & Snyder.

Xystrosus, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 123 (*popoche*).

Body long, compressed; interorbital space low and flat; mouth terminal, oblique; jaws subequal; premaxillaries protractile; no barbel; no pseudobranchia; gill-rakers 6+6, long, slender, crowded on arch; teeth 4-4, hooked, grinding surface oblique, grooved; alimentary canal about twice as long as body; peritoneum dusky; lateral line complete, decurved above pectoral; scales 61, evenly distributed over body; fins falcate; dorsal inserted directly above insertion of ventral; caudal deeply forked; anal basis short. Allied to *Notropis*; differing in the long gill-rakers, the small scales, and the absence of pseudobranchia. ξιστρός, (gill-)raker.

498 (a). XYSTROSUS POPOCHE, Jordan & Snyder.

Head 3½; depth 4½; eye 4½ in head; snout 3½; depth of caudal peduncle 2½; height of dorsal 5 in length; anal 6; length of pectoral 6; ventral 6½; caudal 4; D. 8; A. 7; P. 16; scales 61, 24 between dorsal and occiput, 16 between dorsal and lateral line. Body long and slender; caudal peduncle deep, compressed; head long, its upper contour straight;

interorbital contained at end; not quite slender, thin surface canals crowded the contour decurved 3 times its edge is sharp; color silver more or less tin and thin.

Exact Length of of length, interorbital from snout from snout anal to eye distance & Snyder.

Only th O. Snyder

Xystrosus p Chapala

Falcula, Jo

Body long maxillary surface granular peritoneum lateral scales anal basis closely adjacent.

The g small scales

Head 3½ depth of pectoral 6½ 19 between dorsal and lateral line.

interorbital space broad and flat; eye large; its longitudinal diameter contained 2 times in interorbital space; snout sharp, slightly turned up at end; mouth large, oblique; lower jaw included; lips thin; maxillary not quite extending to orbit; gill-rakers 6+6, on first arch, close together, slender, the longest $\frac{1}{2}$ diameter of orbit. Teeth 4-1, strong, hooked, grinding surface oblique, narrow, grooved, a notch just below the hook; alimentary canal almost twice the length of body; peritoneum dusky. Scales not crowded anteriorly, evenly distributed over body; lateral line shaped like the contour of body, except above pectoral fin, where it is sharply decurved; dorsal inserted directly above ventral, first ray highest, nearly 3 times height of last; when depressed, the fin is falcate; when elevated, its edge is concave; anal similar in shape; caudal deeply forked, the tips sharp; ventrals pointed, not reaching vent; pectoral slightly rounded. Color silvery; darker above, especially on median dorsal area, where a more or less dusky band extends the length of the body; rays of dorsal fin and tips of caudal dusky; lower fins white.

Exact measurements of the only specimen obtained are here given: Length of body in millimeters 92; depth of body, expressed in hundredths of length, 23; depth of caudal peduncle $11\frac{1}{2}$; length of head $28\frac{1}{2}$; width of interorbital space $11\frac{1}{2}$; length of snout 8; diameter of orbit 6; distance from snout to dorsal fin 52; height of longest dorsal rays 20; distance from snout to anal fin 73; height of longest anal rays 17; distance from anal to caudal fin 21; length of caudal fin 25; length of pectoral fin 18; distance from snout to ventral fin 53; length of ventral fin 16. (Jordan & Snyder.)

Only the type known (No. 6151, L. S. Jr. Univ. Mus.) collected by J. O. Snyder in Laguna de Chapala, near Ocotlan, Jalisco, Mexico.

Xystrosus popoche, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 123, Laguna de Chapala, Mexico.

123 (b). **FALCULA**, Jordan & Snyder.

Falcula, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 124 (*chapale*).

Body long, compressed; caudal slender; mouth large, lips thin, premaxillaries protractile; no barbel; teeth in 1 row, 4-4, hooked, grinding surface grooved; gill-rakers few, short, far apart; alimentary canal short; peritoneum silvery; lateral line complete; scales rather small, about 50 in lateral series. Fins high, falcate; dorsal inserted directly over ventrals; anal basis short; first simple rays of dorsal and anal rudimentary and closely adnate to first branched ray; caudal long, deeply notched.

The genus *Falcula* is related to *Notropis*, from which it differs in its small scales and in its very high falcate fins. (*falx*, scythe.)

498 (b). **FALCULA CHAPALE**, Jordan & Snyder.

Head 4; depth $4\frac{1}{2}$; eye $3\frac{1}{2}$ in head; snout $3\frac{1}{2}$; interorbital space $3\frac{1}{2}$; depth of caudal peduncle $9\frac{1}{2}$; height of dorsal 4 in length; anal 5; length of pectoral $4\frac{1}{2}$; ventral $5\frac{1}{2}$; caudal $3\frac{1}{2}$; D. 8; A. 8; P. 17; scales 50, 19 between dorsal and occiput, 16 between dorsal and ventral fins.

Body long, slender, compressed; snout pointed, its length equal to diameter of orbit or to interorbital space; mouth almost horizontal, its cleft extending to vertical from anterior edge of orbit; lips thin; teeth 4-4, strong, slightly hooked, the hook barely evident on lower tooth; three upper teeth with a grooved grinding surface; gill-rakers 3+7, short, pyramidal, and far apart; alimentary canal short; peritoneum silvery; lateral line complete, decurved to a point in a vertical midway between pectoral and ventral fins, from which it extends in a straight line a little below middle of body to the caudal; fins all very high and pointed; dorsal inserted directly over origin of ventrals, falcate when depressed, its first ray longest; anal similar in shape, inserted at a point $\frac{2}{3}$ the distance from tip of snout to base of caudal; ventrals extending to vent, tip of pectoral reaching ventrals; caudal deeply forked. Color silvery; a narrow, dark median band extending down from occiput to base of caudal; dorsal scales with fine dots which give their edges a dusky color.

One specimen known, careful measurements of which are given: Length of body in millimeters 74; depth of body, expressed in hundredths of length, 25; depth of caudal peduncle 11; length of head 25; width of interorbital space 7; length of snout 7; diameter of orbit 7; distance from snout to dorsal fin 47; height of longest dorsal rays 26; distance from snout to anal fin 67; height of longest anal rays 21; distance from anal to caudal fin 27; length of caudal fin 3 $\frac{1}{2}$; length of pectoral fin 22; distance from snout to ventral fin 62; length of ventral fin 21. (Jordan & Snyder.)

Only the type known (No. 6152, L. S. Jr. Univ. Mus.) collected by J. O. Snyder in Laguna de Chapala, near Ocotlan, Jalisco, Mexico.

(Named for the type locality.)

Faleula chapalae, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 124, Laguna de Chapala, Mexico.

Page 311. Before *Agosia nubila* (Girard), insert the following:

514 (a). AGOSIA KLAMATHENSIS, Evermann & Meek.

Head 4; depth 4 $\frac{1}{2}$; eye 4 $\frac{1}{2}$ in head; snout 3 $\frac{1}{2}$; D. I., 8; A. I., 7; scales about 14-78-10 (average, 73 in 49 specimens).

Body robust, subcylindrical, back somewhat elevated; snout rather long, mouth inferior, little oblique, the lower jaw included; maxillary not reaching front of orbit; upper lip without frenum; barbel present, but small. Lateral line incomplete, interrupted in many places, about 30 pores developed. Origin of dorsal fin midway between front of pupil and base of caudal fin; pectoral rather short, reaching about three-fourths the distance to base of ventrals; ventrals reaching vent; anal large, its longest ray 1 $\frac{1}{2}$ in head; caudal widely forked.

Color in alcohol: Olivaceous, mottled and blotched with darker on back and side; under parts pale; an obscure pale streak from eye to base of caudal fin, below which is a broad dark band; dorsal, pectoral, and caudal dusky; other fins plain; a black blotch at base of caudal.

An examination of 48 specimens shows all the important characters to

be fairly absent.

average

different

which the

others the

Close to

Upper

Agosia klaw

fig. 5.

Coll. M.

Page 3

Head 4;
scales 28-

anal mandib

distant b

ventrals;

1 $\frac{1}{2}$ in hea

basal half

gray. L.

Darien; o

Piabucina

Torino,

Page 3

Stations 1

long. 40 $\frac{1}{2}$

and 1,695

Page 3

Head 7;
2.2; inter

not close

than rest

lower ja

Dorsal fi

Color:

longitudi

and bod

head sma

irregular

idle of sid

white sp

more qu

cutting t

be fairly constant. The barbel is in a few cases obscure or possibly absent. The number of scales in a transverse line varies from 68 to 78, the average being 73. The lateral line is in all cases incomplete, though in different degrees. Sometimes it is continuous for only 6 or 8 scales, after which there are several interruptions and only 8 to 10 more pores. In others there are 20 to 30 pores in a continuous series.

Close to *A. nubila*.

Upper Klamath Lake, Oregon.

Ayosia klamathensis, EVERMANN & MEEK, Bull. U. S. Fish Com. 1897 (Jan. 6, 1898), 74, fig. 5. Pelican Bay, Upper Klamath Lake, Oregon. (Type, No. 48225, U. S. N. M. Coll. Meek & Alexander.)

Page 333. After *Piabucina panamensis*, Gill, insert the following:

552 (a). **PIABUCINA FESTE**, Boulenger.

Head $4\frac{1}{2}$; depth 4; eye $4\frac{1}{2}$; snout $4\frac{1}{2}$; interorbital width 9; D. 9; A. 11; scales 28-8; lower jaw longer than snout; teeth 26 above, 32 in external mandibular series; maxillary reaching eye; origin of dorsal equally distant between end of snout and caudal sinus, a little behind base of ventrals; adipose dorsal very small; anal with 8 branched rays; pectoral $1\frac{1}{2}$ in head, longer than ventrals; caudal strongly emarginate, scaled on basal half. Very deep olive brown above, whitish below; fins uniformly gray. Length 160 mm. (Boulenger.) From tributary of Lake Pita, Darien; only the type known. (Named for the collector, Dr. Enrico Festa.)

Piabucina festae, BOULENGER, Bollettino dei Mus. di Zool. ed Anat. Comp. della Univ. de Torino, Vol. XIV, No. 346. April 29, 1899, Rio Sabina, Darien. (Coll. Dr. Enrico Festa.)

Page 367. *Serrivomer beanii* has recently been taken by the *Ingolf* at Stations 12 and 20, lat. $64^{\circ} 38' N.$, long. $32^{\circ} 37' W.$, and lat. $58^{\circ} 20' N.$, long. $40^{\circ} 48' W.$, in Denmark Strait and SSE. of Cape Farewell, in 1,040 and 1,695 fathoms, respectively.

Page 398. After *Lycodontis obscuratus* (Poey), insert:

656 (a). **LYCODONTIS JORDANI**, Evermann & Marsh.

Head 7 in total length; depth about 14; eye 8 in head; snout 5; gape 2.2; interorbital a little less than snout. Teeth uniserial, strong, sharp, not close set, all entire and without basal lobes; tail considerably longer than rest of body; gill-opening smaller than eye; snout rather pointed, lower jaw the shorter, the mouth capable of being completely closed. Dorsal fin high, much higher than anal; nasal tube long, about 3 in eye.

Color: Tawny-ochraceous, paler below; upper jaw gray; iris blue; longitudinal brown stripes on side of head in front of gill-opening; head and body covered with numerous small, round, white spots, those on head smallest; a series of larger ones along upper part of side, and 1 or 2 irregular series of large ones on side of belly; between these on the middle of side the spots are smaller; dorsal with an irregular series of small white spots along the base, and another series of about 16 much larger, more quadrate spots of same color along edge of fin, some of the spots cutting the border, which is black; anal similarly spotted and with black

border. In alcohol the general color is grayish black, yellowish below, the tawny-ochraceous or yellow becoming darker, almost black, and the white spots on body becoming yellowish.

This species seems to be related to *L. obscuratus* (Poey), but differs from it markedly in color. Puerto Rico; only the type known, a specimen 15 inches long collected by the U. S. Fish Commission steamer *Fish Hawk* at Mayagüez, January 20, 1899. (Named for David Starr Jordan.)

Lycodonotis jordani, EVERMANN & MARSH, Rept. U. S. Fish Com. 1899 (Dec. 19), 352. Mayagüez, P. R. (Type, No. 49358, U. S. N. M. Coll. Evermann & Marsh.)

Page 447. After *Stolephorus productus* (Poey), insert:

734 (a). STOLEPHORUS GILBERTI, Evermann & Marsh.

Head 3.25; depth 3.4; eye 4; snout 6; maxillary 1.7; mandible 1.7; interorbital 4.9; D. 15; A. 23; pectoral 2.1; ventral 3.5; caudal 1.3; scales 42-9.

Body comparatively deep and strongly compressed, the belly trenchant, without serrations; snout thick, much projecting; maxillary reaching nearly to root of mandible, scarcely serrate; eye moderate; tip of lower jaw reaching vertical from front of eye; distance from lower posterior angle of cheek to vertical from posterior margin of opercle much less than from same point to eye; dorsal inserted far in advance of anal, just behind insertion of ventrals, midway between anterior edge of eye and base of caudal.

Color in spirits: Back light olivaceous with dark punctulations; rest of body below a line from shoulder to upper base of caudal silvery; faint traces of golden behind eye; no lateral band.

This species is very close to *Stolephorus garmani*, Evermann & Marsh, differing chiefly in the much smaller eye, the more uniform color of the back, the somewhat more sharply compressed belly, and the more nearly entire maxillary.

Puerto Rico; only the type, a specimen 4.5 inches long, known. This was collected by the U. S. Fish Commission expedition to Puerto Rico, at Palo Seco, near San Juan, January 13, 1899, associated with *S. productus*. (Named for Dr. Charles Henry Gilbert.)

Stolephorus gilberti, EVERMANN & MARSH, Rept. U. S. Fish Com. 1899 (Dec. 19, 1899), 352. Palo Seco, P. R. (Type, No. 49359, U. S. N. M. Coll. Evermann & Marsh.)

734 (b.). STOLEPHORUS GARMANI, Evermann & Marsh.

Head 3.2; depth 3.3; eye 3.5; snout 5.5; maxillary 1.7; mandible 1.7; interorbital 5; D. 14; A. 23; pectoral 2; ventral 3.5; caudal 1.3; scales 42-9.

Body comparatively deep and strongly compressed; the belly not strongly trenchant, without serrulations; snout thick, much projecting; maxillary reaching nearly to root of mandible, very finely and weakly serrate; eye large; tip of lower jaw reaching vertical from front of eye; distance from lower posterior angle of cheek to vertical from posterior margin of opercle much less than from same point to eye; dorsal inserted far in advance of anal, just behind insertion of ventrals, midway between anterior edge of pupil and base of caudal.

Color in reddish; silvery;

This sp. unquestionably farther b. It is very in the lar compresses

Puerto Rico was collected Puerto Rico Museum of *Stolephorus* 352. Pue

Page 49

Head 3.3; D. 10; A. 1. elevated, a tip of snout margin of maxillary 1.5; tip of second dorsal ray 2½ in 1. blue above, above lateral line, nearly white, few on canthus, on cheek, opercle; preopercle; pectoral

The genera of Lake C. and the up

Professor "This tr. 'Salmon T. upper outlet style of col. Blue Back these lake moreover, account of can not be

Bu

Color in spirits: Back dark near the median line, below this somewhat reddish; rest of body below a line from shoulder to upper base of caudal silvery; some golden on snout and behind eye; no lateral band.

This species has a general resemblance to *Stolephorus productus*, but is unquestionably distinct from it; the anal is much shorter and inserted farther back, the body is deeper, the eye larger, and the snout longer. It is very close to *Stolephorus gilberti*, Evermann & Marsh, differing chiefly in the larger eye, in the color of the back, and the somewhat less sharply compressed belly.

Puerto Rico; only the type, a specimen 4.5 inches long, known. This was collected by the U. S. Fish Commission expedition to Puerto Rico at Puerto Real, January 27, 1899. (Named for Prof. Samuel Garman, of the Museum of Comparative Zoology.)

stolephorus garmani. EVERMANN & MARSH, Rept. U. S. Fish Com. 1899 (Dec. 19, 1899), 352. Puerto Real, P. R. (Type, No. 49360, U. S. N. M. Coll. Evermann & Marsh.)

Page 497. Before *Salmo gairdneri*, Richardson, insert the following:

780 (k). **SALMO CLARKII DECLIVIFRONS**, Meek.

(SALMON TROUT OF LAKE SOUTHERLAND.)

Head 3.8; depth 4.6; eye 5.14; snout 4.5; maxillary 1.6; scales 148; D. 10; A. 11; branchiostegals 10; gill-rakers 7+10. Body elongate, back elevated, anterior profile much decurved, especially so from nape forward; tip of snout below axis of body; margin of upper lip on a level with lower margin of orbit; gape nearly horizontal, more so than in other trout; maxillary broad, its greatest width 5 in its length, its posterior end reaching beyond eye; dentition strong; posterior margin of dorsal fin straight; tip of second ray reaching middle of last ray in the depressed fin, the last ray $2\frac{1}{2}$ in longest ray; pectoral 1.67 in head; ventral 2.25. Color dark blue above and on side to lateral line anteriorly, and to a short distance above lateral line posteriorly, then becoming abruptly silvery; belly nearly white; no spots on head or body, none on any of the fins, except a few on caudal fin; upper margin of lower jaw black, a dark blue patch on cheek, extending obliquely upward and backward to near top of opercle; pectoral, ventrals, and anal yellowish.

The general color of this species much resembles that of the Blue Back of Lake Crescent. It is some darker, has no spots, except on caudal fin, and the upper anterior profile is much more curved.

Professor Elliot gives the following note on this trout:

"This trout is occasionally taken in Lake Southerland, and is called the 'Salmon Trout.' It is easily recognizable, not only by the sharply curved upper outline of the fore part of the body, but also by its quite different style of coloration, resembling, as stated above, somewhat the style of the Blue Back of Lake Crescent. As there is no water connection between these lakes, and Lake Southerland is 75 feet lower than Crescent, and, moreover, the fish of that lake having no communication with the sea, on account of a very high precipitous fall a short distance from its outlet, it can not be supposed that these two forms are in any way identical. On

of a large number of trout taken by me in Lake Sontherland, only two or three specimens of this form were procured, and they were all of small size, and I did not understand that it was ever obtained of much greater dimensions. This could not be the fault of the lake, which is exceedingly deep, and nearly 3 miles in length. It is a gamey fish, takes the fly, leaps out of water, and is a good fighter for its size." (Meek.)

Length about 10 inches.

Known only from Sontherland Lake, Washington, where 2 or 3 specimens (only 1 of which was preserved) were taken by Professor Elliot in 1898. (*declivus, steep; frons, forehead.*)

Salmo clarki declivifrons, Meek. Notes on a collection of cold-blooded Vertebrates from the Olympic Mountains: Field Columbian Museum Publication 31, Zoological Series. Vol. 1, No. 12, 230, Feb., 1899. **Southerland Lake, Washington.** (Type, No. 2006, Field Columbian Museum.)

780 (d). *Salmo clarkii jordani*, Meek.

(SPOTTED TROUT OF LAKE SOUTHERLAND.)

Head 3.86,* depth 4.79; eye 5.89; snout 4.13; maxillary 1.78; scales 116; D. 10; A. 11; branchiostegals 10 or 11. Body elongate, not much compressed; head short, maxillary rather broad, not extending far behind orbit—its greatest width $4\frac{1}{2}$ in its length, in some specimens about $5\frac{1}{2}$, a slight curve downward under the eye; dentition not so strong as in the Speckled Trout of Lake Crescent; gill-rakers rather long, longer than in the Speckled Trout, but less so than in Blue Back of Lake Crescent; pectoral in head, 1.88; ventral 2.24; origin of dorsal midway between tip of snout and base of caudal, or slightly nearer tip of snout; origin of ventrals under first to third dorsal rays; margin of dorsal fin convex, its base 1.32 in longest ray; snout bluish. Color in alcohol, dark steel-blue above, paler below, becoming nearly white on the belly; back, sides, and head profusely spotted with black; some specimens with black spots on the belly and on all fins; pectorals and ventrals usually without spots; upper half of lower jaw black, red under dentary bones. The life colors are given by Professor Elliot in the following note:

"This beautiful species is exceedingly gamey, takes a fly readily even as late as October, is a great leaper when hooked, and fights à l'entrée. In appearance it resembles *Salmo gairdneri crescentis* of the neighboring lake, being fully as brilliantly colored, but can be at once distinguished by its orange or orange-red fins, red on the jaw, and the number and blackness of its spots, and darker back and top of head. In general appearance there is not the slightest similarity between this species and the specimen from Boulder Creek. At no stage of its existence that I have seen, from fingerlings to fish weighing over 4 pounds, is there any silvery luster, but the colors are all bright-hued, some even metallic. It is one of the most active of its tribe, and I have had them leap after taking the fly in such quick succession, and with such rapid dartings about the lake, that it was impossible to imagine where they would next appear. I be-

*These comparative measurements are the averages of 23 specimens examined by Dr Meek.

leave it
Lake So
and no i
Length

Known
were col

"Nam
more tha
Salmo clarki
Olympic
Vol. 1,
Field C

Page 5

Head 3.
lary 1.66;
9 to 11; 1
terior mar
very long
of the orb
lary; teeth
stronger t
dible very
lunnarou
last ray of
of fin; whe
last dorsal
gairdneri c
with black
has fewer s
dark, dark
gairdneri c
head more
narrower,
that this s
the dentiti

Prof. D.

"This is
Crescent is
much more
other tron
season of t
any kind o
within a fe
in the lake

leave it spawns in the spring, as in the middle of October, when I left Lake Sutherland, the eggs of the females we caught were not enlarged, and no indication of the approach of the spawning season." (Meek.)

Length 1 to 2 feet.

Known only from Sutherland Lake, Washington, where 23 specimens were collected in 1898 by Professor Elliot.

"Named for Dr. D. S. Jordan, president of Stanford University, who, more than anyone else, has studied our Western trout." (Meek.)

Salmo clarkii jordani, MEEK, Notes on a collection of cold-blooded Vertebrates from the Olympic Mountains; Field Columbian Museum Publication 31, Zoological Series, Vol. 1, No. 12, 229, February, 1899, Sutherland Lake, Washington. (Type, No. 2012, Field Columbian Museum.)

Page 500. Before *Salmo iridens*, Gibbons, insert the following:

480 (a). **SALMO BATHYCETOR**, Meek.

(LONG-READED TROUT OF CRESCENT LAKE.)

Head 3.5 to 3.8; depth 5.10 to 5.75; eye 6.75 to 7.60; snout 3.33; maxillary 1.66; scales 150 to 152; gill-rakers 7 or 8 + 11 to 13; branchiostegals 9 to 11; D. 10; A. 11. Body elongate, slender; head much pointed; anterior margin of upper jaw slightly above axis of the body; maxillary very long and very slender, reaching considerably beyond posterior part of the orbit, its greatest width 7 in its length; about 24 teeth on maxillary; teeth on jaws, vomer, and palatines large, the dentition much stronger than in specimens of *Salmo gairdneri crescentis* of same size; mandible very strong; opercle very broad and contains the eye 1½; preoperculum narrow, less than diameter of the eye; gill-rakers very short and thick; last ray of dorsal less than half longest rays, third ray longer than base of fin; when depressed, second and third ray tips reach beyond middle of last dorsal ray; pectoral 1½ in head; ventrals 2½. Color much as in *S. gairdneri crescentis*, except lighter. Head, body, and tail profusely spotted with black spots; ventrals and pectorals dark. No. 2036 (Field Museum) has fewer spots, none on anal and pectoral; anterior part of upper jaw very dark, darker than in the larger specimen. This species differs from *S. gairdneri crescentis* in being much more slender, its back much less elevated, head more slender and pointed, gill-rakers shorter, maxillary straighter, narrower, and longer. The general color pattern is the same, except that this species is less spotted and lighter. No red on the under jaw; the dentition is much stronger in this species than in *S. gairdneri crescentis*. (Meek.)

Prof. D. G. Elliott gives the following interesting note on this trout:

"This is a deep-water fish, keeping always near the bottom. Lake Crescent is of great depth, in some places over 700 feet, and doubtless much more in others not yet ascertained. The present species, unlike other trout, does not come to the surface, as I was informed, at any season of the year, and will not of course take a fly, or indeed a spoon, or any kind of lure. The only way it can be captured is by set-lines sunk within a foot of the bottom, and it seems that there are only a few places in the lake where it can be caught even by this means. The specimens

obtained were procured at a depth of about 200 feet. While it is a brightly colored fish, it lacks some of the iridescent hues of *S. gairdneri crescentis*, and consequently is less attractive in appearance. It is known as the long-nose, or long-headed trout." (Meek.)

Length nearly 2 feet.

Known only from Crescent Lake, Washington, where two specimens were obtained in 1898 by Prof. D. G. Elliot.

Salmo bathaeator, MEEK, Notes on a collection of cold-blooded Vertebrates from the Olympic Mountains: Field Columbian Museum Publication 31, Zoological Series, Vol. I, No. 12, 227, February, 1896, **Crescent Lake, Washington.** (Type, No. 2035, Field Columbian Musm.)

Page 572. *Myctophum gracilis* (Lütken) is reported by Lütken from Denmark Strait, west of Iceland.

Page 583. Before *Yarella*, Goode & Bean, insert the following:

875 (a). CYCLOTHONE MEGALOPS, Lütken.

Together with a great number of *Cyclothona microdon* captured at station 12— $64^{\circ} 38'$ lat. N., $32^{\circ} 37'$ long. W., 1,040 fathoms—there occurred a single specimen of a length of 70 mm., habitually looking much like the said species, but differing by the eyes not being particularly small, and by totally wanting the light glands or "photospheres." It can, therefore, apparently, hardly be referred to the same genus. The dorsal and anal fins are very like those of *C. microdon*, though with the difference that the dorsal fin begins somewhat before the anal fin, while this, on the other hand, ends somewhat farther back than the dorsal fin. Quite black. A somewhat larger specimen (105 mm.) from station 9— $64^{\circ} 18'$ lat. N., and 27° long. W., 295 fathoms—is so badly preserved that it gives only the information that the eyes are not small and that both jaws are armed with small teeth directed obliquely backward, with a few longer ones in the foremost part of the lower jaw and the foremost part of the palate or the intermaxillary. The nearer determination of this specimen must be reserved for a future discovery.

It seems evident that these specimens belong to species else unknown, but as the material is so scanty I shall limit myself to the short preliminary notes made above. (Lütken.)

Cyclothona (?) megalops, LÜTKEN, Ichth. Results Danish *Ingolf* Exped., Vol. II, 10, 1898, **west of Iceland.**

Page 617. *Macdonaldia rostrata* was taken in 1895 by the *Ingolf* expedition west of Iceland.

Page 669. After *Characodon variatus*, Bean, insert:

988 (a). CHARACODON ENCAUSTUS, Jordan & Snyder.

Head 4; depth $3\frac{1}{2}$; depth of caudal peduncle 8; eye 3 in head; snout 4; interorbital space $3\frac{1}{2}$; height of dorsal $4\frac{1}{2}$ in length; anal $6\frac{1}{2}$; length of pectoral $5\frac{1}{2}$; ventral $6\frac{1}{2}$; caudal $4\frac{1}{2}$; D. 16; A. 15; scales 35, 13 transverse series counting upward and forward from origin of anal, 9 on caudal

peduncle, snout to C ventral of eye very large equal to 1 $\frac{1}{2}$ diameter in 2 series attached; not extending half way beyond less than rounded; vent. Space below small scale and at low in alcohol median par caudal; second part of head with a little

This species having more pressed body

Known only where J. C. MUS. (EVR) *Characodon* de Chapeau

Page 682.

Xenendrum, J. C. MUS.

Body deep size. Eye attached, in jaw; the mouth opening not to the diameter; air-bladder and anal in short; anal convex—thin ventral fins

Xenendrum loosely attached

peduncle. Body deep, compressed, dorsal outline almost straight from snout to origin of dorsal, concave from the latter point to base of caudal; ventral outline evenly curved from snout to posterior part of base of anal; eye very large, nearer snout than to posterior edge of opercle by a distance equal to longitudinal diameter of pupil; mouth small, its width equal to $\frac{1}{3}$ diameter of pupil; maxillary protractile; lower jaw projecting; teeth in 2 series, the outer series small, bicuspid, in a single row, rather firmly attached; inner series minute, in small patches; gill-opening restricted, not extending above base of pectoral fin; gill-rakers slender, equal in length to $\frac{1}{2}$ diameter of pupil. Alimentary canal short; air-bladder large, extending posteriorly to a point above origin of anal; dorsal fin inserted halfway between tip of snout and base of caudal, length of base a little less than height of fin; anal inserted below middle of dorsal, its edge rounded; pectoral extending to base of ventrals; ventrals extending to vent. Scales on body large; upper posterior part of head and a narrow space below and posterior to eye with scales; other parts of head naked; small scales on basal part of caudal fin; a row of large pores above eye and at lower edge of suborbital patch of scales; no lateral line. Color in alcohol light, yellowish olive; 9 short and narrow vertical bands on median part of body, the first above base of pectoral, the ninth at base of caudal; scales on dorsal region of body edged with black dots; upper part of head dark; upper half of orbit black; opercles silvery; dorsal fin with a little dusky; other fins without dark color.

This species somewhat resembles *C. eiseni* in appearance. It differs in having more rays in the dorsal and anal fins, smaller scales, a more compressed body, and less dark color on the body. (Jordan & Snyder.)

Known only from Laguna de Chapala, near Ocotlán, Jalisco, Mexico, where J. O. Snyder collected the type, a female, No. 6163, L. S. Jr. Univ. Mus. (*ερχαντός*, branded.)

Characodon encaustus, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 126, Laguna de Chapala, Mexico.

Page 685. After *Goodea atripinnis*, Jordan, insert:

313 (a). **XENENDUM**, Jordan & Snyder.

Xenendum, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 127 (*caliente*).

Body deep, not much compressed; males and females of about the same size. Eye normal; mouth vertical, lower jaw projecting; teeth loosely attached, in 2 series, the first series flat, bicuspid, in 2 or 3 rows on each jaw; the second series minute, in villiform bands, sometimes absent; gill opening not restricted, extending above the pectoral fin a distance equal to the diameter of pupil. Alimentary canal long, with many convolutions; air-bladder present, large. Scales large; no lateral line. Dorsal and anal inserted posteriorly, the one directly over the other, their bases short; anal very slightly modified in male, first rays shorter, edge double-convex—the notch being between smaller and larger groups of rays; ventral fins present; caudal rounded, and not modified in male.

Xenendum differs notably from *Characodon* in having the bicuspid teeth loosely attached and in more than one series. *Characodon* has the bicuspid

teeth firmly attached and in a single series. Its actual affinities are with the genus *Goodea*, and it belongs to the subfamily *Goodeinae*, which has the general characters of the *Paciliinae*, but with bifurcate or trifurcate teeth, and no great differences between the sexes. (ξενός, strange; ἐνδόν, within.)

1008 (b). **XENENDUM CALIENTE**, Jordan & Snyder.

Head $3\frac{1}{2}$; depth $2\frac{3}{4}$; depth of caudal peduncle $5\frac{1}{2}$; eye 4 in head; snout $3\frac{1}{2}$; interorbital space 2; height of dorsal $5\frac{1}{2}$ in length; anal $6\frac{1}{4}$; length of pectoral 5; ventral $7\frac{1}{2}$; caudal 5; D. 13; A. 14; scales 36, in transverse series counting upward and forward from origin of anal, 14; on caudal peduncle, 9. Body large and thick set, deepest at tip of pectoral; width $4\frac{1}{4}$ times in length; head pointed; interorbital space broad, slightly convex; length of snout about equal to diameter of orbit; mouth vertical, its width equal to length of snout; maxillary very protractile; teeth loosely attached, in 2 series, those of the first series larger, flat, and notched, in 2 rows on upper jaw, 3 rows on lower, the individual teeth of each row alternating in position with those of the next, those of the second series very minute, in a villiform band; gill-opening not restricted, extending above the pectoral a distance equal to diameter of pupil; gill-rakers long, slender, and close together, 40 on first arch. Alimentary canal long (in one specimen $4\frac{1}{2}$ times the length of body), coiled many times; peritoneum black; genital opening close to base of anal, covered by a thick, notched pad. Dorsal fin inserted posteriorly, rounded, the base short, its length less than height of fin; anal inserted under dorsal, first 5 rays crowded together and shortened; edge of fin double-convex, the notch being between the shorter and longer sets of rays; pectorals and ventrals with rounded edges; caudal evenly rounded; body and head everywhere, except jaws and preorbital area, with scales; no lateral line. Color light olive, growing darker above; medium dorsal area blackish, each scale with a dark, angular band, those of the sixth series below the dorsal darker, making an indistinct, narrow lateral band; all the fins, except ventrals, dusky.

The females differ but slightly from the males. The body is more thick set, the caudal peduncle a little less deep, the fins a little lower, and the anal evenly rounded. The young are somewhat mottled in color.

Xenendum caliente differs from *X. luitpoldii* in having fewer scales in the lateral and transverse series and on the caudal peduncle, and in a similar way from *X. xaliscone*, besides having villiform teeth, which are absent in *X. xaliscone*. (Jordan & Snyder.)

Known only from Rio Verde, near Aguas Calientes, Mexico. (Type, a female, No. 6147, L. S. Jr. Univ. Mus. Coll. J. O. Snyder.)

Xenendum caliente, JORDAN & SNYDER, Bull U. S. Fish Com. 1899 (1900), 127, Rio Verde, near Aguas Calientes, Mexico.

Characodon luitpoldii, Steindachner (p. 2832) belongs to this new genus, and should stand as

1008 (b). **XENENDUM LUITPOLDII** (Steindachner).

Head $3\frac{1}{2}$; interorbital space 2; height of dorsal $5\frac{1}{2}$; anal $6\frac{1}{4}$; length of caudal peduncle 5; ventral $7\frac{1}{2}$; caudal 5; D. 13; A. 14; scales 36, in transverse series counting upward and forward from origin of anal, 14; on caudal peduncle, 9. Body large and thick set, deepest at tip of pectoral; width $4\frac{1}{4}$ times in length; head pointed; interorbital space broad, slightly convex; length of snout about equal to diameter of orbit; mouth vertical, its width equal to length of snout; maxillary very protractile; teeth loosely attached, in 2 series, those of the first series larger, flat, and notched, in 2 rows on upper jaw, 3 rows on lower, the individual teeth of each row alternating in position with those of the next, those of the second series very minute, in a villiform band; gill-opening not restricted, extending above the pectoral a distance equal to diameter of pupil; gill-rakers long, slender, and close together, 40 on first arch. Alimentary canal long (in one specimen $4\frac{1}{2}$ times the length of body), coiled many times; peritoneum black; genital opening close to base of anal, covered by a thick, notched pad. Dorsal fin inserted posteriorly, rounded, the base short, its length less than height of fin; anal inserted under dorsal, first 5 rays crowded together and shortened; edge of fin double-convex, the notch being between the shorter and longer sets of rays; pectorals and ventrals with rounded edges; caudal evenly rounded; body and head everywhere, except jaws and preorbital area, with scales; no lateral line. Color light olive, growing darker above; medium dorsal area blackish, each scale with a dark, angular band, those of the sixth series below the dorsal darker, making an indistinct, narrow lateral band; all the fins, except ventrals, dusky.

One male known, appearing tentatively as a specimen from Chapala, Mexico.

Xenendum caliente differs from *X. luitpoldii* in having fewer scales in the lateral and transverse series and on the caudal peduncle, and in a similar way from *X. xaliscone*, besides having villiform teeth, which are absent in *X. xaliscone*. (Jordan & Snyder.)

Known only from Rio Verde, near Aguas Calientes, Mexico. (Type, a female, No. 6147, L. S. Jr. Univ. Mus. Coll. J. O. Snyder.)

Xenendum caliente, JORDAN & SNYDER, Bull U. S. Fish Com. 1899 (1900), 127, Rio Verde, near Aguas Calientes, Mexico.

Characodon luitpoldii, Steindachner (p. 2832) belongs to this new genus, and should stand as

1008 (b). **XENENDUM LUITPOLDII** (Steindachner).

Head $3\frac{1}{2}$; interorbital space 2; height of dorsal $5\frac{1}{2}$; anal $6\frac{1}{4}$; length of caudal peduncle 5; ventral $7\frac{1}{2}$; caudal 5; D. 13; A. 14; scales 36, in transverse series counting upward and forward from origin of anal, 14; on caudal peduncle, 9. Body large and thick set, deepest at tip of pectoral; width $4\frac{1}{4}$ times in length; head pointed; interorbital space broad, slightly convex; length of snout about equal to diameter of orbit; mouth vertical, its width equal to length of snout; maxillary very protractile; teeth loosely attached, in 2 series, those of the first series larger, flat, and notched, in 2 rows on upper jaw, 3 rows on lower, the individual teeth of each row alternating in position with those of the next, those of the second series very minute, in a villiform band; gill-opening not restricted, extending above the pectoral a distance equal to diameter of pupil; gill-rakers long, slender, and close together, 40 on first arch. Alimentary canal long (in one specimen $4\frac{1}{2}$ times the length of body), coiled many times; peritoneum black; genital opening close to base of anal, covered by a thick, notched pad. Dorsal fin inserted posteriorly, rounded, the base short, its length less than height of fin; anal inserted under dorsal, first 5 rays crowded together and shortened; edge of fin double-convex, the notch being between the shorter and longer sets of rays; pectorals and ventrals with rounded edges; caudal evenly rounded; body and head everywhere, except jaws and preorbital area, with scales; no lateral line. Color light olive, growing darker above; medium dorsal area blackish, each scale with a dark, angular band, those of the sixth series below the dorsal darker, making an indistinct, narrow lateral band; all the fins, except ventrals, dusky.

1008 (e). **XENENDUM XALISCOME**, Jordan & Snyder.

Head $4\frac{1}{2}$; depth $3\frac{1}{2}$; depth of caudal peduncle $6\frac{1}{2}$; eye $3\frac{1}{2}$ in head; snout 3; interorbital space $1\frac{1}{2}$; height of dorsal $6\frac{1}{2}$ in length; anal 9; length of pectoral $5\frac{1}{2}$; ventral $7\frac{1}{2}$; caudal 5; D. 13; A. 14; scales 42, transverse series, counting upward and forward from origin of anal, 17; on caudal peduncle 12. Body thick set, deepest at origin of ventrals, widest at bases of pectorals; caudal peduncle deep and long; head large and pointed; interorbital space broad, slightly convex; mouth vertical, its width equal to length of snout; maxillary very protractile; teeth loosely attached, broader at distal ends than at bases, bicuspid, in two rows on each jaw; no villiform teeth present; gill openings extending above base of pectorals a distance about equal to diameter of pupil; gill-rakers long, flat, very close together, 56 on first arch. Alimentary canal long, in many folds; peritoneum black. Dorsal fin inserted posteriorly, first ray simple, closely attached to second; edge of fin rounded; anal inserted on a vertical passing through the base of fourth dorsal ray, similar to dorsal in shape; pectoral and ventral fins rounded; edge of caudal a little convex; basal fourth with scales. Scales large, everywhere on body and head, except lower jaw and preorbital area; no lateral line. Color plain, dark above, light below, the dark color leaving off rather abruptly on the head along a line passing through lower edge of eye; on the body, along a line passing from lower edge of base of pectoral to caudal, leaving lower one-fifth of caudal peduncle light; faint traces of a dark spot at base of each scale on dorsal region of body; all the fins, except ventrals, dusky. Laguna de Chapala, Mexico.

One male individual was taken. It resembles the female in general appearance. The anal fin is not advanced nor modified into an intromittent organ. Although it is injured, it shows that the first 5 or 6 rays were close together and shortened.

Xenendum xaliscone differs from *X. caliente* in not having villiform teeth, in having more scales in the lateral and transverse series, and on the caudal peduncle. It differs from *X. luitpoldii* (Steindachner), which is the third known species of the genus, in having a much longer snout, a more pointed head, and in not having villiform teeth. (Jordan & Snyder.)

Known only from Laguna de Chapala, near Ocotlan, Jalisco, Mexico. (Type, a female, No. 6148, L. S. Jr. Univ. Mus. Coll. J. O. Snyder.)

(Name from Jalisco, the type locality.)

Xenendum xaliscone, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 129, Laguna de Chapala, Mexico.

Page 698. Before *Mollienisia*, Le Sueur, insert:

1037 (a). **POECILIA LIMANTOURI**, Jordan & Snyder.

Head $3\frac{1}{2}$ in length; depth 3; depth of caudal peduncle $4\frac{1}{2}$; eye $3\frac{1}{2}$ in head; snout 3; interorbital space 2; height of dorsal $5\frac{1}{2}$ in length; anal $4\frac{1}{2}$; length of pectoral $4\frac{1}{2}$; ventral 6; caudal $3\frac{1}{2}$; D. 9; A. 8; scales 26-9, 8 on caudal peduncle. Body rather deep and compressed; dorsal outline angular, its highest point at insertion of dorsal; lowest point of

ventral outline at base of ventrals. Head pointed, interorbital space wide and flat; eye large, nearer tip of snout than to posterior edge of opercle by a distance equal to diameter of pupil; mouth very oblique, its width equal to 2 times diameter of pupil; premaxillaries protractile; distal end of maxillary visible; lower jaw projecting; teeth in two series on each jaw, the outer series in a single row, small, pointed, loosely attached; second series scarcely discernible, in bands; gill-openings extending above base of pectoral a distance equal to $\frac{1}{2}$ diameter of orbit; gill-rakers on first arch 20, small and slender; alimentary canal very long and slender. Body and entire head, except preorbital area, lips, and lower jaw, covered with large scales; 3 rows of scales on base of caudal; small scales extending on inter-radial membranes of caudal, a distance beyond the basal scales about equal to diameter of the eye. Dorsal fin inserted halfway between base of caudal and anterior edge of pupil, its base contained $6\frac{1}{2}$ times in length of head and body, its height $5\frac{1}{2}$, the last rays a little higher than the first; anal advanced close to base of ventrals, the first and second rays short, closely attached to the next, third ray greatly enlarged and lengthened, a loosely attached ovate, fleshy pad near its tip; fourth and fifth rays slender, as long as third; tips of third and fifth rays bent toward that of the fourth; sixth to eighth rays about $\frac{1}{2}$ as long as third; caudal rounded, its length contained $3\frac{1}{2}$ times in head and body; pectoral rounded, the length contained $1\frac{1}{2}$ times in head; ventrals pointed, extending to middle of longest anal ray. Color in alcohol, light yellowish olive, much lighter on breast and ventral part of head; posterior edges of scales dark; lower jaw, preorbital area, upper part of head, and a narrow, median dorsal stripe, dark; basal $\frac{2}{3}$ of dorsal fin black, distal part of fin white, the boundary between the white and black more definite on the anterior than on posterior part of fin; basal $\frac{2}{3}$ of caudal dusky, distal part without color.

Other male examples have only a few small dark spots on dorsal and caudal. The females have the body more elongate than the males, the depth of the caudal peduncle $5\frac{1}{2}$ in length. The dorsal fin is inserted in advance of the anal, its origin above anal opening, the first rays highest. The ventrals extend to the posterior edge of the vent, but do not reach the anal. The dorsal and caudal have a little dusky coloring.

Lack of material for comparison prevents our commenting on the probable affinities between this and other species of the genus. (Jordan & Snyder.)

Known only from Rio Tamesoe, near Tampico, Tamaulipas, Mexico, where several specimens were collected January 12, 1899, by Mr. Snyder. (Type, a female, No. 6165, U. S. Jr. Univ. Mus.)

(Named for Jose Limantour, the accomplished minister of the "Hacienda" in Mexico.)

Pecilia limantouri, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 129, Rio Tamesoe, near Tampico, Mexico.

Page 702. Before *Xiphophorus guntheri*, Jordan & Evermann, insert:

1043 (n). XIPHOPHORUS MONTEZUMAE, Jordan & Snyder.

Head $4\frac{1}{2}$; depth 3; depth of caudal peduncle $4\frac{1}{2}$; eye $3\frac{1}{2}$ in head; snout $3\frac{1}{2}$; interorbital space 2; height of dorsal $3\frac{1}{2}$ in length; anal 5; length of

pectoral $4\frac{1}{2}$; ventral $4\frac{1}{2}$; upper rays of caudal $3\frac{1}{2}$, lower rays $1\frac{1}{2}$; D. 13; A. 7; scales 29-9, 7 on caudal peduncle. Body deep, compressed, dorsal contour arched, its highest point at insertion of dorsal; caudal peduncle narrow and very deep; head small, pointed; interorbital space wide, slightly convex; eye large, a little nearer tip of snout than to posterior edge of opercle; mouth vertical; teeth in two series, the first in a single row, minute, flat, and pointed, the second in a viiform band, much smaller and narrower than the first, brownish-colored, strongly curved backward; gill opening extending above base of pectoral a distance equal to diameter of pupil; gill-rakers on first arch 19, slender, the length of longest equal to half diameter of eye; intestinal tract slender and long; peritoneum black; scales on head and body large, 1 large, round scale on interorbital space, followed by 2, a row of 11 from the latter to first dorsal ray; 3 rows on base of caudal fin; base of dorsal fin short, $4\frac{1}{2}$ in body, first ray shortest, the others graduated to the eighth, which is longest; ninth, tenth, and eleventh shorter; twelfth and thirteenth longer; the abrupt shortening of the ninth, tenth, and eleventh rays makes a notch in the outline of fin; anal advanced, its origin under third ray of dorsal; first ray greatly enlarged and lengthened; second and third equal^{1/2} lengthened, but more slender, these three with their connecting membranes forming a half tube with a pointed end; other rays half the length of first; upper lobe of caudal rounded, 5 lower rays forming a very long, blunt appendage; ventrals pointed, extending almost to tip of anal; pectoral sharply rounded; color yellowish olive, marked with black; during life there were 4 narrow, longitudinal orange bands, each extending along a row of scales on body; top of head, and a median dorsal band extending to caudal, dusky; a narrow, dusky band on edge of lower jaw; 2 short, vertical bands on snout; 6 upper rows of scales edged with black or dusky; a few black spots irregularly arranged on body above ventrals; a large black spot at base of caudal, its color extending along upper edge of prolongation; a dark line extending along lower edge of caudal peduncle to end of lower caudal rays, the caudal extension with a light central portion bordered with black, the lower border wider; underpart of head and belly without dark color; dorsal fin with black dots and lines; pectorals, ventrals, and anal plain.

Considerable variation in shape of fins and in color is shown among other male examples (cotypes, males and females, No. 6146, L. S. Jr. Univ. Mus.). In some the fins are low and short, the caudal ornament represented only by a slight lengthening of the lower rays. Among individuals, apparently fully grown, there is every graduation from the undeveloped to the very long caudal extension. In every case the scales are conspicuously dark edged. In some examples black spots, crowded together, form a more or less dark line from eye to caudal, while below this line are large, irregular black blotches. Others have no black spots, and the dark caudal patch has almost disappeared.

The females have the fins low or short, and without special modifications, the posterior edge of caudal 2-shaped, the upper part rounded, the lower pointed, the scales dark edged; a narrow, indefinite, dark color band usually present along median line of sides; the dark caudal patch

rarely absent. Fully grown males are scarce, a large catch consisting mostly of females and young.

Xiphophorus montezumae is distinguished from the other known species of the genus by having 7 anal rays, the scales with conspicuous dark edges, a large brown caudal spot, and the caudal appendage not sword-shaped, but with its end enlarged and blunt. (Jordan & Snyder.)

Known only from Rio Verde, near Rascon, San Luis Potosi, Mexico. (Type, a male, No. 6145, L. S. Jr. Univ. Mus. Coll. J. O. Snyder.)

Xiphophorus montezumae, JORDAN & SNYDER, Bull. U. S. Fish Com. 1890 (1900), 131, Rio Verde, near Rascon, Mexico.

Page 702. The recent studies of the eyes of American blind fishes by Dr. C. H. Eigenmann have shown that the species occurring in the caves of southwestern Missouri which has usually been identified as *Typhlichthys subterraneus*, Girard, but which Dr. Eigenmann described as a new species under the name *Typhlichthys rosea* (p. 2835), has had an ancestry quite distinct from that of *Typhlichthys*. It therefore is generically distinct from *Typhlichthys*, and has been made the type of the new genus *Troglomystes* by Eigenmann.

322 (a). *TROGLICHTHYS*, Eigenmann.

Troglomystes, EIGENMANN, Science, N. S., Vol. ix, No. 217, p. 282, Feb. 24, 1899 (*rosea*).

Scleral cartilages present, pigment in the pigment epithelium; vitreal cavity obliterated, no hyaloid membrane; pupil closed, some of the eye muscles developed; no outer reticular layer; outer and inner nuclear layers merged into one; eye in adult not connected with the brain; pigment epithelium developed on the distal face of the eye, rarely over the sides and back; no cones; nuclear layers mere vestiges; ganglionic layer restricted to the anterior face of the eye just within the pigmented epithelium. Maximum diameter of eye about 85 μ . No ventral fins. (*Tρώγλη*, cavern; *ἰχθύς*, fish.)

The genera and species of Amblyopsidae may be determined by the following key (from Eigenmann), based largely on the character of the eyes:

a. Vitreous body and lens normal, the eye functional; no scleral cartilages; eye permanently connected with the brain by the optic nerve; eye muscles normal; no optic fiber layer. Minimum diameter of the eye 0.700 μ . No ventral fins. CHOCOASTER, 321.

b. Eye in adult more than 1 mm. in longitudinal diameter; lens over 0.5 mm. in diameter; retina very simple, its maximum thickness 83.5 μ in the old; the outer and inner nuclear layers consisting of a single series of cells each; the ganglionic layer of isolated cells. Maximum thickness of the outer nuclear layer 5 μ , of the inner nuclear layer 8 μ . CORNUCTUS, 1044.

bb. Eye in adult less than 1 mm. in longitudinal diameter; lens less than 4 mm.; outer nuclear layer composed of at least 2 layers of cells; inner nuclear layer of at least 3 layers of cells, the former at least 10 μ thick, the latter at least 18 μ .

c. Pigment epithelium 65 μ thick in the middle-aged, 102 μ in the old.

PAPILLIFERUS, 1046.

cc. Pigment 49 μ thick in the middle-aged, 74 μ in the old; 24 to 30 per cent thinner than in papilliferous; eye smaller. AGASSIZI, 1045.

aa. The eye a vestige, not functional; vitreous body and lens mere vestiges, the eye collapsed, the inner faces of the retina in contact. Maximum diameter of eye about 200 μ .

dd. No scleral cartilages, no pigment in the pigment epithelium; a minute vitreal cavity; hyaloid membrane with blood vessels; pupil not closed; outer nuclear, outer reticular, inner nuclear, inner reticular, ganglionie, and pigment epithelial layers differentiated; cones probably none; no eye muscles. Maximum diameter of eye 180 μ . Eye probably connected with brain throughout life.

TYPHLICHTHYS, 322.

dd. Scleral cartilages present; pigment in the pigment epithelium; vitreal cavity obliterated, no hyaloid membrane; pupil closed; some of the eye muscles developed; no outer reticular layer; outer and inner nuclear layers merged into one. Eye in adult not connected with the brain.

e. Pigment epithelium well developed; cones well developed; ganglionie cells forming a funnel-shaped mass through the center of the eye; pigmented epithelium over the front of the eye without pigment. Maximum diameter of eye about 200 μ . Ventral fins present.

AMBLVOPSIS, 323.

ee. Pigment epithelium developed over the distal face of the eye, rarely over the sides and back; no cones; nuclear layers mere vestiges; the ganglionie layer restricted to the anterior face of the eye, just within the pigmented epithelium. Maximum diameter of the eye about 85 μ . No ventral fins.

TROGLOCHTHYS, 322 (a).

Page 744. To the synonymy of *Eucalia inconstans* (Kirtland), add the following:

Gasterosteus gymnotus, DAWSON, Canadian Naturalist, Vol. IV, No. 5, October, 1859, 321-324, figs. 1-3, Montreal, Canada. (Coll. Prof. J. W. Dawson.)

Page 789. Recent investigations made by Dr. Jordan in certain freshwater streams and lakes of Mexico resulted in the discovery of several new species of Atherinidae and new facts regarding other species:

1157. *ESLOPSARUM JORDANI* (Woolman).

Locality, Rio Verde, Aguas Calientes, Mexico.

Two specimens of *Eslopsarum jordani* were collected, together with numerous individuals of *E. arge*. *Chirostoma breve*, Steindachner, is probably identical with *E. jordani*, as already supposed by us.

An examination of a number of species of *Eslopsarum* and of *Chirostoma* shows that the number of vertebrae, in addition to the character of the scales, furnishes a distinguishing generic feature.

Species.	Vertebrae.
<i>Eslopsarum jordani</i>	38
<i>arge</i>	37
<i>Chirostoma humboldtianum</i>	44
<i>chapale</i>	45
<i>promelas</i>	45
<i>diasi</i>	44
<i>crystallinum</i>	44
<i>terme</i>	44
<i>oceollane</i>	44

Chirostoma contains two very marked types of species. The one represented by the typical species, *Chirostoma humboldtianum*, has the flesh firm

and opaque in life. The other (*Lethostole*), typified by *Chirostoma estor*, has the flesh thin, translucent, and very pale. There are correlated differences in the firmness of the bones and scales, but thus far we have found no tangible character on which to separate *Lethostole* as a genus from *Chirostoma*. The known species of *Lethostole* are *estor*, *album*, *chapala*, *grandocula*, *promelas*, *diazii*, *crystallinum*, *lermae*, and *oceanum*.

1157 (n). *ESLOPSARUM ARGE*, Jordan & Snyder.

Head $\frac{1}{3}$; depth $4\frac{1}{2}$; depth of caudal peduncle $2\frac{1}{4}$ in head; eye $3\frac{1}{2}$; snout 3; interorbital space $3\frac{1}{2}$; height of spinous dorsal $3\frac{1}{2}$; soft dorsal 2; anal 2; length of pectoral $1\frac{1}{2}$; ventrals $2\frac{1}{2}$; caudal $1\frac{1}{2}$; D. IV-8; A. 15; P. 13; scales 40-41, 5 between the dorsals. Body rather thickset, its deepest part just anterior to base of ventrals; width of body equal to distance from posterior edge of orbit to tip of snout; eye nearer to tip of snout than to posterior edge of opercle by a distance equal to diameter of pupil; interorbital space convex; width of preorbital area equal to diameter of pupil; tip of lower jaw projecting beyond that of upper; mouth large, oblique; lips not much thickened posteriorly, the lower not distinctly folded over the upper at their angle; maxillary extending posteriorly to a perpendicular passing through anterior edge of orbit, its distal end below the level of eye; teeth large, sharp, projecting backwards, in 2 definite rows on each jaw, none on vomer or palatines; gill-rakers on first arch 14, long and slender; air-bladder extending posteriorly to a point a little past insertion of anal; peritoneum black; vertebrae 37. Lateral line represented on the fifth row of scales below the dorsal by a series of partly developed pores; scales large, entire, covering head and body except snout, lower jaw, preorbital area, and a small space around base of pectoral; small scales extending for a short distance on interradial membranes of caudal; first 3 dorsal spines of about the same height, the fourth shorter; first dorsal ray longest, the others gradually shorter; edge of fin straight; anal inserted on a perpendicular, passing halfway between dorsals, its first ray longest; edge of fin slightly concave; caudal notched, the tips and notch rounded, extending to bases of ventrals; ventrals falling short of vent a distance equal to diameter of orbit. Color in life, translucent; a silvery lateral band with its upper edge dark, extending from upper part of base of pectoral to base of caudal, the band less distinct in the region of the pectoral fin; scales of back edged with fine, dark specks; snout, lower jaw, top of head, and upper part of eye dusky; dorsal and caudal fins with a little dusky coloring.

Specimens of *E. arge* were caught in the same seine-haul with *E. jordani*. The former species differs from the latter in having a thicker body, a longer snout, a larger and less oblique mouth, a larger eye, and a wider color band.

In the drawing accompanying the original description of *Eslopsarum jordani* the mouth is wrongly represented. Of the specimens examined, including some of the types, the mouth is much like that of *Chirostoma humboldtianum*. The cleft is not straight in outline. The lower lip folds over the upper at their union. (Jordan & Snyder.) (*arge*, silvery.)

Known
No. 6154

Eslopsarum
near Agu-

Head $\frac{1}{3}$; inter-
1 $\frac{1}{2}$; long
pectoral
its deeper
to poster-
eter of P
eter of 1
ing a lit
thickene
angle of
its dista
tance eq
arranged
first arch
orbit; pe
a point
line exten
dorsal.
crowded
and ante-
membran
2 spines
second d
anterior
dorsals, t
elevated,
pectoral i
to diamet
cent; a s
becoming
scales on
jaws wit
arachnoid

C. chape
having a
the later
has 60 to

Known
(Type, N)

Chirostoma
de Chae

Known only from the Rio Verde, near Aguas Calientes, Mexico. (Type, No. 6154, L. S. Jr. Univ. Mus. Coll. J. O. Snyder.)

Elophus argenteus, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 133, Rio Verde, near Aguas Calientes, Mexico.

1157 (b). *CHIROSTOMA CHAPALE*, Jordan & Snyder.

Head 4; depth $5\frac{1}{2}$; depth of caudal peduncle 3 in head; eye $3\frac{2}{3}$; snout 3; interorbital space 4; height of spinous dorsal 3; soft dorsal $1\frac{1}{2}$; anal 1; length of pectoral $1\frac{1}{2}$; ventrals $2\frac{1}{2}$; caudal $\frac{3}{5}$; D. IV-10; A. 20; pectoral 14; scales 47-13, 6 between dorsals. Body slender, compressed, its deepest part below first dorsal; eye large, nearer tip of snout than to posterior edge of opercle by a distance equal to $1\frac{1}{2}$ times the diameter of pupil; interorbital space convex, its width about equal to diameter of pupil or to preorbital area; symphysis of lower jaw projecting a little beyond tip of upper, the lips meeting; mouth oblique, lips thickened posteriorly, the lower folding over the upper at their union; angle of mouth on a level with center of pupil; maxillary nearly vertical, its distal end in advance of a vertical from anterior edge of orbit, a distance equal to $\frac{2}{3}$ the diameter of pupil. Teeth minute, in bands, not arranged in definite rows; no teeth on vomer or palatines; gill-rakers on first arch 30, very slender, the length of longest equal to diameter of orbit; peritoneum black; air-bladder very large, extending posteriorly to a point above middle of anal fin; vertebrae 45. A well-defined lateral line extending along the body on the eighth row of scales below the first dorsal. Scales large, crenate, not notably reduced in size nor closely crowded together on any part of the body; those on post-occipital region and anterior to pectorals small; scales extending on basal $\frac{2}{3}$ of inter-radial membranes of caudal; lower jaw, snout, and preorbital space naked; first 2 spines of dorsal highest, the following 2 a little shorter; first ray of second dorsal highest, others gradually shorter; anal inserted a little anterior to a perpendicular passing halfway between the origins of dorsals, first ray longest, others successively shorter; when the fin is elevated, its edge is concave; caudal deeply forked, the tips pointed; pectoral notably pointed, extending past base of ventral a distance equal to diameter of orbit; ventrals extending to vent. Color in life, translucent; a silvery lateral band 1 scale wide, bright, and distinct posteriorly, becoming indistinct anteriorly; upper edge of lateral band with dusky; scales on dorsal part of body edged with dark dots; upper and lower jaws with dark dots; upper part of eye black, the dark, pigmented arachnoid showing through the thin skull. (Jordan & Snyder.)

C. chapale is closely related to *C. grandocule*, Steindachner. It differs in having a smaller eye and larger scales. The former has 44 to 51 scales in the lateral series, and 12 to 14 in a transverse series, while *C. grandocule* has 60 to 62 scales in the lateral series and 15 or 16 in a transverse series.

Known only from Laguna de Chapala, near Ocotlán, Jalisco, Mexico. (Type, No. 6165, L. S. Jr. Univ. Mus. Coll. J. O. Snyder.)

Chirostoma chapale, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 135, Laguna de Chapala, Mexico.

1157 (e). *CHIROSTOMA PROMELAS*, Jordan & Snyder.

Head $3\frac{1}{2}$; depth $4\frac{1}{2}$; depth of caudal peduncle $3\frac{1}{2}$ in head; eye $5\frac{1}{2}$; snout $2\frac{1}{2}$; interorbital space $4\frac{1}{2}$; height of spinous dorsal $3\frac{1}{2}$; soft dorsal $2\frac{1}{2}$; anal 2; length of pectoral $1\frac{1}{2}$; ventral $2\frac{1}{2}$; caudal $1\frac{1}{2}$; D. IV-11; A. 19; pectoral 15; scales 53-16, 9 between dorsals. Head slender, triangular; eye small, nearer tip of snout than to edge of opercle by a distance equal to half diameter of pupil; width of preorbital area somewhat greater than diameter of pupil; interorbital space slightly convex; snout pointed; upper jaw projecting a little beyond the lower; cleft of mouth almost horizontal; lips enlarged posteriorly, the lower lip folding over the upper at their junction; angle of mouth on a level with lower part of pupil; maxillary almost vertical in position, its distal end not extending backward as far as anterior edge of orbit; teeth large, curved inward, not arranged in definite rows; none on vomer or palatines; vertebrae 45. A tolerably well-defined lateral line extending along the body about 8 scales below the first dorsal, the pores absent on some of the scales; scales crenate, largest along lateral color band; a post-occipital patch of minute, closely-crowded scales extending backward nearly to a line connecting bases of pectorals; a narrow band of similar scales just posterior to gill-openings, and on base of caudal, the latter extending on inter-radial membranes half their length; scales between dorsals not abruptly smaller than those near by, nor crowded closely together; head with scales except on snout, preorbital region, and on lower jaw; first 3 spines of dorsal nearly equal in length, the fourth a little shorter; first dorsal ray longest, the others gradually shorter; anal inserted on a perpendicular passing midway between dorsals, first ray longest, others successively shorter; edge of fin slightly concave; pectoral pointed; extending beyond base of ventrals a distance equal to diameter of pupil; ventrals not quite reaching vent. Color in alcohol, yellowish olive; a distinct silvery lateral band, the light color of which is underlaid with dark pigment, extending from upper part of base of pectoral to caudal, wider and brighter in color between dorsal and anal, growing narrower on caudal peduncle, widening at its end; scales of upper part of body with dusky coloring on their edges; dorsals, pectoral, and caudal with dark color; eye dusky above, a dark band on interorbital space; snout and jaws black,

Chirostoma promelas is distinguishable from other known species of the genus by the projecting upper jaw and the black snout.

One specimen other than the type was obtained. In it the projection of the upper jaw is more pronounced than in the type. The gill-rakers and abdominal viscera had been removed from both. (Jordan & Snyder.)

Known only from Guadalajara, Jalisco, Mexico, where 2 specimens were obtained in the market. (Type, No. 6156, L. S. Jr. Univ. Mus. Coll. J. O. Snyder.) ($\pi\rho\acute{o}$, before; $\mu\acute{e}\lambda\acute{a}\zeta$, black.)

Chirostoma promelas, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 136, Guadalajara, Mexico.

Head $3\frac{1}{2}$; interorbital $2\frac{1}{2}$; length of body $3\frac{1}{2}$ times standard; scales 69, pressed together, straight, much compressed, lower; in front of mouth, edge of opercle little wider, a horizontal line lower for a short distance, projecting, terminating in a very large and prominent palatines band, groove crowded to the isthmus of the dorsal, a band extending from the snout, ending in a little slight depression, similar to that of another deeply forked dorsal, extended, olive; a single black pigment on the edge of the dorsal.

Our species and viscera probably.

Chirostoma species of the dorsal.

Known No. 6157, 1

(Named Mexico, in)

Chirostoma Mexico

1157 (d). *CHIROSTOMA DIAZI*, Jordan & Snyder.

Head $3\frac{1}{2}$; depth 5; depth of caudal peduncle $3\frac{1}{2}$ in head; eye $5\frac{1}{2}$; snout 2.; interorbital space $4\frac{1}{2}$; height of spinous dorsal $4\frac{1}{2}$; soft dorsal $2\frac{1}{2}$; anal $2\frac{1}{2}$; length of pectoral $1\frac{1}{2}$; ventral 3; caudal 1; D. V-11; A. 20; P. 15; scales 69-20, 22 between dorsals. Body long, deeper and more compressed than that of *C. humboldtiwum*; head large, its dorsal contour straight from tip of snout to occiput; viewed from above, the head is much compressed, the upper jaw is sharply pointed and included by the lower; interorbital space slightly convex; eye nearer tip of snout than to edge of opercle by a distance equal to diameter of pupil; preorbital area a little wider than diameter of orbit; mouth oblique, its cleft extending to a horizontal from lower edge of orbit; lips thickened posteriorly, the lower forming a fold across the upper at angle of mouth; lower jaw projecting, the teeth just passing the edge of the upper; maxillary extending to a vertical from anterior edge of orbit, its distal end angular; teeth large anteriorly, growing gradually smaller posteriorly, canine-like, sharp, projecting backward, not arranged in definite rows, none on vomer or palatines; vertebrae 41. Scales crenate, larger in region of lateral color band, growing smaller dorsally and ventrally, abruptly smaller and closely crowded together in a region anterior to the pectoral fin, extending from the isthmus to the occiput, also between the dorsal fins and along bases of dorsals, anal and caudal; inter-radial membranes of the latter with scales extending $\frac{2}{3}$ its length; head with scales except on jaws, upper part of snout, and on preorbital area; two anterior spines of dorsal longest, others a little shorter; first dorsal ray longest, others gradually shorter; anal similar to soft dorsal in shape, except that its base is much longer; insertion of anal on a perpendicular passing halfway between dorsals; caudal deeply forked, the lobes pointed; pectoral sharp, extending past base of ventrals a distance equal to diameter of orbit; edge of pectoral, when extended, straight. Body, during life, translucent; in alcohol, greenish olive; a silvery lateral band, in which the silver is not underlaid with black pigment, extending from axil to base of caudal; dorsal scales narrowly edged with dusky; upper part of eye dark. Jalisco, Mexico.

Our specimens are all from the market at Guadalajara. The gill-arches and viscera had been removed. The flesh is somewhat shrunken, which probably causes the teeth to appear more prominent than in life.

Chirostoma diazi may be easily distinguished from the other known species of the genus by the small scales crowded closely together between the dorsal fins. (Jordan & Snyder.)

Known only from the market at Guadalajara, Jalisco, Mexico. (Type, No. 6157, L. S. Jr. Univ. Mus. Coll. J. O. Snyder.)

(Named for Porfirio Diaz, the honored President of the Republic of Mexico, in recognition of his interest in the progress of science.)

Chirostoma diazi, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 137, Guadalajara, Mexico.

1157 (e). **CHIROSTOMA CRYSTALLINUM**, Jordan & Snyder.

Head $3\frac{1}{4}$; depth $4\frac{1}{2}$; depth of caudal peduncle $3\frac{1}{2}$ in head; eye $5\frac{1}{2}$; snout $2\frac{1}{2}$; interorbital space $4\frac{1}{2}$; height of spinous dorsal $1\frac{1}{2}$; soft dorsal $2\frac{1}{2}$; anal $2\frac{1}{2}$; length of pectoral $1\frac{1}{2}$; ventral $2\frac{1}{2}$; caudal $1\frac{1}{2}$; D. V-13; A. 24; P. 15; scales 56-58, 10 between dorsals. Body deepest above ventrals, widest at insertion of pectorals; head long and pointed, its upper contour straight; snout, viewed from above, sharply pointed, the lower jaw projecting beyond upper a distance equal to not more than half diameter of orbit; eye nearer tip of snout than to posterior edge of opercle a distance equal to diameter of pupil; mouth oblique, the cleft extending downward to a level with lower edge of pupil; lips thickened posteriorly, the lower folding over the upper at angle of mouth; teeth minute, in wide patches, not arranged in definite rows, on upper and lower jaws; no teeth on vomer or palatines; gill-rakers on first arch 27, slender, the length of longest equal to diameter of pupil; vertebrae 44. An indefinite lateral line extending along the body about 9 scales below first dorsal, the pores absent on many of the scales; scales crenate, largest along color-band, growing smaller above and below, much smaller and closely crowded on the body anterior to the pectoral fin, from the isthmus to the occiput; a few small scales at upper end of opercle, along bases of fins, and on the inter-radial membranes of caudal for half its length; scales between dorsals large and not crowded together; lower jaw and upper part of snout naked; first 3 spines of dorsal about the same length, reaching, when depressed, to $\frac{1}{2}$ their length of insertion of soft dorsal; first dorsal ray longest, others gradually shorter; anal inserted on a perpendicular passing through a point half way between origins of dorsals, base much longer than that of dorsal; first dorsal ray longest, others gradually shorter to middle of fin, all remaining rays of about the same height; caudal deeply forked, the tips rounded; pectoral pointed, extending a little past base of ventrals; tips of ventrals extending to vent. Body, during life, translucent, with a slightly bluish tinge; in alcohol, yellowish; upper scales with dusky edges; top of head dusky; upper part of eye dark; fins, except ventrals and anal, with a little dusky color; a silvery color band extending from upper part of pectoral base to caudal, narrower on caudal, widening at base of caudal.

C. crystallinum closely resembles *C. ocellatum* in general appearance. The much shorter lower jaw and the smaller eye of *C. crystallinum* are at once distinguishing characters. Guadalajara and Laguna de Chapala, Mexico. (Jordan & Snyder.)

Two specimens from the Guadalajara market are shrunken so that the orbit is larger, and the teeth more evident than in those taken at Ocotlan.

Known from Laguna de Chapala, near Ocotlan, Jalisco, Mexico, and the market at Guadalajara. (Type, No. 6158, L. S. Jr. Univ. Mus. Coll. J. O. Snyder.)

Chirostoma crystallinum, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 139, Laguna de Chapala, Mexico.

Head $3\frac{1}{2}$; interorbital $2\frac{1}{2}$; length of body $58-20$, 11 scales between dorsals; eye slightly larger than diameter of orbit; tip of snout slightly raised; upper part of opercle with a horizontal line across it, from anterior to posterior edge; inner row of scales on opercle none on ventral side, along lateral line many of scales much smaller and more numerous, similar scales becoming larger on basal half of body; dorsal rays 44, dorsal rays 27, ratio of anal to dorsal rays 1.33, ratio of origin of dorsal to origin of anal 1.33, one-third of pectoral plus diameter of eye equal to width of silvery band on base of dorsal, dark coloration of body extending to top of head.

C. lemniscatum, Jordan & Snyder, 1890, Pl. 1, Fig. 1, large teeth, larger eye.

Our species distinguished by gill-arches 27, not 28, shrunken, not elongated, Jalisco, Mexico.

Chirostoma,

Mexico.

Head $3\frac{1}{2}$; interorbital $2\frac{1}{2}$; length of body $54-19$, 11 scales between dorsals; eye slightly larger than diameter of orbit; tip of snout slightly raised; upper part of opercle with a horizontal line across it, from anterior to posterior edge; inner row of scales on opercle none on ventral side, along lateral line many of scales much smaller and more numerous, similar scales becoming larger on basal half of body; dorsal rays 44, dorsal rays 27, ratio of anal to dorsal rays 1.33, ratio of origin of dorsal to origin of anal 1.33, one-third of pectoral plus diameter of eye equal to width of silvery band on base of dorsal, dark coloration of body extending to top of head.

By

1157 (d). *CHIROSTOMA LERME*, Jordan & Snyder.

(Head 3); depth 5 $\frac{1}{2}$; depth of caudal peduncle 3 $\frac{1}{2}$ in head; eye 5 $\frac{1}{2}$; snout 2 $\frac{1}{2}$; interorbital space 5 $\frac{1}{2}$; height of spinous dorsal 4 $\frac{1}{2}$; soft dorsal 2 $\frac{1}{2}$; anal 2; length of pectoral 1 $\frac{1}{2}$; ventrals 3; caudal 1 $\frac{1}{2}$; D. IV-11; A. 20; scales 38-40, 11 between dorsals. Body slender, deepest part in region of ventrals; caudal peduncle narrow; snout long and pointed; lower jaw slightly projecting, but not enough to include the upper; eye large, nearer tip of snout than to posterior edge of opercle, a distance equal to diameter of orbit, or to width of preorbital space; cleft of mouth extending to a horizontal through lower edge of orbit; lower lip folded over the upper at their union; maxillary extending posteriorly almost to a perpendicular from anterior edge of orbit, its distal end angular; teeth large and strong, curved backward and inward, arranged in 2 definite rows, those of the inner row of the upper jaw and of the outer row of the lower jaw larger, none on vomer or palatines; vertebrae 44. An indefinite lateral line extending along body about 10 scales below first dorsal, the pores absent on many of the scales; scales crenate, largest along the lateral color-band, much smaller between occiput and first dorsal, those immediately posterior to occiput, minute and very closely crowded; a narrow edging of similar scales along the gill-openings, extending ventrally to the isthmus; scales between dorsals not much reduced in size or crowded together; basal half of inter-radial membranes of caudal with scales; first 2 spines of dorsal longest, third shorter, fourth about $\frac{2}{3}$ as long as the first; first dorsal ray longest, others gradually shorter, edge of fin straight; insertion of anal on a perpendicular passing through a point half way between origins of dorsals, first ray longest, others successively shorter, last ray one-third the length of first; caudal deeply forked, its tips rather pointed; pectoral pointed, extending beyond origin of ventrals a distance equal to diameter of pupil; ventrals reaching vent. Color in alcohol, light olive, a silvery lateral band equal to a scale in width extending from upper part of base of pectoral to base of caudal; edges of upper scales dusky; a little dark color on dorsals, caudal, and pectoral; upper and lower jaws and top of head with minute dark dots; upper part of eye dark.

C. lerme closely resembles *C. crystallinum*. It differs markedly in having large teeth, which are arranged in two rows, a shorter lower jaw and a larger eye. Jalisco, Mexico. (Jordan & Snyder.)

Our specimens of *C. lerme* are all from the market of Guadalajara. The gill-arches and viscera had been removed and the bodies were somewhat shrunken. The specimens are said to have come from Laguna de Chapala, Jalisco, Mexico. (Type, No. 6159, L. S. Jr. Univ. Mus. Coll. J. O. Snyder.)

Chiostoma lerme, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 140, Guadalajara, Mexico.

1157 (g). *CHIROSTOMA OCOTLANE*, Jordan & Snyder.

(Head 3 $\frac{1}{2}$; depth 4 $\frac{1}{2}$; depth of caudal peduncle 3 $\frac{1}{2}$ in head; eye 4 $\frac{1}{2}$; snout 2 $\frac{1}{2}$; interorbital space 5; height of spinous dorsal 3 $\frac{1}{2}$; soft dorsal 2 $\frac{1}{2}$; anal 2; length of pectoral 1 $\frac{1}{2}$; ventral 2 $\frac{1}{2}$; caudal 1 $\frac{1}{2}$; D. V-12; A. 20; scales 31-33, 11 between dorsals. Body long, rather slender, deepest above

Bull. No. 47, pt. 4—IX

ventrals; head long, pointed, its dorsal contour straight from tip of snout to a point a little posterior to the eye, where it curves upward; interorbital space flat; eye high up, nearer tip of snout than edge of opercle by a distance about equal to diameter of pupil; width of preorbital space equal to diameter of pupil; mouth oblique, the cleft extending downward to a point opposite lower edge of pupil; lips growing more fleshy posteriorly, the lower forming a fold across the upper at their junction; distal end of maxillary angular, extending almost to a vertical from anterior edge of orbit; lower jaw very long, projecting beyond upper a distance equal to diameter of pupil; viewed from above, both jaws are a little more pointed than are those of *C. estor* or of *C. humboldtianum*. Teeth on jaws in bands, minute, projecting backward; no teeth on vomer or palatines; gill-rakers slender, close together, length of longest equal to diameter of pupil; vertebræ 44. A rather indefinite lateral line extending along body, 10 scales below the first dorsal, the pores absent on many of the scales; scales crenate, growing smaller dorsally, larger ventrally, those on posterior part of occiput, on nape, on region anterior to pectorals—above and below, and on base of caudal, very small and crowded together, those in the region of dorsals not reduced in size or crowded except at insertion of fins, where there are 4 very small ones; cheek with 5 rows; lower jaw and upper part of snout naked; basal half of interradial membranes of caudal with minute scales; first 3 spines of dorsal highest, extending when depressed within half their length of origin of soft dorsal; first dorsal ray preceded by a shorter, simple, closely attached one; other rays successively shorter than first; edge of fin slightly concave; anal similar to soft dorsal in shape, its basis 1½ times as long as that of soft dorsal; attachment of first ray under a point half way between insertions of dorsals; caudal deeply forked, the lobes equal; pectoral pointed when depressed, upper rays longest, extending to a vertical half way between insertions of first dorsal and ventrals. Body, during life, almost translucent, with a bluish tinge of color; in alcohol, the color is a light olive yellow; a silvery lateral band extending from axil to base of caudal, the band wider and brighter in color between dorsal and anal, growing narrow on caudal peduncle and then widening again at its posterior end. Head and body above, and the lower jaw, dusky; upper part of eye dark; edge of scales above lateral band with small black dots; caudal somewhat dusky on its basal third; other fins with little or no dark color.

Chirostoma ocellane is easily distinguished from all other known species of the genus by its excessively long lower jaw. Except the jaw and somewhat larger eye it resembles *C. estor* in general appearance. Laguna de Chapala, Mexico. (Jordan & Snyder.)

Known only from Laguna de Chapala, near Ocotlan, Mexico. (Type, No. 6160, L. S. Jr. Univ. Mus., Coll. J. O. Snyder.)

Chirostoma ocellane, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 141, Laguna de Chapala, near Ocotlan, Mexico.

We ha
small an
occiput
crowded
and num
almost a

Some c
length o

Length
head 31½
width of
distance
dorsal 12
distance
from ana
tance fro
P. 11; se

The ty
Steindach
species w
localities
large vom
album as p

Page 8

Neomugil, V

General
beyond an
those of ou
small, but
siphonal, v
like; a la
distinct.

This ge
rated, wit
extends no

Head 3;
l, 5; scal
deeply cle
dorsal at 1

1154. CHIROSTOMA ESTOR, Jordan.

We have examined the type specimen of *Chirostoma estor*. The scales are small and closely crowded on the region anterior to the pectoral from the occiput to the isthmus; they are not much reduced in size nor closely crowded together between the dorsals. The teeth on the jaws are large and numerous, not arranged in definite rows; there are 3 vomerine teeth almost as large as those on the jaws.

Some exact measurements of the type, expressed in hundredths of the length of the body, are here given:

Length of body in millimeters 212; depth of caudal peduncle $7\frac{1}{2}$; length of head $31\frac{1}{2}$; distance from snout to occiput 23; tip of lower jaw to occiput 24; width of interorbital space 7; length of snout 12; diameter of orbit $5\frac{1}{2}$; distance from snout to spinous dorsal $56\frac{1}{2}$; insertion of spinous to soft dorsal $12\frac{1}{2}$; height of longest dorsal spines 7; of longest dorsal rays 11; distance from snout to anal $62\frac{1}{2}$; height of longest anal rays 12; distance from anal to caudal 22; length of caudal fin $18\frac{1}{2}$; of pectoral fin 17; distance from snout to ventral $45\frac{1}{2}$; length of ventral fin $10\frac{1}{2}$; D. V-12; A. 18; P. 11; scales 72-19, 9 between dorsals.

1154 (a). CHIROSTOMA ALBUM (Steindachner).

The type of *Chirostoma estor* agrees externally almost perfectly with Steindachner's account of *Chirostoma album* from Lake Pátzenaro, a species which he later places in the synonymy of *C. estor*. But the type localities are widely separated, and Steindachner found no trace of the large vomerine teeth so conspicuous in *C. estor*. We therefore regard *C. album* as probably a valid species, not identical with *C. estor*.

Page 821. After *Joturus picardi*, Poey, add:

368 (a). NEOMUGIL, Vaillant.

Neomugil, VAILLANT, Bull. Soc. Philom., IV, 1893-1894 (June, 1894), 72 (*diguetti*).

General aspect of *Mugil*. Mouth widely cleft, maxillary extending beyond anterior edge of orbit. Fine teeth in many series in each jaw, these of outer series on intermaxillary most developed, all conical, simple, small, but distinct; villiform teeth on vomer and palatines. Stomach siphonal walls membranous, very thin in pyloric region, hence not gizzard-like; a large pyloric ceca and a second smaller one; pseudobranchiae distinct.

This genus resembles *Joturus*, which has the teeth flattened and serrated, with only 1 series on the intermaxillary. The maxillary in *Joturus* extends not so far back. (Vaillant.) (*νέος*, new; *Mugil*.)

1198 (a). NEOMUGIL DIGUETI, Vaillant.

Head 3; depth 4; eye 6; interorbital 2.75. B. 6; D. IV-I, 8; A. I, 9; V. I, 5; scales 34-41; caudal fin 4. Maxillary reaching pupil; mouth deeply cleft; upper jaw the longer; no adipose eyelid. Origin of first dorsal at middle of body, the first spine 2 in depth of body; origin of anal

a little in front of second dorsal. Whole head, except preocular region, scaled like the body, the scales ctenoid.

Known from 20 examples from 99 to 194 mm. long, from Lower California, south of La Paz.

(Named for the collector of the type, M. Léon Dignet.)

Neomugil digueti, VAILLANT, Bull. Soc. Philom., IV, 1883-1894 (June, 1894), 73, torrent in the Sierra de las Cacachilas de Santa Cruz, Lower California.

Page 903. In the description of *Seriola lalandi* for "A. I, 27" read "A. I, 21."

Page 962. *Centrolophidae* can not be maintained as a distinct family. It belongs with the *Stromateidae*.

Page 1038. After *Hadropterus scirurus serrula*, insert:

HADROPTERUS MAXINKUCKIENSIS, EVERMANN

Head 3.75; depth 6; eye 4; snout 4.2; maxillary 3.25; mandible 2.75; interorbital 6; pectoral 1.25; ventral 1.3; D. XIV, 13; A. II, 8; scales 7-61-10.

Body rather long, slender, and subterete; caudal peduncle somewhat compressed, its least width one-half its least depth; head rather long, snout pointed; mouth moderately large, oblique, maxillary reaching past anterior part of eye, lower jaw included; eye rather large, slightly above axis of body; interorbital moderately wide, nearly flat; gill-membranes free from each other and from the isthmus; opercle with a rather long flap and stout spines; premaxillaries not protractile; fins rather large: distance from origin of spinous dorsal to tip of snout slightly greater than base of spinous dorsal or nearly twice base of soft dorsal; longest dorsal spine 2.25 in head; soft dorsal higher than spinous portion, 2 in head, the free edge gently curved; origin of anal under that of soft dorsal, its base 2.1 in head; caudal rounded or slightly emarginate.

Scales firm and strongly ctenoid, lateral line complete and straight, beginning over opercular spine; top of head and an oblong area on nape naked; space in front of spinous dorsal with small embedded scales; opercle with about 7 rows of scales; cheek with a few small embedded scales, breast naked, except 2 or 3 partially embedded scales on median line; one large scale between ventrals; belly naked anteriorly, but with about 10 enlarged, stellate scales posteriorly; space between ventrals broad, equal to width of base of ventral; preopercle smooth.

Color essentially the same as in *H. scirurus*; mottled and vermiculated with light and dark brown or blackish, the middle line of back with about 9 large, roundish, dark, confluent areas, each surrounded by a wavy, whitish line; middle of side with about 7 large, confluent dark spots, the anterior 2 largest and longest, the third small, the fourth large, and the remaining 3 progressively smaller; under parts yellowish white; top of head dark; a narrow whitish line around upper posterior part of orbit; a broad black line downward from eye; upper part of preopercle and nearly whole opercle dark, each dusky on lower part; cheek dusky with fine, dark specks; an irregular pale area at anterior end of lateral

line; sp
the o
dark; so
series of
with wh
ish, wit
lowed by

This sp
ing the
of orbit.
preoper

On t
bee Creel
from the
mann &
Hadropt

Aubee:

Page 10

Head 3,
X-10; A. I

Body ra
rather sh
oblique, lo
above axis
from the i
distinctly, bu
spinous do
outline of
head; long
stronger th
short, abor
rounded, 1
incomplete
usually na
usually ab
naked; 1-4
scaled, 1-9
flap mode
process.

Color in
cal, dark 1
another se
side anteri
head dark
line downw
dark borde

line; spinous dorsal ashy, the first 3 spines black on the middle portion, the other spines dark, but not so distinctly so; tips of last few spines dark; soft dorsal light brownish or grayish, crossed near the base by a series of dark spots and above by 2 series of whitish spots; caudal spotted with white and brown; anal white, dusky with brownish; ventrals whitish, with fine, dark dustings; pectoral whitish, yellowish at base, followed by alternating series of dusky and whitish spots.

This species is related to *H. scirurus*, from which it differs chiefly in having the dorsal fins united, in having the maxillary reaching beyond front of orbit, in the larger scales, the free gill-membranes, and in the smooth preopercle.

On, the type known, an example, 3.5 inches long, taken in Aubeenaukee Creek, the eastern inlet of Lake Maxinkuckee, Indiana, about $\frac{1}{2}$ mile from the lake, August 4, 1899. Type No. 49378, U. S. N. M. Coll. Evermann & Sevall.

Hadropterus maxinkuckensis, EVERMANN, Rept. U. S. Fish Com. 1899 (1900), 366, with plate.
Aubeenaukee Creek, Lake Maxinkuckee, Indiana.

Page 1084. Before *Etheostoma jessiae*, insert:

ETHEOSTOMA AUBEEAUBEI, Evermann.

Head 3.6; depth 5.5; eye 4.5; snout 4.5; maxillary 3; interorbital 5; D. X-10; A. II, 7; scales 4-55 to 58-7, 8 to 24 pores.

Body rather elongate, not much compressed except posteriorly; head rather short; snout short, somewhat decurved; mouth moderate, slightly oblique, lower jaw included, maxillary reaching front of eye; eye small, above axis of body; premaxillaries not protractile; gill-membranes free from the isthmus and each other. Fins not large, the dorsals usually distinctly, but narrowly separated, sometimes scarcely separate; origin of spinous dorsal one-third distance from tip of snout to base of caudal; outline of spinous dorsal gently rounded, the longest spine about 3 in head; longest dorsal ray about 2; first anal spine longer, and slightly stronger than second, 3 to 3.5 in head; longest anal ray about 2; pectoral short, about 1.3 in head; ventrals close together, about 2 in head; caudal rounded, 1.5 in head. Scales rather small, rough-ctenoid; lateral line incomplete, usually developed on only 8 to 24 scales at anterior end; cheek usually naked or with a few small, more or less embedded scales; opercle usually about half scaled, sometimes with but few scales; breast always naked; body with ordinary scales; nape usually densely and regularly scaled, some scales sometimes embedded; preopercle entire; opercular flap moderate, broad; opercular spine rather small; no humeral spot or process.

Color in alcohol, greenish brown above; side with about 12 or 13 vertical, dark blotches, separated by pale orange red areas of similar size; another series of similar but smaller orange blotches along lower part of side anterior to anal fin; under parts whitish; caudal peduncle grayish; head dark above; opercle and cheek dark, with greenish shade; a dark line downward from eye; snout grayish; spinous dorsal with a narrow dark border, below which is a broad orange band, then a broad but irregu-

lar dark band along base of fin; soft dorsal and caudal barred with white and grayish, the latter in spots on the rays; anal and ventrals without markings; pectoral somewhat dusky.

This species is close to *E. iowae*, from which it seems to differ in the almost naked cheek, the less complete scaling of the opercles, the somewhat longer maxillary, more oblique mouth, closer approximation of the dorsal fins, and the coloration. Lake Maxinkuckee, Indiana.

Many examples, each about 2 inches long, taken in Aubeenaubee Creek, the east inlet of Lake Maxinkuckee, August 4, and on other days in August and September, 1899. *E. iowae* is a common species in the lake, but does not occur in the inlet, nor were we able to find any specimens of *E. aubeenaubae* anywhere except in this small creek. Type No. 49379, U.S.N.M. Coll. Evermann & Sevall.

Etheostoma aubeenaubae, EVERMANN, Rept. U. S. Fish Com. 1899 (1900), 307, with plate, Aubeenaubee Creek, Lake Maxinkuckee, Indiana.

Page 1212. Before *Prionodes fasciatus*, Jenyns, insert:

1597 (a). PRIONODES BALDWINI, Evermann & Marsh.

Head 2.5; depth 3.2; eye 4; snout 4.6; maxillary 2.4; mandible 2; interorbital 7; D. x, 12; A. III, 7; pectoral 1.4; ventral 1.3; caudal 1.7; scales 44-42-12.

Body elongate, moderately compressed, not elevated, covered with ctenoid scales; dorsal and ventral outlines alike; head moderate, pointed, naked above and below; eye large, greater than length of snout, high in position; mouth terminal, slightly oblique, the maxillary reaching middle of eye or somewhat beyond; gill-rakers short, 6 developed on lower limb; teeth small, conical, and sharp, on vomer and palatines and in several series in each jaw, with weak canines in front and a few canine-like teeth on middle of side of lower jaw; cheek with about 7 rows of scales; preopercle finely serrate; opercle ending in 3 sharp, flat spines, the middle one largest, a membranous pointed flap projecting beyond; fins all naked, the dorsal continuous, with a slight emargination, the spines slender and pungent, the first 4 or 5 graduated, the rest subequal, 3.2 in head, lower than the soft rays which are contained about 2.5 in head; anal fin short, the second spine longest and strongest, 3 in head, the soft part high, the fifth or sixth ray longest, reaching almost to front of anal, 2 in head; ventral with second ray produced, reaching vent; caudal truncate, or with middle rays very slightly shorter, making the margin slightly concave.

Color in life: Dorsal half of head and trunk and all of caudal peduncle scarlet, ventral portion pale blue, almost white; a yellow longitudinal band nearly as wide as pupil from preopercular margin straight across opercle and along body to lateral line under last dorsal rays; 4 quadrate or oblong black blotches just under this band, the first about under middle of spinous dorsal, second under last spines, third under first rays, fourth under last rays; from each of the first three of these blotches a square, well-defined yellow shade extends downward to belly or base of anal, a similar one from base of pectoral to ventral; 4 smaller black blotches at base of caudal, two others, somewhat larger than the last, just in front of

them on base of dorsal, the other middle under last peduncle by a simi on top of brown spe very defin black blo with their of head us with dark tively fair ing the pr color bord out with 1 is a very ray some pectoral a

In spirit and addit and upper vertical ba round or q longitudin dorsal dus toral, and

Puerto P dredged air tangle, off 15 and 16 f Hawk State light-house expedition artist of the

Prionodes bal
353, off Cu

Page 122

Rhegma, Gmel

Allied, bu large, cteno ian physiog *Rypticine*.

taem on caudal peduncle; a row of 9 round black spots on each side at base of dorsal fin, the first one smallest, opposite membrane of first spine, the other 8 separated somewhat obscurely into pairs, the first pair under middle spines, second under last spines, third under first rays, fourth under last rays; 2 or 3 very small black dots on upper edge of caudal peduncle; 2 or 3 more in front of dorsal on median line, each accompanied by a similar one on either side; in some specimens a few scattered ones on top of head behind eyes, sometimes regularly arranged; a few dark-brown spots behind eye; various dark markings on side of head, without very definite pattern, but usually 2 oblique stripes on cheek, a heavy black blotch on interopercle and 2 on the ramus of the mandible, which, with their fellows of the other side, make distinct crossbars on lower side of head usually extending across maxillary; chin and lower part of opercle with dark spots; lateral line white, with a few broken spots, comparatively faint, just below it; iris red, with an inner ring of white surrounding the pupil; spinous dorsal pale, the edge of the membrane black, this color bordered below with faint yellow; soft dorsal pale, spotted throughout with light orange, with a marginal band of the same, outside of which is a very narrow pale-blue edge; ventral very pale blue, the produced ray somewhat yellow; anal pale blue with some light orange on last rays; pectoral and caudal uniform pale reddish, unmarked.

In spirits all the red and yellow markings disappear, the dark persists, and additional markings are brought out as follows: Along the anterior and upper part of trunk and crossing the lateral line are dark-brown vertical bars, diffuse and running together, or separated and broken into round or quadrate blotches; in the middle part of the course of the yellow longitudinal band appears a row of very small black points; spots on soft dorsal dusky; dark mottlings on caudal; upper and lower base of pectoral, and sometimes axil, dusky.

Puerto Rico. A beautiful and strongly marked species; 2 specimens dredged and 33 others, ranging in size from 0.55 to 2 inches, caught in the tangle, off Culebra and Vieques islands, from coral bottom, in depths of 15 and 16 fathoms; the type, 2 inches long, taken in the tangle at Fish Hawk Station 6093, off Culebra Island, 5.25 miles southwest of Culebritas light-house, February 8, 1899, in 15 fathoms, by the U. S. Fish Commission expedition to Puerto Rico. ("Named for Mr. Albertus H. Baldwin, the artist of the U. S. Fish Commission expedition to Puerto Rico, in recognition of his excellent drawings and paintings of American fishes.")

Priacanthus hablitzini, EVERMANN & MARSH, Report U. S. Fish Comm. 1899 (December 19, 1899), 353, off Culebra Island, P. R. (Type, No. 49361, U.S.N.M. Coll. Evermann & Marsh.)

Page 1229. After *Gramma loreto*, Poey, insert:

518 (a). **RHEGMA**, Gilbert, new genus.

Rhegma, GILBERT, new genus of Serranidae (*thaumasium*).

Allied, but not closely, to *Rypticus*, *Grammistes* and *Gramma*, having the large, ctenoid scales and interrupted lateral line of *Gramma* and the peculiar physiognomy, attachment of gill-membranes, and fin structure of the *Rypticus*.

Scales of moderate size, thin, not embedded, minutely ciliated; lateral lines 2, the upper near base of dorsal, ceasing under origin of posterior fifth of soft dorsal, the lower line beginning slightly in front of the end of the upper line, running along middle of caudal peduncle; the tubes very short, borne on much smaller intercalated scales, and not forming a continuous line; head largely scaled, the snout and jaws naked; gill-membranes united anteriorly, forming a narrow free fold across the isthmus; branchiostegals 7; pseudobranchiae well developed; a wide slit behind last gill-arch; gill-rakers short, broadly triangular, strongly toothed. Upper margin of opercle, above its angle, wholly attached by membrane to the shoulder girdle, as in the *Rypticinae*. Mouth large, protractile, the lower jaw protruding, the maxillary broadly exposed, with a narrow supplemental bone along its upper edge. Teeth all villiform, in broad bands on jaws, vomer and palatines, the inner teeth on jaws slightly longer than the others and depressible. Tongue smooth. Large mucous pores on under side of mandible, and slit-like pores present on edge of preorbital and around front of eye. Anterior nostril near edge of preorbital, provided with a short tube; posterior nostril without tube or raised rim, immediately in front of eye; a short free triangular flap on upper edge of each orbit. Upper portion of preopercle with a single strong plectroid spine, directed backward and downward; bones of the head otherwise unarmed, the preorbital and preopercle with entire edges, the opercle without spines or ridges. Ventrals small and anterior in position, as in the *Rypticinae*, consisting each of one strong spine, and five branched rays, their base being in front of base of pectorals; no enlarged scale behind base of ventrals; vertical fins low, with rounded lobes, their basal portions well scaled; dorsal with 7 low strong spines and 22 profusely branched rays; anal with 3 spines and 18 rays. One species, apparently the type of a distinct subfamily, *Rhegmatisinae*.

(*ρῆγμα*, fracture; referring to the interrupted lateral line.)

1615 (a.) RHEGMA THAUMASIUM, Gilbert, new species.

Head 2 $\frac{1}{2}$ in length; depth 3; eye 5 in head; D. VII, 22; A. III, 18. Scales 45 in a longitudinal series along middle of side. Body elongate, moderately compressed, with very short, deep, caudal peduncle; anterior profile strongly arched, slightly depressed above orbits; interorbital space very narrow, convex, its width two-thirds diameter of orbit. Mouth large, slightly oblique; lower jaw the longer, its tip entering the profile. Dorsal spines low, strong, increasing backward, the last spine a trifle longer than diameter of orbit; anal spines short and strong, the middle spine longest. Lateral line curved strongly upward from its origin to below third dorsal spine, thence running parallel with the back to below middle of second dorsal, from which point it gradually approaches the base of the dorsal, where it terminates under the fifth ray from the last; along its anterior course it is separated from base of dorsal by from 4 to 6 scales (in oblique series); scales minutely ctenoid except on head, breast, and belly, on nape under anterior dorsal spines, and on base of pectoral; top of head scaled forward to interorbital space, the anterior scales here,

as well
outlines
the fo

Total len
Length t
Greatest
Least de
Length o
Distance
Length o
Diameter
Interorbital
Tip of sn
Length o
Tip of sn
Tip of sn
Length o
First anal
Second an
Third ana
First soft
Longest s
First dors
Last dorsa
Longest d
Caudal ...

Color, u
opercular
with whit

One spec
(*Gymnac*
Rhegma tha
Univ. M

Page 13

Head 3 $\frac{1}{2}$
A. III, 12;
Profile v
reaching b
maxillary r
chin; preo
gill rakers
creasing te
creasing to
covered wi

as well as those on cheek, much reduced in size, embedded, so that their outlines can not be distinguished.

The following measurements of the type specimen are given:

	m.m.	One-hun-dreths of length to base of caudal.
Total length.....	85	
Length to base of caudal	70	
Greatest depth.....		33
Least depth caudal peduncle.....		15½
Length of caudal peduncle.....		8
Distance from tip of upper jaw to end of opercular flap.....		37
Length of snout.....		6
Diameter of eye.....		7½
Interorbital width		3½
Tip of snout to end of maxillary.....		19
Length of pectoral.....		27
Tip of snout to base of upper pectoral ray		35
Tip of snout to base of ventrals		31
Length of ventrals.....		13
First anal spine.....		3½
Second anal spine.....		6
Third anal spine.....		4½
First soft anal ray		8½
Longest soft anal ray		15
First dorsal spine.....		4
Last dorsal spine.....		9
Longest dorsal ray		15
Caudal		23

Color, nearly uniform warm brown on head, body, and fins; a dusky opercular blotch; soft dorsal, anal and caudal only narrowly margined with white.

One specimen known, from Panama.

(*θαυμασίος*, wonderful.)

Rhegma thaumasium, GILBERT MS., Panama (Coll. C. H. Gilbert. Type, No. 5978, L. S. Jr. Univ. Mus.)

Page 1300. After *Hemulon scudderii*, Gill, insert the following:

1667 (a). **HEMULON HELEX.E**, Boulenger.

Head 3½; depth 2½; eye 3½ in head; interorbital width 3½. D. XIII, 16; A. III, 12; scales 11-87-20, 67 pores.

Profile very much arched from snout to origin of dorsal; muzzle not reaching beyond lower jaw, little shorter than eye; mouth little oblique; maxillary reaching anterior ¼ of eye; teeth very small; 2 very small pores in chin; preopercle very feebly serrate; head scaled, except snout and chin; gill rakers very short, 15 on lower arm of arch; dorsal spines feeble, increasing to the fourth, which almost equals ½ length of head, then decreasing to the last, which is contained 4½ times in head; soft dorsal low, covered with scales; pectoral fulciform, longer than head, 1½ times ventral;

anal spines small, increasing to the third, which is 5 in head; soft anal low and scaly; caudal almost completely scaled, deeply emarginate, the median rays less than half the external ones; caudal peduncle $\frac{1}{2}$ times longer than deep; scales above lateral line in very oblique series. Grayish above, with oblique brown lines, somewhat undulating; silvery below. Length 22 cm. One specimen from Bay of Santa Elena, Ecuador. (Boulenger.) (Named for the type locality.)

Haemulon heleneae, BOULENGER, Bollettino dei Mus. di Zool. ed Anat. Comp. della Univ. di Torino, Vol. XIV, No. 335, 3, Feb. 15, 1899, Bay of St. Elena, Ecuador. (Coll. Dr. Enrico Festa.)

Page 1334. After *Pomadasys ramosus* (Poey), insert the following:

1706 (a). **POMADASIS LABRACIFORME** (Boulenger).

Head 3; depth $3\frac{1}{2}$; eye 4; snout slightly longer than mandible, by $\frac{1}{2}$ diameter of eye, which is little more than interorbital width; maxillary reaching anterior $\frac{1}{2}$ of eye; D. XIII, 12; A. III, 7; scales 8 or 9-68-22; teeth very small; 2 pores in chin; preopercle strongly serrate, strongest at angle; head scaled, except snout and lips; gill-rakers 13 on lower arm of first arch, the longest equaling $\frac{1}{2}$ length of gill-filaments. Dorsal spine strong, the fifth longest, 2 in head, from which they decrease to next to last, which is $4\frac{1}{2}$ in head, and a little shorter than last; soft dorsal scaled at base, its longest rays a little shorter than longest spine; pectoral $1\frac{1}{2}$ in head, scarcely shorter than ventrals; second anal spine very strong, $3\frac{1}{2}$ in head, third 2 in head; longest anal rays shorter than anal spine; caudal peduncle longer than deep; scales arranged in series scarcely oblique above the lateral line, which has 54 pores. Color uniform silvery. Length 17 cm. One specimen known, from the Bay of St. Elena, Ecuador. (Boulenger.) (*Labrus*; *forma*, *form*.)

Pristipoma labraciforme, BOULENGER, Bollettino dei Mus. di Zool. ed Anat. Comp. della Univ. di Torino, Vol. XIV, No. 335, 3, Feb. 15, 1899, Bay of Santa Elena, Ecuador. (Coll. Dr. Enrico Festa.)

Page 1351. After *Calamus proridens*, Jordan & Gilbert, insert:

1721 (a). **CALAMUS KENDALLI**, Evermann & Marsh.

(PLUMA.)

Head 3.1; depth 2.1; eye 3.5; snout 1.5; maxillary 2.4; interorbital 3.5; preorbital 2.1; D. XII, 12; A. III, 10; pectoral 1; ventral 1.8; caudal 1.3; scales 7-53-16.

Body deep, back strongly elevated, more so than in *C. bajonado*, but less than in *C. calamus* or *C. proridens*, the anterior profile a nearly regular curve, lacking the abrupt unshel elevation of those species; eye large, larger than in *C. proridens*; 7 or 8 rows of scales on cheek; teeth about as in *C. proridens*; molars in 2 or more rows on sides, those of inner row much the largest, those in front becoming more numerous and merging into caridiform teeth, the most anterior of which, in each jaw, are somewhat enlarged; in front of upper jaw are 2 much enlarged antrorse canines, curved slightly upward; highest dorsal spine 2.7 in head, second anal spine 4.6.

Color in
the rows
below; so
numerous
which the
Puerto
Mayagüez
Mayagüez
sion expe
dall, assis
Calamus ke
Mayagt

Page 1
dentalis, V
The syn
Nebri's oceo
Panama
Nebri's zet
America
Coll. D

Page 14

Head 3
D. XI, 21
pores. Sn
scarcely ob
 $\frac{1}{2}$ or $\frac{1}{3}$ of
directed d
arch; thir
first anal s
est soft ray
Grayish ab
below. C
tinguished
month, and
from the p
Corrina mia
Univ. di
Enrico F

Page 15

Head 2
 $2\frac{1}{2}$; inter
 $2\frac{1}{2}$, ray 2;
8; P. 14; s
depth a di

Color in spirits: Silvery, side with bluish longitudinal lines following the rows of scales, plainest above; a pale-blue line bordering the orbit below; some blue lines on preorbital, not evidently reticulated and not as numerous as in *C. proridens*; iris yellow; otherwise as in *C. proridens*, to which this species is very close.

Puerto Rico; known from the type, a specimen 10.5 inches long, from Mayagüez, January 20, 1899, and 2 cotypes, each 8.5 inches long, one from Mayagüez, the other from Arroyo, all collected by the U. S. Fish Commission expedition to Puerto Rico. (Named for Dr. William Converse Kendall, assistant, U. S. Fish Commission.)

Calamus kendalli, EVERMANN & MARSH, Rept. U. S. Fish Com. 1899 (Dec. 19, 1899), 354.
Mayagüez, Puerto Rico. (Type, No. 49362, U.S.N.M. Coll. Evermann & Marsh.)

Page 1417. For *Nebris zetus*, Jordan & Starks, substitute *Nebris occidentalis*, Vaillant.

The synonymy of this species is as follows:

Nebris occidentalis, VAILLANT, Notes from the Leyden Museum, XX, 20, October, 1897,
Panama; after description by Jordan & Eigenmann.

Nebris zetus, JORDAN & STARKS, in JORDAN & EVERMANN, Fishes North & Middle
America, Part II, 1417, October 3, 1898, Panama. (Type, No. 433, L. S. Jr. Univ. Mus.
Coll. Dr. C. H. Gilbert.)

Page 1439. After *Bairdiella chrysoleuca* (Günther), insert the following:

1820 (a). **BAIRDIELLA MIACANTHA** (Boulenger).

Head $3\frac{1}{2}$ to $3\frac{1}{2}$; depth $3\frac{1}{2}$ to $3\frac{1}{2}$; eye $4\frac{1}{2}$ to $4\frac{1}{2}$, 1 to $1\frac{1}{2}$ in interorbital width.
D. XI, 21 or 22; A. II, 7 or 8; scales 5 or 6-45 to 49-51 or 12, 50 to 55
pores. Snout round, scarcely longer than eye; mouth inferior, compressed,
scarcely oblique, with simple subequal teeth, the gape reaching anterior
 $\frac{1}{3}$ or $\frac{1}{2}$ of eye; 2 very strong spines at angle of preopercle, the lower
directed downward; gill-rakers very short, 10 or 11 on lower arm of first
arch; third dorsal spine longest, $\frac{3}{2}$ to $\frac{2}{3}$ length of head; pectoral $1\frac{1}{2}$ in head;
first anal spine very small, the second small, $2\frac{1}{2}$ to 3 in head, or $1\frac{1}{2}$ in long-
est soft ray; caudal acuminate; caudal peduncle a little longer than deep.
Grayish above, with faint brown stripes following series of scales; silvery
below. Close to *B. chrysoleuca*, Günther, from which it is sufficiently dis-
tinguished by the less development of the second anal spine, the smaller
mouth, and the feebler dentition. Length 21 cm. Three specimens known
from the port of Guayaquil. (Boulenger.)

Corina miacanthus, BOULENGER, Bollettino dei Mus. di Zool. ed Anat. Comp. della
Univ. di Torino, Vol. xiv, No. 335, 5, Feb. 15, 1899, port of Guayaqui. (Coll. Dr.
Enrico Festa.)

Page 1526. After *Cichlasoma centrarchus* (Gill & Bransford), insert:

1923 (a). **CICHLASOMA STEINDACHNERI**, Jordan & Snyder.

Head $2\frac{1}{2}$; depth 3; depth of caudal peduncle $7\frac{1}{2}$; eye $3\frac{1}{2}$ in head; snout
 $2\frac{1}{2}$; interorbital space 5; longest dorsal spine 4, ray 2; longest anal spine
 $2\frac{1}{2}$, ray 2; length of pectorals $1\frac{1}{2}$; ventrals $1\frac{1}{2}$; caudal $1\frac{1}{2}$; D. XVI, 10; A. V,
8; P. 14; scales 26-14, 7 on caudal peduncle. Length of head exceeding its
depth a distance equal to diameter of orbit; body elongate, deepest above

ventrals; curve of dorsal outline interrupted by a slight elevation above eye and a rather rapid descent at base of soft dorsal; ventral outline less curved than dorsal; interorbital space convex; orbit somewhat elongate laterally, located slightly nearer tip of snout than to posterior edge of opercle, its lower edge a little above a horizontal from mouth to middle of caudal peduncle; cleft of mouth almost horizontal; maxillary, except distal end, concealed by preorbital; lips thick, the lower with a narrow frenum; jaws equal, the upper moderately protractile; teeth in 2 series on each jaw, outer series in a single row, large, canine-like, far apart; inner series minute, in bands; tips of teeth brown-colored; no teeth on vomer or palatines. Gill-membranes forming a fold across the isthmus; gill-rakers on first arch 10, short and blunt. Body covered with large weakly-ctenoid scales; head with cycloid scales; upper part of head anterior to middle of orbit, snout, preorbital area, and ventral part of head, naked; a single row of small scales along bases of dorsal and anal fins; small scales on basal part of inter-radial membranes of caudal; lateral line interrupted at the fourteenth scale, beginning again 3 scales lower and extending to base of caudal; first dorsal spine very short, others gradually longer to the sixth or seventh, after which the spines are about the same length; fifth and sixth dorsal rays longest, about $1\frac{1}{2}$ times the length of longest spine, depressed fin extending to posterior edge of dark caudal spot; first anal spine shortest, one-fifth as long as fifth spine, third, fourth, and fifth rays longest; depressed fin extending to anterior edge of caudal spot; caudal fin evenly rounded; pectoral rounded, extending to a vertical from vent; outer rays of pectoral longest, extending to vent. Color in alcohol, light olive, darker above than below; an indistinct, dark lateral band extending from snout to caudal; 8 or 9 scarcely distinguishable dark vertical bands on side of body; irregular dark spots at intersection of lateral and vertical bands; a small, dark spot at base of caudal; small, distinct dark dots on anterior dorsal region of head.

The total length of the type is 61 mm. Younger examples measuring about 43 mm. have the body a little deeper and the head shorter. The vertical color bands on posterior half of body are more distinct on the younger individuals. Rio Verde, Mexico. (Jordan & Snyder.)

Cichlasoma steindachneri, JORDAN & SNYDER, Bull. U.S. Fish Com. 1899 (1900), 143, Rio Verde near Rascon, Mexico. (Type, No. 6164, L. S. Jr. Univ. Mus. Coll. J. O. Snyder.)

Page 1540. Before *Theraps*, Günther, insert the following:

1948 (a). HEROS ISLANUS, Jordan & Snyder.

Head $2\frac{2}{3}$; depth $2\frac{2}{3}$; depth of caudal peduncle 7; eye $4\frac{1}{2}$ in head; snout $2\frac{1}{2}$; interorbital space $3\frac{1}{2}$; longest dorsal spine $2\frac{2}{3}$, ray (without filament) $1\frac{1}{2}$; longest anal spine $2\frac{2}{3}$, ray (without filament) $1\frac{1}{2}$; length of pectoral $1\frac{1}{2}$; ventrals $1\frac{1}{2}$; caudal $1\frac{1}{2}$; D. XVI, 10; A. V, 7; P. 14; scales 28-18. Body elongate, compressed, deepest above insertion of ventrals; dorsal outline rising rapidly to origin of dorsal, falling gradually to base of last spine, from which the descent to the caudal peduncle is more abrupt; ventral outline evenly rounded; interorbital space convex; eye large, orbit circular, equally distant from tip of snout and posterior edge of

opercle; mouth oblique, without fangs, protractile; nostrils 2, by the latter connected by a short tube; mouth wide, maxillary reaching to middle of eye; maxillary, except distal end, concealed by preorbital; lips thick, the lower with a narrow frenum; jaws equal, the upper moderately protractile; teeth in 2 series on each jaw, outer series in a single row, large, canine-like, far apart; inner series minute, in bands; tips of teeth brown-colored; no teeth on vomer or palatines. Gill-membranes forming a fold across the isthmus; gill-rakers on first arch 10, short and blunt. Body covered with large weakly-ctenoid scales; head with cycloid scales; upper part of head anterior to middle of orbit, snout, preorbital area, and ventral part of head, naked; a single row of small scales along bases of dorsal and anal fins; small scales on basal part of inter-radial membranes of caudal; lateral line interrupted at the fourteenth scale, beginning again 3 scales lower and extending to base of caudal; first dorsal spine very short, others gradually longer to the sixth or seventh, after which the spines are about the same length; fifth and sixth dorsal rays longest, about $1\frac{1}{2}$ times the length of longest spine, depressed fin extending to posterior edge of dark caudal spot; first anal spine shortest, one-fifth as long as fifth spine, third, fourth, and fifth rays longest; depressed fin extending to anterior edge of caudal spot; caudal fin evenly rounded; pectoral rounded, extending to a vertical from vent; outer rays of pectoral longest, extending to vent. Color in alcohol, light olive, darker above than below; an indistinct, dark lateral band extending from snout to caudal; 8 or 9 scarcely distinguishable dark vertical bands on side of body; irregular dark spots at intersection of lateral and vertical bands; a small, dark spot at base of caudal; small, distinct dark dots on anterior dorsal region of head.

Young individuals have the body a little deeper and the head shorter. The vertical color bands on posterior half of body are more distinct on the younger individuals. Rio Verde, Mexico. (Jordan & Snyder.)

Known only from
No. 6150, L.

Heros istlanus
Morelos, Mexico.

Page 1540.

Head $2\frac{2}{3}$; depth $2\frac{2}{3}$; interorbital space $3\frac{1}{2}$; longest dorsal spine $2\frac{2}{3}$, ray (without filament) $1\frac{1}{2}$; longest anal spine $2\frac{2}{3}$, ray (without filament) $1\frac{1}{2}$; length of pectoral $1\frac{1}{2}$; ventrals $1\frac{1}{2}$; caudal $1\frac{1}{2}$; D. XVI, 10; A. V, 7; P. 14; scales 28-18. Body elongate, compressed, deepest above insertion of ventrals; dorsal outline rising rapidly to origin of dorsal, falling gradually to base of last spine, from which the descent to the caudal peduncle is more abrupt; ventral outline evenly rounded; interorbital space convex; eye large, orbit circular, equally distant from tip of snout and posterior edge of

opercle; mouth horizontal, lower jaw projecting; lips thick, the lower without frenum, folding over the upper at their union; premaxillary protractile; maxillary small, nearly vertical in position, and almost hidden by the large preorbital; teeth on both jaws, in 2 series, the outer a single row of 24 canines, largest in front, growing smaller posteriorly, the inner series villiform; all of the teeth with brown-colored tips; gill-membranes free from isthmus; branchiostegals 5; gill-rakers on first arch 9, short, blunt, far apart. Body covered with large scales; cheek, opercle, interopercle, subopercle, and occipital portion of head with small scales; ventral part of head, preorbital area, snout, and anterior part of interorbital space naked; one row of scales extending on bases of dorsal and anal fins; scales of body weakly-etenoid; scales of head smooth; lateral line interrupted on the nineteenth transverse row of scales, beginning again 3 scales lower down and 2 scales in advance of where it left off and extending to base of caudal; first dorsal spine short and slender, others gradually longer and heavier, posterior spine longest; each spine with a rather stiff distal, ray-like attachment, the anterior edge of which projects above the membrane of fin; tips of fourth and fifth rays of dorsal uniting to form a thread-like filament about as long as the diameter of orbit; dorsal, when depressed, extending on caudal one-third its length; first anal spine shortest, others growing gradually longer and heavier, the last $2\frac{1}{2}$ times as long as the first; spines with distal attachments similar to those of dorsal, third and fourth rays longest, united at their tips, forming a slender filament; tip of anal extending a little farther posteriorly than that of dorsal; caudal rounded; tip of pectoral rounded; ventrals located slightly posterior to base of pectoral, extending to vent, outer ray longest, ending in a filament. Color dark, an oblong brownish-black spot at base of each scale on side of body, the spots growing less distinct above the pectoral; membranes of dorsal, anal, and caudal with small spots, these more distinct and regularly arranged on soft parts of dorsal and anal; pectoral and ventrals without spots.

Young individuals have a brownish-black spot at base of caudal and on side of body at tip of pectoral; a less distinct spot at upper edge of gill-opening and also below posterior end of base of dorsal. The darker of these spots are sometimes faintly indicated on the larger individuals. Morelos, Mexico. (Jordan & Snyder.)

Known only from Rio Ixtla at Puente de Ixtla, Morelos, Mexico. (Type, No. 6150, L. S. Jr. Univ. Mus. Coll. Jordan & Snyder.)

Hechos istlanae, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 144, Rio Ixtla, Morelos, Mexico.

Page 1542. After *Nectroplus nicaraguensis*, Gill & Bransford, insert:

1951 (a). NEETROPLUS CARPINTIS, Jordan & Snyder.

Head $2\frac{1}{2}$; depth 2; depth of caudal peduncle 6; eye 5 in head; snout $2\frac{1}{2}$; interorbital space $2\frac{1}{2}$; longest dorsal spine $2\frac{1}{2}$; ray $1\frac{1}{2}$; longest anal spine 2, ray $1\frac{1}{2}$; length of pectoral $1\frac{1}{2}$; ventrals $1\frac{1}{2}$; caudal $1\frac{1}{2}$; D. XVI, 10; A. V, 8; P. 15; scales 26-17, 7 on caudal peduncle. Body compressed, deepest part above ventrals; dorsal outline straight from tip of snout to a

point above anterior edge of orbit, where it is abruptly curved upward and backward to the origin of dorsal fin; from the latter point it gradually curves downward to base of first dorsal ray, from which the descent to caudal peduncle is abrupt; ventral outline evenly curved from snout to caudal peduncle; interorbital space convex, its middle portion flattened; orbit circular, nearer to posterior edge of opercle than to tip of snout, a distance equal to diameter of pupil; mouth oblique, lower jaw slightly projecting; lips thick; lower jaw with a frenum equal in width to $\frac{1}{2}$ diameter of pupil; upper jaw protractile; maxillary covered by preorbital except at its distal end; teeth in 2 series on each jaw, the outer series in a single row, flat or incisor-like, larger in front, growing much smaller posteriorly, the inner series minute, in narrow bands, all the teeth loosely attached, their tips brown-colored; no teeth on vomer or palatines; gill-membranes forming a fold across the isthmus; gill-rakers on first arch 10, short, far apart. Body covered with large, weakly-ctenoid scales; cheek, opercle, and occipital portion of head with small, cycloid scales; lower jaw, snout, and anterior half of interorbital space naked; bases of dorsal and anal fins with a row of small scales; inter-radial membranes of caudal with very small scales on basal parts; lateral line interrupted on the nineteenth transverse row of scales, beginning again 3 scales lower down on the third row anterior to where it left off, and extending to base of caudal; 2 short rows of mucous tubes on inter-radial caudal scales, one above and the other below end of lateral line; first dorsal spine shortest, others gradually longer and heavier, each spine with a ray-like attachment projecting above and posterior to its tip; first anal spine shortest, others gradually longer and heavier, the fourth three times as long as the first; spines with distal attachments similar to those of the dorsal, third and fourth rays longest, extending posteriorly as far as those of the dorsal; posterior edge of caudal somewhat convex; pectoral rounded; ventrals pointed, the outer ray much the longest, extending a little beyond vent. Color in alcohol, light slate; scales with lighter central spots; posterior parts of dorsal, anal, and caudal lighter; in life the head was covered with round and elongate spots of greenish blue on a background of golden brown; side of body with bluish and brownish spots without regularity of arrangement; pectoral and distal part of soft dorsal with a yellowish tinge.

In the young of this species there are 5 or 6 dark vertical bands, about equal in width to diameter of orbit, on the posterior part of the body; a dark spot sometimes present just below lateral line on a vertical through base of eleventh dorsal spine.

This species differs from *N. nematopus* and *N. nicaraguensis* in having a much deeper body and fewer dorsal and anal spines. Laguna del Carpinte, Mexico. (Jordan & Snyder.)

One specimen (type, No. 6162, L. S. Jr. Univ. Mus.) from Laguna del Carpinte, near Tampico, Tamaulipas, Mexico, collected by J. O. Snyder.

Neetroplus carpintis, JORDAN & SNYDER, Bull. U. S. Fish Com. 1899 (1900), 145, Laguna del Carpinte, near Tampico, Mexico.

Page 1557. *Eupomacentrus flavidatus* (Gill), is the young of *Eupomacentrus rectifrenum* (Gill).

Head 2.0
D. IX, 10;
Body moderately elevated, the outlines nearly straight, profile not slightly concave, and tip of large, high opercle; snout produced, the mouth large, the nostrils with strong folds of skin, of lower; the mouth in front, but not reaching the main mouth and anal edge of dorsal extended, otherwise narrow, slender and graduated to the ninth, which is highest, 2.2 in length, 3.3 in head; greatest, reaching the pointed, reacheing the lateral line of dorsal and anal.

Color in life: lower parts of body obliquely accented with maroon line from eye to snout, shades down to the front of dorsal, blue bars or spots surrounding pale-blue edges, posteriorly just between seventh spine; anal edge of first spine to the seventh rays color bordered with pale transparent.

Page 1612. Before *Nyrula*, Jordan, insert:

2022 (n.) **DORATONOTUS DECORIS**, Evermann & Marsh.

Head 2.6; depth 3.4; eye 4; snout 3.5; maxillary 4; interorbital 4.6; D. IX, 10; A. III, 9; pectoral 1.6; ventral 2.2; caudal 1.6; scales 1-26-6. Body moderately elongate, compressed throughout; the back a little elevated, the caudal peduncle deep and rather long; dorsal and ventral outlines nearly alike, the dorsal somewhat more strongly arched; anterior profile not trenchant, almost straight from snout to front of dorsal, very slightly convex in front of dorsal and very slightly concave between eye and tip of snout; head pointed, interorbital space broad and flat; eye large, high in position, middle of pupil nearer tip of snout than end of opercle; snout long, somewhat longer than diameter of eye, moderately produced, the lips broad in front, characteristically labroid; mouth not large, the maxillary not reaching front of orbit, the jaws equal, armed with strong, sharp teeth, about 4 canines in front of upper jaw, 2 in front of lower; teeth on sides of jaws also canine-like, smaller than those in front, but not distinctly different from them; a few smaller teeth behind the main row of large ones; vomer and palatines toothless; soft dorsal and anal each with a basal sheath of about two rows of large scales, that of dorsal extending over half the fin or more, that of anal lower, the fins otherwise naked; dorsal fin continuous, with a shallow notch, the spines slender and pungent, the second longer than the first, the following ones graduated to the fifth, which is shortest, thence increasing in length to the ninth, which is longest, 2.3 in head; soft dorsal with its middle rays highest, 2.2 in head; anal with three slender, sharp, graduated spines, the third longest, 2.2 in head; the soft part similar to soft dorsal, longest rays 1.3 in head; pectoral large, symmetrical, of 11 rays, the middle ones largest, reaching past tip of ventral nearly to vent; ventral moderate, pointed, reaching half-way to vent; caudal rounded; scales large, cycloid the lateral line on second row below the dorsal interrupted near the end of dorsal and beginning again on the row below, on caudal peduncle.

Color in life: Body chiefly green, darker green on back, lighter below; lower parts of head and breast light yellow; a broad white bar from eye obliquely across cheek and opercle, bordered above by an undulating maroon line and below by a similar but fainter line; a brown bar from eye to snout; 4 dusky spots near base of dorsal extending as fainter shades downward and slightly forward to or beyond lateral line, 1 from in front of dorsal, 2 under spinous dorsal, and 1 under soft rays; short pale-blue bars or spots on breast and about pectoral; iris blue, a pinkish border surrounding pupil; dorsal greenish, the soft part with yellow shade, a pale-blue edging to the whole fin, a maroon border to the green color posteriorly just inside the pale-blue edge, a small dark spot on membrane between seventh and eighth rays and a blue spot on membrane of first spine; anal colored like soft dorsal, the maroon border extending from first spine to last ray inside the pale edging, the dark spot between sixth and seventh rays; ventral green near base, pale blue outwardly, the green color bordered by maroon spots; pectoral plain, pale green; caudal very pale transparent blue, a wedge-shaped maroon spot on the 2 upper rays

near tip and a corresponding one on the 2 lower rays, the base of the wedge on outer ray; base of caudal with a pale undulate vertical bar bordered in front by a black line. In spirits, pale green, the maroon markings faintly persistent, becoming dusky.

Puerto Rico; known only from the type, 1.45 inches long, taken in the seine at Ponce, January 30, 1899, by the United States Fish Commission expedition to Puerto Rico. (*decoris*, beautiful.)

Doratonotus decoris, EVERMANN & MARSH, Rept. U. S. Fish Com. 1899 (Dec. 19, 1899), 355. Mayagüez, P. R. (Type, No. 49363, U.S.N.M. Coll. Evermann & Marsh.)

Page 1720. A specimen of *Alticera monoceros* (Osbeck) has recently been recorded by Dr. H. M. Smith from Woods Hole, in Bull. U. S. Fish Com. 1898 (1899), 273, pl. 64.

Page 1733. *Tetronotus nephelus*, Goode & Bean, is a good species distinct from *T. spengleri*, Bloch, and should stand as

2147 (a). SPHEROIDES NEPHELUS (Goode & Bean).

Page 1968. To the synonymy of *Urauidea gracilis* (Heckel), add:

Cottus gracilis cayuga, MEEK, Ann. N. Y. Ac. Sci., iv, 1888, 315. Cayuga Lake, N. Y.

Page 2017. The recent studies of the tide-pool Cottoids of the California coast by Mr. Arthur W. Greeley have resulted in the discovery of some new species and genera and other additional information regarding this group of fishes. The following new species and genera are to be added:

2384 (a). BLENNICOTTUS RECALVUS, Greeley.

Head $3\frac{1}{2}$; eye $4\frac{1}{2}$ in head; snout 3; D. IX, 15 or 16; A. 11 or 12; P. 14. Body short, stout, broad anteriorly; the head very broad, short and blunt; snout obtuse; interorbital space $5/6$ eye, grooved, the groove leading into a depressed space behind the eyes; mouth distinctly terminal, the maxillary reaching a vertical below the anterior edge of orbit, the lower jaw included; minute conical teeth on jaws, vomer, and palatines; nasal spines very small; no preopercular spine apparent in adults; edge of preopercle rounded; opercle ending in a rounded flap; branchiostegals six, the membranes broadly united, free from the isthmus; gills $3\frac{1}{2}$, a slit behind the last gill. Dorsal fins very long, slightly joined at the base, origin of first dorsal directly over tip of opercle, that of soft dorsal in advance of origin of anal; first dorsal slightly rounded, the middle ray the longest; pectoral reaching origin of anal, membranes of first seven rays deeply emarginate; ventrals reaching vent; anal papilla of male very large; anal low, the membranes of all the rays except the last three deeply emarginate, none of the rays enlarged in the male; caudal short, slightly rounded.

Cirri few and small, those of top of head joined at base in conspicuous bunches, two irregular occipital rows, a few below these on the side of head, and on the margins of preopercle and opercle; a few above origin of pectoral, and a weak row along the anterior third of the lateral line.

Color of body, light brown, vermiculated with white, and marked dorsally with 4 or 5 wedge-shaped spots of dark brown, edged with white,

the more caudal part and along tinged with brown which individual.

The blue mouth margin except *B. gracilis* preopercular its mouth, the young eating that of *Oligocottus* which does not reach *recalvus* is apparently succeeded have been several species of *Oligocottus* *recalvus*. The flap is still in range and can mark, where *recalvus*, b.

Centridermichthys
Oligocottus gracilis
Blennicottus gracilis

MANN, Fish
Girard.
Blennicottus gracilis
Pacific Grove

Blenniculus, G.

This genus
prickly scale,
sharp,
rough-skin-

Head $3\frac{1}{2}$;
V. 1, 3. Bod-
so; head ed-
grooved; tail
blunt, snout
eared with m-

Bull.

and more distinct posteriorly; 2 pinkish spots on the dorsal side of the caudal peduncle, and a faint shading of the same color on side of head and along anterior fourth of lateral line; entire under surface dull brown, tinged with olive; fins indistinctly barred with grayish white; tail faintly tinged with pink. In some specimens the color is an almost uniform dull brown while in others the light markings are prominent. Some young individuals from among the green algae are uniform light green.

The bluntness of the snout and preopercular spines, and the terminal mouth make this species easily distinguishable from all related forms, except *B. globiceps*, from which it is separated by the shape and size of its preopercular and nasal spines, the number of its cirri, 12, and the size of its mouth. The adults of these 2 species can be readily distinguished, but the young of *B. recalvus* is very similar to the young of *B. globiceps*, indicating that *B. globiceps* is probably the ancestral form. Girard's old description of *Oligocottus globiceps* has been erroneously associated with this fish, which does not extend so far north as the type locality of *B. globiceps*. *B. recalvus* is distributed from San Diego to Santa Cruz, where it is immediately succeeded by *B. globiceps* on the north. No specimens of *B. recalvus* have been taken north of the region of Santa Cruz. On the other hand, several specimens of *B. globiceps* have been collected on the coast of Monterey County south of Monterey Bay, therefore within the range of *B. recalvus*. The relations between the two species where their ranges overlap is still to be made out. *B. recalvus* is quite common throughout its range and everywhere inhabits the deep shaded tide-pools, near low water mark, where a large number will often be found in a single pool. (Greeley.) (*recalvus*, bald in front.)

Centridermichthys globiceps, GÜNTHER, Cat., II, 171, 1860; not of Girard.

Oligocottus globiceps, JORDAN & GILBERT, Synopsis, 718, 1883.

Bleonicottus globiceps, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 808; JORDAN & EVERMANN, Fishes of North and Middle America, II, 2017, 1898; not *Oligocottus globiceps* Girard.

Bleonicottus recalvus, GREELEY, Bull. U. S. Fish Com. 1899 (Dec. 13, 1899), 9, fig. 1, Pacific Grove, Cal. (Type, No. 8068, L. S. Jr. Univ. Mus. Coll. Greeley & Cowles.)

746 (a). RUSCICULUS, Greeley.

Rusciculus, GREELEY, Bull. U. S. Fish Com. 1899 (Dec. 13, 1899), 13 (*rimensis*).

This genus is allied to *Oryctocephalus*, differing in the presence of minute prickly scales, which cover dorsal half of body. Preopercular spine simple, sharp. No slit behind the last gill. (*ruscum*, the butcher's broom, a rough-skinned plant.)

2384 (n). RUSCICULUS RIMENSIS, Greeley.

Head $3\frac{1}{2}$; eye 4 in head; snout $3\frac{1}{2}$ in head; D. IX-17 or 18; A. 14; P. 14; V. 1, 3. Body compressed, very slender, the caudal peduncle especially so; head compressed, flat; snout pointed; interorbital space $\frac{5}{6}$ eye, grooved; top of head flat and slightly concave; nasal spines large, blunt, snout abruptly decurved below them. Dorsal half of body covered with minute, embedded, prickly scales partially arranged in obscure

Bull. No. 47, pt. 4—X

oblique rows, none below lateral line. Minute pointed teeth on jaws, vomer, and palatines; jaws subequal, mouth horizontal, the maxillary reaching a vertical below the anterior edge of pupil. Margin of preopercle armed with one sharp spine curved upward, below which are 1 and sometimes 2 very short blunt processes; margin of opercle ending dorsally in a pointed flap. Branchiostegals 6, the membranes broadly united, free from the isthmus; no slit behind the last gill. Dorsal fins not joined, the soft dorsal very large; first dorsal beginning slightly in advance of opercular flap, the upper edge much rounded, the fifth spine being longest; origin of soft dorsal just in front of origin of the anal in the female, directly above it in the male, the fin very long; pectoral large, reaching a vertical below ninth ray of soft dorsal; origin of ventrals posterior to a point midway between anal and base of pectoral in the male, anterior to it in the female, the difference caused by the enlargement of first 2 anal rays in the male; anal fin small, the rays slender, the membranes of all deeply emarginate; the first 2 anal rays of male greatly enlarged, joined by membrane to each other and to the rest of the fin; the posterior edge of tail nearly straight; anal papilla inconspicuous. Cirri small and scarce, always occurring singly, never in bunches or joined at the base, with the exception of a few pairs along the anterior third of the lateral line; one above each orbit, 2 rows of 3 each behind these on top of head, 1 cirrus on the inside of each nasal spine; a cirrus on the end of maxillary, 2 or 3 on the margin of the preopercle below the preopercular spine, and a row along the anterior half of the lateral line. Color, light olive or reddish brown tinged with lavender, marked dorsally with 4 or 5 wedge-shaped, indented spots of black, a broken band of same color along the lateral line, sometimes sending branches below it which show a tendency to inclose round spots; a more or less distinct spot of black on top of the head; a faint postocular line, a spot below the eye, and a preopercular line running from eye to snout, all of same color; pectoral and caudal indistinctly barred with brown, anal tinged with it, and the dorsal covered with fine brown or black spots sometimes very faint; throat and belly pale yellowish white, unspotted.

This species is most closely related to *Orycottas embryum*, with which it agrees in general coloration, but differs decidedly in the presence of scales, the slenderer body, the larger number of soft dorsal and anal rays, the serrated margin of the preopercle, and the arrangement of the ciri. Rare; only 2 other specimens from Point Lobos, California, are known to us. It inhabits the tide-pools lined with corallines, and in its coloration imitates very closely these algae. Length 40 mm. The smallest of our tide-pool fishes. (Greeley.) (*rimus*, a crevice; *rimensis*, living in crevices.)

Ruscarius rimensis, GREELEY, Bull. U. S. Fish Comm. 1890 (Dec. 13, 1890), 13, fig. 3.
Point Lobos, Monterey County, Cal. (Type, No. 6067, L. S. Jr. Univ. Mus. Coll. A. W. Greeley.)

746 (b). *DIALARCHUS*, Greeley.

Dialarchus, GREELEY, Bull. U. S. Fish Com. 1890 (Dec. 13, 1890), 14 (snyderi).

Preopercular spine forked at tip; scales none; first anal ray of male enlarged, joined to the second, the two widely separated from the rest of

the fin,
anal rays
joined

Head 3
to 45; V;
minute co
mouth hor
rior edge
groove lea
spines lar
as eye, fre
pointing a
life; marg
chiostegals
Gills 3½, a
diameter o
pectoral; t
upper edge
spine; orig
well beyon
base of ped
the male t
distinctly s
the last 3
the male o
or 4, those a
above orbit
top of head
head, 2 or 3
preopercula
on its upper
of bunched
defined row
seventeenth
each spine b
scattered bo
body; a cirr

Color, ligh
fine indistin
base of dor
much broke
brown belo
post-ocular
color on top
opercle; thr
bluish green
dorsal fins

the fin. Closely allied to *Oligocottus*, differing only in the character of the anal rays of the male.

($\delta\tau\alpha\lambda\ddot{o}\delta$, divided; $\dot{\alpha}\rho\chi\ddot{o}\zeta$, anus.)

2384 (b). *DIALARCHUS SNYDERI*, Greeley.

Head $3\frac{1}{2}$; eye $4\frac{1}{2}$ in head; snout $3\frac{1}{2}$; D. VIII-18 or 19; A. 13 to 15; P. 13 to 15; V. 1, 3. Body elongate, slender, the snout pointed, compressed; minute conical teeth on jaws, vomer, and front of palatines; jaws equal; mouth horizontal, the maxillary 3 in head, reaching a vertical below anterior edge of pupil. Interorbital space $\frac{6}{7}$ eye, shallowly grooved, the groove leading into a depressed space between the occipital ridges; nasal spines large. Margin of preopercle armed with a strong spine, $\frac{1}{2}$ as long as eye, from the upper border of which at base extends a second spine pointing abruptly upward and inward; both spines covered with skin in life; margin of opercle ending in a pointed flap entirely unarmed. Branchiostegals 6, the membranes broadly united, free from the isthmus. Gills 3½, a slit behind the last gill. Dorsal fins large, separated by $\frac{1}{2}$ diameter of eye, whole length equaling that from caudal to base of pectoral; first dorsal beginning slightly in advance of margin of opercle, upper edge nearly straight, curving abruptly downward from the sixth spine; origin of soft dorsal in advance of anal; pectoral large, reaching well beyond the origin of the anal; ventrals almost midway between base of pectoral and anal; anal fin small, rays all feeble in the female, in the male the first ray only greatly enlarged, joined to the second, the 2 distinctly separated from the rest of the fin, the membranes of all except the last 3 or 4 rays deeply emarginate. Anal papilla small, present in the male only. Cirri very numerous, usually occurring in bunches of 3 or 4, those of head joined at the base, forming a comb; 2 pairs of bunches above orbits, with the rudiments of a third bunch in front of these, 3 on top of head, behind orbits, 2 or 3 bunches just below these on side of head, 2 or 3 single cirri on margin of preopercle, a thick bunch above the preopercular spines, 4 or 5 on lower margin of opercle, with a thick bunch on its upper margin; a short row above the base of the pectoral; a row of bunched cirri along anterior two-thirds of lateral line, another well-defined row along the dorsal fin from the third spine to the sixteenth or seventeenth ray of soft dorsal, this row containing a bunch at the base of each spine and ray, with the occasional exception of the first ray; 5 or 6 scattered bunches between the dorsal and lateral rows on each side of body; a cirrus at the tip of each dorsal spine.

Color, light reddish brown, sometimes almost pink, thickly spotted with fine indistinct white spots; 4 or 5 irregular dark brown spots along the base of dorsal, a band of same color along lateral line, sometimes very much broken and extending ventrally, shading into the uniform reddish brown below, and including 3 or 4 round pinkish spots; a dark brown post-ocular line, another running forward from the eye, a patch of same color on top of head, another on side of head, and 2 or 3 on edge of opercle; throat reddish brown with several distinct white spots; belly bluish green; a silvery white patch between the bases of the pectorals; dorsal fins pale reddish brown with black and clear spots; pectoral

crossed irregularly with white; anal fin pale pink, crossed with dark brown. There are 2 or 3 perfectly distinct types of coloration, as follows: Some specimens from pools containing green algae are pure light green, others from coralline pools are tinged with lavender, as *B. embryum*. This species resembles most closely *O. maculosus*, which name has been erroneously applied to it, but it differs markedly in its slenderer body, more pointed snout, the arrangement of the cirri, and the perfectly distinct coloration, also in the greater length of the dorsal fins, the enlargement of only one anal ray in the male, and the shortness of the maxillary. Coast of California.

Specimens are at hand from Crescent City, Cal., Bolinas Bay, Half Moon Bay, Monterey Bay, and San Luis Obispo, Cal. Found in all kinds of pools, from San Francisco to Monterey Bay, but nowhere common. Length 60 mm. The most beautiful and active of the tide-pool fishes, extremely variable in color. (Greeley.)

Centridermichthys maculosus, GÜNTHER, Cat., II, 171, 1860; not *Oligocottus maculosus*, Girard.

Oligocottus maculosus, JORDAN & GILBERT, Synopsis, 718, 1883; JORDAN & EVERMANN, Fishes of North and Middle America, II, 2013, 1898.

Oligocottus snyderi, GREELEY, in JORDAN & EVERMANN, Fishes of North and Middle America, III, 2871, 1898, Pacific Grove, Cal. (Type, No. 5846, L. S. Jr. Univ. Mus. Coll. Greeley & Maddren.)

Dialarchus snyderi, GREELEY, Bull. U. S. Fish Com., 1899, 15, fig. 4.

946 (c) EXIMIA, Greeley.

Eximia, Greeley, Bull. U. S. Fish Com. 1899 (Dec. 13, 1899), 18 (*rubellio*).

Allied to *Oligocottus*, but differing in the presence of a large three-pointed preopercular spine instead of the simple forked spine of *Oligocottus*. Skin smooth. A slit behind the last gill.

2384 (c). EXIMIA RUBELLIO, Greeley.

Head 2 $\frac{1}{2}$; eye 3 $\frac{1}{2}$ in head; snout 3 $\frac{1}{2}$; D. VII or VIII-15 or 16; A. 12 or 13; P. 13 or 14; V. 1, 3. Body compressed, snout pointed and compressed, head deep, occiput narrow, slightly concave; interorbital space narrow, $\frac{1}{2}$ the large eye, shallowly grooved. Nasal spines prominent, very large and pointed. Teeth small, pointed on jaws, vomer and palatines; jaws equal, mouth horizontal, maxillary 3 in head, reaching a vertical below anterior edge of pupil. Margin of preopercle armed with a very strong spine as long as eye, extending backward and downward, bearing on its upper surface a second and third spine, both pointing back and up; all the spines covered with skin in life; opercle ending in a rounded flap. Branchiostegals 6, not united to the isthmus; gills 3 $\frac{1}{2}$, a slit behind the last gill. Anal papilla inconspicuous. Dorsal fins not joined; first dorsal beginning in advance of margin of opercle, first 2 spines short, the upper margin slightly rounded; soft dorsal beginning in advance of origin of anal, all the rays and spines very slender, pectoral reaching well beyond the origin of the anal; anal fin small, the rays slender, the membranes emarginate between each 2 rays; in the males the first ray enlarged, the second slightly elongated, the 2 united and not separated from the rest

the edge of 1 never jo 2 or 3 cir 3 pairs o side of he of preope scattered anterior l a bunch the last s base of t lateral ro light bro where with more cons dorsal sid green, bee from eye t preopercle spot with very light a brownish trals, light brown, ana is lighter a to *Dialar opercular s the larger o other speci other locali iting only t *Eximia rubell Grove, Ca**

Page 220
For its use i *Gilbertidiae BE*

Page 220

Head 4.4; interorbital longest dors head, longes

Body rath width 1.5 in simple, flexi

the fin, as in *Dialarchus snyderi*; ventrals situated below the upper edge of the base of the pectoral, just reaching the anus. Cirri all distinct, never joined at the base in a comb as in *Dialarchus snyderi*; 3 pairs of 2 or 3 cirri each above the orbits, the first directly above the nasal spines, 3 pairs on top of head behind orbits, a few scattered cirri below these on side of head, a bunch of 2 or 3 on end of maxillary, a row on lower margin of preopercle, a large bunch above the preopercular spines, and several scattered cirri on margin of opercle; a row of bunched cirri along the anterior half of lateral line, a row along the base of the dorsal, including a bunch of 3 or 4 for each spine and ray, the row bending downward at the last spine of the first dorsal, leaving a space between the cirri and the base of the dorsal spines; a few scattered cirri between the dorsal and lateral rows, and below the lateral line behind the pectoral fin. Color light brown to all shades of light red, pink, or lavender, spotted everywhere with white, the spots extremely minute on dorsal half of body, but more conspicuous ventrally; 5 wedge-shaped spots of dark brown along dorsal side of body; head dark brown, sometimes blotched with red or green, becoming lighter on side, leaving a dark postocular line extending from eye to the preopercular spine, and a dark spot on lower margin of preopercle, everywhere very finely marked with white and blue; a white spot with a brown center just in front of first dorsal; throat and belly a very light, bluish green, shading into a faint yellow behind pectoral, and a brownish green on each side of anal; all the fins, excepting the ventrals, light brownish green barred with dark brown; caudal light reddish brown, anal and tip of pectoral tinged with pinkish. A young individual is lighter and more brilliantly colored. This species is most closely allied to *Dialarchus snyderi*, from which it differs in the presence of a third preopercular spine, the greater depth and comparative length of the head, the larger eye and nasal spines, and the arrangement of the cirri. Many other specimens taken at Monterey Bay, but it is not recorded from any other locality. The most brilliantly colored of the tide-pool fishes, inhabiting only the deep pools rich in plant life. (*rubellio*, a rosy one.)

Eximia rubellio, GREELEY, Bull. U. S. Fish Com. 1890 (Dec. 13, 1890), 18, fig. 5, Pacific Grove, Cal. (Type, No. 6066, L. S. Jr. Univ. Mus. Coll. Greeley & Muddren.)

Page 2027. *Gilbertina*, Jordan & Starks, is preoccupied in Lepidoptera. For its use in ichthyology Dr. Berg has substituted *Gilbertidia*.

Gilbertidia BERG, Com. del Museo Nac. de Buenos Aires, 1898 (Dec. 17, 1898), 43 (*sigillatus*).

Page 2207. After *Sicydium rincante*, Jordan & Evermann, insert:

2531 (6). **SICYDIUM CAGUITE**, Evermann & Marsh.

Head 4.4; depth 4.8; eye 5.75; snout 2.5; maxillary 2; mandible 2.75; interorbital width 3; preorbital 3.5; D. VI-I, 10; A. I, 9; scales 83-25; longest dorsal spine 1.5 in head, longest ray 2; longest anal spine 2 in head, longest ray 2; pectoral 1.1; ventral disk 1.75; caudal 1.

Body rather stout, heavy forward; head large, broad; mouth large, its width 1.5 in head; lips very thick; maxillary not greatly produced; teeth simple, flexible; a median cleft in upper lip; pectoral somewhat shorter

than head; dorsal spines without filaments, the longest about 1.5 in depth of body; space between dorsals about equal to orbit; soft rays of dorsal and anal scarcely reaching base of caudal; ventrals united, forming a cup-shaped disk, only about two-fifths posterior edge free from belly; caudal rounded. Scales very small, ctenoid, densely covering entire body except a broad strip on belly; posterior portion of nape with very fine scales; entire head naked.

Color: Dark brown or olivaceous on head, side, and back; under parts pale; fins all pale, the anal with a narrow darkish margin; caudal somewhat dark; no dark vertical bars on body and none at base of pectoral; no H-shaped figure at base of caudal.

This species is close to *S. plumieri*, from which it differs chiefly in the color, the more complete squamation, the shorter pectoral, and the non-filamentous character of the dorsal spines.

Puerto Rico. A single specimen, 3.63 inches long, obtained in the Rio de Caguitas at Caguas, January 9, 1899, by the U. S. Fish Commission expedition to Puerto Rico. (Named for the Rio de Caguitas, from which the type was obtained.)

Sicydium caguitae, EVERMANN & MARSH, Rept. U. S. Fish Com. 1899 (Dec. 19, 1899), 355, Rio de Caguitas, Caguas, P. R. (Type, No. 49364, U.S.N.M. Coll. Evermann & Marsh.)

Page 2230. After *Gobius oceanicus*, Pallas, insert:

2558 (a). *GOBIUS BAYAMONENSIS*, Evermann & Marsh.

Head 4.8; depth 6.4; eye 5; snout 3.2; maxillary 1.8; mandible 1.9; interorbital 7.6; preorbital 4.6; scales 71-19, about 29 before dorsal; D. IV-14, the longest spine about 0.7 in head, the longest ray 1.5; A. 15, the longest ray 1.5; pectoral 1.1; ventrals 1.1; caudal very long and pointed. Body very long and slender; head long; caudal peduncle long; mouth very large, oblique; maxillary long, reaching past posterior border of orbit.

Color as in *G. oceanicus*, which this species closely resembles. The smaller (71 instead of 63 to 65), almost cycloid scales, the longer head, larger mouth, longer maxillary, and the longer and more slender body are differences which we can not reconcile with the descriptions of that species or with the numerous specimens of it which we have from Puerto Rico.

Puerto Rico. Known only from the type, a specimen 9 inches long, obtained in the San Juan market January 14, 1899, by the U. S. Fish Commission expedition to Puerto Rico. It probably came from near the mouth of the Bayamon River at Palo Seco, for which stream the species was named.

Gobius bayamonensis, EVERMANN & MARSH, Rept. U. S. Fish Com. 1899 (Dec. 19, 1899), 355, Mouth of Bayamon River, Palo Seco, P. R. (Type, No. 49365, U.S.N.M.; coll. Evermann & Marsh.)

Page 2218. To the synonymy of *Gobius soporator* add the following:

Gobius arundelii, GARMAN, Proc. N. E. Zool. Club, I, 63, June 9, 1899, Clipperton Island. (Type in M. C. Z.)

Page 2

Head 4
interorbital
longest sp.
in head;

Body long
oblique;
isthmus n.
eyes large
median ke
jaws in na
short, abo
behind ba
space betw
anal reach
pointed, a
of anal; v
reaching o
and breas
scales befo

Color: F
profuse fir
dusky; do
on posterio
what dusky

Length
dredged b
Puerto Rea
type localit

Bolmannia t
1899, 356,
station 60

Page 224

Head 3.75
maxillary 2

Body slen
caudal; hea
high; mo th
lower jaw p
enlarged an
rather narro

Body dens
what reduce

Page 2240. Before, *Aboma*, Jordan & Starks, insert:

2570 (a). **BOLLMANNIA BOQUERONENSIS**, Evermann & Marsh.

Head 4; depth 5.5; eye 3.5; snout 4.4; maxillary 2.2; mandible 2.5; interorbital width 3 in eye; preorbital 6; scales 27-8; D. VII-13, the longest spine 1.5 in head, the longest ray 1.2; A. 12, the longest ray 1.25 in head; pectoral 1; ventrals 1.1; caudal 0.4.

Body long, slender, tapering; head short; snout blunt; mouth large, oblique; jaws subequal, maxillary reaching posterior border of pupil; isthmus narrow, the gill-openings reaching forward to below preopercle; eyes large, high, close together, the interorbital very narrow and without median keel; no fleshy process on inner edge of shoulder girdle; teeth on jaws in narrow bands, those of outer series somewhat enlarged; opercle short, about 3 in head. Fins moderate; origin of spinous dorsal slightly behind base of pectoral, its spines 7 in number, not filamentous; interspace between dorsals less than diameter of eye; soft rays of dorsal and anal reaching, when depressed, beyond base of caudal; caudal long and pointed, as in *Gobius oceanicus*; pectoral pointed, reaching beyond origin of anal; ventral disk moderate, free from belly, the longest rays barely reaching origin of anal. Scales very large, weakly ctenoid; nape, cheek, and breast scaled, the scales somewhat smaller than on body, about 9 scales before the dorsal.

Color: Pale olivaceous or straw-color, back and upper part of head with profuse fine dark punctulations; under parts pale, breast somewhat dusky; dorsal fins barred with white and dark, a large jet-black ocellus on posterior part of spinous dorsal; other fins pale, the ventral disk somewhat dusky in front. The only known Atlantic member of the genus.

Length 2 to 3 inches. Puerto Rico. Known only from 5 specimens dredged by the U. S. Fish Commission expedition to Puerto Rico, off Puerto Real, in Ensenada del Boqueron, January 25, 1899. (Named for the type locality.)

Bollmannia boqueronensis, EVERMANN & MARSH, Rept. U. S. Fish Com. 1899 (Dec. 19, 1899), 356. Ensenada del Boqueron, off Puerto Real, in 8.5 fathoms, at Fish Hawk station 6074. (Type, No. 49366, U.S.N.M. Coll. Evermann & Marsh.)

Page 2245. Before *Microgobius thalassinus*, Jordan & Gilbert, insert:

2575 (a). **MICROGOBIUS MEEKI**, Evermann & Marsh.

Head 3.75; depth 6; eye 3.5; snout 5.5; interorbital 7; preorbital 7; maxillary 2; mandible 1.5; scales 55-52; D. VII-17; A. 16.

Body slender, greatly compressed, tapering regularly from pectorals to caudal; head moderately heavy, interorbital space very narrow; eye large, high; mouth large, oblique; maxillary reaching posterior border of orbit; lower jaw projecting; teeth in bands in each jaw, the outer series greatly enlarged and strongly recurved, those of lower jaw largest; isthmus rather narrow, the gill-openings continue forward.

Body densely scaled, the scales strongly ctenoid, those anteriorly somewhat reduced; nape, breast, and entire head naked. Origin of spinous

dorsal from snout 3.5 in length; dorsals very close together; spines of first dorsal filamentous, exceeding head in length; soft dorsal and anal long, their bases about equal, about 2.5 in body, their last rays reaching past base of caudal when depressed; caudal pointed, its longest rays about equal to head; pectoral about equal to head, reaching origin of anal; ventrals united, almost reaching origin of anal.

Color: Light olivaceous, dusted over uniformly with fine dark punctulations; a large dark shoulder spot between the base of pectoral and origin of spinous dorsal; a few indistinct dark areas on side of head; lower jaw dark at tip; an obscure dark blotch at base of caudal; fins all rather pale except ventrals, which are dark, perhaps bluish in life; caudal somewhat dusky; anal dark-edged. Length 1.5 inches.

This species seems related to *M. eulepis*, Eigenmann & Eigenmann, described from Fortress Monroe, Va., but differs in the smaller and strongly ctenoid scales, greatly compressed body, and in the coloration.

Puerto Rico; known only from the type, 1.5 inches long, dredged by the U. S. Fish Commission expedition to Puerto Rico, February 8, 1899. (Named for Dr. Seth Eugene Meek, assistant curator of zoology, Field Columbian Museum.)

Microgobius meeki, EVERMANN & MARSH, Rept. U. S. Fish Com. 1899 (Dec. 19, 1899), 356. Between Vieques and Culebra islands at Fish Hawk station 6087, in 13.25 fathoms. (Type, No. 49367, U.S.N.M. Coll. Evermann & Marsh.)

Page 2350. After *Enneanectes carinalis* (Jordan & Gilbert), insert:

868 (b). GILLIAS, Evermann & Marsh.

Gillias, EVERMANN & MARSH, Rept. U. S. Fish Com. 1899 (Dec. 19, 1899), 357 (*jordani*).

Body short and stout, tapering rapidly from the short, broad head to the short, compressed caudal peduncle; scales large, rough-ctenoid; lateral line complete, or nearly so, broken under last spines of middle dorsal; a broad, double-pointed tentacle above eye; dorsal fin divided into 3 parts, the first of 3 short spines, the second of 11 longer spines, and the third of 7 rays.

This genus is closely related to *Enneanectes*, Jordan & Evermann, from which it differs in the presence of the orbital tentacle, the more complete development of the lateral line, and the larger scales.

(Named for Dr. Theodore Gill.)

2687 (b). GILLIAS JORDANI, Evermann & Marsh.

Head 3.5; depth 4.3; eye 2.5; snout 3.5; maxillary 2.4; mandible 1.9; scales 2-30-7; D. XII-XII-7; A. II, 15; longest dorsal spine 1.8 in head, longest ray 1.6; longest anal ray 2.3; pectoral 0.8; ventral 1.3; caudal 1.3.

Body short and stout, tapering rapidly to the short, compressed caudal peduncle; head short; snout short; blunt, concave in front of eyes; mouth small, slightly oblique, jaws equal; eye large, high up, interorbital width very narrow; a broad bifid orbital tentacle, none on nape. Scales very large and rough-ctenoid; opercles and entire head rough; lateral line nearly complete, beginning immediately above base of pectoral at upper end of gill-opening and extending parallel with back to posterior

part of mouth; dorsal; rays of caudal; anal; pectoral; dorsal; ventral.

Color in the origin between scales; inky-black; under part narrow light; followed by a fin finally marked an expedition taken on the preceding day.

(Named after Gillias jordani Cordona)

Page 2350

Head 3.3; interorbital 1.3; caudal 1.3; dorsal 1.3; Body slender; vex; mouth thick; lips thick; jaw; a single high, origin in front of last dorsal; rays, no spines anteriorly, with the origin on the

Color in what regular base of dorsal smaller one series at base in color and sometimes forming

part of middle dorsal fin (or for 12 scales) where there is a break, the line dropping down 3 scales, then continuing with one or two interruptions to base of caudal; belly and breast scaled; dorsals 3, the first of 3 short, flexible spines, close to the second, which has 12 longer, rather stiffer, spines, separated from the third by a space one-third diameter of eye; anal long and low, the membranes deeply notched between the rays; pectoral of 15 rays, broad and short, reaching posterior end of second dorsal; ventral 2, slender.

Color in alcohol: Brown, body crossed by 4 broad blackish bars, one at the origin of second dorsal, one under last spines of same fin, the third between second and third dorsals, and the fourth under third dorsal; an inky-black bar across caudal peduncle at base of caudal fin; head and under parts rusty; fins all barred with light and dark; caudal with a narrow light bar at base, then a black one, then a broader white one, followed by a much broader dark bar containing some white areas, the fin finally tipped with white. Puerto Rico. Two specimens of this well-marked and interesting species were obtained by the U. S. Fish Commission expedition to Puerto Rico, the type 1.5 inches long (No. 49368, U.S.N.M.), taken on the Cardona Light-House Reef, at Ponce, February 1, 1899, and another specimen of about the same size taken at the same place the preceding day.

(Named for Dr. David Starr Jordan.)

Gillias jordani, EVERMANN & MARSH, Rept. U. S. Fish Com. 1899 (Dec. 19, 1899), 357,
Cordona Light-House Reef, Ponce, P. R. (coll. Evermann & Marsh.).

Page 2357. After *Malacoctenus lugubris* (Poey), insert:

2696 (a). **MALACOCTENUS CULEBRE**, Evermann & Marsh.

Head 3.35; depth 5; eye 4.2; snout 4.5; maxillary 2.2; mandible 1.8; interorbital 6.5; scales 2-35-11; D. XXI, 8; A. II, 18; pectoral 1.3; ventral 1.3; caudal 1.4.

Body slender, compressed; head rather long, pointed, upper profile convex; mouth large, the maxillary nearly reaching posterior border of orbit; lips thick, jaws equal; teeth very small, conical, a single row in each jaw; a single nasal, ocular, and nuchal filament; dorsal fin moderately high, originating above the origin of lateral line, a shallow notch in front of last two dorsal spines, the membrane free from caudal; anal origin under about tenth dorsal spine; caudal somewhat pointed; pectoral large, reaching anal; ventrals moderate, not reaching anus, of two rays, no spine evident; lateral line distinct throughout, running high anteriorly, where it is slightly curved, turning abruptly downward over the origin of anal, thence median to base of caudal.

Color in spirits: Body everywhere mottled with dark brown, in somewhat regularly arranged blotches, a series of about nine of these at the base of dorsal, barely extending upon the fin; a similar series of much smaller ones at base of anal, not evident on all specimens; below the series at base of dorsal are two other series of the same blotches less deep in color and not so well defined, extending the length of body and sometimes forming, with the upper series, more or less broken vertical bars;

between the blotches a lighter shade of brown is interwoven with pale streaks of ground color; head nearly pale below, save some dark on chin and isthmus; two wide streaks from eye across cheek; opercle dark brown; top of head with the color of body; lips with brown and pale stripes; posterior half of maxillary pale; dorsal rather dark; caudal uniform gray or faintly barred; anal similar to dorsal in color; the rays with pale tips forming a white edge; pectoral like caudal; ventrals pale. Length about 1.5 inches.

A plainly marked species most closely related to *M. lugubris* (Poey). Puerto Rico; only 3 specimens known, all from the coral reefs about Culebra Island, February 9, 1899.

(Named for the type locality.)

Malacoctenus culbrei, EVERMANN & MARSH, Rept. U. S. Fish Com. 1899 (Dec. 19, 1899), 357, Culebra Island. (Type, No. 49369, U.S.N.M. Coll. Evermann & Marsh.)

Page 2358. After *Malacoctenus gilli* (Steindachner), insert:

2697 (a). *MALACOCTENUS MOOREI*, Evermann & Marsh.

Head 3.6; depth 3.7; eye 3.5; snout 3.4; maxillary 4.5; mandible 4.5; interorbital 4; scales 3-45-5. D. xxii, 11; A. ii, 20; pectoral 1 in head; ventral 1.2; caudal 1.2; longest dorsal spine 1.5, ray 1.2; longest anal ray 1.5.

Body short, rather stout, compressed; head short, snout short, but pointed; mouth rather small, little oblique, the gape scarcely reaching orbit; teeth in each jaw in a single series; gill membranes broadly united across the isthmus; eye small, interorbital space wide; dorsal outline rising abruptly to above eye, thence gently curved to origin of dorsal fin, and from there nearly straight to base of caudal fin; ventral outline regularly convex.

Color in alcohol: Light olivaceous, the body crossed by about 9 or 10 broad dark vertical bars, which extend upon dorsal fin, these usually broadest above, the pale interspaces therefore broadest on lower half of body; the fourth from last is a narrow dark line, the one following it is a double spot, the next narrow and indistinct, the last, at base of caudal, more distinct, followed by 3 small irregular white spots; top of head brown; side of head with fine punctulations; a dark line running forward from eye, a dark spot below eye, 2 or 3 dark blotches on anterior edge of opercle; under surface of head crossed by 3 or 4 irregular, indistinct dark lines; caudal and anal with fine dusky punctulations; pectoral and ventrals pale.

This species is close to *M. gilli*, from which it may be distinguished by the larger dorsal and anal fins, the greater depth, wider interorbital, and the coloration.

Puerto Rico; known only from the type, 1.4 inches long, obtained by the U. S. Fish Commission expedition to Puerto Rico, February 11, 1899.

(Named for Dr. H. F. Moore, naturalist on the U. S. Fish Commission steamer *Albatross*.)

Malacoctenus moorei, EVERMANN & MARSH, Rept. U. S. Fish Com. 1899 (Dec. 19, 1899), 358, Culebra Island. (Type, No. 49370, U.S.N.M. Coll. Evermann & Marsh.)

Page 2358. After *Malacoctenus bimaculatus* (Steindachner), Insert:

2697 (b). **MALACOCTENUS PUERTORICENSESIS**, Evermann & Marsh.

Head 3.4; depth 3.4; eye 4; snout 3.5; maxillary 3.4; mandible 2.6; interorbital 7; preorbital 8; scales 4-44-8. D. xx, 10; A. II, 19; P. 14; V. 2; C. 13.

Body short, stout, compressed; head rather long, snout long and pointed; mouth small, little oblique, the maxillary scarcely reaching front of orbit; teeth in a single row in each jaw; gill-membranes broadly united, free from the isthmus; eyes high up, interorbital narrow; caudal peduncle short, compressed, its least depth about 3 in head. Fins rather large; origin of dorsal over upper end of gill-opening, first spine slightly shorter than second, which is somewhat longer than the third, whose length is about 2.2 in head; no notch behind third and fourth spines, all the spines from third to fifteenth being about equal in length, the sixteenth and seventeenth being somewhat shorter, the remaining three progressively longer; soft dorsal higher, its longest ray about 1.7 in head; longest anal ray 1.7; pectoral broad, 1.25 in head, reaching anal; ventral barely reaching origin of anal; a pair of slender oculal cirri, a small supraocular one, a short, slender, nasal cirrus and a few very slender ones at the nape; scales large, not crowded anteriorly; lateral line well arched above the pectoral.

Color in alcohol: Brown, much spotted and vermiculated with darker; top of head brown, sides and under parts pale, crossed by about 5 broad, irregular brown bars; side of body with about 5 or 6 broad, dark cross-bars, broader than the paler interspaces, broadest and darkest above and extending upon dorsal fin; under parts of body paler, more speckled; spinous dorsal with numerous small brown specks, a large, black ocellus on base of 3 anterior spines, and a larger one on base of last 4 dorsal spines, being chiefly on body; soft dorsal, caudal, and anal each crossed by several series of small brown spots; pectoral and ventrals pale, the pectoral with a few brown spots at base.

The above description from the type, a female, 2.5 inches long, obtained at Ilucares, February 14. Three female cotypes gotten at Fajardo, February 17, and one at Culebra, February 9, agree closely with the type; 2 of these, however, show faint traces of narrow horizontal lines along lower part of side.

A male, 2.5 inches long, from Culebra, February 11, taken as one of the cotypes, may be described as follows: Head 3.5; depth 3.7; eye 3.8; snout 3.2; maxillary 3.1; mandible 2.4; interorbital 7; preorbital 6.2; scales 3-45-9; D. xx, 10; A. II, 19; P. 14; V. 2; C. 13; longest dorsal spine 2 in head, longest ray 1.4; longest anal ray 1.5; pectoral 1; ventral 1.1; caudal 1.1. Color in alcohol, tolerably uniform brown; crossbars on side very faint; longitudinal lines more evident than in the female; throat and under parts of head mottled with white and light brown; fins less speckled than in female, the soft dorsal and anal pale, almost without spots.

Another male, 2.25 inches long, from Culebra, February 11, agrees with

the large specimen just described, except that the crossbars on body are more distinct.

This species most closely resembles *M. bimaculatus* Steindachner, from which it differs in the larger head, greater depth, smaller mouth, narrower interorbital, and in the color. The tips of the anal rays are not white, the soft dorsal is spotted like the caudal and anal, and there are no white spots on base of pectoral, as is said to be the case in *M. bimaculatus*.

Puerto Rico; known from the 7 specimens mentioned above, all obtained by the U. S. Fish Commission expedition to Puerto Rico.

Malacoctenus puertoricensis, EVERMANN & MARSH, Rept. U. S. Fish Comm. 1899 (Dec. 19, 1899), 358, Hucares, P. R. (Type, No. 49371, U.S.N.M. Coll. Evermann & Marsh.)

Page 2369. Before *Achenopterus*, Gillther, insert:

878 (a). *AUCHENISTIUS*, Evermann & Marsh.

Auchenistius, EVERMANN & MARSH, Rept. U. S. Fish Comm. 1899 (Dec. 19, 1899), 359 (*stahli*).

This genus has the form of *Achenopterus* and suggests that genus strongly. It differs in the absence of a lateral line, in the much smaller scales, in the absence of a notch at the front of the dorsal fin, and in the union of the membrane of the anal fin with that of the caudal.

(αὐχενίτης, nape; ἵστιος, sail or fin.)

2711 (a). *AUCHENISTIUS STAHLI*, Evermann & Marsh.

Head 5; depth 6.5; eye 4.8; snout 6; maxillary 2.8; scales about 58, about 12 in transverse series; D. XLI or XLII; A. II, 23 or 24; pectoral 2.5; ventral 2.2; caudal 1.3.

Body elongate, somewhat compressed, especially posteriorly, the dorsal and ventral outlines alike; head small, upper profile straight and descending; snout moderate, pointed; mouth large, the maxillary reaching to or beyond middle of eye; the jaws equal, heavy and projecting; teeth in lower jaw conical, short and strong, slightly recurved, in one row; teeth in upper jaw similar to those in lower, but a small patch of smaller teeth in front of jaw behind the main row; teeth on vomer; gill-membranes joined to the isthmus; nostrils with short tubes, a single flap above each eye and one on each side of nape; dorsal fin long, of spines only; last four spines somewhat longer than the preceding, forming a shallow notch, a feature lacking in the other examples; anal origin about midway between tip of snout and tip of caudal, the fin similar to dorsal in shape, but somewhat lower; membrane of dorsal and anal joined to caudal; caudal small, pointed; pectoral small, of 8 rays; ventral small, of 2 rays.

Color in spirits; Body everywhere with a very slight yellowish tinge, in some specimens a faded gray; one specimen has traces of 10 or 12 dark crossbars; fins all pale, in one case with the dorsal and anal dark-edged.

Puerto Rico; known from the type, a specimen 1.2 inches long, obtained by the U. S. Fish Commission expedition to Puerto Rico, February 1, 1899, at Ponce, and 13 cotypes, 8 from the coral and alge on the reefs at the mouth of Culebra harbor, and 5 from Puerto Real.

Name
commonities
sections of
Auchenistius
Ponce, P.

Page 23

Head 3.2
orbital 5.3
branchiost

Body rat
rate, not l
extending t
teeth on eac
teeth on vo
isthmus; e
fringed tu
dorsal spine
scarcely as
anteriorly;
one scale; I

Color: Un
brown, mot
upon anterie
of fin betw
darker, with
white at ba
pectoral bla
at base, the

This speci
opterus integ
species in th

Puerto Ri
obtained by
4, 1899.

(*albus*, wh
Achenopterus
360, Arroyo

271

Head 3.4;
scales 2-32-8

Body slen
above; sno
about reachi
small, conica

(named for Dr. A. Stahl, of Bayamon, Puerto Rico, who, under many difficulties put in his way by Spanish authorities, made important collections of natural history objects of that island.")

Auchenopterus stahli, EVERMANN & MARSH, Rept. U. S. Fish Com. 1899 (Dec. 19, 1899), 359,
1 once, P. R. (Type, No. 49372, U.S.N.M. Coll. Evermann & Marsh.)

Page 2373. Before *Auchenopterus fasciatus* (Steindachner), insert:

2717 (a). **AUCHEOPTERUS ALBICAUDUS**, Evermann & Marsh.

Head 3.2; depth 4; eye 4; snout 4.1; maxillary 2.2; mandible 1.6; interorbital 5.3; D. XXX, 1; A. II, 17; pectoral 1.4; ventral 1.5; caudal 1.6; branchiostegals 6; scales 1-34-6.

Body rather short, compressed; dorsal outline not elevated; head moderate, not broad; snout short, pointed; mouth large, oblique, maxillary extending to below middle of eye; lips broad, prominent; a band of conical teeth on each jaw, those on side somewhat enlarged and recurved; a patch of teeth on vomer, none on palatines; gill-membranes broadly united, free from isthmus; eye large, high up; nasal, supraocular, and nuchal regions with fringed tuft-like cirri; a considerable notch between fourth and fifth dorsal spines, but not reaching base of membrane; longest anterior spine scarcely as long as those of the posterior portion; scales large, reduced anteriorly; lateral line anteriorly separated from the dorsal fin by only one scale; head naked.

Color: Uniform dark brown on head and body, no dark crossbars; dorsal brown, mottled with lighter, narrowly edged with white; a black spot upon anterior 3 or 4 spines and a large black ocellus upon posterior portion of fin between twenty-second and twenty-fourth spines; anal rather darker, with narrow white edge; caudal peduncle black, the fin abruptly white at base, the entire fin being clear white, entirely without specks; pectoral black at base, then barred with white and dark; ventral black at base, the outer two-thirds barred with black and white.

This species seems to be related to the Pacific-coast species, *Auchenopterus integrifinnis*, which it closely resembles, but differs from that species in the larger scales, the deeper body, and the coloration.

Puerto Rico; known only from the type, a specimen 1.5 inches long, obtained by the U. S. Fish Commission expedition to Puerto Rico, February 4, 1899.

(*albus*, white; *cauda*, tail.)

Auchenopterus albicaudus, EVERMANN & MARSH, Rept. U. S. Fish Com. 1899 (Dec. 19, 1899),
360, Arroyo, P. R. (Type, No. 49373, U.S.N.M. Coll. Evermann & Marsh.)

2717 (b). **AUCHEOPTERUS RUBESCENS**, Evermann & Marsh.

Head 3.4; depth 5; eye 5; snout 3.8; maxillary 2.6; interorbital 5.8; scales 2-32-8; D. XXX, 1; A. II, 18; pectoral 1.5; ventral 2; caudal 1.4.

Body slender and compressed; head moderate, somewhat compressed above; snout pointed; mouth moderate, the jaws equal, the maxillary about reaching front of pupil; lips, especially the upper, prominent; teeth small, conical and sharp, in both jaws, in a numerous patch on front of

upper jaw, fewer on sides; in lower jaw less numerous in front, a long single row of somewhat stronger teeth on sides; eye not large; a small nasal flap, and a 3- or 4-branched tentacle over eye and one at nape; scales rather large and regularly arranged; dorsal fin with a notch behind third spine, and with one unbranched soft ray at its end, the membrane joined to caudal; origin of anal under eleventh dorsal spine; lateral line as usual in *Auchenopterus*.

Color in spirits: Everywhere a nearly uniform faded pink, save breast and lower side of head, which are paler; a small, inconspicuous dark round spot on dorsal fin, at twenty-third and twenty-fourth spines, a little nearer base than margin, and made up of very small black punctulations; indications of a yellow tinge on front of dorsal and base of anal in life; fins otherwise all pale.

Puerto Rico; known only from the type, a specimen 1.3 inches long, obtained by the United States Fish Commission expedition to Puerto Rico January 27, 1899. (*rubeocens*, reddening.)

Auchenopterus rubeocens, EVERMANN & MARSH, Rept. U. S. Fish Comm. 1899 (Dec. 19, 1899), 360. Puerto Real, P. R. (Type, No. 49374, U.S.N.M. Coll. Evermann & Marsh.)

2717 (e). **AUCHENOPTERUS CINGULATUS**, Evermann & Marsh.

Head 3; depth 4.4; eye 5; snout 4.2; maxillary 2.2; interorbital 6; scales 2-29-7; D. IV-XXIV, the longest spines 3 in head; A. II, 16, the longest ray 2.25 in head; pectoral 1.3; ventral 1.8; caudal 1.6. Body rather long and slender, strongly compressed; head large, little compressed; snout moderately sharp; mouth large, maxillary reaching posterior border of eye, the lips heavy, the jaws subequal or the lower very slightly projecting; teeth conical and sharp, in more than one row in each jaw, most numerous in front; a patch on vomer; a nasal filament, a 3- or 4-branched supraocular tentacle, and a 4-branched nuchal tentacle, the branches of the latter each with a dark dot on their anterior surface. Dorsal originating over edge of preopercle, of spines only, the second slightly longer than first; second, third, and fourth graduated, the fourth comparatively short, thus forming a notch partly separating the first 4 spines from rest of fin; dorsal membrane joined low with caudal; anal free from caudal, about as high as dorsal, its thirteenth and fourteenth rays longest; first anal spine under tenth or eleventh dorsal spine; caudal rounded, shorter than head, of about 13 rays; pectoral large, reaching anal, of 12 rays; ventrals moderate, of 2 rays, the spine not evident. Lateral line running high to eleventh dorsal spine, here abruptly decurved two rows of scales, thence median to base of caudal.

Color in spirits: Body and head pale yellow; body with 4 heavy dark-brown vertical bars, each about 4 rows of scales wide, extending on the vertical fins; membrane of anterior dorsal spines, opercle, occipital, and scapular region blotched with the same color; a dark bar backward and downward from eye across cheek, rather more than one-half width of eye; top of head between and behind eyes darkened; preorbital, maxillary, lips, and under part of head thickly punctulate with dark; dorsal and anal barred with the extensions of the wide dark body bars, and with the

attenuating with gray sharply separated half of pectoral; 1 the rest bar

A pretty where 1 sp Puerto Rico
(*cingulatus*)
Auchenopterus
361, Ponce

Head 3.27
interorbital
A. II, 17; p.
compressed
the long an
orbic; jaws
front, becom
and weaker
tentacles pr
edge of pred
forming a m
visible unde
anal origin m
line; pectoral
the innermost

Color in sp
posteriorly;
the fins, thei
ward and ba
opercle; max
dark; lower
spots on edge
the extension
dorsal; a dist
fourth dorsa
ground color
lings on the
space without
bar; pectoral

A handsome
long, collecte
dition to Pue

Auchenopterus f
361, Fajardo

and having narrower pale interspaces; caudal mottled or irregularly barred with grayish, its base with the plain pale-yellow ground color, which is sharply separated from the rest of the fin by a curved dark line; posterior half of pectoral barred with dark formed of dots on the rays, the first bar plainest; basal half of pectoral pale; ventral with basal portion dark, the rest barred like pectoral.

A pretty and strongly marked blenny, known only from Puerto Rico, where 4 specimens were obtained on the coral reefs at Ponce and 1 at Puerto Real by the U. S. Fish Commission expedition to Puerto Rico.

(*cinctus*, banded, from the conspicuous vertical bars.)

Auchenipterus cinctus, EVERMANN & MARSH, Rept. U. S. Fish Com. 1899 (Dec. 19, 1899), 361. Ponce, P. R. (Type No. 49375 U.S.N.M. Coll. Evermann & Marsh.)

2717 (a). **AUCHEOPTERUS FAJARDO**, Evermann & Marsh.

Head 3.25; depth 4.8; eye 4.2; snout 4.8; maxillary 1.7; mandible 1.5; interorbital 5.5; scales 2-37-8; B. XXIX, 1, the longest spine 2.3 in head; A. II, 17; pectoral 1.4; ventral 1.7; caudal 1.4. Body elongate, strongly compressed posteriorly; head moderate, little compressed; mouth large, the long and slender maxillary reaching beyond the posterior border of orbit; jaws subequal; teeth of upper jaw conical and sharp, in a patch in front, becoming one row posteriorly; teeth in lower jaw similar, but fewer and weaker; vomerine teeth in two series. Nasal, ocular, and nuchal tentacles present, all but the nasal about 5-branched. Dorsal origin over edge of preopercle, the first 4 spines graduated, the fourth shortest, thus forming a notch; dorsal ending with an unbranched soft ray, the joints visible under a strong lens; membrane of dorsal joined low to caudal; anal origin under eleventh dorsal spine and the decurved portion of lateral line; pectoral reaching past front of anal; ventral moderate, of 3 rays, the innermost shorter and slenderer.

Color in spirits: Body and head light reddish, becoming a little paler posteriorly; body with traces of 6 or 8 dark vertical bars extending on the fins, their margins ill defined; breast pale, 2 dark reddish bars downward and backward from eye across upper and lower edge of cheek to opercle; maxillary blotched with dark; upper lip and tips of both jaws dark; lower part of head spotted with dark; a row of about 5 small dark spots on edge of preopercle; iris pink; dorsal and anal fins gray, except for the extensions of the dark bars of the body and a few white spots on the dorsal; a distinct ocellus on the twenty-second, twenty-third, and twenty-fourth dorsal spines and their membranes; base of caudal gray, like the ground color of dorsal and anal; posterior part of caudal with gray mottlings on the rays only, this portion separated from the basal part by a space without pigment on rays or membrane, making a distinct vertical bar; pectoral and ventrals mottled. Puerto Rico.

A handsomely colored blenny, known only from the type, 1.63 inches long, collected February 17, 1899, by the U. S. Fish Commission expedition to Puerto Rico. (Named for the type locality.)

Auchenipterus fajardo, EVERMANN & MARSH, Rept. U. S. Fish Com. 1899 (Dec. 19, 1899), 361. Fajardo, P. R. (Type, No. 49376, U.S.N.M. Coll. Evermann & Marsh.)

Page 2400. Before *Ophioblennius*, Gill, insert:

893 (a). CORALLIOZETUS, Evermann & Marsh.

Coralliozetus, EVERMANN & MARSH, Rept. U. S. Fish Com. 1899 (Dec. 19, 1899), 362 (*cardone*).

Body slender and strongly compressed, without scales; head large, subcylindrical, bluntly pointed; mouth large; teeth not hooked, about 8 enlarged conical ones in front of each jaw, smaller ones behind; vomer with teeth; dorsal fin with a notch between the rays and spines, the membrane connected with caudal; caudal fin rounded; pectoral large, ventrals small and inserted slightly in advance of pectorals. A strongly marked genus, conspicuous in appearance by its heavy head and thin body, probably related to *Ophioblennius*, from which it is technically separated by the absence of hooked canine teeth, the convex caudal, and the entire absence of a lateral line.

(κοράλλιον, coral; ζητέω, I seek.)

2753 (a). CORALLIOZETUS CARDONE, Evermann & Marsh.

Head 4; depth 5.6; eye 5; snout 4; maxillary 2; D. XVII, 11; A. 21; pectoral 1.3; ventral 1.8; caudal 1.4.

Body scaleless, slender, much compressed; head large and heavy, not compressed nor depressed; snout very short and blunt; mouth large, horizontal, low in position, the maxillary reaching far beyond the eye; eyes small, close together, placed high and well forward; teeth conical, in a patch on the front of each jaw, an outer row of about 8 teeth (4 on a side) in each jaw, much enlarged; a single row of smaller teeth on sides of each jaw; teeth on vomer; a small flap at the nostril and two short filaments above eye, one much the smaller; no appendages at the nape. Dorsal fin long and high, of slender, flexible spines, and longer soft rays, a notch between the soft and spinous portions; anal longer and lower than soft dorsal; anal and dorsal free from caudal; caudal rounded; pectoral large, wide as body, reaching anal or beyond; ventral small, inserted before pectoral, of 3 rays, the innermost very slender.

Color in spirits: Body dark red, much paler in one specimen; head everywhere inky, this color dusted upon the body, particularly on the anterior portion; a pale gray bar downward and backward across cheek; fins pale, except ventrals and front of dorsal, which have color of head; a row of small rosy spots along bases of anal rays, seemingly in the flesh; sometimes a similar fainter row along base of dorsal. Known only from Puerto Rico, where 3 specimens, each about 1 inch long, were obtained in February, 1899, by the U. S. Fish Commission expedition to that island.

(Named for Cardona, a little islet off Playa de Ponce, on the reef of which the type was collected.)

Coralliozetus cardone, EVERMANN & MARSH, Rept. U. S. Fish Com. 1899 (Dec. 19, 1899), 362, *Cardona Islet, P. R.* (Type, No. 49377, U.S.N.M. Coll. Evermann & Marsh.)

Page 2
Index

Head ne
endai, 15
naked; ve
jaws with
trace in it
closely pl:
a curve to
base of an
it ceases.
another la
zontally a
pores here
only by el
each pore
North Atla
Strait.

*Lycodes mure
land, Nor
Lyconchelys n*

Page 257
in Denmark

Page 258

The head
or frontal su
wardly direc
spinose osse
close before
tinned both
prolonged fa
corners of t
middle of t
breadth of t
jaws. The l
isthmus, and
surface. Th
rows of slim
high and its
second dorsa
followed unl
The pectoral
to a fine thre
It may also

Bull

Page 2471. After *Lycenchelys parillus* (Goode & Bean), insert the following:

2833 (a). **LYCENCHELYS MURENA** (Collett).

Head nearly 8; depth about 20; eye 4.5; D. 112 (including one-half of anal, 118); A. 95 (101); P. 13. Body everywhere scaled, head and fins naked; ventrals comparatively close together, 6 times length of pectoral; jaws with teeth in a single series. Lateral line present, but difficult to trace in its entirety; from upper angle of gill-opening a series of rather closely placed, very small pores (connected by a narrow line) descends in a curve to a short distance behind vent; after having reached nearly to base of anal, it runs for a short distance nearly horizontally, whereupon it ceases, or can not be followed in the same direction with certainty; another lateral line, however, begins about over the vent, and runs horizontally along median line, becoming obsolete toward end of tail; the pores herein are very small and more widely separated, and can be seen only by close observation; the short whitish line which runs through each pore forms here no accompanying line. Length 7 to 11 inches. North Atlantic. Recently taken by the *Ingolf* expedition in Denmark Strait.

Lycodes murena, COLLETT, Lorph. Selsk. Christ., No. 14, 15, 1878, off Traenen, in Helgeland, Norway; Lütken, *Ingolf* Expedition, 20, 1898.
Lycenchelys murena, GOODE & BEAN, Oceanic Ichth., 309, 1896.

Page 2578. *Chalinura simula*, Goode & Bean, was taken by the *Ingolf* in Denmark Strait in 912 to 1,236 fathoms.

Page 2587. Before *Caelorhynchus*, Giorna, insert the following:

2963 (a). **MACROURUS INGOLFI**, Lütken.

The head is contained about 5 times in the total length. The superior or frontal surface of the snout is separated from the inferior or more forwardly directed part by a well-developed crest or edge, terminating in 3 spinose osseous tubercles, 1 directly in the middle and 1 on either side close before the naked spot, where the nostrils have their place, and continued both above and below the orbits. The broadly triangular snout is prolonged fairly over and before the mouth, which is relatively little, the corners of the mouth falling in a line with the anterior margin of the middle of the orbits. The eyes are large, their diameter surpassing the breadth of the front between the eyes. The teeth form a fine card in both jaws. The head is scaled with the exception of the gill-membrane, the isthmus, and its foremost superior margin, and almost the whole lower surface. The naked part of the snout is handsomely embroidered with rows of slime glands. The first dorsal, counting 11+9 rays, is singularly high and its longest (second) ray is serrate and as long as the head. The second dorsal begins much forward, its foremost rudimentary rays may be followed until not far from the posterior margin of the first dorsal fin. The pectoral contains 20 rays, and the ventrals, whose external ray tapers to a fine thread and reaches a long stretch beyond the anus, have 8 rays. It may also be remarked that the tail, as in other Macrurans, is really

pointed behind, but in several specimens has lost a shorter or longer part; but the wound has healed, and on the thus truncated point of the tail is developed a distinct caudal fin, a phenomenon which is also observed in some specimens of the preceding species. The scales show distinct rows of thorns, not however so much projecting as in *M. goodei*. (Lütken.)

Several specimens taken by the *Ingolf* about Iceland and in Denmark Strait.

Macrurus ingolfi, LÜTKEN, Ichth. Results Danish *Ingolf* Exped., 27, 1898, south of Iceland.

Page 2622. After *Hippoglossina bollmani*, Gilbert, insert the following:

2988 (a). HIPPOGLOSSINA SABANENSIS, Boulenger.

Head 4; depth 1 $\frac{1}{4}$; eye 5; width of mouth 2 $\frac{1}{4}$ in head; maxillary reaching posterior third of eye; teeth small, more strongly developed on right side; eyes moderately large, the lower a little in advance of the upper; interorbital space tectiform, scarcely half the width of eye. D. 76; A. 63. Gill-rakers very short, tuberculous, 6 on lower arm of first gill-arch; dorsal commencing above anterior third of upper eye, its longest ray 2 in head; left pectoral a little longer than right, 1 $\frac{1}{2}$ in head; left ventral twice as long as right, equaling the head; caudal a little shorter than head, its free edge forming a wide angle, the middle rays more than half longer than the outer; depth of caudal peduncle twice its length; scales 33-110-40, strongly ciliate on both sides; lateral line strongly arched above the pectoral. Deep gray brown; 3 large black ocelli bordered with bluish, 2 superposed on middle of body, equally distant, between lateral line and vertical fin, the third on lateral line a little in front of caudal peduncle; pectoral yellowish, ventrals blackish. Length 230 mm. One specimen from Rio Sabana, Darien. (Boulenger.) (*sabanensis*, from the Rio Sabana.)

Hippoglossina sabanensis, BOULENGER, Boletino del Mus. di Zool. ed. Anat. Comp. della Univ. di Torino, Vol. XIV, No. 346, 4, April 29, 1899, Rio Sabana, Darien. (Coll. Dr. Eurico Festa.)

Page 2789. *Villarius*, Rutter=Hansstor, Jordan & Evermann. All species of *Ameiurus* have villi.

Page 118. Dr. Albert Günther has recently critically studied the Linnean type specimens of American fishes in the possession of the Linnean Society of London, and the results are published in the proceedings of that society for May 24, 1899.

These studies of Dr. Günther necessitate the following changes from the names adopted by us in the earlier parts of this work: The Gaff-Top-sail Catfish becomes

161. FELICHTHYS FELIS (Linnaeus).

The species called by us *Hexanemichthys felis* (Linnaeus) (p. 128) becomes

186. GALEICHTHYS MILBERTI (Cuvier & Valenciennes).

The Red Hind, given by us as *Epinephelus maculosus* (p. 1158), becomes

The 2¹

Page 2

Head 3.
space 2.66
anal 5.75;
line 33; be
10; D. 8; A
dorsal; ca
oblique; j
nostril. T
smallest to
short, with
notably de
as posterio
fourth ray
ing to ven
forked, the

Color silv
end of caud
broader and
band on me
late, except
part of can
edges; dors

This speci
in length of
a little less
with *Notrop*
species may
body, short

Lake Chap

Notropis calien
at Aguas C

Page 965.
1797 for a ge
place in fish

1553. EPINEPHELUS GUTTATUS (Linnaeus).

The spotted Jewfish, given as *Promicrops guttatus* (p. 1162), becomes

1557. PROMICROPS ITAIARA (Lichtenstein).

Page 261. After *Notropis heterodon* (Cope), insert:

406 (a). NOTROPIS CALIENTIS Jordan & Snyder.

Head 3.66 in length; depth 3.33; eye 4 in head; snout 4; interorbital space 2.66; depth of caudal peduncle 2.40; height of dorsal 4.66 in length; anal 5.75; length of pectoral 5.66; ventral 6; caudal 3.83; scales in lateral line 33; between dorsal and occiput 15; between dorsal and ventral fins 10; D. 8; A. 7. Body deep and wide, deepest part anterior to insertion of dorsal; caudal peduncle long and slender; snout blunt, rounded; mouth oblique; jaws equal; maxillary reaching to a vertical from posterior nostril. Teeth 4-4, slender, hooked, grinding surface narrow, absent on smallest tooth. Gill-rakers short, blunt; 9 on first arch. Intestinal canal short, with but 1 convolution; peritoneum white, scales large, not notably deeper than long; lateral line incomplete, extending only as far as posterior edge of pectoral. Dorsal fin rounded, the second, third, and fourth rays longest; anal fin similar in shape; ventral rounded, extending to vent, inserted directly under dorsal; pectoral rounded; caudal forked, the lobes rounded.

Color silvery, an indefinite dark band extending from tip of snout to end of caudal peduncle; the band distinct on snout and caudal peduncle, broader and almost obsolete on middle of body; top of head and a narrow band on median dorsal part of body dark; ventral parts of body immaculate, except a narrow, dark band extending along base of anal and ventral part of caudal peduncle; chin white; scales on upper parts with dark edges; dorsal and caudal fins slightly dusky; other fins lighter.

This species is of small size, the mature males measuring about 34 mm. in length of body; the females 43. The males are more slender and have a little less dark color on the body. The affinities of *Notropis calientis* are with *Notropis cayuga*, *jordani*, and others of the subgenus *Chriope*. The species may be distinguished by its small eye, short, rounded snout, deep body, short lateral line, and rounded fins. (Jordan & Snyder.)

Lake Chapala basin, Mexico.

Notropis calientis, JORDAN & SNYDER, Bull. U. S. Fish Com. 1890 (1900), 122, Rio Verde, at Aguas Calientes, Mexico. (Type, No. —, L. S. Jr. Univ. Mus. Coll. J. O. Snyder.)

Page 965. The generic name *Rhombus* was first used by Humphreys in 1797 for a genus of mollusks, and can not therefore be used in fishes. Its place in fishes will be taken by *Peprilus* Cuvier (1817).

The fol
the "Add
index con

agassizii, Choi
Agosia klamath
albicaudus, A
album, Chirost
Algansea lucus
Alutera monochro
Amblyopsis ...
arge, Eslopsar
arundelii, Gob
aubeanaebei, E
Anchenistius .

^s
Auchenopterus

Bairdiella miaca
baldwini, Prion
balsanus, Istlar
bathecetor, Salu
bayamonensis, C
Blennicottus gle

Bollmannia boqu
boqueronensis, I
breve, Chirostom
caguila, Sleydin
Calamus kendal
caliente, Xenend
calientis, Notrop
cardona, Corallie
carpintis, Nectro
cayuga, Cottus g
Centridermichth

Centrolophidae ...
chapala, Chirost
Falcula
Characodon enea

INDEX TO NAMES OF FISHES MENTIONED IN THE
"ADDITIONAL ADDENDA" IN PART IV.

The following index includes only those names which occur in the "Additional Addenda" found in the present volume. An index complete for the three other parts is given in Part III.

Page.		Page.	
agassizii, Chologaster	3156	Chiostoma album	3165
Agosia klamathensis	3144	breve.....	3157
albicaudus, Auchenopterus.....	3191	chapale	3159
album, Chiostoma	3165	crystallinum	3162
Algansea lacustris.....	3140	diazi.....	3161
Aluterus monoceros	3178	estor.....	3165
Amblyopsis	3157	humboldtianum	3157
arge, Eslopsarum.....	3158	lerme	3163
arundelii, Gobius.....	3184	ocotlane	3163
aubenaupei, Etheostoma.....	3167	prouelas	3160
Auchenistius	3190	Chologaster	3156
stahli	3190	agassizii.....	3156
Auchenopterus albicaudus	3191	cornutus	3156
cingulatus.....	3192	papilliferus.....	3156
fajardo	3193	Cichlasoma steindachneri	3173
rubescens	3191	cingulatus, Auchenopterus	3192
Bairdiella miacantha	3173	clarkii declivifrons, Salmo.....	3147
baldwini, Prionodes	3168	jordani, Salmo	3148
balssami, Istiurus	3139	Coralliozetus.....	3194
batheecotor, Salmo.....	3149	cardone	3194
bayamonensis, Gobius	3184	cornutus, Chologaster	3156
Blennioccottus globiceps	3179	Cottus gracilis cayuga.....	3178
recalvus.....	3178	crystallinum, Chiostoma.....	3162
Bollmannia boqueronensis	3185	culebrae, Malacoctenus.....	3187
boqueronensis, Bollmannia	3185	Cyclothonome megalops	3150
breve, Chiostoma	3157	declivifrons, Salmo clarkii.....	3147
caguata, Sleydium	3183	decoris, Doratonotus	3177
Calamus kendalli	3172	Dialarchus.....	3180
caliente, Xenendum	3152	nyderi	3181
calientis, Notropis.....	3197	diazi, Chiostoma.....	3161
cardona, Coralliozetus	3194	digneti, Neomugil	3165
carpinti, Neetroplus	3175	Doratonotus decoris.....	3177
cayuga, Cottus gracilis	3178	encaustus, Characodon	3150
Centridermichthys globiceps	3179	Epinephelus guttatus.....	3197
maculosus	3182	maculosus.....	3197
Centrolophidae	3166	Eslopsarum arge	3158
chapale, Chiostoma.....	3159	jordani	3157
Falcula.....	3143	estor, Chiostoma.....	3165
Characodon encaustus.....	3150	Etheostoma aubenaupei	3167

Page.	Page.
<i>Eucalia inconstans</i>	3157
<i>Eupomacentrus flavilatus</i>	3176
<i>rectifrenum</i>	3176
<i>Eximia</i>	3182
<i>rubellio</i>	3182
<i>Falcula</i>	3143
<i>chapale</i>	3143
<i>fajardo, Auchenopterus</i>	3193
<i>Felichthys felis</i>	3197
<i>felis, Felichthys</i>	3197
<i>festae, Piabucina</i>	3145
<i>Tachysurus</i>	3138
<i>flavilatus, Eupomacentrus</i>	3176
<i>Galeichthys labiatus</i>	3137
<i>milberti</i>	3197
<i>garmani, Stolephorus</i>	3146
<i>Gasterosteus gymnetes</i>	3157
<i>gilberti, Stolephorus</i>	3146
<i>Gilbertidia</i>	3183
<i>Gilbertina</i>	3183
<i>Gillias</i>	3186
<i>jordani</i>	3186
<i>globiceps, Blennicottus</i>	3179
<i>Centridermicthys</i>	3179
<i>Oligocottus</i>	3179
<i>Gobius arundelii</i>	3184
<i>bayamonensis</i>	3184
<i>soporator</i>	3184
<i>gracile, Myctophum</i>	3150
<i>gracilis cayaga, Cottus</i>	3178
<i>Uranidea</i>	3178
<i>guttatus, Epinephelus</i>	3197
<i>Promicrops</i>	3197
<i>gymnetes, Gasterosteus</i>	3157
<i>Hadropterus maxinkuckiensis</i>	3166
<i>Haemulon belenea</i>	3171
<i>Haustor</i>	3196
<i>helenae, Haemulon</i>	3171
<i>Heros istlarius</i>	3174
<i>Hippoglossina sabanensis</i>	3196
<i>bumboldtianum, Chirostoma</i>	3157
<i>inconstans, Eucalia</i>	3157
<i>ingolfi, Macrourus</i>	3195
<i>istlarius, Heros</i>	3174
<i>Istlarius</i>	3138
<i>balsanus</i>	3139
<i>itaiai, Promicrops</i>	3197
<i>jordani, Eslopsarum</i>	3157
<i>Gillias</i>	3186
<i>Lycodontis</i>	3145
<i>Salmo clarkii</i>	3148
<i>kendalli, Calamus</i>	3172
<i>klamathensis, Agosia</i>	3144
<i>labiatus, Galeichthys</i>	3137
<i>labraciforme, Pomadasys</i>	3172
<i>lacustris, Algansca</i>	3140
<i>lalandi, Seriola</i>	3166
<i>lermei, Chirostoma</i>	3163
<i>limantouri, Poccilia</i>	3153
<i>luitpoldii, Xenendum</i>	3152
<i>Lycenchelys muræna</i>	3195
<i>Lycodontis jordani</i>	3145
<i>Macdonaldia rostrata</i>	3150
<i>Macrourus goli</i>	3195
<i>maculosus, Centridermicthys</i>	3182
<i>Epinephelus</i>	3196
<i>Oligocottus</i>	3182
<i>Malacoctenus cultrifer</i>	3167
<i>moorei</i>	3188
<i>puertoricensis</i>	3189
<i>maxinkuckiensis, Hadropterus</i>	3166
<i>meeki, Microgobius</i>	3185
<i>megalops, Cyclothona</i>	3150
<i>miacantha, Bairdiella</i>	3173
<i>Microgobius meeki</i>	3185
<i>milberti, Galeichthys</i>	3196
<i>monoceros, Aluterus</i>	3178
<i>montezumae, Xiphophorus</i>	3154
<i>moorei, Malacoctenus</i>	3188
<i>muraena, Lycenchelys</i>	3195
<i>muskoka, Notropis</i>	3141
<i>Myctophum gracile</i>	3150
<i>Nebris occidentalis</i>	3173
<i>zestus</i>	3173
<i>Neotroplus carpintis</i>	3175
<i>Neomugil</i>	3165
<i>digneti</i>	3165
<i>nephelus, Spherooides</i>	3178
<i>Tetronodon</i>	3178
<i>Notropis eulentis</i>	317
<i>muskoka</i>	3141
<i>rasconis</i>	3141
<i>occidentalis, Nebris</i>	3173
<i>octolane, Chirostoma</i>	3163
<i>Oligocottus globiceps</i>	3179
<i>maenelosus</i>	3182
<i>snyderi</i>	3182
<i>Orcella</i>	3140
<i>Orcula</i>	3140
<i>papilliferus, Chologaster</i>	3156
<i>Peprius</i>	3197
<i>Piabucina festae</i>	3145
<i>Poccilia limantouri</i>	3153
<i>Pomadasys labraciforme</i>	3172
<i>popoche, Xystrosus</i>	3142
<i>Prionodes baldwini</i>	3168
<i>promelas, Chirostoma</i>	3160
<i>Promicrops guttatus</i>	3197
<i>itaiai</i>	3197
<i>puertoricensis, Malacoctenus</i>	3189
<i>rasconis, Notropis</i>	3141
<i>recalvus, Blennicottus</i>	3178

rectifrenum
Rhegma
Rhombus
rimensis, R.
rose, Trogl.
Typh.
rostrata, Ma.
rubellio, Ex.
rubescent, A.
Ruscarius, I.
sabanensis, I.
Salmo batho
clarki
Seriola laland
Sicydium cap
snyderi, Dial
Oligo
soporator, Ge
spengleri, Te
Spherooides no
stabli, Anche
steindachneri

	Page.		Page.
rectirecum, <i>Eupomacentrus</i>	3176	<i>Stolephorus garmani</i>	3146
<i>Rhegma</i> *	3169	<i>gilberti</i>	3146
<i>thaumasium</i> *	3170	<i>Stromateidae</i>	3166
<i>Rhombus</i>	3197	<i>subterraneus</i> , <i>Typhlichthys</i>	3156
<i>riniensis</i> , <i>Rusciculus</i>	3179	<i>Tachysurus festae</i>	3138
<i>rosie</i> , <i>Troglichthys</i>	3156	<i>Tetronotus nephelus</i>	3178
<i>Typhlichthys</i>	3156	<i>speigleri</i>	3178
<i>rostrata</i> , <i>Macdonaldia</i>	3150	<i>thaumasium</i> , <i>Rhegma</i>	3170
<i>rubellio</i> , <i>Eximia</i>	3182	<i>Troglichthys</i>	3156
<i>rubescens</i> , <i>Auchenopterus</i>	3191	<i>rosie</i>	3156
<i>Rusciculus</i>	3179	<i>Typhlichthys</i>	3157
<i>rimensis</i>	3179	<i>rosie</i>	3156
<i>sabanensis</i> , <i>Hippoglossina</i>	3196	<i>subterraneus</i>	3156
<i>Salmo bathaeceptor</i>	3149	<i>Urandea gracilis</i>	3178
<i>clarkii declivifrons</i>	3147	<i>Villarius</i>	3196
<i>jordani</i>	3148	<i>xaliscone</i> , <i>Xenendum</i>	3153
<i>Seriola lalandi</i>	3166	<i>Xenendum</i>	3151
<i>Sicydium euguitae</i>	3183	<i>caliente</i>	3152
<i>snyderi</i> , <i>Dialarchus</i>	3181	<i>luitpoldii</i>	3152
<i>Oligocottus</i>	3182	<i>xaliscone</i>	3153
<i>soporator</i> , <i>Gobius</i>	3184	<i>Xiphophorus montezumae</i>	3154
<i>spengleri</i> , <i>Tetronotus</i>	3178	<i>Xystrosus</i>	3142
<i>Sphaeroides nephelus</i>	3178	<i>popoche</i>	3142
<i>stabili</i> , <i>Auchenistius</i>	3190	<i>zestus</i> , <i>Nebris</i>	3173
<i>steindachneri</i> , <i>Cichlasoma</i>	3173		

*These names occur as new in this volume.

Abeona mini
Ahomia ethoe
Abramis crys
Abraedilus si
acanthias. Squ
Acantharchus
acepiter. Pod
Achirus fasci
lineatus
Acipenser bre
rus
tra
acipenserinus
ackleyi. Rej
Acrocheilus a
Acrotus willo
aculeatus. Ga
Ste
acus. Tylosur
acutus. Fodiator
Adinla dusgesi
adscensionis.
adspersus. Ta
eglefinus. Ta
eneus. Myoxo
asculapins. A
estivalis mare
Poma

.Ethoprorae luc
.Etobatus mari
affinis. Atherin

Cheilosid
Chimene
Gambus
agassizii. Alepo

Callionymus
Dierc

Lipan
Xenia

aggregatus. Cy
Agonostomus n
Agosia falcati

khamaathi
umatilla

Aldia egmonti
alabama. Alosa

abohunga. Germa
alus. Priomotu

albigutta. Katho
albolineatus. Fr

albus. Haemulo
Alcidea thoburni

Aldrovandia gr
ma

Alepisaurus ase
fer

Alepocephalus a

Aleposomus co

INDEX TO GENERA AND SPECIES ILLUSTRATED.

Genera and species.	Plate and figure.	Text page.
<i>Abeona minima</i>	CCXXVIII, 578	1497
<i>Ahoma ethostoma</i>	CCXXXVIII, 702	3240
<i>Abramis crysoleucus</i>	XLV, 111	259
<i>Abroedafus saxatilis</i>	CCXXXIV, 590	1561
<i>acanthias, Squadius</i>	VII, 24, 24a	54
<i>Acantharchitis pomotis</i>	CLV, 418	089
<i>acipecter, Podoliceus</i>	CCCVIII, 745, 745a	2055
<i>Achirus fasciatus</i>	CCCLXXXVII, 948	2700
<i>lineatus</i>	CCCLXXXVI, 947	2697
<i>Acipenser brevirostrum</i>	XXI, 47	106
<i>rubieundus</i>	XXI, 46	106
<i>sturio oxyrhynchus</i>	XX, 45	105
<i>transmontanus</i>	XX, 44	104
<i>acipenserinus, Podotherina</i>	CCCIX, 746	2061
<i>ackleyi, Raja</i>	X, 31	70
<i>Acrocheilus alutaceus</i>	XXXIX, 97	208
<i>Acorus willoughbyi</i>	CLI, 408	973
<i>aculeatus, Gasterosteus</i>	CNIX, 320	747
<i>Stenotomus</i>	CCXII, 545	1346
<i>acs., Tylosurus</i>	CXVI, 309	716
<i>acutus, Fodiator</i>	CXVII, 315	728
<i>Adinia dugesii</i>	CVIII, 290	661
<i>adscensionis, Epinephelus</i>	CLXXXII, 482	1152
<i>adspersus, Tantogolabrus</i>	CCXXXVI, 595	1577
<i>egleinus, Melanogrammus</i>	CCCLX, 892, 892a	2542
<i>anemus, Myoxocephalus</i>	CCXCV, 715, 715a	1972
<i>asclepias, Alepisaurus</i>	XCVI, 258	595
<i>festivus marconis, Hybopsis</i>	LII, 136	318
<i>Pomolobus</i>	LXXI, 190	426
<i>Ethophrora lucida</i>	XCI, 246	565
<i>Etopterus marinari</i>	XV, XVI, 37, 37a	88
<i>affinis, Atherinops</i>	CXXXVI, 342	807
<i>Cheilodipterus</i>	CXXXIX, 471	1113
<i>Chimara</i>	XIX, 40	95
<i>Gambusia</i>	CXIII, 299, 299a	680
<i>agassizii, Alepocephalus</i>	LXXIV, 197	452
<i>Callionymus</i>	CCXXXIII, 779	2186
<i>Dicromita</i>	CCCLV, 874	2516
<i>Liparis</i>	CCXXVII, 765	2121
<i>Xenichthys</i>	CCIII, 527	1287
<i>aggregatus, Cymatogaster</i>	CCXXVIII, 579a	1498
<i>Agonostomus monticola</i>	CXXVII, 347	819
<i>Agosia falcatata</i>	LII, 135	313
<i>klunzatherensis</i>	LII, 133
<i>umatilla</i>	LII, 134	313
<i>Ahlia egmontis</i>	LX, 158	370
<i>alabama, Alosa</i>	LXXII, 192, 192a	2810
<i>albulungu, Germa</i>	CXXXIV, 367	871
<i>alatus, Prionotus</i>	CCCXIX, 770	2159
<i>albigutta, Kathetostoma</i>	CCXXXIV, 809, 809a	2312
<i>albolineatus, Fundulus</i>	CV, 281	649
<i>Albula vulpes</i>	LXVIII, 179	411
<i>albua, Hamulon</i>	CCIII, 528	1295
<i>Alcidea thoburni</i>	CCLXXXIII, 684	1887
<i>Aldrovandia gracilia</i>	XCVIII, 263	610
<i>macrochir</i>	XCVIII, 262	609
<i>Alepisaurus aesculapius</i>	XCVI, 258	595
<i>ferox</i>	XCV, 257	595
<i>Alepocephalus agassizii</i>	LXXIV, 197	453
<i>Aleposomus copei</i>	LXXV, 199	450

Genera and species.	Plate and figure.	Text page.
<i>aleutensis</i> , <i>Lyconectes</i>	CCCXLVI, 844	244
<i>aleutensis</i> , <i>Cottus</i>	CCXCVI, 711	1957
<i>Alexurus armiger</i>	CCXXXV, 784	2203
<i>Algansen dugesi</i>	XL, 98	211
<i>alletterata</i> , <i>Gymnosarda</i>	CLXXXIV, 366	839
<i>Allopias vulpes</i>	VI, 20	45
<i>Alosa alabamae</i>	LXXII, 192, 192a	280
<i>sapidissima</i>	LXXII, 191	247
<i>Alphesterochloropterus</i>	CLXXXVI, 488, 488a	1161
<i>alpinus aurivulus</i> , <i>Salvelinus</i>	LXXXIII, 220	511
<i>artivellis</i> , <i>Sebastolobus</i>	CLLXIX, 653	1763
<i>altus</i> , <i>Hybopsis</i>	LIV, 138	321
<i>Pseudopriacanthus</i>	CXCV, 512	1239
<i>alutaceus</i> , <i>Aerocheilus</i>	XXXIX, 97	208
<i>Alutera schaeffii</i>	CLLX, 635	1718
<i>scripta</i>	CLLX, 636	1719
<i>alutus</i> , <i>Aponogonichthys</i>	CLLXXVIII, 473	1110
<i>Sebastodes</i>	CCLXXII, 660	179
<i>Ambloplites rupestris</i>	CLVI, 410, 419 a, b, c	990
<i>Amblyopsis spelaea</i>	CXV, 307	706
<i>amblyrhynchus</i> , <i>Hemicaranx</i>	CXL, 386	912
<i>Anelurus dugesi</i>	XXVI, 50, 50a, 50b	138
<i>melas</i>	XXVI, 60	141
<i>platycephalus</i>	XXVI, 61	142
<i>americana</i> , <i>Morone</i>	CXXXI, 470	1134
<i>americanus</i> , <i>Ammodytes</i>	CXXXIX, 351	833
<i>Hemitripterus</i>	CCCV, 738	2023
<i>Menticirrhus</i>	CCXXXV, 572	1474
<i>Polypyron</i>	CLXXXI, 480, 480a	1139
<i>Pseudopleuronectes</i>	CCLXXXIX, 933	2647
<i>Amia calva</i>	XXII, 51, 51a	113
<i>Ammoecrypta beanii</i>	CLXXII, 455	1064
<i>pellucida clara</i>	CLXXII, 454	1063
<i>Ammodytes americanus</i>	CXXXIX, 351	833
<i>ampullaceus</i> , <i>Saccopharynx</i>	LXVI, 173	496
<i>Anableps dovii</i>	CXIII, 300	685
<i>analis</i> , <i>Hypocritichthys</i>	CCXXXIX, 582	1500
<i>Neomenis</i>	CXCVIII, 517	1265
<i>Notacanthus</i>	XCVIII, 264	615
<i>Anarhichas latifrons</i>	CCXLVI, 845	2416
<i>lepturus</i>	CCXLVII, 847	2447
<i>lupus</i>	CCXLVI, 846	2446
<i>ancylodon</i> , <i>Sagenichtys</i>	CCXXXI, 564	1410
<i>Anelyopsetta quadrocincta</i>	CCLXXV, 925	2634
<i>Anglichthys ciliaris</i>	CCLIV, 626, 626a	1084
<i>Anguilla chrysopha</i>	LV, 143	348
<i>anguilla</i> , <i>Ictalurus</i>	XXV, 57	2788
<i>anguillaris</i> , <i>Zonres</i>	CCXLVIII, 850	2457
<i>Anisotremus bilineatus</i>	CCVIII, 538	1349
<i>surinameus</i>	CCVIII, 537	1318
<i>virginicus</i>	CCIX, 539	1322
<i>annularis</i> , <i>Pomoxis</i>	CLIV, 415	987
<i>anomalum</i> , <i>Campostoma</i>	XXXIX, 95	265
<i>Anoplarchus atropurpureus</i>	CCXLII, 835	2422
<i>Anoploma finirostre</i>	CCLXXXIX, 674	1862
<i>Anthias asperlinguis</i>	CXCHI, 507	1227
<i>antillarum</i> , <i>Talismania</i>	LXXIV, 198	455
<i>Antimora violacea</i>	CCCLXII, 893, 893a	2514
<i>Apeltes quadratus</i>	CXX, 322	752
<i>Aploedoderus sayanus</i>	CXXII, 331	786
<i>Aploionotus grunniens</i>	CCXXXVI, 574	1484
<i>Apodichthys flavidus</i>	CCCXL, 830	2411
<i>apodus</i> , <i>Neomenis</i>	CXCVII, 515	1258
<i>Apogon pigmentarius</i>	CXXXVII, 472	1169
<i>retroseilla</i>	CLXXVII, 471	1168
<i>Apogonichthys olutus</i>	CLXXXVII, 473	1110
<i>Aponotis symmetricus</i>	CLIX, 434	998
<i>Aprius macrophthalmus</i>	CCI, 523	1280
<i>Apromon cortezianus</i>	CCCXLIX, 852	2461
<i>Apogon dentatus</i>	CC, 522	1278
<i>araea</i> , <i>Atherina</i>	CXXIII, 333	790
<i>Archoplites interruptus</i>	CLVII, 420	991
<i>Archosargus probatocephalus</i>	CXXVI, 554	1361

Archosargus, Architrichius, C. Arctoscopus, Arctozenus, arctatus, Pardalis, Cardis, arenatus, Paragymnophorus, argentatus, Paragymnophorus, argenteus, Brachygenys, Stenogaster, Argentina, Argyropelecus, argyrosomus, armatus, Leptoclinus, Aleurodilechthys, Ascidiella, ascensionis, asper, Evermanni, aspera, Linnaeus, asperiglauca, Aspidophorus, asprella, Cryphallus, Rutilus, asprellus, Radula, aspro, Hadronemus, A sternopterus, Astroleutes niger, Astronotus, stipes, Astroscopus, atchafalayae, Atheresthes, Atherina, atlantica, Atlanticus, A. Brama, atripinnis, Glaucostegus, atrocaudatus, atronotatus, atropurpureus, Auchenipterus, Aulorhynchus, Aulostomus, auratus, auratus, Mulloidichthys, aurea, Lampetra, aurivulus, Salmo, auritus, Lepidion, aurofrenatum, aurolineatum, aurorubens, austriatum, Axius thazardi, Axius thazardi, Averruncus, avocetta, Neomyxus, axillaris, Myoxocephalus, axinophrys, aya, Neomyxus, Azevia panamensis, Notesthes, azurica, Herichthys

Genera and species.	Plate and figure.	Text page.
<i>Archosargus unimaculatus</i>	CCXVI, 553	1359
<i>arietinus</i> , <i>Calamus</i>	CCXIV, 550	1355
<i>Aritocampus japonicus</i>	CCCXXXIII, 807	2297
<i>Arctozenus cornicans</i>	XCVI, 259	601
<i>arenatus</i> , <i>Pomacanthus</i>	CCLI, 623	1679
<i>ardens</i> , <i>Catostomus</i>	XXXIV, 84	179
<i>arenatus</i> , <i>Priacanthus</i>	CXCV, 511	1237
<i>argentinus</i> , <i>Gairdronopsurus</i>	CCCLXVII, 906	2559
<i>argentea</i> , <i>Bathyclupea</i>	CXXIX, 352	835
<i>Steindachnera</i>	CCCLXVIII, 909	2568
<i>argenteus</i> , <i>Larimus</i>	CCXXI, 565	1421
<i>Argentina silus</i>	LXXXVII, 232	526
<i>Argyropelecus olfersi</i>	XCVII, 261, 261a	604
<i>Argyrosomus</i> , <i>Damalichthys</i>	CCXXXII, 588	1509
<i>nigripinnis</i>	LXXVI, 203	472
<i>armatus</i> , <i>Leptocottus</i>	CCCII, 732	2012
<i>armiger</i> , <i>Alexurus</i>	CCXXXV, 784	2203
<i>Artediellus atlanticus</i>	CCLXXXV, 690	1906
<i>Ascolichthys rhodorus</i>	CCCV, 739	2025
<i>ascensionis</i> , <i>Holocentrus</i>	CXXXI, 358	848
<i>asper</i> , <i>Exuperes</i>	CCCCXXXVII, 818	2367
<i>aspera</i> , <i>Limanda</i>	CCCLXXVII, 930	2645
<i>aspelinguis</i> , <i>Anthias</i>	CXIII, 507	1227
<i>Aspidophoroides guntheri</i>	CCCXII, 755, 755a, 755b	2090
<i>monopterygius</i>	CCCXII, 750, 750a	2091
<i>asprella</i> , <i>Crystallaria</i>	CLXXI, 453	1061
<i>asprellus</i> , <i>Radulinus</i>	CCLXXXVII, 695	1920
<i>astro</i> , <i>Hadropterus</i>	CLXVI, 438	1032
<i>Asternopteryx gunnelliiformis</i>	CCXLIII, 834	2420
<i>Astrolyutes notospilotus</i>	CCLXXXIV, 688, 688a	1899
<i>Astromesisthes gemmifer</i>	XCHI, 251	586
<i>richardsoni</i>	XCIV, 252	587
<i>Astroscopus y griecum</i>	CCXXXIV, 808	2307
<i>atachafalaye</i> , <i>Signalosa</i>	LXIX, 184	2609
<i>Atheresthes stomias</i>	CCCLXXI, 917	2609
<i>Atherina area</i>	CXXIII, 333	790
<i>stipes</i>	CXXII, 322	790
<i>atherinoides</i> , <i>Clitodus</i>	CXVI, 310	719
<i>Atherinops affinis</i>	CXXVI, 342	807
<i>Atherinopsis californiensis</i>	CXXV, 341	806
<i>atlantica</i> , <i>Eblenimaria</i>	CCXXXL, 826	2402
<i>atlanticus</i> , <i>Artediellus</i>	CCLXXXV, 689	1906
<i>Benthodesmus</i>	CXXXVI, 374	887
<i>Rupisartes</i>	CCXXXIX, 825	2397
<i>Tarpon</i>	LXXVII, 177	409
<i>atripinnis</i> , <i>Gooden</i>	CXIV, 301	685
<i>atrocaudalis</i> , <i>Notropis cayuga</i>	XLVI, 114	260
<i>atromaculatus</i> , <i>Semotilus</i>	XI, 100	222
<i>atropurpureus</i> , <i>Anoplarchus</i>	CCXLIII, 895	2422
<i>Anchoaepterus nox</i>	CCCCXXXVII, 819	2373
<i>Athorhynchus flavidus</i>	CXX, 323	754
<i>Autostomus maculatus</i>	CXX, 324	754
<i>aurantiacus</i> , <i>Hypothomus</i>	CLXVIII, 442	1040
<i>auratus</i> , <i>Mullus</i>	CXXXII, 360	856
<i>aurea</i> , <i>Lampræta</i>	II, 5	13
<i>aureolus</i> , <i>Salvelinus alpinus</i>	LXXXIII, 220	511
<i>auritus</i> , <i>Lepomis</i>	CLIX, 425, 425a	1001
<i>australeatum</i> , <i>Sparisoma</i>	CCXLIII, 610	1034
<i>aurolineatum</i> , <i>Bathystoma</i>	CCVII, 535	1310
<i>aurorubens</i> , <i>Rhomboplites</i>	CC, 521	1277
<i>austrinum</i> , <i>Moxostoma</i>	XXXVII, 92	102
<i>Auxis thazard</i>	CXXXIII, 365	867
<i>Averruncus emmeiae</i>	CCCIX, 749, 749a	2069
<i>sterletus</i>	CCCX, 750, 750a	2071
<i>avocetta</i> , <i>Nemichthys</i>	LX, 157	309
<i>Avocettina gilli</i>	LIX, 154, 154a, 154b	367
<i>axillaris</i> , <i>Myoxocephalus</i>	CCXCVII, 721	1980
<i>axinophrys</i> , <i>Xystes</i>	CCCX, 752, 752a	2076
<i>aya</i> , <i>Neomenius</i>	CXCVII, 516	1264
<i>Azovia panamensis</i>	CCCLXXXIV, 942	2677
<i>aztecus</i> , <i>Noropsis</i>	XLV, 112	258
<i>azurea</i> , <i>Hermosilla</i>	CCXIX, 558	1383

Genera and species.	Plate and figure.	Text page.
<i>azureus</i> , <i>Galeichthys</i>	XXIV, 55	275
<i>Azurina hirundo</i>	CXXXIII, 588	154
<i>bahamensis</i> , <i>Teuthis</i>	CCLVI, CCLVII, 629a	1693
<i>Bairdiella chrysura</i>	CCXXII, 566	1433
<i>bairdii</i> , <i>Gastrostomus</i>	LXVII, 176	406
<i>Microspathodon</i>	CCXXXV, 592	1566
<i>bajonado</i> , <i>Calamus</i>	CCXIII, 548	1352
<i>Balistes carolinensis</i>	CCLVIII, 631	1701
<i>balteatus</i> , <i>Leuciscus</i>	XLI, 105, 105a	258
<i>Barathrodemus manatinus</i>	CCCLVII, 889	2517
<i>Barathronus bicolor</i>	CCCLVIII, 883	2524
<i>barbata</i> , <i>Brotula</i>	CCLIV, 871	2500
<i>Pallasina</i>	CCCVIII, 744	2049
<i>barbatulum</i> , <i>Lemonema</i>	CCCLXVI, 904	2536
<i>barbatum</i> , <i>Tyconemus</i>	CCCLII, 863	2474
<i>barraenda</i> , <i>Sphyraena</i>	CXXVIII, 349	823
<i>bartoni</i> , <i>Clethusa</i>	CXXXIII, 587	2841
<i>Bascanichthys peninsulae</i>	LXIII, 166, 166a	1515
<i>seuticaris</i>	LXIII, 165	379
<i>Bassogigas gilli</i>	CCCLVII, 879	378
<i>Bassozetus catena</i>	CCLVI, 876	2509
<i>normalis</i>	CCCLVI, 875	2507
<i>Bathyelupa argentea</i>	CXXIX, 352	845
<i>Bathygadus favosus</i>	CCCLXVII, 908	2565
<i>Bathygonus nigripinnis</i>	CCXXI, 753	2078
<i>Bathylingus benedicti</i>	LXXXVIII, 234	529
<i>Bathymaster signatus</i>	CCCCXXIII, 802	2288
<i>Bathypylhaea ovigerum</i>	CCCVIII, 707	2128
<i>Bathypterois quadrifilis</i>	LXXXIX, 238	544
<i>Bathystoma aurolineatum</i>	CCVII, 535	1310
<i>rimator</i>	CCVI, 534	1308
<i>beanii</i> , <i>Ammocrypta</i>	CLXXII, 455	1064
<i>Lingula</i>	CCCLXXVIII, 962	2646
<i>Serrivomer</i>	LVIII, 158	367
<i>Triglops</i>	CCLXXXVIII, 697	1924
<i>bella</i> , <i>Hypoclydonia</i>	CXXXIX, 475	1115
<i>Bellator egretta</i>	CCXXXI, 773	2174
<i>henedicti</i> , <i>Bathylagus</i>	LXXXVIII, 234	529
<i>Benthodesmus atlanticus</i>	CXXXVI, 374	887
<i>Benthosaurus grallator</i>	LXXXIX, 237	543
<i>beryllina</i> , <i>Menidia gracilis</i>	CCXXIV, 338	797
<i>beryllinus</i> , <i>Cryptotomus</i>	CCXLII, 608	1625
<i>Beryx splendens</i>	CCXXXI, 557	844
<i>bicanalis</i> , <i>Lac toprys</i>	CCLXII, 639, 639a, 639b	1723
<i>bichi</i> , <i>Polypterus</i>	1, 2
<i>bicolor</i> , <i>Barathronus</i>	CCCLVIII, 883	2524
<i>Leuciscus</i>	XI, 102	232
<i>Rondeletia</i>	XC, 240	548
<i>Rutilus</i>	XLIII, 107	241
<i>bilineata</i> , <i>Lepidopsetta</i>	CCCLXXVI, 928	2643
<i>bilineatus</i> , <i>Antistrema</i>	CCVIII, 538	1319
<i>Characodon</i>	CIX, 293	666
<i>billibus</i> , <i>Histioctonus</i>	CCCHII, 735	2018
<i>birostris</i> , <i>Mania</i>	XVII, 39	92
<i>biaon</i> , <i>Enophrys</i>	CCXCII, 705	1958
<i>bistrispinus</i> , <i>Rydiaeus</i>	CXCV, 509	1233
<i>bivittatus</i> , <i>Iridia</i>	CCXXXIX, 601	1505
<i>blackfordi</i> , <i>Yarrella</i>	XCIII, 249	584
<i>Blennioides embryum</i>	CCCHI, 734	2016
<i>bleunioides</i> , <i>Diploion</i>	CLXX, 449	1033
<i>Blennius eriaatus</i>	CCCCXXVIII, 821	2382
<i>favosus</i>	CCCCXXVIII, 820	2380
<i>Blenniopsis cirrhosa</i>	CCCIV, 736, 736a, 736b	2018
<i>Bodianus fulvus punctatus</i>	CLXXXII, 481	1146
<i>Bolechthys fusiformis</i>	CLXXVII, 469	1101
<i>boleoides</i> , <i>Radulinus</i>	CCLXXXVII, 394	1919
<i>Boleosoma canum</i>	CLXXI, 452	1060
<i>nigrum</i>	CLXX, 450	1058
<i>olmstedi</i>	CLXXI, 451	1057
<i>bolmani</i> , <i>Opsopoeodus</i>	XLIV, 110	249
<i>Bolmannia chlamydes</i>	CCCXXVIII, 701	2338

Genera and species.	Plate and figure.	Text page.
bonaci, <i>Mycteroptera</i>	CXXXVII, 492	1174
borealis, <i>feelinus</i>	CCLXXXIV, 687	1896
Borogadus sandia.....	CCLIX, 885	2533
boulengeri, <i>Mycteroptera</i>	CXXXVII, 490	1171
bouvieri, <i>Salmo clarkii</i>	LXXX, 212	496
brachiusculus, <i>Grammicolepis</i>	CLI, 410	974
Brachygenys <i>chrysargyreus</i>	CCVI, 533	1397
Brachystius <i>frenatus</i>	CCXXIX, 580	1499
brachyptera, <i>Remora</i>	CCXXX, 797; 797a	2272
Branchiostoma <i>caribaeum</i>	1, 1	3
brasiliensis, <i>Hemirhamphus</i>	CXVII, 313	722
Narcine	XIII, 35a	78
Scorpaena	CCLXXVII, 660	1842
brevirostris, <i>Hopropnion</i>	V, 18	41
brevirostrum, <i>Acipenser</i>	XXI, 47	106
brevispinis, <i>Sebastodes</i>	CCLXXI, 657	1787
Brevoortia <i>tyrannus</i>	LXXII, 195	434
Brotula <i>barbata</i>	CCCLIV, 871	2500
brunnens, <i>Hyophis</i>	LVI, 145	350
Brycon dentex	LV, 141	337
Bryostema <i>mugator</i>	CCCXL, 820	2410
<i>polyacanthocephalum</i>	CCCXL, 828	2408
caeca, <i>Ericymna</i>	LI, 129	302
balleri, <i>Prionoides</i>	CXCI, 503	1213
cavifrons, <i>Scarus</i>	PCXLIV, 613	1652
Calamus <i>arctifrons</i>	CCXIY, 559	1355
bajonado	CXXIII, 548	1352
calamus	CCXII, 546	1349
penna	CCXIV, 549	1354
proridens	CCXIII, 547	1350
calamus, <i>Calamus</i>	CCXII, 546	1349
Catechelys <i>infrata</i>	I XIII, 164	378
californica, <i>Tetronarce</i>	XII, 34, 34a	77
californiensis, <i>Atherinopsis</i>	CXXV, 341	806
Medialuna	CCXX, 560	1391
Typhlogobius	CCXXIX, 705	2262
callarias, <i>Gadus</i>	CCLXII, 891	2541
Callionymus <i>agassizii</i>	CCXXXIII, 779	2186
callirrhous, <i>Ioglossus</i>	CCXXXIII, 780	2103
callyodon, <i>Neoliparis</i>	CCXIV, 700, 700a	2110
Calotomus <i>xenodon</i>	CCXLIII, 609, 609a	1626
calva, <i>Amia</i>	XXII, 51, 51a	113
Campostoma <i>anomalum</i>	XXXIX, 95	205
canarium, <i>Boleosoma</i>	CLXXI, 452	1060
Etheostoma	CLXXII, 456	1076
canadense, <i>Stizostedion</i>	CLXIV, 434	1022
canadus, <i>Rachycentron</i>	CXLVIII, 401	948
canaliculatus, <i>Icelus</i>	CCLXXXVI, 603	1917
Cantherines <i>carolinus</i>	CCLVIII, 632	1713
capistratus, <i>Chactodon</i>	CCL, 622	1677
caprimus, <i>Otrynter</i>	CCXI, 543	1345
caprodes, <i>Percina</i>	CLV, 436, 436a	1026
Carmus <i>crysos</i>	CXLII, 388	923
hippos	CXLII, 387	920
latius	CXLII, 389	921
medusicola	CXLII, 390	924
carapinnis, <i>Coryphaenoides</i>	CCCLXVIII, 911	2579
Carcharhinus <i>limba</i>	V, 17	38
caribaeum, <i>Branchiostoma</i>	1, 1	3
caribbeus, <i>Colorynchus</i>	CCCLXX, 915	2580
carinatus, <i>Lutjanichthys</i>	LIX, 155	368
carnivalis, <i>Eumeaneutes</i>	CCXXXVI, 814	2350
carnivatus, <i>Celorhynchus</i>	CCCLXIX, 914	2588
carolae, <i>Cantherines</i>	CCLVIII, 632	1713
carolinensis, <i>Balistes</i>	CCLVIII, 631	1701
carolinum, <i>Priacanthus</i>	CCXXVIII, 708	2156
Trachinotus	CXLVII, 398	944
carpio, <i>Cyprinodon</i>	CXII, 297	675
Carpioles <i>cypriinus</i>	XXX, 71	167
caryi, <i>Hystruris</i>	CCXXXI, 585	1508
cataphractus, <i>Gasterosteus</i>	CXIX, 321	749
catenatus, <i>Fundulus</i>	CIV, 278, 278a	648

Genera and species.	Plate and figure.	Text page.
<i>catena</i> , <i>Bassozetus</i>	CCCLVI , 876, 876a	2509
<i>Catostomus ardens</i>	XXXIV , 84	179
<i>catostomus</i>	XXXII , 77	176
<i>commersonii</i>	XXXIV , 80	173
<i>griseus</i>	XXXI , 75	175
<i>latipinnis</i>	XXXI , 74	174
<i>macrochelus</i>	XXXII , 81	178
<i>occidentalis</i>	XXXII , 79	178
<i>pocatus</i>	XXXII , 79	178
<i>tameus</i>	XXXII , 78	177
<i>tsilteoensis</i>	XXXIII , 80	276
<i>catostomus</i> , <i>Catostomus</i>	XXXII , 77	176
<i>Catulus uter</i>	II , 12	25
<i>caudatus</i> , <i>Lepidopus</i>	CXXXVI , 373	866
<i>candillimbatus</i> , <i>Leptocephalus</i>	LVII , 149	355
<i>Caularctus macandrensis</i>	CCCCXXXV , 812	2328
<i>Caulolatilus microps</i>	CCCCXXX , 799	237
<i>Caudophryne jordani</i>	CCXCII , 957	2735
<i>cavrinus</i> , <i>Mylocheilus</i>	XI , 99	219
<i>Sebastodes</i>	CCLXXIV , 663	1820
<i>cavernosus</i> , <i>Hymenocephalus</i>	CCCLXIX , 912	2589
<i>cayorum</i> , <i>Corythoichthys</i>	CX , 322	238
<i>Ogilbia</i>	CCCLV , 873	2503
<i>cavynia atrocaudalis</i> , <i>Notropis</i>	XLVI , 114	266
<i>Centrarchus macropterus</i>	CLV , 417	988
<i>Centrocymnus coelolepis</i>	VIII , 25	35
<i>Centrolophius niger</i>	CXLIX , 403	965
<i>Centropomus undecimalis</i>	CXXIX , 476	1118
<i>Centropristes philadelphicus</i>	CXCI , 501	1201
<i>striatus</i>	CXC , 500	1200
<i>Centroscyllium fabricii</i>	VIII , 26	56
<i>cepediamum</i> , <i>Dorosoma</i>	LXIX , 180	416
<i>Cephalacanthus volitans</i>	CCCXXIV , 778	2100
<i>cephalus</i> , <i>Mugil</i>	CXXXVI , 341	811
<i>Ceratias holboelli</i>	CCCLXXXIX , 954	2729
<i>Ceratocottus diceranus</i>	CCXL , 706	1940
<i>Ceratoscopelus maderiensis</i>	XC , 242	557
<i>cerulea</i> , <i>Scytalinus</i>	CCCXLVIII , 849, 849a, 849b	2454
<i>guttatus</i>	XG , 241	549
<i>Cetorhinus maximus</i>	VII , 23	51
<i>Chenobryttus guulosus</i>	CLVII , 421	999
<i>Chenomugil proboscideus</i>	CXXVII , 346	816
<i>Chaetodipterus faber</i>	CCXLVII , 619	1668
<i>Chactodon capistratus</i>	CCL , 622	1671
<i>nigrirostris</i>	CCXLVIII , 620	1672
<i>ocellatus</i>	CXLIX , 621	1674
<i>chaetodon</i> , <i>Mesogonistius</i>	CLVIII , 423	905
<i>chalceogramma</i> , <i>Theragra</i>	CCCLIX , 887	2535
<i>Chalidura simula</i>	CCCLXVIII , 910	2578
<i>Channamura-mura vitata</i>	LXVI , 174	404
<i>Channa chrysos</i>	LXIX , 182	413
<i>chanos</i> , <i>Chanos</i>	LXIX , 182	414
<i>Characodon bilineatus</i>	CIX , 293	618
<i>varius</i>	CXI , 295, 295a	669
<i>Chasmistes copei</i>	XXXV , 87	2795
<i>bicornis</i>	XXXIV , 85	183
<i>stomias</i>	XXXV , 86	2794
<i>Chasmodes saburrae</i>	CCXXXIX , 824	2392
<i>Chauliodus sloanei</i>	XII , 250	585
<i>Channax pictus</i>	CCXXXIX , 953	2726
<i>Chelidonopterus affinis</i>	CLXIX , 474	1119
<i>cheneyi</i> , <i>Cottogaster</i>	CLXIX , 445	2851
<i>chesteri</i> , <i>Urophycis</i>	CCCLXVI , 903	2536
<i>Chimaodon niger</i>	CCXXXIII , 804	2391
<i>chilannahua</i> , <i>Notropis</i>	XLVI , 116	265
<i>Chilonymetus schopfii</i>	CCXLV , 636	1748
<i>Chimera affinis</i>	XIX , 40	95
<i>Chiostoma humboldtianum</i>	CCXLIV , 334	793
<i>chirurgus</i> , <i>Xiphistes</i>	CCXLIV , 317	2019
<i>Chitonotus pugnax</i>	CCCLXXXIII , 986	1886
<i>chimaenoides</i> , <i>Bolimanina</i>	CCCLXVIII , 701	2238
<i>Chlorichthys grammaticus</i>	CCXLII , 904	1626
<i>chloropterus</i> , <i>Alphestes</i>	CLXXXVI , 388, 388a	1794

Genera and species.	Plate and figure.	Text page.
<i>Chimaera</i> <i>nabrus euryurus</i>	CXLV, 394	938
<i>Chimaera</i> <i>cornutus</i>	CXV, 305	703
<i>Chimaera</i> <i>atherinoides</i>	CXVI, 310	719
<i>Chimaera</i> <i>Pozonias</i>	CCXXV, 573	1482
<i>Chimaera</i> <i>Brachygenys</i>	CCVI, 533	1307
<i>Chimaera</i> <i>serratoris</i> , <i>Pomolobus</i>	LXX, 187	425
<i>Chimaera</i> <i>Roceanus</i>	CLXXX, 477	1132
<i>Stenotomus</i>	CCXI, 544	1346
<i>Chimaera</i> <i>Ornithopristis</i>	CGX, 541	1338
<i>Chimaera</i> <i>Fundulus</i>	CVII, 287	655
<i>Chimaera</i> <i>Bairdiella</i>	CCXXII, 566	1433
<i>Chimaera</i> <i>Cloroseombrus</i>	CLXV, 394	938
<i>Chimaera</i> <i>Micropathodon</i>	CCXXXV, 503	1567
<i>Ocyurus</i>	CXCIX, 520	1275
<i>chrysopsa</i> , <i>Anguilla</i>	LV, 143	348
<i>chrysopsa</i> , <i>Trophycis</i>	CCCLXV, 902	2555
<i>Ciehlasoma</i> <i>bartoni</i>	CCXXXII, 587	1615
<i>ciliatus</i> , <i>Angelicichthys</i>	CCLIV, 626, 626a	1684
<i>ciliatus</i> , <i>Monacanthus</i>	CCLIX, 633	1714
<i>Selmatodes</i>	CCLXX, 655	1783
<i>embrius</i> , <i>Enchelyopus</i>	CCCXLVII, 907	2580
<i>emarginatum</i> , <i>Erheostoma</i>	CLXXIII, 457	1078
<i>Nymphaea</i>	CCXVIII, 536	1372
<i>caratum</i> , <i>Ginglymostoma</i>	IV, 13	26
<i>cirratus</i> , <i>Urophycis</i>	CCCLXIV, 899	2553
<i>Cirrhites</i> <i>rivulatus</i>	CCXXXVII, 576	1491
<i>cirrhosus</i> , <i>Idiopsis</i>	CCCIV, 736, 736a, 736b	2018
<i>Citharichthys</i> <i>marmoratus</i>	CCCLXXXV, 944	2684
<i>sordidus</i>	CCCLXXXIV, 943	2679
<i>clara</i> , <i>Ammocrypta pellucida</i>	CLXXII, 454	1063
<i>clarkii</i> , <i>bouvieri</i> , <i>Salmo</i>	LXXX, 212	496
<i>clarkii</i> , <i>nashawi</i> , <i>Salmo</i>	{ 2819	
<i>clarkii</i> , <i>macdonaldi</i> , <i>Salmo</i>	LXXXIX, 208	493
<i>clarkii</i> , <i>macdonaldi</i> , <i>Salmo</i>	{ 2819	
<i>pleuriticus</i> , <i>Salmo</i>	LXXXI, 214	497
<i>spilurus</i> , <i>Salmo</i>	{ 2819	
<i>stomias</i> , <i>Salmo</i>	LXXX, 211	496
<i>virginalis</i> , <i>Salmo</i>	LXXXIX, 210	495
<i>Cleptodus</i> <i>parva</i>	LXXX, 213	497
<i>Clevelandia</i> <i>los</i>	LXXXIX, 209	495
<i>Clupea</i> <i>harengus</i>	CCXXXVIII, 599	1586
<i>Clupea</i> <i>pulchra</i>	CCXXXVIII, 703	2254
<i>clupeiformis</i> , <i>Coregonus</i>	LXX, 185	421
<i>coccocephalus</i> , <i>Notropis</i>	LXX, 186	422
<i>cocolepis</i> , <i>Centrolymus</i>	LXXVI, 202	405
<i>Celorhynchus</i> <i>caribbaeus</i>	XLIX, 124	284
<i>carminalis</i>	VIII, 25	55
<i>cochlias</i> , <i>Scomerus</i>	CCCLIX, 915	2589
<i>colliei</i> , <i>Hydrolagus</i>	CCCLXIX, 914	2588
<i>Columbia</i> <i>transmontanus</i>	CCCLXIX, 913	2588
<i>commersonii</i> , <i>Catostomus</i>	CXXXIII, 364	866
<i>conger</i> , <i>Leptophthalus</i>	XIX, 41	95
<i>congestum</i> , <i>Moxostoma</i>	C'XII, 330	784
<i>Congrellus</i> <i>davidsoni</i>	XXIV, 83	178
<i>coper</i> , <i>Alepoionus</i>	LVII, 148	354
<i>Chasmistes</i>	XXXVI, 91	192
<i>Copelandellus</i> <i>quiescens</i>	LVIII, 150	357
<i>corallinus</i> , <i>Cryptotrema</i>	LXXV, 199	459
<i>coregonoides</i> , <i>Paralepis</i>	XXXV, 87	2795
<i>Coregonus</i> <i>clupeiformis</i>	CLXXVI, 468	1100
<i>coreoperca</i>	CCXXXVII, 817	2366
<i>coreoperca</i> , <i>williamsoni</i>	XCVI, 260	602
<i>cornuta</i> , <i>Lamna</i>	LXXVI, 202	405
<i>cornuta</i> , <i>Chilogaster</i>	LXXVI, 201	402
<i>cornuta</i> , <i>smilis</i>	LXXV, 200, 200a	403
<i>vertebrata</i> , <i>Aspredon</i>	V, 22	49
	CXV, 305	703
	CCLV, 627	1087
	C'CXLIX, 652	240

Genera and species.	Plate and figure.	Text page.
<i>coruscans</i> , <i>Arctozenus</i>	XCVI , 250	661
<i>Coryphana hippurus</i>	CXLIX , 402	952
<i>Coryphenooides carapinnis</i>	CCCLXVIII , 911	2579
<i>Corythoichthys eayorum</i>	CXX , 326	2838
<i>Cottogaster cheneyi</i>	CLXIX , 445	2851
<i>shumardi</i>	CLXVII , 444	1046
<i>Cottus aleuticus</i>	CCXCIII , 711	1957
<i>evermanni</i>	CCXCIII , 707	1945
<i>klamathensis</i>	CCXCIII , 710	1955
<i>liponius</i>	CCXCIV , 712	1962
<i>perplexus</i>	CCXCIII , 709	1955
<i>princeps</i>	CCXCIV , 713	1962
<i>punctulatus</i>	CCXCII , 708	1948
<i>conesi</i> , <i>Cryptopsaras</i>	CCCNC , 956	2731
<i>coulterii</i> , <i>Coregonus</i>	LXXXVI , 201	492
<i>crassiceps</i> , <i>Plectromus</i>	CXXX , 356	843
<i>crestonius</i> , <i>Tenthis</i>	CCLVI , 628	1692
<i>cretense</i> , <i>Sparisoma</i>	CCXLV , 6166
<i>cristatus</i> , <i>Blennius</i>	CCXXXVIII , 821	2382
<i>Christyomer namayencus</i>	LXXXII , 217	504
<i>eristhleta</i> , <i>Scorpaena</i>	CLXXVI , 668	1841
<i>eroeodilus</i> , <i>Lampanyctus</i>	XCI , 243	558
<i>erossotus</i> , <i>Etropus</i>	CCCLXXXVI , 946	2689
<i>eratulus</i> , <i>Eubryx</i>	CCXLVIII , 851	2458
<i>eremntifer</i> , <i>Pisodonophis</i>	LXII , 163	377
<i>erumenophthalmus</i> , <i>Trachurops</i>	CCXL , 385	911
<i>Cryptacanthodes maculatus</i>	CCXLV , 843	2443
<i>Cryptopsaras conesi</i>	CCCNC , 956	2731
<i>Cryptotomus beryllinus</i>	CCXLII , 608	1625
<i>Cryptotrema coralliphium</i>	CCXXXVII , 817	2366
<i>erysoleucus</i> , <i>Abramus</i>	XLV , 111	250
<i>erysos</i> , <i>Caranx</i>	XLII , 388	923
<i>Crystallaria asprella</i>	CLXXI , 453	1061
<i>culveri</i> , <i>Trachinotus</i>	CXLVII , 397	942
<i>crenema</i> , <i>Mugil</i>	CXXVI , 344	813
<i>cuvieri</i> , <i>Tetragonurus</i>	CLII , 411	976
<i>czanulile</i> , <i>Scarus</i>	CCXLIV , 612	1648
<i>Cycleptus elongatus</i>	XXX , 72	168
<i>Cyclopterus lumpus</i>	CCXXIII , 757	2066
<i>cyclops</i> , <i>Liparis</i>	CCCXV , 764, 764a	2118
<i>Cymatogaster aggregatus</i>	CCXXVIII , 579a	1498
<i>Cynoscion nebulosus</i>	CCXXI , 563	1409
<i>nothus</i>	CCXX , 561	1406
<i>regalis</i>	CCXX , 562	1407
<i>cypho</i> , <i>Xyrauchen</i>	XXXV , 88	184
<i>cyprinella</i> , <i>Ictiobius</i>	XXX , 70	161
<i>Cyprinodon carpio</i>	CXII , 297	675
<i>variegatus</i>	CXL , CXII , 296, 296a	671
<i>cyprioides</i> , <i>Lophogobius</i>	CCCXXVI , 786	2299
<i>Cyprinus</i> , <i>Carpides</i>	XXX , 71	168
<i>Cypsilurus californicus</i>	CXIX , 310	2830
<i>Dallia pectoralis</i>	NXIX , 267	621
<i>Damalichthys argyrosomus</i>	CCXXXII , 586	1509
<i>Dasyatis sabina</i>	XIV , 36, 36a	84
<i>Dasyctotus setiger</i>	CCC , 727	1991
<i>decaenurus</i> , <i>Hexagrammos</i>	CCLXXX , 676, 676a	1867
<i>Decapterus macarellus</i>	XL , 384	909
<i>decurrens</i> , <i>Pleuronichthys</i>	CCLXXV , 926	2637
<i>dekayi</i> , <i>Isurus</i>	VI , 21	48
<i>dennyi</i> , <i>Liparis</i>	CCCXVII , 766a	2124
<i>dentatus</i> , <i>Apisalus</i>	CC , 522	1278
<i>Grammatostomias</i>	XCIV , 254	580
<i>Paralichthys</i>	CCLXXXIII , 922	2629
<i>dentex</i> , <i>Brycon</i>	LV , 141	337
<i>Osmerus</i>	LXXXVI , 229	594
<i>Derichthys serpentinus</i>	LV , 142	343
<i>dermathinus</i> , <i>Lycodapus</i>	CCCLIV , 870	2492
<i>Dermatolepis zauchus</i>	CXXXVI , 489	2854
<i>diaphanus</i> , <i>Fundulus</i>	CII , 275, 275a	645
<i>diapterus</i> , <i>Furcifimus</i>	CCCLI , 861	2472
<i>diceraus</i> , <i>Ceratocottus</i>	CCXCI , 706	1940
<i>Dicrolene intronigra</i>	CCLVIII , 882	2522

Genera and species.	Plate and figure.	Text page.
<i>Dicromita agassizii</i>		
<i>Diodon hystrix</i>		
<i>Diplocrema formosum</i>		
<i>Diplestes blennioides</i>		
<i>Diplodus holbrookii</i>		
<i>dipulus</i> , <i>Iridio</i>		
<i>dodecahedron</i> , <i>Ocea</i>		
<i>delichogaster</i> , <i>Pholis</i>		
<i>dolomieu</i> , <i>Microterus</i>		
<i>Dormitator maculatus</i>		
<i>dormitor</i> , <i>Philypnus</i>		
<i>Dorosoma cepedianum</i>		
<i>dorsalis</i> , <i>Microspathodon</i>		
<i>Seriola</i>		
<i>dovii</i> , <i>Anableps</i>		
<i>drummond-hayi</i> , <i>Epinephelus</i>		
<i>ductor</i> , <i>Xanthurites</i>		
<i>dugesii</i> , <i>Adinia</i>		
<i>Algansia</i>		
<i>Ameturus</i>		
<i>dulcis</i> , <i>Rhinichthys</i>		
<i>duquesnii</i> , <i>Placopharynx</i>		
<i>earli</i> , <i>Trophycis</i>		
<i>Echeneis naucrates</i>		
<i>egmontis</i> , <i>Ahilia</i>		
<i>egrette</i> , <i>Bellator</i>		
<i>eigenmanni</i> , <i>Evarpa</i>		
<i>Schastodes</i>		
<i>Elanura forficata</i>		
<i>elassodon</i> , <i>Hippoglossoides</i>		
<i>Elissoma evergladei</i>		
<i>elegans</i> , <i>evides</i> , <i>Gibbonsia</i>		
<i>Eleginops navaga</i>		
<i>Eleotris pisonis</i>		
<i>elongatus</i> , <i>Cycleptus</i>		
<i>Labichthys</i>		
<i>Ophidion</i>		
<i>Elops saurus</i>		
<i>Endemaria atlantica</i>		
<i>emblematus</i> , <i>Scarus</i>		
<i>embryum</i> , <i>Blennicottus</i>		
<i>Embryx eratilinus</i>		
<i>erilia</i> , <i>Opsopoeodus</i>		
<i>Emmeekia venusta</i>		
<i>enomelane</i> , <i>Averrhoecus</i>		
<i>Emperichthys merriami</i>		
<i>Enchelyopus cimbrinus</i>		
<i>Enneacanthus gloriosus</i>		
<i>Enneanectes carminalis</i>		
<i>Enophrys bison</i>		
<i>Entosphenus tridentatus</i>		
<i>Epinephelus adscensionis</i>		
<i>drummond-hayi</i>		
<i>morio</i>		
<i>striatus</i>		
<i>Epinnula magistralis</i>		
<i>Eques lanceolatus</i>		
<i>erarcha</i> , <i>Eurystole</i>		
<i>Ericynba buekata</i>		
<i>Ermyzon suetetta</i>		
<i>eritacea</i> , <i>Raja</i>		
<i>Erotelis smaragdus</i>		
<i>Esox violaceus</i>		
<i>Eslops armatum</i> , <i>jordani</i>		
<i>Etelis ocellatus</i>		
<i>Etheostoma</i> , <i>Aboma</i>		
<i>Etheostoma canorum</i>		
<i>cineratum</i>		
<i>towae</i>		
<i>lesse</i>		

Genera and species.	Plate and figure.	Text page.
Etheostoma jordani	CLXXXIII, 458	1079
<i>julie</i>	CLXXVI, 466	1092
<i>lepidogenys</i>	CLXXIV, 462	1087
<i>obeyense</i>	CLXXV, 463	1092
<i>pagei</i>	CLXXV, 464	1092
<i>pottsii</i>	CLXXXIII, 459	1082
<i>virgatum</i>	CLXXV, 465	1093
Etropus crossopterus	CCCLXXXVI, 946	2689
<i>rimosus</i>	CCCLXXXV, 945	2688
Eupomacentrus rectifrenum	CCXXXIX, 580	3176
Eupomotis euryurus	CLXI, 428	1008
<i>gibbosus</i>	CLXI, 429	1009
euryurus, Eupomotis	CLXI, 428	1008
Eurystole eriarcha	CXXXV, 230	803
Evarra eigenmanni	LI, 131	304
evergladei, Elassoma	CLIII, 414	982
evermanni, Cottus	CCXCI, 707	1945
<i>Thyrina</i>	CXXXV, 340	804
Evermannia zosterura	CCXXXIX, 794	2256
evides, Gibbonsia elegans	CCCLXXXVI, 814	2352
<i>Hadropterus</i>	CLXVII, 440	1036
evionthas, Quassiremus	LXIV, 167	380
evolans, Prionotus	CCCX, 772	2168
Evoplites viridis	CXCVI, 514	1216
Evoxymetopon tenuitius	CXXXVI, 372	886
Exerpes asper	CXXXXVII, 818	2307
exilis, Schilbeoides	XXVII, 65	147
Excetus volitans	CVI, 318	734
Exoglossum maxillingua	LIV, 140	327
Exonantes exiliens	CXVIII, 316	732
<i>rondeletii</i>	733	2820
exiliens, Exonantes	CXVIII, 317	732
<i>faber, Chaetodipterus</i>	2820	732
<i>fabricii, Centroscyllium</i>	CXVIII, 316	732
<i>falcata, Agonopsis</i>	CCXLVII, 619	166
<i>phenax, Mycteroptera</i>	VIII, 26	56
<i>falcatus, Trachinotus</i>	LIII, 135	313
<i>fuscatus, Achirus</i>	CLXXXVIII, 495	1185
<i>favosus, Bathygadus</i>	CXLVI, 396	941
<i>Bleennius</i>	CCCLXXXVII, 948	2700
<i>Telichthys felis</i>	CCCXLVII, 908	2563
<i>felis, Telichthys</i>	CCXXXVIII, 820	2180
<i>ferox, Alepisaurus</i>	XXIII, 52	118
<i>Stomias</i>	XXIII, 52	118
<i>ferruginea, Limanda</i>	XCV, 257	595
<i>limbra, Anoploponus</i>	XCIV, 253	588
<i>flavineo marianus</i>	CCCLXXVII, 929	2644
<i>flavescens, Pereca</i>	CCLXXXIX, 674	1862
<i>flavidus, Apodichthys</i>	CXXXI, 350	852
<i>Aulorhynchus</i>	CLXV, 435	1029
<i>Congrellus</i>	CCXLII, 850	2411
<i>flavus, Noturus</i>	CXX, 323	751
<i>flora, Neoliparis</i>	LVIII, 150	357
<i>floride, Jordaniella</i>	XXVII, 63	144
<i>Fodiator acutus</i>	CCCXV, 762	2111
<i>fotens, Synodus</i>	CXII, 298	67
<i>fonticola, Miceroperca</i>	CXVII, 315	72
<i>fontinalis, Salvelinus</i>	LXXXVIII, 236	558
<i>forticata, Elanura</i>	CLXXXVII, 470	1104
<i>formosum, Diplectrum</i>	LXXXII, 218	506
<i>francischii, Gyropleurodon</i>	CCLXXXIX, 701	1939
<i>frenatus, Brachyistius</i>	CXIV, 302	657
<i>Sarritor</i>	CXCI, 502	1257
<i>frigidus, Lycones</i>	III, 9	20
<i>fulvus, Physcianus</i>	CCXXXIX, 580	1439
<i>punctatus, Bodianus</i>	CCCX, 751	2073
<i>funduloides, Fundulus</i>	CCCL, 856	2463
<i>Fundulus albo-lineatus</i>	CCCLXIII, 896	2547
<i>Fundulus</i>	CLXXXIII, 481	1146
<i>atratulus</i>	CVI, 282	650
<i>maculatus</i>	CV, 281	649

Fundulus car
cl
di
fu
go
he
jen
luc
ma
no
not
oce
pal
rat
sen
ste
zeb
farcatus, Icti
Phan
furcifer, Param
Furcimarus d
furiosus, Schi
busformis, Bo
Gadus callaris
gairdneri, Saha
Gairdrosaurus
galacturus, No
galeatus, Gym
Galeichthys az
g
m
Galeorhinus zy
Gambusia ahin
Garmannia par
Garra nigrita
Gasterosteus a
c
Gastromonida
gennum, Hypop
genimur, Astr
Germia albulosa
Gerris olithos
Gibbonsia eleg
gibbosus, Enop
gilberti, Galeich
Schilbe
Sebasto
Ulocutu
Gilbertidia sig
gilli, Avocettina
Bassogigas
Cetomimus
Lipogenys
Neobrythi
Ginglymostoma
glacialis, Liops
glauca, Prionace
glauca, Trachin
glesne, Regaleci
globosa, Lycoph
gloriosus, Enne
Gnathopoma ma
gobioidei, Hyps
oceanicus
stigmatica
Goodenia atrilob
goodei, Foudia

Genera and species.	Plate and figure.	Text page.
<i>Fundulus catenatus</i>	CIV, 278, 278a	648
<i>chrysotus</i>	CVII, 287	655
<i>diaphanus</i>	III, 275, 275a	645
<i>funduloides</i>	CVI, 282	650
<i>goodei</i>	CIX, 291	{ 664 2831
<i>heteroclitus</i>	II, 273	640
<i>jenkinsi</i>	CVI, 284	651
<i>luciae</i>	CVII, 286	654
<i>macleodii</i>	CVI, 283	651
<i>majalis</i>	CI, 271, 271a, 271b	639
<i>notatus</i>	CVIII, 289	659
<i>notiti</i>	CVIII, 288	656
<i>ocellaris</i>	II, 274	642
<i>pallidus</i>	II, 272	638
<i>polyurus</i>	CVII, 285	652
<i>rathbuni</i>	CV, 280	649
<i>seminolis</i>	CIV, 277	647
<i>stellifer</i>	CV, 279	648
<i>zebrinus</i>	II, 276	646
<i>fureatus</i> , <i>Ictalurus</i>	XXV, 56	134
<i>Phanerodon</i>	CCXXX, 583	1506
<i>furcifer</i> , <i>Paranthias</i>	CXII, 504	1221
<i>Furcimanus diapterus</i>	CCCLII, 861	2472
<i>furious</i> , <i>Schilbeoides</i>	XXIX, 69, 69a, 69b	149
<i>tusiformis</i> , <i>Boleichthys</i>	CLXXXVII, 469	1101
<i>Gadus callarias</i>	CCCLXI, 891	2541
<i>gairdneri</i> , <i>Salmo</i>	LXXXI, 215	497
<i>Gairdronotus argentatus</i>	CCCLXVII, 906	2559
<i>galacturus</i> , <i>Notropis</i>	XLVII, 122	279
<i>galeatus</i> , <i>Gymnoanthus</i>	CCCI, 730	2010
<i>Galeichthys azureus</i>	XXIV, 55	2775
<i>gilberti</i>	XXIV, 54	2773
<i>milberti</i>	XXIII, 53	128
<i>Galeorhinus zyopterus</i>	IV, 15	32
<i>Gambusia affinis</i>	CXIII, 299, 299a	680
<i>Garmania paradoxa</i>	CCCXXVII, 790	2232
<i>Garupa nigrita</i>	CLXXXIV, 486	1161
<i>Gasterosteus aculeatus</i>	CXIX, 320	747
<i>cataphractus</i>	CXIX, 321	749
<i>Gastrostomus hairii</i>	LXVII, 176	406
<i>gemma</i> , <i>Hypoplectrus</i>	CLXXXIX, 407	1193
<i>genimifer</i> , <i>Astromethes</i>	CXII, 251	586
<i>Germa alalunga</i>	CXXXIV, 367	871
<i>Gerres olishostomus</i>	CCXXVII, 557	1376
<i>Gibbonsia elegans evides</i>	CCXXXVI, 815	2352
<i>gibbosus</i> , <i>Empomotis</i>	CLXI, 429	1069
<i>gilberti</i> , <i>Galeichthys</i>	XXIV, 54	2773
<i>Schilbeoides</i>	XXVIII, 67, 67a, 67b	148
<i>Stenostodes</i>	CCLXXV, 665	1823
<i>Ulocentra</i>	CLXIX, 443	1049
<i>Gilbertidia sanguinea</i>	CCCVI, 741	2028
<i>gilli</i> , <i>Avocettina</i>	LIX, 154, 154a, 154b	{ 267 2801
<i>Bassogigas</i>	CCCLVII, 879	2515
<i>Cetomimus</i>	XC, 241	549
<i>Lipogramys</i>	XCIX, 266	619
<i>Neohylithites</i>	CCCLVII, 877	2512
<i>Ginglymostoma cirratum</i>	IV, 13	26
<i>glacialis</i> , <i>Liopsetta</i>	CCCLXXX, 935	2649
<i>glauca</i> , <i>Prionace</i>	IV, V, 16, 16a	33
<i>glaucus</i> , <i>Trachinops</i>	CXLVI, 395	940
<i>glesne</i> , <i>Regalecus</i>	CCCLXX, 916	2596
<i>globosa</i> , <i>Lyosphara</i>	CCLXVII, 640, 649a	1751
<i>gloriosus</i> , <i>Enneacanthus</i>	CLVIII, 422	993
<i>Gnathopsis maxillosa</i>	CCXXXI, 801	2284
<i>gobiooides</i> , <i>Hypsilemetes</i>	CCXXXIII, 805	2294
<i>Gobius hastatus</i>	CCXXXVI, 788	2229
<i>oceanicus</i>	CCXXXVII, 789, 789a	2230
<i>stigmatus</i>	CCXXXVI, 787	2224
<i>Goodea atripinnis</i>	CXIV, 301	685
<i>goodei</i> , <i>Fundulus</i>	CIX, 291, 664	2831

Genera and species.	Plate and figure.	Text page.
goodei, <i>Ptilichthys</i>	CCCXLVII, 848	2452
gorbuscha, <i>Oncorhynchus</i>	LXXVII, 205	478
gracile, <i>Peristedion</i>	CCXXXII, 776	2179
gracilis, <i>Aldrovandia</i>	XCVIII, 263	610
beryllina, <i>Menidia</i>	CXXIV, 338	797
<i>Photonectes</i>	XCV, 255	591
<i>Platygobio</i>	LIV, 139	326
grallator, <i>Benthosaurus</i>	LXXXIX, 237	543
<i>Gramma</i> loreto.....	CXIII, 508	1229
grammatismus, <i>Chlorichthys</i>	CCXLI, 604	1610
<i>Grammatostomias dentatus</i>	XCIV, 254	590
<i>Grammicolepis brachinsculus</i>	CLII, 410	974
grandicornis, <i>Scorpaena</i>	CCLXXVIII, 671	1850
greeni, <i>Neolamprologus</i>	CCCVI, 763, 763a	2112
griseus, <i>Catostomus</i>	XXXI, 75	175
<i>Hexanchus</i>	II, 8	19
grunniens, <i>Aplidionotus</i>	CCXXVI, 574	1481
guacamaya, <i>Pseudosecarus</i>	CCXLVI, 617	1657
gulosus, <i>Chiobryttus</i>	CLVII, 421	962
gunnelliiformis, <i>Asternopteryx</i>	CCCXIII, 834	2420
gunnelli, <i>Pholidichthys</i>	CCCXLII, 832	2419
guntheri, <i>Aspidophoroides</i>	CCCXII, 755, 755a, 755b	2090
<i>Hadropterus</i>	CLXVI, 439	1633
<i>Hoplopagrus</i>	CXCVI, 513	1244
guttatus, <i>Percopsis</i>	CXXI, 329	784
guttifer, <i>Ophichthus</i>	XLIV, 168	583
Gymnella viridis.....	CCCLII, 864-864c	2479
Gymnophanthus galeatus.....	CCCCI, 730	2010
<i>pistilliger</i>	CCCII, 729, 729a, 729b	2006
Gymnosarda alleterata.....	CXXXIV, 366	869
Gyroleuroodus francisci.....	III, 9	29
<i>Hadropterus aspro</i>	CLXVI, 438	1632
<i>evides</i>	CLXVII, 440	1636
<i>guntheri</i>	CLXVII, 439	1633
<i>macrocephalus</i>	CLXVI, 437	1631
<i>scirurus</i>	CLXVII, 441	1637
<i>Hæmulon album</i>	CCHI, 528	1295
<i>macrostomum</i>	CCIV, 529	1296
<i>parra</i>	CCIV, 530	1297
<i>plumieri</i>	CCV, 532	1304
<i>scirurus</i>	CCV, 531	1303
<i>harengus</i> , <i>Clupea</i>	LXX, 185	421
<i>raleighana</i>	XIX, 42	96
<i>hostatus</i> , <i>Gobius</i>	CCXXXVI, 788	2229
<i>helleri</i> , <i>Xiphophorus</i>	CNV, 304	701
<i>Hemianthias vivanus</i>	CXII, 505	1223
<i>Hemicarax ambylyrhynchus</i>	CLXII, 386	912
<i>hemilepidotus</i> , <i>Hemilepidotus</i>	CCXC, 704, 704a, 704b	1935
<i>hemilepidotus hemilepidotus</i>	CCXC, 704, 704a, 704b	1935
<i>jordani</i>	CCXC, 703	1934
<i>Hemirhamphus brasiliensis</i>	CXVII, 313	722
<i>Hemitripterus americanus</i>	CCCV, 738	2023
<i>henaphilii</i> , <i>Stathmonotus</i>	CCXL, 827	2407
<i>henshawi</i> , <i>Salmo clarkii</i>	LXXXIX, 208	493
<i>bentz</i> , <i>Hypsoblennius</i>	CCXXXIX, 823	2399
<i>hermosilla azurea</i>	CCXIX, 558	1383
<i>Heterandria formosa</i>	CXIV, 302	687
<i>heteroclitus</i> , <i>Fundulus</i>	CII, 273	640
<i>Heterodontus philippinus</i>	III, 10
<i>Hexagrammos decagrammus</i>	CCLXXX, 676, 676a	1867
<i>octogrammus</i>	CCLXXX, 677	1869
<i>otakii</i>	CCLXXXI, 680	1867
<i>stelleri</i>	CCLXXXI, 678	1871
<i>supercliosus</i>	CCLXXXI, 679	1872
<i>Hexanchus griseus</i>	II, 8	19
<i>Hiodon selenops</i>	LXVIII, 181	414
<i>tergisus</i>	LXVIII, 180	413
<i>Hippocampus hudsonius</i>	CXXI, 327	777
<i>zoster</i>	CXXI, 328	778
<i>Hippoglossoides elassodon</i>	CCCLXXII, 920	2615
<i>platessoides</i>	CCCLXXII, 919	2614
<i>Hippoglossus hippoglossus</i>	CCCLXXI, 918	2611

hippoglossus,
hippos, *Caran*,
hippurus, *Cor*,
hirundo, *Azn*,
 Leto,
hippidius, *Mor*,
hoplostethus, *S*,
hoplostethus, *H*,
hoplopterus, *H*,
hoplomimus, *sch*,
hoplomus, *Mugil*,
hudsonius, *Hip*,
hoplostethus, *Seba*,
hoplostethus, *sal*,
humboldtianus,
humeralis, *Para*,
 Sard,
Hybopsis astivis,
 watan,
Hydrolags coll,
hydropholox, *Leu*,
Hymenopthalma,
Hymis hopkinsi,
Hypoclydonis be,
Hypocrichtichys,
Hypohnomus atra,
 spic,
Hypomesus olid,
 pret,
Hypoplectrus ge,
 un,
Hypopriion brevi,
Hyporhamphus n,
 1,
Hypsiconetes go,
Hypsoblennius h,
 ie,
Hypsurus cari,
Hypsypops rubic,
Hystericarpinus tr,
hystrix, *Blodon*,
Iceelius borealis,
Iceelius canaliculat,
leichtys locking,
Ictalurus anguilla,
 toreatus,
Ictiobius punctatus,
Hyphophis cyprinell,
inermis, *Rabirobi*,
internalis, *Histiob*,
inframus, *Novaeul*,
Inopsetta ischrya,
hornatus, *Microle*,
insignis, *Schilbeod*,
interruptus, *Arche*,
intertinctus, *Myst*,
introna, *Dicrol*,
loglossus callirinus,
longus, *Hypsobl*,
los, *Clevelandia*...

Genera and species.	Plate and figure.	Text page.
<i>Hippoglossus</i> , <i>Hippoglossus</i>	CCCLXXI , 918	2611
<i>Hippus</i> , <i>Caranx</i>	CXLI , 387	920
<i>Hippurus</i> , <i>Coryphaena</i>	CXLIX , 402	952
<i>Hirundo</i> , <i>Azurina</i>	CCXXXIII , 588	1544
<i>Leiocotus</i>	CCCI , 731	2011
<i>Hispidus</i> , <i>Monacanthus</i>	CCLIX , 634	1715
<i>Histiobranchus</i> , <i>internalis</i>	LVI , 147	352
<i>Histioctetus</i> , <i>bilobus</i>	CCCIII , 735	2018
<i>Holacanthus</i> , <i>tricolor</i>	CCLII , 626	1684
<i>Holbilla</i> , <i>Ceratias</i>	CCCLXXXIX , 954	2729
<i>Holbrookii</i> , <i>Diphodus</i>	CCXVII , 555, 555a	1362
<i>Holocentrus</i> , <i>ascensionis</i>	CXXXI , 358	848
<i>hopkinsi</i> , <i>Hymnis</i>	CXLII , 391	933
<i>Sébastodes</i>	CCLXII , 659	1789
<i>hoplomystax</i> , <i>Sparisoma</i>	CCXLIV , 611	1632
<i>Hoplopagrus</i> , <i>guntheri</i>	CXCVI , 513	1244
<i>Hoplostethus</i> , <i>mediterraneus</i>	CXXX , 354	837
<i>Hoplmis</i> , <i>schmidti</i>	LVII , 151	361
<i>hospes</i> , <i>Mugil</i>	CXXVII , 345	814
<i>hussonius</i> , <i>Hippocampus</i>	CXXI , 327	777
<i>Notropis</i>	XLVII , 119	269
<i>sardadum</i> , <i>Notropis</i>	XLVIII , 120	270
<i>humboldtianum</i> , <i>Chirostoma</i>	CXXXII , 334	793
<i>humeralis</i> , <i>Paralabrax</i>	CXC , 499	1196
<i>Sardinella</i> *.....	LXXIII , 193	431
<i>Hybopsis</i> , <i>astivalis</i> , <i>marconis</i>	LIII , 136	316
<i>altus</i>	LIV , 138	321
<i>watanga</i>	LII , 137	319
<i>Hydroagus</i> , <i>colliei</i>	XIX , 41	95
<i>Hydrophox</i> , <i>Leuciscus</i>	XLII , 104	238
<i>Hymenocelphalus</i> , <i>cavernosus</i>	CCCLXIX , 912	2580
<i>Hymnis</i> , <i>hopkinsi</i>	CXLIII , 301	933
<i>Hypocephalonia</i> , <i>bella</i>	CLXXXIX , 475	1115
<i>Hypocrichtichthys</i> , <i>analis</i>	CCXXX , 582	1500
<i>Hypohomus</i> , <i>aurantiacus</i>	CLXVIII , 443	1043
<i>spinifer</i>	CLXVIII , 442	1040
<i>Hypomesus</i> , <i>oidius</i>	LXXXVII , 231	525
<i>pretiosus</i>	LXXXVII , 230	525
<i>Hypolectrus</i> , <i>genuala</i>	CLXXXIX , 497	1193
<i>unicolor</i> , <i>nligriceps</i>	CLXXXIX , 496	1193
<i>Hyporhynchus</i> , <i>brevirostris</i>	V , 18	41
<i>Hyporhamphus</i> , <i>roberti</i>	CXVII , 312	721
<i>unifasciatus</i>	CXVI , 311	720
<i>Hypsicometes</i> , <i>gobiooides</i>	CCXXXIII , 805	2294
<i>Hypsobenniuss</i> , <i>beauforti</i>	CCXXXIX , 823	2390
<i>ionthas</i>	CCCXVIII , 822	2388
<i>Hypsurus</i> , <i>euryi</i>	CCXXXI , 585	1508
<i>Hypsypops</i> , <i>rubicundus</i>	CCXXXIV , 591	1564
<i>Hystericarpus</i> , <i>traski</i>	CCXXVII , 577	1496
<i>Ilytrix</i> , <i>Diodon</i>	CCLXVI , 647	1745
<i>Icelinus</i> , <i>borealis</i>	CCLXXXIV , 687	1896
<i>Icelinus</i> , <i>canaliculatus</i>	CCLXXXVI , 693	1917
<i>spiniger</i>	CCLXXXVI , 692	1914
<i>Icichthys</i> , <i>lockingtoni</i>	CII , 406	969
<i>Ictalurus</i> , <i>angustifrons</i>	XXV , 57	2788
<i>fureatus</i>	XXV , 56	134
<i>punctatus</i>	XXV , 58	134
<i>Ictiobus</i> , <i>cypinella</i>	XXX , 70	163
<i>Ilyophis</i> , <i>brunneus</i>	LVI , 145	350
<i>imermis</i> , <i>Rabirinia</i>	CXCIX , 519	1274
<i>internalis</i> , <i>Histiobranchus</i>	LVII , 147	352
<i>infrenatus</i> , <i>Novaculichthys</i>	CCXL , 606	1616
<i>Inopsetta</i> , <i>ischyrus</i>	CCCLXXVI , 927	2641
<i>inhornatus</i> , <i>Microlepidotus</i>	CCX , 542	1341
<i>insignis</i> , <i>Schilbeoides</i>	XXVIII , 66	147
<i>interruptus</i> , <i>Arethoplitess</i>	CLVII , 420	991
<i>interictetus</i> , <i>Mystriophis</i>	LXV , 170	386
<i>intronaigra</i> , <i>Diercoleone</i>	CCCLVIII , 882	2522
<i>loglossus</i> , <i>calliumus</i>	CCXXXIII , 780	2193
<i>ionthas</i> , <i>Hypsoblennius</i>	CCXXXVIII , 822	2388
<i>ios</i> , <i>Clevelandia</i>	CCXXXVIII , 793	2254


* Plate by error labeled *Sardinella sardina*.

Genera and species.	Plate and figure.	Text page.
iovae, <i>Etheostoma</i> .	CLXXXIV, 460	1083
<i>Ipnois murrayi</i> .	LXXXIX, 239	546
<i>Iridens</i> , <i>Salmo</i> .	LXXXI, 216	500
<i>Irido bivittatus</i> .	CXXXIX, 601	1395
<i>dipilus</i> .	CCXL, 602	1397
<i>radiatus</i> .	CCXXXIX, 600	1390
<i>ischyra</i> , <i>Inopsetta</i> .	CCCXLXVI, 927	2611
<i>Istiophorus nigricans</i> .	CXXXVII, 376	891
<i>Istnus dekayi</i> .	VI, 21	48
<i>Itaiara</i> , <i>Promicrops</i> .	CLXXXV, 487, 487a, 487b	1162
<i>jaeki</i> , <i>Myoxocephalus</i> .	CCXCVI, 719	1977
<i>japonicus</i> , <i>Arctoscopus</i> .	CCCXXXIII, 807	2297
<i>Jenkinsi</i> , <i>Fundulus</i> .	CVI, 284	651
<i>Jessiae</i> , <i>Etheostoma</i> .	CLXXIV, 461	1081
<i>Xenocys</i> .	CCH, 526	1285
<i>Jordaniella floridana</i> .	CXII, 208	677
<i>Jordanell</i> , <i>Caulophryne</i> .	CCXCII, 957	2753
<i>Eslopsarum</i> .	CXXXIII, 335	793
<i>Etheostoma</i> .	CLXXXIII, 458	1079
<i>Hemilepidotus</i> .	CCXC, 703	1934
<i>Mycteroptera</i> .	CLXXXVIII, 403	1176
<i>Pantosteus</i> .	XXXI, 73	171
<i>Romquilius</i> .	CCCXXXII, 803	2289
<i>Jordania zonope</i> .	CCLXXXIII, 683	1884
<i>Joturus picturatus</i> .	CCXVIII, 348	821
<i>julie</i> , <i>Etheostoma</i> .	CLXXVI, 466	1063
<i>kanawha</i> , <i>Notropis</i> .	XLVI, 115	264
<i>Kathetostoma alboguttatum</i> .	CCCXXXIV, 809, 809a	2312
<i>kendalli</i> , <i>Vermi</i> .	LX, 159, 159a	375
<i>Kirtlandia vagrans</i> .	CXXIV, 336	794
<i>klamathensis</i> , <i>Alosa</i> .	LII, 133	3144
<i>Cottus</i> .	CCXCIII, 710	1955
<i>Kuhlia rupestris</i> .	CLXIII, 492	432
<i>Kyphosus sectatrix</i> .	CCNIX, 559	1387
<i>Labichthys carinatus</i> .	LIX, 155	368
<i>elongatus</i> .	LIX, 156	369
<i>lacera</i> , <i>Lagochila</i> .	XXVIII, 94, 94a	169
<i>Lachnolaimus maximus</i> .	CCXXXVII, 597	1579
<i>Lactophrys bleekeri</i> .	CCLXI, 638	1721
<i>tricornis</i> .	CCLXIII, 640, 640a	1723
<i>trigonus</i> .	CCLXI, 637	1722
<i>triqueter</i> .	COLXIII, 641	1728
<i>levigatus</i> , <i>Lactophrys</i> .	COLXIII, 641	1728
<i>Lagocephalus levigatus</i> .	CCCVIII, 94, 94a	169
<i>Lagochila lacera</i> .	CCCV, 552	1358
<i>Lagodon rhomboides</i> .	CXL, 382	903
<i>lalandi</i> , <i>Seriola</i> .	v, 17	38
<i>lamna</i> , <i>Carcharhinus</i> .	VI, 22	49
<i>Lammina cornifrons</i> .	XCI, 244	561
<i>Lampadenus speculiger</i> .	XCI, 243	558
<i>Lampanyctus crocodilinus</i> .	II, 5	13
<i>Lampetra aurea</i> .	III, 6	13
<i>spadicæ</i> .	CCCXLIV, 840	2438
<i>lampetraformis</i> , <i>Lumpenus</i> .	CXXVII, 573	1189
<i>lanceolatus</i> , <i>Eques</i> .	CXXVI, 565	1121
<i>Larinus argenteus</i> .	CCXXIX, 725, 725a	1983
<i>laticeps</i> , <i>Megalecottus</i> .	CCXLVI, 845	2416
<i>latifrons</i> , <i>Anarhichas</i> .	XXXI, 74	174
<i>latiphinis</i> , <i>Cottostomus</i> .	CXLI, 389	921
<i>tatus</i> , <i>Caranx</i> .	CCCI, 731	2011
<i>Leiocottus hirundo</i> .	CXXXIV, 712	1662
<i>leiopomus</i> , <i>Cottus</i> .	CXXIII, 569	1458
<i>Leiostomus xanthurus</i> .	CCCLXVI, 904	2556
<i>Lemonema barbatulum</i> .	CCCLXVI, 905	2557
<i>melanurum</i> .	VIII, 28; IX, 28a, 28b	62
<i>lentiginosus</i> , <i>Rhinobatos</i> .	CLXXIV, 462	1087
<i>lepidogenys</i> , <i>Etheostoma</i> .	CCLXXVII, 928	2613
<i>Lepidopsetta bilineata</i> .	CXXXVI, 373	886
<i>Lepidopus caninus</i> .	XXII, 49	119
<i>Lepisosteus platostomus</i> .	XXII, 50	111
<i>fristechus</i> .	CLIX, 425, 425a	1061
<i>Lepomis auritus</i> .		Sal
		macellus, Prior

Genera and species.	Plate and figure.	Text page.
<i>Laemis megalotis</i>	CLX, 426	1002
<i>pallidus</i>	CLX, 427	1005
<i>Lepophidium marmoratum</i>	CCCLIII, 806	2482
<i>profundorum</i>	CCCLIII, 807	2484
<i>Leptocephalus canthilimbatus</i>	LVI, 149	355
<i>conger</i>	LVI, 148	354
<i>Leptocottus armatus</i>	CCCI, 732	2012
<i>Leptops olivaris</i>	XXVII, 62	143
<i>lepturus</i> , <i>Anarhichas</i>	CCCXLVII, 847	2447
<i>Trichirrus</i>	CXXXVII, 375	889
<i>Letharchus velifer</i>	LXI, 160	375
<i>Lethotrema multiceps</i>	CCCXIII, 758	2101
<i>vincentus</i>	CCCXIV, 759	2101
<i>Leuciscus baileyi</i>	XLI, 105, 105a	238
<i>baileyi</i>	XLI, 102	232
<i>hydroploch</i>	XLI, 104	238
<i>lineatus</i>	XLI, 103	232
<i>sinslawi</i>	XLII, 106	2797
<i>Leucoglossus stilbius</i>	LXXXVII, 233	527
<i>Limanda aspera</i>	CCCLXXVII, 930	2645
<i>beamii</i>	CCCLXXVIII, 932	2646
<i>frarginae</i>	CCCLXXVII, 929	2644
<i>proboscidea</i>	CCCLXXVIII, 931	2645
<i>Lineatus</i> , <i>Achirus</i>	CCCLXXXVI, 947	2697
<i>Lineatus</i>	XLI, 103	232
<i>Roccus</i>	CLXXX, 478	1132
<i>Liopsetta glacialis</i>	CCCLXXX, 935	2649
<i>putnami</i>	CCCLXXX, 936	2650
<i>liorus</i> , <i>Chasmistes</i>	XXXIV, 85	183
<i>Liparis agassizii</i>	CCCVII, 765	2121
<i>cyclops</i>	CCCVII, 764, 764a	2118
<i>dearrayi</i>	CCCVII, 766, 766a	2124
<i>Lipogenys gillii</i>	XCIX, 266	619
<i>Lophotes surinamensis</i>	CXCIV, 510	1235
<i>lockingtoni</i> , <i>Teichthys</i>	CLI, 406	969
<i>longispathum</i> , <i>Peristedion</i>	CCXXI, 775	2178
<i>Lophius piscatorius</i>	CCCLXXXVIII, 952	2713
<i>Lophogobius cyprinoides</i>	CCXXXVII, 780	229
<i>Lophosetta maculata</i>	CCCLXXII, 938	2660
<i>loredo</i> , <i>Gramma</i>	CXCH, 508	1229
<i>Lota lota</i>	CCCLXIV, 897	2550
<i>Lotella maxillaris</i>	CCCLXIII, 895	2546
<i>Lucania parva</i>	CIX, 292	665
<i>lucia</i> , <i>Fundulus</i>	CVII, 286	654
<i>lucida</i> , <i>Etheostoma</i>	XCL, 246	565
<i>lucius</i> , <i>Lucius</i>	C, 269, 269a	628
<i>Lucius lucius</i>	C, 269, 269a	628
<i>masquinongy</i>	C, 270	629
<i>Laumentis hompetraeformis</i>	CCXLIV, 840	2478
<i>mackayi</i>	CCXLIV, 839	2436
<i>limpus</i> , <i>Cyclopterus</i>	CCXLII, 757	2006
<i>limpus</i> , <i>Anarhichas</i>	CCXLVII, 846	2446
<i>Lycenchelys paxillns</i>	CCCLI, 860-860a	2471
<i>verrillii</i>	CCCLI, 859	2470
<i>Lycodalepis polaris</i>	CCCLI, 857	2468
<i>turneri</i>	CCCLI, 858	2469
<i>Lycodapus dermatinus</i>	CCCLIV, 870	2492
<i>Lycodes frigidus</i>	CCCL, 856	2465
<i>perspicillum</i>	CCCL, 855	2465
<i>reticulatus</i>	CCXLIX, 854	2465
<i>zonorehus</i>	CCXLIX, 853	2464
<i>Lycodontis moringa</i>	LXV, 171	395
<i>Lycodonus mirabilis</i>	CCCLI, 862	2474
<i>Lycoperca aleutensis</i>	CCXLVI, 844	2444
<i>Lycophena barbatum</i>	CCCLI, 863	2474
<i>Lyosphera globosa</i>	CCLXVII, 649, 649a	1751
<i>Lythrypnus opalescens</i>	CCVII, 536	1312
<i>macarellus</i> , <i>Dicapterus</i>	CXL, 383	909
<i>macdonaldi</i> , <i>Fundulus</i>	CVI, 283	651
<i>Nanobrachium</i>	XCI, 245	563
<i>Notropis</i>	XLIX, 123	284
<i>Salmo clarkii</i>	LXXXI, 214	{ 497 2819
<i>macellus</i> , <i>Prionistius</i>	CCLXXXIX, 700	1928



**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

45
48
28
32
123
36
22
20
18

10

Genera and species.	Plate and figure.	Text page.
mackayi, <i>Lumpenus</i>	CCXLIV, 839	2136
mackenzi, <i>Stenodus</i>	LXXVII, 204	474
macrocephalus, <i>Hadropterus</i>	CLXVI, 437	1031
macrocheilus, <i>Catostomus</i>	XXXIII, 81	158
macrochir, <i>Aldrovandia</i>	XCVIII, 262	609
macrognathum, <i>Ophistognathus</i>	CCCXXXI, 800a	2281
macrolepis, <i>Pontinus</i>	CULXXXVII, 672	1855
macrophtalmus, <i>Aprión</i>	CXI, 523	1280
macrops, <i>Citharichthys</i>	CCLXXXV, 944	2084
macropterus, <i>Centrarchus</i>	CLV, 417	988
macrostomus, <i>Haemulon</i>	CXIV, 529	1296
maculata, <i>Lophopsetta</i>	CCCXXXII, 938	2669
maculatofasciatus, <i>Paralabrax</i>	CXC, 498	1196
maculatus, <i>Aulostomus</i>	CXX, 324	754
<i>Cryptacanthodes</i>	CCXLV, 843	2443
<i>Dormitator</i>	CCCXXIX, 782	2196
<i>Noterichthys</i>	II, 7	17
<i>Scomberomorus</i>	CXXXIV, 368	874
<i>Spherooides</i>	CCLXIV, 644	1733
<i>Upeneus</i>	CXXXII, 362	858
maculosa, <i>Lota</i>	CCCXLIX, 897	2550
maculosus, <i>Oligocottus</i>	CCCI, 733	2033
madeirensis, <i>Ceratoscopelus</i>	XC, 242	557
maeandrius, <i>Caranxchus</i>	CXXXV, 371	2328
magistralis, <i>Epinnula</i>	CCCXXXV, 812	228
magalis, <i>Fundulus</i>	CI, 271; 272, 2, 16	633
malacosteus, <i>niger</i>	XCV, 256	533
malardi, <i>Uraepterus</i>	CCLXXIV, 894	2435
maliger, <i>Sebastodes</i>	CCLXXXIV, 664	1822
Mallotus villosus.....	LXXXV, 225	520
madras, <i>Salvelinus</i>	LXXXII, 219	507
manatus, <i>Barathrodemus</i>	CCCLVII, 880	2317
Mancalias shufeldti.....	CXXEC, 953	2730
Manta birostris.....	XVIII, 39	32
marconis, <i>Hybopsis natalensis</i>	LII, 136	716
marginata, <i>Rissoa</i>	CCCLII, 868	2489
marginatus, <i>Neobathythites</i>	CCLVII, 878	2315
<i>Sympularis</i>	CI XXXVII, 949	2706
mariannus, <i>Flammeo</i>	CXXXI, 350	852
marinus, <i>Petromyzon</i>	1, 3	10
<i>Sebastes</i>	CCLXVII, 652	1760
marmoratum, <i>Lepophidium</i>	CCCLII, 836	2482
marmoratus, <i>Scorpaenichthys</i>	CCLXXXIII, 685	1889
masquinongy, <i>Lucius</i>	C, 270	629
maxillaris, <i>Lotilia</i>	CCCLXIII, 835	2546
maxillingua, <i>Exoglossum</i>	LIV, 140	327
maxillosa, <i>Gnathopis</i>	CCXXXI, 861	2284
maximus, <i>Cetorhinus</i>	VII, 23	51
<i>Lachnolaimus</i>	CCXXXVII, 597	1579
meadie, <i>Ulocentra</i>	CLXIX, 447	2832
meanlyi, <i>Ruscarius</i>	CCLXXXV, 690	1908
Medialuna californiensis.....	CCXXX, 560	1391
medlocris, <i>Pomolobus</i>	LXXI, 188	425
mediterraneus, <i>Hoplostethus</i>	CXXX, 354	837
meduseola, <i>Caranx</i>	CXLIII, 390	924
medusophagus, <i>Schedophilus</i>	CLI, 407	970
Megalocottus laticeps.....	CCXCIX, 725, 725a	1988
<i>platycephalus</i>	CCXCIX, 724, 724a	1987
megulotis, <i>Lepomis</i>	CLX, 420	1002
Melanogrammus nigletinus.....	CCCLXI, 892, 892a	2542
melanops, <i>Minytrema</i>	XXXVI, 90	187
<i>Sebastodes</i>	CCLXIX, 654	1782
melanostictus, <i>Psettichthys</i>	CCCLXXIII, 921	2618
Melmostigma pannulus.....	CCCLII, 865	2479
melanurus, <i>Leuconema</i>	CCCLXV, 905	2557
melas, <i>Amelurus</i>	XXVI, 60	141
Melletes papilio.....	CCLXXXIX, 702	1932
Menidia gracilis beryllina.....	CXXXIV, 338	797
<i>peninsulae</i>	CXXXIV, 337	797
Menticirrhus americanus.....	CCXXXV, 572	1474
Merluccius productus.....	CCCLIX, 884	2531
merriami, <i>Empetrichthys</i>	CX, 294-294d	667

Macrourus
metaphysus
metodon, P.
Microgadus

Microlepidotus
microlepidotid
microtropis, M.
Microptera L.
Micropteron interpus, Cau.
Micropterus

Microspathe

milberti, Gal.
miles, Porop.
mioletta, Po.
minatus, Sel.
minima, Abe.
Minytrema n.
mirabilis, Ly.
mirurus, Schil.
Mola mola...
mola, Mola.
Monacanthus

mona, Stephan.
monopterygin
montanus, Th.
monticola, Ag.
mordax, Osm.
morina, Lyc.
morio, Epinep.
Morone americana
Moxostoma at.
ru

mucoinosus, Neol.
Mugil cephalus
curema
hosipes
mulleri, Penp.
Mulloidichthys
Mullus auratus
multifasciatus
muraena, Caleo.
Murana insula
reticulata

murrayi, Trinop.
muscaram, Rin.
mutiens, Letho.
Myctoperca

1

Myctophium op.
Mylochelius ca.
myops, Trachin.
Myoxocephalus

Genera and species.	Plate and figure.	Text page.
<i>Mesognathus chaetodon</i>	CLVIII, 423	995
<i>metathetus</i> , <i>Neropis</i>	L, 128	297
<i>microdon</i> , <i>Pseudorriakis</i>	IV, 14	27
<i>Microgadus proximus</i>	CCCLX, 889	2539
<i>tomcod</i>	CCCLX, 899	2540
<i>Microlepidotus inornatus</i>	CCX, 542	1341
<i>microlepidotus</i> , <i>Orthodou</i>	XXXIX, 96	207
<i>microlepis</i> , <i>Mycteroptera</i>	CLXXXVIII, 494	1177
<i>Microtia fonticola</i>	CLXXVII, 470	1104
<i>Micropegon undulatus</i>	CCXXXIV, 570	1461
<i>micropterus</i> , <i>Caulolatilus</i>	CCCCXXX, 799	2277
<i>Micropterus dolomieu</i>	CLXII, 430a	1011
<i>salmoideus</i>	CLXIII, 431	1012
<i>Microstethodon bairdii</i>	CCXXXV, 592	1566
<i>chrysurus</i>	CCXXXVI, 593	1567
<i>dorsalis</i>	CCXXXVI, 594	1568
<i>milleri</i> , <i>Galeichthys</i>	XXIII, 53	128
<i>miles</i> , <i>Pterogadus</i>	CCCLVIII, 881	2520
<i>miniatum</i> , <i>Peristedion</i>	CCCXI, 774	2178
<i>minutus</i> , <i>Sebastodes</i>	CCLXXXIII, 662	1794
<i>minima</i> , <i>Abeona</i>	CCXXXVIII, 578	1497
<i>Minytrema melanops</i>	XXXVI, 90	187
<i>mirabilis</i> , <i>Lycodonus</i>	CCCLII, 862	2474
<i>mirinus</i> , <i>Selildodeas</i>	XXIX, 68	148
<i>Mola mola</i>	CCLXVII, 650	1753
<i>mola</i> , <i>Mola</i>	CCLXVII, 650	1753
<i>Monacanthus ciliatus</i>	CCLIX, 633	1714
<i>hispidus</i>	CCLIX, 634	1715
<i>monae</i> , <i>Stephamobery</i>	CXXIX, 353	836
<i>monoptychius</i> , <i>Aspidophoroides</i>	CCCXII, 729, 756a	2091
<i>Pleurogrammus</i>	CCLXXIX, 675	1864
<i>montanus</i> , <i>Thymallus tricolor</i>	LXXXIV, 224a	{ 519 2871
<i>monticola</i> , <i>Agonostomus</i>	CXXVII, 347	819
<i>morlax</i> , <i>Osmernus</i>	LXXXVI, 228	523
<i>moringa</i> , <i>Lycodontis</i>	LXV, 171	395
<i>morio</i> , <i>Epinephelus</i>	CLXXXIV, 485	1160
<i>Morone americana</i>	CLXXXI, 479	1134
<i>Moxostoma austriatum</i>	XXXVII, 92	192
<i>congestum</i>	XXXVI, 91	192
<i>rupicartes</i>	XXXVII, 93	196
<i>mucosus</i> , <i>Neoliparis</i>	CCCXV, 761, 761a	2111
<i>Mugil cephalus</i>	CXXVI, 343	811
<i>enrema</i>	CXXVI, 344	813
<i>bosporus</i>	CXXVII, 345	814
<i>mulieri</i> , <i>Pempheris</i>	CLIII, 412	978
<i>Mullus rathbuni</i>	CXXXII, 361	857
<i>Mullus auratus</i>	CXXXII, 360	856
<i>multifasciatus</i> , <i>Pronotogrammus</i>	CXCH, 506	1226
<i>aurea</i> , <i>Calcechelys</i>	LXIII, 164	378
<i>Murina insularum</i>	LXV, 172	400
<i>reticula</i>	LXVI, 173	401
<i>murrayi</i> , <i>Ipnops</i>	LXXXIX, 239	547
<i>naseatum</i> , <i>Rimicola</i>	CCXXXV, 812	2338
<i>niutiens</i> , <i>Lethotrema</i>	CCCXIII, 758	2101
<i>Mycteroptera bonaci</i>	CLXXXVII, 492	1174
<i>bonlengeri</i>	CLXXXVII, 490	1171
<i>falcata phenax</i>	CLXXXVIII, 495	1185
<i>jordani</i>	CLXXXVIII, 493	1176
<i>microlepis</i>	CLXXXVIII, 494	1177
<i>venenosus</i>	CLXXXVIII, 491	1172
<i>Myctophum opalinum</i>	XCI, 247	571
<i>Mylocheilus caeruleus</i>	XL, 99	219
<i>myops</i> , <i>Trachinocéphalus</i>	LXXXVIII, 235	533
<i>Myoxocephalus anens</i>	CCXCV, 715, 715a	1972
<i>axillaris</i>	CCXCVII, 721	1980
<i>jaak</i>	CCXCVI, 719	1977
<i>niger</i>	CCXCVIII, 723	1985
<i>octodecemspiniferus</i>	CCXCVI, 717	1976
<i>polyacanthocephalus</i>	CCXCVI, 718	1976
<i>scorpius</i>	CCXCV, 710	1974
<i>stelleri</i>	CCXCVIII, 722, 722a	1981
<i>verrucosus</i>	CCXCVII, 720, 720a	1979

Genera and species.	Plate and figure.	Text page.
<i>Myrichthys pantostigmatus</i>	LXII, 162	280
<i>tigrinus</i>	LXI, 161	376
<i>mystes</i> , <i>Scorpaena</i>	CCLXXVII, 670	1849
<i>mystinus</i> , <i>Sebastodes</i>	CCLXX, 656	1784
<i>Mystriophis intertinctus</i>	LXV, 170	386
<i>niamayenii</i> , <i>Cristivomer</i>	LXXXIII, 217	504
<i>Nannobrachium macdonaldi</i>	XCI, 245	503
<i>Narcine brasiliensis</i>	XIII, 35, 35a	78
<i>narmari</i> , <i>Aetobatus</i>	XV, XVI, 37, 37a	88
<i>Nauernes dector</i>	CXXXIX, 379	900
<i>nauernes</i> , <i>Echeneis</i>	CCCXXIX, 796	2299
<i>nayaga</i> , <i>Eleginops</i>	CCCLX, 888	2537
<i>nebulosus</i> , <i>Cynoscion</i>	CCXXI, 563	1409
<i>Nematiatius pectoralis</i>	CXXXVIII, 377	895
<i>Nemichthys avocetta</i>	LX, 157, 157a, 157b	309
<i>Neobythites gilli</i>	CCCLVII, 877	2512
<i>marginalis</i>	CCCLVII, 878	2513
<i>Neoclinus satiricus</i>	CCCXXXVI, 816	2355
<i>Neoliparis calyodon</i>	CCCXIV, 760, 760a	2110
<i>florie</i>	CCCXV, 762	2111
<i>greeni</i>	CCCXVI, 763, 763a	2112
<i>mucosus</i>	CCCXV, 761, 761a	2111
<i>Neouerenis analis</i>	CXCVIII, 517	1265
<i>apodus</i>	CXCVII, 515	1258
<i>aya</i>	CXCVII, 516	1264
<i>synagris</i>	CXCVIII, 518	1266
<i>nerka</i> , <i>Oncorhynchus</i>	LXXXVIII, 207, 207a, 207b	71
<i>niger</i> , <i>Centrolophus</i>	CXLIX, 403	963
<i>Chiasmodon</i>	CCCXXXII, 804	2291
<i>Malacostenus</i>	XCV, 256	593
<i>Myoxocephalus</i>	CXCIVIII, 723	1985
<i>nigricans</i> , <i>Hypoplectrus unicolor</i>	CLXXXIX, 496	1193
<i>Istiophorus</i>	CXXXII, 376	891
<i>nigripinnis</i> , <i>Argyrosomus</i>	LXXXVI, 203	472
<i>Bathygymnus</i>	CCCXI, 753	207
<i>nigrostris</i> , <i>Chetodon</i>	CCXLVIII, 620	1673
<i>nigrita</i> , <i>Garrupa</i>	CLXXXV, 486	1161
<i>nigrocinctus</i> , <i>Sebastodes</i>	CLXXXVI, 667	1827
<i>nigrum</i> , <i>Boleosoma</i>	CLXX, 450	1056
<i>olmstedi</i> , <i>Boleosoma</i>	CLXXI, 451	1057
<i>notomis</i> , <i>Notropis</i>	XLVII, 178	268
<i>nocturnus</i> , <i>Schilbeoides</i>	XXVII, 64	146
<i>normalis</i> , <i>Bassozetus</i>	CCCLVI, 875	2507
<i>Notacanthus analis</i>	XCVIII, 264	615
<i>phasganurus</i>	XCVIII, 265	616
<i>notatus</i> , <i>Fundulus</i>	CVIII, 289	659
<i>notemigonooides</i> , <i>Notropis</i>	L, 127	292
<i>notlus</i> , <i>Cynoscion</i>	CCXX, 561	1406
<i>Notorhynchus maculatus</i>	II, 7	17
<i>notospilotus</i> , <i>Astrolytes</i>	CCLXXXIV, 688, 688a	1899
<i>Notropis acuticeps</i>	XLV, 112	258
<i>cayuga atrocaudalis</i>	XLVI, 114	260
<i>chihualhua</i>	XLVI, 116	265
<i>coccogenis</i>	XLIX, 124	281
<i>galacturus</i>	XLIX, 122	279
<i>hudsonius</i>	XLVII, 119	269
<i>sulidamus</i>	XLVII, 120	270
<i>kanawha</i>	XLVI, 115	264
<i>maedonaldi</i>	XLIX, 123	284
<i>metallifer</i>	XLVI, 116	265
<i>noconis</i>	XLIX, 124	281
<i>notemigonooides</i>	XLVIII, 122	279
<i>nux</i>	XLVII, 119	269
<i>swaini</i>	XLVII, 120	270
<i>telescopus</i>	XLV, 113	2766
<i>welaka</i>	XLVIII, 121	278
<i>whippini</i>	CVIII, 288	656
<i>nottii</i> , <i>Fundulus</i>	XXVII, 63	144
<i>Nothrus flavus</i>	CCXL, 606	1615
<i>Novaculichthys infirmus</i>	CCXL, 605	1615
<i>ventralis</i>	CCCXXXVII, 819	2373
<i>nox</i> , <i>Anchenopterus</i>	CCCXL, 820	2410
<i>nugator</i> , <i>Bryostema</i>		

Genera and species.	Plate and figure.	Text page.
<i>nux</i> , <i>Notropis</i>	<i>NXLVII</i> , 117	267
<i>obeyense</i> , <i>Etheostoma</i>	<i>CLXXV</i> , 463	1092
<i>oblongus</i> , <i>Paralichthys</i>	<i>CCCLXXXIV</i> , 924	2632
<i>ocea</i> , <i>Culorhynchus</i>	<i>CCCLXIX</i> , 913	2588
<i>oreat dodecaedron</i>	<i>CCCVIII</i> , 743	2044
<i>occidentalis</i> , <i>Catostomus</i>	<i>XXXIII</i> , 79	178
<i>Tetronarce</i>	<i>XI</i> , 33	77
<i>oceaniensis</i> , <i>Gobius</i>	<i>CCCXXVII</i> , 789a	2230
<i>ocularis</i> , <i>Fundulus</i>	<i>CI</i> , 274	642
<i>ocellata</i> , <i>Raja</i>	<i>X</i> , 30	68
<i>ocellatus</i> , <i>Chaetodon</i>	<i>CCXLIX</i> , 621	1674
<i>Ophichthus</i>	<i>CCI</i> , 524	1282
<i>Platophrys</i>	<i>LXIV</i> , 169	383
<i>Solenops</i>	<i>CCCLXXXII</i> , 939	2663
<i>Zenopsis</i>	<i>CCXXII</i> , 567	1453
<i>octodecemspinosis</i> , <i>Myoxocephalus</i>	<i>CCXLVI</i> , 618	1660
<i>octogrammus</i> , <i>Hexagrammos</i>	<i>CCXXVI</i> , 717	1976
<i>octonemus</i> , <i>Polydactylus</i>	<i>CCLXXX</i> , 677	1869
<i>oculatus</i> , <i>Etelis</i>	<i>CXXVII</i> , 350	830
<i>Ocyurus chrysurus</i>	<i>CXCIX</i> , 529	1275
<i>ogcocephalus vespertilio</i>	<i>CCCXII</i> , 958a, 958b	2737
<i>Ogilbia cayorum</i>	<i>CCCIV</i> , 873	2503
<i>ventralis</i>	<i>CCCIV</i> , 872	2503
<i>olfersi</i> , <i>Argyropelecus</i>	<i>NCVII</i> , 261, 261a	604
<i>olidus</i> , <i>Hypomesus</i>	<i>LXXXVII</i> , 221	525
<i>oligocottus maculosus</i>	<i>CCCI</i> , 733	2013
<i>Oligoplites saurus</i>	<i>CXXXVIII</i> , 378	898
<i>olmstromus</i> , <i>Gerres</i>	<i>CCXVII</i> , 557	1376
<i>olivaris</i> , <i>Leptos</i>	<i>XXVII</i> , 62	143
<i>olmstedii</i> , <i>Boleosoma nigrum</i>	<i>CLXXI</i> , 451	1057
<i>omostigmum</i> , <i>Otophilium</i>	<i>CCCLIV</i> , 869	2490
<i>Oncoctonus quadricornis</i>	<i>CCC</i> , 728	2001
<i>Oncorhynchus gorbuscha</i>	<i>LXXVII</i> , 205	478
<i>nerka</i>	<i>LXXVII</i> , 207a	481
<i>tschawytscha</i>	<i>LXXVII</i> , 206	479
<i>onitis</i> , <i>Tantoga</i>	<i>CCXXXVII</i> , 596	1578
<i>opalescens</i> , <i>Lythrypnus</i>	<i>CCVII</i> , 536	1312
<i>opalimum</i> , <i>Myctophum</i>	<i>XCI</i> , 247	571
<i>Ophichthus guttifer</i>	<i>LXIV</i> , 168	383
<i>ocellatus</i>	<i>LXIV</i> , 169	383
<i>Ophidion elongatus</i>	<i>CCCLXXXII</i> , 681	1875
<i>Opistognathus macrognathus</i>	<i>CCXXXI</i> , 800, 800a	2281
<i>Opsanus pardus</i>	<i>CCXXXV</i> , 810	2316
<i>Opsopoeodus bellhorni</i>	<i>XLIV</i> , 110	248
<i>emilia</i>	<i>XLIV</i> , 109	248
<i>osculus</i>	<i>XLIV</i> , 108	248
<i>opuissa</i> , <i>Salvelinus</i>	<i>LXXXIII</i> , 221	514
<i>oregonensis</i> , <i>Ptychocheilus</i>	<i>XLI</i> , 101	224
<i>ornatus</i> , <i>Pholis</i>	<i>CCXLII</i> , 833	2419
<i>Orthodon microlepidotus</i>	<i>XXXIX</i> , 96	207
<i>Orthoprists chrysopterus</i>	<i>CCC</i> , 541	1338
<i>reddingi</i>	<i>CCIX</i> , 540	1336
<i>osculus</i> , <i>Opsopoeodus</i>	<i>XLIV</i> , 108	248
<i>Osmerus dentex</i>	<i>LXXXVI</i> , 229	524
<i>mordax</i>	<i>LXXXVI</i> , 228	523
<i>thaleichthys</i>	<i>LXXXV</i> , 227	522
<i>osteobir</i> , <i>Rhombocheirus</i>	<i>CCXXX</i> , 798	2273
<i>otakii</i> , <i>Hexagrammos</i>	<i>CCLXXXI</i> , 680	1867
<i>Otophidium omostigmum</i>	<i>CCCLIV</i> , 869	2490
<i>Otrynter caprimus</i>	<i>CCXI</i> , 543	1345
<i>ovigerum</i> , <i>Bathyphasma</i>	<i>CCCVIII</i> , 767	2128
<i>Ovoides setosus</i>	<i>CCLXV</i> , 646	1739
<i>oxybleius pictus</i>	<i>CCLXXXIII</i> , 682	1878
<i>oxyrhynchus</i> , <i>Aelposser sturio</i>	<i>XX</i> , 45	105
<i>pacificus</i> , <i>Thaleichthys</i>	<i>LXXXV</i> , 226	521
<i>paged</i> , <i>Etheostoma</i>	<i>CLXXV</i> , 464	1092
<i>pagrus</i> , <i>Pagrus</i>	<i>CCXV</i> , 551	1356
<i>Pagrus pagrus</i>	<i>CCXV</i> , 551	1356
<i>pallasi</i> , <i>Chipea</i>	<i>LXX</i> , 186	422
<i>Pallasina barbata</i>	<i>CCCVIII</i> , 744	20, 9
<i>pallidus</i> , <i>Fu duhus</i>	<i>CI</i> , 272	638
<i>Lepomis</i>	<i>CLX</i> , 427	1005
<i>pammelas</i> , <i>Melanostigma</i>	<i>CCCLI</i> , 865	2479

Genera and species.	Plate and figure.	Text page.
<i>panamensis</i> , <i>Azevia</i> .	<i>CCCLXXXIV</i> , 942	2677
<i>Pantosteus jordani</i> .	<i>XXXI</i> , 73	171
<i>pautostigmatus</i> , <i>Myrichthys</i> .	<i>LXII</i> , 162	2802
<i>papilio</i> , <i>Melletes</i> .	<i>CCCLXXXIX</i> , 702	1932
<i>papillosum</i> , <i>Syacium</i> .	<i>CCCLXXXIII</i> , 941	2671
<i>paradoxa</i> , <i>Garmannia</i> .	<i>CCCLXXVII</i> , 700	2332
<i>paradoxus</i> , <i>Psychrolutes</i> .	<i>CXXVI</i> , 740	2026
<i>Paralabrax humeralis</i> .	<i>CXC</i> , 499	1196
<i>maculatofasciatus</i> .	<i>CXC</i> , 498	1196
<i>Paralepis eoregonoides</i> .	<i>XCVI</i> , 260	602
<i>Paralichthys dentatus</i> .	<i>CCCLXXIII</i> , 922	2629
<i>oblongus</i> .	<i>CCCLXXIV</i> , 923	2631
<i>squamiferus</i> .	<i>CXCH</i> , 504	1221
<i>Paranthias furcifer</i> .	<i>LVI</i> , 144	349
<i>parasitens</i> , <i>Simenichelys</i> .	<i>CCXXXV</i> , 810	2516
<i>pardus</i> , <i>Opsanus</i> .	<i>CCXXXIX</i> , 673	1860
<i>parmatum</i> , <i>Setarches</i> .	<i>CCCV</i> , 530	1297
<i>parra</i> , <i>Hemulon</i> .	<i>CCXXXVIII</i> , 599	1586
<i>parra</i> , <i>Clepticus</i> .	<i>CL</i> , 404	965
<i>paru</i> , <i>Peprilus</i> .	<i>CIX</i> , 292	665
<i>parva</i> , <i>Lucania</i> .	<i>CCCLII</i> , 860, 860a, 860b, 860c, 860d,	2471
<i>paxillus</i> , <i>Lycenchelys</i> .	<i>VIII</i> , 27	66
<i>pectinatus</i> , <i>Pristis</i> .	<i>XCIX</i> , 267	621
<i>pectoralis</i> , <i>Dallia</i> .	<i>CCXXXVIII</i> , 377	895
<i>Nematistius</i> .	<i>CLXXII</i> , 454	1063
<i>pellucida clara</i> , <i>Ammocrypta</i> .	<i>CLIII</i> , 412	978
<i>Pempheris mulleri</i> .	<i>CLIII</i> , 413	979
<i>poeyi</i> .	<i>LXIII</i> , 166, 166a	379
<i>peninsulae</i> , <i>Bascanichthys</i> .	<i>CCXXIV</i> , 337	797
<i>Menidia</i> .	<i>CCXIV</i> , 549	1354
<i>penna</i> , <i>Calamus</i> .	<i>CL</i> , 404	965
<i>Peprilus paru</i> .	<i>CLXV</i> , 436, 436a	1026
<i>Perca flavescens</i> .	<i>CXXI</i> , 329	784
<i>Percina caprodes</i> .	<i>CCXXII</i> , 776	2170
<i>Percopsis guttatus</i> .	<i>CCXXI</i> , 775	2178
<i>Peristedion gracile</i> .	<i>CCXXI</i> , 774	2178
<i>longispathum</i> .	<i>CCXXII</i> , 777, 777a	2180
<i>miniatum</i> .	<i>CCXXII</i> , 709	1955
<i>platycephalum</i> .	<i>CCCL</i> , 855	2465
<i>perplexus</i> , <i>Cottus</i> .	<i>LXXXIV</i> , 196	442
<i>perspicillum</i> , <i>Lycodes</i> .	1, 3	10
<i>perthecatus</i> , <i>Stolephorus</i> .	<i>CCXXX</i> , 583	1506
<i>Petromyzon marinus</i> .	<i>XCVIII</i> , 265	616
<i>Phanerodon furcatus</i> .	<i>LI</i> , 130	304
<i>phaeagoros</i> , <i>Notacanthus</i> .	<i>CLXXXVIII</i> , 495	1185
<i>Phenacobius uranops</i> .	<i>CXCI</i> , 501	1291
<i>phenax</i> , <i>Mycteroptera falcata</i> .	III, 10	
<i>philadelphicus</i> , <i>Centropristes</i> .	<i>CCCCXIV</i> , 781	2194
<i>philippii</i> , <i>Heterodontus</i> .	<i>CCXLII</i> , 831	2416
<i>Philypnus dormitor</i> .	<i>CCXLII</i> , 832	2419
<i>Pholidichthys dolichogaster</i> .	<i>CCXLII</i> , 833	2419
<i>gunnellus</i> .	<i>XCV</i> , 255	591
<i>ornatus</i> .	<i>CCCLXIII</i> , 896	2547
<i>Photoneutes gracilis</i> .	<i>CXXXVIII</i> , 348	821
<i>Physciulus fulvus</i> .	<i>CCCLXXXIX</i> , 953	2726
<i>pichardi</i> , <i>Joturus</i> .	<i>CCLXXXII</i> , 682	1878
<i>pictus</i> , <i>Chimaera</i> .	<i>CLXXVII</i> , 472	1169
<i>Oxyblepus</i> .	<i>CCXXXVIII</i> , 598	1585
<i>pigmentarius</i> , <i>Apogon</i> .	<i>CCLXXXVII</i> , 606	1923
<i>Pimeletometaepon pucher</i> .	<i>LVI</i> , 146	351
<i>plugell</i> , <i>Triglops</i> .	<i>CCLXXXIII</i> , 661	1793
<i>pinnatus</i> , <i>Synaphobranchus</i> .	<i>CCCLXXXVIII</i> , 952	2713
<i>pimiliger</i> , <i>Sebastodes</i> .	<i>CCCXXXV</i> , 783	2290
<i>piscatorius</i> , <i>Lophius</i> .	<i>LXII</i> , 163	377
<i>pisounis</i> , <i>Electris</i> .	<i>CCL</i> , 729, 729a, 729b	2096
<i>Pisodonophis erneutifer</i> .	<i>XXXIII</i> , 82	198
<i>pistilliger</i> , <i>Gymnoanthus</i> .	<i>CCCLXXXVIII</i> , 950	2710
<i>Placopharynx duquesnei</i> .	<i>CCCLXXXII</i> , 910	2614
<i>plagiura</i> , <i>Syphurus</i> .	<i>CCCLXXXI</i> , 937, 937a	2652
<i>platessoides</i> , <i>Hippoglossoides</i> .	<i>CCCLXXXII</i> , 939	2663
<i>Platichthys stellatus</i> .	<i>XXI</i> , 48	107
<i>Platophryns ocellatus</i> .		
<i>platyrhynchus</i> , <i>Scaphirhynchus</i> .		

platycephalus
platycephalid
platycephalidae

Platycephalidae

Pleuronectidae

Pleuronichthys

Plumieriidae

Podorhynchidae

Poeciliidae

Poeciliopsis

Pomatomidae

Pomolobus

Pomolobus aspi-

chus

Pomotidae

Pomoxis annularis

Ponticidae

Ponticulus mac-

erulus

Porichthys po-

rigatus

Porogadus mil-

lifer

Poromotus tria-

porosus

Pottsi

Etheostoma

presidium

Priacanthidae

Priacanthus ma-

rcularis

Priacanthus

Genera and species.	Plate and figure.	Text page.
<i>platostomus</i> , <i>Lepisosteus</i>	XXII, 49	110
<i>percephalum</i> , <i>Peristedion</i>	CCCXII, 777, 777a	2180
<i>platycephalus</i> , <i>Amarurus</i>	XXVII, 61	142
<i>Megalocottus</i>	CXCIX, 724, 724a	1987
<i>Platygobio gracilis</i>	LIV, 139	326
<i>Plectrocnemis crassiceps</i>	CXXX, 356	843
<i>suborbitalis</i>	CXXX, 355	841
	LXXX, 211	496
<i>pleuriticus</i> , <i>Salmoneus clarkii</i>		2819
<i>Pleuragrammus monopterygius</i>	CCLXXIX, 675	1864
<i>Pleuronectes quadrituberculatus</i>	CCCLXXIX, 934	2648
<i>Pleuronichthys decurrens</i>	CCCLXXV, 926	2637
<i>pumifer</i> , <i>Hemulon</i>	CCV, 532	1304
<i>pacatello</i> , <i>Catostomus</i>	XXXII, 76	175
<i>Podotrichus acipenserinus</i>	CCCIX, 746	2061
<i>acepiter</i>	CCCVII, 745, 745a	2055
<i>veterinus</i>	CCCIX, 747, 747a	2063
<i>Poecilia presidensis</i>	CXIV, 303	697
<i>poeyi</i> , <i>Pempheris</i>	CLIII, 413	979
<i>Pogonias chrysotaenia</i>	CCXXXV, 573	1482
<i>polaris</i> , <i>Lycodeslepis</i>	CCCL, 857	2468
<i>Pollachius virens</i>	CCCLIX, 886	2534
<i>polyacanthocephalum</i> , <i>Bryostoma</i>	CCCXLII, 828	2408
<i>polyacanthocephalus</i> , <i>Myoxocephalus</i>	CXCVI, 718	1976
<i>Polydactylus octonemus</i>	CXXVIII, 350	830
<i>Polyodon spathula</i>	XX, 43, 43a	101
<i>Polyprion americanus</i>	CLXXXI, 480, 480a	1139
<i>Polypterus bichir</i>	1, 2	...
<i>Pomacanthus arcuatus</i>	CCLI, 623	1679
<i>zonopectus</i>	CCLII, 624	1681
<i>Pomatomus saltatrix</i>	XLVIII, 400	946
<i>Pomolobus testivalis</i>	LXXI, 190	426
<i>chrysochloris</i>	LXX, 187	425
<i>medioocellatus</i>	LXXI, 188	425
<i>pseudoharengus</i>	LXXI, 189	426
<i>pomotis</i> , <i>Acantharchus</i>	CLV, 418	989
<i>Pomoxis annularis</i>	CLIV, 415	987
<i>sparoides</i>	CLIV, 416	987
<i>Ponticus macrolepis</i>	CCLXXVIII, 672	1855
<i>Porichthys porosissimus</i>	CCXXXV, 811	2319
<i>Poregadus miles</i>	CCCLVIII, 881	2520
<i>Poronotus triacanthus</i>	CL, 405	967
<i>porosissimus</i> , <i>Porichthys</i>	CCXXXV, 811	2319
<i>pottsii</i> , <i>Etheostoma</i>	CLXXXIII, 459	1082
<i>predidens</i> , <i>Poecilia</i>	CXIV, 303	697
<i>preciosus</i> , <i>Hypomesus</i>	LXXXVI, 230	525
<i>Priacanthus arenatus</i>	CXCV, 511	1237
<i>princeps</i> , <i>Cottus</i>	CCXCVI, 713	1962
<i>Prionae glauca</i>	IV, V, 16, 16a	33
<i>Prionistius macellus</i>	CCLXXXIX, 700	1028
<i>Prionodes bulleri</i>	CXCI, 503	1213
<i>Prionotus alatus</i>	CCCXIX, 770	2159
<i>carolinus</i>	CCCXVIII, 768	2156
<i>evolans</i>	CCCXX, 772	2168
<i>scintilus</i>	CCCXIX, 769	2157
<i>stejnosi</i>	CCCXN, 771	2166
<i>Pristis pectinatus</i>	VIII, 27	60
<i>probatocephalus</i> , <i>Archosargus</i>	CCCVI, 554	1361
<i>proboscidea</i> , <i>Limanda</i>	CCLXXVIII, 931	2645
<i>proboscideus</i> , <i>Chacromugil</i>	CXXVII, 346	816
<i>procera</i> , <i>Venefica</i>	LVIII, 152	365
<i>productus</i> , <i>Merluccius</i>	CCCLIX, 884	2531
<i>profundorum</i> , <i>Leptodipterus</i>	CCCLXII, 867	2484
<i>Seydlorhinus</i>	III, 11	22
<i>Zesticelus</i>	CCC, 726	1960
<i>Promicrops itaiara</i>	CLXXXV, 487, 487a, 487b	1162
<i>Pronotogrammus multifasciatus</i>	CXII, 506	1226
<i>proximus</i> , <i>Calamus</i>	CCXIII, 547	1350
<i>proximus</i> , <i>Microgadus</i>	CCCLX, 880	2539
<i>Psettichthys melanostictus</i>	CCCLXXIII, 921	2618
<i>pseudoharengus</i> , <i>Pomolobus</i>	LXXI, 189	425
<i>Pseudopleronectes americanus</i>	CCCLXXIX, 933	2647
<i>Pseudopriacanthus altus</i>	CXCV, 512	1239

Genera and species.	Plate and figure.	Text page.
<i>Pseudoscars guacamai</i>	CCXLVI, 617	1637
<i>Pseudotriakis microdon</i>	IV, 14	27
<i>psittacus</i> , <i>Xyrichtys</i>	CCXLII, 607	1618
<i>Psychrolutes paradoxus</i>	CCCVI, 730	2026
<i>Psychromaster inscumbia</i>	CLXXVI, 467	1100
<i>Ptilichthys goodei</i>	CCXLVII, 848	2452
<i>Ptychocheilus oregonensis</i>	XLI, 101	224
<i>pugetensis</i> , <i>Chitonotus</i>	CCLXXXIII, 686	1890
<i>pulcher</i> , <i>Pimelomelopon</i>	CXXXVIII, 598	1585
<i>pulvereus</i> , <i>Fundulus</i>	CVI, 285	652
<i>punctatus</i> , <i>Bodianus fulvus</i>	CLXXXII, 481	1146
<i>Ictalurus</i>	XXV, 58	134
<i>Stichaeus</i>	CXXXLV, 841	2439
<i>Xeasurus</i>	CCLVII, 630	1694
<i>punctulatus</i> , <i>Cottus</i>	CXCII, 708	1948
<i>putnami</i> , <i>Liosetta</i>	CCCLXXX, 936	2650
<i>pygmaea</i> , <i>Umbrina</i>	CXIX, 268	624
<i>quadricornis</i> , <i>Oncorhynchus</i>	CXX, 322	752
<i>quadrifilis</i> , <i>Bathypterois</i>	CCC, 728	2001
<i>quadriradiatus</i> , <i>Pleuronectes</i>	LXXXIX, 298	545
<i>quadrocellata</i> , <i>Aneylopetta</i>	CCLXXXIX, 934	2648
<i>Quassiremus evionthas</i>	CCCLXXV, 925	2634
<i>quiescens</i> , <i>Copelandellus</i>	LXIV, 107	380
<i>Rabiania inermis</i>	CLXXVI, 468	1100
<i>Rachycentron canadum</i>	CXCIX, 519	1274
<i>radiatus</i> , <i>Iridio</i>	CXLVIII, 401	948
<i>Ralidinus asprellus</i>	CXXXXIX, 600	1590
<i>boleoides</i>	CCLXXXVII, 695	1920
<i>Raja ackleyi</i>	CCLXXXVII, 694	1919
<i>erinacea</i>	X, 31	70
<i>ocellata</i>	IX, 29	68
<i>stellulata</i>	X, 30	68
<i>raleighana</i> , <i>Harriota</i>	XI, 32	75
<i>Ranzania truncata</i>	XIX, 42	96
<i>raphidoma</i> , <i>Tylosurus</i>	CCLXVIII, 651	1735
<i>Rastrinus scutiger</i>	CXVI, 308	715
<i>rathbuni</i> , <i>Fundulus</i>	CCLXXXVI, 691	1949
<i>Mulloidies</i>	CV, 280	649
<i>rectifrenum</i> , <i>Eupomacentrus</i>	CXXXI, 361	857
<i>reddingi</i> , <i>Orthopristis</i>	CCXXXIII, 589	1757
<i>Regalecus glesne</i>	CCIX, 540	1336
<i>regalis</i> , <i>Cynoscion</i>	CCLXX, 910	2506
<i>Scomberomorus</i>	CCXX, 562	1407
<i>regius</i> , <i>Urophycis</i>	CXXXV, 369	875
<i>Remora brachiptera</i>	CCCLXIV, 898	2553
<i>reticulatus</i> , <i>Lycodes</i>	CCXXX, 797, 797a	2272
<i>retifera</i> , <i>Muricena</i>	CCXLIX, 854	2465
<i>retroscella</i> , <i>Apogon</i>	LXVI, 173	401
<i>Rhaeoichthys toxotes</i>	CLXXVII, 471	1108
<i>Rhamphocottus richardsoni</i>	CCXXXI, 584	1507
<i>Rhinichthys dracis</i>	CCCVII, 742, 742a, 742b, 742c	2030
<i>Rhinobatos lentiginosus</i>	LH, 132	306
<i>Rhinoptera steindachneri</i>	VIII, IX, 28, 28a, 28b	62
<i>rhodorus</i> , <i>Ascelichthys</i>	XVII, 38, 38a	91
<i>Rhombochirus osteochir</i>	CCCV, 739	2025
<i>rhomboidea</i> , <i>Lagodon</i>	CCCCXXX, 798	2273
<i>Rhombopterus aurorubens</i>	CCXV, 552	1358
<i>richardsoni</i> , <i>Astronesthes</i>	CC, 521	1277
<i>Rhamphocottus</i>	CXIV, 252	587
<i>rimator</i> , <i>Bathystoma</i>	CCCVII, 742, 742a, 742b, 742c	2030
<i>Rimicola muscarum</i>	CCVI, 534	1368
<i>rimosus</i> , <i>Etropus</i>	CCXXXV, 813	2338
<i>Rissola marginata</i>	CCCLXXXV, 945	2688
<i>rivulatus</i> , <i>Cirrhites</i>	CCCLIII, 868	2489
<i>roberti</i> , <i>Hyporhamphus</i>	CCXXXVII, 576	1491
<i>Roecus chrysops</i>	CXVII, 312	721
<i>lineatus</i>	CLXXX, 477	1132
<i>Roucadour stearnsi</i>	CLXXX, 478	1132
<i>Rondeletia bicolor</i>	CCXXIII, 568	1457
<i>rondeletii</i> , <i>Exonantes</i>	XC, 240	548
<i>Ronquilus jordani</i>	CXVII, 317	733
	CCXXXII, 803	2289

Genera and species.	Plate and figure.	Text page.
rosaceus, Zalembius.....	CCXXXIX, 581	1500
ruficundus, Acpenser.....	XXI, 46	106
Hypsypops.....	CCXXXIV, 501	1564
rupestris, Xiphidion.....	CXCXLIV, 838	2426
rupestris, Ambloplites.....	CLVI, 419, 419a, 419b, 419c	990
Kuhlia.....	CLXIII, 432
Rupisartes atlanticus.....	CCCCXXXIX, 825	2397
Rupisartes, Moxostoma.....	XXXVII, 93	196
Rusearius meanyi.....	CCLXXXV, 690	1908
Rutilus bicolor.....	XLII, 107	244
Rypticus bistrispinus.....	CXCIV, 509	1233
sabina, Dasyatis.....	XIV, 36a	84
suburri, Chasmodes.....	CCCCXXXIX, 824	2392
Saccopharynx ampullaceus.....	LXVI, 175	406
Sagenichthys aencyclodon.....	CCXXI, 564	1416
saita, Borogadus.....	CCCLIX, 885	2533
Salmo clarkii honvieri.....	LXXX, 212	496
hemshawi.....	LXXXIX, 208	2819
maedonaldi.....	LXXXI, 214	497
pleuritiens.....	LXXX, 211	2819
spilurus.....	LXXXIX, 210	495
stomias.....	LXXX, 213	2819
virginalis.....	LXXXIX, 209	495
gardineri.....	LXXXI, 215	497
iridens.....	LXXXII, 216	500
salmoides, Micropterus.....	CLXII, 431	1012
saltatrix, Pomatomus.....	CXLVIII, 400	946
saludanus, Notropis ludovicianus.....	XLVII, 120	270
Salvelinus alpinus aureolus.....	LXXXIII, 220	511
fontinalis.....	LXXXII, 218	506
malma.....	LXXXII, 219	507
oquassa.....	LXXXIII, 221	514
sapidissima, Alosa.....	LXXII, 191	427
Sardinella himmalaris*.....	LXXIII, 193	431
stolifera.....	LXXIII, 194	431
Sarrtor frenatus.....	CCCX, 751	2073
sartorius, Neoclinus.....	CCCCXXXVI, 816	2355
saurus, Elops.....	LXVII, 178	410
Oligoplites.....	CXXXVIII, 378	898
Scomberesox.....	CXVII, 314	725
saxatilis, Abulidesdus.....	CCXXXIV, 590	1561
sayanensis, Aphredoderus.....	CXXII, 331	786
Scaphirhynchus platorhynchus.....	XXI, 48	107
Scarus ceruleus.....	CCXLIV, 613	1652
cuzamiae.....	CCXLIV, 612	1648
emblematicus.....	CCXLV, 614	1654
strongylocephalus.....	CCLXV, 615a, 615a
septicus, Triglops.....	CCLXXXVIII, 698	1925
Schedophilus medusophagus.....	CLI, 407	970
Schilbeoides exilis.....	XXVIII, 65	147
turiosus.....	XXIX, 69, 69a, 69b	149
gilberti.....	XXVIII, 67a, 67b	148
insignis.....	XXVIII, 66	147
minor.....	XXIX, 68	148
nocturnus.....	XXVII, 64	146
schmidti, Hoplunnis.....	LVIII, 151	361
schepeti, Chilomycterus.....	CCLXVI, 648	1748
schepeti, Aluterus.....	CCLX, 635	1718
Sciænops ocellatus.....	CCXXXI, 567	1453
scierus, Hadropterus.....	CLXVII, 441	1037
seitudus, Prionotus.....	CCCXIX, 768	2157
seurus, Hamulon.....	CCV, 531	1303
Scouber collis.....	CXXXIII, 304	866
scombrus.....	CXXXIII, 303	865
Scomberomorus maculatus.....	CXXXIV, 308	874
regalis.....	CXXXV, 309	875

* Plate by error labeled *Sardinella sardina*.

Genera and species.	Plate and figure.	Text page.
<i>Scomberesox saurus</i>	CXVII, 314	725
<i>scombrus</i> , <i>Scomber</i>	CXXXIII, 363	865
<i>Scorpaena brasiliensis</i>	CCLXXXVII, 669	1842
<i>cristulata</i>	CCLXXVI, 668	1841
<i>grandicornis</i>	CCLXXVIII, 671	1850
<i>mystes</i>	CCLXXVII, 670	1849
<i>Scorpenichthys marmoratus</i>	CCLXXXIII, 685	1889
<i>scorpius</i> , <i>Myoxocephalus</i>	CCXCV, 716	1974
<i>scripta</i> , <i>Aluterus</i>	CCLX, 686	1719
<i>scutellaris</i> , <i>Bascanichthys</i>	LXIII, 165	378
<i>sentiger</i> , <i>Rastrinus</i>	CCLXXXVI, 691	1909
<i>Seydlorhinus profundorum</i>	III, 11	21
<i>Scyatlina cerdale</i>	CCCXLVIII, 810, 849a, 849b	2454
<i>Sebastes marinus</i>	CCLXVII, 652	1760
<i>Sebastodes alutus</i>	CCLXVII, 660	1790
<i>brevispinis</i>	CCLXXI, 657	1787
<i>caurinus</i>	CCLXXIV, 663	1820
<i>ellatus</i>	CCLXX, 655	1783
<i>eigenmanni</i>	CCLXXI, 658	1789
<i>gilberti</i>	CCLXXV, 665	1823
<i>hopkinsi</i>	CCLXXII, 659	1789
<i>maliger</i>	CCLXXIV, 664	1822
<i>melanops</i>	CCLXIX, 654	1782
<i>miniatu</i>	CCLXXII, 662	1794
<i>mystinus</i>	CCLXX, 656	1784
<i>nigrocinetus</i>	CCLXXVI, 667	1827
<i>pinniger</i>	CCLXXIII, 661	1793
<i>sericeps</i>	CCLXXV, 666	1827
<i>Sebastolobus altivelis</i>	CCLXIX, 653	1763
<i>sectatrix</i> , <i>Kyphosus</i>	CCXIX, 559	1387
<i>Selene vomer</i>	CXLIV, CXLV, 393, 393a	936
<i>seleneops</i> , <i>Hiodon</i>	LXVIII, 181	414
<i>seminalis</i> , <i>Fundulus</i>	CIV, 277	647
<i>Semotilus atromaculatus</i>	XL, 100	322
<i>Seriola dorsalis</i>	CXXXIX, 380	902
<i>lalandi</i>	CXL, 382	903
<i>zonata</i>	CXXXIX, 381	902
<i>serpentinus</i> , <i>Derlichthys</i>	LV, 142	343
<i>sericeps</i> , <i>Sebastodes</i>	CCLXXV, 666	1827
<i>Serrivomer beani</i>	LVIII, 153	367
<i>Setarches parvatus</i>	CCLXXIX, 673	1860
<i>setiger</i> , <i>Dasyctettus</i>	CCC, 727	1991
<i>setipinnis</i> , <i>Vomer</i>	CXLIV, 392	934
<i>setosus</i> , <i>Ovoides</i>	CCLV, 646	1739
<i>shufeldti</i> , <i>Maneblas</i>	CCCXC, 955	2730
<i>shumardi</i> , <i>Cottogaster</i>	CLXVIII, 444	1046
<i>Signalosa atchafalaya</i>	LXIX, 184	2809
<i>signatus</i> , <i>Bathymaster</i>	CCXXXII, 802	2288
<i>signifer</i> , <i>Thymallus</i>	LXXXIII, 229	515
<i>sigillatus</i> , <i>Gilbertidius</i>	CCCVI, 741	2028
<i>sileneus</i> , <i>Zaprora</i>	CLU, 409	2550
<i>silus</i> , <i>Argentina</i>	LXXXVII, 232	526
<i>Simenichelys parasitica</i>	LVI, 144	349
<i>simotera</i> , <i>Ulocentra</i>	CLXX, 448	1053
<i>simula</i> , <i>Chalinura</i>	CCCLXVIII, 910	2578
<i>simiole</i> , <i>Umbrina</i>	CCXXXIV, 571	1468
<i>Siphonotoma starksii</i>	CXX, 325	771
<i>siniawii</i> , <i>Leuciscus</i>	XLIID, 706	2797
<i>stoma</i> , <i>Chauliodus</i>	XCHI, 250	585
<i>smaragdus</i> , <i>Eretelis</i>	CCXXXV, 785	2204
<i>sordidus</i> , <i>Citharichthys</i>	CCCLXXXIV, 943	2679
<i>Verilus</i>	CCII, 525	1284
<i>spadicea</i> , <i>Lampetra</i>	II, 6	13
<i>Sparisoma aurofrenatum</i>	CCXLIII, 610	1634
<i>cretense</i>	CCXLIV, 611	1632
<i>hoplomyctax</i>	CLIV, 416	987
<i>sparoides</i> , <i>Pomoxis</i>	XX, 43, 43a	101
<i>spatula</i> , <i>Polydectes</i>	XCI, 244	561
<i>speculifera</i> , <i>Lampadena</i>	CXY, 307	706
<i>speleus</i> , <i>Amblyopsis</i>	CCLXIV, 643	1732
<i>spengleri</i> , <i>Spherooides</i>	CCLXIV, 644	1733
<i>Spherooides maculatus</i>	CCLXIV, 642	1732
<i>nephelus</i>		

Genera and species.	Plate and figure.	Text page.
<i>Sphoeroides</i> — <i>steindleri</i>	CCLXIV, 643	1732
..... <i>testudineus</i>	CCLXV, 645, 645a	1734
<i>Sphyraena</i> — <i>barracuda</i>	CXXXVII, 349	{ 823 2841
<i>Sphyraena</i> — <i>tourou</i>	V, 19	41
<i>Sphyraena</i> — <i>sphyraena</i>	CLXVIII, 443	{ 1043 496
<i>Sphyraena</i> — <i>salmo</i>	LXXIX, 219	2819
<i>spilurus</i> , <i>clarkii</i>	CCLXXXVI, 602	1914
<i>spinniger</i> , <i>Icelus</i>	CXXXI, 357	844
<i>splendens</i> , <i>Beryx</i>	VII, 24, 24a	54
<i>Squatinus</i> — <i>acanthias</i>	CCLXXIV, 923	2631
<i>squamiferus</i> , <i>Parachaetichthys</i>	CXN, 325	771
<i>starksii</i> , <i>Siphonotoma</i>	CCCXL, 827	2407
<i>stethmonotus</i> , <i>hemophilus</i>	CCCXXI, 771	2166
<i>stearnsi</i> , <i>Prionotus</i>	CXXXIII, 568	1457
..... <i>Roncador</i>	XVII, 38, 38a	91
<i>steindachneri</i> , <i>Rhinoptera</i>	CCLXVIII, 909	2568
<i>steindachneria</i> , <i>argentea</i>	CCCIX, 748, 748a	2087
<i>Stellifer</i> — <i>vulgaris</i>	CCLXXXVI, 937, 937a	2632
<i>stellatus</i> , <i>Platichthys</i>	CCLXXXI, 678	1871
<i>stelleri</i> , <i>Hexagrammos</i>	CCXC VIII, 722a, 722b	1981
..... <i>Myoxocephalus</i>	CV, 279	648
<i>stellifer</i> , <i>Fundulus</i>	XI, 32	75
<i>stellifera</i> , <i>Raja</i>	LXXVII, 204	474
<i>Steindodus</i> — <i>mackenzii</i>	CXXII, 545	1346
<i>stenotomus</i> — <i>aculeatus</i>	CCXI, 544	1346
..... <i>chrysops</i>	CXXIX, 353	836
<i>Stephanoberryx</i> — <i>mona</i>	CCCX, 750, 750a	2071
<i>sterletus</i> , <i>Avermanni</i>	CCLXXXVIII, CCLXXXIX, 699, 699a	1927
<i>Sternias</i> — <i>xenostethus</i>	CCCXLV, 841	2439
<i>Stichaeus</i> — <i>punctatus</i>	CXXXVI, 787	2234
<i>stigmatus</i> , <i>Gobius</i>	CXLVII, 399	2848
<i>stigbe</i> , <i>Zalocys</i>	LXXXVII, 233	527
<i>stilbius</i> , <i>Leucoglossus</i>	CXII, 332	790
<i>stipes</i> , <i>Atherinum</i>	CLXIV, 434	1022
<i>stizostedion</i> — <i>canadense</i>	CLXIV, 433, 433a	1021
..... <i>vitreum</i>	LXXIV, 196	442
<i>stolephorus</i> — <i>pertheatus</i>	LXXIII, 194	431
<i>stolifera</i> , <i>Sardinella</i>	CCLXXXI, 917	2609
<i>stomias</i> , <i>Atheresthes</i>	XXXV, 86	2794
..... <i>Chasmistes</i>	LXXX, 213	{ 497 2819
..... <i>Salmo</i> — <i>clarkii</i>	XCIV, 253	588
<i>Stomias</i> — <i>ferox</i>	CXC, 500	1199
<i>striaatus</i> , <i>Centropristes</i>	CCLXXXIII, 483	1157
<i>Epinephelus</i>	CCXLV, 615, 615a	—
<i>strongylocephalus</i> , <i>Scarus</i>	XX, 45	105
<i>sturio oxyrinchus</i> , <i>Actipenser</i>	CCCXLV, 841	2440
<i>subbifurcata</i> , <i>Ulvaria</i>	CXXX, 355	841
<i>suborbitalis</i> , <i>Plectromus</i>	CXV, 306	704
<i>subterraneus</i> , <i>Typhlechthys</i>	XXXVI, 89	185
<i>suetta</i> , <i>Erimyzon</i>	CCLXXXI, 679	1872
<i>superciliosus</i> , <i>Hexagrammos</i>	CCVIII, 537	1318
<i>surinamensis</i> , <i>Anisotremus</i>	CXCIV, 510	1235
..... <i>Lobotes</i>	XLIX, 125	290
<i>waini</i> , <i>Notropis</i>	CCCLXXXIX, 941	2671
<i>sciacium</i> , <i>papillosum</i>	CLIX, 424	908
<i>symmetrius</i> , <i>Apionotis</i>	CCCLXXXVII, 949	2706
<i>synaphirus</i> — <i>marginatus</i>	CCCLXXXVIII, 950	2710
..... <i>plagiusa</i>	CCCLXXXVIII, 951	2711
..... <i>williamsi</i>	CXCVII, 518	1270
<i>synagris</i> , <i>Neomanus</i>	LVI, 146	351
<i>Synaphobranchus</i> — <i>pinnatus</i>	LXXXVII, 236	598
<i>Synodus</i> — <i>fetens</i>	CXXXVI, 372	886
<i>tentatus</i> , <i>Erythrometopon</i>	XXXII, 78	177
<i>tahitiensis</i> , <i>Catostomus</i>	LXXIV, 198	455
<i>Talismania</i> — <i>antillarum</i>	XCI, 248	575
<i>Tarletonbeania</i> — <i>tenua</i>	LXVII, 177	409
<i>Tarpon</i> — <i>atlanticus</i>	CCXXXVII, 596	1578
<i>Tautogolabrus</i> — <i>adspersus</i>	CCXXXVII, 595	1577
<i>telescopus</i> , <i>Notropis</i>	L, 126	292
<i>temna</i> , <i>Tarletonbeania</i>	XCI, 248	575

Genera and species.	Plate and figure.	Text page.
<i>tenulus</i> , <i>Uranidea</i>	CCXCIV, 714	1966
<i>tenuis</i> , <i>Urophycis</i>	CCCLXV, 901	2555
<i>tergisus</i> , <i>Hiodon</i>	LXVII, 180	413
<i>testudinens</i> , <i>Spherooides</i>	CCLXV, 645, 645a	1734
<i>Tetragonurus cuvieri</i>	CLII, 411	976
<i>Tetronarce californica</i>	XII, 34, 34a	77
<i>occidentalis</i>	XI, 33	77
<i>Tenthitis bahamensis</i>	CCLVI, CCLVII, 629a	1693
<i>crestonis</i>	CCLVI, 628	1692
<i>thaleichthys</i> , <i>Osmerus</i>	LXXXV, 227	592
<i>Thaleichthys pacificus</i>	LXXXV, 226	591
<i>thazard</i> , <i>Auxis</i>	CXXXIII, 363	867
<i>Theragra chalcogramma</i>	CCCXLX, 887	2335
<i>thoburni</i> , <i>Alcidia</i>	CCLXXXIII, 684	1887
<i>Thymallus tricolor</i>	LXXXIV, 223	518
<i>montanus</i>	LXXXIV, 224, 224a	519
<i>signifer</i>	LXXXIII, 222	515
<i>Thyrina overmanni</i>	CXXV, 340	994
<i>tiburo</i> , <i>Sphyraena</i>	V, 19	44
<i>tigrinus</i> , <i>Myrichthys</i>	LXI, 161	376
<i>tomeod</i> , <i>Microgadus</i>	CCLIX, 890	2540
<i>toxotes</i> , <i>Rhacocelus</i>	CCXXXI, 584	1567
<i>Trachinocephalus myops</i>	LXXXVIII, 233	533
<i>Trachinotus carolinus</i>	CXLVII, 398	944
<i>elveri</i>	CXLVII, 397	942
<i>falcatus</i>	CXLVI, 396	941
<i>glauces</i>	CXLVI, 395	940
<i>Trachurops crumenophthalmus</i>	CXLIX, 385	911
<i>Trachurus trachurus</i>	CXLIX, 384	910
<i>trachurus</i> , <i>Trachurus</i>	CXLIX, 384	910
<i>transmontana</i> , <i>Columbia</i>	CXXII, 330	784
<i>transmontanus</i> , <i>Acipenser</i>	XX, 44	104
<i>traski</i> , <i>Hystericarpus</i>	CCXXVII, 577	1496
<i>triangularis</i> , <i>Poromotus</i>	CL, 405	967
<i>Trichium lepturus</i>	CXXXVII, 373	889
<i>Trichodus trichodon</i>	CCCXXXIII, 806	2265
<i>trichodon</i> , <i>Trichodon</i>	CCCCXXXIII, 806	2265
<i>Trichopsetta ventralis</i>	CCLXXXIII, 940	2669
<i>tricolor</i> , <i>Holacanthus</i>	CCLIII, 625	1634
<i>montanus</i> , <i>Thymallus</i>	LXXXIV, 224, 224a	519
<i>Thymallus</i>	LXXXIV, 223	518
<i>tricornis</i> , <i>Lactophrys</i>	CCLXI, 638	1731
<i>tridentatus</i> , <i>Entosphenus</i>	1, 4	32
<i>Triglops beani</i>	CCLXXXVIII, 697	1924
<i>pingoli</i>	CCLXXXVIII, 696	1923
<i>scepticus</i>	CCLXXXVIII, 698	1923
<i>trigonus</i> , <i>Lactophrys</i>	CCLXIII, 640, 640a	1723
<i>triqueter</i> , <i>Lactophrys</i>	CCLXII, 637	1722
<i>tristichthus</i> , <i>Lepisosteus</i>	XXI, 50	111
<i>truncata</i> , <i>Ranania</i>	CCLXVIII, 651	1735
<i>tachawytscha</i> , <i>Oneirolynchus</i>	LXXXVII, 206	479
<i>tsiltcoosenensis</i> , <i>Catostomus</i>	XXXII, 80	2793
<i>tunneri</i> , <i>Lycodespis</i>	CCCL, 858	2469
<i>tuscumbia</i> , <i>Psychromaster</i>	CLXXVI, 467	1100
<i>Tylosurus acus</i>	CXVI, 309	716
<i>raphidoma</i>	CXVI, 308	715
<i>Typhlichthys subterraneus</i>	CXV, 306	704
<i>Typhlogobius californiensis</i>	CCCCXIX, 795	2262
<i>tyrannus</i> , <i>Brevoortia</i>	LXXXII, 195	433
<i>Ulocentra gibberti</i>	CLXIX, 446	1049
<i>meadue</i>	CLXIX, 447	1052
<i>sinotera</i>	CLXX, 448	1053
<i>ulvae</i> , <i>Xiphistes</i>	CCCCXLI, 836	2423
<i>Ulvaria subfibrata</i>	CCCLXV, 842	2410
<i>umatilla</i> , <i>Agosia</i>	LI, 134	313
<i>Umbräa pygmaea</i>	XCIX, 268	624
<i>Umbrina sinaloae</i>	CCXXXIV, 571	1463
<i>undecimialis</i> , <i>Centropomus</i>	CLXXXIX, 476	1118
<i>undulatus</i> , <i>Micropogon</i>	CCXXXIV, 570	1461

Genera and species.	Plate and figure.	Text page.
<i>maculatus nigricans</i> , <i>Hypoplectrus</i> .	CXXXIX, 496	1193
<i>maculatus</i> , <i>Hyporhamphus</i> .	CXVI, 311	720
<i>maculatus</i> , <i>Archosargus</i> .	CCXVI, 553	1350
<i>maculatus</i> .	CXXXII, 362	858
<i>maculatus</i> .	CCCLXII, 894	2545
<i>malabarensis</i> .	CCXCIV, 714	1966
<i>malabaricus</i> .	LI, 130	304
<i>malabaricus</i> .	CCCLXVI, 903	2550
<i>malabaricus</i> .	CCCLXV, 902	2555
<i>malabaricus</i> .	CCCLXIV, 890	2553
<i>malabaricus</i> .	CCCLXV, 900	2554
<i>malabaricus</i> .	CCCLXIV, 899	2553
<i>malabaricus</i> .	CCCLXV, 901	2555
<i>malabaricus</i> .	III, 12	25
<i>malabaricus</i> .	CXXIV, 336	704
<i>malabaricus</i> .	CXI, 295, 295a	600
<i>malabaricus</i> .	CXI, CXII, 296, 296a	671
<i>malabaricus</i> .	LXI, 160	375
<i>malabaricus</i> .	LVIII, 152	305
<i>malabaricus</i> .	CLXXXVII, 491	1172
<i>malabaricus</i> .	CCXLII, 605	1615
<i>malabaricus</i> .	CCCLV, 872	2503
<i>malabaricus</i> .	CCCLXXXIII, 940	2669
<i>malabaricus</i> .	CCXL, 803	1602
<i>malabaricus</i> .	CCHI, 525	1284
<i>malabaricus</i> .	LX, 159, 150a	375
<i>malabaricus</i> .	CCCLI, 850	2479
<i>malabaricus</i> .	CCXCIII, 720, 720a	1979
<i>malabaricus</i> .	CCCXII, 958-958b	2727
<i>malabaricus</i> .	CCCIX, 747, 747a	2063
<i>malabaricus</i> .	LXXXV, 225	520
<i>malabaricus</i> .	CCXIV, 759	2101
<i>malabaricus</i> .	CCCLXII, 893, 893a	2544
<i>malabaricus</i> .	CXXXV, 370	878
<i>malabaricus</i> .	CCCLIX, 886	2524
<i>malabaricus</i> .	CLXXV, 465	1093
<i>malabaricus</i> .	LXXXIX, 209	405
<i>malabaricus</i> .	CCIX, 539	1322
<i>malabaricus</i> .	CXCVI, 514	1246
<i>malabaricus</i> .	CCCLI, 864-864c	2479
<i>malabaricus</i> .	CLXIV, 433, 433a	1021
<i>malabaricus</i> .	LXVI, 174	404
<i>malabaricus</i> .	CXCH, 505	1223
<i>malabaricus</i> .	CCXXXIII, 778	2183
<i>malabaricus</i> .	CXVIII, 318	734
<i>malabaricus</i> .	CXLIV, 393, 393a	936
<i>malabaricus</i> .	CXLIV, 392	934
<i>malabaricus</i> .	LXVIII, 179	411
<i>malabaricus</i> .	VI, 20	45
<i>malabaricus</i> .	CCCIX, 748, 748a	2067
<i>malabaricus</i> .	LH, 137	319
<i>malabaricus</i> .	XLV, 113	2799
<i>malabaricus</i> .	XLVIII, 121	278
<i>malabaricus</i> .	CCCLXXXVIII, 951	2711
<i>malabaricus</i> .	CLI, 408	973
<i>malabaricus</i> .	CCXXIII, 569	1458
<i>malabaricus</i> .	CCHI, 527	1287
<i>malabaricus</i> .	CCCXI, 754, 754a	2084
<i>malabaricus</i> .	CCHI, 528	1285
<i>malabaricus</i> .	CCXLIII, 609, 609a	1626
<i>malabaricus</i> .	CCXLVI, 618	1660
<i>malabaricus</i> .	CCLXXXVIII, 699, 699a	1927
<i>malabaricus</i> .	CCLVII, 630	1694
<i>malabaricus</i> .	CCCXLIV, 838	2426
<i>malabaricus</i> .	CCCXLIV, 837	2424
<i>malabaricus</i> .	CCCXLIII, 830	2423
<i>malabaricus</i> .	CXV, 304	701
<i>malabaricus</i> .	XXXV, 88	184
<i>malabaricus</i> .	CCXLII, 607	1618
<i>malabaricus</i> .	CCXVIII, 556	1372
<i>malabaricus</i> .	CCCX, 752, 752a	2076

Genera and species.	Plate and figure.	Text page.
<i>Varrella blackfordi</i>	XCHI, 249	584
<i>y-gracum, Astroscopus</i>	CCCXXXIV, 808	2367
<i>Zalembius rosaceus</i>	CXXXIX, 581	1500
<i>Zalocys stilbe</i>	CXLVIII, 399	2848
<i>zanculus, Dermatolepis</i>	CLXXXVI, 489	2834
<i>Zanclus cornutus</i>	CCLV, 627	1687
<i>Zaprora silenus</i>	CLI, 409	2850
<i>zebrinus, Fundulus</i>	CIII, 276	646
<i>Zesticelus profundorum</i>	CCC, 726	1990
<i>Zoarces anguillaris</i>	CCXLVIII, 850	2457
<i>zorichthys, Lycoodes</i>	CCCXLIX, 853	2464
<i>zonata, Seriola</i>	CXXXIX, 381	902
<i>zonipactus, Pomaanthus</i>	CCLI, 624	1681
<i>zonope, Jordania</i>	CCLXXXII, 683	1884
<i>zosteræ, Hippocampus</i>	CXXI, 328	778
<i>zosterura, Evermannia</i>	CCCXXXI, 794	2256
<i>zyopterus, Galeorhinus</i>	IV, 15	32

NOTE.—
illustration
by the us
the photo
same prop

1. Branchiostomæ
Drawing
Stat.
2. Polycarpa
Drawing
3. Petromyzon
Drawing
Wood
4. Entosphenus
Drawing
lecter

5. Lampetra
Drawing
Lucie
6. Lampetra
Drawing
U.S.N.M.
7. Notorhynchus
Drawing
in Hu
8. Hexanchus
Drawing
in Cu

9. Gyropleuron
Drawing
California
10. Heterodontus
Drawing

EXPLANATION OF PLATES AND FIGURES.

NOTE.—The actual size of the specimen from which each illustration was drawn may, in most instances, be determined by the use of the inch mark beneath the engraving, which, in the photographic reduction of the drawing, is reduced in the same proportion as the drawing itself.

PLATE I.

Text page.

1. <i>Branchiostoma caribaeum</i> Snndeval	3
Drawing by A. H. Baldwin from a specimen in the United States National Museum.	
2. <i>Polypterus bichir</i> Geoff. St. L'Paire	
Drawing by Anna L. Brown.	
3. <i>Petromyzon marinus</i> Linnaeus	10
Drawing by H. L. Todd from No. 10654, U.S.N.M., collected at Woods Hole, Massachusetts, by V. N. Edwards.	
4. <i>Eutrophenus tridentatus</i> (Gairdner)	12
Drawing by A. H. Baldwin from No. 48204, U.S.N.M., collected in Kamchatka by Dr. L. Stejneger.	

PLATE II.

5. <i>Lampetra aurea</i> (Bean)	13
Drawing by H. L. Todd from No. 21524, U.S.N.M., collected by Lucien M. Turner in the Yukon River, Alaska.	
6. <i>Lampetra spadicea</i> Bean	13
Drawing by H. L. Todd from the type specimen, No. 38005, U.S.N.M., collected by Prof. A. Dugès at Guanajato, Mexico.	
7. <i>Notorhynchus maculatus</i> Ayres	17
Drawing by H. L. Todd from No. 27191, U.S.N.M., collected in Humboldt Bay, California, by Dr. Jordan.	
8. <i>Hexanchus griseus</i> (Gmelin)	19
Drawing by H. L. Todd from No. 37790, U.S.N.M., collected in Currinnek Inlet, North Carolina, by D. M. Etheridge.	

PLATE III.

9. <i>Gyropleurodon francisci</i> (Girard)	20
Drawing from No. 24977, U.S.N.M., collected at Wilmington, California.	
10. <i>Heterodontus philippii</i>	
Drawing by Anna L. Brown.	

	Text page
11. <i>Scylliorhinus profundorum</i> Goode and Bean.....	22
Drawing by M. M. Smith from the type, No. 35646, U.S.N.M., collected by the <i>Albatross</i> at 39° 9' N. and 72° 3' 15" W.	
12. <i>Catulus uter</i> Jordan and Gilbert.....	25
Drawing by A. H. Baldwin from No. 26866, U.S.N.M., col- lected at Santa Barbara, California.	

PLATE IV.

13. <i>Ginglymostoma cirratum</i> (Gmelin).....	26
Drawing by H. L. Todd from No. 2983, U.S.N.M., collected at Fort Jefferson, Florida, by Dr. Whitehurst.	
14. <i>Pseudotriakis microdon</i> Capello	27
Drawing by H. L. Todd from No. 32516, U.S.N.M., collected at Amagansett Island by J. B. Edwards.	
15. <i>Galeorhinus zyopterus</i> Jordan and Gilbert.....	32
16. <i>Prionace glauca</i> (Linnaeus).....	33
Drawing by H. L. Todd from No. 38001, U.S.N.M., collected by schooner <i>Grampus</i> .	

PLATE V.

16a. <i>Prionace glauca</i> (Linnaeus).....	33
Drawing by H. L. Todd from No. 38001, U.S.N.M., collected by the schooner <i>Grampus</i> .	
17. <i>Carcharhinus lamia</i> (Rafinesque).....	38
Drawing by H. L. Todd.	
18. <i>Hypoprion brevirostris</i> Poey.....	41
Drawing by W. S. Haines from No. 38497, U.S.N.M., collected by the <i>Albatross</i> at Watlings Island.	
19. <i>Sphyrna tiburo</i> (Linnaeus).....	44
Drawing by H. L. Todd from No. 26582, U.S.N.M., collected by Silas Stearns at Cedar Keys, Florida.	

PLATE VI.

20. <i>Alopias vulpes</i> (Gmelin)	45
Drawing by H. L. Todd from No. 25962, U.S.N.M., collected off Newport, Rhode Island, by Captain Rockliff.	
21. <i>Isurus dekayi</i> (Gill).....	48
Drawing by H. L. Todd from No. 37665, U.S.N.M., collected by Silas Stearns at Pensacola, Florida.	
22. <i>Lamna cornubica</i> (Gmelin).....	49
Drawing by H. L. Todd from No. 27368, U.S.N.M., collected at Santa Barbara, California, by Dr. Jordan.	

PLATE VII.

23. <i>Cetorhinus maximus</i> (Gunner).....	51
Drawing by S. F. Denton, copied from Annales du Muséum d'Histoire Naturelle, vol. 18, pl. 6.	

24, 25a. Sq.

25. Centros
 Drawi
 the A26. Centros
 Drawi
 Geor27. Pristis p
 Drawi
 Pens28. Rhinoba
 Drawi
 Pensa28a, 28b. Rhin
 v
 Dr29. Raja erin
 Drawing
 Woods30. Raja ocell
 Drawing
 Provini31. Raja ackle
 Drawing
 the Alb32. Raja stellu
 Drawing33. Tetronarce
 Drawing
 Hole, M34, 34a. Tetro
 Drawi
 lect

	Text page.
24, 24a. <i>Squalus acanthias</i> Linnaeus	54

Drawings by S. F. Denton from No. 22316, U.S.N.M., collected at Gloucester, Massachusetts, by the U. S. F. C.

PLATE VIII.

25. <i>Centroscymnus coelolepis</i> Bocage and Capello	55
Drawing by H. L. Todd from No. 26219, U.S.N.M., collected by the <i>Fish Hawk</i> off Newport, Rhode Island.	
26. <i>Centroscyllium fabricii</i> (Reinhardt)	56
Drawing by H. L. Todd from No. 22879, U.S.N.M., collected by George W. Scott at $44^{\circ} 23' N.$, $53^{\circ} 25' W.$, in 200 fathoms.	
27. <i>Pristis pectinatus</i> Linnaeus	60
Drawing by H. L. Todd from No. 30678, U.S.N.M., collected at Pensacola, Florida, by Silas Stearns.	
28. <i>Rhinobatus lentiginosus</i> Garman; lateral view	62
Drawing by H. L. Todd from No. 30175, U.S.N.M., collected at Pensacola, Florida, by Silas Stearns.	

PLATE IX.

28a, 28b. <i>Rhinobatus lentiginosus</i> Garman; dorsal and ventral views.....	62
Drawings by H. L. Todd from No. 30175, U.S.N.M., collected at Pensacola, Florida, by Silas Stearns.	
29. <i>Raja erinacea</i> Mitchell	68

Drawing by H. L. Todd from No. 10428, U.S.N.M., collected at Woods Hole, Massachusetts, by S. F. Baird.

PLATE X.

30. <i>Raja ocellata</i> Mitchell	68
Drawing by H. L. Todd from No. 24228, U.S.N.M., collected at Provincetown, Massachusetts, by U.S.F.C.	
31. <i>Raja ackleyi</i> Garman	70

Drawing by H. L. Todd from No. 2684, U.S.N.M., collected by the *Albatross* at $39^{\circ} 35' N.$, and $70^{\circ} 54' W.$.

PLATE XI.

32. <i>Raja stellulata</i> Jordan and Gilbert	75
Drawing by H. L. Todd.	
33. <i>Tetronarce occidentalis</i> (Storer)	77

Drawing by H. L. Todd from a specimen collected at Woods Hole, Massachusetts, in the summer of 1881 by the U.S.F.C.

PLATE XII.

34, 34a. <i>Tetronarce californica</i> (Ayres)	77
Drawings by H. L. Todd from No. 27212, U.S.N.M., collected by Dr. Jordan at Soquel, California.	

PLATE XIII.

	Text page.
35, 35a. <i>Narcine brasiliensis</i> (Ölfers).....	78
Drawings by H. L. Todd from No. 30178, U.S.N.M., collected by Silas Stearns at Pensacola, Florida.	

PLATE XIV.

36, 36a. <i>Dasyatis sabina</i> (Le Sueur).....	81
Drawings by H. L. Todd from No. 31043, U.S.N.M., collected at Galveston, Texas, by Dr. Jordan.	

PLATE XV.

37. <i>Aetobatus narinari</i> (Euphrasen).....	88
Drawing by H. L. Todd from No. 37196, U.S.N.M., collected by E. F. Denechaud near Cedar Keys, Florida.	

PLATE XVI.

37a. <i>Aetobatus narinari</i> (Euphrasen).....	88
Drawing by H. L. Todd from No. 37196, U.S.N.M., collected by E. F. Denechaud near Cedar Keys, Florida.	

PLATE XVII.

38, 38a. <i>Rhinoptera steindachneri</i> Evermann and Jenkins.....	91
Drawings by A. H. Baldwin from the type, No. 43235, U.S.N.M., collected at Guaymas, on the Gulf of California, by Drs. Evermann and Jenkins in 1887.	

PLATE XVIII.

39. <i>Manta birostris</i> (Walbaum)	92
Drawing by A. H. Baldwin.	

PLATE XIX.

40. <i>Chimæra affinis</i> Capello; female.....	95
Drawing by H. L. Todd from a specimen collected on the southeastern portion of Le Have Bank, in $42^{\circ} 40' N.$, $63^{\circ} 23' W.$.	
41. <i>Hydrolagus colliei</i> (Lay and Bennett).....	95
Drawing by H. L. Todd, from No. 26976, U.S.N.M., collected at Monterey, California, by Dr. Jordan.	
42. <i>Harriota raleighana</i> Goode and Bean.....	96
Drawing by A. H. Baldwin from the type, No. 35520, U.S.N.M., collected by the <i>Albatross</i> at station 2210, lat. $39^{\circ} 37' 45'' N.$, long. $71^{\circ} 18' 45'' W.$, at a depth of 991 fathoms.	

PLATE XX.

43, 43a. <i>Polyodon spathula</i> (Walbaum).....	101
Drawings by H. L. Todd from No. 12235, U.S.N.M., collected at Cincinnati, Ohio, by J. W. Milner.	

44. *Heterodontus* Dr.
45. *Acipenser* Dr.
46. *Acipenser* Dra.
47. *Acipenser* Dra.
48. *Scaphirhynchus* Dra.
49. *Lepisosteus* Dra.
50. *Lepisosteus* Dra.
51, 51a. *Alosa* Dra.
52. *Felichthys* Dra.
53. *Galeichthys* Dra.
54. *Galeichthys* Draw.
55. *Galeichthys* Draw.

	Text page.
44. Heterocercal tail of <i>Acipenser transmontanus</i> Richardson..	104
Drawing by Anna L. Brown.	
45. <i>Acipenser sturio oxyrhynchus</i> (Mitchill)	105
Drawing by H. L. Todd from No. 22495, U.S.N.M., collected by William Woltz in the Potomac River.	

PLATE XXI.

46. <i>Acipenser rubicundus</i> Le Sueur.....	106
Drawing by H. L. Todd from No. 10252, U.S.N.M., collected at Ecorse, Michigan, by J. W. Milner.	
47. <i>Acipenser brevirostrum</i> Le Sueur	106
Drawing by H. L. Todd from photograph of specimen taken at Woods Hole by the U.S.F.C.	
48. <i>Scaphirhynchus platorhynchus</i> (Rafinesque)	107
Drawing by H. L. Todd.	

PLATE XXII

49. <i>Lepisosteus platostomus</i> Rafinesque.....	110
Drawing by H. L. Todd from No. 3241, U.S.N.M., collected at Cleveland, Ohio, by Spencer F. Baird.	
50. <i>Lepisosteus tristoechus</i> (Bloch and Schneider).....	111
Drawing by A. H. Baldwin from No. 24794, U.S.N.M., col- lected by Professor Poey in Cuba.	
51, 51a. <i>Amia calva</i> Linnaeus, female.....	113
Drawings by S. F. Denton.	

PLATE XXIII.

52. <i>Felichthys felis</i> (Linnaeus)	118
Drawing by H. L. Todd from No. 10422, U.S.N.M., collected at Woods Hole, Massachusetts, by Spencer F. Baird.	
53. <i>Galeichthys milberti</i> (Cuvier and Valenciennes)	128
Drawing by H. L. Todd from a specimen collected at Pensa- cola, Florida, by Silas Stearns.	

PLATE XXIV.

54. <i>Galeichthys gilberti</i> Jordan and Williams.....	2773
Drawing by Anna L. Brown from the type, No. 1667, L. S. Jr. Univ. Mus., collected by the Hopkins expedition at Mazat- lan, Mexico.	
55. <i>Galeichthys azureus</i> Jordan and Williams.....	2775
Drawing by Anna L. Brown from the type, No. 1575, L. S. Jr. Univ. Mus., collected by the Hopkins expedition at Mazat- lan, Mexico.	

PLATE XXV.

	Text page.
56. <i>Ictalurus furcatus</i> (Le Sueur).....	134
Drawing by M. M. Smith from No. 838, U.S.N.M., type of <i>Pimelodus affinis</i> , collected at Brownsville, Texas, by Capt. Van Vliet.	
57. <i>Ictalurus anguilla</i> Evermann and Kendall.....	2788
Drawing by A. H. Baldwin from the type, No. 48788, U.S.N.M., collected by Dr. Evermann in the Atchafalaya River, Morgan City, Louisiana.	
58. <i>Ictalurus punctatus</i> (Rafinesque).....	131
Drawing by H. L. Todd from No. 27846, U.S.N.M., collected at Pekin, Illinois, by Dr. S. A. Forbes.	

PLATE XXVI.

59, 59a, 59b. <i>Ameiurus dugesii</i> Bean.....	138
Drawings by H. L. Todd from the type, No. 23123, U.S.N.M., collected by Prof. A. Dugès in Rio Turbio, Mexico.	
60. <i>Ameiurus melas</i> (Rafinesque)	141
Drawing by H. L. Todd from No. 1497, U.S.N.M., collected in Aux Plaines River, Illinois, by Robert Kennicott.	

PLATE XXVII.

61. <i>Ameiurus platycephalus</i> (Girard).....	142
Drawing by H. L. Todd from No. 1534, U.S.N.M., collected by Mrs. E. Daniel at Anderson, South Carolina.	
62. <i>Leptops olivaris</i> (Rafinesque)	143
Drawing by H. L. Todd from No. 27873, U.S.N.M., collected by S. A. Forbes in the Illinois River.	
63. <i>Noturus flavus</i> Rafinesque.....	144
Drawing by H. L. Todd from No. 1478, U.S.N.M., collected by Professor Baird in Lake Champlain, Westport, New York.	
64. <i>Schilbeodes nocturnus</i> (Jordan and Gilbert).....	146
Drawing by H. L. Todd from the type, No. 36461, U.S.N.M., collected by Jordan and Gilbert in the Sabine River near Benton, Arkansas.	

PLATE XXVIII.

65. <i>Schilbeodes exilis</i> (Nelson).....	147
Drawing by H. L. Todd from No. 36261, U.S.N.M., collected by Gilbert and Swain in the Ozark Fork of the Gasconade River, Marshfield, Missouri.	
66. <i>Schilbeodes insignis</i> (Richardson)	147
Drawing by H. L. Todd from No. 18016, U.S.N.M., collected by T. H. Bean in Consy Creek, Bainbridge, Pennsylvania.	
67, 67a, 67b. <i>Schilbeodes gilberti</i> (Jordan and Evermann).....	148
Drawings by S. F. Denton from the type, No. 39931, U.S.N.M., collected by D. S. Jordan and party in Roanoke River, Salem, Virginia.	

68. *Schilbe*
 Drawn by S.
 69, 69a, 69b
 Drawn coll. Nor
70. *Ictiobu*
 Drawn by S.
71. *Carpio*
 Drawn by Bea
72. *Cyclop*
 Drawn by J.
73. *Pantost*
 Drawn colle near
74. *Catosto*
 Drawn the C
75. *Catosto*
 Drawn by J.
76. *Catosto*
 Drawn colle Rive
77. *Catosto*
 Drawn by E.
78. *Catosto*
 Drawn Lake
79. *Catosto*
 Drawn Sacra

PLATE XXIX.

- | | Text page. |
|---|------------|
| 68. <i>Schilbeodes miurus</i> (Jordan) | 148 |
| Drawing by H. L. Todd from No. 20926, U.S.N.M.; collected
by J. W. Milner in Tar River, North Carolina. | |
| 69. 69a. 69b. <i>Schilbeodes furcatus</i> (Jordan and Meek). | 149 |
| Drawings by S. F. Denton from the type, No. 39932, U.S.N.M.,
collected by D. S. Jordan and party in Neuse River, Raleigh,
North Carolina. | |

PLATE XXX.

- | | |
|---|-----|
| 70. <i>Ictiobus cyprinella</i> (Cuvier and Valenciennes) | 162 |
| Drawing by H. L. Todd from No. 20774, U.S.N.M., collected
by S. A. Forbes at Normal, Illinois. | |
| 71. <i>Carpoides cyprinus</i> (Le Sueur) | 167 |
| Drawing by H. L. Todd from a specimen collected by Dr.
Bean at Havre de Grace, Maryland. | |
| 72. <i>Cycleptus elongatus</i> (Le Sueur)..... | 168 |
| Drawing by H. L. Todd from No. 10790, U.S.N.M., collected
by J. W. Milner in the Ohio River at Cincinnati, Ohio. | |

PLATE XXXI.

- | | |
|--|-----|
| 73. <i>Pantosteus jordani</i> Evermann | 171 |
| Drawing by A. H. Baldwin from the type, No. 43963, U.S.N.M.,
collected by Evermann and McCormick, in Whitewood Creek,
near Deadwood, South Dakota. | |
| 74. <i>Catostomus latipinnis</i> Baird and Girard..... | 174 |
| Drawing by H. L. Todd from No. 20078, U.S.N.M., collected in
the Gila River, Arizona. | |
| 75. <i>Catostomus griseus</i> (Girard)..... | 175 |
| Drawing by S. F. Denton from No. 21197, U.S.N.M., collected
by Jordan and Evermann. | |

PLATE XXXII.

- | | |
|--|-----|
| 76. <i>Catostomus pocatello</i> Gilbert and Evermann..... | 175 |
| Drawing by A. H. Baldwin from the type, No. 45385, U.S.N.M.,
collected by Evermann and Gilbert in Ross Fork of Snake
River, near Pocatello, Idaho. | |
| 77. <i>Catostomus catostomus</i> (Forster)..... | 176 |
| Drawing by H. L. Todd from No. 32888, U.S.N.M., collected
by E. W. Nelson, in the Yukon River, Alaska. | |
| 78. <i>Catostomus tahoensis</i> Gill and Jordan | 177 |
| Drawing by H. L. Todd from No. 5240, U.S.N.M., collected in
Lake Tahoe. | |

PLATE XXXIII.

- | | |
|---|-----|
| 79. <i>Catostomus occidentalis</i> Ayres | 178 |
| Drawing by A. H. Baldwin from a specimen collected in the
Sacramento River, by Dr. Jordan. | |

	Text page
80. <i>Catostomus tsiltcoosensis</i> Evermann and Meek	2793
Drawing by A. H. Baldwin from the type, No. 48479, U.S.N.M., collected by Dr. Meek in Tsilteos Lake, Oregon.	
81. Pharyngeal teeth of <i>Catostomus macrocheilus</i> Girard.....	178
82. Pharyngeal teeth of <i>Placopharynx duquesnii</i> (Le Sueur)....	198

PLATE XXXIV.

83. <i>Catostomus commersonii</i> (Lacépède)	178
Drawing by H. L. Todd from No. 10458, U.S.N.M., collected by J. W. Milner at Ecorse, Michigan.	
84. <i>Catostomus ardens</i> Jordan and Gilbert.....	179
Drawing by H. L. Todd from the type, No. 27363, U.S.N.M., collected by Dr. Jordan in Utah Lake, Utah.	
85. <i>Chasmistes liorus</i> Jordan	183
Drawing by S. F. Denton from No. 3042, U.S.N.M., collected by Dr. Jordan in Utah Lake.	

PLATE XXXV.

86. <i>Chasmistes stomias</i> Gilbert	2791
Drawing by A. H. Baldwin from the type, No. 48223, U.S.N.M., collected by Gilbert and Cramer in Upper Klamath Lake, Oregon.	
87. <i>Chasmistes copei</i> Evermann and Meek	2795
Drawing by A. H. Baldwin from the type, No. 48224, U.S.N.M., collected by Meek and Alexander in Upper Klamath Lake, Oregon.	
88. <i>Xyrauchen cypho</i> (Lockington).....	184
Drawing by S. F. Denton from a specimen collected by Dr. Jordan in Green River, Blake City, Utah.	

PLATE XXXVI.

89. <i>Erimyzon suetta</i> Lacépède	185
Drawing by H. L. Todd from No. 27867, U.S.N.M., collected by Dr. Forbes in Nipisink Lake, Illinois.	
90. <i>Minytrema melanops</i> (Rafinesque)	187
Drawing by H. L. Todd from No. 17800, U.S.N.M., collected by Dr. Bean in Round Lake, near Montgomery, Alabama.	
91. <i>Moxostoma congestum</i> (Baird and Girard).....	192
Drawing by H. L. Todd from No. 36510, U.S.N.M., collected in the Rio Lampasas, Belton, Texas.	

PLATE XXXVII.

92. <i>Moxostoma austrinum</i> Bean.....	192
Drawing by H. L. Todd from the type, No. 23121, U.S.N.M., collected at Piedad, in Morelia (Michoacan), Mexico, by Prof. A. Dugès.	

93. *Moxostoma*
 Drawing
 coll
 lina
- 94, 94a. *Lag*
 19
95. *Campos*
 Drawing
 by J
96. *Orthodo*
 Drawing
 by D
97. *Acroche*
 Drawing
 by C
98. *Algancea*
 Drawing
 by P
99. *Myloche*
 Drawing
100. *Semotilu*
 Drawing
 by Re
101. *Ptychoch*
 Drawing
102. *Leuciscu*
 Drawing
 Upper
 Otaki
103. *Leuciscu*
 Drawing
 stone
104. *Leuciscu*
 Drawing
 stone

Text page.

93. *Moxostoma rupisartes* Jordan and Jenkins 196
 Drawing by S. F. Denton from the type, No. 39927, U.S.N.M.,
 collected in the Catawba River, Morganton, North Caro-
 lina, by Jordan, Jenkins, and Meek.

PLATE XXXVIII.

- 94, 94a. *Lagochila lacera* Jordan and Brayton 199
 Drawing by H. L. Todd from No. 25238, U.S.N.M., collected
 by J. B. McElwin at Fairview, Tennessee.

PLATE XXXIX.

95. *Campostoma anomalum* (Rafinesque) 205
 Drawing by W. S. Haines from No. 36328, U.S.N.M., collected
 by Jordan and Gilbert in White River, near Eureka Springs,
 Arkansas.
 96. *Orthodon microlepidotus* (Ayer) 207
 Drawing by H. L. Todd from No. 27139, U.S.N.M., collected
 by Dr. Jordan in the Sacramento River, California.
 97. *Acrocheilus alutaceus* Agassiz and Pickering 208
 Drawing by H. L. Todd from No. 30298, U.S.N.M., collected
 by Capt. Charles Bendire in John Day River, Oregon.

PLATE XL.

98. *Algansea dugesi* Bean 211
 Drawing by A. H. Baldwin from No. 43764, U.S.N.M., collected
 by Professor Dugès in Mexico.
 99. *Mylocheilus caurinus* (Richardson) 219
 Drawing by A. H. Baldwin.
 100. *Semotilus atromaculatus* (Mitchill) 222
 Drawing by H. L. Todd from No. 19163, U.S.N.M., collected
 by Robert Kennicott in Aux Plaines River, Illinois.

PLATE XLI.

101. *Ptychocheilus oregonensis* (Richardson) 224
 Drawing by A. H. Baldwin.
 102. *Leuciscus bicolor* (Girard) 232
 Drawing by Chloe Lesley Starks from a specimen collected in
 Upper Klamath Lake, Oregon, by Gilbert, Cramer, and
 Otaki.
 103. *Leuciscus lineatus* (Girard) 232
 Drawing by S. F. Denton from a specimen collected in Yellow-
 stone Park by Dr. Jordan.

PLATE XLII.

104. *Leuciscus hydrophlox* (Cope) 238
 Drawing by S. F. Denton from a specimen collected in Yellow-
 stone Park by Dr. Jordan.

	Text page.
105. <i>Leuciscus balteatus</i> (Richardson).....	238
Drawing by H. L. Todd from No. 36869, U.S.N.M., collected in the Columbia River, Oregon, by J. H. Clarke.	
105a. <i>Leuciscus balteatus</i> (Richardson).....	238
Drawing by A. H. Baldwin from No. 43953, U.S.N.M., type of <i>Leuciscus gilli</i> Evermann, collected in Browns Gulch Creek, Silver Bow, Montana, by Evermann, Jenkins, and Clapham.	

PLATE XLIII.

106. <i>Leuciscus siuslawi</i> Evermann and Meek	279
Drawing by A. H. Baldwin from the type, No. 48480, U.S.N.M., collected by Dr. Meek in the Siuslaw River, Mapleton, Oregon.	
107. <i>Rutilus bicolor</i> (Girard).....	241
Drawing by Anna L. Brown from a specimen collected by Gilbert, Cramer, and Otaki in Upper Klamath Lake, Oregon.	

PLATE XLIV.

108. <i>Opsopoeodus osculus</i> Evermann.....	248
Drawing by A. H. Baldwin from the type, No. 45560, U.S.N.M., collected by Evermann, Scovell, and Gurley in Buffalo Bayou, near Houston, Texas.	
109. <i>Opsopoeodus emiliae</i> Hay	248
Drawing by A. H. Baldwin.	
110. <i>Opsopoeodus bollmani</i> Gilbert	248
Drawing by S. F. Denton from the type, collected by Bollman and Fesler in Buckland Creek, Millen, Georgia.	

PLATE XLV.

111. <i>Aramis crysoleucas</i> (Mitchill)	250
Drawing from No. 20243, U.S.N.M., collected by Professor Baird in the Hackensack River, New Jersey.	
112. <i>Notropis aztecus</i> Woolman	258
Drawing by A. H. Baldwin from the type, No. 45569, U.S.N.M., collected by A. J. Woolman at the City of Mexico.	
113. <i>Notropis welaka</i> Evermann and Kendall	279
Drawing by A. H. Baldwin from the type, No. 48786, U.S.N.M., collected by Dr. Kendall in St. Johns River, Welaka, Florida.	

PLATE XLVI.

114. <i>Notropis cayuga atrocaudalis</i> Evermann.....	260
Drawing by A. H. Baldwin from the type, No. 45537, U.S.N.M., collected by Evermann, Scovell, and Gurley in the Neches River, near Palestine, Texas.	

115. *Notropis*
Drawing
coll.
W.
116. *Notropis*
Drawing
coll.
Ch.

117. *Notropis*
Drawing
coll.
Riv.

118. *Notropis*
Drawing
coll.
Riv.

119. *Notropis*
Drawing
D. H.

120. *Notropis*
Drawing
Huds.
Cree.

121. *Notropis*
Drawing
Whit.

122. *Notropis*
Drawing
Ever.
Plain.

123. *Notropis*
Drawing
collect.
Luray.

124. *Notropis*
Drawing
Ever.
Mount.

125. *Notropis*
Drawing
Ever.
Marco.

Text page.

115. *Notropis kanawha* Jordan and Jenkins 264
 Drawing by S. F. Denton from the type, No. 39928, U.S.N.M.,
 collected by Jordan, Evermann, and Jenkins, in Reed Creek,
 Wytheville, Virginia.
 116. *Notropis chihuahua* Woolman 265
 Drawing by A. H. Baldwin from the type, No. 44151, U.S.N.M.,
 collected by Woolman and Cox in the Rio de las Conchos,
 Chihuahua, Mexico.

PLATE XLVII.

117. *Notropis nux* Evermann 267
 Drawing by A. H. Baldwin from the type, No. 45555, U.S.N.M.,
 collected by Evermann, Seovell, and Gurley, in Trinity
 River, near Palestine, Texas.
 118. *Notropis nocomis* Evermann 268
 Drawing by A. H. Baldwin from the type, No. 45556, U.S.N.M.,
 collected by Evermann, Scovell, and Gurley in Trinity
 River, Magnolia Point, Texas.
 119. *Notropis hudsonius* (Clinton) 269
 Drawing by A. H. Baldwin from a specimen collected by Geo.
 D. Head in Kilpatrick Lake, Minnesota.

PLATE XLVIII.

120. *Notropis hudsonius saludanus* (Jordan and Brayton) 270
 Drawing by H. L. Todd from No. 23513, U.S.N.M., the type of
Hudsonius euryopa Bean, collected by A. Graves in McBean
 Creek, Georgia.
 121. *Notropis whipplii* (Girard) 278
 Drawing by H. L. Todd from No. 36748, U.S.N.M., collected in
 White River, Indiana, by Dr. Gilbert.
 122. *Notropis galacturus* (Cope) 279
 Drawing by A. H. Baldwin from a specimen collected by
 Evermann, Scovell, and Gurley in Tellico River, Tellico
 Plains, Tennessee.

PLATE XLIX.

123. *Notropis macdonaldi* Jordan and Jenkins 284
 Drawing by S. F. Denton from the type, No. 39859, U.S.N.M.,
 collected by Jordan and Jenkins in the Shenandoah River,
 Luray, Virginia.
 124. *Notropis coccogenis* (Cope) 284
 Drawing by A. H. Baldwin from a specimen collected by
 Evermann, Scovell, and Gurley in Aruwine Spring Creek,
 Mount Verd, Tennessee.
 125. *Notropis swaini* Jordan 290
 Drawing by A. H. Baldwin from a specimen collected by
 Evermann, Scovell, and Gurley in San Marcos River, San
 Marcos, Texas.

PLATE L.

Text page.

126. *Notropis telescopus* (Cope) 292
 Drawing by A. H. Baldwin from a specimen collected by Evermann, Sevall, and Gurley in Tellico River, Tellico Plains, Tennessee.
127. *Notropis notemigonoides* Evermann 292
 Drawing by A. H. Baldwin from the type, No. 45559, U.S.N.M., collected by Evermann, Scovell, and Gurley in Neches River, near Palestine, Texas.
128. *Notropis metallicus* Jordan and Meek 297
 Drawing by A. H. Baldwin from a specimen collected by A. J. Woolman in the Withlacoochee River, near Richland, Florida.

PLATE LI.

129. *Ericympba buccata* Cope 302
 Drawing by A. H. Baldwin from No. 36803, U.S.N.M., collected by Dr. Meek in Lost Creek, near Defiance, Ohio.
130. *Phenacobius iranops* Cope 304
 Drawing by A. H. Baldwin from a specimen collected by Evermann, Scovell, and Gurley in Chickamanga Creek, Lee & Gordon's mill, Georgia.
131. *Evarra eigenmanni* Woolman 304
 Drawing by A. H. Baldwin from the type, No. 45571, U.S.N.M., collected by Woolman and Cox at the City of Mexico.

PLATE LII.

132. *Rhinichthys dulcis* (Girard). 306
 Drawing by S. F. Denton from a specimen collected by Dr. Jordan in Yellowstone Park.
133. *Agosia klamathensis* Evermann and Meek 314
 Drawing by A. H. Baldwin from the type, No. 48225, U.S.N.M., collected by Meek and Alexander in Upper Klamath Lake, Oregon.
134. *Agosia umatilla* Gilbert and Evermann 313
 Drawing by A. H. Baldwin from the type, No. 45390, U.S.N.M., collected by Gilbert and Rutter in the Columbia River at Umatilla, Oregon.

PLATE LIII.

135. *Agosia falcata* Eigenmann and Eigenmann 313
 Drawing by A. H. Baldwin from the type collected by C. H. Eigenmann in Boise River near Caldwell, Idaho.
136. *Hybopsis aestivalis marconis* Jordan and Gilbert 316
 Drawing by A. H. Baldwin from a specimen collected in the San Marcos River, San Marcos, Texas, by Evermann, Scovell, and Gurley.
137. *Hybopsis watauga* Jordan and Evermann 319
 Drawing by S. F. Denton from the type, No. 39929, U.S.N.M., collected by Jordan and Evermann in the Watauga River near Elizabethtown, Tennessee.

138. *Hybop-**Draw-**col-**A.*139. *Platyg-**Draw-**Eve-**lain*140. *Exoglo-**Draw-*141. *Brycon-**Drawi-**by L.*142. *Derichto-**Drawin-**tress*143. *Anguilla-**Drawin-**at Ho*144. *Simench-**Drawin-**by Ca*145. *Ilyophis-**Drawin-**Albat*146. *Synapho-**Drawin-**by Ca*147. *Histiobra-**Drawin-**by the*148. *Leptocep-**Drawin-**at No*149. *Leptocep-**Drawin-**Bean,**Pensac**Bull. No. 4*

PLATE LIV.

	Text page.
138. <i>Hybopsis altus</i> (Jordan)	321
Drawing by H. L. Todd from the type, No. 23125, U.S.N.M., collected in Lake Tupataro, Guanajuato, Mexico, by Prof. A. Duges.	
139. <i>Platygobio gracilis</i> (Richardson)	326
Drawing by A. H. Baldwin from a specimen collected by Evermann, Cox, and Rutter in White River near Chamberlain, South Dakota.	
140. <i>Exoglossum maxillingua</i> (Le Sueur)	327
Drawing by Anna L. Brown.	

PLATE LV.

141. <i>Brycon dentex</i> Giinther	337
Drawing by A. H. Baldwin from No. 39909, U.S.N.M., collected by Dr. L. E. II. Best in Nicaragua.	
142. <i>Derichthys serpentinus</i> Gill.....	343
Drawing by H. L. Todd from the type collected by the <i>Albatross</i> in the Gulf Stream at Station 2094 in 1,022 fathoms.	
143. <i>Anguilla chrysypa</i> Rafinesque.....	348
Drawing by H. L. Todd from No. 20470, U.S.N.M., collected at Holyoke, Massachusetts.	

PLATE LVI.

144. <i>Simenichelys parasiticus</i> Gill	349
Drawing by H. L. Todd from No. 21673, U.S.N.M., collected by Capt. N. McPhee near Sable Island Bank.	
145. <i>Ilyophis brunneus</i> Gilbert.....	350
Drawing by A. H. Baldwin from the type collected by the <i>Albatross</i> at Chatham Island.	
146. <i>Synaphobranchus pinnatus</i> (Gronow)	351
Drawing by H. L. Todd from No. 21681, U.S.N.M., collected by Captain Olsen on Le Have Bank.	

PLATE LVII.

147. <i>Histiobranchus infernalis</i> Gill	352
Drawing by H. L. Todd from No. 38205, U.S.N.M., collected by the <i>Albatross</i> in lat. $36^{\circ} 35' N.$, long. $74^{\circ} 08' 30'' W.$	
148. <i>Leptocephalus conger</i> (Linnaeus)	354
Drawing by H. L. Todd from No. 16027, U.S.N.M., collected at Noank, Connecticut, by J. W. Latham.	
149. <i>Leptocephalus caudilimbatus</i> (Poey)	355
Drawing by H. L. Todd from the type of <i>Conger caudicula</i> Beau, No. 30709, U.S.N.M., collected by Silas Stearns at Pensacola, Florida.	

PLATE LVIII.

	Text page.
150. Congrellus flavus (Goode and Bean)	357
Drawing by H. L. Todd from the type, No. 44612, U.S.N.M., collected by the <i>Albatross</i> in the Gulf Stream.	
151. Hoplunnis schmidtii Kaup	361
Drawing by J. C. Van Hook from No. 44240, U.S.N.M., col- lected by the <i>Albatross</i> at Station 2402.	
152. Venefica proceria Goode and Bean	365
Drawing by H. L. Todd from a specimen collected in the Gulf Stream.	
153. Serrivomer beanii Gill and Ryder	367
Drawing by A. H. Baldwin from the type, No. 33383, U.S.N.M., collected by the <i>Albatross</i> in the Gulf Stream.	

PLATE LIX.

154. Avocettina gillii Bean	367; 2801
Drawing by A. H. Baldwin from the type, No. 44239, U.S.N.M., collected by the <i>Albatross</i> east of Prince of Wales Island, Alaska, at Station 2859.	
154a, 154b. Avocettina gillii Bean	2801
Drawings by Anna L. Brown from No. 679, L. S. Jr. Univ. Mus., collected by the <i>Albatross</i> at Station 2860.	
155. Labichthys carinatus Gill and Ryder	368
Drawing by A. H. Baldwin from the type, No. 33239, U.S.N.M., collected by the <i>Albatross</i> at Station 2076 in the Gulf Stream.	
156. Labichthys elongatus Gill and Ryder	369
Drawing by A. H. Baldwin from the type, No. 33755, U.S.N.M., collected by the <i>Albatross</i> at Station 2100 in the Gulf Stream.	

PLATE LX.

157, 157a, 157b. Nervichthys avocetta Jordan and Gilbert	369
Drawings by Anna L. Brown from a specimen in the Pro- vincial Museum at Victoria, British Columbia, collected by M. W. Norgate, at Beacon Hill, near Victoria.	
158. Ahlia egmontis (Jordan)	370
Drawing by H. L. Todd from the type, No. 35086, U.S.N.M., collected by Dr. Jordan at Egmont Key, Florida.	
159. Verma kendalli (Gilbert)	375
Drawings by A. H. Baldwin from the type, No. 44304, U.S.N.M., collected by W. C. Kendall off west coast of Florida.	

PLATE LXI.

160. Letharchus velifer Goode and Bean	375
Drawing by H. L. Todd from the type, No. 31458, U.S.N.M., collected by Kaiser and Martin, on west coast of Florida.	
161. Myrichthys tigrinus Girard	376
Drawing by A. H. Baldwin.	

162. **Myripristis** Dr. Univ.
163. **Pisodonos** Draw col.
164. **Calechelys** Draw col.
165. **Bascanichthys** Draw Goo and
166. **166a. Bathyraja** Draw
167. **Quassiremus** Draw
168. **Ophichthus** Draw col.
169. **Ophichthoides** Draw by S.
170. **Mystrius** Draw by K.
171. **Lycodes** Draw at Ga.
172. **Muraena** Drawin
173. **Muraena** Drawin collect
174. **Channichthys** Drawin
175. **Saccopharynx** Drawin

PLATE LXII.

	Text page.
162. <i>Myrichthys pantostigmus</i> Jordan and McGregor	2802
Drawing by W. S. Atkinson from the type, No. 5710, L. S. Jr. Univ. Mus., collected by R. C. McGregor, at Clarion Island.	
163. <i>Pisodonophis cruentifer</i> Goode and Bean	377
Drawing by A. H. Baldwin from the type, No. 28938, U.S.N.M., collected by the <i>Fish Hawk</i> .	

PLATE LXIII.

164. <i>Calechelys muræna</i> Jordan and Evermann	378
Drawing by A. H. Baldwin from the type, No. 37996, U.S.N.M., collected by Silas Stearns at the Pensacola Snapper Banks.	
165. <i>Bascanichthys acuticaris</i> (Goode and Bean)	378
Drawing by H. L. Todd from the type of <i>Sphagebranchus teres</i> Goode and Bean, No. 31457, U.S.N.M., collected by Kaiser and Martin on west coast of Florida.	
166, 166a. <i>Bascanichthys peninsulæ</i> (Gilbert)	379
Drawings by A. H. Baldwin from the type, No. 44297, U.S.N.M., collected by the <i>Albatross</i> in La Paz Bay, Lower California.	

PLATE LXIV.

167. <i>Quassiremus evionthas</i> (Jordan and Bollman)	380
Drawing by A. H. Baldwin.	
168. <i>Ophichthus guttifer</i> (Bean and Dresel)	383
Drawing by H. L. Todd from the type, No. 32647, U.S.N.M., collected by Silas Stearns on the Pensacola Snapper Banks.	
169. <i>Ophichthus ocellatus</i> (Le Sueur)	383
Drawing by H. L. Todd from No. 22289, U.S.N.M., collected by Silas Stearns at Pensacola, Florida.	

PLATE LXV.

170. <i>Mystriophis intertinctus</i> (Richardson)	386
Drawing by H. L. Todd from No. 22865, U.S.N.M., collected by Kaiser and Martin on west coast of Florida.	
171. <i>Lycodontis moringa</i> (Cuvier)	395
Drawing by H. L. Todd from No. 6994, U.S.N.M., collected at Garden Key, Florida.	
172. <i>Muraena insularum</i> Jordan and Davis	400
Drawing by A. H. Baldwin.	

PLATE LXVI.

173. <i>Muraena retifera</i> Goode and Bean	401
Drawing by H. L. Todd from the type, No. 31393, U.S.N.M., collected by C. C. Leslie, at Charleston, South Carolina.	
174. <i>Channomuraena vittata</i> (Richardson)	404
Drawing by A. H. Baldwin.	
175. <i>Saccopharynx ampullaceus</i> (Harwood)	406
Drawing from Günther, Challenger Report, Vol. xxii, Pl. LXVI.	

PLATE LXVII.

	Text page.
176. <i>Gastrostomus bairdii</i> Gill and Ryder	406
Drawing by H. L. Todd from No. 33386, U.S.N.M., collected by the <i>Albatross</i> at Station 2074, in N. lat. $41^{\circ} 43'$, W. long. $65^{\circ} 21' 50''$, at a depth of 1,309 fathoms.	
177. <i>Tarpon atlanticus</i> (Cuvier and Valenciennes)	409
Drawing by H. L. Todd.	
178. <i>Elops saurus</i> Linnaeus	410
Drawing by H. L. Todd.	

PLATE LXVIII.

179. <i>Albula vulpes</i> (Linnaeus)	411
Drawing by H. L. Todd.	
180. <i>Hiodon tergisus</i> Le Sueur	413
Drawing by H. L. Todd from No. 8710, U.S.N.M., collected by George Clark, at Ecorse, Michigan.	
181. <i>Hiodon selenops</i> Jordan and Bean	411
Drawing by H. L. Todd from the type, No. 19844, U.S.N.M., collected by Daniel J. Duffy, at Chattanooga, Tennessee.	

PLATE LXIX.

182. <i>Chanos chanos</i> (Forskål).....	414
Drawing by A. H. Baldwin from No. 28240, U.S.N.M., col- lected by Dr. Gilbert at Mazatlan, Mexico.	
183. <i>Dorosoma cepedianum</i> (Le Sueur)	416
Drawing by H. L. Todd.	
184. <i>Signalosa atchafalayæ</i> Evermann and Kendall	2809
Drawing by A. H. Baldwin from the type, No. 48790, U.S.N.M., collected by Fred M. Chamberlain in the Atchafalaya River at Melville, Louisiana.	

PLATE LXX.

185. <i>Clupea harengus</i> Linnaeus.....	421
Drawing by H. L. Todd.	
186. <i>C'upea pallasi</i> Cuvier and Valenciennes.....	422
Drawing by H. L. Todd from No. 27718, U.S.N.M., collected by L. Bailey at Unalaska.	
187. <i>Pomolobus chrysichloris</i> Rafinesque	425
Drawing by H. L. Todd from No. 30159, U.S.N.M., collected by Silas Stearns at Pensacola, Florida.	

PLATE LXXI.

188. <i>Pomolobus mediocris</i> (Mitchill)	425
Drawing by H. L. Todd from No. 25132, U.S.N.M., collected at Washington market by Dr. Bean.	

189. *Pomacanthus*
190. *Pomacanthus*
191. *Alosa*
192. 192a. *Alosa*
193. *Sardinella*
194. *Sardinella*
195. *Brevoclinus*
196. *Stolephorus*
197. *Alepocephalus*
198. *Talismanus*
199. *Aleposoma*

	Text page.
189. <i>Pomolobus pseudoharengus</i> (Wilson).....	426
Drawing by H. L. Todd from No. 25197, U.S.N.M., collected at Washington market.	

	Text page.
190. <i>Pomolobus æstivalis</i> (Mitchill)	426
Drawing by H. L. Todd from No. 32639, U.S.N.M., collected at Washington market.	

PLATE LXXII.

191. <i>Alosa sapidissima</i> (Wilson).....	427
Drawing by H. L. Todd from No. 25099, U.S.N.M., collected by U. S. F. C. at Norfolk, Virginia.	
192. 192a. <i>Alosa alabamae</i> Jordan and Evermann.....	2810
Drawings by A. H. Baldwin from the types, Nos. 47690, male, and 47689, female, U.S.N.M., collected by J. N. Fitts in Black Warrior River, at Tuscaloosa, Alabama.	

PLATE LXXIII.

193. <i>Sardinella humeralis</i> (Cuvier & Valenciennes)	431
Drawing by H. L. Todd from the type of <i>Harengula pensacola</i> Goode and Bean, No. 22831, U.S.N.M., collected by Silas Stearns at Pensacola, Florida.	
194. <i>Sardinella stolifera</i> (Jordan and Gilbert).....	431
Drawing by Anna L. Brown from No. 2693, L. S. Jr. Univ. Mus., collected by the Hopkins Expedition at Mazatlan, Mexico.	
195. <i>Brevoortia tyrannus</i> (Latrobe).....	433
Drawing by H. L. Todd from No. 20666, U.S.N.M., collected at Woods Hole, Massachusetts.	

PLATE LXXIV.

196. <i>Stolephorus perthecatus</i> Goode and Bean	442
Drawing by H. L. Todd from No. 30483, U.S.N.M., collected by Silas Stearns in south Florida.	
197. <i>Alepocephalus agassizii</i> Goode and Bean	453
Drawing by H. L. Todd from No. CCCXXXVIII, M. C. Z., collected by A. Agassiz in N. lat. $38^{\circ} 18' 40''$, W. long. $73^{\circ} 18' 10''$, at a depth of 922 fathoms.	
198. <i>Talismania antillarum</i> Goode and Bean.....	455
Drawing by M. M. Smith from the type, No. 43739, U.S.N.M., collected by the <i>Albatross</i> at Station 2304, in N. lat. $28^{\circ} 38' 30''$, W. long. $87^{\circ} 02'$, at a depth of 420 fathoms.	

PLATE LXXV.

199. <i>Aleposomus copei</i> Gill.....	459
Drawing by H. L. Todd from the type, No. 33551, U.S.N.M., collected by the <i>Albatross</i> at Station 2099, in N. lat. $37^{\circ} 12' 20''$, W. long. $69^{\circ} 39'$, at a depth of 2,949 fathoms.	

	Text page
200. <i>Coregonus williamsoni</i> Girard	463
Drawing by A. H. Baldwin from a breeding male collected by B. A. Bean in Little Spokane River, near Clarks Spring, Washington.	
200a. <i>Coregonus williamsoni</i> Girard.....	463
Drawing by A. H. Baldwin from a young individual col- lected by Dr. Evermann in Montana.	

PLATE LXXVI.

201. <i>Coregonus coulteri</i> Eigenmann and Eigenmann.....	462
Drawing by A. H. Baldwin from the type, No. 44875, U.S. N.M., collected by Dr. Eigenmann in the Kicking Horse River at Field, British Columbia.	
202. <i>Coregonus clupeiformis</i> (Mitchill).....	465
Drawing by H. L. Todd from No. 10300, U.S.N.M., collected by George Clark at Ecorse, Michigan.	
203. <i>Argyrosomus nigripinnis</i> Gill.....	472
Drawing by A. H. Baldwin from a nearly ripe male collected in Lake Michigan, off Sheboygan, Wisconsin, in 90 fathoms.	

PLATE LXXVII.

204. <i>Stenodus mackenzii</i> (Richardson)	474
Drawing from No. 29889, U.S.N.M., collected at Nulato, Alaska, by E. W. Nelson.	
205. <i>Oncorhynchus gorbuscha</i> (Walbaum).....	478
Drawing by H. L. Todd from No. 27743, U.S.N.M., collected by Dr. Bean in Cooks Inlet, Alaska.	
206. <i>Oncorhynchus tshawytscha</i> (Walbaum).....	479
Drawing by H. L. Todd.	

PLATE LXXVIII.

207. <i>Oncorhynchus nerka</i> (Walbaum)	481
Drawing by A. H. Baldwin from a male specimen of the large form collected by T. M. Williams at Big Payette Lake, Idaho.	
207a. <i>Oncorhynchus nerka</i> (Walbaum).....	481
Drawing by A. H. Baldwin from a female specimen of the small form collected by Dr. Evermann at Alturas Lake, Idaho.	
207b. <i>Oncorhynchus nerka</i> (Walbaum).....	481
Drawing by A. H. Baldwin from a mutilated male specimen of the small form collected by Dr. Evermann at Alturas Lake, Idaho.	

PLATE LXXIX.

208. <i>Salmo clarkii henshawi</i> (Gill and Jordan).....	493; 2819
Drawing by S. F. Denton from No. 17086, U.S.N.M., collected by H. W. Henshaw in Lake Tahoe, California.	

209. *Salmo*
Draw
Ma
210. *Salmo*
Draw
an

211. *Salmo*
Draw
Draw
an

212. *Salmo*
Draw
Ch

213. *Salmo*
Draw
Draw
dan

214. *Salmo*
Draw
coll

215. *Salmo* g

216. *Salmo* i
Draw
exam
Mis

217. *Cristivo*
Drawi

218. *Salvelin*

Drawi

by E

219. *Salvelin*

Drawi

by T

220. *Salvelin*

Drawi

cole

shire

221. *Salvelin*

Drawi

by E

222. *Thymall*

Drawi

Nels

- Text page.
209. *Salmo clarkii virginalis* (Girard) 495; 2819
 Drawing by S. F. Denton from a specimen collected by Peter
 Madsen in Utah Lake, Utah.
210. *Salmo clarkii spilurus* (Cope) 495; 2819
 Drawing by S. F. Denton from a specimen collected by Jordan
 and Evermann at Del Norte, Colorado.

PLATE LXXX.

211. *Salmo clarkii pleuriticus* (Cope) 496; 2819
 Drawing by S. F. Denton from specimen collected by Jordan
 and Evermann at Trappers Lake, Colorado.
212. *Salmo clarkii bouvieri* (Bendire) 496; 2819
 Drawing by S. F. Denton from the type, collected by Capt.
 Charles Bendire at Waha Lake, Idaho.
213. *Salmo clarkii stomias* (Cope) 497; 2819
 Drawing by S. F. Denton from a specimen collected by Jor-
 dan and Evermann at Twin Lakes, Colorado.

PLATE LXXXI.

214. *Salmo clarkii macdonaldi* Jordan and Evermann 497; 2819
 Drawing by S. F. Denton from the type, No. 41730, U.S.N.M.,
 collected by George Fisher in Twin Lakes, Colorado.
215. *Salmo gairdneri* Richardson 497
216. *Salmo irideus* Gibbons 500
 Drawing by S. F. Denton from No. 37782, U.S.N.M., a male
 example, collected by William Montgomery at Verona,
 Missouri, where the species had been introduced.

PLATE LXXXII.

217. *Cristivomer namaycush* (Walbaum) 504
 Drawing by S. F. Denton.
218. *Salvelinus fontinalis* (Mitchill) 506
 Drawing by H. L. Todd from No. 28651, U.S.N.M., collected
 by E. G. Blackford.
219. *Salvelinus malma* (Walbaum) 507
 Drawing by H. L. Todd from No. 27740, U.S.N.M., collected
 by T. H. Bean in Cooks Inlet.

PLATE LXXXIII.

220. *Salvelinus alpinus aureolus* (Bean) 511
 Drawing by H. L. Todd from No. 39899, U.S.N.M., a female,
 collected by E. B. Hodge at Sunapee Lake, New Hamp-
 shire.
221. *Salvelinus oxyessa* (Girard) 514
 Drawing by H. L. Todd from No. 20688, U.S.N.M., collected
 by E. G. Blackford at Oquassa Lake, Maine.
222. *Thymallus signifer* (Richardson) 517
 Drawing by H. L. Todd from a specimen collected by E. W.
 Nelson at Nulato, Alaska.

PLATE LXXXIV.

- Text page.
23
23
223. *Thymallus tricolor* Cope 518; 281
 Drawing by H. L. Todd from No. 11115, U.S.N.M., collected
 by J. W. Milner in Ansable River, Michigan.
224. *Thymallus tricolor montanus* (Milner) 519; 281
 Drawing by S. F. Denton from a specimen collected by Dr.
 Jordan at Horsethief Springs, Montana.
- 224a. *Thymallus tricolor montanus* (Milner) 519; 281
 Drawing by A. H. Baldwin from a young example collected
 in the Yellowstone Park by Dr. Evermann.

PLATE LXXXV.

225. *Mallotus villosus* (Miller) 520
 Drawing by H. L. Todd from No. 3418, U.S.N.M., collected
 by Dr. E. Cones in Grosswater Bay.
226. *Thaleichthys pacificus* (Richardson) 521
 Drawing by H. L. Todd from No. 28001, U.S.N.M., collected
 by Marcus Baker in Chilkat River, Alaska.
227. *Osmerus thaleichthys* Ayres 522
 Drawing by H. L. Todd from No. 28087, U.S.N.M., collected
 by Dr. Bean at San Francisco, California.

PLATE LXXXVI.

228. *Osmerus mordax* (Mitchill) 523
 Drawing by H. L. Todd from No. 21435, U.S.N.M., collected
 by V. N. Edwards at Woods Hole, Massachusetts.
229. *Osmerus dentex* Steindachner 524
 Drawing by H. L. Todd from No. 27914, U.S.N.M., collected
 at Port Clarence, Alaska, by Dr. Bean.
230. *Hypomesus pretiosus* (Girard) 525
 Drawing by H. L. Todd from No. 27995, U.S.N.M., collected
 in Yakutat Bay, Alaska, by Dr. Bean.

PLATE LXXXVII.

231. *Hypomesus olidus* (Pallas) 525
 Drawing by H. L. Todd from No. 23973, U.S.N.M., collected
 by L. M. Turner at St. Michaels, Alaska.
232. *Argentina silus* Ascanius 526
 Drawing by H. L. Todd from No. 37801, U.S.N.M., collected
 by E. H. Bunker at Biddeford Pool, Maine.
233. *Leuroglossus stilbius* Gilbert 527
 Drawing by A. H. Baldwin from the type, No. 44283, U.S.N.M.,
 collected by the *Albatross* at Station 2997 in 221 fathoms.

PLATE LXXXVIII.

234. *Batryllagus benedicti* Goode and Bean 529
 Drawing by A. H. Baldwin from a specimen collected by the
Albatross at Station 2711, in N. lat. $38^{\circ} 59'$, W. long. $70^{\circ} 07'$, in 1,344 fathoms.

23
23
23
23238
238
239
239240
241
241242
243
243244. 1
245. 1

	Text page.
235. <i>Thachinocephalus myops</i> (Forster).....	533
Drawing by H. L. Todd.	
236. <i>Synodus icetens</i> (Linnaeus).....	538
Drawing by H. L. Todd from No. 25998, U.S.N.M., collected by C. C. Leslie at Charleston, South Carolina.	

PLATE LXXXIX.

237. <i>Benthosaurus grallator</i> Goode and Bean.....	543
Drawing by H. L. Todd from a specimen collected by the <i>Blake</i> at Station CLXXIV, in N. lat. $24^{\circ} 23'$, W. long. 84° $23'$, in 1,850 fathoms.	
238. <i>Bathypterois quadrifilis</i> Günther.....	545
Drawing by A. H. Baldwin from a specimen collected by the <i>Blake</i> at Station XCVIII, off St. Vincent.	
239. <i>Ipnotops murrayi</i> Günther.....	547
Drawing by H. L. Todd from a specimen collected by the <i>Blake</i> at Station CCXXXIII, in N. lat. $24^{\circ} 36'$, W. long. 84° $5'$, in 955 fathoms.	

PLATE XC.

240. <i>Rondeletia bicolor</i> Goode and Bean	548
Drawing by H. L. Todd from the type, No. 38202, U.S.N.M., collected by the <i>Albatross</i> at Station 2206, in N. lat. $36^{\circ} 47'$, W. long. $73^{\circ} 25'$, in 1,641 fathoms.	
241. <i>Cetomimus gilli</i> Goode and Bean	549
Drawing by M. M. Smith from the type, No. 35529, U.S.N.M., collected by the <i>Albatross</i> at Station 2206, in N. lat. $39^{\circ} 35'$, W. long. $71^{\circ} 24' 30''$, in 1,043 fathoms.	
242. <i>Ceratoscopelus madeirensis</i> (Lowe)	557
Drawing by J. C. Van Hook from No. 43776, U.S.N.M., col- lected by the <i>Albatross</i> at Station 2328, in N. lat. $41^{\circ} 47'$ W. long. $65^{\circ} 37' 30''$, in 677 fathoms.	

PLATE XCI.

243. <i>Lampanyctus crocodilus</i> (Risso).....	558
Drawing by A. H. Baldwin from a specimen collected at Nice, and obtained by the U.S.N.M. through the Royal Zoolog- ical Museum at Florence, Italy.	
244. <i>Lampadema speculigera</i> Goode and Bean	561
Drawing by J. C. Van Hook from the type, No. 43797, U.S.N.M., collected by the <i>Fish Hawk</i> at Station 797, off Newport, Rhode Island, in 16.5 fathoms.	
245. <i>Nannobrachium macdonaldi</i> Goode and Bean	563
Drawing by S. F. Denton from No. 35445, U.S.N.M., collected by the <i>Albatross</i> at Station 2182, in N. lat. $39^{\circ} 25' 30''$, W. long. $71^{\circ} 44'$, in 861 fathoms.	

PLATE XCII.

- Text page.
246. *Aethophrora lucida* Goode and Bean 565
 Drawing by A. H. Baldwin from No. 44084, U.S.N.M., collected by the *Albatross* at Station 2127, in N. lat. $19^{\circ} 45'$, W. long. $75^{\circ} 04'$, in 1639 fathoms.
247. *Myctophum opalinum* Goode and Bean 571
 Drawing by J. C. Van Hook from No. 43798, U.S.N.M., collected by the *Albatross* at Station 2585, in N. lat. $39^{\circ} 08' 30''$, W. long. $72^{\circ} 17'$, in 542 fathoms.
248. *Tarletonbeania tenua* Eigenmann and Eigenmann 575
 Drawing by A. H. Baldwin from No. 41882, U.S.N.M., collected by C. H. Eigenmann off Point Loma, near San Diego, California.

PLATE XCIII.

249. *Varrella blackfordii* Goode and Bean 584
 Drawing by A. H. Baldwin from the type, No. 44242, U.S.N.M., collected by the *Albatross* at Station 2276, in N. lat. $29^{\circ} 03' 15''$, W. long. $88^{\circ} 16'$, in 321 fathoms.
250. *Chauliodus sloanei* Bloch and Schneider 585
 Drawing by H. L. Todd from No. 23420, U.S.N.M., collected by Capt. Charles Anderson and crew, of the Gloucester fishing fleet, in N. lat. $42^{\circ} 08'$, W. long. $65^{\circ} 35'$, in 185 fathoms.
251. *Astronesthes gemmifer* Goode and Bean 586
 Drawing by A. H. Baldwin from the type, No. 24645, U.S.N.M., collected by the schooner *Polar Ware* from the stomach of a halibut, in N. lat. $44^{\circ} 25'$, W. long. $53^{\circ} 12'$, in 300 fathoms.

PLATE XCIV.

252. *Astronesthes richardsoni* Poey 587
 Drawing by M. M. Smith from No. 35540; U.S.N.M., collected by the *Albatross* at Station 2202, in N. lat. $39^{\circ} 38'$, W. long. $71^{\circ} 39' 45''$, in 515 fathoms.
253. *Stomias ferox* Reinhardt 588
 Drawing by H. L. Todd from No. 23360, U.S.N.M., collected by Capt. David Cammel and crew, of the Gloucester fishing fleet, at East Banquereux.
254. *Grammatostomias dentatus* Goode and Bean 590
 Drawing by H. L. Todd from the type, No. 37370, U.S.N.M., collected by the *Albatross*, at Station 2565, in N. lat. $38^{\circ} 19' 20''$, W. long. $69^{\circ} 02' 30'$, in 2069 fathoms.

PLATE XCV.

255. *Photonectes gracilis* Goode and Bean 591
 Drawing by M. M. Smith from the type, collected by the *Blake* at Station XL, off Martinique, in 472 fathoms.

256. *Malacoctenus* Dray by G. 127
257. *Alepis* Drawn 127
258. *Alepis* Drawn 127
259. *Arctozenus* Drawn 127
260. *Paralepidotus* Drawn Nat
- 261, 261a. *Aldrovanda* Drawn Nat
262. *Aldrovanda* Drawn Blak
263. *Aldrovanda* Drawn Blak
264. *Notacanthus* Drawn coll. W. L.
265. *Notacanthus* Drawn coll. ever,
266. *Lipogenys* Drawn coll. 30%

	Text page.
256. <i>Malacosteus niger</i> Ayres.....	593
Drawing by H. L. Todd, from No. 32169, U.S.N.M., collected by Capt. Charles Anderson and crew, of the schooner <i>Alice</i> <i>G. Wanson</i> , on the northeastern edge of Georges Bank, in 125 fathoms.	
257. <i>Alepisaurus ferox</i> Lowe.....	595
Drawing by H. L. Todd from No. 20593, U.S.N.M., ob- tained in the New York market by E. G. Blackford.	

PLATE XCVI.

258. <i>Alepisaurus æsculapius</i> (Bean).....	595
Drawing by H. L. Todd from the type, No. 27705, U.S.N.M., collected by Robert King at Unalaska, Alaska.	
259. <i>Arctozenus coruscans</i> (Jordan & Gilbert).....	601
Drawing by A. H. Baldwin from the type, No. 27171, U.S.N.M., collected by Mr. Brown, at Port Townsend, Washington.	
260. <i>Paralepis coregonoides</i> Risso.....	602
Drawing by H. L. Todd from a specimen in the Academy of Natural Sciences of Montreal.	

PLATE XCVII.

261, 261a. <i>Argyropelecus olfersi</i> (Cuvier).....	604
Drawings by H. L. Todd from No. 33393, U. S. N. M., collected by the <i>Albatross</i> in N. lat. $41^{\circ} 40' 30''$, W. long. $65^{\circ} 35'$, in 855 fathoms.	

PLATE XCVIII.

262. <i>Aldrovandia macrochir</i> (Günther).....	609
Drawing by H. L. Todd from a specimen collected by the <i>Blake</i> at Station LIII, off Havana, in 242 fathoms.	
263. <i>Aldrovandia gracilis</i> Goode and Bean.....	610
Drawing by S. F. Denton, from a specimen collected by the <i>Blake</i> at Station LXX, off Guadalupe, in 789 fathoms.	
264. <i>Notacanthus analis</i> Gill	615
Drawing by H. L. Todd from the type, No. 37856, U.S.N.M., collected by the <i>Albatross</i> at Station 2677, in N. lat. $32^{\circ} 39'$, W. long. $76^{\circ} 50' 30''$, in 478 fathoms.	
265. <i>Notacanthus phasganorus</i> Goode.....	616
Drawing by H. L. Todd from the type, No. 25972, U.S.N.M., collected by Capt. Briggs Gilpatrick, of the schooner <i>Gatherer</i> , from the stomach of a ground shark, on the Grand Bank.	

PLATE XCIX.

266. <i>Lipogenys gillii</i> Goode and Bean	619
Drawing by H. L. Todd from the type, No. 39212, U.S.N.M., collected by the <i>Albatross</i> at Station 2742, in N. lat. $37^{\circ} 46'$ 30'', W. long. $73^{\circ} 53' 30''$, in 865 fathoms.	

	Text page
267. <i>Dallia pectoralis</i> Bean.....	621
Drawing by H. L. Todd from the type, No. 23498, U.S.N.M., collected by L. M. Turner, at St. Michaels, Alaska.	
268. <i>Umbrä pygmæa</i> (De Kuy).....	624
Drawing by H. L. Todd from No. 34886, U.S.N.M., collected in New Jersey.	

PLATE C.

269, 269a. <i>Lucius lucius</i> (Linnaeus)	628
Drawings by H. L. Todd from No. 9389, U.S.N.M., collected at Ecorse, Michigan.	
270. <i>Lucius masquinongy</i> (Mitchill).....	629
Drawing by H. L. Todd from No. 10607, U.S.N.M., collected by Geo. Clark at Ecorse, Michigan.	

PLATE CI.

271, 271a. <i>Fundulus majalis</i> (Walbaum)	639
Drawings by H. L. Todd from No. 13788, U.S.N.M. (female and male), collected by the U. S. Fish Commission at Woods Hole, Massachusetts.	
271b. <i>Fundulus majalis</i> (Walbaum).....	639
Drawing by A. H. Baldwin from a young example collected by H. M. Smith at St. George Island, Maryland.	

PLATE CII.

272. <i>Fundulus pallidus</i> Evermann	638
Drawing by A. H. Baldwin from the type, No. 45564, U.S.N.M., collected by Evermann, Scovell, and Gurley in Galveston Bay, near Swan Lake, Galveston, Texas.	
273. <i>Fundulus heteroclitus</i> (Linnaeus).....	640
Drawing by A. H. Baldwin from a male example collected by H. M. Smith at St. George Island, Maryland.	
274. <i>Fundulus ocellaris</i> Jordan and Gilbert.....	642
Drawing by A. H. Baldwin from a specimen collected in the Withlacoochee River, Florida, by A. J. Woolman.	

PLATE CIII.

275. <i>Fundulus diaphanus</i> (Le Sueur).....	645
Drawing by A. H. Baldwin from a male example collected by H. M. Smith at St. George Island, Maryland.	
275a. <i>Fundulus diaphanus</i> (Le Sueur)	645
Drawing by A. H. Baldwin from a female example collected by H. M. Smith at St. George Island, Maryland.	
276. <i>Fundulus zebra</i> Jordan and Gilbert.....	646
Drawing by H. L. Todd from No. 36610, U.S.N.M., collected at Ellis, Kansas, by F. W. Cragin.	

277. *Fundulus* Drawn by Ch.
 278. *Fundulus* Drawn by Dr.
 278a. *Fundulus* Drawn by Dr.
 279. *Fundulus* Drawn by Dr.
 280. *Fundulus* Drawn by Dr.
 281. *Fundulus* Drawn by Dr.
 282. *Fundulus* Drawn by Dr.
 283. *Fundulus* Drawn by Dr.
 284. *Fundulus* Drawn by Dr.
 285. *Fundulus* Drawn by Dr.
 286. *Fundulus* Drawn by Dr.
 287. *Fundulus* Drawn by Dr.

PLATE CIV.

	Text page.
277. <i>Fundulus seminolis</i> Girard.....	647
Drawing by A. H. Baldwin from a specimen collected in Charlie Apopka Creek, Florida, by A. J. Woolman.	
278. <i>Fundulus catenatus</i> (Storer).....	648
Drawing by A. H. Baldwin from a male example collected by Dr. Gurley in Ball Creek, near Tazewell, Tennessee.	
278a. <i>Fundulus catenatus</i> (Storer).....	648
Drawing by H. L. Todd from No. 36456, U.S.N.M., collected by Jordan and Gilbert in Saline River, Benton, Arkansas.	

PLATE CV.

279. <i>Fundulus stellifer</i> (Jordan).....	648
Drawing by H. L. Todd from No. 17888, U.S.N.M., collected by Dr. Jordan in Etowah River, Rome, Georgia.	
280. <i>Fundulus rathbuni</i> Jordan and Meek.....	649
Drawing by S. F. Denton from the type, No. 39860, U.S.N.M., collected in Allemane Creek, Greensboro, North Carolina, by Jordan, Jenkins, and Meek.	
281. <i>Fundulus albolineatus</i> Gilbert.....	649
Drawing by S. F. Denton from the type, collected by P. H. Kirsch in Spring Creek, Huntsville, Alabama.	

PLATE CVI.

282. <i>Fundulus funduloides</i> (Evermann)	650
Drawing by A. H. Baldwin from the type, No. 45563, U.S.N.M., collected by Evermann, Seovell, and Gurley in Dickinson Bayou, near Galveston, Texas.	
283. <i>Fundulus macdonaldi</i> (Meek).....	651
Drawing by S. F. Denton from the type collected by Meek, Drew, and Rettger in Jones Creek, Dixon, Missouri.	

284. <i>Fundulus jenkinsi</i> Evermann	651
Drawing by A. H. Baldwin from the type, No. 45562, U.S.N.M., collected by Evermann, Scovell, and Gurley in Dickinson Bayou, near Galveston, Texas.	

PLATE CVII.

285. <i>Fundulus pulvereus</i> (Evermann).....	652
Drawing by A. H. Baldwin from the type, No. 45561, U.S.N.M., collected in Dickinson Bayou, near Galveston, Texas, by Evermann, Scovell, and Gurley.	
286. <i>Fundulus luciae</i> (Baird).....	654
Drawing by A. H. Baldwin from a specimen collected by H. M. Smith at St. George Island, Maryland.	

287. <i>Fundulus chrysotus</i> Holbrook	655
Drawing by A. H. Baldwin from a specimen collected by A. J. Woolman in Florida.	

PLATE CVIII.

	Text page
288. <i>Fundulus notti</i> (Agassiz).....	656
Drawing by H. L. Todd from No. 31439, U.S.N.M., collected in Elbow Creek, Florida, by J. A. Henshall.	
289. <i>Fundulus notatus</i> (Rafinesque).....	659
Drawing by H. L. Todd from No. 36324, U.S.N.M., collected in White River at Eureka Springs, Arkansas, by Jordan and Gilbert.	
290. <i>Adinia dugesii</i> (Bean)	661
Drawing by H. L. Todd from the type, No. 37831, U.S.N.M., collected by Prof. A. Dugès, in Guanajuato, Mexico.	

PLATE CIX.

291. <i>Fundulus goodei</i> (Jordan).....	664; 2831
Drawing by A. H. Baldwin, from a specimen collected by A. J. Woolman in Florida.	
292. <i>Lucania parva</i> (Baird and Girard).....	665
Drawing by W. S. Haines from a specimen collected by Dr. Bean in New Jersey.	
293. <i>Characodon bilineatus</i> Bean.....	668
Drawing by W. S. Haines from the type, No. 37832, U.S.N.M., collected by Professor Dugès in Guanajuato, Mexico.	

PLATE CX.

294, 294a, 294b, 294c, 294d. <i>Empetrichthys merriami</i> Gilbert	667
Drawings from the type, No. 46101, U.S.N.M., collected by Merriam and Bailey in Ash Meadows, Amargosa Desert, on the boundary between California and Nevada.	

PLATE CXI.

295, 295a. <i>Characodon variatus</i> Bean	669
Drawings by H. L. Todd from No. 37810, U.S.N.M., the type of <i>C. ferrugineus</i> Bean, collected by Professor Dugès in Guanajuato, Mexico.	
296. <i>Cyprinodon variegatus</i> Lacépède.....	671
Drawing by A. H. Baldwin from a male example collected by H. M. Smith at St. George Island, Maryland.	

PLATE CXII.

296a. <i>Cyprinodon variegatus</i> Lacépède	671
Drawing by A. H. Baldwin from a young example collected by H. M. Smith at St. George Island, Maryland.	
297. <i>Cyprinodon carpio</i> Günther	675
Drawing by H. L. Todd from No. 32151, U.S.N.M., collected by Dr. J. W. Velie at Boca Ciega Bay, Florida.	
298. <i>Jordanella floridae</i> Goode and Bean	677
Drawing by A. H. Baldwin from a specimen collected by A. J. Woolman in Florida.	

299. *Gambusia* Drawn
 collected
299a. *Gambusia* Drawn
 by
300. *Anableps* Drawn
 lecter
301. *Goodea* Drawn
 Prof
302. *Heteropneustes* Drawn
303. *Poecilia* Drawn
 L. S.
 Maz
304. *Xiphophorus* Drawn
 lecter
305. *Chologlanis* Drawn
 and J.
 Meek
306. *Typhlichthys* Drawn
 in Ma
307. *Amblyopus* Drawn
 Manu
308. *Tylosurus* Drawing
309. *Tylosurus* Drawing
 from I
310. *Chriodorus* Drawing
 collect
311. *Hyperoplus* Drawing
 by Ott

PLATE CXIII.

	Text page.
299. <i>Gambusia affinis</i> (Baird and Girard)	680
Drawing by A. H. Baldwin from No. 37839, U.S.N.M., a male, collected by Professor Dugès in Mexico.	
299a. <i>Gambusia affinis</i> (Baird and Girard)	680
Drawing by A. H. Baldwin from a female example collected by H. M. Smith at St. George Island, Maryland.	
300. <i>Anableps dovii</i> Gill.....	685
Drawing by A. H. Baldwin from No. 48214, U.S.N.M., col- lected by E. W. Nelson in Tehuantepec, Mexico.	

PLATE CXIV.

301. <i>Goodea atripinnis</i> Jordan.....	685
Drawing from the type, No. 23137, U.S.N.M., collected by Professor Dugès at Leon in Guanajuato, Mexico.	
302. <i>Heterandria formosa</i> Agassiz.....	687
Drawing by A. H. Baldwin.	
303. <i>Pœcilia presidionis</i> Jordan and Culver	697
Drawing by Anna L. Brown from the type, a female, No. 2687, L. S. Jr. Univ. Mns., collected by the Hopkins Expedition at Mazatlan, Mexico, in the Rio Presidio.	

PLATE CXV.

304. <i>Xiphophorus helleri</i> Heckel.....	701
Drawing by A. H. Baldwin from No. 44948, U.S.N.M., col- lected by A. L. Herrera at Vera Cruz, Mexico.	
305. <i>Chologaster cornutus</i> Agassiz.....	703
Drawing by S. F. Denton from the type of <i>C. aritus</i> Jordan and Jenkins, No. 39864, U.S.N.M., collected by Jenkins and Meek in the outlet of the Dismal Swamp, Virginia.	
306. <i>Typhlichthys subterraneus</i> Girard.....	704
Drawing by H. L. Todd from No. 36806, U.S.N.M., collected in Mammoth Cave, Kentucky.	
307. <i>Amblyopsis spelæus</i> DeKay	706
Drawing by H. L. Todd from No. 5863, U.S.N.M., collected in Mammoth Cave, Kentucky.	

PLATE CXVI.

308. <i>Tylosurus raphidoma</i> (Ranzani)	715
Drawing by H. L. Todd.	
309. <i>Tylosurus acus</i> (Lacépède).....	716
Drawing by H. L. Todd from No. 21422, U.S.N.M., received from E. G. Blackford.	
310. <i>Chirodorus atherinoides</i> Goode and Bean	719
Drawing by H. L. Todd from the type, No. 26593, U.S.N.M., collected by Silas Stearns at Key West, Florida.	
311. <i>Hyporhamphus unifasciatus</i> (Ranzani)	720
Drawing by H. L. Todd from No. 16944, U.S.N.M., collected by Otto Lugger in Chesapeake Bay.	

PLATE CXVII.

	Text page.
312. <i>Hyporhamphus roberti</i> (Cuvier and Valenciennes)	721
Drawing by W. S. Haines from a specimen collected at Longport, New Jersey, by Dr. Bean.	
313. <i>Hemirhamphus brasiliensis</i> (Linnaeus)	722
Drawing by H. L. Todd from No. 26400, U.S.N.M., collected at Hungers Wharf, Virginia, by J. T. Wilkins.	
314. <i>Scombrexox saurus</i> (Walbaum)	725
Drawing by H. L. Todd from No. 19853, U.S.N.M., collected by V. N. Edwards at Woods Hole.	
315. <i>Fodiator acutus</i> (Cuvier and Valenciennes)	728
Drawing by A. H. Baldwin from No. 43427, U.S.N.M., collected by the <i>Albatross</i> at Panama.	

PLATE CXVIII.

316. <i>Exonauta exsiliens</i> (Miller)	732; 2830
Drawing by H. L. Todd from No. 25187, U.S.N.M., collected by E. G. Blackford.	
317. <i>Exonauta rondeletii</i> (Cuvier and Valenciennes)	733; 2830
Drawing by H. L. Todd from No. 21871, U.S.N.M., collected by Capt. N. McPhee in N. lat. $41^{\circ} 40'$, W. long. $62^{\circ} 28'$.	
318. <i>Exocetus volitans</i> Linnaeus	734
Drawing by H. L. Todd from No. 20636, U.S.N.M., collected by V. N. Edwards at Woods Hole.	

PLATE CXIX.

319. <i>Cypsilurus californicus</i> (Cooper)	740; 2830
Drawing by H. L. Todd from No. 26907, U.S.N.M., collected by Dr. Jordan at Santa Barbara, California.	
320. <i>Gasterosteus aculeatus</i> Linnaeus	747
Drawing by H. L. Todd from No. 20875, U.S.N.M., collected at Woods Hole, Massachusetts, by V. N. Edwards.	
321. <i>Gasterosteus cataphractus</i> (Pallas)	749
Drawing by H. L. Todd from a specimen collected by Dr. Bean at St. Paul Island, Alaska.	

PLATE CXX.

322. <i>Apeltes quadratus</i> (Mitchill)	752
Drawing by A. H. Baldwin from No. 13409, U.S.N.M., collected by V. N. Edwards at Woods Hole, Massachusetts.	
323. <i>Aulorhynchus flavidus</i> Gill	754
Drawing by A. H. Baldwin from No. 21585, U.S.N.M., collected by J. G. Swan in Puget Sound.	
324. <i>Aulostomus maculatus</i> Valenciennes	754
Drawing by H. L. Todd from No. 34838, U.S.N.M., collected by U. S. Fish Commission, Potomac River, Washington.	

325. *Siphonognathus*
 Drawing by H. L. Todd from No. 21585, U.S.N.M., collected by U. S. Fish Commission, Potomac River, Washington.
 326. *Coryphaenoides*
 Drawing by H. L. Todd from No. 21585, U.S.N.M., collected by U. S. Fish Commission, Potomac River, Washington.
 327. *Hippocampus*
 Drawing by H. L. Todd from No. 21585, U.S.N.M., collected by U. S. Fish Commission, Potomac River, Washington.
 328. *Hippocampus*
 Drawing by H. L. Todd from No. 21585, U.S.N.M., collected by U. S. Fish Commission, Potomac River, Washington.
 329. *Percopis*
 Drawing by H. L. Todd from No. 21585, U.S.N.M., collected by U. S. Fish Commission, Potomac River, Washington.
 330. *Columbus*
 Drawing by H. L. Todd from No. 21585, U.S.N.M., collected by U. S. Fish Commission, Potomac River, Washington.
 331. *Aphredoderus*
 Drawing by H. L. Todd from No. 21585, U.S.N.M., collected by U. S. Fish Commission, Potomac River, Washington.
 332. *Atherinops*
 Drawing by H. L. Todd from No. 21585, U.S.N.M., collected by U. S. Fish Commission, Potomac River, Washington.
 333. *Atherina*
 Drawing by H. L. Todd from No. 21585, U.S.N.M., collected by U. S. Fish Commission, Potomac River, Washington.
 334. *Chiurostomus*
 Drawing by H. L. Todd from No. 21585, U.S.N.M., collected by U. S. Fish Commission, Potomac River, Washington.
 335. *Eslopsaris*
 Drawing by H. L. Todd from No. 21585, U.S.N.M., collected by U. S. Fish Commission, Potomac River, Washington.
 336. *Kirklandia*
 Drawing by H. L. Todd from No. 21585, U.S.N.M., collected by U. S. Fish Commission, Potomac River, Washington.

Text page.

325. *Siphostoma starksii* Jordan & Culver..... 771
 Drawing by Anna L. Brown from the type, No. 2686, U. S. N. M.
 Univ. Mus., collected by the Hopkins Expedition in the Rio
 Presidio, Mazatlan, Mexico.
326. *Corythoichthys cayorum* Evermann and Kendall..... 2838
 Drawing by A. H. Baldwin from the type, No. 48784, U. S. N. M.,
 a male, collected by Evermann and Kendall at Key West,
 Florida.

PLATE CXXI.

327. *Hippocampus hudsonius* De Kay 777
 Drawing by H. L. Todd.
328. *Hippocampus zosteræ* Jordan and Gilbert 778
 Drawing by W. S. Atkinson from a specimen collected by Dr.
 Jordan at Pensacola, Florida.
329. *Percopsis guttatus* Agassiz..... 784
 Drawing by A. H. Baldwin from a specimen collected by Dr.
 Evermann in East Okoboji Lake, Iowa.

PLATE CXXII.

330. *Columbia transmontana* Eigenmann and Eigenmann 784
 Drawing by A. H. Baldwin from the type, collected at the
 mouth of the Umatilla River, Oregon, by Dr. Eigenmann.
331. *Aphredoderus sayanus* (Gilliams) 786
 Drawing from No. 34401, U. S. N. M., collected by William
 McAdams in the Illinois River.
332. *Atherina stipes* Müller and Troschel..... 790
 Drawing by A. H. Baldwin from No. 38456, U. S. N. M., col-
 lected by the *Albatross* in the Bahamas.

PLATE CXXIII.

333. *Atherina aræa* Jordan and Gilbert..... 790
 Drawing by W. S. Haines from the type, No. 34937, U. S. N. M.,
 collected by Dr. Jordan at Key West, Florida.
334. *Chiostoma humbolditanum* (Cuvier and Valenciennes) 793
 Drawing by A. H. Baldwin from No. 44156, U. S. N. M., col-
 lected by Seovell and Woolman in Mexico.
335. *Eslopsarum jordani* (Woolman) 793; 2840
 Drawing by A. H. Baldwin from the type, collected by A. J.
 Woolman in the canals at Salamanca, Mexico.

PLATE CXXIV.

336. *Kirklandia vagrans* (Goode and Bean) 794
 Drawing by H. L. Todd from No. 22864, U. S. N. M., collected
 by Silas Stearns at Pensacola, Florida.
 Bull. No. 47, pt. 4—xv

	Text page.
337. <i>Menidia peninsulae</i> (Goode and Bean).....	797
Drawing by H. L. Todd from No. 21481, U.S.N.M., collected at Pensacola, Florida., by Silas Stearns.	
338. <i>Menidia gracilis beryllina</i> (Cope).....	797
Drawing by A. H. Baldwin from a specimen collected by H. M. Smith in the Potomac River.	

PLATE CXXV.

339. <i>Eurystole eriarcha</i> (Jordan and Gilbert).....	803
Drawing by Anna L. Brown from No. 2689, L. S. Jr. Univ. Mus., collected by the Hopkins expedition at Mazatlan, Mexico.	
340. <i>Thyrima evermanni</i> Jordan and Culver	804
Drawing by Anna L. Brown from the type, No. 2688, L. S. Jr. Univ. Mus., collected by the Hopkins Expedition at Mazatlan, Mexico.	
341. <i>Atherinopsis californiensis</i> Girard	806
Drawing from No. 26764, U.S.N.M., collected by Dr. Jordan at San Diego, California.	

PLATE CXXVI.

342. <i>Atherinops affinis</i> (Ayres).....	807
Drawing by Anna L. Brown.	
343. <i>Mugil cephalus</i> Linnaeus	811
Drawing by H. L. Todd from No. 24456, U.S.N.M., collected at Woods Hole, Massachusetts, by V. N. Edwards.	
344. <i>Mugil curema</i> Cuvier and Valenciennes.....	813
Drawing by H. L. Todd.	

PLATE CXXVII.

345. <i>Mugil hospes</i> Jordan and Culver	811
Drawing by Anna L. Brown from the type, No. 2890, L. S. Jr. Univ. Mus., collected by the Hopkins Expedition at Mazatlan, Mexico.	
346. <i>Chænomugil proboscideus</i> (Günther).....	816
Drawing by A. H. Baldwin from No. 46563, U.S.N.M., collected by the <i>Albatross</i> at Clarion Island.	
347. <i>Agonostomus monticola</i> (Bancroft).....	819
Drawing by A. H. Baldwin from No. 45482, U.S.N.M., collected by E. W. Nelson in Mexico.	

PLATE CXXVIII.

348. <i>Joturus pichardi</i> Poey	821
Drawing by A. H. Baldwin from No. 31010, U.S.N.M., collected by Captain Dow at Panama.	

349. *Sphyraena*
Drawing
fig.
350. *Polydactylus*
Drawing
S...

351. *Anniellula*
Drawing
the
352. *Bathygobius*
Drawing
Bl...
353. *Stephanius*
Drawing
coll.
Str...

354. *Hoplostethus*
Drawing
lect...
355. *Plectrocanthus*
Drawing
coll.
W. L...
356. *Plectrocanthus*
Drawing
by t...
fath...

357. *Beryx*
Drawing
Albula
in 44...
358. *Elocentrus*
Drawing
Barb...
Drawing
Univ...
Jama...

360. *Mullus*
Drawing
361. *Mulloidichthys*
Drawing
coll...

- Text page.
349. *Sphyraena barracuda* (Walbaum)..... 823; 2841
 Drawing by H. L. Todd from No. 14978, U.S.N.M., received
 from E. G. Blackford, collected in Florida.
350. *Polydactylus octonemus* (Girard)..... 830
 Drawing by H. L. Todd from No. 22821, U.S.N.M., collected by
 Silas Stearns at Pensacola, Florida.

PLATE CXXIX.

351. *Ammodytes americanus* De Kay..... 833
 Drawing by H. L. Todd from No. 16500, U.S.N.M., collected by
 the U. S. Fish Commission at Nantucket.
352. *Bathyclupea argentea* Goode and Bean..... 835
 Drawing by A. H. Baldwin from the type collected by the
Blake, off Neris.
353. *Stephanoberyx monæ* Gill..... 836
 Drawing by H. L. Todd from the type, No. 33445, U.S.N.M.,
 collected by the *Albatross* at Station 2077, in the Gulf
 Stream.

PLATE CXXX.

354. *Hoplostethus mediterraneus* Cuvier and Valenciennes..... 837
 Drawing by A. H. Baldwin from No. 43624, U.S.N.M., col-
 lected by the *Albatross* at Station 2059.
355. *Plectromus suborbitalis* Gill 841
 Drawing by H. L. Todd from the type, No. 33271, U.S.N.M.,
 collected by the *Albatross* at Station 2036, in N. lat. $38^{\circ} 52' 40''$,
 W. long. $69^{\circ} 24' 40''$, in 1,735 fathoms.
356. *Plectromus crassiceps* Günther 843
 Drawing by S. F. Denton from No. 33378, U.S.N.M., collected
 by the *Albatross* in N. lat. $41^{\circ} 40' 30''$, W. long. $65^{\circ} 35'$, in 855
 fathoms.

PLATE CXXXI.

357. *Beryx splendens* Lowe 844
 Drawing by M. M. Smith from a specimen collected by the
Albatross at Station 2413, in N. lat. $30^{\circ} 44'$, W. long. $79^{\circ} 26'$,
 in 440 fathoms.
358. *Holocentrus ascensionis* (Osbeck)..... 848
 Drawing by H. L. Todd.
- Lammeo marianus* (Cuvier and Valenciennes)..... 852; 2871
 Drawing by W. S. Atkinson from specimen No. 4972, L. S. Jr.
 Univ. Mus., collected by Rev. J. S. Roberts at Kingston,
 Jamaica.

PLATE CXXXII.

360. *Mullus auratus* Jordan and Gilbert 856
 Drawing by A. H. Baldwin.
361. *Mulloides rathbuni* (Evermann and Jenkins) 857
 Drawing by A. H. Baldwin from the type, No. 43241, U.S.N.M.,
 collected by Evermann and Jenkins at Guaymas, Mexico.

	Text page
362. <i>Upeneus maculatus</i> (Bloch)	858
Drawing by H. L. Todd from No. 21910, U.S.N.M., collected by V. N. Edwards at Woods Hole, Massachusetts.	

PLATE CXXXIII.

363. <i>Scomber scombrus</i> Linnaeus	865
Drawing by H. L. Todd from No. 25256, U.S.N.M., obtained in the Washington market.	
364. <i>Scomber colias</i> Gmelin	866
Drawing by H. L. Todd from No. 23480, U.S.N.M., collected by the U. S. Fish Commission at Provincetown, Massachusetts.	
365. <i>Auxis thazard</i> (Lacépède)	867
Drawing by H. L. Todd from No. 35136, U.S.N.M., collected by the U. S. Fish Commission at Woods Hole, Massachusetts.	

PLATE CXXXIV.

366. <i>Gymnosarda alleterata</i> (Rafinesque)	869
Drawing by H. L. Todd from No. 10436, U.S.N.M., collected by S. F. Baird at Woods Hole, Massachusetts.	
367. <i>Gerimo alalunga</i> (Gmelin)	871
Drawing by H. L. Todd from No. 21844, U.S.N.M., collected by Capt. William Thompson of the schooner <i>Magic</i> , in 300 fathoms.	
368. <i>Scomberomorus maculatus</i> (Mitchill)	874
Drawing by H. L. Todd from No. 15582, U.S.N.M., obtained by E. G. Blackford in the New York market.	

PLATE CXXXV.

369. <i>Scomberomorus regalis</i> (Bloch)	875
Drawing by H. L. Todd from No. 12527, U.S.N.M., collected by E. G. Blackford at Key West.	
370. <i>Escolar violaceus</i> (Bean)	878; 2843
Drawing by S. F. Denton from the type, No. 39287, U.S.N.M., collected by Capt. Thomas Thompson on Le Have Bank in 125 fathoms.	
371. <i>Epinnula magistralis</i> Poey	880
Drawing by H. L. Todd from No. 37238, U.S.N.M., collected by the <i>Albatross</i> in the Caribbean Sea.	

PLATE CXXXVI.

372. <i>Evoxymetopon tæniatus</i> Poey	886
Drawing by H. L. Todd from No. 5735, U.S.N.M., collected by Professor Poey in Cuba.	
373. <i>Lepidopus caudatus</i> (Euphrasen)	886
Drawing by A. H. Baldwin from No. 10115, U.S.N.M., col- lected by John Xantus at Cape San Lucas, Lower California.	

374. Bent
Dra
ta
or
B375. Trich
Dra
by
376. Istiop
Dra
by377. Nem
:Draw378. Oligo
Draw379. Naucr
Draw
by V380. Seriola
Draw
and381. Seriola
Draw
by V382. Seriola
Drawin
by th383. Decapter
Drawin384. Trachur
Drawin
at Ne385. Trachur
Drawin
by V

Text page.

374. *Benthodesmus atlanticus* Goode and Bean 887
 Drawing by H. L. Todd from the type, No. 29116, U.S.N.M.,
 taken from the stomach of a halibut by Capt. R. Morrison,
 of the schooner *Laura Nelson*, on the west edge of the Grand
 Bank of Newfoundland, in 80 fathoms.

PLATE CXXXVII.

375. *Trichiurus lepturus* Linnaeus 889
 Drawing by H. L. Todd from No. 18028, U.S.N.M., collected
 by Dr. Janeway at St. Augustine, Florida.
 376. *Istiophorus nigricans* (Lacépède) 891
 Drawing by H. L. Todd from No. 37923, U.S.N.M., collected
 by U. S. Fish Commission at Woods Hole, Massachusetts.

PLATE CXXXVIII.

377. *Nematistius pectoralis* Gill 895
 Drawing by A. H. Baldwin.
 378. *Oligoplites saurus* (Bloch and Schneider) 898
 Drawing by H. L. Todd from No. 16354, U.S.N.M., collected
 by the U. S. Fish Commission at Memensha Bight.

PLATE CXXXIX.

379. *Nauprates ductor* Linnaeus 900
 Drawing by H. L. Todd from No. 23197, U.S.N.M., collected
 by William A. Bansett at New Bedford, Massachusetts.
 380. *Seriola dorsalis* (Gill). 902
 Drawing by H. L. Todd from a specimen collected by Jordan
 and Gilbert at San Pedro, California.
 381. *Seriola zonata* (Mitchill). 902
 Drawing by H. L. Todd from No. 18720, U.S.N.M., collected
 by V. N. Edwards at Woods Hole, Massachusetts.

PLATE CXL.

382. *Seriola lalandi* Cuvier and Valenciennes. 903
 Drawing by H. L. Todd from No. 37918, U.S.N.M., collected
 by the U. S. Fish Commission at Woods Hole, Massachusetts.
 383. *Decapterus macarellus* (Cuvier and Valenciennes). 909
 Drawing by H. L. Todd.
 384. *Trachurus trachurus* (Linnaeus). 910
 Drawing by H. L. Todd from No. 23421, U.S.N.M., collected
 at Newport, Rhode Island.

PLATE CXLI.

385. *Trachurops crumenophthalmus* (Bloch) 911
 Drawing by H. L. Todd from No. 20681, U.S.N.M., collected
 by V. N. Edwards, at Woods Hole, Massachusetts.

	Text page.
386. Hemicaranx amblyrhynchus (Cuvier and Valenciennes).....	912
Drawing by H. L. Todd from No. 30177, U.S.N.M., collected by Silas Stearns, at Pensacola, Florida.	
387. Caranx hippos (Linnaeus)	920
Drawing by H. L. Todd from No. 10431, U.S.N.M., collected by V. N. Edwards, at Woods Hole, Massachusetts.	
PLATE CXLII.	
388. Caranx latus Agassiz.....	921
Drawing by H. L. Todd from No. 20247, U.S.N.M., collected by V. N. Edwards, at Woods Hole, Massachusetts.	
389. Caranx cryos (Mitchill).....	923
Drawing by H. L. Todd from No. 16512, U.S.N.M., collected by the U.S. Fish Commission at Woods Hole, Massachu- sets.	
PLATE CXLIII.	
390. Caranx medusicola Jordan and Starks	924
Drawing by Anna L. Brown from the type, No. 2645, L. S. Jr. Univ. Mus., collected by the Hopkins Expedition at Mazatlan, Mexico.	
391. Hynnis hopkinsi Jordan and Starks.....	933
Drawing by Anna L. Brown from the type, No. 1563, L. S. Jr. Univ. Mus., collected by the Hopkins Expedition at Mazatlan, Mexico.	
PLATE CXLIV.	
392. Vomer setipinnis (Mitchill)	934
Drawing by H. L. Todd from No. 16252, U.S.N.M., obtained by E. G. Blackford in the Fulton Market, New York.	
393. Selene vomer (Linnaeus)	936
Drawing by H. L. Todd from No. 22007, U.S.N.M., a young example, collected by C. J. Copley at Stapleton, Staten Island.	
PLATE CXLV.	
393a. Selene vomer (Linnaeus).....	936
Drawing by H. L. Todd from No. 22279, U.S.N.M., an adult.	
394. Chloroscombrus chrysurus (Linnaeus)	938
Drawing by H. L. Todd from No. 21286, U.S.N.M., collected by Dr. Goode in the St. Johns River, Florida.	
PLATE CXLVI.	
395. Trachinotus glaucus (Bloch)	940
Drawing by H. L. Todd from No. 30176, U.S.N.M., collected by Silas Stearns at Pensacola, Florida.	
396. Trachinotus falcatus (Linnaeus)	941
Drawing by H. L. Todd from No. 26585, U.S.N.M., collected by Silas Stearns in Florida.	

397. **Trachinus** Dravida
Jr. Mar
398. **Trachinus** Draw
by
399. **Zaloces** Draw
Un
400. **Pomatomus** Draw
401. **Rachycentron** Draw
402. **Coryphaenoides** Draw
by
403. **Centrolophus** Draw
Mas
404. **Peprilus** Draw
405. **Poronotus** Draw
S. F.
406. **Icichthys** Draw
at P
407. **Schedophilus** Draw
Alba
408. **Acrotus** Draw
colle
west

PLATE CXLVII.

	Text page.
397. <i>Trachinotus culveri</i> Jordan and Starks.....	942
Drawing by Anna L. Brown from the type, No. 2691, L. S. Jr. Univ. Mus., collected by the Hopkins Expedition at Mazatlan, Mexico.	

398. <i>Trachinotus carolinus</i> (Linnaeus)	944
Drawing by H. L. Todd from No. 10427, U.S.N.M., collected by S. F. Baird at Woods Hole, Massachusetts.	

PLATE CXLVIII.

399. <i>Zalocys stilbe</i> Jordan and McGregor.....	2848
Drawing by Anna L. Brown from the type, No. 11996, L. S. Jr. Univ. Mus., collected by R. C. McGregor, at Clarion Island.	
400. <i>Pomatomus saltatrix</i> (Linnaeus)	946
Drawing by H. L. Todd, from a cast.	
401. <i>Rachycentron canadus</i> (Linnaeus)	948
Drawing by H. L. Todd from No. 18563, U.S.N.M.	

PLATE CXLIX.

402. <i>Coryphaena hippurus</i> Linnaeus.....	952
Drawing by H. L. Todd from No. 16482, U.S.N.M., obtained by E. G. Blackford in Fulton Market.	
403. <i>Centrolophus niger</i> (Gmelin).....	963
Drawing by S. F. Denton from a specimen obtained at Dennis, Massachusetts.	

PLATE CL.

404. <i>Peprilus paru</i> (Linnaeus)	965
Drawing by H. L. Todd from No. 15234, U.S.N.M.	
405. <i>Poronotus triacanthus</i> (Peck).....	967
Drawing by H. L. Todd from No. 787, U.S.N.M., collected by S. F. Baird at Beesleys Point, New Jersey.	

PLATE CLI.

406. <i>Icichthys lockingtoni</i> Jordan and Gilbert	969
Drawing by A. H. Baldwin from No. 27397, U.S.N.M., collected at Point Reyes, California.	
407. <i>Schedophilus medosophagus</i> Cocco.....	970
Drawing from H. L. Todd from a specimen collected by the Albatross in lat. 27° 49' N., 76° 12' W.	
408. <i>Acrotus willoughbyi</i> Bean.....	973
Drawing by H. L. Todd from the type, No. 39340, U.S.N.M., collected by Chas. Willoughby at the Quinault Agency, west coast of Washington.	

PLATE CLII.

	Text page.
409. <i>Zaprora silenus</i> Jordan.....	2850
Drawing by Anna L. Brown from the type, in the Provincial Museum at Victoria, British Columbia, collected by H. T. Stainton, at Nanaimo, Vancouver Island.	
410. <i>Grammicolepis brachiusculus</i> Poey.....	974
Drawing by A. H. Baldwin from a drawing by R. W. Shufeldt, Jour. Morph., Vol. II.	
411. <i>Tetragonurus cuvieri</i> Risso	976
Drawing by Mary Hildebrandt from a specimen collected by V. N. Edwards at Woods Hole.	

PLATE CLIII.

412. <i>Pempheris mulleri</i> Poey.....	978
Drawing by M. M. Smith from No. 37111, U.S.N.M., collected by the <i>Albatross</i> at Cozumel.	
413. <i>Pempheris poeyi</i> Bean	979
Drawing by M. M. Smith from the type, No. 37184, U.S.N.M., collected by Prof. Poey in Cuba.	
414. <i>Elassoma evergladei</i> Jordan	982
Drawing by A. H. Baldwin from a specimen collected by A. J. Woolman in Florida.	

PLATE CLIV.

415. <i>Pomoxis annularis</i> Rafinesque.....	987
Drawing by H. L. Todd from No. 10387, U.S.N.M.	
416. <i>Pomoxis sparoides</i> (Lacépède).....	987
Drawing by H. L. Todd from No. 10077, U.S.N.M.	

PLATE CLV.

417. <i>Centrarchus macropterus</i> (Lacépède).....	988
Drawing by H. L. Todd from No. 20397, U.S.N.M., collected by W. S. Hyatt at Kingston, North Carolina.	
418. <i>Acantharchus pomotis</i> (Baird).....	989
Drawing by H. L. Todd from No. 6475, U.S.N.M., collected at Tarboro, North Carolina.	

PLATE CLVI.

419. <i>Ambloplites rupestris</i> (Rafinesque)	990
Drawing by H. L. Todd from No. 9401, U.S.N.M., collected by George Clark at Ecorse, Michigan.	
419, A, B, C. Skull of Ambloplites rupestris (Rafinesque)	990
Drawings from Boulenger's Catalogue of Perciform Fishes, Vol. I, p. 3.	

420. Archon
 Drawn
 by
421. Chæno
 Drawn
 by
 Ala
422. Enmead
 Drawn
 by
423. Mesoge
 Drawn
 at T
424. Apomo
 Drawn
 Eve
 Tex
425. Lepomi
 Drawn
425a. Skull o
 Drawn
 Vol
426. Lepomi
 Drawn
 by Je
427. Lepomi
 Drawn
428. Eupom
 Drawn
 collec
429. Eupom
 Drawn
 by Pr
430. Micropte
 Drawn
 in the
430a. Vertebr
 From Ba

PLATE CLVII.

- | | Text page. |
|--|------------|
| 420. <i>Archoplites interruptus</i> (Girard)..... | 991 |
| Drawing by H. L. Todd from No. 27137, U.S.N.M., collected
by Dr. Jordan at San Francisco. | |
| 421. <i>Chænobryttus gulosus</i> (Cuvier and Valenciennes) | 992 |
| Drawing by H. L. Todd from No. 17803, U.S.N.M., collected
by Knmlien and Bean in Jackson Lake, near Montgomery,
Alabama. | |

PLATE CLVIII.

- | | |
|--|-----|
| 422. <i>Enneacanthus gloriosus</i> (Holbrook)..... | 993 |
| Drawing by H. L. Todd from No. 20356, U.S.N.M., collected
by C. C. Abbott at Trenton, New Jersey. | |
| 423. <i>Mesogonistius chaetodon</i> (Baird) | 995 |
| Drawing from No. 20354, U.S.N.M., collected by C. C. Abbott
at Trenton, New Jersey. | |

PLATE CLIX.

- | | |
|--|------|
| 424. <i>Apomotis symmetricus</i> (Forbes)..... | 998 |
| Drawing by A. H. Baldwin from a specimen collected by
Evermann, Seovell, and Gurley in Kilpers Pond, Houston,
Texas. | |
| 425. <i>Lepomis auritus</i> (Linnaeus) | 1001 |
| Drawing by H. L. Todd. | |
| 425a. <i>Skull of Lepomis auritus</i> (Linnaeus)..... | 1001 |
| Drawing from Boulenger's Catalogue of Perciform Fishes,
Vol. I, p. 24. | |

PLATE CLX.

- | | |
|---|------|
| 426. <i>Lepomis megalotis</i> (Rafinesque) | 1002 |
| Drawing by H. L. Todd from No. 36465, U.S.N.M., collected
by Jordan and Gilbert in Saline River, Benton, Arkansas. | |
| 427. <i>Lepomis pallidus</i> (Mitchill) | 1005 |
| Drawing by H. L. Todd from No. 8448, U.S.N.M. | |

PLATE CLXI.

- | | |
|--|------|
| 428. <i>Eupomotis euryorus</i> (McKay) | 1008 |
| Drawing by H. L. Todd from the type No. 4109, U.S.N.M.,
collected in Lake Huron, at Fort Gratiot, Michigan. | |
| 429. <i>Eupomotis gibbosus</i> (Linnaeus) | 1009 |
| Drawing by H. L. Todd from No. 4163, U.S.N.M., collected
by Professor Baird in Root River, Wisconsin. | |

PLATE CLXII.

- | | |
|---|------|
| 430. <i>Micropterus dolomieu</i> Lacépède | 1011 |
| Drawing by H. L. Todd from No. 14143, U.S.N.M., obtained
in the Washington market. | |
| 430a. <i>Vertebræ of Micropterus dolomieu</i> Lacépède | 1011 |
| From Boulenger's Catalogue Perciform Fishes, Vol. I, p. 4. | |

PLATE CLXIII.

- Text page.
- | | |
|--|------|
| 431. <i>Micropterus salmoides</i> (Lacépède) | 1012 |
| Drawing by H. L. Todd. | |
| 432. <i>Kuhlia rupestris</i> (Lacépède) | |
| Drawings from Boulenger's Catalogue of Perciform Fishes,
Vol. I, fig. 4, p. 35. | |

PLATE CLXIV.

- | | |
|--|------|
| 433. <i>Stizostedion vitreum</i> (Mitchill) | 1021 |
| Drawing by H. L. Todd from No. 10072, U.S.N.M., collected by
J. W. Milner at Rochester, New York. | |
| 433a. Tail of <i>Stizostedion vitreum</i> (Mitchill) | 1021 |
| Drawing by Anna L. Brown. | |
| 434. <i>Stizostedion canadense</i> (Smith) | 1022 |
| Drawing by H. L. Todd from No. 10555, U.S.N.M., collected
by J. W. Milner at Ecorse, Michigan. | |

PLATE CLXV.

- | | |
|---|------|
| 435. <i>Perca flavescens</i> (Mitchill) | 1023 |
| Drawing by H. L. Todd. | |
| 436. <i>Percina caprodes</i> (Rafinesque) | 1026 |
| Drawing by H. L. Todd from No. 34744, U.S.N.M., collected by
H. G. Dresel in Licking County reservoir, Ohio. | |
| 436a. Skull of <i>Percina caprodes</i> (Rafinesque) | 1026 |
| Drawing from Boulenger's Catalogue of Perciform Fishes,
Vol. I, fig. 7, p. 57. | |

PLATE CLXVI.

- | | |
|---|------|
| 437. <i>Hadropterus macrocephalus</i> (Cope) | 1031 |
| Drawing by A. H. Baldwin from a specimen collected by Dr.
Gurley in Indian Creek, near Cumberland Gap, Tennessee. | |
| 438. <i>Hadropterus aspro</i> (Cope and Jordan) | 1032 |
| Drawing by A. H. Baldwin from a specimen collected by
Evermann, Sevell, and Gurley in Chickamauga Creek, at
Lee and Gordon's mill, Georgia. | |

PLATE CLXVII.

- | | |
|---|------|
| 439. <i>Hadropterus gruntheri</i> Eigenmann and Eigenmann | 1033 |
| Drawing by A. H. Baldwin from a specimen collected by A.
J. Woolman in the Red River of the North, Moorehead,
Minnesota. | |
| 440. <i>Hadropterus evides</i> (Jordan and Copeland) | 1036 |
| Drawing by A. H. Baldwin from a specimen collected by
Evermann, Sevell, and Gurley in Clinch River, at Walkers
Ford, Tennessee. | |
| 441. <i>Hadropterus scierus</i> Swain | 1037 |
| Drawing by A. H. Baldwin from a specimen collected by
Evermann, Sevell, and Gurley in Clinch River, at Walkers
Ford, Tennessee. | |

442. Hypo
Drawing
Ev
Fe
443. Hypo
Drawing
444. Cottog
Drawing
col
Ne
445. Cottog
Drawing
col
Wa
446. Ulocen
Drawing
coll
Wa
447. Ulocen
Drawing
coll
Ten
448. Ulocen
Drawing
Eve
Ver
449. Diplesio
Drawing
Eve
For
450. Boleosco
Drawing
Gur
451. Boleoso
Drawing
lecte
Gur
452. Boleoso
Drawing
colle
453. Crystall
Drawing
lecte
mony

PLATE CLXVIII.

	Text page.
442. <i>Hypohomus aurantiacus</i> (Cope)	1040
Drawing by A. H. Baldwin from a specimen collected by Evermann, Sevall, and Gurley in Clinch River, at Walkers Ford, Tennessee.	
443. <i>Hypohomus spilotus</i> (Gilbert).....	1043
Drawing by A. H. Baldwin.	
444. <i>Cottogaster shumardi</i> (Girard).....	1046
Drawing by H. L. Todd from No. 17852, U.S.N.M., collected by Dr. Jordan in the Wabash River, Indiana.	

PLATE CLXIX.

445. <i>Cottogaster cheneyi</i> Evermann and Kendall.....	2851
Drawing by A. H. Baldwin from the type No. 48781, U.S.N.M., collected by Evermann and Bean in Racket River, Norfolk, New York.	
446. <i>Ulocentra giberti</i> Evermann and Thoburn.....	1049
Drawing by A. H. Baldwin from the type No. 47531, U.S.N.M., collected by Evermann, Sevall, and Gurley in Clinch River, Walkers Ford, Tennessee.	
447. <i>Ulocentra meadiae</i> Jordan and Evermann.....	2852
Drawing by A. H. Baldwin from the type No. 48903, U.S.N.M., collected by Dr. Gurley in Indian Creek, Cumberland Gap, Tennessee.	

PLATE CLXX.

448. <i>Ulocentra simotera</i> (Cope)	1051
Drawing by A. H. Baldwin from a specimen collected by Evermann, Sevall, and Gurley in Arnwine Sprung, Mount Verd, Tennessee.	
449. <i>Diplesion blennioides</i> (Rafinesque).....	1053
Drawing by A. H. Baldwin from a specimen collected by Evermann, Sevall, and Gurley in Clinch River, at Walkers Ford, Tennessee.	
450. <i>Boleosoma nigrum</i> (Rafinesque)	1056
Drawing by A. H. Baldwin from a specimen collected by R. R. Gurley in Ball Creek, near Tazewell, Tennessee.	

PLATE CLXXI.

451. <i>Boleosoma nigrum olmstedi</i> (Storer).....	1057
Drawing by H. L. Todd from No. 30243, U.S.N.M., collected by Col. Marshall McDonald in the Potomac River, Gaithersburg, Virginia.	
452. <i>Boleosoma camurum</i> Forbes	1060
Drawing by H. L. Todd from the type No. 23455, U.S.N.M., collected by Dr. Jordan in the Illinois River.	

453. *Crystallaria asprella* (Jordan)

 Drawing by A. H. Baldwin from No. 45455, U.S.N.M., collected by Dr. Evermann in the Wabash River, New Harmony, Indiana.

1061

PLATE CLXXII.

- Text page.
454. *Ammocrypta pellucida clara* (Jordan and Meek) 1063
 Drawing by S. F. Denton from the type No. 35828, U.S.N.M., collected by Jordan and Meek in the Des Moines River, Ottumwa, Iowa.
455. *Ammocrypta beanti* Jordan 1064
 Drawing by A. H. Baldwin from No. 17833, U.S.N.M., collected by Bean and Maxson in Notalbany River, Louisiana.
456. *Etheostoma camurum* (Cope) 1076
 Drawing by A. H. Baldwin from a specimen collected by R. R. Gurley in Indian Creek, near Cumberland Gap, Tennessee.

PLATE CLXXIII.

457. *Etheostoma cinereum* Storer 1078
 Drawing by A. H. Baldwin from a specimen collected by P. H. Kirsch in Wolf River, Byrdstown, Tennessee.
458. *Etheostoma jordani* Gilbert 1079
 Drawing by S. F. Denton.
459. *Etheostoma pottsii* (Girard) 1082
 Drawing by H. L. Todd from No. 38245, U.S.N.M., the type of *Etheostoma micropterus* Gilbert, collected by E. Wilkinson in Chihuahua, Mexico.

PLATE CLXXIV.

460. *Etheostoma iowae* Jordan and Meek 1083
 Drawing by A. H. Baldwin from a specimen collected by A. J. Woolman in Pomme de Terre River, Appleton, Wisconsin.
461. *Etheostoma jessiae* (Jordan and Brayton) 1084
 Drawing by H. L. Todd from No. 27896, U.S.N.M., collected by Dr. Forbes, in Lake Peoria, Illinois.
462. *Etheostoma lepidogenys* Evermann and Kendall 1087
 Drawing by A. H. Baldwin from the type, No. 44840, U.S.N.M., collected by Evermann, Seovell, and Gurley, in Comal Spring, New Braunfels, Texas.

PLATE CLXXV.

463. *Etheostoma obeyense* Kirsch 1082
 Drawing by A. H. Baldwin from the type, No. 45565, U.S.N.M., collected by Dr. Kirsch, in tributary of Obeys River, Clinton County, Kentucky.
464. *Etheostoma pagei* Meek 1092
 Drawing by A. H. Baldwin from the type, No. 45556, U.S.N.M., collected by Dr. Meek in Spring Branch at Neosho, Missouri.
465. *Etheostoma virgatum* (Jordan) 1093
 Drawing by A. H. Baldwin from No. 36644, U.S.N.M., collected by Gilbert and Swain in Rock Castle River, Livingston, Kentucky.

466. *Etheostoma* Drawn
 467. *Psychrolutes* Drawn
 468. *Copelatus* Drawn
469. *Boleichthys* Drawn
 470. *Micropogonias* Drawn
 471. *Apogonichthys* Drawn

472. *Apogonichthys* Drawn
 473. *Apogonichthys* Drawn
 by J.

474. *Cheilodipterus* Drawn
 by Pr.
 475. *Hypoclymene* Drawn
 by th.
 long.

476. *Centropogon* Drawn

477. *Roccus* Drawn
 478. *Roccus* Drawn
 by Pr.

PLATE CLXXVI.

	Text page.
466. <i>Etheostoma juliae</i> Meek.....	1093 Drawing by S. F. Denton from the type collected by Meek, Drew, and Rettger in James River, near Springfield, Missouri.
467. <i>Psychromaster tuscumbia</i> (Gilbert and Swain)	1100 Drawing by A. H. Baldwin from the type, No. 36154, U.S.N.M., collected by Gilbert and Swain in Spring Creek, Tuscum- bia, Alabama.
468. <i>Copelandellus quiescens</i> (Jordan)	1100 Drawing by A. H. Baldwin.

PLATE CLXXVII.

469. <i>Boleichthys fusiformis</i> (Girard).....	1101
Drawing by H. L. Todd from No. 36415, U.S.N.M., collected by Jordan and Gilbert in Washita River, Arkadelphia, Ar- kansas.	
470. <i>Micropogonias fonticola</i> (Jordan and Gilbert)	1104
Drawing by Ernest Copeland from the type, No. 36523, U.S.N.M., collected by Jordan and Gilbert in San Marcos River, San Marcos, Texas.	
471. <i>Apogon retrosella</i> (Gill)	1108
Drawing by Anna L. Brown from No. 2917, L. S. Jr. Univ. Mus., collected by the Hopkins expedition at Mazatlan, Mexico.	

PLATE CLXXVIII.

472. <i>Apogon pigmentarius</i> (Poey)	1109
473. <i>Apogonichthys alatus</i> (Jordan & Gilbert).....	1110

 Drawing by A. H. Baldwin from No. 30874, U.S.N.M., collected
 by Jordan and Stearns at Pensacola, Florida.

PLATE CLXXIX.

474. <i>Cheilodipterus affinis</i> Poey.....	1113
Drawing by A. H. Baldwin from No. 37416, U.S.N.M., collected by Professor Poey in Cuba.	
475. <i>Hypoclydonia bella</i> Goode and Bean	1115
Drawing by S. F. Denton from No. 39338, U.S.N.M., collected by the <i>Albatross</i> at Station 2426, in N. lat. $36^{\circ} 01' 30''$, W. long. $74^{\circ} 47' 30''$, in 93 fathoms.	
476. <i>Centropomus undecimalis</i> (Bloch).....	1118
Drawing by H. L. Todd from No. 19907, U.S.N.M.	

PLATE CLXXX.

477. <i>Roccus chrysops</i> (Rafinesque)	1132
Drawing by H. L. Todd from No. 10326, U.S.N.M., collected by J. W. Milner at Sandusky, Ohio.	
478. <i>Roccus lineatus</i> (Bloch).....	1132
Drawing by H. L. Todd from No. 25219, U.S.N.M., obtained by Professor Baird in the Washington market.	

PLATE CLXXXI.

- Text page.
479. *Moro americana* (Gmelin)..... 1134
 Drawing by H. L. Todd from No. 15861, U.S.N.M., received
 from the New York market.
- 480, 480a. *Polyprion americanus* (Bloch & Schneider)..... 1139
 Drawings from Boulenger's Catalogue of Perciform Fishes
 Vol. I, fig. 12, p. 149.

PLATE CLXXXII.

481. *Bodianus fulvus punctatus* (Linnaeus) 1146
 Drawing by A. H. Baldwin from No. 33717, U.S.N.M., collected
 by J. C. Brevoort in the West Indies.
482. *Epinephelus adscensionis* (Osbeck)..... 1152
 Drawing by H. L. Todd from No. 26574, U.S.N.M., collected
 by Silas Stearns, at Key West, Florida.

PLATE CLXXXIII.

483. *Epinephelus striatus* (Bloch)..... 1157
 Drawing by H. L. Todd from No. 31910, U.S.N.M., collected
 by Silas Stearns at Pensacola, Florida.
484. *Epinephelus drummond-hayi* Goode and Bean 1159
 Drawing by H. L. Todd from No. 31719, U.S.N.M., collected
 by Silas Stearns at Pensacola, Florida.

PLATE CLXXXIV.

485. *Epinephelus morio* (Cuvier and Valenciennes) 1160
 Drawing by H. L. Todd from No. 22129, U.S.N.M., collected
 in Florida and obtained in the Washington market by Dr.
 Goode.
486. *Garrupa nigrita* (Holbrook)..... 1161
 Drawing by H. L. Todd from No. 37207, U.S.N.M., collected
 by the *Albatross*.

PLATE CLXXXV.

487. *Promicrops itaiara* (Lichtenstein) 1162
 Drawing by H. L. Todd from No. 22306, U.S.N.M., collected
 at mouth of St. Johns River, Florida.
- 487a, 487b. *Promicrops itaiara* (Lichtenstein)..... 1162
 Drawings from Boulenger's Catalogue of Perciform Fishes,
 Vol. I, fig. 13, p. 164.

PLATE CLXXXVI.

488. *Alphestes chloropterus* (Cuvier and Valenciennes)..... 1164
 Drawing by A. H. Baldwin from No. 9821, U.S.N.M., collected
 by Professor Poey in Cuba.
- 488a. *Alphestes chloropterus* (Cuvier and Valenciennes)..... 1164
 Drawing from Boulenger's Catalogue of Perciform Fishes,
 Vol. I, fig. 18, p. 254.

489. Dern
Dra
e490. Myct
Dra
U
la491. Myct
Dra
by492. Skull
Dra
Ve493. Myct
Dra
U
Ev494. Myct
Dra
by495. Myct
Dra
a*496. Hypop
Draw
Ga497. Hypop
Draw
coll498. Paralal
Draw
lect499. Skull c
Draw
Vol500. Centrop
Draw
by t

Text page.

489. *Dermatolepis zanclus* Evermann and Kendall 2854
 Drawing by A. H. Baldwin from the type, No. 48843, U.S.N.M.,
 collected by Evermann and Kendall at Key West, Florida.

PLATE CLXXXVII.

490. *Mycteroperca boulengeri* Jordan and Starks 1171
 Drawing by Anna L. Brown from the type, No. 1691, L. S. Jr.
 Univ. Mus., collected by the Hopkins expedition to Mazat-
 lan, Mexico.
491. *Mycteroperca venenosa* (Linnaeus) 1172
 Drawing by H. L. Todd from No. 35103, U.S.N.M., collected
 by Dr. Jordan at Havana, Cuba.
492. Skull of *Mycteroperca bonaci* (Poey) 1174
 Drawing from Boulenger's Catalogue of Perciform Fishes,
 Vol. I, fig. 19, p. 258.

PLATE CLXXXVIII.

493. *Mycteroperca jordani* (Jenkins and Evermann) 1176
 Drawing by A. H. Baldwin from the type, No. 39628,
 U.S.N.M., collected at Guaymas, Mexico, by Jenkins and
 Evermann.
494. *Mycteroperca microlepis* (Goode and Bean) 1177
 Drawing by H. L. Todd from No. 26587, U.S.N.M., collected
 by Silas Stearns at Key West, Florida.
495. *Mycteroperca falcata phenax* Jordan and Swain 1185
 Drawing by H. L. Todd from No. 34992, U.S.N.M., collected
 a^t Key West, Florida, by Dr. Jordan.

PLATE CLXXXIX.

496. *Hypoplectrus unicolor nigricans* (Poey) 1193
 Drawing by H. L. Todd from No. 3423, U.S.N.M., collected at
 Garden Key, Florida.
497. *Hypoplectrus gemma* Goode and Bean 1193
 Drawing by H. L. Todd from the type, No. 3422, U.S.N.M.,
 collected at Garden Key, Florida.

PLATE CXC.

498. *Paralabrax maculatusfasciatus* (Steindachner) 1196
 Drawing by A. H. Baldwin from No. 34754, U.S.N.M., col-
 lected by Rosa Smith at San Diego, California.
499. Skull of *Paralabrax humeralis* (Cuvier and Valenciennes) ... 1196
 Drawing from Boulenger's Catalogue of Perciform Fishes,
 Vol. I, fig. 20, p. 275.
500. *Centropristes striatus* (Linnaeus) 1199
 Drawing by H. L. Todd from No. 14838, U.S.N.M., collected
 by the U. S. Fish Commission at Noank, Connecticut.

PLATE CXCI.

- | | Text page. |
|--|------------|
| 501. <i>Centropristes philadelphicus</i> (Linnaeus)..... | 1201 |
| Drawing by A. H. Baldwin from No. 33161, U.S.N.M., collected by Dr. Gilbert at Charleston, South Carolina. | |
| 502. <i>Diplectrum formosum</i> (Linnaeus) | 1205 |
| Drawing by H. L. Todd from No. 21543, U.S.N.M., collected by C. C. Leslie at Charleston, South Carolina. | |
| 503. <i>Prionodes bulleri</i> (Boulenger)..... | 1213 |
| Drawing from Boulenger's Catalogue of Perciform Fishes, Vol. I, pl. x. | |

PLATE CXII.

- | | |
|--|------|
| 504. <i>Paranthias furcifer</i> (Cuvier and Valenciennes) | 1221 |
| Drawing by A. H. Baldwin from No. 12540, U.S.N.M., collected by Professor Poey in Cuba. | |
| 505. <i>Hemianthias vivanus</i> (Jordan and Swain)..... | 1223 |
| Drawing by W. S. Atkinson from specimen No. 1655, L. S. Jr. Univ. Mus., collected by Dr. Jordan at Pensacola, Florida. | |
| 506. <i>Pronotogrammus multifasciatus</i> Gill..... | 1226 |
| Drawing by H. L. Todd from the type, No. 2762, U.S.N.M., collected by John Xantus at Cape San Lucas, Lower California. | |

PLATE CXIII.

- | | |
|--|------|
| 507. <i>Anthias asperilinguis</i> Günther..... | 1227 |
| Drawing from Boulenger's Catalogue of Perciform Fishes, Vol. I, pl. XIII. | |
| 508. <i>Gramma loreto</i> Poey | 1229 |
| Drawing by J. W. Folsom from the type, No. 10031, M.C.Z., collected by Professor Poey at Matanzas, Cuba. | |

PLATE CXCIV.

- | | |
|---|------|
| 509. <i>Rypticus bistrispinus</i> (Mitchill)..... | 1233 |
| Drawing by H. L. Todd. | |
| 510. <i>Lobotes surinamensis</i> (Bloch)..... | 1235 |
| Drawing by H. L. Todd. | |

PLATE CXCV.

- | | |
|--|------|
| 511. <i>Priacanthus arenatus</i> Cuvier and Valenciennes | 1237 |
| Drawing by H. L. Todd from No. 20680, U.S.N.M., collected by V. N. Edwards at Woods Hole, Massachusetts. | |
| 512. <i>Pseudopriacanthus altus</i> Gill | 1239 |
| Drawing by H. L. Todd from No. 2317, U.S.N.M., collected by the Albatross off Key West, Florida. | |

- | | |
|---|------|
| 513. <i>Hoplostethus mediterraneus</i> (Linnaeus) | 1241 |
| Drawing by H. L. Todd. | |
| 514. <i>Evoplosurus maculatus</i> (Günther) | 1245 |
| Drawing by H. L. Todd. | |
| 515. <i>Neomyxus maculatus</i> (Günther) | 1249 |
| Drawing by H. L. Todd. | |
| 516. <i>Neomyxus maculatus</i> (Günther) | 1253 |
| Drawing by H. L. Todd. | |
| 517. <i>Neomyxus maculatus</i> (Günther) | 1257 |
| Drawing by H. L. Todd. | |
| 518. <i>Neomyxus maculatus</i> (Günther) | 1261 |
| Drawing by H. L. Todd. | |

- | | |
|--|------|
| 519. <i>Rabibus maculatus</i> (Günther) | 1265 |
| Drawing by H. L. Todd. | |
| 520. <i>Ocyurus maculatus</i> (Günther) | 1269 |
| Drawing by H. L. Todd. | |
| 521. <i>Rhombopteryx maculatus</i> (Günther) | 1273 |
| Drawing by H. L. Todd. | |
| 522. <i>Apsilus maculatus</i> (Günther) | 1277 |
| Drawing by H. L. Todd. | |
| 523. <i>Aprion maculatus</i> (Günther) | 1281 |
| Drawing by H. L. Todd. | |
| 524. <i>Etellis maculatus</i> (Günther) | 1285 |
| Drawing by H. L. Todd. | |

PLATE CXCVI.

Text page.

513. *Hoplopagrus guntheri* Gill..... 1244
 Drawing by A. H. Baldwin from No. 28367, U.S.N.M., collected
 by Dr. Gilbert at Mazatlan, Mexico.
 514. *Evoplites viridis* (Valenciennes) 1246
 Drawing by W. S. Atkinson from a specimen collected by
 Alfonso Forrer at Tres Marias Islands, Mexico.

PLATE CXCVII.

515. *Neomænis apodus* (Walbaum) 1258
 Drawing by H. L. Todd from No. 33220, U.S.N.M., collected
 by Dr. Henshall at Indian River Inlet, Florida.
 516. *Neomænus aya* (Bloch) 1264
 Drawing by A. H. Baldwin from a specimen obtained in the
 Washington Market.

PLATE CXCVIII.

517. *Neomænis analis* (Cuvier and Valenciennes)..... 1265
 Drawing by H. L. Todd from No. 35030, U.S.N.M.
 518. *Neomænis synagris* (Linnæus)..... 1266
 Drawing by H. L. Todd from No. 36922, U.S.N.M., collected
 by the U. S. Fish Commission at Key West, Florida.

PLATE CXCIX.

519. *Rabirubia inermis* (Peters) 1274
 Drawing by Anna L. Brown from No. 4983, L. S. Jr. Univ.
 Mus., collected by the *Albatross* at Panama.
 520. *Ocyurus chrysurus* (Bloch) 1275
 Drawing by S. F. Denton from No. 26583, U.S.N.M., collected
 by Silas Stearns at Key West, Florida.

PLATE CC.

521. *Rhomboplites aurorubens* (Cuvier and Valenciennes) 1277
 Drawing by H. L. Todd from No. 21338, U.S.N.M., collected
 by Silas Stearns at Pensacola, Florida.
 522. *Apsilus dentatus* Guichenot 1278
 Drawing by A. H. Baldwin from No. 9809, U.S.N.M., collected
 by Professor Poey in Cuba.

PLATE CCI.

523. *Aprion macrophthalmus* (Miller and Troschel)..... 1280
 Drawing by M. M. Smith from a specimen collected by the
Blake in N. lat. $23^{\circ} 13'$, W. long. $89^{\circ} 10'$, in 84 fathoms.
 524. *Etelis oculatus* (Cuvier and Valenciennes) 1282
 Drawing by A. H. Baldwin from No. 35049, U.S.N.M., collected
 by Dr. Jordan at Havana, Cuba.
 Bull. No. 47, pt. 4—xvi

PLATE CCII.

- Text page.
525. *Verilus sordidus* Poey 1284
 Drawing by A. H. Baldwin from No. 12565, U.S.N.M., collected
 by Professor Poey in Cuba.
526. *Xenocys jessiae* Jordan and Bollman 1285
 Drawing by A. H. Baldwin from the type, No. 41166, U.S.N.M.,
 collected by the *Albatross* at Charles Island, Galapagos
 Group.

PLATE CCIII.

527. *Xenichthys agassizii* Steindachner 1287
 Drawing by A. H. Baldwin from No. 26671, U.S.N.M., collected
 in the Galapagos Islands.
528. *Hæmulon album* Cuvier and Valenciennes 1295
 Drawing by H. L. Todd from No. 35035, U.S.N.M., collected at
 Key West, Florida, by Dr. Jordan.

PLATE CCIV.

529. *Hæmulon macrostomum* Günther 1296
 Drawing by H. L. Todd from No. 30057, U.S.N.M., received
 from the Kingston Public Museum, Jamaica.
530. *Hæmulon parra* (Desmarest) 1297
 Drawing by H. L. Todd.

PLATE CCV.

531. *Hæmulon sciurus* (Shaw) 1303
 Drawing by H. L. Todd from No. 32603, U.S.N.M., collected
 at Key West, Florida, by J. A. Henshall.
532. *Hæmulon plumieri* (Lacépède) 1301
 Drawing by H. L. Todd from No. 20980, U.S.N.M., collected
 by Dr. Goode at Charleston, South Carolina.

PLATE CCVI.

533. *Brachygenys chrysargyreus* (Günther) 1307
 Drawing by H. L. Todd from No. 35150, U.S.N.M., collected at
 Havana, Cuba, by Dr. Jordan.
534. *Bathystoma rimator* (Jordau and Swain) 1308
 Drawing by H. L. Todd from No. 34957, U.S.N.M., collected
 by the *Albatross* at St. Thomas, West Indies.

PLATE CCVII.

535. *Bathystoma aurolineatum* (Cuvier and Valenciennes) 1310
 Drawing by H. L. Todd from No. 26567, U.S.N.M., collected
 by Silas Stearns at Key West, Florida.
536. *Lythrylon opalescens* Jordan and Starks 1312
 Drawing by Anna L. Brown from the type, No. 2963, L. S. Jr.
 Univ. Mus., collected by the Hopkins Expedition at Mazat-
 lan, Mexico.

537. *Aniso-*
Drav.

Ev.

538. *Aniso-*
Draw.

Dr.

539. *Aniso-*
Draw.

by

540. *Orthop-*
Draw.

Uni-

bor-

541. *Orthop-*
Draw.

Pen-

542. *Microle-*
Drawin-

lect-

543. *Otrynte-*
Drawin-

by S-

544. *Stenoto-*
Drawin-

by P-

545. *Stenoto-*
Drawin-

by R-

546. *Calamus*
Drawin-

by D-

547. *Calamus*
Drawin-

by Dr-

548. *Calamus*
Drawin-

by Dr-

PLATE CCVIII.

- Text page.
537. *Anisotremus surinamensis* (Bloch) 1318
 Drawing by A. H. Baldwin from a specimen collected by Evermann and Bean in Indian River at Fort Pierce, Florida.
538. *Anisotremus bilineatus* (Cuvier and Valenciennes) 1319
 Drawing by H. L. Todd from No. 30878, U.S.N.M., collected by Dr. J. A. Henshall in Indian River Inlet, Florida.

PLATE CCIX.

539. *Anisotremus virginicus* (Linnaeus) 1322
 Drawing by H. L. Todd from No. 33189, U.S.N.M., collected by Dr. J. A. Henshall at Key West, Florida.
540. *Orthopristis reddingi* Jordan and Richardson 1336
 Drawing by Anna L. Brown from the type, No. 3458, L. S. Jr. Univ. Mus., collected by J. A. Richardson in La Paz Harbor, Lower California.

PLATE CCX.

541. *Orthopristis chrysopterus* (Linnaeus) 1338
 Drawing by H. L. Todd from No. 3113, U.S.N.M., collected at Pensacola, Florida.
542. *Microlepidotus inornatus* Gill 1341
 Drawing by A. H. Baldwin, from No. 43267, U.S.N.M., collected by Evermann and Jenkins at Guaymas, Mexico.

PLATE CCXI.

543. *Otrynter caprinus* (Bean) 1345
 Drawing by H. L. Todd from No. 32683, U.S.N.M., collected by Silas Stearns at Pensacola, Florida.
544. *Stenotomus chrysops* (Linnaeus) 1346
 Drawing by H. L. Todd from No. 10425, U.S.N.M., collected by Professor Baird at Woods Hole, Massachusetts.

PLATE CCXII.

545. *Stenotomus aculeatus* (Cuvier and Valenciennes) 1346
 Drawing by H. L. Todd from No. 24694, U.S.N.M., collected by R. E. Earll near Charleston, South Carolina.
546. *Calamus calamus* (Cuvier and Valenciennes) 1349
 Drawing by H. L. Todd from No. 35040, U.S.N.M., collected by Dr. Jordan at Key West, Florida.

PLATE CCXIII.

547. *Calamus proridens* Jordan and Gilbert 1350
 Drawing by H. L. Todd from No. 35056, U.S.N.M., collected by Dr. Jordan at Key West, Florida.
548. *Calamus bajonado* (Bloch and Schneider) 1352
 Drawing by H. L. Todd from No. 35042, U.S.N.M., collected by Dr. Jordan at Key West, Florida.

PLATE CCXIV.

- Text page.
549. **Calamus pennia** (Cuvier and Valenciennes) 1354
 Drawing by H. L. Todd from No. 6134, U.S.N.M., collected
 in Charlotte Harbor, Florida.
550. **Calamus arctifrons** Goode and Bean 1355
 Drawing by H. L. Todd from the type, No. 30163, U.S.N.M.,
 collected by Silas Stearns at Pensacola, Florida.

PLATE CCXV.

551. **Pagrus pagrus** (Linnaeus) 1356
 Drawing by H. L. Todd from a specimen collected by Silas
 Stearns at Pensacola, Florida.
552. **Lagodon rhomboides** (Linnaeus) 1358
 Drawing by H. L. Todd from No. 21280, U.S.N.M., collected
 by Dr. Goode in St. Johns River, Florida.

PLATE CCXVI.

553. **Archosargus unimaculatus** (Bloch) 1359
 Drawing by H. L. Todd from No. 35031, U.S.N.M., collected
 by Dr. Jordan at Key West, Florida.
554. **Archosargus probatocephalus** (Walbaum) 1361
 Drawing by H. L. Todd from No. 19647, U.S.N.M., collected
 by Dr. H. C. Yarrow at Beaufort, North Carolina.

PLATE CCXVII.

555. **Diplodus holbrookii** (Bean) 1362
 Drawing by H. L. Todd from No. 32632, U.S.N.M., collected
 at Pensacola, Florida.
- 555a. **Diplodus holbrookii** (Bean) 1362
 Drawing by H. L. Todd from a young example, No. 4693,
 U.S.N.M.

PLATE CCXVIII.

556. **Xystoma cinereum** (Walbaum) 1372
 Drawing by H. L. Todd from No. 35054, U.S.N.M., collected
 by Dr. Jordan at Key West, Florida.
557. **Gerres olistostomus** Goode and Bean 1376
 Drawing by H. L. Todd from the type, No. 25118, U.S.N.M.,
 collected by R. E. Earll in Indian River, Florida.

PLATE CCXIX.

558. **Hermosilla azurea** Jenkins and Evermann 1383
 Drawing by A. H. Baldwin from the type, No. 36269, U.S.N.M.,
 collected by Jenkins and Evermann at Guaymas, Mexico.
559. **Kyphosus sectatrix** (Linnaeus) 1387
 Drawing by H. L. Todd from No. 20635, U.S.N.M., collected
 at Woods Hole, Massachusetts.

560. **Meda**
 Drawing
561. **Cynoscion**
 Drawing
562. **Cynoscion**
 by

563. **Cynoscion**
 Drawing

564. **Sagenarius**
 Drawing

565. **Larimus**
 Drawing

- Museum

566. **Bairdiella**
 Drawing

567. **Sciæno**
 Drawing

- Dr. J.

568. **Roncador**
 Drawing

- Dr. J.

569. **Leiostomus**
 Drawing

- Sauvage

570. **Micropogonias**
 Drawing

- by Sa

571. **Umbrina**
 Drawing

- Jr. U.
 Mazatlan

572. **Menticirrhus**
 Drawing

- by Sila

573. **Pogonias**
 Drawing

- J. C. W.

PLATE CCXX.

	Text page.
560. <i>Medialuma californiensis</i> (Steindachner)	1391
Drawing by H. L. Todd.	
561. <i>Cynoscion nothus</i> (Holbrook)	1406
Drawing by H. L. Todd from No. 34921, U.S.N.M., collected by the <i>Albatross</i> at Trinidad, West Indies.	
562. <i>Cynoscion regalis</i> (Bloch and Schneider).....	1407

PLATE CCXXI.

563. <i>Cynoscion nebulosus</i> (Cuvier and Valenciennes)	1409
Drawing by H. L. Todd.	
564. <i>Sagenichthys aencyclodon</i> (Bloch and Schneider).....	1416
Drawing by A. H. Baldwin from No. 29722, U.S.N.M., collected by Dr. Gilbert at Panama.	
565. <i>Larimus argenteus</i> (Gill).....	1421
Drawing by Anna L. Brown from a specimen in L. S. Jr. Univ. Mus.	

PLATE CCXXII.

566. <i>Bairdiella chrysura</i> (Lacépède)	1433
Drawing by H. L. Todd.	
567. <i>Sciænops ocellatus</i> (Linnaeus)	1453
Drawing by H. L. Todd from No. 622, U.S.N.M., collected by Dr. Kennerly at Indianola, Texas.	

PLATE CCXXIII.

568. <i>Roncador stearnsi</i> (Steindachner).....	1557
Drawing by H. L. Todd from No. 26864, U.S.N.M., collected by Dr. Jordan at Santa Barbara, California.	
569. <i>Leiostomus xanthurus</i> Lacépède	1458
Drawing by H. L. Todd from No. 20222, U.S.N.M., collected by Samuel Powell at Newport, Rhode Island.	

PLATE CCXXIV.

570. <i>Micropogon undulatus</i> (Linneus)	1461
Drawing by H. L. Todd from No. 20742, U.S.N.M., collected by Samuel Powell at Newport, Rhode Island.	
571. <i>Umbrina sinaloæ</i> Scofield	1468
Drawing by Anna I. Brown from the type, No. 1632, L. S. Jr. Univ. Mus., collected by the Hopkins Expedition at Mazatlan, Mexico.	

PLATE CCXXV.

572. <i>Menticirrhus americanus</i> (Linneus).....	1474
Drawing by H. L. Todd from No. 22832, U.S.N.M., collected by Silas Stearns at Pensacola, Florida.	
573. <i>Pogonias chromis</i> (Linneus)	1482
Drawing by H. L. Todd from No. 18036, U.S.N.M., collected by J. C. Willetts at Matanzas River Inlet, Florida.	

PLATE CCXXVI.

- Text page.
574. *Aplidonotus grunniens* Rafinesque 1484
 Drawing by H. L. Todd from No. 10542, U.S.N.M., collected
 by J. W. Milner at Ecorse, Michigan.
575. *Eques lanceolatus* (Linnaeus) 1489
 Drawing by H. L. Todd from No. 32097, U.S.N.M., collected
 by Professor Poey in Cuba.

PLATE CCXXVII.

576. *Cirrhitus rivulatus* Valenciennes 1491
 Drawing by A. H. Baldwin from a specimen collected by the
Albatross in the Pacific.
577. *Hysteroxarpus traski* Gibbons 1496
 Drawing by J. C. Van Hook.

PLATE CCXXVIII.

578. *Abeona minima* (Gibbons) 1497
 Drawing by A. H. Baldwin from No. 26913, U.S.N.M., col-
 lected by Dr. Jordan at Santa Barbara, California.
579. *Cymatogaster aggregatus* Gibbons 1498
 Drawing by M. M. Smith from No. 31971, U.S.N.M., a male,
 collected at Fort Wrangell, Alaska, by Capt. H. E. Nichols.
- 579a. *Cymatogaster aggregatus* Gibbons 1498
 Drawing by W. S. Atkinson from a specimen in L. S. Jr.
 Univ. Mus.

PLATE CCXXIX.

580. *Brachyistius frenatus* Gill 1499
 Drawing by A. H. Baldwin from No. 26990, U.S.N.M., col-
 lected by Dr. Jordan in Monterey Bay, California.
581. *Zalembius rosaceus* (Jordan and Gilbert) 1500
 Drawing by W. S. Atkinson from a specimen in L. S. Jr.
 Univ. Mus.

PLATE CCXXX.

582. *Hypocritichthys anal'* (A. Agassiz) 1500
 Drawing by A. H. Baldwin from No. 27075, U.S.N.M., col-
 lected by Dr. Jordan in Monterey Bay, California.
583. *Phanerodon furcatus* Girard 1506
 Drawing by A. H. Baldwin from No. 24905, U.S.N.M., col-
 lected by Dr. Jordan at San Diego, California.

PLATE CCXXXI.

584. *Rhacochilus toxotes* Agassiz 1507
 Drawing from No. 27015, U.S.N.M., collected by Dr. Jordan
 at Monterey, California.
585. *Hypsurus caryi* (Agassiz) 1508
 Drawing by W. S. Atkinson from a specimen in L. S. Jr.
 Univ. Mus.

586. *Dama*
Draw
by587. *Cichla*
Draw
col
Lu588. *Azurin*
Draw
McD589. *Eupom*
Draw
Mus
Mex590. *Abudef*
Draw
leete591. *Hypsop*
Draw
lecte592. *Microspa*
Drawin
Mus.,
Mexic593. *Microspa*
Drawin
lected594. *Microspa*
Drawing
Mus.,
by the
Mexic595. *Tautogola*
Drawing
by V.

PLATE CCXXXII.

	Text page.
586. <i>Damalichthys argyrosomus</i> (Girard).....	1509
Drawing by H. L. Todd from No. 29811, U.S.N.M., collected by Capt. H. E. Nichols in Friendly Cove, British Columbia.	
587. <i>Cichlasoma bartoni</i> (Bean).....	1515
Drawing by A. H. Baldwin from the type, No. 43765, U.S.N.M., collected by Professor Dugès at Huazteca Potosina, San Luis Potosí, Mexico.	

PLATE CCXXXIII.

588. <i>Azurina hirundo</i> Jordan and McGregor.....	1544
Drawing by Anna L. Brown from the type, collected by R. C. McGregor at Guadalupe Island.	
589. <i>Eupomacentrus rectifrenum</i> (Gill)	1557; 3176
Drawing by Anna L. Brown from No. 3460, L. S. Jr. Univ. Mus., collected by the Hopkins Expedition at Mazatlan, Mexico.	

PLATE CCXXXIV.

590. <i>Abudefduf saxatilis</i> (Linnaeus).....	1561
Drawing by A. H. Baldwin from No. 33688, U.S.N.M., col- lected by J. C. Brevoort in the West Indies.	
591. <i>Hypsopops rubicundus</i> (Girard).....	1564
Drawing by W. S. Atkinson from No. 43080, U.S.N.M., col- lected by Rosa Smith at San Diego, California.	

PLATE CCXXXV.

592. <i>Microspathodon bairdii</i> (Gill)	1566
Drawing by Anna L. Brown from No. 2940, L. S. Jr. Univ. Mus., collected by the Hopkins Expedition at Mazatlan, Mexico.	
593. <i>Microspathodon chrysurus</i> (Cuvier and Valenciennes)	1567
Drawing by A. H. Baldwin from No. 13042, U.S.N.M., col- lected by Professor Poey in Cuba.	

PLATE CCXXXVI.

594. <i>Microspathodon dorsalis</i> (Gill)	1568
Drawing by Anna L. Brown from No. 2895, L. S. Jr. Univ. Mus., type of <i>M. azurissimus</i> Jordan and Starks, collected by the Hopkins Expedition at Venados Island, Mazatlan, Mexico.	
595. <i>Tautogolabrus adspersus</i> (Walbaum)	1577
Drawing by H. L. Todd from No. 17745, U.S.N.M., collected by V. N. Edwards at Woods Hole, Massachusetts.	

PLATE CCXXXVII.

- Text page.
596. *Tautoga onitis* (Linnaeus) 1578
 Drawing by H. L. Todd from No. 17738, U.S.N.M., collected by
 V. N. Edwards at Woods Hole, Massachusetts.
 597. *Lachnolaimus maximus* (Walbaum) 1579
 Drawing by H. L. Todd.

PLATE CCXXXVIII.

598. *Pimelometopon pulcher* (Ayres) 1580
 Drawing by H. L. Todd from No. 24890, U.S.N.M., collected
 by Dr. Jordan at San Diego, California.
 599. *Clepticus parvus* (Bloch and Schneider) 1586
 Drawing by A. H. Baldwin from No. 9797, U.S.N.M., collected
 by Professor Poey in Cuba.

PLATE CCXXXIX.

600. *Iridio radiatus* (Linnaeus) 1590
 Drawing by H. L. Todd from No. 31168, U.S.N.M., collected at
 Key West, Florida.
 601. *Irido bivittatus* (Bloch) 1595
 Drawing by A. H. Baldwin from No. 35168, U.S.N.M., collected
 at Key West, Florida.

PLATE CCXL.

602. *Iridio dispilus* (Günther) 1597
 Drawing by Anna L. Brown from No. 2904, L. S. Jr. Univ.
 Mus., collected by the Hopkins Expedition in the Astillero
 at Mazatlan, Mexico.
 603. *Emmeekia venusta* (Jenkins & Evermann) 1602
 Drawing by A. H. Baldwin from the type, No. 39631, U.S.N.M.,
 collected by Jenkins & Evermann at Guaymas, Mexico.

PLATE CCXLI.

604. *Chlorichthys grammaticus* (Gilbert) 1610
 Drawing by A. H. Baldwin from No. 46934, U.S.N.M., collected
 by the *Albatross* at Socorro Island.
 605. *Novaculichthys ventralis* (Bean) 1615
 Drawing by S. F. Denton from the type, No. 37077, U.S.N.M.,
 collected by Dr. Bean at Cozumel Island.
 606. *Novaculichthys infirmus* (Bean) 1616
 Drawing by H. L. Todd from the type, No. 37076, U.S.N.M.,
 collected by Dr. Bean at Cozumel Island.

PLATE CCXLII.

607. *Xyrichtys psittacus* (Linnaeus) 1618
 Drawing by H. L. Todd from No. 5815, U.S.N.M., collected by
 Dr. Whitehurst at Garden Key, Florida.
 608. *Cryptotomus beryllinus* (Jordan and Swain) 1625
 Drawing by H. L. Todd from a specimen collected by the
Albatross.

609. 609a. Dra
le
610. Spari
Dra
U

611. Spari
Draw
S.
Dy

612. Scaru
Draw
col
Draw
Un

614. Scaru
Draw
Mua
615, 615a. Sp
Lower
L. I
616, 616a. Sp
Lower
L. I

617. Pseudor
Drawi
618. Zenopsi
Drawin
Histe

619. Chaetodi
Drawin
coast

620. Chaetodi
Drawin
Mus.,

PLATE CCXLIII.

- | | Text page. |
|--|------------|
| 609, 609a. <i>Calotomus xenodon</i> Gilbert | 1626 |
| Drawings by A. H. Baldwin from No. 46935, U.S.N.M., collected by the <i>Albatross</i> at Socorro Island. | |
| 610. <i>Sparisoma aurofrenatum</i> (Cuvier and Valenciennes) | 1634 |
| Drawing by Anna L. Brown from a specimen in L. S. Jr. Univ. Mus. | |

PLATE CCXLIV.

- | | |
|--|------|
| 611. <i>Sparisoma hoplomystax</i> (Cope)..... | 1632 |
| Drawing by H. L. Todd from No. 35173, U.S.N.M., type of <i>S. cyanoleuca</i> Jordan and Swain, collected by Jordan and Dye at Key West, Florida. | |
| 612. <i>Scarus cuzamilae</i> Bean..... | 1648 |
| Drawing by S. F. Denton from the type, No. 37128, U.S.N.M., collected by Dr. Bean at Cozumel Island. | |
| 613. Jaws of <i>Scarus cæruleus</i> (Bloch) | 1652 |
| Drawing by Anna L. Brown from a specimen in L. S. Jr. Univ. Mus. | |

PLATE CCXLV.

- | | |
|--|------|
| 614. <i>Scarus emblematicus</i> Jordan and Rutter..... | 1654 |
| Drawing by W. S. Atkinson from the type in L. S. Jr. Univ. Mus., collected by Rev. J. S. Roberts in Jamaica. | |
| 615, 615a. <i>Scarus strongylocephalus</i> | |
| Lower and upper pharyngeal bones. Drawings by Anna L. Brown. | |
| 616, 616a. <i>Sparisoma cretense</i> | |
| Lower and upper pharyngeal bones. Drawings by Anna L. Brown. | |

PLATE CCXLVI.

- | | |
|---|------|
| 617. <i>Pseudoscarus guacamai</i> (Cuvier)..... | 1657 |
| Drawing by H. L. Todd. | |
| 618. <i>Zenopsis ocellatus</i> (Storer) | 1660 |
| Drawing from a specimen in the Boston Society of Natural History. | |

PLATE CCXLVII.

- | | |
|---|------|
| 619. <i>Chætodipterus faber</i> (Broussonet)..... | 1668 |
| Drawing by H. L. Todd from No. 22921, U.S.N.M., from east coast of United States. | |

PLATE CCXLVIII. *

- | | |
|---|------|
| 620. <i>Chætodon nigrirostris</i> (Gill)..... | 1673 |
| Drawing by W. S. Atkinson from No. 5981, L. S. Jr. Univ. Mus., collected by R. C. McGregor at Clarion Island. | |

PLATE CCXLIX.

- Text page.
621. *Chætodon ocellatus* (Bloch) 1674
 Drawing by W. S. Haines. Specimen collected by Dr. Bean
 at Beesley's Point, New Jersey.

PLATE CCL.

622. *Chætodon capistratus* Linnæus 1677
 Drawing by M. M. Smith from No. 30143, U.S.N.M., received
 from the Public Museum of Kingston, Jamaica.

PLATE CCLI.

623. *Pomacanthus arcuatus* (Linnæus) 1679
 Drawing by H. L. Todd from No. 33197, U.S.N.M., collected
 by Lewis G. Mitchell at Barnegat, New Jersey.

PLATE CCLII.

624. *Pomacanthus zonipectus* (Gill). 1681
 Drawing from No. 31482, U.S.N.M., collected by Kaiser and
 Martin in West Florida.

PLATE CCLIII.

625. *Holacanthus tricolor* (Bloch) 1684
 Drawing by A. H. Baldwin from a specimen collected by Dr.
 Evermann at Arroyo, Puerto Rico.

PLATE CCLIV.

626. *Angelichthys ciliaris* (Linnæus) 1684
 Drawing by A. H. Baldwin from No. 30023, U.S.N.M., received
 from the Public Museum at Kingston, Jamaica.

- 626a. *Angelichthys ciliaris* (Linnæus), skeleton showing vertebrae
 reduced in number and enlarged 1684
 Drawing by W. S. Atkinson from a specimen in L. S. Jr.
 Univ. Mus.

PLATE CCLV.

627. *Zanclus cornutus* (Linnæus) 1687
 Drawing by W. S. Atkinson from a specimen in L. S. Jr.
 Univ. Mus.

629a. Tail
630. Xesur
Draw
le

631. Baliste
Draw
at N

632. Canthe
Draw
Uni

633. Monaca
Drawi
634. Monaca
Drawi

635. Alutera
Drawi
by J

636. Alutera
Drawing
by V

637. Lactoph
Drawing
at Ga

638. Lactoph
Drawing
by C.

639, 639a, 639b.

628. *Teuthis crestonis* Jordan and Starks 1692
 Drawing by Anna L. Brown from the type, No. 2899, L. S. Jr.
 Univ. Mus., collected by the Hopkins Expedition in the
 Astillero at Mazatlan, Mexico.

629. *Teuthis bahianus* (Castelnau) 1693
 Drawing by H. L. Todd from No. 3424, U.S.N.M., collected at
 Garden Key, Florida.

640, 640a. Lacto
Dra
le

641. Lagoceph
Drawing

PLATE CCLVII.

	Text page.
629a. Tail of <i>Teuthis bahianus</i> (Castelnau).....	1693
630. <i>Xesurus punctatus</i> (Gill).....	1694

Drawing by Anna L. Brown from No. 2911, U.S.N.M., collected by the Hopkins Expedition at Mazatlan, Mexico.

PLATE CCLVIII.

631. <i>Balistes carolinensis</i> Gmelin	1701
Drawing by H. L. Todd from No. 15233, U.S.N.M., collected at New York.	
632. <i>Cantherines carolæ</i> Jordan and McGregor	1713
Drawing by Anna L. Brown from the type, No. 11995, L. S. Jr. Univ. Mus., collected by R. C. McGregor at Socorro Island.	

PLATE CCLIX.

633. <i>Monacanthus ciliatus</i> (Mitchill).....	1714
Drawing by W. S. Haines.	
634. <i>Monacanthus hispidus</i> (Linnaeus)	1715
Drawing by W. S. Haines.	

PLATE CCLX.

635. <i>Alutera schœppii</i> (Walbaum)	1718
Drawing by W. S. Haines from No. 6068, U.S.N.M., collected by Judge Steele at Cedar Keys, Florida.	
636. <i>Alutera scripta</i> (Osbeck)	1719
Drawing by W. S. Haines from No. 34397, U.S.N.M., collected by V. N. Edwards at Woods Hole, Massachusetts.	

PLATE CCLXI.

637. <i>Lactophrys triqueter</i> (Linnaeus)	1722
Drawing by W. S. Haines from No. 5989, U.S.N.M., collected at Garden Key, Florida, by Dr. Whitehurst.	
638. <i>Lactophrys tricornis</i> (Linnaeus)	1724
Drawing by W. S. Haines from No. 21548, U.S.N.M., collected by C. C. Leslie at Charleston, South Carolina.	

PLATE CCLXII.

639, 639a, 639b. <i>Lactophrys bicaudalis</i> (Linnaeus).....	1723
Drawings by H. L. Todd from No. 37130, U.S.N.M., collected at Cozumel Island by the <i>Albatross</i> ; 640a, ventral view; 640b, front view.	

PLATE CCLXIII.

640, 640a. <i>Lactophrys trigonus</i> (Linnaeus)	1723
Drawing by H. L. Todd from No. 7987, U.S.N.M., collected at Garden Key, Florida, by Dr. Whitehurst.	
641. <i>Lagocephalus leavigatus</i> (Linnaeus)	1728
Drawing by H. L. Todd from No. 35135, U.S.N.M.	

PLATE CCLXIV.

- Text page.
642. **Sphercides nephelus** Goode and Bean 1732
 Drawing by H. L. Todd from No. 31428, U.S.N.M., the type.
 collected by R. E. Earll in Indian River, Florida.
643. **Spheroides spengleri** (Bloch) 1732
 Drawing from No. 20608, U.S.N.M., collected by V. N. Edwards
 at Woods Hole, Massachusetts.
644. **Spheroides maculatus** (Bloch and Schneider) 1733
 Drawing by W. S. Haines from No. 14827, U.S.N.M., collected
 by the U. S. Fish Commission at Noank, Connecticut.

PLATE CCLXV.

- 645, 645a. **Spheroides testudineus** (Linnaeus) 1734
 Drawings by H. L. Todd from No. 30060, U.S.N.M.,
 received from the museum at Kingston, Jamaica.
646. **Ovoides setosus** (Rosa Smith) 1739
 Drawing by W. S. Atkinson from No. 319, L. S. Jr. Univ. Mus.,
 collected by the *Albatross* at Clarion Island.

PLATE CCLXVI.

647. **Diodon hystrix** Linnaeus 1745
 Drawing by H. L. Todd from No. 5985, U.S.N.M., collected by
 H. E. Wright at Tortugas Islands.
648. **Chilomycterus schœpfi** (Walbaum) 1748
 Drawing from a specimen collected at Noank, Connecticut.

PLATE CCLXVII.

649. **Lyosphæra globosa** Evermann and Kendall 1751
 Drawing by A. H. Baldwin from the type, No. 48794, U.S.N.M.,
 collected by W. C. Kendall in the Rappahannock River,
 Virginia.
- 649a. **Lyosphæra globosa** Evermann and Kendall 1751
 Drawing by A. H. Baldwin from the cotype, a young example
 collected by Evermann and Kendall at Cape Florida,
 Florida.
650. **Mola mola** (Linnaeus) 1753
 Drawing taken from a washed drawing taken from a cast.

PLATE CCLXVIII.

651. **Ranzania truncata** (Retzius) 1755
 Drawing from a specimen collected in the Bermudas.
652. **Sebastodes marinus** (Linnaeus) 1760
 Drawing by H. L. Todd.

653. **Sebastodes**
 Drawing
 Alba
654. **Sebastodes**
 Drawing
 by L

655. **Sebastodes**
 Drawing
 in To
656. **Sebastodes**
 Drawing
 by Dr

657. **Sebastodes**
 Drawing
 collect
658. **Sebastodes**
 Drawing
 Univ.
 Grove,

659. **Sebastodes**
 Drawing
 Jr. Un
 Grove,
660. **Sebastodes**
 Drawing
 collec

661. **Sebastodes**
 Drawing
 by J. G.
662. **Sebastodes**
 Drawing
 Univ. M
 tively in

PLATE CCLIX.

	Text page.
633. <i>Sebastolobus altivelis</i> Gilbert	1763
Drawing by A. H. Baldwin from the type collected by the <i>Albatross</i> at Station 3338, south of the Alaskan Peninsula, in 625 fathoms.	
634. <i>Sebastodes melanops</i> (Girard).....	1782
Drawing by H. L. Todd from No. 27628, U.S.N.M., collected by Dr. Bean at Sitka, Alaska.	

PLATE CCLXX.

635. <i>Sebastodes ciliatus</i> (Tilesius).....	1783
Drawing by H. L. Todd from No. 32014, U.S.N.M., collected in Tolstoi Bay, Alaska, by Capt. H. E. Nichols.	
636. <i>Sebastodes mystinus</i> (Jordan and Gilbert).....	1784
Drawing by H. L. Todd from No. 27031, U.S.N.M., collected by Dr. Jordan at Monterey, California.	

PLATE CCLXXI.

637. <i>Sebastodes brevispinis</i> (Bean).....	1787
Drawing by W. S. Haines from the type, No. 32004, U.S.N.M., collected by Capt. H. E. Nichols in Hassler Harbor, Alaska.	
638. <i>Sebastodes eigenmanni</i> Cramer.....	1789
Drawing by Anna L. Brown from the type, No. 4046, L. S. Jr. Univ. Mus., collected by Dr. W. W. Thoburn at Pacific Grove, California.	

PLATE CCLXXII.

639. <i>Sebastodes hopkinsi</i> Cramer.....	1789
Drawing by Anna L. Brown from the type, No. 2282, L. S. Jr. Univ. Mus., collected by Gilbert and Starks at Pacific Grove, California.	
640. <i>Sebastodes alutus</i> (Gilbert)	1790
Drawing by A. H. Baldwin from the type, No. 48244, U.S.N.M., collected by the <i>Albatross</i> at Station 3339, Santa Barbara Islands.	

PLATE CCLXXIII.

641. <i>Sebastodes pinniger</i> (Gill).....	1793
Drawing by H. L. Todd from No. 27488, U.S.N.M., collected by J. G. Swan at Neah Bay, Washington.	
642. <i>Sebastodes miniatus</i> (Jordan and Gilbert)	1794
Drawing by W. S. Atkinson from a specimen in L. S. Jr. Univ. Mus., showing vertebrae in moderate number, rela- tively increased in size.	

PLATE CCLXXIV.

- | | Text page. |
|---|------------|
| 663. <i>Sebastodes caurinus</i> (Richardson)..... | 1820 |
| Drawing by H. L. Todd from No. 31999, U.S.N.M., collected
by Capt. H. E. Nichols in Departure Bay, British Columbia. | |
| 664. <i>Sebastodes maliger</i> (Jordan and Gilbert) | 1822 |
| Drawing by H. L. Todd from No. 27713, U.S.N.M., collected
by Dr. Bean at Sitka, Alaska. | |

PLATE CCLXXV.

- | | |
|--|------|
| 665. <i>Sebastodes gilberti</i> Cramer | 1823 |
| Drawing by Anna L. Brown from the type, No. 3893, L. S. Jr.
Univ. Mus., collected by Dr. Jordan in the San Francisco
market. | |
| 666. <i>Sebastodes serriceps</i> (Jordan and Gilbert)..... | 1827 |
| Drawing by H. L. Todd from No. 27041, U.S.N.M., collected
by Dr. Jordan at Monterey, California. | |

PLATE CCLXXVI.

- | | |
|---|------|
| 667. <i>Sebastodes nigrocinctus</i> (Ayres)..... | 1827 |
| Drawing by H. L. Todd from No. 27285, U.S.N.M., collected by
Dr. Jordan in Puget Sound. | |
| 668. <i>Scorpaena cristulata</i> Goode and Bean..... | 1841 |
| Drawing by H. L. Todd from the type, No. 39326, U.S.N.M.,
collected by the <i>Albatross</i> at Station 2415, in N. lat. $30^{\circ} 44'$,
W. long. $79^{\circ} 26'$, off Georgia, in 440 fathoms. | |

PLATE CCLXXVII.

- | | |
|--|------|
| 669. <i>Scorpaena brasiliensis</i> Cuvier and Valenciennes | 1842 |
| Drawing by H. L. Todd from No. 30169, U.S.N.M., collected by
Silas Stearns at Pensacola, Florida. | |
| 670. <i>Scorpaena mystes</i> Jordan and Starks..... | 1849 |
| Drawing by Anna L. Brown from the type, No. 2919, L. S. Jr.
Univ. Mus., collected by the Hopkins Expedition at Mazat-
lan, Mexico. | |

PLATE CCLXXVIII.

- | | |
|--|------|
| 671. <i>Scorpaena grandicornis</i> Cuvier and Valenciennes..... | 1850 |
| Drawing by H. L. Todd from No. 35101, U.S.N.M., collected by
Dr. Jordan at Key West, Florida. | |
| 672. <i>Pontinus macrolepis</i> Goode and Bean | 1855 |
| Drawing by A. H. Baldwin from a specimen collected by the
<i>Brake</i> off Barbados. | |

PLATE CCLXXIX.

- | | |
|---|------|
| 673. <i>Setarches parmatus</i> Goode | 1860 |
| Drawing by H. L. Todd from a specimen collected by the
<i>Albatross</i> at Station 2397. | |

674. *Anoplo-*
Drawing675. *Pleuro-*
Drawing676. *Hexagra-*
Drawing677. *Skeleto-*
A mail

incre-

W. S.

678. *Hexagra-*
Drawing679. *Hexagra-*
St. Mi680. *Hexagra-*
Drawing

Univ.

681. *Ophidion*
Drawing

by Dr.

682. *Oxylebius*
Drawing

Mus.,

Franci

683. *Jordania*
Drawing

Univ.

Hubbe

684. *Alcidea th*
Drawing*Albatro*685. *Skull of S*

Text page.

674. *Anoplopoma fimbria* (Pallas) 1862
 Drawing by H. L. Todd.
675. *Pleurogrammus monopterygius* (Pallas) 1864
 Drawing by A. H. Baldwin from a specimen collected by Dr. Evermann at Atka Island, Alaska.

PLATE CCLXXX.

676. *Hexagrammos decagrammus* (Pallas) 1867
 Drawing by W. S. Haines from No. 27710, U.S.N.M., collected at Old Sitka by L. A. Beardslee.
- 676a. *Skeleton of Hexagrammos decagrammus* 1867
 A mail-cheeked fish from northern waters, showing vertebrae increased in number and diminished in size. Drawing by W. S. Atkinson from a specimen in L. S. Jr. Univ. Mus.
677. *Hexagrammos octogrammus* (Pallas) 1869
 Drawing by W. S. Haines from No. 27975, U.S.N.M., collected at Unalaska by Dall and Bean.

PLATE CCLXXXI.

678. *Hexagrammos stelleri* Tilesius 1871
 Drawing by H. L. Todd from No. 21530, U.S.N.M., collected at St. Michaels, Alaska, by L. M. Turner.
679. *Hexagrammos superciliosus* (Pallas) 1872
 Drawing by M. M. Smith from No. 27934, U.S.N.M., collected at Chernoffsky, Unalaska Island, by T. H. Bean.
680. *Hexagrammos otakii* Jordan and Starks 1867
 Drawing by Anna L. Brown from the type, No. 3704, L. S. Jr. Univ. Mus., collected by Keinosuke Otaki at Tokio, Japan.

PLATE CCLXXXII.

681. *Ophidion elongatus* Girard 1875
 Drawing by H. L. Todd from No. 27657, U.S.N.M., collected by Dr. Bean at Sitka, Alaska.
682. *Oxylebius pictus* Gill 1878
 Drawing by Anna L. Brown from No. 2381, L. S. Jr. Univ. Mus., collected by the California Fish Commission at San Francisco, California.
683. *Jordania zonope* Starks 1884
 Drawing by Anna L. Brown from the type, No. 3124, L. S. Jr. Univ. Mus., collected by Miss Maude Parker and Adam Hubert at Point Orebard, Puget Sound, Washington.

PLATE CCLXXXIII.

684. *Alcidea thoburni* (Gilbert) 1887
 Drawing by A. H. Baldwin from a specimen collected by the Albatross at Station 3350.
685. *Skull of Scorpænichthys marmoratus* (Ayres) 1889

- Text page.
1890
686. *Chitonotus pugetensis* (Steindachner)
 Drawing by A. H. Baldwin from No. 27238, U.S.N.M., collected by Dr. Jordan in the Straits of Juan de Fuca, Washington.

PLATE CCLXXXIV.

687. *Icelinus borealis* Gilbert 1896
 Drawing by A. H. Baldwin from the type collected by the *Albatross* at Station 3223.
688. *Astrolytes notospilotus* (Girard) 1899
 Drawing by A. H. Baldwin from No. 27146, U.S.N.M., collected by Dr. Jordan in Puget Sound.
- 688a. *Astrolytes notospilotus* (Girard) 1899
 Drawing from No. 23936, U.S.N.M., a young example, collected by W. H. Dall in the Shumagin Islands.

PLATE CCLXXXV.

689. *Arteidiellus atlanticus* Jordan and Evermann 1906
 Drawing by H. L. Todd from the type, No. 21069, U.S.N.M., collected in Massachusetts Bay in 90 fathoms.
690. *Ruscarius meanyi* Jordan and Starks 1908
 Drawing by Anna L. Brown from the type, No. 3127, L. S. Jr. Univ. Mus., collected by E. C. Starks at Point Orchard, Puget Sound, Washington.

PLATE CCLXXXVI.

691. *Rastrinus scutiger* (Bean) 1909
 Drawing by W. S. Atkinson from a specimen in L. S. Jr. Univ. Mus.
692. *Icelus spiniger* Gilbert 1914
 Drawing by A. H. Baldwin from the type, collected by the *Albatross* at Station 3278.
693. *Icelus canaliculatus* Gilbert 1917
 Drawing by A. H. Baldwin from the type, collected by the *Albatross* at Station 3329, off Unalaska.

PLATE CCLXXXVII.

694. *Radulinus boleoides* Gilbert 1919
 Drawing by A. H. Baldwin from the type, No. 48795, U.S.N.M., collected by the *Albatross* at Station 3664, off Santa Catalina Island, California, in 59 fathoms.
695. *Radulinus aspiellus* Gilbert 1920
 Drawing by Anna L. Brown from No. 3781, L. S. Jr. Univ. Mus., collected by E. C. Starks in Puget Sound, near Seattle, Washington.
696. *Triglops pingeli* Reinhardt 1923
 Drawing by H. L. Todd from a specimen taken in trawl 8 miles off Chebucto in 521 fathoms.

697. *Triglops* Drawing
Albatros
 698. *Triglops* Drawing
Albatros
 699. *Sternias* Drawing
Albatros
 699a. *Sternias* Drawing
Albatros

700. *Prionistius* Drawing
 Drawing by collected
 Columbia
 701. *Elanura* Drawing
 Drawing by
Albatross
 702. *Melletes* Drawing
 Drawing by
 collected
 Group, Ba

703. *Hemilepidot* Drawing
 Drawing by
 collected
 704. 704a, 704b. *H* Drawing by
 Drawings by
 at Sitka b

705. *Enophrys bis* Drawing
 Drawing by
 Jordan in
 706. *Ceratocottus* Drawing
 Drawing by
 by Capt. H

707. *Cottus everma* Drawing
 Drawing by
 U.S.N.M., e
 River, near
 Bull. No. 47, p

PLATE CCLXXXVIII.

Text page.

697. *Triglops beani* Gilbert 1924
 Drawing by H. L. Todd from the type, collected by the
Albatross at Station 3220.
698. *Triglops scepticus* Gilbert 1925
 Drawing by A. H. Baldwin from the type, collected by the
Albatross at Station 3339.
699. *Sternias xenostethus* (Gilbert) 1927
 Drawing by A. H. Baldwin from the type, collected by the
Albatross at Station 3220, north of Unalaska, in 34 fathoms.
- 699a. *Sternias xenostethus* (Gilbert) 1927
 Drawing by A. H. Baldwin from the type, collected by the
Albatross at Station 3220, north of Unalaska, in 34 fathoms.

PLATE CCLXXXIX.

700. *Prionistius macellus* Bean 1928
 Drawing by H. L. Todd from the type, No. 31958, U.S.N.M.,
 collected by Capt. H. E. Nichols in Carter Bay, British
 Columbia.
701. *Elanura forficata* Gilbert 1930
 Drawing by Anna L. Brown from the type, collected by the
Albatross at Station 3214, in the Aleutian Islands.
702. *Melletes papilio* Bean 1932
 Drawing by H. L. Todd from the type, No. 23751, U.S.N.M.,
 collected by H. W. Elliott at St. Paul Island, Pribilof
 Group, Bering Sea.

PLATE CCXC.

703. *Hemilepidotus jordani* Bean 1934
 Drawing by H. L. Todd from the type, No. 27598, U.S.N.M.,
 collected by Sylvanus Bailey at Unalaska.
- 704, 704a, 704b. *Hemilepidotus hemilepidotus* (Tilesius) 1935
 Drawings by S. F. Denton from No. 27609, U.S.N.M., collected
 at Sitka by T. H. Bean.

PLATE CCXCI.

705. *Enophrys bison* (Girard) 1738
 Drawing by H. L. Todd from a specimen collected by Dr.
 Jordan in Puget Sound.
706. *Ceratocottus diceraus* (Pallas) 1940
 Drawing by W. S. Haines, from No. 32007, U.S.N.M., collected
 by Capt. H. E. Nichols in Tolstoi Bay, Alaska.

PLATE CCXII.

707. *Cottus evermanni* Gilbert 1945
 Drawing by Chloe Lesley Starks from the type, No. 48228,
 U.S.N.M., collected by Gilbert, Cramer, and Otaki in Lost
 River, near Lostine, Oregon.
 Bull. No. 47, pt. 4—xvii

708. *Cottus punctulatus* (Gill) 1948
 Drawing by S. F. Denton from a specimen collected in Yellowstone Park by Dr. Jordan.

PLATE CCXCIII.

709. *Cottus perplexus* Gilbert and Evermann 1955
 Drawing by A. H. Baldwin from the type, No. 45387, U.S.N.M., collected by Drs. Gilbert and Jenkins in the Skookumchuck River, near Chehalis, Washington.
710. *Cottus klamathensis* Gilbert 1955
 Drawing by Anna L. Brown from the type, No. 48226, U.S.N.M., collected by Gilbert, Cramer, and Otaki, in Upper Klamath Lake, Oregon.
711. *Cottus aleuticus* Gilbert 1957
 Drawing from No. 26922, U.S.N.M., the type of *Uranidea microstoma* Lockington, collected by W. J. Fischer at St. Paul, Kadiak Island, Alaska.

PLATE CCXCIV.

712. *Cottus leiopomus* Gilbert and Evermann 1962
 Drawing by A. H. Baldwin from the type, No. 45389, U.S.N.M., collected by H. H. Kinsey in Little Wood River, near Shoshone, Idaho.
713. *Cottus princeps* Gilbert 1962
 Drawing by Anna L. Brown from the type, No. 48227, U.S.N.M., collected by Gilbert, Cramer, and Otaki in Upper Klamath Lake, Oregon.
714. *Uranidea tenuis* Evermann and Meek 1966
 Drawing by A. H. Baldwin from the type, No. 48229, U.S.N.M., collected by Meek and Alexander in Upper Klamath Lake, Oregon.

PLATE CCXCV.

- 715, 715a. *Myoxocephalus seneus* (Mitchill) 1972
 Drawings by H. L. Todd from No. 15093, U.S.N.M., collected by Mr. Copley at Tompkinsville, New York.
716. *Myoxocephalus scorpius* (Linneaus) 1974
 Drawing by H. L. Todd from No. 21989, U.S.N.M., collected by L. Kumlein in Cumberland Gulf.

PLATE CCXCVI.

717. *Myoxocephalus octodecimspinosis* (Mitchill) 1976
 Drawing by H. L. Todd from No. 4552, U.S.N.M., collected by Professor Baird at Beesley Point, New Jersey.
718. *Myoxocephalus polyacanthocephalus* Pallas 1976
 Drawing by H. L. Todd from No. 23499, U.S.N.M., collected by W. H. Dall at Unalaska, Alaska.
719. *Myoxocephalus jack* (Cuvier and Valenciennes) 1977
 Drawing by H. L. Todd from No. 21522, U.S.N.M., collected by L. M. Turner at St. Michaels, Alaska.

720, 720a. **M**
 Drawn
 by I.721. **Myoxo**
 Drawn
 U. S.
 Stra722, 722a, 722
 Drawin
 by U.723. **Myoxoce**
 Drawin
 collec724, 724a. **Meg**
 Drawing
 lected725, 725a. **Meg**
 Drawing
*Albatro*726. **Zesticelus**
 Drawing
*Albatro*727. **Dasycottu**
 Drawing
 Mus., e728. **Oncocottu**
 Drawing
 by E. W.729, 729a, 729b. **G**730. **Gymnocant**
 Drawing b
 collected

PLATE CCXCVII.

Text page.

- 720, 720a. *Myoxocephalus verrucosus* (Bean) 1979
 Drawing by H. L. Todd from No. 27847, U.S.N.M., collected
 by Dall and Bean in Plover Bay, Alaska.
721. *Myoxocephalus axillaris* (Gill) 1980
 Drawing by H. L. Todd from No. 32442, U.S.N.M., collected by
 U. S. Signal Service Point Barrow Expedition at Bering
 Straits, Alaska.

PLATE CCXCVIII.

- 722, 722a, 722b. *Myoxocephalus stelleri* Tilesius 1981
 Drawings by H. L. Todd from No. 32442, U.S.N.M., collected
 by U. S. Signal Service at Bering Straits, Alaska.
723. *Myoxocephalus niger* (Bean) 1985
 Drawing by H. L. Todd from the type, No. 27971, U.S.N.M.,
 collected by Dr. Bean at St. Paul Island, Bering Sea.

PLATE CCXCIX.

- 724, 724a. *Megalocottus platycephalus* (Pallas) 1987
 Drawings by A. H. Baldwin from No. 21522, U.S.N.M., col-
 lected at St. Michaels, Alaska, by L. M. Turner.
- 725, 725a. *Megalocottus laticeps* (Gilbert) 1988
 Drawings by A. H. Baldwin from the type, collected by the
Albatross in the Nushagak River, Alaska.

PLATE CCC.

726. *Zesticelus profundorum* (Gilbert) 1990
 Drawing by A. H. Baldwin from the type, collected by the
Albatross at Station 3329 in Bering Sea, north of Unalaska.
727. *Dasycottus setiger* Bean 1991
 Drawing by Anna L. Brown from No. 3761, L. S. Jr. Univ.
 Mus., collected by E. C. Starks in Puget Sound, near
 Seattle, Washington.
728. *Oncocottus quadricornis* (Linnaeus) 2001
 Drawing by W. S. Haines from No. 32962, U.S.N.M., collected
 by E. W. Nelson at St. Michaels, Alaska.

PLATE CCCI.

- 729, 729a, 729b. *Gymnophanthus pistilliger* (Pallas) 2006
 Drawings by H. L. Todd from No. 21743, U.S.N.M.,
 collected at Niantilik Harbor, Cumberland Gulf,
 by Ludwig Kunlien.
730. *Gymnophanthus galeatus* Bean 2010
 Drawing by W. S. Haines from the type, No. 28097, U.S.N.M.,
 collected by Dr. Bean at Unalaska, Alaska.

PLATE CCCII.

731. *Leiocottus hirundo* Girard 2011
 Drawing by A. H. Baldwin from No. 313, U.S.N.M., collected by
 W. P. Trowbridge at San Miguel, California.
732. *Leptocottus armatus* Girard 2012
 Drawing by H. L. Todd from No. 27968, U.S.N.M., collected at
 Sitka, Alaska, by Dr. Bean.
733. *Oligocottus maculosus* Girard 2013
 Drawing by H. L. Todd from No. 27514, U.S.N.M., collected by
 Dr. Bean at Sitka, Alaska.

PLATE CCCIII.

734. *Blennicottus embryum* (Jordan and Starks) 2016; 2864
 Drawing by Anna L. Brown from the type, No. 3128, L. S. Jr.
 Univ. Mus., collected by E. C. Starks, Neah Bay, Washington.
735. *Histiocottus bilobus* (Cuvier and Valenciennes) 2018
 Drawing by W. S. Haines from No. 30307, U.S.N.M., collected
 by W. J. Fischer at St. Paul, Kadiak Island, Alaska.

PLATE CCCIV.

736. *Blepsias cirrhosus* (Pallas) 2018
 Drawing by A. H. Baldwin from No. 27118, U.S.N.M., collected
 by Dr. Jordan in the Straits of Fuca.
- 736a, 736b. *Blepsias cirrhosus* (Pallas) 2018
 Drawings by H. L. Todd from No. 30307, U.S.N.M., collected
 by W. J. Fischer at St. Paul, Kadiak Island, Alaska.

PLATE CCCV.

738. *Hemitripterus americanus* (Gmelin) 2023
 Drawing by H. L. Todd from No. 23199, U.S.N.M., collected by
 the U. S. Fish Commission at Halifax, Nova Scotia, in 16
 fathoms.
739. *Ascelichthys rhodorus* Jordan and Gilbert 2025
 Drawing by A. H. Baldwin from No. 28488, U.S.N.M., collected
 by J. G. Swan at Neah Bay, Washington.

PLATE CCCVI.

740. *Psychrolutes paradoxus* Günther 2026
 Drawing by Anna L. Brown from No. 3371, L. S. Jr. Univ. Mus.,
 collected by E. C. Starks in Puget Sound near Point
 Orchard, Washington.
741. *Gilbertidia sigulata* (Jordan and Starks) 2028
 Drawing by Anna L. Brown from the type, No. 3129, L. S. Jr.
 Univ. Mus., collected by E. C. Starks in Puget Sound near
 Point Orchard, Washington.

- 742, 742a. *Rh* 1
 Drawing by W. J. Fischer at Port Clarence, Alaska.
- 742b, 742c. *F* 1
 Drawing by W. J. Fischer at Port Clarence, Alaska.
743. *Occa do* 1
 Drawing by W. J. Fischer at Port Clarence, Alaska.
744. *Pallasina* 1
 Drawing by W. J. Fischer at Port Clarence, Alaska.
- 745, 745a. *Poda* 1
 Drawing by W. J. Fischer at Port Clarence, Alaska.
746. *Podothecu* 1
 Drawing by W. J. Fischer at Port Clarence, Alaska.
- 747, 747a. *Podo* 1
 Drawing by W. J. Fischer at Port Clarence, Alaska.
- 748, 748a. *Stelg* 1
 Drawing by W. J. Fischer at Port Clarence, Alaska.
- 749, 749a. *Averr* 1
 Drawing by W. J. Fischer at Port Clarence, Alaska.
- 750, 750a. *Averru* 1
 Drawing by W. J. Fischer at Port Clarence, Alaska.
751. *Sarritor frenata* 1
 Drawing by W. J. Fischer at Port Clarence, Alaska.

PLATE CCCVII.

	Text page.
742, 742a. <i>Rhamphocottus richardsoni</i> Günther	2030
Drawings by H. L. Todd from a specimen collected by W. N. Lockington, in California.	
742b, 742c. <i>Rhamphocottus richardsoni</i> Günther	2030
Drawings by Anna L. Brown from No. 3385, L. S. Jr. Univ. Mus., collected by E. C. Starks in Puget Sound near Point Orchard, Washington.	

PLATE CCCVIII.

743. <i>Occa dodecaedron</i> (Tilesius)	2044
Drawing by Anna L. Brown from No. 14936, U.S.N.M., col- lected by Gordon Prince in Kamchatka.	
744. <i>Pallasina barbata</i> (Steindachner)	2049
Drawing by H. L. Todd from No. 28052, U.S.N.M., collected at Port Mulgrave, Alaska, by Dr. Bean.	

745, 745a. *Podothecus accipiter* Jordan and Starks

Drawing by Anna L. Brown from the type, No. 3835, L. S. Jr. Univ. Mus., collected by Capt. J. C. Blair at Robben Island.	2055
--	------

PLATE CCCIX.

746. <i>Podothecus acipenserinus</i> (Tilesius)	2061
Drawing by M. M. Smith from No. 32481, U.S.N.M., collected by W. J. Fischer at Kadiak Island, Alaska.	
747, 747a. <i>Podothecus veterinus</i> Jordan and Starks	2063
Drawing by Anna L. Brown from the type, No. 4823, L. S. Jr. Univ. Mus., collected by Capt. J. C. Blair at Robben Island.	
748, 748a. <i>Stelgis vulsus</i> (Jordan and Gilbert)	2067
Drawing by Anna L. Brown from the type in the U.S.N.M., collected by Jordan and Gilbert at Point Reyes, California.	
749, 749a. <i>Averruncus emmelane</i> Jordan and Starks	2069
Drawing by Anna L. Brown from the type, No. 3135, L. S. Jr. Univ. Mus., collected by E. C. Starks in Puget Sound near Point Orchard, Washington.	

PLATE CCCX.

750, 750a. <i>Averruncus sterletus</i> Gilbert	2071
Drawings by Anna L. Brown from the type in the U.S.N.M., collected by the <i>Albatross</i> at Station 3662, off Avalon, Coronado Island, California, in 47 fathoms.	
751. <i>Sarritor frenatus</i> (Gilbert)	2073
Drawing by A. H. Baldwin from the type collected by the <i>Albatross</i> at Station 3229.	

- Text page.
- 752, 752a. **Xystes axinophrys** Jordan and Starks 2076
 Drawings by Anna L. Brown from the type, No. 3130,
 L. S. Jr. Univ. Mus., collected by E. C. Starks in
 Puget Sound near Seattle, Washington.

PLATE CCCXI.

753. **Bathygonus nigripinnis** Gilbert 2078
 Drawing by A. H. Baldwin from No. 46614, U.S.N.M., col-
 lected by the *Albatross* in the Aleutian Islands.
- 754, 754a. **Xenochirus triacanthus** Gilbert 2084
 Drawings by Anna L. Brown from No. 3760, L. S. Jr.
 Univ. Mus., collected by E. C. Starks in Puget
 Sound near Seattle, Washington.

PLATE CCCXII.

- 755, 755a, 755b. **Aspidophoroides guntheri** Beau 2090
 Drawings by W. S. Haines from No. 37032,
 U.S.N.M., collected by G. M. Stoney in Alaska.
- 756, 756a. **Aspidophoroides monoptygius** (Bloch) 2091
 Drawings by H. L. Todd from No. 21761, U.S.N.M.,
 collected by the U. S. Fish Commission at Sandwich
 Point, Halifax, Nova Scotia, in 18 fathoms.

PLATE CCCXIII.

757. **Cyclopterus lumpus** Linnaeus 2096
 Drawing by H. L. Todd from No. 14795, U.S.N.M., collected
 by the U. S. Fish Commission at Eastport, Maine.
758. **Lethotremus muticus** Gilbert 2101
 Drawing by A. H. Baldwin from the type, collected by the
Albatross at Station 3223, in Unimak Pass, Alaska.

PLATE CCCXIV.

759. **Lethotremus vinolentus** Jordan and Starks 2101
 Drawing by Anna L. Brown from the type, No. 3131, L. S. Jr.
 Univ. Mus., collected by E. C. Starks in Puget Sound, near
 Seattle, Washington.
- 760, 760a. **Neoliparis callyodon** (Pallas) 2110
 Drawings by M. M. Smith from No. 30317, U.S.N.M., col-
 lected by W. J. Fischer at St. Paul, Kadiak Island,
 Alaska.

PLATE CCCXV.

- 761, 761a. **Neoliparis mucosus** (Ayres) 2111
 Drawings by Anna L. Brown from No. 360, Cal. Acad.
 Sci., collected by H. D. Dunn off San Francisco, Califor-
 nia.

762. **Neoliparis**
 Drawi
 Univ
 Neal

763, 763a. **Ne**
 I

764, 764a. **Lip**
 Dr

765. **Liparis** a
 Drawing
 Bean,
 ing Se

766, 766a. **Lipa**
 Dr
 I
 1

767. **Bathyphas**
 Drawing
 collecte
 lotte Is

768. **Prionotus** c
 Drawing
 Professo

769. **Prionotus** a
 Drawing
 Wiirden

770. **Prionotus** a
 Drawing f
 ton, Son

771. **Prionotus** s
 Drawing f
 Banks.

772. **Prionotus** e
 Drawing b
 Professor

Text page.

762. *Neoliparis floræ* Jordan and Starks 2111
 Drawing by Anna L. Brown from the type, No. 3126, L. S. Jr.
 Univ. Mus., collected by E. C. Starks at Waadda Island,
 Neah Bay, Washington.

PLATE CCCXVI.

- 763, 763a. *Neoliparis greeni* Jordan and Starks 2112
 Drawings by Anna L. Brown from the type, No. 3019,
 L. S. Jr. Univ. Mus., collected by A. N. Green in Vic-
 toria Harbor, Victoria, British Columbia.
 764, 764a. *Liparis cyclopus* Günther 2118
 Drawings by H. L. Todd from No. 24007, U.S.N.M., col-
 lected at Port Muller, Alaska, by W. H. Dall.

PLATE CCCXVII.

765. *Liparis agassizii* Putnam 2121
 Drawing by H. L. Todd from one of the types of *L. gibbus*
 Bean, No. 24047, U.S.N.M., collected at St. Paul Island, Ber-
 ing Sea, by H. W. Elliott.
 766, 766a. *Liparis dennyi* Jordan and Starks 2124
 Drawings by Anna L. Brown from the type, No. 3703,
 L. S. Jr. Univ. Mus., collected by E. C. Starks in Ad-
 miralty Inlet, near Seattle, Washington.

PLATE CCCXVIII.

767. *Bathyphasma ovigerum* Gilbert 2128
 Drawing by A. H. Baldwin from the type, No. 48622, U.S.N.M.,
 collected by the *Albatross* at Station 3342, off Queen Char-
 lotte Island, in 1,588 fathoms.
 768. *Prionotus carolinus* (Linnaeus) 2156
 Drawing by H. L. Todd from No. 774, U.S.N.M., collected by
 Professor Baird at Beesleys Point, New Jersey.

PLATE CCCXIX.

769. *Prionotus scitulus* Jordan 2157
 Drawing by H. L. Todd from a specimen collected by G.
 Würdemann in Florida.
 770. *Prionotus alatus* Goode and Bean 2159
 Drawing from the type, collected in deep water off Charles-
 ton, South Carolina.

PLATE CCCXX.

771. *Prionotus stearnsi* Jordan and Swain 2166
 Drawing from a specimen obtained on the Pensacola Snapper
 Banks.
 772. *Prionotus evolans* (Linnaeus) 2168
 Drawing by H. L. Todd from No. 5556, U.S.N.M., collected by
 Professor Baird at Woods Hole, Massachusetts.

PLATE CCCXXI.

- | | Text page |
|---|-----------|
| 773. <i>Bellator egretta</i> (Goode and Bean)..... | 2171 |
| Drawing by M. M. Smith from the type in the M. C. Z., collected by the <i>Blake</i> at Station LXIV, off Barbados. | |
| 774. <i>Peristedion miniatum</i> Goode | 2178 |
| Drawing by H. L. Todd from the type, No. 26023, U.S.N.M., collected by the <i>Fish Hawk</i> at Station 869, in N. lat. $40^{\circ} 02' 18''$, W. long. $70^{\circ} 23' 06''$, in 192 fathoms. | |
| 775. <i>Peristedion longispathum</i> Goode and Bean | 2178 |
| Drawing by H. L. Todd from the type, collected by the <i>Blake</i> at Station LVIII, off Havana, in 242 fathoms. | |

PLATE CCCXXII.

- | | |
|---|------|
| 776. <i>Peristedion gracile</i> Goode and Bean | 2179 |
| Drawing by H. L. Todd from the type, collected by the <i>Albatross</i> at Station 2401, in N. lat. $28^{\circ} 38' 30''$, W. long. $85^{\circ} 52' 30''$, in 142 fathoms. | |
| 777, 777a. <i>Peristedion platycephalum</i> Goode and Bean..... | 2180 |
| Drawings by H. L. Todd from the type, collected by the <i>Blake</i> at Station LX, off Barbados, in 123 fathoms. | |

PLATE CCCXXIII.

- | | |
|---|------|
| 778. <i>Cephalacanthus volitans</i> (Linnaeus) | 2183 |
| Drawing by W. S. Haines. | |
| 779. <i>Callionymus agassizii</i> Goode and Bean..... | 2186 |
| Drawing by H. L. Todd from a specimen collected by the <i>Blake</i> at Station XXX, off Barbados, in 209 fathoms. | |
| 780. <i>Ioglossus calliurus</i> Bean | 2193 |
| Drawing from No. 30797, U.S.N.M., collected by Silas Stearns at the Pensacola Snapper Banks. | |

PLATE CCCXXIV.

- | | |
|---|------|
| 781. <i>Philypnus dormitor</i> (Lacépède) | 2191 |
| Drawing by A. H. Baldwin from a specimen collected by Dr. Evermann at Palo Seco, Puerto Rico. | |
| 782. <i>Dormitator maculatus</i> (Bloch)..... | 2196 |
| Drawing by A. H. Baldwin from a specimen collected by Dr. Evermann at Hucares, Puerto Rico. | |

PLATE CCCXXV.

- | | |
|---|------|
| 783. <i>Eleotris pisonis</i> (Gmelin) | 2200 |
| Drawing by A. H. Baldwin from No. 5757, U.S.N.M., collected by Dr. Whitelhurst at Garden Key, Florida. | |
| 784. <i>Alexurus armiger</i> Jordan and Richardson..... | 2203 |
| Drawing by Anna L. Brown from the type, No. 3455, L. S. Jr. Univ. Mus., collected by J. A. Richardson in La Paz Harbor, Lower California. | |
| 785. <i>Erotelis smaragdus</i> (Cuvier and Valenciennes)..... | 2204 |
| Drawing by A. H. Baldwin. | |

- | | |
|---|--|
| 786. <i>Lophogobius</i> Drawing by F. P. ... | |
| 787. <i>Gobius</i> Drawing by S. ... | |
| 788. <i>Gobius</i> Drawing by D. ... | |
| 789. <i>Gobius</i> Drawing by Dr. ... | |
| 789a. <i>Gobius</i> Drawing by Evermann ... | |
| 790. <i>Garmanni</i> Drawing Mus., co. Mexico ... | |
| 791. <i>Bollmanni</i> Drawing Mus., co. ... | |
| 792. <i>Aboma ethiopicum</i> Drawing Univ. M. Astilleri ... | |
| 793. <i>Clevelandia</i> Drawing Mus., co. Washington ... | |
| 794. <i>Evermannia</i> Drawing Mus., co. Mexico ... | |
| 795. <i>Typhlogobius</i> Drawing collected by ... | |
| 796. <i>Echeneis</i> Drawing by ... | |

PLATE CCCXXVI.

	Text page.
786. <i>Lophogobius cyprinoides</i> (Pallas)	2209
Drawing by A. H. Baldwin from No. 37509, U.S.N.M., collected by Professor Poey in Cuba.	
787. <i>Gobius stigmaticus</i> (Poey)	2224
Drawing by W. S. Haines from No. 30469, U.S.N.M., collected by Silas Stearns in south Florida.	
788. <i>Gobius hastatus</i> Girard	2229
Drawing by H. L. Todd from No. 35155, U.S.N.M., collected by Dr. Jordan at Key West, Florida.	

PLATE CCCXXVII.

789. <i>Gobius oceanicus</i> Pallas	2230
Drawing by A. H. Baldwin from No. 35155, U.S.N.M., collected by Dr. Jordan at Key West, Florida.	
789a. <i>Gobius oceanicus</i> Pallas	2230
Drawing by A. H. Baldwin from a specimen collected by Dr. Evermann at Palo Seco, Puerto Rico.	
790. <i>Garmannia paradoxa</i> (Giluther).	2232
Drawing by Anna L. Brown from No. 3765, L. S. Jr. Univ. Mus., collected by the Hopkins Expedition at Mazatlan, Mexico.	

PLATE CCCXXVIII.

791. <i>Bollmannia chlamydes</i> Jordan	2238
Drawing by W. S. Atkinson from a specimen in L. S. Jr. Univ. Mus., collected by the <i>Albatross</i> off the coast of Colombia.	
792. <i>Aboma etheostoma</i> Jordan and Starks	2240
Drawing by Anna L. Brown from the type, No. 3459, L. S. Jr. Univ. Mus., collected by the Hopkins Expedition in the Astillero at Mazatlan, Mexico.	
793. <i>Clevelandia ios</i> (Jordan and Gilbert).	2254
Drawing by Anna L. Brown from No. 3666, L. S. Jr. Univ. Mus., collected by E. C. Starks in Elliot Bay, near Seattle, Washington.	

PLATE CCCXXIX.

794. <i>Evermannia zosterura</i> (Jordan and Gilbert).	2256
Drawing by Anna L. Brown from No. 2927, L. S. Jr. Univ. Mus., collected by the Hopkins Expedition at Mazatlan, Mexico.	
795. <i>Typhlogobius californiensis</i> Steindachner.	2262
Drawing by A. H. Baldwin from No. 34747, U.S.N.M., col- lected by Rosa Smith at San Diego, California.	
796. <i>Echeneis naucrates</i> Linnaeus.	2269
Drawing by H. L. Todd.	

PLATE CCCXXX.

- Text page.
- 797, 797a. *Remora brachyptera* (Lowe) 2272
 Drawing by H. L. Todd from No. 23374, U.S.N.M., collected by Daniel McEachran in N. lat. $42^{\circ} 40'$, W. long. $63^{\circ} 6'$, in 250 fathoms.
798. *Rhombochirus osteochir* (Cuvier) 2273
 Drawing by H. L. Todd from No. 19022, U.S.N.M., collected by the U. S. Fish Commission at Woods Hole, Massachusetts.
799. *Caulolatilus microps* Goode and Bean 2277
 Drawing by H. L. Todd from No. 20971, U.S.N.M., collected by Silas Stearns at the Pensacola Snapper Banks.

PLATE CCCXXXI.

800. *Opistognathus macrognathum* Poey 2281
 Drawing by H. L. Todd from No. 5936, U.S.N.M., the type of *O. scaphiurus* Goode and Bean, collected by Dr. Whitehurst at Garden Key, Florida.
- 800a. *Opistognathus macrognathum* Poey 2281
 Drawing by A. H. Baldwin from No. 5936, U.S.N.M., the type of *O. scaphiurus* Goode and Bean, collected by Dr. Whitehurst at Garden Key, Florida.
801. *Gnathopops maxillosa* Poey 2284
 Drawing by H. L. Todd from No. 5866, U.S.N.M., collected at Garden Key, Florida.

PLATE CCCXXXII.

802. *Bathymaster signatus* Cope 2288
 Drawing by H. L. Todd from No. 27646, U.S.N.M., collected by Capt. E. P. Herendeen at the Shumagin Islands, Alaska.
803. *Eonquilus jordani* (Gilbert) 2289
 Drawing by Anna L. Brown from No. 3410, L. S. Jr. Univ. Mus., collected by the Young Naturalist Society of Seattle in Elliot Bay, near Seattle, Washington.
804. *Chiasmodon niger* Johnson 2291
 Drawing by H. L. Todd from No. 25632, U.S.N.M., collected by Capt. Thomas F. Hodgdon at the surface on Le Have Bank.

PLATE CCCXXXIII.

805. *Hypsicometes gobiooides* Goode 2294
 Drawing by A. H. Baldwin from a specimen collected by the Albatross at Station 2377.
806. *Trichodon trichodon* (Tilesius) 2295
 Drawing by M. M. Smith from No. 27980, U.S.N.M., collected by Marcus Baker at the Shumagin Islands.
807. *Arctoscopus japonicus* (Steindachner) 2297
 Drawing by W. S. Atkinson from No. 5656, L. S. Jr. Univ. Mus., collected by the Albatross at Station 3652.

808. *Astr*
 Dra
 by

810. *Opsa*
 Dra
 co
 FI
 811. *Poric*
 Dra
 Al

812. *Caula*
 Dra
 813. *Rimic*
 Draw
 Mn
 Cal

814. *Ennea*
 Draw
 coll
 815. *Gibbo*
 Draw
 by
 816. *Neoclin*
 Draw
 coll

PLATE CCCXXXIV.

	Text page.
808. <i>Astroscopus y-græcum</i> (Cuvier and Valenciennes)	2307
Drawing by M. M. Smith from No. 18029, U.S.N.M., collected by J. C. Willets at Matanzas River Inlet, Florida.	
809, 809a. <i>Kathetostoma alboguttata</i> Bean	2312
Drawings by M. M. Smith from No. 2403, U.S.N.M., col- lected by the <i>Albatross</i> .	

PLATE CCCXXXV.

810. <i>Opsanus pardus</i> (Goode and Bean)	2316
Drawing by H. L. Todd from the type, No. 22237, U.S.N.M., collected by Silas Stearns at the Pensacola Snapper Banks, Florida.	
811. <i>Porichthys porosissimus</i> (Cuvier and Valenciennes).....	2319
Drawing by A. H. Baldwin from a specimen collected by the <i>Albatross</i> at Station 2121.	
812. <i>Caularchus mæandricus</i> (Girard)	2328
Drawing by H. L. Todd.	
813. <i>Rimicola muscarum</i> (Meek and Pierson).....	2338
Drawing by A. H. Baldwin from the type in L. S. Jr. Univ. Mus., collected by Meek and Pierson in Monterey Bay, California.	

PLATE CCCXXXVI.

814. <i>Enneanectes carminalis</i> (Jordan and Gilbert)	2350
Drawing by Anna L. Brown from No. 3854, L. S. Jr. Univ. Mus., collected by the Hopkins Expedition at Mazatlan, Mexico.	
815. <i>Gibbonsia elegans evides</i> (Jordan and Gilbert).....	2352
Drawing by A. H. Baldwin from No. 34784, U.S.N.M., collected by Rosa Smith at San Diego, California.	

816. <i>Neoclinus satiricus</i> Girard	2355
Drawing by W. S. Atkinson from No. 2288, L. S. Jr. Univ. Mus., collected by Dr. Gilbert at Pacific Grove, California.	

PLATE CCCXXXVII.

817. <i>Cryptotrema corallinum</i> Gilbert	2266
Drawing by W. S. Atkinson from a specimen in L. S. Jr. Univ. Mus.	
818. <i>Exerpes asper</i> (Jenkins and Evermann)	2367
Drawing by A. H. Baldwin from the type, No. 39643, U.S.N.M., collected by Jenkins and Evermann at Guaymas, Mexico.	

819. <i>Auchenopterus nox</i> (Jordan and Gilbert)	2373
Drawing by Anna L. Brown from the type collected by Dr. Jordan at Key West, Florida.	

PLATE CCCXXXVIII.

	Text page
820. <i>Blennius favosus</i> Goode and Bean	2380
Drawing from the type, No. 2020, U.S.N.M., collected by G. Wiedemann at Garden Key, Florida.	
821. <i>Blennius cristatus</i> Linnaeus	2382
822. <i>Hypsoblennius hentz</i> (Jordan and Gilbert)	2388

PLATE CCCXXXIX.

823. <i>Hypsoblennius hentz</i> (Le Sueur)	2390
Drawing by M. M. Smith from No. 26308, U.S.N.M., collected at Charleston, South Carolina, by C. C. Leslie.	
824. <i>Chasmodes sabra</i> \varnothing Jordan and Gilbert	2392
Drawing by H. L. Todd from the type, No. 30824, U.S.N.M., collected by Jordan and Stearns at Pensacola, Florida.	
825. <i>Rupisartes atlanticus</i> (Cuvier and Valenciennes)	2397
Drawing by A. H. Baldwin from No. 30916, U.S.N.M., collected by C. H. Townsend at San Cristobal, Lower California.	

PLATE CCCXL.

826. <i>Enchlemaria atlantica</i> Jordan and Evermann	2402
Drawing by A. H. Baldwin from the type, No. 33915, U.S.N.M., collected by Silas Stearns at the Pensacola Snapper Banks, Florida.	
827. <i>Stathmonotus hemphilli</i> Bean	2407
Drawing by H. L. Todd from the type, No. 37193, U.S.N.M., collected by Henry Hemphill at Key West, Florida.	

PLATE CCCXLI.

828. <i>Bryostema polyactocephalum</i> (Pallas)	2408
Drawing by H. L. Todd from No. 36889, U.S.N.M., collected by J. G. Swan in Puget Sound, Washington.	
829. <i>Bryostema nigrator</i> Jordan and Williams	2410
Drawing by Anna L. Brown from the type, No. 3134, L. S. Jr., Univ. Mus., collected by the Young Nat. Soc. of Seattle in Elliot Bay, near Seattle, Washington.	
830. <i>Apodichthys flavidus</i> Girard	2411
Drawing by W. S. Haines from No. 100065, U.S.N.M., collected by J. G. Swan at Port Townsend, Washington.	

PLATE CCCXLII.

831. <i>Pholis dellichogaster</i> (Pallas)	2416
Drawing by H. L. Todd from No. 32881, U.S.N.M., collected by Dr. L. Stejneger at Bering Island.	
832. <i>Pholis gunnellus</i> (Linnaeus)	2419
Drawing by H. L. Todd from No. 23198, U.S.N.M., collected by R. E. Earll at Gloucester, Massachusetts.	

833. *Pholis* \varnothing Drawing by P.
 834. *Asterocottus* Drawing by S.
 835. *Anoplarchus* Drawing by I.
 836. *Xiphister* Drawing Univ. Neah.
 837. *Xiphister* Drawing W. H.
 838. *Xiphidium* Drawing Sylva.
 839. *Lumpenus* Drawing Nusha.
 840. *Lumpenus* Drawing
 841. *Stichaeus* Drawing by L.
 842. *Ulvaria* Drawing of Nat.
 843. *Cryptacanthodes* Drawing from F.
 844. *Lyconectes* Drawing *Ulnare*.
 845. *Aharichthys* Drawing by Tap.

	Text page.
833. <i>Pholis ornatus</i> (Girard)	2419
Drawing by M. M. Smith from No. 27996, U.S.N.M., collected by Dr. Bean at Sitka, Alaska.	

PLATE CCCXLIII.

834. <i>Asternopteryx gunnelliformis</i> Riippell	2420
Drawing by A. H. Baldwin from No. 48153, U.S.N.M., collected by Schuchert and White in Omanak Fjord, Karsak, Nour- sook Peninsula, Greenland.	
835. <i>Anoplarchus atropurpureus</i> (Kittlitz)	2422
Drawing by H. L. Todd from No. 29820, U.S.N.M., collected by L. M. Turner at Atka Island, Alaska.	
836. <i>Xiphistes ulvae</i> Jordan and Starks	2423
Drawing by Anna L. Brown from the type, No. 3132 L. S. Jr. Univ. Mus., collected by E. C. Starks at Waadda Island, Neah Bay, Washington.	

PLATE CCCXLIV.

837. <i>Xiphistes chiru</i> (Jordan and Gilbert)	2424
Drawing by H. L. Todd from No. 23964, U.S.N.M., collected by W. H. Dall at Amchitka Island, Alaska.	
838. <i>Xiphidion rupestre</i> (Jordan and Gilbert)	2426
Drawing by H. L. Todd from a specimen collected by Sylvanus Bailey at Sitka, Alaska.	
839. <i>Lumpenus mackayi</i> Gilbert	2436
Drawing by A. H. Baldwin from the type collected in the Nushagak River, Alaska, by the <i>Albatross</i> .	
840. <i>Lumpenus lampetraeformis</i> (Walbaum)	2438
Drawing by H. L. Todd from No. 13852, U.S.N.M.	

PLATE CCCXLV.

841. <i>Stichaeus punctatus</i> (Fabricius)	2439
Drawing by H. L. Todd from No. 21748, U.S.N.M., collected by L. M. Turner at St. Michaels, Alaska.	
842. <i>Ulvaria subbifurcata</i> (Storer)	2440
Drawing by H. L. Todd from a specimen from the Academy of Natural Sciences of Montreal.	
843. <i>Cryptacanthodes maculatus</i> Storer	2443
Drawing by H. L. Todd from No. 22309, U.S.N.M., obtained from E. G. Blackford.	

PLATE CCCXLVI.

844. <i>Lyconectes aleutensis</i> Gilbert	2444
Drawing by A. H. Baldwin from the type, collected by the <i>Albatross</i> at Station 3H12, north of Unalaska, in 15 fathoms.	
845. <i>Anarhichas latifrons</i> Steenstrup and Hallgrímsson ..	2446
Drawing by H. L. Todd from No. 24373, U.S.N.M., collected by Capt. J. W. Collins in N. lat. 43° 56', W. long. 170° 04'.	

PLATE CCCXLVII.

- | | Text page. |
|---|------------|
| 846. <i>Anarhichas lupus</i> Linnaeus..... | 2446 |
| Drawing by H. L. Todd from No. 21846, U.S.N.M., collected
by Capt. John Gourville at Georges Bank. | |
| 847. <i>Anarhichas lepturus</i> Bean..... | 2447 |
| Drawing from the type, collected by L. M. Turner at St. Mi-
chael's, Alaska. | |
| 848. <i>Ptilichthys goodei</i> Bean | 2452 |
| Drawing by H. L. Todd from the type, No. 26619, U.S.N.M.,
collected at Iliniuk (Unalaska), Alaska. | |

PLATE CCCXLVIII.

- | | |
|--|------|
| 849, 849a. 849b. <i>Scyatlina cerdale</i> Jordan and Gilbert | 2451 |
| Drawings by Anna L. Brown from No. 3389, L. S.
Jr. Univ. Mus., collected by E. C. Starks at
Waadda Island, Neah Bay, Washington. | |
| 850. <i>Zoarces anguillaris</i> (Peck)..... | 2457 |
| Drawing by H. L. Todd from No. 10438, U.S.N.M., collected
by the U. S. Fish Commission at Eastport, Maine. | |
| 851. <i>Embryx crotalinus</i> (Gilbert) | 2458 |
| Drawing by A. H. Baldwin from the type, collected by the
Albatross at Station 2980, off Santa Barbara Islands. | |

PLATE CCCXLIX.

- | | |
|--|------|
| 852. <i>Aprodon cortezianus</i> Gilbert | 2461 |
| Drawing by A. H. Baldwin from the type, No. 46457, U.S.N.M.,
collected by the Albatross on Cortez Banks, off San Diego,
California. | |
| 853. <i>Lycodes zoarchus</i> Goode and Bean..... | 2464 |
| Drawing by H. L. Todd from the type, No. 39298, U.S.N.M.,
collected by the Albatross in N. lat. $44^{\circ} 46' 30''$, W. long. $59^{\circ} 53' 45''$, off Nova Scotia, in 130 fathoms. | |
| 854. <i>Lycodes reticulatus</i> Reinhardt | 2465 |
| Drawing by H. L. Todd from a specimen collected by Capt.
R. Markuson at S. W. Banquereau, in 300 fathoms. | |

PLATE CCCL.

- | | |
|---|------|
| 855. <i>Lycodes perspicillum</i> Kröyer..... | 2465 |
| Drawing by H. L. Todd from No. 39336, U.S.N.M., collected
by the Albatross at Station 2456, in N. lat. $47^{\circ} 29'$, W. long.
$52^{\circ} 18'$, in 86 fathoms. | |
| 856. <i>Lycodes frigidus</i> Collett..... | 2465 |
| Drawing by H. L. Todd from No. 32995, U.S.N.M., collected
by the Albatross at Station 2018, in N. lat. $37^{\circ} 12' 22''$, W.
long. $74^{\circ} 20' 04''$, in 788 fathoms. | |

- | | |
|--|------|
| 857. <i>Lycod</i>
Draw
Ly
me | 857. |
| 858. <i>Lycoda</i>
Draw
coll | 858. |
| 859. <i>Lycenc</i>
Drawi
male
sontl | 859. |
| 860, 860a, 860b
D | 860, |
| 861. <i>Furcinia</i>
Drawin
Univ. | 861. |
| 862. <i>Lycodon</i>
Drawing
by the
long. 7 | 862. |
| 863. <i>Lyconema</i>
Drawing
Albatro | 863. |
| 864, 864a, 864b, 8 | 864, |
| 865. <i>Melanosti</i>
Drawing
Albatro | 865. |
| 866. <i>Lepophidiu</i>
Drawing
U.S.N.M.
W. long | 866. |
| 867. <i>Lepophidiu</i>
Drawing
Albatro | 867. |
| 868. <i>Rissola mar</i> | 868. |

	Text page.
857. <i>Lycodalepis polaris</i> (Sabine).....	2468
Drawing by H. L. Todd from No. 27748, U.S.N.M., the type of <i>Lycodes coccineus</i> Bean, collected by Dr. Bean at Big Dio- mede Island, Bering Strait.	
858. <i>Lycodalepis turneri</i> (Bean).....	2469
Drawing by H. L. Todd from the type, No. 21529, U.S.N.M., collected by L. M. Turner at St. Michaels, Alaska.	

PLATE CCCLI.

859. <i>Lycenchelys verrillii</i> (Goode and Bean)	2470
Drawing by H. L. Todd from No. 21015, U.S.N.M., an old male, collected by the U. S. Fish Commission, 27 miles southwest of Chebucto, Nova Scotia.	
860, 860a, 860b, 860c, 860d. <i>Lycenchelys paxillus</i> (Goode and Bean).	2471
Drawings by H. L. Todd from the type, No. 22177, U.S.N.M., collected by Capt. J. W. Collins in N. lat. $42^{\circ} 48'$, W. long. $63^{\circ} 07'$.	
861. <i>Furcinitanus diapterus</i> (Gilbert).....	2472
Drawing by W. S. Atkinson from a specimen in L. S. Jr. Univ. Mus.	

PLATE CCCLII.

862. <i>Lycodonus mirabilis</i> (Goode and Bean)	2474
Drawing by S. F. Denton from No. 39207, U.S.N.M., collected by the <i>Albatross</i> at Station 2742, in N. lat. $37^{\circ} 46' 30''$, W. long. $73^{\circ} 56' 30''$.	
863. <i>Lyconema barbatum</i> Gilbert.....	2474
Drawing by A. H. Baldwin from the type, collected by the <i>Albatross</i> at Station 3129, in 204 fathoms.	
864, 864a, 864b, 864c. <i>Gymnelis viridis</i> (Fabricius).....	2479
Drawing by H. L. Todd from No. 21999a, U.S.N.M., col- lected at Niantilik Harbor, Cumberland Gulf, by W. A. Mintzer.	
865. <i>Melanostigma pammelas</i> Gilbert.....	2479
Drawing by A. H. Baldwin from the type, collected by the <i>Albatross</i> at Station 3202, on coast of California.	

PLATE CCCLIII.

866. <i>Lepophidium marmoratum</i> (Goode and Bean)	2482
Drawing by M. M. Hildebrandt from the type, No. 37237, U.S.N.M., collected by the <i>Albatross</i> in N. lat. $23^{\circ} 10' 39''$, W. long. $82^{\circ} 20' 21''$, in 213 fathoms.	
867. <i>Lepophidium profundorum</i> (Gill)	2484
Drawing by A. H. Baldwin from a specimen collected by the <i>Albatross</i> at Station 2042.	
868. <i>Rissola marginata</i> (De Kay)	2489

PLATE CCCLIV.

- Text page.
869. *Otophidium omostigmum* (Jordan and Gilbert) 2490
 Drawing by H. L. Todd from the type, No. 29670, U.S.N.M.,
 collected by Jordan and Stearns at the Pensacola Snapper
 Banks.
870. *Lycodapus dermatinus* Gilbert 2492
 Drawing by A. H. Baldwin from the type, collected by the
Albatross at Station 3162, off Lower California.
871. *Brotula barbata* (Bloch and Schneider) 2500
 Drawing by A. H. Baldwin from No. 5337, U.S.N.M., collected
 by Professor Poe in Cuba.

PLATE CCCLV.

872. *Ogilbia ventralis* (Gill) 2503
 Drawing by Anna L. Brown from No. 2903, L. S. Jr. Univ.
 Mus., collected by the Hopkins Expedition at Mazatlan,
 Mexico.
873. *Ogilbia cayorum* Evermann and Kendall 2503
 Drawing by A. H. Baldwin from the type, No. 48792, U.S.N.M.,
 collected by Evermann and Kendall at Key West, Florida.
874. *Dicromita agassizii* Goode and Bean 2506
 Drawing by H. L. Todd from the type, collected by the *Blake*
 off Grenada.

PLATE CCCLVI.

875. *Bassozetus normalis* Gill. 2507
 Drawing by H. L. Todd from No. 39416, U.S.N.M., collected
 by the *Albatross* in N. lat. $28^{\circ} 02' 30''$, W. long. $87^{\circ} 43' 45''$.
876. *Bassozetus catena* Goode and Bean 2509
 Drawn by A. H. Baldwin from the type, No. 37341, U.S.N.M.,
 collected by the *Albatross* in N. lat. $28^{\circ} 00' 15''$, W. long.
 $87^{\circ} 42'$, in 1,467 fathoms.
- 876a. *Bassozetus catena* Goode and Bean 2509
 Drawn by H. L. Todd from same specimen.

PLATE CCCLVII.

877. *Neobythites gilli* Goode and Bean 2512
 Drawing by A. H. Baldwin from the type, No. 37340, U.S.N.M.,
 collected by the *Albatross* in N. lat. $28^{\circ} 33'$, W. long. $85^{\circ} 33'$,
 in 111 fathoms.
878. *Neobythites marginatus* Goode and Bean 2513
 Drawing by H. L. Todd from the type collected by the *Blake*
 off Barbados, in 209 fathoms.
879. *Bassogigas gilli* Goode and Bean 2515
 Drawing by A. H. Baldwin from the type, No. 39417, U.S.N.M.,
 collected by the *Albatross* off Cape Henlopen, Delaware, in
 N. lat. $39^{\circ} 35'$, W. long. $70^{\circ} 54'$, in 1,106 fathoms.
880. *Barathrodesmus manatinus* Goode and Bean 2517
 Drawing by H. L. Todd from the type in M. C. Z., collected by
 the *Blake* in N. lat. $33^{\circ} 35' 20''$, W. long. 76° , in 647 fathoms.

927. *Inopsetta* Drawing at U
928. *Lepidopsetta* Drawing St. I
929. *Limanda* Drawing at Gl
930. *Limanda* Drawing Sitka

931. *Limanda* Drawing *Albatr*
932. *Limanda* Drawing collect coast o
933. *Pseudopleuronectes* Drawing
934. *Pleuronectes* Drawing St. Pan

935. *Liopsetta* Drawing Kotzebue
936. *Liopsetta* Drawing Salem, M

- 937, 937a. *Platichthys* Drawing lect

PLATE CCCLXXVI.

	Text page.
927. <i>Inopsetta ischyra</i> (Jordan and Gilbert).....	2641
Drawing by H. L. Todd from No. 32913, U.S.N.M., collected at Unalaska by E. W. Nelson.	
928. <i>Lepidopsetta bilineata</i> (Ayres).....	2643
Drawing from No. 27602, U.S.N.M., collected by Dr. Bean at St. Paul, Kadiak Island, Alaska.	

PLATE CCCLXXVII.

929. <i>Limanda ferruginea</i> (Storer).....	2644
Drawing by H. L. Todd from No. 21903, U.S.N.M., collected at Gloucester, Massachusetts, by the U. S. Fish Commission.	
930. <i>Limanda aspera</i> (Pallas).....	2645
Drawing from No. 27944, U.S.N.M., collected by Dr. Bean at Sitka, Alaska.	

PLATE CCCLXXVIII.

931. <i>Limanda proboscidea</i> Gilbert.....	2645
Drawing by A. H. Baldwin from the type, collected by the <i>Albatross</i> in Bristol Bay, Alaska.	
932. <i>Limanda beanii</i> Goode	2646
Drawing by H. L. Todd from the type, No. 26102, U.S.N.M., collected by the <i>Fish Hawk</i> at Station 875, off the south coast of New England.	

PLATE CCCLXXIX.

933. <i>Pseudopleuronectes americanus</i> (Walbaum).....	2647
Drawing by H. L. Todd from a specimen in the U.S.N.M.	
934. <i>Pleuronectes quadrifasciatus</i> Pallas	2648
Drawing by H. L. Todd from No. 28025, U.S.N.M., collected at St. Paul, Kadiak, by W. J. Fischer.	

PLATE CCCLXXX.

935. <i>Liopsetta glacialis</i> (Pallas).....	2649
Drawing by H. L. Todd from No. 27947, U.S.N.M., collected in Kotzebue Sound by Dall and Bean.	
936. <i>Liopsetta putnami</i> (Gill).....	2650
Drawing by H. L. Todd from No. 5368, U.S.N.M., collected at Salem, Massachusetts, by F. W. Putnam.	

PLATE CCCLXXXI.

937, 937a. <i>Platichthys stellatus</i> (Pallas).....	2652
Drawing by H. L. Todd from No. 27640, U.S.N.M., col- lected by Dr. Bean in Alaska.	

PLATE CCCLXXXII.

938. *Lophopsetta maculata* (Mitchill) 2660
 Drawing by H. L. Todd.
939. *Platophrys ocellatus* (Agassiz) 2663
 Drawing by H. L. Todd from No. 34972, U.S.N.M., type of
 P. nebularis Jordan and Gilbert, collected by Dr. Jordan
 at Key West, Florida.

PLATE CCCLXXXIII.

940. *Trichopsetta ventralis* (Goode and Bean) 2669
 Drawing by H. L. Todd from No. 37372, U.S.N.M., collected
 by the *Albatross* at Station 2378, in N. lat. $29^{\circ} 14' 30''$, W.
 long. $88^{\circ} 09' 30''$, in 68 fathoms.
941. *Syacium papillosum* (Linnaeus) 2671
 Drawing by H. L. Todd from the type of *Hemirhombus patulus*
 Bean, No. 30180, U.S.N.M., collected by Silas Stearns at
 Pensacola, Florida.

PLATE CCCLXXXIV.

942. *Azevia panamensis* (Steindachner) 2677
 Drawing by A. H. Baldwin from No. 15, Mus. Comp. Zool.
943. *Citharichthys sordidus* (Girard) 2679
 Drawing by H. L. Todd from No. 31991, U.S.N.M., collected
 by Capt. H. E. Nichols in Johnstons Straits, British
 Columbia.

PLATE CCCLXXXV.

944. *Citharichthys macrops* Dresel 2684
 Drawing by H. L. Todd from the type, No. 21500, U.S.N.M.,
 collected by Silas Stearns at Pensacola, Florida.
945. *Etropus rimosus* Goode and Bean 2688
 Drawing by H. L. Todd from the type, collected by the
 Albatross at Station 2408, in N. lat. $28^{\circ} 28'$, W. long. $84^{\circ} 25'$,
 in 21 fathoms.

PLATE CCCLXXXVI.

946. *Etropus crossotus* Jordan and Gilbert 2689
 Drawing by H. L. Todd from No. 26571, U.S.N.M., collected
 by Silas Stearns at Cedar Keys, Florida.
947. *Achirus lineatus* (Linnaeus) 2697
 Drawing by H. L. Todd.

PLATE CCCLXXXVII.

948. *Achirus fasciatus* Lacépède 2700
 Drawing by H. L. Todd.
949. *Syphurus marginatus* (Goode and Bean) 2706
 Drawing by H. L. Todd from a specimen collected by the
 Albatross at Station 2376, in N. lat $29^{\circ} 03' 15''$, W. long. 88°
 $16'$, in 324 fathoms.

950. *Syphu-*
Drawing
by L.951. *Syphu-*
Drawing
Univ.952. *Lophins-*
Drawing953. *Chaunax-*
Drawing
by th.954. *Ceratias-*
Drawing
Poiss.955. *Mancalia-*
Drawing
collect.956. *Cryptops-*
Drawing
by the
long. 6957. *Caulophry-*
Drawing
collected
15', in 1

958, 958a, 958b. C.

PLATE CCCLXXXVIII.

	Text page.
950. <i>Sympodus plagiusa</i> (Linnaeus)	2710
Drawing by H. L. Todd from No. 15017, U.S.N.M., collected by Dr. H. C. Yarrow at Beaufort, North Carolina.	
951. <i>Sympodus williamsi</i> Jordan and Culver	2711
Drawing by Anna L. Brown from the type, No. 2943, L. S. Jr. Univ. Mus., collected by the Hopkins Expedition at Mazat- lan, Mexico.	
952. <i>Lophius piscatorius</i> Linnaeus	2713
Drawing by H. L. Todd from a specimen in the U.S.N.M.	

PLATE CCCLXXXIX.

953. <i>Chaunax pictus</i> Lowe	2726
Drawing by H. L. Todd from No. 26021, U.S.N.M., collected by the <i>Albatross</i> at Station 869.	
954. <i>Ceratias holboelli</i> Kröyer	2729
Drawing by A. H. Baldwin from Gaimard, <i>Voy. Skand.</i> , Poissons, pl. ix.	

PLATE CCCXC.

955. <i>Mancalias shufeldti</i> (Gill)	2730
Drawing by H. L. Todd from the type, No. 33552, U.S.N.M., collected by the <i>Albatross</i> at Station 2099, in N. lat. $37^{\circ} 12'$ $20''$, W. long. $69^{\circ} 39'$, in 2,949 fathoms.	
956. <i>Cryptopsaras couesii</i> Gill	2731
Drawing by H. L. Todd from No. 33558, U.S.N.M., collected by the <i>Albatross</i> at Station 2101, in N. lat. $38^{\circ} 13' 30''$, W. long. $68^{\circ} 24'$, in 1,686 fathoms.	

PLATE CCCXCI.

957. <i>Caulophryne jordani</i> Goode and Bean	2735
Drawing by S. F. Denton from the type, No. 39265, U.S.N.M., collected by the <i>Albatross</i> in N. lat. $39^{\circ} 27'$, W. long. 71° $15'$, in 1,276 fathoms.	

PLATE CCCXCII.

958, 958a, 958b. <i>Ogcocephalus vespertilio</i> (Linnaeus)	2737
Drawing by H. L. Todd from No. 2316, U.S.N.M., collected by the <i>Albatross</i> in N. lat. $24^{\circ} 25' 30''$, W. long. $81^{\circ} 47' 45''$.	

881. *Porogadus*
Drawing
W. I.

882. *Dicrolene*
Drawing
in the

883. *Barathra*
Drawing
at Sta.

884. *Merluccius*
Drawing
W. F.

885. *Borogadus*
Drawing
L. M. T.

886. *Pollachium*
Drawing
Profess.

887. *Theragra c*
Drawing
W. H. D.

888. *Eleginus na*
Drawing
H. Bann.

889. *Microgadus*
Drawing b
Yakutat

890. *Microgadus*
Drawing b
V. N. Ed.

891. *Gadus callaris*
Drawing by
Professor

892. *Melanogrammus*
Drawing by
Professor J.

892a. *Melanogrammus*
Drawing of

Bull. No. 47, 1

PLATE CCCLVIII.

	Text page.
881. <i>Porogadus miles</i> Goode and Bean	2520
Drawing by A. H. Baldwin from the type, No. 35625, U.S.N.M., collected by the <i>Albatross</i> at Station 2230, in N. lat. $38^{\circ} 27'$, W. long. $73^{\circ} 02'$, in 1,168 fathoms.	
882. <i>Dicrolene intronigra</i> Goode and Bean	2522
Drawing by H. L. Todd from a specimen collected by the <i>Blake</i> in the Gulf Stream.	
883. <i>Barathronus bicolor</i> Goode and Bean	2524
Drawing by M. M. Smith from the type collected by the <i>Blake</i> at Station LXXI, off Guadeloupe, in 769 fathoms.	

PLATE CCCLIX.

884. <i>Merluccius productus</i> (Ayres)	2531
Drawing by H. L. Todd from No. 26638, U.S.N.M., collected by W. F. Prosser at Seattle, Washington.	
885. <i>Borogadus saida</i> (Lepechin)	2533
Drawing by H. L. Todd from No. 24031, U.S.N.M., collected by L. M. Turner at St. Michaels, Alaska.	
886. <i>Pollachius virens</i> (Linnaeus)	2534
Drawing by H. L. Todd from No. 10443, U.S.N.M., collected by Professor Baird at Eastport, Maine.	

PLATE CCCLX.

887. <i>Theragra chalcogramma</i> (Pallas)	2535
Drawing by H. L. Todd from No. 27637, U.S.N.M., collected by W. H. Dall in Pirate Cove, Shumagin Island, Alaska.	
888. <i>Eleginus navaga</i> (Kölreuter)	2537
Drawing by H. L. Todd from No. 9286, U.S.N.M., collected by H. Bannister at St. Michaels, Alaska.	
889. <i>Microgadus proximus</i> (Girard)	2539
Drawing by H. L. Todd from No. 27982, U.S.N.M., collected at Yakutat Bay, Alaska, by Dr. T. H. Bean.	
890. <i>Microgadus tomcod</i> (Walbaum)	2540
Drawing by H. L. Todd from No. 17733, U.S.N.M., collected by V. N. Edwards at Woods Hole, Massachusetts.	

PLATE CCCLXI.

891. <i>Gadus callarias</i> Linnaeus	2541
Drawing by H. L. Todd from No. 10444, U.S.N.M., collected by Professor Baird at Eastport, Maine.	
892. <i>Melanogrammus æglefinus</i> Linnaeus	2542
Drawing by H. L. Todd from No. 10440, U.S.N.M., collected by Professor Baird at Eastport, Maine.	

892a. <i>Melanogrammus æglefinus</i> Linnaeus	2542
Drawing of skull by H. L. Todd.	







IMAGE EVALUATION TEST TARGET (MT-3)





6"



Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503



EE
2.8
2.2
2.2
2.2
2.0
1.8

IT
Oil

PLATE CCCLXII.

	Text page.
893, 893a. <i>Antimora viola</i> (Goode and Bean)	2544
Drawings by H. L. Todd from the type, No. 21837.	
U.S.N.M., collected by Capt. J. W. Collins on Le Havre Bank, in 400 to 500 fathoms.	

894. <i>Uraleptus malardi</i> (Risso)	2545
Drawing by H. L. Todd from a specimen collected by the <i>Blake</i> at Station LXXXI, off Neris.	

PLATE CCCLXIII.

895. <i>Lotella maxillaris</i> Bean	2546
Drawing by H. L. Todd from the type, No. 29832, U.S.N.M., collected by the <i>Fish Hawk</i> in N. lat. $39^{\circ} 55'$, W. long. $70^{\circ} 28'$, in 396 fathoms.	
896. <i>Physiculus fulvus</i> Bean	2547
Drawing by H. L. Todd from the type, No. 28466, U.S.N.M., collected by the <i>Fish Hawk</i> in N. lat. $40^{\circ} 01'$, W. long. $69^{\circ} 56'$, in 79 fathoms.	

PLATE CCCLXIV.

897. <i>Lota maculosa</i> (Le Sueur)	2550
Drawing by H. L. Todd.	
898. <i>Urophycis regius</i> (Walbaum)	2553
Drawing by H. L. Todd from No. 20923, U.S.N.M., obtained by E. G. Blackford from the New York Aquarium.	
899. <i>Urophycis cirratus</i> (Goode and Bean)	2553
Drawing by H. L. Todd from the type, No. 39059, U.S.N.M., collected by the <i>Albatross</i> at Station 2376, in N. lat. $29^{\circ} 03' 15''$, W. long. $88^{\circ} 16'$.	

PLATE CCCLXV.

900. <i>Urophycis earlii</i> (Bean)	2554
Drawing by H. L. Todd from the type, No. 25207, U.S.N.M., collected by R. E. Earll at Charleston, South Carolina.	
901. <i>Urophycis tenuis</i> (Mitchill)	2555
Drawing by H. L. Todd from No. 21029, U.S.N.M., collected by the <i>Speedwell</i> in Halifax Harbor.	
902. <i>Urophycis chuss</i> (Walbaum)	2555
Drawing by H. L. Todd from No. 28707, U.S.N.M., collected by the <i>Fish Hawk</i> at Station 918, in N. lat. $40^{\circ} 20' 24''$, W. long. $70^{\circ} 41' 30''$, in 245 fathoms.	

PLATE CCCLXVI.

903. <i>Urophycis chesteri</i> Goode and Bean	2556
Drawing by H. L. Todd from No. 21840, U.S.N.M., collected by the <i>Speedwell</i> at Station 174, off Cape Ann, in 140 fathoms.	

904. *Len...*
D...905. *Len...*
Dr...
V...906. *Gair...*
Dr...
N...907. *Euch...*
Dra...
by...908. *Bathy...*
Dra...
of...909. *Steind...*
Draw...
col...
88...910. *Chalin...*
Draw...
by...
70...911. *Coryp...*
Draw...
Bl...912. *Hymer...*
Draw...
col...
W...913. *Cælorh...*
Draw...
cell...
W...914. *Cælorh...*
Draw...
by...
372...

	Text page.
904. <i>Lemonema barbatulum</i> Goode and Bean	2556
Drawing by W. S. Haines from No. 38331, U.S.N.M., collected by the <i>Albatross</i> at Station 2397 in N. lat. $28^{\circ} 42'$, W. long. $86^{\circ} 36'$, in 280 fathoms.	
905. <i>Lemonema melanurum</i> Goode and Bean	2557
Drawing by W. S. Haines from the type, No. 38270, U.S.N.M., collected by the <i>Albatross</i> at Station 2415 in N. lat. $30^{\circ} 44'$, W. long. $79^{\circ} 26'$, in 440 fathoms.	

PLATE CCCLXVII.

906. <i>Gairdropsarus argentatus</i> (Rheinhardt)	2559
Drawing by H. L. Todd from No. 7212, U.S.N.M., collected at Nahant, Massachusetts, by F. W. Putnam.	
907. <i>Euchelyopus cimbrinus</i> (Linneus)	2560
Drawing by H. L. Todd from No. 21721, U.S.N.M., collected by Edward Brown in Bay Chaleur.	
908. <i>Bathygadus favosus</i> Goode and Bean	2565
Drawing by H. L. Todd from the type collected by the <i>Blake</i> off Martinique, in 472 fathoms.	

PLATE CCCLXVIII.

909. <i>Steindachneria argentea</i> Goode and Bean	2568
Drawing by H. L. Todd from the type, No. 37350, U.S.N.M., collected by the <i>Albatross</i> in N. lat. $39^{\circ} 14' 30''$, W. long. $88^{\circ} 9' 30''$, in 68 fathoms.	
910. <i>Chalinura simula</i> Goode and Bean	2578
Drawing by H. L. Todd from No. 39152, U.S.N.M., collected by the <i>Albatross</i> at Station 2095 in N. lat. $39^{\circ} 29'$, W. long. $70^{\circ} 58' 40''$, in 1,342 fathoms.	
911. <i>Coryphaenoides carapinus</i> Goode and Bean	2579
Drawing by H. L. Todd from a specimen collected by the <i>Blake</i> in N. lat. $39^{\circ} 43'$, W. long. $70^{\circ} 55' 25''$.	

PLATE CCCLXIX.

912. <i>Hymenocephalus cavernosus</i> (Goode and Bean)	2580
Drawing by S. F. Denton from the type, No. 37337, U.S.N.M., collected by the <i>Albatross</i> at Station 2398 in N. lat. $28^{\circ} 45'$ W. long. $86^{\circ} 26'$, in 227 fathoms.	
913. <i>Cælorhynchus occa</i> (Goode and Bean)	2588
Drawing by H. L. Todd from the type No. 37334, U.S.N.M., collected by the <i>Albatross</i> at Station 2396 in N. lat. $28^{\circ} 34'$, W. long. $86^{\circ} 48'$, in 335 fathoms.	
914. <i>Cælorhynchus carminatus</i> (Goode)	2588
Drawing by H. L. Todd from No. 26187, U.S.N.M., collected by the <i>Fish Hawk</i> at Station 893, off Martha's Vineyard, in 372 fathoms.	

PLATE CCCLXX.

- Text page.
915. *Coelorhynchus caribbaeus* (Goode and Bean)..... 2589
 Drawing by H. L. Todd from the type, No. 37333, U.S.N.M.,
 collected by the *Albatross* at Station 2377, in the northern
 part of the Gulf of Mexico, in N. lat. $29^{\circ} 7' 30''$, W. long.
 $88^{\circ} 8'$, in 210 fathoms.
916. *Regalecus glesne* (Ascanius)..... 2596
 Drawing from Day, Fishes of Great Britain and Ireland,
 pl. 64.

PLATE CCCLXXI.

917. *Atheresthes stomias* (Jordan and Gilbert)..... 2600
 Drawing by A. H. Baldwin from No. 27186, U.S.N.M., col-
 lected by Dr. Jordan at Point Reyes, California.
918. *Hippoglossus hippoglossus* (Linnaeus)..... 2611
 Drawing by H. L. Todd from No. 27605, U.S.N.M., collected
 at Marmot Island, Alaska, by Dr. Bean.

PLATE CCCLXXII.

919. *Hippoglossoides platessoides* (Fabricius) 2614
 Drawing by H. L. Todd from No. 21002, U.S.N.M., collected
 at Le Have Bank.
920. *Hippoglossoides elassodon* Jordan and Gilbert..... 2615

PLATE CCCLXXIII.

921. *Psettichthys melanostictus* Girard..... 2618
 Drawing by H. L. Todd from No. 24167, U.S.N.M., collected
 at San Francisco, California, by Dr. Jordan.
922. *Paralichthys dentatus* (Linnæus)..... 2629
 Drawing by A. H. Baldwin from a specimen collected by
 Dr. H. M. Smith at St. George Island, Maryland.

PLATE CCCLXXIV.


923. *Paralichthys squamilentus* Jordan and Gilbert..... 2631
 Drawing by H. L. Todd from No. 30862, U.S.N.M., collected
 at Pensacola, Florida, by Jordan and Stearns.
924. *Paralichthys oblongus* (Mitchill) 2632
 Drawing by H. L. Todd.

PLATE CCCLXXV.


925. *Ancylopsetta quadrocellata* Gill..... 2634
 926. *Pleuronichthys decurrens* Jordan and Gilbert..... 2637
 Drawing by A. H. Baldwin from No. 27115, U.S.N.M., col-
 lected by Dr. Jordan at San Francisco, California.




1



2



3





4

1. BRANCHIOSTOMA CARIBEUM. (P. 3.)


2. TAIL OF POLYPTERUS BICHIR.

3. PETROMYZON MARINUS. (P. 10.)

4. ENTOSPHENUS TRIDENTATUS. (P. 12.)



5. LAM.
6. LAM.
7. NOT.
8. HEX.




5

63




6





7

4




8


- * 5. *LAMPETRA AUREA*. (P. 13.) *Yellow Lamprey*
6. *LAMPETRA SPADICEA*. (P. 13.)
7. *NOTORHYNCHUS MACULATUS*. (P. 17.) *Cow Head*
8. *HEXANCHUS GRISEUS*. (P. 19.)




9. GYROPI
10. HETERO
11. SCYLLO
12. CATUL




9



10



11



12


9. GYROPLEURODUS FRANCISCI. (P. 20.)

10. HETEROERCAL TAIL OF HETERODONTUS PHILIPPI.


11. SCYLLIORHINUS PROFUNDORUM. (P. 22.) *francisci*

12. CATULUS UTER. (P. 25.)







13



14



15






16

13. GINGLYMOSSTOMA CIRRATUM. (P. 26.)


14. PSEUDOTRIAKIS MICRODON. (P. 27.)

15. GALEORHINUS ZYOPTERUS. (P. 32.) *Ogilby's Blue shark*16. PRIONACE GLAUCA. (P. 33.) *Blue shark*







16a



17




18




19

- 16a. *PRIONACE GLAUCA*. (P. 33.)
17. *CARCHARHINUS LAMIA*. (P. 38.)
18. *HYPOPRION BREVIROSTRIS*. (P. 41.)
19. *SPIYRNA TIBURO*. (P. 44.)






20




21




22

- 16 20. ALOPIAS VULPES. (P. 45.) *Zanclorhynchus vulpes*
17 21. ISURUS DEKAYI. (P. 48.) *Porbeagle*.
17 22. LAMNA CORNUBICA. (P. 49.) *Mackerel*
Shark






23



24




24a

23. *CETORHINUS MAXIMUS*. (P. 51.) *Great White*
24, 24a. *SQUALUS ACANTHIAS*. (P. 54.) *Black Dog*



25
26
27
28



25



26





27




28

25. *CENTROSCYMNUS COELOLEPIS*. (P. 55.) *Ahi* - Dogfish
26. *CENTROSCYLLIUM FABRICII*. (P. 56.) *Spiri* - Dogfish
27. *PRISTIS PECTINATUS*. (P. 60.) *Corna de la muerte*
28. *RHINOBATUS LENTIGINOSUS*. (P. 62.)






28a





28b



29


28a, 28b. *RHINOBATUS LENTIGINOSUS*. (P. 62.)
29. *RAJA ERINACEA*. (P. 68.) *Common Skate*






30


15




31

27 30. *RAJA OCCELLATA*. (P. 68.) *Big Skate*
31. *RAJA ACKLEYI*. (P. 70.)






32




33

32. RAJA STELLULATA. (P. 75.)
33. TETRONARCE OCCIDENTALIS. (P. 77.)





34




34a


34, 34a. *TETRONARCE CALIFORNICA*. (P. 77.)

U.S NATIONAL





35




35a


35, 35a. *NARCINE BRASILIENSIS.* (P. 78.)

U.S. NATION





36




36a

36, 36a. *DASYATIS SABINA*. (P. 84.)

U.S. NATION






37


37. *AETORATUS NARINAЕ.* (P. 88.)





37a. *AETOBATUS KARNIARI.* (P. 88.)

38. I
38a. E





38

38a

38. RHINOPTERA STEINDACHNERI. (P. 91.)
38a. PAVED TEETH OF RHINOPTERA STEINDACHNERI. (P. 91.)


U.S. NATIONAL







39

39. *MANTA BIROSTRIS.* (P. 92.)






40



41

19




42

40. CHIMAERA AFFINIS. (P. 95.) *Leptorhynchidae*.
41. HYDROLAGUS COLLIEI. (P. 95.) *Lophotidae*.
42. HARRIOTTA RALEIGHANA. (P. 96.)




43 43, 43
44, 11
45, A




43

20




43a

20



44

21



45

22

- 43, 43a. *POLYODON SPATHULA*. (P. 101.) *Paddlefish*
44. HETEROERCERAL TAIL OF *ACIPENSER TRANSMONTANUS*. (P. 104.) *White Sturgeon*
45. *ACIPENSER STURIO OXYRHYNCHUS*. (P. 105.) *Common Sturgeon*





46


12
62

47


12
64

48

46. *ACIPENSER RUBICUNDUS*. (P. 106.) *Lake Sturgeon*.
47. *ACIPENSER BREVIROSTRUM*. (P. 106.) *Short-nosed Sturgeon*.
48. *SCAPHIRHYNCHUS PLATIRYNCHUS*. (P. 107.)




4
5
5
5
5
4
4




49

25-




50

25



51

1




51a


25

49. *LEPISTOEVUS PLATOSTOMUS*. (P. 110.) *Short-nosed Garpike*
50. *LEPISTOEVUS TRISTECHUS*. (P. 111.)
51. *AMIA CALVA*; female. (P. 113.) *Bowfin; Dogfish*
51a. HETEROERCAL TAIL OF *AMIA CALVA*. (P. 113.)





52





53

52. *FELICHTHYS MARINUS*. (P. 118.)
53. *GALEICHTHYS FELIS*. (P. 128.)


U.S. NATIONAL






54. *GALEICHTHYS GILBERTI*. (P. 2773.)
55. *GALEICHTHYS AZUREUS*. (P. 2775.)






56



57




58


56. *ICTALURUS FURCATUS.* (P. 134.)57. *ICTALURUS ANGUILLA.* (P. 2788.)58. *ICTALURUS PUNCTATUS.* (P. 134.)

blue catfish







59



59a




59b




60


28

59, 59a, 59b. *AMEIURUS DUGESII*. (P. 138.)
60. *AMEIURUS MELAS*. (P. 141.) *Black Bullhead*







61



62




63




64

61. *AMEIURUS PLATYCEPHALUS*. (P. 142.)
62. *LEPTOPS OLIVARIS*. (P. 143.)
63. *NOTURUS FLAVUS*. (P. 144.) *Stone batfish*
64. *SCHILBEODES NOCTURNUS*. (P. 146.)




67a


65,
66,
67,




65




66



67



67a




67b

65. SCHILBEODES EXILIS. (P. 147.)


66. SCHILBEODES INSIGNIS. (P. 147.)

67, 67a, 67b. SCHILBEODES GILBERTI. (P. 148.)






68




69




68. *SCHILBEODES MURUS*. (P. 148.)
69, 69a, 69b. *SCHILBEODES FURIOSUS*. (P. 149.)






70





71



72


1.2 70. *ICTIOBUS CYPRINELLA.* (P. 163.) Common Buffalo Fish.
71. *CARPIOIDES CYPRINUS.* (P. 168.)
72. *CYCLEPTUS ELONGATUS.* (P. 168.)






73

31




74






75


22

73. *PANTOSTEUS JORDANI*. (P. 171.) Mountain Sucker.
74. *CATOSTOMUS LATIPINNIS*. (P. 174.)
75. *CATOSTOMUS GRISEUS*. (P. 175.) Gray Sucker.







76. *CATOSTOMUS POCATELLO.* (P. 175.)
77. *CATOSTOMUS CATOSTOMUS.* (P. 176.) *Northern Sucker*
78. *CATOSTOMUS TAOENSIS.* (P. 177.)




79. CATO
80. CATO
81. PHA
82. PHA



79



80




81



82

79. *CATOSTOMUS OCCIDENTALIS.* (P. 178.)80. *CATOSTOMUS TSILTCOENSIS.* (P. 279.)81. PHARYNGEAL TEETH OF *CATOSTOMUS MACROCHEILUS.* (P. 178.) *Giving River*82. PHARYNGEAL TEETH OF *PLACOPHARYNX DUQUESNII.* (P. 198.)





35




83. *CATOSTOMUS COMMERSONII*. (P. 178.)


84. *CATOSTOMUS ARDENSI*. (P. 179.)

85. *CHASMISTES LIORUS*. (P. 183.)






86






87





88

86. CHASMISTES STOMIAS. (P. 2794.)
87. CHASMISTES COPEI. (P. 2795.)
88. XYRAUCHEN CYPHO. (P. 184.)




89. *ERIMYZON SUCETTA*. (P. 185.)90. *MINYTREMA MELANOPS*. (P. 187.)91. *MONOSTOMA CONGESTUM*. (P. 192.)






92




93

92. MOXOSTOMA AUSTRINUM. (P. 192.)
93. MOXOSTOMA RUPISARTES. (P. 196.)







94



94a


94, 94a. *LAGOCHILA LACERA.* (P. 199.)






95

38



96


39




97

95. *CAMPOSTOMA ANOMALUM*. (P. 205.) *Stone Perch*; *Atom Logger*
96. *ORTHODON MICROLEPIDOTUS*. (P. 207.)
97. *ACROCHEILUS ALUTACEUS*. (P. 208.) *Eelinch-mont*






98




99




100

98. ALGANSEA DUGESI. (P. 211.)
99. MYLOCHEILUS CAURINUS. (P. 219.) *Lobes. Rio. black.*
100. SEMOTILUS ATROMACULATUS. (P. 222.) *Breath. black. Horned. 2 nose.*






101



102




103


101. *Ptychocheilus oregonensis*. (P. 224.)
102. *Leuciscus bicolor*. (P. 232.)
103. *Leuciscus lineatus*. (P. 232.)




D1 L 105
105 L 105
105 L 105



104



105




105a

104. *LEUCISCUS HYDROPHILUS*. (P. 238.)


105. *LEUCISCUS BALTEATUS*. (P. 238.)

105^a. *LEUCISCUS BALTEATUS*, FROM TYPE OF *LEUCISCUS GILLI*. (P. 238.)





106




107


106. *LEUCISCUS SIUSLAWI*. (P. 2797.)

107. *RUTILUS BICOLOR*. (P. 244.)






108






109



110

108. OPSOPOEODUS OSCULUS. (P. 248.)
109. OPSOPOEODUS EMILLE. (P. 248.)
110. OPSOPOEODUS BOLLMANI. (P. 249.)







111. *ABRAMIS CRYSOLEUCAS*. (P. 250.)


112. *NOTROPIS AZTECUS*. (P. 258.)

113. *NOTROPIS WELAKA*. (P. 2799.)






114



115



116

114. NOTROPIS CAYUGA ATROCAUDALIS. (P. 260.)

115. NOTROPIS KANAWHA. (P. 264.)

116. NOTROPIS CHIHUAHUA. (P. 265.)









IMAGE EVALUATION TEST TARGET (MT-3)



6"





Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503


EE
44 28
43 25
42 32
41 36
40 2.2
39 2.0
38 1.8

EE
11 10
10 9






117



118




119


117. NOTROPIS NUX. (P. 267.)

118. NOTROPIS NOOCOMIS. (P. 268.)


119. NOTROPIS HUDSONIUS. (P. 269.)

116 *sharpened : by tail*
more






120



121





122

120. NOTROPIS HODSONIUS SALUDANUS. (P. 270.)


121. NOTROPIS WHIPPLEI. (P. 278.)

122. NOTROPIS GALACTURUS. (P. 279.)






123



124





125

123. *NOTROPIS MACDONALDI*. (P. 284.)


124. *NOTROPIS COCCOGENIS*. (P. 284.)

125. *NOTROPIS SWAINI*. (P. 290.)






126



127





128

126. *NOTROPIS TELESCOPUS*. (P. 292.)


127. *NOTROPIS NOTEMIGONOIDES*. (P. 292.)

128. *NOTROPIS METALLICUS*. (P. 297.)






129





130




131

129. ERICYMBIA BUCCATA. (P. 302.)
130. PHENACOBIA URANOPS. (P. 304.)
131. EVARRA EIGENMANNI. (P. 304.)






132







133




134


132. RHINICHTHYS DULCIS. (P. 306.)
133. AGOSIA KLAMATHENSIS. (P. ——.)
134. AGOSIA UMATILLA. (P. 313.)






135. *AGOSIA FALCATA.* (P. 313.)
136. *HYBOPSIS AESTIVALIS MARCONIS.* (P. 316.)
137. *HYBOPSIS WATAUGA.* (P. 319.)






138



139

51




140


52

138. HYBOPSIS ALTUS. (P. 321.)


139. PLATYGOBIO GRACILIS. (P. 326.)

140. EXOGLOSSUM MAXILLINGUA. (P. 327.)






141





142



143


141. *BRYCON DENTEX*. (P. 337.)
142. *DERICHTHYS SERPENTINUS*. (P. 343.)
143. *ANGUILA CHRYSOPA*. (P. 348.)





144

54




145




146


55

144. *SIMENICHELYS PARÁSITICUS.* (P. 349.)
145. *ECHIODON BRUNNEUS.* (P. 350.)
146. *SYNAPOBRANCHUS PINNATUS.* (P. 351.)






147




148




149

147. *HISTIOBRANCHUS INFERNALIS.* (P. 352.)148. *LEPTOCEPHALUS CONGER.* (P. 354.)149. *LEPTOCEPHALUS CAUDILIMBATUS.* (P. 355.)







150



151





152



153

150. *CONGRELLUS FLAVUS*. (P. 357.)
151. *HOPLUUNIS SCHMIDTI*. (P. 361.)
152. *VENEFICA PROCERA*. (P. 365.)
153. *SERRIVOMER BEANII*. (P. 367.)






154




154a



154b



155




156

154. *AVOCETTINA GILLI.* (Pp. 367, 2801.)
154a, 154b. *AVOCETTINA GILLI.* (P. 2801.)
155. *LABICHTHYS CARINATUS.* (P. 368.)
156. *LABICHTHYS ELONGATUS.* (P. 369.)







157




157a



157b




158




159


157, 157a, 157b. *NEMICHTHYS AVOCETTA*. (P. 369.)
158. *AHLIA EG Montis*. (P. 370.)

159, 159a. *VERMA KENDALLI*. (P. 375.)







160




161

160. *LETHARCHUS VELIFER.* (P. 375.)
161. *MYRICHTHYS TIGRINUS.* (P. 376.)







162




163

162. *MYRICHTHYS PANTOSTIGMIUS.* (P. 2802.)
163. *PISOODONOPHIS CRUENTIFER.* (P. 377.)






164




165




166a


166

164. CALECHELYS MURENA. (P. 378.)
165. BASCANICHTHYS SCUTICARIS. (P. 378.)
166, 166a. BASCANICHTHYS PENINSULÆ. (P. 379.)






167





168




169

167. *QASSIREMUS EVIONTHAS.* (P. 380.)168. *OPHICHTHUS GUTTIFER.* (P. 383.)169. *OPHICHTHUS OCELLATUS.* (P. 383.)






170





171




172

170. *MYSTRIOPHIS INTERTINCTUS.* (P. 386.)
171. *LYCODONTIS MORINGA.* (P. 395.)
172. *MURENA INSULARUM.* (P. 400.)






173




174




175

173. MURENA RETIFERA. (P. 401.)
174. CHANOMURENA VITTATA. (P. 404.)
175. SACCOPHARYNX AMPULLACEUS. (P. 406.)






176




177




178

176. *GASTROSTOMUS BAIRDII*. (P. 406.)
177. *TARPON ATLANTICUS*. (P. 409.)
178. *ELOPS SAURUS*. (P. 410.)






179





180




181

179. *ALBULA VULPES.* (P. 411.)
180. *HIODON TERGISUS.* (P. 413.)
181. *HIODON SELENOPOD.* (P. 414.)






182





183




184

182. *CHANOS CHANOS*. (P. 414.)
183. *DOROSOMA CEPEDIANUM*. (P. 416.)
184. *SIGNALOSA ATCHAFALAY.E.* (P. 2809.)






185





186




187

185. *CLUPEA HARENGUS.* (P. 421.)
186. *CLUPEA PALLASII.* (P. 422.)
187. *POMOLOBUS CHRYSICHLORIS.* (P. 425.)






188





189




190

188. *POMOLOBUS MEDIOCRISS.* (P. 425.)
189. *POMOLOBUS PSEUDOHARENGUS.* (P. 426.)
190. *POMOLOBUS FESTIVALIS.* (P. 426.)






191





192




192a

191. *ALOSA SAPIDISSIMA*. (P. 427.)
192. *ALOSA ALABAMÆ*; male. (P. 2810.)
192a. *ALOSA ALABAMÆ*; female. (P. 2810.)






193




194




195

193. *SARDINELLA SARDINA*. (P. 430.)
194. *SARDINELLA STOLIFERA*. (P. 431.)
195. *BREVOORTIA TYRANNUS*. (P. 433.)






196





197




198

196. *STOLEPHORUS PERTHECATUS*. (P. 442.)
197. *ALEPOCEPHALUS AGASSIZII*. (P. 453.)
198. *TALISMANIA ANTILLARUM*. (P. 455.)






199





200




200a

199. *ALEPOSOMUS COPEI*. (P. 459.)200. *COREGONUS WILLIAMSONI*. (P. 463.)200a. *COREGONUS WILLIAMSONI*; young. (P. 463.)






201





202




203

201. *COREGONUS COULTERII*. (P. 462.)
202. *COREGONUS CLUPEIFORMIS*. (P. 465.)
203. *ARGYROSOMUS NIGRIPINNIS*. (P. 472.)






204





205




206

204. *STENODUS MACKENII.* (P. 474.)
205. *ONCORHYNCHUS GORBUSCHA.* (P. 478.)
206. *ONCORHYNCHUS TSCHAWYTSCHA.* (P. 479.)






207





207a




207b

207. *ONCORHYNCHUS NERKA*; adult male, large form. (P. 481.)
207a. *ONCORHYNCHUS NERKA*; adult female, small form. (P. 481.)
207b. *ONCORHYNCHUS NERKA*; adult mutilated male, small form. (P. 481.)






208





209




210

208. *SALMO CLARKII HENSHAWI*. (Pp. 493, 2819.)
209. *SALMO CLARKII VIRGINALIS*. (Pp. 495, 2819.)
210. *SALMO CLARKII SPILURUS*. (Pp. 495, 2819.)






211



212





213

211. *SALMO CLAEKII PLEURITICUS.* (Pp. 496, 2819.)


212. *SALMO CLARKII BOUVIERI.* (Pp. 496, 2819.)

213. *SALMO CLARKII STOMIAS.* (Pp. 497, 2819.)






214



215





216

214. *SALMO CLARKII MACDONALDI*. (Pp. 497, 2819.)


215. *SALMO GAIARDNERI*. (P. 497.)

216. *SALMO IRIDEUS*. (P. 500.)






217





218




219

217. *CRISTIVOMER NAMAYCUSH.* (P. 504.)
218. *SALVELINUS FONTINALIS.* (P. 506.)
219. *SALVELINUS MALMA.* (P. 507.)






220




221




222

220. *SALVELINUS ALPINUS AUREOLUS*. (P. 511.)
221. *SALVELINUS OQUASSA*. (P. 514.)
222. *THYMALLUS SIGNIFER*. (P. 517.)






223



224





224a

223. THYMALLUS TRICOLOR. (Pp. 518, 2871.)


224. THYMALLUS TRICOLOR MONTANUS. (Pp. 519, 2871.)

224a. THYMALLUS TRICOLOR MONTANUS; young. (Pp. 519, 2871.)






225





226




227

225. *MALLOTUS VILLOSUS*. (P. 520.)
226. *THALEICHTHYS PACIFICUS*. (P. 521.)
227. *OSMERUS THALEICHTHYS*. (P. 522.)






228





229




230

228. *OSMERUS MORDAX*. (P. 523.)
229. *OSMERUS DENTEX*. (P. 524.)
230. *HYPOMESUS PRETIOSUS*. (P. 525.)






231




232




233

231. HYPOMESUS OLIDUS. (P. 525.)
232. ARGENTINA SILUS. (P. 526.)
233. LEUROGLOSSUS STILBIUS. (P. 527.)






234





235




236

234. *BATHYLAGUS BENEDICTI*. (P. 529.)
235. *TRACHINOCEPHALUS MYOPS*. (P. 533.)
236. *SYNODUS FOETENS*. (P. 538.)






237



238





239

237. *BENTHOSAURUS OBALJATOR*. (P. 543.)

238. *BATHYPTEROIS QUADRIFILIS*. (P. 545.)

239. *IPNOTOPS MURRAYI*. (P. 547.)






240



241





242

240. RONDELETIA BICOLOR. (P. 548.)


241. CETOMIMUS GILLII. (P. 549.)

242. CERATOSCOPELUS MADEIRENSIS. (P. 557.)






243



244





245

243. LAMPANYCTUS CROCODILUS. (P. 558.)


244. LAMPADENA SPECULIGERA. (P. 561.)

245. NANNOBRACHIUM MACDONALDI. (P. 563.)






246





247




248

246. *AETHOPRORA LUCIDA*. (P. 565.)
247. *MYCTOPHUM OPALINUM*. (P. 571.)
248. *TARLETONBEANIA TENUA*. (P. 575.)






249



250




251

249. *YARRELLA BLACKFORDI*. (P. 584.)
250. *CHAULIODUS SLOANEI*. (P. 585.)
251. *ASTRONESTHES GEMMIFER*. (P. 586.)



**IMAGE EVALUATION
TEST TARGET (MT-3)**




**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

45
28
32
25
22
20
18

oil





252




253




254

252. *ASTRONESTHES RICHARDSONI*. (P. 587.)253. *STOMIAS FEROX*. (P. 588.)254. *GRAMMATOSTOMIAS DENTATUS*. (P. 590.)






255





256




257

255. PHOTONECTES GRACILIS. (P. 591.)
256. MALACOSTEUS NIGER. (P. 593.)
257. ALEPISaurus FEROX. (P. 595.)






258



259




260

258. *ALEPISaurus ESCULAPIUS.* (P. 595.)


259. *ARCTOZENUS CORUSCANS.* (P. 601.)

260. *PARALEPIS COREGONOIDES.* (P. 602.)






261




261a

261, 261a. ARGYROPELECUS OLFFERSI. (P. 604.)







262



263



264



265


262. ALDROVANDIA MACROCHIR. (P. 609.)

263. ALDROVANDIA GRACILIS. (P. 610.)


264. NOTACANTHUS ANALIS. (P. 615.)

265. NOTACANTHUS PHASGANORUS. (P. 616.)






266




267




268

266. LIOPENYS GILLII. (P. 619.)
267. DALLIA PECTORALIS. (P. 621.)
268. UMBRA PYMÆA. (P. 624.)






269





269a




270

269, 269a. *LUCIUS LUCIUS*. (P. 628.)
270. *LUCIUS MASQUINONGY*. (P. 629.)






271



271a





271b

271. *FUNDULUS MAJALIS*; female. (P. 639.)


271a. *FUNDULUS MAJALIS*; male. (P. 639.)

271b. *FUNDULUS MAJALIS*; young. (P. 639.)






272





273




274

272. *FUNDULUS PALLIDUS.* (P. 638.)273. *FUNDULUS HETEROCLITUS*; male. (P. 640.)274. *FUNDULUS OCCELLARIS.* (P. 642.)






275



275a




276

275. *FUNDULUS DIAPHANUS*; male. (P. 645.)


275a. *FUNDULUS DIAPHANUS*; female. (P. 645.)

276. *FUNDULUS ZEBRINUS*. (P. 646.)






277



278





278a

277. *FUNDULUS SEMINOLIS*. (P. 647.)


278. *FUNDULUS CATENATUS*; male. (P. 648.)

278a. *FUNDULUS CATENATUS*; female. (P. 648.)






279




280



281


279. *FUNDULUS STELLIFER.* (P. 648.)280. *FUNDULUS RATHBUNI.* (P. 649.)281. *FUNDULUS ALBOLINEATUS.* (P. 649.)






1/5 in.

282



283




284


1/5 in.

282. *FUNDULUS FUNDULOIDES*. (P. 650.)
283. *FUNDULUS MACDONALDI*. (P. 651.)
284. *FUNDULUS JENKINSI*. (P. 651.)






285




286




287

285. *FUNDULUS PULVEREUS*. (P. 652.)
286. *FUNDULUS LUCLE*. (P. 654.)
287. *FUNDULUS CHRYSOTUS*. (P. 655.)






288





289




290

288. *FUNDULUS NOTATUS*. (P. 656.)
289. *FUNDULUS NOTATUS*. (P. 659.)
290. *ADINIA DUGESII*. (P. 661.)






291









292



293


291. *FUNDULUS GOODEI*. (Pp. 664, 2831.)292. *LUCANIA PARVA*. (P. 665.)293. *CHARACODON BILINEATUS*. (P. 668.)






294, 294a, 294b, 294c, 294d. *EMPETRICHTHYS MERRIAMI.* (P. 667.)






295





295a




296

295. CHARACODON VARIATUS. (P. 669.)
295a. TEETH OF CHARACODON VARIATUS. (P. 669.)
296. CYPRINODON VARIEGATUS. (P. 671.)






296a



297







298

296a. *CYPRINODON VARIEGATUS*; young. (P. 671.)

297. *CYPRINODON CARPIO*. (P. 675.)

298. *JORDANELLA FLORIDE*. (P. 677.)






299. GAMBUSIA AFFINIS; male. (P. 680.)


299a. GAMBUSIA AFFINIS; female. (P. 680.)

300. ANABLEPS DOVI. (P. 685.)






301







302




303

301. *GOODEA ATRIPINNIS*. (P. 685.)
302. *HETERANDRIA FORMOSA*. (P. 687.)
303. *POECILIA PRESIDIONIS*; female. (P. 697.)






304. *XIPHOPHORUS HELLERI*. (P. 701.)
305. *CHOLOGASTER CORNUTUS*. (P. 703.)
306. *TYPHLICHTHYS SUBTERRANEUS*. (P. 704.)
307. *AMBLYOPSIS SPELEUS*. (P. 706.)







308



309



310




311


308. *TYLOSURUS RAPHIDOMA.* (P. 715.)

309. *TYLOSURUS ACUS.* (P. 716.)


310. *CHRIODORUS ATERINOIDES.* (P. 719.)

311. *HYPORHAMPHUS UNIFASCIATUS.* (P. 720.)







312



313





314




315

312. *HYPORHAMPHUS ROBERTI*. (P. 721.)
313. *HEMIRAMPHUS BRASILIENSIS*. (P. 722.)
314. *SOMBRESOX SAURUS*. (P. 725.)
315. *FODIATOR ACUTUS*. (P. 728.)






316





317




318

316. *EXONAUTES EXSILIENS*. (Pp. 732, 2830.)
317. *EXONAUTES RONDELETHII*. (Pp. 733, 2830.)
318. *EXOCETUS VOLITANS*. (P. 734.)






319




320



321

319. *CYP SILURUS CALIFORNICUS*. (Pp. 740, 2830.)
320. *GASTEROSTEUS ACULEATUS*. (P. 747.)
321. *GASTEROSTEUS CATAPHRACTUS*. (P. 749.)






322




323



324




325




326

322. *APELTES QUADRACUS*. (P. 752.)
323. *AULORHYNCHUS FLAVIDUS*. (P. 754.)
324. *AULOSTOMUS MACULATUS*. (P. 754.)
325. *SIPHOSTOMA STARKII*. (P. 771.)
326. *CORYTHROICHTHYS CAYORUM*. (P. 2838.)






327




328






329


1/2 in.


327. *Hippocampus hudsonius*. (P. 777.)
328. *Hippocampus zosterae*. (P. 778.)
329. *Percopsis guttatus*. (P. 784.)






330. *COLUMBIA TRANSMONTANA*. (P. 784.)
331. *APHREDODERUS SAYANUS*. (P. 786.)
332. *ATHERINA STIPES*. (P. 790.)






333





334




335

333. *ATHERINA AREA.* (P. 790.)334. *CHIROSTOMA HUMBOLDTIANUM.* (P. 793.)335. *ESLOPSARUM JORDANI.* (Pp. 793, 2840.)






336




337




338

336. *KIRTLANDIA VAGRANS.* (P. 794.)337. *MENIDIA PENINSULE.* (P. 797.)338. *MENIDIA GRACILIS BERYLLINA.* (P. 797.)






339



340




341

339. EURYSTOLE ERIARCHA. (P. 803.)


340. THYRINA EVERMANNI. (P. 804.)

341. AATHERINOPSIS CALIFORNIENSIS. (P. 806.)






342




343




344

342. *ATHERINOPS AFFINIS.* (P. 807.)343. *MUGIL CEPHALUS.* (P. 811.)344. *MUGIL CUREMA.* (P. 813.)






345






346





347

345. *MUGIL HOSPES.* (P. 814.)
346. *CHÆNOMUGIL PROBOSCIDEUS.* (P. 816.)
347. *AGONOSTOMUS MONTICOLA.* (P. 819.)




348. *JOTURUS PICHARDI.* (P. 821.)349. *SPHYRENA BARRACUDA.* (Pp. 823, 2841.)350. *POLYDACTYLUS OCTONEMUS.* (P. 830.)






351




352




353

351. *AMMODYTES AMERICANUS.* (P. 833.)
352. *BATHYCLUPEA ARGENTEA.* (P. 835.)
353. *STEPHANOBERYX MONÆ.* (P. 836.)






354




355




356

354. *HOPLOSTETHUS MEDITERRANEUS.* (P. 837.)
355. *PLECTROMUS SUBORBITALIS.* (P. 841.)
356. *PLECTROMUS CRASSICEPS.* (P. 843.)






357




358




359

357. *BERYX SPLENDENS.* (P. 844.)358. *HOLOCENTRUS ASCENSIONIS.* (P. 848.)359. *FLAMMEO MARIANUS.* (Pp. 852, 2871.)






360




361




362

360. *MULLUS AURATUS.* (P. 856.)
361. *MULLOIDES RATHBUNI.* (P. 857.)
362. *UPENEUS MACULATUS.* (P. 858.)






363






364



365


363. SCOMBER SCOMBRUS. (P. 865.)
364. SCOMBER COLIAS. (P. 866.)
365. AUXIS THAZARD. (P. 867.)






366. *GYMNO SARD A. ALLETERATA.* (P. 869.)
367. *GERMO ALALUNGA.* (P. 871.)
368. *SCOMBEROMORUS MACULATUS.* (P. 874.)






369




370



371

369. *SCOMBEROMORUS REGALIS.* (P. 875.)
370. *ESCOLAR VIOLACEUS.* (Pp. 878, 2843.)
371. *EPINNULA MAGISTRALIS.* (P. 880.)






372




373




374

372. *EVOXYMETOPON TENIATUS*. (P. 886.)
373. *LEPIDOPUS CAUDATUS*. (P. 886.)
374. *BENTHODESMUS ATLANTICUS*. (P. 887.)

4. 110




375




376

375. *TRICHIURUS LEPTURUS*. (P. 889.)
376. *ISTIOPHORUS NIGRICANS*. (P. 891.)






377




378

377. *NEMATISTIUS PECTORALIS.* (P. 895.)
378. *OLIGOPLITES SAURUS.* (P. 898.)






379




380




381

379. NAUCRATES DUCTOR. (P. 900.)
380. SERIOLA DORSALIS. (P. 902.)
381. SERIOLA ZONATA. (P. 902.)






382



383



384

382. *SERIOLA LALANDI.* (P. 903.)
383. *DECAPTERUS MACARELLUS.* (P. 909.)
384. *TRACHURUS TRACHURUS.* (P. 910.)









IMAGE EVALUATION TEST TARGET (MT-3)



6"




Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503


45
28
34
34
22
25
20
18

Oil






385




386




387

385. *TRACHUROPS CRUMENOPHTHALMUS*. (P. 911.)
386. *HEMICARANX AMBLYRHYNCHUS*. (P. 912.)
387. *CARANX HIPPOS*. (P. 920.)






388




389

388. *CARANX CRYOS.* (P. 921.)
389. *CARANX LATUS.* (P. 923.)






390




391

390. *CARANX MEDUSICOLA*. (P. 924.)
391. *HYNNIS HOPKINSI*. (P. 933.)






392




393

392. *VOMER SETIPINNIS*. (P. 934.)393. *SELENE VOMER*; young. (P. 936.)






393a




394

393a. *SELENE VOMER*; adult. (P. 936.)
394. *CHLOROSCOMBRUS CHRYSURUS*. (P. 938.)






395




396

395. *TRACHINOTUS GLAUCUS.* (P. 940.)
396. *TRACHINOTUS FALCATUS.* (P. 941.)






397




398

397. *TRACHINOTUS CULVERI.* (P. 942.)398. *TRACHINOTUS CAROLINUS.* (P. 944.)






399




400




401

399. *ZALOCYS STILBE*. (P. 284.)400. *POMATOMUS SALTATRIX*. (P. 946.)401. *RACHYCENTRON CANADUS*. (P. 948.)






402




403

402. *CORYPHENA HIPPURUS.* (P. 952.)
403. *CENTROLOPHUS NIGER.* (P. 963.)






404




405

404. *RHOMBUS PARU*. (P. 965.)405. *PORONOTUS TRIACANTHIUS*. (Pp. 967, 2849.)






406




407




408

406. *ICICHTHYS LOCKINGTONI*. (P. 969.)
407. *SCHEDOPHILUS MEDUSOPHAGUS*. (P. 970.)
408. *ACROTUS WILLOUGHBYI*. (P. 973.)






409



410





411

409. *ZAPRORA SILENUS*. (P. 2850.)


410. *GRAMMICOLEPIS BRACHIUSCUS*. (P. 974.)

411. *TETRAGONURUS CUVIERI*. (P. 976.)






412



413




414

412. PEMPHERIS MULLERI. (P. 978.)


413. PEMPHERIS POEYI. (P. 979.)

414. ELASSOMA EVERG. ADEI. (P. 982.)






415




416


415. POMOXIS ANNULARIS. (P. 987.)

416. POMOXIS SPAZOIDES. (P. 987.)







417




418

417. *CENTRARCHUS MACROPTERUS*. (P. 988.)
418. *ACANTHARCHUS POMOTIS*. (P. 989.)







419




419. AMBLOPLITES RUPESTRIS. (P. 990.)
A, B, C. SKULL OF AMBLOPLITES RUPESTRIS. (P. 990.)






420




421


420. *ARCHOPLITES INTERRUPTUS*. (P. 991.)

421. *CHENOBRYTTUS GULOSUS*. (P. 992.)







422




423

422. ENNEACANTHUS GLORIOSUS. (P. 993.)
423. MESOGONISTIUS CHAETODON. (P. 995.)






424



425





425a

424. APOMOTIS SYMMETRICUS. (P. 998.)


425. LEPOMIS AURITUS. (P. 1001.)

425a. SKULL OF LEPOMIS AURITUS. (P. 1001.)







426




427

426. LEPOMIS MEGALOTIS. (P. 1002.)
427. LEPOMIS PALLIDUS. (P. 1005.)







428




429

428. *EUPOMOTIS EURYORUS*. (P. 1008.)
429. *EUPOMOTIS GIBBOSUS*. (P. 1009.)






430




430a

430. *MICROPTERUS DOLOMIEU.* (P. 1011.)430a. VERTEBRAE OF *MICROPTERUS DOLOMIEU.* (P. 1011.)






431




432

431. MICROPTERUS SALMOIDES. (P. 1012.)
432. SKULL OF KUHLIA RUPESTRIS.






433



433a





434

433. STIZOSTEDION VITREUM. (P. 1021.)


433a. TAIL OF STIZOSTEDION VITREUM. (P. 1021.)

434. STIZOSTEDION CANADENSE. (P. 1022.)






435



436





436a

435. PERCA FLAVESCENS. (P. 1023.)


436. PERCINA CAPRODES. (P. 1026.)

436a. SKULL OF PERCINA CAPRODES. (P. 1026.)




 $\frac{1}{4} \text{ in.}$


437

 $\frac{1}{3} \text{ in.}$


438

437. *HADROPTERUS MACROCEPHALUS*. (P. 1031.)
438. *HADROPTERUS ASPRO*. (P. 1032.)






439




440




441

439. *HADROPTERUS GUNTHERI.* (P. 1033.)
440. *HADROPTERUS EVIDES.* (P. 1036.)
441. *HADROPTERUS SCIERUS.* (P. 1037.)






442



443




444

442. HYPOHOMUS AURANTIACUS. (P. 1040.)


443. HYPOHOMUS SPILOTUS. (P. 1043.)

444. COTTOGASTER SHUMARDI. (P. 1046.)






445





446




447

445. *COTTOGASTER CHENEYI*. (P. 2851.)
446. *ULOCENTRA GILBERTI*. (P. 1049.)
447. *ULOCENTRA MEADIE*. (P. 2852.)






448





449




450

448. *ULOCENTRA SIMOTERA*. (P. 1051.)
449. *DIPLESION BLENNIOIDES*. (P. 1053.)
450. *BOLEOSOMA NIGRUM*. (P. 1056.)






451




452




453

451. *BOLEOSOMA NIGRUM OLMISTEDI.* (P. 1057.)
452. *BOLEOSOMA CAMURUM.* (P. 1060.)
453. *CRYSTALLARIA ASPRELLA.* (P. 1061.)






454





455




456

454. AMMOCRYPTA PELLUCIDA CLARA. (P. 16.)
455. AMMOCRYPTA BRANII. (P. 1064.)
456. ETHEOSTOMA CAMURUM. (P. 1076.)






457



458




459

457. *ETHEOSTOMA CINEREUM*. (P. 1078.)


458. *ETHEOSTOMA JORDANI*. (P. 1079.)

459. *ETHEOSTOMA POTTERI*. (P. 1082.)






460




461




462

460. *ETHEOSTOMA IOW.E.* (P. 1083.)
461. *ETHEOSTOMA JESSE.E.* (P. 1084.)
462. *ETHEOSTOMA LEPIDOGENYS.* (P. 1087.)






463



464




465

463. *Etheostoma obeyense*. (P. 1092.)


464. *Etheostoma pagei*. (P. 1092.)

465. *Etheostoma virgatum*. (P. 1093.)






466



467

1/4 in.




468


1/5 in.

466. *ETHEOSTOMA JULIE.* (P. 1093.)
467. *PSYCHROMASTER TUSCUMBIA.* (P. 1100.)
468. *COPELANDELLUS QUIESCENS.* (P. 1100.)






469





470




471

469. *BOLEICHTHYS FUSIFORMIS.* (P. 1101.)470. *MICROPERCA FONTIOLA.* (P. 1104.)471. *APOGON RETROSELLA.* (P. 1108.)







472




473

472. APOGON PIGMENTARIUS. (P. 1109.)
473. APOGONICHTHYS ALUTUS. (P. 1110.)






474




475




476

474. *CHEILODIPTERUS AFFINIS.* (P. 1113.)475. *HYPOCLYDONIA BELLA.* (P. 1115.)476. *CENTROPOMUS UNDECIMALIS.* (P. 1118.)






477




478

477. *ROCCUS CHRYSOPS.* (P. 1132.)
478. *ROCCUS LINEATUS.* (P. 1132.)






479





480




480a

479. *MORONE AMERICANA*. (P. 1134.)480, 480a. SKULL OF *POLYPRION AMERICANUS*. (P. 1139.)






481




482

481. *BODIANUS FULVUS PUNCTATUS.* (P. 1146.)
482. *EPINEPHELUS ADSCENSIONIS.* (P. 1152.)







483




484

483. *EPINEPHELUS STRIATUS*. (P. 1157.)484. *EPINEPHELUS DRUMMOND-HAYI*. (P. 1159.)







485




486

485. *EPINEPHELUS MORIO.* (P. 1160.)
486. *GARRUPA NIGRITA.* (P. 1161.)






487




487a




487b

487, 487a, 487b. *PROMICROPS GUTTATUS.* (P. 1162.)






488





488a




489

488. *ALPIESTES AFER*. (P. 1164.)488a. SKULL OF *ALPIESTES AFER*. (P. 1164.)489. *DERMATOLEPIS ZANCLUS*. (P. 2854.)






490



491



492

490. *MYCTEROOPERCA BOULEGERI*. (P. 1171.)491. *MYCTEROOPERCA VENENOSA*. (P. 1172.)492. SKULL OF *MYCTEROOPERCA BONACI*. (P. 1174.)










IMAGE EVALUATION TEST TARGET (MT-3)



6"





Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503


EE
28
25
22
20
18

IT
oil






493





494




495

493. *MYCTEROPERCA JORDANI*. (P. 1176.)
494. *MYCTEROPERCA MICROLEPIIS*. (P. 1177.)
495. *MYCTEROPERCA FALCATA PHENAX*. (P. 1185.)







496




497

496. *HYPPOLECTRUS UNICOLOR NIGRICANS.* (P. 1193.)
497. *HYPPOLECTRUS GEMMA.* (P. 1193.)






498





499




500

498. *PARALABRAX MACULATOFASCIATUS.* (P. 1196.)
499. *PARALABRAX HUMERALIS.* (P. 1196.)
500. *CENTROPRISTES STRIATUS.* (P. 1199.)






501





502




503

501. *CENTROPRISTES PHILADELPHICUS*. (P. 1201.)
502. *DIPLECTRUM FORMOSUM*. (P. 1207.)
503. *PRIONODES BULLERI*. (P. 1213.)






504





505




506

504. *PARANTHIAS FURCIFER*. (P. 1221.)505. *HEMIANTHIAS VIVANUS* (P. 1223.)506. *PRONOTOGRAMMUS MULTIFASCIATUS*. (P. 1226.)







507




508

507. ANTHIAS ASPERILINGUIS. (P. 1227.)
508. GRAMMA LORETO. (P. 1229.)






509




510

509. *RYPTICUS BISTRISPINUS*. (P. 1233.)
510. *LOBOTES SURINAMENSIS*. (P. 1235.)






511




512


511. *PRIACANTHUS ARENATUS*. (P. 1237.)

512. *PSEUDOPRIACANTHUS ALTUS*. (P. 1239.)






513




514

513. *HOPLOPAGRUS GUNTHERI.* (P. 1244.)
514. *EVOPLITES VIRIDIS.* (P. 1246.)






515




516

515. *NEOMENIS APODUS.* (P. 1258.)
516. *NEOMENIS AYA.* (P. 1264.)





517




518


517. NEOMENIS ANALIS. (P. 1265.)

518. NEOMENIS SYNAGRIS. (P. 1270.)







519




520

519. *RABIRUBIA INERMIS.* (P. 1274.)
520. *OXYURUS CHRYSURUS.* (P. 1275.)






521




522

521. *RHOMBOPLITES AURORUBENS*. (P. 1277.)
522. *APSILUS DENTATUS*. (P. 1278.)






523




524

523. APRION MACROPIHTHALMUS. (P. 1280.)
524. ETELIS OCULATUS. (P. 1282.)






525




526

525. *VERILUS SORDIDUS*. (P. 1284.)
526. *XENOCYS JESSIE*. (P. 1285.)







527




528

527. *XENICHTHYS AGASSIZII*. (P. 1287.)
528. *HÆMULON ALBUM*. (P. 1295.)






529




530

529. HÆMULON MACROSTOMUM. (P. 1296.)
530. HÆMULON PARRA. (P. 1297.)






531




532

531. *HEMULON SCIURUS.* (P. 1303.)
532. *HEMULON PLUMIERI.* (P. 1304.)






533




534

533. *BRACHYGENYS CHRYSARGYREUS.* (P. 1307.)
534. *BATHYSTOMA RIMATOR.* (P. 1308.)






535




536

535. *BATHYSTOMA AUROLINEATUM*. (P. 1310.)
536. *LYTHRULON OPALESCENS*. (P. 1312.)






537




538

537. *ANISOTREMUS SURINAMENSIS*. (P. 1318.)
538. *ANISOTREMUS BILINEATUS*. (P. 1319.)






539




540

539. *ANISOTREMUS VIRGINICUS*. (P. 1322.)
540. *ORTHOPRISTIS REDDINGI*. (P. 1336.)






541




542

541. *ORTHOPRISTIS CHRYSOPTERUS.* (P. 1338.)
542. *MICROLEPIDOTUS INORNATUS.* (P. 1341.)







543




544

543. OTRYNTER CAPRINUS. (P. 1345.)
544. STENOTOMUS CHRYSOPS. (P. 1346.)







545




546

545. *STENOTOMUS ACULEATUS*. (P. 1346.)
546. *CALAMUS CALAMUS*. (P. 1349.)







547




548

547. *CALAMUS PRORIDENS*. (P. 1350.)
548. *CALAMUS BAJONADO*. (P. 1352.)







549




550

549. *CALAMUS PENNA.* (P. 1354.)
550. *CALAMUS ARCTIFRONS.* (P. 1355.)







551




552

551. *PAGRUS PAGRUS*. (P. 1356.)
552. *LAGODON RHOMBOIDES*. (P. 1358.)






553




554

553. *ARCHOSARGUS UNIMACULATUS.* (P. 1359.)
554. *ARCHOSARGUS PROBATOCEPHALUS.* (P. 1361.)







555




555a

555. *DIPLODUS HOLBROOKII*. (P. 1362.)
555a. *DIPLODUS HOLBROOKII*; young. (P. 1362.)






556




557

556. *XYSTEMA CINEREUM*. (P. 1372.)
557. *GERRES OLISTOSTOMUS*. (P. 1376.)






558




559

558. HERMOSILLA AZUREA. (P. 1383.)
559. KYPHOSUS SECTATRIX. (P. 1387.)






560





561




562

560. *MEDIALUNA CALIFORNIENSIS*. (P. 1391.)
561. *CYNOSCION NOTHUS*. (P. 1406.)
562. *CYNOSCION REGALIS*. (P. 1407.)






563




564




565

563. *CYNOSCIUS NEBULOSUS*. (P. 1409.)
564. *SAGENICHTHYS ANCYLODON*. (P. 1416.)
565. *LARIMUS ARGENTEUS*. (P. 1421.)






566




567

566. BAIRDIELLA CHRYSURA. (P. 1433.)
567. SCLÆNOPS OCCELLATUS. (P. 1453.)






568




569

568. RONCADOR STEARNSI. (P. 1457.)
569. LEIOSTOMUS XANTHURUS. (P. 1458.)






570




571

570. *MICROPOGON UNDULATUS*. (P. 1461.)
571. *UMBIRINA SINALOÆ*. (P. 1468.)






572




573

572. *MENTICIRRUS AMERICANUS.* (P. 1474.)
573. *POGONIAS CHROMIS.* (P. 1482.)






574




575

574. *APLODINOTUS GRUNNIENS*. (P. 1484.)
575. *EQUES LANCEOLATUS*. (P. 1489.)






576




577

576. CIRRHIITES RIVULATUS. (P. 1491.)
577. HYSTEROXANTHUS TRASKI. (P. 1496.)






578



579




579a

578. ABEONA MINIMA. (P. 1497.)


579. CYMATOGASTER AGGREGATUS. (P. 1498.)

579a. CYMATOGASTER AGGREGATUS; female. (P. 1498.)






580




581

580. *BRACHYISTIUS FRENATUS.* (P. 1499.)
581. *ZALEMBIUS ROSACEUS.* (P. 1500.)






582




583

582. HYPORHAMPHUS ANALIS. (P. 1500.)
583. PHANERODON FURCATUS. (P. 1506.)







584




585

584. *RHACOCHILUS TOXOTES*. (P. 1507.)
585. *HYPSURUS CARYI*. (P. 1508.)






586




587

586. *DAMALICHTHYS ARGYROSOMUS*. (P. 1509.)
587. *CICHLASOMA BARTONI*. (P. 1515.)






588




589

588. AZURINA HIRUNDO. (P. 1544.)
589. EUPOMACENTRUS FLAVILATUS. (P. 1557.)





590




591

590. *ABUDEFDUF SAXATILIS.* (P. 1561.)
591. *HYPSYPOPS RUBICUNDUS.* (P. 1564.)



IMAGE EVALUATION TEST TARGET (MT-3)




23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503


Photographic
Sciences
Corporation








592




593

592. *MICROSPATHODON BAIRDII.* (P. 1566.)
593. *MICROSPATHODON CHRYSURUS.* (P. 1567.)





594




595


594. *MICROSPATHODON DORSALIS*. (P. 1568.)

595. *TAUTOGOLABRUS ADSPERSUS*. (P. 1577.)





596




597


596. TAUTOGA ONITIS. (P. 1578.)

597. LACHNOLAIMUS MAXIMUS. (P. 1579.)

U S N




598




599

598. *PIMELOMETOPON PULCHER*. (P. 1585.)
599. *CLEPTICUS PARVUS*. (P. 1586.)



600




601


600. *IRIDIO RADIATUS.* (P. 1590.)
601. *IRIDIO BIVITTATUS.* (P. 1595.)

U S N





602




603


602. IRIDIO DISPILUS. (P. 1597.)

603. EMMEEKIA VENUSTA. (P. 1602.)






604




605




606

604. *CHLORICHTHYS GRAMMATICUS*. (P. 1610.)
605. *NOVACULICHTHYS VENTRALIS*. (P. 1615.)
606. *NOVACULICHTHYS INFIRMUS*. (P. 1616.)



607




608


607. XYRICHTHYS PSITTACUS. (P. 1618.)
608. CRYPTOTOMUS BERYLLINUS. (P. 1625.)

U.S.






609



609a




610


609, 609a. *CALOTOMUS XENODON*. (P. 1626.)

610. *SPARISOMA AUROFRENATUM*. (P. 1634.)






611




612




613

611. SPARISOMA HOPLOMYSTAX. (P. 1632.)
612. SCARUS CUZAMILÆ. (P. 1648.)
613. JAWS OF SCARUS CÆRULEUS. (P. 1652.)


614, SCA
615, 615a
616, 616a




614



615



615a



616




616a

614. SCARUS EMBLEMATICUS. (P. 1654.)


615, 615a. LOWER AND UPPER PHARYNGEAL BONES OF SCARUS STRONGYLOCEPHALUS.

616, 616a. LOWER AND UPPER PHARYNGEAL BONES OF SPARISOMA CRETENSE.







617



618


617. *PSEUDOSCARUS GUACAMAIA*. (P. 1657.)
618. *ZENOPSIS OCCELLATUS*. (P. 1660.)





619


619. **CHÆTODIPTERUS FABER.** (P. 1668.)



620


620. *CHAETODON NIGRIROSTRIS.* (P. 1673.)






621


621. *CHEETODON OCCELLATUS.* (P. 1674.)






622

622. **CHETODON CAPISTRATUS.** (P. 1677.)



623


623. **POMACANTHUS ARCUATUS.** (P. 1679.)



624

624. *POMACANTHUS ZONIPECTUS*; YOUNG. (P. 1681.)







625

625. **HOLACANTHUS TRICOLOR.** (P. 1684.)






626



626a


626, 626a. ANGELICHTHYS CILIARIS. (P. 1684.)

U.S.




627

627. *ZANCLUS CORNUTUS.* (P. 1687.)




628




629

628. *TEUTHIS CRESTONIS*. (P. 1692.)
629. *TEUTHIS BAHIANUS*. (P. 1693.)




630




631

630. TAIL OF *TEUTHIS BAHIANUS*. (P. 1693.)
631. *XESURUS PUNCTATUS*. (P. 1694.)





632




633


632. *BALISTES CAROLINENSIS*. (P. 1701.)
633. *CANTHERINES CAROLEE*. (P. 1713.)

U.S. NAT






634




635

634. *MONOCANTHUS CILIATUS.* (P. 1714.)
635. *MONOCANTHUS HISPIDUS.* (P. 1715.)



636




637


636. *ALUTERA SCHCEPFII*. (P. 1718.)
637. *ALUTERA SCRIPTA*. (P. 1719.)

U.S.N.A.







638




639

638. *LACTOPHYRS TRIQUETER*. (P. 1722.)
639. *LACTOPHYRS TRICORNIS*. (P. 1724.)






640




640a




640b


640, 640a, 640b. *LACTOPHYRYS BICAUDALIS.* (P. 1723.)



641




641a




642

641, 641a, *LACTOPHYRYS TRIGONUS*. (P. 1723.)
642. *LAGOCEPHALUS LEVIGATUS*. (P. 1728.)






643



644




645

643. SPHEROIDES NEPHELUS. (P. 1732.)


644. SPHEROIDES SPENGLERI. (P. 1732.)

645. SPHEROIDES MACULATUS. (P. 1733.)






646




646a




647

646, 646a. SPHEROIDES TESTUDINEUS. (P. 1734.)
647. OVOIDES SETOSUS. (P. 1739.)





648




649


648. *DIODON HYSTRIX*. (P. 1745.)

649. *CHILOMYCTERUS SCHEPFI*. (P. 1748.)






650




650a




651

650. LYOSPHERA GLOBOSA. (P. 1751.)
650a. LYOSPHERA GLOBOSA; young. (P. 1751.)
651. MOLA MOLA. (P. 1753.)

11




652




653

652. *RANZANIA TRUNCATA*. (P. 1755.)
653. *SEBASTES MARINUS*. (P. 1760.)







654




655

654. *SEBASTOLOBUS ALTIVELIS.* (P. 1763.)
655. *SEBASTODES MELANOPS.* (P. 1782.)






656




657

656. *SEBASTODES CILIATUS*. (P. 1783.)
657. *SEBASTODES MYSTINUS*. (P. 1784.)




658




659

658. *SEBASTODES BREVISPINIS.* (P. 1787.)
659. *SEBASTODES EIGENMANNI.* (P. 1789.)






660




661

660. *SEBASTODES HOPKINSI.* (P. 1789.)
661. *SEBASTODES ALUTUS.* (P. 1790.)

RP



662




663


662. *SEBASTODES PINNIGER*. (P. 1793.)
663. *SEBASTODES MINIATUS*. (P. 1794.)

10

11





664




665

664. *SEBASTODES CAURINUS.* (P. 1820.)
665. *SEBASTODES MALIGER.* (P. 1822.)





666




667

666. *SEBASTODES GILBERTI*. (P. 1823.)667. *SEBASTODES SERRICEPS*. (P. 1827.)


U.S.

10

AVC




668




669

668. *SEBASTODES NIGROCINCTUS.* (P. 1827.)
669. *SCORPENA CRISTULATA.* (P. 1841.)





670




671


670. *SCORPENA BRASILIENSIS.* (P. 1842.)
671. *SCORPENA MYSTES.* (P. 1849.)

U.S.

11



672



673


672. SCORPÆNA GRANDICORNIS. (P. 1850.)
673. PONTINUS MACROLEPIS. (P. 1855.)

U.S.


Alton

Alton


Alton



674



675




676

674. SETARCHEΣ PARMATUS. (P. 1860.)


675. ANOPLOPOMA FIMBRIA. (P. 1862.)

676. PLEUROGRAMMUS MONOPTERYGIUS. (P. 1864.)


U.S.



677



677a




678


677, 677a. *HEXAGRAMMOS DECAGRAMMUS.* (P. 1867.)
678. *HEXAGRAMMOS OCTOGRAMMUS.* (P. 1869.)

U.S.






679



680



681

679. *HEXAGRAMMOS STELLERI*. (P. 1871.)

680. *HEXAGRAMMOS SUPERCILIOSUS*. (P. 1872.)

681. *HEXAGRAMMOS OTAKII*. (P. 1867.)









IMAGE EVALUATION TEST TARGET (MT-3)



6"



Photographic
Sciences
Corporation


23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

45 2.8
43 3.2
42 3.5
38 2.2
32 2.0
18


10
11

U S NATION






682



683



684


682. *OPHIDION ELONGATUS.* (P. 1875.)

683. *OXYLEBIUS PICTUS.* (P. 1878.)


684. *JORDANIA ZONOPE.* (P. 1884.)




683
684
687



685



686




687

685. *ALCIDEA THOBURNI*. (P. 1887.)


686. SKULL OF *SCORPENICHTHYS MARMORATUS*. (P. 1889.)

687. *CHITONOTUS PUGETENSIS*. (P. 1890.)






688



689



689a


688. ICELINUS BOREALIS. (P. 1896.)

689. ASTROLYTES NOTOSPILOTUS. (P. 1899.)


689a. ASTROLYTES NOTOSPILOTUS; young† (P. 1899.)

U.S. NATION






690




691

690. *ARTEDIELLUS ATLANTICUS.* (P. 1906.)
691. *RUSCARIUS MEANYI.* (P. 1908.)






692



693




694

692. *Rastrinus scutiger*. (P. 1909.)


693. *Icelus spiniger*. (P. 1914.)

694. *Icelus canaliculatus*. (P. 1917.)






695





696




697

695. *RADULINUS BOLEOIDES*. (P. 1919.)
696. *RADULINUS ASPRELLUS*. (P. 1920.)
697. *TRIGLOPS PINGELI*. (P. 1923.)







698



699




700




700a

698. *TRIGLOPS BEANI*. (P. 1924.)699. *TRIGLOPS SCEPTICUS*. (P. 1925.)700, 700a. *STERNIAS XENOSTETHUS*. (P. 1927.)






701



702



703


701. *PRIONISTIUS MACELLUS*. (P. 1928.)

702. *ELANURA FORFICATA*. (P. 1930.)


703. *MELLETES PAPILIO*. (P. 1932.)




704. HEMI
705, 705a,




704



705



705a




705b


704. *HEMILEPIDOTUS JORDANI.* (P. 1934.)

705, 705a, 705b. *HEMILEPIDOTUS HEMILEPIDOTUS.* (P. 1935.)






706




707

706. ENOPHRYS BISON. (P. 1938.)
707. CERATOCOTTUS DICERAUS. (P. 1940.)







708




709

708. *COTTUS EVERMANNI*. (P. 1945.)
709. *COTTUS PUNCTULATUS*. (P. 1948.)






710





711




712

710. *COTTUS PERPLEXUS*. (P. 1955.)
711. *COTTUS KLAMATHENSIS*. (P. 1955.)
712. *COTTUS ALEUTICUS*. (P. 1957.)






713




714




715

713. *COTTUS LEIOPOMUS*. (P. 1962.)714. *COTTUS PRINCEPS*. (P. 1962.)715. *URANIDEA TENUIS*. (P. 1966.)






716



716a




717


716, 716a. *MYOXOCEPHALUS AENEUS.* (P. 1972.)
717. *MYOXOCEPHALUS SCORPIUS.* (P. 1974.)




718. MYO
719. MYO
720. MYO



718



719



720


718. *MYOXOCEPHALUS OCTODECIMSPINOSUS*. (P. 1976.)

719. *MYOXOCEPHALUS POLYACANTHOCEPHALUS*. (P. 1976.)


720. *MYOXOCEPHALUS JAOK*. (P. 1977.)

U.S. NATION






721



721a




722


721, 721a. *MYOXOCEPHALUS VERRUCOSUS.* (P. 1979.)
722. *MYOXOCEPHALUS AXILLARIS.* (P. 1980.)




723,
724.




723



723a



723b



724


723, 723a, 723b. *MYOXOCEPHALUS STELLERI.* (P. 1981.)
724. *MYOXOCEPHALUS NIGER.* (P. 1985.)




725a




725, 726
726, 727




725



725a



726a




726


725, 725a. *MEGALOCOTTUS PLATYCEPHALUS*. (P. 1987.)
726, 726a. *MEGALOCOTTUS LATICEPS*. (P. 1988.)

U.S. NATIONAL






727



728



729


727. *ZESTICELUS PROFUNDORUM*. (P. 1990.)

728. *DASYCOTTUS SETIGER*. (P. 1991.)


729. *ONCOCOTTUS QUADRICORNIS*. (P. 2001.)




730a




730, 731. G




730



730a




730b




731

730, 730a, 730b. *Gymnophthalmus pistilliger.* (P. 2006.)
731. *Gymnophthalmus galeatus.* (P. 2010.)






732



733




734

732. *LEIOCOTTUS HIRUNDO*. (P. 2011.)


733. *LEPTOCOTTUS ARMATUS*. (P. 2012.)

734. *OLIGOCOTTUS MACULOSUS*. (P. 2013.)






735




736

735. *BLENNICOTTUS EMBRYUM*. (Pp. 2016, 2864.)
736. *HISTIOCOTTUS BILOBUS*. (P. 2018.)






737





737a




737b

737, 737a, 737b. BLEPSIAS CIRRHOUS. (P. 2018.)





738




739


1/2 in.

738. *Hemitripterus americanus*. (P. 2023.)
739. *Ascelichthys rhodorus*. (P. 2025.)






740




741


740. *PSYCHROLUTES PARADOXUS*. (P. 2026.)
741. *GILBERTIDIA SIGOLUTES*. (P. 2028.)




742, 742a




742



742a




742b



742c

742, 742a, 742b, 742c. *RHAMPHOCOTTUS RICHARDSONI*. (P. 2030.)






743



744



745



745a

743. *OCCA DODECAEDRON*. (P. 2044.)

744. *PALLASINA BARBATA*. (P. 2049.)

745, 745a. *PODOTHECUS ACCIPITER*. (P. 2055.)




54a

746. P.


747. 74

748. 74


749. 74




746



747




748




749



746a




747a




748a


746. *PODOTHECUS ACIPENSERINUS*. (P. 2061.)747, 747a. *PODOTHECUS VETERNUS*. (P. 2063.)748, 748a. *STELGIS YULSUS*. (P. 2067.)749, 749a. *AVERRUNCUS EMMELANE*. (P. 2069.)




73
74
75




750




751



752



750a




752a

750, 750a. *AVERRUNCUS STERLETUS*. (P. 2071.)


751. *SARRITOR FRENATUS*. (P. 2073.)

752, 752a. *XYSTES AXINOPHYRS*. (P. 2076.)





753



754



754a


753. *BATHYAGONUS NIGRIPINNIS*. (P. 2078.)
754, 754a. *XENOCHIRUS TRIACANTHUS*. (P. 2084.)




755




755, 756
756, 757



755



755a



755b



756




756a


755, 755a, 755b. *ASPIDOPHOROIDES GUNTHERI.* (P. 2090.)

756, 756a. *ASPIDOPHOROIDES MONOPTERYGIUS.* (P. 2091.)





757




758


757. CYCLOPTERUS LUMPUS. (P. 2096.)

758. LETHOTREMUS MUTICUS. (P. 2101.)






759




760




760a

759. *LETHOTREMUS VIOLENATUS*. (P. 2101.)
760, 760a. *NEOLIPARIS CALYODON*. (P. 2110.)






761




761a





762

761, 761a. *NEOLIPARIS MUCOSUS*. (P. 2111.)
762. *NEOLIPARIS FLORE*. (P. 2111.)




764

763



763a






764a

763, 763a. *NEOLIPARIS GREENI.* (P. 2112.)


764, 764a. *LIPARIS CYCLOPUS.* (P. 2118.)






765. *LIPARIS AGASSIZII*. (P. 2121.)
766, 766a. *LIPARIS DENNYI*. (P. 2124.)






767




768

767. *BATHYPHASMA OVIGERUM*. (P. 2128.)
768. *PRIONOTUS CAROLINUS*. (P. 2156.)





769




770


769. *PRIONOTUS SCITULUS*. (P. 2157.)

770. *PRIONOTUS ALATUS*. (P. 2159.)





771




772


771. PRIONOTUS STEARNSI. (P. 2166.)

772. PRIONOTUS EVOLANS. (P. 2168.)






773



774





775

773. *BELLATOR EGRETTA*. (P. 2174.)


774. *PERISTEDION MINIATUM*. (P. 2178.)

775. *PERISTEDION LONGISPATHUM*. (P. 2178.)






776



777




777a


776. *PERISTEDION GRACILE.* (P. 2179.)777, 777a. *PERISTEDION PLATYCEPHALUM.* (P. 2180.)

U.S. NATIONAL






778



779



780


778. *CEPHALACANTHUS VOLITANS.* (P. 2183.)

779. *CALLIONYMUS AGASSIZII.* (P. 2186.)


780. *IOGLOSSUS CALLIURUS.* (P. 2193.)

U.S. NATION






781




782

781. *PHILYPNUS DORMITOR.* (P. 2194.)
782. *DORMITATOR MACULATUS.* (P. 2196.)






783

 $\frac{1}{2}$ in.

784




785


783. ELEOTRIS PISONIS. (P. 2200.)
784. ALEXURUS ARMIGER. (P. 2203.)
785. ERETELIS SMARAGDUS. (P. 2204.)

U.S. NATIONAL






786



787



788


786. *LOPHOGOBIUS CYPRINOIDES*. (P. 2209.)

787. *GOBIUS STIGMATICUS*. (P. 2224.)


788. *GOBIUS HASTATUS*. (P. 2229.)

U.S. NATION






789

 $\frac{1}{2}10$ 


789a




790

789 789a. GOBIUS OCEANICUS. (P. 2230.)
790 GARMANNIA PARADOXA. (P. 2232.)






791



792




793

791. *BOLLMANNIA CHLAMYDES*. (P. 2238.)

792. *ABOMA ETHEOSTOMA*. (P. 2240.)

793. *CLEVELANDIA IOS*. (P. 2254.)








IMAGE EVALUATION TEST TARGET (MT-3)



6"





Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-45C3


EE
128
2.5
32
34
2.2
20
8

II
10
EE






794



795

1/2 in.




796

794. EVERMANNIA ZOSTERURA. (P. 2256.)


795. TYPHLOGOBius CALIFORNIENSIS. (P. 2262.)

796. ECHENEUS NAUCRATES. (P. 2269.)







797



797a



798




799

797, 797a. REMORA BRACHYPTERA. (P. 2272.)


798. RHOMBOSOLEA OSTEOCRIR. (P. 2273.)

799. CAULOLATILUS MICROPS. (P. 2277.)






800





800a




801

800, 800a. *OPISTHOGNATHUS MACROGNATHUM*. (P. 2281.)
801. *GNATHYPOPS MAXILLOSA*. (P. 2284.)






802




803




804

802. *BATHYMASTER SIGNATUS.* (P. 2288.)
803. *RONQUILUS JORDANI.* (P. 2289.)
804. *CHIASMODON NIGER.* (P. 2291.)






805



806




807

805. *HYPSCOMETES GOBIOIDES*. (P. 2294.)


806. *TRICHODON TRICHOON*. (P. 2295.)

807. *ARCTOSCOPUS JAPONICUS*. (P. 2297.)






808



809




809a


808. ASTROSCOPUS Y-GRÆCUM. (P. 2307.)

809, 809a. KATHETOSTOMA ALBIGUTTA. (P. 2312.)







810



811




812




813

810. *OPSANUS PARDUS.* (P. 2316.)811. *PORICHTHYS POROSISSIMUS.* (P. 2319.)812. *CAULARCHUS MEANDRICUS.* (P. 2328.)813. *RIMICOLA MUSCARUM.* (P. 2338.)






814



815

1/2 in.





816

814. ENNEANECTES CARMINALIS. (P. 2350.)


815. GIBBONSIA ELEGANS EVIDES. (P. 2352.)

816. NEOCLINUS SATIRICUS. (P. 2355.)






817



818





819

817. *Cryptotrema corallinum*. (P. 2366.)


818. *Exerpes asper*. (P. 2367.)

819. *Auchenopterus nox*. (P. 2373.)






820



821




822

820. *BLENNIUS FAVOSUS*. (P. 2380.)


821. *BLENNIUS CRISTATUS*. (P. 2382.)

822. *HYPSOBLENNIUS IONTHIAS*. (P. 2388.)






823




824




825

823. *HYPSOBLENNIUS HENTZ.* (P. 2390.)824. *CHASMODES SABURRI.* (P. 2392.)825. *RUPISCARTES ATLANTICUS.* (P. 2397.)





826



827


826. EMBLEMARIA ATLANTICA. (P. 2402.)

827. STATHMONOTUS HEMPHILLII. (P. 2407.)






828






829



830

828. *BRYOSTEMA POLYACTOCEPHALUM*. (P. 2408.)
829. *BRYOSTEMA NUGATOR*. (P. 2410.)
830. *APODICHTHYS FLAVIDUS*. (P. 2411.)






831



832




833

831. *PHOLIS DOLICHOGASTER.* (P. 2416.)

832. *PHOLIS GUNNELLUS.* (P. 2419.)

833. *PHOLIS ORNATUS.* (P. 2419.)






834





835



836

834. *ASTEROPTERYX GUNNELLIFORMIS*. (P. 2420.)
835. *ANOPLARCHUS ATROPURPUREUS*. (P. 2422.)
836. *XIPHISTES ULV.E.* (P. 2423.)





837



838





839




840

837. *XIPHISTES CHIRUS*. (P. 2424.)
838. *XIPHIODON RUPESTRE*. (P. 2426.)
839. *LUMENUS MACKAYI*. (P. 2436.)
840. *LUMENUS LAMPETRÆFORMIS*. (P. 2438.)






841



842



843


841. *STICHLÆUS PUNCTATUS*. (P. 2439.)842. *ULVARIA SUBBIFURCATA*. (P. 2440.)843. *CRYPTACANTHODES MACULATUS*. (P. 2443.)

U.S. NATION






844




845

844. LYCONECTES ALEUTENSIS. (P. 2444.)
845. ANARHICHAS LATIFRONS. (P. 2446.)






846



847



848

846. ANARHICHAS LUPUS. (P. 2446.)

847. ANARHICHAS LEPTURUS. (P. 2447.)

848. PTILICHTHYS GOODEI. (P. 2452.)





849




849a



849b



850




851

849, 849a, 849b. *SCYTALINA CERDALE*. (P. 2454.)

850. *ZOARCES ANGUILLARIS*. (P. 2457.)

851. *EMBRYX CROTALINUS*. (P. 2458.)






852



853






854

852. *APRODON CORTEZIANUS.* (P. 2461.)

853. *LYCOPTERUS ZOARCHUS.* (P. 2464.)

854. *LYCODES RETICULATUS.* (P. 2465.)






855



856



857



858

855. LYCODES PERSPICILLUM. (P. 2465.)

856. LYCODES FRIGIDUS. (P. 2465.)


857. LYCODALEPIS POLARIS. (P. 2468.)

858. LYCODALEPIS TURNERI. (P. 2468.)



—





859



860



860a




860c



860d



860b




861

859. *LYCENCHELYS VERRILLII*. (P. 2470.)

860, 860a, 860b, 860c, 860d. *LYCENCHELYS PAXILLUS*. (P. 2471.)

861. *FURCIMANUS DIAPTERUS*. (P. 2472.)






862




863



864



864a



864b



864c



865


862. *LYCODONUS MIRABILIS*. (P. 2474.)

863. *LYCONEMA BARBATUM*. (P. 2474.)


864, 864a, 864b, 864c. *GYMNELIS VIRIDIS*. (P. 2477.)

865. *MELANOSTIGMA PAMMELAS*. (P. 2479.)






866



867





868

866. *LEPOPHIDIUM MARMORATUM*. (P. 2482.)


867. *LEPOPHIDIUM PROFUNDORUM*. (P. 2484.)

868. *RISSOLA Marginata*. (P. 2489.)






869



870





871

869. *OTOPHIDIUM OMOSTIGMUM*. (P. 2490.)


870. *LYCODAPUS DERMATINUS*. (P. 2492.)

871. *BROTULA BARBATA*. (P. 2500.)






872




873




874

872. *OGILBIA VENTRALIS*. (P. 2503.)
873. *OGILBIA CAYORUM*. (P. 2503.)
874. *DICROMITA AGASSIZII*. (P. 2506.)






875




876




876a

875. *BASSOZETUS NORMALIS.* (P. 2507.)
876, 876a. *BASSOZETUS CATENA.* (P. 2509.)







877



878



879



880

877. *NEORYTHITES GILLII.* (P. 2512.)


878. *NEORYTHITES MARGINATUS.* (P. 2513.)

879. *BASSOGIGAS GILLII.* (P. 2515.)


880. *BARATHRODEMUS MANATINUS.* (P. 2517.)

U. S. NATI






881



882




883


881. *POROGADUS MILES.* (P. 2520.)
882. *DICROLENE INTRONIGRA.* (P. 2522.)
883. *BARATHRONUS BICOLOR.* (P. 2524.)

U. S. NATION






884





885




886

884. *MERLUCCIUS PRODUCTUS.* (P. 2531.)885. *BOROGADUS SAIDA.* (P. 2533.)886. *POLLACHIUS VIRENS.* (P. 2534.)







887



888



889



890

887. *THERAGRA CHALCOGRAMMA*. (P. 2535.)


888. *ELEGINUS NAVAGA*. (P. 2537.)

889. *MICROGADUS PROXIMUS*. (P. 2539.)


890. *MICROGADUS TOMCOD*. (P. 2540.)




t



891



892





892a

891. *GADUS COLLARIAS*. (P. 2541.)

892. *MELANOGRAMMUS EGLEFINUS*. (P. 2542.)

892a. *MELANOGRAMMUS EGLEFINUS*; SKULL. (P. 2542.)






893



893a




894


893, 893a. ANTIMORA VIOLA. (P. 2544.)

894. URALEPTUS MALARDI. (P. 2545.)





895




896

895. *LOTELLA MAXILLARIS.* (P. 2546.)


896. *PHYSICULUS FULVUS.* (P. 2547.)

U. S. NAT






897




898




899

897. *Lota maculosa*. (P. 2550.)
898. *Urophycis regius*. (P. 2553.)
899. *Urophycis cirratus*. (P. 2553.)






900



901



902


900. *UROPHYCIS EARLLI*. (P. 2554.)

901. *UROPHYCIS TENUIS*. (P. 2555.)


902. *UROPHYCIS CHUSS*. (P. 2555.)

U. S. NAT






903



904




905


903. *UROPHYCIS CHESTERI*. (P. 2556.)
904. *LEMONEMA BARBATULUM*. (P. 2556.)
905. *LEMONEMA MELANURUM*. (P. 2557.)

U. S. NAT.






906



907





908

906. *GAIIRDROPSARUS ARGENTATUS*. (P. 2559.)


907. *ENCHELYOPUS CIMRIUS*. (P. 2560.)

908. *BATHYGADUS FAVOSUS*. (P. 2565.)






909



910



911


909. STEINDACHNERIA ARGENTEA. (P. 2568.)

910. CHALINURA SIMULA. (P. 2578.)


911. CORYPHLEONOOIDES CARAPINUS. (P. 2579.)

U. S. NATI






912



913




914

912. HYMENOCEPHALUS CAVERNOSUS. (P. 2580.)


913. CÆLORHYNCHUS OCCA. (P. 2588.)

914. CÆLORHYNCHUS CARMINATUS. (P. 2588.)

U. S. N.



915




916


915. *COLOMHNCHUS CARIBBEUS*. (P. 2589.)
916. *REGALECUS GLESNE*. (P. 2596.)

U. S. N





917




918


917. *ATHERESTHES STOMIAS.* (P. 2609.)
918. *HIPPOGLOSSUS HIPPOGLOSSUS.* (P. 2611.)

U. S. NA





919




920


919. HIPPOGLOSSOIDES PLATESSOIDES. (P. 2614.)
920. HIPPOGLOSSOIDES ELASSODON. (P. 2615.)

U. S. NA





921




922


921. *PSETTICHTHYS MELANOSTICTUS.* (P. 2618.)
922. *PARALICHTHYS DENTATUS.* (P. 2629.)

U. S. NAT





923




924

923. *PARALICHTHYS SQUAMILEATUS.* (P. 2631.)


924. *PARALICHTHYS OBLONGUS.* (P. 2632.)

U.S. NATION





925



926

925. *ANCYLOSETTA QUADROCELLATA*. (P. 2634.)
926. *PLEURONICHTHYS DECURRENS*. (P. 2637.)










IMAGE EVALUATION TEST TARGET (MT-3)




Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503


1.8
2.0
2.2
2.5
2.8
3.2
3.5
4.0

0.1
0.5
1.0
1.5
2.0
2.5
3.0

U. S. N




927




928

927. *INOPSETTA ISCHYRA*. (P. 2641.)
928. *LEPIDOPSETTA BILINEATA*. (P. 2643.)

U.S.




929




930

929. *LIMANDA FERRUGINEA*. (P. 2644.)
930. *LIMANDA ASPERA*. (P. 2645.)





931




932

931. *LIMANDA PROBOSCIDEA*. (P. 2645.)
932. *LIMANDA BEANII*. (P. 2646.)






933




934

933. *PSEUDOPLEURONECTES AMERICANUS*. (P. 2647.)
934. *PLEURONECTES QUADRITUBERCULATUS*. (P. 2648.)




935




936

935. LIOPSETTA GLACIALIS. (P. 2649.)
936. LIOPSETTA PUTNAMI. (P. 2650.)



937




937a


937, 937a. *PLATICHTHYS STELLATUS.* (P. 2652.)

W. H. G.

A



938




939


938. *LOPHOPSETTA MACULATA*. (P. 2660.)
939. *PLATOPHRYS OCELLATUS*. (P. 2663.)

U. S. N.






940




941


940. *TRICHOPSETTA VENTRALIS.* (P. 2669.)

941. *SYACIUM PAPILLOSUM.* (P. 2671.)





942




943


942. *AZEVIA PANAMENSIS*. (P. 2677.)

943. *CITHARICHTHYS SORDIDUS* (P. 2679.)

U.S.




944




945

944. *CITHARICHTHYS MACROPS.* (P. 2684.)945. *ETROPUS RIMOSUS.* (P. 2688.)





946




947


946. ETROPUS CROSSOTUS. (P. 2689.)

947. ACHIRUS LINEATUS. (P. 2697.)

U. S. N.




948




949


948. *ACHIRUS FASCIATUS*. (P. 2700.)
949. *SYMPHURUS MARGINATUS*. (P. 2706.)



950




951




952

950. *SYMPHURUS PLAGIUSA*. (P. 2710.)
951. *SYMPHURUS WILLIAMSI*. (P. 2711.)
952. *LOPHIUS PISCATORIUS*. (P. 2713.)

I.J. S. N




953




954

953. *CHAUNAX PICTUS.* (P. 2726.)
954. *CERATIAS HOLBOLLI.* (P. 2729.)

U. S.




955





956

955. *MANCALIAS SHUFELDTI*. (P. 2730.)
956. *CRYPTOPSARAS COUESII*. (P. 2731.)




957. CAULOPHRYNE JORDANI. (P. 2735.)






958



958a



958b

958, 958a, 958b. *Ogcocephalus vespertilio*. (P. 27²⁷)

