

**CIHM  
Microfiche  
Series  
(Monographs)**

**ICMH  
Collection de  
microfiches  
(monographies)**



**Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques**

**© 1997**

## Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming are checked below.

- ☐ Coloured covers / Couverture de couleur
- ☐ Covers damaged / Couverture endommagée
- ☐ Covers restored and/or laminated / Couverture restaurée et/ou pelliculée
- ☐ Cover title missing / Le titre de couverture manque
- ☐ Coloured maps / Cartes géographiques en couleur
- ☐ Coloured ink (i.e. other than blue or black) / Encre de couleur (i.e. autre que bleue ou noire)
- ☐ Coloured plates and/or illustrations / Planches et/ou illustrations en couleur
- ☒ Bound with other material / Relié avec d'autres documents
- ☐ Only edition available / Seule édition disponible
- ☒ Tight binding may cause shadows or distortion along interior margin / La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure.
- ☐ Blank leaves added during restorations may appear within the text. Whenever possible, these have been omitted from filming / Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- ☐ Additional comments / Commentaires supplémentaires:

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- ☐ Coloured pages / Pages de couleur
- ☐ Pages damaged / Pages endommagées
- ☐ Pages restored and/or laminated / Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed / Pages décolorées, tachetées ou piquées
- ☐ Pages detached / Pages détachées
- ☒ Showthrough / Transparence
- ☐ Quality of print varies / Qualité inégale de l'impression
- ☐ Includes supplementary material / Comprend du matériel supplémentaire
- ☐ Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image / Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.
- ☐ Opposing pages with varying colouration or discolourations are filmed twice to ensure the best possible image / Les pages s'opposant ayant des colorations variables ou des décolorations sont filmées deux fois afin d'obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below /  
Ce document est filmé au taux de réduction indiqué ci-dessous.

|     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 10x | 12x | 14x | 16x | 18x | 20x | 22x | 24x | 26x | 28x | 30x | 32x |
|     |     |     |     |     |     |     |     | ✓   |     |     |     |

The copy filmed here has been reproduced thanks to the generosity of:


Library  
Agriculture Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol  $\rightarrow$  (meaning "CONTINUED"), or the symbol  $\nabla$  (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:


Bibliothèque  
Agriculture Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.


Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole  $\rightarrow$  signifie "A SUIVRE", le symbole  $\nabla$  signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.



# MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART Na. 2)



**APPLIED IMAGE Inc**

1653 East Main Street  
Rochester, New York 14609 USA  
(716) 482 - 0300 - Phone  
(716) 286 - 5989 - Fax

PROVINCE OF BRITISH COLUMBIA.

DEPARTMENT OF AGRICULTURE  
(LIVE-STOCK BRANCH).

PRACTICAL POULTRY-RAISING.

BULLETIN No. 26  
*(Third Edition).*



PRINTED BY AUTHORITY OF  
THE LEGISLATIVE ASSEMBLY OF BRITISH COLUMBIA.

VICTORIA, B.C.:  
Printed by WILLIAM H. CULLIN, Printer to the King's Most Excellent Majesty.  
1912.

*Hon. Price Ellison,*

*Minister of Agriculture,*

*Victoria, B.C.*

SIR,—I have the honour to submit for your approval the third edition of Bulletin No. 26, as revised by J. R. Terry, Chief Poultry Instructor, containing practical information on the care and management of poultry.

I have the honour to be,

Sir,

Your obedient servant,

WM. E. SCOTT,

*Deputy Minister of Agriculture.*

*Victoria, B.C., June, 1912.*

PROVINCE OF BRITISH COLUMBIA.

---

DEPARTMENT OF AGRICULTURE.

---

LIVE-STOCK BRANCH.

---

*Minister of Agriculture*—HON. PRICE ELLISON.

*Deputy Minister of Agriculture*—W. E. SCOTT.

*Live-stock Commissioner*—W. T. McDONALD, B.S.A., M.S.

*Dairy Instructor*—H. RIVE, B.S.A.

*Chief Poultry Instructor*—I. R. TERRY.

*Poultry Instructor*—H. E. UPTON.

*Chief Veterinary Inspector*—A. KNIGHT, V.S.

*Veterinary Inspectors*—

S. A. K. WHITE, V.S., W. W. ALTON, V.S., B. R. ILSLEY, V.S.

*Secretary of the Department*—W. J. BONAVIA.



Fig. (1.) Feeding growing stock. "The man behind the feed-pail" is an important factor in the successful management of poultry.



# PRACTICAL POULTRY-RAISING.

---

By J. R. TERRY, CHIEF POULTRY INSTRUCTOR.

THE business of poultry-raising is one of the most attractive pursuits of all the agricultural industries. More than ever the poultry industry is being regarded as a national industry. Its importance is being realized by the public and considerable increased interest is being shown because of prevailing conditions. The product of the domestic hen is demanded more than any other product of agricultural or animal industry, with the exception of milk and flour. There is a constant and increasing demand particularly for the fresh egg, and within the past few years the number marketed, and consumed, has increased tremendously. The prices have increased also, proving that the supply is not keeping pace with the demand and indicating very positively that the business will not be overdone or become unprofitable. Few departments of agricultural industry offer greater returns on equal capital invested to those who work diligently and consistently. The hen to-day may be considered the most profitable condenser of the raw materials of the land into a finished product. In other words, the egg is manufactured at less cost to the country than almost any other staple food product.

Any farm is improved by a poultry department, and few businesses can be operated in as wide a range of climate and in as varied conditions. Poultry-raising, then, is more independent than almost any other business and few branches can be carried on with as inexpensive an outlay. The returns on the money invested in a poultry plant are immediate and regular.

It is the purpose of the author to present in this bulletin such information as may be followed by the poultryman with his small or large flock. No hard-and-fast rules are laid down, since it is a matter of record that no one method can be described which will prove adequate and suitable under all conditions with which the poultryman on a small or large scale may be confronted. Through emphasis of the general underlying principles in raising poultry the reader is left to exercise individual judgment in applying the principles to his own particular conditions.

---

## THE POULTRY INDUSTRY.

---

Until recently very little attention has been given to poultry-raising in this Province, the industry having been largely considered as a side-issue of but little importance. The attention of the rural community has been absorbed almost entirely in fruit-growing and dairying. These are the two leading industries of the Province, but it is surprising how the poultry industry has been neglected when the proportional values of the horticulture, dairy, and poultry products are compared.

Poultry husbandry, then, is seen to be deserving of more attention than it has received in the past.

### THE IMPORTANCE OF THE INDUSTRY.

That the poultry industry is an important one can be fully realized by a study of the statistics of production and importations. In 1901 there were in the neighbourhood of 17,000,000 head of poultry on Canadian farms. The value of stock and products was over \$10,000,000—being about \$10,500,000 for eggs alone. This amount points to the magnitude of the industry in Canada. Of this enormous sum, British Columbia produced 420,020 dollars' worth of eggs and 200,717 dollars' worth of other produce. It is readily seen what an insignificant part the British Columbia hen plays in the production of Canadian poultry products. In 1905 there were laid in Canada 90,000,000 dozen eggs, and in the same year about 10,000,000 dollars' worth of dressed poultry was consumed. Although comparatively small quantities are produced within the Province, large importations are made. The extent of the importations is convincing that the markets of the Province are good.

### POULTRY AND MIXED HUSBANDRY.

The system of mixed farming as carried on in Canada is ideal for the rearing of poultry. In some of the Provinces, especially Ontario, encouragement is given to general farmers to keep a few fowls on each farm. The establishment of exclusive poultry plants is not encouraged. In this Province, because of our highly specialized industries and because of other conditions, we find that the keeping of fowls in large numbers is a good investment, when combined with expert knowledge, favourable surroundings, and intelligence. Quite a few poultry plants, where eggs and other poultry produce form the basis of returns, are quite successful. Conditions seem to allow of intensive specialization. At the same time it must not be forgotten that excellent profits are possible on farms of mixed husbandry. The returns from the flock should add materially to the income of the fruit-grower, the dairyman, or even the man who is merely in the process of clearing his land. The orchard makes an admirable run for poultry, especially growing stock, as does also the cattle-yard. In the orchard, many insect pests will be destroyed and the manure from the fowls will greatly enrich the soil. By-products in the orchard and in the fields are converted into profit. Fowls are economizers, and when a poultry department is carried on with other branches of farming a system of rotation could be adopted.

### THE MARKET.

Poultry produce finds a more ready sale with better prices on our home markets than elsewhere in Canada. There is no market like the "home" market, but in the past it has been largely supplied by poultrymen of the East and South. During 1911 there was imported 1,024,875 dollars' worth of poultry produce from the Eastern Provinces, as well as 117,800 dollars' worth of produce from the United States. The importations of produce in 1911 amounted to \$1,142,735. A study of these figures is convincing that British Columbia is annually losing enormous sums of money. Such a condition is most undesirable, especially when we consider the adaptability of the Province for increased production.

In 1906 the average price paid per dozen for eggs was 25 cents.

In 1907 the average price was 30 cents.

In 1908 the average price was 35 cents.

In 1909 the average price was 37 cents.

In 1910 the average price reached 45 cents in some parts of the Province.

## THE CLIMATE.

The poultry industry is more independent of climate than other branches of agriculture. The climate of the Province as a whole is quite favourable. Some sections of the Province have slight advantages in climate over other sections, but the average temperature is not low. Climate conditions are such that fowls may be kept under the most natural conditions and therefore the best. This means healthier flocks with less cost in maintaining than in other parts of Canada where artificial methods must be adopted.

## THE SOIL.

The nature of the soil is a more important factor than many poultrymen realize. The soil need not be very fertile, but above all it should be clean. A sandy soil suits best, but it should be able to grow a sufficient amount of green food. Loams and clays will often grow abundance of green food, but are sometimes difficult to drain. Then they are apt to become contaminated and this is one thing that poultrymen must avoid. Considering our comparatively moist climate in some parts of the Province, the question of soil contamination becomes one of our most important problems. The runs and yards should be kept sweet and clean by frequent ploughing and sowing down to some form of green food. Land that is well drained and porous should be selected if possible. Danger of soil contamination must be avoided. On a soil of a porous nature the droppings are deodorized by absorption of their liquids, and the rains wash the remaining solids below the surface. This tends to prevent contamination and disease is kept down; whereas in very heavy clay soil, especially where the top soil tends to bake, the surface soon becomes foul and outbreaks of disease occur. If possible, enough land should be secured so that not more than 300 fowls be placed on an acre. Adult birds should be given double yards, so that green food is growing in one run while the fowls are in the other.

It is not advisable to start a poultry ranch on less than 5 acres. Where 500 head of layers are kept on this acreage, the growing stock can be provided with free range. Chicks so raised can be grown much cheaper, with less mortality, than chicks reared on restricted range.

A preferable site is on a gentle slope facing south or south-east. Such an exposure admits of the greatest amount of sunlight, which is an important factor in the rearing of poultry.

## THE BREEDS.

Of the many breeds of poultry, only the more popular commercial varieties can be given here.

**PLYMOUTH ROCKS.**—The most popular varieties of this breed are the Barred, White, and Buff. The Barred Plymouth Rock is the oldest of the three varieties. All three varieties are noted for their qualities as general-purpose fowls, being excellent flesh producers and good layers. They have single combs, yellow legs and flesh, are fairly hardy, and layers of brown or tinted eggs.

The standard weights are:—Cock, 9½ lb.; cockerel, 8 lb.; hen, 7½ lb.; and pullet, 6½ lb.

**WYANDOTTES.**—There are several varieties of this breed, including the Silver, Golden, White, Buff, Black, Partridge, Silver Pencilled, and Columbian.



Fig. (2.) Barred Plymouth Rock.

The last variety is the newest of all, while the White is the most popular. The Wyandottes bear the same general characteristics, from a commercial standpoint, as the Plymouth Rocks. They have rose combs and possess a different shape. The White Wyandotte is bred by many market poultrymen.



Fig. (3.) Single-comb White Leghorn.

The standard weights are:—Cock,  $8\frac{1}{2}$  lb.; cockerel,  $7\frac{1}{2}$  lb.; hen,  $6\frac{1}{2}$  lb.; and pullet,  $5\frac{1}{2}$  lb.

**RHODE ISLAND RED.**—The only two varieties of this breed are the Single and Rose Comb and they have no material differences except in combs.

Although a breed of comparatively recent origin, the Rhode Island Red is quite popular as a utility fowl. As with all American breeds, the Reds have yellow flesh and legs, and lay brown or fluted eggs.

The standard weights are:—Cock,  $8\frac{1}{2}$  lb.; cockerel,  $7\frac{1}{2}$  lb.; hen,  $6\frac{1}{2}$  lb.; and pullet, 5 lb.

**ORPINGTONS.**—The most popular varieties of this breed are the Buff, Black, and White, the Buff being the favourite.

They are hardy, good layers, and flesh well for the market; having a white skin. They are single combed, and lay brown or fluted eggs.

The standard weights are:—Cock, 10 lb.; cockerel,  $8\frac{1}{2}$  lb.; hen, 8 lb.; and pullet, 7 lb.

**LEGHORNS.**—This breed composes a group characterized by great activity and sprightliness. The most popular varieties are the Single-comb White, Brown, and Buff, while the Rose-comb varieties receive favour in some parts. The Single-comb White Leghorn is the most numerous fowl in the Province.

All Leghorns are essentially egg breeds, the egg being white in colour. The young cockerels make splendid squab broilers and broilers.

**MINORCAS.**—The Single-comb Black is the most popular variety. The Minorcas have much of the same characteristics of the Leghorns, except that they are larger, have larger combs, and lay larger eggs.

The standard weights for the Single-comb Black Minorca are:—Cock, 9 lb.; cockerel,  $7\frac{1}{2}$  lb.; hen,  $7\frac{1}{2}$  lb.; and pullet,  $6\frac{1}{2}$  lb.

**GAMES.**—The Indian Game is the chief variety of interest to the farmer. They are a good meat fowl, having short, thick legs, large thighs, and a deep and broad breast. However, they are very poor layers generally.

The standard weights are:—Cock, 9 lb.; cockerel,  $7\frac{1}{2}$  lb.; hen,  $6\frac{1}{2}$  lb.; and pullet,  $5\frac{1}{2}$  lb.

**BRONZE TURKEYS.**—The Bronze turkey is the largest of all varieties and is also the most popular. They are hardy, good foragers, and dress a good proportion of flesh.

The standard weights are:—Adult cock, 36 lb.; yearling cock, 33 lb.; cockerel, 25 lb.; hen, 20 lb.; and pullet, 16 lb.

**PEKIN DUCKS.**—The marketable value as table poultry has made the Pekin duck a favourite with the majority of duck-breeders. In "the making of the duck industry of America" the Pekin was used almost exclusively, and its popularity is exceeded by no other variety, because of its utilitarian qualities. As layers and rapid producers of flesh they are unexcelled.

The standard weights are:—Adult drake, 8 lb.; young drake, 7 lb.; adult duck, 7 lb.; and young duck, 6 lb.

**ROUEN DUCKS.**—The Rouen is quite hardy, of large size, and dresses well for the table.

The standard weights are:—Adult drake, 9 lb.; young drake, 8 lb.; adult duck, 8 lb.; and young duck, 7 lb.

**INDIAN RUNNER DUCKS.**—The variety is noted for its peculiar shape and for its productivity. They are splendid layers of white transparent eggs, but are too small to be classed as a meat fowl.

The standard weights are:—Drake,  $4\frac{1}{2}$  lb., and duck, 4 lb.

**TOULOUSE GESE.**—There is very little difference between the Toulouse and Embden geese as far as weight is concerned, both being classed as the heavyweights of the goose family. The plumage is grey and white. The Toulouse are better layers, but not such good setters as the Embden.

The standard weights are:—Adult gander, 20 lb.; young gander, 18 lb.; adult goose, 18 lb.; and young goose, 15 lb.

**EMILDEN GEES.**—Having white plumage, the Emlden is sometimes preferred for the white feathers, feathers being an important item in goose-culture.

The standard weights are:—Adult gander, 20 lb.; young gander, 18 lb., adult goose, 18 lb.; and young goose, 16 lb.

#### STARTING IN BUSINESS.

To the man who wishes to make a business of poultry-raising, to raise superior birds from a commercial standpoint, British Columbia offers inducements such as few other Provinces in the Dominion. There is room for great development. There is a large and stable home market, which is expanding very rapidly. Markets are opening up in the North, and Alaska also has a market which British Columbia should be supplying. The increase in prices with the increase in importations gives some idea of the great demand for poultry products. The supply is not nearly keeping pace with the constant and increasing demand. There is little doubt but that the output of poultry products by the Province could be more than trebled without seriously affecting market prices. In fact, British Columbia, in poultry produce at least, should be an exporter rather than an importer.



Fig. (4.) Farm scene in British Columbia. A pretty rural scene, although the picture reveals a method of poultry-keeping that it is not wise to follow. Turkeys should be rooted away from all buildings or surroundings over which ordinary fowls roam. Fifth and successful turkey-raising will not combine. Sooner or later "black-head" will follow, and once it makes its appearance, turkey-raising generally has to be discontinued owing to its ravages.

Profits are not dependent entirely upon market prices; when markets are good the egg yield also should be good. It is essential to start with the best strains of some of the commercial breeds. Strain probably has more to do with the success than breed. Also for those who intend taking up poultry-raising, experience is necessary; without experience the importance of the details of the business will not be realized and success is almost impossible. This one fact must be borne in mind, that successful poultry-keeping depends upon the regular and accurate performance of many small tasks.

#### INCUBATION.

Since the principles of incubation and brooding have been dealt with at length in Bulletin No. 39, only a few of the more important points will be discussed here.



Fig. (5.) Colony brooder where "round" hay is used.

Both natural and artificial means of incubation are practised the world over. On small farms where only a few fowls are kept the buying of an incubator would not be justified. Here the chicks are hatched by hens altogether. On large plants where extensive methods are employed incubators must be used. The degree of success in artificial incubation depends to some extent upon the machine selected to do the work; therefore care should be

exercised in making the selection. Nearly all of the different machines used are constructed on similar principles, and there is little difference in the hatching power of the hot-air incubator as compared with the hot-water make. Both makes are giving good satisfaction in the Province. Whatever machine is purchased, it should be a good one. A low-grade, cheap machine should never be experimented with. The purchase of the best machine on the market is the best investment in the long run.



Fig. (6.) Chicks of the same age. An illustration of what sometimes happens when pullets' eggs are used for hatching; eggs from forced stock; fabred fowls; lice-infested chicks; improperly fed or improperly housed breeding stock.

The best place for an incubator is one where there are no draughts of air, and yet where the air of the room is pure at all times. It is desirable to have an even temperature in the room, although throughout the greater part of the Province such precautions need not be taken as in the East, since the climatic conditions at hatching time are not as changeable. Incubator houses which give best results are usually those built half in the ground and half out. The best cellar is one with a 4- or 5-foot basement with the rest of the house above ground. Cement is often used in the construction of the basement of the house.

The eggs require to be turned often, twice daily being generally found sufficient. Usually two tests are made, on the seventh and sixteenth day. It is generally advisable to supply moisture during incubation. The object of supplying moisture is to prevent the eggs from losing too much weight. Many fully-formed chicks die in the shell because of lack of moisture. The chicks are weak and the shells are very hard, consequently the little chicks are unable to force their way out.






Fig. (7.) Sketch of first-class hen or A coop. The wire at top admits sufficient fresh air, and yet it is not too cool for any but very severe weather. Early in spring it is advisable to set the coop on a board bottom, attached from the coop to allow of easy cleaning. When no bottom is used, the coop should be either moved forward or to one side, never backwards, because of the danger from feeding chicks amongst filth. A little air-slaked lime should be sprinkled over portion of soil used.



Fig. (8.) Four-compartment setting-coop. After hatching can be made into one broody-coop.

At hatching time the incubator chamber should be kept dark. The chicks should not be removed from the incubator until about thirty-six hours after hatching.

#### **BROODING.**

One of the main things in brooding is the temperature. The chicks' lungs are located along the spinal column, and are not protected very well anatomically. They have a thin membranous lining over them over which a few feathers grow. This is very meagre protection, so meagre that the lungs may become readily chilled, or readily overheated. Then an even temperature, with as few variations as possible is most desirable. On the other hand, the "fireless" brooder is being used with success by many poultrymen in different parts of the Province. It should not be taken up too readily by amateurs. Artificial heat may be supplied the chicks for the first week or ten days, and then the "fireless" system may be found to carry them through for the remainder of the brooding season.

#### **FEEDING.**

Some good commercial "chick feed" is good to start them on, and after they learn to scratch they can be fed dry mash from a hopper. The drier the food for the first few days the better. The dry mash should not

be given them all at once, rather let them have a little from time to time, as in this way they will become accustomed to it gradually, and will not overeat. A good mash may be composed of wheat, bran, shorts, oatmeal, cornmeal, equal parts by measure, and from 5 to 10 per cent. beef scraps. The beef scraps will go towards the formation of bone, which is so essential in the development of the chicken. Another excellent food which serves the same purpose is skim-milk. Green food in some form is necessary. It is surprising what large quantities of green food little chicks will consume. Lettuce, grass-tops, and kale are relished. Charcoal, fine grit, and oyster-shells are other requisites for the growing chicks. The premises should be kept in the most sanitary condition and the food should also be kept sweet and clean. Sanitation is one of the most important factors in the poultry industry, and the most careful consideration should be given while the chicks are young.



Fig. (9) An expensive way of raising chicks. Chicks kept in such small quarters cost more to rear, inasmuch as everything they consume has to be provided for them, and in addition the land room becomes "chicken-tainted," causing disease and mortality. In such quarters chicks need to be supplied with green food. There is no comparison as regards constitution, vigour, and size, between chicks reared in such quarters and those raised on free range, such as orchard, corn, or pasture land.

#### FEEDING THE GROWING STOCK.

All of the chicks are usually in the same flock until about ten or twelve weeks old. About this time the cockerels are separated from the pullets, and all cull pullets which would not develop into suitable breeders are taken out. Different methods of feeding should be employed with the broilers, those intended for roasters, and the females which are being kept for laying and breeding purposes. On all poultry plants there are always a certain number of male birds to be disposed of. With the lighter breeds it is most profitable to sell the cockerels as broilers, while with the heavier breeds it may be as well to hold them, and later make roasters of them. The method of feeding the broilers, roasters, or capons would not vary much. The principal object is to secure as rapid flesh development as possible. A chicken ten or twelve

weeks old will not tend to fatten very rapidly, but if penned up and fed heavily for a few days will add considerably to its weight. Those intended for roasters should develop as large frames as possible, so that when they become full-grown they will put on plenty of flesh.




Fig. (10.) A-shape colony house, suitable for growing stock. The writer has found that this type of house affords too much surface to the sun's rays during the day. However, when placed under trees, it is satisfactory.

Wet mash along with whole grains tends to give quick growth. Variety should be used, the principal grains being wheat, corn, and oats. A mash composed of two parts cornmeal, two parts middlings, two parts beef scraps, one part bran, and one part alfalfa meal may be used. This may be moistened with water or skim-milk, and fed once or twice a day. Different ground grains may be used, but the important thing is to have a forcing mash. Broilers may be sold as squabs when weighing from 12 oz. to 1 lb., or held a few weeks longer, when they weigh about 2 lb.

#### FEEDING THE ROASTERS.

There is a great scarcity of properly dressed poultry on the markets of the Province. The broiler trade is developing and roasters are in great demand, also dressed turkeys, ducks, and geese. Prices for all forms of dressed poultry are exceptionally good.

In fowls to be fattened for roasters the first point we should look for is a good constitution in order that they may be fattened early and economically.

A good market bird should have much the same characteristics as a choice beef animal. It should be fairly low set, broad-breasted, and, as a rule, should be rather short in the bill, bright in the eye, and broad in the head.




Fig. (11.) B or secondary coop. Dimensions: Length, 6 feet; width, 2 feet 6 inches; height, front 2 feet 4 inches, back 18 inches. A hen's brood of chicks, at about eight weeks of age, should be placed in here from smaller quarters, or the B coop itself makes a splendid coop for hen and chicks from the start. When using the coop for this purpose, the roosts can be removed, and roost-holes in ends covered temporarily with a shingle tacked over to keep out draughts and vermin.

Crate fattening is more economical than pen fattening, although the latter practice is often resorted to because of the lack of proper facilities for crate fattening. Birds fattened in crates, if properly fed, will be ready for market about three weeks after being put in the crates. A very easily made and efficient fattening-crate can be made of laths, and a few light boards which are used for ends and partitions. The crate should be 6 feet 6 inches long, 20 inches high, and 16 inches wide. The top, back, and bottom are formed of laths, running lengthwise. The laths on the top and back should be about  $1\frac{1}{2}$  inches apart. The slats on the front run up and down and are placed about 2 inches apart, so that the birds may eat from the V-shaped trough in front of the crate. The laths on the bottom are usually placed  $\frac{3}{4}$  inch apart. Care should be taken to have the back slat on the bottom at least  $\frac{3}{4}$  to 1 inch from the back, so that the droppings may pass through and not accumulate.

A V-shaped trough, 2 inches deep and  $2\frac{1}{2}$  inches wide at the top, inside measurements, is placed in front of the crate on brackets, the trough being raised about 2 inches from the level of the bottom of the crate. The crate when finished should stand on legs about  $2\frac{1}{2}$  to 3 feet high.

For the first few days after the birds are placed in the fattening-crate they should not be fed too heavily. The success of fattening depends to a large extent upon the feeding of the birds for the first few days. The crops should never be too well filled as the appetites should be kept keen. If this method is practised, it will be found that the birds will feed better throughout the fattening period than if fed too heavily at first. No more food should be given at any time than the birds will eat up clean. If all food is not eaten the troughs should be taken away or the feed removed, and the troughs turned over. Feeds should be, if possible, twelve hours apart, and the food should be mixed at least twelve hours before feeding.

The following are good mashes which may be used for fattening:—

1. Two parts oat-chop, one part cornmeal or barley-chop, and one part low-grade flour.
2. Equal parts low-grade flour, barley-meal, and wheat middlings.

These mashes may be mixed with buttermilk or skim-milk. Milk is an excellent food for fattening chickens, as it tends to develop the tissues and apparently improves the quality of the meat. The proportion of milk to dry mash is about double the quantity of milk by weight or measure.



Fig. (12.) This type of colony coop could be improved by having more space open in front, and lowering the opening. As depicted, the rain can obtain easy ingress and the sun's rays can only reach a very small space at the back. Would place wire door in front, fitted with muslin-cloth frame.

#### FEEDING THE LAYING STOCK.

As soon as possible the pullets should be placed in colony houses and allowed free range. The principal object is to have them fully matured into healthy fowls with plenty of constitutional vigour. The stamina should be kept up so that they may be able to keep in the best of health when heavy egg production is demanded of them. The health of the laying stock has a great deal to do with success in poultry-raising, for heavy egg production is principally a problem of how to maintain laying hens in a high state of health. Before any marked difference may be expected from

adopting improved methods of feeding, the fowls must necessarily have plenty of constitutional vigour, with inherited stamina. They must be supplied with an abundance of fresh air, day and night, and must also be induced to take plenty of exercise.



Fig. (13.) Small house, suited to Coast conditions. The kerosene can was inserted in the wire screen to allow birds egress and ingress, and as a protection against vermin.

The hen is simply a manufacturer, manufacturing the raw materials, the food, into the finished product, the egg. The food serves various purposes, part of it to furnish energy to carry on the various activities of the body, and to keep the body warm; other purposes being that of building up the tissues and organs, and keeping them in repair, and to supply material for egg production. For these various purposes different classes of nutrients are demanded. The heat and energy required by a fowl are derived mainly from the fat and a number of other carbonaceous materials in the food termed carbohydrates. The organic part of the bone, the tendons, the internal organs, and the muscles in the body of a hen are derived from the nitrogenous constituents of the food, commonly termed protein. Then to give substance to the bones, and other various organs, and also for the formation of the shell of the egg, ash constituents are required. Then there are three classes of nutrients, carbohydrates, protein, and ash, which are used in feeding for egg production. The egg is also composed of these three classes of nutrients. The object in feeding is to give the hen such foods which are composed of the same constituents as the egg—for it is the egg we are after—and in as near the same proportion as possible. The nutritive ratio of a food is the proportion of digestible protein to digestible carbohydrates, and in the body of the hen and in the egg it is found to be as two to one. Now, the




Fig. (14.) Illustration shows colony system of breeding pens. The advantages of this type of house are that they can be moved to any part of the ranch thus enabling the owner to practice rotation of crops, at the same time sweetening the soil.



majority of the more staple grains have much wider nutritive ratios, which shows that they are deficient in protein. To make up this deficiency in protein, foods rich in nitrogen must be used. Speaking practically, this means that laying hens must have a variety of grain, and the better the proportion in which the grains are fed the more economically can the hen convert the food into the finished product.

### DIGESTIBLE CONSTITUENTS OF VARIOUS FOODSTUFFS.

| Food.              | Dry Matter, per cent. | Ash or Mineral Matter, per cent. | Protein or Muscle Formers, per cent. | Equivalent Carbohydrates or Fat Formers, per cent. | Nutritive Ratio. | Manorial Value in pounds. |
|--------------------|-----------------------|----------------------------------|--------------------------------------|----------------------------------------------------|------------------|---------------------------|
| Wheat .....        | 86.0                  | 1.8                              | 10.2                                 | 73.0                                               | 1:7.2            | .420                      |
| Oats .....         | 86.0                  | 3.0                              | 9.2                                  | 56.8                                               | 1:9.2            | .378                      |
| Corn .....         | 86.0                  | 1.5                              | 7.0                                  | 70.4                                               | 1:9.7            | .326                      |
| Barley .....       | 86.0                  | 2.4                              | 8.7                                  | 69.2                                               | 1:7.0            | .288                      |
| Penn. ....         | 90.0                  | 2.0                              | 10.8                                 | 53.4                                               | 1:3.2            | .548                      |
| Buckwheat .....    | 87.0                  | 2.0                              | 7.7                                  | 53.3                                               | 1:3.9            | .247                      |
| Rice .....         | 87.0                  | 0.4                              | 4.8                                  | 72.0                                               | 1:15.2           | .175                      |
| Kaffir corn .....  | 81.8                  | 1.5                              | 7.8                                  | 69.2                                               | 1:8.1            | ...                       |
| Sunflower seed ..  | 92.5                  | 2.0                              | 12.1                                 | 85.8                                               | 1:7.1            | .428                      |
| Wheat bran .....   | 88.0                  | 1.0                              | 9.0                                  | 70.0                                               | 1:7.1            | .320                      |
| Wheat middlings .. | 88.0                  | 3.8                              | 12.2                                 | 65.3                                               | 1:3.7            | .618                      |
| Gluten meal .....  | 92.0                  | 0.8                              | 25.8                                 | 60.7                                               | 1:4.7            | .471                      |
| Red clover hay ..  | 85.0                  | 0.2                              | 6.8                                  | 66.0                                               | 1:2.5            | .774                      |
| Alfalfa hay .....  | 92.0                  | 7.4                              | 11.0                                 | 42.3                                               | 1:3.8            | .431                      |
| Linseed meal ..... | 92.1                  | ...                              | 11.3                                 | 63.0                                               | 1:5.5            | ...                       |
| Dried scraps ..... | 80.3                  | 5.3                              | 27.2                                 | 37.0                                               | 1:14.0           | ...                       |
| Dried blood .....  | ...                   | 4.1                              | 68.4                                 | 30.7                                               | 1:0.4            | .175                      |
| Green bone .....   | ...                   | 4.7                              | 52.3                                 | 5.0                                                | 1:0.1            | 2.128                     |
| Skin-milk .....    | ...                   | 11.5                             | 18.0                                 | 43.0                                               | 1:5.3            | ...                       |
| Pontones .....     | ...                   | 0.7                              | 2.0                                  | 5.0                                                | 1:4.6            | .103                      |
| Mangel .....       | 21.0                  | 1.0                              | 0.9                                  | 10.5                                               | 1:18.3           | .075                      |
| Cabbage .....      | 13.0                  | 1.1                              | 1.1                                  | 10.4                                               | 1:5.1            | .051                      |
| Onion .....        | 15.0                  | 1.4                              | 1.8                                  | 9.1                                                | 1:5.1            | .082                      |
| Rape .....         | 11.0                  | 2.0                              | 1.5                                  | 8.0                                                | 1:5.4            | .062                      |
| Red clover (seed)  | 20.0                  | 2.1                              | 2.0                                  | 10.4                                               | 1:5.6            | .107                      |

### COMPARISON- COMPOSITION OF RAW MATERIALS AND FINISHED PRODUCTS.

#### Raw Materials.

|                   | Water per cent. | Ash, per cent. | Protein per cent. | Carbohydrates per cent. | Fat, per cent. |
|-------------------|-----------------|----------------|-------------------|-------------------------|----------------|
| Wheat .....       | 1               | 1.8            | 11.0              | 71.0                    | 2.1            |
| Oats .....        | 11.0            | 3.0            | 10.8              | 56.7                    | 5.0            |
| Corn .....        | 10.0            | 1.5            | 10.4              | 70.3                    | 5.0            |
| Barley .....      | 10.0            | 2.4            | 12.4              | 69.8                    | 1.8            |
| Rice .....        | 12.1            | 0.4            | 7.1               | 79.2                    | 0.4            |
| Beef scraps ..... | 1.3             | 8.0            | 58.0              | ...                     | 32.9           |
| Green bone .....  | 0.9             | 24.5           | 22.3              | ...                     | 16.5           |
| Skin-milk .....   | 99.6            | ...            | 2.0               | 5.2                     | 0.3            |
| Grass .....       | 76.4            | 2.4            | 2.3               | 13.8                    | 1.0            |
| Beets .....       | 90.9            | 1.1            | 1.4               | 5.5                     | 0.2            |

#### Finished Product.

| Hen .....       | 55.8 | 3.8  | 21.6 | ... | 17.0 |
|-----------------|------|------|------|-----|------|
| Pullet .....    | 55.4 | 3.4  | 21.2 | ... | 18.0 |
| Capon .....     | 41.6 | 3.7  | 19.4 | ... | 33.9 |
| Fresh egg ..... | 65.7 | 12.2 | 11.4 | ... | 8.9  |

Among the more staple grains are wheat, oats, barley, and corn. Wheat is one of the best grains for the poultry plant, though some poultrymen claim it to be an expensive food. However, when the net results are balanced up, poultrymen in British Columbia can readily afford to use this valuable grain. Since it is a staple of human food, the price of good wheat usually runs high as compared with other grains. Broken wheat, frozen wheat, and good wheat screenings are as good for poultry-feeding as the very finest milling wheats.

Oats make a very good poultry food, but when poor in quality the percentage of hull is sometimes very large and such oats are not very palatable or digestible, hence should not be used. When fed dry or in a mash, they are more relished by the birds. Sprouting the oats by soaking them for about twenty-four hours increases their palatability. Oatmeal or rolled oats make a good food for young chicks, and ground oats without the hull is excellent for fattening fowls.

Barley, like oats, has a large percentage of hull, but it is a valuable food.

Corn is one of the most commonly used of all grains, especially in the great corn belts. Corn is rich in fat and forms an excellent food when used with other grains. Cracked corn is good as a chick food, while cornmeal is a good fattening food, though it tends to produce the undesirable yellow flesh.

Besides the whole grains, there are the mashes, green foods, grits, oyster-shells, and other foods.

The dry-mash system is used much more than the wet-mash system. In feeding dry mash, different ground grains are used and mixed together. This mixture is kept before the fowls all of the time, in hoppers.

Some good mashes which are largely used are as follows:—

1. By measure, four parts bran, two middlings, two ground oats, one cornmeal, one ground barley, one-half flaxseed-meal, and one-half beef scraps.
2. By measure, four parts bran, four ground oats, two cornmeal, one middlings, one cut alfalfa (well cured), and one part beef scraps.
3. Equal parts by measure of bran, low-grade flour, and barley chop or meal.
4. By weight, two bran, one cornmeal, one middlings, one gluten-meal, one flaxseed-meal, and one beef scraps.

In wet-mash feeding many by-products may be economically disposed of, though this method involves more labour, especially with large flocks, than dry-mash feeding. Small potatoes or other refuse may be boiled and mixed with the mash and fed. Probably the best time to feed a wet mash is at noon.

The following makes a good mash:—Bran, shorts, ground oats, cornmeal, 10 per cent, beef scraps, and green food, such as clover or alfalfa. Soak the alfalfa-meal over night in boiling water and mix with mash next afternoon, then feed.

Alfalfa, clover, and cut-grass or lawn clippings are two sources of green food. Mangel form one of the cheapest foods for the laying hens and are easily grown. If the flock is to produce maximum results it must be liberally supplied with nutritious food.

It is also necessary that animal food be supplied. If beef scraps or green bone cannot be obtained, then blood-meal should be used. Experience has proved, however, that breeders should be careful in feeding concentrated material. An overabundance will result in ovarian trouble, and digestive diseases. Grit, oyster or clam shells, and pure water are essentials.

One of the most successful methods in feeding laying hens is to feed them some whole grain, wheat, in the early morning as the fowls come from the roosts, and at noon give them some green food, sprouted oats, and at night a good feed of corn or corn and oats is given. The whole grains should be scattered in litter 12 or 14 inches deep, as in this way the fowls are induced to take plenty of exercise. Dry mash is kept before them all the time and they are given liberal quantities of green food in the form of alfalfa or mangels. Clean water, grit, and shells are never wanting.

#### EGG PRODUCTION.

However careful the poultryman may be in feeding, he must not expect appreciable results unless he has good laying stock. The best strains of the proper variety should be selected. The strain of a variety is more important than the variety itself, although the variety has an important commercial aspect.

At present the average number of eggs per hen is not over seventy, but the average hen in every flock should produce at least 150 eggs per year.

A poor layer is a poor payer. The unprofitable hens should be weeded out, thus saving loss in the amount of feed consumed, and at the same time giving the good layer a better show. Different methods of selection may be employed. Up to the present, trap-nesting cannot be recommended for commercial plants on account of the labour involved. Trap-nesting is followed by many poultrymen, and each hen that is being trap-nested wears a leg band which is stamped with her number. A record is kept of the number of eggs she lays, and by this method the unprofitable hen may be weeded out. It requires considerable time in keeping the record and in releasing the birds from the nest, though it has many distinct advantages. In the hands of those who understand the principles which must be kept in view, this system is an excellent one. It needs to be very carefully applied. Some breeders test the laying qualities of their hens by means of the pelvic-bone test. If this test is to be satisfactory it is necessary to test the fowls during the months of November, December, and January. Any fowl showing good width between the pelvic bones during these months can generally be relied upon to do her share in filling the egg-basket.

The thing of importance is to breed a heavy egg-laying strain that will reproduce its egg-laying characteristics in the offspring, at the same time maintaining the highest possible degree of vitality.

Finally, as we have seen, the question of egg production involves many factors. The principal problem is how to maintain the heavy-laying hens in a high state of health. Inherited stamina, fresh air, exercise, and food are necessities if constitutional vigour is to be maintained. All of these factors must be considered when endeavouring to improve the qualities of any strain of birds.

#### POULTRY-HOUSES.

The chief factors in poultry-house construction are the cost of the house, the comfort of the fowls, and the proper provision of light and ventilation. Expensive houses are unnecessary and often are not as convenient as they should be. The cost of the building is influenced by its shape. It costs more to house flocks in colony houses than in a continuous house. On the other




Fig. (15.) Continuous open front house.

hand, on the colony system the fowls are on free range, and can be fed cheaper than when they are kept in large flocks. Naturally the cost of labour is greater when fowls are kept in small flocks, yet notwithstanding the amount of labour required to do the extra work in connection with the colony plan, there are many advantages. Expense of fencing and congestion of flocks is avoided. Colony houses may be built on cleared land and even in wooded areas, and the fowls can be hopper-fed. Where a large number of fowls are being kept the continuous house will give good results with much less labour, but proper precautions must be taken to prevent the outbreak of disease through soil contamination and filth.

It is very necessary that the house be built on a dry location, a location where there is ample drainage, and where, if possible, the subsoil is of a gravelly or open nature. The floor of the house should be somewhat elevated above the level of the land, so as to be sure of drainage. The houses should be faced to the south or south-east, with as much as possible of the front open. We do not advise the use of drop curtains in any part of the building, except, possibly, during cold snaps in winter. The house must be free from draughts at all times. Ventilation and abundance of fresh air are two of the most essential requirements in the hen-house, if the flock is to be kept in a healthy condition. The old idea that the house must, during cold weather, be shut up close and kept warm has long since died out, and it has been proven that hens give best results when allowed to run and work in the fresh air during the day. One essential factor is to have the birds roosting in comfortable quarters at night, although the air should be pure. For this reason, we do not advocate the use of a drop curtain before the roosts, except in districts where the temperature falls to a very low degree; but we do maintain that the roosts should be so placed in the house as to avoid all danger from draughts and other detrimental influences.




Fig. (16.) Woods fresh-air house. A type of house eminently suitable for small breeding-pens, or small laying flocks. Its construction—low-down front—makes it somewhat difficult and expensive to clean out when used as a long-continuous house.

Another satisfactory house is found in the "Woods" fresh-air house. (Fig. 16.) This house is of the semi-monitor top type and bears out the principle of the "Tolman" house, providing some advantages. Each section is about 7 feet deep, making the house, in all, 14 feet deep. If built on the continuous plan the partitions may be placed every 15 or 20 feet. The actual

opening in front may vary in different sections of the Province. A board 10 or 12 inches wide is necessary at the bottom to prevent the fowls from scratching the litter out and also to act as a wind-break. The house is  $4\frac{1}{2}$  feet high in front, and where the top and bottom boards are used the actual front opening would be  $2\frac{1}{2}$  feet, whereas it would be  $3\frac{1}{2}$  feet where the top protection board is not necessary. A row of windows is placed along the top part of the back section, and this permits of the sun's rays reaching the roosting quarters. These should be on hinges, so that they can be opened when desirable. When opened during warm weather a free circulation of air is obtained and the construction of the house provides for comfort in winter. When properly constructed it is serviceable and durable.




Fig. (17.) Inside arrangements of hen-house, showing roosts, droppings-boards, nests. The nests, roosts, and droppings-boards should be removable, so as to allow of disinfection.

#### General Requirements.

In brief, a cheaply built house with an open front will give equally as good results as, if not better than, a more expensive house. Not only will fowls lay more eggs if the house is supplied with plenty of fresh air, but the hatchability of the eggs from such houses will be greater, and a stronger and more thrifty bunch of chickens will be the result.

There are many plans of houses which receive a wide use, and no one can truthfully say one plan is the best. The house must suit the local conditions, but it is advisable to get as much sunlight and fresh air into the house as possible. Interior arrangements should be simple and as convenient as possible. The nests may be under the droppings-board, so as to give the fowl all the floor space possible. The roosts should not be too high from the floor, or there is danger that the fowl may injure its feet in jumping from the

perches to the floor. The dropping-board for a single roost should be from 18 inches to 20 inches wide, and 3 feet wide for a double roost. Roosts may be made of 2 by 3 scantlings, and the edges may be rounded. For conveniences in cleaning the dropping-board, it is well to have it fairly low down.

#### LITTER.

The number as well as the fertility of the eggs depends largely upon the vitality of the flock. The vitality, in turn, largely depends upon the exercise the hens are given. It is only natural for a hen to work for most of her living; consequently, the reproductive system will be better nourished than if the hen were fed large quantities of food only a few times a day and induced to take but little exercise. Exercise helps to start egg production. The litter should be quite deep, from about 8 inches to 12 inches. It should be kept in such a condition that the hens may readily scratch down to the floor, thus being able to pick up every grain that is in the litter. The fowls should keep the litter free of grain. If, some time after feeding, any grain except a few coarse oats are left in the litter, there is something wrong with the method of feeding. The flock should not be fed too heavy in the morning or throughout the day; but at night the fowls should be given all they will pick up clean. Through vigorous exercise the fowls will become hardier, and when cold weather comes on they will be better able to resist the change and will make better breeders and layers.



Fig. (18.) A good type of colony house. Measurements, 6 feet long, 8 feet deep, 6 feet 6 inches in front, 2 feet 6 inches rear. Colony houses can be hauled to any part of the ranch, thus giving young or old stock opportunity to range on clean soil. Can be used as cockerel, pullet, or breeding houses after chickens are matured. The top shutter or board window makes a splendid hood or rain-shield and can be changed to a muslin-curtain screen if desired.

### CLEANING THE HOUSE.

Cleanliness is one of the most important factors to be considered in the poultry industry. The house should be cleaned occasionally and the roosts painted with kerosene or some other disinfectant. The interior of the house should be white-washed frequently with about 5% carbolic acid used. The nests must be kept clean, since eggs readily become tainted.

### MOVABLE HOUSES.

Conditions in this Province are quite favourable for the use of movable houses. The most satisfactory house is one built on runners, so that the house may be moved with a horse or team. Movable houses are used principally for growing stock and breeders. They may be drawn from place to place over the farm, giving the fowls new ground. Stump lands and orchards are excellent fields for poultry. Usually some form of open-front house is used.

### POULTRY RUNS.

For success with poultry, they should not be allowed to run year after year on the same land without the soil being turned over and a green crop sown. It cannot be too strongly emphasized that some form of soil cultivation should be linked with the poultry industry. It has been proven that many diseases which break out in the flock arise from a contaminated soil. On commercial plants the whole plant may be divided into two parts by one wire fence running through about the centre of the plant. This will allow of the cultivation of one part and the sowing of some crop, while the fowls have the range of the other part. The land should be top-dressed with lime frequently and a solution of sulphuric acid may be used. Keeping the soil free from contamination has a great deal to do with the success of the industry. Wherever possible, shade should be provided in the runs. Fruit-trees, sun-flowers, kale, or corn is excellent.

---

### MARKETING POULTRY PRODUCTS.

A large part of the profits in poultry-keeping depends on the marketing of the products. The poultryman must be a good salesman as well as a good raiser of poultry. He must have good markets to begin with, or know how to make them. With high-grade products that somebody wants, and that most people will not take the trouble to supply, a high-class trade can be built up through skill and tact. The nearer the producer can come to the consumer in selling his products the higher will be the price realised. The producer should strive to eliminate the middleman. The extra price per dozen which is paid for quality is almost clear gain. There is fair profit only in producing eggs and other produce at regular market prices. The largest profits must come from superior marketing and from special market advantages in selling eggs and other produce. A special market advantage has been introduced into British Columbia at Duncan, where a co-operative egg station has been established. This is only one instance of the many co-operative establishments which might be advantageously introduced into this Province.



### MARKETING EGGS.

Fig. 21 shows eggs from a leading grocery store. The eggs were in no way selected. Dirty eggs are in evidence. Poultrymen should never send dirty eggs to the market. The shells of the eggs are very porous and allow gases and bacteria to enter the egg. If the egg is laid in filthy nests or is packed in unclean cases, it soon becomes tainted and unfit for use.



Fig. (19.) A method sometimes employed when shipping high-grade eggs in large consignments.

The egg is at its best when just laid, and the longer it takes to place it in the consumer's hands, the less it is worth. Eggs, when prepared for market, should be spotlessly clean. They should be packed in clean cases, of which the thirty-dozen size is preferable. Absolutely clean fillers and cardboards should be used. The quality of the eggs should be unquestionable. It is only in this way that a good trade can be worked up.

The egg industry in Denmark has been enormously improved owing to the application of the principle of co-operation to the collecting, grading, packing, and export of the eggs. The great progress of Denmark during the past fifty years may be at once attributed to the wonderful system of education available for its people, and to the extent to which the principle of co-operation has been adopted in every part of the country and in every phase of its national industry.

There has lately been formed through the instrumentality of Prof. F. C. Elford, of Macdonald Agricultural College, a Poultry Producers' Association in Eastern Canada, based on the co-operative system. They "do not hope at first to revolutionize or change the whole poultry industry," but they do, nevertheless, hope to remedy certain conditions which are greatly to the disadvantage of the industry as a whole. The Egg Station at Duncan was

established with this end in view. The principal object of the station is to eliminate the "middleman" and thus increase the profits of the producer. A good market is sought where the best quality of produce is offered. Quality sells itself.




Fig. (20.) Thirty-dozen egg-crates, as commonly used in the Province. When eggs are kept for any length of time in these crates, it is important to see that the cardboard fillers or partitions are dry and clean. Mouldy or damp fillers will deteriorate the eggs very quickly.

At present thousands of dollars are lost annually in British Columbia through bad eggs. Consequently, it is of vital importance that the poultrymen throughout British Columbia produce sufficient quantities to supply the home trade; and, finally, to solve this vital problem, we must have, first, a campaign of education among egg producers; and, second, a system of buying eggs that will guarantee a price according to quality.

#### MARKETING DRESSED POULTRY.

The same principles hold good for dressed poultry as for eggs. A little care and skill in preparing poultry for market will often make a large increase

in the selling price. The expense of killing, picking, and packing poultry is small in comparison with the first cost of raising. All poultry should be carefully assorted according to size, colour of shanks, age, and condition. Each case should be uniform in quality.



Fig. (21.) Basket of eggs as generally exhibited in stores—all ages, all sizes, all colours, clean, dirty, cracked, and broken. A basket of fertile eggs placed in the sun for about six hours at a temperature exceeding 70 degrees will start hatching. Eggs for sale should be kept in shaded parts of store or dry cellar, and cards inserted in window announcing the fact.



Fig. (22.) One dozen cartons. These are of cardboard and cost in some localities from \$12 to \$16 per thousand, the latter price including printing.

**Market Requirements.**—Fattened chickens when prepared for the consumer should conform to the following market requirements:—

The dressed chicken should present a neat, clean, plump appearance.

The preferable weight is about 4 lb. Plump, crate-fed chickens up to the weight of 5 lbs. are more readily disposed of than heavier ones.

The shape of the breast should be long and broad, to give a well-mented appearance.

The legs should be as short as possible, indicating the low, blocky type.

The colour of the flesh should be white, and the grain fine.

The minimum of bone and offal are two important factors in the best grade of marketable products.

Do not market lean chickens. It is an unprofitable business both to the producer and the consumer.

**Killing, Plucking, and Packing.**—Before killing, the birds should be fasted at least twenty-four hours, which will prevent food remaining in the crop and the intestines which would decompose and spoil the flavour of the meat. Several hours after the last feed allow the birds all the clean water they want, and then give them a complete fast until killed. The best method of killing is to stick the bird in the mouth, braining it, and at the same time bleeding the bird by cutting the arteries.




Fig. (23.) Illustration of shaper in frame.

Suspend the bird by a cord from the ceiling of the killing-room so that it will be as low as the operator's shoulder. To bleed, place the left arm over the bird, grasp the bird by the neck with the thumb on the throat. With the right hand insert the knife and sever the arteries by drawing the knife down, first on one side and then by a quick half-turn down the other side of the mouth. Then pierce the brain by passing the knife through the roof of the mouth towards the top of the head and give it half a turn. To be successfully

bled, both arteries must be severed. These arteries are exposed on each side of the mouth. If the fowl squawks when stuck, the operator will know that the operation has been successful.

Plucking should be commenced at once, pulling the body feathers first, then the wing and tail feathers. As soon as the fowl is rough-plucked the plu-feathering and finishing can be done more quickly in a sitting position.

Returns are greater when shipped in a compact, plump condition than in a rough, unprepared condition. All birds for market should be properly shaped.

The shaper is shown in Fig. 23, and is made by nailing two ¾ by 6-inch plain boards together at right angles. The trough should be nailed into a frame and inclined slightly backwards, as shown in Fig. 35.




Fig. (24.) Showing fowl in shaper. Should weights be used to compress bird, it is advisable to place clean paper next to flesh. Dressed fowls should not be shipped until all animal heat has escaped. Warm fowls will show blackened or green flesh upon unpacking after a short journey during hot weather.

With legs placed alongside the breast and with the breast downward, force the bird into the angle of the shaper, cover with paper and lay a brick on the back and one against it, to hold it in position. The shaping should be done in as cool a temperature as possible without freezing, and should be continued for about twelve hours. When thoroughly cooled the birds are placed in shipping cases, as shown in Fig. 25. Cases of basswood or spruce, with corners lock-joined, are most satisfactory. The box should be lined with parchment paper.

There are different methods of packing. The general method employed in shipments into British Columbia from the East is by packing the birds in a single row. Packing the birds in two rows is employed for short shipping distances. The case should be properly stencilled, giving the name and address of producer, number and weight of birds, and whether they are chicken or fowl.



Fig. (25.) Showing method of packing dressed fowls for shipment. It is important that the mouth and feet of birds should be washed or cleansed with moist rag or cloth. Filth clinging to shanks or feet will generally adhere to breast of fowl when unpacked, causing an unsightly appearance.

## PESTS, PARASITES, AND DISEASES.

### PESTS.

**RATS.**—The rat is the worst mammalian pest known to the poultryman. Its depredations everywhere result in losses amounting to thousands of dollars annually, and though thousands are killed, other thousands soon replace the slain. The most important step toward the suppression of the rat is to adopt some method to prevent it from increasing in numbers so rapidly. Restricted supplies of food means fewer rats. Their bill of fare includes seeds and grass of all kinds, flour, meal, eggs, chicks, young turkeys, ducklings, andlings, and other foods. It has been found that it costs at least 25 cents every year to feed one rat, and this on the cheaper foods alone. The failure of efforts to exterminate this rodent has not been due to lack of effective methods so much as to negligence and the absence of concerted action.

The more important means of fighting rats are: (1) Natural enemies of the rat; (2) Rat-proof construction of buildings; (3) Keeping food from rats; (4) Driving away rats; (5) Destroying rats. Trapping, if persistently followed, is one of the most effective ways of destroying rats. Another method is by the use of strychnine sulphate. The dry crystals may be inserted in small pieces of raw meat or toasted cheese, and these placed in rat runs and burrows.

The following method is recommended for poisoning rats where the rats are among chickens. Take two wooden boxes, one larger than the other, and each having two or more holes in the sides large enough to admit rats. The poisoned bait should be placed on the bottom and near the middle of the smaller box, and the larger box should then be inverted over the other. Rats have thus free access to the bait, but fowls are excluded.

**HAWKS, CROWS, ETC.**—In some parts of the Province it is almost impossible to raise poultry without very heavy losses, because of the depredation of these birds. An ordinary milk-trap (rusty ones are best) placed on top of a pole at least 15 feet high, and securely chained, will help to rid the flocks of these enemies. The trap need not be baited. Festoons of tape or white strings strung across rearing yards near the coops and second-hand fish-nets have proven very satisfactory in warding off birds of prey. Poisoned infertile or worthless eggs, placed out in a field frequented by crows, skuks, milk, etc., are also successful remedies. Crystallized form of strychnine is one of the best poisons to use. The eggs should be placed in the soft, large end up, and the top removed, and a few crystals of the poison inserted and mixed with the contents of egg. These should be placed out over-night, and removed before chicks are released in the morning.

### PARASITES.

**LICE.**—Lice probably exist in small numbers wherever there are fowls, but as long as fowls are healthy and active lice cannot increase rapidly enough to seriously annoy the birds. They can only become troublesome from gross neglect, either of the fowls or of their habitations. In the one case the remedy is some good common insecticide placed in the dust-bath, which may be mixed powdered tobacco or a little sulphur; in the other an energetic lime-washing of the houses and interior arrangements, with a free use of a solution of carbolic acid.

Lice in the poultry-houses and on the fowls means loss of money through decreased egg production. The interior arrangements of the houses should be simple and convenient, so that spraying and disinfecting may be easily done, giving as little chance as possible for vermin to increase.

**SCALY LEG.**—This is caused by a mite which burrows under the scales of the feet and shanks. Wash every morning with strong carbolic soap, after soaking bird's legs in warm water for at least ten minutes on the first day, and apply sulphur ointment or carbollized vaseline at night.

#### DISEASES.

Disease does more injury than any other factor. "Prevention of disease" should be the motto for the poultryman. Fresh, air, abundance of sunlight, clean water, and clean quarters are necessary for success. The poultryman should watch his flock carefully, and on noting the first symptoms of sickness among any of his birds should at once remove that bird to quarters where it cannot come in contact with any healthy birds. This bird should be treated according to the disease. The general remedy for all such birds is the axe, and the carcase burnt, not buried. Neither does one gain anything by trying to cure deformed or diseased chicks.

#### COLDS.

Colds affect fowls variously, and if neglected often lead to something serious. Changeable weather and continuous damp weather often make colds epidemic in the early fall. A common cause of colds and distempers is closing the houses too close at night. Often colds will be spread among young pullets by bringing them in from the range to close, dark buildings.

Simple remedies should be used at once; give neonite in the drinking-water. A good remedy to keep on hand for colds is equal parts of cayenne pepper, ginger, and mustard, mixed as stiff as possible in lard, then flour worked in to make a stiff dough. Form in small pellets and give by opening the mouth and dropping down the throat. Disinfect drinking water by means of permanganate of potash. Feed all sick, especially runpy, fowls on soft feed, and see that Epsom salts are given every second day: 1/2 teaspoonful being ordinary dose for adult fowls.

#### WHITE DIARRHOEA.

At present information pertaining to white diarrhoea is rather indefinite; the latest investigations disclose two diseases under the one name, "Diarrhoea." The case characterized by cheesy feces in the lungs are caused by the mold *Aspergillus fumigatus*. The cases characterized by distended and cheesy ceca invariably show the presence of cecidia. This disease is receiving a great deal of attention from poultrymen and scientists in the East, though it seems to be on the increase in this Province. The health of the flock and the cleanliness of the incubation quarters will determine to a large degree the prevalence (or not) of this disease. The soil should never be allowed to become contaminated.

#### EGG-BOUND.

Treatment is rarely successful. Give dose of castor oil, wash the vent with warm water and pass in a feather dipped in sweet oil, or hold hen over a pail of boiling water for about twenty minutes.



**GAPES.**

This is caused by gape-worms in the wind-pipe, which finally kill by either wasting or actual suffocation. Place the birds in a large box covered with a coarse cloth and dust air-slaked lime on the cloth. The lime breathed in by the birds causes the worms to relax their hold and they are coughed up. Destroy the worms.

**LEG WEAKNESS.**

In chicks it is often due to excessive heat in brooder, and through chicks being kept indoors too much during the first few weeks. In mature stock it is generally due to forelug or to lack of bone-forming material. Discontinue high feeding, giving more green food, fresh air, and where board floors are used, place several inches of earth on, and then litter scattered on this.

**BLACKHEAD.**

Blackhead has been found in chickens, ducks, and turkeys as well as other birds, but is most fatal to turkeys. Many flocks have been killed off through the ravages of this disease. The nature of the disease makes it very difficult of eradication since the disease may reach a very advanced stage before any noticeable symptoms are exhibited. The victim then begins to mope about with drooping wings and lowered head. It may die very shortly or it may linger on for a long time. Naturally the head is discoloured and the bird presents a very unhealthy appearance. The disease was first called infectious entero-hepatitis, and it has been found to be due to a minute parasite called *Amoeba meleagridis* or the amoeba of the turkey. Of late, however, the parasite is claimed to be a *Coccidia*. The disease begins in the ceca and generally affects both organs. Once in the tissues the organisms multiply rapidly. The wall of the affected cecum is thickened and the inflammation may extend to other organs of the body. Associated with this condition of the ceca, there is in nearly all cases a serious disease of the liver. The liver is very much enlarged, and over its surface yellow spots appear, usually being slightly depressed below the surface. In the liver the parasites begin to multiply and spread in all directions.

During the course of affection the parasites multiply in the ceca and become mixed with the intestinal contents, many of them being discharged with the contents. This gives rise to a rapid spread of the disease. The food especially, where the turkeys are continually fed in the same place, becomes contaminated, and healthy birds devour the parasites which pass through the digestive canal into the ceca where they multiply.

Treatment of diseased birds is useless. The most obvious precaution is immediate isolation of affected birds. All dead birds should be burned and the greatest care should be exercised to prevent the spread of the disease to healthy birds. Yards and houses which have been exposed to blackhead should be thoroughly disinfected. Poults should be raised on entirely new ground, the old ground having been sprinkled with lime and ploughed. Where the disease is present the whole flock should be killed and burned, and no more turkeys should be raised for a few years. It is next to impossible to rear chickens, ducks, or turkeys together satisfactorily. Sooner or later the turkeys will suffer.

### TUBERCULOSIS.

Tuberculosis in fowls is a more serious disease than poultrymen realize. The disease exists extensively among poultry in the Province, but seldom kills enough fowls at any one time to excite the alarm of the poultryman. The peculiar nature of the disease and its method of infection greatly increases the necessity for preventive measures being taken.

There seems to be no noticeable symptoms until the disease has progressed far enough to cause emaciation or weakness, lameness often occurs, and the fowls usually begin to lose weight. In the last stages of the disease, the affected bird crouches, apparently from weakness. The comb becomes pale, the feathers ruffled, and in some stages of the disease some fowls may show tuberculous nodules on the skin of various parts of the body. Frequently these nodules appear on the skin of the face and the throat. The liver is nearly always affected, being studded with cheesy spots throughout. The spleen may be affected to such an extent that it is abnormally enlarged. The peritoneum or serous membrane of the abdominal cavity may be covered with tubercles the size of a pin-head or smaller, and when this occurs the abdomen contains more or less liquid. The intestinal walls are often thickened or ulcerated, and contain tubercles of various sizes. As with blackhead and enteritis (a very common disease amongst poultry and often mistaken for blackhead or tuberculosis) tuberculous fowls pass manure of a greenish cast, turning to a yellowish diarrhoea as the case becomes chronic. Such manure is very contagious, and should be cleaned up and burnt.

Tuberculosis of birds is caused by a bacillus, *Bacillus tuberculosis*. There is no known remedy for this disease. Eradication from infected premises can only be attempted with fair prospects of success when all the birds are affected. Any birds that are preserved are liable to have ulceration of the intestines from which the bacilli are constantly distributed. Consequently there should be no attempt to save any birds from an infected flock. The disease must be controlled by measures designed to keep the healthy fowls separated from the diseased ones. All affected birds should be removed immediately from the rest of the flock, and should be killed at once, the bodies being burnt.

The further spread of the disease through a flock where it already exists may be impeded by sanitary measures. The roosting-houses should be cleaned and disinfected as often as practicable. As with human beings, the disease can be avoided by means of good house ventilation, direct sunlight, and good wholesome food.

### ROUP.

Roup may be expected to cause trouble, especially where the houses are of the closed type. A few years ago it was the most prevalent disease in Southern Ontario, but by the introduction of the fresh-air house, it has been controlled to a very large extent. It causes very little, if any, trouble on Vancouver Island, but it must be carefully watched and guarded against in some sections of the Province where there is a considerable rainfall. It is particularly prevalent in the fall and spring of the year. Do not allow the fowls to catch cold under any consideration, but keep them dry and comfortable and they will give good returns.

In the majority of cases the first symptoms appear in the nostrils, while sometimes it may be localised in the eyes. It usually appears first in weak fowls as catarrh of the nose. A serous fluid may be observed at one or both nostrils. This secretion soon becomes streaked with grey; a silmy matter forms and dries quickly to dirty crusts, which often completely close the nostrils. The secretion decomposes and emits an offensive odour from the diseased membranes of the nostrils. The inflammation easily spreads to the mucous membrane of the mouth, pharynx, and larynx; yet those parts may be the first seat of the infection. When the disease affects the eyes there is often formed in the inner corner of the eye a vicious secretion. Soon the secretion is changed into a firm, smooth, yellowish-white, cheese-like body, which fills up the whole eye. This cheesy mass may be easily removed, but the eye is soon completely filled again. Swellings occur in the membrane of the eye, mouth, and larynx; oftentimes the cheesy mass develops in the mouth, forcing the mandibles apart so that the bird is unable to eat or drink. The fowls thus infected become weak, separate themselves from the other birds, and sit around with the head drawn close to the body. The eyes are kept shut; in fact, in many cases the cheesy matter fills the eye, destroying the sight. The head is sometimes drawn under the wings so that the feathers are smeared with secretions and some fall out. Usually the birds die, though they may live in this apathetic condition for a long time.

Healthy fowls may become affected if the diseased ones are allowed to run with them. It is usually brought into the flock by a diseased fowl from another place or the bacteria transmitted from one flock to another by means of person's clothing, coops, etc.

Poultry-keepers know how predisposed chickens and all other fowls are to colds caused by sudden changes in the weather. They must be kept in as natural a condition as possible; hence the wisdom of the adoption of the fresh-air house.

Experienced practical poultrymen do not doctor fowls which have roup; they kill and burn them.

A simple and fairly efficient remedy is swabbing the nostrils and throat out twice a day with a feather which has been dipped in coal-oil. The same feather should not be used more than once. Epsom salts should be given  $\frac{1}{2}$  teaspoonful to each affected bird and the bird fed on soft food entirely.

It will be seen, then, that the majority of poultry diseases arise through negligence on the part of the poultrymen. Health is the normal condition for a fowl, and it is of more importance to know how to keep fowls in good health than it is to know how to cure them after they are sick. However, under the best conditions sickness sometimes makes its appearance in the flocks. It is then important to know what is the cause of the sickness, and how to remove the source of trouble.

**BULLETINS AT PRESENT AVAILABLE FOR DISTRIBUTION BY THE  
DEPARTMENT OF AGRICULTURE.**

- 
- No. 7.—Flax.  
 .. 8.—Feeding Farm Animals.  
 .. 20.—Varieties of Fruit recommended. (Revised.)  
 .. 25.—Orchard Cleansing.  
 .. 26.—Practical Poultry-raising.  
 .. 28.—Production of Eggs.  
 .. 29.—Poultry Industry on the Pacific Coast.  
 .. 30.—Guide to Bee-keeping in British Columbia.  
 .. 32.—Control of Bovine Tuberculosis in British Columbia.  
 .. 33.—Fruit-growing Possibilities, Skeena River and Porcher Island Districts.  
 .. 34.—Fruit-trees and Black-spot Canker.  
 .. 35.—The Place and Purpose of Family Life.  
 .. 36.—The Preparation of Food.  
 .. 37.—The Preservation of Food.  
 .. 38.—The Construction of Silos.  
 .. 39.—Natural and Artificial Incubation and Brooding.  
 .. 40.—Alfalfa.  
 .. 41.—Labour-saving Devices in Household.

Applications for bulletins should be addressed to the Secretary, Department of Agriculture, Victoria, B.C.

---

VICTORIA, B.C.:

Printed by WILLIAM H. CULLEN, Printer to the King's Most Excellent Majesty.  
1912.

