
The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below. L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

Coloured covers/	Coloured pages/
Couverture de couleur	Pages de couleur
Covers damaged/	Pages damaged/
Couverture endommagée	Pages endommagées
Covers restored and/or laminated/	Pages restored and/or laminated/
Couverture restaurée et/ou pelliculée	Pages restaurées et/ou pelliculées
Cover title missing/	Pages discoloured, stained or foxed/
Le titre de couverture manque	Pages décolorées, tachetées ou piquées
Coloured maps/	Pages detached/
Cartes géographiques en couleur	Pages détachées
Coloured ink (i.e. other than blue or black)/	Showthrough/
Encre de couleur (i.e. autre que bleue ou noire)	Transparence
Coloured plates and/or illustrations/	Quality of print varies/
Planches et/ou illustrations en couleur	Qualité inégale de l'impression
Bound with other material/	Continuous pagination/
Relié avec d'autres documents	Pagination continue
Tight binding may cause shadows or distortion	Includes index(es)/
along interior margin/	Comprend un (des) index
La reliure serrée peut causer de l'ombre ou de la	Title on header taken from:/
distorsion le long de la marge intérieure	Le titre de l'en-tête provient:
Blank leaves added during restoration may appear within the text. Whenever possible, these have been omitted from filming/ Il se peut que certaines pages blanches ajoutées	Title page of issue/ Page de titre de la livraison
lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.	Caption of issue/ Titre de départ de la livraison
	Masthead/ Générique (périodiques) de la livraison
Additional comments:/ Commentaires supplémentaires:	
This item is filmed at the reduction ratio checked below/ Ce document est filmé au taux de réduction indiqué ci-dessous.	



August, 1887. PPE. VOLUME I. NO. V. The .TRADSACTIODS. Öf ilje \* Oftavla · Rield=Paturalists' · Club \* (Organized March, 1879. Incorporated March, 1884.) COPTEPTS. The Great Ice Age and Subsequent Formations at Ottawa, Ontario. H. M. Ami, M.A., F.G.S. 65 Report of the Botanical Branch..... .... 75 Additions to the "Flora Ottawaensis" ...... 77 Excursion to Buckingham..... Sub-Excursions. . . . OFFAWA, CADADA: Eilizen Printing and Publishing Company, Metcalfe St. 1887. 6 serv Published Monthly at \$1.00 per annum.

Patron :

### HIS EXCELLENCY THE MARQUIS OF LANSDOWNE, GOVERNOR GENERAL OF CANADA. President: R. B. WHYTE.

Vice-Presidents : 1st, PROF. J. MACOUN | 2ND, PROF. S. WOODS.

Secretary : W. H. HARBINGTON. (Post Office Dept.)

Treasurer : JAMES FLETCHEB. (Library of Parliament.)

Librarian: T. J. MACLAUGHLIN. (Dept. Public Works.)

Committee: Dr. H. B. SMALL, | H. M. AMI, | Dr. GEO. BAPTIE.

#### Standing Committees of Conncil:

Publishing-Prof. S. Woods, W. H. HARRINGTON, JAMES FLETCHER. Excursions-T. J. MACLAUGHLIN, H. M. AMI, DR. BAPTIE. Sources-Prof. J. MACOUN, PROF. S. WOODS, DR. SMALL.

#### Leaders :

Geology--H. M. AMI, REV. PROF. MARSAN, JOHN STEWART. Botany-J. FLETCHER, DR. SMALL, DR. BAPTIE. Conchology-F. R. LATCHFORD, H. B. SMALL. Entomology-W. H. HARRINGTON, J. FLETCHER, T. J. MACLAUGHLIN. Ornithology-G. R. WHITE, PROF. MACOUN.

Editor: W. H. HARBINGTON.

The Librarian will furnish the Publications of the Club at the following rates :--

Transactions, Part 1, Not sold singly )

66	2,	25 (	Cents.	PLOD C. TALT		
"	3,		<b>6</b>	\$1.00 for Vol. I.		
**	4,	25	<b>66</b>	) - Let status		
66	5,	30	"	A second seco		
"	9,	40	66	\$1.00 for Vol. II.		
H H	7.	30	#	No. State State State		

# (Less 10% to members.)

The Ottawa Naturalist, \$1.00 per annum. Extra copy to members, .75 " Monthly parts, 10 cents each, \$1.00 per doz. To members, 8 " .75 "

# THE GREAT ICE AGE AND SUBSEQUENT FORMATIONS AT OTTAWA, ONTARIO.

#### By H. M. Ami, M.A., F.G.S.

Among the most interesting and captivating subjects which attract the attention of even a casual observer in the realms of geological science, few are so full of interest and afford so much information as researches amongst the most recently deposited strata. Besides this interest, there is carried with it the fact of its practical importance, so that the economic aspects of the question have likewise to be taken into consideration.

At what period in the Earth's History did the Glacial Epoch or the Great Ice Age make its appearance?

What were the causes which led to it, what phenomena characterized it, what was its duration and what traces did it leave behind? And again, specially, to what extent was this continent submerged, for how long, and what traces of animal life has that period left behind it; and, further, are there unquestionable proofs of a period of subsidence followed by another of elevation carrying us on to the present day, during which time numerous and varied lacustrine or alluvial deposits were laid down, and in the lapse of which man made his appearance.

Then, in which of the newer deposits are traces of the existence of certain tribes of the American Indians to be found? What are these traces? To what extent do they assist in forming an estimate of the degree of civilization to which these aborigines attained? What customs and modes of life are exemplified by the implements of various kinds found in what has been very appropriately termed the Human . Period? At what time and for how long did these inhabitants occupy the land before the intrusion of the whites, and what was their history?

Е

It may not be thought amiss to note first what has already been done in the field of research with which we have to deal. In the "Geology of Canada," 1863, a report by Sir Wm. Logan and staffthere is a chapter on "superficial geology" in which a number of interesting notes are recorded from Ottawa and its environs, an examination of which had been entrusted to Dr. R. Bell. Then comes the work done in the Post-Pliocene geology of Ottawa by Sir J. A. Grant who produced a number of valuable papers, some of which were published in the United States and others here in Canada. At the mouth of and along Green's Creek, six miles distant from the city, and a favourite resort for students of Post-Tertiary geology, Sir J. A. Grant and Sir W. Dawson made important discoveries. The collections of the late Dr E. Van Cortland show that he also devoted considerable attention 'to these interesting deposits, whilst the I te Mr. E. Billings in his Canadian "Naturalist and Geologist" published notes on the same subject. The above mentioned work was prior to the organization of the Field-Naturalists' Club which has since vigorously pushed investigation in this Nearly a score of members, have taken a more or less direction. active part in these researches, whilst the abundance of work and material make it probable that greater attention will continue to be paid to the deposits in question. The work done already is considerable; but there remains a hundred-fold more to do. Mr. Surtees, the City Engineer, has been carrying on an extensive series of excavations in all parts of the city. These excavations or trenches are dug or blasted out to a depth ranging from eleven feet to eighteen feet six inches; so that deep and interesting sections have been exposed.

For the description of the Post-Tertiary or Post-Pliocene (Pleistocene) deposits it is first necessary to ascertain whence the material came which composes them, and in order to do this it is obviously necessary to examine the older rocks of the district, and to see of what their measures consist, and know the stratigraphical relations existing. between the various members of these older underlying series.

Just as we have a great diversity of formations about Ottawa, so also have we a great diversity of substances in the materials which make up the rock of the Post-Tertiary deposits, (and let it be borne in mind, that in using the word "rock" it is used in its true geological sense, so that a handful of sand, a lump of clay, a mass of publies, cemented or not by finer detritus, are all as much "rock" as a pillar of freestone, a block of limestone or a column of granite). Most of the materials which are found in these newer deposits were derived from the older formations of the district, whilst erratic blocks, and the like, may have come from great distances. To go into details regarding all the kinds of rocks met with, would necessitate a protracted study of a great quantity of material which years of labour could not exhaust, and which would form the constituent elements of all the formations from the Laurentian to the Hudson River as they are developed in the "Ottawa Basin;" from the gneissoid, granitic and hornblendic rocks of the former, to the shaly magnesian and arenaceous measures of the latter. There would be the gueisses, pegmeties, crystalline limestones, serpentines, dolomites and diorites of the Lower Laurentian, occurring at Chelsea in the Laurentide Hills, there would also be included portions of the conglomerates and quartzites and calc-bearing sandrock materials peculiar to the unconformably overlying Potsdam and Calciferous formations, whilst the sandstones, shales and limestones of the Chazy, followed upwards without a break by the impure calcareous strata of the Black River and Trenton formations would all be mixed together with the likewise conformably overlying bituminous schists of the Utica.

The materials which compose the series of formations just mentioned and newer than the Laurentian were themselves derived from the Laurentian System, for this latter contains all the elements necessary for the formation of the sandstones, shales and limestones of the newer overlying Cambro-Silurian or Ordoviciau strata.

Having ascertained the series of strata whence the material was obtained which constitutes the various beds in the Post Tertiary deposits, let us consider the condition of affairs previous to and at the coming in of the Glacial Epoch.

67

This portion of the American Continent, which, during the earlier palæozoic period had alternately been submerged and elevated, remained in this latter state a long period of time, during which denuding agencies, such as atmospheric erosion, rain and other solvents carried away a great deal of material. This is a lapse of time, which, in other parts of Canada and elsewhere, is marked by a regular ascending series of newer formations deposited, for the most part, beneath the level of the then existing oceans, a period embracing within itself the whole of the Silurian and Devonian systems, together with the Carboniferous age or the coal measures. The Palæozoic Era thus ended Mesozoic times came in and the Triassic, Jurassic, and Cretaceous systems followed, overlying which all the Laramie and Tertiaries were laid, a'l of which are entirely absent in our district marking a great unconformity between the Glacial deposits and the Hudson River rocks about Ottawa.

The Glacial Epoch or the Great Ice Age, then, is the first of the series of Post Tertiary times, with which we have to deal, as it rests immediately upon, though with discordance of stratification (if that term may be employed here), and overlies the Cambro-Silurian and older formations in this district. Just previous to this period of glaciation, and. whilst it lasted, there must have taken place a great elevation in this part of the North American continent, so that an extreme Alpine or Arctic climate was the natural result. Nor was this part of America the only one which enjoyed this particular state of affairs, but throughout the greater portion of North America as far west as the Great Missouri Coteau, in Europe, and in other continents, evidence of extreme cold, the result of great elevation, has been ascertained beyond Prof. Favre, of Geneva, whose admirable researches in Alpine doubt. , geology have made him so famous the world over, in the "Résumé" of his "Geological Researches in Savoie and the neighbourhood of Mt. Blanc," points out clearly what was the origin of the glacial epoch in that part of Europe. "The amount of moisture or humidity," he says, "with which the atmosphere of Europe was filled on account of , the elevation of land subsequent to the deposition of the tertiary deposits-the cooling effect of the neighbouring mountains, then more elevated than now-a-days-together with many other causes, led to a reduction in the temperature of the atmosphere resulting in an abundant

precipitation of snow on the leading peaks of the vegion." So in Canada, and in the Ottawa district, a great reduction in the temperature followed the great elevation, and immense quantities of snow, ice and water followed and glaciers were formed all over the district-a vastmer-de-glace covered this portion of Canada, whose height above the ocean level of that period was considerable. These glaciers, like modern ones, were characterized by many interesting particulars which a study of the latter can afford. The number, direction, movements, thickness, erosive or denuding power and the constituent parts of a glacier, or a system of glaciers, are questions full of interest. It has been ascertained: that over four hundred glaciers can be seen in the central portion of the Alps, from Mont Blanc to the Tyrol, some of which are only three miles in length, whilst others exceed twenty miles from head to foot or from the point of origin to the snout. There is abundant evidence to show that the number of glaciers which must have existed here about Ottawa is very considerable. Perhaps the greater number, were subordinate or small ones and may, at length, have been absorbed in and formed part of "a great glacier." The direction in which they moved depended of course on the nature of the district, its physical or orographical character. The general trend of the great mer-de-glace in Canada during this epoch has been ascertained to be approximately N. E. and S. W. With regard to the direction of some of the glaciers, the striations or grooves on the rocks about Ottawa show that in some cases they travelled almost due east and west, as may be seen along Park Avenue, on Nicholas street and in other parts of the city, at other times they appear at a considerable angle to this direction, bearing almost due north and south, as at Buckingham on the Lièvre River. Regarding their movements and the speed with which glaciers travel, we consult Agassiz and find that he obtained the following results in 1841 and 1842 on some of the Aar glaciers :---

I. FINSTER AAR-	ANNUAL NOTION. 		
Stake near centre of glacier			
{ " side "	160 "		
II. LAUTER AAR-			
(Stake nearest centre of glacier	.245 "		
Stake nearest centre of glacier	.124 "		

۰.

:

whilst at *Chamonix* the ice near the shore of the *mer-de-glace* was found to move as follows, from June 29th to June 8th of the following year :

From	June	29	to	Sept.	28	132	feet.
"	Sept.	<b>28</b>	"	Dec.	28	70	"
	Dec.	12	"	Feb.	17	76	"
					4		
					8		
	•				•		

Total (in less than one year)..... 432 feet This would average over five hundred feet or about one-tenth of a mile in twelve months. The rapidity in the motion of a glacier, of course, depends upon the nature of the obstacles to be surmounted, as well as to a great extent upon the time or month of the year, different portions of the same glacier moving at different rates. A glacier which decends into a valley below, or discharges itself into a sea or arm of an ocean, does not necessarily lose any of its length, for whilst its snout is being melted and carried away to warmer portions, the head or initial point is ever receiving additional snow and ice to supply it constantly, and only a subsidence of the continent could produce a change in the climate of such an ice bound district.

We have no data existing here cr traces left by means of which we can calculate the rate of motion of the glaciers about Ottawa during the great ice age, suffice it to say that as in the case of modern glaciers their rate of travelling varied at different times. Then as to the thickness of the great ice-mass which then invaded this district, that is a problem which to a great extent, has yet to be solved with us, nevertheless, let us examine the data at our disposal in reference to this interesting phenomenon. Taking the Ottawa Valley, in and around the city, as a typical example of a valley of erosion with subordinate branches, we see that facing the river and the north, there occurs a series of high "buffs" or cliffs where the strata are clearly seen along their sides to be throughout nearly horizontal.\*

That these beds could not have been deposited in such a position is beyond question, so that the prolongation of them northward must at one time have existed. What was it then, which removed all these and

70

Ŧ

<sup>\*</sup>There are but few exceptions to this, due to dislocations, faults and folds in the strata, of purely local origin, but not of general significance in this problem.

to such a depth ? This very question gives us a clue to one factor at least in the estimate of the powerful ice-mass which, coming from the West or W. N. W. struck down upon the shales and limestones of the formations here to be found. It also gives us data to estimate its thickness. The occurrence of striated rocks at the top of old Barrack Hill, where the Parliament Buildings now stand, shows that as that cliff is one hundred and eighty-seven feet above the level of the river, and over two hundred feet above the level of the bed of the river, the mass must have been much over two hundred feet. Further, in order that a mass of ice or a glacier carrying boulders and detritus-moraineprofonde-can groove and polish the rocks of a district to such an extent as was the case here, the superincumbent weight and attending pressures must have been enormous, and from what is known of present glaciers, whether in alpine or arctic regions, we know that its thickness must have been very great. A fair estimate, we believe, of the thickness of the glacier or mer-de-glace extending over our city and its environs during the glacial epoch must have been very little short of one thousand feet, if indeed that number is not too small. The erosive or denuding force of glaciers has as yet only casually been touched upon, for when we take into consideration the millions of tons of material which have been removed from even the small area about our city, it is marvellous to know where it all went. You can hardly find a loose rock or boulder in the fields without seeing written upon it indubitable marks of scratching and grooving, v nich, along with • millions of others were held firm in a mixture of cementing clay and sand (to a small extent) carried forward upon the floor of the glacier and ground one against the other, at times, to such an extent that all angularities and rough points were removed and the boulders left smooth and polished. The striations, grooves and polished surfaces of rocks which up to this date attest clearly to the fact of the existence of those glaciers, besides the boulders themselves, may be seen not only in the places already mentioned, but at the corner of Sussex and Rideau streets, where there is an interesting exposure.

The effect of these glaciers upon the softer shaly strata of our neighbourhood is clearly shown in such a deposit of the Utica shales as is met with at Cumming's Bridge, on the Rideau River, or at the corner of Maria and O'Connor streets, some 12 feet below the surface of the roadway. At these two places, whilst the shales of the Utica formation also occur *in situ* and undisturbed at a greater depth than is visible in either section, the uppermost measures of the section exposed and examined cannot certainly be said to be strictly *in situ*, as the beds are tilted at every conceivable angle, crushed and broken, and in the overlying glacial deposits are to be found some of the boulders themselves which assisted intilting and disturbing these once horizontal measures.

There occur a vast number of faults and dislocations in the measures. of the Trenton and other formations about Ottawa, great breaks, which at times, run more or less parallel to each other and were the result of great pressure brought to bear upon the beds in question. Whether these faults and breaks are due to disturbances which took place about the close of the Silurian Age, or at the introduction of the Devonian, when Rigaud and Montreal mountains, and other similar volcanic or intrusive masses, were ejected amidst great perturbation; or whether some of these faults were not in part due to the enormous pressure which the great ice-mass exerted upon the strata in later glacial times are questions which, though readily suggested to one's mind by the phenomena examined, do not find so ready a solution. Having now examined the number, direction, movements, thickness and the erosive power of the glaciers during this Great Ice Age, having very cursorily glanced at the results which were effected in giving to the country the general appearance which it possesses at the present day, there remains to find out what are the materials and under what conditions they were - deposited.

The masses of boulders, also termed "boulder clay," "moraineprofonde," &c., unlike both the underlying older and overlying newer deposits are not stratified, *i. e.* they have no divisional planes of stratification or true bedding. Pebbles of various sizes and of every kind of rock in the district, usually rounded and smooth, held together or cemented by an argillaceous paste or clay with a certain admixture of arenaceous material derived from the more finely crushed *detritus* and *debris* at the bottom of the glacier, form the lowest division of our Post-Tertiary deposits. These "boulder clays," as they are appropriately termed, have a very large percentage of boulders in them, the finer material being scanty and in a finely-divided or comminuted state. Such a deposit is one which "land ice" alone produces and one which resembles wonderfully the "moraine-profonde" of the ancient Rhône glacier as they may be seen near the Western extremity of Lake Geneva (a mile and a-half below) and in the adjoining districts. In further corroboration of these boulder clays being due to land ice is the fact that none of the organisms which would be expected to characterize marine clays are present therein. The total absence of organic remains (so far as ascertained) in these glacial clays, coupled with the fact of their occurrence in abundance in the Leda clays above, points clearly to a wide difference in the mode and condition of deposition of both, the one being laid at a great elevation above the sea level, the other below the level of an ocean or arm of a sea.

In examining the surface geology of Ottawa, one is struck withthe diversity in the distribution and extent of this "boulder clay formation:" In some places, the only indexes present, which point to its existence at one time, are the striæ and grooves over the bare rocks, such as are exposed principally about Hull and Ottawa in the vicinity of the Grand River, whilst there are also numerous fields and tracts. of country which exhibit that formation very clearly. In such post-glacial valleys and districts, from which the Leda clay, and Saxicava sand and overlying strata, have been removed by denudation, there occurs a large quantity of these boulders. Amongst these are no. doubt included, at the present day, the erratics which were dropped by ice-bergs at a period subsequent to the Great Ice Age. The Rideau River Valley, of Post-Tertiary Age, and very recent, geologically speaking, presents numerous points of interest from its mouth at the falls in New Edinburgh to the Hog's Back. Nearly the whole of the Post-Tertiary formations were carried away by the once wide stream which flowed there, and even the glacial clays suffered not a little, as the materials. cementing the pebbles are to a great extent entirely wanting. The Rideau Rifle Range extends, for the most part, over this formation, whilst the southern portion of the range, as well as its northern limit (at the 600 yard butte) are on the outskirts of the newer overlying marine clays. We have already spoken of moraines. These vary very much in extent and distribution just as the "boulder clay" or "till,"-

as it is also sometime called, varies from next to nothing to twenty feet or more in thickness in different places. They are extensively developed about Ashburnham Hill, Gilmour's Mills, near the Hog's Back, etc., occupying their lower and regular position at the bottom of the Post-Tertiary series in Upper Town, Centre Town and Stewarton, cropping out in the rear of the City Hall, on the east side of the canal, and ever keeping in a normal position. This "till" is thence very generally distributed in beds of varying thickness, in the area included in a curve drawn from the New Militia Stores on the canal, along Sussex street up to St. Patrick street, then produced on to the bridges over the Rideau river, pretty nearly in a line with the curve which Sussex street there describes, and across to New Edinburgh through the Rideau Hall grounds, to a small extent, where these deposits thin out markedly, and continuing the line through Beechwood, in Gloucester, on in a southeasterly direction, we have a horse-shoe curve, in which are included vast accumulations of material left us by this Ice Period. But to come back to the moraines :-- there are, besides these hard coherent masses of "boulder clay," large aggregations of more or less uniformly sized boulders held very loosely together, and form a prominent feature of the county. At Gilmour's Mills there is a good example of one of these whilst there has already been pointed out in the "Geology of Canada," 1863-already cited-that a number of zones or belts of boulders cross the Ottawa at different places near the city, one of which a few miles below Ottawa produces a shoal on account of which the navigability of the river at this point becomes dangerous and has obliged the Government to erect a lighthouse. These are what are termed "morainic belts," and are prominent features to consider.

There are a number of other particulars respecting which the detailed notes obtained from the excavations that have been carried on in our streets, though bearing immediately on this subject, cannot be included in the present exposition of the subject.

(To be Continued.)

### REPORT OF THE BOTANICAL BRANCH.

# To the Council of the Ottawa Field-Naturalists' Club:

The same plan of work, found so successful in former years, has been adhered to during the past season with quite satisfactory results. The attendance of botanists at the sub-excursions has been particularly gratifying to the leaders, and there is a decided increase in the amount of individual work being done by members. The Flora of the district is now pretty well worked up and the number of additions to the "Flora Ottawaensis" is smaller than usual. The leaders desire to call the attention of the Council to this list which was prepared by Mr. Fletcher for Part I. of our Transactions, and would suggest that, as most of the plants in the district have probably now been discovered, the time has arrived for a more complete list to be issued, in which all the additions of subsequent years should be included as well as the special localities and the period of inflorescence of each species. We hope that the Council will be able to order such a republication as a supplement to our present Transactions.

We think it well in the interests of the Club to mention the important work done by some of our members in connection with the Colonial and Indian Exhibition, held in London, England, during the past summer. Our worthy President, Prof. Macoun, the Dominion Botanist, spent the whole summer in England, where he was able, by lecturing and in conversation at the Exhibition, to disseminate very valuable information concerning Canada and its natural products. То Mr. James Fletcher, one of the active members of this branch, was entrusted the laying out and arrangement of a Botanical Garden in connection with the Exhibition, where were shown growing specimens of a large number of our Canadian plants. A good representative collection was made of our forest trees, and plants of special interest, whether economic, medicinal or scientific, were shown. In making the collection of roots Mr. Fletcher was materially assisted in his work by several members of the Club, particularly by Mr. H. M. Ami, of this city, and Rev. G. W. Taylor, of Victoria, B.C. It is gratifying to see from the report of the High Commissioner of Canada that this garden

was a decided success, and those who visited the exhibition pronounced it a very interesting feature of the Canadian Exhibit.

We consider it our duty to refer to the valuable work which has been done for the Club by Mr. R. B. Whyte, our senior leader, who, from April to June, gave a systematic course of instruction in Botany to a large class in the Central School West. We believe that in no way will the Club be made more useful and popular than in action similar to this.

Of the plants added to the list during the past season none are of very special interest. New localities for already recorded species have been noted, and add much to our knowledge of the district. *Bidens Beckii*, a plant which seldom flowers freely, this season perfected its flowers in profusion in many of our waters, as Brigham's Creek, Hull, and all along the Rideau River. *Cellis occidentalis*, a tree of rare occurrence here, was observed at Britannia and the Little Chaudiere.

Dipsacus sylvestris, Mill. "The Wild Teazle," although possibly indigenous in Western Canada is not so here, and the specimen collected by Mr. Ami in one of our streets was undoubtedly from seed, accidentally introduced.

Caucalis daucoides, L. "Beaked Parsley." One specimen of this plant was found in a wheat field where it had probably been introduced with the seed.

Symplocarpus foetidus, "The Skunk Cabbage," and Podophylliumpellatum, L., "The May Apple," were both found by Mr. R. B. Whyte at Perth, Ont., but this is at too great a distance to be included in our ' local list.

The leaders again express a hope that the study of the Mosses may be taken up with greater enthuciasm by our members. There is no branch of Botany which presents a more fruitful field for research, nor in which there is so little trouble in preparing the specimens.

During the past season Mr. Fletcher has continued his studies in growing from the seed, and cultivating here, plants indigenous to other parts of the Dominion, and has succeeded in flowering the following species in addition to those already mentioned: Silene Drummondii (Nor.-West Ter.), Claytonia perfoliata, Donn. (Brit. Col.), Geranium incisum, Nutt. (B.C.), Lupinus polyphyllus, Lindl. (B.C.), Petalostemon

candidus, Mx. (N.-W.T.), Rubus Nutkanus, Moçino (B.C.), Liatris cylindracea, Michx. (W. Ont.), L. scariosa (Manitoba), L. punctata, Hook. (N.-W.T.), Soldago rigida, L. (N.-W.T.), Gaillardia aristata, Pursh. (B.C.), Nemophila parviflora, Doug. (B.C.), Phacelia Menziesii, Torr. (B.C.), Convolvulus spithamœus, L. (L. Superior), Solanum triflorum, Nutt. (N.-W.T.), Allium siellatum (B.C.) The following grow from the bulb and stand our winters well: Lilium parviflorum, Camassia esculenta and Fritillaria lanceolata. All of the above with the exception of the first two and Solanum triflorum are worthy of a place in our gardens as ornamental plants.

(Signed)

R. B. WHYTE. SAML. WOODS. H. B. SMALL. (M.D.)

#### ADDITIONS TO THE "FLORA OTTAWAENSIS."

#### EXCURSION:

The third regular monthly Excursion was held on Thursday, 21st July, to Buckingham, P.Q. The morning opened dull and cloudy with an occasional drop or two of rain. This dullness, undoubtedly, affected very seriously the attendance, for when the whistle of the "Maude" sounded at 7.20, only 24 members were present. The day turned out a perfect success, the overcast sky during part of the day, accompanied by a light breeze rendered the labours of the Collectors a much pleasanter occupation than has been the case during the greater part of this hot dry month. Rich collections were made in all branches. At noon the party reassen bled for luncheon at the grist

1

mill over the rapids, after which the Phosphate Crushing Works were visited. The Excursionists received every attention from Mr. Taylor, who, in the absence of the Manager, showed them over the mill and explained the machinery by which the raw material is crushed and prepared for the manufacture of super-phosphate. There were in the mill over 150 tons of ground phosphate (finer than wheat flour) ready for shipment. Having returned to the wharf the usual elementary lectures were delivered by the leaders Mr. Fletcher was first called upon by the President to speak on Entomology. He stated that rare insects had been collected, and announced the capture of a female of Colias Interior, a very rare butterfly, and one he was particularly anxious to breed from the egg. His remarks were directed towards showing the advantages which are to be derived from students working out the lifehistories of different species of insects, and showed that it was one of the most important factors in the success of Economic Entomologists.

Mr. Henry M. Ami then explained the different rock-formations which were to be seen in the locality. The Laurentian and Potsdam formations were the two which occurred of the older Geologic Epochs at Buckingham. In the former were found inexhaustible supplies of iron, apatite or phosphate of lime, plumbago, mica, baryta and other mineral products, and the Potsdam formation afforded building materials of first quality. Mr. Ami then drew attention to the traces of glaciation which were well exposed in several places and gave a succinct and lucid account of the Great Ice Age. The President, Mr. R. B. Whyte, gave the address upon the Botanical specimens collected, he said the locality was an exceptionally rich one as was evidenced by the many rarities which had rewarded the collectors. Mr. Whyte took advantage of some of the familiar plants to explain in a clear and attractive manner a few of the rudiments of structural botany. He used the Nymphaa tuberosa to show the gradation of all the floral organs from the leaf-form. Impatiens fulve, the common balsam or "Jewel Weed" of our swamps, was exhibited and everyone was advised to take special notice of it, as it was reputed to be the antidote for the Poison Ivy (Rhus toxicodendron). Many other rare and beautiful · plants were spoken of and made interesting by the remarks of the speaker. The party reached the city again about 9 o'clock, when the

much needed rain, which had kept off during the day, began to fall and continued to do so all night.

:

# SUB-EXCURSIONS.

The Sub-Excursion arranged by the Botanical and Entomological Leaders for the 2nd July had to be postponed on account of wet weather.

SEVENTH .-- On the 9th of July, notwithstanding the excessive heat of the day, about lifteen members of the Club, including several ladies, turned up at the usual rendezvous-the General Post Office. At 2 o'clock the party started in the Billings Bridge stage for Dow's swamp. This locality, which lies about two miles south of the city, is one possessing special attractions to the Entomologist and Botanist. The ·Botanists were under the leadership of the President, Mr. R. B. Whyte, who also delivered the address upon Botany, none of the regular leaders being able to attend upon this occasion. Many plants of interest were collected and their important features were drawn attention to in a pleasing and simple manner, by the President. The magnificent white panicles of the Meadow Rue, Thalictrum Cornuti, were conspicuous objects in the cleared spaces of the bog, and beneath the shade of the cedars the rare little orchids, Microstylis monophyllos and M. ophioglossoides, were found. These treasures only reward the patient searcher or the Botanist with a well-frained eve. Another orchid of interest, but like the last named, possibly more inconspicuous than scarce, was Liparis Laselii, of which a few specimens were secured. In all about forty species of plants were collected in flower. Mr. Fred. Magee represented the Entomologists and exhibited his collections of the afternoon. Dow's Swamp is one of our few localities for the beautiful butterfly Melitæa Phaeton, which may still be found in an open glade to the south east of the lake, where it was noted by the late Mr. B. Billings many years ago. Its bright colours, rich velvety black spotted with red and white, make it a conspicuous object as it flits slowly over the low bushes. Its food plant, Chelone glabra, grows here in abundance.

EIGHTH.-Rockcliffe was the point decided upon for investigation for the Sub-Excursion of 16th July. The heat was excessive, but over a dozen enthusiastic members turned up. As usual this locality produced much of interest. Here, curiously enough, is the only locality, yet discovered in this neighbourhood for the New Jersey Tea (Ceanothus Americanus). Here too occur Pinus resinosa and Woodsia Ilvensis, both uncommon in the neighbourhood, and on the occasion of the Excursion it was discovered by Mr. A. J. Forward that the rare Ericaceous plant, Pterospora andromedea, also grew here. An address was delivered by Mr. Fletcher, leader in Botany and Entomology, upon those subjects. He spoke of the peculiar features of the Flora of the rocky point upon which they stood, and mentioned the plants which had been found. Pterospora was probably a parasite upon the roots of pines, the only other place that he knew of, where it could be found about Ottawa, was upon the Chelsea Road. Comandra, Monotropa, and other parasites were referred to. Astragalus Canadensis and A. Cooperi were exhibited, and it was shown how they might be known The latter is a much earlier plant, the seed being ripe by the apart. time the former is in flower. The fruit of Prunus pumila was exhibited. This shrub is quite rare at Ottawa, one bush only having been previously observed, which grows upon the Chaudiere Islands. In speaking of the insects collected, the different kinds of Cicadæ were described, their "song" was treated of and their life-history was sketched. The different kinds of mosquitoes also formed an interesting topic which was simply explained. Notwithstanding the heat the Sub-Excursion was a great success and the members were well pleased.

#### ANNOUNCEMENTS.

EXCURSION.—The fourth Excursion will be held on Saturday, 13th August, to Britannia on the Canadian Pacific Railway. The trains leave at 10 a.m. and 12.30 p.m. and the party will return by the 6.45 train in the evening. Return tickets : Members, 25 cents; non-members, 30 cents children, 15 cents.

SUB-EXCURSIONS.—August 6, Gatineau Point; August 20, Beechwood; August 27, Skead's Gravel Road.

# International Tent and Awning Company, 184 Sparks Street, Ottawa.

Manufactuters of Tents, Camp Furniture, Awnings, Window Shades, Plain and Decotated Curtain Poles, Curtain Chains, Flags of all Nations. Waterproof Horse and Wag gon Covers, &c., on hand and made to order in any size or style on short notice.

NOTICE—Special Discount to Lumbermen, Contractors, and other large Buyers.

Send for Prices. Catalogues Free on Application

# Address, A. G. FORGIE, Manager International Tent & Awning Co. Telephone Connection.

e de la competitione

