




**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**Photographic  
Sciences  
Corporation**

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503

**CIHM  
Microfiche  
Series  
(Monographs)**

**ICMH  
Collection de  
microfiches  
(monographies)**



**Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques**

**© 1992**

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured covers/  
Couverture de couleur
- Covers damaged/  
Couverture endommagée
- Covers restored and/or laminated/  
Couverture restaurée et/ou pelliculée
- Cover title missing/  
Le titre de couverture manque
- Coloured maps/  
Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black)/  
Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations/  
Planches et/ou illustrations en couleur
- Bound with other material/  
Relié avec d'autres documents
- Tight binding may cause shadows or distortion  
along interior margin/  
La reliure serrée peut causer de l'ombre ou de la  
distorsion le long de la marge intérieure
- Blank leaves added during restoration may appear  
within the text. Whenever possible, these have  
been omitted from filming/  
Il se peut que certaines pages blanches ajoutées  
lors d'une restauration apparaissent dans le texte,  
mais, lorsque cela était possible, ces pages n'ont  
pas été filmées.
- Additional comments: /  
Commentaires supplémentaires:

- Coloured pages/  
Pages de couleur
- Pages damaged/  
Pages endommagées
- Pages restored and/or laminated/  
Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed/  
Pages décolorées, tachetées ou piquées
- Pages detached/  
Pages détachées
- Showthrough/  
Transparence
- Quality of print varies/  
Qualité inégale de l'impression
- Continuous pagination/  
Pagination continue
- Includes index(es)/  
Comprend un (des) index
- Title on header taken from: /  
Le titre de l'en-tête provient:
- Title page of issue/  
Page de titre de la livraison
- Caption of issue/  
Titre de départ de la livraison
- Masthead/  
Générique (périodiques) de la livraison

This item is filmed at the reduction ratio checked below/  
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
12X	16X	20X	24X	28X	32X

The copy filmed here has been reproduced thanks to the generosity of:



National Library of Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol  $\rightarrow$  (meaning "CONTINUED"), or the symbol  $\nabla$  (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

Bibliothèque nationale du Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole  $\rightarrow$  signifie "A SUIVRE", le symbole  $\nabla$  signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

qu'il  
cet  
de vue  
ge  
ation  
ués



32x

IN

Optic

SO

[Re-print

PRI

AN  
INTRODUCTION TO THE STUDY  
OF THE  
**Optical Defects of the Eye,**  
AND  
THEIR TREATMENT  
BY THE  
SCIENTIFIC USE OF SPECTACLES.

---

BY  
A. M. ROSEBURGH, M.D.,  
TORONTO.

---

*[Re-printed from the Canadian Journal for January, 1866.]*

TORONTO :  
PRINTED BY LOVELL & GIBSON, 67 YONGE STREET.  
1866.

OPT

S

(R

THE f  
of lectur  
have not  
lishing th  
only to  
medical

In thei  
to the el  
berg Wel  
treatise of  
New Syde

*From the Canadian Journal for January, 1866.*

---

THE  
OPTICAL DEFECTS OF THE EYE,  
AND  
THEIR TREATMENT  
BY THE  
SCIENTIFIC USE OF SPECTACLES.

BY A. M. ROSEBRUGH, M.D.

*(Read before the Canadian Institute, February 3rd, 1866.)*

THE following pages were written as an introduction to a course of lectures recently delivered by me on the diseases of the eye. I have not thought it necessary to alter the form, as I propose publishing them as a pamphlet, hoping that they may be useful, not only to the members of my ophthalmic class, but to Canadian medical students generally.

In their preparation, I must here acknowledge my indebtedness to the elaborate works of Mr. J. Z. Laurence and Mr. J. Soelberg Wells, of London, and especially to the very comprehensive treatise of Professor Donders, of Utrecht, published in 1864 by the New Sydenham Society.



## CHAPTER I.—OPTICAL CONSIDERATIONS.

The eye is pre-eminently an optical instrument, and the phenomena of vision all depend upon the laws of optics. Hence, a knowledge of some, at least, of the elementary principles of light is essential to a correct appreciation of the physiology of the eye. The diagnosing of optical defects of the eye,—long and short sight, &c. &c., and their treatment with the scientific use of spectacles, require some knowledge of the laws of refraction, and the properties of convex and concave lenses.

The philosophy of the ophthalmoscope can hardly be understood unless the principles of both refraction and reflection are thoroughly mastered.

You will therefore, I hope, not consider the time ill spent if, before proceeding with the investigation of diseases of the eye—you review with me some of the elementary principles of optics which lie at the foundation of all ophthalmic science.

The nature of light is not known. I can no more tell you what light is, than your professor of physiology can tell you what life is. We know that the sun shines, but how it shines we cannot tell.

“Two different theories have been advanced of the more intimate nature of light.” “One, the *Newtonian (corpuscular)* conceives that each luminous point is constantly giving off a succession of luminous corpuscles which follow each other in uninterrupted succession on an imaginary line or axis like a string of beads on a rigid thread.”

The *undulatory* theory (Christian Huychens') on the other hand considers space as pervaded by a subtle gaseous fluid or ether; that luminous bodies have the power of communicating to this ether a wave motion which affects the retina the same as vibrations of the air affect the auditory nerve.

Sir John Herschel, speaking of the great ingenuity of the undulatory theory says, “if it is not true it deserves to be.”

The sun is the great natural source of light; as it shines by its own light it is called *self-luminous*. The fixed stars are also self-luminous; so is a lighted lamp and bodies in a state of ignition. But most bodies by which we are surrounded, are seen only by reflected

light  
befor  
and  
E  
infin  
alone  
light  
diver  
are c

-Fig  
passin  
rays p  
the po  
The  
rays a  
The po  
a focus

Stric  
the ne  
from th  
ever, w  
pupil o  
tance g  
twenty

A g  
lighted  
a small  
upon th  
the oper

light The light from an object seen by moonlight is reflected twice before it reaches the eye. The moon reflects the light from the sun, and the object, the light which it receives from the moon.

Every luminous object gives off, or radiates, in every direction, an infinite number of straight lines of light. Each of these lines taken alone is called a *ray* of light. A bundle of rays is called a *beam* of light when the rays run *parallel* to each other. When the rays *diverge* from a luminous point or are made to *converge* to a focus they are called a *pencil* of rays, thus :



Fig. 1 represents a pencil of rays diverging from a flame F, after passing a convex lens they are rendered parallel and these parallel rays passing the second convex lens B, the rays are converged to the point (focus) P.

The parallel rays may be called a *parallel pencil*; the diverging rays a *divergent pencil*, and the convergent rays a *convergent pencil*. The point where rays of light meet is called the *focal point* or simply a *focus*.

Strictly speaking, there is no such thing in nature as parallel rays; the nearest approach we have to it are the rays of light we receive from the sun and the fixed stars. Practically, for our purpose however, we may consider rays of light parallel that are received by the pupil of the eye from objects that are twenty feet distant or any distance greater than that. Pencils of light from objects less than twenty feet distant are more decidedly divergent.

A good illustration of a divergent pencil can be obtained from a lighted lamp or candle in a dark room. If a piece of card board, with a small circular opening in it, be held near the lamp, you will have, upon the opposite wall, an illuminated spot of the same shape as the opening in the card, but very much larger.

This will prove not only that the rays *diverge*, but also that the rays proceed in straight lines.\*

*Convex lenses* :—We shall now proceed to the consideration of convex lenses, which, for our purpose, is the most important part of the subject. Lenses are made of various transparent substances as amber, alum, quartz, glass, diamond, and even of ice. Those in ordinary use are made of glass. When the two surfaces of a convex lens have the same degree of curvature, the lens is said to be equi-convex. When one of the surfaces is flat or plane, the lens is called a plano-convex lens. Glass spectacles used by old persons for reading, &c., are commonly made double convex.

In order to simplify the subject as much as possible, let us confine our attention to lenses that are equi-convex.



In fig. 2 let A be the centre of the circle B, C, D, of which A, B, is the radius, and let E be the centre of the circle F, G, H, of which the radius E, F, is equal to the radius A, B. The circle F, G, H, will be equal to the circle B, C, D. The part D, H, common to both circles, represent a section of an equi-convex lens. The line A, E, is called the *axis* of the lens, and the line D, H is called the *diameter*. The centre of the diameter (where it is intersected by the axis) is the optical centre of the lens.

Reading glasses, and burning glasses, are examples of a double convex lens. Many of you have, doubtless, seen the experiment of

(\* Convergent pencils of light do not exist in nature. Parallel pencils or divergent pencils of rays can be rendered convergent by means of a convex lens. Thus in fig. 1, the rays diverging from F, are made to converge to P by the convex lenses, A. and B.)

sett  
The  
prop  
the

In  
focus

Th  
alway  
The  
equal

Th  
the le  
of one  
inch l  
centre  
receiv  
an inc  
strong

they  
remov  
The w  
inches  
the let

Let  
princip

Sup  
one of  
the gla  
words-  
would  
ascerta  
brough  
inches

, but also that the

consideration of con-  
sistent part of the  
ent substances as  
of ice. Those in  
surfaces of a convex  
is said to be equi-  
e, the lens is called  
l persons for read-

ible, let us confine

of which A, B, is  
G, H, of which  
circle F, G, H, will  
mon to both cir-  
The line A, E, is  
led the *diameter*.  
d by the axis) is

cles of a double  
ne experiment of

el pencils or diver-  
convex lens. Thus  
the convex lenses,

setting fire to wood, paper, &c., by means of a burning or sun glass. The explanation of this is simply that the convex lens possesses the property of converging a portion of the sun's rays to a point called the focus.



In Fig. 3, P, P, represent a pencil of parallel rays converged to a focus at F by means of the double convex lens, L.

The focus for parallel rays is called the *principal focus*. It is always the same distance from the optical centre in the same lens. The length of the focus for parallel rays is, in equi-convex lenses, equal to the length of the radius of curvature.

The shorter the focus, the greater is the "power" or "strength" of the lens. A lens that can bring parallel rays to a focus at a distance of one inch from the optical centre of the lens, would be called a *one inch* lens. Another lens whose focus is two inches from the optical centre, is called a *two inch* lens, and so on. Convex lenses therefore receive their names according to the number of inches or fraction of an inch, the principal focus is distant from the centre of the lens. The strongest lenses used for spectacles are what are called cataract glasses; they are worn by patients who have had their crystalline lenses removed. Their strength ranges from 2 to 4 inches focal length. The weakest spectacles that are ordinarily used have a focus of 36 inches. Convex lenses having a focus of 36 inches do not enlarge the letters of a book at the ordinary reading distance.

Let us now see what practical application we can make of this principle of convex lenses.

Supposing that a person accustomed to using convex spectacles, gets one of the glasses broken, and applies to you to learn the strength of the glass that would be necessary to replace the broken one, or in other words—to learn the strength of the glass that is still whole. How would you proceed? One method is to use the lens as a sun glass, and ascertain by measurement, how far from the glass, the sun's rays are brought to a focus. If you find, for instance, that the focus is 10 inches from the lens, you will have ascertained that the person has

been wearing glasses of 10 inch focus, or as they are sometimes called No. 10 convex, or simply + 10 (plus 10).

The method, however, that is usually adopted, depends upon a property of convex lenses that will be more fully explained further on.

If, for instance, you hold up a 10 inch convex lens at a distance of 10 inches from a white wall—the wall being about 20 feet from an open window, opposite—there will appear, behind the lens, upon the wall, an inverted, miniature picture of the window, and trees or buildings, &c., in front of the window. If the lens be held at a greater or less distance from the wall than the focal length of the lens, the inverted picture will be indistinct. Measuring the distance therefore that the lens must be held from the wall, to produce the sharpest picture, will give the focal length of the lens.

Suppose, now, that we bring the lens to within, say 5 feet of the window, and hold a sheet of white paper at the principal focal distance behind the lens, viz., at ten inches, we will find a change in the inverted picture, there will still appear distant buildings, trees, &c. but the sash of the window will be very indistinct. If, however, we move the sheet of paper 12 inches from the lens—that is, two inches farther from the lens, we will again see the image of the sash but scarcely any trace of the buildings, trees, &c. This experiment is an illustration of the fact that the nearer an object approaches the front of a convex lens, the farther will be its image behind the lens; thus, when an object is 5 feet or rather 60 inches from the front of a 10 inch convex lens the inverted image is found to be 12 inches behind the lens; when 30 inches, it will be 15 in.; when 20, that is, double the length of the focus, the image will be double the length of the focus behind the lens; viz., 20 inches; when 15 inches, the image behind the lens will be removed to 30 inches. As the object approaches the principal focal distance of the lens the image recedes much more rapidly; thus when at 12 inches, the image will be 60 inches; when at 11, the image will be 110 inches behind the lens. When however we bring the object to within 10 inches of the lens—that is, at its principal focus there will be no image formed behind the lens, as the rays after passing the lens will be parallel.

(I would strongly urge you, gentlemen, to perform all these experiments for yourselves, as in that way only can you become familiar with these important principles. These latter experiments can be performed best in a dark room—taking for an object the flame of a lamp or candle).

Fr  
the U  
proac  
will t  
lens.  
(the r  
distan  
Qu  
length  
tance  
will b  
nically  
and th  
the foc  
follow  
far bel  
This  
but th  
The  
the len  
object  
the di  
50=12  
The  
to men  
Wh  
the obj  
focus, t  
focus o  
inch len  
inch len  
The 1  
inch len  
inch, &  
The  
focal le  
the "p

From the above we can easily understand the principle, that the *less* divergent the rays of a pencil (that is, the nearer they approach parallel rays,) incident or falling upon a convex lens, the nearer will the focus of the convergent pencil be to the principal focus of the lens. 2nd. The *more* divergent the incident pencil, the less convergent (the more nearly parallel) will be the refracted pencil, and the more distant will its focus be from the principal focus of the lens.

Questions of the following nature very often arise in optics, viz., the length of the principal focus of a convex lens being given, and the distance a certain object is in front of it;—to find how far behind the lens will be the inverted image of the object. Or to express it more technically, the length of the principal focus of a convex lens being given and the length of the divergent incident pencil, to find the length of the focus of convergent refracted pencil. Thus: Suppose you had the following question: A 10 inch lens is 60 inches from an object; how far behind the lens will be the inverted image?

This could be solved immediately, by actual trial, and measurement, but this is not always practical.

The rule given in some text books on optics is as follows: multiply the length of the divergent incident pencil, that is, the distance the object is from the lens, by the focal length of the lens, and divide by the difference; thus:  $60 \times 10 = 600$ ,  $60 - 10 = 50$ ,  $600$  divided by  $50 = 12$ ; or  $\frac{60 \times 10}{60 - 10} = \frac{600}{50} = 12 =$  the distance behind the lens.

There is another property of convex lenses which I must not omit to mention; namely, what is called it magnifying power.

When a convex lens is placed between the eye and an object,—the object being at a less distance from the lens than its principal focus, the object will appear enlarged or magnified. The shorter the focus of the lens, the greater is its magnifying power. Thus, a 4 inch lens has a greater magnifying power than an 8 inch lens; a 2 inch lens greater than a 4, and a 1 inch greater than a 2 inch lens. The 1 inch lens has, in fact, double the magnifying power of a 2 inch lens; a 2, double that of 4 inch; a 4 inch, double that of an 8 inch, &c.

The "power" of a lens is therefore inversely proportional to its focal length. For this reason a different form is used in expressing the "power" or strength of a lens. A 1 inch lens is taken as unity,

and as a 2 inch lens is just half the strength, it is simply expressed  $\frac{1}{2}$ , and as a 3 inch lens has just one-third the strength of a 1 inch, it is written  $\frac{1}{3}$ ; a 4 inch is  $\frac{1}{4}$  &c. We will find that this nomenclature is not only very convenient, but scientifically correct.

For example, suppose we have two lenses of 4 inch focus each, and we wish to know their combined "power" when used as one lens; we simply add their reciprocals thus  $\frac{1}{4} + \frac{1}{4} = \frac{2}{4} = \frac{1}{2}$ . The two lenses have, therefore, the magnifying power of  $\frac{1}{2}$ , which is the reciprocal of 2, and are consequently, together, equal to a 2 inch lens, which can be proved by actual measurement. Again, suppose we have a 6 inch lens, and a 12 inch lens, and we wish to know their combined strength,  $\frac{1}{6} + \frac{1}{12} = \frac{2}{12} = \frac{1}{6}$  which represents the power of a 4 inch lens; the 6 and the 12 inch lenses taken together being equal to one lens having a focus of 4 inches.

To save repetition, I may here state that when a *concave* lens enters into combination with a *convex* lens, it has a neutralizing effect upon the convex lens. If we have a convex 6 and a concave 6 the one would neutralize the other,—thus  $\frac{1}{6} - \frac{1}{6} = 0$ . But if the convex lens has the higher power, the concave lens simply weakens it—that is, lengthens its focus—thus, if we have a convex 6 and a concave 9 the result will be  $\frac{1}{6} - \frac{1}{9} = \frac{3}{18} - \frac{2}{18} = \frac{1}{18}$ , which represents the strength of one lens having a focus of 18 inches. If, however, the concave lens has the higher "power" it will simply be weakened by the concave lens,—the combination will be equal to a concave lens having a lower "power," or a longer focus than the concave lens taken,—thus reversing the last example, suppose we have a *concave* 6 and a *convex* 9, we will then have  $-\frac{1}{6} + \frac{1}{9}$  or simply  $\frac{1}{9} - \frac{1}{6} = \frac{2}{18} - \frac{3}{18} = -\frac{1}{18}$ , which represents the strength of a *concave* lens having a focus of 18 inches.

This fractional nomenclature (taking 1 for numerator and the focal length of the lens for denominator) will assist us also in understanding the principle of the formation of images at different distances behind a convex lens, according to the distance of objects in front of it.

Let me remind you that when an object, for instance the flame of a candle, is placed in the focus of a convex lens, the diverging rays of light from the object are rendered parallel by the lens. Thus, a lens having a focus of 20 inches will render parallel pencils of light diverging from an object 20 inches from the lens. Bearing this in mind let us again try the solution of the following question, pro-

pound  
a 10 in  
of the  
light e  
what d

Now  
of the  
from t  
to divi  
60 in  
would l  
Deduct  
portion  
are equ  
parallel  
falling  
principl  
illustrat  
inches i  
know h  
divergin  
this  $\frac{1}{18}$   
bring th

Fig. 4  
ten inch  
one havin  
The 30 in  
so as to  
parallel r  
are conve  
lens, viz.,

is simply expressed  
length of a 1 inch, it  
at this nomenclature  
rect.

inch focus each, and  
used as one lens;  
 $\frac{1}{2}$ . The two lenses  
is the reciprocal of  
inch lens, which can  
we have a 6 inch  
combined strength, the 6  
4 inch lens; the 6  
to one lens having

when a *concave* lens  
neutralizing effect  
and a concave 6  
 $=0$ . But if the  
s simply weakens  
a convex 6 and  
 $\frac{1}{3}$ , which repre-  
sents 3 inches. If, how-  
ever, it will simply be  
equal to  
or focus than the  
example, suppose we  
 $-\frac{1}{3} + \frac{1}{3}$  or simply  
of a *concave* lens

generator and the  
also in under-  
different distances  
acts in front of it.  
hence the flame of  
the diverging rays  
lens. Thus, a  
pencils of light  
Bearing this in  
question, pro-

pounded not long since, viz. :—When an object is 60 inches in front of a 10 inch convex lens, how far behind the lens will be the inverted image of the object? Or, to express it differently, when a divergent pencil of light emanates from a point 60 inches from a 10 inch convex lens, at what distance behind the lens will the pencil be converged to a focus?

Now, we know that a lens of 60 inches focus, placed in the position of the 10 inch lens, would render the rays parallel that fall upon it from the object 60 inches distant. Were it possible, therefore, to divide the 10 inch lens into two lenses, one having a focus of 60 inches to render the rays parallel, the remaining portion would bring these parallel rays to a focus at its principle focus. Deducting then  $\frac{1}{60}$  from  $\frac{1}{10}$  will give the strength of the remaining portion of the lens  $\frac{1}{10} - \frac{1}{60} = \frac{5}{60} = \frac{1}{12}$ ; the two parts then  $\frac{1}{60}$  and  $\frac{1}{12}$  are equal to the one lens  $\frac{1}{10}$ . And, as the  $\frac{1}{60}$  will render the rays parallel from the object 60 inches distant, and these parallel rays falling upon the other part  $\frac{1}{12}$ , they will be brought to a focus at the principle focus of this part, viz. : at 12 inches from the lens. Let us illustrate this with another example. Suppose that an object is 30 inches in front of a convex lens of 10 inch focus, and we wish to know how far behind the lens will be the focus of a pencil of rays diverging from a point in the object. We will have  $\frac{1}{10} - \frac{1}{30} = \frac{2}{30} = \frac{1}{15}$ ; this  $\frac{1}{15}$  represents the power of a 15 inch lens, which we know will bring the parallel rays to a focus at 15 inches behind the lens.



Fig. 4 illustrates this; O represents an object 30 inches from a ten inch convex lens, the lens supposed to be divided into two parts, one having a focus of 30 inches, and the other a focus of 15 inches. The 30 inch lens refracts the rays of the divergent pencil  $d, d, d, d,$  so as to render them parallel, as shown at P, P, P, P, P. These parallel rays, meeting the 15 inch lens, are again refracted and are converged to a focus at F, which is the principle focus of the lens, viz., at 15 inches.



Fig. 1, page 3, represents a 10 inch lens, at a distance of 20 inches from an object, F. The lens is supposed to be divided into two equal parts, of 20 inch focus each: the first half renders the diverging pencil parallel, and the second half converges the parallel pencil to a focus, at 20 inches from the lens;  $\frac{1}{10} - \frac{1}{20} = \frac{1}{20}$ .

(Dr. Giraud-Teulon, of Paris, has ascribed the origination of the above theory to Mr. J. Z. Laurence, of London, to whom we are very much indebted, for his praiseworthy efforts to popularize this hitherto neglected, field of Physiological and Pathological Optics.)

Let me next direct your attention to certain optical considerations which have a most important application, in the treatment of optical defects of the eye.

You may remember that in a former experiment, a 10 inch lens was held ten inches from a white wall, so as to show the miniature inverted picture of the window, &c., 20 ft. distant; and that when the lens was brought to a distance of 60 inches from the window, it was found that the image of the window was formed 12 inches behind the lens, instead of 10 inches, and that at 10 inches, the image was so indistinct as to be scarcely recognizable.

Now suppose that a 12 inch lens be immovably fixed 12 inches from the same wall, it will then be in a proper position to bring parallel rays to a focus on the wall, where it will form an inverted picture of the window, and objects at a distance beyond the window.

If we now bring the flame of a lamp, for instance, to a distance of 60 inches from the lens, no distinctly defined image of the flame will appear upon the wall; but if, by any means, we can render the pencil parallel that diverges from the flame, the 12 inch lens will then converge it accurately to a focus upon the wall, where we will have an inverted image of the flame.

From the knowledge that we have now obtained, we know that a 60 inch lens placed in front of the 12 inch lens will render these rays parallel. All that we have to do then is to combine a 60 inch lens with the 12 inch lens: the 60 inch lens to render the rays parallel that diverge from the flame, 60 inches distant, and the 12 inch lens to converge these rays to a focus, at the principal focal length of the lens. This is exactly what we do in supplying old people with convex spectacles. Their eyes are constructed to bring parallel rays to a focus, on the retina; but the rays from near objects are too divergent to be focussed upon the retina without artificial aid: this deficiency is what we supply with suitable glasses.

Before  
subject  
mation

Many  
is admit  
pears up  
picture  
also be a  
few inch

The p

Let A  
short dis  
of diverg  
direction  
minate th  
small ori  
box, and

Rays c  
of these  
back of t  
pencils o  
one ray f  
tion of t  
back of t  
of the bo  
The reas  
the same  
flame, aft  
part of t  
flame mu  
upper pa

distance of 20 inches  
be divided into two  
half renders the diver-  
erges the parallel pen-  
 $\frac{1}{20} = \frac{1}{20}$ .

the origination of the  
ndon, to whom we  
rts to popularize this  
Pathological Optics,  
ptical considerations  
treatment of optica

ment, a 10 inch lens  
show the miniature  
ant; and that when  
from the window, i  
formed 12 inches be  
10 inches, the imag

ably fixed 12 inches  
sition to bring par  
orm an inverted pic  
ond the window.  
tance, to a distance  
image of the flame  
we can render the  
e 12 inch lens will  
wall, where we will

ned, we know that  
s will render these  
combine a 60 inch  
nder the rays par-  
t, and the 12 inch  
ncipal focal length  
plying old people  
d to bring parallel  
a near objects are  
out artificial aid-  
ses.

Before leaving the consideration of optical lenses, there is one subject to which I wish to direct your attention; namely, the formation of an inverted image behind a convex lens.

Many of you are, probably, familiar with the fact, that when light is admitted into a darkened room, through a small orifice, there appears upon the opposite wall of the room, an inverted, dim, shadowy picture of buildings, trees, &c., in front of the aperture. This can also be seen, on a smaller scale, by holding a sheet of white paper a few inches from the key-hole of a darkened hall.

The philosophy of this is seen in Fig. 5.



Let A, B, represent the position of a flame of a lamp that is a short distance in front of an aperture of a darkened box. Pencils of divergent rays of light radiate from the apex of the flame in every direction; one of these pencils is represented in the figure to illuminate the end of the box, and one of the rays escaping through the small orifice *c*; this ray passes in a straight line to the back of the box, and strikes the point *a*, which it illuminates.

Rays of light diverge from the lower part of the flame, also; one of these rays is shown to enter the aperture *c*, and to pass to the back of the box at *b*. In a similar way it might be illustrated that pencils of light radiate from every point in the flame A, B, and that one ray from each point passes into the box and illuminates a portion of the back. In this way we get an illuminated spot at the back of the box, which is an exact counterpart of the flame in front of the box, but *inverted*, the apex of the flame pointing downwards. The reason that the picture is reversed is that, as rays of light (in the same medium) pass in straight lines, a ray from the top of the flame, after passing the aperture, must necessarily pass to the lower part of the back of the box; and a ray from the lower part of the flame must necessarily (in moving in a straight line) pass to the upper part of the back of the box. You will observe, also, that the

size of the image depends upon its distance behind the aperture; if the image is as far behind the aperture, as the object is in front, the image will be of the same size as the object, if half the distance, half the size, as seen at f, g.



If, in the above experiment, the aperture be enlarged, it will be found that the image at the back of the box will become much less distinct; the more the aperture is enlarged, the more indistinct will be the image. The reason of this indistinctness in the image is this: when the aperture is enlarged, a number of diverging rays from every point in the flame pass through the aperture, and each one repeats the image, so that the parts of the image overlap each other.

This is shown in Fig. 6. A, B, represents the flame of the lamp, and C, E, D, F, the image behind an aperture. The aperture is supposed to be just large enough to admit two divergent rays, each of these rays produces a separate image; thus, the point A is repeated twice at D and F, and the point B is repeated at C and E. The larger the aperture, the more light is admitted, but the more indistinct is the image.

If now, a convex lens be inserted in the enlarged aperture, the divergent rays that enter the aperture (from every point of the object) are converged to a focus; thus in

Fig. 7.




Fig. 7.  
 o the in  
 point A a  
 rays from  
 manner, c  
 enter the  
 We will t  
 of the obj  
 paper we  
 upon thin  
 behind; an  
 glass be re  
 pertion) o  
 pried—bein  
 called an a  
 able circum  
 line with th  
 behind the

The hum  
 diameter.  
 crystalline l  
 convexity o  
 about one i

When a r  
 of rest), par  
 and a very  
 upon that r  
 back of the  
 decreas  
 opposite sid  
 in front of t

The impr  
 through the  
 municates to  
 more than v  
 "God has w  
 We are a

behind the aperture  
the object is in front  
if half the distance

Fig. 7. A C represents an object in front of a convex lens, and  $a c$  the inverted image behind the lens. Rays diverging from the point A and falling upon the lens L are brought to a focus at  $a$ ; rays from B are similarly focussed at  $b$ , and so on. In a similar manner, diverging rays from every point in the object A C that enter the lens are brought to a focus in the image between  $a$  and  $c$ . We will then have in the position of  $a c$  a distinct inverted image of the object A C. If this image is received upon a sheet of white paper we can see it only upon its front surface; but if it is received upon thin oiled paper, or upon ground glass, we can see it from behind; and if, while viewing the image from behind, the ground glass be removed, we can still see the inverted image (or at least a portion) occupying the same position as the ground glass just occupied—being suspended, as it were, in the air, and forming what is called an ærial image. In order to see this ærial image under favourable circumstances, one eye only should be used, and should be in a line with the lens and the object, and should be at least ten inches behind the position of the inverted lens.

enlarged, it will  
will become much  
more indistinct  
in the image is  
diverging rays from  
and each one repe  
up each other.

the flame of the lac  
e. The aperture  
divergent rays, ca  
the point A is  
repeated at C and  
mitted, but the m

enlarged aperture, the  
every point of

## CHAPTER II.—OPTICS OF NORMAL EYE.

The human eye, from before backwards, is about one inch in diameter. Its transparent media are the cornea, aqueous humour, crystalline lens, and vitreous humour. This combination, with the convexity of the cornea, is equal to a convex lens having a focus of about one inch (more accurately  $\frac{1}{8}$  of an inch.)

When a normal eye is directed to a distant object (*i. e.* in a state of rest), parallel rays of light are brought to a focus upon the retina, and a very minute inverted picture of the object is sharply defined upon that membrane. If the sclerotic coat be removed from the back of the eye of an ox, and the eye be placed in an aperture of a darkened room, with the cornea looking, for instance, towards the opposite side of the street, an inverted image of the buildings, &c., in front of the aperture will be seen at the back of the eye.

The impression that objects make upon the retina, is conveyed through the optic nerve to the brain, but in what manner this communicates to the mind a knowledge of the appearance of objects, is more than we can tell. We can simply say with Potterfield, that "God has willed it so."

We are aware, however, that although the eye may be free from

disease, and the connection between the retina and brain in every way perfect, if the optical mechanism of the eye be in any way defective so as to produce ill defined images upon the retina,—vision will be indistinct, and that the distinctness or indistinctness of vision will be in exact proportion to the distinctness or indistinctness of the inverted picture. Hence the necessity of understanding the optics of the eye in order to comprehend the pathology and treatment of the numerous optical defects to which it is liable.

CASE 1. Let me here take an example. A few weeks ago a physician of this city sent a patient for my advice, fearing that he was losing the sight of his left eye. Upon examination, I found that he had what we call "paralysis of accommodation" of that eye.

He could see distant objects with perfect distinctness, but near objects he was unable to define; he could not read large type unless the letters were very large, and several feet from the eye. The eye was, in fact, simply passive, like a convex lens, or a camera-obscura, with the screen to receive the image immovably fixed at the principal focus of the lens, and could only bring parallel rays to a focus upon the retina.

I found that by rendering the diverging rays parallel, by means of a convex lens, he could see near objects distinctly; by placing a 6 inch convex lens before that eye, he could read fine type at six inches, with a 10 inch lens at ten inches, with an 18 inch lens at eighteen inches, &c. &c. The 6 inch lens rendered the rays parallel that diverged from the letters six inches distant, and these parallel rays falling upon the eye were brought to a focus upon the retina. A 6 inch lens does not increase the apparent size of letters one-half, whereas this patient could not see letters ten times the ordinary size at six inches, or any distance less than about two feet from the eye. The 10 inch lens rendered the rays parallel from objects ten inches distant, and the 18 inch lens from objects eighteen inches distant.

The eye was unable to bring diverging rays to a focus upon the retina; in other words it had lost the power of "accommodation." (We can temporarily paralyse the accommodation of the eye by applying a strong solution of Atropine.)

A normal eye differs from the glass lenses we have been described in the fact that it can, not only focus parallel rays upon the retina, but also rays that diverge from objects as near as from four to six inches from the eye. When parallel rays fall upon a 1 inch convex

lens, they form an object, for instance, 6 inches from the lens, will be inverted behind the lens upon the lamp upon the screen behind the lens.

Now when the eye has no accommodation, it has no power, as to the eye, that has it, rays parallel

Fig. 8 represents the eye accommodated for distant vision. The rays from F, when they reach the eye, are parallel. The manner in which they are effected is shown in the diagram. The nature,—a t

The accommodation of the eye is effected by the contraction of the ciliary muscles. The iris is contracted, and the lens is brought into a more convex shape. The power of the eye is increased, and it is able to focus near objects upon the retina. When the eye is accommodated for distant vision, the ciliary muscles are relaxed, and the lens is brought into a less convex shape. The power of the eye is decreased, and it is able to focus parallel rays upon the retina.

ina and brain in eve  
 re be in any way defe  
 the retina,—vision w  
 undistinctness of visio  
 or indistinctness of t  
 nderstanding the opt  
 ology and treatment  
 ble.  
 few weeks ago a ph  
 , fearing that he w  
 nation, I found that  
 " of that eye.  
 distinctness, but he  
 read large type unlik  
 om the eye. The  
 s, or a camera-observ  
 y fixed at the principl  
 l parallel rays to a focus

lens, they are brought to a focus one inch behind the lens, but if an object, for instance the flame of a lamp, be brought to within four inches of the lens, we know that the focus will fall farther than one inch behind the lens. If we wish to receive the inverted image of the lamp upon a screen, the screen must be held one inch and a third behind the lens.

Now when an object is brought to within, say four inches of the eye, it has no power to move the retina backwards to receive the image that would be formed behind that membrane, but, what answers the same purpose, it has the property of so far increasing its refractive power, as to be able not only to render parallel, these diverging rays, but also to focus them upon the retina. This increase in the power of the eye, is equal to the addition of a 4 inch lens in front of an eye that has its "accommodation" paralysed, as a 4 inch lens renders rays parallel that diverge from objects four inches distant.



s parallel, by means  
 nctly; by placing a  
 fine type at six inch  
 3 inch lens at eight  
 the rays parallel th  
 and these parallel r  
 s upon the retina.  
 size of letters one-h  
 times the ordinary  
 wo feet from the ey  
 om objects ten inch  
 teen inches distant  
 s to a focus upon  
 of "accommodatio  
 on of the eye by app

Fig. 8 represents the section of a normal eye. When it is accommodated for distant objects parallel rays P, P, are focussed upon the retina at F, while diverging rays from O, would form a focus at *fd*. When, however, the eye is accommodated for the near object O, these diverging rays are focussed upon the retina at F.

The manner in which this increase in the refractive power of the eye is effected is still a disputed point. Most physiologists however are now inclined to the theory that it is caused by an increase in the curvature,—a thickening from before backwards, of the crystalline lens.\*

e have been describ  
 ys upon the retina,  
 om four to six or  
 upon a 1 inch cou

\* The accommodation of the eye was at one time believed to be produced by the external muscles, but it is now ascertained that the accommodation can remain perfect with all the external muscles paralysed.

The iris was thought, by others, to have the power of increasing the refractive power of the eye, but it was proved by a case that occurred in Dr. Von Graefe's practice that accommodation can still be effected with entire absence of the iris.

Helmholtz and Cramer have proved by means of the ophthalmometre, that when the eye is accommodated for a near object it undergoes the following changes:—

The "near" and "far" point.—The nearest point to which objects can be brought to an eye and be seen with perfect distinctness, is called the "near" point, and the farthest point of distinct vision is called the "far" point.

In a normal eye the "near" point is about seven inches from the front of the cornea, and the "far" point is at an unlimited distance. In childhood, however, the "near" point is about  $3\frac{1}{2}$  inches from the eye and recedes as age advances. At the age of forty the "near" point of a normal eye is nearly eight inches from the eye.

When the "near" point recedes to a greater distance than eight inches from the eye it becomes inconvenient; such an eye is called *presbyopic* or long-sighted.

When the "far" point is not unlimited, but is at a definite distance from the eye, as for instance from six inches to four or five feet from the eye—such an eye is called *myopic* or short-sighted.

*Range of Accommodation.*—The distance between the "near" and "far" point in any eye, is called the "range of accommodation." If a person can read distinctly very fine type at four inches from the eye, and can also see clearly at an infinite distance the range of accommodation would be said to equal  $\frac{1}{4}$  because, when such an eye is directed to objects at an infinite distance, (accommodated for parallel rays) in order to see clearly objects only four inches distant, it is necessary to increase the curvature of the crystalline lens, or in other words the "power" of the eye to an extent equal to the addition of a 4 inch convex lens; the power of which is expressed by  $\frac{1}{4}$ . If a person's "near" point is at eight inches from the eye, and his "far" point at an infinite distance, his range of accommodation would be said to equal  $\frac{1}{8}$ .

If the "near" point of a myopic eye be 3 inches, and the "far" point be 12 inches, we get the range of accommodation by the equation  $\frac{1}{3} - \frac{1}{12} = \frac{1}{4}$ .

### CHAPTER III.—MYOPIA.

CONCAVE LENSES.—Before proceeding to the consideration of Myopia, it will be well for us to glance at some of the properties of concave lenses; and, in order to simplify the subject, we will confine

1st. The pupil contracts; 2nd. The pupillary edge of the iris moves forward; 3rd. The peripheral portion of the iris moves backwards; 4th. The anterior surface of the lens becomes more convex (arched); 5th. The lens does not change its position; 6th. The cornea retains the same degree of curvature.

course  
two s  
which

Fig  
centre  
rays, l  
leading  
centre  
princip  
In a co  
lens, it  
lens p  
When  
to a fo  
diverge  
rent si  
behind  
therefo  
a conve  
lens. ( (-), a  
To asce  
convex

1. In  
fair deg  
inverted  
the pict  
object c

Patier  
objects i

t point to which objects  
perfect distinctness,  
point of distinct vision is

seven inches from the  
an unlimited distance  
out  $3\frac{1}{2}$  inches from the  
forty the "near" point  
eye.

r distance than eight  
such an eye is called

s at a definite distance  
four or five feet from  
sighted.

een the "near" and  
accommodation." If  
inches from the eye,  
the range of accommoda-  
ch an eye is directed  
for parallel rays) in  
nt, it is necessary to  
in other words the  
tion of a 4 inch con-  
f a person's "near"  
"far" point at an  
d be said to equal  $\frac{1}{2}$   
nes, and the "far"  
accommodation by the

e consideration of  
f the properties of  
ect, we will confine

moves forward; 3rd.  
he anterior surface of  
not change its posi-

ourselves to equi-concave lenses. An equi-concave lens is bounded by two surfaces, which are portions of the concave side of two circles which have equal radii.



Fig. 9. A, B, one of the concave surfaces of the lens. C is the centre of curvature, and C, R the radius of curvature. When parallel rays, P, P, strike one surface of the lens, they have a divergence upon leaving the second surface of the lens, as if they proceeded from the centre of curvature, C, which, in an equi-concave lens, is also the principal focus of the lens. C, R, is the focal length of the lens. In a convex lens, the focus is measured *behind* the lens; in a concave lens, it is measured *in front* of it. If we call the focus of the convex lens positive, we must call the focus of the concave lens negative. When parallel rays of light fall upon a convex lens, they are converged to a focus. When they fall upon a concave lens, they are made to diverge. A convex lens enlarges, and a concave diminishes the apparent size of objects. The focal length of a convex lens is measured behind; and that of a concave lens, in front of the lens. They are, therefore, entirely opposite in all their properties; and, for this reason, a convex lens is called a positive lens; and a concave one, a negative lens. Or, shorter still, they are indicated by the plus (+) and minus (-), algebraic symbols; thus, + 5, and - 5; or,  $+\frac{1}{5}$ , and  $-\frac{1}{5}$ . To ascertain the focal length of a concave lens, we ascertain what convex lens it will neutralize.

1. In a myopic eye, parallel rays, as well as those that have a certain degree of divergence, are focussed *in front* of the retina; and, the inverted image of distant objects being formed in the same position, the picture upon the retina will be ill-defined, and vision for distant object consequently indistinct.

Patients with myopia complain that, although their vision for near objects is perfect, they cannot see objects at a distance with any dis-



tinctness. They can read the smallest type, when brought near the eyes, even better than persons with normal vision, but they are unable to recognize their friends at a distance of fifteen or twenty feet.

In order to enable such persons to see distinctly at a distance, it is necessary for them to wear concave spectacles of such a strength, that the parallel rays from distant objects may have such a degree of divergence, that, falling upon the myopic eye, they may form a focus upon the retina. Theoretically, we should prescribe concave glasses of such a strength that their focus will correspond with the patient's "far" point. Thus, if the "far" point be at 12 inches, we should prescribe  $-12$ , as a twelve inch concave lens, placed before such an eye, will give parallel rays from distant objects the same degree of divergence as if they proceeded from the "far" point of the eye, namely, at 12 inches from the eye. Thus, in Fig. 9, P. P. represent parallel rays falling upon the concave lens, A. B.; they are made to diverge, as if coming from the focus, C., and falling upon the eye, divergingly, they are focussed upon the retina at F. Practically, however, we would find that  $-12$  would be rather too strong, and that  $-15$ , or  $-16$  would probably answer better. As a rule, the weaker glasses should be worn that will enable the patient to see distant objects with distinctness.

In testing the degree of myopia, we use a series of test types that are so constructed that No. I (smallest) can be distinctly seen and read by a person having normal vision, at a distance of 1 foot; No. II, at 2 feet; No. V, at 5 feet; No. XX, at 20 feet; and so on. A series of specimens of these types will be annexed to this paper. The types are also used in testing the acuteness of vision in Presbyopia, Hypermetropia, Amblyopia, &c.

2. In determining the degree of myopia in any case, we ascertain the greatest distance at which No. I test types can be read distinctly. If at 10 inches, the "far" point will be at 10 inches, and the myopia would be called  $\frac{1}{10}$ ; if at 6 inches, the myopia would be called  $\frac{1}{6}$ . From this we can, as stated above, get a proximate knowledge of the strength of the concave lens necessary to relieve the myopia.

3. A myopic eye, when in a state of rest, is adjusted for divergent rays. To enable such an eye to see distant objects, that is, to give parallel rays to a focus on the retina, it is necessary to give the parallel rays a preliminary degree of divergence by the interposition of the proper concave lens.

4. Myopia by the far method: test type that a patient at 6 inches, or paired the can see a the contra (much sm even No.

This distant near object frequent cause

Short-sighted of spectacles will enable changed to more frequent should be that short-writing, &c purpose, for

1st. "Be paired with ated action, existing with increase of upon the eye convergent, sure cannot staphyloma.

2d. "On regarding the cause an increase within the eye oeditis poste

4. Myopia can be distinguished from every other defect of vision, by the fact that concave glasses improve vision for distant objects. If we have no concave glasses convenient, we can diagnose it from Amphyopia, (insensibility of the retina) by the following ready method:—A person with normal vision can read distinctly, No. I test type at 12 inches, and even a little farther. We will suppose that a patient's vision is so impaired, that he can only read No. II at 6 inches; if he is *not* also myopic, he can also read No. IV at 12 inches, or No. LX at 180 inches—that is at 15 feet. However impaired then a person's vision may be, unless he be also myopic, he can see as well proportionately, at one distance as at another. On the contrary, a person with myopia, say  $\frac{1}{2}$ , can see the smallest type (much smaller than No. I,) at 6 inches, but he cannot see No. II, or even No. V, at 12 inches.

This disease is often hereditary. Over exertion of the eyes upon near objects at the age of puberty, (about 14 or 15) is a very frequent cause of myopia.

Short-sighted persons often inquire if we would advise the use of spectacles. There can be no objection to wearing glasses that will enable them to see distant objects; for their eyes are thus changed to normal ones, but as most persons use their eyes much more frequently upon near than upon distant objects; the glasses should be no stronger than necessary. Some contend, however, that short-sighted persons should dispense with glasses for reading, writing, &c. Prof. Donders, however, recommends their use for this purpose, for the following reasons:—

1st. "Because strong convergence of the optic axes is necessarily paired with tension of the accommodation. The latter is an associated action, not arising from the mechanism of the convergence, but existing within the eye itself, and may consequently easily lead to an increase of the myopia. Besides this, the pressure of the muscles upon the eye ball appears to be greater when the optic axes are convergent, than when they are parallel, and this increase of pressure cannot but tend to give rise to the development of posterior staphyloma.

2d. "On account of the habit which short-sighted persons have of bending their head forwards during reading or writing. This must cause an increased flow of blood to the eye, and an increased tension within the eye itself. Owing to this development of sclerotic—choroiditis posterior, effusions of blood and detachment of the retina

which are so apt to occur in short-sighted persons, are undoubtedly greatly promoted. For this reason, we should always tell the patients to read with their head well thrown back, and to write at sloping desk. But it may, on the other hand, be urged that it is just in looking at near objects that myopic persons have an advantage, for they can see them remarkably distinctly. And the great danger is, that after reading for a short time with spectacles, the patient, on getting somewhat fatigued will, instead of laying the book aside, approach it nearer to the eye, in order to gain greater retinal images, and thus strain and tax his power of accommodation too much. If we, for instance, give a patient whose far point lies at 12 inches, a pair of spectacles which enable him to read at 12 inches, he will, if not very careful, after a short time almost insensibly bring the book nearer to his eyes, and thus have to make use of a great amount of accommodation. If he does this frequently, he will so increase his myopia. The greater the range of accommodation the less harm will spectacles do, and *vice versa*. Spectacles may also be used for near objects in those cases of myopia in which asthenopia (depending upon insufficiency of the internal recti muscles) shows itself as soon as the patient has read or worked at near objects for a short time. Whilst these forms of myopia may be furnished with spectacles for near objects, it is very dangerous to permit them in patients whose range of accommodation is very limited, and who moreover, suffer perhaps from such an amount of amblyopia (generally depending upon sclerotic—choroiditis posterior) that they cannot read No. 4 or 5 Jäger even with the most accurately chosen glasses. Such patients will bring the object very close to the eye in order to obtain large retinal images, the accommodation will be greatly strained, the intra-ocular tension be increased, and great mischief will be sure to ensue. If there is much amblyopia, spectacles should not be permitted at all for near objects.\*

In cases where the myopia is extreme, there usually co-exists a posterior staphyloma of the sclerotic. Von Graefe says it is present in all cases of myopia where the "far" point is less than five inches, the myopia being less than  $\frac{1}{2}$ . Out of sixty cases of myopia examined by J. Z. Laurence, forty-four had posterior staphyloma.

The presence of this disease can be easily diagnosed with the ophthalmoscope. (See Hulke or Zander on the ophthalmoscope

\*Mr. J. Z. Laurence, of London, recommends that deeply concave lenses be fitted in order to obviate their "dazzling" effect.—(Med. Times and Gazette, Oct. 22nd, 1864.)

Pos  
sensib  
of the  
detach  
preven  
tened

Don  
clo e  
and cr

The  
1st.  
2nd.  
3rd.  
rays.

4th.

You  
and dir  
upon th  
parallel  
howeve  
(if con  
is, ther  
of the e  
When t  
power o  
hyperme  
verging  
at a def  
metropi  
improve  
one.

This i  
within t  
of Glasg

From

persons, are undoubtedly should always tell the back, and to write at and, be urged that it persons have an adva distinctly. And the gra me with spectacles, t instead of laying the bo to gain greater retin of accommodation t whose far point lies a to read at 12 inches, almost insensibly br make use of a great frequently, he will se of accommodation t Spectacles may also pia in which asthenic al recti muscles) she d at near objects fo may be furnished wous to permit their very limited, and w at of amblyopia (ge s posterior) that t most accurately cho t very close to the e accommodation will o increased, and gr uch amblyopia, spee ects."\*

usually co-exists p te says it is present less than five inch cases of myopia ex or staphyloma.

y diagnosed with the ophthalmoscop

Posterior staphyloma is a serious complication in myopia, as the sensibility of the retina becomes more or less impaired in the position of the bulging of the sclerotic, and in some cases the retina becomes detached from the choroid. It is the existence of this disease that prevents improvement in cases of myopia, as the eye becomes flattened with advancing age.

Donders considers that in myopia, the antero-posterior diameter is alone at fault; that is, it is too much elongated, and that the cornea and crystalline lens have usually a normal curvature.

The characteristics of a myopic eye, are\*

- 1st. Parallel rays are focussed in front of the retina.
- 2nd. The "far" point is at a definite distance and positive.
- 3rd. When the eye is in a state of rest it is adapted for divergent rays.
- 4th. Concave glasses improve vision.

#### CHAPTER IV.—HYPERMETROPIA.

You will remember that when a normal eye is in a state of rest, and directed to a distant object, parallel rays are brought to a focus upon the retina, and that when a myopic eye is in a state of rest, parallel rays are brought to a focus in front of the retina. When, however, a hypermetropic eye is in a state of rest, parallel rays would (if continued) form a focus behind the retina. Hypermetropia is, therefore, the reverse of myopia. In myopia, the refractive power of the eye is excessive, and in hypermetropia it is not strong enough. When the accommodation of a myopic eye is paralysed, it has the power of focussing none but diverging rays upon the retina, but a hypermetropic eye under the same circumstances can focus only converging rays upon the retina. The "far" point of a myopic eye is at a definite distance and positive, but the "far" point of a hypermetropic eye is at a definite distance and negative. Concave glasses improve the vision for a myopic eye, and convex for a hypermetropic one.

This is an affection which has received very little attention until within the last ten years. It was indeed noticed by Dr. McKenzie of Glasgow, in 1841, but it was not until about five years ago that

\* From Donders' system of classification.

Prof. Donders, of Utrecht, from his elaborate researches on this subject, first pointed out how common this affection is, and how frequently it is the sole cause of that peculiar weakness of sight (formerly so little understood) called asthenopia.

Donders believes that this condition of the eye depends more upon a shortening of the antero-posterior diameter of the eye, than upon a too low degree of its refractive power; that the cornea and crystalline lens have a normal degree of curvature, and that parallel rays would form a focus at the normal distance behind the lens, were the retina far enough back to receive it.

A very good illustration of a hypermetropic eye is one in which the crystalline lens has been removed in the operation for cataract. To enable such an eye to see distinctly, even distant objects, it is necessary to place in front of it a strong convex lens of about four inches focus, called a cataract glass. The eye having too low a refractive power to converge rays to a focus, on the retina, it is necessary to give rays falling upon the eye, a preliminary degree of convergence; the eye having sufficient power to complete their refraction to a point upon the retina. We do the same thing in relieving cases of hypermetropia.




Fig. 10 represents a hypermetropic eye in a state of rest.  $PP$  are parallel rays which are focussed behind the retina at  $f$ .  $L$ , Fig. 11, is a convex lens which changes the parallel rays to convergent ones, at  $c$ ,  $e$ , as if they came from the direction  $ab$  and  $de$ , which again are refracted by the eye, and brought to a focus upon the retina at  $F$ .

When a hypermetropic eye is in a state of rest, and directed to distant objects, it is adjusted for convergent rays; images upon the retina will consequently be ill defined, and vision will be indistinct. To remedy this, it is necessary for the eye to increase its refractive power by increasing the antero-posterior diameter of the crystalline lens, so as to bring parallel rays to a focus on the retina.

Wh  
or acc  
tax hi  
diverg  
comm  
than a  
"near  
becom  
arise w  
Diag  
his eye  
xxx.,  
xx. at  
his visi  
same ty  
&c., un  
distan  
dation  
them;  
presum  
complai  
pronoun  
lens he  
Again  
distan  
at 20 fe  
convex  
hyper  
In or  
in any c  
power c  
cases of  
dation c  
distant  
their ow  
former  
accomm  
of his e  
retina, a

researches on this defect is, and how weakness of sight

depends more upon the eye, than upon the cornea and crystalline lens, and that parallel rays behind the lens,

is one in which the eye is not fit for cataract. To see distant objects, it is necessary to use a lens of about four inches focal length, or too low a refractive power. It is necessary to give the eye a degree of convergence; the eye is not adapted to a point of convergence in ordinary cases of hyper-

of rest. P P are at f. L, Fig. 11, are convergent ones, and d e, which again converge on the retina at F. If the rays are directed to distant objects, the images upon the retina will be indistinct. To increase its refractive power, the curvature of the crystalline lens is increased, and the retina.

When a person with hypermetropia, attempts to read or write, or accommodate his eyes to short distances, it is necessary for him to tax his accommodation to its utmost extent, in order to bring the diverging rays to a focus on the retina. This excessive effort at accommodating the eye for short distances, can not be kept up for more than a few minutes, when the ciliary muscle begins to relax,—the “near” point commences to recede, and (if he is reading) the letters become indistinct. The eye also feels fatigued, and other symptoms arise which will be referred to when speaking of Asthenopia.

*Diagnosis.*—When we suspect a patient has hypermetropia, we test his eyes as follows:—We place a series of test-types, No. xv., xx., xxx., &c., at a distance of about 20 feet. If he can read No. xv. or xx. at this distance, his acuteness of vision is normal. We then try his vision with weak convex glasses, say No. 50, and if he can read the same type, at the same distance, we try successively No. 40, 36, 30, 24, &c., until we reach the glasses that render the test type indistinct at that distance. Some persons may possibly be able to relax their accommodation so as to see as well at a distance, with convex 50 lenses, as without them; and not be hypermetropic; it would, however, be very strong presumptive evidence of its presence; and if, in addition, the patient complains of the symptoms of Asthenopia, we would be generally safe in pronouncing it a case of hypermetropia. The shorter the focus of the lens he can use, the stronger is the presumptive evidence of the disease.

Again, if another patient be tested with the same type, at the same distance, and we find that he can not read a smaller type than No. xl. at 20 feet without spectacles, and that he can read No. xv. or xx. with convex glasses, say + 10 or + 12, his would be called a case of hypermetropia *absolute*.

In order, however, to test accurately the degree of hypermetropia in any case, it is necessary to neutralize one element in the refractive power of the eye; namely, the power of accommodation. In most cases of hypermetropia, particularly in young subjects, the accommodation of the eye is so constantly exercised, even when directed to distant objects, that it is quite impossible for them, by any effort of their own, to completely relax that accommodation. I related in a former chapter, the case of a patient who had lost the power of accommodating his eye to different distances. As the refraction of his eye was normal, parallel rays were brought to a focus upon the retina, and vision for distant objects remained perfect.

Had his eye been hypermetropic, parallel rays would not have been sufficiently converged by the refractive power of the eye, to form a focus upon the retina; vision would, consequently, have been indistinct. By placing, however, the proper convex lens in front of such an eye, the requisite preliminary convergence would be given to the rays, to enable the eye, with its low refractive power, to focus the rays upon the retina, and thus render vision distinct.

The lens used in such a case would indicate the degree of hypermetropia. If the lens were a + 15 inch, the hypermetropia would be equal to  $\frac{1}{15}$ ; if a + 10, the hypermetropia would be  $\frac{1}{10}$ , and so forth.

We have, however, the means of temporarily producing this condition of the eye by artificial means. By applying a four-grain solution of atropine to the eye, within two hours the action of the ciliary muscle will be completely paralysed. A solution of one grain of atropine to an ounce of pure water (also a solution of the extract of belladonna) will dilate the pupil widely, and in some cases, render the eye slightly presbyopic, but it will not paralyse the accommodation.

If we test, in this manner, the case of suspected hypermetropia mentioned above, and find that after his accommodation is paralysed, he is not able to read No. xxx. even with + 50, and that the only glass with which he can read No. xv. and No. xx. 20 feet is + 20; his hypermetropia is therefore  $\frac{1}{20}$ . But as he could see as well with + 50 as without them, before his accommodation was paralysed; he had a manifest hypermetropia of  $\frac{1}{33}$ . The difference between his total hypermetropia and his manifest hypermetropia will give the amount of the latent hypermetropia which he overcame with the exercise of his accommodation, namely  $\frac{1}{33}$ , thus  $\frac{1}{20} - \frac{1}{33} = \frac{1}{66}$ .\*

*Asthenopia*, according to Donders, depends almost invariably upon hypermetropia. He describes it as follows: "The power of vision is usually acute,—and nevertheless, in reading, writing, and other close work, especially by artificial light, or in a gloomy place, the objects after a short time, become indistinct and confused, and a feeling of fatigue and tension comes on in, and especially above the eyes, necessitating a suspension of work. The person affected now often involuntarily closes his eyes, and rubs his hand over the forehead

\* Hypermetropia can easily be diagnosed with the ophthalmoscope.

eyelids.

the same

Accor

tion cor

give of t

gio pains

cases ext

year ago,

symptom

even on

forehead

the lid o

object for

Others, a

that, afte

that they

close the

same sym

In rega

when it is

of the acc

glasses to

cases), a c

will proba

As age

a normal c

Treatm

for the pat

to enable

the accom

the patient

approxima

degree of r

ary to par

strength, i

When we a

that we mu

lasts about

glasses. H

s would not have been of the eye, to form, nly, have been ind, x lens in front of su would be given to th e power, to focus the stinct.

the degree of hypermetropia would be  $\frac{1}{10}$ , and so forth.

ly producing this co applying a four gra ours the action of th solution of one gra solution of the extr and in some cases, r ot paralyse the acco

spected hypermetrop accommodation is pa with + 50, and th xv. and No. xx. fore  $\frac{1}{10}$ . But as em, before his acco hypermetropia of, pia and his mainit latent hypermetrop accommodation, name

almost invariably The power of vision writing, and other ch my place, the obje used, and a feeling above the eyes, ne detected now often in ver the forehead

hthalmoscope.

eyelids. After some moments rest, he once more sees distinctly, but the same phenomena are again developed more rapidly than before."

According to my own experience with these cases, the above description corresponds very closely with the description that most patients give of their symptoms. Some give more prominence to the neuralgic pains which they experience in and around the eye, and in some cases extending to the back of the head. I was consulted, about a year ago, by a lady from the town of Simcoe, C.W., who had all these symptoms in the most aggravated form. If she attempted to read even one line, it gave her so much pain in her eyes and forehead that, for several years, she had scarcely dared to even raise the lid of a book. She was unable to keep her eyes upon any one object for more than an instant at a time, without causing her pain. Others, again, do not speak of any pain or fatigue of the eye; but that, after reading a short time, the letters become indistinct, so that they are obliged to stop or look away at something distant, or close the eyes for a short time, when they can again proceed, the same symptoms recurring.

In regard to the *prognosis* in hypermetropia, Donders thinks that when it is once developed it never gives way. All the inconvenience of the accompanying Asthenopia can be relieved by wearing the proper glasses to relieve the hypermetropia; but the cause, namely (in most cases), a congenital flattening of the eye-ball from before, backwards, will probably remain through life.

As age advances, the "near" point recedes from the eye, as in a normal eye, so that in time it becomes complicated with presbyopia.

*Treatment.*—In order to correct this optical defect, it is necessary for the patient to wear a pair of convex spectacles of sufficient strength to enable him to see distant objects distinctly, without any effort of the accommodation. In cases where the hypermetropia is absolute, and the patients are not able to see distinctly at any distance, they can, approximately, by trial, select the glasses that will remedy the low degree of refraction of their eyes. But, in all other cases, it is necessary to paralyse the accommodation, and test with lenses of different strength, in order accurately to ascertain the degree of hypermetropia. When we ascertain this fact, we also know the number of the glasses that we must prescribe for them. The effect of the atropine usually lasts about a week, after which the patient can commence wearing glasses. Before, however, he use the spectacles that he is to wear



permanently, his accommodation must first be gradually relaxed by the use of weaker lenses. Donders' rule is to prescribe first that glass that will neutralize his manifest hypermetropia, and  $\frac{1}{4}$  of his latent hypermetropia, and every two or three weeks change them for a stronger pair, as he becomes accustomed to their use, until the glasses are reached that we found to be necessary to correct his hypermetropia. Thus, if a patient has a total amount of hypermetropia equal to  $\frac{1}{10}$ , and a manifest hypermetropia of  $\frac{1}{30}$ , his latent hypermetropia ( $\frac{1}{10} - \frac{1}{30} = \frac{2}{30}$ ), would equal  $\frac{1}{15}$ ; one fourth of  $\frac{1}{15}$  is  $\frac{1}{60}$ ; this added to  $\frac{1}{30}$  ( $\frac{1}{30} + \frac{1}{60} = \frac{2}{60} = \frac{1}{30}$ ), equals  $\frac{1}{30}$ . We would therefore prescribe, at first, 20 inch convex spectacles, which we would afterwards change successively for + 18, + 16, + 14, &c., until he has relaxed his accommodation that he can, with ease, wear + 10. It will not be until he becomes accustomed to this last pair that all his symptoms of Asthenopia will disappear.

*Strabismus.*—Prof Donders was the first to direct attention to the fact, that nearly all cases of convergent strabismus arise from the presence of hypermetropia. We know that when both eyes are directed to a near object, they are very much converged,—the optic axes cross at the point to which they are directed. If one eye is covered, and the opposite eye be accommodated for its "near" point, the converged eye will be found to be very decidedly converged toward the nose,—to have, in fact, a temporary convergent squint. This arises from the constant association of the act of accommodating the eye for short distances, with the act of contracting the internal recti muscles. The hypermetropic, however, being obliged to exert the accommodation of their eyes, even when looking at distant objects, it is easy to understand that they would be inclined to contract their internal recti-muscles unduly, so as to increase this power of accommodation. This converges the eyes to a point at a nearer distance than the object looked at, and causes one of the eyes to turn inward while the other is fixed upon the object. When, therefore, they wish to see distinctly with one eye, they instinctively turn in the other. At first the convergent strabismus is seen occasionally only, and in this stage may be prevented by using the proper spectacles to correct the hypermetropia. After the squint has existed sometime, it becomes confirmed and cannot be cured without an operation.

If the convergence exceeds three lines, a partial tenotomy, upon each eye, should be performed, and the effect controlled by a conjunc-

tival s  
operat

Wh  
withou  
very so

To  
they sh  
when r

The

1st.

2nd.

3rd.

4th.

5th.

nopia a

This

45. Mo

vision,

beginni

can see

In re

and per

Reading

frequent

In ch

from  $3\frac{1}{2}$

ited dist

$3\frac{1}{2}$  to 4 i

being in

tance.

40 the "

"near"

calls it

should n

10 inches

gradually relaxed by the  
 prescribe first that glau-  
 opia, and  $\frac{1}{4}$  of his later  
 weeks change them for  
 their use, until the glasse  
 correct his hypermetr  
 of hypermetropia equ  
 his latent hypermetrop  
 urch of  $\frac{1}{15}$  is  $\frac{1}{80}$ ; this  
 $\frac{1}{80}$ . We would therefor  
 which we would after  
 14, &c., until he has a  
 case, wear + 10. I  
 his last pair that all his

direct attention to the  
 strabismus arise from the  
 when both eyes are  
 converged,—the optic  
 rected. If one eye be  
 for its "near" point  
 dly converged toward  
 vergent squint. This  
 of accommodating the  
 icting the internal rec  
 obliged to exert the  
 g at distant objects, i  
 ined to contract the  
 this power of accom  
 at a nearer distance  
 eyes to turn inward  
 , therefore, they wish  
 ely turn in the other  
 sionally only, and in  
 r spectacles to correc  
 existed sometime, in  
 an operation.

partial tenotomy, upon  
 controlled by a conjunc-

tival suture, by which means we have the power of regulating our operation, in proportion to the effect we wish to produce.

When Strabismus shows itself in childhood, it should be treated without delay, for, if not corrected, the vision of the "cross-eye" will very soon become impaired.

To get the full benefit of spectacles, in cases of hypermetropia, they should be used both on the street, and at church, as well as when reading or writing,—in fact whenever the eyes are used.

The characteristics of a hypermetropic eye then are :

1st. Parallel rays form a focus behind the retina.

2nd. The "far" point is at an definite distance and negative.

3rd. The eye, in a state of rest, is adjusted for convergent rays.

4th. Convex glasses improve vision.

5th. This affection is usually accompanied by symptoms of Asthenopia and Amblyopia, and frequently by convergent strabismus.

#### CHAPTER V.—PRESBYOPIA.

This affection usually develops itself between the ages of 40 and 45. Most persons at this age, although previously enjoying excellent vision, complain that their sight, particularly in the evening, is beginning to fail for near objects, as small print, &c., although they can see distant objects as well as ever.

In reading they will hold the book or paper at nearly arm's length and perhaps bring the lamp almost between their eyes and the page. Reading in this manner soon fatigues them, and they are obliged frequently to rest,—or to resort to spectacles.

In childhood, when the vision is normal, the "near" point is from  $3\frac{1}{2}$  to 4 inches from the eye, and the "far" point at an unlimited distance; that is, we can see objects distinctly as near as from  $3\frac{1}{2}$  to 4 inches from the eye, and we can see objects clearly (the size being in proportion to the distance) from that to an indefinite distance. As age advances the "near" point recedes. At the age of 40 the "near" point is about eight inches from the eyes. When the "near" point recedes to a greater distance than 8 inches, Donders calls it a case of presbyopia; Laurence, however, thinks that it should not be called presbyopia unless the "near" point is at least 10 inches from the eye.

Presbyopia, then, is not an optical defect of the nature of myopia or hypermetropia, but is simply a lessening of the accommodative power of the eye.

It is supposed to depend upon, or to be caused by, the crystalline lens becoming hardened as age advances, so that it does not yield sufficiently to the contraction of the ciliary muscle.

In a case of pure presbyopia where, for instance, the "near" point is 12 inches from the eye, vision will remain normal for all points beyond that distance. When the "near" point is 12 inches distant and the "far" point at an infinite distance, the accommodation is only  $\frac{1}{12}$ . Taking eight inches as the normal "near" point,  $\frac{1}{8}$  would represent the normal accommodation. Deducting  $\frac{1}{12}$  from  $\frac{1}{8}$  gives the degree of presbyopia thus:  $\frac{1}{8} - \frac{1}{12} = \frac{1}{24}$ . The degree of presbyopia in this case would then be  $\frac{1}{24}$ . This fraction  $\frac{1}{24}$  also represents the strength of the glasses necessary to correct the presbyopia, namely 24 inch convex. Practically, we would probably find that a pair of 30 inch convex would answer better, as the weakest glass that can be worn with comfort, is the one that should be prescribed. Again, if a person's "near" point be at 16 inches, his presbyopia ( $\frac{1}{8} - \frac{1}{16} = \frac{1}{16}$ ) will be  $\frac{1}{16}$ , and a 16 inch convex lens would enable him to read at 8 inches.

"There can be no question as to the advisability and necessity of affording far-sighted persons the use of spectacles. They should be furnished with them as soon as they are in the slightest degree annoyed or inconvenienced by the presbyopia. Some medical men think that presbyopic patients should do without spectacles as long as possible, for fear the eye should, even at an early period, get used to them as soon to find them indispensable. This is, however, an error, for if such persons are permitted to work without glasses we observe that the presbyopia soon rapidly increases.\*"

If, however, we call all cases presbyopia, where the "near" point recedes to a greater distance than eight inches from the eye, it will follow that we may have presbyopia in cases of myopia and hypermetropia. If a person's far point be at 20 inches from the eye, he would be called *near-sighted* and if his near point recedes to 10 inches from the eye, he would be also *far-sighted*.

In some persons, as age advances, the "far" point also recedes

\* J. Soelberg Wells.

as to re  
seldom  
presbyop  
reading  
a pair o  
and as  
with th  
for rea  
Thus  $\frac{1}{8}$   
his hyp

In th  
enable  
usually

The  
general  
the pres

At 4

" 4

" 5

" 5

" 5

" 6

" 6

Prof.

we shou  
the pers  
of 12 in

There  
called as  
cal lines  
crystallin  
case, th  
astigmat

A very  
Times ar  
M.B., of  
The p  
referred

the nature of myopia  
of the accommodative

ed by, the crystalline  
that it does not yield  
ucle.

ce, the "near" point  
normal for all points  
t is 12 inches distant  
ne accommodation is  
near" point,  $\frac{1}{8}$  would  
ting  $\frac{1}{2}$  from  $\frac{1}{8}$  give  
r. The degree of  
s fraction  $\frac{1}{2}$  also re  
o correct the presby  
we would probably  
nswer better, as the  
s the one that should  
at be at 16 inches, h  
h convex lens would

ility and necessity  
les. They should be  
the slightest degree

Some medical men  
out spectacles as long  
an early period, get  
e. This is, however,  
work without glasses  
reases."\*

re the "near" point  
from the eye, it w  
f myopia and hyper  
ches from the eye  
point recedes to  
l.

point also recedes

as to render the person hypermetropic; this form of hypermetropia seldom exceeds  $\frac{1}{4}$ . When a person has both hypermetropia and presbyopia, it is necessary for him to use a stronger pair of glasses for reading, &c., than for ordinary use. If a person for instance, wears a pair of 18 inch convex spectacles to correct a hypermetropia of  $\frac{1}{15}$ , and as age advances his "near" point recedes to 12 inches, even with the addition of his glasses, it will be necessary for him to wear, for reading, a pair of glasses having a focus of about  $10\frac{1}{2}$  inches. Thus  $\frac{1}{8} - \frac{1}{15} = \frac{1}{24} =$  presbyopia, this added to the lens to correct his hypermetropia, ( $\frac{1}{15} + \frac{1}{24} = \frac{1}{10\frac{1}{2}}$  nearly) equals  $10\frac{1}{2}$  nearly.

In the very aged, it is necessary to prescribe glasses, that will enable them to read at 5 or 7 inches from the eye, as their vision is usually somewhat impaired.

The following table constructed by Dr. Kitchener may give a general idea of the glasses required at different periods of life when the presbyopia is unaccompanied by hypermetropia or amblyopia.

At 40 years,—36 inch focus.				At 70 years,—12 inch focus.			
"	45	"	30	"	75	"	10
"	50	"	24	"	80	"	9
"	55	"	20	"	85	"	8
"	58	"	18	"	90	"	7
"	60	"	16	"	100	"	6
"	65	"	14	"		"	

Prof. Donders thinks that when there is no hypermetropia present we should generally advise those glasses to be worn that will enable the person to read distinctly No. 1 (smallest) test type at a distance of 12 inches.

There is an optical defect of the eye that is occasionally met with called astigmatism (from *a* and *στῆγμα*) in which horizontal and vertical lines are not brought to a focus at the same distance behind the crystalline lens. It is relieved by glasses specially ground for each case, these glasses are cylindrical. I have seen but one case of astigmatism.

A very comprehensive article on this subject appears in the Medical Times and Gazette, Nov., 1864, from the pen of J. Zachariah Laurence, M.B., of London.

The paralysis of the accommodation of the eye I have already referred to in a case on page 14.

## SPECIMENS OF JÄGER'S TEST TYPES.

*No. I.—Brilliant, omitted for want of type.*

*No. II.—Pearl.*

A person with normal vision should be able to read No. II at any distance from eight inches to two feet by the eyes.  
 temperance was virtue. They wrought with cheerfulness on days of labour; but observed festivals as intervals of idleness and pleasure. They kept up the Christmas carol, sent true-love knots on Valentine morning, pancakes on Shrove-tide, shewed their wit on the first of April, and religiously cracked nuts on Michaelmas.

*No. III.—Nonpareil.*

Being apprised of our approach, the whole neighbourhood came out to meet their minister dressed in their fine cloths, and preceded by a pipe and tabor; a feast also was provided for our reception, at which we sat cheerfully down; and what the conversation wanted in

*No. VI.—Bourgeois.*

was made up in laughter. Our little habitation was situated at the foot of a sloping hill, sheltered with a beautiful underwood behind, and pratt

*No. VIII.—Small Pica.*

ling river before; on one side a meadow, on the other a green  
 My farm consisted of about twenty acres of excellent land

*No. X.—Pica.*

having given a hundred pounds for my predecessor's  
 good will. Nothing could exceed the neatness of my

*No. XII.—Great Primer.*

little enclosure; the elms and hedge-  
 rows appearing with an inexpressible

*No. XVI.—2-line Great Primer.*

and was covered with

*Cannon. No. XX.—Snellen.*

thatch, which

THE EYE.

BEST TYPES.

ant of type.

nce from eight inches to two feet h

our; but observed festivals as inter  
love knots on Valentine morning,  
usly cracked nuts on Michaelmas

me out to meet their ministr  
; a feast also was provided in  
the conversation wanted in w

was situated at the foot  
wood behind, and prais

n the other a green  
s of excellent land

my predecessor's  
neatness of my

and hedge-  
expressible

r.  
d with

nich

OPTICAL DEFECTS OF THE EYE.

31

4-line Roman. No. XXX.—Snellen.

**gave it an**

8-line Roman. No. XL.—Snellen.

**air of**

ADVERTISEMENTS.

CHARLES POTTER,



OPTICIAN,

20, King Street East, Toronto,

SIGN OF THE SPECTACLES.

ESTABLISHED, 18

Spectacles, Convex and Concave Lenses, Opera Glasses, Microscopes, Telescopes, Steroscopes, &c., always on hand.

Manufacturer of Dr. Rosebrugh's DEMONSTRATING OPHTHALMOSCOPE.

R. A. WOOD & BRO.,

DISPENSING CHEMISTS

dc., dc.,

280, YONGE STREET,

TORONTO.

DR. ROSEBRUGH'S SOLUTION OF ATROPINE, AND CALABAR BEAN PAPER, SUPPLIED TO THE PROFESSION.

W. C. CHEWETT & CO.,

Booksellers, Stationers, Printers,

Lithographers, Bookbinders, &c., &c.

ALL THE STANDARD WORKS AND TEXT BOOKS IN LAW AND MEDICINE, KEPT IN STOCK.

Books imported to Order, at Lowest Rates.

Agents for New Sydenham Society, and Braithwaite's Retrospect.

TTER,



A N,

Toronto,

ESTABLISHED, 186

Opera Glasses, Microscopes,  
always on hand.

PHALMOSCOPE.

BRO.,

EMISTS

REET.

AND CALABAR BEAN  
SESSION.

CO.,

, Printers,

&c., &c.

BOOKS IN LAW AND

Best Rates.

ect.



