

**CIHM  
Microfiche  
Series  
(Monographs)**

**ICMH  
Collection de  
microfiches  
(monographies)**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

**© 1997**

## Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming are checked below.

- Coloured covers / Couverture de couleur
- Covers damaged / Couverture endommagée
- Covers restored and/or laminated / Couverture restaurée et/ou pelliculée
- Cover title missing / Le titre de couverture manque
- Coloured maps / Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black) / Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations / Planches et/ou illustrations en couleur
- Bound with other material / Relié avec d'autres documents
- Only edition available / Seule édition disponible
- Tight binding may cause shadows or distortion along interior margin / La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure.
- Blank leaves added during restorations may appear within the text. Whenever possible, these have been omitted from filming / Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- Additional comments / Commentaires supplémentaires:

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured pages / Pages de couleur
- Pages damaged / Pages endommagées
- Pages restored and/or laminated / Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed / Pages décolorées, tachetées ou piquées
- Pages detached / Pages détachées
- Showthrough / Transparence
- Quality of print varies / Qualité inégale de l'impression
- Includes supplementary material / Comprend du matériel supplémentaire
- Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image / Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.
- Opposing pages with varying colouration or discolourations are filmed twice to ensure the best possible image / Les pages s'opposant ayant des colorations variables ou des décolorations sont filmées deux fois afin d'obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below / Ce document est filmé au taux de réduction indiqué ci-dessous.

10x	14x	18x	22x	26x	30x
12x	16x	20x	24x	✓	28x

**The copy filmed here has been reproduced thanks  
to the generosity of:**


**Stauffer Library  
Queen's University**

**The images appearing here are the best quality  
possible considering the condition and legibility  
of the original copy and in keeping with the  
filming contract specifications.**

**Original copies in printed paper covers are filmed  
beginning with the front cover and ending on  
the last page with a printed or illustrated impres-  
sion, or the back cover when appropriate. All  
other original copies are filmed beginning on the  
first page with a printed or illustrated impres-  
sion, and ending on the last page with a printed  
or illustrated impression.**

**The last recorded frame on each microfiche  
shall contain the symbol → (meaning "CON-  
TINUED"), or the symbol ▽ (meaning "END"),  
whichever applies.**

**Maps, plates, charts, etc., may be filmed at  
different reduction ratios. Those too large to be  
entirely included in one exposure are filmed  
beginning in the upper left hand corner, left to  
right and top to bottom, as many frames as  
required. The following diagrams illustrate the  
method:**



**L'exemplaire filmé fut reproduit grâce à la générosité de:**



**Stauffer Library  
Queen's University**

**Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.**

**Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminent soit par le dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plié, selon le cas. Tous les autres exemplaires origineux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminent par la dernière page qui comporte une telle empreinte.**


**Un des symboles suivants apparaîtra sur le dernière image de chaque microfiche, selon le cas: le symbole → signifie "A SUIVRE", le symbole ▽ signifie "FIN".**

**Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.**



MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)



APPLIED IMAGE Inc

1651 East Main Street  
Rochester, New York 14609 USA  
(716) 482-0300 - Phone  
(716) 288-5989 - Fax

REPORT  
OF THE  
**CANADIAN ARCTIC EXPEDITION**  
**1913-18**

VOLUME VII: CRUSTACEA

PART F: PYCNOGONIDA

By LEON J. COLE

SOUTHERN PARTY 1913-1916

G  
670  
1913  
02t  
v.007  
pt.F  
c.3



OTTAWA  
THOMAS MULVEY  
PREFEDE TO THE KING'S MOST EXCELLENT MAJESTY  
1911

Issued Jan. 1921

# Report of the Canadian Arctic Expedition, 1913-18.

## VOLUME VII: CRUSTACEA

- Part A: DECAPOD CRUSTACEANS. By Miss Mary J. Rathbun. (*Issued August 18, 1919*)
- Part B: SCHIZOPOD CRUSTACEANS. By Waldo L. Schmitt. (*Issued September 22, 1919*)
- Part C: CERAMACEA. By W. T. Calman. (*Issued October 15, 1920*)
- Part D: ISOPODA. By Miss P. L. Boone. (*Issued November 10, 1920*)
- Part E: AMPHIPODA. By Clarence L. Shoemaker. (*Issued September 5, 1920*)
- Part F: PYCNOGONIDA. By Leon J. Cole. (*In press*)
- Part G: EUPHYLLOPODA. By Frits Johnsen. (*In preparation*)
- Part H: CLADOCERA. By Chaney Juday. (*Issued June 23, 1920*)
- Part I: OSTRACODA. By R. W. Sharpe. (*In preparation*)
- Part J: FRESHWATER COPEPODA. By C. Dwight Marsh. (*Issued April 21, 1920*)
- Part K: MARINE COPEPODA. By Arthur Willey. (*Issued June 25, 1920*)
- Part L: PARASITIC COPEPODA. By Charles B. Wilson. (*Issued August 6, 1920*)
- Part M: CIRRIPEDIA. By H. H. Pilsbry. (*In preparation*)

REPORT  
OF THE  
**CANADIAN ARCTIC EXPEDITION**  
**1913-18**

VOLUME VII: CRUSTACEA

PART F: PYCNOGONIDA

By LEON J. COLE

SOUTHERN PARTY—1913-1916



OTTAWA  
THOS. MULVEY  
PRINTER TO THE KING     AT THE EXCELSIOR MAJESTY

# Report on Pycnogonida Collected by the Canadian Arctic Expedition, 1913-1918.

By LEON J. COLE

*University of Wisconsin*

Arctic explorations extending back more than a century have started in what may perhaps be considered a fairly complete knowledge of the pycnogonid fauna in one half of the Arctic ocean, while that of the other half has remained almost wholly unknown. Numerous collections have been reported from the Kara sea (limited by about the 70th meridian) on the east to Baffin bay and Smith sound (to nearly 80° W. long.) on the west. These include collections from the Kara sea, Barents sea, Franz Josef land, Spitzbergen, the north coast of Norway and the Norwegian sea, the north Atlantic, the shores of Greenland, Davis strait, Baffin bay and Smith sound. In contrast to this two or three early records and the reporting of two species of *Nymphon* at point Barrow (Murdoch, 1885)<sup>1</sup> apparently stood for many years as the only records for this group on the arctic coasts of Canada (to the westward of Baffin bay), Alaska and Siberia. The Russian Polar Expedition, 1900-1903 (Junkewitsch, 1907a), added considerably to our knowledge of the Siberian pycnogonid fauna as far east as the New Siberian islands (to longitude 150° E.). The collections of the Canadian Arctic Expedition, while they contain only three species, help to fill in another gap, as they come from Dolphin and Union strait, approximately midway between Baffin bay and point Barrow.

There were transmitted to me through the Smithsonian Institution of Washington, for purposes of identification, two lots of Pycnogonida collected by the Canadian Arctic Expedition. These lots both came from the same station (43a) and comprise five specimens belonging to three different species of the Nymphonidae. The labels give the following data for this station:

"Off Cockburn Point, Dolphin and Union Str., Arctic Can. Sta. 43a. Depth about 100 meters, Gray mud with pebbles, 4 ft. beam-trawl, about 1 hour C.A.E.—F. Johansen."

Following are the species represented:

## 1. *Nymphon sluteri* Hock.

One specimen, adult.

The shape of the eye-tubercle and the shape and relative proportions of the terminal claw to the second tarsal joint are diagnostic.

According to the tabular summary of the distribution of temperate and arctic Pycnogonida given by Norman (1908) this species has previously been reported from the following regions:

- a. British area.
- b. East Arctic—Siberia to East Finnmark.
- c. High Arctic—Spitzbergen, Franz Josef land &c.
- d. West Arctic—Jan Mayen, Iceland, east Greenland.

It has also been reported from west Greenland (Coutts inlet, Rodger, 1893)<sup>2</sup>, while the records of the Russian Polar Expedition from Kara sea, Taimur bay,

<sup>1</sup>Through the kindness of Dr. Paul Bartsch, Curator of Marine Invertebrates in the U.S. National Museum, I have recently had the privilege of examining the point Barrow specimens and have been able to verify Murdoch's identification of them as *Nymphon longitarse* and *Nymphon grossipes*.

<sup>2</sup>Mr. Fritz Johansen has called my attention to the record of two pycnogonids reported in Sutherland's (1852) "Journal of a Voyage in Baffin's Bay and Barrow Straits, in the years 1850-1851" (Vol. II, Appendix, pages cxxvii and cxxviii), which appears to have been overlooked by the authors dealing with this region. The descriptions, by Mr. Adam White, are very inadequate, and the illustrations are little better for specific determination. As Mr. Johansen suggests, his *Nymphon crassipes* is probably a *Chatonympion* (though the hairiness is not mentioned except on the palps, and is not represented in the figure); and might perhaps be *Chatonympion hirtipes*. The other species, which he describes as a species of *Nymphon* similar to the *Pycnogonum grossipes* of Otto Fabricius but smaller and "more slim" would appear to be very close to *Nymphon sluteri* or *Nymphon longitarse*, though the proportionate lengths of the joints of the legs as represented do not agree with the latter species. It would, however, be worse than useless to attempt to assign these forms definitely to known species on the basis of the descriptions furnished. The specimens came from Union bay (about 75° N., 92° W.). At our suggestion, Dr. W. T. Calman has kindly made search for these specimens in the collections of the British Museum, but has been unable to find them.

and north of the New Siberian islands (Schimkewitsch, 1907a) add it to the Siberian arctic area. The present record, from what may be called the American Arctic area<sup>1</sup> completes the circle, making this a truly circumpolar species.

### 2. *Nymphon longitarse* Kröyer.

One specimen, adult.

The known distribution of *N. longitarse* is even more extensive than that of the preceding species. Norman (1908) has collected records from:

- a. British area.
- b. Scandinavian—Norway, South and West.
- c. East Arctic.
- d. High Arctic.
- e. West Greenland<sup>2</sup>.
- f. North-east American to lat. 35° N. (Cape Hatteras).

The American records for this species range from the region of Smith sound (Ortmann, 1901), Baffin bay and Davis Strait (Meinert, 1899; Vanhoffen, 1907), to extreme south Greenland (Stephensen, 1913, 1916), and according to Wilson (1878, 1880) it extends as far south along the east coast of North America as Massachusetts bay. The present record form a link to the westward with that from point Barrow (Murdoch, 1885). Schimkewitsch (1907 b) reported it from the Okhotsk sea,<sup>3</sup> but it does not appear to have been taken by the Russian Polar Expedition, though its otherwise circumpolar distribution would lead to the presumption that it occurs also in the Siberian Arctic.

### 3. *Chaetonymphon hirtipes* (Bell).

One adult male; one adult female; one immature.

The present record extends the range of this well-known species to the westward. Norman (1908) lists it from:

- a. British Area.
- b. Scandinavian.
- c. East Arctic.
- d. High Arctic.
- e. Faroe Channel.
- f. West Arctic.
- g. West Greenland.<sup>2</sup>
- h. North-East American

It has been found widely distributed on the East and West coast of Greenland (see Stephensen, 1913 and 1916, for detailed localities) and according to Wilson (1878, 1880), like *Nymphon longitarse* extends south to Massachusetts bay on the American coast. Carpenter (1898) has said of this form: "*C. hirtipes* is one of the most familiar Arctic pycnogons; and from the numerous localities from which it has been dredged, it would appear to have a complete circumpolar range." It should be noted, however, that in this last respect it does not yet equal either of the foregoing species as it does not appear up to this time to have been found between Dolphin and Union strait and the Kara sea.

The Canadian Arctic Expedition specimens had a number of Foraminifera attached to them, identified by Dr. J. A. Cushman of Boston as *Truncatulina lobatula* (Walker and Jacob). See Report of Canad. Arct. Exped., Vol. IX, Part M, p. 94.

<sup>1</sup>This term is suggested, rather than Canadian Arctic, to include the whole Arctic area adjacent to the North American continent from Baffin bay to Bering strait. Should there be reason for doing so it can be subdivided into the Canadian Arctic and Alaskan Arctic, the division point being practically Mackenzie bay, at the mouth of the Mackenzie river. These two sub-areas differ strikingly in that one is composed of the straits and sounds among the islands of the Arctic Archipelago, while the other is an open coast.

<sup>2</sup>See also footnote 2, p. 3r.

<sup>3</sup>Schimkewitsch (1913) has later described a new variety (*N. longitarse* var. *minus*) from this region, which he says is only about half the size of the European form.

**Other Records for the North American (west of Long. 80°W.) and East Siberian (east of Long. 70° E.) Arctic.**

At least three other species have been reported from the half of the Arctic ocean here considered, making a total of six in all. This is a great contrast to the forty-eight species recorded by Norman (1908) as occurring in the East Arctic, High Arctic, Faroe Channel, West Arctic and West Greenland areas combined. This difference is doubtless due in part to the great discrepancy in the amount of marine exploration of the two regions, but the half of the Arctic above the Atlantic Ocean is more varied and more open to southern influence, and may accordingly have a more varied fauna.

The species referred to are as follows:

**Nymphon grossipes** O. Fabr.

Reported by Murdoch (1885) as "rather plenty but small off Point Franklin [near Pt. Barrow] in 13½ fathoms," and from Norton sound in Bering sea. Aside from these records this species has a wide distribution, corresponding roughly to that of *Chaetonymphon hirtipes*.

**Boreonymphon robustum** (Bell).

Bell (1855) described this species as *Nymphon robustum* from specimens collected by the Belcher Expedition at their winter quarters in Northumberland sound, 76° 52' N., 97° W.

**Colossendeis proboscidea** (Sabine).


This species was described as *Phoxichilus proboscideus* by Sabine (1824) from specimens collected on the Parry Expedition, at North Georgia island (one of the Parry islands, about lat. 75° N., long. 100° W.).

In the narrative of the voyage of the "Vega" (Nordenskiöld, 1881, Vol. 1, p. 349) is a figure of an unnamed specimen of *Colossendeis* which resembles *C. proboscidea* very closely in appearance, particularly in the shape and size of the proboscis. It was taken off the northeast coast of the Taimur peninsula. Lönnberg (1902, p. 359) evidently refers to this record in discussing the distribution of this species, but does not cite his authority for its identification<sup>1</sup>. If the identification is correct it adds another known species to the little-known half of the Arctic previously referred to.

<sup>1</sup> "The original report of the "Vega" expedition has not been available to me, but Dr. Bartsch, who has been so kind as to look it up, states that on page 709 of the "Vega-Expeditiones Vetenskapliga Iakttagelser," volume I (1882) is a picture labelled *Colossendeis gigantea*. This is given as a synonym for *C. proboscidea* by Sars (Pycnogonida of the Norwegian North-Atlantic Expedition, 1891, p. 138). The figure is the same one used in the English translation of the "Vega" narrative.

## REFERENCES.

- Bell, Thomas. 1855. Account of the Crustacea. In The last of the Arctic voyages; being a narrative of the expedition in H.M.S. *Assistance*, under the command of Captain Sir Edward Belcher, C.B., in search of Sir John Franklin, during the years 1852-53-54. Lond., 2 vols. Vol. 2, pp. 400-411, pls. 34, 35.
- Carpenter, G. II. 1898. On Pantopoda collected by Mr. W. S. Bruce in the neighbourhood of Franz-Joseph Land, 1896-97. *Journ. Linn. Soc.-Zool.*, Vol. 24, pp. 626-634, pl. 46.
- Meinert, Fr. 1899. The Danish Ingolf-Expedition, Vol. III, 1, Pyenogonida. Copenhagen.
- Murdoch, John. 1885. Marine invertebrates. (Exclusive of mollusks.) Report of the International Polar Expedition to Point Barrow, Alaska. Washington. Pp. 136-176, pls. 1, 2.
- Lönnberg, Einar. 1902. List of pyenogonids collected by the Swedish zoological expedition to Spitzbergen and East Greenland 1900. *Kongl. Vetensk.-Akad. Förhandl.* 1902, pp. 353-359.
- Nordenskiöld, A. E. 1881. The voyage of the *Vega* around Asia and Europe, with a historical review of previous journeys along the north coast of the world. (Translated by Alexander Leslie.) London: Macmillan and Co., 2 vols.
- Norman, A. M. 1908. The Podosomata (=Pyenogonida) of the Temperate Atlantic and Arctic Oceans. *Journ. Linn. Soc.-Zool.*, Vol. 30, pp. 198-238, pls. 29, 30.
- Ortmann, A. E. 1901. Crustacea and Pyenogonida collected during the Princeton Expedition to North Greenland. *Proc. Acad. Nat. Sci. Phila.* 1901, pp. 144-168.
- Rodger, A. M. 1893-95. Preliminary account of natural history collections made on a voyage to the Gulf of St. Lawrence and Davis Straits. *Proc. Roy. Soc. Edinb.*, Vol. 20, pp. 154-163.
- Sabine, E. 1824. Supplement to the Appendix of Capt. Parry's voyage for the discovery of the North-West Passage in the years 1819-20, containing the zoological and botanical notices. *Marine Invertebrate Animals*, by Captain Edward Sabine. London.
- Schimkewitsch, W. 1907a. Zur Pantopoden-Fauna des sibirischen Eismeeeres. Résultats scientifiques de l'Expédition Polaire Russe en 1900-1903, sous la direction du Baron E. Toll. Section E: Zoologie. Vol. 1, livr. 6., Mém. de l'Acad. Imp. d. Sci. de St.-Pétersb., VIII<sup>e</sup> Scire. Classe Physico-Math., Vol. XVIII, No. 6, pp. 1-10, pl. 1
- 1907 b. Übersicht der von P. Schmidt und W. Broschnikow in den ostasiatischen Ufergewässern gesammelten Pantopoden. *Annuaire du Musée Zoologique de l'Acad. Imp. d. Sci. de St.-Pétersb.*, T. XI, 1906, pp. 246-252, pl. 1
- 1913. Einige neue Pantopoden. *Ibid.*, T. XVIH, 1913, pp. 240-251, pl. 3a.
- Stephensen, K. 1913. Grönlands Krebsdyr og Pyenogonider. (Conspicuum Crustaceorum et Pyenogonidorum Groenlandiae). "Meddelelser om Grönland," XXII. Copenhagen, 479 pp.
- 1916. Zoogeographical investigation of certain fjords in Southern Greenland, with special reference to Crustacea, Pyenogonida and Echinodermata; including a list of Aleyonaria and Pisces. "Meddelelser om Grönland," LIH. Copenhagen, pp. 229-378, 31 figs.
- Sutherland, Peter C. 1852. Journal of a voyage in Baffin's Bay and Barrow Straits, in the years 1850-1851, performed by H.M. Ships "Lady Franklin" and "Sophia," under the command of Mr. William Penny..., London: Longman, Brown, Green and Longmans, 2 vols.
- Vanhöffen, E. 1897. Crustaceen; Drygalski, Grönland-Exped. d. Gesellsch. f. Erdkunde zu Berlin, 1891-93, Vol. 2.
- Wilson, E. B. 1878. A synopsis of the Pyenogonida of New England. *Trans. Conn. Acad. Arts. and Sci.*, Vol. 5, pp. 1-26, pls. 1-7.
- 1880. The Pyenogonida of New England and adjacent waters. *Rept. U.S. Comm. Fish and Fisheries* for 1878, Part VI, pp. 461-506, pls. 1-7.



## Report of the Canadian Arctic Expedition, 1913-1918.

### Volume I: General Introduction, Narrative, Etc.

Part A: Northern Party, 1913-18.

Part B: Southern Party, 1913-16. By Rudolph Martin Anderson. (*In preparation*).

### Volume II: Mammals and Birds.

Part A: Mammals. By Rudolph Martin Anderson. (*In preparation*).

Part B: Birds. By R. M. Anderson and P. A. Taverner. (*In preparation*).

### Volume III: Insects.

Introduction. By C. Gordon Hewitt. (*Issued*).

Part A: Collembola. By Justus W. Folsom. (*Issued*).

Part B: Neuropteroid Insects. By Nathan Banks. (*Issued*).

Part C: Diptera. By Chas. W. Alexander, Harrison G. Dyar, and J. R. Malloch. (*Issued*).

Part D: Mallophaga and Anoplura. By A. W. Baker, G. F. Ferris, and G. H. F. Nuttall. (*Issued*).

Part E: Coleoptera. By J. M. Swaine, H. C. Fall, C. W. Leng, and J. D. Sherman, Jr. (*Issued*).

Part F: Hemiptera. By E. P. Van Duzee. (*Issued*).

Part G: Hymenoptera and Plant Galls. By Alex. D. MacGillivray, Charles T. Brues, L. W. T. Shantz, and E. Porter Felt. (*Issued*).

Part H: Spiders, Mites, and Myriapoda. By J. H. Emerton, Nathan Banks, and Rudolph V. Chamberlin. (*Issued*).

Part I: Lepidoptera. By Arthur Gibson. (*Issued*).

Part J: Orthoptera. By E. M. Walker. (*Issued*).

Part K: General Observations on Insect Life in the Arctic. By Eric Johansen. (*In preparation*).

### Volume IV: Botany.

Part A: Freshwater Algae and Freshwater Diatoms. By Charles W. Faxon. (*In preparation*).

Part B: Marine Algae. By F. S. Collins. (*In preparation*).

Part C: Fungi. By John Durness. (*In preparation*).

Part D: Lichens. By G. K. Merrill. (*In preparation*).

Part E: Mosses. By R. S. Williams. (*In press*).

### Volume V: Botany.

Part A: Vascular Plants. By James M. Macoun and Theo. Holm. (*In preparation*).

Part B: Contributions to Morphology, Synonymy and General Distribution of Arctic Plants. By Theo. Holm. (*In press*).

Part C: General Notes on Arctic Vegetation. By Eric Johansen. (*In preparation*).

### Volume VI: Fishes, Trilobites, Etc.

Part A: Fishes. By E. Johansen. (*In preparation*).

Part B: Ascidians, etc. By A. G. Huntsman. (*In preparation*).

### Volume VII: Crustacea.

Part A: Decapod Crustacea. By Mary T. Rathbun. (*Issued*).

Part B: Schizopod Crustaceans. By Waldo L. Schmitt. (*Issued*).

Part C: Ciliaceæ. By W. T. Calman. (*Issued*).

Part D: Isopoda. By Miss P. L. Hoops. (*Issued*).

Part E: Amphipoda. By Oskar R. Showmaker. (*In preparation*).

Part F: Pyemozonida. By Leon J. Cole. (*In preparation*).

Part G: Euphylllopoda. By E. Johansen. (*In preparation*).

Part H: Cladocera. By Chauncey Juday. (*Issued*).

Part I: Ostracoda. By R. W. Shantz. (*In preparation*).

Part J: Freshwater Copepoda. By C. Dwight Marsh. (*Issued*).

Part K: Marine Copepoda. By A. Willey. (*Issued*).

Part L: Parasitic Copepoda. By Claus. B. Wilson. (*Issued*).

Part M: Cirripedia. By H. A. Pilsbry. (*In preparation*).

### Volume VIII: Mollusks, Echinoderms, Coelenterates, Etc.

Part A: Mollusks, Recent and Pleistocene. By Wm. H. Dall. (*Issued*).

Part B: Cephalopoda and Pteropoda. By S. S. Berry and W. F. Clapp. (*In preparation*).

Part C: Echinoderms. By Austin H. Clark. (*Issued*).

Part D: Bryozoa. By R. C. Ostara. (*In preparation*).

Part E: Rotatoria. By H. K. Harring. (*In preparation*).

Part F: Chaetognatha. By A. G. Huntsman. (*In preparation*).

Part G: Actinozoa and Aleyrodomilia. By A. E. Verrell. (*In preparation*).

Part H: Medusae and Ctenophora. By H. B. Bigelow. (*Issued*).

Part I: Hydrozoa. By McLean Fraser. (*In preparation*).

Part J: Porifera. (*In preparation*).

### Volume IX: Annelids, Parasitic Worms, Protozoans, Etc.

Part A: Oligochaeta. By Frank Smith and Paul S. Welch. (*Issued*).

Part B: Polychaeta. By Ralph V. Chamberlin. (*Issued*).

Part C: Hirudinea. By J. P. Morris. (*In press*).

Part D: Gephyrea. By Ralph V. Chamberlin. (*Issued*).

Part E: Acanthocephala. By H. J. Van Cleave. (*Issued*).

Part F: Nemertea. By N. A. Cobb. (*In preparation*).

Part G: H. Trematoda and Cestoda. By A. R. Cooper. (*In preparation*).

Part H: Turbellaria. By A. Hassell. (*In preparation*).

Part I: Gordiacea.

Part K: Nematini.

Part L: Sporozoa. By J. W. Mayor. (*In preparation*).

Part M: Foraminifera. By J. A. Cushman. (*Issued*).

### Volume X: Plankton, Hydrography, Tides, etc.

Part A: Plankton. By Albert Mann. (*In preparation*).

Part B: Marine Diatoms. By L. W. Bailey. (*In preparation*).

Part C: Tidal Observations and Results. By W. Bell Davis. (*In preparation*).

Part D: Hydrography. (*In preparation*).



