




**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**CIHM/ICMH  
Microfiche  
Series.**

**CIHM/ICMH  
Collection de  
microfiches.**



Technical Notes / Notes techniques

The Institute has attempted to obtain the best original copy available for filming. Physical features of this copy which may alter any of the images in the reproduction are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Certains défauts susceptibles de nuire à la qualité de la reproduction sont notés ci-dessous.

- |                                     |   |                          |   |
|-------------------------------------|---|--------------------------|---|
| <input type="checkbox"/>            | Coloured covers/<br>Couvertures de couleur  | <input type="checkbox"/> | Coloured pages/<br>Pages de couleur     |
| <input type="checkbox"/>            | Coloured maps/<br>Cartes géographiques en couleur   | <input type="checkbox"/> | Coloured plates/<br>Planches en couleur |
| <input checked="" type="checkbox"/> | Pages discoloured, stained or foxed/<br>Pages décolorées, tachetées ou piquées  | <input type="checkbox"/> | Show through/<br>Transparence           |
| <input type="checkbox"/>            | Tight binding (may cause shadows or distortion along interior margin)/<br>Reliura serré (peut causer de l'ombre ou de la distortion le long de la marge intérieure) | <input type="checkbox"/> | Pages damaged/<br>Pages endommagées     |
| <input type="checkbox"/>            | Additional comments/<br>Commentaires supplémentaires  |                          |   |
- 

Bibliographic Notes / Notes bibliographiques

- |                          |   |                          |  |
|--------------------------|---|--------------------------|--|
| <input type="checkbox"/> | Only edition available/<br>Seule édition disponible         | <input type="checkbox"/> | Pagination incorrect/<br>Erreurs de pagination     |
| <input type="checkbox"/> | Bound with other material/<br>Relié avec d'autres documents | <input type="checkbox"/> | Pages missing/<br>Des pages manquent               |
| <input type="checkbox"/> | Cover title missing/<br>Le titre de couverture manque       | <input type="checkbox"/> | Maps missing/<br>Des cartes géographiques manquent |
| <input type="checkbox"/> | Plates missing/<br>Des planches manquent                    |                          |  |
| <input type="checkbox"/> | Additional comments/<br>Commentaires supplémentaires        |                          |  |



The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

The last recorded frame on each microfiche shall contain the symbol → (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

The original copy was borrowed from, and filmed with, the kind consent of the following institution:

National Library of Canada

Maps or plates too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole → signifie "A SUIVRE", le symbole ∇ signifie "FIN".

L'exemplaire filmé fut reproduit grâce à la générosité de l'établissement prêteur suivant :

Bibliothèque nationale du Canada

Les cartes ou les planches trop grandes pour être reproduites en un seul cliché sont filmées à partir de l'angle supérieure gauche, de gauche à droite et de haut en bas, en prenant le nombre d'images nécessaire. Le diagramme suivant illustre la méthode :



NOTES on the OCCURRENCE of MAMMOTH-REMAINS in the YUKON  
DISTRICT of CANADA and in ALASKA. By GEORGE M. DAWSON,  
C.M.G., LL.D., F.R.S., F.G.S., Assistant Director of the Geolo-  
gical Survey of Canada.

THESE notes, relating primarily to the occurrence of remains of the Mammoth in the geographical valley of the Yukon River, are the result of a correspondence between Mr. H. Moody of the Canadian Pacific Railway Co., the Assistant Secretary of the Geological Society, and the writer, respecting statements which had reached Mr. Moody from a friend resident in the extreme north-western part of the Dominion of Canada. It has been suggested that a brief notice of the facts in this connexion, so far as these are known, may be of some interest to the Geological Society.

The original discovery of bones of the Mammoth in the Yukon region is due to Mr. Robert Campbell, an officer in the service of the Hudson's Bay Company, who between 1840 and 1852 travelled through and established trading-posts in the upper valley of the Yukon, and was the first white man to penetrate this remote part of North America.

In a brief account of his explorations, printed at Winnipeg in 1885, Campbell writes:—"I saw the bones, heads, and horns of Buffaloes [Musk-Oxen?]; but this animal had become extinct before our visit, as had also some species of Elephant, whose remains were found in various swamps. I forwarded an Elephant's thigh-bone to the British Museum, where it may still be seen".<sup>1</sup>

<sup>1</sup> 'The Discovery and Exploration of the Yukon (Pelly) River,' Winnipeg, 1885.

As Campbell's posts on the Upper Yukon were finally abandoned in 1852, the bone thus referred to by him must have been sent out before this date. It was a tibia, not a thigh-bone, and was described by Sir John Richardson in 1855 as referable to *Elephas primigenius*. Richardson states that it was identical in form with, though larger than, a corresponding bone of the same animal brought back by Capt. Beechey from Eschscholtz Bay. The skeleton of which it formed part was said to be complete when found; but most of the bones were lost by the Indians who extracted them for Campbell. According to a statement subsequently obtained from Campbell, these bones were found at some place not far from the former site of Fort Selkirk, at the confluence of the Lewes and Pelly Rivers.<sup>1</sup>

Dr. W. H. Dall in 1866-67, during his connexion with the Western Union Telegraph Expedition (abandoned on the completion of the Atlantic Cable), visited a number of places in the lower valley of the Yukon, within what is now the Territory of Alaska. In the volume which resulted from his explorations, and in other publications, he frequently mentions the occurrence of Mammoth-remains in this region, writing in one place as follows:—

“Wild and exaggerated stories have found a place, even in official documents, in regard to fossil ivory. This is not uncommon in many parts of the valley of the Yukon and Kuskoquim. It is usually found on the surface, not buried as in Siberia, and all that I have seen has been so much injured by the weather that it was of little commercial value. It is usually blackened, split, and so fragile as to break readily to pieces. A lake near Nushergak, the Inglutalik River, and the Kóto River are noted localities for this ivory.”<sup>2</sup>

In 1886 the Geological Survey of Canada acquired from Mr. F. Mercier, who had spent many years as a trader in the Yukon region, a number of bones, tusks, and teeth of the Mammoth. These were chiefly obtained by Mr. Mercier near the mouth of the Tananá River, one of the main feeders of the Yukon on the south side. Mr. J. F. Whiteaves, F.G.S., Paleontologist to the Geological Survey of Canada, has kindly furnished the subjoined note on these remains:—

“In my judgment all the Elephantine remains collected by Mr. Mercier in the Yukon region, and now in our Museum, are clearly *Elephas* (sub-genus *Euclephas*) and not *Mastodon*.

“Four of the specimens collected by Mr. Mercier are perfect molars, essentially similar to those from Burlington Heights, near Hamilton, Ontario, which E. Billings referred to *Elephas Jacksoni* of Briggs and Foster, but which Dr. Falconer subsequently identified with *E. primigenius*, Blumenbach.

“The specific relations of the North American fossil Elephants

<sup>1</sup> ‘Zoology of the Voyage of H.M.S. ‘Herald,’ (1854) p. 142; Am. Journ. Sci. ser. 2, vol. xix. (1855) p. 132; Annual Report, Geol. Surv. Canada, 1887, p. 41 B.

<sup>2</sup> ‘Alaska and its Resources,’ 1870, pp. 238, 460, 479; Am. Journ. Sci. ser. 2, vol. xlv. (1868) p. 99.

(as distinguished from Mastodons) are treated of at considerable length in vol. ii. pp. 234-238 of the 'Palaontological Memoirs and Notes of the late Dr. Hugh Falconer,' under the heading 'Synonymy of American Fossil Elephants.'

"It is there stated that there are but two species of fossil Elephant in North America. The first of these is the *Elephas (Euclaphus) primigenius*, Blumenbach, of which *E. Jacksoni*, of Briggs and Foster, and *E. americanus*, Leidy, are synonyms. According to Dr. Falconer, all the specimens from the Yukon, Alaska, and Burlington Heights are *E. primigenius*. The second species is *E. Columbi*, Falconer, of the southern part of the United States and Mexico."

The writer, in 1887, carried out an extended reconnaissance-survey in the Yukon District, in the valleys of the Pelly and Lewes branches of the main stream, but not going below the confluence of these two rivers.<sup>1</sup> In the whole region thus traversed no Mammoth-remains were met with, nor was their presence reported by such of the gold-miners as had worked in parts of these valleys; though some of the same men had frequently noted Mammoth-bones farther down the Yukon valley, particularly in the vicinity of Forty-Mile Creek, where rather important placer-mining has been carried on.

The above notes refer particularly to the occurrence of Mammoth-remains in the inland region of Alaska, and in parts of the adjacent Yukon District of the North-west Territory of Canada—the International boundary following the 141st meridian. The existence of similar remains, as well as those of other animals not now inhabiting the region, has long been known at various places on the coast, both to the south and north of Bering Straits. The most notable and the first discovered of these localities is Kotzebue Sound, where bones were collected by Kotzebue in 1816, Capt. Beechey, of H.M.S. 'Blossom,' in 1826, Capt. Kellett, of H.M.S. 'Herald,' in 1848, Dr. W. H. Dall in 1880, and Mr. Nelson in 1881. The specimens brought back by the three first-named expeditions were described by Eschscholtz, Buckland, Forbes, and Richardson in appendices or auxiliary works to the narratives of the several voyages.

Dall has recently given a summary of what is known respecting these localities, with full references to the published accounts of them.<sup>2</sup> The bones found at Kotzebue Sound and at other places on the coast are associated with what he calls the 'ground-ice formation.' The localities are indicated in a general manner on the map accompanying Dall's work; but, so far as these are described or the writer is aware, no information exists to show that such bones are associated with 'ground-ice' anywhere south of Kotzebue Sound.

The following list of species obtained in Kotzebue Sound is given

<sup>1</sup> Annual Report, Geol. Surv. Canada, 1887-88, Part B.

<sup>2</sup> Bull. U.S. Geol. Survey, no. 84, 1892, pp. 260-267.



by Dall, chiefly from Richardson's report, but with revised nomenclature<sup>1</sup> :—

*Elephas primigenius*, Blumenbach

*Elephas Columbi*, Falconer [?].<sup>2</sup>

*Equus major*, De Kay.

*Alces americanus*, Jardine = *Muchlis*, Ogilby.

*Rangifer Caribou*, Baird.

*Ovibos moschatus*, Flainville.

*Ovibos maximus*, Richardson = *O. cavifrons*, Leidy.

*Bison crassicornis*, Richardson = *B. antiquus*, Leidy.

No *Mastodon*-bones appear to have been found in any portion of the extreme north-west of North America.

Of particular interest in connexion with the general question of the distribution of Mammoth-remains in the Alaskan region is the occurrence of such remains (a tooth) on St. George Island of the Pribilof group, in Bering Sea, and on Unalaska Island of the Aleutian Chain.<sup>3</sup> Mr. J. Stanley-Brown further notes the discovery of a Mammoth-tusk on St. Paul Island of the Pribilof group, but appears at the same time to throw doubt on the means by which these remains reached the Pribilof Islands, writing—"As there is not a foot of earth upon either island, save that which has resulted from the decomposition of the native rock and the decay of vegetation, the value of such testimony is questionable."<sup>4</sup>

The precise intention of the cautionary remark just quoted is not clear to the writer. The finding of the bones upon St. George and St. Paul Islands does not appear to be doubtful. Both islands were uninhabited previous to their discovery by the Russians; they show neither traces of glacial action nor erratics; and in what way the Mammoth can be supposed to have reached these islands, except by means of a former connexion with the mainland, it is difficult to understand. We have, moreover, the Mammoth-bones already mentioned on Unalaska Island, vouchered for by Dr. Stein, and a like explanation must be found for all these cases. This does not appear to be difficult, for the whole eastern part of Bering Sea is rather notably shallow, nearly everywhere less than 50 fathoms in depth. An elevation of the land by about 300 feet would thus suffice to unite the islands mentioned, with a number of others, to the American Continent, and it appears scarcely to admit of doubt that it was across such a practicable plain that the Mammoth found its way to these places.

The most important observation to be based on the foregoing notes is that the remains of the Mammoth, with those of other associated animals, are, in the north-western part of the North American Continent, abundant in, if not strictly confined to the

<sup>1</sup> *Op. cit.* p. 264.

<sup>2</sup> I have ventured to place a mark of interrogation against this species, for Falconer gives its range as being from Mexico to Georgia and perhaps farther south. See 'Paleontological Memoirs and Notes,' vol. ii. pp. 230-231. See also Howorth, 'The Mammoth and the Flood,' pp. 274-276.

<sup>3</sup> Bull. U.S. Geol. Survey, no. 84, p. 266.

<sup>4</sup> Bull. Geol. Soc. Am. vol. iii. (1892) p. 499.

limits of, a great unglaciated area there existing. With the exception of the southern mountainous sea-margin of Alaska, and doubtless also that of certain local inland ranges, this unglaciated area may be described as comprising nearly the whole of Alaska, together with a considerable portion of the adjacent Yukon District of Canada.

As the result of his explorations in this part of the continent, the writer has been able to determine the fact that during the glacial period the Rocky Mountain or Cordilleran region, from about the 48th to the 63rd degree of latitude North, was at one time buried beneath a great confluent ice-mass some 1200 miles in greatest length in a north-west by south-east bearing, with an average width of about 400 miles.<sup>1</sup>

This Greenland-like ice-cap was distinct from the still greater Laurentide Glacier of Eastern North America, and, because of the trend of the mountain-ranges which it covered, it moved principally in two directions—south-eastward and north-westward. The south-easterly motion of one part of this ice-mass the writer had demonstrated in 1877,<sup>2</sup> but it was not till 1887, and then as a result of the Yukon expedition, that he was enabled to ascertain the north-westerly movement of its northern part, and to show that there was a definite limit to its extent in both directions. Being thus clearly distinct from any extension of polar ice, as well as from the great Laurentide ice-mass, it became appropriate to designate it as the Cordilleran Glacier.<sup>3</sup> Further evidence respecting the northern limit of glaciation in this region has since been obtained by Mr. R. G. McConnell, of the Canadian Geological Survey (1888), Mr. I. C. Russell, of the U. S. Geological Survey (1889), and Mr. C. W. Hayes, of the same Survey (1891).<sup>4</sup> The area covered by, and the directions of movement of, the Cordilleran ice-mass have been approximately mapped in one of the papers above referred to,<sup>5</sup> and the later observations of the above-named gentlemen have not in any material degree changed the indications there given.

Within the area which was covered by the great Cordilleran Glacier, remains of the Mammoth are either entirely wanting or are very scarce. The reported finding of a tooth on the southern part of Vancouver Island, and that of a portion of a large bone (which, though not determinable, may have belonged to such an animal) in gravels worked for gold on Cherry Creek,<sup>6</sup> are the only possible exceptions known to the writer, and the deposits from which the last-mentioned bone was obtained may be of pre-Glacial age.

<sup>1</sup> 'On the later Physiographical Geology of the Rocky Mountain Region in Canada,' *Trans. Royal Soc. Canada*, vol. viii. (1890) sect. iv. p. 27.

<sup>2</sup> Report of Progress, *Geol. Surv. Canada*, 1877-78, pp. 136 B, 151 B; *Quart. Journ. Geol. Soc.* vol. xxxiv. (1878) p. 119, vol. xxxvii. (1881) p. 283.

<sup>3</sup> 'American Geologist,' vol. vi. (1890) p. 162.

<sup>4</sup> Annual Report, *Geol. Surv. Canada*, 1888-89, p. 28 D; *Bull. Geol. Soc. Am.* vol. i. (1890) p. 144; *National Geogr. Mag.*, Washington, vol. iv. p. 157.

<sup>5</sup> *Trans. Royal Soc. Canada*, *op. cit.* pl. ii. map no. 4.

<sup>6</sup> Okanagan District, British Columbia.

The likeness of the non-glaciated north-western portion of North America, with its abundant Mammoth-remains, to the similarly characterized northern part of Asia has already been recognized. The purport of the foregoing remark is to indicate the existence of a south-eastern boundary to the Mammoth-inhabited portion of Alaska and the Yukon District; nor can it be reasonably doubted that the North American and Asiatic land was continuous at the time of the existence of the Mammoth, or for some portion of that time; for an elevation of the land sufficient to enable the Mammoth to reach the islands in Bering Sea, already referred to, would result in the obliteration of Bering Straits.

Many conjectures have been advanced as to the mode of occurrence and origination of the 'ground-ice formation,' in association with which the bones of the Mammoth and other animals are found along the northern coasts of Alaska. Dall summarizes these in his work previously cited,<sup>1</sup> and it may now be confidently assumed that the descriptions of Kotzebue and his party, of Capt. Kellett and others on the 'Herald,' of Dall and Lieut. Cantwell,<sup>2</sup> correctly indicate the facts of the case. The clearest descriptions of the phenomena are those of Seemann and Dall.<sup>3</sup> From these it appears that the lower parts of cliffs which have some extent on Kotzebue Sound are composed of solid ice, somewhat discoloured and impure, and showing indications of stratification. Above this ice rests a layer of clay, in which the bones occur, and capping the whole is a peaty layer supporting the vegetation of the region. It is further apparent that this or a very similar formation occurs at a number of points along the northern coast of Alaska, but nothing has been adduced to show that it is absolutely continuous over any great area;—there is, in fact, some reason to believe that it is confined to limited tracts, even in the vicinity of Kotzebue Sound.<sup>4</sup>

In the present connexion, the 'ground-ice formation' is of interest only in so far as its existence and the mode of its origination may throw light on the date and method of entombment of the Mammoth-remains associated with it. With respect to the origin of the deposits, the writer ventures to offer the following suggestions.

The country in which the 'ground-ice formation' occurs is low in its relief, and the formation occupies its lower tracts. The ice itself must undoubtedly have been produced upon a land-surface, and since the time of its production this surface can never have been covered by the sea; for this would inevitably have reduced the frozen condition of the overlying clays, and have resulted in the destruction of the icy sub-stratum as well.

With an elevation of the land by an amount of 300 feet or more (such as appears to be required by the Mammoth-remains on islands already mentioned) the warmer waters connecting with the Pacific

<sup>1</sup> Bull. U.S. Geol. Survey, No. 84, pp. 260-264.

<sup>2</sup> 'American Geologist,' vol. vi. (1890) p. 51.

<sup>3</sup> 'Voyage of H.M.S. 'Herald,'' vol. ii. pp. 33 *et seqq.*; Bull. U.S. Geol. Surv. No. 84, pp. 261 *et seqq.*

<sup>4</sup> 'American Geologist,' vol. vi. (1890) p. 52.

would be confined to the deeper western portion of what is now Bering Sea, forming there a limited gulf, without outlet to the north, from which the region where the 'ground-ice formation' is now found would be so far removed as to greatly reduce its mean annual temperature. Snow falling upon this nearly level, northern land, and only in part removed during the summer, would naturally tend to accumulate in *nevé*-like masses in the valleys and lower tracts, and the underlying layers of such accumulations would pass into the condition of ice, though without the necessary slope or head to produce moving glaciers. The evidence does not seem to imply that the Mammoth resorted to this extreme northern portion of the region during the actual time of ice-accumulation, but this animal may be supposed to have passed between Asia and America along the southern parts of the wide land-bridge then existing.

At a later date, when the land became depressed to about its present level, Bering Sea extended itself far to the eastward, and Bering Straits were opened. The perennial accumulation of snow upon the lowlands ceased, and in the southern parts of Alaska such masses as had been formed may have been entirely removed. Farther to the north and at a greater distance from the Pacific waters, while the total precipitation would probably be increased, a greater proportion would fall as rain, and floods resulting from this and the melting of snow on the higher tracts would be frequent. Thus it may be supposed that deposits of clay and soil from adjacent highlands and from the overflow of rivers covered large parts of the remaining ice of the lowlands, and that wherever so covered it has since remained; the winter temperature being still sufficiently low to ensure the persistence of a layer of frozen soil between the surface annually thawed and the subjacent ice. Over the new land thus formed the Mammoth and associated animals appear to have roamed and fed, and wherever local areas of decay of the ice may have arisen, bottomless bogs and sink-holes must have been produced which served as veritable traps.

It will be observed that this hypothesis requires a rather abrupt passage from the conditions under which the ice accumulated to those in which, before it had time to disappear, it began to be covered up by soil, but the change may nevertheless have extended over a considerable number of years. The association of the Mammoth with an animal so essentially Arctic as the Musk-Ox requires—as has frequently been pointed out—the admission that the Mammoth was capable of living in a rigorous climate, though it may be that the southern limit of the migration-range of one animal merely overlapped the northern limit of the migration-range of the other. The occurrence of the Moose (*Alces americanus*) implies the existence at that time of woodland, or at least of well-grown thickets.

In the Cordilleran region generally, the Pliocene and Glacial periods were characterized by several important changes in elevation and depression of land;<sup>1</sup> but it is unsafe to assume that these

<sup>1</sup> Trans. Royal Soc. Canada, vol. viii. (1890) sect. iv. p. 54; Bull. U.S. Geol. Survey, no. 84, p. 278.

changes equally affected the northern region here particularly treated of; for it is not only very distant from the localities which have so far been specially studied, but the physical features of the Cordilleran belt become diffuse and ill-marked to the north, and such mountain-ridges as remain assume new trends. It may, however, be taken for granted that this region shared to some extent in these great movements of elevation and depression, and as the very existence of the 'ground-ice' shows that the area where it is found has not since the date of its formation been materially lower than at present, it may reasonably be argued that it dates from a period approaching the conclusion of the series of changes in level, or subsequent to the last well-marked epoch of depression of the land.

Thus, without entering into any details respecting the sequence of these great earth-movements in the Cordilleran region of British Columbia,<sup>1</sup> it may be stated as probable that the uprising of the land which led to the accumulation of the 'ground-ice' was coincident with the second (and latest) epoch of maximum glaciation, which was followed by an important subsidence in British Columbia.

#### DISCUSSION.

The PRESIDENT said that many interesting points had been brought forward by the Author. The differentiation of the glaciated from the unglaciated area, and the clear recognition of a north-western as well as a south-eastern boundary to the Cordilleran ice-mass, struck him as being of great importance.

Sir HENRY HOWORTH remarked upon the long and careful survey of N.W. America which has been made by the Author, and upon the value of the conclusions to which he has come: firstly, in regard to the absence of ancient glaciation in Alaska and its borders; secondly, in regard to the existence of a great glacier in the Cordilleras, whose products are quite independent of and have nothing to do with the Laurentian drift; and thirdly, in regard to the distribution of the Mammoth. It was a new fact to him, and one of great importance, that Mammoth-remains had occurred in Unalashka and the Pribilof Islands in Bering Sea, proving that in the Mammoth age there was a land-bridge here, as many inquirers had argued. It would be very interesting to have the western frontier defined, where the Mammoth-remains cease to be found. It would also be very interesting to know how far south on the west of the Cordilleras the true Mammoth, as distinguished from *Elephas Columbi*, has occurred.

Regarding one conclusion of Dr. Dawson's, the speaker could not agree with his friend, namely, about the age of the strata of ice sometimes found under the Mammoth-beds in Alaska as they have been found in Siberia. The speaker was of opinion that this ice has accumulated since the beds were laid down, and was not there when the Mammoth roamed about in the forests where he and his com-

<sup>1</sup> For a discussion of which see Trans. Royal Soc. Canada, vol. viii. (1890) sect. iv. pp. 40-55.

panions lived. Humus and soil cannot accumulate upon ice except as a moraine, and there are no traces of moraines or of great surface-glaciation in Alaska and Siberia. Nor could either the flora or fauna of the Mammoth age have survived conditions consistent with the accumulation of these beds of ice almost immediately below the surface, or consistent with their presence there. The speaker considered that these beds are due to the filtration of water in the summer down to the point where there is a stratum of frozen soil, through which it cannot pass and where it consequently accumulates, freezes, raises the ground, and in the next season grows by the same process until a thick bed of ice has been formed. The evidence goes to show that the present is the coldest period known in recent geological times in Siberia and Alaska, and that the period of the Mammoth and its companions was followed and not preceded by an Arctic climate where its remains occur.

Dr. HENRY WOODWARD mentioned that in 1850 Capt. Kellett and Lieut. Wood brought remains of Musk-Ox and Mammoth to the British Museum from Kotzebue Sound, Alaska; and in 1873 the Rev. R. McDonald (one of the Hudson's Bay Company's Chaplains) from Fort McPherson, Mackenzie River, Arctic America, gave to the National Collection, from the Poreupine River, remains of Mammoth, Musk-Ox, *Bison priscus*, and Horse. The *Mastodon* has lately been found in Kent County, Ontario, Canada. These instances prove the former abundance of the land Mammalia in high latitudes in North America. The most interesting point in Dr. Dawson's paper is the mention by him of the remains of Mammoth on the Aleutian Islands, proving that this was the old high road for this and other mammals from Asia into North America in Pleistocene times.

Prof. HULL observed that, with reference to the requirements of the large animals referred to in Dr. Dawson's interesting paper, he had seen it stated that one had been discovered in N.W. America nearly entire, and in its stomach were about seven bushels of vegetable matter. However that might be, it seemed clear that the climate of the circumpolar regions had undergone a great change since the Mammoth had become extinct; in consequence of which the vegetation had materially fallen off. He also desired to call attention to the clear evidence which the Author's paper afforded of the former wider extension of land in the Arctic regions during the Mammoth period.